

IDENTIFYING PHASE TRANSITION THRESHOLDS OF PERMUTED LINEAR REGRESSION VIA MESSAGE PASSING

Anonymous authors

Paper under double-blind review

ABSTRACT

This paper considers the permuted linear regression, i.e., $\mathbf{Y} = \mathbf{\Pi}^\dagger \mathbf{X} \mathbf{B}^\dagger + \mathbf{W}$, where $\mathbf{Y} \in \mathbb{R}^{n \times m}$, $\mathbf{\Pi}^\dagger \in \mathbb{R}^{n \times n}$, $\mathbf{X} \in \mathbb{R}^{n \times p}$, $\mathbf{B}^\dagger \in \mathbb{R}^{p \times m}$, and $\mathbf{W} \in \mathbb{R}^{n \times m}$ represent the observations, missing (or incomplete) information about ordering, sensing matrix, signal of interests, and additive sensing noise, respectively. As is shown in the previous work, there exists phase transition phenomena in terms of the *signal-to-noise ratio* (snr), number of permuted rows, etc. While all existing works only concern the convergence rates without specifying the associate constants in front of them, we give a precise identification of the phase transition thresholds via the message passing algorithm. Depending on whether the signal \mathbf{B}^\dagger is known or not, we separately identify the corresponding critical points around the phase transition regimes. Moreover, we provide numerical experiments and show the empirical phase transition points are well aligned with theoretical predictions.

1 INTRODUCTION

This paper considers the permuted linear regression

$$\mathbf{Y} = \mathbf{\Pi}^\dagger \mathbf{X} \mathbf{B}^\dagger + \sigma \mathbf{W},$$

where $\mathbf{Y} \in \mathbb{R}^{m \times n}$ denotes the sensing result, $\mathbf{\Pi}^\dagger \in \mathbb{R}^{n \times n}$ represents the permutation matrix, $\mathbf{X} \in \mathbb{R}^{n \times p}$ is the sensing matrix, \mathbf{B}^\dagger is the signal of interests, \mathbf{W} denotes the additive noise, and σ is the noise variance. The research on this problem dates back at least to 1970s under the name ‘broken sample problem’ (Goel, 1975; Bai & Hsing, 2005; DeGroot et al., 1971; DeGroot & Goel, 1976; 1980). In recent years, we have witnessed a revival of this problem due to its board spectrum of applications in privacy protection, data integration, etc (Pananjady et al., 2018; Unnikrishnan et al., 2015; Slawski et al., 2020; Slawski & Ben-David, 2019; Pananjady et al., 2017; Zhang et al., 2022; Zhang & Li, 2020).

Associated with this problem comes a phase transition phenomenon: the error rate for the permutation recovery suddenly drops to zero once some parameters reach certain thresholds. Despite previous work such as Slawski et al. (2020); Slawski & Ben-David (2019); Pananjady et al. (2017); Zhang et al. (2022); Zhang & Li (2020) can all explain this phenomenon, the precise positions of the phase transition thresholds are never studied but rather their statistical order. In this work, we would like to leverage *message passing* (MP) algorithm to identify their precise location. As a byproduct, we also come up with an algorithm to partially recover the permutation matrix.

Related work. The line of research starts with the literature in permuted linear regression. Among all the works mentioned above, the most related works include Slawski et al. (2020); Slawski & Ben-David (2019); Pananjady et al. (2017); Zhang et al. (2022); Zhang & Li (2020), in which almost the same settings as ours are used. Pananjady et al. (2018); Slawski & Ben-David (2019) consider the single observation model ($m = 1$) and proved the snr for the correct permutation recovery will be $\mathbb{O}_p(n^c)$, where $c > 0$ is some positive constant. Later, Slawski et al. (2020); Zhang et al. (2022); Zhang & Li (2020) investigate the multiple observations model ($m > 1$) and suggest the snr requirement can be significantly decreased, to put it more specifically, from $\mathbb{O}_p(n^c)$ to $\mathbb{O}_p(n^{c/m})$. In particular, Zhang & Li (2020) develop an estimator which we will analyze as part of our contributions. Although they obtained the correct convergence rate to restore the correspondence,

which are minimax-optimal in certain regimes, their results fail to specify the leading coefficients, or equivalently, the precise location of the phase transition threshold. Moreover, their analysis does not consider the intertwined influence among the parameters n, p, m , etc. One example would be the impact of the p/n ratio on the maximum allowed number of permuted rows, which has not been studied before this work.

Another line of research comes from the field of statistical physics, which begins with Mézard & Parisi (1986; 1985). Using the replica method, they study the *linear assignment problem* (LAP), i.e., $\min_{\Pi} \sum_{i,j} \Pi_{ij} E_{ij}$ where Π denotes a permutation matrix and E_{ij} is i.i.d random variable uniformly distributed within the regime $[0, 1]$. Martin et al. (2005) then generalize the LAP to multi-index matching and presented a investigation based on MP algorithm. And Caracciolo et al. (2017); Malatesta et al. (2019) extend the distribution of E_{ij} to a broader class. However, all the above works exhibit no phase transition. In Chertkov et al. (2010), this method is extended to the particle tracking problem, where a phase transition phenomenon is first observed. Later, Semerjian et al. (2020) modify it to fit the graph matching problem, which paves way for our work in studying the permuted linear regression.

Our technical **contributions** are summarized as follows

- We propose the first framework that can identify the precise location of phase transition thresholds associated with permuted linear regression. In the oracle case where \mathbf{B}^{\natural} is known, our scheme is able to determine the phase transition snr. In the non-oracle case where \mathbf{B}^{\natural} is not given, our scheme can further predict the maximum allowed permuted rows and uncover its dependence on the ratio n/p .
- We generalize the full permutation estimator and first obtain a partial permutation estimator. Consider the example where the correspondence for a single index is desired. By removing all function nodes except that corresponds to that index, we exploit the MP algorithm and design an algorithm that converge in one step. Moreover, we show its performance almost match the estimator for the full permutation recovery.

In addition, we would like to briefly mention the technical challenges. Compared with the previous works (Mezard & Montanari, 2009; Talagrand, 2010; Linusson & Wästlund, 2004; Mézard & Parisi, 1987; 1986; Parisi & Ratiéville, 2002; Semerjian et al., 2020), where the edge weights are relatively simple, our edge weights usually involve high-order interactions across Gaussian random variables and are densely correlated. To tackle this issue, our proposed approximation method to compute the phase transition thresholds consists of three parts: (i) perform Taylor expansion; (ii) modify the leave-one-out technique; and (iii) size correction scheme. A detailed explanation can be found in Section 5. Hopefully, it will serve independent technical interests for researchers in the machine learning community.

Notations. We use $a \xrightarrow{\text{a.s.}} b$ to suggest a converges almost surely to b . We denote $f(n) \approx g(n)$ when $\lim_{n \rightarrow \infty} f(n)/g(n) = 1$. We denote $f(n) = \mathcal{O}_{\mathbb{P}}(g(n))$ if the sequence $f(n)/g(n)$ is bounded in probability; while we denote $f(n) = o_{\mathbb{P}}(g(n))$ if the sequence $f(n)/g(n)$ converges to zero in probability. The inner product between two vectors (resp. matrices) are denoted as $\langle \cdot, \cdot \rangle$. In addition, for two distributions d_1 and d_2 , we write $d_1 \cong d_2$ if they are the equal up to some normalization. Moreover, we define \mathcal{P}_n as the set of all possible permutation matrices, i.e., $\mathcal{P}_n \triangleq \{\mathbf{\Pi} \in \{0, 1\}^{n \times n}, \sum_i \mathbf{\Pi}_{ij} = 1, \sum_j \mathbf{\Pi}_{ij} = 1\}$; and associate each permutation matrix $\mathbf{\Pi} \in \mathcal{P}_n$ with a mapping π of $\{1, 2, \dots, n\}$, where $\pi(i)$ denotes the correspondence of index i permuted by $\mathbf{\Pi}$, $1 \leq i \leq n$. The *signal-to-noise-ratio* (snr) is written as $\|\mathbf{B}^{\natural}\|_{\text{F}}^2 / (m \cdot \sigma^2)$, where $\|\cdot\|_{\text{F}}$ is the Frobenius norm and σ^2 denotes the variance of the sensing noise.

2 PROBLEM SETTING

In this paper, we consider the linear regression with permuted labels reading as

$$\mathbf{Y} = \mathbf{\Pi}^{\natural} \mathbf{X} \mathbf{B}^{\natural} + \sigma \mathbf{W},$$

where $\mathbf{Y} \in \mathbb{R}^{n \times m}$ represents the sensing result, $\mathbf{\Pi}^{\natural} \in \mathcal{P}_n$ denotes a permutation matrix awaiting to be reconstructed, $\mathbf{X} \in \mathbb{R}^{n \times p}$ is the sensing matrix with each entry X_{ij} following the i.i.d standard

normal distribution, $\mathbf{B}^\natural \in \mathbb{R}^{p \times m}$ is the signal of interests, and $\mathbf{W} \in \mathbb{R}^{n \times m}$ represents the additive sensing noise and its entries \mathbf{W}_{ij} are i.i.d standard normal random variables. In addition, we denote h as the Hamming distance between the identity matrix and the permutation matrix $\mathbf{\Pi}^\natural$, i.e., $h \triangleq \sum_i \mathbb{1}(\pi^\natural(i) \neq i)$.

The goal is to reconstruct the permutation matrix $\mathbf{\Pi}^\natural$ from the pair (\mathbf{Y}, \mathbf{X}) . As is well known in the previous works (Zhang & Li, 2020; Pananjady et al., 2018; Zhang et al., 2022), there exists a phase transition phenomenon inherent in this problem. However, all these work only present the statistical order without specifying the constants. In this work, we would like to identify the precise position of the phase transition points in the large-system limit, i.e., n, m, p , and h all approach to infinity with $m/n \rightarrow \tau_m, p/n \rightarrow \tau_p$ and $h/n \rightarrow \tau_h$.¹

Inspired by the Mezard & Montanari (2009); Semerjian et al. (2020); Chertkov et al. (2010), we borrow the tools from the statistical physics to identify the precise location of the phase transition threshold. In the following context, we separately study the phase transition phenomenon in (i) the oracle case where \mathbf{B}^\natural is given as a prior and (ii) the non-oracle case where \mathbf{B}^\natural is unknown.

3 BACKGROUND KNOWLEDGE ON GRAPHICAL MODELS

To begin with, we briefly review the *linear assignment problem* (LAP), which is defined as

$$\hat{\mathbf{\Pi}} = \operatorname{argmin}_{\mathbf{\Pi} \in \mathcal{P}_n} \langle \mathbf{\Pi}, -\mathbf{E} \rangle, \quad (1)$$

where $\mathbf{E} \in \mathbb{R}^{n \times n}$ is a fixed matrix and \mathcal{P}_n denotes the set of all possible permutation matrices. The following context investigates the behavior of $\hat{\mathbf{\Pi}}$ with the *message-passing* (MP) algorithm. First, we follow the approach in Semerjian et al. (2020); Mezard & Montanari (2009) and introduce a probability measure over the permutation matrix $\mathbf{\Pi}$, which reads as

$$\mu(\mathbf{\Pi}) = Z^{-1} \prod_i \mathbb{1}\left(1 - \sum_j \Pi_{ij}\right) \prod_j \mathbb{1}\left(1 - \sum_i \Pi_{ij}\right) e^{-\beta \sum_{i,j} \Pi_{ij} E_{ij}}, \quad (2)$$

where $\mathbb{1}(\cdot)$ is the indicator function, Z is the normalization constant of the probability measure $\mu(\mathbf{\Pi})$, and $\beta > 0$ is an auxiliary parameter. We can verify that the solution to the ML estimator maximizes the probability measure $\mu(\mathbf{\Pi})$, which means we can study the properties of $\hat{\mathbf{\Pi}}$ via the configuration $\operatorname{argmax}_{\mathbf{\Pi}} \mu(\mathbf{\Pi})$. In addition, we notice the probability measure $\mu(\mathbf{\Pi})$ concentrates on $\hat{\mathbf{\Pi}}$ when letting $\beta \rightarrow \infty$. Then we identify the phase transition thresholds by studying the marginals of $\mu(\mathbf{\Pi})$.

3.1 CONSTRUCTION OF GRAPHICAL MODEL

To start with, we construct the factor graph associated with the probability measure in (2). Adopting the same strategy as in Chapter 16 in Mezard & Montanari (2009), we (i) associate each variable Π_{ij} a variable node v_{ij} ; (ii) connect the variable node v_{ij} a function node representing the term $e^{-\beta \Pi_{ij} E_{ij}}$; and (iii) associate each constraint $\sum_i \Pi_{ij} = 1$ one function node and similarly for the constraint $\sum_j \Pi_{ij} = 1$. A graphical representation is put in Figure 1.

Then we briefly review the MP algorithm. Informally speaking, MP is local algorithm to compute the marginal probabilities over the graphical model. In each iteration, the variable node v transmits the message to its incident function node f by multiplying all incoming messages except that along the edge (v, f) . And the function




Figure 1: Illustration of the constructed graphical model. The circle icons represents the variable node; while the square icons represent the function node: the blue square icon represents the constraints for the rows of $\mathbf{\Pi}$, the green square icon represents the constraints for the columns of $\mathbf{\Pi}$, and the red square icon denotes the function $e^{-\beta \pi E_{ij}}$.

¹Although our analysis concerns the large-system limit, numerical results matches our predicted results to a good extent even when n, m, p , and h are a few hundreds.

node f transmits the message to its incident variable node v by computing the weighted summary of all incoming messages except that along the edge (f, v) . For a detailed introduction to MP, we refer readers to Kschischang et al. (2001); Mezard & Montanari (2009).

MP is able to obtain the exact marginals (Mezard & Montanari, 2009) for the tree-like graphical models. While for graphs with a lot of short loops, which happens to be our case, this claim may become invalid. However, past works all suggest that MP can still obtain meaningful results when applying to LAP (Mezard & Montanari, 2009; Semerjian et al., 2020).

3.2 MESSAGE PASSING (MP) ALGORITHM

Next, we turn to the permutation recovery via MP. The following derivation follows the standard procedure, which can be found in the previous works (Semerjian et al., 2020; Mezard & Montanari, 2009). Denote the message flow from the node i^L to the variable node (i^L, j^R) as $\widehat{m}_{i^L \rightarrow (i^L, j^R)}(\cdot)$ and that from the edge (i^L, j^R) to node i^L as $m_{(i^L, j^R) \rightarrow i^L}(\cdot)$. Similarly, we define $\widehat{m}_{j^R \rightarrow (i^L, j^R)}(\cdot)$ and $\widehat{m}_{(i^L, j^R) \rightarrow j^R}(\cdot)$ as the message flow transmitted between the functional node j^R and the variable node (i^L, j^R) . Here the superscripts L and R are used to indicate the positions of the node (left and right). Roughly speaking, these transmitted messages can be viewed as (unnormalized) conditional probability $\mathbb{P}(\Pi_{i,j} = \{0, 1\} | (\cdot))$ with the joint pdf being defined in (2). And the message transmission process is to iteratively compute these conditional probabilities. For a more detailed introduction of the MP algorithm, we refer to Mezard & Montanari (2009); MacKay et al. (2003).

First, we consider the message flows transmitted between the functional node i^L and the variable node (i^L, j^R) , which are written as

$$\begin{aligned} m_{(i^L, j^R) \rightarrow i^L}(\pi) &\cong \widehat{m}_{j^R \rightarrow (i^L, j^R)}(\pi) e^{-\beta \pi E_{i^L, j^R}}; \\ \widehat{m}_{i^L \rightarrow (i^L, j^R)}(\pi) &\cong \sum_{\pi_{i^L, k^R}} \prod_{k^R \neq j^R} \widehat{m}_{k^R \rightarrow (i^L, k^R)}(\pi_{i^L, k^R}) e^{-\beta \pi_{i^L, k^R} E_{i^L, k^R}} \mathbb{1}(\pi + \sum_k \pi_{i^L, k^R} = 1), \end{aligned} \quad (3)$$

where $\pi \in \{0, 1\}$ is a binary data. Similarly, we can write the message flows between the functional node j^R and the variable node, which are denoted as (i^L, j^R) as $m_{(i^L, j^R) \rightarrow j^R}(\pi)$ and $\widehat{m}_{j^R \rightarrow (i^L, j^R)}(\pi)$, respectively. With the parametrization method, we define

$$h_{i^L \rightarrow (i^L, j^R)} \triangleq \frac{1}{\beta} \log \frac{\widehat{m}_{i^L \rightarrow (i^L, j^R)}(1)}{\widehat{m}_{i^L \rightarrow (i^L, j^R)}(0)}; \quad h_{j^R \rightarrow (i^L, j^R)} \triangleq \frac{1}{\beta} \log \frac{\widehat{m}_{j^R \rightarrow (i^L, j^R)}(1)}{\widehat{m}_{j^R \rightarrow (i^L, j^R)}(0)}.$$

Define ζ as $h_{i^L \rightarrow (i^L, j^R)} + h_{j^R \rightarrow (i^L, j^R)} - E_{i^L, j^R}$, we select the edge (i^L, j^R) according to the probability $m_{(i^L, j^R)}(\pi) \triangleq \frac{\exp(\beta \pi \zeta_{i^L, j^R})}{1 + \exp(\beta \zeta_{i^L, j^R})}$, $\pi \in \{0, 1\}$. Provided $m_{(i^L, j^R)}(1) > m_{(i^L, j^R)}(0)$, or equivalently,

$$\zeta_{i^L, j^R} > 0, \quad (4)$$

we pick $\widehat{\pi}(i^L) = j^R$; otherwise, we have $\widehat{\pi}(i^L) \neq j^R$. Due to the fact that $\mu(\Pi)$ concentrates on $\widehat{\Pi}$ when β is sufficiently large, we can thus simplify the MP update equation as

$$h_{i^L \rightarrow (i^L, j^R)} = \min_{k^R \neq j^R} E_{i^L, k^R} - h_{k^R \rightarrow (i^L, k^R)}; \quad h_{j^R \rightarrow (i^L, j^R)} = \min_{k^L \neq i^L} E_{k^L, j^R} - h_{k^L \rightarrow (k^L, j^R)}, \quad (5)$$

which is obtained by letting $\beta \rightarrow \infty$.

4 ANALYSIS OF ORACLE CASE

As a warm-up example, we first consider the oracle scenario, where \mathbf{B}^\natural is given a prior. To reconstruct the permutation matrix Π^\natural , we adopt the *maximum-likelihood* (ML) estimator reading as

$$\widehat{\Pi}^{\text{oracle}} = \operatorname{argmax}_{\Pi} \langle \Pi, \mathbf{Y} \mathbf{B}^{\natural \top} \mathbf{X}^\top \rangle, \quad \text{s.t.} \quad \sum_i \Pi_{ij} = 1, \sum_j \Pi_{ij} = 1, \Pi \in \{0, 1\}^{n \times n}. \quad (6)$$

Denote the variable $\mathbf{E}_{ij}^{\text{oracle}}$ as $-\mathbf{X}_{\pi^\natural(i)}^\top \mathbf{B}^\natural \mathbf{B}^{\natural \top} \mathbf{X}_j + \sigma \mathbf{W}_i^\top \mathbf{B}^{\natural \top} \mathbf{X}_j$, ($1 \leq i, j \leq n$), we can transform the objective function in (6) as the canonical form of LAP, i.e., $-\sum_{i,j} \Pi_{ij} \mathbf{E}_{ij}^{\text{oracle}}$.

4.1 IDENTIFYING THE PHASE TRANSITION THRESHOLD

This subsection studies the phase transition phenomenon inherent in the MP update equation (5). Following the same strategy as in Semerjian et al. (2020), we divide all edges (i^L, j^R) into two categories based on whether the edge (i^L, j^R) corresponds to the ground-truth permutation matrix $\mathbf{\Pi}^{\natural}$ or not. Within each category, we assume the edges' weights and the message flows along them are independent identically distributed. For the edge $(i^L, \pi^{\natural}(i^L))$ corresponding to the ground-truth correspondence, we represent its weight as a random variable called Ω and the associated message flow as a random variable called H (both $h_{i^L \rightarrow (i^L, j^R)}$ and $h_{j^R \rightarrow (i^L, j^R)}$). Similarly, we define random variables $\widehat{\Omega}$ and \widehat{H} for other edges. Then we can rewrite (5) as

$$\widehat{H}^{(t+1)} = \min \left(\Omega - H^{(t)}, H'^{(t)} \right), \quad H^{(t+1)} = \min_{1 \leq i \leq n-1} \widehat{\Omega}_i - \widehat{H}_i^{(t)}, \quad (7)$$

where $(\cdot)^{(t)}$ denotes the update in the t th iteration, H' is an independent copy of H , and $\{H_i^{(t)}\}_{1 \leq i \leq n-1}$ and $\{\widehat{\Omega}_i\}_{1 \leq i \leq n-1}$ denote the i.i.d. copies of random variables $H_{(\cdot)}^{(t)}$ and $\widehat{\Omega}_{(\cdot)}$.

Then we turn to computing the critical point where the permutation matrix can be perfectly reconstructed. According to (4), this means the event $H + H' > \Omega$ holds with probability one. Conditional on this event, we can simplify (7) to be

$$H^{(t+1)} = \min_{1 \leq i \leq n-1} H_i^{(t)} + \Xi_i, \quad (8)$$

where the random variable Ξ is defined as the difference between $\widehat{\Omega}$ and Ω , i.e., $\Xi \triangleq \widehat{\Omega} - \Omega$; and $\{H_i^{(t)}\}_{1 \leq i \leq n-1}$ and $\{\Xi_i\}_{1 \leq i \leq n-1}$ denote the i.i.d. copies of random variables $H_{(\cdot)}^{(t)}$ and $\Xi_{(\cdot)}$.

This equation can be viewed as the analogous version of the *density evolution* and *state evolution*, which are used to analyze the convergence of the message passing and approximate message passing algorithm, respectively (Chung, 2000; Richardson & Urbanke, 2001; 2008; Maleki, 2010; Donoho et al., 2009; Bayati & Montanari, 2011; Rangan, 2011). Adopting the same viewpoint of Semerjian et al. (2020), we treat (8) as a *branching random walk* (BRW) process, which satisfies

Theorem 1 ((Biggins, 1977; Hammersley, 1974; Kingman, 1975; Semerjian et al., 2020)). *Consider the recursive distributional equation $K^{(t+1)} = \min_{1 \leq i \leq n} K_i^{(t)} + \Xi_i$, where $K_i^{(t)}$ and Ξ_i are i.i.d. copies of random variables $K_{(\cdot)}^{(t)}$ and $\Xi_{(\cdot)}$, we have $\frac{K^{(t+1)}}{t} \xrightarrow{\text{a.s.}} -\inf_{\theta > 0} \frac{1}{\theta} \log \left[\sum_{i=1}^n \mathbb{E} e^{-\theta \Xi_i} \right]$, conditional on the event $\lim_{t \rightarrow \infty} K^{(t)} \neq \infty$.*

With Theorem 1, we conclude the critical point for the correct permutation recovery, i.e., $H + H' > \Omega$, can be computed by letting $\inf_{\theta > 0} \frac{1}{\theta} \log \left[\sum_{i=1}^n \mathbb{E} e^{-\theta \Xi_i} \right] \leq 0$, since otherwise the condition in (4) will be violated. In the oracle case where \mathbf{B}^{\natural} is known, we have random variable Ξ be written as

$$\Xi = \mathbf{x}^{\top} \mathbf{B}^{\natural} \mathbf{B}^{\natural \top} (\mathbf{x} - \mathbf{y}) + \sigma \mathbf{w} \mathbf{B}^{\natural \top} (\mathbf{x} - \mathbf{y}), \quad (9)$$

where \mathbf{x} and \mathbf{y} follow the distribution $\mathcal{N}(\mathbf{0}, \mathbf{I}_{p \times p})$, and \mathbf{w} follows the distribution $\mathcal{N}(\mathbf{0}, \mathbf{I}_{m \times m})$.

For the convenience of computation, we consider the simple case where \mathbf{B}^{\natural} is $\lambda \mathbf{I}_{p \times p}$ ($m = p$).

Proposition 1. *Consider the case where \mathbf{B}^{\natural} is a re-scaled version of the identity matrix, i.e., $\lambda \mathbf{I}_{p \times p}$, we can write the expectation $\mathbb{E} e^{-\theta \Xi}$, which is defined in (9), as*

$$\mathbb{E} e^{-\theta \Xi} = \left(1 + 2\theta\lambda^2 - \theta^2\lambda^2 (\lambda^2 + 2\sigma^2) \right)^{-\frac{m}{2}} \quad (10)$$

provided that $\theta^2\sigma^2\lambda^2 < 1$, and $\theta^2\lambda^2 (\lambda^2 + 2\sigma^2) \leq 1 + 2\theta\lambda^2$. (11)

Provided the conditions in (11) is violated, we have the expectation $\mathbb{E} e^{-\theta \Xi}$ to diverge to infinity, which suggests the optimal θ_* for $\inf_{\theta > 0} \frac{\log(n \mathbb{E} e^{-\theta \Xi})}{\theta}$ cannot be achieved. Using (10), we can compute the optimal θ_* as $\theta_* \cong \sqrt{\frac{2 \log n}{\|\mathbf{B}^{\natural \top} \mathbf{B}^{\natural}\|_F^2 + 2\sigma^2 \|\mathbf{B}^{\natural}\|_F^2}}$. The corresponding $\text{snr}_{\text{oracle}}$ is written as

$$\text{snr}_{\text{oracle}} \cong \frac{4 \log n}{m - 2 \log n}. \quad (12)$$

The comparison between the theoretical values of the phase transition threshold and the numerical values are put in Table 1, from which we conclude the phase transition threshold snr can be predicted to a good extent. As m increases, we believe the gap between the theoretical values and the numerical values will keep shrinking.

Table 1: Comparison between the predicted value of the phase transition threshold $\text{snr}_{\text{oracle}}$ and its numerical value when $n = 500$. **P** denotes the predicted value while **N** denotes the numerical value. **N** value corresponds to the $\text{snr}_{\text{oracle}}$ when the error rate drops below 0.05.)

m	20	30	40	50	60	70
P	3.283	1.415	0.902	0.662	0.523	0.432
N	2.466	1.290	0.862	0.644	0.513	0.426

4.2 APPROXIMATED COMPUTATION OF PHASE TRANSITION THRESHOLDS

The computation of phase transition threshold is by setting $\inf_{\theta > 0} \log(\mathbb{E}Z \cdot \mathbb{E}e^{-\theta\Xi})/\theta$ be zero. However, in certain scenarios, it can be extremely difficult or impossible to obtain a closed-formula of $\mathbb{E}e^{-\theta\Xi}$, let alone the optimal solution θ . To handle such difficulties, we propose to approximate $\mathbb{E}e^{-\theta\Xi}$ with Taylor expansion, which proceeds as

$$\mathbb{E}e^{-\theta\Xi} = e^{-\theta\mathbb{E}\Xi} \cdot \mathbb{E}e^{-\theta(\Xi - \mathbb{E}\Xi)} \stackrel{\textcircled{1}}{=} e^{-\theta\mathbb{E}\Xi} \cdot \left[1 + \frac{\theta^2}{2} \cdot \mathbb{E}(\Xi - \mathbb{E}\Xi)^2 + \mathcal{O}_{\mathbb{P}}\left(\theta^4 \mathbb{E}(\Xi - \mathbb{E}\Xi)^4\right) \right],$$

where $\textcircled{1}$ is due to the fact $\mathbb{E}(\Xi - \mathbb{E}\Xi)^3 = 0$. To simplify the computation, we adopt one widely-used assumption, stating as

Assumption 1. We assume $\theta^4 \cdot \mathbb{E}(\Xi - \mathbb{E}\Xi)^4$ to be negligible.

Then we obtain

$$\mathbb{E}e^{-\theta\Xi} \approx e^{-\theta\mathbb{E}\Xi} \cdot \left(1 + \frac{\theta^2}{2} \text{Var}\Xi \right) \stackrel{\textcircled{2}}{\approx} \exp\left(-\theta\mathbb{E}\Xi + \frac{\theta^2}{2} \text{Var}\Xi\right), \quad (13)$$

where in $\textcircled{2}$ we use the approximation $1 + x \approx e^x$ when x is near zero. In this way, the rather complicated computation of $\mathbb{E}e^{-\theta\Xi}$ is replaced by the computation of mean $\mathbb{E}\Xi$ and variance $\text{Var}\Xi$, which is still complex but manageable. With this approximation, the optimal θ_* for $\log(n\mathbb{E}e^{-\theta\Xi})/\theta$ is computed as $\sqrt{2 \log n / \text{Var}\Xi}$ and hence the critical point corresponding to the phase transition as

$$2(\log n)\text{Var}\Xi = (\mathbb{E}\Xi)^2. \quad (14)$$

To verify that this approximation can yield meaningful results, we revisit the oracle case and have

$$\mathbb{E}\Xi = \|\mathbf{B}^{\natural}\|_{\text{F}}^2; \quad \text{Var}\Xi = 3\|\mathbf{B}^{\natural}\mathbf{B}^{\natural\top}\|_{\text{F}}^2 + 2\sigma^2\|\mathbf{B}^{\natural}\|_{\text{F}}^2. \quad (15)$$

Plugging (15) into (14) then yields the relation

$$6 \log n \|\mathbf{B}^{\natural}\mathbf{B}^{\natural\top}\|_{\text{F}}^2 + 4\sigma^2(\log n)\|\mathbf{B}^{\natural}\|_{\text{F}}^2 = \|\mathbf{B}^{\natural}\|_{\text{F}}^4,$$

from which we can determine the critical point of snr .

Discussion. As a comparison, we first revisit the simple case where \mathbf{B}^{\natural} is $\lambda\mathbf{I}_{p \times p}$ ($p = m$). We can compute the phase transition snr as $4 \log n / (m - 6 \log n)$. This solution is almost identical to (12) in the large-system limit as $\text{snr}_{\text{oracle}} \approx \widetilde{\text{snr}}_{\text{oracle}} \approx n^{\frac{4}{m}} - 1$. Moreover, we should stress that (i) our approximation method applies to other types of matrices as well, rather than limited to the identity matrix; and (ii) our approximation method can predict the phase transition thresholds even when the entries \mathbf{X}_{ij} are sub-gaussian. An illustration is given in Table 2: in **(Case I)**, half of eigenvalues are with Ener while the other half are with $\text{Ener}/2$; in **(Case II)**, half of the eigenvalues are with Ener while the other half are with $3\text{Ener}/4$.

5 ANALYSIS OF NON-ORACLE CASE

Having presented the oracle case as a warm-up example, we now extend the analysis to the non-oracle case, where the value of \mathbf{B}^{\natural} is not given a priori. To begin with, we need to recast the permutation recovery problem as a LAP. As shown in Zhang et al. (2022), the ML estimator yields a *quadratic assignment problem* (QAP), which is NP-hard to solve and fails to meet this requirement. Fortunately,

Table 2: Comparison between the predicted value of the phase transition threshold $\text{snr}_{\text{oracle}}$ and its numerical value when $n = 800$. **Gauss** refers to $\mathbf{X}_{ij} \stackrel{\text{i.i.d}}{\sim} \mathcal{N}(0, 1)$ while **Unif** refers to $\mathbf{X}_{ij} \stackrel{\text{i.i.d}}{\sim} \text{Unif}[-1, 1]$. We averaged over 20 experiments.

m	100	110	120	130
(Case I) P	0.63	0.52	0.43	0.38
(Gauss) N	0.6 ~ 0.65	0.55 ~ 0.6	0.55 ~ 0.6	0.5 ~ 0.55
(Unif) N	0.6 ~ 0.65	0.6 ~ 0.65	0.55 ~ 0.6	0.55 ~ 0.6
(Case II) P	0.47	0.40	0.35	0.31
(Gauss) N	0.5 ~ 0.55	0.5 ~ 0.55	0.45 ~ 0.5	0.45 ~ 0.5
(Unif) N	0.5 ~ 0.55	0.5	0.45 ~ 0.5	0.45 ~ 0.5

the proposed estimator in Zhang & Li (2020) is able to fill the gap. Define the edge weight $\mathbf{E}_{ij}^{\text{non-oracle}}$ as $\mathbf{E}_{ij}^{\text{non-oracle}} \triangleq -\mathbf{Y}_i \mathbf{Y}_j^\top \mathbf{X} \mathbf{X}_j^\top$, we can reconstruct the permutation matrix $\mathbf{\Pi}^{\natural}$ as (1). Before proceeding, we would like to justify using the estimator in Zhang & Li (2020) to analyze the permuted linear regression in the non-oracle case: first, this estimator is proved to achieve both the computational and statistical optimality; second, this estimator also exhibits a phase transition phenomenon, which behaves similarly to that in the oracle case. Naturally, we should expect this estimator will incorporate some inherent properties of the permutation recovery problem (with unknown \mathbf{B}^{\natural}), from which we can gain meaningful insights.

5.1 ANALYSIS OF NON-ORACLE CASE

Having illustrated soundness of the approximation method in (13), we apply it to the non-oracle case, where the random variable is written as

$$\Xi = \Xi_1 + \sigma(\Xi_2 + \Xi_3) + \sigma^2 \Xi_4, \quad (16)$$

where Ξ_i ($1 \leq i \leq 4$) are defined as

$$\begin{aligned} \Xi_1 &\triangleq \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{B}^{\natural} \mathbf{B}^{\natural\top} \mathbf{X}^\top \mathbf{\Pi}^{\natural\top} \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j); & \Xi_2 &\triangleq \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{B}^{\natural} \mathbf{W}^\top \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j); \\ \Xi_3 &\triangleq \mathbf{W}_i^\top \mathbf{B}^{\natural\top} \mathbf{X}^\top \mathbf{\Pi}^{\natural\top} \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j); & \Xi_4 &\triangleq \mathbf{W}_i^\top \mathbf{W}^\top \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j), \end{aligned}$$

respectively. Then we conclude

Theorem 2. *The mean $\mathbb{E}\Xi$ of Ξ in (39) and its variance $\text{Var}\Xi$ are computed as*

$$\begin{aligned} \mathbb{E}\Xi &\cong n(1 - \tau_h) \left[(1 + \tau_p) \|\mathbf{B}^{\natural}\|_{\text{F}}^2 + n\tau_m\tau_p\sigma^2 \right]; \\ \text{Var}\Xi &\cong n^2\tau_h(1 - \tau_h)\tau_p^2 \left[\|\mathbf{B}^{\natural}\|_{\text{F}}^2 + m\sigma^2 \right]^2 + n^2 \left[2\tau_p + 3(1 - \tau_h)^2 \right] \|\mathbf{B}^{\natural\top} \mathbf{B}^{\natural}\|_{\text{F}}^2 \\ &\quad + n^2 \left[6\tau_p(1 - \tau_h)^2 + (3 - \tau_h)\tau_p^2 \right] \|\mathbf{B}^{\natural\top} \mathbf{B}^{\natural}\|_{\text{F}}^2, \end{aligned}$$

respectively, where the definitions of τ_p , τ_m and τ_h can be found in Section 2.

For the clarity of presentation, we only present the proof outlines.

Computation of mean $\mathbb{E}\Xi$. For the computation of the mean $\mathbb{E}\Xi$, easily we can verify that $\mathbb{E}\Xi_2$ and $\mathbb{E}\Xi_3$ are both zero, which is due to the independence between \mathbf{X} and \mathbf{W} . Regarding the computation of $\mathbb{E}\Xi_1$ and $\mathbb{E}\Xi_4$, we adopt Wick's theorem and obtain

$$\mathbb{E}\Xi_1 = n(1 - \tau_h)(1 + \tau_p) [1 + o_{\text{P}}(1)] \|\mathbf{B}^{\natural}\|_{\text{F}}^2; \quad \mathbb{E}\Xi_4 = n^2\tau_m\tau_p(1 - \tau_h)(1 + o_{\text{P}}(1)).$$

Computation of variance $\text{Var}\Xi$. With the relation $\text{Var}(\Xi) = \mathbb{E}\Xi^2 - (\mathbb{E}\Xi)^2$, our goal becomes computing $\mathbb{E}\Xi^2$, which consists the calculation of the following six terms

$$\mathbb{E}\Xi^2 = \mathbb{E}\Xi_1^2 + \sigma^2\mathbb{E}\Xi_2^2 + \sigma^2\mathbb{E}\Xi_3^2 + \sigma^4\mathbb{E}\Xi_4^2 + 2\sigma^2\mathbb{E}\Xi_1\Xi_4 + 2\sigma^2\mathbb{E}\Xi_2\Xi_3.$$

The computation of above terms turns to be quite complex due to the high order Gaussian chaos. For example, term $\mathbb{E}\Xi_1^2$ involves the eighth-order Gaussian chaos; terms $\mathbb{E}\Xi_2^2$, $\mathbb{E}\Xi_3^2$, $\mathbb{E}\Xi_1\Xi_4$ and $\mathbb{E}\Xi_2\Xi_3$ all involves the sixth-order Gaussian variables. To alleviate the computational burden, we compute the expectation $\mathbb{E}\Xi^2$ in the following three phases.

Table 3: Comparison between the predicted value of the phase transition threshold τ_h and its numerical value when $n = 500$. \mathbf{P} denotes the predicted value while \mathbf{N} denotes the numerical value. (The numerical value of τ_h is the minimum τ_h when the correct permutation rate drops below 0.05.)

p	75	100	125	150	175	200
\mathbf{P}	0.82	0.73	0.68	0.62	0.56	0.52
\mathbf{N}	0.77	0.74	0.7	0.66	0.61	0.57

- **Phase I.** The solution in this phase comes from a modification of the so-called *leave-one-out* technique (Sur et al., 2019; El Karoui, 2013; 2018; Bai & Silverstein, 2010). Notice that the major technical difficulty comes from the correlation between the product $\mathbf{X}^\top \mathbf{\Pi}^{\natural} \mathbf{X}$ and the difference $\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j$. We decompose this correlation by first rewriting the matrix $\mathbf{X}^\top \mathbf{\Pi}^{\natural} \mathbf{X}$ as the sum $\sum_{\ell} \mathbf{X}_{\ell} \mathbf{X}_{\pi^{\natural}(\ell)}^\top$. Then we collect all terms $\mathbf{X}_{\ell} \mathbf{X}_{\pi^{\natural}(\ell)}^\top$ independent of $\mathbf{X}_{\pi^{\natural}(i)}$ and \mathbf{X}_j in the matrix $\mathbf{\Sigma}$ and leave the rest terms to matrix $\mathbf{\Delta}$, which means $\mathbf{\Delta} \triangleq \mathbf{X}^\top \mathbf{\Pi}^{\natural} \mathbf{X} - \mathbf{\Sigma}$. This decomposition is in the same spirit of the leave-one-out technique.

With this method, we divide all terms involved in the computation of $\mathbb{E}\Xi^2$ into three categories: (i) those only containing matrix $\mathbf{\Sigma}$; (ii) those containing both $\mathbf{\Sigma}$ and $\mathbf{\Delta}$; and (iii) those only containing $\mathbf{\Delta}$. Easily we can see that the first two categories contain most vectors' outer products while the last category only contains a finite number of such terms.

- **Phase II.** Concerning the terms in the first two categories, which contains majority of terms, we can exploit the independence among rows in the sensing matrix \mathbf{X} and reduce the order of Gaussian random variables by separately taking expectation w.r.t $\mathbf{\Sigma}$ and w.r.t vectors $\mathbf{X}_{\pi^{\natural}(i)}$ and \mathbf{X}_j .
- **Phase III.** For the few terms in the third category which contains high-order Gaussian chaos, we compute their expectations by iterative applying of Wick's Theorem and Stein's Lemma, which filters out the zero terms to reduce higher-order interactions between Gaussian random variables to lower-order interactions.

For more technical details, we refer the interested readers to the supplementary material.

5.2 IDENTIFYING THE PHASE TRANSITION THRESHOLD

Having explained the computation of $\mathbb{E}\Xi$ and $\text{Var}\Xi$, we turn to identifying the phase transition threshold. Different from the oracle case, we notice the edge weight E_{ij} are strongly correlated in this case especially when $j = \pi^{\natural}(j)$, which corresponds to the non-permuted rows. To factor out these independence, we only take the permuted rows into account and correct the sample size from n to $\tau_h n$. Thus, we have

Proposition 2. *The critical point for the phase transition phenomenon transforms from (14) to $2(\log \tau_h n) \text{Var}\Xi = (\mathbb{E}\Xi)^2$.*

Example 1. *We consider the case where $\mathbf{B}^{\natural} = \lambda \mathbf{I}_{p \times p}$ as an illustration. With Theorem 2 and Proposition 2, we obtain the solution*

$$\text{snr}_{\text{non-oracle}} = \eta_1 / \eta_2, \quad (17)$$

where η_1 and η_2 are defined as

$$\begin{aligned} \eta_1 &\triangleq 2\tau_h \tau_p^2 \log(n\tau_h) - \tau_p(\tau_p + 1)(1 - \tau_h) + \sqrt{2}\tau_p \sqrt{(1 - \tau_h)\tau_h(\log(n\tau_h))}; \\ \eta_2 &\triangleq 2\tau_h \tau_p^2 \log(n\tau_h) - (1 - \tau_h)(\tau_p + 1)^2. \end{aligned}$$

Notice that the negative solution has been abandoned due to the non-negativity requirement of snr.

Discussion. *For the accuracy of the predicted phase transition $\text{snr}_{\text{non-oracle}}$, we notice a increasing gap between the theoretical value and the numerical value when compared with that in the oracle case. Possible reasons include strong correlation across the edge weights $\{E_{ij}\}_{1 \leq i, j \leq n}$ and the error within the approximation relation $\mathbb{E}e^{-\theta\Xi} \approx \mathbb{E}\exp(\theta\mathbb{E}\Xi - \theta^2\text{Var}\Xi/2)$.*

In addition, we observe a singularity point, i.e., τ_h is approximately 0.73 in Figure 2, which suggests a phase transition phenomenon. To validate the predicted phenomenon, we consider the noiseless

case, i.e., $\text{snr} = \infty$, and reconstruct the permutation matrix $\Pi^{\mathfrak{b}}$ with (1). Numerical experiments confirm our prediction by showing that the correct rate of the permutation recovery exceeds 0.2 when $h/n \leq 0.73$ and drops below 0.05 when $h/n > 0.74$. Additional experiments are put in Table 3, from which we conclude the solution (17) can predict the critical points w.r.t. τ_h to a good extent.

Remark 1. Compared with the prior work (Zhang & Li, 2020) which only yields the statistical order, i.e. $h/n \leq c$ (c is a positive constant), our framework can (i) specify the positive constant c , and (ii) uncover the dependence of τ_h on the ratio τ_p . In addition, Zhang & Li (2020) requires $n \gg p$ while our work allows n to be the same order of p , i.e., $p/n = \tau_p$ as $n, p \rightarrow \infty$. Notice that this is consistent with numerical experiments such that $n = \mathcal{O}_p(p)$ is sufficient for the permutation recovery.

6 PARTIAL PERMUTATION RECOVERY

Apart from the phase transition thresholds, we would like to exploit MP to design algorithms for a partial permutation recovery. As an illustration, we consider recovering the correspondence $\pi^{\mathfrak{b}}(i)$ for a single index i . In an effortless way, we can generalize it to recovering the correspondences for multiple indices by iterative applying the following procedure.

To start with, we modify the graphical model in Figure 1 by removing the unnecessary function nodes and their incident edge. Then we can rewrite the MP update equation

$$\widehat{m}_{i^L \rightarrow (i^L, j^R)}(\pi) \simeq \sum_{\pi_{i^L, k^R}} \prod_{k^R \neq j^R} \exp(-\beta \pi_{i^L, k^R} E_{i^L, k^R}) \mathbb{1}(\pi + \sum_k \pi_{i^L, k^R} = 1).$$

Compared with the MP for the full permutation recovery, MP for the partial permutation recovery can reach convergence in one iteration. Letting $\beta \rightarrow \infty$, we obtain the edge selection criteria as

$$\widehat{\pi}(i^R) = \operatorname{argmin}_j E_{ij},$$

which turns out to be a greedy selection scheme.

Following the same procedure as in Sections 4 and 5, we can analyze its statistical properties, which are similar (although degraded) to the estimator for the full permutation recovery thereof (both the oracle case and non-oracle case). This claim is also confirmed by numerical experiments in Figure 3.

7 CONCLUSION

This is the first work that can identify the precise location of phase transition thresholds of permuted linear regressions. For the oracle case where the signal $\mathbf{B}^{\mathfrak{b}}$ is given a prior, our analysis can predict the phase transition threshold $\text{snr}_{\text{oracle}}$ to a good extent. For the non-oracle case where $\mathbf{B}^{\mathfrak{b}}$ is not given, we modified the leave-one-out technique to approximately compute the phase critical $\text{snr}_{\text{non-oracle}}$ value for the phase transition, as the precise computation becomes significantly complicated as the high-order interaction between Gaussian random variables is involved. Moreover, we associated the singularity point in $\text{snr}_{\text{non-oracle}}$ with a phase transition point w.r.t the maximum allowed number of permuted rows. In the end, we generalized the full permutation recovery and obtained a partial permutation recovery algorithm. Following the same analytical procedure, we argued it would have a similar although degraded performance compared with the full permutation recovery algorithm, which is later confirmed by our numerical experiments. In the future, we will incorporate the *replica symmetry breaking* scheme into our framework and extend this framework to broad areas.




Figure 2: $\text{snr}_{\text{non-oracle}}$ when $n = 500$ and $p = 100$.




Figure 3: Comparison between the full permutation recovery and partial permutation recovery. We set $n = 500$, $p = 100$, and $m = 75$. The partial permutation recovery estimator also exhibits a phase transition, similar to the full permutation recovery estimator. However, their phase transition points are larger when compared with the corresponding points for the full permutation recovery.

REFERENCES

- Zhidong Bai and Tailen Hsing. The broken sample problem. *Probab. Theory Relat. Fields*, 131(4): 528–552, 2005.
- Zhidong Bai and Jack W Silverstein. *Spectral analysis of large dimensional random matrices*, volume 20. Springer, 2010.
- Mohsen Bayati and Andrea Montanari. The dynamics of message passing on dense graphs, with applications to compressed sensing. *IEEE Transactions on Information Theory*, 57(2):764–785, 2011.
- JD464415 Biggins. Chernoff’s theorem in the branching random walk. *Journal of Applied Probability*, 14(3):630–636, 1977.
- Sergio Caracciolo, Matteo P D’Achille, Enrico M Malatesta, and Gabriele Sicuro. Finite-size corrections in the random assignment problem. *Physical Review E*, 95(5):052129, 2017.
- Michael Chertkov, Lukas Kroc, F Krzakala, M Vergassola, and L Zdeborová. Inference in particle tracking experiments by passing messages between images. *Proceedings of the National Academy of Sciences*, 107(17):7663–7668, 2010.
- Sae-Young Chung. *On the construction of some capacity-approaching coding schemes*. PhD thesis, Massachusetts Institute of Technology, 2000.
- Morris H. DeGroot and Prem K. Goel. The matching problem for multivariate normal data. *Sankhyā: The Indian Journal of Statistics, Series B (1960-2002)*, 38(1):14–29, 1976.
- Morris H. DeGroot and Prem K. Goel. Estimation of the correlation coefficient from a broken random sample. *The Annals of Statistics*, 8(2):264–278, 03 1980.
- Morris H. DeGroot, Paul I. Feder, and Prem K. Goel. Matchmaking. *Annals of Mathematical Statistics*, 42(2):578–593, 04 1971.
- David L Donoho, Arian Maleki, and Andrea Montanari. Message-passing algorithms for compressed sensing. *Proceedings of the National Academy of Sciences*, 106(45):18914–18919, 2009.
- Noureddine El Karoui. Asymptotic behavior of unregularized and ridge-regularized high-dimensional robust regression estimators: rigorous results. *arXiv preprint arXiv:1311.2445*, 2013.
- Noureddine El Karoui. On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. *Probab. Theory Relat. Fields*, 170(1-2): 95–175, 2018.
- Prem K. Goel. On re-pairing observations in a broken random sample. *The Annals of Statistics*, 3(6): 1364–1369, 11 1975.
- John M Hammersley. Postulates for subadditive processes. *The Annals of Probability*, 2(4):652–680, 1974.
- Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge university press, 1990.
- Svante Janson. *Gaussian Hilbert Spaces*. Cambridge Tracts in Mathematics. Cambridge University Press, 1997. doi: 10.1017/CBO9780511526169.
- John Frank Charles Kingman. The first birth problem for an age-dependent branching process. *The Annals of Probability*, pp. 790–801, 1975.
- Frank R Kschischang, Brendan J Frey, and H-A Loeliger. Factor graphs and the sum-product algorithm. *IEEE Transactions on information theory*, 47(2):498–519, 2001.
- Svante Linusson and Johan Wästlund. A proof of parisi’s conjecture on the random assignment problem. *Probability theory and related fields*, 128(3):419–440, 2004.
- David JC MacKay, David JC Mac Kay, et al. *Information theory, inference and learning algorithms*. Cambridge university press, 2003.

- Enrico M Malatesta, Giorgio Parisi, and Gabriele Sicuro. Fluctuations in the random-link matching problem. *Physical Review E*, 100(3):032102, 2019.
- Mohammad Ali Maleki. *Approximate message passing algorithms for compressed sensing*. Stanford University, 2010.
- OC Martin, M Mézard, and Olivier Rivoire. Random multi-index matching problems. *Journal of Statistical Mechanics: Theory and Experiment*, 2005(09):P09006, 2005.
- Marc Mezard and Andrea Montanari. *Information, physics, and computation*. Oxford University Press, 2009.
- Marc Mézard and Giorgio Parisi. Replicas and optimization. *Journal de Physique Lettres*, 46(17):771–778, 1985.
- Marc Mézard and Giorgio Parisi. Mean-field equations for the matching and the travelling salesman problems. *EPL (Europhysics Letters)*, 2(12):913, 1986.
- Marc Mézard and Giorgio Parisi. On the solution of the random link matching problems. *Journal de Physique*, 48(9):1451–1459, 1987.
- Heinz Neudecker and Tom Wansbeek. Fourth-order properties of normally distributed random matrices. *Linear Algebra and Its Applications*, 97:13–21, 1987.
- Ashwin Pananjady, Martin J Wainwright, and Thomas A Courtade. Denoising linear models with permuted data. In *Proceedings of the 2017 IEEE International Symposium on Information Theory (ISIT)*, pp. 446–450, Aachen, Germany, 2017.
- Ashwin Pananjady, Martin J. Wainwright, and Thomas A. Courtade. Linear regression with shuffled data: Statistical and computational limits of permutation recovery. *IEEE Trans. Inf. Theory*, 64(5):3286–3300, 2018.
- Giorgio Parisi and Matthieu Ratiéville. On the finite size corrections to some random matching problems. *The European Physical Journal B-Condensed Matter and Complex Systems*, 29(3):457–468, 2002.
- Sundeep Rangan. Generalized approximate message passing for estimation with random linear mixing. In *Information Theory Proceedings (ISIT), 2011 IEEE International Symposium on*, pp. 2168–2172. IEEE, 2011.
- Thomas J Richardson and Rüdiger L Urbanke. The capacity of low-density parity-check codes under message-passing decoding. *IEEE Transactions on information theory*, 47(2):599–618, 2001.
- Tom Richardson and Ruediger Urbanke. *Modern coding theory*. Cambridge university press, 2008.
- Guilhem Semerjian, Gabriele Sicuro, and Lenka Zdeborová. Recovery thresholds in the sparse planted matching problem. *Physical Review E*, 102(2):022304, 2020.
- Martin Slawski and Emanuel Ben-David. Linear regression with sparsely permuted data. *Electron. J. Statist.*, 13(1):1–36, 2019.
- Martin Slawski, Emanuel Ben-David, and Ping Li. Two-stage approach to multivariate linear regression with sparsely mismatched data. *J. Mach. Learn. Res.*, 21(204):1–42, 2020.
- Pragya Sur, Yuxin Chen, and Emmanuel J Candès. The likelihood ratio test in high-dimensional logistic regression is asymptotically a rescaled chi-square. *Probab. Theory Relat. Fields*, 175(1-2):487–558, 2019.
- Michel Talagrand. *Mean field models for spin glasses: Volume I: Basic examples*, volume 54. Springer Science & Business Media, 2010.
- Jayakrishnan Unnikrishnan, Saeid Haghighatshoar, and Martin Vetterli. Unlabeled sensing: Solving a linear system with unordered measurements. In *Proceedings of the 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pp. 786–793, Monticello, IL, 2015.

Hang Zhang and Ping Li. Optimal estimator for unlabeled linear regression. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, pp. 11153–11162, Virtual Event, 2020.

Hang Zhang, Martin Slawski, and Ping Li. The benefits of diversity: Permutation recovery in unlabeled sensing from multiple measurement vectors. *IEEE Trans. Inf. Theory*, 68(4):2509–2529, 2022.

A APPENDIX: ANALYSIS OF ORACLE CASE

Denote random variable Ξ as

$$\Xi = \mathbf{x}^\top \mathbf{B}^\natural \mathbf{B}^{\natural\top} (\mathbf{x} - \mathbf{y}) + \sigma \mathbf{w} \mathbf{B}^{\natural\top} (\mathbf{x} - \mathbf{y}).$$

A.1 EXACT COMPUTATION

We would like to compute $\mathbb{E}e^{-\theta\Xi}$. Due to the assumption such that \mathbf{B}^\natural is proportional to the identity matrix, we can rewrite Ξ as

$$\Xi = \sum_{i=1}^m \lambda^2 x_i (x_i - y_i) + \sigma \sum_{i=1}^m \lambda w_i (x_i - y_i),$$

where λ is the principal singular value of \mathbf{B}^\natural . Due to the independent across \mathbf{x} , \mathbf{x} , and \mathbf{y} , we compute $\mathbb{E}e^{-\theta\Xi}$ as

$$\begin{aligned} \mathbb{E}e^{-\theta\Xi} &= \left[\mathbb{E}_{\mathbf{x}, \mathbf{y}, \mathbf{w}} \exp \left[-\theta \lambda^2 \mathbf{x} (\mathbf{x} - \mathbf{y}) - \theta \sigma \lambda \mathbf{w} (\mathbf{x} - \mathbf{y}) \right] \right]^m \\ &= \left[\mathbb{E}_{\mathbf{x}, \mathbf{y}} \exp \left(\frac{1}{2} \theta \lambda^2 (\mathbf{x} - \mathbf{y}) (\theta \sigma^2 (\mathbf{x} - \mathbf{y}) - 2\mathbf{x}) \right) \right]^m \\ &\stackrel{\textcircled{1}}{=} \left[\mathbb{E}_{\mathbf{x}} \frac{\exp \left(\frac{\theta \lambda^2 \mathbf{x}^2 (\theta (\lambda^2 + \sigma^2) - 2)}{2 - 2\theta^2 \lambda^2 \sigma^2} \right)}{\sqrt{1 - \theta^2 \lambda^2 \sigma^2}} \right]^m \\ &\stackrel{\textcircled{2}}{=} (1 + 2\theta \lambda^2 - \theta^2 \lambda^2 (\lambda^2 + 2\sigma^2))^{-\frac{m}{2}}, \end{aligned}$$

where in $\textcircled{1}$ we use the fact $\theta^2 \sigma^2 \lambda^2 < 1$ and in $\textcircled{2}$ we use the fact $\theta^2 \lambda^2 (\lambda^2 + 2\sigma^2) \leq 1 + 2\theta \lambda^2$. Then we relax the function

$$\begin{aligned} \frac{\log [n \mathbb{E}e^{-\theta\Xi}]}{\theta} &\stackrel{\textcircled{3}}{\approx} \frac{\log n - \frac{m}{2} (2\theta \lambda^2 - \theta^2 \lambda^2 (\lambda^2 + \sigma^2))}{\theta} \\ &= \frac{\log n}{\theta} - \|\mathbf{B}^\natural\|_F^2 + \frac{\theta \left[\|\mathbf{B}^{\natural\top} \mathbf{B}^\natural\|_F^2 + 2\sigma^2 \|\mathbf{B}^\natural\|_F^2 \right]}{2}, \end{aligned}$$

where in $\textcircled{3}$ we use the relation $\log(1 + x) \approx x$ when x is close to zero.

A.2 TAYLOR APPROXIMATION

Our goal is to compute the mean $\mathbb{E}\Xi$ and $\text{Var}\Xi$. Since their computations follow the same method as that in the non-oracle case, we omit the detailed computation for the conciseness of presentation.

B APPENDIX: ANALYSIS OF NON-ORACLE CASE

B.1 NOTATIONS

Note that our analysis can involve the terms containing $(\mathbf{X}_{\pi^\natural(i)} - \mathbf{X}_j)$ and $\mathbf{X}^\top \mathbf{\Pi}^\natural \mathbf{X}$ simultaneously. To decouple the dependence between $(\mathbf{X}_{\pi^\natural(i)} - \mathbf{X}_j)$ and $\mathbf{X}^\top \mathbf{\Pi}^\natural \mathbf{X}$, we first rewrite the matrix $\mathbf{X}^\top \mathbf{\Pi}^\natural \mathbf{X}$ as the sum $\sum_\ell \mathbf{X}_\ell \mathbf{X}_{\pi^\natural(\ell)}^\top$ and then collect all terms $\mathbf{X}_\ell \mathbf{X}_{\pi^\natural(\ell)}^\top$ independent of $\mathbf{X}_{\pi^\natural(i)}$ and \mathbf{X}_j in the matrix $\mathbf{\Sigma}$, which is written as

$$\mathbf{\Sigma} \triangleq \sum_{\ell, \pi^\natural(\ell) \neq \pi^\natural(i), j} \mathbf{X}_\ell \mathbf{X}_{\pi^\natural(\ell)}^\top. \quad (18)$$

The rest terms are then put in the matrix $\mathbf{\Delta}$ such that $\mathbf{X}^\top \mathbf{\Pi}^\natural \mathbf{X} = \mathbf{\Sigma} + \mathbf{\Delta}$. Notice that the expression of $\mathbf{\Delta}$ varies under different cases such that

- **Case (s, s) :** $i = \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. We have

$$\Delta = \Delta^{(s,s)} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top. \quad (19)$$

- **Case (s, d) :** $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. We have

$$\Delta = \Delta^{(s,d)} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top. \quad (20)$$

- **Case (d, s) :** $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. We have

$$\Delta = \Delta^{(d,s)} = \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top + \mathbf{X}_j \mathbf{X}_j^\top. \quad (21)$$

- **Case (d, d) :** $i \neq \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. We have

$$\Delta = \Delta^{(d,d)} = \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top. \quad (22)$$

In addition, we define the matrix \mathbf{M} as $\mathbf{B}^{\natural} \mathbf{B}^{\natural \top}$, and define the index sets \mathcal{S} , \mathcal{D} , and $\mathcal{D}_{\text{pair}}$ as

$$\mathcal{S} \triangleq \{\ell \mid \ell \neq i \text{ or } j, \ell = \pi^{\natural}(\ell)\}; \quad (23)$$

$$\mathcal{D} \triangleq \{\ell \mid \ell, \pi^{\natural}(\ell) \neq i \text{ or } j, \ell \neq \pi^{\natural}(\ell)\}; \quad (24)$$

$$\mathcal{D}_{\text{pair}} \triangleq \{(\ell_1, \ell_2) : \ell_1 = \pi^{\natural}(\ell_2), \ell_2 = \pi^{\natural}(\ell_1), \ell_1, \ell_2 \in \mathcal{D}\}, \quad (25)$$

respectively.

B.2 SUPPORTING LEMMAS

First, we study the higher order expectations of Gaussian random vectors' inner product, which hopefully will serve independent interests.

Lemma 1. *Assume $\mathbf{x} \in \mathbb{R}^p$ and $\mathbf{y} \in \mathbb{R}^p$ are Gaussian distributed random vectors whose entries follow the i.i.d. standard normal distribution, then we have*

$$\mathbb{E} \text{Tr}(\mathbf{y} \mathbf{y}^\top \mathbf{x} \mathbf{x}^\top \mathbf{M}) = \text{Tr}(\mathbf{M}); \quad (26)$$

$$\mathbb{E} \text{Tr}(\mathbf{y} \mathbf{y}^\top \mathbf{x}^\top \mathbf{M} \mathbf{x}) = \mathbb{E} \|\mathbf{y}\|_2^2 \text{Tr}(\mathbf{x}^\top \mathbf{M} \mathbf{x}) = p \text{Tr}(\mathbf{M}); \quad (27)$$

$$\mathbb{E}(\mathbf{x}^\top \mathbf{M} \mathbf{x})^2 = [\text{Tr}(\mathbf{M})]^2 + \text{Tr}(\mathbf{M} \mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}); \quad (28)$$

$$\mathbb{E} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) = (p+2) \text{Tr}(\mathbf{M}); \quad (29)$$

$$\mathbb{E} \|\mathbf{x}\|_2^4 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) = (p+2)(p+4) \text{Tr}(\mathbf{M}), ; \quad (30)$$

$$\mathbb{E} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})^2 = (p+4) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M} \mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right]; \quad (31)$$

$$\mathbb{E} \|\mathbf{x}\|_2^4 (\mathbf{x}^\top \mathbf{M} \mathbf{x})^2 = (p+4)(p+6) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M} \mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right]; \quad (32)$$

$$\mathbb{E}(\mathbf{x}^\top \mathbf{y})^2 \mathbf{y}^\top \mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2 \mathbf{y} = 2 \text{Tr}(\mathbf{M}_1) \text{Tr}(\mathbf{M}_2) + (p+4) \text{Tr}(\mathbf{M}_1 \mathbf{M}_2) + 2 \text{Tr}(\mathbf{M}_1 \mathbf{M}_2^\top) \quad (33)$$

where $\mathbf{M} \in \mathbb{R}^{p \times p}$ is a fixed matrix.

Remark 2. *If we assume $\mathbf{M} = \mathbf{I}_{p \times p}$, we can get $\mathbb{E} \|\mathbf{x}\|_2^4 = p(p+2)$, $\mathbb{E} \|\mathbf{x}\|_2^6 = p(p+2)(p+4)$, and $\mathbb{E} \|\mathbf{x}\|_2^8 = p(p+2)(p+4)(p+6)$.*

Proof. • **Proof of (26) and (27).** The proof can be conducted easily with the property such that $\text{Tr}(\mathbf{u} \mathbf{v}^\top) = \mathbf{u}^\top \mathbf{v} = \text{Tr}(\mathbf{v} \mathbf{u}^\top)$ holds for arbitrary vectors \mathbf{u} and \mathbf{v} .

- **Proof of (28).** This property is a direct consequence of Neudecker & Wansbeek (1987) (Equation (3.2)), which is attached in Lemma 18 for the sake of self-containing.

- **Proof of (29).** Invoking the Stein's lemma, we have

$$\mathbb{E} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) = \mathbb{E} [\nabla_{\mathbf{x}} (\mathbf{x}^\top \mathbf{M} \mathbf{x}) \mathbf{x}].$$

Then our goal transforms to computing the trace of the Hessian matrix $\text{Tr} [\nabla_{\mathbf{x}} \text{Tr}(\mathbf{x}^\top \mathbf{M} \mathbf{x})]$. For the i th entry of the gradient, we have

$$\frac{d}{dx_i} \mathbf{x}^\top \mathbf{M} \mathbf{x} = \langle \mathbf{M}_i, \mathbf{x} \rangle + \langle (\mathbf{M}^\top)_i, \mathbf{x} \rangle,$$

where \mathbf{M}_i is the i th row (or column) of \mathbf{M} . Then we obtain

$$\frac{d}{dx_i} [x_i \text{Tr}(\mathbf{x}^\top \mathbf{M} \mathbf{x})] = \mathbf{x}^\top \mathbf{M} \mathbf{x} + x_i [\langle \mathbf{M}_i, \mathbf{x} \rangle + \langle (\mathbf{M}^\top)_i, \mathbf{x} \rangle],$$

and hence

$$\begin{aligned} \mathbb{E} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) &= \sum_{i=1}^p \mathbb{E}(\mathbf{x}^\top \mathbf{M} \mathbf{x}) + \sum_{i=1}^p \mathbb{E} [x_i (\langle \mathbf{M}_i, \mathbf{x} \rangle + \langle (\mathbf{M}^\top)_i, \mathbf{x} \rangle)] \\ &= p \text{Tr}(\mathbf{M}) + 2 \sum_i \mathbf{M}_{ii} = (p+2) \text{Tr}(\mathbf{M}). \end{aligned}$$

- **Proof of (30).** Following the same strategy as in proving (29), we have

$$\mathbb{E} \|\mathbf{x}\|_{\mathbb{F}}^4 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) = \mathbb{E} [\nabla_{\mathbf{x}} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) \mathbf{x}].$$

Then our goal transforms to computing the trace of the Hessian matrix $\text{Tr} [\nabla_{\mathbf{x}} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})]$. For the i th entry of the gradient, we obtain

$$\begin{aligned} \frac{d}{dx_i} [x_i \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})] &= \|\mathbf{x}\|_2^2 \cdot (\mathbf{x}^\top \mathbf{M} \mathbf{x}) + x_i \frac{d}{dx_i} [\|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})] \\ &= \|\mathbf{x}\|_2^2 \cdot (\mathbf{x}^\top \mathbf{M} \mathbf{x}) + 2x_i^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x}) + x_i \|\mathbf{x}\|_2^2 [\langle \mathbf{M}_i, \mathbf{x} \rangle + \langle (\mathbf{M}^\top)_i, \mathbf{x} \rangle], \end{aligned}$$

whose expectation reads as

$$\begin{aligned} &\mathbb{E} \|\mathbf{x}\|_2^2 \cdot (\mathbf{x}^\top \mathbf{M} \mathbf{x}) + 2\mathbb{E} [x_i^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})] + \mathbb{E} x_i \|\mathbf{x}\|_2^2 [\langle \mathbf{M}_i, \mathbf{x} \rangle + \langle (\mathbf{M}^\top)_i, \mathbf{x} \rangle] \\ &= (p+2) \text{Tr}(\mathbf{M}) + 2\mathbb{E} \left[x_i^2 M_{ii} x_i^2 + x_i^2 \left(\sum_{j \neq i} M_{jj} x_j^2 \right) \right] + 2M_{ii} \mathbb{E} [x_i^2 \|\mathbf{x}\|_2^2] \\ &= (p+2) \text{Tr}(\mathbf{M}) + 2M_{ii} (\mathbb{E} x_i^4) + 2 \sum_{j \neq i} M_{jj} (\mathbb{E} x_i^2) (\mathbb{E} x_j^2) + 2M_{ii} \left[\mathbb{E} (x_i^4) + \sum_{j \neq i} (\mathbb{E} x_i^2) (\mathbb{E} x_j^2) \right] \\ &= (p+2) \text{Tr}(\mathbf{M}) + 6M_{ii} + 2 \sum_{j \neq i} M_{jj} + 2M_{ii} (3+p-1) \\ &= (p+4) \text{Tr}(\mathbf{M}) + 2(p+4)M_{ii}. \end{aligned}$$

Then we conclude

$$\begin{aligned} \mathbb{E} \text{Tr} [\nabla_{\mathbf{x}} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})] &= p(p+2) \text{Tr}(\mathbf{M}) + 2(p+4) \sum_i M_{ii} + 2p \text{Tr}(\mathbf{M}) \\ &= (p+2)(p+4) \text{Tr}(\mathbf{M}). \end{aligned}$$

- **Proof of (31).** Invoking the Stein's lemma, we have

$$\mathbb{E} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M} \mathbf{x})^2 = \sum_i \frac{d}{dx_i} [x_i (\mathbf{x}^\top \mathbf{M} \mathbf{x})^2] = p (\mathbf{x}^\top \mathbf{M} \mathbf{x})^2 + 4 \sum_i x_i (\mathbf{x}^\top \mathbf{M} \mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle.$$

The proof is then completed by invoking Lemma 2.

- **Proof of (32).** Following the same strategy as in proving (31), we consider the i th gradient w.r.t x_i , which can be written as

$$\begin{aligned}
\mathbb{E}\|\mathbf{x}\|_2^4 (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 &= \sum_i \frac{d}{dx_i} \left[x_i \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 \right] \\
&= \sum_i \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 + 2 \sum_i x_i^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 + 4 \sum_i x_i \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}^{(\text{sym})}, \mathbf{x} \rangle \\
&= \sum_i \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 + 2 \sum_i \mathbb{E} \frac{d}{dx_i} \left[x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 \right] + 4 \sum_i \mathbb{E} \frac{d}{dx_i} \left[\|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle \right].
\end{aligned} \tag{34}$$

Noticing the following relations

$$\frac{d}{dx_i} \left[x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 \right] = (\mathbf{x}^\top \mathbf{M}\mathbf{x})^2 + 4x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle; \tag{35}$$

$$\begin{aligned}
\frac{d}{dx_i} \left[\|\mathbf{x}\|_2^2 \text{Tr}(\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle \right] &= 2x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle \\
&\quad + 2\|\mathbf{x}\|_2^2 \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle^2 + M_{ii} \|\mathbf{x}\|_2^2 (\mathbf{x}^\top \mathbf{M}\mathbf{x}),
\end{aligned} \tag{36}$$

we can conclude the proof by combining (31), (34), (35), (36), and Lemma 2.

- **Proof of (33).** Due to the independence between \mathbf{x} and \mathbf{y} , we first condition on \mathbf{x} and have

$$\begin{aligned}
&\mathbb{E}(\mathbf{x}^\top \mathbf{y})^2 \mathbf{y}^\top \mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2 \mathbf{y} = \mathbb{E}_{\mathbf{x}} \mathbb{E}_{\mathbf{y}} \mathbf{y}^\top \mathbf{x} \mathbf{x}^\top \mathbf{y} \mathbf{y}^\top \mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2 \mathbf{y} \\
&\stackrel{\textcircled{1}}{=} \mathbb{E}_{\mathbf{x}} \text{Tr}(\mathbf{x} \mathbf{x}^\top) \text{Tr}(\mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2) + \mathbb{E}_{\mathbf{x}} \text{Tr}(\mathbf{x} \mathbf{x}^\top \mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2) + \mathbb{E}_{\mathbf{x}} \text{Tr}(\mathbf{x} \mathbf{x}^\top \mathbf{M}_2^\top \mathbf{x} \mathbf{x} \mathbf{M}_1^\top) \\
&= \mathbb{E}_{\mathbf{x}} \|\mathbf{x}\|_2^2 \mathbf{x}^\top \mathbf{M}_2 \mathbf{M}_1 \mathbf{x} + \mathbb{E}_{\mathbf{x}} \mathbf{x}^\top \mathbf{M}_1 \mathbf{x} \mathbf{x}^\top \mathbf{M}_2 \mathbf{x} + \mathbb{E}_{\mathbf{x}} \mathbf{x}^\top \mathbf{M}_1^\top \mathbf{x} \mathbf{x}^\top \mathbf{M}_2^\top \mathbf{x} \\
&\stackrel{\textcircled{2}}{=} 2 \text{Tr}(\mathbf{M}_1) \text{Tr}(\mathbf{M}_2) + (p+4) \text{Tr}(\mathbf{M}_1 \mathbf{M}_2) + 2 \text{Tr}(\mathbf{M}_1 \mathbf{M}_2^\top),
\end{aligned}$$

where in $\textcircled{1}$ and $\textcircled{2}$ we both use Lemma 18. □

Lemma 2. For a fixed matrix $\mathbf{M} \in \mathbb{R}^{p \times p}$, we associate it with a symmetric matrix $\mathbf{M}^{(\text{sym})}$ defined as $\mathbf{M} + \mathbf{M}^\top / 2$. Consider the Gaussian distributed random vector $\mathbf{x} \sim \mathcal{N}(\mathbf{0}, \mathbf{I})$, we have

$$\mathbb{E} \sum_i x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle = (\text{Tr}(\mathbf{M}))^2 + \|\mathbf{M}\|_{\text{F}}^2 + \text{Tr}(\mathbf{M}\mathbf{M})$$

Proof. This lemma is a direct application of Wick's theorem, which is completed by showing

$$\begin{aligned}
&\mathbb{E} x_i (\mathbf{x}^\top \mathbf{M}\mathbf{x}) \langle \mathbf{M}_i^{(\text{sym})}, \mathbf{x} \rangle = \mathbb{E} \sum_j \sum_{\ell_1, \ell_2} M_{i,j}^{(\text{sym})} M_{\ell_1, \ell_2} x_i x_j x_{\ell_1} x_{\ell_2} \\
&= \underbrace{\mathbb{E} \sum_j \sum_{\ell_1, \ell_2} \mathbb{1}_{\ell_1=i} \mathbb{1}_{\ell_2=j} M_{i,j}^{(\text{sym})} M_{\ell_1, \ell_2} x_i x_j x_{\ell_1} x_{\ell_2}}_{\sum_j M_{i,j}^{(\text{sym})} M_{i,j}} + \underbrace{\mathbb{E} \sum_j \sum_{\ell_1, \ell_2} \mathbb{1}_{\ell_2=i} \mathbb{1}_{\ell_1=j} M_{i,j}^{(\text{sym})} M_{\ell_1, \ell_2} x_i x_j x_{\ell_1} x_{\ell_2}}_{\sum_j M_{i,j}^{(\text{sym})} M_{j,i}} \\
&\quad + \underbrace{\mathbb{E} \sum_j \sum_{\ell_1, \ell_2} \mathbb{1}_{i=j} \mathbb{1}_{\ell_1=\ell_2} M_{i,j}^{(\text{sym})} M_{\ell_1, \ell_2} x_i x_j x_{\ell_1} x_{\ell_2}}_{\sum_{\ell} M_{i,i}^{(\text{sym})} M_{\ell, \ell}} \\
&= \sum_j 2 \left[M_{i,j}^{(\text{sym})} \right]^2 + M_{i,i} \text{Tr}(\mathbf{M}),
\end{aligned}$$

where $\mathbf{M}^{(\text{sym})}$ is defined as $(\mathbf{M} + \mathbf{M}^\top) / 2$. □

Then we study the properties of Σ , which is defined as $\mathbf{X}^\top \Pi^h \mathbf{X} - \Delta$.

Lemma 3. For a fixed matrix \mathbf{M} , we have

$$\mathbb{E} \text{Tr} \left(\Sigma \mathbf{M} \Sigma^\top \right) = n^2 \left[\frac{p}{n} + \left(1 - \frac{h}{n} \right)^2 + o(1) \right] \text{Tr}(\mathbf{M}),$$

where matrix Σ is defined in (18).

Proof. We conclude the proof by showing

$$\begin{aligned} \mathbb{E} \text{Tr} \left(\Sigma \mathbf{M} \Sigma^\top \right) &\stackrel{\textcircled{1}}{=} \sum_{\ell_1, \ell_2 \in \mathcal{S}} \mathbb{E} \text{Tr} \left[\mathbf{X}_{\ell_1} \mathbf{X}_{\ell_1}^\top \mathbf{M} \mathbf{X}_{\ell_2} \mathbf{X}_{\ell_2}^\top \right] + \sum_{\ell_1, \ell_2 \in \mathcal{D}} \mathbb{E} \text{Tr} \left[\mathbf{X}_{\ell_1} \mathbf{X}_{\pi^h(\ell_1)}^\top \mathbf{M} \mathbf{X}_{\pi^h(\ell_2)} \mathbf{X}_{\ell_2}^\top \right] \\ &= \sum_{\ell \in \mathcal{S}} \mathbb{E} \text{Tr} \left(\mathbf{X}_\ell \mathbf{X}_\ell^\top \mathbf{M} \mathbf{X}_\ell \mathbf{X}_\ell^\top \right) + \sum_{\ell_1, \ell_2 \in \mathcal{S}, \ell_1 \neq \ell_2} \mathbb{E} \text{Tr} \left[\mathbf{X}_{\ell_1} \mathbf{X}_{\ell_1}^\top \mathbf{M} \mathbf{X}_{\ell_2} \mathbf{X}_{\ell_2}^\top \right] \\ &\quad + \sum_{\ell \in \mathcal{D}} \underbrace{\mathbb{E} \text{Tr} \left[\mathbf{X}_\ell \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \mathbf{X}_{\pi^h(\ell)} \mathbf{X}_\ell^\top \right]}_{p \text{Tr}(\mathbf{M})} + \sum_{(\ell_1, \ell_2) \in \mathcal{D}_{\text{pair}}} \underbrace{\mathbb{E} \text{Tr} \left[\mathbf{X}_{\ell_1} \mathbf{X}_{\ell_2}^\top \mathbf{M} \mathbf{X}_{\ell_1} \mathbf{X}_{\ell_2}^\top \right]}_{\text{Tr}(\mathbf{M})} \\ &= (n-h)(p+2) \text{Tr}(\mathbf{M}) + (n-h)(n-h-1) \text{Tr}(\mathbf{M}) + hp \text{Tr}(\mathbf{M}) + |\mathcal{D}_{\text{pair}}| \text{Tr}(\mathbf{M}) \\ &\stackrel{\textcircled{2}}{=} n^2 \left[\frac{p}{n} + \left(1 - \frac{h}{n} \right)^2 + o(1) \right] \text{Tr}(\mathbf{M}), \end{aligned}$$

where $\textcircled{1}$ is due to the definitions of index sets \mathcal{S} and \mathcal{D} (Equation (23) and in Equation (24)), and $\textcircled{2}$ is because $|\mathcal{D}_{\text{pair}}| \leq h$. \square

Lemma 4. For a fixed matrix \mathbf{M} , we have

$$\mathbb{E} \text{Tr}(\Sigma \mathbf{M} \Sigma \mathbf{M}) = (n-h + |\mathcal{D}_{\text{pair}}|) [\text{Tr}(\mathbf{M})]^2 + (n-h)^2 \text{Tr}(\mathbf{M} \mathbf{M}) + n \text{Tr}(\mathbf{M} \mathbf{M}^\top)$$

Proof. Following the same strategy as in proving Lemma 3, we complete the proof by showing

$$\begin{aligned} \mathbb{E} \text{Tr}(\Sigma \mathbf{M} \Sigma \mathbf{M}) &= \sum_{\ell = \pi^h(\ell)} \mathbb{E} \text{Tr} \left(\mathbf{X}_\ell^\top \mathbf{M} \mathbf{X}_\ell \mathbf{X}_\ell^\top \mathbf{M} \mathbf{X}_\ell \right) + \sum_{\ell_1, \ell_2 \in \mathcal{S}, \ell_1 \neq \ell_2} \mathbb{E} \text{Tr} \left(\mathbf{X}_{\ell_1} \mathbf{X}_{\ell_1}^\top \mathbf{M} \mathbf{X}_{\ell_2} \mathbf{X}_{\ell_2}^\top \mathbf{M} \right) \\ &\quad + \sum_{\ell \in \mathcal{D}} \mathbb{E} \text{Tr} \left(\mathbf{X}_\ell \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \mathbf{X}_\ell \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \right) + \sum_{\ell \in \mathcal{D}_{\text{pair}}} \mathbb{E} \text{Tr} \left(\mathbf{X}_\ell \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \mathbf{X}_{\pi^h(\ell)} \mathbf{X}_\ell^\top \mathbf{M} \right) \\ &= (n-h + |\mathcal{D}_{\text{pair}}|) [\text{Tr}(\mathbf{M})]^2 + (n-h)^2 \text{Tr}(\mathbf{M} \mathbf{M}) + n \text{Tr}(\mathbf{M} \mathbf{M}^\top). \end{aligned}$$

\square

Lemma 5. For a fixed matrix \mathbf{M} , we have

$$\mathbb{E} [\text{Tr}(\Sigma \mathbf{M})]^2 = (n-h)^2 [\text{Tr}(\mathbf{M})]^2 + n \text{Tr}(\mathbf{M}^\top \mathbf{M}) + (n-h + |\mathcal{D}_{\text{pair}}|) \text{Tr}(\mathbf{M} \mathbf{M}).$$

Proof. We complete the proof by showing

$$\begin{aligned} \mathbb{E} (\text{Tr}(\Sigma \mathbf{M}))^2 &= \sum_{\ell \in \mathcal{S}} \mathbb{E} (\mathbf{X}_\ell^\top \mathbf{M} \mathbf{X}_\ell)^2 + \sum_{\ell_1, \ell_2 \in \mathcal{S}, \ell_1 \neq \ell_2} (\text{Tr}(\mathbf{M}))^2 + \sum_{\ell \in \mathcal{D}} \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \mathbf{X}_\ell \mathbf{X}_\ell^\top \mathbf{M}^\top \mathbf{X}_{\pi^h(\ell)}}_{\text{Tr}(\mathbf{M}^\top \mathbf{M})} \\ &\quad + \sum_{\ell \in \mathcal{D}_{\text{pair}}} \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(\ell)}^\top \mathbf{M} \mathbf{X}_\ell \mathbf{X}_\ell^\top \mathbf{M} \mathbf{X}_{\pi^h(\ell)}}_{\text{Tr}(\mathbf{M} \mathbf{M})} \\ &= (n-h)^2 [\text{Tr}(\mathbf{M})]^2 + n \text{Tr}(\mathbf{M}^\top \mathbf{M}) + (n-h + |\mathcal{D}_{\text{pair}}|) \text{Tr}(\mathbf{M} \mathbf{M}). \end{aligned}$$

\square

Lemma 6. For a fixed matrix \mathbf{M} , we have

$$\mathbb{E} \sum_{\ell=\pi^{\natural}(\ell)} \mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{M} \mathbf{X}_{\ell} \mathbf{X}_{\ell}^{\top} \mathbf{X}_{\pi^{\natural}(i)} = (n-h) \text{Tr}(\mathbf{M}) + (p+1) \mathbb{1}_{i=\pi^{\natural}(i)} \text{Tr}(\mathbf{M}).$$

Proof. Provided that $i = \pi^{\natural}(i)$, we have

$$\begin{aligned} \mathbb{E} \Xi_{1,1} &= \mathbb{E} \mathbf{X}_i^{\top} \mathbf{M} \mathbf{X}_i \mathbf{X}_i^{\top} \mathbf{X}_i + \sum_{\ell \neq i, \ell=\pi^{\natural}(\ell)} \mathbb{E} \mathbf{X}_i^{\top} \mathbf{M} \mathbf{X}_{\ell} \mathbf{X}_{\ell}^{\top} \mathbf{X}_i \\ &= (p+2) \text{Tr}(\mathbf{M}) + \sum_{\ell \neq i, \ell=\pi^{\natural}(\ell)} \text{Tr}(\mathbf{M}) = (n-h+p+1) \text{Tr}(\mathbf{M}) \mathbb{1}_{i=\pi^{\natural}(i)}. \end{aligned} \quad (37)$$

Provided that $i \neq \pi^{\natural}(i)$, we have

$$\mathbb{E} \Xi_{1,1} = \sum_{\ell=\pi^{\natural}(\ell)} \mathbb{E} \mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{M} \mathbf{X}_{\ell} \mathbf{X}_{\ell}^{\top} \mathbf{X}_{\pi^{\natural}(i)} = (n-h) \text{Tr}(\mathbf{M}) \mathbb{1}_{i \neq \pi^{\natural}(i)}. \quad (38)$$

Combining (37) and (38) then completes the proof. \square

Lemma 7. For a fixed \mathbf{M} , we have

$$\mathbb{E} \sum_{\ell} \mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{M} \mathbf{X}_{\pi^{\natural}(\ell)} \mathbf{X}_{\ell}^{\top} \mathbf{X}_j = (p \mathbb{1}_{i=j} \mathbb{1}_{i \neq \pi^{\natural}(i)} + \mathbb{1}_{j=\pi^{\natural}(i)}) \text{Tr}(\mathbf{M}).$$

We omit its proof as it is a direct application of Wick's theorem (Theorem 3).

Lemma 8. We have

$$\begin{aligned} \mathbb{E} \mathbb{1}(i = \pi^{\natural}(i)) &\cong \frac{n-h}{n}; \\ \mathbb{E} \mathbb{1}_{i=j} &\cong \frac{h}{n^2} \\ \mathbb{E} \mathbb{1}_{j=\pi^{\natural}(i)} &\cong \frac{h}{n^2}; \\ \mathbb{E} \mathbb{1}_{i=j} \mathbb{1}_{i=\pi^{\natural}(i)} &\cong \frac{|\mathcal{D}_{\text{pair}}|}{n^2} \end{aligned}$$

This lemma can be easily proved by assuming the indices $i, j, \pi^{\natural}(i)$, and $\pi^{\natural}(j)$ are uniformly sampled from the set $\{1, 2, \dots, n\}$

B.3 MAIN COMPUTATION

In this case, we can write Ξ as

$$\Xi = \underbrace{\mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{B}^{\natural} \mathbf{B}^{\natural\top} \mathbf{X}^{\top} \mathbf{\Pi}^{\natural\top} \mathbf{X} [\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j]}_{\triangleq \Xi_1} + \sigma \underbrace{\mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{B}^{\natural} \mathbf{W}^{\top} \mathbf{X} [\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j]}_{\triangleq \Xi_2} \quad (39)$$

$$+ \sigma \underbrace{\mathbf{W}_i^{\top} \mathbf{B}^{\natural\top} \mathbf{X}^{\top} \mathbf{\Pi}^{\natural\top} \mathbf{X} [\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j]}_{\triangleq \Xi_3} + \sigma^2 \underbrace{\mathbf{W}_i^{\top} \mathbf{W}^{\top} \mathbf{X} [\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j]}_{\triangleq \Xi_4}. \quad (40)$$

The following context separately computes its expectation $\mathbb{E} \Xi$ and its variance $\text{Var} \Xi$.

Expectation Easily we can verify that both $\mathbb{E} \Xi_2$ and $\mathbb{E} \Xi_3$ are zero. Then our goal turns to calculating the expectation of $\mathbb{E} \Xi_1$ and $\mathbb{E} \Xi_4$. First, we have

$$\mathbb{E} \Xi_1 = \mathbb{E} \sum_{\ell=\pi^{\natural}(\ell)} \mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{M} \mathbf{X}_{\ell} \mathbf{X}_{\ell}^{\top} \mathbf{X}_{\pi^{\natural}(i)} - \mathbb{E} \sum_{\ell} \mathbf{X}_{\pi^{\natural}(i)}^{\top} \mathbf{M} \mathbf{X}_{\pi^{\natural}(\ell)} \mathbf{X}_{\ell}^{\top} \mathbf{X}_j.$$

With Lemma 6 and Lemma 7, we conclude

$$\begin{aligned}\mathbb{E}\Xi_1 &= (n-h)\text{Tr}(\mathbf{M}) + (p+1)\mathbb{E}\mathbb{1}_{i=\pi^{\natural}(i)}\text{Tr}(\mathbf{M}) - (p\mathbb{E}\mathbb{1}_{i=j} + \mathbb{E}\mathbb{1}_{j=\pi^{\natural 2}(i)})\text{Tr}(\mathbf{M}) \\ &= (n+p-h-hp/n)[1+o(1)]\text{Tr}(\mathbf{M}).\end{aligned}\quad (41)$$

Meanwhile, we have

$$\begin{aligned}\mathbb{E}\Xi_4 &= \mathbb{E}[\mathbf{W}_i^\top \mathbf{W}_1 \cdots \mathbf{W}_i^\top \mathbf{W}_i \cdots \mathbf{W}_i^\top \mathbf{W}_n] \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) \\ &= m\mathbb{E}\mathbf{X}_i^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) = mp(\mathbb{E}\mathbb{1}_{i=\pi^{\natural}(i)} - \mathbb{E}\mathbb{1}_{i=j}) = \frac{mp(n-h)\sigma^2}{n}[1+o(1)].\end{aligned}\quad (42)$$

Combining (41) and (42) then yields

$$\mathbb{E}\Xi \simeq (n+p)(1-h/n)\|\mathbf{B}^{\natural}\|_F^2 + \frac{mp(n-h)\sigma^2}{n}.$$

Variance Then we study the variance of Ξ . With the relation $\text{Var}(\Xi) = \mathbb{E}\Xi^2 - (\mathbb{E}\Xi)^2$, our goal reduces to computing $\mathbb{E}\Xi^2$, which can be written as

$$\mathbb{E}\Xi^2 = \mathbb{E}\Xi_1^2 + \sigma^2\mathbb{E}\Xi_2^2 + \sigma^2\mathbb{E}\Xi_3^2 + \sigma^4\mathbb{E}\Xi_4^2 + 2\sigma^2\mathbb{E}\Xi_1\Xi_4 + 2\sigma^2\mathbb{E}\Xi_2\Xi_3.$$

The following context separately computes each terms

$$\begin{aligned}\mathbb{E}\Xi_1^2 &\approx (n-h)^2 \left(1 + \frac{2p}{n} + \frac{p^2}{n(n-h)}\right) [\text{Tr}(\mathbf{M})]^2 \\ &\quad + n^2 \left[\frac{2p}{n} + 3\left(1 - \frac{h}{n}\right)^2 + \frac{6(n-h)^2p}{n^3} + \frac{(3n-h)p^2}{n^3}\right] \text{Tr}(\mathbf{M}\mathbf{M}); \\ \mathbb{E}\Xi_2^2 &\approx 2np(1+p/n)\text{Tr}(\mathbf{M}); \\ \mathbb{E}\Xi_3^2 &\approx 2n^2 \left(\frac{p}{n} + \left(1 - \frac{h}{n}\right)^2 + \frac{p^2}{n^2} + \frac{4p(n-h)^2}{n^3}\right) \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Xi_4^2 &\approx \frac{(n-h)m^2p^2}{n}; \\ \mathbb{E}\Xi_1\Xi_4 &\approx \frac{mp(n-h)(n+p-h)}{n}\text{Tr}(\mathbf{M}); \\ \mathbb{E}\Xi_2\Xi_3 &\approx \frac{p(n-h)(n+p-h)}{n}\text{Tr}(\mathbf{M}).\end{aligned}$$

The detailed computation is attached as follows.

Lemma 9. *We have*

$$\begin{aligned}\mathbb{E}\Xi_1^2 &= (n-h)^2 \left(1 + \frac{2p}{n} + \frac{p^2}{n(n-h)} + o(1)\right) [\text{Tr}(\mathbf{M})]^2 \\ &\quad + n^2 \left[\frac{2p}{n} + 3\left(1 - \frac{h}{n}\right)^2 + \frac{6(n-h)^2p}{n^3} + \frac{(3n-h)p^2}{n^3} + o(1)\right] \text{Tr}(\mathbf{M}\mathbf{M}),\end{aligned}$$

where Ξ_1 is defined in (39).

Proof. We begin the proof by decomposing Ξ_1^2 as

$$\begin{aligned}\mathbb{E}\Xi_1^2 &= \mathbb{E} \underbrace{(\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \Sigma^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)}_{\Lambda_1} \\ &\quad + 2\mathbb{E} \underbrace{(\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \Delta^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)}_{\Lambda_2}\end{aligned}$$

$$+ \underbrace{\mathbb{E} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \boldsymbol{\Delta} \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)}_{\Lambda_3},$$

and separately bound each term as in Lemma 10, Lemma 11, and Lemma 12. \square

Lemma 10. *We have*

$$\begin{aligned} & \mathbb{E} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \boldsymbol{\Sigma} \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Sigma}^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) \\ &= (n-h)^2 (1+o(1)) [\text{Tr}(\mathbf{M})]^2 + n^2 \left[\frac{2p}{n} + 3 \left(1 - \frac{h}{n}\right)^2 + o(1) \right] \text{Tr}(\mathbf{M}\mathbf{M}). \end{aligned}$$

Proof. Due to the independence among different rows of the sensing matrix \mathbf{X} , we condition on $\boldsymbol{\Sigma}$ and take expectation w.r.t. $\mathbf{X}_{\pi^h(i)}$ and \mathbf{X}_j , which leads to

$$\mathbb{E} \Lambda_1 = \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \boldsymbol{\Sigma} \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Sigma}^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{1,1}} + \underbrace{\mathbb{E} \mathbf{X}_j^\top \boldsymbol{\Sigma} \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Sigma}^\top \mathbf{X}_j}_{\Lambda_{1,2}}.$$

For $\mathbb{E} \Lambda_{1,1}$, we obtain

$$\begin{aligned} \mathbb{E} \Lambda_{1,1} &\stackrel{\textcircled{1}}{=} \mathbb{E} [\text{Tr}(\boldsymbol{\Sigma} \mathbf{M}) \text{Tr}(\boldsymbol{\Sigma} \mathbf{M})] + \mathbb{E} \text{Tr}(\boldsymbol{\Sigma} \mathbf{M} \mathbf{M}^\top \boldsymbol{\Sigma}^\top) + \mathbb{E} \text{Tr}(\boldsymbol{\Sigma} \mathbf{M} \boldsymbol{\Sigma} \mathbf{M}) \\ &\stackrel{\textcircled{2}}{=} (n-h)^2 [1+o(1)] [\text{Tr}(\mathbf{M})]^2 + n^2 \left[\frac{p}{n} + 2 \left(1 - \frac{h}{n}\right)^2 + o(1) \right] \text{Tr}(\mathbf{M}\mathbf{M}), \end{aligned}$$

where $\textcircled{1}$ is due to (28), and $\textcircled{2}$ is due to Lemma 3, Lemma 4, and Lemma 5. As for $\mathbb{E} \Lambda_{1,2}$, we have

$$\mathbb{E} \Lambda_{1,2} = \mathbb{E} \text{Tr}(\boldsymbol{\Sigma} \mathbf{M} \mathbf{M}^\top \boldsymbol{\Sigma}^\top) = n^2 \left[\frac{p}{n} + \left(1 - \frac{h}{n}\right)^2 + o(1) \right] \text{Tr}(\mathbf{M}^\top \mathbf{M}),$$

and hence complete the proof. \square

Lemma 11. *We have*

$$\mathbb{E} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \boldsymbol{\Sigma} \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) \simeq \frac{(n-h)^2 p}{n} \left[(\text{Tr}(\mathbf{M}))^2 + 3 \text{Tr}(\mathbf{M}\mathbf{M}) \right].$$

Proof. Similar as above, we first expand Λ_2 as

$$\begin{aligned} \mathbb{E} \Lambda_2 &= (n-h) \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{2,1}} + (n-h) \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top \mathbf{X}_j}_{\Lambda_{2,2}} \\ &\quad - (n-h) \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top \mathbf{X}_j}_{\Lambda_{2,3}} - (n-h) \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \boldsymbol{\Delta}^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{2,4}}. \end{aligned}$$

Case (s, s) : $i = \pi^h(i)$ and $j = \pi^h(j)$. We first compute $\Lambda_{2,1}$ as

$$\begin{aligned} \mathbb{E} \Lambda_{2,1} &= \underbrace{\mathbb{E} \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_{\mathbb{E} \|\mathbf{X}_i\|_2^2 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E} \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_i}_{\mathbb{E} (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} \\ &= (p+5) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right]. \end{aligned}$$

We consider $\Lambda_{2,2}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{2,2} &= \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)}_{\mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}(\mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j)}_{\mathbb{E}\|\mathbf{X}_j\|_2^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} \\
&= (\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) + (p+2) \text{Tr}(\mathbf{M}\mathbf{M}).
\end{aligned}$$

As for $\Lambda_{2,3}$ and $\Lambda_{2,4}$, we can verify that they are both zero, which gives

$$\mathbb{E}\Lambda_2 = (n-h)p[1+o(1)][\text{Tr}(\mathbf{M})]^2 + 3(n-h)p[1+o(1)]\text{Tr}(\mathbf{M}\mathbf{M}). \quad (43)$$

Case (s, d): $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$.

We can compute $\Lambda_{2,1}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i + \underbrace{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)}}_0 \\
&\quad + \underbrace{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_{\pi^{\natural}(i)}}_0 \\
&= (p+4) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right].
\end{aligned}$$

We consider $\Lambda_{2,2}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j}_0 \\
&\quad + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_j}_0 \\
&= \mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2 = (\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}).
\end{aligned}$$

Similarly, we can verify that both $\mathbb{E}\Lambda_{2,3}$ and $\mathbb{E}\Lambda_{2,4}$ are zero and hence have

$$\mathbb{E}\Lambda_2 = (n-h)p[1+o(1)][\text{Tr}(\mathbf{M})]^2 + 2(n-h)p[1+o(1)]\text{Tr}(\mathbf{M}\mathbf{M}). \quad (44)$$

Case (d, s): $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. We compute $\Lambda_{2,1}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \underbrace{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)}}_0 + \underbrace{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)}}_0 \\
&\quad + \underbrace{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)}}_{\mathbb{E}\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}} = (\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}).
\end{aligned}$$

We consider $\Lambda_{2,2}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{2,2} &= \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_j}_0 + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j}_0 \\
&\quad + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_j\|_2^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} = (p+2) \text{Tr}(\mathbf{M}\mathbf{M}).
\end{aligned}$$

As for $\mathbb{E}\Lambda_{2,3}$ and $\Lambda_{2,4}$, we can follow the same strategy and prove they are both zero, which yields

$$\mathbb{E}\Lambda_2 = (n-h)[\text{Tr}(\mathbf{M})]^2 + (n-h)p[1+o(1)]\text{Tr}(\mathbf{M}\mathbf{M}). \quad (45)$$

Case (d, d): $i \neq \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. Contrary to the previous cases, we have $\mathbb{E}\Lambda_{2,1}$ and $\mathbb{E}\Lambda_{2,2}$ to be zero in this case rather than $\mathbb{E}\Lambda_{2,3}$ and $\mathbb{E}\Lambda_{2,4}$.

Hence our focus turns to the calculation of $\mathbb{E}\Lambda_{2,3}$ and that of $\mathbb{E}\Lambda_{2,4}$. For $\Lambda_{2,3}$, we have

$$\begin{aligned}\mathbb{E}\Lambda_{2,3} &= \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_j}_{p \mathbb{1}_{i=j} \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} + \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{h2}(i)} \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} \\ &+ \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_j}_{p \mathbb{1}_{i=j} \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} + \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{h2}(i)} \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} \\ &= 2(p \mathbb{1}_{i=j} + \mathbb{1}_{j=\pi^{h2}(i)}) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right].\end{aligned}$$

Then we turn to the calculation of $\mathbb{E}\Lambda_{2,4}$, which proceeds as

$$\begin{aligned}\mathbb{E}\Lambda_{2,4} &= \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}(i=j) \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{j=\pi^{h2}(i)} \mathbb{E} \left\| \mathbf{X}_{\pi^h(i)} \right\|_2^2 \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{M} \mathbf{X}_{\pi^h(i)}} \\ &+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}(i=j) \mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)}} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{j=\pi^{h2}(i)} \mathbb{E} \left\| \mathbf{X}_{\pi^h(i)} \right\|_2^2 \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{M} \mathbf{X}_{\pi^h(i)}} \\ &= 2 \mathbb{1}_{j=\pi^{h2}(i)} (p+2) \text{Tr}(\mathbf{M}\mathbf{M}) + 2 \mathbb{1}(i=j) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right].\end{aligned}$$

Then we conclude

$$\begin{aligned}\mathbb{E}\Lambda_2 &= -2(n-h) \left[(p+1) \mathbb{1}_{i=j} + \mathbb{1}_{j=\pi^{h2}(i)} \right] \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right] \\ &\quad - 2(n-h)(p+2) \mathbb{1}_{j=\pi^{h2}(i)} \text{Tr}(\mathbf{M}\mathbf{M}).\end{aligned}\tag{46}$$

The proof is thus completed by combining (43), (44), (45), and (46). \square

Lemma 12. *We have*

$$\begin{aligned}&\mathbb{E}(\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \Delta \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \Delta^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) \\ &= \left(1 - \frac{h}{n} + o(1)\right) p^2 [\text{Tr}(\mathbf{M})]^2 + \left(3 - \frac{h}{n} + o(1)\right) p^2 \text{Tr}(\mathbf{M}\mathbf{M}).\end{aligned}$$

Proof. We begin the proof by expanding Λ_3 as

$$\begin{aligned}\mathbb{E}\Lambda_3 &= \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \Delta \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \Delta^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{3,1}} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \Delta \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \Delta^\top \mathbf{X}_j}_{\Lambda_{3,2}} \\ &\quad - \underbrace{\mathbb{E}\mathbf{X}_{\pi^h(i)}^\top \Delta \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \Delta^\top \mathbf{X}_j}_{\Lambda_{3,3}} - \underbrace{\mathbb{E}\mathbf{X}_j^\top \Delta \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \Delta^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{3,4}}.\end{aligned}$$

Case (s, s): $i = \pi^h(i)$ and $j = \pi^h(j)$. First we compute $\Lambda_{3,1}$ as

$$\begin{aligned}\mathbb{E}\Lambda_{3,1} &= \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^4 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} \\ &\quad + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_i}_{\mathbb{E}(\mathbf{X}_j^\top \mathbf{X}_i)^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j} \\ &= (p+4)(p+8) \left[(\text{Tr}(\mathbf{M}))^2 + 2 \text{Tr}(\mathbf{M}\mathbf{M}) \right] + 2 (\text{Tr}(\mathbf{M}))^2 + (p+6) \text{Tr}(\mathbf{M}\mathbf{M}).\end{aligned}$$

We consider $\Lambda_{3,2}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{3,2} &= \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j}_{(p+2)\mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} \\
&+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j}_{(p+2)\mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_j\|_2^4 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} \\
&= (3p+8) \left[(\text{Tr}(\mathbf{M}))^2 + \text{Tr}(\mathbf{M}\mathbf{M}) + \text{Tr}(\mathbf{M}^\top \mathbf{M}) \right] + (p+2)(p+4) \text{Tr}(\mathbf{M}\mathbf{M}).
\end{aligned}$$

In addition, we can verify that $\mathbb{E}\Lambda_{3,3}$ and $\mathbb{E}\Lambda_{3,4}$ are both zero. Hence we conclude

$$\mathbb{E}\Lambda_3 = (p^2 + 15p + 42) [\text{Tr}(\mathbf{M})]^2 + (3p^2 + 37p + 94) \text{Tr}(\mathbf{M}\mathbf{M}). \quad (47)$$

Case (s, d) : $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$.

We can compute $\Lambda_{3,1}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{3,1} &= \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^4 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_i}_0 \\
&+ \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_i}_0 + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_0 \\
&+ \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_i}_{\mathbb{1}_{j=\pi^{\natural 2}(j)} \mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} \\
&+ \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i}_0 + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_i}_{\mathbb{1}_{j=\pi^{\natural 2}(j)} \mathbb{E}(\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} \\
&+ \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_i}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{M} \mathbf{X}_i} \\
&= (p^2 + 10p + 24 + 2\mathbb{1}_{j=\pi^{\natural 2}(j)}) \left[(\text{Tr}(\mathbf{M}))^2 + 2 \text{Tr}(\mathbf{M}\mathbf{M}) \right] + 2(p+2) \text{Tr}(\mathbf{M}\mathbf{M}).
\end{aligned}$$

We consider $\Lambda_{3,2}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{3,2} &= \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_i\|_2^2 (\mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i)^2} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j}_0 \\
&+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j}_0 \\
&+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j}_0 + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_i\|_2^4 \text{Tr}(\mathbf{M}\mathbf{M})} \\
&+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{\natural 2}(j)} \mathbb{E}\|\mathbf{X}_j\|_2^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j}_0 \\
&+ \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{\natural 2}(j)} \mathbb{E}\|\mathbf{X}_j\|_2^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} + \underbrace{\mathbb{E}\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j}_{\mathbb{E}\|\mathbf{X}_j\|_2^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{M} \mathbf{X}_j} \\
&= (p+4) \left[(\text{Tr}(\mathbf{M}))^2 + 2 \text{Tr}(\mathbf{M}\mathbf{M}) \right] + (p+2) (p+1 + 2\mathbb{1}_{j=\pi^{\natural 2}(j)}) \text{Tr}(\mathbf{M}\mathbf{M}).
\end{aligned}$$

As for $\Lambda_{3,3}$ and $\Lambda_{3,4}$, we can prove that they are both zero in this case. Hence we conclude

$$\mathbb{E}\Lambda_3 = (p^2 + 11p + 28 + 2\mathbb{1}_{j=\pi^{\natural 2}(j)}) (\text{Tr}(\mathbf{M}))^2 + [3p^2 + 27p + 62 + 2(p+4)\mathbb{1}_{j=\pi^{\natural 2}(j)}] \text{Tr}(\mathbf{M}\mathbf{M}). \quad (48)$$

$$\begin{aligned}
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^{h+2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^{h+2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{h+3}(i)} \mathbb{E} \left\| \mathbf{X}_{\pi^h(i)} \right\|_{\text{F}}^2 \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{M} \mathbf{X}_{\pi^h(i)}} \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_j}_{\mathbb{1}_{i=\pi^h(j)} \mathbb{E} \left[\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \right]^2} \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h+2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h+2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_j}_{\mathbb{1}_{j=\pi^{h+3}(i)} \mathbb{E} \left[\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \right]^2} \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_j}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_j}_0 \\
& = \left[(p+1) \mathbb{1}_{i=\pi^h(j)} + \mathbb{1}_{j=\pi^{h+3}(i)} \right] \left[\text{Tr}(\mathbf{M}) \right]^2 \\
& + \left[2(p+1) \mathbb{1}_{i=\pi^h(j)} + (p+4) \mathbb{1}_{j=\pi^{h+3}(i)} \right] \text{Tr}(\mathbf{M} \mathbf{M}). \tag{52}
\end{aligned}$$

Then we consider the term $\mathbb{E} \Lambda_{3,4}$, which can be written as

$$\begin{aligned}
\mathbb{E} \Lambda_{3,4} & = \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h+2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{i=\pi^h(j)} \mathbb{E} \left[\mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \right]^2} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^{h+2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^{h+2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{h+2}(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^{h+2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_0
\end{aligned}$$

$$\begin{aligned}
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)}^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_{\mathbb{1}_{j=\pi^{\mathfrak{h}3}(i)} \mathbb{E} \left[\mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right]^2} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}}(j)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_{\mathbb{1}_{i=\pi^{\mathfrak{h}}(j)} p \mathbb{E} \left[\mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right]^2} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}}(j)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}2}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}}(j)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}}(j)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)}^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}2}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_{\mathbb{1}_{j=\pi^{\mathfrak{h}3}(i)} \mathbb{E} \left\| \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right\|_{\mathbb{F}}^2 \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\mathfrak{h}-1}(j)}^\top \mathbf{X}_{\pi^{\mathfrak{h}}(i)}}_0 \\
& = \left[(p+1) \mathbb{1}_{i=\pi^{\mathfrak{h}}(j)} + \mathbb{1}_{j=\pi^{\mathfrak{h}3}(i)} \right] [\text{Tr}(\mathbf{M})]^2 \\
& + \left[2(p+1) \mathbb{1}_{i=\pi^{\mathfrak{h}}(j)} + (p+4) \mathbb{1}_{j=\pi^{\mathfrak{h}3}(i)} \right] \text{Tr}(\mathbf{M}\mathbf{M}). \tag{53}
\end{aligned}$$

Combing (50), (51), (52), and (53) together then yields

$$\mathbb{E} \Xi_3 = \frac{4hp(p+2)}{n^2} [1 + o(1)] [\text{Tr}(\mathbf{M})]^2 + 2p^2 [1 + o(1)] \text{Tr}(\mathbf{M}\mathbf{M}). \tag{54}$$

The proof is thus completed by summarizing the computations thereof. \square

Lemma 13. *We have*

$$\mathbb{E} \Xi_2^2 = 2[(p+2)(p+3) + (n-2)(p+1)] \left\| \mathbf{B}^{\mathfrak{h}} \right\|_{\mathbb{F}}^2 = 2np(1 + p/n + o(1)) \left\| \mathbf{B}^{\mathfrak{h}} \right\|_{\mathbb{F}}^2,$$

where Ξ_2 is defined in (39).

Proof. We have

$$\begin{aligned}
\mathbb{E} \Xi_2^2 & = \mathbb{E} \left[\mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{B}^{\mathfrak{h}} \mathbf{W}^\top \mathbf{X} (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j)^\top \mathbf{X}^\top \mathbf{W} \mathbf{B}^{\mathfrak{h}\top} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right] \\
& = \mathbb{E} \left[\mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{B}^{\mathfrak{h}} \text{Tr} \left[\mathbf{X} (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j)^\top \mathbf{X}^\top \right] \mathbf{B}^{\mathfrak{h}\top} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right] \\
& = \mathbb{E} \left[\left\| \mathbf{X} (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j) \right\|_2^2 \mathbf{X}_{\pi^{\mathfrak{h}}(i)}^\top \mathbf{B}^{\mathfrak{h}} \mathbf{B}^{\mathfrak{h}\top} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right] \\
& = \mathbb{E} \left[\left\| \mathbf{X} (\mathbf{X}_{\pi^{\mathfrak{h}}(i)} - \mathbf{X}_j) \right\|_2^2 \times \left\| \mathbf{B}^{\mathfrak{h}\top} \mathbf{X}_{\pi^{\mathfrak{h}}(i)} \right\|_2^2 \right].
\end{aligned}$$

For the conciseness of notation, we assume $\pi^{\mathfrak{h}}(i) = 1$ and $j = 2$ w.l.o.g. Decomposing the term $\left\| \mathbf{X} (\mathbf{X}_1 - \mathbf{X}_2) \right\|_2^2$ as

$$\|\mathbf{X}(\mathbf{X}_1 - \mathbf{X}_2)\|_F^2 = \underbrace{[\mathbf{X}_1^\top (\mathbf{X}_1 - \mathbf{X}_2)]^2}_{\mathcal{T}_1} + \underbrace{[\mathbf{X}_2^\top (\mathbf{X}_1 - \mathbf{X}_2)]^2}_{\mathcal{T}_2} + \underbrace{\sum_{i=3}^n [\mathbf{X}_i^\top (\mathbf{X}_1 - \mathbf{X}_2)]^2}_{\mathcal{T}_3},$$

we then separately bound the above three terms. For the first term $\mathbb{E}\mathcal{T}_1 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2$, we have

$$\begin{aligned} \mathbb{E}\mathcal{T}_1 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 &= \mathbb{E} \left[\left(\|\mathbf{X}_1\|_2^4 + (\mathbf{X}_1^\top \mathbf{X}_2)^2 \right) \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 \right] \\ &= \underbrace{\mathbb{E} \|\mathbf{X}_1\|_2^4 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2}_{(p+2)(p+4)\|\mathbf{B}^\natural\|_F^2} + \underbrace{\mathbb{E} (\mathbf{X}_1^\top \mathbf{X}_2)^2 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2}_{(p+2)\|\mathbf{B}^\natural\|_F^2} \stackrel{\textcircled{1}}{=} (p+2)(p+5)\|\mathbf{B}^\natural\|_F^2, \end{aligned} \quad (55)$$

where $\textcircled{1}$ is due to (29) and (30).

Similarly, for term $\mathbb{E}\mathcal{T}_2 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2$, we invoke (29) and (30), which gives

$$\begin{aligned} \mathbb{E}\mathcal{T}_2 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 &= \underbrace{\mathbb{E} \left[(\mathbf{X}_1^\top \mathbf{X}_2)^2 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 \right]}_{(p+2)\|\mathbf{B}^\natural\|_F^2} + \underbrace{\mathbb{E} \|\mathbf{X}_2\|_2^4 \times \mathbb{E} \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2}_{p(p+2)\|\mathbf{B}^\natural\|_F^2} \\ &\stackrel{\textcircled{2}}{=} (p+1)(p+2)\|\mathbf{B}^\natural\|_F^2, \end{aligned} \quad (56)$$

where $\textcircled{2}$ is due to (29).

For the last term $\mathbb{E}\mathcal{T}_3 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2$, we exploit the independence among the rows of matrix \mathbf{X} and have

$$\begin{aligned} \mathbb{E}\mathcal{T}_3 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 &= \sum_{i \geq 3} \mathbb{E} \left[(\mathbf{X}_i^\top (\mathbf{X}_1 - \mathbf{X}_2))^2 \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 \right] \\ &= \sum_{i \geq 3} \mathbb{E} \left[\|\mathbf{X}_1 - \mathbf{X}_2\|_2^2 \cdot \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 \right] \\ &= \sum_{i \geq 3} \mathbb{E} \left[\left(\|\mathbf{X}_1\|_2^2 + \|\mathbf{X}_2\|_2^2 \right) \cdot \|\mathbf{B}^{\natural\top} \mathbf{X}_1\|_F^2 \right] \\ &= 2 \sum_{i \geq 3} (p+1) \|\mathbf{B}^\natural\|_F^2 = 2(n-2)(p+1) \|\mathbf{B}^\natural\|_F^2. \end{aligned} \quad (58)$$

The proof is then completed by combining (55), (57), and (58). \square

Lemma 14. *We have*

$$\mathbb{E}\Xi_3^2 = 2n^2 \left[\frac{p}{n} + \left(1 - \frac{h}{n}\right)^2 + \frac{p^2}{n^2} + \frac{4p(n-h)^2}{n^3} + o(1) \right] \text{Tr}(\mathbf{M}),$$

where Ξ_3 is defined in (39).

Proof. To begin with, we decompose the term $\mathbb{E}\Xi_3^2$ as

$$\begin{aligned} \mathbb{E}\Xi_3^2 &= \mathbb{E} \left[\underbrace{(\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)^\top \Sigma \mathbf{M} \Sigma^\top (\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)}_{\triangleq \Lambda_1} \right] + 2\mathbb{E} \left[\underbrace{(\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)^\top \Sigma \mathbf{M} \Delta^\top (\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)}_{\triangleq \Lambda_2} \right] \\ &\quad + \mathbb{E} \left[\underbrace{(\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)^\top \Delta \mathbf{M} \Delta^\top (\mathbf{X}_{\pi^{\natural(i)}} - \mathbf{X}_j)}_{\triangleq \Lambda_3} \right]. \end{aligned} \quad (59)$$

Step I. First we consider $\mathbb{E}\Lambda_1$, which can be written as

$$\mathbb{E}\Lambda_1 = 2\mathbb{E} \text{Tr} \left(\Sigma \mathbf{M} \Sigma^\top \right) \stackrel{\textcircled{1}}{=} 2n^2 \left[\frac{p}{n} + \left(1 - \frac{h}{n}\right)^2 + o(1) \right] \text{Tr}(\mathbf{M}), \quad (60)$$

where ① is due to Lemma 3.

Step II. Then we turn to $\mathbb{E}\Lambda_2$, which can be written as

$$\begin{aligned}\mathbb{E}\Lambda_2 &= (n-h)\mathbb{E}\underbrace{\left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M}\Delta^\top \mathbf{X}_{\pi^{\natural}(i)}\right]}_{\Lambda_{2,1}} + (n-h)\mathbb{E}\underbrace{\left[\mathbf{X}_j^\top \mathbf{M}\Delta^\top \mathbf{X}_j\right]}_{\Lambda_{2,2}} \\ &\quad - (n-h)\mathbb{E}\underbrace{\left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M}\Delta^\top \mathbf{X}_j\right]}_{\Lambda_{2,3}} - (n-h)\mathbb{E}\underbrace{\left[\mathbf{X}_j^\top \mathbf{M}\Delta^\top \mathbf{X}_{\pi^{\natural}(i)}\right]}_{\Lambda_{2,4}}.\end{aligned}$$

Case (s, s): $i = \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. We have

$$\begin{aligned}\mathbb{E}\Lambda_{2,1} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_i = (p+3) \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_j^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_j = (p+3) \text{Tr}(\mathbf{M}).\end{aligned}$$

In addition, we can verify that $\mathbb{E}\Lambda_{2,2}$ and $\Lambda_{2,3}$ are both zero, which suggests that

$$\mathbb{E}\Lambda_2 = 2(n-h)(p+3) \text{Tr}(\mathbf{M}). \quad (61)$$

Case (s, d): $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. We have

$$\begin{aligned}\mathbb{E}\Lambda_{2,1} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{M} \left(\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \right) \mathbf{X}_i = (p+2) \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_j^\top \mathbf{M} \left(\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \right) \mathbf{X}_j = \text{Tr}(\mathbf{M}).\end{aligned}$$

Moreover, we have both $\mathbb{E}\Lambda_{2,3}$ and $\mathbb{E}\Lambda_{2,4}$ be zero, which suggests that

$$\mathbb{E}\Lambda_2 = (n-h)(p+3) \text{Tr}(\mathbf{M}). \quad (62)$$

Case (d, s): $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. We have

$$\begin{aligned}\mathbb{E}\Lambda_{2,1} &= \mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right) = \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j = (p+2) \text{Tr}(\mathbf{M}).\end{aligned}$$

Similar as above, we can verify both $\mathbb{E}\Lambda_{2,3}$ and $\mathbb{E}\Lambda_{2,4}$ are zero, which suggests that

$$\mathbb{E}\Lambda_2 = (n-h)(p+3) \text{Tr}(\mathbf{M}). \quad (63)$$

Case (d, d): $i \neq \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. Different from the above three cases, we have $\mathbb{E}\Lambda_{2,1}$ and $\mathbb{E}\Lambda_{2,2}$ be zero and focus on the calculation of $\mathbb{E}\Lambda_{2,3}$ and $\mathbb{E}\Lambda_{2,4}$, which proceeds as

$$\begin{aligned}\mathbb{E}\Lambda_{2,3} &= \underbrace{\mathbb{E} \left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_j \right]}_{p \mathbb{1}_{i=j} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} + \underbrace{\mathbb{E} \left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \right]}_{\mathbb{1}_{j=\pi^{\natural 2}(i)} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} \\ &\quad + \underbrace{\mathbb{E} \left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right]}_{p \mathbb{1}_{i=j} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} + \underbrace{\mathbb{E} \left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j \right]}_{\mathbb{1}_{\pi^{\natural}(i)=\pi^{\natural-1}(j)} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} \\ &= 2 [p \mathbb{1}_{i=j} + \mathbb{1}_{j=\pi^{\natural 2}(i)}] \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Lambda_{2,4} &= \underbrace{\mathbb{E} \left[\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right]}_{\mathbb{1}_{i=j} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} + \underbrace{\mathbb{E} \left[\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right]}_{p \mathbb{1}_{j=\pi^{\natural 2}(i)} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} \\ &\quad + \underbrace{\mathbb{E} \left[\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right]}_{\mathbb{1}_{i=j} \|\mathbf{B}^{\natural}\|_{\text{F}}^2} + \underbrace{\mathbb{E} \left[\mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right]}_{p \mathbb{1}_{j=\pi^{\natural 2}(i)} \|\mathbf{B}^{\natural}\|_{\text{F}}^2}\end{aligned}$$

$$= 2 [p\mathbb{1}_{j=\pi^{h^2}(i)} + \mathbb{1}_{i=j}] \text{Tr}(\mathbf{M}),$$

which suggests that

$$\mathbb{E}\Lambda_2 = -2(n-h)(p+1) (\mathbb{1}_{j=\pi^{h^2}(i)} + \mathbb{1}_{i=j}) \text{Tr}(\mathbf{M}). \quad (64)$$

Combing (61), (62), (63), and (64), we conclude

$$\mathbb{E}\Lambda_2 = \frac{2p(n-h)^2}{n} \text{Tr}(\mathbf{M}) [1 + o(1)]. \quad (65)$$

Step III. Then we turn to the calculation of $\mathbb{E}\Lambda_3$. First we perform the following decomposition

$$\Lambda_3 = \underbrace{\mathbf{X}_{\pi^h(i)}^\top \Delta \mathbf{M} \Delta^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{3,1}} + \underbrace{\mathbf{X}_j^\top \Delta \mathbf{M} \Delta^\top \mathbf{X}_j}_{\Lambda_{3,2}} - \underbrace{\mathbf{X}_{\pi^h(i)}^\top \Delta \mathbf{M} \Delta^\top \mathbf{X}_j}_{\Lambda_{3,3}} - \underbrace{\mathbf{X}_j^\top \Delta \mathbf{M} \Delta^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{3,4}}.$$

Case (s, s): $i = \pi^h(i)$ and $j = \pi^h(j)$. We have

$$\begin{aligned} \mathbb{E}\Lambda_{3,1} &= \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i)}_{\mathbb{E}\|\mathbf{X}_i\|_2^4 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_i)}_{(p+2)\|\mathbf{B}^h\|_F^2} \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i)}_{(p+2)\|\mathbf{B}^h\|_F^2} + \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_i)}_{(p+2)\|\mathbf{B}^h\|_F^2} = (p+2)(p+7) \text{Tr}(\mathbf{M}); \end{aligned}$$

$$\mathbb{E}\Lambda_{3,2} = \mathbb{E}\mathbf{X}_j^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_j = (p+2)(p+7) \text{Tr}(\mathbf{M}).$$

As for $\mathbb{E}\Lambda_{3,3}$ and $\mathbb{E}\Lambda_{3,4}$, easily we can verify that they are both zero and hence have

$$\mathbb{E}\Lambda_3 = 2(p+2)(p+7) \text{Tr}(\mathbf{M}) = 2p^2 \text{Tr}(\mathbf{M}) [1 + o(1)]. \quad (66)$$

Case (s, d): $i = \pi^h(i)$ and $j \neq \pi^h(j)$. We can write $\Lambda_{3,1}$ as

$$\begin{aligned} \mathbb{E}\Lambda_{3,1} &= \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i)}_{\mathbb{E}\|\mathbf{X}_i\|_2^4 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_i)}_0 \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_i)}_0 \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i)}_0 + \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_i)}_{p\|\mathbf{B}^h\|_F^2} \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_i)}_{\mathbb{1}_{j=\pi^{h^2}(j)} \text{Tr}(\mathbf{M})} \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i)}_0 + \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_i)}_{\mathbb{1}_{j=\pi^{h^2}(j)} \text{Tr}(\mathbf{M})} \\ &+ \underbrace{\mathbb{E}(\mathbf{X}_i^\top \mathbf{X}_{\pi^{h-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{h-1}(j)}^\top \mathbf{X}_i)}_{p\|\mathbf{B}^h\|_F^2} \\ &= (p^2 + 8p + 8 + 2\mathbb{1}_{j=\pi^{h^2}(j)}) \text{Tr}(\mathbf{M}). \end{aligned}$$

Mean $\Lambda_{3,2}$ can be written as

$$\mathbb{E}\Lambda_{3,2} = \underbrace{\mathbb{E}(\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j)}_{(p+2)\|\mathbf{B}^h\|_F^2} + \underbrace{\mathbb{E}(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_j)}_0$$

$$\begin{aligned}
& + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j \right)}_0 \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j \right)}_0 + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right)}_{\mathbb{E} \|\mathbf{X}_j\|_{\mathbb{F}}^4 \text{Tr}(\mathbf{M})} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j \right)}_{\mathbb{1}_{\pi^{\natural-1}(j)=\pi^{\natural}(j)}(p+2) \|\mathbf{B}^{\natural}\|_{\mathbb{F}}^2} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j \right)}_0 + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right)}_{\mathbb{1}_{\pi^{\natural-1}(j)=\pi^{\natural}(j)}(p+2) \|\mathbf{B}^{\natural}\|_{\mathbb{F}}^2} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j \right)}_{(p+2) \|\mathbf{B}^{\natural}\|_{\mathbb{F}}^2} \\
& = (p+2) (p+2 + 2 \mathbb{1}_{j=\pi^{\natural 2}(j)}) \text{Tr}(\mathbf{M}).
\end{aligned}$$

And for $\mathbb{E}\Lambda_{3,3}$ and $\mathbb{E}\Lambda_{3,4}$, easily we can verify that they are both zero. Then we conclude

$$\mathbb{E}\Lambda_3 = 2(p^2 + 6p + 6 + (p+3) \mathbb{1}_{j=\pi^{\natural 2}(j)}) \text{Tr}(\mathbf{M}) = 2p^2 \text{Tr}(\mathbf{M}) [1 + o(1)]. \quad (67)$$

Case (d, s) : $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$.

In this case, we can write $\Lambda_{3,1}$ as

$$\begin{aligned}
\mathbb{E}\Lambda_{3,1} & = \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=\pi^{\natural 2}(i)} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=\pi^{\natural 2}(i)} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^4 \text{Tr}(\mathbf{M})} \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 \\
& + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\
& = (p+2) (p+2 + 2 \mathbb{1}_{i=\pi^{\natural 2}(i)}) \text{Tr}(\mathbf{M}).
\end{aligned}$$

$$\begin{aligned}
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{i=\pi^h(j)} \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^h-1(j)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h-2(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_0 + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h-2(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h-2(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h-2(i)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_0 + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)} \mathbf{X}_{\pi^h-2(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^h-1(j)}^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{j=\pi^h-3(i)} \text{Tr}(\mathbf{M})} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_{p \mathbb{1}_{i=\pi^h(j)} \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h-2(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_0 + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_j \mathbf{X}_{\pi^h(j)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^h-1(j)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h-1(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^h(i)}}_0 + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h-1(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h-2(i)} \mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_{\pi^h(i)}}_{p \mathbb{1}_{j=\pi^h-3(i)} \text{Tr}(\mathbf{M})} \\
& + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h-1(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_{\pi^h(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^h(i)}}_0 + \underbrace{\mathbb{E} \mathbf{X}_j^\top \mathbf{X}_{\pi^h-1(j)} \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^h-1(j)}^\top \mathbf{X}_{\pi^h(i)}}_0 \\
& = (p+1) (\mathbb{1}_{i=\pi^h(j)} + \mathbb{1}_{j=\pi^h-3(i)}) \text{Tr}(\mathbf{M}).
\end{aligned}$$

In summary, we have

$$\begin{aligned}
\mathbb{E} \Lambda_3 & = 2(p^2 + 5p + 2) \text{Tr}(\mathbf{M}) + 2(p+3) [\mathbb{1}_{i=\pi^h-2(i)} + \mathbb{1}_{j=\pi^h-2(j)}] \text{Tr}(\mathbf{M}) \\
& + (3p^2 + 7p + 6) (\mathbb{1}_{i=j} + \mathbb{1}_{j=\pi^h-2(i)}) \text{Tr}(\mathbf{M}) + \mathbb{1}_{i=j} \mathbb{1}_{i=\pi^h-2(i)} 4(3p+5) \text{Tr}(\mathbf{M}) \\
& - 2(p+1) [\mathbb{1}_{i=\pi^h(j)} + \mathbb{1}_{j=\pi^h-3(i)}] \text{Tr}(\mathbf{M}) = 2p^2 \text{Tr}(\mathbf{M}) [1 + o(1)].
\end{aligned} \tag{69}$$

Combining (66), (67), (68), and (69) then yields

$$\mathbb{E} \Lambda_3 = 2p^2 \text{Tr}(\mathbf{M}) [1 + o(1)]. \tag{70}$$

The proof is then completed by (59), (60), (65), and (70). \square

Lemma 15. *We have*

$$\begin{aligned}
\mathbb{E} \Xi_4^2 & = m(m+1) [p(p+2) (\mathbb{1}_{i=\pi^h(i)} + \mathbb{1}_{i=j}) + p (\mathbb{1}_{i \neq \pi^h(i)} + \mathbb{1}_{i \neq j})] + 2mp(n+p+1) \\
& = \frac{(n-h)m^2 p^2}{n} [1 + o(1)],
\end{aligned}$$

where Ξ_4 is defined in (39).

Proof. For the conciseness of notation, we define $\mathbf{\Gamma}$ as $\mathbf{X} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \mathbf{X}^\top$ and hence have

$$\mathbb{E} \Xi_4^2 = \mathbb{E} \mathbf{W}_i^\top \mathbf{W}^\top \mathbf{\Gamma} \mathbf{W} \mathbf{W}_i.$$

We begin the discussion by expanding $\mathbf{W} \mathbf{W}_i$ as

$$\begin{bmatrix} \mathbf{W}_1^\top \\ \mathbf{W}_2^\top \\ \dots \\ \mathbf{W}_n^\top \end{bmatrix} \mathbf{W}_i = \begin{bmatrix} \mathbf{W}_1^\top \mathbf{W}_i \\ \mathbf{W}_2^\top \mathbf{W}_i \\ \dots \\ \mathbf{W}_n^\top \mathbf{W}_i \end{bmatrix}.$$

Then we obtain

$$\mathbb{E} \Xi_4^2 = \sum_{s=1}^n \sum_{t=1}^n \Gamma_{ij} \mathbb{E} [(\mathbf{W}_s^\top \mathbf{W}_i) (\mathbf{W}_t^\top \mathbf{W}_i)]$$

$$\begin{aligned}
&= \Gamma_{ii} \mathbb{E} (\mathbf{W}_i^\top \mathbf{W}_i)^2 + \underbrace{\sum_{s \neq i} \sum_{t \neq i} \Gamma_{st} \mathbb{E} (\mathbf{W}_s^\top \mathbf{W}_i \mathbf{W}_t^\top \mathbf{W}_i)}_{\sum_{s \neq i} \Gamma_{ss} \mathbb{E} (\mathbf{W}_s^\top \mathbf{W}_i)^2} \\
&= \Gamma_{ii} \mathbb{E} \left(\sum_{j=1}^m W_{ij}^2 \right)^2 + \sum_{s \neq i} \Gamma_{ss} \cdot m = \Gamma_{ii} [3m + m(m-1)] + m \sum_{s \neq i} \Gamma_{ss} \\
&= m(m+1) \mathbb{E} \Gamma_{ii} + m \mathbb{E} \text{Tr}(\mathbf{\Gamma}). \tag{71}
\end{aligned}$$

We can thus complete the proof by separately computing $\mathbb{E} \text{Tr}(\mathbf{M})$ and $\mathbb{E} \Gamma_{ii}$. First we compute $\mathbb{E} \Gamma_{ii}$, which proceeds as

$$\begin{aligned}
\mathbb{E} \Gamma_{ii} &= \mathbb{E} (\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)})^2 + \mathbb{E} (\mathbf{X}_i^\top \mathbf{X}_j)^2 \\
&= \mathbb{1}_{i=\pi^{\natural}(i)} p(p+2) + \mathbb{1}_{i \neq \pi^{\natural}(i)} p + \mathbb{1}_{i=j} p(p+2) + \mathbb{1}_{i \neq j} p \\
&= p(p+2) [\mathbb{1}_{i=\pi^{\natural}(i)} + \mathbb{1}_{i=j}] + p [\mathbb{1}_{i \neq \pi^{\natural}(i)} + \mathbb{1}_{i \neq j}]. \tag{72}
\end{aligned}$$

Then we turn to the computation of $\mathbb{E} \text{Tr}(\mathbf{M})$, which proceeds as

$$\begin{aligned}
\mathbb{E} \text{Tr}(\mathbf{\Gamma}) &= \|\mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)\|_{\text{F}}^2 \\
&= \mathbb{E} \left\| \mathbf{X}_{\pi^{\natural}(i)}^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) \right\|_2^2 + \mathbb{E} \left\| \mathbf{X}_j^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) \right\|_2^2 + \sum_{s \neq \pi^{\natural}(i), j} \mathbb{E} \left\| \mathbf{X}_s^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) \right\|_2^2 \\
&= 2 \mathbb{E} \left\| \mathbf{X}_{\pi^{\natural}(i)} \right\|_2^4 + 2 \mathbb{E} (\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j)^2 + 2 \sum_{s \neq \pi^{\natural}(i), j} \mathbb{E} \left\| \mathbf{X}_s \right\|_2^2 \\
&= 2p(p+3) + 2(n-2)p = 2p(n+p+1). \tag{73}
\end{aligned}$$

The proof is thus completed by combing (71), (73), and (72). \square

Lemma 16. *We have*

$$\mathbb{E} \Xi_1 \Xi_4 = \frac{mp(n-h)(n+p-h)}{n} [1 + o(1)] \text{Tr}(\mathbf{M}),$$

where Ξ_1 and Ξ_4 are defined in (39).

Proof. We have

$$\mathbb{E} \Xi_1 \Xi_4 = \underbrace{\mathbb{E} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}^\top \mathbf{\Pi}^{\natural} \mathbf{X} (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)^\top \mathbf{X}^\top \mathbf{W} \mathbf{W}_i}_{\triangleq \mathbf{v}^\top}.$$

First we conditional on \mathbf{X} . Expanding the product $\mathbf{W} \mathbf{W}_i$ as

$$\mathbb{E} \begin{bmatrix} \mathbf{W}_1^\top \mathbf{W}_i \\ \mathbf{W}_2^\top \mathbf{W}_i \\ \dots \\ \mathbf{W}_i^\top \mathbf{W}_i \\ \dots \\ \mathbf{W}_n^\top \mathbf{W}_i \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ \dots \\ m \\ \dots \\ 0 \end{bmatrix},$$

we can compute $\mathbb{E} \Xi_1 \Xi_2$ w.r.t. \mathbf{W} as

$$\mathbb{E} (\Xi_1 \Xi_2) = \mathbb{E} \mathbf{v}^\top \mathbf{W} \mathbf{W}_i = m \mathbb{E} \mathbf{v}_i,$$

where \mathbf{v}_i denotes the i th entry of \mathbf{v} and can be written as

$$\mathbf{v}_i = \underbrace{\mathbf{X}_i^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)^\top \mathbf{\Sigma} \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_1} + \underbrace{\mathbf{X}_i^\top (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j) (\mathbf{X}_{\pi^{\natural}(i)} - \mathbf{X}_j)^\top \mathbf{\Delta} \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_2}.$$

For Λ_1 , we conclude

$$\begin{aligned}
\mathbb{E}\Lambda_1 &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} + \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \\
&\quad - \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_j^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} - \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(i)}^\top \Sigma \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \\
&= \mathbb{1}_{i=\pi^{\natural}(i)}(p+3)\mathbb{E}\text{Tr}(\Sigma \mathbf{M}) - \mathbb{1}_{i=j}(p+1)\mathbb{E}\text{Tr}(\Sigma \mathbf{M}) \\
&= (n-h)(\mathbb{1}_{i=\pi^{\natural}(i)}(p+3) - \mathbb{1}_{i=j}(p+1))\text{Tr}(\mathbf{M}) \\
&= \frac{p(n-h)^2}{n} [1 + o(1)] \text{Tr}(\mathbf{M}).
\end{aligned}$$

Then we turn to $\mathbb{E}\Lambda_2$ and obtain

$$\begin{aligned}
\mathbb{E}\Lambda_2 &= \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \Delta \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_{2,1}} + \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top \Delta \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_{2,2}} \\
&\quad - \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_j^\top \Delta \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_{2,3}} - \underbrace{\mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(i)}^\top \Delta \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)}}_{\Lambda_{2,4}}.
\end{aligned}$$

We compute the value of $\mathbb{E}\Lambda_2$ under the four different cases.

Case (s, s) : $i = \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. In this case, we have Δ be

$$\Delta = \Delta^{(s,s)} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top.$$

We have

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = (p+2)(p+5)\text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = 2(p+2)\text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,3} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_j^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = 0; \\
\mathbb{E}\Lambda_{2,4} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_i^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = 0,
\end{aligned}$$

which implies

$$\mathbb{E}\Lambda_2 = (p+2)(p+7)\text{Tr}(\mathbf{M}).$$

Case (s, d) : $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. First we write Δ as

$$\Delta^{(s,d)} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top.$$

Then we conclude

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = (p+2)(p+4)\text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,2} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_j^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = (p+2)\text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,3} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_j^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = 0; \\
\mathbb{E}\Lambda_{2,4} &= \mathbb{E}\mathbf{X}_i^\top \mathbf{X}_j \mathbf{X}_i^\top (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}(j)}^\top + \mathbf{X}_{\pi^{\natural-1}(j)} \mathbf{X}_j^\top) \mathbf{M} \mathbf{X}_i = 0,
\end{aligned}$$

which suggests that

$$\mathbb{E}\Lambda_2 = (p+2)(p+5)\text{Tr}(\mathbf{M}).$$

Case (d, s) : $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. In this case, Δ reduces to

$$\Delta^{(d,s)} = \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural 2}(i)}^\top + \mathbf{X}_j \mathbf{X}_j^\top.$$

We have

$$\mathbb{E}\Xi_2\Xi_3 = \frac{p(n-h)(n+p-h)}{n} \text{Tr}(\mathbf{M}) [1 + o(1)],$$

where Ξ_2 and Ξ_3 are defined in (39).

Proof. To start with, we write the expectation as $\mathbb{E}\Xi_2\Xi_3$

$$\begin{aligned} \mathbb{E}\Xi_2\Xi_3 &= \mathbb{E} \underbrace{\mathbf{X}_{\pi^h(i)}^\top \mathbf{B}^h \mathbf{W}^\top}_{\mathbf{u}^\top} \underbrace{\mathbf{X} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)}_{\mathbf{p} \in \mathbb{R}^{n \times 1}} \underbrace{\mathbf{W}_i^\top \mathbf{B}^{h^\top} \mathbf{X}^\top \mathbf{\Pi}^h \mathbf{X}}_{\mathbf{v}} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) \\ &= \mathbb{E} \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{W}_i^\top \mathbf{v} = \mathbb{E} \langle \mathbf{W}_i, \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v} \rangle. \end{aligned}$$

Exploiting the independence among \mathbf{X} and \mathbf{W} , we condition on \mathbf{X} and have

$$\mathbb{E}_{\mathbf{W}} \langle \mathbf{W}_i, \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v} \rangle = \mathbb{E}_{\mathbf{W}} \text{Tr} (\nabla_{\mathbf{W}_i} \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v}).$$

Notice that only the diagonal entries of the Hessian matrix $\nabla_{\mathbf{W}_i} \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v}$ matters. For an arbitrary index s , we can compute the gradient of the s th entry of $\mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v}$ w.r.t. $\mathbf{W}_{i,s}$ as

$$\frac{d}{dW_{i,s}} (\mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v}_s) = \mathbf{v}_s \frac{d}{dW_{i,s}} (\mathbf{u}^\top \mathbf{W}^\top \mathbf{p}) = \mathbf{v}_s \sum_{t=1}^n \frac{d}{dW_{i,s}} (\mathbf{p}_t \mathbf{W}_t^\top \mathbf{u}) = \mathbf{v}_s \frac{d}{dW_{i,s}} \mathbf{p}_i \mathbf{W}_i^\top \mathbf{u} = \mathbf{p}_i \mathbf{v}_s \mathbf{u}_s.$$

Invoking the definitions of \mathbf{p} , \mathbf{v} and \mathbf{u} , we have

$$\begin{aligned} \mathbb{E}_{\mathbf{W}, \mathbf{X}} \langle \mathbf{W}_i, \mathbf{u}^\top \mathbf{W}^\top \mathbf{p} \mathbf{v} \rangle &= \mathbb{E}_{\mathbf{X}} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \mathbf{X}_i \sum_{s=1}^m \left[\mathbf{X}_{\pi^h(i)}^\top (\mathbf{B}^{h^\top})_s (\mathbf{B}^{h^\top})_s^\top \mathbf{X}^\top \mathbf{\Pi}^h \mathbf{X} (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j) \right] \\ &\stackrel{\textcircled{1}}{=} \mathbb{E} \left[\underbrace{(\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)}_{\Lambda_1} \right] + \mathbb{E} \left[\underbrace{(\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Delta}^\top (\mathbf{X}_{\pi^h(i)} - \mathbf{X}_j)}_{\Lambda_2} \right], \end{aligned}$$

where in $\textcircled{1}$ we use the relation $\sum_{s=1}^m (\mathbf{B}^{h^\top})_s (\mathbf{B}^{h^\top})_s^\top = \mathbf{B}^h \mathbf{B}^{h^\top} = \mathbf{M}$.

For the first term Λ_1 , we obtain

$$\begin{aligned} \mathbb{E}\Lambda_1 &= \mathbb{E} \left(\underbrace{\mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_{\pi^h(i)}}_{\mathbb{1}_{i=\pi^h(i)} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_i} \right) + \mathbb{E} \left(\underbrace{\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_j}_{\mathbb{1}_{i=\pi^h(i)} \mathbb{E} \mathbf{X}_i^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_i} \right) \\ &\quad - \mathbb{E} \left(\underbrace{\mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_j}_{\mathbb{1}_{i=j} \mathbb{1}_{i \neq \pi^h(i)} \mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_{\pi^h(i)}} \right) - \mathbb{E} \left(\underbrace{\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_{\pi^h(i)}}_{p \mathbb{1}_{i=j} \mathbb{1}_{i \neq \pi^h(i)} \mathbb{E} \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Sigma}^\top \mathbf{X}_{\pi^h(i)}} \right) \\ &= (n-h) \left[\mathbb{1}_{i=\pi^h(i)} (p+3) - (p+1) \mathbb{1}_{i=j} \mathbb{1}_{i \neq \pi^h(i)} \right] \text{Tr}(\mathbf{M}). \end{aligned}$$

Then we consider the second term Λ_2 , which can be decomposed further into four sub-terms reading as

$$\begin{aligned} \mathbb{E}\Lambda_2 &= \mathbb{E} \left(\underbrace{\mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Delta}^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{2,1}} \right) + \mathbb{E} \left(\underbrace{\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Delta}^\top \mathbf{X}_j}_{\Lambda_{2,2}} \right) \\ &\quad - \mathbb{E} \left(\underbrace{\mathbf{X}_{\pi^h(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Delta}^\top \mathbf{X}_j}_{\Lambda_{2,3}} \right) - \mathbb{E} \left(\underbrace{\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^h(i)}^\top \mathbf{M} \mathbf{\Delta}^\top \mathbf{X}_{\pi^h(i)}}_{\Lambda_{2,4}} \right); \end{aligned}$$

Case (s, s) : $i = \pi^h(i)$ and $j = \pi^h(j)$. In this case, we have $\mathbf{\Delta}$ be

$$\mathbf{\Delta}^{(s,s)} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top.$$

Hence we conclude

$$\mathbb{E}\Lambda_{2,1} = \mathbb{E} \left[\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_i \right] = \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^4 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i + \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i$$

$$\begin{aligned}
&= (p+2)(p+4) \text{Tr}(\mathbf{M}) + (p+2) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,2} &= \mathbb{E} \left[\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_j \right] = \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i + \mathbb{E} \|\mathbf{X}_j\|_{\mathbb{F}}^2 \mathbf{X}_j^\top \mathbf{M} \mathbf{X}_j \\
&= 2\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i = 2(p+2) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,3} &= \mathbb{E} \left[\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_j^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_j \right] = 0; \\
\mathbb{E}\Lambda_{2,4} &= \mathbb{E} \left[\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} (\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_j \mathbf{X}_j^\top) \mathbf{X}_i \right] = 0,
\end{aligned}$$

which suggests $\mathbb{E}\Lambda_2 = (p+2)(p+7) \text{Tr}(\mathbf{M})$.

Case (s, d): $i = \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. First we write Δ as

$$\Delta^{(s,d)\top} = \mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top.$$

Then we conclude

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \underbrace{\mathbb{E} \left(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_i \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^4 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E} \left(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_i \right)}_0 \\
&\quad + \underbrace{\mathbb{E} \left(\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_i \right)}_0 = \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^4 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i = (p+2)(p+4) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,2} &= \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i \mathbf{X}_i^\top \mathbf{X}_j \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right)}_0 \\
&\quad + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \mathbf{X}_j \right)}_0 = \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i = (p+2) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,3} &= \mathbb{E} \left[\mathbf{X}_i^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \left(\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \right) \mathbf{X}_j \right] = 0; \\
\mathbb{E}\Lambda_{2,4} &= \mathbb{E} \left[\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_i^\top \mathbf{M} \left(\mathbf{X}_i \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural-1}(j)}^\top \right) \mathbf{X}_i \right] = 0,
\end{aligned}$$

which suggests $\mathbb{E}\Lambda_2 = (p+2)(p+5) \text{Tr}(\mathbf{M})$.

Case (d, s): $i \neq \pi^{\natural}(i)$ and $j = \pi^{\natural}(j)$. In this case, Δ reduces to

$$\Delta^{(d,s)\top} = \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_j \mathbf{X}_j^\top.$$

Then we obtain

$$\begin{aligned}
\mathbb{E}\Lambda_{2,1} &= \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=\pi^{\natural 2}(i)} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\
&\quad + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 = (1 + \mathbb{1}_{i=\pi^{\natural 2}(i)}) (p+2) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,2} &= \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=j} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^4 \text{Tr}(\mathbf{M}) + \mathbb{1}_{i \neq j} p \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=\pi^{\natural 2}(i)} [\mathbb{1}_{i=j} \mathbb{E} \|\mathbf{X}_i\|_{\mathbb{F}}^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i + \mathbb{1}_{i \neq j} \text{Tr}(\mathbf{M})]} \\
&\quad + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_j^\top \mathbf{X}_j \right)}_0 = (p + \mathbb{1}_{i=\pi^{\natural 2}(i)}) \text{Tr}(\mathbf{M}); \\
\mathbb{E}\Lambda_{2,3} &= \mathbb{E} \left[\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \left(\mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_j \mathbf{X}_j^\top \right) \mathbf{X}_j \right] = 0; \\
\mathbb{E}\Lambda_{2,4} &= \mathbb{E} \left[\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \left(\mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural 2}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_j \mathbf{X}_j^\top \right) \mathbf{X}_{\pi^{\natural}(i)} \right] = 0,
\end{aligned}$$

which suggests

$$\mathbb{E}\Lambda_2 = 2(p+1) \text{Tr}(\mathbf{M}) + (p+3) \mathbb{1}_{i=\pi^{\natural}(i)} \text{Tr}(\mathbf{M}).$$

Case (d, d): $i \neq \pi^{\natural}(i)$ and $j \neq \pi^{\natural}(j)$. In this case, Δ is written as

$$\Delta^{(d,d)\top} = \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top + \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top + \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top + \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top.$$

Then we have

$$\begin{aligned} \mathbb{E}\Lambda_{2,1} &= \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{E} \|\mathbf{X}_i\|_F^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=\pi^{\natural}(i)} \mathbb{E} \|\mathbf{X}_i\|_F^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\ &+ \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=j} \mathbb{E} \|\mathbf{X}_i\|_F^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=j} \mathbb{1}_{i=\pi^{\natural}(i)} \mathbb{E} \|\mathbf{X}_i\|_F^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\ &= (p+2) [1 + \mathbb{1}_{i=\pi^{\natural}(i)} + \mathbb{1}_{i=j} + \mathbb{1}_{i=\pi^{\natural}(i)} \mathbb{1}_{i=j}] \text{Tr}(\mathbf{M}); \\ \mathbb{E}\Lambda_{2,2} &= \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=j} \mathbb{E} \|\mathbf{X}_i\|_F^4 \text{Tr}(\mathbf{M}) + p \mathbb{1}_{i \neq j} \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=\pi^{\natural}(i)} [\mathbb{1}_{i=j} (p+2) \text{Tr}(\mathbf{M}) + \mathbb{1}_{i \neq j} \text{Tr}(\mathbf{M})]} \\ &+ \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=j} \mathbb{E} \|\mathbf{X}_i\|_F^4 \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_j \right)}_{\mathbb{1}_{i=j} \mathbb{1}_{i=\pi^{\natural}(i)} \mathbb{E} \|\mathbf{X}_i\|_F^2 \mathbf{X}_i^\top \mathbf{M} \mathbf{X}_i} \\ &= 2\mathbb{1}_{i=j} p(p+2) \text{Tr}(\mathbf{M}) + p \mathbb{1}_{i \neq j} \text{Tr}(\mathbf{M}) + \mathbb{1}_{i=\pi^{\natural}(i)} [\mathbb{1}_{i=j} 2(p+2) \text{Tr}(\mathbf{M}) + \mathbb{1}_{i \neq j} \text{Tr}(\mathbf{M})] \\ \mathbb{E}\Lambda_{2,3} &= \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_j \right)}_0 + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_j \right)}_0 \\ &+ \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_j \right)}_{p \mathbb{1}_{i=\pi^{\natural}(j)} \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \left(\mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_j \right)}_0 \\ \mathbb{E}\Lambda_{2,4} &= \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_i^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(i)} \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0 \\ &+ \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_{\pi^{\natural}(j)} \mathbf{X}_j^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_{\mathbb{1}_{i=\pi^{\natural}(j)} \text{Tr}(\mathbf{M})} + \underbrace{\mathbb{E} \left(\mathbf{X}_j^\top \mathbf{X}_i \mathbf{X}_{\pi^{\natural}(i)}^\top \mathbf{M} \mathbf{X}_j \mathbf{X}_{\pi^{\natural}-1(j)}^\top \mathbf{X}_{\pi^{\natural}(i)} \right)}_0, \end{aligned}$$

which gives

$$\mathbb{E}\Lambda_2 = (p+1) [2 + 2(p+1) \mathbb{1}_{i=j} + \mathbb{1}_{i=\pi^{\natural}(j)}] \text{Tr}(\mathbf{M}) + \mathbb{1}_{i=\pi^{\natural}(i)} [p+3 + (3p+5) \mathbb{1}_{i=j}] \text{Tr}(\mathbf{M}).$$

□

C USEFUL FACTS

This section collects some useful facts for the sake of self-containing.

Theorem 3 (Wick's theorem (Theorem 1.28 in Janson (1997))). *Considering the centered jointly normal variables g_1, g_2, \dots, g_n , we conclude*

$$\mathbb{E}(g_1 g_2 \cdots g_n) = \sum_{\substack{\text{all possible disjoint} \\ \text{pairs } (i_k, j_k) \in \{1, 2, \dots, n\}}} \prod_k \mathbb{E}(g_{i_k} g_{j_k}).$$

With Wick's theorem, we can reduce the computation of high-order Gaussian chaos to calculating the expectations of a series of low-order Gaussian chaos.

Lemma 18. For a normally distributed random matrix $\mathbf{G} \in \mathbb{R}^{n \times p}$ which satisfies $\mathbb{E}\mathbf{G} = \mathbf{0}$ and $\mathbb{E}\text{vec}(\mathbf{G})\text{vec}(\mathbf{G})^\top = \mathbf{U} \otimes \mathbf{V}$, we have

$$\begin{aligned} \mathbb{E}(\mathbf{G}^\top \mathbf{A} \mathbf{G} \mathbf{C} \mathbf{G}^\top \mathbf{B} \mathbf{G}) &= \text{Tr}(\mathbf{A} \mathbf{U}) \text{Tr}(\mathbf{B} \mathbf{U}) \mathbf{V} \mathbf{C} \mathbf{V} \\ &+ \text{Tr}(\mathbf{A} \mathbf{U} \mathbf{B}^\top \mathbf{U}) \mathbf{V} \mathbf{C}^\top \mathbf{V} \\ &+ \text{Tr}(\mathbf{A} \mathbf{U} \mathbf{B} \mathbf{U}) \text{Tr}(\mathbf{C} \mathbf{V}) \mathbf{V}, \end{aligned}$$

where $\text{vec}(\cdot)$ is the vector operation; \otimes is the Kronecker product (Horn & Johnson, 1990); and \mathbf{A} , \mathbf{B} and \mathbf{C} are arbitrary fixed matrices.

Lemma 19 (Stein’s Lemma (cf. Section 1.3 in Talagrand (2010))). Let $g \sim \mathcal{N}(0, 1)$. Then for any differentiable function $f : \mathbb{R} \mapsto \mathbb{R}$ we have

$$\mathbb{E}[gf(g)] = \mathbb{E}f'(g),$$

where $\lim_{\|g\| \rightarrow \infty} f(g)e^{-a\|g\|_2^2} = 0$ for any $a > 0$.

REFERENCES

- Zhidong Bai and Tailen Hsing. The broken sample problem. *Probab. Theory Relat. Fields*, 131(4): 528–552, 2005.
- Zhidong Bai and Jack W Silverstein. *Spectral analysis of large dimensional random matrices*, volume 20. Springer, 2010.
- Mohsen Bayati and Andrea Montanari. The dynamics of message passing on dense graphs, with applications to compressed sensing. *IEEE Transactions on Information Theory*, 57(2):764–785, 2011.
- JD464415 Biggins. Chernoff’s theorem in the branching random walk. *Journal of Applied Probability*, 14(3):630–636, 1977.
- Sergio Caracciolo, Matteo P D’Achille, Enrico M Malatesta, and Gabriele Sicuro. Finite-size corrections in the random assignment problem. *Physical Review E*, 95(5):052129, 2017.
- Michael Chertkov, Lukas Kroc, F Krzakala, M Vergassola, and L Zdeborová. Inference in particle tracking experiments by passing messages between images. *Proceedings of the National Academy of Sciences*, 107(17):7663–7668, 2010.
- Sae-Young Chung. *On the construction of some capacity-approaching coding schemes*. PhD thesis, Massachusetts Institute of Technology, 2000.
- Morris H. DeGroot and Prem K. Goel. The matching problem for multivariate normal data. *Sankhyā: The Indian Journal of Statistics, Series B (1960-2002)*, 38(1):14–29, 1976.
- Morris H. DeGroot and Prem K. Goel. Estimation of the correlation coefficient from a broken random sample. *The Annals of Statistics*, 8(2):264–278, 03 1980.
- Morris H. DeGroot, Paul I. Feder, and Prem K. Goel. Matchmaking. *Annals of Mathematical Statistics*, 42(2):578–593, 04 1971.
- David L Donoho, Arian Maleki, and Andrea Montanari. Message-passing algorithms for compressed sensing. *Proceedings of the National Academy of Sciences*, 106(45):18914–18919, 2009.
- Noureddine El Karoui. Asymptotic behavior of unregularized and ridge-regularized high-dimensional robust regression estimators: rigorous results. *arXiv preprint arXiv:1311.2445*, 2013.
- Noureddine El Karoui. On the impact of predictor geometry on the performance on high-dimensional ridge-regularized generalized robust regression estimators. *Probab. Theory Relat. Fields*, 170(1-2): 95–175, 2018.
- Prem K. Goel. On re-pairing observations in a broken random sample. *The Annals of Statistics*, 3(6): 1364–1369, 11 1975.

- John M Hammersley. Postulates for subadditive processes. *The Annals of Probability*, 2(4):652–680, 1974.
- Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge university press, 1990.
- Svante Janson. *Gaussian Hilbert Spaces*. Cambridge Tracts in Mathematics. Cambridge University Press, 1997. doi: 10.1017/CBO9780511526169.
- John Frank Charles Kingman. The first birth problem for an age-dependent branching process. *The Annals of Probability*, pp. 790–801, 1975.
- Frank R Kschischang, Brendan J Frey, and H-A Loeliger. Factor graphs and the sum-product algorithm. *IEEE Transactions on information theory*, 47(2):498–519, 2001.
- Svante Linusson and Johan Wästlund. A proof of parisi’s conjecture on the random assignment problem. *Probability theory and related fields*, 128(3):419–440, 2004.
- David JC MacKay, David JC Mac Kay, et al. *Information theory, inference and learning algorithms*. Cambridge university press, 2003.
- Enrico M Malatesta, Giorgio Parisi, and Gabriele Sicuro. Fluctuations in the random-link matching problem. *Physical Review E*, 100(3):032102, 2019.
- Mohammad Ali Maleki. *Approximate message passing algorithms for compressed sensing*. Stanford University, 2010.
- OC Martin, M Mézard, and Olivier Rivoire. Random multi-index matching problems. *Journal of Statistical Mechanics: Theory and Experiment*, 2005(09):P09006, 2005.
- Marc Mezard and Andrea Montanari. *Information, physics, and computation*. Oxford University Press, 2009.
- Marc Mézard and Giorgio Parisi. Replicas and optimization. *Journal de Physique Lettres*, 46(17):771–778, 1985.
- Marc Mézard and Giorgio Parisi. Mean-field equations for the matching and the travelling salesman problems. *EPL (Europhysics Letters)*, 2(12):913, 1986.
- Marc Mézard and Giorgio Parisi. On the solution of the random link matching problems. *Journal de Physique*, 48(9):1451–1459, 1987.
- Heinz Neudecker and Tom Wansbeek. Fourth-order properties of normally distributed random matrices. *Linear Algebra and Its Applications*, 97:13–21, 1987.
- Ashwin Pananjady, Martin J Wainwright, and Thomas A Courtade. Denoising linear models with permuted data. In *Proceedings of the 2017 IEEE International Symposium on Information Theory (ISIT)*, pp. 446–450, Aachen, Germany, 2017.
- Ashwin Pananjady, Martin J. Wainwright, and Thomas A. Courtade. Linear regression with shuffled data: Statistical and computational limits of permutation recovery. *IEEE Trans. Inf. Theory*, 64(5):3286–3300, 2018.
- Giorgio Parisi and Matthieu Ratiéville. On the finite size corrections to some random matching problems. *The European Physical Journal B-Condensed Matter and Complex Systems*, 29(3):457–468, 2002.
- Sundeep Rangan. Generalized approximate message passing for estimation with random linear mixing. In *Information Theory Proceedings (ISIT), 2011 IEEE International Symposium on*, pp. 2168–2172. IEEE, 2011.
- Thomas J Richardson and Rüdiger L Urbanke. The capacity of low-density parity-check codes under message-passing decoding. *IEEE Transactions on information theory*, 47(2):599–618, 2001.
- Tom Richardson and Ruediger Urbanke. *Modern coding theory*. Cambridge university press, 2008.

- Guilhem Semerjian, Gabriele Sicuro, and Lenka Zdeborová. Recovery thresholds in the sparse planted matching problem. *Physical Review E*, 102(2):022304, 2020.
- Martin Slawski and Emanuel Ben-David. Linear regression with sparsely permuted data. *Electron. J. Statist.*, 13(1):1–36, 2019.
- Martin Slawski, Emanuel Ben-David, and Ping Li. Two-stage approach to multivariate linear regression with sparsely mismatched data. *J. Mach. Learn. Res.*, 21(204):1–42, 2020.
- Pragya Sur, Yuxin Chen, and Emmanuel J Candès. The likelihood ratio test in high-dimensional logistic regression is asymptotically a rescaled chi-square. *Probab. Theory Relat. Fields*, 175(1-2): 487–558, 2019.
- Michel Talagrand. *Mean field models for spin glasses: Volume I: Basic examples*, volume 54. Springer Science & Business Media, 2010.
- Jayakrishnan Unnikrishnan, Saeid Haghighatshoar, and Martin Vetterli. Unlabeled sensing: Solving a linear system with unordered measurements. In *Proceedings of the 53rd Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pp. 786–793, Monticello, IL, 2015.
- Hang Zhang and Ping Li. Optimal estimator for unlabeled linear regression. In *Proceedings of the 37th International Conference on Machine Learning (ICML)*, pp. 11153–11162, Virtual Event, 2020.
- Hang Zhang, Martin Slawski, and Ping Li. The benefits of diversity: Permutation recovery in unlabeled sensing from multiple measurement vectors. *IEEE Trans. Inf. Theory*, 68(4):2509–2529, 2022.