
A Faster Decentralized Algorithm for Nonconvex Minimax Problems

Wenhan Xian, Feihu Huang, Yanfu Zhang, Heng Huang

Electrical and Computer Engineering, University of Pittsburgh, Pittsburgh, PA 15213
wex37@pitt.edu, huangfeihu2018@gmail.com, yaz91@pitt.edu, heng.huang@pitt.edu

Abstract

In this paper, we study the nonconvex-strongly-concave minimax optimization problem on decentralized setting. The minimax problems are attracting increasing attentions because of their popular practical applications such as policy evaluation and adversarial training. As training data become larger, distributed training has been broadly adopted in machine learning tasks. Recent research works show that the decentralized distributed data-parallel training techniques are specially promising, because they can achieve the efficient communications and avoid the bottleneck problem on the central node or the latency of low bandwidth network. However, the decentralized minimax problems were seldom studied in literature and the existing methods suffer from very high gradient complexity. To address this challenge, we propose a new faster decentralized algorithm, named as DM-HSGD, for nonconvex minimax problems by using the variance reduced technique of hybrid stochastic gradient descent. We prove that our DM-HSGD algorithm achieves stochastic first-order oracle (SFO) complexity of $O(\kappa^3\epsilon^{-3})$ for decentralized stochastic nonconvex-strongly-concave problem to search an ϵ -stationary point, which improves the existing best theoretical results. Moreover, we also prove that our algorithm achieves linear speedup with respect to the number of workers. Our experiments on decentralized settings show the superior performance of our new algorithm.

1 Introduction

Minimax optimization has enormous applications in machine learning tasks such as Generative Adversarial Net (GAN) [8], adversarial training [26] and multi-agent reinforcement learning [43]. Specifically, in minimax optimization, variable x aims to minimize a payoff loss function $f(x, y) : \mathbb{R}^{d_1} \times \mathbb{R}^{d_2} \rightarrow \mathbb{R}$ while variable y tries to maximize the loss, which can be formulated as

$$\min_{x \in \mathcal{X}} \max_{y \in \mathcal{Y}} f(x, y), \quad (1)$$

where $\mathcal{X} \subseteq \mathbb{R}^{d_1}$ and $\mathcal{Y} \subseteq \mathbb{R}^{d_2}$. In the past a few decades, there are plenty of works to study minimax optimization problem in a variety of research fields and many methods have been developed. The most intuitive solution is Gradient Descent Ascent (GDA) algorithm [6, 29] with equal stepsize $\eta_x = \eta_y$. Asymptotic and nonasymptotic convergence analysis has been provided when f is convex in x and concave in y . Recently, many deterministic and stochastic gradient algorithms for nonconvex-strongly-concave and nonconvex-concave problems were proposed. Some algorithms improve the performance of vanilla GDA method by adopting different stepsize on x and y , such as [10, 19], where the stepsize of y is typically larger than the stepsize of x . Some algorithms update x and y at different frequency, such as [14, 25, 32]. These kind of algorithms usually involve a nested loop structure that updates y more frequently than x to make $f(x, y)$ close to function $\Phi(x)$, which is defined by

$$\Phi(x) = \max_{y \in \mathcal{Y}} f(x, y). \quad (2)$$

As more large-scale machine learning problems are arising, distributed training becomes a popular and crucial framework because of its ability and efficiency to deal with large data. It is desired to generalize minimax optimization to distributed training to solve large-scale minimax problems. In distributed optimization, the original centralized optimization suffers from a bottleneck communication problem, *i.e.* the communication traffic on the busiest central node, especially when the network is large [18, 51]. To tackle this communication issue, decentralized optimization was proposed and has emerged as a promising technique. It is a kind of distributed machine learning training paradigm that does not rely on the centralized network topology. Different worker nodes collaboratively utilize their own local data to implement large-scale training tasks and at each iteration they only have to communicate with their neighbors. Decentralized algorithms have been shown to enhance the communication efficiency by avoiding the communication overhead problem. Decentralized methods are also advantageous when the network suffers from communication restriction or has low bandwidth between some nodes and the central node. Besides, it is also an essential method in some situations where data are geographically distributed and centralized data processing is not available or there are concerns to preserve data privacy [48].

Recently many works were proposed to improve the performance of decentralized training. D-PSGD [18] theoretically justifies the potential advantage of decentralized algorithm. D^2 [38] improves the convergence rate to outperform D-PSGD by eliminating the influence of data variance among different workers. D-SPIDER-SFO [33] incorporates D^2 and SPIDER [7, 44], which is a kind of variance reduction technique [15], to further reduce the gradient complexity. DQSFWD [45] studies decentralized constrained problem with Frank-Wolfe method. GT-HSGD [46] extends hybrid stochastic gradient descent to decentralized setting, which is a variance-reduced approach that does not compute mega batch periodically. However, the decentralized minimax optimization is still very limited and existing methods suffer from very high gradient complexity [21, 41]. Thus, we are motivated to design an accelerated decentralized algorithm for minimax problems.

In this paper, thus, we propose a faster Decentralized Minimax Hybrid Stochastic Gradient Descent (DM-HSGD) algorithm to solve the following decentralized stochastic minimax optimization problem:

$$\min_{x \in \mathbb{R}^d} \max_{y \in \mathcal{Y}} f(x, y) = \frac{1}{n} \sum_{i=1}^n f_i(x, y), \quad f_i(x, y) := \mathbb{E}_{\xi^{(i)} \sim D_i} F_i(x, y; \xi^{(i)}) \quad (3)$$

where n is the number of worker nodes, \mathcal{Y} is a convex set. Here the local component objective function $F_i(x, y; \xi^{(i)})$ is L -smooth, nonconvex in x , and strongly-concave in y . D_i is the data distribution on the i -th node. In this paper, the data distribution can be non-identical. Random variable $\xi^{(i)}$ is an index sampled from the local data. We summarize our contributions as follows:

- (1) In this paper, we propose a new accelerated decentralized stochastic first-order algorithm, named as DM-HSGD, to solve the decentralized nonconvex-strongly-concave minimax optimization problems. Our algorithm is the first stochastic gradient algorithm to solve general decentralized minimax problem on non-identical distributed data with theoretical guarantees. Besides, our algorithm does not require large batch size or nested loop which makes it more practical and efficient to implement.
- (2) We provide a completed proof to guarantee the convergence of our algorithm to solve decentralized stochastic minimax optimization. Under nonconvex-strongly-concave condition, our algorithm obtains SFO complexity of $O(\kappa^3 \epsilon^{-3})$ to search an ϵ -stationary point of function $\Phi(x) = \max_{y \in \mathcal{Y}} f(x, y)$. This result is faster than the complexity of previous decentralized minimax algorithms [21, 41]. Moreover, we also prove that our method achieves linear speedup as the number of workers n increases, which verifies its ability to solve large-scale problems.

The rest of this paper will be organized as follows. In Section 2, we will introduce related works. In Section 3, we will introduce our new DM-HSGD algorithm. In Section 4, we will show the main theorems of convergence and complexity analysis. In Section 5, we will discuss our experimental results, and Section 6 will conclude the paper.

Table 1: Comparison of Related Algorithms for Minimax Optimization

Name	SFO	Decentralized	Stochastic	Implementation	Reference
SGDA	$O(\kappa^3 \epsilon^{-4})$	×	✓	single-loop	[19]
SGDmax	$O(\kappa^3 \epsilon^{-4} \log(\frac{1}{\epsilon}))$	×	✓	double-loop	[19]
SREDA	$O(\kappa^3 \epsilon^{-3})$	×	✓	double-loop	[25]
Acc-MDA	$O(\kappa^3 \epsilon^{-3})$	×	✓	single-loop	[11]
DPOSG	$O(\epsilon^{-12})$	✓ (iid)	✓	single-loop	[21]
GT/DA	$O(N \epsilon^{-2} \log(\frac{1}{\epsilon}))$	✓ (non-iid)	×	double-loop	[41]
DM-HSGD	$O(\kappa^3 \epsilon^{-3})$	✓ (non-iid)	✓	single-loop	Ours

2 Related Works

2.1 Centralized Minimax Optimization

In recent years, many algorithms for solving minimax optimization were proposed, and the majority of them were studied under the nonconvex-strongly-concave condition. SGDmax [14] is a double loop algorithm that achieves SFO complexity of $O(\kappa^3 \epsilon^{-4} \log(1/\epsilon))$ where $\kappa = L/\mu$ is the condition number. Proximally Guided Stochastic Mirror Descent and Variance Reduction (PGSMD/PGSVRG) [34] are double loop algorithms that achieve SFO complexity of $O(\kappa^3 \epsilon^{-4})$ for stochastic problem and $O(\kappa^2 N \epsilon^{-2})$ for finite-sum problem where N is the number of samples. Multistep GDA (MGDA) [32] is a double loop algorithm and HiBSA [23] is a single loop algorithm. Both MGDA and HiBSA are deterministic hence they can only solve finite-sum problems. Both of them achieve SFO complexity of $O(\kappa^4 N \epsilon^{-2})$. Proximal Dual Implicit Accelerated Gradient (ProxDIAG) is a deterministic triple loop algorithm whose SFO complexity for finite-sum problem is $O(\kappa^{1/2} N \epsilon^{-2})$.

SGDA [19], Stochastic Recursive gradiEnt Descent Ascent (SREDA) [25], and Hybrid Variance-Reduced SGD [40] are more related to our work. SGDA is a single loop algorithm to solve nonconvex-strongly-concave and nonconvex-concave minimax problems. For nonconvex-strongly-concave problem, it requires $O(\kappa^3 \epsilon^{-4})$ SFO complexity to find an ϵ -stationary point of $\Phi(x)$. In this paper, we will prove that our method achieves a better SFO complexity.

SREDA [25] is a double loop algorithm that achieves $O(\kappa^3 \epsilon^{-3})$ SFO complexity. It accelerates SGDA by using SPIDER, which is a variance reduction technique and utilizes the newest gradient information [7, 30]. SREDA also involves a separated initialization algorithm called PiSARAH [31] to ensure the convergence. More recently, [13] proposed an efficient mirror descent ascent algorithm for nonconvex-strongly-concave minimax optimization with nonsmooth regularization based on Bregman distance and variance reduced technique of SPIDER. In our paper, we use another variance-reduced technique named STORM or hybrid stochastic gradient descent [3] to accelerate the algorithm. We will discuss the challenges of using SPIDER on decentralized settings in Section 3. Different from SREDA, our method only requires a large batch at the first iteration. Except the first iteration, we can use either a single sample or a mini-batch to calculate the stochastic gradient. However, SREDA loads a mega-batch with size $O(\epsilon^{-2})$ periodically (every q iterations) and needs $O(\epsilon^{-1})$ gradient oracles at each iteration, which is not practical for large-scale problems. Besides, the maximizer in SREDA is a nested loop to update variable y and if we count the loop of SPIDER then SREDA is actually a triple algorithm. On the contrary, there is no nested loop in our DM-HSGD, which makes our method more efficient and convenient to implement. Moreover, unlike SREDA, our method does not require a separated initialization algorithm to calculate a precise initial value for y .

Hybrid Variance-Reduced SGD algorithm also takes advantage of hybrid stochastic gradient descent to accelerate minimax optimization. For example, [40, 11] applied the Hybrid Variance-Reduced SGD to minimax problems. More recently, [9, 12] proposed some efficient adaptive gradient descent ascent methods for nonconvex-strongly-concave minimax optimization based on momentum techniques including Hybrid Variance-Reduced SGD.

2.2 Decentralized Minimax Optimization

At decentralized setting, most minimax algorithms were proposed for convex-concave problem [17, 28]. In [22] a nonconvex-nonconcave algorithm DPPSP was proposed. However, it is not

gradient-based and the closed-form solution to the subproblem is not ensured in our problem. Hence we will not discuss it in this paper. Decentralized Parallel Optimistic Stochastic Gradient (DPOSG) [21] is the first algorithm applicable to a general decentralized minimax problem with theoretical guarantees. It is a single loop minimax algorithm that generalizes Optimistic Stochastic Gradient (OSG) [2] to decentralized training. However, DPOSG has some obvious drawbacks. The first one is that the gradient complexity $O(\epsilon^{-12})$ is too high and we are motivated to design a faster algorithm. The second one is that DPOSG only works in the case where the data distribution is identical. When the data distribution is non-identical, the Lemma 3 in [21] is not satisfied. Actually the assumption of identical data distribution is not satisfied at most decentralized training tasks. Thus, in this paper, we do not use this assumption.

More recently, [41] studied decentralized nonconvex-strongly-concave minimax problems and proposed a double loop deterministic Gradient Tracking/Descent-Ascent algorithm which extends the vanilla GDA to decentralized setting and combines it with gradient tracking. It achieves a gradient complexity of $O(\epsilon^{-2})$. However, in large-scale machine learning tasks such as deep neural network, generally the full gradient is unavailable and the application of deterministic algorithms is very restricted. If we convert Gradient Tracking/Descent-Ascent to stochastic gradient version, the SFO complexity should be at least $O(\epsilon^{-4})$, which is the same result as SGD in nonconvex optimization. Under the same conditions, our new algorithm achieves a better SFO complexity of $O(\epsilon^{-3})$.

[24] studied decentralized reinforcement learning problem based on distributed constrained Markov decision process model and proposed a decentralized policy gradient optimization method named Safe Dec-PG, which achieves SFO complexity of $O(\epsilon^{-4})$. However, the problem studied in [24] has a special form that is linear in y . In this paper, we focus on general minimax problem. [1] is a simultaneous work of our work that studies a more general decentralized variational inequality problem with higher complexity. We summarize the comparison of related algorithms for general minimax optimization in Table 1. For decentralized algorithms DPOSG, GT/DA, and DM-HSGD, we also discuss whether they can converge on non-identical distributed data.

3 Proposed New Algorithm

3.1 Preliminaries

Before we propose our algorithms, we will introduce the notations used in this paper and some important concepts. We use lower case $x_t^{(i)}$ and $y_t^{(i)}$ to represent the column vector parameters on i -th worker node. We use upper case X_t and Y_t to represent the n -column matrix formed by $x_t^{(i)}$ and $y_t^{(i)}$ respectively, which means $X_t = [x_t^{(1)}, x_t^{(2)}, \dots, x_t^{(n)}]$ and $Y_t = [y_t^{(1)}, y_t^{(2)}, \dots, y_t^{(n)}]$. Column vectors $u_t^{(i)}$, $v_t^{(i)}$, $g_t^{(i)}$ and $h_t^{(i)}$ are gradient estimators used in our algorithms. Upper case U_t , V_t , G_t and H_t are matrices of which the i -th column is $u_t^{(i)}$, $v_t^{(i)}$, $g_t^{(i)}$ and $h_t^{(i)}$ respectively. Lower case with a bar represents the mean vector. Upper case with a bar represents the matrix that each column is the mean vector. For example, $\bar{x}_t = \frac{1}{n} \sum_{i=1}^n x_t^{(i)}$ and $\bar{X}_t = [\bar{x}_t, \bar{x}_t, \dots, \bar{x}_t]$. We define the optimal maximum value of y as:

$$y^*(\cdot) = \arg \max_{y \in \mathcal{Y}} f(\cdot, y), \quad \hat{y}_t = \arg \max_{y \in \mathcal{Y}} f(\bar{x}_t, y) \quad (4)$$

Note that when f is strongly-concave in y , \hat{y}_t is unique. We also define:

$$\delta_t = \|\hat{y}_t - \bar{y}_t\|^2 \quad (5)$$

Bold number $\mathbf{0}$ and $\mathbf{1}$ are $n \times 1$ column vectors that each entry is 0 and 1, respectively. For matrices, we use $\|\cdot\|_F$ to denote Frobenius norm and $\|\cdot\|_2$ to denote spectral norm. We use ∇_x and ∇_y to denote the partial derivative with respect to x and y .

Mixing matrix W represents the weights of averaging among the communication network topology. It is doubly stochastic which satisfies:

$$W\mathbf{1} = W^T\mathbf{1} = \mathbf{1} \quad (6)$$

We should notice that here matrix W is not assumed to be symmetric so that the communication network is not restricted to undirected graph.

Algorithm 1 DM-HSGD

Input: mixing matrix W , initial value $x_0^{(i)} = x_0, y_0^{(i)} = y_0, v_{-1}^{(i)} = g_{-1}^{(i)} = \mathbf{0}, u_{-1}^{(i)} = h_{-1}^{(i)} = \mathbf{0}$

Parameter: stepsize η_x, η_y , weight β_x, β_y , batch size b_0 , iteration T

Output: \bar{x}_ζ , where ζ is chosen randomly from $\{1, 2, \dots, T\}$

```
1: On i-th node:
2: for  $t = 0, 1, \dots, T - 1$  do
3:   if  $t = 0$  then
4:      $g_t^{(i)} = \nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_{x,t}^{(i)})$ ,  $|\xi_{x,t}^{(i)}| = b_0$ 
5:      $h_t^{(i)} = \nabla_y F_i(x_t^{(i)}, y_t^{(i)}; \xi_{y,t}^{(i)})$ ,  $|\xi_{y,t}^{(i)}| = b_0$ 
6:   else
7:      $g_t^{(i)} = \nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) + (1 - \beta_x)(g_{t-1}^{(i)} - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}))$ 
8:      $h_t^{(i)} = \nabla_y F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) + (1 - \beta_y)(h_{t-1}^{(i)} - \nabla_y F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}))$ 
9:   end if
10:  Communicate with neighbors and update gradient estimator as follows
11:   $v_t^{(i)} = \sum_{j=1}^n w_{ij}(v_{t-1}^{(j)} + g_t^{(j)} - g_{t-1}^{(j)})$ 
12:   $u_t^{(i)} = \sum_{j=1}^n w_{ij}(u_{t-1}^{(j)} + h_t^{(j)} - h_{t-1}^{(j)})$ 
13:  Communicate with neighbors and update model parameter as follows
14:   $x_{t+1}^{(i)} = \sum_{j=1}^n w_{ij}(x_t^{(j)} - \eta_x v_t^{(j)})$ 
15:   $y_{t+\frac{1}{2}}^{(i)} = \sum_{j=1}^n w_{ij}(y_t^{(j)} + \eta_y u_t^{(j)})$ ,  $y_{t+1}^{(i)} = P_{\mathcal{Y}}(y_{t+\frac{1}{2}}^{(i)})$ 
16: end for
```

3.2 Decentralized Minimax Hybrid Stochastic Gradient Descent

In this subsection, we introduce our new Decentralized Minimax Hybrid Stochastic Gradient Descent (DM-HSGD) algorithm. Our algorithm is a single loop minimax algorithm (summarized in Algorithm 1) which does not contain a nested loop structure.

The initial points of different nodes are the same, *i.e.* $x_0^{(i)} = x_0$ and $y_0^{(i)}$. $g_t^{(i)}$ and $h_t^{(i)}$ are the gradient estimators with respect to x and y on i -th node. $g_t^{(i)}$ and $h_t^{(i)}$ are computed in the same way as STORM [3]. When $t = 0$, we load a large batch with size b_0 to calculate the stochastic gradient (lines 4 and 5 in Algorithm 1). When $t > 0$, we can use either a single sample or a mini-batch to calculate the gradient (lines 7 and 8 in Algorithm 1). $g_t^{(i)}$ can also be written as

$$g_t^{(i)} = \beta_x \nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) + (1 - \beta_x) \left(g_{t-1}^{(i)} - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}) + \nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) \right) \quad (7)$$

which is a linear combination of the gradient estimators of stochastic gradient descent (the first part) and SPIDER (the second part). As we have mentioned, SPIDER is a variance-reduced method that utilizes the newest gradient information. Thus, estimator Eq. (7) is also called hybrid stochastic gradient descent. It is the same with $h_t^{(i)}$. Then each worker communicates with their neighbors to compute gradient estimator $v_t^{(i)}$ and $u_t^{(i)}$. Here we use gradient tracking [5, 47] to reduce the consensus error (lines 11 and 12 in Algorithm 1). We will discuss why gradient tracking is necessary in our method at next subsection. After we obtain $u_t^{(i)}$ and $v_t^{(i)}$, each worker communicates with their neighbors again and updates the model parameters x and y . Here $P_{\mathcal{Y}}(\cdot)$ represents the projection onto convex set \mathcal{Y} . In the theoretical analysis, we define $Y_{-\frac{1}{2}} = Y_0$.

3.3 Discussions on STORM and Gradient Tracking

In this subsection, we will discuss the intuition of our algorithm and explain why we choose STORM and gradient tracking rather than generalizing SREDA for decentralized setting. The first reason is that SREDA requires large batch or full batch periodically, which is expensive and even unavailable. Besides, there are too many nested loops in SREDA and it is not efficient or convenient. From the view of theoretical analysis, normalization or projection are likely to cause divergence in decentralized training on non-identical data distribution, which is indicated by the following Example 1. Therefore, in the circumstance of this paper, SPIDER will probably not converge to a stationary point. Besides,

SREDA adopts smaller stepsize at the beginning and larger stepsize at the end when $\|v_t\|$ becomes small enough. However, when the data distribution is non-identical, $\|v_t\|$ may not tend to 0 and the stepsize of SREDA will probably always keep small. In contrast, STORM can avoid these issues and we use STORM to accelerate the decentralized minimax algorithm.

In the standard decentralized framework D-PSGD [18], the consensus error satisfies $\|X_t - \bar{X}_t\|_F \leq O(\epsilon)$ when the stepsize η is $O(\epsilon)$ and t is large enough. The following Example 2 is a simple example to show that this bound is tight and there are cases where consensus error $\|X_t - \bar{X}_t\|_F$ is exactly $\Theta(\eta)$ when the data distribution is non-identical. However, according to the analysis of STORM [3] without gradient tracking, the error term $e_t = \bar{g}_t - \nabla_x f(\bar{x}_t, \bar{y}_t)$ between the averaged update direction and the correct direction is supposed to satisfy:

$$\|e_t\|^2 \leq (1 - \beta_x)\|e_{t-1}\|^2 + O(\eta_x^4). \quad (8)$$

Nevertheless, the consensus error $\|X_t - \bar{X}_t\|_F^2$ is only $O(\eta_x^2)$ and cannot be as small as $O(\eta_x^4)$ if there is no gradient tracking. Therefore, to inherit the analysis framework of STORM, the gradient tracking in our algorithm is essential.

Example 1. Assume $f(x) = f_1(x) + f_2(x)$, where $x = (a, b) \in \mathbb{R}^2$. $f_1(x) = a$ and $f_2(x) = \sqrt{3}b$ are defined on two different nodes. Let W be the uniform weighted mixing matrix. We can compute $v_1 = (1, 0)$ and $v_2 = (0, \sqrt{3})$. The ideal averaged gradient direction is $v^* = (1/2, \sqrt{3}/2)$. However, if we do normalization before making consensus, the obtained gradient estimator is $v = (1/2, 1/2)$, which is deviated from v^* .

Example 2. Suppose there are two sequences $\{p_t\}$ and $\{q_t\}$ defined on two different nodes with $p_0 = q_0$. They are updated by $p_{t+\frac{1}{2}} = p_t - \eta a$ and $q_{t+\frac{1}{2}} = q_t - \eta b$ at each iteration respectively where a and b are fixed gradient directions. As data distribution is non-identical, we have $a \neq b$. Assume the mixing matrix is

$$W = \begin{bmatrix} 2/3 & 1/3 \\ 1/3 & 2/3 \end{bmatrix}$$

Then we have

$$p_{t+1} - q_{t+1} = \frac{1}{3}(p_t - q_t) - \frac{\eta}{3}(a - b) = \frac{1}{3^{t+1}}(p_0 - q_0) - \eta \left(\sum_{s=1}^{t+1} \frac{1}{3^s} \right) (a - b) = \frac{\eta}{2} \left(1 - \frac{1}{3^{t+1}} \right) (b - a) \quad (9)$$

Therefore, $\lim_{t \rightarrow \infty} \|p_t - q_t\| = \frac{\eta}{2} \|a - b\|$.

4 Convergence Analysis

In this section, we will show the main theorems of our convergence analysis. The theoretical results show that the SFO complexity of our algorithm is $O(\kappa^3 \epsilon^{-3})$, which is the same as the best result in centralized minimax problem [25]. First we will introduce the following assumptions.

Assumption 1. (Lipschitz Gradient). Each component function $F_i(x, y; \xi)$ is L -smooth, which means there exists a constant L such that for any (x, y) and (x', y') , we have

$$\|\nabla F_i(x, y; \xi) - \nabla F_i(x', y'; \xi)\|^2 \leq L^2(\|x - x'\|^2 + \|y - y'\|^2)$$

Assumption 2. (Bounded Variance). The gradient of each component function $F_i(x, y; \xi)$ is an unbiased estimator of $\nabla f_i(x, y)$ and has bounded variance, i.e.,

$$\mathbb{E}\|\nabla F_i(x, y; \xi) - \nabla f_i(x, y)\|^2 \leq \sigma < +\infty$$

Assumption 3. (Lower Bound). The function $\Phi(\cdot)$ is lower bounded, i.e., $\inf_x \Phi(x) = \Phi^* > -\infty$.

Assumption 4. (Spectral Gap). The doubly stochastic matrix W satisfies $\|W - \frac{\mathbf{1}\mathbf{1}^T}{n}\|_2 = \lambda \in [0, 1)$.

Assumption 5. (Strongly Concave). The function $f_i(x, y)$ is μ -strongly-concave in y . That is, there exists a constant $\mu > 0$, for any x, y and y' , we have

$$f_i(x, y) \leq f_i(x, y') + \langle \nabla_y f_i(x, y'), y - y' \rangle - \frac{\mu}{2} \|y - y'\|^2$$

These are very common and mild assumptions that are frequently assumed in previous works. Assumptions 1, 2 and 3 are also used in minimax methods [25] and [19]. Assumption 4 is used in [46]. Typically, the spectral gap assumption is stated as W is symmetric and $|\lambda_2| < 1$, $|\lambda_n| < 1$ where $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ are the eigenvalues of W [16, 18, 51]. Our Assumption 4 is automatically satisfied if the typical spectral gap assumption holds (see Lemma 16 in [16]). Assumption 5 is the definition of strong concavity.

In nonconvex-strongly-concave problem, we use ϵ -stationary point of $\Phi(x)$, *i.e.* $\|\nabla\Phi(x)\| \leq \epsilon$ as the convergence criterion. From Lemma 4.3 in [19], we know $\Phi(x)$ is differentiable and $(L + \kappa L)$ -smooth and $y^*(\cdot)$ is κ -Lipschitz, which means $\|y^*(x_1) - y^*(x_2)\| \leq \kappa\|x_1 - x_2\|$ for any $x_1, x_2 \in \mathbb{R}^{d_1}$. Furthermore, we have:

$$\nabla\Phi(\bar{x}_t) = \nabla_x f(\bar{x}_t, \hat{y}_t) + \nabla_y f(\bar{x}_t, \hat{y}_t) \cdot \partial y^*(\bar{x}_t) = \nabla_x f(\bar{x}_t, \hat{y}_t) \quad (10)$$

since $\nabla_y f(\bar{x}_t, \hat{y}_t) = 0$. This criterion is broadly used in the analysis of nonconvex-strongly-concave minimax optimization [19, 39]. Now we will provide the main theorems of our convergence analysis. Completed proof can be found in the Supplementary Material.

Theorem 1. *Let Assumptions 1 to 5 hold. When parameters $\beta_x = \frac{\epsilon \min\{1, n\epsilon\}}{20}$, $\beta_y = \frac{\epsilon \min\{1, n\epsilon\}}{500\kappa^2}$, $\eta_x = \frac{(1-\lambda)^2 \min\{1, n\epsilon\}}{2000\kappa^3 L}$, $\eta_y = \frac{(1-\lambda)^2 \min\{1, n\epsilon\}}{500\kappa L}$, $b_0 = \frac{400}{\min\{1, n\epsilon\}}$, $T = \frac{4000\kappa^3 \epsilon^{-2}}{(1-\lambda)^2 \min\{1, n\epsilon\}}$, our Algorithm 1 satisfies*

$$\begin{aligned} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi(\bar{x}_t)\|^2 &\leq L(\Phi(x_0) - \Phi^*)\epsilon^2 + \sigma^2\epsilon^2 + L^2\delta_0\epsilon^2 + \frac{\epsilon^2}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_x f_i(x_0, y_0)\|^2 \\ &\quad + \frac{\epsilon^2}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_y f_i(x_0, y_0)\|^2 \end{aligned} \quad (11)$$

Corollary 1. *When the parameters are defined as Theorem 1, we can see $\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi(\bar{x}_t)\|^2 \leq O(\epsilon^2)$. Therefore, if $n \leq O(\epsilon^{-1})$, the SFO complexity of Algorithm 1 is $O(\kappa^3 \epsilon^{-3})$. If $n > O(\epsilon^{-1})$, the SFO complexity is $O(\kappa^3 n \epsilon^{-2})$. Besides, from the proof of Theorem 1 we can see error $\|\bar{y}_t - y^*(\bar{x}_t)\|^2$ is also bounded by the right side of Eq. (11).*

Theorem 1 is the theoretical result when T is determined by ϵ . If the number of iteration T is not fixed, we have the following conclusion.

Theorem 2. *Let Assumptions 1 to 5 hold. We set the parameters as $T = \frac{4000\kappa^3 T_0}{(1-\lambda)^2}$, $\beta_x = \frac{n^{1/3}}{20T_0^{2/3}}$, $\beta_y = \frac{n^{1/3}}{500\kappa^2 T_0^{2/3}}$, $\eta_x = \frac{(1-\lambda)^2 n^{2/3}}{2000\kappa^3 T_0^{1/3} L}$, $\eta_y = \frac{(1-\lambda)^2 n^{2/3}}{500\kappa T_0^{1/3} L}$, $b_0 = \frac{T_0^{1/3}}{n^{2/3}}$, where we suppose $T_0 \geq 10n^2$. Then our algorithm satisfies*

$$\begin{aligned} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi(\bar{x}_t)\|^2 &\leq \frac{L(\Phi(x_0) - \Phi^*) + \sigma^2 + L^2\delta_0}{(nT_0)^{2/3}} + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_x f_i(x_0, y_0)\|^2}{T_0} \\ &\quad + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_y f_i(x_0, y_0)\|^2}{T_0} \end{aligned} \quad (12)$$

Corollary 2. *From Theorem 2, we know $\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla\Phi(\bar{x}_t)\|^2 \leq O(\frac{1}{(nT_0)^{2/3}}) + O(\frac{1}{T_0})$ when parameters are defined as above. As we suppose $T_0 \geq O(n^2)$, the dominating term in the convergence rate is $O(\frac{1}{(nT_0)^{2/3}})$, which indicates the linear speedup of our algorithm.*

5 Experiments

5.1 Robust Logistic Regression

We conduct the experiment of decentralized robust logistic regression¹ task as the first experiment, which was proposed in [49] and was also conducted in the related work [25]. Given dataset

¹<https://github.com/Trashzzz/DM-HSGD>

$\{(a_i, b_i)\}_{i=1}^n$, where $a_i \in \mathbb{R}^d$ is the feature and $b_i \in \{-1, 1\}$ is the label, the robust logistic regression problem is formulated as follows:

$$\min_{x \in \mathbb{R}^d} \max_{y \in \Delta_n} f(x, y) = \sum_{i=1}^n y_i l_i(x) - V(y) + g(x) \quad (13)$$

where y_i is the i -th component of variable y . $l_i(x)$ is the logistic loss function which is defined by $l_i(x) = \log(1 + \exp(-b_i a_i^T x))$. $V(y)$ is a divergence measure defined by $V(y) = \frac{1}{2} \lambda_1 \|ny - \mathbf{1}\|^2$. Δ_n represents the simplex in \mathbb{R}^n , which means

$$\Delta_n = \{y \in \mathbb{R}^n | 0 \leq y_i \leq 1, \sum_{i=1}^n y_i = 1\} \quad (14)$$

$g(x)$ is a nonconvex regularization with form $g(x) = \lambda_2 \sum_{i=1}^d \frac{\alpha x_i^2}{1 + \alpha x_i^2}$. Following the experimental settings in [25, 49], we let $\lambda_1 = \frac{1}{n^2}$, $\lambda_2 = 0.001$ and $\alpha = 10$ in our experiment.

Table 2: Descriptions of datasets used in our experiment

Name	a9a	covtype	ijcnn1	phishing	rcv1	w8a
N	32561	581012	49990	11055	20242	49749
d	123	54	22	68	47236	300

We conduct our experiment on six real-world training datasets ‘‘a9a’’, ‘‘covtype’’, ‘‘ijcnn1’’, ‘‘phishing’’, ‘‘rcv1’’ and ‘‘w8a’’, which can be downloaded from LIBSVM² repository. The description of datasets is listed in Table 2 where N is the number of samples and d is the number of features. We implement our code on an MPI cluster where each node is equipped with 12-core Intel Xeon E5-2620 v3 2.40 GHz processor.




Figure 1: Results of our decentralized robust logistic regression task. Figure (a) to (f) show the value of $\Phi(x)$ with respect to the number of gradient oracles divided by 10^3 . Figure (a), (b), (c), (d), (e) and (f) are experimental results on ‘‘a9a’’, ‘‘covtype’’, ‘‘ijcnn1’’, ‘‘phishing’’, ‘‘rcv1’’ and ‘‘w8a’’ respectively.

We compare our DM-HSGD algorithm with baseline algorithms: SGDA [19], SREDA [25], DPOSG [21], and stochastic Gradient Tracking/Descent Ascent (SGTDA) [41]. We consider the algorithms

²<https://www.csie.ntu.edu.tw/~cjlin/libsvmtools/datasets>

for solving stochastic problem. We set the number of worker nodes to $n = 20$ and use the ring-based topology as the communication network. For each algorithm, we grid search the learning rates η_x and η_y from $\{0.1, 0.01, 0.001, 0.0001\}$. The mini-batch size is set to 20. The number of iterations in the nested loop for double-loop algorithms is set to $K = 5$. For DM-HSGD, we set the batch size of the first iteration to $b_0 = 10000$. β_x and β_y are set to 0.01. For SREDA, we set $\epsilon = 0.1$ in the factor $\frac{\epsilon}{\|v_t\|}$, period $q = 50$ and large batch size $S_1 = 1000$. We compare the value of $\Phi(x)$ with respect to the number of gradient oracles among different algorithms, which can also be calculated by the projection onto simplex Δ_n . The experimental results are shown in Figure 1. From the experimental results in Figure 1, we can see our new DM-HSGD algorithm converges faster than other baseline algorithms, which verifies the performance of our method.

5.2 Policy Evaluation

Our second experiment is the decentralized policy evaluation (PE) task. PE is an important task in reinforcement learning, which aims to estimate the value function of a given policy. The most intuitive and frequently used method for PE is temporal-difference (TD) method that relies on the Bellman equation [4]. However, traditional TD method, which is probably not true gradient descent method as pointed out in [20] and [37], are shown to be unstable in the case of off-policy sampling or nonlinear function approximation. [36] first proposed a method to optimize the objective function of mean-squared projected Bellman error (MSPBE) and MSPBE is proven to achieve asymptotic convergence with arbitrary nonlinear smooth function approximation in [27]. In [42], the MSPBE objective function with nonlinear approximation is converted into a nonconvex-strongly-concave minimax problem by Fenchel’s duality. The problem can be formulated as:

$$\min_{\theta} \max_w L(\theta, w) = \frac{1}{nN_i} \sum_{i=1}^n \sum_{j=1}^{N_i} L_j^{(i)}(\theta, w),$$

$$L_j^{(i)}(\theta, w) = \langle w, [R_i(s_j, a_j) + \gamma V_{\theta}(s_{j+1}) - V_{\theta}(s_j)] g_{\theta}(s_j) \rangle - \frac{1}{2} (w^T g_{\theta}(s_j))^2 \quad (15)$$

where s_j is a state and a_j is an action. R_i represents the reward and $\gamma \in (0, 1)$ is the discount factor. V is a value function that maps the state space to a real number. θ is the parameter to estimate the value function. Function g_{θ} is the gradient of V_{θ} and parameter w is yield by Fenchel’s duality.

Mountaincar [35] is a preliminary task in reinforcement learning. [42] and [43] ran offline PE task of this problem with primal-dual MSPBE, where the objective function is formulated as Eq. (15). Following the experimental settings in [42], we use Sarsa [35] to generate trajectories of transitions (s_i, a_i, s_{i+1}, r_i) with d features and $N = 5000$ samples on each worker node. We parameterize value function V_{θ} as a 2-layer neural network with H hidden neurons. We use Sigmoid function as activation and set discount factor to $\gamma = 0.95$. This experiment is run on an MPI cluster where each node is equipped with 12-core Intel Xeon E5-2620 v3 2.40 GHz processor.




Figure 2: Results of our policy evaluation task. Figures (a), (b) and (c) show the value of $\Phi(\theta)$ with respect to the number of gradient oracles divided by 10^3 . In Figures, (a) $d = 200$, $H = 50$; (b) $d = 300$, $H = 100$; (c) $d = 400$, $H = 200$.

We compare our DM-HSGD algorithm with baseline algorithms: SGDA [19], SREDA [25], DPOSG [21], and stochastic Gradient Tracking/Descent Ascent (SGTDA) [41]. We also consider algorithms

for solving stochastic problem. We set the number of worker nodes to $n = 20$. We also use a ring-based topology with uniform weights as the communication network in this task. For each algorithm, we grid search the learning rates η_x and η_y from $\{0.1, 0.01, 0.001, 0.0001\}$. The mini-batch size is set to 20. The number of iterations in the nested loop for double-loop algorithms is set to $K = 5$. For DM-HSGD, we set the batch size of the first iteration to $b_0 = 2500$. β_x and β_y are set to 0.01. For SREDA, we set $\epsilon = 0.1$ in the factor $\frac{\epsilon}{\|v_t\|}$, period $q = 50$ and large batch size $S_1 = 1000$. We compare the value of $\Phi(\theta)$ with respect to the number of gradient oracles among different algorithms, which can be calculated by quadratic optimization. The experimental results are shown in Figure 2.

Figure 2 (a), (b) and (c) show that our DM-HSGD algorithm achieves the fastest convergence regarding the number of gradient oracles. From the experimental result, we can also see that nested loop algorithm for minimax optimization usually consumes more gradient complexity during the training process than single-loop algorithm.

6 Conclusion

In this paper, we proposed a novel accelerated decentralized minimax algorithm, Decentralized Minimax Hybrid Stochastic Gradient Descent (DM-HSGD), to solve the stochastic nonconvex-strongly-concave minimax optimization problems. We prove that our new method obtains SFO complexity of $O(\kappa^3\epsilon^{-3})$ which outperforms the existing results in decentralized minimax optimization and matches state-of-the-art in centralized minimax optimization. Our method also achieves linear speedup with respect to the number workers, which shows its ability to solve large-scale problems. We also conduct experiments on two machine learning tasks, decentralized robust logistic regression and policy evaluation to validate the superior performance of our algorithm. In our future work, we will explore the decentralized nonconvex-concave minimax optimization without the strong concavity so that it can solve a broader range of problems including the loss functions that are linear in y . We will probably consider the methods that add a perturbation such as Catalyst [50].

Acknowledgments and Disclosure of Funding

This work was partially supported by NSF IIS 1845666, 1852606, 1838627, 1837956, 1956002, OIA 2040588.

References

- [1] Aleksandr Beznosikov, Pavel Dvurechensky, Anastasia Koloskova, Valentin Samokhin, Sebastian U Stich, and Alexander Gasnikov. Decentralized local stochastic extra-gradient for variational inequalities. *arXiv preprint arXiv:2106.08315*, 2021.
- [2] Chao-Kai Chiang, Tianbao Yang, Chia-Jung Lee, Mehrdad Mahdavi, Chi-Jen Lu, Rong Jin, and Shenghuo Zhu. Online optimization with gradual variations. *Conference on Learning Theory*, 2012.
- [3] Ashok Cutkosky and Francesco Orabona. Momentum-based variance reduction in non-convex sgd. *Neural Information Processing Systems (NeurIPS)*, 2019.
- [4] Christoph Dann, Gerhard Neumann, and Jan Peters. Policy evaluation with temporal differences: A survey and comparison. *Journal of Machine Learning Research*, 2014.
- [5] Paolo Di Lorenzo and Gesualdo Scutari. Next: In-network nonconvex optimization. *IEEE Transactions on Signal and Information Processing over Networks*, 2(2):120–136, 2016.
- [6] Simon S. Du and Wei Hu. Linear convergence of the primal-dual gradient method for convex-concave saddle point problems without strong convexity. *International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2019.
- [7] Cong Fang, Chris Junchi Li, Zhouchen Lin, and Tong Zhang. Spider: Near-optimal non-convex optimization via stochastic path integrated differential estimator. *Neural Information Processing Systems (NeurIPS)*, 2018.

- [8] Ian J. Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron Courville, and Yoshua Bengio. Generative adversarial nets. *Neural Information Processing Systems (NeurIPS)*, 2014.
- [9] Zhishuai Guo, Yi Xu, Wotao Yin, Rong Jin, and Tianbao Yang. On stochastic moving-average estimators for non-convex optimization. *arXiv preprint arXiv:2104.14840*, 2021.
- [10] Martin Heusel, Hubert Ramsauer, Thomas Unterthiner, Bernhard Nessler, and Sepp Hochreiter. Gans trained by a two time-scale update rule converge to a local nash equilibrium. *Neural Information Processing Systems (NeurIPS)*, 2017.
- [11] Feihu Huang, Shangqian Gao, Jian Pei, and Heng Huang. Accelerated zeroth-order and first-order momentum methods from mini to minimax optimization. *Journal of Machine Learning Research*, 22:1–70, 2021.
- [12] Feihu Huang and Heng Huang. Adagda: Faster adaptive gradient descent ascent methods for minimax optimization. *arXiv preprint arXiv:2106.16101*, 2021.
- [13] Feihu Huang, Xidong Wu, and Heng Huang. Efficient mirror descent ascent methods for nonsmooth minimax problems. *Advances in Neural Information Processing Systems*, 34, 2021.
- [14] Chi Jin, Praneeth Netrapalli, and Michael I. Jordan. What is local optimality in nonconvex-nonconcave minimax optimization? *arXiv:1902.00618v2*, 2019.
- [15] R. Johnson and T. Zhang. Accelerating stochastic gradient descent using predictive variance reduction. *Neural Information Processing Systems (NeurIPS)*, 2013.
- [16] Anastasia Koloskova, Sebastian Stich, and Martin Jaggi. Decentralized stochastic optimization and gossip algorithms with compressed communication. In *International Conference on Machine Learning*, pages 3478–3487. PMLR, 2019.
- [17] Alec Koppel, Felicia Y. Jakubiec, and Alejandro Ribeiro. A saddle point algorithm for networked online convex optimization. *IEEE Transactions on Signal Processing*, 2015.
- [18] Xiangru Lian, Ce Zhang, Huan Zhang, Cho-Jui Hsieh, Wei Zhang, and Ji Liu. Can decentralized algorithms outperform centralized algorithms? a case study for decentralized parallel stochastic gradient descent. *Neural Information Processing Systems (NeurIPS)*, 2017.
- [19] Tianyi Lin, Chi Jin, and Michael I. Jordan. On gradient descent ascent for nonconvex-concave minimax problems. *arXiv:1906.00331v4*, 2019.
- [20] Bo Liu, Ji Liu, Mohammad Ghavamzadeh, Sridhar Mahadevan, and Marek Petrik. Finite-sample analysis of proximal gradient algorithms. *Conference on Uncertainty in Artificial Intelligence*, 2015.
- [21] Mingrui Liu, Wei Zhang, Youssef Mroueh, Xiaodong Cui, Jerret Ross, Tianbao Yang, and Payel Das. A decentralized parallel algorithm for training generative adversarial nets. *Smooth Games Optimization and Machine Learning Workshop (NeurIPS)*, 2019.
- [22] Weijie Liu, Aryan Mokhtari, Asuman Ozdaglar, Sarath Pattathil, Zebang Shen, and Nenggan Zheng. A decentralized proximal point-type method for saddle point problems. *arXiv:1910.14380v1*, 2019.
- [23] Songtao Lu, Ioannis Tsaknakis, Mingyi Hong, and Yongxin Chen. Hybrid block successive approximation for one-sided non-convex min-max problems: Algorithms and applications. *arXiv:1902.08294v1*, 2019.
- [24] Songtao Lu, Kaiqing Zhang, Tianyi Chen, Tamer Basar, and Lior Horesh. Decentralized policy gradient descent ascent for safe multi-agent reinforcement learning. *Association for the Advancement of Artificial Intelligence (AAAI)*, 2021.
- [25] Luo Luo, Haishan Ye, and Tong Zhang. Stochastic recursive gradient descent ascent for stochastic nonconvex-strongly-concave minimax problems. *Neural Information Processing Systems (NeurIPS)*, 2020.

- [26] Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. *International Conference on Learning Representations (ICLR)*, 2018.
- [27] Hamid Reza Maei, Csaba Szepesvári, Shalabh Bhatnagar, Doina Precup, David Silver, and Richard S. Sutton. Convergent temporal-difference learning with arbitrary smooth function approximation. *Neural Information Processing Systems (NeurIPS)*, 2010.
- [28] David Mateos-Núñez and Jorge Cortés. Distributed subgradient methods for saddle-point problems. *IEEE Conference on Decision and Control*, 2015.
- [29] Arkadi Nemirovski. Prox-method with rate of convergence $o(1/t)$ for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM Journal on Optimization*, 2004.
- [30] Lam M. Nguyen, Jie Liu, Katya Scheinberg, and Martin Takáč. Sarah: A novel method for machine learning problems using stochastic recursive gradient. *arXiv:1703.00102v2*, 2017.
- [31] Lam M. Nguyen, Katya Scheinberg, and Martin Takáč. Inexact sarah algorithm for stochastic optimization. *arXiv:1811.10105*, 2018.
- [32] Maher Nouiehed, Maziar Sanjabi, Tianjian Huang, and Jason D. Lee. Solving a class of non-convex min-max games using iterative first order methods. *Neural Information Processing Systems (NeurIPS)*, 2019.
- [33] Taoxing Pan, Jun Liu, and Jie Wang. D-spider-sfo: A decentralized optimization algorithm with faster convergence rate for nonconvex problems. *Association for the Advancement of Artificial Intelligence (AAAI)*, 2020.
- [34] Hassan Rafique, Mingrui Liu, Qihang Lin, and Tianbao Yang. Non-convex min-max optimization: Provable algorithms and applications in machine learning. *arXiv:1810.02060v3*, 2018.
- [35] Richard S. Sutton and Andrew G. Barto. *Reinforcement Learning: An Introduction*. MIT Press, 1998.
- [36] Richard S. Sutton, Hamid Reza Maei, Doina Precup, Shalabh Bhatnagar, David Silver, Csaba Szepesvári, and Eric Wiewiora. Fast gradient-descent methods for temporal-difference learning with linear function approximation. *International Conference on Machine Learning (ICML)*, 2009.
- [37] Csaba Szepesvári. *Algorithms for Reinforcement Learning*. Synthesis Lectures on Artificial Intelligence and Machine Learning, 2010.
- [38] Hanlin Tang, Xiangru Lian, Ming Yan, Ce Zhang, and Ji Liu. d^2 : Decentralized training over decentralized data. *International Conference on Machine Learning (ICML)*, 2018.
- [39] Kiran Koshy Thekumprampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Efficient algorithms for smooth minimax optimization. *Neural Information Processing Systems (NeurIPS)*, 2019.
- [40] Q. Tran-Dinh, D. Liu, and L.M. Nguyen. Hybrid variance-reduced sgd algorithms for minimax problems with nonconvex-linear function. *Neural Information Processing Systems (NeurIPS)*, 2020.
- [41] Ioannis Tsaknakis, Mingyi Hong, and Sijia Liu. Decentralized min-max optimization: Formulations, algorithms and applications in network poisoning attack. In *ICASSP 2020-2020 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 5755–5759. IEEE, 2020.
- [42] Hoi-To Wai, Mingyi Hong, Zhuoran Yang, Zhaoran Wang, and Kexin Tang. Variance reduced policy evaluation with smooth function approximation. *Neural Information Processing Systems (NeurIPS)*, 2019.

- [43] Hoi-To Wai, Zhuoran Yang, Zhaoran Wang, and Mingyi Hong. Multi-agent reinforcement learning via double averaging primal-dual optimization. *Neural Information Processing Systems (NeurIPS)*, 2018.
- [44] Zhe Wang, Kaiyi Ji, Yi Zhou, Yingbin Liang, and Vahid Tarokh. Spiderboost: A class of faster variance-reduced algorithms for nonconvex optimization. *arXiv:1810.10690v2*, 2018.
- [45] Wenhan Xian, Feihu Huang, and Heng Huang. Communication-efficient frank-wolfe algorithm for nonconvex decentralized distributed learning. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, pages 10405–10413, 2021.
- [46] Ran Xin, Usman A. Khan, and Soumya Kar. A hybrid variance-reduced method for decentralized stochastic non-convex optimization. *arXiv:2102.06752*, 2021.
- [47] Jinming Xu, Shanying Zhu, Yeng Chai Soh, and Lihua Xie. Augmented distributed gradient methods for multi-agent optimization under uncoordinated constant stepsizes. In *2015 54th IEEE Conference on Decision and Control (CDC)*, pages 2055–2060. IEEE, 2015.
- [48] Feng Yan, Shreyas Sundaram, S. Vishwanathan, and Yuan Qi. Distributed autonomous online learning: Regrets and intrinsic privacy-preserving properties. *IEEE Transactions on Knowledge and Data Engineering*, 2013.
- [49] Yan Yan, Yi Xu, Qihang Lin, Lijun Zhang, and Tianbao Yang. Stochastic primal-dual algorithms with faster convergence than $O(1/\sqrt{T})$ for problems without bilinear structure. *arXiv:1904.10112*, 2019.
- [50] Junchi Yang, Siqi Zhang, Negar Kiyavash, and Niao He. A catalyst framework for minimax optimization. In *NeurIPS*, 2020.
- [51] Kun Yuan, Qing Ling, and Wotao Yin. On the convergence of decentralized gradient descent. *SIAM Journal on Optimization*, 2016.

A Proof of Convergence Analysis

A.1 Basic Lemmas

First, we introduce following basic lemmas, which are broadly used in the convergence analysis of optimization algorithms.

Lemma 1. *Let vector X be a stochastic variable. Then we have*

$$0 \leq \mathbb{E}\|X - \mathbb{E}X\|^2 = \mathbb{E}\|X\|^2 - \|\mathbb{E}X\|^2 \leq \mathbb{E}\|X\|^2 \quad (16)$$

Lemma 2. *Let X_1, X_2, \dots, X_n be n independent stochastic variables of which the means are 0. Then we have*

$$\mathbb{E}\left\|\sum_{i=1}^n X_i\right\|^2 = \sum_{i=1}^n \mathbb{E}\|X_i\|^2 \quad (17)$$

Lemma 3. *Suppose A and B are two matrices. Then it satisfies*

$$\|AB\|_F \leq \|A\|_2 \|B\|_F \quad (18)$$

A.2 Important Conclusions

Next, we will propose and prove some conclusions that are important to the proof our main theorems.

Lemma 4. *(Lemma 4.3 in paper [19]) $\Phi(x)$ is $(L + \kappa L)$ -smooth and $y^*(\cdot)$ is κ -Lipschitz, which means $\|y^*(x_1) - y^*(x_2)\| \leq \kappa\|x_1 - x_2\|$ for any $x_1, x_2 \in \mathbb{R}^{d_1}$.*

Proof. As $y^*(x_1)$ and $y^*(x_2)$ achieve the maximum, we have $\nabla_y f(x_1, y^*(x_1)) = \mathbf{0}$ and $\nabla_y f(x_2, y^*(x_2)) = \mathbf{0}$. Then we have

$$\begin{aligned} & \|y^*(x_1) - y^*(x_2)\| \\ & \leq \frac{1}{\mu} \|\nabla_y f(x_1, y^*(x_1)) - \nabla_y f(x_1, y^*(x_2))\| \\ & = \frac{1}{\mu} \|\nabla_y f(x_2, y^*(x_2)) - \nabla_y f(x_1, y^*(x_2))\| \leq \frac{L}{\mu} \|x_1 - x_2\| = \kappa \|x_1 - x_2\| \end{aligned} \quad (19)$$

where the first inequality is derived from μ -strong concavity and the second inequality is derived from L -smoothness. Since $\nabla\Phi(x) = \nabla_x f(x, y^*(x))$, from Assumption 1 we get

$$\|\nabla\Phi(x_1) - \nabla\Phi(x_2)\| \leq L\|x_1 - x_2\| + L\|y^*(x_1) - y^*(x_2)\| \leq (L + \kappa L)\|x_1 - x_2\| \quad (20)$$

which implies $\Phi(x)$ is $(L + \kappa L)$ -smooth. \square

Lemma 5. *When $\eta_y \leq \frac{1}{5L}$ we have following estimation for δ_t .*

$$\begin{aligned} \sum_{t=0}^{T-1} \delta_t & \leq \frac{4\kappa}{L\eta_y} \delta_0 + \frac{18\eta_y}{\mu} \sum_{t=1}^{T-1} \left(1 - \frac{\mu\eta_y}{4}\right)^{T-t-1} \sum_{s=0}^{t-1} \|\bar{u}_s - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{72\kappa^2}{n} \sum_{t=0}^{T-1} (\|X_t - \bar{X}_t\|_F^2 \\ & + \|Y_t - \bar{Y}_t\|_F^2) + \frac{20\kappa^4 \eta_x^2}{L^2 \eta_y^2} \sum_{t=0}^{T-1} \|\bar{v}_t\|^2 - \frac{12}{5\mu^2} \sum_{t=0}^{T-1} \left(1 - \left(1 - \frac{\mu\eta_y}{4}\right)^{T-t}\right) \|\bar{u}_t\|^2 \end{aligned} \quad (21)$$

Proof. Define $z_t = \bar{y}_t + \theta \bar{u}_t$ for some constant θ . As function f is strongly-concave in y we have

$$\begin{aligned} f(\bar{x}_t, \hat{y}_t) & \leq f(\bar{x}_t, \bar{y}_t) + \langle \nabla_y f(\bar{x}_t, \bar{y}_t), \hat{y}_t - \bar{y}_t \rangle - \frac{\mu}{2} \|\hat{y}_t - \bar{y}_t\|^2 \\ & = f(\bar{x}_t, \bar{y}_t) + \langle \bar{u}_t, \hat{y}_t - z_t \rangle + \langle \nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t, \hat{y}_t - z_t \rangle \\ & \quad + \theta \langle \nabla_y f(\bar{x}_t, \bar{y}_t), \bar{u}_t \rangle - \frac{\mu}{2} \|\hat{y}_t - \bar{y}_t\|^2 \end{aligned} \quad (22)$$

By Assumption 1, we also have

$$-\frac{L\theta^2}{2} \|\bar{u}_t\|^2 \leq f(\bar{x}_t, z_t) - f(\bar{x}_t, \bar{y}_t) - \theta \langle \nabla_y f(\bar{x}_t, \bar{y}_t), \bar{u}_t \rangle \quad (23)$$

Add Eq. (22) and Eq. (23) together we obtain

$$0 \leq \langle \bar{u}_t, \hat{y}_t - z_t \rangle + \langle \nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t, \hat{y}_t - z_t \rangle - \frac{\mu}{2} \|\hat{y}_t - \bar{y}_t\|^2 + \frac{L\theta^2}{2} \|\bar{u}_t\|^2 \quad (24)$$

where we also use the definition of \hat{y}_t so that $f(\bar{x}_t, \hat{y}_t) \geq f(\bar{x}_t, z_t)$.

$$\langle \bar{u}_t, \hat{y}_t - z_t \rangle = -\theta \|\bar{u}_t\|^2 + \langle \bar{u}_t, \hat{y}_t - \bar{y}_t \rangle \quad (25)$$

Combining Eq. (24) and Eq. (25) we have

$$0 \leq \langle \bar{u}_t, \hat{y}_t - \bar{y}_t \rangle - \frac{\mu}{2} \|\hat{y}_t - \bar{y}_t\|^2 + \langle \nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t, \hat{y}_t - z_t \rangle - (\theta - \frac{L\theta^2}{2}) \|\bar{u}_t\|^2 \quad (26)$$

By Cauchy-Schwartz inequality we have

$$\langle \nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t, \hat{y}_t - z_t \rangle \leq \frac{4}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 + \frac{\mu}{8} \|\hat{y}_t - \bar{y}_t\|^2 + \frac{\mu\theta^2}{8} \|\bar{u}_t\|^2 \quad (27)$$

Therefore, we obtain

$$\begin{aligned} 0 &\leq \langle \bar{u}_t, \hat{y}_t - \bar{y}_t \rangle - \frac{\mu}{4} \|\hat{y}_t - \bar{y}_t\|^2 + \frac{4}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 - (\theta - \frac{L\theta^2}{2} - \frac{\mu\theta^2}{8}) \|\bar{u}_t\|^2 \\ &\leq \langle \bar{u}_t, \hat{y}_t - \bar{y}_t \rangle - \frac{\mu}{4} \|\hat{y}_t - \bar{y}_t\|^2 + \frac{4}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 - \frac{2}{5\mu} \|\bar{u}_t\|^2 \end{aligned} \quad (28)$$

where we let $\theta = \frac{4}{5\mu}$. As we have

$$2\eta_y \langle \bar{u}_t, \hat{y}_t - \bar{y}_t \rangle = \|\bar{y}_t - \hat{y}_t\|^2 + \|\bar{y}_{t+1} - \bar{y}_t\|^2 - \|\bar{y}_{t+1} - \hat{y}_t\|^2 \quad (29)$$

Eq. (28) is equivalent to

$$\|\bar{y}_{t+1} - \hat{y}_t\|^2 \leq (1 - \frac{\mu\eta_y}{2}) \|\bar{y}_t - \hat{y}_t\|^2 + \|\bar{y}_{t+1} - \bar{y}_t\|^2 + \frac{8\eta_y}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 - \frac{4\eta_y}{5\mu} \|\bar{u}_t\|^2 \quad (30)$$

When $L\eta_y \leq \frac{1}{5}$, from Eq. (30) we know

$$\|\bar{y}_{t+1} - \hat{y}_t\|^2 \leq (1 - \frac{\mu\eta_y}{2}) \|\bar{y}_t - \hat{y}_t\|^2 + \frac{8\eta_y}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 - \frac{3\eta_y}{5\mu} \|\bar{u}_t\|^2 \quad (31)$$

According to Young's inequality we have

$$\begin{aligned} \|\bar{y}_{t+1} - \hat{y}_{t+1}\|^2 &\leq (1 + \frac{\mu\eta_y}{4}) \|\bar{y}_{t+1} - \hat{y}_t\|^2 + (1 + \frac{4}{\mu\eta_y}) \|\hat{y}_{t+1} - \hat{y}_t\|^2 \\ &\leq (1 - \frac{\mu\eta_y}{4}) \|\bar{y}_t - \hat{y}_t\|^2 + \frac{9\eta_y}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 + \frac{5\kappa}{L\eta_y} \|\hat{y}_{t+1} - \hat{y}_t\|^2 - \frac{3\eta_y}{5\mu} \|\bar{u}_t\|^2 \\ &\leq (1 - \frac{\mu\eta_y}{4}) \|\bar{y}_t - \hat{y}_t\|^2 + \frac{9\eta_y}{\mu} \|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 + \frac{5\kappa^3\eta_x^2}{L\eta_y} \|\bar{v}_t\|^2 - \frac{3\eta_y}{5\mu} \|\bar{u}_t\|^2 \end{aligned} \quad (32)$$

In the second inequality we use Eq. (31) and $L\eta_y \leq \frac{1}{5}$. The last inequality is because function $y^*(\cdot)$ is κ -Lipschitz. By Cauchy-Schwartz inequality and Assumption 1 we also have

$$\|\nabla_y f(\bar{x}_t, \bar{y}_t) - \bar{u}_t\|^2 \leq 2 \|\bar{u}_t - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_t^{(i)}, y_t^{(i)})\|^2 + \frac{2L^2}{n} (\|X_t - \bar{X}_t\|_F^2 + \|Y_t - \bar{Y}_t\|_F^2) \quad (33)$$

Using the definition of δ_t and the recursion in Eq. (32) we obtain

$$\begin{aligned} \delta_t &\leq (1 - \frac{\mu\eta_y}{4})^t \delta_0 + \frac{9\eta_y}{\mu} \sum_{s=0}^{t-1} (1 - \frac{\mu\eta_y}{4})^{t-s-1} \|\bar{u}_s - \nabla_y f(\bar{x}_s, \bar{y}_s)\|^2 + \frac{5\kappa^3\eta_x^2}{L\eta_y} \sum_{s=0}^{t-1} (1 - \frac{\mu\eta_y}{4})^{t-s-1} \|\bar{v}_s\|^2 \\ &\quad - \frac{3\eta_y}{5\mu} \sum_{s=0}^{t-1} (1 - \frac{\mu\eta_y}{4})^{t-s-1} \|\bar{u}_s\|^2 \end{aligned} \quad (34)$$

Summing above equation we have

$$\begin{aligned} \sum_{t=0}^{T-1} \delta_t &\leq \frac{4\kappa}{L\eta_y} \delta_0 + \frac{18\eta_y}{\mu} \sum_{t=1}^{T-1} (1 - \frac{\mu\eta_y}{4})^{T-t-1} \sum_{s=0}^{t-1} \|\bar{u}_s - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{72\kappa^2}{n} \sum_{t=0}^{T-1} (\|X_t - \bar{X}_t\|_F^2 \\ &\quad + \|Y_t - \bar{Y}_t\|_F^2) + \frac{20\kappa^4\eta_x^2}{L^2\eta_y^2} \sum_{t=0}^{T-1} \|\bar{v}_t\|^2 - \frac{12}{5\mu^2} \sum_{t=0}^{T-1} (1 - (1 - \frac{\mu\eta_y}{4})^{T-t}) \|\bar{u}_t\|^2 \end{aligned} \quad (35)$$

where Eq. (33) is used. \square

Lemma 6. For all $t \in \{0, 1, \dots, T\}$ we have $\bar{v}_t = \bar{g}_t$ and $\bar{u}_t = \bar{h}_t$.

Proof. As matrix W is doubly stochastic, we have

$$\bar{v}_t = \bar{v}_{t-1} + \bar{g}_t - \bar{g}_{t-1} \quad (36)$$

which is equivalent to $\bar{v}_t - \bar{g}_t = \bar{v}_{t-1} - \bar{g}_{t-1}$. Since $\bar{u}_{-1} = \bar{g}_{-1}$, we have $\bar{v}_t = \bar{g}_t$ for all $t \in \{0, 1, \dots, T\}$. Similarly, we have $\bar{u}_t = \bar{h}_t$. \square

Lemma 7. Let A_t, B_t be positive sequences satisfying

$$A_{t+1} \leq (1-c)A_t + B_t \quad (37)$$

for some constant $c \in (0, 1)$. Then for any positive integer T we have

$$\sum_{t=0}^T A_t \leq \frac{1}{c}A_0 + \frac{1}{c} \sum_{t=0}^{T-1} B_t \quad (38)$$

Proof. Using recursion on Eq. (37) we can obtain

$$A_t \leq (1-c)^t A_0 + \sum_{s=0}^{t-1} (1-c)^{t-s-1} B_s \quad (39)$$

for $\forall t \geq 0$. Sum above inequality and we achieve the desired conclusion Eq. (38), where we use the condition A_t, B_t are positive and the fact that $\sum_{t=0}^{\infty} (1-c)^t = \frac{1}{c}$. \square

Lemma 8. We can prove the following bound for gradient estimator \bar{v}_t and \bar{u}_t .

$$\begin{aligned} \sum_{s=0}^{t-1} \mathbb{E} \|\bar{v}_s - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 &\leq \frac{\sigma^2}{n\beta_x b_0} + \frac{2\beta_x \sigma^2 t}{n} + \frac{12L^2}{n^2 \beta_x} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\ &\quad + \frac{6L^2}{n\beta_x} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \end{aligned} \quad (40)$$

$$\begin{aligned} \sum_{s=0}^{t-1} \mathbb{E} \|\bar{u}_s - \frac{1}{n} \sum_{i=1}^n \nabla_y f_i(x_s^{(i)}, y_s^{(i)})\|^2 &\leq \frac{\sigma^2}{n\beta_y b_0} + \frac{2\beta_y \sigma^2 t}{n} + \frac{12L^2}{n^2 \beta_y} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\ &\quad + \frac{6L^2}{n\beta_y} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \end{aligned} \quad (41)$$

for all $t \in \{1, 2, \dots, T\}$.

Proof. By the definition of $g_t^{(i)}$ and Lemma 6 we have

$$\begin{aligned} &\bar{v}_t - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_t^{(i)}, y_t^{(i)}) \\ &= (1-\beta_x)(\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})) + \frac{\beta_x}{n} \sum_{i=1}^n (\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})) \\ &\quad + (1-\beta_x) \frac{1}{n} \sum_{i=1}^n \left(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}) + \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) \right. \\ &\quad \left. - \nabla_x f_i(x_t^{(i)}, y_t^{(i)}) \right) \end{aligned} \quad (42)$$

Taking expectation on $\xi_t^{(i)}$ the last two terms of Eq. (42) are 0. Therefore,

$$\begin{aligned} &\mathbb{E} \|\bar{v}_t - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\ &= (1-\beta_x)^2 \mathbb{E} \|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + \mathbb{E} \left\| \frac{\beta_x}{n} \sum_{i=1}^n (\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})) \right\|^2 \end{aligned}$$

$$\begin{aligned}
& -\nabla_x f_i(x_t^{(i)}, y_t^{(i)}) + (1 - \beta_x) \frac{1}{n} \sum_{i=1}^n \left(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}) \right. \\
& \quad \left. + \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)}) \right) \|^2 \\
& \leq (1 - \beta_x)^2 \mathbb{E} \|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + \frac{2\beta_x^2}{n^2} \sum_{i=1}^n \mathbb{E} \|\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) \\
& \quad - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 + \frac{2(1 - \beta_x)^2}{n^2} \sum_{i=1}^n \mathbb{E} \|\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)}) \\
& \quad + \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& \leq (1 - \beta_x)^2 \mathbb{E} \|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + \frac{2\beta_x^2 \sigma^2}{n} + \frac{2L^2(1 - \beta_x)^2}{n^2} (\mathbb{E} \|X_t - X_{t-1}\|_F^2 \\
& \quad + \mathbb{E} \|Y_t - Y_{t-1}\|_F^2) \tag{43}
\end{aligned}$$

The first inequality is obtained by Cauchy-Schwartz inequality. In the last inequality we use Lemma 2 on the last two terms and then use Assumption 2, Lemma 1 and Assumption 1. By Cauchy-Schwartz inequality we have estimations

$$\|X_t - X_{t-1}\|_F^2 \leq 3\|X_t - \bar{X}_t\|_F^2 + 3n\eta_x^2 \|\bar{v}_{t-1}\|^2 + 3\|X_{t-1} - \bar{X}_{t-1}\|_F^2 \tag{44}$$

$$\|Y_t - Y_{t-1}\|_F^2 \leq 3\|Y_t - \bar{Y}_t\|_F^2 + 3n\eta_y^2 \|\bar{u}_{t-1}\|^2 + 3\|Y_{t-1} - \bar{Y}_{t-1}\|_F^2 \tag{45}$$

Combining above two inequalities with Eq. (43) and Lemma 7 we have

$$\begin{aligned}
& \sum_{s=0}^{t-1} \mathbb{E} \|\bar{v}_s - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 \\
& \leq \frac{1}{\beta_x} \mathbb{E} \|\bar{v}_0 - \nabla_x f(x_0, y_0)\|^2 + \frac{2\beta_x \sigma^2 t}{n} + \frac{12L^2}{n^2 \beta_x} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{6L^2}{n\beta_x} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \\
& \leq \frac{\sigma^2}{n\beta_x b_0} + \frac{2\beta_x \sigma^2 t}{n} + \frac{12L^2}{n^2 \beta_x} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{6L^2}{n\beta_x} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \tag{46}
\end{aligned}$$

for all $t \in \{1, 2, \dots, T\}$. In the first inequality we use the fact $\frac{1}{1 - (1 - \beta_x)^2} \leq \frac{1}{\beta_x}$ when $\beta_x \leq 1$. The second inequality is because $\mathbb{E} \|\bar{v}_0 - \nabla_x f(x_0, y_0)\|^2 \leq \frac{\sigma^2}{nb_0}$ by Assumption 2 and Lemma 2. Note that if we do not use Lemma 7 on the last term we will get

$$\begin{aligned}
& \sum_{s=0}^{t-1} \mathbb{E} \|\bar{v}_s - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 \\
& \leq \frac{\sigma^2}{n\beta_x b_0} + \frac{2\beta_x \sigma^2 t}{n} + \frac{12L^2}{n^2 \beta_x} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{6L^2}{n\beta_x} \sum_{s=0}^{t-2} (1 - (1 - \beta_x)^{t-s-1}) (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \tag{47}
\end{aligned}$$

Mimic above steps we can also prove the second conclusion in Lemma 8. \square

Lemma 9. *The consensus error satisfies the following recursive relation*

$$\|X_{t+1} - \bar{X}_{t+1}\|_F^2 \leq \frac{1 + \lambda^2}{2} \|X_t - \bar{X}_t\|_F^2 + \frac{2\lambda^2\eta_x^2}{1 - \lambda^2} \|V_t - \bar{V}_t\|_F^2 \quad (48)$$

$$\|Y_{t+1} - \bar{Y}_{t+1}\|_F^2 \leq \frac{1 + \lambda^2}{2} \|Y_t - \bar{Y}_t\|_F^2 + \frac{2\lambda^2\eta_y^2}{1 - \lambda^2} \|U_t - \bar{U}_t\|_F^2 \quad (49)$$

Proof. Let $J = \frac{\mathbf{1}\mathbf{1}^T}{n}$. According to the update rule we have

$$\begin{aligned} & \|X_{t+1} - \bar{X}_{t+1}\|_F^2 \\ &= \|(X_t - \eta_x V_t)W - (\bar{X}_t - \eta_x \bar{V}_t)\|_F^2 = \|(X_t - \bar{X}_t)(W - J) - \eta_x(V_t - \bar{V}_t)(W - J)\|_F^2 \\ &\leq \lambda^2 \|X_t - \bar{X}_t\|_F^2 + \lambda^2 \eta_x^2 \|V_t - \bar{V}_t\|_F^2 - 2\langle (X_t - \bar{X}_t)(W - J), \eta_x(V_t - \bar{V}_t)(W - J) \rangle \\ &\leq (\lambda^2 + \theta\lambda^2) \|X_t - \bar{X}_t\|_F^2 + \left(\frac{\lambda^2\eta_x^2}{\theta} + \lambda^2\eta_x^2\right) \|V_t - \bar{V}_t\|_F^2 \\ &\leq \frac{1 + \lambda^2}{2} \|X_t - \bar{X}_t\|_F^2 + \frac{2\lambda^2\eta_x^2}{1 - \lambda^2} \|V_t - \bar{V}_t\|_F^2 \end{aligned} \quad (50)$$

In the first inequality we use Assumption 4 and Lemma 3. In the second inequality we use Young's inequality and θ is an arbitrary positive constant. Let $\theta = \frac{1 - \lambda^2}{2\lambda^2}$ and we can get the last inequality. Similar to Eq. (50), we can obtain the following estimation

$$\begin{aligned} \|Y_{t+1} - \bar{Y}_{t+1}\|_F^2 &= \|(Y_t + \eta_y U_t)W - (\bar{Y}_t + \eta_y \bar{U}_t)\|_F^2 \\ &\leq (\lambda^2 + \theta\lambda^2) \|Y_t - \bar{Y}_t\|_F^2 + \left(\frac{\lambda^2\eta_y^2}{\theta} + \lambda^2\eta_y^2\right) \|U_t - \bar{U}_t\|_F^2 \\ &\leq \frac{1 + \lambda^2}{2} \|Y_t - \bar{Y}_t\|_F^2 + \frac{2\lambda^2\eta_y^2}{1 - \lambda^2} \|U_t - \bar{U}_t\|_F^2 \end{aligned} \quad (51)$$

□

Lemma 10. *For all $t \in \{0, 1, \dots, T - 1\}$ we have*

$$\begin{aligned} \sum_{s=0}^t \mathbb{E} \|V_s - \bar{V}_s\|_F^2 &\leq \frac{2}{1 - \lambda^2} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{48\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\ &\quad + \frac{24n\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2 + \frac{24n\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 \\ &\quad + \frac{8\lambda^2 \beta_x^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \sum_{i=1}^n \mathbb{E} \|g_s^{(i)} - \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{6n\lambda^2 \beta_x^2 \sigma^2 t}{1 - \lambda^2} \end{aligned} \quad (52)$$

$$\begin{aligned} \sum_{s=0}^t \mathbb{E} \|U_s - \bar{U}_s\|_F^2 &\leq \frac{2}{1 - \lambda^2} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 + \frac{48\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\ &\quad + \frac{24n\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2 + \frac{24n\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 \\ &\quad + \frac{8\lambda^2 \beta_y^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t-1} \sum_{i=1}^n \mathbb{E} \|h_s^{(i)} - \nabla_y f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{6n\lambda^2 \beta_y^2 \sigma^2 t}{1 - \lambda^2} \end{aligned} \quad (53)$$

Proof. By the definition of V_t , Assumption 4 and Lemma 3, we have

$$\begin{aligned} & \|V_{t+1} - \bar{V}_{t+1}\|_F^2 \\ &= \|(V_t + G_{t+1} - G_t)W - (\bar{V}_t + \bar{G}_{t+1} - \bar{G}_t)\|_F^2 \\ &= \|(V_t - \bar{V}_t)(W - J) + (G_{t+1} - G_t)(W - J)\|_F^2 \\ &\leq \lambda^2 \|V_t - \bar{V}_t\|_F^2 + \lambda^2 \|G_{t+1} - G_t\|_F^2 + 2\langle (V_t - \bar{V}_t)(W - J), (G_{t+1} - G_t)(W - J) \rangle \end{aligned} \quad (54)$$

Review the definition of $g_t^{(i)}$

$$g_{t+1}^{(i)} - g_t^{(i)} = \nabla_x F_i(x_{t+1}, y_{t+1}; \xi_{t+1}^{(i)}) - \nabla_x F_i(x_t, y_t; \xi_{t+1}^{(i)}) - \beta_x (g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})) + \beta_x (\nabla_x F_i(x_t, y_t; \xi_{t+1}^{(i)}) - \nabla_x f_i(x_t, y_t^{(i)})) \quad (55)$$

and take expectation on $\xi_{t+1}^{(i)}$, then we have

$$\mathbb{E}[g_{t+1}^{(i)} - g_t^{(i)}] = \nabla_x f_i(x_{t+1}, y_{t+1}^{(i)}) - \nabla_x f_i(x_t, y_t^{(i)}) - \beta_x (g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})) \quad (56)$$

Taking expectation on $\xi_{t+1}^{(i)}$ the last term of Eq. (54) can be bounded by

$$\begin{aligned} & \mathbb{E} \langle (V_t - \bar{V}_t)(W - J), (G_{t+1} - G_t)(W - J) \rangle \\ &= \langle (V_t - \bar{V}_t)(W - J), \mathbb{E}[G_{t+1} - G_t](W - J) \rangle \leq \lambda \|V_t - \bar{V}_t\|_F \cdot \lambda \|\mathbb{E}[G_{t+1} - G_t]\|_F \\ &\leq \frac{1 - \lambda^2}{4} \|V_t - \bar{V}_t\|_F^2 + \frac{\lambda^4}{1 - \lambda^2} \|\mathbb{E}[G_{t+1} - G_t]\|_F^2 \\ &\leq \frac{1 - \lambda^2}{4} \|V_t - \bar{V}_t\|_F^2 + \frac{2\lambda^4}{1 - \lambda^2} \sum_{i=1}^n \|\nabla_x f_i(x_{t+1}, y_{t+1}^{(i)}) - \nabla_x f_i(x_t, y_t^{(i)})\|^2 \\ &\quad + \frac{2\lambda^4 \beta_x^2}{1 - \lambda^2} \sum_{i=1}^n \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 \\ &\leq \frac{1 - \lambda^2}{4} \|V_t - \bar{V}_t\|_F^2 + \frac{2\lambda^4 L^2}{1 - \lambda^2} (\|X_{t+1} - X_t\|_F^2 + \|Y_{t+1} - Y_t\|_F^2) \\ &\quad + \frac{2\lambda^4 \beta_x^2}{1 - \lambda^2} \sum_{i=1}^n \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 \end{aligned} \quad (57)$$

where the second inequality is resulted from Young's inequality, the third inequality is resulted from Cauchy-Schwartz inequality and the last inequality is resulted from Assumption 1. Besides, applying Cauchy-Schwartz inequality to Eq. (55) we have

$$\begin{aligned} & \mathbb{E} \|g_{t+1}^{(i)} - g_t^{(i)}\|^2 \\ &\leq 3\mathbb{E} \|\nabla_x F_i(x_{t+1}, y_{t+1}^{(i)}; \xi_{t+1}^{(i)}) - \nabla_x F_i(x_t, y_t^{(i)}; \xi_{t+1}^{(i)})\|^2 + 3\beta_x^2 \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 \\ &\quad + 3\beta_x^2 \mathbb{E} \|\nabla_x F_i(x_t, y_t^{(i)}; \xi_{t+1}^{(i)}) - \nabla_x f_i(x_t, y_t^{(i)})\|^2 \\ &\leq 3L^2 (\mathbb{E} \|x_{t+1}^{(i)} - x_t^{(i)}\|^2 + \mathbb{E} \|y_{t+1}^{(i)} - y_t^{(i)}\|^2) + 3\beta_x^2 \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 + 3\beta_x^2 \sigma^2 \end{aligned} \quad (58)$$

where in the last inequality we use Assumption 1 and Assumption 2. Combining Eq. (54), (57) and (58) we can obtain

$$\begin{aligned} \mathbb{E} \|V_{t+1} - \bar{V}_{t+1}\|_F^2 &\leq \frac{1 + \lambda^2}{2} \mathbb{E} \|V_t - \bar{V}_t\|_F^2 + \frac{4\lambda^2 L^2}{1 - \lambda^2} (\mathbb{E} \|X_{t+1} - X_t\|_F^2 + \mathbb{E} \|Y_{t+1} - Y_t\|_F^2) \\ &\quad + \frac{4\lambda^2 \beta_x^2}{1 - \lambda^2} \sum_{i=1}^n \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 + 3n\lambda^2 \beta_x^2 \sigma^2 \end{aligned} \quad (59)$$

Then using Eq. (44) and (45) in above inequality we have

$$\begin{aligned} & \mathbb{E} \|V_{t+1} - \bar{V}_{t+1}\|_F^2 \\ &\leq \frac{1 + \lambda^2}{2} \mathbb{E} \|V_t - \bar{V}_t\|_F^2 + \frac{12\lambda^2 L^2}{1 - \lambda^2} (\mathbb{E} \|X_{t+1} - \bar{X}_{t+1}\|_F^2 + \mathbb{E} \|Y_{t+1} - \bar{Y}_{t+1}\|_F^2) \\ &\quad + \frac{12\lambda^2 L^2}{1 - \lambda^2} (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2) + \frac{12n\lambda^2 L^2 \eta_y^2}{1 - \lambda^2} \mathbb{E} \|\bar{u}_t\|^2 \\ &\quad + \frac{12n\lambda^2 L^2 \eta_y^2}{1 - \lambda^2} \mathbb{E} \|\bar{v}_t\|^2 + \frac{4\lambda^2 \beta_x^2}{1 - \lambda^2} \sum_{i=1}^n \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t, y_t^{(i)})\|^2 + 3n\lambda^2 \beta_x^2 \sigma^2 \end{aligned} \quad (60)$$

By Lemma 7, we can further achieve

$$\sum_{s=0}^{t'} \mathbb{E} \|V_s - \bar{V}_s\|_F^2 \leq \frac{2}{1 - \lambda^2} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{48\lambda^2 L^2}{(1 - \lambda^2)^2} \sum_{s=0}^{t'} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2)$$

$$\begin{aligned}
& + \frac{24n\lambda^2 L^2}{(1-\lambda^2)^2} \sum_{s=0}^{t'-1} \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2 + \frac{24n\lambda^2 L^2}{(1-\lambda^2)^2} \sum_{s=0}^{t'-1} \eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 \\
& + \frac{8\lambda^2 \beta_x^2}{(1-\lambda^2)^2} \sum_{s=0}^{t'-1} \sum_{i=1}^n \mathbb{E} \|g_s^{(i)} - \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{6n\lambda^2 \beta_x^2 \sigma^2 t'}{1-\lambda^2} \quad (61)
\end{aligned}$$

for all $t' \in \{0, 1, \dots, T-1\}$. Here we should notice that term $\mathbb{E} \|X_{t+1} - \bar{X}_{t+1}\|_F^2$ in Eq. (60) is summed from $\mathbb{E} \|X_1 - \bar{X}_1\|_F^2$ to $\mathbb{E} \|X_{t'} - \bar{X}_{t'}\|_F^2$, while term $\mathbb{E} \|X_t - \bar{X}_t\|_F^2$ is summed from $\mathbb{E} \|X_0 - \bar{X}_0\|_F^2$ to $\mathbb{E} \|X_{t'-1} - \bar{X}_{t'-1}\|_F^2$. As $X_0 = \bar{X}_0$, these two terms can be merged together. And it is the same with term $\mathbb{E} \|Y_{t+1} - \bar{Y}_{t+1}\|_F^2$. Mimic above steps and we can prove the conclusion for $\sum_{s=0}^{t'} \mathbb{E} \|U_s - \bar{U}_s\|_F^2$ in the similar way. \square

Lemma 11. *We can prove the gradient estimators \bar{g}_t and \bar{h}_t satisfy the following conclusion*

$$\begin{aligned}
\sum_{s=0}^t \sum_{i=1}^n \mathbb{E} \|g_s^{(i)} - \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 & \leq \frac{n\sigma^2}{\beta_x b_0} + 2n\beta_x \sigma^2 t + \frac{12L^2}{\beta_x} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& + \frac{6nL^2}{\beta_x} \sum_{s=0}^{t-1} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \quad (62)
\end{aligned}$$

$$\begin{aligned}
\sum_{s=0}^t \sum_{i=1}^n \mathbb{E} \|h_s^{(i)} - \nabla_y f_i(x_s^{(i)}, y_s^{(i)})\|^2 & \leq \frac{n\sigma^2}{\beta_y b_0} + 2n\beta_y \sigma^2 t + \frac{12L^2}{\beta_y} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& + \frac{6nL^2}{\beta_y} \sum_{s=0}^{t-1} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \quad (63)
\end{aligned}$$

for all $t \in \{0, 1, \dots, T-1\}$.

Proof. According to the definition of $g_t^{(i)}$ we have

$$\begin{aligned}
& g_t^{(i)} - \nabla_x f_i(x_t^{(i)}, y_t^{(i)}) \\
& = (1-\beta_x)(g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})) + \beta_x(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})) \\
& + (1-\beta_x)(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)})) \\
& + \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)}) \quad (64)
\end{aligned}$$

The last two terms of Eq. (64) is 0 after taking expectation of $\xi_t^{(i)}$. Hence we have

$$\begin{aligned}
& \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& = (1-\beta_x)^2 \mathbb{E} \|g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + \mathbb{E} \|\beta_x(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})) \\
& + (1-\beta_x)(\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)})) \\
& + \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& \leq (1-\beta_x)^2 \mathbb{E} \|g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + 2\beta_x^2 \mathbb{E} \|\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& + 2(1-\beta_x)^2 \mathbb{E} \|\nabla_x F_i(x_t^{(i)}, y_t^{(i)}; \xi_t^{(i)}) - \nabla_x F_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}; \xi_t^{(i)})\|^2 \\
& \leq (1-\beta_x)^2 \mathbb{E} \|g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + 2\beta_x^2 \sigma^2 + 2(1-\beta_x)^2 L^2 (\mathbb{E} \|x_t^{(i)} - x_{t-1}^{(i)}\|^2 \\
& + \mathbb{E} \|y_t^{(i)} - y_{t-1}^{(i)}\|^2) \quad (65)
\end{aligned}$$

where we use Cauchy-Schwartz inequality and Lemma 1 in the first inequality and use Assumption 1 and Assumption 2 in the last inequality. Sum above inequality from $i = 1$ to n and we have

$$\begin{aligned}
\sum_{i=1}^n \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 & \leq (1-\beta_x)^2 \sum_{i=1}^n \mathbb{E} \|g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + 2n\beta_x^2 \sigma^2 \\
& + 2(1-\beta_x)^2 L^2 (\mathbb{E} \|X_t - X_{t-1}\|^2 + \mathbb{E} \|Y_t - Y_{t-1}\|^2) \quad (66)
\end{aligned}$$

Then by Eq. (44) and (45) we have

$$\begin{aligned}
& \sum_{i=1}^n \mathbb{E} \|g_t^{(i)} - \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& \leq (1 - \beta_x)^2 \sum_{i=1}^n \mathbb{E} \|g_{t-1}^{(i)} - \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + 2n\beta_x^2\sigma^2 \\
& \quad + 6(1 - \beta_x)^2 L^2 (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2 + \mathbb{E} \|X_{t-1} - \bar{X}_{t-1}\|_F^2 \\
& \quad + \mathbb{E} \|Y_{t-1} - \bar{Y}_{t-1}\|_F^2) + 6n(1 - \beta_x)^2 L^2 (\eta_x^2 \mathbb{E} \|\bar{v}_{t-1}\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_{t-1}\|^2)
\end{aligned} \tag{67}$$

Applying Lemma 7 to Eq. (67), similar to Eq. (61), we can obtain

$$\begin{aligned}
& \sum_{s=0}^t \sum_{i=1}^n \mathbb{E} \|g_s^{(i)} - \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 \\
& \leq \frac{1}{\beta_x} \sum_{i=1}^n \mathbb{E} \|g_0^{(i)} - \nabla_x f_i(x_0^{(i)}, y_0^{(i)})\|^2 + \frac{12L^2}{\beta_x} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{6nL^2}{\beta_x} \sum_{s=0}^{t-1} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) + 2n\beta_x\sigma^2 t \\
& \leq \frac{n\sigma^2}{\beta_x b_0} + 2n\beta_x\sigma^2 t + \frac{12L^2}{\beta_x} \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{6nL^2}{\beta_x} \sum_{s=0}^{t-1} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2)
\end{aligned} \tag{68}$$

for all $t \in \{0, 1, \dots, T-1\}$. Here the last inequality is derived by $\mathbb{E} \|g_0^{(i)} - \nabla_x f_i(x_0^{(i)}, y_0^{(i)})\|^2 \leq \frac{\sigma^2}{b_0}$ due to Lemma 2. The estimation of $h_t^{(i)}$ can be achieved in the same way as above. \square

Lemma 12. Let $\eta_x \leq \frac{(1-\lambda)^2}{500L}$ and $\eta_y \leq \frac{(1-\lambda)^2}{500L}$. The consensus error can be bounded by

$$\begin{aligned}
& \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \leq \frac{16\lambda^2\eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{16\lambda^2\eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 + \frac{576n\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 \\
& \quad + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) + \frac{64n\lambda^4 (\beta_x \eta_x^2 + \beta_y \eta_y^2) \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{196n\lambda^4 (\beta_x^2 \eta_x^2 + \beta_y^2 \eta_y^2) \sigma^2 t}{(1-\lambda^2)^4}
\end{aligned} \tag{69}$$

for all $t \in \{0, 1, \dots, T\}$.

Proof. Combining Lemma 7 and Lemma 9, for all $t \in \{0, 1, \dots, T\}$ we have

$$\sum_{s=0}^t \|X_s - \bar{X}_s\|_F^2 \leq \frac{4\lambda^2\eta_x^2}{(1-\lambda^2)^2} \sum_{s=0}^{t-1} \|V_s - \bar{V}_s\|_F^2 \tag{70}$$

Substitute the right side with Lemma 10 we have

$$\begin{aligned}
\sum_{s=0}^t \mathbb{E} \|X_s - \bar{X}_s\|_F^2 & \leq \frac{8\lambda^2\eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{192\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{96n\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2 + \frac{96n\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} \eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 \\
& \quad + \frac{32\lambda^4 \beta_x^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} \sum_{i=1}^n \mathbb{E} \|g_s^{(i)} - \nabla_x f_i(x_s^{(i)}, y_s^{(i)})\|^2 + \frac{24n\lambda^4 \beta_x^2 \eta_x^2 \sigma^2 (t-1)}{(1-\lambda^2)^3}
\end{aligned} \tag{71}$$

Apply Lemma 11 and we get

$$\begin{aligned}
& \sum_{s=0}^t \mathbb{E} \|X_s - \bar{X}_s\|_F^2 \\
& \leq \frac{8\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{192\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{96n\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) + \frac{32n\lambda^4 \beta_x \eta_x^2 \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{64n\lambda^4 \beta_x^3 \eta_x^2 \sigma^2 (t-2)}{(1-\lambda^2)^4} \\
& \quad + \frac{24n\lambda^4 \beta_x^2 \eta_x^2 \sigma^2 (t-1)}{(1-\lambda^2)^3} + \frac{384\lambda^4 \beta_x L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{192n\lambda^4 \beta_x L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-3} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \\
& \leq \frac{8\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{576\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{288n\lambda^4 L^2 \eta_x^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) + \frac{32n\lambda^4 \beta_x \eta_x^2 \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{98n\lambda^4 \beta_x^2 \eta_x^2 \sigma^2 t}{(1-\lambda^2)^4} \tag{72}
\end{aligned}$$

where we use $\beta_x \leq 1$ to simplify the equation. Similarly, we have

$$\begin{aligned}
& \sum_{s=0}^t \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2 \\
& \leq \frac{8\lambda^2 \eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 + \frac{576\lambda^4 L^2 \eta_y^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \quad + \frac{288n\lambda^4 L^2 \eta_y^2}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) + \frac{32n\lambda^4 \beta_y \eta_y^2 \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{98n\lambda^4 \beta_y^2 \eta_y^2 \sigma^2 t}{(1-\lambda^2)^4} \tag{73}
\end{aligned}$$

Add Eq. (72) and (73) together. Then we have

$$\begin{aligned}
& \sum_{s=0}^t (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) \\
& \leq \frac{8\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{8\lambda^2 \eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 + \frac{576\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \sum_{s=0}^{t-1} (\mathbb{E} \|X_s - \bar{X}_s\|_F^2 \\
& \quad + \mathbb{E} \|Y_s - \bar{Y}_s\|_F^2) + \frac{288n\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E} \|\bar{v}_s\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_s\|^2) \\
& \quad + \frac{32n\lambda^4 (\beta_x \eta_x^2 + \beta_y \eta_y^2) \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{98n\lambda^4 (\beta_x^2 \eta_x^2 + \beta_y^2 \eta_y^2) \sigma^2 t}{(1-\lambda^2)^4} \tag{74}
\end{aligned}$$

As $\lambda < 1$, when $\eta_x \leq \frac{(1-\lambda)^2}{500L}$ and $\eta_y \leq \frac{(1-\lambda)^2}{500L}$ it satisfies

$$\frac{576\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \leq \frac{1}{2} \tag{75}$$

Therefore, Eq. (74) implies

$$\begin{aligned}
& \sum_{s=0}^t (\mathbb{E}\|X_s - \bar{X}_s\|_F^2 + \mathbb{E}\|Y_s - \bar{Y}_s\|_F^2) \\
& \leq \frac{16\lambda^2\eta_x^2}{(1-\lambda^2)^3} \mathbb{E}\|V_0 - \bar{V}_0\|_F^2 + \frac{16\lambda^2\eta_y^2}{(1-\lambda^2)^3} \mathbb{E}\|U_0 - \bar{U}_0\|_F^2 + \frac{576n\lambda^4L^2(\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \sum_{s=0}^{t-2} (\eta_x^2 \mathbb{E}\|\bar{v}_s\|^2 \\
& \quad + \eta_y^2 \mathbb{E}\|\bar{u}_s\|^2) + \frac{64n\lambda^4(\beta_x\eta_x^2 + \beta_y\eta_y^2)\sigma^2}{(1-\lambda^2)^4b_0} + \frac{196n\lambda^4(\beta_x^2\eta_x^2 + \beta_y^2\eta_y^2)\sigma^2t}{(1-\lambda^2)^4} \tag{76}
\end{aligned}$$

which reaches the conclusion of Lemma 12. \square

A.3 Proof for main Theorems

Now we will move forward to the main Theorems in our paper. Here we revise some constant coefficients in the statement, but it does not actually affect the result in our convergence analysis.

Theorem 3. (Restatement of Theorem 1) *Let Assumptions 1 to 5 hold. When parameters $\beta_x = \frac{\epsilon \min\{1, n\epsilon\}}{20}$, $\beta_y = \frac{\epsilon \min\{1, n\epsilon\}}{500\kappa^2}$, $\eta_x = \frac{(1-\lambda)^2 \min\{1, n\epsilon\}}{15000\kappa^3L}$, $\eta_y = \frac{(1-\lambda)^2 \min\{1, n\epsilon\}}{1500\kappa L}$, $b_0 = \frac{400\kappa}{\min\{1, n\epsilon\}}$, $T = \frac{30000\kappa^3\epsilon^{-2}}{(1-\lambda)^2 \min\{1, n\epsilon\}}$, our Algorithm 1 satisfies*

$$\begin{aligned}
\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}\|\nabla\Phi(\bar{x}_t)\|^2 & \leq L(\Phi(x_0) - \Phi^*)\epsilon^2 + \sigma^2\epsilon^2 + L^2\delta_0\epsilon^2 + \frac{\epsilon^2}{n} \sum_{i=1}^n \|\nabla_x f_i(x_0, y_0)\|^2 \\
& \quad + \frac{\epsilon^2}{n} \sum_{i=1}^n \|\nabla_y f_i(x_0, y_0)\|^2 \tag{77}
\end{aligned}$$

Proof. Since $\Phi(x)$ is $(\kappa L + L)$ -smooth we have

$$\begin{aligned}
\Phi(\bar{x}_t) & \leq \Phi(\bar{x}_{t-1}) - \eta_x \langle \bar{v}_{t-1}, \nabla\Phi(\bar{x}_{t-1}) \rangle + \eta_x^2 \kappa L \|\bar{v}_{t-1}\|^2 \\
& = \Phi(\bar{x}_{t-1}) - \frac{\eta_x}{2} \|\bar{v}_{t-1}\|^2 - \frac{\eta_x}{2} \|\nabla\Phi(\bar{x}_{t-1})\|^2 + \frac{\eta_x}{2} \|\bar{v}_{t-1} - \nabla\Phi(\bar{x}_{t-1})\|^2 + \eta_x^2 \kappa L \|\bar{v}_{t-1}\|^2 \\
& \leq \Phi(\bar{x}_{t-1}) - \frac{\eta_x}{2} \|\nabla\Phi(\bar{x}_{t-1})\|^2 - \left(\frac{\eta_x}{2} - \eta_x^2 \kappa L\right) \|\bar{v}_{t-1}\|^2 + \eta_x \|\bar{v}_{t-1} - \nabla_x f(\bar{x}_{t-1}, \bar{y}_{t-1})\|^2 \\
& \quad + \eta_x \|\nabla\Phi(\bar{x}_{t-1}) - \nabla_x f(\bar{x}_{t-1}, \bar{y}_{t-1})\|^2 \tag{78}
\end{aligned}$$

where the last inequality is caused by Cauchy-Schwartz inequality. As we have $\nabla\Phi(\bar{x}_{t-1}) = \nabla_x f(\bar{x}_{t-1}, \hat{y}_{t-1})$, by Assumption 1 the last term satisfies

$$\|\nabla\Phi(\bar{x}_{t-1}) - \nabla_x f(\bar{x}_{t-1}, \bar{y}_{t-1})\|^2 \leq L^2 \|\hat{y}_{t-1} - \bar{y}_{t-1}\|^2 = L^2 \delta_{t-1} \tag{79}$$

Besides, according to Cauchy-Schwartz inequality we also have

$$\begin{aligned}
& \|\bar{v}_{t-1} - \nabla_x f(\bar{x}_{t-1}, \bar{y}_{t-1})\|^2 \\
& \leq 2\|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + 2\|\frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)}) - \nabla_x f(\bar{x}_{t-1}, \bar{y}_{t-1})\|^2 \\
& \leq 2\|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 + \frac{2L^2}{n} (\|X_{t-1} - \bar{X}_{t-1}\|_F^2 + \|Y_{t-1} - \bar{Y}_{t-1}\|_F^2) \tag{80}
\end{aligned}$$

Combine Eq. (78), (79), (80) and rearrange the inequality

$$\begin{aligned}
\|\nabla\Phi(\bar{x}_{t-1})\|^2 & \leq \frac{2(\Phi(\bar{x}_{t-1}) - \Phi(\bar{x}_t))}{\eta_x} - (1 - 2\kappa L\eta_x) \|\bar{v}_{t-1}\|^2 + 2L^2\delta_{t-1} + \frac{4L^2}{n} (\|X_{t-1} - \bar{X}_{t-1}\|_F^2 \\
& \quad + \|Y_{t-1} - \bar{Y}_{t-1}\|_F^2) + 4\|\bar{v}_{t-1} - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_{t-1}^{(i)}, y_{t-1}^{(i)})\|^2 \tag{81}
\end{aligned}$$

Telescoping and taking expectation on Eq. (81) we have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \mathbb{E}\Phi(\bar{x}_T))}{\eta_x T} - \frac{(1 - 2\kappa L \eta_x)}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 + \frac{2L^2}{T} \sum_{t=0}^{T-1} \mathbb{E} \delta_t + \frac{4L^2}{nT} \sum_{t=0}^{T-1} (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 \\
& \quad + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2) + \frac{4}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2
\end{aligned} \tag{82}$$

Applying Assumption 3, Lemma 5 and Lemma 8 we have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} - (1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2}) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 + \frac{8\kappa L^2 \delta_0}{TL \eta_y} \\
& \quad + \frac{148\kappa^2 L^2}{nT} \sum_{t=0}^{T-1} (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2) + \frac{4}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t - \frac{1}{n} \sum_{i=1}^n \nabla_x f_i(x_t^{(i)}, y_t^{(i)})\|^2 \\
& \quad + \frac{36\kappa L \eta_y}{T} \sum_{t=1}^{T-1} (1 - \frac{\mu \eta_y}{4})^{T-t-1} \sum_{s=0}^{t-1} \|\bar{u}_s - \frac{1}{n} \sum_{i=1}^n \nabla f_i(x_s^{(i)}, y_s^{(i)})\|^2 \\
& \quad - \frac{24\kappa^2}{5T} \sum_{t=0}^{T-1} (1 - (1 - \frac{\mu \eta_y}{4})^{T-t}) \mathbb{E} \|\bar{u}_t\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} - (1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2}) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 + \frac{8\kappa L^2 \delta_0}{TL \eta_y} + \frac{4\sigma^2}{nb_0 T} (\frac{1}{\beta_x} + \frac{36\kappa^2}{\beta_y}) \\
& \quad + \frac{8\sigma^2}{n} (\beta_x + 36\kappa^2 \beta_y) + \frac{4L^2}{nT} (47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y}) \sum_{t=0}^{T-1} (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2) \\
& \quad + \frac{24L^2}{n\beta_x T} \sum_{t=0}^{T-1} (1 - (1 - \beta_x)^{T-t}) (\eta_x^2 \mathbb{E} \|\bar{v}_t\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_t\|^2) + \frac{864\kappa^2 L^2}{n\beta_y T} \sum_{t=0}^{T-1} (1 - (1 - \frac{\mu \eta_y}{4})^{T-t}) \\
& \quad \cdot (\eta_x^2 \mathbb{E} \|\bar{v}_t\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_t\|^2) - \frac{24\kappa^2}{5T} \sum_{t=0}^{T-1} (1 - (1 - \frac{\mu \eta_y}{4})^{T-t}) \mathbb{E} \|\bar{u}_t\|^2
\end{aligned} \tag{83}$$

where we use Eq. (47) in the last inequality. As

$$\frac{1}{\beta_x} (1 - (1 - \beta_x)^{T-t}) = \sum_{s=0}^{T-t-1} (1 - \beta_x)^s \tag{84}$$

we know Eq. (84) is increasing when β_x is decreasing. Hence $\frac{1}{\beta_x} (1 - (1 - \beta_x)^{T-t}) \leq \frac{300\kappa^2}{(1-\lambda)^2 \beta_x} (1 - (1 - \frac{(1-\lambda)^2 \beta_x}{300\kappa^2})^{T-t})$. According to the definition of β_x and η_y , we have $\frac{(1-\lambda)^2 \beta_x}{300\kappa^2} \leq \frac{\mu \eta_y}{4}$ and

$$\frac{24L^2}{n\beta_x T} (1 - (1 - \beta_x)^{T-t}) \leq \frac{7200L^2 \kappa^2}{n(1-\lambda)^2 \beta_x T} (1 - (1 - \frac{\mu \eta_y}{4})^{T-t}) \tag{85}$$

Therefore, using the definition of β_x , β_y and η_y we obtain

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} - (1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2}) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 + \frac{8\kappa L^2 \delta_0}{TL \eta_y} + \frac{4\sigma^2}{nb_0 T} \left(\frac{1}{\beta_x} + \frac{36\kappa^2}{\beta_y} \right) \\
& \quad + \frac{8\sigma^2}{n} (\beta_x + 36\kappa^2 \beta_y) + \frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \sum_{t=0}^{T-1} (\mathbb{E} \|X_t - \bar{X}_t\|_F^2 + \mathbb{E} \|Y_t - \bar{Y}_t\|_F^2) \\
& \quad + \left(\frac{24L^2 \eta_x^2}{n\beta_x} + \frac{864\kappa^2 L^2 \eta_x^2}{n\beta_y} \right) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 - \frac{\kappa L \eta_y}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{u}_t\|^2 \tag{86}
\end{aligned}$$

Besides, according to Lemma 12 we have

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} - (1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2}) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 + \frac{8\kappa L^2 \delta_0}{TL \eta_y} + \frac{4\sigma^2}{nb_0 T} \left(\frac{1}{\beta_x} + \frac{36\kappa^2}{\beta_y} \right) \\
& \quad + \frac{8\sigma^2}{n} (\beta_x + 36\kappa^2 \beta_y) + \frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \left(\frac{16\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 \right. \\
& \quad \left. + \frac{16\lambda^2 \eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 + \frac{64n\lambda^4 (\beta_x \eta_x^2 + \beta_y \eta_y^2) \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{196n\lambda^4 (\beta_x^2 \eta_x^2 + \beta_y^2 \eta_y^2) \sigma^2 T}{(1-\lambda^2)^4} \right) \\
& \quad + \frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \frac{576n\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \sum_{t=0}^{T-1} (\eta_x^2 \mathbb{E} \|\bar{v}_t\|^2 + \eta_y^2 \mathbb{E} \|\bar{u}_t\|^2) \\
& \quad + \left(\frac{24L^2 \eta_x^2}{n\beta_x} + \frac{864\kappa^2 L^2 \eta_x^2}{n\beta_y} \right) \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{v}_t\|^2 - \frac{\kappa L \eta_y}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\bar{u}_t\|^2 \tag{87}
\end{aligned}$$

When β_x , β_y , η_x and η_y are defined as Theorem 3, we have

$$\frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \frac{576n\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \eta_y^2 \leq \frac{\kappa L \eta_y}{2T} \tag{88}$$

and

$$\begin{aligned}
& 1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2} - \frac{24L^2 \eta_x^2}{n\beta_x} - \frac{864\kappa^2 L^2 \eta_x^2}{n\beta_y} \\
& \quad - \frac{4L^2}{n} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \frac{576n\lambda^4 L^2 (\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \eta_x^2 \geq \frac{2}{5} \tag{89}
\end{aligned}$$

Thus, we obtain

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} + \frac{8\kappa L^2 \delta_0}{TL \eta_y} + \frac{4\sigma^2}{nb_0 T} \left(\frac{1}{\beta_x} + \frac{36\kappa^2}{\beta_y} \right) + \frac{8\sigma^2}{n} (\beta_x + 36\kappa^2 \beta_y) \\
& \quad + \frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \left(\frac{16\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{16\lambda^2 \eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 \right. \\
& \quad \left. + \frac{64n\lambda^4 (\beta_x \eta_x^2 + \beta_y \eta_y^2) \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{196n\lambda^4 (\beta_x^2 \eta_x^2 + \beta_y^2 \eta_y^2) \sigma^2 T}{(1-\lambda^2)^4} \right) \tag{90}
\end{aligned}$$

By Assumption 4 and Cauchy-Schwartz inequality we also have

$$\mathbb{E}\|V_0 - \bar{V}_0\|_F^2 = \mathbb{E}\|G_0(W - J)\|_F^2 \leq \lambda^2 \mathbb{E}\|G_0\|_F^2 \leq \frac{2n\lambda^2\sigma^2}{b_0} + 2\lambda^2 \sum_{i=1}^n \|\nabla_x f_i(x_0, y_0)\|^2 \quad (91)$$

Similarly, we have

$$\mathbb{E}\|U_0 - \bar{U}_0\|_F^2 \leq \frac{2n\lambda^2\sigma^2}{b_0} + 2\lambda^2 \sum_{i=1}^n \|\nabla_y f_i(x_0, y_0)\|^2 \quad (92)$$

Combine above three inequalities and substitute the parameters with their definitions. We achieve

$$\begin{aligned} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}\|\nabla\Phi(\bar{x}_t)\|^2 &\leq L(\Phi(x_0) - \Phi^*)\epsilon^2 + L^2\delta_0\epsilon^2 + \sigma^2\epsilon^2 + \frac{\epsilon^2}{n} \sum_{i=1}^n \|\nabla_x f_i(x_0, y_0)\|^2 \\ &\quad + \frac{\epsilon^2}{n} \sum_{i=1}^n \|\nabla_y f_i(x_0, y_0)\|^2 \end{aligned} \quad (93)$$

where we use following inequalities for simplification.

$$\begin{aligned} \beta_x \geq \beta_y, \quad \frac{144\kappa^2}{n\beta_y b_0 T} &\leq \frac{144\kappa^2 \cdot 500\kappa^2 (\min\{1, n\epsilon\})^2 \epsilon^2}{n\epsilon \min\{1, n\epsilon\} 400\kappa \cdot 30000\kappa^3} \leq \frac{3\epsilon^2}{500} \\ 4L^2(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y}) &\leq 200L^2\kappa^2 + \frac{1800L^2\kappa^2}{n\beta_y} \\ \frac{8\beta_x}{n} \leq \frac{8\epsilon \cdot n\epsilon}{20n} = \frac{2\epsilon^2}{5}, \quad \frac{288\kappa^2\beta_y}{n} &\leq \frac{288\kappa^2\epsilon \cdot n\epsilon}{500n\kappa^2} \leq \frac{288\epsilon^2}{500} \\ \frac{L^2\beta_x\eta_x^2}{(1-\lambda)^4 b_0 T} &\leq \frac{\epsilon(\min\{1, n\epsilon\})^5 \epsilon^2}{20 \cdot 400\kappa \cdot 30000\kappa^3 (15000\kappa^3)^2}, \quad \frac{L^2\beta_y\eta_y^2}{(1-\lambda)^4 b_0 T} \leq \frac{\epsilon(\min\{1, n\epsilon\})^5 \epsilon^2}{500 \cdot 400\kappa \cdot 30000\kappa^3 (1500\kappa)^2} \\ \frac{L^2\beta_x^2\eta_x^2}{(1-\lambda)^4} &\leq \frac{\epsilon^2(\min\{1, n\epsilon\})^4}{400(15000\kappa^3)^2}, \quad \frac{L^2\beta_y^2\eta_y^2}{(1-\lambda)^4} \leq \frac{\epsilon^2(\min\{1, n\epsilon\})^4}{(500\kappa^2)^2(1500\kappa)^2} \end{aligned} \quad (94)$$

□

Theorem 4. (Restatement of Theorem 2) Let Assumptions 1 to 5 hold. We set the parameters as $T = \frac{30000\kappa^3 T_0}{(1-\lambda)^2}$, $\beta_x = \frac{n^{1/3}}{20T_0^{2/3}}$, $\beta_y = \frac{n^{1/3}}{500\kappa^2 T_0^{2/3}}$, $\eta_x = \frac{(1-\lambda)^2 n^{2/3}}{15000\kappa^3 T_0^{1/3} L}$, $\eta_y = \frac{(1-\lambda)^2 n^{2/3}}{1500\kappa T_0^{1/3} L}$, $b_0 = \frac{400\kappa T_0^{1/3}}{n^{2/3}}$, where we suppose $T_0 \geq 10n^2$. Then our algorithm satisfies

$$\begin{aligned} \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E}\|\nabla\Phi(\bar{x}_t)\|^2 &\leq \frac{L(\Phi(x_0) - \Phi^*) + \sigma^2 + L^2\delta_0}{(nT_0)^{2/3}} + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E}\|\nabla_x f_i(x_0, y_0)\|^2}{T_0} \\ &\quad + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E}\|\nabla_y f_i(x_0, y_0)\|^2}{T_0} \end{aligned} \quad (95)$$

Proof. When the parameters are defined as Theorem 4, the conditions in Lemma 5 and Lemma 12 are also satisfied. Hence we can prove Eq. (83) and (87) still hold. When β_x , β_y , η_x and η_y are defined as Theorem 4, we also have

$$\frac{4L^2}{nT} (47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y}) \frac{576n\lambda^4 L^2(\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \eta_y^2 \leq \frac{\kappa L \eta_y}{2T} \quad (96)$$

and

$$\begin{aligned} 1 - 2\kappa L \eta_x - \frac{40\kappa^4 \eta_x^2}{\eta_y^2} - \frac{24L^2 \eta_x^2}{n\beta_x} - \frac{864\kappa^2 L^2 \eta_x^2}{n\beta_y} \\ - \frac{4L^2}{n} (47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y}) \frac{576n\lambda^4 L^2(\eta_x^2 + \eta_y^2)}{(1-\lambda^2)^4} \eta_x^2 \geq \frac{2}{5} \end{aligned} \quad (97)$$

Similar to Theorem 3, we can also obtain

$$\begin{aligned}
& \frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 \\
& \leq \frac{2(\Phi(x_0) - \Phi^*)}{\eta_x T} + \frac{8\kappa L^2 \delta_0}{TL\eta_y} + \frac{4\sigma^2}{nb_0 T} \left(\frac{1}{\beta_x} + \frac{36\kappa^2}{\beta_y} \right) + \frac{8\sigma^2}{n} (\beta_x + 36\kappa^2 \beta_y) \\
& \quad + \frac{4L^2}{nT} \left(47\kappa^2 + \frac{12}{n\beta_x} + \frac{432\kappa^2}{n\beta_y} \right) \left(\frac{16\lambda^2 \eta_x^2}{(1-\lambda^2)^3} \mathbb{E} \|V_0 - \bar{V}_0\|_F^2 + \frac{16\lambda^2 \eta_y^2}{(1-\lambda^2)^3} \mathbb{E} \|U_0 - \bar{U}_0\|_F^2 \right) \\
& \quad + \frac{64n\lambda^4 (\beta_x \eta_x^2 + \beta_y \eta_y^2) \sigma^2}{(1-\lambda^2)^4 b_0} + \frac{196n\lambda^4 (\beta_x^2 \eta_x^2 + \beta_y^2 \eta_y^2) \sigma^2 T}{(1-\lambda^2)^4} \tag{98}
\end{aligned}$$

Substitute the parameters with their definitions and we have

$$\begin{aligned}
\frac{1}{T} \sum_{t=0}^{T-1} \mathbb{E} \|\nabla \Phi(\bar{x}_t)\|^2 & \leq \frac{L(\Phi(x_0) - \Phi^*) + \sigma^2 + L^2 \delta_0}{(nT_0)^{2/3}} + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_x f_i(x_0, y_0)\|^2}{T_0} \\
& \quad + \frac{\frac{1}{n} \sum_{i=1}^n \mathbb{E} \|\nabla_y f_i(x_0, y_0)\|^2}{T_0} \tag{99}
\end{aligned}$$

which achieves the conclusion of Theorem 4. \square