The Many Faces of Adversarial Risk

Abstract

Adversarial risk quantifies the performance of classifiers on adversarially perturbed data. Numerous definitions of adversarial risk—not all mathematically rigorous and differing subtly in the details—have appeared in the literature. In this paper, we revisit these definitions, make them rigorous, and critically examine their similarities and differences. Our technical tools derive from optimal transport, robust statistics, functional analysis, and game theory. Our contributions include the following: generalizing Strassen's theorem to the unbalanced optimal transport setting with applications to adversarial classification with unequal priors; showing an equivalence between adversarial robustness and robust hypothesis testing with ∞-Wasserstein uncertainty sets; proving the existence of a pure Nash equilibrium in the two-player game between the adversary and the algorithm; and characterizing adversarial risk by the minimum Bayes error between a pair of distributions belonging to the ∞-Wasserstein uncertainty sets. Our results generalize and deepen recently discovered connections between optimal transport and adversarial robustness and reveal new connections to Choquet capacities and game theory.

1 Introduction

Neural networks are known to be vulnerable to *adversarial attacks*, which are imperceptible perturbations to input data that maximize loss [39, 15, 5]. Developing algorithms resistant to such attacks has received considerable attention in recent years [8, 30, 25, 20], motivated by safety-critical applications such as autonomous driving [18, 28], medical imaging [17, 24, 22] and law [21, 6].

A classification algorithm with high accuracy (low risk) in the absence of an adversary may have poor accuracy (high risk) when an adversary is present. Thus, a modified notion known as *adversarial risk* is used to quantify the adversarial robustness of algorithms. Algorithms that minimize adversarial risk are deemed robust. Procedures for finding them have been effective in practice [25, 42, 29], spurring numerous theoretical investigations into adversarial risk and its minimizers.

There is no universally agreed upon definition of adversarial risk. Even the simplest setting of binary classification in $\mathbb{R}^d$ with an ℓ_2 adversary admits various definitions involving set expansions [9, 16], transport maps [31], Markov kernels [32], and couplings [27]. These works broadly interpret adversarial risk as a measure of robustness to small perturbations, but their definitions differ in subtle details such as the class of adversaries and algorithms considered, budget constraints placed on the adversary, assumptions on the loss function, and the geometry of decision boundaries.

Optimal adversarial risk is most commonly defined as the minimax risk under adversarial contamination [25, 34]. Other notable characterizations include an optimal transport cost between data generating distributions in [32, 2, 10, 11], the optimal value of a distributionally robust optimization problem [37, 36, 41], and the value of a two-player zero-sum game [27, 31, 3, 4].

The diversity of definitions for adversarial risk makes it challenging to compare approaches. Moreover, not all approaches are rigorous. For instance, the classes of adversarial strategies and classifier

algorithms are often unclear, and issues of measurability are ignored. Although this may be harmless for applied research, it has led to incorrect proofs and insufficient assumptions in some theoretical works; a mathematically rigorous foundation for adversarial risk is essential for future research.

In this paper, we examine various notions of adversarial risk for binary classification. We present rigorous definitions of adversarial risk and identify conditions under which these definitions are equivalent. We consider the general setting of Polish spaces (complete, separable metric spaces), and present stronger results for the Euclidean space ($\mathbb{R}^d$). Our contributions are as follows:

- We examine the definition of adversarial risk based on set expansions. For Polish spaces, we observe that adversarial risk is not well-defined when the decision region is an arbitrary Borel measurable set. We show well-definedness when the decision region is both F_σ and G_δ (classes of measurable sets; see section 2.1). For the Euclidean space with the Lebesgue σ-algebra, we show that adversarial risk is well-defined for any Lebesgue measurable decision region. Our key lemma (Lemma 4.2) shows that the Lebesgue σ-algebra is preferred over the Borel σ-algebra because set expansions are Lebesgue measurable but not necessarily Borel measurable. These results are contained in Section 4.

- We show that the definition of adversarial risk using set expansions is identical to a notion of risk that appears in robust hypothesis testing with ∞-Wasserstein uncertainty sets. We prove this result in Polish spaces using the theory of measurable selections for multi-valued mappings [44]. In $\mathbb{R}^d$, we are able to use the powerful theory of Choquet capacities [7] (in particular, Huber and Strassen's 2-alternating capacities [19]) to establish results of a similar nature. These results are contained in Section 5.

- We consider the binary classification setup with unequal priors and show (under suitable assumptions) that the optimal adversarial risk from the above definitions is characterized by an unbalanced optimal transport cost between data-generating distributions. For both Polish spaces and $\mathbb{R}^d$, the main tool we use is Theorem 6.1 in which we generalize a classical result of Strassen on excess-cost optimal transport [38, 43] from probability measures to finite measures with possibly unequal mass. This generalizes results of [32, 2] on binary classification, which were only for equal priors. These results are contained in Section 6.

- We consider the setup of a zero-sum game between the adversary and the algorithm. We show that the value of this game (adversarial risk) is equal to the minimum Bayes error between a pair of distributions belonging to the ∞-Wasserstein uncertainty sets centered around true data-generating distributions. We prove the existence of a pure Nash equilibrium in this game for $\mathbb{R}^d$ and for Polish spaces with a *midpoint property*. This extends/strengthens the results of [27, 31, 3] to non-parametric classifiers. These results are contained in Section 7.

The paper is organized as follows: In Section 2, we present preliminary definitions from optimal transport and metric space topology. In Section 3, we discuss various definitions of adversarial risk and present more related work. Sections 4, 5, 6 and 7 contain our main contributions summarized above. We conclude the paper in Section 8 and discuss future research directions.

Notation: Throughout the paper, we use $\mathcal{X}$ to denote a Polish space (a complete, separable metric space) with metric d and Borel sigma algebra $\mathcal{B}(\mathcal{X})$. For $x \in \mathcal{X}$ and $r \geq 0$, let $B_r(x)$ denote the closed ball of radius r centered at x. We use $\mathcal{P}(\mathcal{X})$ and $\mathcal{M}(\mathcal{X})$ to denote the space of probability measures and finite measures defined on the measure space $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$, respectively. For $\mu, \nu \in \mathcal{M}(\mathcal{X})$, we say ν *dominates* μ if $\mu(A) \leq \nu(A)$ for all $A \in \mathcal{B}(\mathcal{X})$ and write $\mu \preceq \nu$. When $\mathcal{X}$ is $\mathbb{R}^d$, we use $\mathcal{L}(\mathcal{X})$ to denote the Lebesgue σ-algebra and λ to denote the d-dimensional Lebesgue measure. We use $\overline{\mathcal{P}}(\mathcal{X})$ to denote the set of probability measures defined over the complete measure space $(\mathcal{X}, \mathcal{L}(\mathcal{X}))$. For a positive integer n, we use $[n]$ to denote the finite set $\{1, \ldots, n\}$.

2 Preliminaries

2.1 Metric Space Topology

We introduce three different notions of set expansions. For $\epsilon \geq 0$ and $A \in \mathcal{B}(\mathcal{X})$, the ϵ-*Minkowski expansion* of A is given by $A^{\oplus \epsilon} := \cup_{a \in A} B_\epsilon(a)$. The ϵ-*closed expansion* of A is defined as $A^\epsilon := \{x \in \mathcal{X} : d(x, A) \leq \epsilon\}$, where $d(x, A) = \inf_{a \in A} d(x, a)$. The ϵ-*open expansion* of A is defined

as $A^{\epsilon)} := \{x \in \mathcal{X} : d(x, A) < \epsilon\}$. We use the notation $A^{-\epsilon}$ to denote $((A^c)^\epsilon)^c$. Similarly, $A^{\ominus\epsilon} := ((A^c)^{\oplus\epsilon})^c$. For example, consider the set $A = (0, 1]$ in the space $(\mathcal{X}, d) = (\mathbb{R}, |\cdot|)$ and $\epsilon > 0$. Then $A^{\oplus\epsilon} = (-\epsilon, 1 + \epsilon]$, $A^\epsilon = [-\epsilon, 1 + \epsilon]$ and $A^{\epsilon)} = (-\epsilon, 1 + \epsilon)$. For any $A \in \mathcal{B}(\mathcal{X})$, A^ϵ is closed and $A^{\epsilon)}$ is open. Hence, $A^\epsilon, A^{\epsilon)} \in \mathcal{B}(\mathcal{X})$. Moreover, $A^{\epsilon)} \subseteq A^{\oplus\epsilon} \subseteq A^\epsilon$. However, $A^{\oplus\epsilon}$ may not be in $\mathcal{B}(\mathcal{X})$ (see Example 2.4 in [23]). In general, the Minkowski sum of two Borel sets need not be Borel [13], and that of two Lebesgue measurable sets need not be Lebesgue measurable [35].

2.2 Optimal Transport

Let $\mu, \nu \in \mathcal{P}(\mathcal{X})$. A *coupling* between μ and ν is a joint probability measure $\pi \in \mathcal{P}(\mathcal{X}^2)$ with marginals μ and ν. The set $\Pi(\mu, \nu) \subseteq \mathcal{P}(\mathcal{X}^2)$ denotes the set of all couplings between μ and ν. The *optimal transport cost* between μ and ν under a cost function $c : \mathcal{X} \times \mathcal{X} \to [0, \infty)$ is defined as $T_c(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X}^2} c(x, x') d\pi(x, x')$. For a positive integer p, the *p-Wasserstein distance* between μ and ν is defined as, $W_p(\mu, \nu) = (T_{d^p}(\mu, \nu))^{\frac{1}{p}}$. The ∞-*Wasserstein metric* is defined as $W_\infty(\mu, \nu) = \lim_{p \to \infty} W_p(\mu, \nu)$. It can also be expressed in the following ways:

$$W_\infty(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \operatorname*{ess\,sup}_{(x, x') \sim \pi} d(x, x') = \inf\{\delta > 0 : \mu(A) \leq \nu(A^\delta) \forall A \in \mathcal{B}(\mathcal{X})\}. \tag{1}$$

Given a $\mu \in \mathcal{P}(\mathcal{X})$ and a measurable function $f : \mathcal{X} \to \mathcal{X}$, the *push-forward* of μ by f is defined as a probability measure $f_{\sharp\mu} \in \mathcal{P}(\mathcal{X})$ given by $f_{\sharp\mu} = \mu(f^{-1}(A))$ for all $A \in \mathcal{B}(\mathcal{X})$.

3 Adversarial Risk: Definitions and Related Work

We consider a binary classification setting on feature space $\mathcal{X}$. Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ be the data-generating distributions for labels 0 and 1, respectively. Let the prior probabilities for labels 0 and 1 be in the ratio $T : 1$ where we assume $T \geq 1$ without loss of generality. For any set $A \in \mathcal{B}(\mathcal{X})$, we may consider a classifier $f_A(x) := \mathbb{1}\{x \in A\}$ which labels any point in the set A as 1 and any point in A^c as 0. We say that such a classifier has a decision region A. The error (standard risk) incurred by such a classifier under the 0-1 loss function is, $R_0(\ell_{0/1}, A) = \frac{T}{T+1} p_0(A) + \frac{1}{T+1} p_1(A^c)$.

An adversary of budget $\epsilon > 0$ can perturb any $x \in \mathcal{X}$ to $x' \in B_\epsilon(x)$. It follows that any $x \in A$ can be perturbed to $x' \in \cup_{a \in A} B_\epsilon(a) = A^{\oplus\epsilon}$. Hence, adversarial risk could be defined as

$$R_{\oplus\epsilon}(\ell_{0/1}, A) = \frac{T}{T+1} p_0(A^{\oplus\epsilon}) + \frac{1}{T+1} p_1((A^c)^{\oplus\epsilon}). \tag{2}$$

The above formulation has been studied in [9, 16], and is a special case of the widely used expected supremum formulation of adversarial risk for general loss functions [25, 34]. A problem with this formulation is the ambiguity over the measurability of the sets $A^{\oplus\epsilon}$ and $(A^c)^{\oplus\epsilon}$. Even when $A \in \mathcal{B}(\mathcal{X})$, it is not guaranteed that $A^{\oplus\epsilon}, (A^c)^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})$ [23, Example 2.4]. Hence, $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is not well-defined for all $A \in \mathcal{B}(\mathcal{X})$. It is shown in [32] that $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined when A is either closed or open, but its validity beyond that is unknown.

An alternate formulation [26] replaces the Minkowski set expansion by closed set expansion:

$$R_\epsilon(\ell_{0/1}, A) = \frac{T}{T+1} p_0(A^\epsilon) + \frac{1}{T+1} p_1((A^c)^\epsilon). \tag{3}$$

The above definition is well-defined for any $A \in \mathcal{B}(\mathcal{X})$ because A^ϵ and $(A^c)^\epsilon$ are both closed and hence, measurable. However, under the above definition, a point $x \in A$ may be perturbed to $x' \in A^\epsilon$ such that $d(x, x') > \epsilon$. For example, when $A = (0, 1)$, we have $A^\epsilon = [-\epsilon, \epsilon]$ and an adversary may transport $x = \delta > 0$ to $x' = -\epsilon$, violating the budget constraint at x.

Another approach to defining adversarial risk is by explicitly defining the class of adversaries of budget ϵ as measurable transport maps $f : \mathcal{X} \to \mathcal{X}$ that push-forward the true data distribution such that no point is transported by more than a distance of ϵ; i.e., $d(x, f(x)) \leq \epsilon$. The transport map-based adversarial risk [31] is formally defined as follows:

$$R_{F_\epsilon}(\ell_{0/1}, A) = \sup_{\substack{f_0, f_1 : \mathcal{X} \to \mathcal{X} \\ \forall x \in \mathcal{X}, d(x, f_i(x)) \leq \epsilon}} \frac{T}{T+1} f_{0\sharp p_0}(A) + \frac{1}{T+1} f_{1\sharp p_1}((A^c)). \tag{4}$$

128 Yet another definition uses the robust hypothesis framework with W_∞ uncertainty sets. In this
129 approach, an adversary perturbs the true distribution p_i to a corrupted distribution p_i' such that
130 $W_\infty(p_i, p_i') \le \epsilon$. From (1), this is equivalent to the existence of a coupling $\pi \in \Pi(p_i, p_i')$ such that
131 $\operatorname{ess\,sup}_{(x,x')\sim\pi} d(x,x') \le \epsilon$. The adversarial risk with such an adversary is given by

$$R_{\Gamma_\epsilon}(\ell_{0/1}, A) = \sup_{W_\infty(p_1, p_1'), W_\infty(p_0, p_0') \le \epsilon} \frac{T}{T+1} p_0'(A) + \frac{1}{T+1} p_1'((A^c)). \tag{5}$$

132 Clearly, $R_{F_\epsilon}(\ell_{0/1}, A) \le R_{\Gamma_\epsilon}(\ell_{0/1}, A)$, but conditions for equality were not studied in prior work.
133 Moreover, their relation to set expansion-based definitions in (2) and (3) was also unknown.

134 Next we discuss some characterizations of optimal adversarial risk. In [32, 2], it is shown that
135 $R_\epsilon^* := \inf_{A \in \mathcal{B}(\mathcal{X})} R_\epsilon(\ell_{0/1}, A) = \frac{1}{2}[1 - D_\epsilon(p_0, p_1)]$ for equal priors ($T = 1$), where D_ϵ is an optimal
136 transport cost defined as follows.

137 **Definition 1** (D_ϵ cost). Let $\mu, \nu \in \mathcal{P}(\mathcal{X})$. Let $\epsilon \ge 0$. Let $c_\epsilon : \mathcal{X}^2 \to \{0,1\}$ be such that $c_\epsilon(x, x') =$
138 $\mathbb{1}\{(x, x') \in \mathcal{X} \times \mathcal{X} : d(x, x') > 2\epsilon\}$. Then for $\mu, \nu \in \mathcal{P}(\mathcal{X})$ and $\epsilon \ge 0$, $D_\epsilon(\mu, \nu) = T_{c_\epsilon}(\mu, \nu)$.

139 For $\epsilon = 0$, D_ϵ reduces to the total variation distance. While D_0 is a metric on $\mathcal{P}(\mathcal{X})$, D_ϵ (for $\epsilon > 0$)
140 is neither a metric nor a pseudometric [32].

141 Other formulations of optimal adversarial risk are inspired from game theory [31, 27, 3]. Consider a
142 game between two players: (1) The adversary whose action space is pairs of distributions $p_0', p_1' \in$
143 $\mathcal{P}(\mathcal{X})$, and (2) The algorithm whose action space is the space of decision regions of the form
144 $A \in \mathcal{A} := \{A \in \mathcal{B}(\mathcal{X}) : A^{\ominus\epsilon}, A^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})\}$ (or a subset thereof). For $T > 0$, define $r :$
145 $\mathcal{A} \times \mathcal{P}(\mathcal{X}) \times \mathcal{P}(\mathcal{X}) \to [0,1]$ as, $r(A, \mu, \nu) = \frac{T}{T+1}\mu(A) + \frac{1}{T+1}\nu((A^c))$. The payoff when the
146 algorithm plays first is given by $\inf_{A \in \mathcal{A}} \sup_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \le \epsilon} r(A, p_0', p_1')$, and this quantity is
147 interpreted as the optimal adversarial risk in this setup.

4 Well-Definedness of Adversarial Risk

149 As stated in Section 3, $R_{\oplus\epsilon}(\ell_{0/1}, A)$ may not be well-defined for some decision regions $A \in \mathcal{B}(\mathcal{X})$
150 because of the non-measurability of the sets $A^{\oplus\epsilon}$ and $(A^c)^{\oplus\epsilon}$. In this section, we lay down the
151 conditions under which the ambiguity can be resolved.

152 We begin by introducing the concept of G_δ and F_σ sets in a topological space. A subset of the
153 topological space $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$ is called a G_δ set if it is a countable intersection of open sets, and an
154 F_σ set if it is a countable union of closed sets. Note that the complement of a G_δ set is a F_σ set and
155 vice versa. The following lemma shows that Minkowski set expansions of F_σ sets are measurable. It
156 generalizes a similar result of [32] for closed and open sets.

157 **Lemma 4.1.** Let $A \in \mathcal{B}(\mathcal{X})$. If A is an F_σ set, then $A^{\oplus\epsilon}$ is also an F_σ set and hence $A^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})$.

158 In the following theorem, we use Lemma 4.1 to show the well-definedness of $R_{\oplus\epsilon}(\ell_{0/1}, A)$.

159 **Theorem 4.1.** Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$. Let $A \in \mathcal{B}(\mathcal{X})$ be G_δ and F_σ. Then $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined.

160 *Proof.* Since A is a G_δ set, A^c is a F_σ set. By Lemma 4.1, both $A^{\oplus\epsilon}$ and $(A^c)^{\oplus\epsilon}$ are F_σ sets and
161 hence measurable. Therefore $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined. $\qquad\square$

162 In a Polish space, every closed set is G_δ and every open set is F_σ [1]. Hence, any countable union of
163 closed or open sets is both G_δ and F_σ. Theorem 4.1 establishes that $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined for
164 a much broader class of decision regions than closed and open sets, strengthening the result of [32].

165 Now, we will show that if $\mathcal{X} = \mathbb{R}^d$, then $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined for any Lebesgue measurable
166 set A as long as the data generating distributions are defined on the complete measure space $(\mathcal{X}, \mathcal{L}(\mathcal{X}))$
167 instead of the default Borel measure space $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$. For this, we use the concept of porous sets.

168 **Definition 2** (Porous set). A set $E \subseteq \mathcal{X}$ is said to be porous if there exists $\alpha \in (0,1)$ and $r_0 > 0$
169 such that for every $r \in (0, r_0]$ and every $x \in \mathcal{X}$, there is an $x' \in \mathcal{X}$ such that $B_{\alpha r}(x') \subseteq B_r(x) \backslash E$.

170 Porous sets are a subclass of nowhere dense sets. Importantly, $\lambda(E) = 0$ for any porous set $E \subseteq \mathbb{R}^d$
171 [48]. By the following lemma, the set difference between the closed/open set expansions is porous.

172 **Lemma 4.2.** Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$ and $A \in \mathcal{L}(\mathcal{X})$. Then $E = A^\epsilon \backslash A^{\epsilon)}$ is porous.

173 Lemma 4.2 plays a crucial role in proving that $A^{\oplus \epsilon} \in \mathcal{L}(\mathcal{X})$ whenever $A \in \mathcal{L}(\mathcal{X})$. We recall that
174 $A^{\oplus \epsilon}$ is the Minkowski sum of A with the closed ϵ-ball. In general, the Minkowski sum of two
175 Lebesgue measurable sets is not always Lebesgue measurable [35, 14]. So the fact that one of them
176 is a closed ball in case of $A^{\oplus \epsilon}$ is important. In the following theorem, we use Lemma 4.2 to prove the
177 measurability of $A^{\oplus \epsilon}$ and in turn prove that $R_{\oplus \epsilon}(\ell_{0/1}, A)$ is well-defined for any $A \in \mathcal{L}(\mathcal{X})$.

178 **Theorem 4.2.** Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$. Let $p_0, p_1 \in \overline{\mathcal{P}}(\mathcal{X})$ and let $\epsilon \geq 0$. Then for any $A \in \mathcal{L}(\mathcal{X})$,
179 $R_{\oplus \epsilon}(\ell_{0/1}, A)$ is well-defined. If, in addition, p_0 and p_1 are absolutely continuous with respect to the
180 Lebesgue measure, then $R_{\oplus \epsilon}(\ell_{0/1}, A) = R_\epsilon(\ell_{0/1}, A)$.

181 *Proof.* By Lemma 4.2 $A^\epsilon \backslash A^{\epsilon)}$ is porous, and so $\lambda(A^\epsilon \backslash A^{\epsilon)}) = 0$. Hence, $\lambda(A^\epsilon) = \lambda(A^{\epsilon)})$. Using
182 the fact that $A^{\epsilon)} \subseteq A^{\oplus \epsilon} \subseteq A^\epsilon$, we have $A^{\oplus \epsilon} \backslash A^{\epsilon)} \subseteq A^\epsilon \backslash A^{\epsilon)}$. Hence, $\lambda(A^{\oplus \epsilon} \backslash A^{\epsilon)}) = 0$. Therefore,
183 $A^{\oplus \epsilon} \in \mathcal{L}(\mathcal{X})$ and $\lambda(A^{\oplus \epsilon}) = \lambda(A^\epsilon) = \lambda(A^{\epsilon)})$.

184 Since $A^{\oplus \epsilon}, (A^c)^{\oplus \epsilon} \in \mathcal{L}(\mathcal{X})$, $R_{\oplus \epsilon}(\ell_{0/1}, A)$ is well-defined. If p_0 and p_1 are absolutely continuous
185 with respect to the Lebesgue measure, the equation $R_{\oplus \epsilon}(\ell_{0/1}, A) = R_\epsilon(\ell_{0/1}, A)$ follows from the
186 previous conclusion that $\lambda(A^{\oplus \epsilon}) = \lambda(A^\epsilon)$. $\qquad\square$

187 Now that we have established the conditions for which $R_{\oplus \epsilon}(\ell_{0/1}, A)$ is well-defined, in the next
188 section, we explore its relation to other notions of adversarial risk.

5 Equivalence with ∞-Wasserstein Robustness

190 In this section, we show the conditions under which $R_{\oplus \epsilon}(\ell_{0/1}, A)$ is equivalent to other notions of
191 adversarial risk based on transport maps and W_∞ robustness. In subsection 5.1, we consider general
192 Polish spaces and in subsection 5.2, we consider the Euclidean space.

5.1 W_∞ Robustness in Polish Spaces via Measurable Selections

194 We begin by presenting a lemma that links the measure of ϵ-Minkowsi set expansion to the worst
195 case measure over a W_∞ probability ball of radius ϵ.

196 **Lemma 5.1.** Let $\mu \in \mathcal{P}(\mathcal{X})$. If $A \in \mathcal{B}(\mathcal{X})$ is both a G_δ set and an F_σ set, then
197 $\sup_{W_\infty(\mu,\mu') \leq \epsilon} \mu'(A) = \mu(A^{\oplus \epsilon})$. Moreover, the supremum in the previous equation is achieved by
198 a $\mu^* \in \mathcal{P}(\mathcal{X})$ that is induced from μ via a measurable transport map $f : \mathcal{X} \to \mathcal{X}$ (i.e. $\mu^* = f_{\sharp \mu}$)
199 satisfying $d(x, f(x)) \leq \epsilon$ for all $x \in \mathcal{X}$.

200 The proof of Lemma 5.1 proceeds by defining a multivalued mapping of the action space of the
201 adversary given by $x \mapsto B_\epsilon(x) \cap A$, and then using the Kuratowski-Ryll-Nardzewski measurable
202 selection theorem ([44]) to find a measurable transport map. We need the set A in Lemma 5.1 to
203 be G_δ for the measurable selection theorem to be applicable. We need A to be F_σ for $A^{\oplus \epsilon}$ to be
204 measurable (by Lemma 4.1) so that the term $\mu(A^{\oplus \epsilon})$ is well-defined.

205 In the following theorem, we use Lemma 5.1 to establish the equivalence between three different
206 notions of adversarial risk introduced in section 3 for the class of sets that are both G_δ set and F_σ.

207 **Theorem 5.1.** Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$. Let $A \in \mathcal{B}(\mathcal{X})$ be both a G_δ set and a F_σ set. Then
208 $R_{\oplus \epsilon}(\ell_{0/1}, A) = R_{F_\epsilon}(\ell_{0/1}, A) = R_{\Gamma_\epsilon}(\ell_{0/1}, A)$. In addition, the supremum over f_0 and f_1 in
209 $R_{F_\epsilon}(\ell_{0/1}, A)$ is attained. Similarly, the supremum over p'_0 and p'_1 in $R_{\Gamma_\epsilon}(\ell_{0/1}, A)$ is attained.

5.2 W_∞ Robustness in $\mathbb{R}^d$ via 2-Alternating Capacities

211 In this subsection, we extend Theorem 5.1 from sets that are both G_δ set and F_σ set to the much
212 broader class of Lebesgue measurable sets in $\mathbb{R}^d$ using the theory of Choquet capacities [7]. We
213 begin with the following definitions.

214 **Definition 3** (Capacity). A set function $v : \mathcal{B}(\mathcal{X}) \to [0, 1]$ is a *capacity* if it satisfies the following
215 conditions: (1) $v(\varnothing) = 0$ and $v(\mathcal{X}) = 1$; (2) For $A, B \in \mathcal{B}(\mathcal{X})$, $A \subseteq B \implies v(A) \leq v(B)$; (3)
216 $A_n \uparrow A \implies v(A_n) \uparrow v(A)$; and (4) $F_n \downarrow F, F_n$ closed $\implies v(F_n) \downarrow v(F)$.

Definition 4 (2-Alternating Capacity). A capacity v defined on the measure space $(\mathcal{X}, \mathcal{B}(\mathcal{X}))$ is called 2-alternating if $v(A \cup B) + v(A \cap B) \leq v(A) + v(B)$ for all $A, B \in \mathcal{B}(\mathcal{X})$.

For any compact set of probability measures $\Xi \subseteq \mathcal{P}(\mathcal{X})$, the upper probability $v(A) = \sup_{\mu \in \Xi} \mu(A)$ is a capacity [19]. The upper probability of ϵ-neighborhoods of a $\mu \in \mathcal{P}(\mathcal{X})$ defined using either the total variation metric or the Levy-Prokhorov metric can be shown to be a 2-alternating capacity [19]. The following lemma shows that $A \mapsto \mu(A^{\oplus \epsilon})$ is a 2-alternating capacity under some conditions.

Lemma 5.2. Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$. Let $\mu \in \overline{\mathcal{P}}(\mathcal{X})$ and let $\epsilon \geq 0$. Define a set function v on $\mathcal{X}$ such that for any $A \in \mathcal{L}(\mathcal{X})$, $v(A) := \mu(A^{\oplus \epsilon})$. Then v is a 2-alternating capacity.

Now we relate the capacity defined in Lemma 5.2 to the W_∞ metric. Since the ϵ-neighborhood of a $\mu \in \mathcal{P}(\mathcal{X})$ in W_∞ metric is a compact set of probability measures [47], the upper probability over this W_∞ ϵ-ball is a capacity. The following lemma shows that it is a 2-alternating capacity, and identifies it with the capacity defined in Lemma 5.2.

Lemma 5.3. Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$. Let $\mu \in \overline{\mathcal{P}}(\mathcal{X})$. Then for any $A \in \mathcal{L}(\mathcal{X})$, $\sup_{W_\infty(\mu, \mu') \leq \epsilon} \mu'(A) = \mu(A^{\oplus \epsilon})$. Moreover, the supremum in the previous equation is attained.

Lemma 5.3 plays a similar role to Lemma 5.1 in proving the following equivalence between adversarial robustness and W_∞ robustness.

Theorem 5.2. Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$. Let $p_0, p_1 \in \overline{\mathcal{P}}(\mathcal{X})$ and let $\epsilon \geq 0$. Then for any $A \in \mathcal{L}(\mathcal{X})$, $R_{\oplus \epsilon}(\ell_{0/1}, A) = R_{\Gamma_\epsilon}(\ell_{0/1}, A)$, and the supremum over p_0' and p_1' in $R_{\Gamma_\epsilon}(\ell_{0/1}, A)$ is attained.

The proof follows by converting the expression for R_{Γ_ϵ} into one for $R_{\oplus \epsilon}$ using Lemma 5.3. Unlike Theorem 5.1, Theorem 5.2 does not show the equivalence of $R_{F_\epsilon}(\ell_{0/1}, A)$ with the other definitions under the relaxed assumption of $A \in \mathcal{L}(\mathcal{X})$. This is because Lemma 5.3 does not provide a push-forward map f such that $\mu^* = f_{\sharp \mu}$ with μ^* attaining the supremum over the W_∞ ball.

6 Optimal Adversarial Risk via Generalized Strassen's Theorem

In section 5, we analyzed adversarial risk for a specific decision region $A \in \mathcal{B}(\mathcal{X})$. In this section, we analyze infimum of adversarial risk over all possible decision regions; i.e., the optimal adversarial risk. We show that optimal adversarial risk in binary classification with unequal priors is characterized by an unbalanced optimal transport cost between data-generating distributions. Our main technical lemma generalizes Strassen's theorem to unbalanced optimal transport. We present this result in subsection 6.1 and present our characterization of optimal adversarial risk in subsection 6.2.

6.1 Unbalanced Optimal Transport & Generalized Strassen's Theorem

Recall from Section 3 that the optimal transport cost D_ϵ characterizes the optimal adversarial risk in binary classification for equal priors. The following result gives an alternative characterization of D_ϵ.

Proposition 6.1 (Strassen's theorem). [Corollary 1.28 in [43]] Let $\mu, \nu \in \mathcal{P}(\mathcal{X})$. Let $\epsilon \geq 0$. Then

$$\sup_{A \in \mathcal{B}(\mathcal{X})} \mu(A) - \nu(A^{2\epsilon}) = D_\epsilon(\mu, \nu). \tag{6}$$

Proposition 6.1 is a special case of Kantorovich-Rubinstein duality [43] applied to $\{0, 1\}$-valued cost functions. We now generalize this result to measures with unequal masses. We begin with some definitions that generalize the concepts we introduced in subsection 2.2.

Let $\mu, \nu \in \mathcal{M}(\mathcal{X})$ be such that $\mu(\mathcal{X}) \leq \nu(\mathcal{X})$. A *coupling* between μ and ν is a measure $\pi \in \mathcal{M}(\mathcal{X}^2)$ such that for any $A \in \mathcal{B}(\mathcal{X})$, $\pi(A \times \mathcal{X}) = \mu(A)$ and $\pi(\mathcal{X} \times A) \leq \nu(A)$. The set $\Pi(\mu, \nu)$ is defined to be the set of all couplings between μ and ν. For a cost function $c : \mathcal{X}^2 \to [0, \infty)$, the optimal transport cost between μ and ν under c is defined as $T_c(\mu, \nu) = \inf_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X}^2} c(x, x') d\pi(x, x')$.

Theorem 6.1 (Generalized Strassen's theorem). Let $\mu, \nu \in \mathcal{M}(\mathcal{X})$ be such that $0 < M = \mu(\mathcal{X}) \leq \nu(\mathcal{X})$. Let $\epsilon > 0$. Define $c_\epsilon : \mathcal{X}^2 \to \{0, 1\}$ as $c_\epsilon(x, x') = \mathbb{1}\{(x, x') \in \mathcal{X}^2 : d(x, x') > 2\epsilon\}$. Then

$$\sup_{A \in \mathcal{B}(\mathcal{X})} \mu(A) - \nu(A^{2\epsilon}) = T_{c_\epsilon}(\mu, \nu) = M \inf_{\nu' \in \mathcal{P}(\mathcal{X}): \nu' \preceq \nu/M} D_\epsilon(\mu/M, \nu'). \tag{7}$$

Moreover, the infimum on the right hand side is attained. (Equivalently, there is a coupling $\pi \in \Pi(\mu, \nu)$ that attains the unbalanced optimal transport cost $T_{c_\epsilon}(\mu, \nu)$.)

Our proof of Theorem 6.1 leverages strong duality in linear programming. We first establish (7) for discrete measures on a finite support. We then apply the discrete result on a sequence of measures supported on a countable dense subset of the Polish space $\mathcal{X}$. Using the tightness of finite measures on $\mathcal{X}$, we construct an optimal coupling that achieves the cost $T_{c_\epsilon}(\mu, \nu)$ in (7). We then show that the constructed coupling satisfies (7). This proof strategy is adapted from the works of [12] and [33].

6.2 Optimal Adversarial Risk for Unequal Priors

Generalized Strassen's theorem involves closed set expansions. The following lemma allows us to switch to Minkowski set expansions.

Lemma 6.1. Let $\mu, \nu \in \mathcal{M}(\mathcal{X})$ and let $\epsilon \geq 0$. Then $\sup_{A \in \mathcal{B}(\mathcal{X})} \mu(A) - \nu(A^{2\epsilon}) = \sup_{A:\ A^{\ominus\epsilon}, A^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})} \mu(A^{\ominus\epsilon}) - \nu(A^{\oplus\epsilon})$. Moreover, the supremum in the right hand side of the above equality can be replaced by a supremum over closed sets.

Using the above lemma and the generalized Strassen's theorem, we show the following result on optimal adversarial risk for unequal priors, generalizing the result of [32, 2].

Theorem 6.2. Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ and let $\epsilon \geq 0$. Then

$$\inf_{A:\ A^{\ominus\epsilon}, A^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})} R_{\oplus\epsilon}(\ell_{0/1}, A) = \frac{1}{T+1}\left[1 - \inf_{q \in \mathcal{P}(\mathcal{X}): q \preceq Tp_0} D_\epsilon(q, p_1)\right]. \tag{8}$$

Moreover, the infimum on the left hand side can be replaced by an infimum over closed sets.

The proof follows by using Lemma 6.1 to convert the expression with Minkowski expansion to one with closed expansions, followed by an application of Theorem 6.1 to arrive at the final optimal transport-based expression.

The above theorem extends the result of [32] in two ways: (1) the infimum is taken over all sets for which $R_{\oplus\epsilon}(\ell_{0/1}, A)$ is well-defined, instead of restricting to closed sets, and (2) the priors on both labels can be unequal. We also note that for $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$, if $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ in Theorem 6.2 are in $\overline{\mathcal{P}}(\mathcal{X})$, then (8) holds with the infimum on the left hand side taken over all $A \in \mathcal{L}(\mathcal{X})$.

7 Minimax Theorems and Nash Equilibria

In this section, we revisit the zero-sum game between the adversary and the algorithm introduced in section 3. Recall that for $A \in \mathcal{A} := \{A \in \mathcal{B}(\mathcal{X}) : A^{\ominus\epsilon}, A^{\oplus\epsilon} \in \mathcal{B}(\mathcal{X})\}$ and $p_0', p_1' \in \mathcal{P}(\mathcal{X})$, the payoff function is given by

$$r(A, p_0', p_1') = \frac{T}{T+1}p_0'(A) + \frac{1}{T+1}p_1'((A^c)). \tag{9}$$

The max-min inequality gives us

$$\sup_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} \inf_{A \in \mathcal{A}} r(A, p_0', p_1') \leq \inf_{A \in \mathcal{A}} \sup_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} r(A, p_0', p_1'). \tag{10}$$

If the inequality in (10) is an equality, we say that the game has zero duality gap, and admits a value equal to either expression in (10). Then there is no advantage to a player making the first move. Our minimax theorems establish such an equality. If in addition to having an equality in (10), there exist $p_0^*, p_1^* \in \mathcal{P}(\mathcal{X})$ that achieve the supremum on the left-hand side and $A^* \in \mathcal{A}$ that achieves the infimum on the right-hand side, we say that $((p_0^*, p_1^*), A^*)$ is a pure Nash equilibrium of the game.

In Section 7.1, we prove the minimax theorem and the existence of a pure Nash equilibrium in $\mathbb{R}^d$ using the theory of 2-alternating capacities [19] and the relation to adversarial risk from Section 5.2. Section 7.2 extends these results to more general Polish spaces with a "midpoint property."

7.1 Minimax Theorem in $\mathbb{R}^d$ via 2-Alternating Capacities

The following theorem proves the minimax equality and the existence of a Nash equilibrium for the adversarial robustness game in $\mathbb{R}^d$.

Theorem 7.1 (Minimax theorem in $\mathbb{R}^d$)**.** Let $(\mathcal{X}, d) = (\mathbb{R}^d, \|\cdot\|)$. Let $p_0, p_1 \in \overline{\mathcal{P}}(\mathcal{X})$ and let $\epsilon \geq 0$. Define r as in (9). Then,

$$\sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} \inf_{A \in \mathcal{L}(\mathcal{X})} r(A, p_0', p_1') = \inf_{A \in \mathcal{L}(\mathcal{X})} \sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} r(A, p_0', p_1'). \quad (11)$$

Moreover, there exist $p_0^*, p_1^* \in \overline{\mathcal{P}}(\mathcal{X})$ and $A^* \in \mathcal{A}$ that achieve the supremum and infimum on the left and right hand sides of the above equation.

Proof. By Lemma 5.2, the set-valued maps $A \mapsto p_0(A^{\oplus \epsilon})$ and $A^c \mapsto p_1((A^c)^{\oplus \epsilon})$ are 2-alternating capacities. Hence, the existence of $A^* \in \mathcal{L}(\mathcal{X})$ that attains the infimum on the right in (11) follows from Lemma 3.1 in [19] and the equality $R_{\oplus \epsilon}(\ell_{0/1}, A) = R_{\Gamma_\epsilon}(\ell_{0/1}, A)$ proved in Theorem 5.2. By Theorem 4.1 in [19], there exist $q_0, q_1 \in \mathcal{P}(\mathcal{X})$ such that $W_\infty(p_i, q_i) \leq \epsilon$ for $i = 0, 1$ and,

$$\inf_{A \in \mathcal{L}(\mathcal{X})} \sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} r(A, p_0, p_1) = \inf_{A \in \mathcal{L}(\mathcal{X})} r(A, q_0, q_1).$$

Hence,

$$\inf_{A \in \mathcal{L}(\mathcal{X})} \sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} r(A, p_0, p_1) = \inf_{A \in \mathcal{A}} r(A, q_0, q_1)$$
$$\leq \sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} \inf_{A \in \mathcal{L}(\mathcal{X}} r(A, p_0, p_1).$$

The desired result follows from combining the above inequality with the max-min inequality (10). Clearly, $q_0 = p_0^*$ and $q_1 = p_1^*$. □

The above proof technique is not applicable in general Polish spaces because the map $A \mapsto \mu(A^{\oplus \epsilon})$ is not a capacity for a general $\mu \in \mathcal{P}(\mathcal{X})$. This is because $A^{\oplus \epsilon}$ is not measurable for all $A \in \mathcal{B}(\mathcal{X})$.

7.2 Minimax Theorem in Polish Spaces via Optimal Transport

We now extend the minimax theorem from $\mathbb{R}^d$ to general Polish spaces with the following property.

Definition 5 (Midpoint property)**.** A metric space $(\mathcal{X}, d)$ is said to have the midpoint property if for every $x_1, x_2 \in \mathcal{X}$, there exists $x \in \mathcal{X}$ such that, $d(x_1, x) = d(x, x_2) = d(x_1, x_2)/2$.

Any normed vector space with distance defined as $d(x, x') = \|x - x'\|$ satisfies the midpoint property. An example of a metric space without this property is the discrete metric space where $d(x, x') = \mathbb{1}\{x \neq x'\}$. The midpoint property plays a crucial role in proving the following theorem, which shows that the D_ϵ transport cost between two distributions is the shortest total variation distance between their ϵ-neighborhoods in W_∞ metric. A similar result was also presented in [11].

Theorem 7.2 (D_ϵ as shortest D_{TV} between W_∞ balls)**.** Let $(\mathcal{X}, d)$ have the midpoint property. Let $\mu, \nu \in \mathcal{P}(\mathcal{X})$ and let $\epsilon \geq 0$. Then $D_\epsilon(\mu, \nu) = \inf_{W_\infty^\epsilon(\mu, \mu'), W_\infty^\epsilon(\nu, \nu') \leq \epsilon} D_{TV}(\mu', \nu')$. Moreover, the infimum over D_{TV} in the above equation is attained.

The following theorem uses Theorem 7.2 to prove the minimax equality and the existence of a Nash equilibrium for any Polish space with the midpoint property for the case of equal priors.

Theorem 7.3 (Minimax theorem for equal priors)**.** Let $(\mathcal{X}, d)$ have the midpoint property. Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ and let $\epsilon \geq 0$. Let $\mathcal{A}$ denote the set of all subsets of $\mathcal{X}$ that are both F_σ and G_δ. Define r as in (9) with $T = 1$. Then

$$\sup_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} \inf_{A \in \mathcal{A}} r(A, p_0', p_1') = \inf_{A \in \mathcal{A}} \sup_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} r(A, p_0', p_1'). \quad (12)$$

Moreover, there exist $p_0^*, p_1^* \in \mathcal{P}(\mathcal{X})$ and $A^* \in \mathcal{A}$ that achieve the supremum and infimum on the left and right hand sides of the above equation.

Proof. For $\mu \in \mathcal{P}(\mathcal{X})$, let $WB(\mu)$ denote the set of all $\mu' \in \mathcal{P}(\mathcal{X})$ such that $W_\infty(\mu, \mu') \leq \epsilon$.

$$\inf_{A \in \mathcal{A}} \sup_{\substack{p_0' \in WB(p_0) \\ p_1' \in WB(p_1)}} r(A, p_0', p_1') = \inf_{A \in \mathcal{A}} R_{\Gamma_\epsilon}(\ell_{0/1}, A) \overset{(i)}{=} \inf_{A \in \mathcal{A}} R_{\oplus \epsilon}(\ell_{0/1}, A) \overset{(ii)}{=} \frac{1}{2}\left[1 - D_\epsilon(p_0, p_1)\right]$$

$$\sup_{\substack{p_0' \in WB(p_0) \\ p_1' \in WB(p_1)}} \inf_{A \in \mathcal{A}} r(A, p_0', p_1') \overset{(iv)}{=} \sup_{\substack{p_0' \in WB(p_0) \\ p_1' \in WB(p_1)}} \frac{1}{2}\left[1 - D_{TV}(p_0', p_1')\right] = \frac{1}{2}\left[1 - \inf_{\substack{p_0' \in WB(p_0) \\ p_1' \in WB(p_1)}} D_{TV}(p_0', p_1')\right],$$

where (i) follows from Theorem 5.1, (ii) from Theorem 6.2, and (iii) again from Theorem 6.2 with $\epsilon = 0$. The expressions on the right extremes of the above equations are equal by Theorem 7.2. The existence of $A^* \in \mathcal{A}$ and $p_0^*, p_1^* \in \mathcal{P}(\mathcal{X})$ follow from Theorem 6.2 and Theorem 7.2 respectively. $\qquad \square$

To prove the minimax theorem for unequal priors, we need the following generalization of Theorem 7.2 to finite measures of unequal mass.

Lemma 7.1. Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ and let $\epsilon \geq 0$. Then for $T \geq 1$,

$$
\begin{aligned}
\inf_{q \in \mathcal{P}(\mathcal{X}): q \preceq Tp_0} D_\epsilon(q, p_1) &= \inf_{q \in \mathcal{P}(\mathcal{X}): q \preceq Tp_0} \inf_{W_\infty^\epsilon(q,q'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} D_{TV}(q', p_1') \\
&= \inf_{W_\infty^\epsilon(p_0, p_0'), W_\infty^\epsilon(p_1, p_1') \leq \epsilon} \inf_{q' \in \mathcal{P}(\mathcal{X}): q' \preceq Tp_0'} D_{TV}(q', p_1') \quad (13)
\end{aligned}
$$

Now, we prove the minimax equality for unequal priors.

Theorem 7.4 (Minimax theorem for unequal priors)**.** Let $(\mathcal{X}, d)$ have the midpoint property. Let $p_0, p_1 \in \mathcal{P}(\mathcal{X})$ and let $\epsilon \geq 0$. Let $\mathcal{A}$ denote the set of all subsets of $\mathcal{X}$ that are both F_σ and G_δ. For $T > 0$, define r as in (9). Then

$$
\sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} \inf_{A \in \mathcal{A}} r(A, p_0', p_1') = \inf_{A \in \mathcal{A}} \sup_{p_0' \in W_\infty^\epsilon(p_0), p_1' \in W_\infty^\epsilon(p_1)} r(A, p_0', p_1'). \quad (14)
$$

The proof of Theorem 7.4 uses the following ingredients: (1) the characterization of inf-sup payoff in terms of unbalanced optimal transport using Theorem 5.1; (2) Lemma 7.1; and (3) the minimax equality of Theorem 7.3 for equal priors.

8 Discussion

In this paper, we rigorously examine different notions of adversarial risk in a binary classification setting with 0-1 loss function, and lay down the conditions under which these definitions are equivalent. By verifying the conditions presented in Sections 4 and 5, researchers can now fluidly move from one definition of adversarial risk to another. A variety of definitions have also been proposed for adversarial risk under general loss functions [32, 27] using analogous constructions like transport maps, couplings and suprema over ϵ-neighborhoods. Extending our equivalence results to more general loss functions is left for future work.

We analyze optimal adversarial risk over non-parametric classifiers identified by their decision regions. Using a formulation of optimal transport between finite measures of unequal mass, we extend the optimal transport based characterization of adversarial risk of [32, 2] to unequal priors. This extension is nontrivial. We prove it through a generalization of Strassen's theorem which may find applications in the study of excess cost optimal transport [46, 45]. A recent work [40] obtains a different characterization of optimal adversarial risk using optimal transport on the product space $\mathcal{X} \times \mathcal{Y}$ where $\mathcal{Y}$ is the label space. Further, they show the evolution of the optimal classifier A^* as ϵ grows, in terms of a mean curvature flow. This raises an interesting question on the evolution of the optimal adversarial distributions $p_0^*, p_1^* \in \mathcal{P}(\mathcal{X})$ with ϵ.

Using our unbalanced optimal transport characterization adversarial risk, we prove the minimax theorem for adversarial robustness game and the existence of a Nash equilibrium. In this process we give a construction for the worst-case pair of distributions $p_0^*, p_1^* \in \mathcal{P}(\mathcal{X})$ in terms of true data distributions, and show that their total variation distance gives the optimal adversarial risk. Newer robust learning algorithms may benefit from this result by focusing on identifying the worst-case distributions first, and then applying standard learning algorithms to obtain a robust classifier.

To prove our results, we use the tools from the theory of Choquet capacities in $\mathbb{R}^d$ and the theory of measurable selections in general Polish spaces. Specifically, we show that the measure of ϵ-Minkowski expansion is a 2-alternating capacity. This connection may facilitate results from adversarial robustness to be applied to more general perturbation models defined by upper probabilities.

Limitations: While our results for $\mathbb{R}^d$ hold for any Lebesgue-measurable sets, some of our results for general Polish spaces are restricted to sets that are G_δ and F_σ, since we guarantee the measurability of ϵ-Minkowski set expansions only for such sets. For proving minimax theorems, we assume that the Polish space has a midpoint property. It is not clear if these assumptions can be relaxed further.

References

[1] C. Aliprantis and K.C. Border. Infinite dimensional analysis: A hitchhiker's guide. 1994.

[2] A.N. Bhagoji, D. Cullina, and P. Mittal. Lower bounds on adversarial robustness from optimal transport. In *Advances in Neural Information Processing Systems*, pages 7496–7508, 2019.

[3] A.J. Bose, G. Gidel, H. Berrard, A. Cianflone, P. Vincent, S. Lacoste-Julien, and W.L. Hamilton. Adversarial example games. *Advances in Neural Information Processing Systems*, 2020.

[4] S.R. Bulò, B. Biggio, I. Pillai, M. Pelillo, and F. Roli. Randomized prediction games for adversarial machine learning. *IEEE Transactions on Neural Networks and Learning Systems*, 28(11):2466–2478, 2016.

[5] N. Carlini and D. Wagner. Towards evaluating the robustness of neural networks. In *IEEE Symposium on Security and Privacy*, pages 39–57. IEEE, 2017.

[6] I. Chalkidis and D. Kampas. Deep learning in law: early adaptation and legal word embeddings trained on large corpora. *Artificial Intelligence and Law*, 27(2):171–198, 2019.

[7] G. Choquet. Theory of capacities. In *Annales de l'institut Fourier*, volume 5, pages 131–295, 1954.

[8] J.M. Cohen, E. Rosenfeld, and J. Z. Kolter. Certified adversarial robustness via randomized smoothing. In *International Conference on Machine Learning*, 2019.

[9] D. I. Diochnos, S. Mahloujifar, and M. Mahmoody. Adversarial risk and robustness: General definitions and implications for the uniform distribution. *Advances in Neural Information Processing Systems*, 31:10359–10368, 2018.

[10] E. Dohmatob. Generalized no free lunch theorem for adversarial robustness. In *International Conference on Machine Learning*, pages 1646–1654. PMLR, 2019.

[11] E. Dohmatob. Universal lower-bounds on classification error under adversarial attacks and random corruption. *arXiv preprint arXiv:2006.09989*, 2020.

[12] R.M. Dudley. Distances of probability measures and random variables. In *Selected Works of RM Dudley*, pages 28–37. Springer, 2010.

[13] P. Erdős and A. Stone. On the sum of two Borel sets. *Proceedings of the American Mathematical Society*, 25(2):304–306, 1970.

[14] R. Gardner. The Brunn-Minkowski inequality. *Bulletin of the American Mathematical Society*, 39(3):355–405, 2002.

[15] I. J. Goodfellow, J. Shlens, and C. Szegedy. Explaining and harnessing adversarial examples. *International Conference on Learning Representations*, 2015.

[16] Pascale Gourdeau, Varun Kanade, Marta Kwiatkowska, and James Worrell. On the hardness of robust classification. *Advances in Neural Information Processing Systems*, 2019.

[17] H. Greenspan, B. Van Ginneken, and R.M. Summers. Guest editorial deep learning in medical imaging: Overview and future promise of an exciting new technique. *IEEE Transactions on Medical Imaging*, 35(5):1153–1159, 2016.

[18] S. Grigorescu, B. Trasnea, T. Cocias, and G. Macesanu. A survey of deep learning techniques for autonomous driving. *Journal of Field Robotics*, 37(3):362–386, 2020.

[19] P.J. Huber and V. Strassen. Minimax tests and the Neyman-Pearson lemma for capacities. *The Annals of Statistics*, pages 251–263, 1973.

[20] A. Jalal, A. Ilyas, C. Daskalakis, and A.G. Dimakis. The robust manifold defense: Adversarial training using generative models. *arXiv preprint arXiv:1712.09196*, 2017.

[21] R.S.S. Kumar, D.R. O'Brien, K. Albert, and S. Vilojen. Law and adversarial machine learning. *NeurIPS Workshop on Security in Machine Learning*, 2018.

[22] Z. Liu, J. Zhang, V. Jog, P. Loh, and A.B. McMillan. Robustifying deep networks for image segmentation. *arXiv preprint arXiv:1908.00656*, 2019.

[23] H. Luiro, M. Parviainen, and E. Saksman. On the existence and uniqueness of p-harmonious functions. *Differential and Integral Equations*, 27(3/4):201 – 216, 2014.

[24] X. Ma, Y. Niu, L. Gu, Y. Wang, Y. Zhao, J. Bailey, and F. Lu. Understanding adversarial attacks on deep learning based medical image analysis systems. *Pattern Recognition*, 110:107332, 2021.

[25] A. Madry, A. Makelov, L. Schmidt, D. Tsipras, and A. Vladu. Towards deep learning models resistant to adversarial attacks. *International Conference on Learning Representations*, 2018.

[26] S. Mahloujifar, D. I. Diochnos, and M. Mahmoody. The curse of concentration in robust learning: Evasion and poisoning attacks from concentration of measure. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pages 4536–4543, 2019.

[27] L. Meunier, M. Scetbon, R. Pinot, J. Atif, and Y. Chevaleyre. Mixed Nash equilibria in the adversarial examples game. *arXiv preprint arXiv:2102.06905*, 2021.

[28] K. Muhammad, A. Ullah, J. Lloret, J. Del Ser, and V.H.C. de Albuquerque. Deep learning for safe autonomous driving: Current challenges and future directions. *IEEE Transactions on Intelligent Transportation Systems*, 2020.

[29] N. Papernot, F. Faghri, N. Carlini, I. Goodfellow, R. Feinman, A. Kurakin, C. Xie, Y. Sharma, T. Brown, and A. Roy. Technical report on the cleverhans v2. 1.0 adversarial examples library. *arXiv preprint arXiv:1610.00768*, 2016.

[30] N. Papernot, P. McDaniel, X. Wu, S. Jha, and A. Swami. Distillation as a defense to adversarial perturbations against deep neural networks. In *2016 IEEE Symposium on Security and Privacy*, pages 582–597. IEEE, 2016.

[31] R. Pinot, R. Ettedgui, G. Rizk, Y. Chevaleyre, and J. Atif. Randomization matters how to defend against strong adversarial attacks. In *International Conference on Machine Learning*, pages 7717–7727. PMLR, 2020.

[32] M. S. Pydi and V. Jog. Adversarial risk via optimal transport and optimal couplings. In *International Conference on Machine Learning*, pages 7814–7823. PMLR, 2020.

[33] G. Schay. Nearest random variables with given distributions. *The Annals of Probability*, pages 163–166, 1974.

[34] Uri Shaham, Yutaro Yamada, and Sahand Negahban. Understanding adversarial training: Increasing local stability of supervised models through robust optimization. *Neurocomputing*, 307:195–204, 2018.

[35] W. Sierpiński. Sur la question de la mesurabilité de la base de M. Hamel. *Fundamenta Mathematicae*, 1(1):105–111, 1920.

[36] A. Sinha, H. Namkoong, and J. C. Duchi. Certifying some distributional robustness with principled adversarial training. In *International Conference on Learning Representations*, 2017.

[37] M. Staib and S. Jegelka. Distributionally robust deep learning as a generalization of adversarial training. In *NIPS workshop on Machine Learning and Computer Security*, 2017.

[38] V. Strassen. The existence of probability measures with given marginals. *The Annals of Mathematical Statistics*, 36(2):423–439, 1965.

[39] C. Szegedy, W. Zaremba, I. Sutskever, J. Bruna, D. Erhan, I. Goodfellow, and R. Fergus. Intriguing properties of neural networks. *International Conference on Learning Representations*, 2014.

[40] N.G. Trillos and R. Murray. Adversarial classification: Necessary conditions and geometric flows. *arXiv preprint arXiv:2011.10797*, 2020.

[41] Z. Tu, J. Zhang, and D. Tao. Theoretical analysis of adversarial learning: A minimax approach. In *Advances in Neural Information Processing Systems*, pages 12280–12290, 2019.

[42] J. Uesato, B. Oadonoghue, P. Kohli, and A. Oord. Adversarial risk and the dangers of evaluating against weak attacks. In *International Conference on Machine Learning*, pages 5025–5034. PMLR, 2018.

[43] C. Villani. *Topics in Optimal Transportation*. American Mathematical Society, 2003.

[44] D. H. Wagner. Survey of measurable selection theorems. *SIAM Journal on Control and Optimization*, 15(5):859–903, 1977.

[45] L. Yu. Asymptotics of Strassen's optimal transport problem. *arXiv preprint arXiv:1912.02051*, 2019.

[46] L. Yu and V. Tan. Asymptotic coupling and its applications in information theory. *IEEE Transactions on Information Theory*, 65(3):1321–1344, 2018.

[47] M-C Yue, D. Kuhn, and W. Wiesemann. On linear optimization over Wasserstein balls. *arXiv preprint arXiv:2004.07162*, 2020.

[48] L. Zajíček. On σ-porous sets in abstract spaces. In *Abstract and Applied Analysis*, volume 2005, pages 509–534. Hindawi, 2005.