LARGE-WIDTH FUNCTIONAL ASYMPTOTICS FOR DEEP GAUSSIAN NEURAL NETWORKS

ABSTRACT

In this paper, we consider fully connected feed-forward deep neural networks where weights and biases are independent and identically distributed according to Gaussian distributions. Extending previous results (Matthews et al., 2018a;b; Yang, 2019) we adopt a function-space perspective, i.e. we look at neural networks as infinite-dimensional random elements on the input space $\mathbb{R}^I$. Under suitable assumptions on the activation function we show that: i) a network defines a continuous Gaussian process on the input space $\mathbb{R}^I$; ii) a network with re-scaled weights converges weakly to a continuous Gaussian process in the large-width limit; iii) the limiting Gaussian process has almost surely locally γ-Hölder continuous paths, for $0 < \gamma < 1$. Our results contribute to recent theoretical studies on the interplay between infinitely wide deep neural networks and Gaussian processes by establishing weak convergence in function-space with respect to a stronger metric.

1 INTRODUCTION

The interplay between infinitely wide deep neural networks and classes of Gaussian processes has its origins in the seminal work of Neal (1995), and it has been the subject of several theoretical studies. See, e.g., Der & Lee (2006), Lee et al. (2018), Matthews et al. (2018a;b), Yang (2019) and references therein. Let consider a fully connected feed-forward neural network with re-scaled weights composed of $L \geq 1$ layers of widths $n_1, \ldots, n_L$, i.e.

$$
\begin{aligned}
f_i^{(1)}(x) &= \sum_{j=1}^{I} w_{i,j}^{(1)} x_j + b_i^{(1)} & i &= 1, \ldots, n_1 \\
f_i^{(l)}(x) &= \frac{1}{\sqrt{n_{l-1}}} \sum_{j=1}^{n_{l-1}} w_{i,j}^{(l)} \phi(f_j^{(l-1)}(x)) + b_i^{(l)} & l &= 2, \ldots, L, \ \ i = 1, \ldots, n_l
\end{aligned}
\tag{1}
$$

where ϕ is a non-linearity and $x \in \mathbb{R}^I$ is a real-valued input of dimension $I \in \mathbb{N}$. Neal (1995) considered a finite number $k \in \mathbb{N}$ of fixed distinct inputs $(x^{(1)}, \ldots, x^{(k)})$, with each $x^{(r)} \in \mathbb{R}^I$, and weights $w_{i,j}^{(l)}$ and biases $b_i^{(l)}$ independently and identically distributed (iid) as Gaussian distributions. Under appropriate assumptions on the activation ϕ Neal (1995) showed that: i) for each $l \geq 2$, and for a fixed unit i, the k-dimensional random vector $(f_i^{(l)}(x^{(1)}), \ldots, f_i^{(l)}(x^{(k)}))$ converges in distribution, as the widths $n_1, \ldots, n_L$ go to infinity sequentially over network's layers, to a k-dimensional Gaussian random vector; ii) the large-width convergence holds jointly over finite collections of i's and the limiting k-dimensional Gaussian random vectors are independent across the index i. Matthews et al. (2018a;b) extended the work of Neal (1995) by assuming that the width n grows to infinity jointly over network's layers, instead of sequentially, and by establishing joint convergence over all i and countable distinct inputs. The joint growth over the layers is certainly more realistic than the sequential growth, since the infinite Gaussian limit is considered as an approximation of a very wide network. We operate in the same setting, hence from here onward $n \geq 1$ denotes the common layer width, i.e. $n_1, \ldots, n_L = n$. Finally, similar large-width limits have been established for a great variety of neural network architectures, see for instance Yang (2019).

The assumption of a finite (or countable) number of fixed distinct inputs is the common trait of the literature on large-width asymptotics for deep neural networks. Under this assumption, the

large-width limit of a network boils down to the study of the large-width asymptotic behavior of the k-dimensional random vector $(f_i^{(l)}(x^{(1)}), \ldots, f_i^{(l)}(x^{(k)}))$ over $i \geq 1$ for finite k. Such limiting finite-dimensional distributions describe the large-width distribution of a neural networks a priori over any dataset, which is finite by definition. When the limiting distribution is Gaussian, as it often is, this immediately paves the way to Bayesian inference for the limiting network. Such an approach is competitive with the more standard stochastic gradient descent training for the fully-connected architectures object of our study (Lee et al., 2020). However, knowledge of the limiting finite-dimensional distributions is not enough to infer properties of the limiting neural network which are inherently uncountable such as the continuity of the limiting neural network, or the distribution of its maximum over a bounded interval. Results in this direction give a more complete understanding of the assumptions being made a priori, and hence whether a given model is appropriate for a specific application. For instance, Van Der Vaart & Van Zanten (2011) shows that the for Gaussian Processes the function smoothness under the prior should match the smoothness of the target function for satisfactory inference performance.

In this paper we thus consider a novel, and more natural, perspective to the study of large-width limits of deep neural networks. This is an infinite-dimensional perspective where, instead of fixing a finite number of distinct inputs, we look at $f_i^{(l)}(x, n)$ as a stochastic process over the input space $\mathbb{R}^I$. Under this perspective, establishing large-width limits requires considerable care and, in addition, it requires to show the existence of both the stochastic process induced by the neural network and its large-width limit. We start by proving the existence of: i) a continuous Gaussian process, indexed by the network's width n, corresponding to the fully connected feed-forward deep neural network; ii) a continuous Gaussian process corresponding to the infinitely-wide limit of the deep neural network. Then, we prove that the Gaussian process i) converges weakly, as the width n goes to infinity, to the Gaussian process ii) jointly over all units i. As a by-product of our results, we show that the limiting Gaussian process have almost surely locally γ-Hölder continuous paths, for $0 < \gamma < 1$. To make the exposition self-contained we include a compact proof of the main result of Matthews et al. (2018a;b), i.e. the finite-dimensional limit for full connected neural networks.

The paper is structured as follows. In Section 2 we introduce the setting under which we operate, whereas in Section 3 we present a high-level overview of the approach taken to establish our results. Section 4 contains the core arguments of the proof of our large-width functional limit for deep Gaussian neural networks, which are spelled out in detail in the supplementary material (SM). We conclude in Section 5.

2 SETTING

Let $(\Omega, \mathcal{H}, \mathbb{P})$ be the probability space on which all random elements of interest are defined. Furthermore, let $N(\mu, \sigma^2)$ denote a Gaussian distribution with mean $\mu \in \mathbb{R}$ and variance $\sigma^2 \in \mathbb{R}_+$, and let $N_k(\mu, \Sigma)$ be a k-dimensional Gaussian distribution with mean $\mu \in \mathbb{R}^k$ and covariance matrix $\Sigma \in \mathbb{R}^{k \times k}$. In particular, $\mathbb{R}^k$ is equipped with $\| \cdot \|_{\mathbb{R}^k}$, the euclidean norm induced by the inner product $\langle \cdot, \cdot \rangle_{\mathbb{R}^k}$, and $\mathbb{R}^\infty = \times_{i=1}^\infty \mathbb{R}$ is equipped with $\| \cdot \|_{\mathbb{R}^\infty}$, the norm induced by the distance $d(\mathbf{a}, \mathbf{b})_\infty = \sum_{i \geq 1} (|a_i - b_i|)/2^i(1 + |a_i - b_i|)$ for $\mathbf{a}, \mathbf{b} \in \mathbb{R}^\infty$. Note that $(\mathbb{R}, |\cdot|)$ and $(\mathbb{R}^\infty, \| \cdot \|_{\mathbb{R}^\infty})$ are Polish spaces, i.e. separable and complete metric spaces. Let $\omega_{i,j}^{(l)}$ be the random weights of the l-th layer, and assume that they are iid as $N(0, \sigma_\omega^2)$, i.e.

$$\varphi_{\omega_{i,j}^{(l)}}(t) = \mathbb{E}[e^{\mathrm{i}t\omega_{i,j}^{(l)}}] = e^{-\frac{1}{2}\sigma_\omega^2 t^2} \tag{2}$$

is the characteristic function of $\omega_{i,j}^{(l)}$, for $i \geq 1$, $j = 1, \ldots, n$ and $l \geq 1$. Let $b_i^{(l)}$ be the random biases of the l-th layer, and assume that they are iid as $N(0, \sigma_b^2)$, i.e.

$$\varphi_{b_i^{(l)}}(t) = \mathbb{E}[e^{\mathrm{i}tb_i^{(l)}}] = e^{-\frac{1}{2}\sigma_b^2 t^2} \tag{3}$$

is the characteristic function of $b_i^{(l)}$, for for $i \geq 1$ and $l \geq 1$. Weights $\omega_{i,j}^{(l)}$ are independent of biases $b_i^{(l)}$, for any $i \geq 1$, $j = 1, \ldots, n$ and $l \geq 1$. Let $\phi : \mathbb{R} \to \mathbb{R}$ denote a continuous non-linearity. For the finite-dimensional limit we will assume the polynomial envelop condition

$$|\phi(s)| \leq a + b|s|^m, \tag{4}$$

for any $s \in \mathbb{R}$ and some real values $a, b > 0$ and $m \geq 1$. For the functional limit we will use a stronger assumption on ϕ, assuming ϕ to be Lipschitz on $\mathbb{R}$ with Lipschitz constant L_ϕ.

Let $Z(x)$ be a stochastic process on $\mathbb{R}^I$, i.e. $x \in \mathbb{R}^I$, such that $Z(x)$ is defined on $(\Omega, \mathcal{H}, \mathbb{P})$ and it takes values in $\mathbb{R}$. For any $k \in \mathbb{N}$ and $x_1, \ldots, x_k \in \mathbb{R}^I$, let $P_{x_1, \ldots, x_k} = \mathbb{P}(Z(x_1) \in A_1, \ldots, Z(x_k) \in A_k)$, with $A_1, \ldots, A_k \in \mathcal{H}$. Then, the family of finite-dimensional distributions of $Z(x)$ is defined as the family of distributions $\{P_{x_1, \ldots, x_k} : x_1, \ldots, x_k \in \mathbb{R}^I \text{ and } k \in \mathbb{N}\}$. See, e.g., Billingsley (1995). In Definition 1 and Definition 2 we look at the deep neural network (1) as a stochastic process on input space $\mathbb{R}^I$, that is a stochastic process whose finite-dimensional distributions are determined by a finite number $k \in \mathbb{N}$ of fixed distinct inputs $(x^{(1)}, \ldots, x^{(k)})$, with each $x^{(r)} \in \mathbb{R}^I$. The existence of the stochastic processes of Definition 1 and Definition 2 will be thoroughly discussed in Section 3.

Definition 1. *For any fixed $l \geq 2$ and $i \geq 1$, let $(f_i^{(l)}(n))_{n \geq 1}$ be a sequence of stochastic processes on $\mathbb{R}^I$. That is, $f_i^{(l)}(n) : \mathbb{R}^I \to \mathbb{R}$, with $x \mapsto f_i^{(l)}(x, n)$, is a stochastic process on $\mathbb{R}^I$ whose finite-dimensional distributions are the laws, for any $k \in \mathbb{N}$ and $x^{(1)}, \ldots, x^{(k)} \in \mathbb{R}^I$, of the k-dimensional random vectors*

$$f_i^{(1)}(\boldsymbol{X}, n) = f_i^{(1)}(\boldsymbol{X}) = [f_i^{(1)}(x^{(1)}, n), \ldots, f_i^{(1)}(x^{(k)}, n)]^T = \sum_{j=1}^{I} \omega_{i,j}^{(1)} \boldsymbol{x}_j + b_i^{(1)} \boldsymbol{1} \qquad (5)$$

$$f_i^{(l)}(\boldsymbol{X}, n) = f_i^{(l)}(\boldsymbol{X}, n) = [f_i^{(l)}(x^{(1)}, n), \ldots, f_i^{(l)}(x^{(k)}, n)]^T = \frac{1}{\sqrt{n}} \sum_{j=1}^{n} \omega_{i,j}^{(l)}(\phi \bullet f_j^{(l-1)}(\boldsymbol{X}, n)) + b_i^{(l)} \boldsymbol{1} \qquad (6)$$

where $\boldsymbol{X} = [x^{(1)}, \ldots, x^{(k)}] \in \mathbb{R}^{I \times k}$ is a $I \times k$ input matrix of k distinct inputs $x^{(r)} \in \mathbb{R}^I$, $\boldsymbol{1}$ denote a vector of dimension $k \times 1$ of $1s$, $\boldsymbol{x}_j$ denote the j-th row of the input matrix and $\phi \bullet \boldsymbol{X}$ the element-wise application of ϕ to the matrix $\boldsymbol{X}$.

Remark: in contrast to (1), we have defined (5)-(6) over an infinite number of units $i \geq 1$ over each layer l, but the dependency on each previous layer $l - 1$ remains limited to the first n components.

Definition 2. *For any fixed $l \geq 2$, let $(\boldsymbol{F}^{(l)}(n))_{n \geq 1}$ be a sequence of stochastic processes on $\mathbb{R}^I$. That is, $\boldsymbol{F}^{(l)}(n) : \mathbb{R}^I \to \mathbb{R}^\infty$, with $x \mapsto \boldsymbol{F}^{(l)}(x, n)$, is a stochastic process on $\mathbb{R}^I$ whose finite-dimensional distributions are the laws, for any $k \in \mathbb{N}$ and $x^{(1)}, \ldots, x^{(k)} \in \mathbb{R}^I$, of the k-dimensional random vectors*

$$\begin{cases} \boldsymbol{F}^{(1)}(\boldsymbol{X}) = \left[f_1^{(1)}(\boldsymbol{X}), f_2^{(1)}(\boldsymbol{X}), \ldots \right]^T \\ \boldsymbol{F}^{(l)}(\boldsymbol{X}, n) = \left[f_1^{(l)}(\boldsymbol{X}, n), f_2^{(l)}(\boldsymbol{X}, n), \ldots \right]^T. \end{cases}$$

where $f_{r,i}^{(l)}(\boldsymbol{X}, n) = \boldsymbol{I}_r^T f_i^{(l)}(\boldsymbol{X}, n) = f_i^{(l)}(x^{(r)}, n)$ denote the r-th component of the $k \times 1$ vector $f_i^{(l)}(\boldsymbol{X}, n)$ being $\boldsymbol{1}_r$ a vector of dimension $k \times 1$ with 1 in the r-the entry and 0 elsewhere.

Remark: for k inputs, the vector $\mathbf{F}^{(l)}(\mathbf{X}, n)$ is an $\infty \times k$ array, and for a single input $x^{(r)}$, $\mathbf{F}^{(l)}(x^{(r)}, n)$ can be written as $[f_1^{(l)}(x^{(r)}, n), f_2^{(l)}(x^{(r)}, n), \ldots]^T \in \mathbb{R}^{\infty \times 1}$. We define $\mathbf{F}_r^{(l)}(\mathbf{X}, n) = \mathbf{F}^{(l)}(x^{(r)}, n)$ the r-th column of $\mathbf{F}^{(l)}(\mathbf{X}, n)$. When we write $\langle \mathbf{F}^{(l-1)}(x, n), \mathbf{F}^{(l-1)}(y, n) \rangle_{\mathbb{R}^n}$ (see (9)) we treat $\mathbf{F}^{(l)}(x, n)$ and $\mathbf{F}^{(l)}(y, n)$ as elements in $\mathbb{R}^n$ and not in $\mathbb{R}^\infty$, i.e. we consider only the first n components of $\mathbf{F}^{(l)}(x, n)$ and $\mathbf{F}^{(l)}(y, n)$.

3 PLAN SKETCH

We start by recalling the notion of convergence in law, also referred to as convergence in distribution or weak convergence, for a sequence of stochastic processes. See Billingsley (1995) for a comprehensive account.

Definition 3 (convergence in distribution). *Suppose that f and $(f(n))_{n \geq 1}$ are random elements in a topological space C. Then, $(f(n))_{n \geq 1}$ is said to converge in distribution to f, if $\mathbb{E}[h(f(n))] \to \mathbb{E}[h(f)]$ as $n \to \infty$ for every bounded and continuous function $h : C \to \mathbb{R}$. In that case we write $f(n) \xrightarrow{d} f$.*

In this paper, we deal with continuous and real-valued stochastic processes. More precisely, we consider random elements defined on $C(\mathbb{R}^I; S)$, namely the space of continuous functions on the input space $\mathbb{R}^I$ into a Polish space (S, d). Our aim is to study in $C(\mathbb{R}^I; S)$ the convergence in distribution as the width n goes to infinity for:

i) the sequence $(f_i^{(l)}(n))_{n \geq 1}$ for a fixed $l \geq 2$ and $i \geq 1$ with $(S, d) = (\mathbb{R}, |\cdot|)$, i.e. the neural network process for a single unit;

ii) the sequence $(\mathbf{F}^{(l)}(n))_{n \geq 1}$ for a fixed $l \geq 2$ with $(S, d) = (\mathbb{R}^\infty, \|\cdot\|_\infty)$, i.e. the neural network process for all units.

Since applying the Definition 3 in a function space is not easy we need the following proposition.

Proposition 1 (convergence in distribution in $C(\mathbb{R}^I; S)$, (S, d) Polish). *Suppose that f and $(f(n))_{n \geq 1}$ are random elements in $C(\mathbb{R}^I; S)$ with (S, d) Polish space. Then, $f(n) \xrightarrow{d} f$ if: i) $f(n) \xrightarrow{f_d} f$ and ii) the sequence $(f(n))_{n \geq 1}$ is uniformly tight.*

We denoted with $\xrightarrow{f_d}$ the convergence in law of the finite-dimensional distributions of a sequence of stochastic processes. The notion of tightness formalizes the concept that the probability mass is not allowed to "escape at infinity": a single random element f in a topological space C is said to be tight if for each $\epsilon > 0$ there exists a compact $T \subset C$ such that $\mathbb{P}[f \in C \setminus T] < \epsilon$. If a metric space (C, ρ) is Polish any random element on the Borel σ-algebra of C is tight. A sequence of random elements $(f(n))_{n \geq 1}$ in a topological space C is said to be uniformly tight[1] if for every $\epsilon > 0$ there exists a compact $T \subset C$ such that $\mathbb{P}[f(n) \in C \setminus T] < \epsilon$ for all n.

Proof of Proposition 1. By Proposition 16.6 of Kallenberg (2002) $f(n) \xrightarrow{d} f$ in $C(\mathbb{R}^I; S)$ iff $f(n) \xrightarrow{d} f$ in $C(K; S)$ for any $K \subset \mathbb{R}^I$ compact. By Lemma 16.2 of of Kallenberg (2002) the latter holds iff $f(n) \xrightarrow{f_d} f$ and $(f(n))_{n \geq 1}$ is relatively compact in distribution in $C(K; S)$. Note that converge of the finite dimensional distributions holds in $\mathbb{R}^I$ iif it holds in the restriction K for any compact $K \subset \mathbb{R}^I$. The space $(C(K; S), \rho)$, namely the space of continuous functions from a generic compact $K \subset \mathbb{R}^I$ to a Polish space S and $C(K; S)$ endowed with the uniform metric $\rho(f, g) = \sup_{x \in K} d(f(x), g(x))$, is itself a Polish space (Aliprantis & Border, 2006, Lemma 3.97 and Lemma 3.99). Thus by Proposition 16.3 of Kallenberg (2002), i.e. Prohorov Theorem, on $C(K, S)$ $(f(n))_{n \geq 1}$ is relatively compact in distribution iif $(f(n))_{n \geq 1}$ is uniformly tight. Thus, so far we have shown that $f(n) \xrightarrow{d} f$ in $C(\mathbb{R}^I; S)$ iff: i) $f(n) \xrightarrow{f_d} f$ and ii) the sequence $(f(n))_{n \geq 1}$ is uniformly tight on $C(K, S)$ for a generic compact $K \subset \mathbb{R}^I$. It remain to show that the latter holds if $(f(n))_{n \geq 1}$ is uniformly tight on $C(\mathbb{R}^I; S)$. It directly follows by proposition 16.4 of Kallenberg (2002) noting that $C(\mathbb{R}^I; S)$ is a subspace of the metric space $(C(K; S), \rho)$. $\qquad\square$

According to Proposition 1, to achieve convergence in distribution in function spaces we need the following Steps A-D:

Step A) to establish the existence of the finite-dimensional weak-limit f on $\mathbb{R}^I$. We will rely on Theorem 5.3 of Kallenberg (2002), known as Levy theorem.

Step B) to establish the existence of the stochastic processes f and $(f(n))_{n \geq 1}$ as elements in $C(\mathbb{R}^I; S)$. We make use of Daniell-Kolmogorov criterion (Kallenberg, 2002, Theorem 6.16): given a family of multivariate distributions $\{\mu_\mathcal{I}$ probability measure on $\mathbb{R}^{\dim(\mathcal{I})} \mid \mathcal{I} \subset \{x^{(1)}, \ldots, x^{(k)}\}_{x^{(z)} \in \mathbb{R}^I, k \in \mathbb{N}}\}$ there exists a stochastic process with $\{\mu_\mathcal{I}\}$ as finite-dimensional distributions if $\{\mu_\mathcal{I}\}$ satisfies the projective property: $\mu_J(\cdot \times \mathbb{R}_{J \setminus \mathcal{I}}) = \mu_\mathcal{I}(\cdot), \mathcal{I} \subset J \subset \{x^{(1)}, \ldots, x^{(k)}\}_{x^{(z)} \in \mathbb{R}^I, k \in \mathbb{N}}$. That is, it is required consistency with respect to the marginalization over arbitrary components.

Step C) to show that the stochastic process $(f(n))_{n \geq 1}$ and f belong to $C(\mathbb{R}^I; S)$. With regards to $(f(n))_{n \geq 1}$ this is a direct consequence of (5)-(6) and the continuity of ϕ. With regards to the limiting

---

[1] Kallenberg (2002) uses the same term "tightness" for both cases of a single random element and of sequences of random elements; we find that the introduction of "uniform tightness" brings more clarity.

process f, with an additional Lipschitz assumption on ϕ, we rely on the following Kolmogorov-Chentsov criterion (Kallenberg, 2002, Theorem 3.23):

Proposition 2 (continuous version and local-Hölderianity, (S, d) complete). *Let f be a process on $\mathbb{R}^I$ with values in a complete metric space (S, d), and assume that there exist $a, b, H > 0$ such that,*

$$\mathbb{E}[d(f(x), f(y))^a] \leq H \|x - y\|^{(I+b)}, \quad x, y \in \mathbb{R}^I$$

Then f has a continuous version (i.e. f belongs to $C(\mathbb{R}^I; S)$), and the latter is a.s. locally Hölder continuous with exponent c for any for any $c \in (0, {}^b/a)$.

Step D) the uniform tightness of $(f(n))_{n \geq 1}$ in $C(\mathbb{R}^I; S)$. We rely on an extension of the Kolmogorov-Chentsov criterion (Kallenberg, 2002, Corollary 16.9), which is stated in the following proposition.

Proposition 3 (uniform tightness in $C(\mathbb{R}^I; S)$, (S, d) Polish). *Suppose that $(f(n))_{n \geq 1}$ are random elements in $C(\mathbb{R}^I; S)$ with (S, d) Polish space. Assume that $f(0_{\mathbb{R}^I}, n)_{n \geq 1}$ (i.e. $f(n)$ evaluated at the origin) is uniformly tight in S and that there exist $a, b, H > 0$ such that,*

$$\mathbb{E}[d(f(x, n), f(y, n))^a] \leq H \|x - y\|^{(I+b)}, \quad x, y \in \mathbb{R}^I, n \in \mathbb{N}$$

uniformly in n. Then $(f(n))_{n \geq 1}$ is uniformly tight in $C(\mathbb{R}^I; S)$.

4 LARGE-WIDTH FUNCTIONAL LIMITS

4.1 LIMIT ON $C(\mathbb{R}^I; S)$, WITH $(S, d) = (\mathbb{R}, |\cdot|)$, FOR A FIXED UNIT $i \geq 1$ AND LAYER l

Lemma 1 (finite-dimensional limit). *If ϕ satisfies (4) then there exists a stochastic process $f_i^{(l)} : \mathbb{R}^I \to \mathbb{R}$ such that $(f_i^{(l)}(n))_{n \geq 1} \xrightarrow{f_d} f_i^{(l)}$ as $n \to \infty$.*

Proof. Fix $l \geq 2$ and $i \geq 1$. To study the weak limit of the finite-dimensional distributions of $f_i^{(l)}(n)$, fix k inputs $\mathbf{X} = [x^{(1)}, \ldots, x^{(k)}]$. We show that as $n \to +\infty$

$$f_i^{(l)}(\mathbf{X}, n) \xrightarrow{d} N_k(\mathbf{0}, \Sigma(l)), \tag{7}$$

where $\Sigma(l)$ denotes the $k \times k$ covariance matrix, which can be computed through the recursion:

$$\begin{cases} \Sigma(1)_{i,j} = \sigma_b^2 + \sigma_\omega^2 \langle x^{(i)}, x^{(j)} \rangle_{\mathbb{R}^I} \\ \Sigma(l)_{i,j} = \sigma_b^2 + \sigma_\omega^2 \int \phi(f_i)\phi(f_j) q^{(l-1)}(\mathrm{d}f), \end{cases}$$

where $q^{(l-1)} = N_k(\mathbf{0}, \Sigma(l-1))$. See SM A for details. By means of (2), (3), (5) and (6),

$$f_i^{(1)}(\mathbf{X}) \stackrel{d}{=} N_k(\mathbf{0}, \Sigma(1)), \quad \Sigma(1)_{i,j} = \sigma_b^2 + \sigma_\omega^2 \langle x^{(i)}, x^{(j)} \rangle_{\mathbb{R}^I}, \tag{8}$$

and for $l \geq 2$,

$$f_i^{(l)}(\mathbf{X}, n)|f_{1,\ldots,n}^{(l-1)} \stackrel{d}{=} N_k(\mathbf{0}, \Sigma(l, n)), \tag{9}$$

$$\Sigma(l, n)_{i,j} = \sigma_b^2 + \frac{\sigma_\omega^2}{n} \left\langle (\phi \bullet \mathbf{F}_i^{(l-1)}(\mathbf{X}, n)), (\phi \bullet \mathbf{F}_j^{(l-1)}(\mathbf{X}, n)) \right\rangle_{\mathbb{R}^n}.$$

Since (8) does not depends on n we will take the limit as $n \to \infty$ only for (9). From (6) we get that for every fixed $l \geq 2$ and n the sequence $\left(f_i^{(l)}(\mathbf{X}, n)\right)_{i \geq 1}$ is exchangeable. Let $p_n^{(l)}$ denote the de Finetti random probability measure of the exchangeable sequence $\left(f_i^{(l)}(\mathbf{X}, n)\right)_{i \geq 1}$. That is, by de Finetti representation theorem, conditionally to $p_n^{(l)}$ the $f_i^{(l)}(\mathbf{X}, n)$'s are iid as $p_n^{(l)}$. Now, let consider the induction hypothesis that, $p_n^{(l-1)} \xrightarrow{d} q^{(l-1)}$ as $n \to +\infty$, where $q^{(l-1)} = N_k(\mathbf{0}, \Sigma(l-1))$. To establish the convergence in distribution we rely on Theorem 5.3 of Kallenberg (2002), known as Levy theorem, taking into account the point-wise convergence of the characteristic functions. Therefore,

$$\varphi_{f_i^{(l)}(\mathbf{X}, n)}(\mathbf{t}) = \mathbb{E}[\mathbb{E}[e^{\mathrm{i} \mathbf{t}^T f_i^{(l)}(\mathbf{X}, n)}|f_{1,\ldots,n}^{(l-1)}]] = \mathbb{E}\left[\exp\left\{ -\frac{1}{2} \mathbf{t}^T \Sigma(l, n) \mathbf{t} \right\} \right]$$

$$= e^{-\frac{1}{2}\sigma_b^2(\mathbf{t}^T \mathbf{1})^2} \mathbb{E}\left[\mathbb{E}\left[\exp\left\{ -\frac{\sigma_\omega^2}{2n} \sum_{j=1}^n \left(\mathbf{t}^T (\phi \bullet f_j^{(l-1)}(\mathbf{X}, n)) \right)^2 \right\} | p_n^{(l-1)} \right] \right]$$

Conditionally to the random probability measure $p_n^{(l-1)}$ the $f_j^{(l)}(\mathbf{X}, n)$'s are iid as $p_n^{(l-1)}$. Then, we can write

$$\varphi_{f_i^{(l)}(\mathbf{X},n)}(\mathbf{t}) = e^{-\frac{1}{2}\sigma_b^2(\mathbf{t}^T\mathbf{1})^2}\mathbb{E}\Big[\Big(\int \exp\Big\{-\frac{\sigma_\omega^2}{2n}\big(\mathbf{t}^T(\phi\bullet f)\big)^2\Big\}p_n^{(l-1)}(\mathrm{d}f)\Big)^n\Big],$$

where $f = [f_1,\dots,f_k]^T \in \mathbb{R}^k$ and $\mathrm{d}f = (\mathrm{d}f_1,\dots,\mathrm{d}f_k)$. Let $\xrightarrow{p}$ denote the convergence in probability. As it is proved in SM A, the following three auxiliary lemmas hold true:

L1) for each $l \geq 2$ and $s \geq 1$, $\mathbb{P}[p_n^{(l-1)} \in Y_s] = 1$, where $Y_s = \{p : \int \|\phi\bullet f\|_{\mathbb{R}^k}^s p(\mathrm{d}f) < +\infty\}$;

L2) $\int \|\phi\bullet f\|_{\mathbb{R}^k}^2 p_n^{(l-1)}(\mathrm{d}f) \xrightarrow{p} \int \|\phi\bullet f\|_{\mathbb{R}^k}^2 q^{(l-1)}(\mathrm{d}f)$, as $n \to +\infty$;

L3) $\int \|\phi\bullet f\|_{\mathbb{R}^k}^2 [1 - \exp\{-\frac{\sigma_\omega^2}{2n}(\mathbf{t}^T(\phi\bullet f))^2\}]p_n^{(l-1)}(\mathrm{d}f) \xrightarrow{p} 0$, as $n \to +\infty$.

By Lagrange theorem, for each n fixed, there exists a real value $\theta_n \in (0,1)$ such that the following equality holds:

$$\varphi_{f_i^{(l)}(\mathbf{X},n)}(\mathbf{t}) = e^{-\frac{1}{2}\sigma_b^2(\mathbf{t}^T\mathbf{1})^2}\mathbb{E}\Big[\mathbb{I}_{\{p_n^{(l-1)}\in Y_2\}}\Big(1 - \frac{\sigma_\omega^2}{2n}\int\big(\mathbf{t}^T(\phi\bullet f)\big)^2 p_n^{(l-1)}(\mathrm{d}f)+$$
$$+ \frac{\sigma_\omega^2}{2n}\int\big(\mathbf{t}^T(\phi\bullet f)\big)^2\Big[1 - \exp\Big\{-\theta_n\frac{\sigma_\omega^2}{2n}\big(\mathbf{t}^T(\phi\bullet f)\big)^2\Big\}\Big]p_n^{(l-1)}(\mathrm{d}f)\Big)^n\Big].$$

Since for $y > 0$ and $\theta \in (0,1)$ $-e^{-\theta_n y} \leq -e^{-y}$ and since $\big(\mathbf{t}^T(\phi\bullet f)\big)^2 \leq \|\mathbf{t}\|^2\|\phi\bullet f\|^2$ we get that the last integral tends to zero as $n \to +\infty$ by L3. For the remaining part, using the definition of exponential function, i.e. $e^x = \lim_{n\to\infty}(1+\frac{x}{n})^n$, and using L1, L2 and L3 we get the following

$$\varphi_{f_i^{(l)}(\mathbf{X},n)}(\mathbf{t}) \to \exp\Big\{-\frac{1}{2}\Big[\sigma_b^2(\mathbf{t}^T\mathbf{1})^2 + \sigma_\omega^2\int\big(\mathbf{t}^T(\phi\bullet f)\big)^2 q^{(l-1)}(\mathrm{d}f)\Big]\Big\} = \exp\Big\{-\frac{1}{2}\mathbf{t}^T\Sigma(l)\mathbf{t}\Big\},$$

where $\Sigma(l)$ is a $k \times k$ matrix with elements $\Sigma(l)_{i,j} = \sigma_b^2 + \sigma_\omega^2\int\phi(f_i)\phi(f_j)q^{(l-1)}(\mathrm{d}f)$ with $q^{(l-1)} = N_k(\mathbf{0}, \Sigma(l-1))$. The limiting distribution of $f_i^{(l)}(\mathbf{X})$ is a k-dimensional Gaussian distribution with parameter $(\mathbf{0}, \Sigma(l))$. $\qquad\square$

Lemma 1 proves **Step A**. This proof gives an alternative and self-contained proof of the main result of Matthews et al. (2018b), under the more general assumption that the activation function ϕ satisfies the polynomial envelop (4). Now we prove **Step B**, i.e. the existence of the stochastic processes $f_i^{(l)}(n)$ and $f_i^{(l)}$ on the space $C(\mathbb{R}^I; \mathbb{R})$, for each layer $l \geq 1$, unit $i \geq 1$ and $n \in \mathbb{N}$. In SM E.1 we show that the finite-dimensional distributions of $f_i^{(l)}(n)$ satisfies Daniell-Kolmogorov criterion (Kallenberg, 2002, Theorem 6.16), and hence the stochastic process $f_i^{(l)}(n)$ exists. In SM E.2 we prove a similar result for the finite-dimensional distributions of the limiting process $f_i^{(l)}$. In order to prove **Step C** note that $f_i^{(1)}(x) = \sum_{j=1}^I \omega_{i,j}^{(1)}x_j + b_i^{(1)}$ is continuous by construction, thus by induction on l, if $f_i^{(l-1)}(n)$ are continuous for each $i \geq 1$ and n, then $f_i^{(l)}(x,n) = \frac{1}{\sqrt{n}}\sum_{j=1}^n \omega_{i,j}^{(l)}\phi(f_j^{(l-1)}(x,n)) + b_i^{(l)}$ is continuous being composition of continuous functions. For the limiting process $f_i^{(l)}$ we assume ϕ to be Lipschitz with Lipschitz constant L_ϕ. In particular we have the following:

Lemma 2 (continuity). *If ϕ is Lipschitz on $\mathbb{R}$ then $f_i^{(l)}(1), f_i^{(l)}(2),\dots$ are $\mathbb{P}$-a.s. Lipschitz on $\mathbb{R}^I$, while the limiting process $f_i^{(l)}$ is $\mathbb{P}$-a.s. continuous on $\mathbb{R}^I$ and locally γ-Hölder continuous for each $0 < \gamma < 1$.*

Proof. Here we present a sketch of the proof, and we defer to SM B.1 and SM B.2 for the complete proof. For $(f_i^{(l)}(n))_{n\geq 1}$ it is trivial to show that for each n

$$|f_i^{(l)}(x,n) - f_i^{(l)}(y,n)| \leq H_i^{(l)}(n)\|x-y\|_{\mathbb{R}^I}, \quad x,y \in \mathbb{R}^I, \mathbb{P}-a.s. \tag{10}$$

where $H_i^{(l)}(n)$ denotes a suitable random variable, which is defined by the following recursion over l

$$\begin{cases} H_i^{(1)}(n) = \sum_{j=1}^{I} |\omega_{i,j}^{(1)}| \\ H_i^{(l)}(n) = \frac{L_\phi}{\sqrt{n}} \sum_{j=1}^{n} |\omega_{i,j}^{(l)}| H_j^{(l-1)}(n) \end{cases} \tag{11}$$

To establish the continuity of the limiting process $f_i^{(l)}$ we rely on Proposition 2. Take two inputs $x, y \in \mathbb{R}^I$. Defining $\mathbf{a}^T = [1, -1]$, from (7) we know that $f_i^{(l)}(y) - f_i^{(l)}(x) \sim N(\mathbf{a}^T\mathbf{0}, \mathbf{a}^T\Sigma(l)\mathbf{a})$. Thus $|f_i^{(l)}(y) - f_i^{(l)}(x)|^{2\theta} \sim |\sqrt{\mathbf{a}^T\Sigma(l)\mathbf{a}}N(0,1)|^{2\theta} \sim (\mathbf{a}^T\Sigma(l)\mathbf{a})^\theta |N(0,1)|^{2\theta}$. We proceed by induction over the layers. For $l = 1$, we get $\mathbb{E}\left[|f_i^{(1)}(y) - f_i^{(1)}(x)|^{2\theta}\right] = C_\theta(\sigma_\omega^2)^\theta \|y - x\|_{\mathbb{R}^I}^{2\theta}$, where $C_\theta = \mathbb{E}[|N(0,1)|^{2\theta}]$. By hypothesis induction there exists a constant $H^{(l-1)} > 0$ such that $\int |u - v|^{2\theta} q^{(l-1)}(\mathrm{d}u, \mathrm{d}v) \leq H^{(l-1)}\|y - x\|_{\mathbb{R}^I}^{2\theta}$. Then, we write

$$|f_i^{(l)}(y) - f_i^{(l)}(x)|^{2\theta} \sim |N(0,1)|^{2\theta}\left(\sigma_\omega^2 \int [|\phi(u)|^2 - 2\phi(u)\phi(v) + |\phi(v)|^2]q^{(l-1)}(\mathrm{d}u, \mathrm{d}v)\right)^\theta$$

$$\leq |N(0,1)|^{2\theta}(\sigma_\omega^2 L_\phi^2)^\theta \int |u - v|^{2\theta} q^{(l-1)}(\mathrm{d}u, \mathrm{d}v)$$

$$\leq |N(0,1)|^{2\theta}(\sigma_\omega^2 L_\phi^2)^\theta H^{(l-1)}\|y - x\|_{\mathbb{R}^I}^{2\theta},$$

where we used $|\phi(u)|^2 - 2\phi(u)\phi(v) + |\phi(v)|^2 = |\phi(u) - \phi(v)|^2 \leq L_\phi|u - v|^2$ and the Jensen inequality. Thus,

$$\mathbb{E}\left[|f_i^{(l)}(y) - f_i^{(l)}(x)|^{2\theta}\right] \leq H^{(l)}\|y - x\|_{\mathbb{R}^I}^{2\theta}, \tag{12}$$

where $H^{(l)}$ is a suitable constant, which can be explicitly derived by solving the following system of equations

$$\begin{cases} H^{(1)} = C_\theta(\sigma_\omega^2)^\theta \\ H^{(l)} = C_\theta(\sigma_\omega^2 L_\phi^2)^\theta H^{(l-1)}. \end{cases}$$

Then, it is easy to get $H^{(l)} = C_\theta^l (\sigma_\omega^2)^{l\theta}(L_\phi^2)^{(l-1)\theta}$. Observe that $H^{(l)}$ does not depend on i. By Proposition 2, setting $\alpha = 2\theta$, and $\beta = 2\theta - I$ (since β needs to be positive, it is sufficient to choose $\theta > I$) we get that $f_i^{(l)}$ has a continuous version and the latter is $\mathbb{P}$-a.s locally γ-Hölder continuous for every $0 < \gamma < 1 - \frac{I}{2\theta}$, for each $\theta > I$. Taking the limit as $\theta \to +\infty$ we conclude the proof. $\quad\square$

Lemma 3 (uniform tightness). *If ϕ is Lipschitz on $\mathbb{R}$ then $(f_i^{(l)}(n))_{n\geq 1}$ is uniformly tight in $C(\mathbb{R}^I; \mathbb{R})$.*

Proof. The proof requires to verify the conditions of Proposition 3. We defer the proof to SM B.3 since it applies the same arguments that have been applied in the proof of Lemma 2. $\quad\square$

Note that Lemma 3 provides the last **Step D** that allows us to prove the desired result which is explained in the theorem that follows:

Theorem 1 (functional limit). *If ϕ is Lipschitz on $\mathbb{R}$ then $f_i^{(l)}(n) \xrightarrow{d} f_i^{(l)}$ on $C(\mathbb{R}^I; \mathbb{R})$.*

Proof. We apply Proposition 1 to $(f_i^{(l)}(n))_{n\geq 1}$. By Lemma 2, we have that $f_i^{(l)}, (f_i^{(l)}(n))_{n\geq 1}$ belong to $C(\mathbb{R}^I; \mathbb{R})$. From Lemma 1 we have the convergence of the finite-dimensional distributions of $(f_i^{(l)}(n))_{n\geq 1}$, and form Lemma 3 we have the uniform tightness of $(f_i^{(l)}(n))_{n\geq 1}$. $\quad\square$

4.2 LIMIT ON $C(\mathbb{R}^I; S)$, WITH $(S, d) = (\mathbb{R}^\infty, \|\cdot\|_{\mathbb{R}^\infty})$, FOR A FIXED LAYER l

As in the previous section we prove Steps A-D for the sequence $(\mathbf{F}^{(l)}(n))_{n\geq 1}$. Remark that each stochastic process $\mathbf{F}^{(l)}, \mathbf{F}^{(l)}(1), \mathbf{F}^{(l)}(2), \dots$ defines on $C(\mathbb{R}^I; \mathbb{R}^\infty)$ a joint measure whose marginals are the measures introduced in SM E.2 and SM E.1. Let $\mathbf{F}^{(l)} \stackrel{d}{=} \bigotimes_{i=1}^{\infty} f_i^{(l)}$, where $\bigotimes$ denotes the product measure.

Lemma 4 (finite-dimensional limit). *If ϕ satisfies (4) then $\boldsymbol{F}^{(l)}(n) \overset{f_d}{\to} \boldsymbol{F}^{(l)}$ as $n \to \infty$.*

Proof. The proof follows by Lemma 1 and Cramér-Wold theorem for finite-dimensional projection of $\mathbf{F}^{(l)}(n)$: it is sufficient to establish the large n asymptotic of linear combinations of the $f_i^{(l)}(\mathbf{X}, n)$'s for $i \in \mathcal{L} \subset \mathbb{N}$. In particular, we show that for any choice of inputs elements $\mathbf{X}$, as $n \to +\infty$

$$\mathbf{F}^{(l)}(\mathbf{X}, n) \overset{d}{\to} \bigotimes_{i=1}^{\infty} N_k(\mathbf{0}, \Sigma(l)), \tag{13}$$

where $\Sigma(l)$ is defined in (7). The proof is reported in SM C. $\qquad\square$

Lemma 5 (continuity). *If ϕ is Lipschitz on $\mathbb{R}$ then $\boldsymbol{F}^{(l)}, (\boldsymbol{F}^{(l)}(n))_{n \geq 1}$ belong to $C(\mathbb{R}^I; \mathbb{R}^\infty)$. More precisely $\boldsymbol{F}^{(l)}(1), \boldsymbol{F}^{(l)}(2), \ldots$ are $\mathbb{P}$-a.s. Lipschitz on $\mathbb{R}^I$, while the limiting process $\boldsymbol{F}^{(l)}$ is $\mathbb{P}$-a.s. continuous on $\mathbb{R}^I$ and locally γ-Hölder continuous for each $0 < \gamma < 1$.*

Proof. It derives immediately from Lemma 2. We defer to SM D.1 and SM D.2. $\qquad\square$

Theorem 2 (functional limit). *If ϕ is Lipschitz on $\mathbb{R}$ then $(\boldsymbol{F}^{(l)}(n))_{n \geq 1} \overset{d}{\to} \boldsymbol{F}^{(l)}$ as $n \to \infty$ on $C(\mathbb{R}^I; \mathbb{R}^\infty)$.*

Proof. This is Proposition 1 applied to $(\mathbf{F}^{(l)}(n))_{n \geq 1}$. From Lemma 4 and Lemma 5 it remains to show the uniform tightness of the sequence $(\mathbf{F}^{(l)}(n))_{n \geq 1}$ in $C(\mathbb{R}^I; \mathbb{R}^\infty)$. This is immediate by the uniform tightness of each component (Lemma 3). Indeed we know that for each $i \in \mathbb{N}$, for each $\epsilon > 0$ there exists a compact $K_i \subset C(\mathbb{R}^I; \mathbb{R})$ such that $\mathbb{P}[f_i^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}) \setminus K_i] < \epsilon$ for each $n \in \mathbb{N}$. Set $K = \times_{i=1}^{\infty} K_i$ and fix $j \in \mathbb{N}$. Since $\bigcap_{i=1}^{\infty} \{f_i^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}) \setminus K_i\} \subset \{f_j^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}) \setminus K_j\}$, we obtain that $\mathbb{P}\left[\mathbf{F}^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}^\infty) \setminus K\right] = \mathbb{P}\left[\bigcap_{i=1}^{\infty} \{f_i^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}) \setminus K_i\}\right] \leq \mathbb{P}\left[f_j^{(l)}(n) \in C(\mathbb{R}^I; \mathbb{R}) \setminus K_j\right] \leq \epsilon$ which concludes the proof. $\qquad\square$

5 DISCUSSION

We looked at deep Gaussian neural networks as stochastic processes, i.e. infinite-dimensional random elements, on the input space $\mathbb{R}^I$, and we showed that: i) a network defines a Gaussian process on the input space $\mathbb{R}^I$; ii) under suitable assumptions on the activation function, a network with re-scaled weights converges weakly to a Gaussian process in the large-width limit. These results extend previous works (Neal, 1995; Der & Lee, 2006; Lee et al., 2018; Matthews et al., 2018a;b; Yang, 2019) that investigate the limiting distribution of neural network over a finite (or countable) number of distinct inputs. From the point of view of applications, the convergence in distribution is the starting point for the convergence of expectations. Let consider a continuous function $g : C(\mathbb{R}^I; \mathbb{R}^\infty) \to \mathbb{R}$. By the continuous function mapping theorem (Billingsley, 1999, Theorem 2.7), we have $g((f_i^{(l)}(n))_{i \geq 1}) \overset{d}{\to} g((f_i^{(l)})_{i \geq 1})$ as $n \to +\infty$, and under uniform integrability (Billingsley, 1999, Section 3), we have (Billingsley, 1999, Theorem 3.5) $\mathbb{E}[g((f_i^{(l)}(n))_{i \geq 1})] \to \mathbb{E}[g((f_i^{(l)})_{i \geq 1})]$ as $n \to +\infty$. See also Dudley (2002) and references therein.

As a by-product of our results we showed that, under a Lipschitz activation function, the limiting Gaussian process has almost surely locally γ-Hölder continuous paths, for $0 < \gamma < 1$. This raises the question on whether it is possible to strengthen our results to cover the case $\gamma = 1$, or even the case of local Lipschitzianity of the paths of the limiting process. In addition, if the activation function is differentiable, does this property transfer to the limiting process? We leave these questions to future research. Finally, while fully connected deep neural networks represents an ideal starting point for theoretical analysis, modern neural network architectures are composed of a much richer class of layers which includes convolutional, residual, recurrent and attention components. The technical arguments followed in this paper are amenable to extensions to more complex network architectures. Providing a mathematical formulation of network's architectures and convergence results in a way that it allows for extensions to arbitrary architectures, instead of providing an ad-hoc proof for each specific case, is a fundamental research problem.

REFERENCES

Charalambos D. Aliprantis and Kim Border. *Infinite Dimensional Analysis: A Hitchhiker's Guide*. Springer-Verlag Berlin and Heidelberg GmbH & Company KG, 2006.

Patrick Billingsley. *Probability and measure*. John Wiley & Sons, 3rd edition, 1995.

Patrick Billingsley. *Convergence of Probability Measures*. Wiley-Interscience, 2nd edition, 1999.

Ricky Der and Daniel D Lee. Beyond gaussian processes: On the distributions of infinite networks. In *Advances in Neural Information Processing Systems*, pp. 275–282, 2006.

RM Dudley. *Real Analysis and Probability*. Cambridge University Press, 2002.

Olav Kallenberg. *Foundations of modern probability*. Springer Science & Business Media, 2nd edition, 2002.

Jaehoon Lee, Jascha Sohl-dickstein, Jeffrey Pennington, Roman Novak, Sam Schoenholz, and Yasaman Bahri. Deep neural networks as gaussian processes. In *International Conference on Learning Representations*, 2018.

Jaehoon Lee, Samuel S Schoenholz, Jeffrey Pennington, Ben Adlam, Lechao Xiao, Roman Novak, and Jascha Sohl-Dickstein. Finite versus infinite neural networks: an empirical study. *arXiv preprint arXiv:2007.15801*, 2020.

Alexander G. de G. Matthews, Jiri Hron, Mark Rowland, Richard E. Turner, and Zoubin Ghahramani. Gaussian process behaviour in wide deep neural networks. In *International Conference on Learning Representations*, 2018a.

Alexander G. de G. Matthews, Mark Rowland, Jiri Hron, Richard E Turner, and Zoubin Ghahramani. Gaussian process behaviour in wide deep neural networks. *arXiv preprint arXiv:1804.11271*, 2018b.

Radford M Neal. *Bayesian Learning for Neural Networks*. PhD thesis, University of Toronto, 1995.

Aad Van Der Vaart and Harry Van Zanten. Information rates of nonparametric gaussian process methods. *Journal of Machine Learning Research*, 12(6), 2011.

Greg Yang. Wide feedforward or recurrent neural networks of any architecture are gaussian processes. In *Advances in Neural Information Processing Systems*, pp. 9951–9960, 2019.