Stochastic $L^\natural$-convex Function Minimization

Abstract

We study an extension of the stochastic submodular minimization problem, namely, the stochastic $L^\natural$-convex minimization problem. We develop the first polynomial-time algorithms that return a near-optimal solution with high probability. We design a novel truncation operation to further reduce the computational complexity of the proposed algorithms. When applied to a stochastic submodular function, the computational complexity of the proposed algorithms is lower than that of the existing stochastic submodular minimization algorithms. In addition, we provide a strongly polynomial approximate algorithm. The algorithm execution also does not require any prior knowledge about the objective function except the $L^\natural$-convexity. A lower bound on the computational complexity that is required to achieve a high probability error bound is also derived. Numerical experiments are implemented to demonstrate the efficiency of our theoretical findings.

1 Introduction

As an important branch of optimization, discrete stochastic optimization problems (Futschik & Pflug, 1995; Gutjahr & Pflug, 1996; Futschik & Pflug, 1997; Kleywegt et al., 2002; Semelhago et al., 2020) appear in a wide range of applications such as image processing (Jegelka & Bilmes, 2011), pricing optimization (Ito & Fujimaki, 2016) and simulation-based optimization (Altman et al., 2003; Freund et al., 2017). Since a discrete optimization problem is generally $\mathcal{NP}$-hard to solve, specific structures are necessary to guarantee the existence of polynomial-time algorithms. In the special case when the feasible set is $\{0,1\}^d$ for some positive integer d, submodular functions are viewed as the discrete analogy of convex functions in the field of combinatorial optimization. The submodular function minimization (SFM) problem has wide applications in computer vision, economics, game theory and is well-studied in literature (Lee et al., 2015; Axelrod et al., 2020; Zhang et al., 2020). However, for certain problems, because of the stochastic structure, the exact value of the objective function is not available but only noisy samples of the objective function value are available. These scenarios, for example, include the optimization of expected performances for complex stochastic systems. In contrast to the deterministic problem, the stochastic SFM problem is less understood (Ito, 2019; Blais et al., 2019).

In this work, we consider the stochastic minimization problem of a class of generalized submodular functions, named the $L^\natural$-*convex functions* (Murota, 2003). More concretely, a function $f : \{1, \ldots, N\}^d \mapsto \mathbb{R}$ is called a $L^\natural$-convex function if it satisfies the *discrete midpoint convexity*:

$$f(x) + f(y) \geq f(\lceil (x+y)/2 \rceil) + f(\lfloor (x+y)/2 \rfloor), \quad \forall x, y \in \{1, \ldots, N\}^d,$$

where N, d are positive integers and $\lceil \cdot \rceil, \lfloor \cdot \rfloor$ are the ceiling and flooring functions applied component-wisely to vectors. We note that $L^\natural$-convex functions are reduced to submodular functions if $N = 2$. As an extension to the stochastic SFM problem, the stochastic $L^\natural$-convex minimization problem also appears in a variety of decision making problems. For example, the user satisfaction function of a bike-sharing system (Freund et al., 2017) and the expected total waiting time of a service network (Altman et al., 2003) are proved to be multimodular functions, which is equivalent to a $L^\natural$-convex function under a linear transformation. Since the evaluation of both functions includes the simulation of random user behaviors, we only have access to noisy objective function value oracles and thus both

Table 1: Computational complexity for stochastic $L^\natural$-convex function minimization. Here, M is an absolute constant. Constants except for $d, N, \epsilon, \delta, c, M$ are omitted in the $O(\cdot)$ notation.

Algorithms	Unknown Sub-optimality Gap	Known Sub-optimality Gap c
Truncated Subgradient	$O(d^2 N^2 \epsilon^{-2} \log(1/\delta))$	$O(d^2 \log(N) c^{-2} \log(1/\delta))$
Multi-dim Uniform Sampling	$O(C^d \epsilon^{-2} \log(1/\delta))$	$O(C^d c^{-2} \log(1/\delta))$
Stochastic Cutting-plane	$O(d^3 N^2 (d + \log(N))$ $\epsilon^{-2} \log(1/\delta))$	$O(d^3 \log(N)(d + \log(N))$ $c^{-2} \log(1/\delta))$

40 problems can be formulated as stochastic $L^\natural$-convex minimization problems. We refer the readers to
41 the appendix and Murota (2003) for more concrete examples of $L^\natural$-convex minimization problems.

42 The contributions of this work are two-fold. For the algorithm design, we provide the first polynomial-
43 time algorithms to achieve high-probability bounds for the stochastic $L^\natural$-convex function minimiza-
44 tion problem. Besides the $L^\natural$-convexity, no additional structure of the function such as the shape of
45 the landscape is needed for the algorithm execution. We note that we only consider *the number of*
46 *objective function value evaluations* in the computational complexity, since for problems with com-
47 plex stochastic structures, the function value evaluation oracle is usually the major contributor to the
48 running time of algorithms. The first algorithm is based on the observation that the Lovász extension
49 of a $L^\natural$-convex function transforms the original problem to an equivalent continuous convex opti-
50 mization problem. Therefore, we can apply the stochastic subgradient method to solve the continuous
51 convex problem and then round the solution of the new problem to an integral solution. The idea of
52 combining the Lovász extension and the stochastic subgradient method has already been considered in
53 Hazan & Kale (2012); Ito (2019) for stochastic online convex optimization and stochastic submodular
54 minimization, respectively. However, the direct application of the stochastic subgradient method
55 will lead to a running time of $O(d^3 N^2 \epsilon^{-2} \log(1/\delta))$. To reduce the computational complexity in
56 the high-probability regime, we add an extra truncation step to the stochastic subgradient and the
57 computational complexity is reduced by $O(d)$. We also implement numerical experiments to show
58 the significance of the truncation step. In the special case when $N = 2$, our problem reduces to the
59 stochastic SFM problem and the computational complexity of the truncated subgradient method is
60 $O(d^2)$ smaller than the stochastic cutting-plane method in Blais et al. (2019). In addition, we extend
61 the dimension reduction method in Jiang (2020) to the stochastic case and get a strongly polynomial
62 approximate algorithm with computational complexity $O(d^3 N^2 (d + \log(N)) \epsilon^{-2} \log(1/\delta))$. Finally,
63 the uniform sampling method in Zhang et al. (2021) is extended to the multi-dimensional case and
64 the computational complexity is independent of N. In summary, we propose several polynomial-time
65 algorithms for the stochastic $L^\natural$-convex minimization problem. The dimension reduction algorithm
66 is suitable for the case when the Lipschitz constant is hard to estimate or the dimension is not high,
67 while the stochastic subgradient method is suitable for the high-dimensional case. Table 1 summarizes
68 the upper bounds on the computational complexities to find a solution whose objective value is at
69 most ϵ larger than the optimum with probability at least $1 - \delta$.

70 On the other hand, we derive lower bounds on the computational complexity for achieving a high-
71 probability error bound. We construct a set of $L^\natural$-convex functions whose landscapes are close
72 to each other while their optima are different with each other. Intuitively, the algorithm requires
73 sufficiently many evaluations to differentiate those "similar" $L^\natural$-convex functions, where the similarity
74 between functions is characterized by the KL-divergence. More rigorously, we utilize the information
75 inequality in Kaufmann et al. (2016) and get the lower bound $O(d\epsilon^{-2} \log(1/\delta))$ on the complexity.

1.1 Notation

77 For $N \in \mathbb{N}$, we define $[N] := \{1, 2, \ldots, N\}$. For a given set $\mathcal{S}$ and an integer $d \in \mathbb{N}$, the product
78 set $\mathcal{S}^d$ is defined as $\{(x_1, x_2 \ldots, x_d) : x_i \in \mathcal{S}, i \in [d]\}$. For two vectors $x, y \in \mathbb{R}^d$, we use
79 $(x \wedge y)_i := \min\{x_i, y_i\}$ and $(x \vee y)_i := \max\{x_i, y_i\}$ to denote the component-wise minimum and
80 maximum. Similarly, the ceiling function $\lceil \cdot \rceil$ and the flooring function $\lfloor \cdot \rfloor$ round each component to
81 an integer when applied to vectors. The failing probability of algorithms is denoted as δ. The notation
82 $f = O(g)$ (resp. $f = \Theta(g)$) means that there exist absolute constants $c_1, c_2 > 0$ such that $f \leq c_1 g$
83 (resp. $c_1 g \leq f \leq c_2 g$). Similarly, the notation $f = \tilde{O}(g)$ (resp. $f = \tilde{\Theta}(g)$) means that there exist

absolute constants $c_1, c_2 > 0$ and constant $c_3 > 0$ independent of δ such that $f \leq c_1 g + c_3$ (resp. $c_1 g \leq f \leq c_2 g + c_3$).

2 Problem Formulation

We have in mind a complex stochastic system and consider the following discrete stochastic optimization problem

$$\min_x \ f(x) := \mathbb{E}[F(x, \xi_x)] \quad \text{s.t. } x \in \mathcal{X}, \tag{1}$$

in which $\mathcal{X}$ is the discrete feasible set, ξ_x summarizes all the randomness supported on some proper probability space $(\mathsf{Y}, \mathcal{B}_\mathsf{Y})$ and $F : \mathcal{X} \times \mathsf{Y} \mapsto \mathbb{R}$ is a deterministic function. Specifically, ξ_x summarizes all the random variables and stochastic processes involved in the system for the decision variable x. The function F maps the decision and associated randomness into a random system performance. For example, in service systems such as the bike-sharing service, this function F captures the full operational logic regarding how services are arranged. We write $\xi_{x,1}, \xi_{x,2}, \ldots, \xi_{x,n}$ as independent and identically distributed (i.i.d.) copies of ξ_x. We use $\hat{F}_n(x) := \frac{1}{n} \sum_{j=1}^n F(x, \xi_{x,j})$ to denote the empirical mean of the n independent evaluations for the choice of decision variables x. The triad of the set $\mathcal{X}$, the space of randomness $(\mathsf{Y}, \mathcal{B}_\mathsf{Y})$ and the function $F(\cdot, \cdot)$ is called the *model* of problem (1). In this work, we assume the $L^\natural$-convexity of the objective function. For the ease of presentation, we assume the feasible set $\mathcal{X} = [N]^d$, where d denotes the dimension and N represents the number of choices on each dimension for the decision variable x.

Assumption 1. The objective function $f(\cdot)$ is a $L^\natural$-convex function on $\mathcal{X} = [N]^d$, i.e.,

$$f(x) + f(y) \geq f(\lceil (x+y)/2 \rceil) + f(\lfloor (x+y)/2 \rfloor), \quad \forall x, y \in \mathcal{X},$$

Remark 1. In general, the definition of $L^\natural$-convexity also requires the feasible set $\mathcal{X}$ to be a $L^\natural$-convex set, namely, the indicator function of $\mathcal{X}$ should be $L^\natural$-convex on $\mathbb{Z}^d$. In our case, the feasible set $\mathcal{X} = [N]^d$ is a $L^\natural$-convex set. Hence, we only need the discrete mid-point convexity property to define $L^\natural$-convex functions on $\mathcal{X}$.

Moreover, we assume that the distribution of the stochastic output $F(x, \xi_x)$ is sub-Gaussian.

Assumption 2. The distribution of $F(x, \cdot)$ is sub-Gaussian with a *known* parameter σ^2 for all $x \in \mathcal{X}$.

The knowledge of σ^2 represents the maximum level of stochasticity involved in the system performance. For scenarios where this σ^2 is not known, it has to be adaptively learned and adds additional complexity to the problem, but is not the focus of this work. We further note that our algorithm design and analysis can be generalized to any probability distribution with provable concentration inequalities. The set of all models satisfying both Assumption 1 and Assumption 2 is denoted as $\mathcal{MC}(\mathcal{X})$, or simply $\mathcal{MC}$. The main goal is to design algorithms that guarantee the selection of a near-optimal solution with high probability. Formally, this criterion is called the (ϵ, δ)-*Probably Approximately Correct (PAC)* guarantee; see Even-Dar et al. (2002); Kaufmann et al. (2016); Ma & Henderson (2017). We say that a solution $\hat{x}$, as the output of an algorithm, satisfies the (ϵ, δ)-PAC guarantee if

$$\mathbb{P}\left[f(\hat{x}) - \min_{x \in \mathcal{X}} f(x) \geq \epsilon\right] \leq \delta,$$

where the probability is calculated over the distribution of $\hat{x}$.

An algorithms is called an $[(\epsilon, \delta)\text{-}PAC, \mathcal{MC}]$-algorithm if it always returns an (ϵ, δ)-PAC solution for any model in $\mathcal{MC}$. The *computational complexity* of an algorithm is defined as the number of evaluations of $F(\cdot, \cdot)$ before termination. The difficulty of satisfying the PAC guarantee for any model in $\mathcal{MC}$ is characterized by the worst-case performance of an optimal algorithm.

Definition 1. The **computational complexity** of the (ϵ, δ)-PAC guarantee and class $\mathcal{MC}$ is defined as

$$T(\epsilon, \delta, \mathcal{MC}) := \inf_{\mathbf{A} \text{ is } ((\epsilon,\delta)\text{-}PAC,\mathcal{MC})} \sup_{M \in \mathcal{MC}} \mathbb{E}\left[\tau_\mathbf{A}\right],$$

where the random variable $\tau_\mathbf{A}$ is the computational complexity for each implementation of $\mathbf{A}$.

The reminder of this paper is focused on providing upper bounds on $T(\epsilon, \delta, \mathcal{MC})$ by designing algorithms and providing lower bounds by an information-theoretical analysis. In this paper, we consider the behavior of algorithms in the high-probability regime, namely, in the setting when the failing probability δ is small enough.

130 **Assumption 3.** The failing probability δ is sufficiently small.

131 In general, the sub-Gaussian randomness leads to

$$T(\epsilon, \delta, \mathcal{M}) = C(\epsilon, \mathcal{M}) \cdot \log(1/\delta) \quad 0 < \delta \ll 1,$$

132 where $C(\epsilon, \mathcal{M})$ is some function only depending on the precision ϵ and the model class $\mathcal{M}$. For
133 example, it is known that the sample complexity is $\Theta(N^d \epsilon^{-2}) \log(1/\delta)$ for the general best-arm
134 selection problem (Even-Dar et al., 2002) when no convex structure is in presence.

3 Truncated Stochastic Subgradient Algorithms

136 In this section, we develop gradient-based algorithms for problem (1). We first show that the properties
137 of $L^\natural$-convex functions make it possible to construct a continuous convex function on $[1, N]^d$ that
138 has the same set of optima as the original objective function. Then, a rounding process is designed to
139 round the solution of the continuous problem to an integral solution. Finally, the truncated stochastic
140 subgradient algorithm is proposed and its computational complexity is analyzed. Moreover, in this
141 section, we make the assumption that the ℓ_∞-Lipschitz constant of the objective function can be
142 estimated a priori. For example, this estimation is possible for service systems optimization, where
143 it represents an upper bound on how much impact an additional server over a certain time range
144 contributes to the system performance.

145 **Assumption 4.** The ℓ_∞-Lipschitz constant of $f(x)$ is known to be L a priori. Namely, we know
146 beforehand that
$$|f(x) - f(y)| \le L, \quad \forall x, y \in \mathcal{X} \quad \text{s.t. } \|x - y\|_\infty \le 1.$$

3.1 Reduction to a continuous convex optimization problem

148 We first show that the original discrete minimization problem (1) can be equivalently transformed
149 to a continuous convex optimization problem. Based on the fact that the restriction of a $L^\natural$-convex
150 function to a hypercube is a submodular function (Murota, 2003), we show that we can piece together
151 the Lovász extension of the objective function in each hypercube to form a continuous convex function
152 on $\text{conv}(\mathcal{X}) = [1, N]^d$. More importantly, it can be shown that the continuous convex function has
153 exactly the same global minima as the original $L^\natural$-convex function and an unbiased subgradient
154 estimator is available at each point at the cost of $2d$ evaluations to $F(\cdot, \cdot)$.

155 **Theorem 1.** *Suppose that function $f(\cdot)$ is a $L^\natural$-convex function on $[N]^d$. Then, there exists a*
156 *continuous convex function $\tilde{f}(\cdot)$ on $[1, N]^d$, which we also name as the Lovász extension, such that*

157   • *It holds $\tilde{f}(x) = f(x)$ for any $x \in \mathcal{X}$.*

158   • *The minimizers of $\tilde{f}$ satisfy $\arg\min_{y \in [1,N]^d} \tilde{f}(y) = \arg\min_{y \in [N]^d} f(y)$.*

159   • *For any point $x \in [1, N]^d$, there exists an unbiased estimator $\hat{g}_x$ to a subgradient of $\tilde{f}(\cdot)$ at*
160   *point x, which can be evaluated via $2d$ evaluations to $F(\cdot, \cdot)$.*

161 The proof along with a detailed discussion on $L^\natural$-convex functions is provided in the appendix.

162 Next, we describe the rounding process that takes an approximate solution to the relaxed problem

$$\min_{x \in [1,N]^d} \tilde{f}(x) \tag{2}$$

163 to an approximate integral solution to the original problem (1). We list the pseudo-code in Algorithm
164 1. The following theorem proves the correctness of Algorithm 1 and estimates the computational
165 complexity. Note that all upper bounds on computational complexity in this paper are proved to hold
166 almost surely and thus also in expectation.

167 **Theorem 2.** *Suppose that Assumptions 1-4 hold. The solution returned by Algorithm 1 satisfies the*
168 *(ϵ, δ)-PAC guarantee. The computational complexity of Algorithm 1 is at most*

$$O\left[\frac{d}{\epsilon^2} \log\left(\frac{d}{\delta}\right) + d\right] = \tilde{O}\left[\frac{d}{\epsilon^2} \log\left(\frac{1}{\delta}\right)\right].$$

Algorithm 1 Rounding process to a feasible solution

Input: Model $\mathcal{X}, \mathcal{B}_Y, F(x, \xi_x)$, optimality guarantee parameters ϵ, δ, $(\epsilon/2, \delta/2)$-PAC solution $\bar{x}$ to problem (2).
Output: An (ϵ, δ)-PAC solution x^* to problem (1).
 1: Compute the neighbouring points (defined in the appendix) of $\bar{x}$, denoted as $S^0, \ldots, S^d$.
 2: Simulate at S^i until the $1 - \delta/(4d)$ confidence half-width of $\hat{F}_n(S^i)$ is smaller than $\epsilon/4$ for all i.
 3: Return the optimal point $x^* \leftarrow \arg\min_{S \in \{S^0, \ldots, S^d\}} \hat{F}_n(S)$.

Algorithm 2 Projected and truncated SSGD method for the PAC guarantee

---

Input: Model $\mathcal{X}, \mathcal{B}_Y, F(x, \xi_x)$, optimality guarantee parameters ϵ, δ, number of iterations T, step size η, truncation threshold M.
Output: An (ϵ, δ)-PAC solution x^* to problem (1).
 1: Choose the initial point $x^0 \leftarrow ((N+1)/2, \ldots, (N+1)/2)^T$.
 2: **for** $t = 0, \ldots, T-1$ **do**
 3:     Generate a stochastic subgradient $\hat{g}^t$ at x^t.
 4:     Truncate the stochastic subgradient $\tilde{g}^t \leftarrow \mathcal{T}_M(\hat{g}^t)$.
 5:     Update $x^{t+1} \leftarrow \mathcal{P}(x^t - \eta \tilde{g}^t)$.
 6: **end for**
 7: Compute the averaging point $\bar{x} \leftarrow \left(\sum_{t=0}^{T-1} x^t \right) / T$.
 8: Round $\bar{x}$ to an integral point by Algorithm 1.

3.2 Algorithm and complexity analysis

Utilizing the tools derived in the last subsection, we can first apply the stochastic subgradient (SSGD) method to find an approximate solution to problem (2) and then round the solution to get an approximate solution to problem (1). The main difficulty of giving sharp upper bounds lies in the fact that the Lovász extension is neither smooth nor strongly-convex. This property of the Lovász extension prohibits the application of Nesterov's acceleration method (Nesterov, 1983) and variance reduction methods (Johnson & Zhang, 2013; Defazio et al., 2014).

Now, we propose the projected and truncated SSGD method for the (ϵ, δ)-PAC guarantee. The orthogonal projection onto the feasible set, which is defined as

$$\mathcal{P}(x) := (x \wedge N\mathbf{1}) \vee \mathbf{1}, \quad \forall x \in \mathbb{R}^d,$$

is applied after each iteration to ensure the feasibility of iteration point. In addition to the projection, the component-wise truncation of stochastic subgradient is critical in reducing expected computational complexities. The truncation operator with threshold $M > 0$ is defined as

$$\mathcal{T}_M(g) := (g \wedge M\mathbf{1}) \vee (-M\mathbf{1}), \quad \forall g \in \mathbb{R}^d.$$

The pseudo-code of projected and truncated SSGD method is listed in Algorithm 2. With a suitable choice of the step size, the truncation threshold and the number of iterations, Algorithm 2 returns an (ϵ, δ)-PAC solution and the computational complexity scales as $O(d^2 N^2)$.

Theorem 3. *Suppose that Assumptions 1-4 hold. If we choose*

$$T = \tilde{\Theta}\left[\frac{dN^2}{\epsilon^2} \log(\frac{1}{\delta}) \right], \quad M = \tilde{\Theta}\left[\sqrt{\log(\frac{dNT}{\epsilon})} \right], \quad \eta = \frac{N}{M\sqrt{T}},$$

then Algorithm 2 returns an (ϵ, δ)-PAC solution. Furthermore, we have

$$T(\epsilon, \delta, \mathcal{MC}) = O\left[\frac{d^2 N^2}{\epsilon^2} \log(\frac{1}{\delta}) + \frac{d^3 N^2}{\epsilon^2} \log\left(\frac{d^2 N}{\epsilon^3}\right) + \frac{d^3 N^2 L^2}{\epsilon^2} \right] = \tilde{O}\left[\frac{d^2 N^2}{\epsilon^2} \log(\frac{1}{\delta}) \right].$$

We can prove that the computational complexity without the truncation step (i.e., $M = \infty$) is

$$\tilde{O}\left[\frac{d^3 N^2}{\epsilon^2} \log(\frac{1}{\delta}) \right].$$

Hence, the truncation of stochastic subgradient is necessary for reducing the asymptotic expected computational complexity. This relation has also been observed in our numerical experiments. While the error of the normal SSGD method only contains the optimization residual and the variance terms, the error of the truncated SSGD method has an extra bias term. We note that the bias term can be made arbitrarily small with high probability by choosing large enough M and utilizing the tail bound for sub-Gaussian random variables, and therefore the total error can be controlled similarly as the normal SSGD method. We mention that Algorithm 2 also applies when $\mathcal{X}$ is a general $L^\natural$-convex set.

Next, we consider a special case with additional condition that the norm of stochastic subgradients is bounded almost surely. We then show that this additional condition, whenever available, can contribute to further complexity reduction of our proposed algorithm.

Assumption 5. There exist a constant G and an unbiased subgradient estimator $\hat{g}$ such that

$$\mathbb{P}\left[\|\hat{g}\|_1 \leq G\right] = 1.$$

Moreover, the cost of generating each $\hat{g}$ is at most β evaluations of $F(\cdot, \cdot)$.

We note that G and β may depend on d and N. This assumption is common when analyzing the high-probability error bound of stochastic subgradient methods (Hazan & Kale, 2011; Xu et al., 2016). We first give two examples where Assumption 5 holds.

Example 1. Suppose that the randomness of each feasible $x \in \mathcal{X}$ shares the same measure space, i.e., there exists a measure space $(Z, \mathcal{B}_Z)$ such that ξ_x can be any element in the measure space for all $x \in \mathcal{X}$. Moreover, for any fixed $\xi \in \mathcal{B}_Z$, the function $F(\cdot, \xi)$ is also $L^\natural$-convex and has ℓ_∞-Lipschitz constant $\tilde{L}$. Then, the properties of $L^\natural$-convex functions imply that there exists an unbiased subgradient estimator $\hat{g}$ such that the cost of evaluating $\hat{g}$ is $d + 1$ and $\|\hat{g}\|_1 \leq 3\tilde{L}/2$. Therefore, the Assumption 5 holds with $G = 3\tilde{L}/2$ and $\beta = d + 1$. We note that this example is similar to the case when $k = d$ in Ito (2019) while the noise of our problem can be unbounded.

When the distribution at each feasible point is bounded, we show that Assumption 5 also holds.

Example 2. Suppose that the distribution at each point $x \in \mathcal{X}$ is bounded, namely, there exists a constant $C \geq 0$ such that

$$\mathbb{P}_{\xi_x}[|F(x, \xi_x)| \leq C] = 1, \quad \forall x \in \mathcal{X}.$$

We note that bounded distributions are sub-Gaussian with parameter $2C^2$. In this case, there exists an unbiased subgradient estimator such that Assumption 5 holds with $G = 2dC$ and $\beta = 2$.

More details about these two examples are provided in the appendix. Since the stochastic subgradient is bounded, the truncation step is unnecessary in Algorithm 2. The computational complexity of Algorithm 2 is estimated in the following theorem. The proof is similar to Lemma 10 in Hazan & Kale (2011) and, since the feasible set is the hypercube $[1, N]^d$, we replace the ℓ_2-norm with the ℓ_∞-norm to bound distances between points.

Theorem 4. *Suppose that Assumptions 1-5 hold. If we skip the truncation step in Algorithm 2 (i.e., set $M = \infty$) and choose*

$$T = \tilde{\Theta}\left[\frac{N^2(L+G)^2}{\epsilon^2}\log(\frac{1}{\delta})\right], \quad \eta = \sqrt{\frac{dN^2}{TG^2}},$$

then Algorithm 2 returns an (ϵ, δ)-PAC solution. Furthermore, we have

$$T(\epsilon, \delta, \mathcal{MC}) = O\left[\frac{\beta N^2(L+G)^2 + d}{\epsilon^2}\log(\frac{1}{\delta}) + \frac{d^2G^2}{\epsilon^2}\right] = \tilde{O}\left[\frac{\beta N^2(L+G)^2 + d}{\epsilon^2}\log(\frac{1}{\delta})\right].$$

In the case of Example 1, we have $\beta = d + 1, G = 3\tilde{L}/2$ and then the computational complexity of Algorithm 2 is at most $\tilde{O}[dN^2(L+\tilde{L})^2\epsilon^{-2}\log(1/\delta)]$. If both Lipschitz constants are independent of d and N, the computational complexity becomes $\tilde{O}[dN^2\epsilon^{-2}\log(1/\delta)]$, which is $O(d)$ better than the general case without Assumption 5. Moreover, in the case of Example 2, we have $G = d$ and $\beta = 2$. Hence, the computational complexity is at most $\tilde{O}[d^2N^2\epsilon^{-2}\log(1/\delta)]$, even without the truncation.

Finally, we consider the case when the global optimum is unique and the sub-optimality gap is known to be at least $c > 0$. In this case, the Lovász extension satisfies the *weak sharp minimum* condition (Burke & Ferris, 1993) and we can design an acceleration scheme similar to that in Xu et al. (2016). The acceleration scheme reduces the dependence on N from N^2 to $\log(N)$ and we provide detailed discussions in the appendix.

4 Stochastic Localization Algorithms

Although the algorithms developed in Section 3 are guaranteed to find an approximate solution in polynomial time, gradient-based algorithms require the prior knowledge about the Lipschitz constant of the objective function to choose the optimal step size. In addition, the computational complexity of subgradient methods has a quadratic dependence on the Lipschitz constant. However, for certain applications, the upper bound on the Lipschitz constant may be difficult to estimate a priori or is very large so that the computational complexity is prohibitive for large-scale problems. To deal with those difficult scenarios, we utilize the discrete natural of problem (1) to design algorithms that do not require the prior knowledge about the objective function except the $L^\natural$-convexity and whose computational complexity has an upper bound that is independent of the Lipschitz constant. More specifically, we propose two approaches to deal with this problem. The first approach is to generalize the uniform sampling algorithm in Zhang et al. (2021) to the multi-dimensional case. In the second approach, we propose a strongly polynomial approximate algorithm by extending a recently proposed dimension reduction algorithm (Jiang, 2020) to the stochastic case.

4.1 Multi-dimensional uniform sampling algorithm

We first generalize the uniform sampling algorithm in Zhang et al. (2021) to the multi-dimensional case. The basic idea is that if the computational complexity for finding an (ϵ, δ)-PAC solution has the form $C\epsilon^{-2}\log(1/\delta)$, we can view the algorithm as a (possibly biased) estimator with sub-Gaussian tails to the optimal value, where $C > 0$ is a constant. For example, the uniform sampling algorithm provides a sub-Gaussian estimator for the optimal value of problem (1) in the one-dimensional case. Now suppose that we have derived a sub-Gaussian estimator for the optimal value for the $(d-1)$-dimensional case. Consider the function

$$f^1(x_1) := \min_{x_2,\ldots,x_d \in [N]} f(x_1,\ldots,x_d), \quad \forall x_1 \in [N], \tag{3}$$

which is an one-dimensional convex function. Since the $(d-1)$-dimensional algorithm has a sub-Gaussian tail, we can view each implementation of the algorithm as a noisy function value oracle to $f^1(\cdot)$ and the uniform sampling algorithm can be modified to solve the convex problem (3). By a similar method as Zhang et al. (2021), we can prove that the computational complexity of the resulting d-dimensional algorithm is a constant multiple of that of the $(d-1)$-dimensional algorithm. By the induction method, we get an approximate algorithm for the multi-dimensional problem (1) whose computational complexity is independent of the Lipschitz constant L and the scale N in the high-probability regime. A rigorous proof to the following result is provided in the appendix.

Theorem 5. *Suppose that Assumptions 1-3 hold. There exists an $[(\epsilon, \delta)\text{-PAC}, \mathcal{MC}]$-algorithm with computational complexity*

$$T(\epsilon, \delta, \mathcal{MC}) = \tilde{O}\left[\frac{C^d}{\epsilon^2}\log(\frac{1}{\delta})\right],$$

where C is an absolute constant.

The exponential dependence on the dimension d makes it impossible to use the generalized algorithm even for moderate dimension problems. However, this result implies that the computational complexity almost does not grow with N in the regime $N \gg d$ in the asymptotic regime with the failing probability δ is very small. Therefore this algorithm can be preferable when d is small and N is large.

4.2 Dimension reduction algorithm

Next, we take an alternative approach and derive a strongly polynomial approximate algorithm for problem (1). We observe that a subgradient of a convex function provides a separation oracle to the global minima, which can be used to gradually shrink the set that contains global minima. This idea leads to the deterministic cutting-plane methods (Vaidya, 1996; Bertsimas & Vempala, 2004; Lee et al., 2015). In the stochastic case, we can view stochastic subgradients as approximate separation oracles. We prove that if we average $\tilde{O}[dN^2\epsilon^{-2}\log(1/\delta)]$ stochastic subgradients, we can get a separation oracle for the set of ϵ-approximate solutions with probability at least $1 - \delta$. Utilizing this stochastic separation oracle, we may extend the deterministic cutting-plane methods to the stochastic case to find PAC solutions.

However, the stochastic cutting-plane methods still require the knowledge of the Lipschitz constant of the objective function. To design a strongly polynomial algorithm, we observe that if a convex body

$P \subset \mathbb{R}^d$ has a volume smaller than $(d!)^{-1} = O[\exp(-d\log(d))]$, then all integral points in P must lie on a hyperplane. Otherwise, if there exist $d+1$ points $x_0, \ldots, x_d \in P$ that are not on the same hyperplane, then the convex body P contains the polytope $\text{conv}\{x_0, \ldots, x_d\}$, which has the volume

$$\frac{1}{d!}|\det(x_1 - x_0, \ldots, x_d - x_0)| \geq \frac{1}{d!} > \text{vol}(P),$$

where $\text{vol}(P)$ is the volume of the polytope P. This leads to a contradiction. Hence, we may use the stochastic version of Vaidya's method (Vaidya, 1996) or the random walk-based (Bertsimas & Vempala, 2004) cutting-plane method to reduce the volume of the search polytope P to $O[\exp(-d\log(d))]$ and then reduce the problem dimension by projecting the polytope onto the hyperplane. After $d-1$ dimension reductions, we have an one-dimensional convex problem and algorithms in Zhang et al. (2021) can be applied.

To make the dimension reduction algorithm practical, the most important step is to design a polynomial-time algorithm to identify the hyperplane given the volume of P is small enough. Recently, this algorithm was developed in Jiang (2020) via the LLL algorithm (Lenstra et al., 1982). We show that the results can be naturally extended to the stochastic case to achieve the PAC guarantee.

Theorem 6. *Suppose that Assumptions 1-3 hold. Then, the dimension reduction algorithm is an $[(\epsilon, \delta)\text{-PAC}, \mathcal{MC}]$-algorithm with computational complexity*

$$T(\epsilon, \delta, \mathcal{MC}) = \tilde{O}\left[\frac{d^3 N^2 (d + \log(N))}{\epsilon^2} \log(\frac{1}{\delta})\right].$$

We note that if we allow exponentially many arithmetic operations, the LLL algorithm is not necessary and the computational complexity can be reduced to $\tilde{O}[d^4 N^2 \epsilon^{-2} \log(1/\delta)]$. Similar to the gradient-based method, the acceleration scheme can reduce the number of required objective function value evaluations if we have a positive lower bound to the sub-optimality gap.

5 Lower Bounds

In this section, we provide a lower bound result on $[(\epsilon, \delta)\text{-PAC}, \mathcal{MC}]$-algorithms. This result shows the limit of algorithms that achieve the PAC guarantee for any stochastic $L^\natural$-convex functions. To prove lower bounds, basically, we construct several convex models that are "similar" to each other but they have distinct optimal solutions, where the similarity between two models is characterized by the KL-divergence between their distributions. Hence, algorithms need a large number of evaluations to differentiate these models. More rigorously, the information-theoretical inequality in Kaufmann et al. (2016) is utilized to get the following lower bound:

Theorem 7. *Suppose Assumptions 1-3 hold. We have*

$$T(\epsilon, \delta, \mathcal{MC}) \geq \Theta\left[\frac{d}{\epsilon^2} \log(\frac{1}{\delta})\right].$$

The results in Sections 3-4 imply that

$$T(\epsilon, \delta, \mathcal{MC}) = \tilde{O}\left[\frac{\min\{d^2 N^2, M^d\}}{\epsilon^2} \log(\frac{1}{\delta})\right].$$

Therefore, we conjecture that the optimal computational complexity is asymptotically independent of N and leave the design of optimal algorithms for future works.

6 Numerical Experiments

In this section, we present a brief account of numerical experiments to illustrate several of our theoretical findings. First, we show that the extra new truncation step in the gradient-based method can achieve a better performance, compared to the gradient-based method without the truncation step. We implement Algorithm 2 with and without the truncation operation on a staffing allocation stochastic optimization problem with staffing cost constraint, which can be formulated into the regularized and the constrained forms. Details about the problem setup are provided in the appendix. We summarize the computational complexities and the objective values in Table 2. We can see that the two algorithms achieve similar object values but the "Truncated" version observes an advantage in the computational cost compared to the "Not Truncated" version. We note that our algorithms do not require the feasible region to be a hypercube and in fact the problems considered in this section

Table 2: Complexity and returned objective value of Algorithm 2 on the optimal allocation problem.

Params.		Regularized				Constrained			
		Truncated		Not truncated		Truncated		Not truncated	
d	N	Cost	Obj.	Cost	Obj.	Cost	Obj.	Cost	Obj.
4	10	2.99e5	2.10e2	6.56e5	2.11e2	3.00e5	4.76e1	4.99e5	4.97e1
4	20	1.21e5	3.53e2	2.61e5	3.53e2	1.14e5	5.23e1	1.77e5	5.38e1
4	30	8.85e4	4.75e2	1.68e5	4.76e2	7.38e4	5.24e1	1.23e5	5.21e1
4	40	6.25e4	5.91e2	1.34e5	6.07e2	5.28e4	5.31e1	9.24e4	5.28e1
4	50	5.34e4	7.07e2	1.08e5	7.07e2	4.66e4	5.64e1	6.61e4	5.51e1
8	10	1.19e6	1.75e2	3.80e6	1.76e2	1.20e6	3.11e1	2.23e6	3.02e1
12	10	2.68e6	1.59e2	9.48e6	1.59e2	2.69e6	1.87e1	5.36e6	1.86e1
16	10	6.35e6	1.49e2	1.31e7	1.50e2	4.78e6	1.49e1	1.08e7	1.41e1
20	10	9.91e6	1.43e2	2.09e7	1.46e2	9.43e6	1.17e1	1.70e7	1.28e1
24	10	1.50e7	1.35e2	3.09e7	1.41e2	1.36e7	9.43e0	2.10e7	1.17e1

Table 3: Complexity and coverage rate of different algorithms on separable convex functions.

Params.		Truncated subgradient		Dimension reduction	
d	N	Cost(10^7)	Rate(%)	Cost(10^7)	Rate(%)
2	50	0.39	100.0	0.19	100.0
2	100	0.44	100.0	0.22	100.0
2	150	0.51	100.0	0.25	100.0
6	50	0.96	100.0	1.51	100.0
6	100	1.07	100.0	1.76	100.0
6	150	1.24	100.0	1.92	100.0
10	50	1.26	100.0	3.47	100.0
10	100	1.43	100.0	4.02	100.0
10	150	1.64	100.0	4.33	100.0
15	50	1.52	100.0	4.14	100.0
20	50	1.69	100.0	4.28	99.00

have more general feasible set with constraints. Especially, we can see that the truncation plays an important role in reducing the computational complexity, especially when the dimension is high.

Next, we compare the performances of the gradient-based methods and stochastic localization algorithms on multi-dimensional problems. To facilitate the comparison, we construct a separable stochastic convex function, which is a $L^\natural$-convex function by the results in Murota (2003). We have observed that using $(N\epsilon/4, \delta/4)$-$\mathcal{SO}$ oracles is enough for producing high-probability guarantees using the dimension reduction method within the range $1 \leq N \leq 150$. More details about the experiment setup are provided in the appendix. We summarize the results in Table 3, where the coverage rate refers to the percentage of implementations that produce an ϵ-optimal solution. Since the coverage rates of the algorithms are 100%, the PAC guarantee is satisfied by all of the algorithms except the $(d, N) = (20, 50)$ case for the dimension reduction method. This failure is because of our loosened choice of the separation oracles. When the dimension d is not large, the dimension reduction method has the best performance and has the advantage of not requiring any knowledge about the Lipschitz constant; when the dimension d is large, the higher order dependence on d makes the dimension reduction method inferior to the stochastic subgradient method. In addition, the computational complexity of the dimension reduction method has little dependence on N. This is because we choose the separation oracle to be $(N\epsilon/4, \delta/4)$-$\mathcal{SO}$ and the computational complexity of each evaluation is independent of N.

7   Conclusions

In this work, we provide the first polynomial-time algorithms for the stochastic $L^\natural$-convex minimization problem, which serves as an important extension of the stochastic submodular minimization problem. Comparing to existing stochastic submodular minimization algorithms, our algorithms either achieve a lower computational complexity or do not require extra information about the objective function. We also derive a lower bound on the computational complexity to achieve the PAC guarantee. Numerical results are implemented to verify our theoretical findings.

References

Altman, E., Gaujal, B., and Hordijk, A. *Discrete-event control of stochastic networks: Multimodularity and regularity*. springer, 2003.

Axelrod, B., Liu, Y. P., and Sidford, A. Near-optimal approximate discrete and continuous submodular function minimization. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pp. 837–853. SIAM, 2020.

Bach, F. Submodular functions: from discrete to continuous domains. *Mathematical Programming*, 175(1-2):419–459, 2019.

Balasubramanian, K. and Ghadimi, S. Zeroth-order nonconvex stochastic optimization: Handling constraints, high-dimensionality and saddle-points. *arXiv preprint arXiv:1809.06474*, 2018.

Bertsimas, D. and Vempala, S. Solving convex programs by random walks. *Journal of the ACM (JACM)*, 51(4):540–556, 2004.

Blais, E., Canonne, C. L., Eden, T., Levi, A., and Ron, D. Tolerant junta testing and the connection to submodular optimization and function isomorphism. *ACM Transactions on Computation Theory (TOCT)*, 11(4):1–33, 2019.

Borjesson, P. and Sundberg, C.-E. Simple approximations of the error function q (x) for communications applications. *IEEE Transactions on Communications*, 27(3):639–643, 1979.

Burke, J. V. and Ferris, M. C. Weak sharp minima in mathematical programming. *SIAM Journal on Control and Optimization*, 31(5):1340–1359, 1993.

Defazio, A., Bach, F., and Lacoste-Julien, S. Saga: A fast incremental gradient method with support for non-strongly convex composite objectives. In *Advances in neural information processing systems*, pp. 1646–1654, 2014.

Duchi, J. C., Jordan, M. I., Wainwright, M. J., and Wibisono, A. Optimal rates for zero-order convex optimization: The power of two function evaluations. *IEEE Transactions on Information Theory*, 61(5):2788–2806, 2015.

Even-Dar, E., Mannor, S., and Mansour, Y. Pac bounds for multi-armed bandit and markov decision processes. In *International Conference on Computational Learning Theory*, pp. 255–270. Springer, 2002.

Favati, P. Convexity in nonlinear integer programming. *Ricerca operativa*, 53:3–44, 1990.

Freund, D., Henderson, S. G., and Shmoys, D. B. Minimizing multimodular functions and allocating capacity in bike-sharing systems. In *International Conference on Integer Programming and Combinatorial Optimization*, pp. 186–198. Springer, 2017.

Fujishige, S. Theory of submodular programs: A fenchel-type min-max theorem and subgradients of submodular functions. *Mathematical programming*, 29(2):142–155, 1984.

Fujishige, S. *Submodular functions and optimization*. Elsevier, 2005.

Futschik, A. and Pflug, G. Confidence sets for discrete stochastic optimization. *Annals of Operations Research*, 56(1):95–108, 1995.

Futschik, A. and Pflug, G. C. Optimal allocation of simulation experiments in discrete stochastic optimization and approximative algorithms. *European Journal of Operational Research*, 101(2): 245–260, 1997.

Ghadimi, S. and Lan, G. Stochastic first-and zeroth-order methods for nonconvex stochastic programming. *SIAM Journal on Optimization*, 23(4):2341–2368, 2013.

Graur, A., Pollner, T., Ramaswamy, V., and Weinberg, S. M. New query lower bounds for submodular function minimization. In *11th Innovations in Theoretical Computer Science Conference (ITCS 2020)*. Schloss Dagstuhl-Leibniz-Zentrum für Informatik, 2020.

Gutjahr, W. J. and Pflug, G. C. Simulated annealing for noisy cost functions. *Journal of global optimization*, 8(1):1–13, 1996.

Hassani, H., Soltanolkotabi, M., and Karbasi, A. Gradient methods for submodular maximization. In *Proceedings of the 31st International Conference on Neural Information Processing Systems*, pp. 5843–5853, 2017.

Hassidim, A. and Singer, Y. Submodular optimization under noise. In *Conference on Learning Theory*, pp. 1069–1122. PMLR, 2017.

Hazan, E. and Kale, S. Beyond the regret minimization barrier: an optimal algorithm for stochastic strongly-convex optimization. In *Proceedings of the 24th Annual Conference on Learning Theory*, pp. 421–436, 2011.

Hazan, E. and Kale, S. Online submodular minimization. *Journal of Machine Learning Research*, 13 (10), 2012.

Homem-de Mello, T. and Bayraksan, G. Monte carlo sampling-based methods for stochastic optimization. *Surveys in Operations Research and Management Science*, 19(1):56–85, 2014.

Ito, S. Submodular function minimization with noisy evaluation oracle. In *Advances in Neural Information Processing Systems*, pp. 12103–12113, 2019.

Ito, S. and Fujimaki, R. Large-scale price optimization via network flow. In *NIPS*, pp. 3855–3863, 2016.

Jegelka, S. and Bilmes, J. Submodularity beyond submodular energies: coupling edges in graph cuts. In *CVPR 2011*, pp. 1897–1904. IEEE, 2011.

Jiang, H. Minimizing convex functions with integral minimizers. *arXiv preprint arXiv:2007.01445*, 2020.

Johnson, R. and Zhang, T. Accelerating stochastic gradient descent using predictive variance reduction. In *Advances in neural information processing systems*, pp. 315–323, 2013.

Karimi, M. R., Lucic, M., Hassani, H., and Krause, A. Stochastic submodular maximization: the case of coverage functions. In *Proceedings of the 31st International Conference on Neural Information Processing Systems*, pp. 6856–6866, 2017.

Kaufmann, E., Cappé, O., and Garivier, A. On the complexity of best-arm identification in multi-armed bandit models. *The Journal of Machine Learning Research*, 17(1):1–42, 2016.

Kleywegt, A. J., Shapiro, A., and Homem-de Mello, T. The sample average approximation method for stochastic discrete optimization. *SIAM Journal on Optimization*, 12(2):479–502, 2002.

Larson, J., Menickelly, M., and Wild, S. M. Derivative-free optimization methods. *arXiv preprint arXiv:1904.11585*, 2019.

Lee, Y. T., Sidford, A., and Wong, S. C.-w. A faster cutting plane method and its implications for combinatorial and convex optimization. In *2015 IEEE 56th Annual Symposium on Foundations of Computer Science*, pp. 1049–1065. IEEE, 2015.

Lenstra, A. K., Lenstra, H. W., and Lovász, L. Factoring polynomials with rational coefficients. *Mathematische annalen*, 261(ARTICLE):515–534, 1982.

Lovász, L. Submodular functions and convexity. In *Mathematical programming the state of the art*, pp. 235–257. Springer, 1983.

Ma, S. and Henderson, S. G. An efficient fully sequential selection procedure guaranteeing probably approximately correct selection. In *2017 Winter Simulation Conference (WSC)*, pp. 2225–2236. IEEE, 2017.

Mokhtari, A., Hassani, H., and Karbasi, A. Conditional gradient method for stochastic submodular maximization: Closing the gap. In *International Conference on Artificial Intelligence and Statistics*, pp. 1886–1895. PMLR, 2018.

Murota, K. Discrete convex analysis. In *Society for Industrial and Applied Mathematics*. Citeseer, 2003.

Nemirovsky, A. S. and Yudin, D. B. *Problem complexity and method efficiency in optimization.* 1983.

Nesterov, Y. *Lectures on convex optimization*, volume 137. Springer, 2018.

Nesterov, Y. and Spokoiny, V. Random gradient-free minimization of convex functions. *Foundations of Computational Mathematics*, 17(2):527–566, 2017.

Nesterov, Y. E. A method for solving the convex programming problem with convergence rate o (1/k^ 2). In *Dokl. akad. nauk Sssr*, volume 269, pp. 543–547, 1983.

Sekar, S., Vojnovic, M., and Yun, S.-Y. A test score-based approach to stochastic submodular optimization. *Management Science*, 67(2):1075–1092, 2021. doi: 10.1287/mnsc.2020.3585. URL https://doi.org/10.1287/mnsc.2020.3585.

Semelhago, M., Nelson, B. L., Song, E., and Wächter, A. Rapid discrete optimization via simulation with Gaussian Markov random fields. *INFORMS journal on Computing*, Articles in Advance, 2020.

Shamir, O. A variant of azuma's inequality for martingales with subgaussian tails. *arXiv preprint arXiv:1110.2392*, 2011.

Vaidya, P. M. A new algorithm for minimizing convex functions over convex sets. *Mathematical programming*, 73(3):291–341, 1996.

Xu, Y., Lin, Q., and Yang, T. Accelerated stochastic subgradient methods under local error bound condition. *arXiv preprint arXiv:1607.01027*, 2016.

Yang, T. and Lin, Q. Rsg: Beating subgradient method without smoothness and strong convexity. *The Journal of Machine Learning Research*, 19(1):236–268, 2018.

Zhang, H., Zheng, Z., and Lavaei, J. Selection of the best under convexity. *SSRN preprint 3848865*, 2021.

Zhang, J., Lin, H., Sra, S., and Jadbabaie, A. On complexity of finding stationary points of nonsmooth nonconvex functions. *arXiv preprint arXiv:2002.04130*, 2020.

Zinkevich, M. Online convex programming and generalized infinitesimal gradient ascent. In *Proceedings of the 20th international conference on machine learning (icml-03)*, pp. 928–936, 2003.