Nonlinear Functions of Gaussian Random Vectors Are Subgaussian: An AI-Assisted Note

Guangyi Zou [1] **Roman Vershynin** [1]

Abstract

This short note presents a dimension-independent subgaussian concentration bound for Gaussian vectors under coordinate-wise nonlinear mappings. Discovered by Gemini 3.5 Flash, this result applies to any bounded function under a well-conditioned covariance. We apply this tool to resolve an open question posed by Simone Bombari on sign-quantized linear maps $Y = \text{sgn}(Wx)$. By partitioning the square matrix into rectangular blocks, we bypass the typical singularity issues of square random matrices to establish the desired concentration.

1. Introduction: An AI-for-Math Perspective

Simone Bombari asked us whether the 1-bit quantized random vector $Y = \text{sgn}(Wx)$ has subgaussian norm bounded by a universal constant. Here W is an $n \times n$ random Gaussian matrix, and x is an independent standard normal random vector in $\mathbb{R}^n$. The question is nontrivial since the coordinates of Y are not independent. We give a strong positive answer to this question – for any bounded map instead of $\text{sgn}(\cdot)$ – using AI:

- **AI Discovery and Generalization (Theorem 2.1):** To handle coordinate dependence, Gemini 3.5 Flash[2] proposed decomposing the Gaussian vector into independent parts, using one part to "smooth" the sign function,

[1]Department of Mathematics, University of California, Irvine, CA, USA. Correspondence to: Guangyi Zou <gzou3@uci.edu>, Roman Vershynin <rvershyn@uci.edu>.

The 3^{rd} *AI for Math Workshop* at the 43^{rd} International Conference on Machine Learning (ICML), Seoul, South Korea, 2026.

[2]The original Chinese dialogue is archived at https://gemini.google.com/share/daca36d5cfbc; see https://github.com/The8ravo/conversation-history-with-Gemini/blob/main/conversation_history_with_gemini.pdf for an AI-generated English translation with a timeline. The reproduced interactive dialogue and full derivations are archived at https://gemini.google.com/share/cefbb71da6d2.

and then applying Gaussian concentration for Lipschitz functions. It subsequently generalized the result for the sign function to an arbitrary bounded functions. However, this idea works for well-conditioned matrices W only.

- **Human Design (Section 3):** To apply this general tool to square random matrices W – which are typically ill conditioned – the human authors used a simple row-partitioning trick. This breaks the square matrix into two well-conditioned rectangular blocks, bypassing the singularity issue.

2. Main Result: Subgaussianity under Well-conditioned Covariance

The subgaussian norm of a random vector is defined as the largest subgaussian norm of its 1-dimensional marginals:

$$\|Y\|_{\psi_2} = \sup_{v \in S^{n-1}} \|\langle v, Y \rangle\|_{\psi_2},$$

(Vershynin, 2026, Definition 3.4.1). The condition number of a positive-definite matrix Σ is the ratio of the largest to smallest eigenvalues:

$$\kappa(\Sigma) := \frac{\lambda_{\max}(\Sigma)}{\lambda_{\min}(\Sigma)}.$$

The following theorem, giving dimension-independent subgaussian bound, was proved by Gemini 3.5 Flash.

Theorem 2.1. *Let $X \sim \mathcal{N}(0, \Sigma)$ be an n-dimensional normal random vector with a nonsingular covariance matrix Σ. Let $\phi : \mathbb{R} \to \mathbb{R}$ satisfy $\|\phi\|_{\infty} \leq 1$. Define the random vector $Y = \phi(X) \in \mathbb{R}^n$ by applying ϕ to each coordinate of X. Then*

$$\|Y - \mathbb{E}Y\|_{\psi_2} \leq C\sqrt{\kappa(\Sigma)},$$

where $C > 0$ is an absolute constant.

Proof. By definition, we need to show that for any direction $v \in S^{n-1}$, the random variable

$$\langle v, Y - \mathbb{E}Y \rangle = \sum_{i=1}^{n} v_i(Y_i - \mathbb{E}Y_i)$$

has a uniformly bounded subgaussian norm.

Step 1: Covariance Splitting. Let $a = \lambda_{\min}(\Sigma)$ and $b = \lambda_{\max}(\Sigma)$, so that $\kappa(\Sigma) = b/a$. Defining the residual covariance matrix $\Sigma_G := \Sigma - aI_n \geq 0$, we can decompose X into the sum of two independent Gaussian vectors:

$$X = \sqrt{a}Z + G, \quad \text{where } Z \sim \mathcal{N}(0, I_n) \text{ and } G \sim \mathcal{N}(0, \Sigma_G).$$

Step 2: Conditional Hoeffding Bound. Conditioned on G, the coordinates $Y_i = \phi(\sqrt{a}Z_i + G_i)$ are mutually independent. Let

$$\mu_i(x) := \mathbb{E}\,\phi(\sqrt{a}Z_i + x), \quad x \in \mathbb{R}.$$

The conditional expectation satisfies $\mathbb{E}[Y_i \mid G] = \mu_i(G_i)$. Since $\|\phi\|_\infty \leq 1$, each centered coordinate is deterministically bounded: $|v_i(Y_i - \mu_i(G_i))| \leq 2|v_i|$. By Hoeffding's inequality (see e.g. (Vershynin, 2026, Theorem 2.2.6)), a random variable bounded within an interval of length $4|v_i|$ has MGF bounded by $\exp(\lambda^2(4|v_i|)^2/8) = \exp(2v_i^2\lambda^2)$. Summing over the independent coordinates gives:

$$\mathbb{E}\left[\exp\left(\lambda \sum_{i=1}^n v_i(Y_i - \mu_i(G_i))\right) \;\middle|\; G\right] \\ \leq \exp\left(\sum_{i=1}^n 2v_i^2\lambda^2\right) = \exp\left(2\lambda^2\right). \tag{1}$$

Step 3: Concentration of the Conditional Mean. Define the function

$$F(x_1, \dots, x_n) := \sum_{i=1}^n v_i\mu_i(x_i).$$

Although ϕ is not assumed to be differentiable, the Gaussian density smooths the expectation via convolution. By writing $\mu_i(x) = \int_{\mathbb{R}} \phi(u)\frac{1}{\sqrt{2\pi a}}e^{-\frac{(u-x)^2}{2a}}\,du$ and differentiating the Gaussian kernel directly with respect to x, a simple change of variables yields:

$$|\mu_i'(x)| = \left|\frac{1}{\sqrt{a}}\mathbb{E}\,Z_i\,\phi(\sqrt{a}Z_i + x)\right| \leq \frac{1}{\sqrt{a}}\mathbb{E}|Z_i| = \sqrt{\frac{2}{\pi a}}.$$

Thus, the gradient satisfies $\|\nabla F\|_2 \leq \max_i |\mu_i'(x_i)| \cdot \|v\|_2 \leq \sqrt{\frac{2}{\pi a}}$, meaning F is L_μ-Lipschitz with $L_\mu = \sqrt{\frac{2}{\pi a}}$. Note also that

$$\mathbb{E}F(G) = \sum_{i=1}^n v_i\mathbb{E}Y_i.$$

By standard Gaussian concentration for the Lipschitz function of $G \sim \mathcal{N}(0, \Sigma_G)$ (see e.g. (Vershynin, 2026, Theorem

5.2.11) or (Boucheron et al., 2013, Theorem 5.5)), we have:

$$\mathbb{E}\left[\exp\left(\lambda(F(G) - \mathbb{E}F(G))\right)\right] \\ \leq \exp\left(\frac{\lambda^2 L_\mu^2\|\Sigma_G\|_{\mathrm{op}}}{2}\right) \\ \leq \exp\left(\frac{\lambda^2}{\pi a}(b - a)\right) = \exp\left(\frac{\lambda^2}{\pi}(\kappa(\Sigma) - 1)\right). \tag{2}$$

Step 4: Total MGF Bound. We decompose the target variable into conditional fluctuations and the variation of the conditional mean:

$$\langle v, Y - \mathbb{E}Y \rangle = \sum_{i=1}^n v_i(Y_i - \mu_i(G_i)) + (F(G) - \mathbb{E}F(G)).$$

By the law of total expectation, we combine (1) and (2):

$$\mathbb{E}\left[\exp\left(\lambda\langle v, Y - \mathbb{E}Y \rangle\right)\right] \\ = \mathbb{E}_G\left[\mathbb{E}\left[\exp\left(\lambda\sum_{i=1}^n v_i(Y_i - \mu_i(G_i))\right) \;\middle|\; G\right] \\ \cdot \exp\left(\lambda(F(G) - \mathbb{E}F(G))\right)\right] \\ \leq \exp(2\lambda^2) \cdot \exp\left(\frac{\lambda^2}{\pi}(\kappa(\Sigma) - 1)\right) \\ = \exp\left(\frac{\sigma^2\lambda^2}{2}\right),$$

where

$$\sigma^2 = 4 + \frac{2}{\pi}(\kappa(\Sigma) - 1) \leq 4\kappa(\Sigma),$$

since $\kappa(\Sigma) \geq 1$.

By the standard equivalence of subgaussian properties (see e.g. (Vershynin, 2026, Proposition 2.6.1)), this implies $\|\langle v, Y - \mathbb{E}Y \rangle\|_{\psi_2} \leq C\sqrt{\kappa(\Sigma)}$, where C is an absolute constant. Taking the supremum over $v \in S^{n-1}$ completes the proof. $\qquad\square$

Remark 2.2 (Independent and simultaneous work of Roth). After completing the initial version of this note, we learned that Valentin Roth, working independently and simultaneously, obtained a dependent version of McDiarmid's inequality by a different method based on approximate tensorization of entropy; see Roth (2026). In the Gaussian case, Roth's approach gives a Hamming-Lipschitz concentration inequality for $X \sim \mathcal{N}(\mu, \Sigma)$ with dependence controlled by an ATE constant. This constant is of the order of $\kappa(\Sigma)$ and can be sharpened to

$$\kappa_*(\Sigma) = \lambda_{\max}(D^{1/2}\Sigma D^{1/2}), \qquad D = \mathrm{diag}(\Sigma^{-1}),$$

which may be smaller than the usual condition number. Roth's approach also applies more broadly to measures

satisfying approximate tensorization of entropy, including strongly log-concave and log-smooth distributions. Our proof is elementary and complementary; in particular, the same Gaussian smoothing idea also yields the Hölder-continuous extension described below.

Remark 2.3 (Extension to Continuous Functions). If ϕ is α-Hölder continuous instead of globally bounded, the same decomposing trick still works, and the subgaussian norm becomes bounded by $O(\sqrt{\lambda_{\max}(\Sigma)}\,\lambda_{\min}(\Sigma)^{\frac{\alpha-1}{2}})$. See the conversation history for Gemini 3.5 Flash's proof.

Furthermore, the well-conditioning of Σ is strictly necessary. A natural counterexample is the rank-one matrix $\Sigma = \mathbf{1}\mathbf{1}^\top$ with $\mathbf{1} = (1,\ldots,1)^\top \in \mathbb{R}^n$, which yields $\|\operatorname{sgn}(X)\|_{\psi_2} \geq c\sqrt{n}$ for an absolute constant $c > 0$.

3. Application: Sign-Quantized Square Linear Maps

We now apply Theorem 2.1 to resolve the question posed by Simone Bombari.

Corollary 3.1. *Let $W \in \mathbb{R}^{n \times n}$ be a random matrix with i.i.d. standard normal entries $W_{ij} \sim \mathcal{N}(0,1)$. Let $x \sim \mathcal{N}(0, I_n)$ be an n-dimensional standard normal random vector. Define the random vector $Y = \operatorname{sgn}(Wx) \in \{-1, 1\}^n$ by applying the sign function to each coordinate of x. Then*

$$\mathbb{E}\|Y\|_{\psi_2 | W} \leq C,$$

where $C > 0$ is an absolute constant. Here, the conditional norm $\|Y\|_{\psi_2 | W}$ is with respect to the random vector x for a fixed realization of W, and the expectation is with respect to the random matrix W.

Proof. Directly applying Theorem 2.1 to W fails because the Wishart matrix $\Sigma = WW^\top$ typically has large condition number.

To fix this, we split W horizontally into two rectangular blocks $W_1 \in \mathbb{R}^{m \times n}$ and $W_2 \in \mathbb{R}^{(n-m) \times n}$, where $m = \lfloor n/2 \rfloor$:

$$W = \begin{pmatrix} W_1 \\ W_2 \end{pmatrix}. \tag{3}$$

This induces a corresponding split on the output vector into

$$Y^{(1)} = \operatorname{sgn}(W_1 x) \in \{-1, 1\}^m$$

and

$$Y^{(2)} = \operatorname{sgn}(W_2 x) \in \{-1, 1\}^{n-m}.$$

Applying the triangle inequality yields $\|Y\|_{\psi_2 | W} \leq \|Y^{(1)}\|_{\psi_2 | W_1} + \|Y^{(2)}\|_{\psi_2 | W_2}$. By symmetry, it suffices to show that $\mathbb{E}\|Y^{(1)}\|_{\psi_2 | W_1}$ is bounded by a universal constant.

Since the aspect ratio satisfies $m/n \leq 1/2$, the rectangular block W_1 is well-conditioned. Specifically, its sample covariance $\Sigma_1 = W_1 W_1^\top$ satisfies $nc_1 I \preceq \Sigma_1 \preceq nc_2 I$, and therefore

$$\kappa(\Sigma_1) \leq c_2/c_1,$$

with high probability $1 - c_3 e^{-c_4 n}$ for universal constants $c_1, c_2, c_3, c_4 > 0$ (see e.g., (Vershynin, 2026, Corollary 7.3.2 and Exercise 7.13)). Let $\mathcal{E} := \{\kappa(\Sigma_1) \leq c_2/c_1\}$ denote this well-conditioned event.

Since $W_1 x$ is a centered Gaussian vector, its distribution is symmetric about the origin, which implies $\mathbb{E}[Y^{(1)} \mid W_1] = 0$. Applying Theorem 2.1 directly to $Y^{(1)}$ yields:

$$\|Y^{(1)}\|_{\psi_2 | W_1} \leq C_0 \sqrt{\kappa(\Sigma_1)} \leq C_0 \sqrt{\frac{c_2}{c_1}} =: C_1. \tag{4}$$

To bound the expectation over W, we decompose

$$\mathbb{E}\|Y^{(1)}\|_{\psi_2 | W_1} = \mathbb{E}\|Y^{(1)}\|_{\psi_2 | W_1} \mathbb{1}_{\mathcal{E}} + \mathbb{E}\|Y^{(1)}\|_{\psi_2 | W_1} \mathbb{1}_{\mathcal{E}^c}.$$

On the event $\mathcal{E}$, we use (4). On the rare complement $\mathcal{E}^c$, we use the trivial deterministic bound $|\langle v, Y^{(1)} \rangle| \leq \sqrt{m} \leq \sqrt{n}$ for all $v \in S^{m-1}$, which implies $\|Y^{(1)}\|_{\psi_2 | W_1} \leq c_5 \sqrt{n}$. Plugging in the bounds for each case and using the tail probability $\mathbb{P}(\mathcal{E}^c) \leq c_3 e^{-c_4 n}$ gives:

$$\mathbb{E}_W[\|Y^{(1)}\|_{\psi_2}]$$
$$\leq C_0 \sqrt{\frac{c_2}{c_1}} \mathbb{P}(\mathcal{E}) + c_5 \sqrt{n} \mathbb{P}(\mathcal{E}^c)$$
$$\leq C_0 \sqrt{\frac{c_2}{c_1}} + c_5 c_3 \sqrt{n} e^{-c_4 n}.$$

Summing the bounds for both blocks via the triangle inequality completes the proof. $\square$

Remark 3.2 (What AI could do, and what it couldn't). It is instructive to contrast the AI model's capabilities across these results. While Gemini 3.5 Flash autonomously proved the initial query for the sign function and generalized the argument to arbitrary bounded mappings (yielding Theorem 2.1), it could not prove Corollary 3.1 outright, via direct prompting. Transitioning from well-conditioned covariances to square random matrices required a human-designed row-partitioning trick (3) to bypass the singularity issues.

Impact Statement

This paper presents work whose goal is to advance the field of Machine Learning. There are many potential societal consequences of our work, none which we feel must be specifically highlighted here.

Acknowledgments

The authors thank Simone Bombari for posing the question that motivated this work and for pointing out its connection

with McDiarmid's inequality. We thank Valentin Roth for communicating his independent and simultaneous work on McDiarmid-type inequalities under dependence. We also acknowledge the mathematical collaboration with Google Gemini 3.5 Flash. A reproduced interactive dialogue and derivation transcripts are archived at https://gemini.google.com/share/cefbb71da6d2. The authors are supported by NSF Grant DMS 2451011 and U.S. Air Force Grant FA9550-25-1-0294.

References

Boucheron, S., Lugosi, G., and Massart, P. *Concentration inequalities*. Oxford University Press, Oxford, 2013. ISBN 978-0-19-953525-5. doi: 10.1093/acprof:oso/9780199535255.001.0001. URL https://doi.org/10.1093/acprof:oso/9780199535255.001.0001. A nonasymptotic theory of independence, With a foreword by Michel Ledoux.

Roth, V. On mcdiarmid's inequality under dependence via approximate tensorization of entropy. *arXiv preprint arXiv:2606.12720*, 2026.

Vershynin, R. *High-Dimensional Probability: An Introduction with Applications in Data Science*. Cambridge Series in Statistical and Probabilistic Mathematics. Cambridge University Press, Cambridge, 2 edition, 2026.