Split-kl and PAC-Bayes-split-kl Inequalities

Abstract

We present a new concentration of measure inequality for sums of independent bounded random variables, which we name a split-kl inequality. The inequality combines the combinatorial power of the kl inequality with ability to exploit low variance. While for Bernoulli random variables the kl inequality is tighter than the Empirical Bernstein, for random variables taking values inside a bounded interval and having low variance the Empirical Bernstein inequality is tighter than the kl. The proposed split-kl inequality yields the best of both worlds. We discuss an application of the split-kl inequality to bounding excess losses. We also derive a PAC-Bayes-split-kl inequality and use a synthetic example and several UCI datasets to compare it with the PAC-Bayes-kl, PAC-Bayes Empirical Bernstein, PAC-Bayes Unexpected Bernstein, and PAC-Bayes Empirical Bennett inequalities.

1 Introduction

Concentration of measure inequalities for sums of independent random variables are the most fundamental analysis tools in statistics and many other domains [Boucheron et al., 2013]. Their history stretches almost a century back, and inequalities such as Hoeffding's [Hoeffding, 1963] and Bernstein's [Bernstein, 1946] are the main work horses of learning theory.

We restrict our attention to bounded random variables. Without loss of generality we can talk about random variables bounded in the $[0, 1]$ interval. For Bernoulli random variables, which take values at the endpoints of the interval, $\{0, 1\}$, one of the tightest concentration of measure inequalities is the kl inequality [Maurer, 2004, Langford, 2005, Foong et al., 2021, 2022], which is based on combinatorial properties of a sum of n independent random variables. However, while being extremely tight for Bernoulli random variables, the kl inequality is not necessarily a good choice for sums of random variables that take values both at the boundary and inside the interval. In the latter case the Empirical Bernstein [Maurer and Pontil, 2009] and the Unexpected Bernstein [Mhammedi et al., 2019] inequalities can be significantly tighter due to their ability to exploit low variance, as shown by Tolstikhin and Seldin [2013].

Our study was motivated by an attempt to derive tighter concentration inequalities for excess losses. More specifically, we were aiming to improve concentration inequalities for ternary random variables originating from the difference between the zero-one losses of a prediction rule h and a reference prediction rule h^*, $Z = \ell(h(X), Y) - \ell(h^*(X), Y) \in \{-1, 0, 1\}$. This kind of excess losses was used, for example, by Mhammedi et al. [2019] in the context of PAC-Bayesian generalization bounds for classification. In their work h^* was pre-trained on part of the data, and then bounds on the excess loss were used to construct a randomized classifier based on the remaining part of the data. Mhammedi et al. were seeking to exploit low variance of excess losses by applying the Unexpected Bernstein inequality. However, the challenge is that if the predictions of h and h^* are similar, then Z is frequently zero and the Empirical Bernstein or Unexpected Bernstein are indeed tighter than the kl inequality (where kl is applied after rescaling Z to the $[0, 1]$ interval). But, if the predictions differ a lot, then Z primarily takes values at the endpoints $\{-1, 1\}$, and the kl inequality is tighter.

The aim of exploiting low variance while staying competitive with the kl inequality at high variance has challenged all prior work on Empirical Bernstein inequalities [Tolstikhin and Seldin, 2013, Mhammedi et al., 2019, Wu et al., 2021].

We present the split-kl inequality, which simultaneously matches the tightness of Empirical/Unexpected Bernstein and kl, and outperforms both for certain distributions of Z. It works for any bounded random variables, not just the ternary ones that we used to motivate the problem. The idea behind the split-kl inequality is to write $Z = \mu + Z^+ - Z^-$, where μ is a constant, $Z^+ = \max\{0, Z - \mu\}$, and $Z^- = \max\{0, \mu - Z\}$. Then $\mathbb{E}[Z] = \mu + \mathbb{E}[Z^+] - \mathbb{E}[Z^-]$ and, given an i.i.d. sample $Z_1, \ldots, Z_n$, we can bound the distance between $\frac{1}{n}\sum_{i=1}^n Z_i$ and $\mathbb{E}[Z]$ by using kl upper and lower bounds on the distances between $\frac{1}{n}\sum_{i=1}^n Z_i^+$ and $\mathbb{E}[Z^+]$, and $\frac{1}{n}\sum_{i=1}^n Z_i^-$ and $\mathbb{E}[Z^-]$, respectively. This approach provides the best of both worlds: the combinatorial tightness of the kl bound and exploitation of low variance, as in Empirical Bernstein inequalities. We further elevate the idea to the PAC-Bayes domain and derive a PAC-Bayes-split-kl inequality.

We present an extensive set of experiments, where we first compare the basic kl, Empirical Bernstein, Unexpected Bernstein, and split-kl on simulated data, and then compare PAC-Bayes-kl, PAC-Bayes-Unexpected-Bersnstein, PAC-Bayes-split-kl, and, in some of the setups, PAC-Bayes-Empirical-Bennett, for several prediction models on several UCI datasets. In particular, we evaluate the bounds in the linear classification setup studied by Mhammedi et al. [2019] and in the weighted majority prediction setup studied by Wu et al. [2021]. To the best of our knowledge, this is also the first time when Empirical Bernstein and Unexpected Bernstein are directly compared, with and without the PAC-Bayesian extension.

2 Basic Concentration of Measure Inequalities

We start with the most basic question in probability theory and statistics: how far can an average of an i.i.d. sample $Z_1, \ldots, Z_n$ deviate from its expectation? We cite the major existing inequalities, the kl, Empirical Bernstein, and Unexpected Bernstein, then derive the new split-kl inequality, and then provide a numerical comparison.

2.1 Background

We use $\mathrm{KL}(\rho\|\pi)$ to denote the Kullback-Leibler divergence between two probability distributions, ρ and π. We further use $\mathrm{kl}(p\|q)$ as a shorthand for the Kullback-Leibler divergence between two Bernoulli distributions with biases p and q, namely $\mathrm{kl}(p\|q) = \mathrm{KL}((1-p, p)\|(1-q, q))$. For $\hat{p} \in [0, 1]$ and $\varepsilon \geq 0$ we define the upper and lower inverse of kl, respectively, as $\mathrm{kl}^{-1,+}(\hat{p}, \varepsilon) := \max\{p \leq 1 : \mathrm{kl}(\hat{p}\|p) \leq \varepsilon\}$ and $\mathrm{kl}^{-1,-}(\hat{p}, \varepsilon) := \min\{p \geq 0 : \mathrm{kl}(\hat{p}\|p) \leq \varepsilon\}$.

The first inequality that we cite is the kl inequality.

Theorem 1 (kl Inequality [Langford, 2005, Foong et al., 2021, 2022]). *Let $Z_1, \cdots, Z_n$ be i.i.d. random variables bounded in the $[0, 1]$ interval and with $\mathbb{E}[Z_i] = p$ for all i. Let $\hat{p} = \frac{1}{n}\sum_{i=1}^n Z_i$ be the empirical mean. Then, for any $\delta \in (0, 1)$:*

$$\mathbb{P}\left(p \geq \mathrm{kl}^{-1,+}\left(\hat{p}, \frac{1}{n}\ln\frac{1}{\delta}\right)\right) \leq \delta, \tag{1}$$

$$\mathbb{P}\left(p \leq \mathrm{kl}^{-1,-}\left(\hat{p}, \frac{1}{n}\ln\frac{1}{\delta}\right)\right) \leq \delta. \tag{2}$$

We note that the PAC-Bayes-kl inequality (Theorem 5 below) is based on the inequality $\mathbb{E}\left[e^{n\,\mathrm{kl}(\hat{p}\|p)}\right] \leq 2\sqrt{n}$ [Maurer, 2004], which gives $\mathbb{P}\left(\mathrm{kl}(\hat{p}\|p) \geq \frac{\ln\frac{2\sqrt{n}}{\delta}}{n}\right) \leq \delta$. Foong et al. [2021, 2022] reduce the logarithmic factor down to $\ln\frac{1}{\delta}$ by basing the proof on Chernoff's inequality, but this proof technique cannot be combined with PAC-Bayes. Therefore, when we move on to PAC-Bayes we pay the extra $\ln 2\sqrt{n}$ factor in the bounds.

Next we cite two versions of the Empirical Bernstein inequality.

Theorem 2 (Empirical Bernstein Inequality [Maurer and Pontil, 2009]). *Let $Z_1, \cdots, Z_n$ be i.i.d. random variables bounded in the $[a, b]$ interval for some $a, b \in \mathbb{R}$, and with $\mathbb{E}[Z_i] = p$ for all i. Let*

84    $\hat{p} = \frac{1}{n} \sum_{i=1}^{n} Z_i$ *be the empirical mean and let* $\hat{\sigma} = \frac{1}{n-1} \sum_{i=1}^{n} (Z_i - \hat{p})^2$ *be the empirical variance.*

85   *Then for any* $\delta \in (0,1)$:

$$\mathbb{P}\left(p \geq \hat{p} + \sqrt{\frac{2\hat{\sigma} \ln \frac{2}{\delta}}{n}} + \frac{7(b-a) \ln \frac{2}{\delta}}{3(n-1)} \right) \leq \delta. \tag{3}$$

86 **Theorem 3** (Unexpected Bernstein Inequality [Fan et al., 2015, Mhammedi et al., 2019])**.** *Let*
87 $Z_1, \cdots, Z_n$ *be i.i.d. random variables bounded from above by* b *for some* $b > 0$, *and with* $\mathbb{E}[Z_i] = p$
88 *for all* i. *Let* $\hat{p} = \frac{1}{n} \sum_{i=1}^{n} Z_i$ *be the empirical mean and let* $\hat{\sigma} = \frac{1}{n} \sum_{i=1}^{n} Z_i^2$ *be the empirical mean*
89 *of the second moments. Let* $\psi(u) := u - \ln(1+u)$ *for any* $u \in \mathbb{R}$. *Then, for any* $\gamma \in (0, 1/b)$ *and*
90 *any* $\delta \in (0,1)$:

$$\mathbb{P}\left(p \geq \hat{p} + \frac{\psi(-\gamma b)}{\gamma b^2} \hat{\sigma} + \frac{\ln \frac{1}{\delta}}{\gamma n} \right) \leq \delta. \tag{4}$$

91 To facilitate a comparison with other bounds, Theorem 3 provides a slightly different form of the
92 Unexpected Bernstein inequality than the one used by Mhammedi et al. [2019]. We provide a proof
93 of the theorem in Appendix A, which is based on the Unexpected Bernstein Lemma [Fan et al., 2015].

2.2 The Split-kl Inequality

95 Let Z be a random variable bounded in a $[a, b]$ interval for some $a, b \in \mathbb{R}$ and let $\mu \in [a, b]$
96 be a constant. We decompose $Z = \mu + Z^+ - Z^-$, where $Z^+ = \max(0, Z - \mu)$ and $Z^- = $
97 $\max(0, \mu - Z)$. Let $p = \mathbb{E}[Z]$, $p^+ = \mathbb{E}[Z^+]$, and $p^- = \mathbb{E}[Z^-]$. For an i.i.d. sample $Z_1, \ldots, Z_n$ let
98 $\hat{p}^+ = \frac{1}{n} \sum_{i=1}^{n} Z_i^+$ and $\hat{p}^- = \frac{1}{n} \sum_{i=1}^{n} Z_i^-$.

99 With these definitions we present the split-kl inequality.

100 **Theorem 4** (Split kl inequality)**.** *Let* $Z_1, \ldots, Z_n$ *be i.i.d. random variables in a* $[a, b]$ *interval for*
101 *some* $a, b \in \mathbb{R}$, *then for any* $\mu \in [a, b]$ *and* $\delta \in (0,1)$:

$$\mathbb{P}\left(p \geq \mu + (b - \mu)\, \mathrm{kl}^{-1,+}\left(\frac{\hat{p}^+}{b-\mu}, \frac{1}{n} \ln \frac{2}{\delta} \right) - (\mu - a)\, \mathrm{kl}^{-1,-}\left(\frac{\hat{p}^-}{\mu-a}, \frac{1}{n} \ln \frac{2}{\delta} \right) \right) \leq \delta. \tag{5}$$

Proof.

$$\mathbb{P}\left(p \geq \mu + (b - \mu)\, \mathrm{kl}^{-1,+}\left(\frac{\hat{p}^+}{b-\mu}, \frac{1}{n} \ln \frac{2}{\delta} \right) - (\mu - a)\, \mathrm{kl}^{-1,-}\left(\frac{\hat{p}^-}{\mu-a}, \frac{1}{n} \ln \frac{2}{\delta} \right) \right)$$

$$\leq \mathbb{P}\left(p^+ \geq (b - \mu)\, \mathrm{kl}^{-1,+}\left(\frac{\hat{p}^+}{b-\mu}, \frac{1}{n} \ln \frac{2}{\delta} \right) \right) + \mathbb{P}\left(p^- \leq (\mu - a)\, \mathrm{kl}^{-1,-}\left(\frac{\hat{p}^-}{\mu-a}, \frac{1}{n} \ln \frac{2}{\delta} \right) \right)$$

$$\leq \delta,$$

102 where the last inequality follows by application of the kl upper and lower bounds from Theorem 1 to
103 the first and the second terms in the middle line, respectively.

∎

2.3 Empirical Comparison

105 We present an empirical comparison of the tightness of the above four concentration inequalities:
106 the kl, the Empirical Bernstein, the Unexpected Bernstein, and the split-kl. We take n i.i.d. samples
107 $Z_1, \ldots, Z_n$ taking values in $\{-1, 0, 1\}$. The choice is motivated both by instructiveness of presenta-
108 tion and by subsequent applications to excess losses. We let $p_{-1} = \mathbb{P}(Z = -1)$, $p_0 = \mathbb{P}(Z = 0)$, and
109 $p_1 = \mathbb{P}(Z = 1)$, where $p_{-1} + p_0 + p_1 = 1$. Then $p = \mathbb{E}[Z] = p_1 - p_{-1}$. We also let $\hat{p} = \frac{1}{n} \sum_{i=1}^{n} Z_i$.

110 In Figure 1 we plot the difference between the bounds on p given by the inequalities (1), (3), (4), and
111 (5), and $\hat{p}$. Lower values in the plot correspond to tighter bounds. To compute the kl bound we first
112 rescale the losses to the $[0, 1]$ interval, and then rescale the bound back to the $[-1, 1]$ interval. For
113 the Empirical Bernstein bound we take $a = -1$ and $b = 1$. For the Unexpected Bernstein bound we
114 take a grid of $\gamma \in \{1/(2b), \cdots, 1/(2^k b)\}$ for $k = \lceil \log_2(\sqrt{n/\ln(1/\delta)}/2) \rceil$ and a union bound over

(a) Comparison of the concentration bounds with the number of sample $n = 100$, and the probabilities $p_{-1} = p_1 = 0.5(1 - p_0)$.

(b) Comparison of the concentration bounds with the number of sample $n = 100$ and the probabilities $p_1 = 0.99(1 - p_0)$, $p_{-1} = 0.01(1 - p_0)$.

Figure 1: Empirical comparison of concentration bounds.

the grid, as proposed by Mhammedi et al. [2019]. For the split-kl bound we take μ to be the middle value of the ternary random variable 0. In the experiments we take $\delta = 0.05$, and cut the bounds to 1.

In the first experiment, presented in Figure 1a, we take $p_{-1} = p_1 = (1 - p_0)/2$ and plot the difference between the values of the bounds and $\hat{p}$ as a function of p_0. For $p_0 = 0$ the random variable Z becomes Bernoulli and, as expected, the kl inequality performs the best, followed by split-kl, and then Unexpected Bernstein. As p_0 grows closer to 1, the variance of Z decreases and, also as expected, the kl inequality falls behind, whereas split-kl and Unexpected Bernstein go closely together. Empirical Bernstein falls behind all other bounds throughout most of the range, except slightly outperforming kl when p_0 gets very close to 1.

In the second experiment, presented in Figure 1b, we take a skewed random variable with $p_1 = 0.99(1 - p_0)$ and $p_{-1} = 0.01(1 - p_0)$, and again plot the difference between the values of the bounds and $\hat{p}$ as a function of p_0. This time the kl also starts well for p_0 close to zero, but then falls behind due to its inability of properly handling the values inside the interval. Unexpected Bernstein exhibits the opposite trend due to being based on uncentered second moment, which is high when p_0 is close to zero, even though the variance is small in this case. Empirical Bernstein lags behind all other bounds for most of the range due to poor constants, whereas split-kl matches the tightest bounds, kl and Unexpected Bernstein, at the endpoints of the range of p_0 and outperforms all other bounds in the middle of the range, around $p_0 = 0.6$, due to being able to exploit the combinatorics of the problem.

The experiments demonstrate that split-kl is a powerful alternative to existing concentration of measure inequalities. To the best of our knowledge, this is also the first empirical evaluation of the Unexpected Bernstein inequality, and it shows that in many cases it is also a powerful inequality. We also observe that in most settings the Empirical Bernstein is weaker than the other three inequalities we consider, and it is always weaker than the split-kl inequality in our experiments. More numerical evaluations are provided in Appendix D.

3 PAC-Bayesian Inequalities

Now we elevate the basic concentration of measure inequalities to the PAC-Bayesian domain. We start with the supervised learning problem setup, then provide a background on existing PAC-Bayesian inequalities, and finish with presentation of the PAC-Bayes-split-kl inequality.

3.1 Supervised Learning Problem Setup and Notations

Let $\mathcal{X}$ be a sample space, $\mathcal{Y}$ be a label space, and let $S = \{(X_i, Y_i)\}_{i=1}^{n}$ be an i.i.d. sample drawn according to an unknown distribution $\mathcal{D}$ on the product-space $\mathcal{X} \times \mathcal{Y}$. Let $\mathcal{H}$ be a hypothesis space containing hypotheses $h : \mathcal{X} \to \mathcal{Y}$. The quality of a hypothesis h is measured using the zero-one loss $\ell(h(X), Y) = \mathbb{1}(h(X) \neq Y)$, where $\mathbb{1}(\cdot)$ is the indicator function. The expected loss of h is

denoted by $L(h) = \mathbb{E}_{(X,Y)\sim\mathcal{D}}\left[\ell(h(X), Y)\right]$, and the empirical loss of h on a sample S is denoted by $\hat{L}(h, S) = \frac{1}{|S|}\sum_{(X,Y)\in S}\ell(h(X), Y)$. We use $\mathbb{E}_{\mathcal{D}}[\cdot]$ as a shorthand for $\mathbb{E}_{(X,Y)\sim\mathcal{D}}[\cdot]$.

PAC-Bayesian bounds bound the generalization error of Gibbs prediction rules. For each input $X \in \mathcal{X}$, Gibbs prediction rule associated with a distribution ρ on $\mathcal{H}$ randomly draws a hypothesis $h \in \mathcal{H}$ according to ρ and predicts $h(X)$. The expected loss of the Gibbs prediction rule is $\mathbb{E}_{h\sim\rho}[L(h)]$ and the empirical loss is $\mathbb{E}_{h\sim\rho}[\hat{L}(h, S)]$. We use $\mathbb{E}_{\rho}[\cdot]$ as a shorthand for $\mathbb{E}_{h\sim\rho}[\cdot]$.

3.2 PAC-Bayesian Analysis Background

Now we present a brief background on the relevant results from the PAC-Bayesian analysis.

PAC-Bayes-kl **Inequality**   The PAC-Bayes-kl inequality cited below is one of the tightest known generalization bounds on the expected loss of the Gibbs prediction rule.

Theorem 5 (PAC-Bayes-kl Inequality, Seeger, 2002, Maurer, 2004)**.** *For any probability distribution π on $\mathcal{H}$ that is independent of S and any $\delta \in (0, 1)$:*

$$\mathbb{P}\left(\exists\rho \in \mathcal{P} : \mathrm{kl}\left(\mathbb{E}_{\rho}[\hat{L}(h, S)]\middle\|\mathbb{E}_{\rho}\left[L(h)\right]\right) \geq \frac{\mathrm{KL}(\rho\|\pi) + \ln(2\sqrt{n}/\delta)}{n}\right) \leq \delta, \tag{6}$$

where $\mathcal{P}$ is the set of all possible probability distributions on $\mathcal{H}$ that can depend on S.

The following relaxation of the PAC-Bayes-kl inequality based on Refined Pinsker's relaxation of the kl divergence helps getting some intuition about the bound [McAllester, 2003]. With probability at least $1 - \delta$, for all $\rho \in \mathcal{P}$ we have

$$\mathbb{E}_{\rho}[L(h)] \leq \mathbb{E}_{\rho}[\hat{L}(h, S)] + \sqrt{2\mathbb{E}_{\rho}[\hat{L}(h, S)]\frac{\mathrm{KL}(\rho\|\pi) + \ln(2\sqrt{n}/\delta)}{n}} + \frac{2\left(\mathrm{KL}(\rho\|\pi) + \ln(2\sqrt{n}/\delta)\right)}{n}.$$

If $\mathbb{E}_{\rho}[\hat{L}(h, S)]$ is close to zero, the middle term in the inequality above vanishes, leading to so-called "fast convergence rates" (convergence of $\mathbb{E}_{\rho}[\hat{L}(h, S)]$ to $\mathbb{E}_{\rho}[L(h)]$ at the rate of $1/n$). However, achieving low $\mathbb{E}_{\rho}[\hat{L}(h, S)]$ is not always possible [Dziugaite and Roy, 2017, Zhou et al., 2019]. Subsequent research in PAC-Bayesian analysis has focused on two goals: (1) achieving fast convergence rates when the variance of prediction errors is low (and not necessarily the errors themselves), and (2) reducing the $\mathrm{KL}(\rho\|\pi)$ term, which may be quite large for large hypothesis spaces. For the first goal Tolstikhin and Seldin [2013] developed the PAC-Bayes-Empirical-Bernstein inequality and Mhammedi et al. [2019] proposed to use excess losses and also derived the alternative PAC-Bayes-Unexpected-Bernstein inequality. For the second goal Ambroladze et al. [2007] suggested to use informed priors and Mhammedi et al. [2019] perfected the idea by proposing to average over "forward" and "backward" construction with informed prior. Next we explain the ideas behind the excess losses and informed priors in more details.

Excess Losses   Let h^* be a reference prediction rule that is independent of S. We define the excess loss of a prediction rule h with respect to the reference h^* by

$$\Delta_{\ell}(h(X), h^*(X), Y) = \ell(h(X), Y) - \ell(h^*(X), Y).$$

If ℓ is the zero-one loss, the excess loss naturally gives rise to ternary random variables, but it is well-defined for any real-valued loss function. We use $\Delta_L(h, h^*) = \mathbb{E}_{\mathcal{D}}[\Delta_{\ell}(h(X), h^*(X), Y)] = L(h) - L(h^*)$ to denote the expected excess loss of h relative to h^* and $\Delta_{\hat{L}}(h, h', S) = \frac{1}{|S|}\sum_{(X,Y)\in S}\Delta_{\ell}(h(X), h^*(X), Y) = \hat{L}(h) - \hat{L}(h^*)$ to denote the empirical excess loss of h relative to h^*. The expected loss of a Gibbs prediction rule can then be written as

$$\mathbb{E}_{\rho}[L(h)] = \mathbb{E}_{\rho}[\Delta_L(h, h^*)] + L(h^*).$$

A bound on $\mathbb{E}_{\rho}[L(h)]$ can thus be decomposed into a summation of a PAC-Bayes bound on $\mathbb{E}_{\rho}[\Delta_L(h, h^*)]$ and a bound on $L(h^*)$. When the variance of the excess loss is small we can use tools that exploit small variance, such as PAC-Bayes-Empirical-Bernstein, PAC-Bayes-Unexpected-Bernstein, or PAC-Bayes-Split-kl proposed below, to achieve fast convergence rates for the excess loss. Bounding $L(h^*)$ involves just a single prediction rule and does not depend on the value of $\mathrm{KL}(\rho\|\pi)$. We note that it is essential that the variance and not just the magnitude of the excess loss is small. For example, if the excess losses primarily take values in $\{-1, 1\}$ and average up to zero, fast convergence rates are impossible.

Informed Priors The idea behind informed priors is to split the data into two subsets $S = S_1 \cup S_2$ and to use S_1 to learn a prior π_{S_1}, and then use it to learn a posterior on S_2 Ambroladze et al. [2007]. Note that since the size of S_2 is smaller than the size of S, this approach gains in having potentially smaller $\mathrm{KL}(\rho\|\pi_{S_1})$, but loses in having a smaller sample size in the denominator of PAC-Bayes bounds. The balance between the advantage and disadvantage depends on the data: for some data sets it strengthens the bounds, but for some it weakens them. Mhammedi et al. [2019] perfected the approach by proposing to use it in the "forward" and "backward" direction and average over the two. Let S_1 and S_2 be of equal size. The "forward" part uses S_1 to train π_{S_1} and then computes a posterior on S_2, while the "backward" part uses S_2 to train π_{S_2} and then computes a posterior on S_1. Finally, the two posteriors are averaged with equal weight and the KL term becomes $\frac{1}{2}\left(\mathrm{KL}(\rho\|\pi_{S_1}) + \mathrm{KL}(\rho\|\pi_{S_2})\right)$. See [Mhammedi et al., 2019] for the derivation.

Excess Losses and Informed Priors Excess losses and informed priors make an ideal combination. If we split S into two equal parts $S = S_1 \cup S_2$, we can use S_1 to train both a reference prediction rule h_{S_1} and a prior π_{S_1}, and then learn a PAC-Bayes posterior on S_2, and the other way around. By combining the "forward" and "backward" approaches we can write

$$\mathbb{E}_\rho[L(h)] = \frac{1}{2}\mathbb{E}_\rho[\Delta_L(h, h_{S_1})] + \frac{1}{2}\mathbb{E}_\rho[\Delta_L(h, h_{S_2})] + \frac{1}{2}\left(L(h_{S_1}) + L(h_{S_2})\right) \tag{7}$$

and we can use PAC-Bayes to bound the first term using the prior π_{S_1} and the data in S_2 and to bound the second term using the prior π_{S_2} and the data in S_1, and we can bound $L(h_{S_1})$ and $L(h_{S_2})$ using "complementary" data in S_2 and S_1, respectively.

PAC-Bayes-Empirical-Bernstein Inequalities The excess losses are ternary random variables taking values in $\{-1, 0, 1\}$ and, as we have already discussed, the kl inequality is not well-suited for them. PAC-Bayesian inequalities tailored for non-binary random variables were derived by Seldin et al. [2012], Tolstikhin and Seldin [2013], Wu et al. [2021], and Mhammedi et al. [2019]. Seldin et al. [2012] derived the PAC-Bayes-Bernstein oracle bound, which assumes knowledge of the variance. Tolstikhin and Seldin [2013] made it into an empirical bound by deriving the PAC-Bayes-Empirical-Bernstein bound for the variance and plugging it into the PAC-Bayes-Bernstein bound of Seldin et al.. Wu et al. [2021] derived an oracle PAC-Bayes-Bennett inequality, which again assumes oracle knowledge of the variance, and showed that it is always at least as tight as the PAC-Bayes-Bernstein, and then also plug in the PAC-Bayes-Empirical-Bernstein bound for the variance. Mhammedi et al. [2019] derived the PAC-Bayes-Unexpected-Bernstein inequality, which directly uses the empirical second moment. Since we have already shown that the Unexpected Bernstein inequality is tighter than the Empirical Bernstein, and since the approach of Wu et al. requires a combination of two inequalities, PAC-Bayes-Empirical-Bernstein for the variance and PAC-Bayes-Bennett for the loss, whereas the approach of Mhammedi et al. only makes a single application of PAC-Bayes-Unexpected-Bernstein, we only compare our work to the latter.

We cite the inequality of Mhammedi et al. [2019], which applies to an arbitrary loss function. We use $\tilde{\ell}$ and matching tilde-marked quantities to distinguish it from the zero-one loss ℓ. For any $h \in \mathcal{H}$, let $\tilde{L}(h) = \mathbb{E}_D[\tilde{\ell}(h(X), Y)]$ be the expected tilde-loss of h and let $\hat{\tilde{L}}(h, S) = \frac{1}{|S|}\sum_{(X,Y)\in S}\tilde{\ell}(h(X), Y)$ be the empirical tilde-loss of h on a sample S.

Theorem 6 (PAC-Bayes-Unexpected-Bernstein inequality [Mhammedi et al., 2019]). *Let $\tilde{\ell}(\cdot, \cdot)$ be an arbitrary loss function bounded from above by b for some $b > 0$, and assume that $\hat{\tilde{\mathbb{V}}}(h, S) = \frac{1}{|S|}\sum_{(X,Y)\in S}\tilde{\ell}(h(X), Y)^2$ is finite for all h. Let $\psi(u) := u - \ln(1 + u)$ for any $u \in \mathbb{R}$. Then for any distribution π on $\mathcal{H}$ that is independent of S, any $\gamma \in (0, 1/b)$, and any $\delta \in (0, 1)$:*

$$\mathbb{P}\left(\exists \rho \in \mathcal{P} : \mathbb{E}_\rho[\tilde{L}(h)] \geq \mathbb{E}_\rho[\hat{\tilde{L}}(h, S)] + \frac{\psi(-\gamma b)}{\gamma b^2}\mathbb{E}_\rho[\hat{\tilde{\mathbb{V}}}(h, S)] + \frac{\mathrm{KL}(\rho\|\pi) + \ln\frac{1}{\delta}}{\gamma n}\right) \leq \delta,$$

where $\mathcal{P}$ is the set of all possible probability distributions on $\mathcal{H}$ that can depend on S.

In optimization of the bound, we take the same grid of $\gamma \in \{1/(2b), \cdots, 1/(2^k b)\}$ for $k = \lceil\log_2(\sqrt{n/\ln(1/\delta)}/2)\rceil$ and a union bound over the grid, as we did for Theorem 3.

3.3 PAC-Bayes-Split-kl Inequality

Now we present our PAC-Bayes-Split-kl inequality. For an arbitrary loss function $\tilde{\ell}$ taking values in a $[a, b]$ interval for some $a, b \in \mathbb{R}$, let $\tilde{\ell}^+ := \max\{0, \tilde{\ell} - \mu\}$ and $\tilde{\ell}^- := \max\{0, \mu - \tilde{\ell}\}$ for some $\mu \in [a, b]$. For any $h \in \mathcal{H}$, let $\tilde{L}^+(h) = \mathbb{E}_D[\tilde{\ell}^+(h(X), Y)]$ and $\tilde{L}^-(h) = \mathbb{E}_D[\tilde{\ell}^-(h(X), Y)]$. The corresponding empirical losses are denoted by $\hat{\tilde{L}}^+(h, S) = \frac{1}{n}\sum_{i=1}^n \tilde{\ell}^+(h(X_i), Y_i)$ and $\hat{\tilde{L}}^-(h, S) = \frac{1}{n}\sum_{i=1}^n \tilde{\ell}^-(h(X_i), Y_i)$.

Theorem 7 (PAC-Bayes-Split-kl Inequality)**.** *Let $\tilde{\ell}(\cdot, \cdot)$ be arbitrary loss function taking values in a $[a, b]$ interval for some $a, b \in \mathbb{R}$. Then for any distribution π on $\mathcal{H}$ that is independent of S, any $\mu \in [a, b]$, and any $\delta \in (0, 1)$:*

$$
\mathbb{P}\left[\exists \rho \in \mathcal{P} : \mathbb{E}_\rho[\tilde{L}(h)] \geq \mu + (b - \mu)\,\mathrm{kl}^{-1,+}\left(\frac{\mathbb{E}_\rho[\hat{\tilde{L}}^+(h, S)]}{b - \mu}, \frac{\mathrm{KL}(\rho\|\pi) + \ln\frac{4\sqrt{n}}{\delta}}{n}\right)\right.
$$
$$
\left. - (\mu - a)\,\mathrm{kl}^{-1,-}\left(\frac{\mathbb{E}_\rho[\hat{\tilde{L}}^-(h, S)]}{\mu - a}, \frac{\mathrm{KL}(\rho\|\pi) + \ln\frac{4\sqrt{n}}{\delta}}{n}\right)\right] \leq \delta,
$$

where $\mathcal{P}$ is the set of all possible probability distributions on $\mathcal{H}$ that can depend on S.

Proof. We have $\mathbb{E}_\rho[\tilde{L}(h)] = \mu + \mathbb{E}_\rho[\tilde{L}^+(h)] - \mathbb{E}_\rho[\tilde{L}^-(h)]$. Similar to the proof of Theorem 4, we take a union bound of PAC-Bayes-kl upper bound on $\mathbb{E}_\rho[\tilde{L}^+(h)]$ and PAC-Bayes-kl lower bound on $\mathbb{E}_\rho[\tilde{L}^-(h)]$. ∎

3.4 PAC-Bayes-split-kl with Excess Loss and Informed Prior

Looking back at the expected loss decomposition in equation (7), we can use PAC-Bayes-split-kl to bound the first two terms and a bound on the binomial tail distribution to bound the last term. For n i.i.d. Bernoulli random variables $Z_1, \ldots, Z_n$ with bias $p \in (0, 1)$, we define the binomial tail distribution $\mathsf{Bin}(n, k, p) = \mathbb{P}(\sum_{i=1}^n X_i \leq k)$ and its inverse $\mathsf{Bin}^{-1}(n, k, \delta) = \max\{p \leq 1 : \mathsf{Bin}(n, k, p) \geq \delta\}$. The following theorem relates $\hat{p} = \frac{1}{n}\sum_{i=1}^n Z_i$ and p.

Theorem 8 (Test Set Bound [Langford, 2005])**.** *Let $Z_1, \ldots, Z_n$ be n i.i.d. Bernoulli random variables with bias $p \in (0, 1)$ and let $\hat{p} = \frac{1}{n}\sum_{i=1}^n Z_i$ be the empirical mean. Then for any $\delta \in (0, 1)$:*

$$
\mathbb{P}\left(p \geq \mathsf{Bin}^{-1}(n, n\hat{p}, \delta)\right) \leq \delta.
$$

By applying Theorems 7 and 8 to equation (7) we obtain the following result.

Theorem 9. *For any $\mu \in [-1, 1]$ and any $\delta \in (0, 1)$:*

$$
\mathbb{P}\left(\exists \rho \in \mathcal{P} : \mathbb{E}_\rho[L(h)] \geq \mu + (1 - \mu)(a) - (\mu + 1)(b) + \frac{1}{2}(c)\right) \leq \delta,
$$

where $\mathcal{P}$ is the set of all possible probability distributions on $\mathcal{H}$ that can depend on S,

$$
(a) = \mathrm{kl}^{-1,+}\left(\frac{1}{2}\frac{\mathbb{E}_\rho[\Delta_{\hat{L}}^+(h, h_{S_1}, S_2)]}{1 - \mu} + \frac{1}{2}\frac{\mathbb{E}_\rho[\Delta_{\hat{L}}^+(h, h_{S_2}, S_1)]}{1 - \mu}, \frac{\mathrm{KL}(\rho\|\pi) + \ln\frac{8\sqrt{n/2}}{\delta}}{n/2}\right),
$$

$$
(b) = \mathrm{kl}^{-1,-}\left(\frac{1}{2}\frac{\mathbb{E}_\rho[\Delta_{\hat{L}}^-(h, h_{S_1}, S_2)]}{\mu + 1} + \frac{1}{2}\frac{\mathbb{E}_\rho[\Delta_{\hat{L}}^-(h, h_{S_2}, S_1)]}{\mu + 1}, \frac{\mathrm{KL}(\rho\|\pi) + \ln\frac{8\sqrt{n/2}}{\delta}}{n/2}\right),
$$

in which $\pi = \frac{1}{2}\pi_{S_1} + \frac{1}{2}\pi_{S_2}$, and

$$
(c) = \mathsf{Bin}^{-1}\left(\frac{n}{2}, \frac{n}{2}\hat{L}(h_{S_1}, S_2), \frac{\delta}{4}\right) + \mathsf{Bin}^{-1}\left(\frac{n}{2}, \frac{n}{2}\hat{L}(h_{S_2}, S_1), \frac{\delta}{4}\right).
$$

The proof is postponed to Appendix C.

Figure 2: Comparison of the bounds and the test losses of the optimized Gaussian posterior ρ^* generated by PBkl with informed priors, PBUB with excess losses and informed priors, and PBSkl with excess losses and informed priors. The test losses of the corresponding bounds are shown in black. We report the mean and the standard deviation over 20 runs of the experiments.

4 Experiments

We investigate the performance of the PAC-Bayes-split-kl inequality under two settings on several data sets from UCI and LibSVM repositories [Dua and Graff, 2019, Chang and Lin, 2011]. An overview of the data sets is provided in Appendix E.1. We reproduce the experimental setups of Mhammedi et al. [2019] and Wu et al. [2021].

4.1   The Experimental Setup of Mhammedi et al. [2019]: Linear Classifiers

In the first experiment we follow the experimental setup of Mhammedi et al. [2019], who consider binary classification problems with linear classifiers in $\mathbb{R}^d$ and Gaussian priors and posteriors. A classifier h_w associated with a vector $w \in \mathbb{R}^d$ makes a prediction on an input X by $h_w(X) = \mathbb{1}\big(\phi(w^\top X) > 0.5\big)$, where $\phi(x) := 1/(1 + e^{-x})$ for $x \in \mathbb{R}$ is the sigmoid function. The posteriors have the form of Gaussian distribution centered at $w_S \in \mathbb{R}^d$ with a covariance Σ_S that depend on a sample S, $\rho = \mathcal{N}(w_S, \Sigma_S)$. The informed priors $\pi_{S_1} = \mathcal{N}(w_{S_1}, \Sigma_{S_1})$ and $\pi_{S_2} = \mathcal{N}(w_{S_2}, \Sigma_{S_2})$ are also taken to be Gaussian distributions centered at w_{S_1} and w_{S_2}, with covariance Σ_{S_1} and Σ_{S_2}, respectively. We take the classifier associated with w_{S_1} as the reference classifier h_{S_1} and the classifier associated with w_{S_2} as the reference classifier h_{S_2}. More details on the construction are provided in Appendix E.2.

Figure 2 compares the PAC-Bayes-Unexpected-Bernstein bound PBUB and the PAC-Bayes-split-kl bound PBSkl with excess losses and informed priors. The ternary random variable in this setup takes a value in $\{-1, 0, 1\}$ and we select μ to be the middle value 0. Since the PAC-Bayes-kl bound PBkl is one of the tightest known generalization bounds, we take PBkl with informed priors as a baseline. The details on bound calculation and optimization are provided in Appendix E.2.

4.2   The Experimental Setup of Wu et al. [2021]: Weighted Majority Vote

In the second experiment we reproduce the experimental setup of Wu et al. [2021], who consider multiclass classification by a weighted majority vote. Given an input $X \in \mathcal{X}$, a hypothesis space $\mathcal{H}$, and a distribution ρ on $\mathcal{H}$, a ρ-weighted majority vote classifier predicts $\mathrm{MV}_\rho(X) = \arg\max_{y \in \mathcal{Y}} \mathbb{E}_\rho[\mathbb{1}(h(X) = y)]$. One of the tightest bound for the majority vote is the tandem bound (TND) proposed by Masegosa et al. [2020], which is based on tandem losses for pairs of hypotheses, $\ell(h(X), h'(X), Y) = \mathbb{1}(h(X) \neq Y)\mathbb{1}(h'(X) \neq Y)$, and the second order Markov's inequality. Wu et al. [2021] proposed two improved forms of the bound, both based on a parametric form of the Chebyshev-Cantelli inequality. The first, CCTND, using Chebyshev-Cantelli with the tandem losses

Figure 3: Comparison of the bounds and the test losses of the weighted majority vote on ensembles of heterogeneous classifiers with optimized posterior ρ^* generated by TND, CCTND, CCPBB, CCPBUB, and CCPBSkl. The test losses of the corresponding bounds are shown in black. We report the mean and the standard deviation over 10 runs of the experiments.

and the PAC-Bayes-kl bound for bounding the tandem losses. The second, CCPBB, using tandem losses with an offset, defined by $\ell_\alpha(h(X), h'(X), Y) = (\mathbb{1}(h(X) \neq Y) - \alpha)(\mathbb{1}(h'(X) \neq Y) - \alpha)$ for $\alpha < 0.5$, and PAC-Bayes-Empirical-Bennett inequality for bounding the tandem losses with an offset. We note that while the tandem losses are binary random variables, tandem losses with an offset are ternary random variables taking values in $\{\alpha^2, -\alpha(1 - \alpha), (1 - \alpha)^2\}$ and, therefore, application of Empirical Bernstein type inequalities makes sense. However, in the experiments of Wu et al. CCPBB lagged behind TND and CCTND. We replaced PAC-Bayes-Empirical-Bennett with PAC-Bayes-Unexpected-Bernstein (CCPBUB) and PAC-Bayes-split-kl (CCPBSkl) and showed that the weakness of CCPBB was caused by looseness of PAC-Bayes-Empirical-Bernstein, and that CCPBUB and CCPBSkl lead to tighter bounds that are competitive and sometimes outperforming TND and CCTND. For the PAC-Bayes-split-kl bound we took μ to be the middle value of the tandem loss with an offset, namely, for $\alpha \geq 0$ we took $\mu = \alpha^2$, and for $\alpha < 0$ we took $\mu = -\alpha(1 - \alpha)$.

In Figure 3 we present a comparison of the TND, CCTND, CCPBB, CCPBUB, and CCPBSkl bounds on weighted majority vote of heterogeneous classifiers (Linear Discriminant Analysis, k-Nearest Neighbors, Decision Tree, Logistic Regression, and Gaussian Naive Bayes), which adds the two new bounds, CCPBUB and CCPBSkl to the experiment done by Wu et al. [2021]. A more detailed description of the experiment and results for additional data sets are provided in Appendix E.3. In Appendix E.4 we replicate another experiment of Wu et al., where we use the bounds to reweigh trees in a random forest classifier. The results are similar to the results for heterogeneous classifiers.

5 Discussion

We have presented the split-kl and PAC-Bayes-split-kl inequalities. In our empirical evaluation the split-kl inequality was always competitive with the kl and Unexpected Bernstein inequalities and outperformed both in certain regimes, whereas Empirical Bernstein typically lagged behind. In the PAC-Bayesian setting PAC-Bayes-split-kl was always comparable to PAC-Bayes-Unexpected-Bernstein, whereas PAC-Bayes-Empirical-Bennett most often lagged behind. The first two inequalities were usually comparable to PAC-Bayes-kl, although in some cases the attempt to exploit low variance did not pay off and PAC-Bayes-kl outperformed, which is also the trend observed earlier by Mhammedi et al. [2019]. To the best of our knowledge, this is the first time when the various approaches to exploitation of low variance were directly compared, and the proposed split-kl emerged as a clear winner in the basic setting, whereas in the PAC-Bayes setting we have shown that PAC-Bayes-Unexpected-Bernstein and PAC-Bayes-split-kl are comparable, and preferable over the more commonly known and used PAC-Bayes-Empirical-Bernstein and PAC-Bayes-Empirical-Bennett.

References

Amiran Ambroladze, Emilio Parrado-Hernández, and John Shawe-Taylor. Tighter PAC-Bayes bounds. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2007.

Sergei N. Bernstein. *Probability Theory*. Moscow-Leningrad, 4^{th} edition, 1946. In Russian.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration Inequalities: A Nonasymptotic Theory of Independence*. Oxford University Press, 2013.

Chih-Chung Chang and Chih-Jen Lin. LIBSVM: A library for support vector machines. *ACM Transactions on Intelligent Systems and Technology*, 2, 2011.

Dheeru Dua and Casey Graff. UCI machine learning repository, 2019. URL http://archive.ics.uci.edu/ml.

Gintare Karolina Dziugaite and Daniel M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In *UAI*, 2017.

Xiequan Fan, Ion Grama, and Quansheng Liu. Exponential inequalities for martingales with applications. *Electronic Journal of Probability*, 20:1–22, 2015.

Andrew Foong, Wessel Bruinsma, David Burt, and Richard Turner. How tight can pac-bayes be in the small data regime? 2021.

Andrew Y. K. Foong, Wessel P. Bruinsma, and David R. Burt. A note on the chernoff bound for random variables in the unit interval. *arXiv preprint arXiv.2205.07880*, 2022.

W. Hoeffding. Probability inequalities for sums of bounded random variables. *Journal of the American Statistical Association*, 58(301):13–30, 1963.

John Langford. Tutorial on practical prediction theory for classification. *Journal of Machine Learning Research*, 6, 2005.

Andrés R. Masegosa, Stephan S. Lorenzen, Christian Igel, and Yevgeny Seldin. Second order PAC-Bayesian bounds for the weighted majority vote. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2020.

Andreas Maurer. A note on the PAC-Bayesian theorem. arXiv preprint cs/0411099, 2004.

Andreas Maurer and Massimiliano Pontil. Empirical Bernstein bounds and sample variance penalization. In *Proceedings of the Conference on Learning Theory (COLT)*, 2009.

David McAllester. PAC-Bayesian stochastic model selection. *Machine Learning*, 51, 2003.

Zakaria Mhammedi, Peter Grünwald, and Benjamin Guedj. PAC-Bayes un-expected Bernstein inequality. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2019.

Matthias Seeger. PAC-Bayesian generalization error bounds for Gaussian process classification. *Journal of Machine Learning Research*, 3, 2002.

Yevgeny Seldin, François Laviolette, Nicolò Cesa-Bianchi, John Shawe-Taylor, and Peter Auer. PAC-Bayesian inequalities for martingales. *IEEE Transactions on Information Theory*, 58, 2012.

Ilya Tolstikhin and Yevgeny Seldin. PAC-Bayes-Empirical-Bernstein inequality. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2013.

Yi-Shan Wu, Andres Masegosa, Stephan Lorenzen, Christian Igel, and Yevgeny Seldin. Chebyshev-cantelli pac-bayes-bennett inequality for the weighted majority vote. In *Advances in Neural Information Processing Systems (NeurIPS)*, 2021.

Wenda Zhou, Victor Veitch, Morgane Austern, Ryan P. Adams, and Peter Orbanz. Non-vacuous generalization bounds at the imagenet scale: a PAC-bayesian compression approach. In *ICLR*, 2019.