Near-Optimal Bounds for
Testing Histogram Distributions

Clément L. Canonne
University of Sydney
clement.canonne@sydney.edu.au

Ilias Diakonikolas*
University of Wisconsin-Madison
ilias@cs.wisc.edu

Daniel M. Kane†
University of California, San Diego
dakane@cs.ucsd.edu

Sihan Liu
University of California, San Diego
sil046@ucsd.edu

Abstract

We investigate the problem of testing whether a discrete probability distribution over an ordered domain is a histogram on a specified number of bins. One of the most common tools for the succinct approximation of data, *k-histograms* over $[n]$ are probability distributions that are piecewise constant over a set of k intervals. Given samples from an unknown distribution $\mathbf{p}$ on $[n]$, we want to distinguish between the cases that $\mathbf{p}$ is a k-histogram versus far from any k-histogram, in total variation distance. Our main result is a sample near-optimal and computationally efficient algorithm for this testing problem, and a nearly-matching (within logarithmic factors) sample complexity lower bound.

1 Introduction

1.1 Background and Motivation

A classical approach for the efficient exploration of massive datasets involves the construction of succinct data representations, see, e.g., the survey [CGHJ12]. One of the most useful and commonly used compact representations are *histograms*. For a dataset S, whose elements are from the universe $[n] := \{1, \dots, n\}$, a k-histogram is a function that is piecewise constant over k interval pieces. Histograms constitute the oldest and most popular synopsis structure in databases and have been extensively studied in the database community since their introduction in the 1980s [Koo80], see, e.g., [GMP97, JKM+98, CMN98, TGIK02, GGI+02, GKS06, ILR12, ADH+15, Can16], for a partial list of references. In both the statistics and computer science literatures, several methods have been proposed to estimate histogram distributions in a range of natural settings [Sco79, FD81, DL04, LN96, Kle09, CDSS14, ADH+15, ADLS17, DLS18].

In this work, we study the algorithmic task of deciding whether a (potentially very large) dataset S over the domain $[n]$ is a k-histogram (i.e., it has a succinct histogram representation with k interval pieces) or is "far" from *any* k-histogram representation (in a well-defined technical sense). Our focus is on *sublinear time* algorithms [Rub06]. Instead of reading the entire dataset S, which could be highly impractical, one can instead use randomness to sample a small subset of the dataset. Note that

---

*Supported by NSF Medium Award CCF-2107079, NSF Award CCF-1652862 (CAREER), a Sloan Research Fellowship, and a DARPA Learning with Less Labels (LwLL) grant. Some of this work was performed while the author was visiting the Simons Institute for the Theory of Computing.

†Supported by NSF Medium Award CCF-2107547, NSF Award CCF-1553288 (CAREER), and a Sloan Research Fellowship.

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

sampling a (uniformly) random element from S is equivalent to drawing a sample from the underlying probability distribution $\mathbf{p}$ of relative empirical frequencies. This observation brings our algorithmic problem of "histogram testing" in the framework of distribution property testing (statistical hypothesis testing) [BFR$^+$00, BFR$^+$13], see, e.g., [Can20] for a survey.

Formally, we study the following task: for an integer $1 \leq k \leq n$, denote by $\mathcal{H}_k^n$ the set of k-histogram distributions over $\{1, 2, \ldots, n\}$, i.e., the set of all distributions $\mathbf{p}$ such that there exists a partition of $[n]$ into k consecutive intervals (not necessarily of the same size) with $\mathbf{p}$ being uniform on each interval. Given access to i.i.d. samples from an unknown distribution $\mathbf{p}$ on $[n]$ and a desired error tolerance $0 < \varepsilon < 1$, we want to correctly distinguish (with high probability) between the cases that $\mathbf{p}$ is a k-histogram versus ε-far from any k-histogram, in total variation distance (or, equivalently, ℓ_1-norm). It should be noted that the histogram testing problem studied here is not new. Prior work within the algorithms and database theory community has investigated the complexity of the problem in the past ten years (see, e.g., [ILR12, ADH$^+$15, Can16] and Section 1.4 for a detailed summary of prior work). However, known algorithms for this task are sub-optimal, and in particular there is a polynomial gap between the best known upper and lower bounds on the sample complexity of the problem. At a high level, the difficulty of our histogram testing problem in the sub-linear regime lies in the fact that the location and "length" of the k intervals defining the histogram representation (if one exists) is a priori unknown to the algorithm.

We believe that the histogram testing problem is natural and interesting in its own right. Moreover, a sample-efficient algorithm for this testing task can be used as a key primitive in the context of *model selection*, where the goal is to identify the "most succinct" data representation. Indeed, various algorithms are known for learning k-histograms from samples whose sample complexities (and running times) scale proportionally to the succinctness parameter k (and are completely independent of the domain size n) [CDSS14, ADH$^+$15, ADLS17]. Combined with an efficient tester for the property of being a k-histogram (used to identify the smallest possible value of k such that $\mathbf{p}$ is a k-histogram, e.g., via binary search), one can obtain a sketch of the underlying dataset. See Appendix C for a detailed description.

1.2 Our Results

Our main contribution is a near-characterization of the sample complexity of the histogram testing problem. Specifically, we provide (1) a sample near-optimal and computationally efficient testing algorithm for the problem, and (2) a nearly-matching sample complexity lower bound (within logarithmic factors). In particular, we establish the following theorem:

Theorem 1 (Main Result). *There exists a testing algorithm for the class of k-histograms on $[n]$ with sample complexity $m = \widetilde{O}(\sqrt{nk}/\varepsilon + k/\varepsilon^2 + \sqrt{n}/\varepsilon^2)$ and running time $\mathrm{poly}(m)$. Moreover, for any $k \in [n]$ and $0 < \varepsilon < 1$, any testing algorithm for the class of k-histograms on $[n]$ requires at least $\widetilde{\Omega}(\sqrt{nk}/\varepsilon + k/\varepsilon^2 + \sqrt{n}/\varepsilon^2)$ samples.*

(The $\tilde{O}(\cdot)$ and $\tilde{\Omega}(\cdot)$ notation hides polylogarithmic factors in the argument.) Theorem 1 characterizes the complexity of the histogram testing problem within polylogarithmic factors. Note that there are three terms in the sample complexity; namely, $\sqrt{nk}/\varepsilon$, k/ε^2, and $\sqrt{n}/\varepsilon^2$. The sample complexity of the problem is dominated by one of these three different terms, depending on the relative sizes of n, k and $1/\varepsilon$. An illustration is given in Figure 1.

Prior to our work, the best previous histogram testing algorithm had sample complexity $\widetilde{O}(\sqrt{kn}/\varepsilon^3)$ [CDGR18], while the best known lower bound was $\widetilde{\Omega}(\sqrt{n}/\varepsilon^2 + k/\varepsilon)$ [Can16].[3]

We note that previous upper and lower bounds exhibit a polynomial gap, even for constant values of ε or k. For example, in the "large-k" regime where $k = n^c$ for some constant $0 < c < 1$, there was a gap between $\widetilde{O}(n^{1/2+c/2})$ and $\tilde{\Omega}(n^{1/2})$ in the sample complexity. In this regime, however, Theorem 1 results in the near-optimal bound of $\widetilde{\Theta}(n^{1/2+c/2})$. Similarly, in the "high-accuracy" regime where $\varepsilon = 1/n^c$ for some constant $c > 0$ (and, say, constant k), previous bounds only established that the sample complexity lies between $\widetilde{O}(n^{1/2+3c})$ and $\widetilde{\Omega}(n^{1/2+2c})$, while our result

---

[3]As discussed in Section 1.4, while an upper bound of $\widetilde{\Omega}(\sqrt{n}/\varepsilon^2 + k/\varepsilon^3)$ is claimed in [Can16], the analysis of their algorithm is flawed; and, indeed, our work shows that the sample complexity bound stated in [Can16] *cannot* hold, as it would contradict our lower bound.

Figure 1: The x-axis, y-axis are $\log(k)/\log(n)$ and $\log(1/\varepsilon)/\log(n)$ respectively. Each point in the graph corresponds to a setting of n, k, ε, and is colored based on the corresponding dominating term.

shows the (nearly-tight) bound is $\widetilde{\Theta}(n^{1/2+c})$. These are only two specific examples: more generally, the previously known bounds are suboptimal by polynomial factors in $1/\varepsilon$ when $\varepsilon \geq \sqrt{k/n}$; and by polynomial factors in all parameters $k, n, 1/\varepsilon$ when $\varepsilon \leq \sqrt{k/n}$. Theorem 1 settles the sample complexity of the problem, up to logarithmic factors, for *every* parameter setting.

At a technical level, our sample complexity lower bound construction conceptually differs from previous work in distribution testing, drawing instead from sophisticated techniques from the distribution *estimation* literature. Our upper bound follows from the "Testing-via-Learning" framework proposed in [ADK15]. The main technical innovation is a sample- and time- efficient *adaptive* algorithm which can simultaneously learn an unknown histogram *with unknown interval structure* distribution and identify a domain where the learned result is accurate. We elaborate on these aspects next.

1.3 Overview of Techniques

Sample Complexity Lower Bound. We follow the typical high-level approach in proving sample complexity lower bound. Namely, we define two ensembles of distributions D_{YES} and D_{NO} such that, with high probability, the following conditions are satisfied: (1) a random distribution from D_{YES} is a k-histogram, (2) a random distribution from D_{NO} is ε-far from any k-histogram, and (3) given a sample of appropriate size, it is information-theoretically impossible to distinguish a random distribution drawn from D_{YES} from a random distribution drawn from D_{NO}.

We start by describing our hard instances for the case that the accuracy parameter ε is a small universal constant. On the one hand we define D_{YES} so that all $\mathbf{p}_i$'s are the same except for a "small" number of domain elements i.e., $c \cdot k$ for a small constant $c \in (0, 1)$. On the other hand, for a distribution $\mathbf{p}$ drawn from D_{NO}, $\mathbf{p}_i$ will be randomly 0 or roughly $2/n$, except for at most a constant fraction of the elements. It is not hard to see that, with high probability, a distribution drawn from D_{YES} (resp. D_{NO}) will be a k-histogram (resp. far from being a k-histogram).

To ensure that the underlying distributions are indistinguishable using a small sample size, we want to guarantee that, for all small values of t, the number of elements with exactly t samples will be roughly the same for D_{YES} and D_{NO}, as this rules out any test statistic relying on counting the number of t-way collisions among the sample. Following [Val11, VV13, JVYHW15, WY16] this is essentially equivalent to showing that distributions drawn from D_{YES} and D_{NO} *match their low-degree moments*.

In particular, for a random pair of distributions $\mathbf{p}, \mathbf{p}'$ drawn from D_{YES} and D_{NO} respectively, we want that $\sum_i \mathbf{p}_i^t$ and $\sum_i \mathbf{p}'^t_i$ are roughly the same for all small values t. We note that the non-exceptional elements of a distribution $\mathbf{p}'$ drawn from D_{NO} — which have probability mass either 0 or roughly $2/n$ — will have second moment larger than the non-exceptional elements of a distribution $\mathbf{p}$ drawn from D_{YES} — which have probability mass roughly $1/n$ — by approximately $1/n$. To counteract this discrepancy, the (fewer than k) exceptional elements in D_{YES} must have average mass at least $1/\sqrt{kn}$. Fortunately, using techniques from [VV13, WY$^+$19], we are able to construct distributions that match $t = \widetilde{\Theta}(\log n)$ moments in which no individual bin has mass more than $\widetilde{O}(1/\sqrt{kn})$. Combining this construction with basic information-theoretic arguments gives us an $\widetilde{\Omega}(\sqrt{kn})$ sample complexity lower bound. We note that this lower bound is tight in the sense that with more than $\widetilde{\Omega}(\sqrt{kn})$ samples one can reliably identify the exceptional elements, as they will each have relatively large numbers of samples with high probability, allowing us to distinguish D_{YES} from D_{NO} simply based on the sub-distributions over these elements.

Given the aforementioned construction (for constant ε), it is easy to obtain a sample lower bound of $\widetilde{\Omega}(\sqrt{kn}/\varepsilon)$ by mixing our hard instances with the uniform distribution (with mixing weights ε and $1 - \varepsilon$ respectively). In fact, even if the testing algorithm knows in advance which samples come from the uniform part and which samples come from the original hard instance, distinguishing would require $\widetilde{\Omega}(\sqrt{kn})$ samples from the original hard instance, and therefore $\widetilde{\Omega}(\sqrt{kn}/\varepsilon)$ samples overall. This sample size lower bound turns out to be tight for ε relatively large, as one can still reliably identify the exceptional bins with only $\widetilde{\Omega}(\sqrt{kn}/\varepsilon)$ samples. However, when ε becomes sufficiently small, identifying the exceptional bins becomes more challenging. Indeed, if we take m samples, we expect that an exceptional bin has roughly $m\varepsilon/\sqrt{kn}$ more samples than a non-exceptional bin. On the other hand, a non-exceptional bin will have roughly $\mathrm{Poi}(m/n)$ samples with standard deviation $\sqrt{m/n}$. When $m/n \gg m\varepsilon/\sqrt{kn}$ (which happens in the regime $\varepsilon \ll \sqrt{k/n}$), in order for the exceptional bins to be distinguishable, we would need that $m\varepsilon/\sqrt{kn} \gg \sqrt{m/n}$ or $m \gg k/\varepsilon^2$ many samples. Using a careful information-theoretic argument, we formalize this intuition to show that $\widetilde{\Omega}(k/\varepsilon^2)$ is indeed a sample lower bound in this regime.

Sample-Efficient Tester. The starting point of our efficient tester is the *Testing-via-Learning* approach of [ADK15]. Very roughly speaking, we first design a learning procedure which outputs a distribution $\hat{\mathbf{p}}$ that would be close to $\mathbf{p}$ in χ^2 divergence, assuming that $\mathbf{p}$ was in fact a k-histogram. Then we use a χ^2/ℓ_1 tolerant tester, in the spirit of the one introduced in [ADK15], to distinguish between the cases that $\mathbf{p}$ is close to $\hat{\mathbf{p}}$ in χ^2 divergence versus far from $\hat{\mathbf{p}}$ in ℓ_1-distance. This step is however significantly harder than this simple outline suggests, as it turns out challenging to perform the first step exactly. Instead, we design a specific learning algorithm with an implicit "hybrid" learning guarantee, (see Lemma 5) which in turns requires us to considerably generalize and adapt the "tolerant testing part" to avoid spurious discrepancies (introduced in the imperfect learning stage) which may lead to false negatives.

To implement the first step, we follow the general "learn-and-sieve" idea suggested in [Can16], with important modifications to address the flaw in their approach and its analysis. In particular, suppose that $\mathbf{p}$ is a k-histogram. Then, if we *knew* the corresponding k intervals (that make up the partition for the k-histogram), it suffices to learn the mass of $\mathbf{p}$ on each interval, and let $\hat{\mathbf{p}}$ be uniform on each interval (with the appropriate total mass). Of course, a key source of difficulty arises from the fact that we do not know the partition a priori. To circumvent this issue, we divide $[n]$ into (roughly) $K = \Theta(k)$ intervals and try to detect if $\mathbf{p}$ is far in χ^2 divergence from being uniform on any of these intervals. If an interval from our partition incurs large χ^2 error (we call such an interval *bad*), we know that $\mathbf{p}$ must not be constant within this interval. Therefore, we proceed to subdivide these bad intervals into roughly equal parts, and recurse on the $\Theta(k)$ intervals in our new partition. Assuming $\mathbf{p}$ is a k-histogram, we subdivide at most k intervals in each iteration, since there could be at most k intervals from any interval partition of $[n]$ where $\mathbf{p}$ is not constant. Hence, in each iteration, we decrease the mass of the bad intervals by at least a constant factor. We repeat the process for at most $O(\log(1/\varepsilon))$ many iterations; after this many iterations, the total mass of the bad intervals will become $O(\varepsilon)$, and thus they may be safely ignored.

A significant difference between our method and the approach from [Can16] lies in the method of *sieving*. In [Can16], it was only said that the algorithm would filter out a subset of *breakpoint* intervals based on the χ^2 statistics ([ADK15]) with the goal of reducing discrepancy; this is where

the main gap in their analysis lies, and the particular (flawed) approach they suggested does not seem to be fixable [Can22]. On the contrary, we characterize the exact set of intervals that need to (and can) be removed with the formal definition of *bad* intervals with respect to a given partition $\mathcal{I}$ of $[n]$ (See Definition 2). Based on that, our approach is to search for *any* sub-intervals J (not necessarily an interval in $\mathcal{I}$) on which the χ^2 divergence between $\mathbf{p}$ and $\hat{\mathbf{p}}$, an approximation of $\mathbf{p}$ assuming $\mathbf{p}$ is uniform over intervals within the given partition, is more than $\widetilde{\Omega}(\varepsilon^2/k)$. For an interval I from the partition $\mathcal{I}$, we show the inclusion of such "bad sub-interval" $J \subseteq I$ then certifies the "badness" of I itself. To find such a J, we need a technique for accurately approximating $\mathbf{p}(J)$ simultaneously for all intervals $J \subseteq [n]$, in both absolute and relative error; a notion of approximation much stronger that what classical tools from statistical learning theory such as the VC inequality or the Dvoretzky–Kiefer–Wolfowitz (DKW) inequality provide. Notice that, for a *fixed* interval $J \subseteq [n]$, taking the empirical distribution over b samples gives an estimate $\mathbf{q}$ of $\mathbf{p}$ such that $|\mathbf{q}(J) - \mathbf{p}(J)| < \sqrt{\mathbf{p}(J)/b}$ with constant probability. By taking $\Theta(\log(n))$ batches of samples (each containing b i.i.d. samples from $\mathbf{p}$), and computing the median value of all of the $\mathbf{q}(J)$'s, with high probability for each J, we then obtain an estimate $\hat{\varphi}(J)$ [4] for which the above condition holds. Using the sub-routine, as long as b is at least $\Omega(k/\varepsilon^2)$, we can ensure that $|\hat{\varphi}(J) - \mathbf{p}(J)|^2/\mathbf{p}(J) \ll \varepsilon^2/k$, and we can then safely use our estimate $\hat{\varphi}(J)$ as a proxy for $\mathbf{p}(J)$ for the detection of those "bad sub-intervals" for which $|\mathbf{p}(J) - \hat{\mathbf{p}}(J)|^2/\hat{\mathbf{p}}(J)$ is large, which in turns certify the bad intervals from a given partition. This suffices *unless* $\mathbf{p}(J)$ is substantially larger than our estimate $\hat{\mathbf{p}}(J)$.

Unfortunately, this bad case where $\mathbf{p}(J) \gg \hat{\mathbf{p}}(J)$ can happen if $\mathbf{p}(J)$ is smaller than $1/b$. In such a case, in a collection of b samples from $\mathbf{p}$, we are likely to see no samples in J, and thus our empirical estimate $\hat{\mathbf{p}}(J)$ will be 0. We can fix this issue (i.e., the case where $\hat{\mathbf{p}}(J)$ is actually 0) by mixing both $\mathbf{p}$ and $\hat{\mathbf{p}}$ with the uniform distribution, thus allowing us to assume that $\hat{\mathbf{p}}(J) \geq |J|/2n \geq 1/(2n)$. Yet, this still leaves a potential gap of roughly n/b between the ratio of $\mathbf{p}(J)$ and $\hat{\mathbf{p}}(J)$. Fortunately, if we select $b \gg \sqrt{nk}/\varepsilon$, we will have that $|\hat{\varphi}(J) - \mathbf{p}(J)|^2/\mathbf{p}(J) \ll \varepsilon/\sqrt{nk}$, and even accounting for losing a factor of b/n, we will still have that $|\hat{\varphi}(J) - \mathbf{p}(J)|^2/\mathbf{p}(J) \ll \varepsilon^2/k$. This implies that we will successfully detect any bad intervals and achieve our learning guarantees.

1.4 Prior Work

Motivated by the question of building provably good succinct representations of a dataset from only a small sub-sample of the data, [ILR12] first introduced histogram testing as a preliminary, ultra-efficient decision subroutine to find the best parameter k for the number of bins. They provided an algorithm for this task which required $\widetilde{O}(\sqrt{kn}/\varepsilon^5)$ samples from the dataset, a sample complexity which beats the naïve approach (reading and processing the whole dataset) for small values of k and relatively large values of the accuracy parameter ε. Subsequent work [CDGR18] reduced the dependence on ε from quintic to cubic, giving an algorithm with sample complexity $\widetilde{O}(\sqrt{kn}/\varepsilon^3)$. This bound was, however, still quite far from the "trivial" lower bound of $\Omega(\sqrt{n}/\varepsilon^2)$, which follows from a reduction to uniformity testing (i.e., the case $k = 1$) [Pan08].

Prior to the current work, an $\widetilde{O}(\sqrt{n}/\varepsilon^2 + k/\varepsilon^3)$ upper bound and an $\widetilde{\Omega}(\sqrt{n}/\varepsilon^2 + k/\varepsilon)$ lower bound were obtained in [Can16]. While the lower bound is theoretically sound (albeit, as it turns out, suboptimal), as pointed out in [Can22], the upper bound does not hold due to a technical flaw in the analysis, leaving the optimal sample complexity of the problem open for even constant ε. Moreover, the lower bound of [Can16], based on a reduction of histogram testing to the well-studied problem of support size estimation, provably cannot be improved to provide either (i) a quadratic dependence on ε, i.e., $\widetilde{\Omega}(k/\varepsilon^2)$ or (ii) coupling between the two domain parameters k, n, i.e., $\widetilde{\Omega}(\sqrt{nk}/\varepsilon)$. Our work remedies all those issues, fully resolving the question of histogram testing, for the whole parameter range, with logarithmic factors.

Finally, we note that a number of works have obtained algorithms and lower bounds for related, yet significantly different, testing problems. Specifically, [DK16] gave a sample-optimal testing algorithm for the special case of our problem where the k intervals are known *a priori*. Moreover, a number of works [DKN15b, DKN15a, DKN17] have obtained identity and equivalence testers *under the assumption* that the input distributions are k-histograms.

---

[4]Notice that $\hat{\varphi}$ is neither a distribution nor a measure, but just a map from intervals to positive real values.

Preliminaries. We denote by $\mathrm{TV}(\mathbf{p}, \mathbf{q})$ the total variation (TV) distance between probability distributions $\mathbf{p}, \mathbf{q}$ over $[n] := \{1, 2, \ldots, n\}$, defined as $\mathrm{TV}(\mathbf{p}, \mathbf{q}) := \sup_{S \subseteq [n]}(\mathbf{p}(S) - \mathbf{q}(S)) = \frac{1}{2} \sum_{i=1}^{n} |\mathbf{p}(i) - \mathbf{q}(i)|$, where $\mathbf{p}(S) := \sum_{i \in S} \mathbf{p}(i)$. We will make essential use of the χ^2-divergence of $\mathbf{p}$ with respect to $\mathbf{q}$, defined as $d_{\chi^2}(\mathbf{p}\|\mathbf{q}) := \sum_{i=1}^{n} (\mathbf{p}_i - \mathbf{q}_i)^2 / \mathbf{q}_i$. We will also require generalizations of these definitions on restrictions of the domain. In particular, given $S \subseteq [n]$, we use the notation $\mathrm{TV}^S(\mathbf{p}, \mathbf{q}) := (1/2) \sum_{i \in S} |\mathbf{p}(i) - \mathbf{q}(i)|$ and $d_{\chi^2}^S(\mathbf{p}\|\mathbf{q}) := \sum_{i \in S} (\mathbf{p}_i - \mathbf{q}_i)^2 / \mathbf{q}_i$. We note that for any $S \subseteq [n]$, it holds that $\mathrm{TV}^S(\mathbf{p}, \mathbf{q})^2 \leq \frac{1}{4} d_{\chi^2}^S(\mathbf{p}\|\mathbf{q})$.

The asymptotic notation $\tilde{O}$ (resp. $\tilde{\Omega}$) suppresses logarithmic factors in its argument, i.e., $\tilde{O}(f(n)) = O(f(n) \log^c f(n))$ and $\tilde{\Omega}(f(n)) = \Omega(f(n) / \log^c f(n))$, where $c > 0$ is a universal constant. The notations $\ll$ and $\gg$ intuitively mean "much less than" and "much greater than" respectively. Formally, we write $f(n) \ll g(n)$ to denote that $f(n) < c \cdot g(n)$, for some universal constant $0 < c$.

2 An efficient testing algorithm

A preliminary simplification. Without loss of generality, we will assume that $\mathbf{p}(i) \geq \frac{1}{2n}$ for every $i \in [n]$. Indeed, this can be ensured by mixing the unknown distribution with the uniform distribution $\mathbf{u}_n$ on $[n]$ beforehand i.e. $\mathbf{p}' := \frac{1}{2}(\mathbf{p} + \mathbf{u}_n)$ (See Fact 3 in Appendix for how to sample from $\mathbf{p}'$ efficiently). It is easy to see that $\mathbf{p}'$ remains a histogram after mixing: $\mathbf{p}' \in \mathcal{H}_k^n$ if $\mathbf{p} \in \mathcal{H}_k^n$, and $\mathbf{p}'$ is at least $(\varepsilon/2)$-far away from every histogram if $\mathbf{p}$ is ε-far from every histogram.

Testing via Learning. The main approach is to follow the **Testing-via-Learning** framework proposed in [ADK15]. In particular, suppose we have a learning algorithm capable of constructing $\hat{\mathbf{p}}$ that is close to $\mathbf{p}$ in χ^2 divergence when $\mathbf{p} \in \mathcal{H}_n^k$. Then, (1) if $\mathbf{p} \in \mathcal{H}_k^n$, we will have that $\mathbf{p}$ and $\hat{\mathbf{p}}$ are close *and* (as a consequence of this) that $\hat{\mathbf{p}}$ is close to being a k-histogram. Yet, (2) if $\mathbf{p}$ is far from being a k-histogram, then by the triangle inequality we must have either that $\hat{\mathbf{p}}$ is far from being a k-histogram, or that $\mathbf{p}$ and $\hat{\mathbf{p}}$ are far from each other in ℓ_1 distance. We can use dynamic programming to check the explicit description is indeed close to a k-histogram in ℓ_1 distance efficiently (See Lemma 4.11 of [CDGR18]). To verify $\mathbf{p}$ and $\hat{\mathbf{p}}$ are close, we will use a result of [ADK15] on tolerant identity testing. In particular, given an explicit description $\hat{\mathbf{p}}$, the tester takes sample from the unknown distribution $\mathbf{p}$ and decides whether $\mathbf{p}$ and $\hat{\mathbf{p}}$ are closed in χ^2 divergence or far in ℓ_1 distance. We remark that $\hat{\mathbf{p}}$ can be relaxed to be a positive measure.

Lemma 1 (Adapted from Lemmas 2 and 3 [ADK15]). *Let $\mathbf{p}$ and $\hat{\mathbf{p}}$ be a distribution and a positive measure defined on $[n]$ respectively. Fix $\varepsilon \in (0, 1)$ and let $\mathcal{A} = \{i \in [n] : \hat{\mathbf{p}}(i) \geq \varepsilon/(50n)\}$. There exists a tester **Tolerance-Identity-Test**, which takes $\mathrm{Poi}(m)$ for $m = \Theta\left(\sqrt{n}/\varepsilon^2\right)$ i.i.d. samples from $\mathbf{p}$ and outputs **Accept** if $d_{\chi^2}^{\mathcal{A}}(\mathbf{p}\|\hat{\mathbf{p}}) \leq \varepsilon^2/500$ and **Reject** if $\mathrm{TV}^{\mathcal{A}}(\mathbf{p}, \hat{\mathbf{p}}) \geq \varepsilon$ with constant probability.*

Outline for Learning. If $\mathbf{p} \in \mathcal{H}_k^n$ and we know the partition of $\mathbf{p}$ in advance, one can learn $\mathbf{p}$ up to ε^2 in χ^2 divergence with $\Theta(k/\varepsilon^2)$ samples (following the analysis of Laplace estimator from [KOPS15]). Without the partition information, we can nonetheless achieve a weaker guarantee. That is, we can output a fully specified measure $\hat{\mathbf{p}}$ on $[n]$, together with a sub-domain $\mathcal{G} \subseteq [n]$, such that $d_{\chi^2}^{\mathcal{G}}(\mathbf{p}\|\hat{\mathbf{p}})$ is small. In particular, we can achieve the guarantee in three steps. (i) Divide the domain $[n]$ into $K \gg k$ many intervals obliviously (Lemma 2). (ii) Output a succinct measure $\hat{\mathbf{p}}$ that is constant on each interval specified by Step (i) (Section 2.1). (iii) Identify the intervals I where $d_{\chi^2}^{I}(\mathbf{p}\|\hat{\mathbf{p}})$ is large (Section 2.2). Denote $\mathcal{B} = [n]\backslash\mathcal{G}$. The fact that we only have $\mathbf{p}$ and $\hat{\mathbf{p}}$ close in χ^2 divergence on a sub-domain $\mathcal{G}$ is a reasonable compromise as long as $\mathbf{p}(\mathcal{B}), \hat{\mathbf{p}}(\mathcal{B}) \ll \varepsilon$: if $\mathbf{p}$ is ε-far away from $\hat{\mathbf{p}}$ in ℓ_1 distance on $[n]$, $\mathbf{p}$ is at least $(\varepsilon - \mathbf{p}(\mathcal{B}) - \hat{\mathbf{p}}(\mathcal{B}))$-far away from $\hat{\mathbf{p}}$ on $[n]\backslash\mathcal{B}$. Otherwise, we may take more samples from $\mathbf{p}$ restricted to $\mathcal{B}$ and sub-divide the problematic intervals identified in Step (iii). Repeating the above steps iteratively then brings us to the case $\mathbf{p}(\mathcal{B}) \ll \varepsilon$.

Equitable Partition. The first step is to divide the domain into $\Theta(k)$ many intervals over which the masses of $\mathbf{p}$ are approximately equal. As shown in [ADK15], this can be done with $\widetilde{\Theta}(k)$ many samples through a routine we denote as **Approx-Divide**. We also need a routine for sub-dividing a set of disjoint intervals into even lighter sub-intervals. Nonetheless, one can reduce the sub-dividing task to domain partitioning by running **Approx-Divide** on the sub-distribution restricted to the set of disjoint intervals. Proofs are provided in Appendix A.1.

Lemma 2. *There exists an algorithm* **Approx-Sub-Divide** *that, given parameters $\delta \in (0, 1]$ and integer $B > 1$, as well as a set of disjoint intervals $\mathcal{I} = \{I_1, I_2, \cdots, I_q\}$, given sample access to $\mathbf{p}$ on $[n]$, outputs a list of partitions $\mathcal{S}_1, \ldots, \mathcal{S}_q$, where $\mathcal{S}_i$ is the partition of the interval $I_i \in \mathcal{I}$, such that the following holds with probability at least $1 - \delta$. (i) The algorithm uses $O\left(B/\mathbf{p}(\mathcal{I}) \cdot \log\left(B/\delta\right)\right)$ samples. (ii) The output contains at most $(8B + q)$ intervals in total. (iii) Every non-singleton interval $S \in \bigcup_{j=1}^{q} \mathcal{S}_j$ satisfies $\mathbf{p}(S) \leq \mathbf{p}(\mathcal{I}) \cdot 16/B$.*

2.1 Simultaneously Estimating Mass of Intervals

In this section, we first introduce **Interval-Mass-Estimate**, a sub-routine that can accurately approximate the mass of $\mathbf{p}(J)$ for all intervals $J \subseteq [n]$ simultaneously and then show how we can use it to learn $\mathbf{p}$ (assuming $\mathbf{p} \in \mathcal{H}_n^k$).

Interval-Mass-Estimate first divides the number of samples drawn into $\Theta(\log(n/\delta))$ batches. For an interval I, we compute the estimate (number of samples falling in I divided by the batch size) for each batch separately and compute the median over the statistics. This is often referred as the "Median Trick" and is crucial in achieving the learning guarantees with high probability. Pseudo-code and analysis are provided in Appendix A.2.

Lemma 3. *Let be $\mathbf{p}$ be supported on $[n]$ such that $\mathbf{p}(i) \geq 1/(2n)$. Fix $b \in \mathbb{Z}^+$ and $\delta \in (0, 1]$. The algorithm* **Interval-Mass-Estimate** *takes $3b\log(n/\delta)$ i.i.d. samples from $\mathbf{p}$ and outputs $\hat{\varphi}$, a map from sub-intervals of $[n]$ to real values, such that, with probability at least $1 - \delta$, for every sub-interval $I \subseteq [n]$ it holds that $\mathbf{p}(I)/\hat{\varphi}(I) \leq \max(2, 8n/b)$, $\hat{\varphi}(I)/\mathbf{p}(I) \leq 3$ and $|\hat{\varphi}(I) - \mathbf{p}(I)| \leq \sqrt{\mathbf{p}(I)/b}$.*

Let $\mathcal{I}$ be a partition of $[n]$. We try to learn $\mathbf{p}$ pretending that $\mathbf{p}$ is constant over each interval within $\mathcal{I}$ with the routine **Empirical-Learning**. In particular, the algorithm uses **Interval-Mass-Estimate** to obtain estimations of the mass of $I \in \mathcal{I}$ and then flatten the mass uniformly among elements $i \in I$. Notice that, due to the application of the median trick, the output is not necessarily a distribution but rather a positive measure[5] $\hat{\mathbf{p}}$ on $[n]$ which is constant over each interval within $\mathcal{I}$.

If $\mathbf{p}$ is indeed a k-histogram, errors are only incurred on a special type of intervals (of which there are at most k) which we refer to as the *breakpoint intervals*.

Definition 1 (Breakpoint Intervals). *Given a k-histogram $\mathbf{p}$ on $[n]$, we say that $i \in [n]$ is a* breakpoint *with respect to $\mathbf{p}$ if $\mathbf{p}(i) \neq \mathbf{p}(i + 1)$; and that an interval $I \subseteq [n]$ is a* breakpoint interval *(with respect to $\mathbf{p}$) if I contains at least one breakpoint.*

With Definition 1 in mind, we now specify the formal learning guarantees. Pseudo-code and proofs are provided in Appendix A.3.

Lemma 4. *Suppose $\mathbf{p} \in \mathcal{H}_k^n$. Let $\mathcal{I}$ a partition of $[n]$ into K intervals. Let $b \in \mathbb{Z}^+$, $\delta \in (0, 1]$ and $T := 3\log(K/\delta)$. There exists an algorithm* **Empirical-Learning**, *given (Tb) i.i.d. samples from $\mathbf{p}$, outputs a positive measure $\hat{\mathbf{p}}$, which satisfies the following with probability at least $1 - \delta$. (i) $\hat{\mathbf{p}}$ is constant within each interval in $\mathcal{I}$. (ii) For every sub-intervals $J \subseteq I$ where $I \in \mathcal{I}$ is a* non-breakpoint *interval with respect to $\mathbf{p}$, we have $\mathbf{p}(J)/\hat{\mathbf{p}}(J) \leq \max(2, 8 \cdot n/b)$ and $|\hat{\mathbf{p}}(J) - \mathbf{p}(J)| \leq \sqrt{\mathbf{p}(J)/b}$.*

By combining the two guarantees in Lemma 4, one can see the χ^2 divergence between $\mathbf{p}$ and $\hat{\mathbf{p}}$, restricted to the non-breakpoint intervals, will be at most ε^2 with high probability if taking $\Theta(KT/\varepsilon^2)$ many samples. However, following a result from [KOPS15, Can16], one only need $\Theta(K/\varepsilon^2)$ samples to learn a K-histogram up to ε^2 error in this restricted notion of χ^2 divergence. One may wonder whether this is enough for us, and if the stronger (but less natural) guarantees provided by Lemma 4, which end up increasing the number of samples required, are necessary. As we will see in the next section, we indeed need not only that the χ^2 divergence is small, but also that the ratio $\mathbf{p}(I)/\hat{\mathbf{p}}(I)$ is bounded for all non-breakpoint intervals. In particular, this latter property enables us to compute relatively accurate estimates of the χ^2 divergence restricted to sub-intervals and (consequently) to tell whether $\mathbf{p}$ is constant or from far from being constant on an interval.

2.2 Bad Interval Detection

While large contributions to the χ^2 divergence (assuming the learning phase was successful) will only come from breakpoint intervals, not all of them will necessarily contribute significantly to the

---

[5]That is, $\hat{\mathbf{p}}$ might not sum to one, and thus is not itself a probability distribution.

χ^2 divergence. In particular, a breakpoint interval is only considered "bad" and needs to be filtered out if the error incurred is proportional to the number of breakpoints within.

Definition 2 (ε-Bad-Interval). *Fix a partition $\mathcal{I}$ of $[n]$ containing K intervals. Let $I \in \mathcal{I}$ be a breakpoint interval of $\mathbf{p}$. Furthermore, suppose I contains $j-1$ breakpoints i.e. $\mathbf{p}$ is j-piece-wise uniform in I. We say $I \in \mathcal{I}$ is an ε-bad interval with respect to $\hat{\mathbf{p}}$ and $\mathcal{I}$ if $d_{\chi^2}^I \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \geq j \cdot \varepsilon^2/K$.*

The definition suits our purpose for two reasons. (i) The total χ^2 error between $\mathbf{p}$ and $\hat{\mathbf{p}}$ on the set of "good" intervals (complement of the set of "bad" intervals) is small. Indeed, let $\mathcal{G} \in \mathcal{I}$ be a set containing no ε-bad intervals. Since there are at most K intervals contained in $\mathcal{G}$ and k breakpoints contained in the intervals in $\mathcal{G}$, it is easy to see that $d_{\chi^2}^{\mathcal{G}} \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \leq O(\varepsilon^2)$. (ii) One can reliably separate bad intervals from non-breakpoint intervals assuming the learning phase was successful. To see why, note that in that case every non-breakpoint interval I satisfies $d_{\chi^2}^J \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \ll \varepsilon^2/K$ for all $J \subseteq I$ with high probability. On the contrary, for any bad interval I, we claim there must be a sub-interval $Q \subseteq I$ where $d_{\chi^2}^Q \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \geq \varepsilon^2/K$ and both $\mathbf{p}$ and $\hat{\mathbf{p}}$ are constant within. In particular, if I is an ε-bad interval that contains $(j-1)$ breakpoints, we then have a partition $\{Q_1, \cdots, Q_j\}$ of I over which $\mathbf{p}$ is piece-wise constant and at least one of them will have χ^2 error at least ε^2/K.

Our next step is to show how we can leverage the separating condition to design an efficient bad interval detection mechanism. This is where our method *significantly differs* from [Can16]. At a high level, we take another set of independent samples to get an estimate $\hat{\varphi}(Q)$ of $\mathbf{p}(Q)$ for all $Q \subseteq [n]$ simultaneously. Then, we compare $\hat{\varphi}(Q)$ with $\hat{\mathbf{p}}(Q)$ to see whether we have $d_{\chi^2}^Q \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \geq \varepsilon^2/K$, which would in turns imply the interval $I \supseteq Q$ from the given partition is ε-bad. We next provide the pseudo-code for **Learn-And-Sieve**, which finds a positive measure $\hat{\mathbf{p}}$ on $[n]$ and a domain $\mathcal{B}$ such that $d_{\chi^2}^{[n] \setminus \mathcal{B}} \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \leq O(\varepsilon^2)$ provided $\mathbf{p} \in \mathcal{H}_k^n$. Its detailed analysis can be found in Appendix A.4.

Algorithm 1 Learn-And-Sieve

Require: Sample access to $\mathbf{p}$; a partition $\mathcal{I}$ of $[n]$ containing K intervals; accuracy ε; failure probability δ.
1: Let $m = C \cdot (K/\varepsilon^2 + \sqrt{Kn}/\varepsilon) \cdot \log(n/\delta)$ for a sufficiently large constant C.
2: Draw $2m$ i.i.d. samples from $\mathbf{p}$ and split the samples evenly into $\mathcal{S}_1, \mathcal{S}_2$.
3: $\hat{\mathbf{p}} \leftarrow$ **Empirical-Learning**$(\mathcal{S}_1, \mathcal{I}, \delta/4)$, $\hat{\varphi} \leftarrow$ **Interval-Mass-Estimate**$(\mathcal{S}_2, \delta/4)$, $\mathcal{B} \leftarrow \{\}$.
4: **for** all intervals $Q \subseteq I$ for some $I \in \mathcal{I}$ **do**
5:      **if** $\hat{\varphi}(Q)/\hat{\mathbf{p}}(Q) > 6 \cdot \max(1, \varepsilon \cdot \sqrt{n/K})$ **or** $|\hat{\varphi}(Q) - \hat{\mathbf{p}}(Q)| > 0.5\sqrt{\hat{\mathbf{p}}(Q)\varepsilon^2/K}$ **then**
6:          Add I to $\mathcal{B}$.
7: Output Reject if $\mathcal{B}$ contains more than k intervals. Otherwise, **return** $\mathcal{B}, \hat{\mathbf{p}}$.

Lemma 5 (Sieving Lemma). *Given a partition $\mathcal{I}$ containing K intervals, sample access to $\mathbf{p}$ on $[n]$ and $\delta \in (0, 1)$. Then, the output of **Learn-And-Sieve** (Algorithm 1) satisfies the following. (i) Suppose $\mathbf{p} \in \mathcal{H}_k^n$. Then the algorithm returns a positive measure $\hat{\mathbf{p}}$ and $\mathcal{B}$ such that $d_{\chi^2}^{[n] \setminus \mathcal{B}} \left(\mathbf{p} \| \hat{\mathbf{p}} \right) \leq \varepsilon^2$ with probability at least $1 - \delta$. (ii) The output $\mathcal{B}$ contains at most k intervals (if the algorithm does not reject). (iii) At most $O((K/\varepsilon^2 + \sqrt{Kn}/\varepsilon) \cdot \log(n/\delta))$ samples are used.*

Proof Sketch. We claim that, if $\mathbf{p} \in \mathcal{H}_k^n$, $\mathcal{B}$ contains all the ε-bad intervals and no non-breakpoint intervals with high probability. Let I be a non-breakpoint interval. For $b = \Theta(m/\log(n/\delta)) = \Theta(K/\varepsilon^2 + \sqrt{Kn}/\varepsilon)$, we have, with high probability, $|\hat{\varphi}(Q) - \mathbf{p}(Q)| \leq \sqrt{\mathbf{p}(Q)/b}$, $|\hat{\mathbf{p}}(Q) - \mathbf{p}(Q)| \leq \sqrt{\mathbf{p}(Q)/b}$ and $\mathbf{p}(Q)/\hat{\mathbf{p}}(Q) \leq \max(2, 8 \cdot n/b)$ which follow from Lemmas 3 and 4. Combining this with triangle inequality and our choice of b implies the second condition of Line 5 will be false. The first condition can be shown to be false by rewriting $\hat{\varphi}(Q)/\hat{\mathbf{p}}(Q)$ as $\hat{\varphi}(Q)/\mathbf{p}(Q) \cdot \mathbf{p}(Q)/\hat{\mathbf{p}}(Q)$, which are themselves bounded, with high probability, by 3 and $\Theta(1) \cdot \max(1, \varepsilon\sqrt{n/K})$ again by Lemmas 3 and 4 and our choice of b.

Let I be a breakpoint interval. We then have $|\mathbf{p}(Q) - \hat{\mathbf{p}}(Q)| \geq \sqrt{\hat{\mathbf{p}}(Q) \cdot \varepsilon^2/K}$ for some sub-interval $Q \subset I$. If $\mathbf{p}(Q)$ is light ($\mathbf{p}(Q) \leq 2\varepsilon/\sqrt{Kn}$), we can show $\mathbf{p}(Q)/b \leq 1/4 \cdot \hat{\mathbf{p}}(Q) \cdot \varepsilon^2/K$, making $\hat{\varphi}(Q)$, our estimation for $\mathbf{p}(Q)$, sufficiently accurate such that the second condition of Line 5 will be true. Otherwise, as $b \gg \sqrt{Kn}/\varepsilon$, the estimation $\hat{\varphi}(Q)$ will be within multiplicative factors of $\mathbf{p}(Q)$. If $\hat{\mathbf{p}}(Q)$ is not much lighter than $\mathbf{p}(Q)$, we can again show $\mathbf{p}(Q)/b \leq 1/4 \cdot \hat{\mathbf{p}}(Q) \cdot \varepsilon^2/K$.

Otherwise, the first condition of Line 5 will be true. Conditioned on that $\mathcal{B}$ includes all ε-bad intervals and no non-breakpoint intervals, it is easy to see that $\mathcal{B}$ will contain no more than k intervals and $d_{\chi^2}^{\mathcal{I}\setminus\mathcal{B}}\left(\mathbf{p}\big\|\hat{\mathbf{p}}\right) \leq O(\varepsilon^2)$. We note that points (i) and (iii) follow from the definition of the algorithm. □

Learn-and-Sieve (Algorithm 1) outputs a fully specified description $\hat{\mathbf{p}}$ and a sub-domain $\mathcal{G} := [n]\setminus\mathcal{B}$ such that $d_{\chi^2}^{\mathcal{G}}\left(\mathbf{p}\big\|\hat{\mathbf{p}}\right)$ is small given $\mathbf{p} \in \mathcal{H}_n^k$. For testing purposes, this is a reasonable divergence from the ideal guarantee that $d_{\chi^2}\left(\mathbf{p}\big\|\hat{\mathbf{p}}\right)$ is small *as long as* $\mathbf{p}(\mathcal{B})$ *is also small*. If so, we can set $\hat{\mathbf{p}}(i) = 0$ for $i \in \mathcal{B}$ and invoke **Tolerant-Identity-Test** with $\mathbf{p}$ and $\hat{\mathbf{p}}$. If the test passes, we then know that $\mathrm{TV}^{\mathcal{G}}\left(\mathbf{p}, \hat{\mathbf{p}}\right) \leq \varepsilon/2$: this together with $\mathbf{p}(\mathcal{B}) \leq \varepsilon/2$ then gives $\mathrm{TV}\left(\mathbf{p}, \hat{\mathbf{p}}\right) \leq \varepsilon$.

Unfortunately, running **Learn-and-Sieve** only once we may have $\mathbf{p}(\mathcal{B}) = \Omega(1)$. To handle this, we will need more fine-grained sieving procedure, which uses **Approx-Sub-Divide** to further partition the bad intervals detected and invokes **Learn-and-Sieve** *iteratively*. In each iteration, the total mass of the bad intervals shrinks by a constant factor, allowing us to reach $\mathbf{p}(\mathcal{B}) \ll \varepsilon$ in at most $O(\log(1/\varepsilon))$ iterations. The pseudo-code (Algorithm 4) and detailed analysis are provided in Appendix A.5.

3 Histogram Lower Bound

In this section, we describe the hard instance of histogram testing, which leads to an $\widetilde{\Omega}(\sqrt{kn}/\varepsilon + k/\varepsilon^2)$ lower bound. We will apply the so-called Poissonization trick: we will relax P, the unknown object being tested, to be a positive measure with total mass $\Theta(1)$. We denote such a measure as an approximate probability vector and give the corresponding notion of histogram.

Definition 3 (Approximate Probability Vector). *We define the set of ν-approximate probability vectors (APV) on the domain $[n]$ by $\tilde{\mathcal{P}}^n(\nu) := \{P : P_i \in [0,\infty) \,\forall i \in [n], |\|P\|_1 - 1| \leq \nu\}$. Accordingly, the set of* histogram APV *is given by $\tilde{\mathcal{H}}_k^n(\nu) := \{P \in \tilde{\mathcal{P}}^n(\nu) : P/\|P\|_1 \in \mathcal{H}_k^n\}$.*

Under the Poisson sampling model, given an unknown $P \in \tilde{\mathcal{P}}^n(\nu)$, the goal it to decide whether $P \in \tilde{\mathcal{H}}_k^n(\nu)$ or P is at least $\varepsilon(1+\nu)$-far[6] from any $P' \in \tilde{\mathcal{H}}_k^n(\nu)$ in ℓ_1 distance when given the vector $\{M_1, M_2, \cdots M_n\}$ where $M_i \sim \mathrm{Poi}(m \cdot P_i)$. We denote the sample complexity of the problem by $m_{\mathrm{hist}}^{\mathrm{poi}}(n, k, \varepsilon, \nu)$ and provide its formal definition in Appendix B.

To lower bound $m_{\mathrm{hist}}^{\mathrm{poi}}(n, k, \varepsilon, \nu)$, we follow the idea of *moment matching* illustrated in [Val11, VV13, WY16]. In particular, one first constructs two discrete non-negative random variables U, U' whose first few moments are identical. Moreover, U and U' will be designed to have different properties such that one can use i.i.d. copies of U (and U') to generate random measures that are histograms (and far-away from histograms respectively).

Our construction of such a pair of random variables is based on *Chebyshev's polynomials*, a standard tool in approximation theory and the parameter estimation literature. The two variables will be supported on the roots of the polynomial $p(x) = x\left(x - \frac{1}{n}\right)\left(x - \frac{2}{n}\right)T_d\left(1 - \frac{\sqrt{kn}}{C\cdot\log^2 n}\cdot x\right)$, where $T_d(\cdot)$ is the *Chebyshev's polynomial* (of the first kind) and C is a sufficiently large constant. More precisely, U will be supported on roots r where the derivatives $p'(r) < 0$, U' will be on roots where $p'(r) > 0$, and the probabilities will be proportional to $p'(r)$ accordingly. Consequently, U will most likely be $1/n$ (hence useful for histogram construction) and U' will most likely be 0 or $2/n$, each with non-trivial probabilities (hence appropriate for non-histogram construction). Besides, they will have their maximums bounded by $\widetilde{O}(1/\sqrt{kn})$, which is crucial to achieve the nearly optimal lower bounds. The detailed construction and analysis are provided in Appendix B.1.

Lemma 6. *Given positive integers k, n where $k < n$, there exists a pair of non-negative random variable U, U' supported on $[0, 1)$ and absolute constants $c, c' > 0$ satisfying (i) $\Pr\left[U \neq \frac{1}{n}\right] \ll \frac{k}{n}$. (ii) $\Pr\left[U' = 0\right] > 1/3$ and $\Pr\left[U' = \frac{2}{n}\right] > 1/3$. (iii) $U, U' \leq c'\log^2 n/\sqrt{kn}$. (iv) $\mathbf{E}[U] = \mathbf{E}[U'] = \frac{1}{n}(1 + O(\sqrt{k/n}))$. (v) $\mathbf{E}[U^t] = \mathbf{E}[U'^t]$ for $1 \leq t \leq c \cdot \log n$.*

We the proceed to construct two families of Approximate Probability Vectors, one of which belongs to $\tilde{\mathcal{H}}_k^n$ and the other far from it using the random variables stated in Lemma 6. To do so, we

---

[6]The extra $(1+\nu)$ factor is to accommodate the fact that P may not be a distribution i.e. $1 \leq \|P\|_1 < (1+\nu)$.

define $H = \left(1/n + \varepsilon U^{(1)}, \cdots, 1/n + \varepsilon U^{(n)}\right)$, $H' = \left(1/n + \varepsilon U'^{(1)}, \cdots, 1/n + \varepsilon U'^{(n)}\right)$ where $U^{(i)}, U'^{(i)}$ are n i.i.d. copies of U, U' in Lemma 6.

We address the two regimes $\sqrt{k/n} \leq \varepsilon \log^2 n$ and $\sqrt{k/n} \geq \varepsilon \log^2 n$ separately. In the former case, the heaviest element among H and H' are roughly $\widetilde{\Theta}(\varepsilon/\sqrt{kn})$. Hence, when the algorithm takes $\widetilde{o}(\sqrt{kn}/\varepsilon)$ samples, it rarely sees any element appearing a large number of times. By the moment-matching property of U and U', the probabilities of seeing some elements appearing for t times for $t \leq \log n$ are almost identical under H and H', therefore making H and H' indistinguishable. In the latter case, we have $\varepsilon U \ll \frac{1}{n}$, implying that no elements in the measures are significantly heavier than the rest. As a result, H and H' are both almost uniform except with a different number of "bumps" (elements that are slightly heavier). Subsequently, the algorithm needs more samples (about $\widetilde{\Omega}(k/\varepsilon^2)$) to tell whether a certain element is heavier than the rest, leading to a phase transition in the sample complexity of the problem. We remark that whether $\widetilde{\Omega}(k/\varepsilon^2)$ or $\widetilde{\Omega}(\sqrt{nk}/\varepsilon)$ dominates depends exactly on the relationship between $\sqrt{k/n}$ and ε (omitting poly-logarithmic factors). Combining the two regimes then gives us the following lower bound, whose proof is provided in Appendix B.2.

Proposition 2. *There exists a constant $\nu \in (0, 1)$ such that for any sufficiently large n and $\varepsilon \in (0, 1/10)$, it holds $m_{\mathrm{hist}}^{\mathrm{poi}}(n, k, \varepsilon, \nu) \geq \Omega(\max(\sqrt{kn}/(\varepsilon \log n), k/(\varepsilon^2 \log^3 n)))$.*

Finally, we can easily translate our lower bound result in the Poissonized sampling model to the Multinomial (standard fixed-size) sampling model by a standard reduction. Combining it with the known $\Omega(\sqrt{n}/\varepsilon^2)$ bound (see [Can16, Proposition 4.1]) then concludes our lower bound argument. Formal proofs are given in Appendix B.3.

References

[ADH+15] J Acharya, I. Diakonikolas, C. Hegde, J. Li, and L. Schmidt. Fast and near-optimal algorithms for approximating distributions by histograms. In Tova Milo and Diego Calvanese, editors, *Proceedings of the 34th ACM Symposium on Principles of Database Systems, PODS 2015*, pages 249–263. ACM, 2015.

[ADK15] J. Acharya, C. Daskalakis, and G. Kamath. Optimal testing for properties of distributions. In *NeurIPS*, pages 3591–3599, 2015.

[ADLS17] J. Acharya, I. Diakonikolas, J. Li, and L. Schmidt. Sample-optimal density estimation in nearly-linear time. In *Proceedings of the Twenty-Eighth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2017*, pages 1278–1289, 2017. Full version available at https://arxiv.org/abs/1506.00671.

[BFR+00] T. Batu, L. Fortnow, R. Rubinfeld, W. D. Smith, and P. White. Testing that distributions are close. In *IEEE Symposium on Foundations of Computer Science*, pages 259–269, 2000.

[BFR+13] T. Batu, L. Fortnow, R. Rubinfeld, W. D. Smith, and P. White. Testing closeness of discrete distributions. *J. ACM*, 60(1):4, 2013.

[Can] C. L. Canonne. A short note on poisson tail bounds.

[Can16] C. L. Canonne. Are few bins enough: Testing histogram distributions. In *Proceedings of the 35th ACM SIGMOD-SIGACT-SIGAI Symposium on Principles of Database Systems*, PODS '16, page 455–463, New York, NY, USA, 2016. Association for Computing Machinery.

[Can20] C. L. Canonne. *A Survey on Distribution Testing: Your Data is Big. But is it Blue?* Number 9 in Graduate Surveys. Theory of Computing Library, 2020.

[Can22] C. L. Canonne, 2022. Personal communication. Corrigendum for [Can16] sent to the conference.

[CDGR18] C. L. Canonne, I. Diakonikolas, T. Gouleakis, and R. Rubinfeld. Testing shape restrictions of discrete distributions. *Theory Comput. Syst.*, 62(1):4–62, 2018. Invited issue for STACS'16.

[CDSS14] S. Chan, I. Diakonikolas, R. Servedio, and X. Sun. Near-optimal density estimation in near-linear time using variable-width histograms. In *NIPS*, pages 1844–1852, 2014.

[CGHJ12] G. Cormode, M. Garofalakis, P. J. Haas, and C. Jermaine. Synopses for massive data: Samples, histograms, wavelets, sketches. *Found. Trends databases*, 4:1–294, 2012.

[CMN98] S. Chaudhuri, R. Motwani, and V. R. Narasayya. Random sampling for histogram construction: How much is enough? In *SIGMOD Conference*, pages 436–447, 1998.

[DK16] I. Diakonikolas and D. M. Kane. A new approach for testing properties of discrete distributions. In *2016 IEEE 57th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 685–694. IEEE, 2016.

[DKN15a] I. Diakonikolas, D. M. Kane, and V. Nikishkin. Optimal algorithms and lower bounds for testing closeness of structured distributions. In *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015*, pages 1183–1202, 2015.

[DKN15b] I. Diakonikolas, D. M. Kane, and V. Nikishkin. Testing identity of structured distributions. In *Proceedings of the Twenty-Sixth Annual ACM-SIAM Symposium on Discrete Algorithms, SODA 2015*, pages 1841–1854, 2015.

[DKN17] I. Diakonikolas, D. M. Kane, and V. Nikishkin. Near-optimal closeness testing of discrete histogram distributions. In *44th International Colloquium on Automata, Languages, and Programming, ICALP 2017*, pages 8:1–8:15, 2017.

[DL04] L. Devroye and G. Lugosi. Bin width selection in multivariate histograms by the combinatorial method. *Test*, 13(1):129–145, 2004.

[DLS18] I. Diakonikolas, J. Li, and L. Schmidt. Fast and sample near-optimal algorithms for learning multidimensional histograms. In *Conference On Learning Theory, COLT 2018*, volume 75 of *Proceedings of Machine Learning Research*, pages 819–842. PMLR, 2018.

[FD81] D. Freedman and P. Diaconis. On the histogram as a density estimator:l2 theory. *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 57(4):453–476, 1981.

[GGI+02] A. C. Gilbert, S. Guha, P. Indyk, Y. Kotidis, S. Muthukrishnan, and M. Strauss. Fast, small-space algorithms for approximate histogram maintenance. In *STOC*, pages 389–398, 2002.

[GKS06] S. Guha, N. Koudas, and K. Shim. Approximation and streaming algorithms for histogram construction problems. *ACM Trans. Database Syst.*, 31(1):396–438, 2006.

[GMP97] P. B. Gibbons, Y. Matias, and V. Poosala. Fast incremental maintenance of approximate histograms. In *VLDB*, pages 466–475, 1997.

[Han19] Y. Han. Mixture vs. mixture and moment matching, 2019.

[ILR12] P. Indyk, R. Levi, and R. Rubinfeld. Approximating and Testing k-Histogram Distributions in Sub-linear Time. In *PODS*, pages 15–22, 2012.

[JKM+98] H. V. Jagadish, N. Koudas, S. Muthukrishnan, V. Poosala, K. C. Sevcik, and T. Suel. Optimal histograms with quality guarantees. In *VLDB*, pages 275–286, 1998.

[JVYHW15] J. Jiao, K. Venkat, Yanjun Y. Han, and T. Weissman. Minimax estimation of functionals of discrete distributions. *IEEE Trans. Inform. Theory*, 61(5):2835–2885, 2015.

[Kle09] J. Klemela. Multivariate histograms with data-dependent partitions. *Statistica Sinica*, 19(1):159–176, 2009.

[Koo80] R. P. Kooi. *The Optimization of Queries in Relational Databases*. PhD thesis, 1980.

[KOPS15] S. Kamath, A. Orlitsky, D. Pichapati, and A. T. Suresh. On learning distributions from their samples. In Peter Grünwald, Elad Hazan, and Satyen Kale, editors, *Proceedings of The 28th Conference on Learning Theory*, volume 40 of *Proceedings of Machine Learning Research*, pages 1066–1100, Paris, France, 03–06 Jul 2015. PMLR.

[LN96] G. Lugosi and A. Nobel. Consistency of data-driven histogram methods for density estimation and classification. *Ann. Statist.*, 24(2):687–706, 04 1996.

[Pan08] L. Paninski. A coincidence-based test for uniformity given very sparsely sampled discrete data. *IEEE Transactions on Information Theory*, 54(10):4750–4755, 2008.

[Rub06] R. Rubinfeld. Sublinear time algorithms. Proceedings of the International Congress of Mathematicians (ICM), 2006.

[Sco79] D. W. Scott. On optimal and data-based histograms. *Biometrika*, 66(3):605–610, 1979.

[TGIK02] N. Thaper, S. Guha, P. Indyk, and N. Koudas. Dynamic multidimensional histograms. In *SIGMOD Conference*, pages 428–439, 2002.

[Val11] P. Valiant. Testing symmetric properties of distributions. *SIAM J. Comput.*, 40(6):1927–1968, 2011.

[VV13] G. Valiant and P. Valiant. Estimating the unseen: improved estimators for entropy and other properties. *Advances in Neural Information Processing Systems*, 26, 2013.

[WY16] Y. Wu and P. Yang. Minimax rates of entropy estimation on large alphabets via best polynomial approximation. *IEEE Trans. Inform. Theory*, 62(6):3702–3720, 2016.

[WY+19] Y. Wu, P. Yang, et al. Chebyshev polynomials, moment matching, and optimal estimation of the unseen. *Annals of Statistics*, 47(2):857–883, 2019.

A Upper Bound

We provide in this appendix the proofs of some technical lemmas, which were omitted from the main paper due to space constraints.

Fact 3. *Let* $\mathbf{u}$ *be the uniform distribution and* $\mathbf{p}$ *be an arbitrary distribution among* $[n]$. *We can sample from* $\mathbf{p}' := 1/2(\mathbf{p} + \mathbf{u})$ *efficiently with sample access to* $\mathbf{p}$.

A.1 Proof of Equitable Partition

Lemma 7 (Claim 1 in [ADK15]). *There exists an algorithm **Approx-Divide** that, given parameters* $\delta \in (0, 1]$ *and integer* $B > 1$, *takes* $O\big(B \log(B/\delta)\big)$ *samples from a distribution* $\mathbf{p}$ *on* $[n]$ *and, with probability at least* $1 - \delta$, *outputs a partition of* $[n]$ *into* $K \leq 8B$ *intervals* $I_1, I_2, \ldots, I_K$ *such that the following holds:*

1. *For each* $i \in [n]$ *such that* $\mathbf{p}(i) > 1/B$, *there exists* $\ell \in [K]$ *such that* $I_\ell = \{i\}$. *(All "heavy" elements are left as singletons)*

2. *Every other interval is such that* $\mathbf{p}(I) \leq 16/B$.

Proof. Let $B > 1$, and consider an arbitrary (unknown) probability distribution $\mathbf{p}$ over $[n]$. Let $\hat{\mathbf{p}}$ be the empirical distribution obtained by taking $m = \left\lceil 18B \ln \frac{12B}{\delta} \right\rceil = O(B \log(B/\delta))$ i.i.d. samples from $\mathbf{p}$, where $C > 0$ is an absolute constant to be chosen later in the proof.

Denote by $i_1 < \cdots < i_T$ the elements i such that $\mathbf{p}(i) > 1/B$; note that $0 \leq T \leq B$ since there can be at most B elements with probability mass greater than $1/B$.

Consider the following (deterministic, and unknown to the algorithm) partition of the domain into consecutive intervals:

- each of $i_1 < \cdots < i_T$ is a singleton interval $J_{\ell_j} = \{i_j\}$;

- setting for convenience $i_0 = -1$ and $i_{T+1} = n + 1$, define the remaining intervals greedily as follows. For each $1 \leq j \leq T$, to partition $\{i_j + 1, i_j + 2, \ldots, i_{j+1} - 1\}$, do

  1. set $J \leftarrow \emptyset, t \leftarrow i_j + 1$
  2. while $t < i_{j+1}$
     - if $\mathbf{p}(J \cup \{t\}) < 3/(2B)$, set $J \leftarrow J \cup \{t\}$
     - else add J as an interval to the partition, then start a new interval: $J \leftarrow \emptyset$
     - consider the next element: $t \leftarrow t + 1$

That is, every remaining interval ("in-between" two heavy elements i_j, i_{j+1}) is greedily divided from left to right into maximal subintervals of probability mass at most $3/(2B)$. Note that since every element $i \in \{i_j + 1, \ldots, i_{j+1} - 1\}$ has mass at most $1/B$, this is indeed possible and leads to a partition of $\{i_j + 1, \ldots, i_{j+1} - 1\}$ in intervals, where all but at most one (the rightmost one) has probability mass $\mathbf{p}(J) \in [1/(2B), 3/(2B)]$.

Overall, this process (which the algorithm cannot directly run, not knowing $\mathbf{p}$) guarantees the existence of a partition $J_1, \ldots J_S$ of the domain in at most

$$S \leq T + T + 2B \leq 4B$$

consecutive, disjoint intervals (the T singleton intervals, the at most T "rightmost, in-between" intervals which may have probability mass at most $1/(2B)$, and the remaining "in-between" intervals which all have mass at least $1/(2B)$, and of which there can thus be at most $2B$ in total).

Now, consider the following partition $\hat{J}_1, \ldots \hat{J}_K$ defined by the algorithm from $\hat{\mathbf{p}}$:

- every element $i \in [n]$ such that $\hat{\mathbf{p}}(i) > 1/(2B)$ is added as a singleton interval; let their number be T';

- then the algorithm proceeds as in the above process (which defined $J_1, \ldots J_S$), but using $\hat{\mathbf{p}}$ instead of $\mathbf{p}$.

The same analysis (but with $T' \leq 2B$ instead of $T \leq B$, since the threshold for "singletons" is now $1/(2B)$ shows that this will result in a partition of the domain into K consecutive disjoint intervals $\hat{J}_1, \ldots \hat{J}_K$, with

$$K \leq 8B.$$

It remains to argue about the properties of $\hat{J}_1, \ldots \hat{J}_K$, by using those of (the unknown) $J_1, \ldots J_S$. First, we have that, for every $1 \leq j \leq T$, by a Chernoff bound,

$$\Pr\left[\hat{\mathbf{p}}(i_j) \leq \frac{1}{2B} \right] \leq e^{-(1/2)^2 \frac{m}{2B}} \leq \frac{\delta}{3B}$$

from our setting our m. Now, for any interval J_j (where $1 \leq j \leq S$) such that $1/(2B) \leq \mathbf{p}(J_j) \leq 3/(2B)$ (there are at most $2B$ of them), we also have, all by a Chernoff bound,

$$\Pr\left[\hat{\mathbf{p}}(i_j) \notin \left[\frac{1}{4B}, \frac{2}{B} \right] \right] \leq 2e^{-\frac{m}{18B}} \leq \frac{\delta}{6B}$$

while for any J_j such that $\mathbf{p}(J_j) < 1/(2B)$ (there are at most T of them)

$$\Pr[\hat{\mathbf{p}}(i_j) \geq 1/B] \leq e^{-\frac{m}{6B}} \leq \frac{\delta}{3B}.$$

This means, by a union bound, that with probability at least $1 - \left(T \cdot \frac{\delta}{3B} + 2B \cdot \frac{\delta}{6B} + T \cdot \frac{\delta}{3B} \right) \geq 1 - \delta$:

- all of the $(1/B)$-heavy elements for $\mathbf{p}$ will also be $(1/(2B))$-heavy elements for $\hat{\mathbf{p}}$, and thus constitute their own singleton interval, as desired;

- all of the intervals J_j such that $1/(2B) \leq \mathbf{p}(J_j) \leq 3/(2B)$ (call those "heavy") satisfy $1/(4B) \leq \hat{p}(J_j) \leq 2/B$;

- all of the intervals J_j such that $\mathbf{p}(J_j) < 1/(2B)$ (call those "light") satisfy $\hat{p}(J_j) \leq 2/B$.

Conditioning on this happening, we can analyze the properties of $\hat{J}_1, \ldots \hat{J}_K$. Specifically, fix any $\hat{J}_j$ which is not a singleton of the form $\{i_\ell\}$ for one of the T $(1/B)$-heavy elements for $\mathbf{p}$. We want to argue that $\mathbf{p}(\hat{J}_j) \leq 16/B$.

- If $\hat{J}_j$ intersects with at most 8 $J_{j'}$'s (which must then all be consecutive, and at least 7 are "heavy", and the rightmost one is either heavy or light), we are done, since then $\mathbf{p}(\hat{J}_j) \leq 8 \cdot 2/B = 16/B$.

- Moreover, $\hat{J}_j$ cannot intersect with more than 8 $J_{j'}$'s, as otherwise those are consecutive, and thus $\hat{J}_j$ must contain at least 6 "heavy" $J_{j'}$'s. But in that case $\hat{\mathbf{p}}(\hat{J}_j) \geq 6 \cdot 1/(4B) = 3/(2B)$, while by our greedy construction we ensured that $\hat{\mathbf{p}}(\hat{J}_j) < 3/(2B)$.

This concludes the proof. $\qquad\square$

Proof of Lemma 2. Reduction from Lemma 7. $\qquad\square$

A.2 Proof of Interval Mass Estimate

Algorithm 2 Interval-Mass-Estimate

Require: m i.i.d. samples from distribution $\mathbf{p}$ on $[n]$; failure probability δ.

1: Let $T := 3\log(n/\delta)$.
2: Split the samples into T batches $S^{(1)}, S^{(2)}, \cdots S^{(T)}$, each of m/T samples.
3: Initialize a Map $\hat{\varphi} : \mathcal{I} \mapsto \mathbb{R}$ which maps sub-intervals of $[n]$ to real values.
4: **for** all intervals $I \subseteq [n]$ **do**
5:    Let $X_j^{(i)}$ be the number of samples falling in interval I_j from samples $S^{(i)}$.
6:    Compute the median $X_j = \text{Median}_{1\leq i \leq T}\left(X_j^{(i)}\right)$. Accordingly, set

$$\hat{\varphi}(I) = \max\left(\frac{X_j}{m}, \frac{|I|}{2n}\right).$$

7: Output the map $\hat{\varphi}$.

Claim 4. *Let I be an interval (or subset) over $[n]$ and $\mathbf{p}$ a distribution over $[n]$. For $\delta \in (0,1)$, assume one takes $T := 3\log(1/\delta)$ batches of i.i.d. samples from $\mathbf{p}$ where each batch is of size $b \in \mathbb{Z}^+$. Denote by $X_I^{(i)}$ the number of samples falling in I from the i-th batch, for $i \leq [T]$. Then, the median $X_I := \text{Median}_i(X_I^{(i)})$ satisfies*

$$\left|\frac{X_I}{b} - \mathbf{p}(I)\right| \leq 3 \cdot \sqrt{\frac{\mathbf{p}(I)}{b}} \quad \text{and} \quad \frac{X_I}{b} \leq 3 \cdot \mathbf{p}(I),$$

with probability at least $1 - \delta$.

Proof of Lemma 3. Let $I \subseteq [n]$ be an arbitrary interval and X_I be the median of the numbers of samples falling in from each batch. By Claim 4, we have that

$$\left|\frac{X_I}{m} - \mathbf{p}(I)\right| \leq \sqrt{\mathbf{p}(I)/b}, \tag{1}$$

$$\frac{X_I}{m} \leq 3\mathbf{p}(I). \tag{2}$$

with probability at least $1 - \delta/(2n)$. Since $\mathbf{p}(i) > 1/(2n)$ for any $i \in [n]$, the max operation decreases the distance between $\mathbf{p}$ and $\hat{\mathbf{p}}$ pointwise. Hence, with probability at least $1 - \delta/2K$, we have

$$|\hat{\varphi}(I) - \mathbf{p}(I)| \leq \sqrt{\mathbf{p}(I)/b}. \tag{3}$$

Since there are at most n^2 sub-intervals $I \subseteq [n]$ overall, by a union bound over all sub-intervals, we have that Equation (3) holds simultaneously with probability at least $1 - \delta/2$ for every sub-interval I.

To show that the ratio $\mathbf{p}(I)/\hat{\varphi}(I)$ is bounded, we first consider the case $\mathbf{p}(I) \leq 4/b$. Since $\hat{\varphi}(I) > 1/(2n)$, it is easy to see that $\mathbf{p}(I)/\hat{\varphi}(I) \leq 8 \cdot b/n$. Otherwise, we have $\mathbf{p}(I) > 4/b$. Following Equation (3). it holds

$$|\mathbf{p}(I) - \hat{\varphi}(I)| \leq \sqrt{\mathbf{p}(I)/b} \leq \frac{1}{2}\mathbf{p}(I),$$

This then implies that

$$\frac{\mathbf{p}(I)}{\hat{\varphi}(I)} \leq 2.$$

Lastly, since $\mathbf{p}(I) \geq |I|/(2n)$, together with Equation (2), we have

$$\frac{\hat{\varphi}(I)}{\mathbf{p}(I)} = \frac{\max\left(|I|/(2n), X_I/m\right)}{\mathbf{p}(I)} \leq 3.$$

$\square$

A.3 Proof of Empirical Learning

Algorithm 3 Empirical-Learning

Require: m i.i.d. samples from distribution $\mathbf{p}$ on [n]; partition $\mathcal{I} = \{I_1, I_2, \cdots, I_K\}$ of the domain $[n]$; error rate δ
1: Denote the m i.i.d samples as $\mathcal{S}$.
2: Construct the Interval Estimator $\mathcal{P} \leftarrow$ **Interval-Mass-Estimate**$(\mathcal{S}, \delta)$.
3: **for** $j = 1 \cdots K$ **do**
4:     Compute the estimate $\hat{\mathbf{p}}(i) = \frac{\mathcal{P}(I_j)}{|I_j|}$ $1 \leq j \leq K, i \in I_j$.
5: Output the measure $\hat{\mathbf{p}}$.

A.4 Detailed Analysis of Learn-And-Sieve

Proof of Lemma 4. Let I be a non-breakpoint interval. It is easy to see that $\hat{\mathbf{p}}(I) = \hat{\varphi}(I)$. By Lemma 3, it holds that

$$\frac{\mathbf{p}(I)}{\hat{\mathbf{p}}(I)} \leq \max(2, 8 \cdot n/b),$$

$$|\hat{\mathbf{p}}(I) - \mathbf{p}(I)| \leq \sqrt{\mathbf{p}(I)/b}.$$

with probability at least $1 - \delta$. Condition on that, we easily have, for any sub-interval $J \subseteq I$,

$$\frac{\mathbf{p}(J)}{\hat{\mathbf{p}}(J)} = \frac{\mathbf{p}(I)}{\hat{\mathbf{p}}(I)} \leq \max(2, 8 \cdot n/b)$$

since both $\mathbf{p}$ and $\hat{\mathbf{p}}$ are uniform within I. Furthermore, we also have

$$|\mathbf{p}(J) - \hat{\mathbf{p}}(J)| = |\mathbf{p}(I) - \hat{\mathbf{p}}(I)| \cdot \frac{|J|}{|I|} \leq \sqrt{\frac{\mathbf{p}(I)}{m}} \cdot \frac{|J|}{|I|} \leq \sqrt{\frac{\mathbf{p}(J)}{m}},$$

where the first equality follows from the fact that $\mathbf{p}$ and $\hat{\mathbf{p}}$ are both uniform within I, the second inequality follows from our conditioning, the third inequality follows from $|J| \leq |I|$. □

Proof of Lemma 5. The first point is easy to see since otherwise Line 7 would reject. We claim that the following holds with probability at least $1 - \delta$. If $\mathbf{p} \in \mathcal{H}_k$, $\mathcal{B}$ contains all the ε-bad intervals and no non-breakpoint intervals. Condition on that, it is easy to see $\mathcal{B}$ would contain no more than k intervals since there are at most k break-point intervals. Besides, for any $I \in \mathcal{G} := \mathcal{I} \backslash \mathcal{B}$, we have $d_{\chi^2}^I (\mathbf{p} \| \hat{\mathbf{p}}) \leq \varepsilon^2/K$. Overall, it then holds $d_{\chi^2}^{\mathcal{I} \backslash \mathcal{B}} (\mathbf{p} \| \hat{\mathbf{p}}) \leq \varepsilon^2$. This concludes the proof of the second point.

It remains to show why $\mathcal{B}$ separates the bad intervals from the rest. We will separately establish that with high probability $\mathcal{B}$ will not include any non-breakpoint intervals and that it will include all bad intervals separately, and conclude by a union bound. The following statement, which follows from Lemma 3, will be useful in both aspects of the analysis. In particular, it holds

$$|\hat{\varphi}(Q) - \mathbf{p}(Q)| \leq \sqrt{\mathbf{p}(Q)/b} \tag{4}$$

$$\hat{\varphi}(Q) \leq 3 \cdot \mathbf{p}(Q) \tag{5}$$

for all sub-intervals Q with probability at least $1 - \delta/2$. We will condition on that in the analysis.

Exclusion of non-breakpoint intervals By Lemma 4, if I is not a break-point interval, then for every sub-interval Q, with probability at least $1 - \delta/3$, we have

$$|\hat{\mathbf{p}}(Q) - \mathbf{p}(Q)| \leq \sqrt{\mathbf{p}(Q)/b} \tag{6}$$

$$\frac{\mathbf{p}(Q)}{\hat{\mathbf{p}}(Q)} \leq \max(2, 8 \cdot n/b), \tag{7}$$

As a result, we have

$$(\hat{\varphi}(Q) - \hat{\mathbf{p}}(Q))^2 \leq 4 \cdot \mathbf{p}(Q)/b \leq O(1/b + n/b^2) \cdot \hat{\mathbf{p}}(Q) \leq \frac{1}{4}\hat{\mathbf{p}}(Q) \cdot \varepsilon^2/K\,,$$

where the first inequality follows Equations (6), (4) and the triangle inequality, the second inequality follows from the bounded ratio between $\hat{\mathbf{p}}(Q)$ and $\mathbf{p}(Q)$ (Equation (7)) and the last inequality follows from $b \gg \sqrt{nK}/\varepsilon + K/\varepsilon^2$. This asserts that second condition of Line 5 will be false.

Next, we shift our focus to the first condition of Line 5. Together with Equation (7), which is the condition in Lemma 4, and Equation (5), the guarantee obtained from Claim 3, we have

$$\frac{\hat{\varphi}(Q)}{\hat{\mathbf{p}}(Q)} = \frac{\hat{\varphi}(Q)}{\mathbf{p}(Q)} \cdot \frac{\mathbf{p}(Q)}{\hat{\mathbf{p}}(Q)} \leq \max\left(6\,, 24 \cdot n/b\right). \leq \max\left(6\,, \varepsilon\sqrt{n/K}\right).$$

where the last inequality follows from $b \gg \sqrt{nK}/\varepsilon$. This asserts that Line 5 won't pass. Hence, with probability at least $1 - \delta$, $\mathcal{B}$ won't include any non-breakpoint intervals.

Inclusion of ε-bad intervals Let I be an ε-bad interval. Then, there exists a sub-interval Q such that $\mathbf{p}$ is constant within Q and $\frac{(\mathbf{p}(Q) - \hat{\mathbf{p}}(Q))^2}{\hat{\mathbf{p}}(Q)} \geq \varepsilon^2/K$, which implies that

$$|\mathbf{p}(Q) - \hat{\mathbf{p}}(Q)| \geq \sqrt{\hat{\mathbf{p}}(Q) \cdot \frac{\varepsilon^2}{K}}. \tag{8}$$

We then proceed to analyze the cases $\mathbf{p}(Q) \leq 2 \cdot \frac{\varepsilon}{\sqrt{Kn}}$ and $\mathbf{p}(Q) > 2 \cdot \frac{\varepsilon}{\sqrt{Kn}}$ separately.

In the former case, we have

$$\frac{\mathbf{p}(Q)}{b} = \frac{\hat{\mathbf{p}}(Q)}{b} \cdot \frac{\mathbf{p}(Q)}{\hat{\mathbf{p}}(Q)} \leq \hat{\mathbf{p}}(Q) \cdot O\left(\varepsilon \cdot \sqrt{\frac{n}{K}}\right) \cdot \frac{1}{b} \leq \frac{1}{4}\hat{\mathbf{p}}(Q) \cdot \frac{\varepsilon^2}{K}\,,$$

where the first inequality follows from that $\mathbf{p}(Q) \leq 2\varepsilon/\sqrt{nK}$ and $\hat{\mathbf{p}}(Q) \geq \frac{1}{2n}$, and the second inequality follows from $b \gg \sqrt{nK}/\varepsilon$. This together with Equations (4), (8) and triangle inequality gives

$$|\hat{\mathbf{p}}(Q) - \hat{\varphi}(Q)| \geq \frac{1}{2}\sqrt{\hat{\mathbf{p}}(Q) \cdot \frac{\varepsilon^2}{K}}\,,$$

which asserts that the second if condition of Line 5 is true.

In the latter case, since $\mathbf{p}(Q) > 2\frac{\varepsilon}{\sqrt{nK}} \gg \frac{1}{b}$, it holds $\frac{\mathbf{p}(Q)}{b} \ll \mathbf{p}(Q)^2$. From Equation (4) we have

$$\hat{\varphi}(Q) \geq \mathbf{p}(Q) - \sqrt{\mathbf{p}(Q)/b} \geq \frac{1}{2}\mathbf{p}(Q). \tag{9}$$

If $\hat{\varphi}(Q)/\hat{\mathbf{p}}(Q) \geq \max(6, \varepsilon\sqrt{n/K})$, the first condition of Line 5 is true, implying I will be included in $\mathcal{B}$. Otherwise, we could argue that

$$\frac{\mathbf{p}(Q)}{b} = \frac{\hat{\mathbf{p}}(Q)}{b} \cdot \frac{\mathbf{p}(Q)}{\hat{\varphi}(Q)} \cdot \frac{\hat{\varphi}(Q)}{\hat{\mathbf{p}}(Q)} \leq \frac{\hat{\mathbf{p}}(Q)}{b} \cdot 2 \cdot \max(6, \varepsilon\sqrt{n/K}) \leq \frac{1}{4}\hat{\mathbf{p}}(Q)\varepsilon^2/K\,,$$

where the first inequality follows from Equation (9) and the fact that the first condition of Line 5 is false, and the second inequality follows from $b \gg \sqrt{nK}/\varepsilon + K/\varepsilon^2$. In other words, we have

$$\sqrt{\frac{\mathbf{p}(Q)}{b}} \leq \frac{1}{2}\sqrt{\hat{\mathbf{p}}(Q) \cdot \frac{\varepsilon^2}{K}}.$$

This together with Equations (4), (8) and triangle inequality allows us to conclude that

$$|\hat{\mathbf{p}}(Q) - \hat{\varphi}(Q)| \geq \frac{1}{2}\sqrt{\hat{\mathbf{p}}(Q) \cdot \frac{\varepsilon^2}{K}}\,,$$

which asserts that the second condition of Line 5 is true. $\qquad\square$

A.5 Main Testing Algorithm

Algorithm 4 Divide-And-Learn-And-Sieve

Require: Sample access to the distribution $\mathbf{p}$; domain size $[n]$; accuracy ε.
1: Set $\mathcal{I}^{(0)} = \mathcal{B}^{(0)} = \{[n]\}$.
2: Set $T \leftarrow 3\log(1/\varepsilon)$, $\delta \leftarrow \frac{1}{100T}$, $t \leftarrow 0$, $r = 1$.
3: **while** $r > \varepsilon/8$ **do**
4:      $\mathcal{S}^{(t+1)} \leftarrow$ **Approx-Sub-Divide** $\left(32k, \mathcal{B}^{(t)}, \delta\right)$.
5:      Set $\mathcal{I}^{(t+1)} = (\mathcal{I}^{(t)} \setminus \mathcal{B}^{(t)}) \cup \mathcal{S}^{(t+1)}$. $\triangleright$ Note that $\mathcal{I}^{(t+1)}$ is still a partition of $[n]$.
6:      $\mathcal{Q}^{(t+1)}, \hat{\mathbf{p}}^{(t+1)} \leftarrow$ **Learn-And-Sieve** $\left(\mathcal{I}^{(t+1)}, \frac{\varepsilon}{4\sqrt{T}}, \delta\right)$.
7:      Reject if **Learn-And-Sieve** outputs Reject.
8:      Set $\mathcal{B}^{(t+1)} = \mathcal{S}^{(t+1)} \cap \mathcal{Q}^{(t+1)}$.
9:      Take $\ell = \Theta\left(\log(1/\delta)/\varepsilon\right)$ i.i.d. samples from $\mathbf{p}$.
10:      Denote X as the number of samples falling in $\mathcal{B}^{(t+1)}$. Set $r = X/\ell$.
11:      $t \leftarrow t + 1$
12: Denote $\mathcal{G}^{(j)} = \mathcal{S}^{(j)} \setminus \mathcal{B}^{(j)}$ for all $j \geq 1$. $\triangleright$ Note that the union of $\mathcal{G}^{(1)}, \cdots \mathcal{G}^{(t)}, \mathcal{B}^{(t)}$ forms a partition of the domain $[n]$.
13: We will consider the measure $\bar{\mathbf{p}}$ such that on intervals $I \in \mathcal{G}^{(j)}$, $\bar{\mathbf{p}}(i) = \hat{\mathbf{p}}^{(j)}(i)$; and on intervals $I \in \mathcal{B}^{(t)}$, $\bar{\mathbf{p}}(i) = 0$.
14: Use Dynamic Programming to check whether there is a k-histogram that is $\varepsilon/2$-close to $\bar{\mathbf{p}}$ in ℓ_1 distance. Otherwise Reject. $\triangleright$ Can be done in time $\text{poly}(k, 1/\varepsilon, n)$ as shown in Lemma 4.11 of [CDGR18].
15: Output Accept if **Tolerant-Identity-Test** $(\mathbf{p}, \hat{\mathbf{p}}, \varepsilon)$ outputs Accept. Otherwise Reject.

Proof of Upper Bound in Theorem 1. We first argue that the algorithm terminates in $T = O(\log(1/\varepsilon))$ rounds with high probability. By Lemma 7, we have $\mathbf{p}(I) \leq \frac{16}{32k} \cdot \mathbf{p}(\mathcal{B}^{(t)}) = \frac{1}{2k} \cdot \mathbf{p}(\mathcal{B}^{(t)})$ for every non-singleton interval $I \in \mathcal{S}^{(t+1)}$ with probability at least $1 - \delta$. By Lemma 5, the subroutine **Learn-And-Sieve** selects (removes) at most k intervals if it does not output reject. Notice that $\mathcal{B}^{(t+1)}$ will not include any singleton as singletons cannot be break-point intervals. Then, it holds that $\mathbf{p}(\mathcal{B}^{(t+1)}) \leq \mathbf{p}(\mathcal{Q}^{(t+1)}) \leq \frac{1}{2}\mathbf{p}(\mathcal{B}^{(t)})$. Hence, the mass of $\mathcal{B}^{(t)}$ will drop below $\varepsilon/100$ after at most $T := 3 \cdot \log(1/\varepsilon)$ iterations with probability at least

$$(1 - \delta)^T = (1 - \frac{1}{100T})^T \geq \frac{9}{10}, \tag{10}$$

where the second inequality holds when T is sufficiently large. On the other hand, in the end of iteration, we always $r \leq 10 \cdot \mathbf{p}(\mathcal{B}^{(t)}) \leq \varepsilon/10$ with probability at least $\frac{9}{10}$ by Markov's Inequality. Combining the two facts then gives the algorithm exits the while loop in at most $T := 3 \cdot \log(1/\varepsilon)$ iterations with probability at least $0.9 \cdot 0.9 > 8/10$.

Furthermore, we claim, with probability at least $9/10$, it holds $\mathbf{p}(\mathcal{B}^{(t)}) \leq \varepsilon/4$ when the algorithm exits the while loop. Suppose at the j-th iteration, we have $\mathbf{p}(\mathcal{B}^{(j)}) > \varepsilon/4$. Then, by the multiplicative Chernoff Bound, we have

$$\Pr\left[r \leq \varepsilon/8\right] \leq \Pr\left[r \leq 1/2 \cdot \mathbf{p}(\mathcal{B}^{(j)})\right] \leq 1/\delta.$$

Hence, following the same calculation as Equation (10), our claim holds with probability at least $9/10$. Conditioning on (i) the algorithm terminates in T iterations and (ii)

$$\mathbf{p}(\mathcal{B}^{(t)}) \leq \varepsilon/4 \tag{11}$$

when the algorithm exits the loop, we proceed to argue it outputs the correct testing result with probability at least $2/3$.

Completeness   Suppose we have $\mathbf{p} \in \mathcal{H}_k^n$. At the t-th iteration, we claim that, with probability at least $1 - \delta$, it holds

$$d_{\chi^2}^{\mathcal{G}^{(t)}}\left(\mathbf{p}\|\hat{\mathbf{p}}^{(t)}\right) \leq \frac{\varepsilon^2}{16T}, \tag{12}$$

where $\mathcal{G}^{(t)} = \mathcal{S}^{(t)} \backslash \mathcal{B}^{(t)}$ as defined in 12 and $\hat{\mathbf{p}}^{(t)}$ is the learned distribution in the t-th iteration. By Lemma 5, it holds

$$d_{\chi^2}^{\mathcal{I}^{(t)} \backslash \mathcal{Q}^{(t)}} \left(\mathbf{p} \| \hat{\mathbf{p}}^{(t)} \right) \leq \frac{\varepsilon^2}{16T}.$$

Since $\mathcal{G}^{(t)} = \mathcal{S}^{(t)} \backslash \mathcal{B}^{(t)}$ is a subset of $\mathcal{I}^{(t)} \backslash \mathcal{Q}^{(t)}$, the claim in Eq. (12) follows. Recall that we condition on the algorithm runs for at most T iterations. Combining this with Eq. (12), if we denote $\mathcal{G} = \bigcup \mathcal{G}^t$ for $1 \leq t \leq T$, it holds

$$d_{\chi^2}^{\mathcal{G}} \left(\mathbf{p} \| \bar{\mathbf{p}} \right) \leq \sum_{t=1}^{T} d_{\chi^2}^{\mathcal{G}^{(t)}} \left(\mathbf{p} \| \hat{\mathbf{p}}^{(t)} \right) \leq \varepsilon^2/16 \qquad (13)$$

with probability at least $(1 - \delta)^T \geq \frac{9}{10}$. Notice that $\mathcal{G}$ is precisely the sub-domain $\mathcal{A} = \{i \in [n] : \bar{\mathbf{p}}(i) \geq \varepsilon/(50n)\}$ that will be used to compute the χ^2 statistic since $\bar{\mathbf{p}}(i) \geq \frac{1}{2n}$ for $i \in \mathcal{G}$ and $\bar{\mathbf{p}}(i) = 0$ for $i \notin \mathcal{G}$. Conditioning on Equation (13), by Proposition 1, line 15 will output accept with probability at least $9/10$ by Chebyshev's Inequality. Then, Equation (13) together with the conditioning $\mathbf{p}(\mathcal{B}^{(t)}) \leq \varepsilon/4$ also implies $\mathrm{TV}(\mathbf{p}, \bar{\mathbf{p}}) \leq \varepsilon/2$ and $\mathbf{p} \in \mathcal{H}_k^n$, line 14 will also pass. Overall, the algorithm accepts with probability at least $2/3$.

Soundness Suppose now that $\mathrm{TV}(\mathbf{p}, H) > \varepsilon$ for every $H \in \mathcal{H}_k^n$. For the sake of contradiction, assume that line 14 and line 15 both pass. By line 15 and the contra-positive of Proposition 1, we have $\mathrm{TV}^{\mathcal{G}}(\mathbf{p}, \bar{\mathbf{p}}) \leq \varepsilon/4$ with probability at least $\frac{9}{10}$.

By definition we have $\bar{\mathbf{p}}(\mathcal{B}^{(t)}) = 0$. Since $\mathbf{p}(\mathcal{B}^{(t)}) \leq \varepsilon/4$ by our conditioning, it then holds that $\mathrm{TV}(\mathbf{p}, \bar{\mathbf{p}}) \leq \varepsilon/2$. Then, by line 14, there exists a k-histogram $\mathbf{h}$ satisfying $\mathrm{TV}(\mathbf{h}, \bar{\mathbf{p}}) \leq \varepsilon/2$. By the triangle inequality, we have $\mathrm{TV}(\mathbf{h}, \mathbf{p}) \leq \varepsilon$. This contradicts the assumption that $\mathbf{p}$ is more than ε-far from any k-histogram. Hence, at least one of the two lines will output reject with probability at least $2/3$.

Finally, the samples are consumed by five different types of routines – dividing (Line 4), learning, sieving (Line 6), testing (Line 15) and mass estimation (Line 10). By Lemma 2, in one iteration, the dividing phase consumes $O\left(k \log k / \mathbf{p}\left(\mathcal{B}^{(t)}\right)\right)$ samples where $\mathbf{p}(\mathcal{B}^{(t)})$ is the mass of the to-be-divided intervals at the t-th iteration. Since $\mathbf{p}(\mathcal{B}^{(t)})$ shrinks exponentially in every iteration, the samples consumed are dominated by the last iteration. Hence, at most $O\left(k \log k / \varepsilon\right)$ samples are consumed by the dividing phase in total.

At the t-th iteration, the partition size $K := \left|\mathcal{I}^{(t+1)}\right|$ is upper bounded by $O(T \cdot k)$. Hence, by Lemma 5, the **Learn-And-Sieve** procedure consumes in total

$$O\left(\left(\frac{Tk}{\varepsilon^2} + \frac{\sqrt{Tkn}}{\varepsilon}\right) \cdot \log\left(n \cdot \log(1/\varepsilon)\right) \cdot T\right) \qquad (14)$$

samples. The process of testing the mass of $\mathcal{B}^{(t)}$ consumes $\Theta(T \cdot \log(1/T)/\varepsilon)$ samples in total.

After the algorithm exits the for loop, the chi-square tester consumes $\Theta(\sqrt{n}/\varepsilon^2)$ samples. Thus, in total, the algorithm consumes

$$O\left(\frac{\sqrt{n}}{\varepsilon^2} + \left(\frac{k}{\varepsilon^2} \cdot \log^2(1/\varepsilon) + \frac{\sqrt{kn}}{\varepsilon} \cdot \log^{3/2}(1/\varepsilon)\right) \cdot \log\left(n \cdot \log(1/\varepsilon)\right)\right) \qquad (15)$$

if we sum up the samples consumed by all different routines and substitute $T = 3 \log(1/\varepsilon)$. $\qquad \square$

B Lower Bound

Definition 4 (Histogram Testing under Poisson Model). *For $0 < \nu < 1$, $0 < \varepsilon < 1$, define the sample complexity of histogram testing (under Poisson model) as*

$$m_{\mathrm{hist}}^{\mathrm{poi}}(n, k, \varepsilon, \nu) := \min\left\{m \geq 0 : \exists \hat{I} \text{ such that } \forall P \in \tilde{\mathcal{P}}^n\right.$$

$$\left.\Pr[\hat{I} = 1, \mathbb{1}\{\inf_{P' \in \tilde{\mathcal{H}}_k^n(\nu)} \mathrm{TV}(P', P) \geq \varepsilon(1 + \nu)\}] + \Pr[\hat{I} = 0, \mathbb{1}\{P \in \tilde{\mathcal{H}}_k^n(\nu)\}] \leq 1/10\right\},$$

where $\hat{I}$ is a binary indicator measurable with respect to $M = (M_1, M_2, \ldots) \sim \mathrm{Poi}(mP_i)$.[7]

B.1 Construction of Moment-Matching Random Variables

Fact 5. *Suppose p is a degree-d polynomial with distinct roots $r_1, \cdots r_d$. Then, for every $0 \leq k \leq d - 2$, $\sum_{i=1}^{d} \frac{r_i^k}{p'(r_i)} = 0$.*

Proof of Lemma 6. We construct such a pair explicitly, relying on properties of Chebyshev polynomials. For every integer $d \geq 0$, recall that the corresponding Chebyshev polynomial (of the first kind) T_d is given by $T_d(x) = \cos(d \cdot \arccos(x))$ for $x \in [-1, 1]$. (and can be shown to be a polynomial of degree d). Then, let $\Delta := \frac{\sqrt{kn}}{C \cdot \log^2 n}$ and $d := c \cdot \log n$ (where $c, C > 0$ are two absolute constants, suitably large), and consider the polynomial

$$p(x) = x \left(x - \frac{1}{n} \right) \left(x - \frac{2}{n} \right) T_d \left(1 - \Delta \cdot x \right), \tag{16}$$

for $x \in [0, 1/\Delta]$. This is a degree-$(d+3)$ polynomial, whose roots are listed in the table below along with the corresponding values of its derivative.

Roots	p'
$r_0 = 0$	$\frac{2}{n^2} \cdot T_d(1) = \frac{2}{n^2}$
$r_1 = \frac{1}{n}$	$-\frac{1}{n^2} \cdot T_d \left(1 - \frac{\Delta}{n} \right)$
$r_2 = \frac{2}{n}$	$\frac{2}{n^2} \cdot T_d \left(1 - \frac{2\Delta}{n} \right)$
$r_{2+m} = \frac{1}{\Delta} \left(1 - \cos \left(\frac{2m-1}{2d} \pi \right) \right)$ for $1 \leq m \leq d$	$|p'(r_{2+m})| = \Theta \left(\frac{m^5}{\Delta^2 d^4} \right)$ for $1 \leq m \leq d$.

The roots in the last row are those associated with the Chebyshev Polynomial T_d, and the last line relies on the facts that, for r such that $T(1 - \Delta r) = 0$, we have

$$p'(r) = -\Delta r (r - 1/n)(r - 2/n) T_d'(1 - \Delta r)$$

and that $T_d'(\cos \theta) = \frac{d \sin(d\theta)}{\sin \theta}$. This latter identity implies that $|T_d'(1 - \Delta r_{2+m})| = \frac{d}{|\sin \frac{2m-1}{2d}|} = \Theta \left(\frac{d^2}{m} \right)$, and since $r_{2+m} = \Theta(\frac{m^2}{\Delta d^2})$ we get the claimed bound:

$$|p'(r_{2+m})| = \Theta \left(\Delta \cdot \frac{m^2}{\Delta d^2} \cdot \max \left(\frac{m^2}{\Delta d^2}, \frac{1}{n} \right)^2 \cdot \frac{d^2}{m} \right) = \Theta \left(\frac{m^5}{\Delta^2 d^4} \right) \tag{17}$$

Some of these $d + 3$ roots have positive derivatives, while others have negative derivatives: this will tell us which ones to use for our construction of U, and which ones for U'. Namely, for root r, we set (1) $\Pr(U = r) = 0$ and $\Pr(U' = r) \propto \frac{1}{p'(r)}$ if $p'(r) > 0$, and (2) $\Pr(U = r) \propto \frac{1}{p'(r)}$ and $\Pr(U' = r) = 0$ otherwise.

We now derive some bounds on the derivatives. Firstly, for sufficiently large choice of C (compared to c), notice that the weights on the Chebyshev polynomial's roots are overall bounded by

$$\sum_{m=1}^{d} \frac{1}{p'(r_{m+2})} = \Theta(1) \cdot \Delta^2 d^4 \sum_{m=1}^{d} \frac{1}{m^5} \ll nk, \tag{18}$$

since $\Delta^2 d^4 = kn \cdot (c^2/C)^2$ and the series $\sum_m \frac{1}{m^5}$ is convergent. Secondly, we claim that

$$T_d \left(1 - \Delta/n \right), T_d \left(1 - 2\Delta/n \right) \geq \cos(\pi/3) = \frac{1}{2}. \tag{19}$$

---

[7] We remark that the choice of the constant $1/10$ for error rate is arbitrary and our argument can easily be adapted to show a lower bound on the sample complexity under any constant error rate $\delta \in (0, 1/2)$.

This is because $\cos(\pi/3d) \leq 1 - \frac{2}{d^2}$; while $1 - \Delta/n \geq 1 - \frac{1}{C\log^2 n} \geq 1 - \frac{2}{d^2}$, the latter inequality again for a sufficiently large choice of C (with respect to c). The claim then follows by noticing that T_d is monotonically increasing in the region $[\pi/3d, 1]$ and $T_d(\cos(\pi/3d)) = \cos(\pi/3)$. We then proceed to verify each property claimed in the lemma.

1. By our construction, U is only supported on roots with negative derivatives. Hence, it can only takes values from $r_1 = \frac{1}{n}$ and some of the roots of the Chebyshev polynomials r_{2+m}. Moreover, by Equations (18), (19), the probabilities are bounded respectively by (before renormalization) $\Pr[U = \frac{1}{n}] \propto \Omega(n^2)$ and $\Pr[U \neq \frac{1}{n}] \propto \sum_{m=1}^{d} \mathbb{1}\{p'(r_{2+m}) < 0\}|p'(r_{m+2})| \ll nk$. It then follows that $\Pr[U \neq \frac{1}{n}] \ll \frac{nk}{nk+n^2} \leq \frac{k}{n}$.

2. Similarly, we have $\Pr[U' = 0], \Pr[U' = 2/n] \propto \Omega(n^2)$ and by (19) are within a factor 2 of each other. Combined with the fact that

$$\Pr[U' \notin \{0, 2/n\}] \propto \sum_{m=1}^{d} \mathbb{1}\{p'(r_{2+m}) > 0\}p'(r_{2+m}) \ll nk\,,$$

this immediately yields after renormalization that $\Pr[U' = 0], \Pr[U' = 2/n] \geq \frac{1}{3}$.

3. The largest values U, U' can take are the largest root of the Chebyshev polynomial, which is at most $\max_{m \in [d]} \frac{1}{\Delta}\left(1 - \cos\left(\frac{2m-1}{2d}\pi\right)\right) \leq \frac{2}{\Delta} = \frac{2C\log^2 n}{\sqrt{kn}}$.

4. First, as we used earlier, recall that by Taylor expansion of $\cos$

$$r_{m+2} = \frac{1}{\Delta}\left(1 - \cos\left(\frac{2m-1}{2d}\pi\right)\right) = \Theta\left(\frac{m^2}{\Delta d^2}\right)$$

Then, it holds

$$\mathbf{E}[U] = \Pr\left[U = \frac{1}{n}\right] \cdot \frac{1}{n} + \sum_{m=1}^{d} \Pr[U = r_{m+2}] \cdot r_{m+2}$$

$$\leq \frac{1}{n} + \Theta(1) \cdot \sum_{m=1}^{d} \frac{\Delta^2 d^4}{n^2 m^5} \cdot \frac{m^2}{\Delta d^2}$$

$$\leq \frac{1}{n} + \Theta\left(\frac{\Delta d^2}{n^2}\right) \cdot \sum_{m=1}^{d} \frac{1}{m^3} \leq \frac{1}{n}\left(1 + O(\sqrt{k/n})\right).$$

The argument for $\mathbf{E}[U'] = \frac{1}{n}(1 + O(\sqrt{k/n}))$ is similar.

5. The claim follows from Fact 5.

$\square$

B.2 Lower Bounds under Poisson Sampling

Proof of Proposition 2. Our goal is to argue H and H', specified in Equation (20) below, satisfy the following properties (i) H and H' are positive measures with total mass $1 + \Theta(\varepsilon)$ with probability at least $99/100$; (ii) H is a k-histogram with probability at least $99/100$ and H' is $\Omega(\varepsilon)$-far away from any k-histogram with probability at least $99/100$; and (iii) the distributions of the vectors $M = \{M_1, M_2, \cdots M_n\}$ and $M' = \{M'_1, M'_2, \cdots M'_n\}$, where $M_i \sim \mathrm{Poi}(m \cdot H_i)$ and $M'_i \sim \mathrm{Poi}(m \cdot H'_i)$, are $1/4$-close to each other in TV distance as long as $m = o(\sqrt{kn}/\log^2 n)$. If all these properties are satisfies, the result follows by applying Le Cam's Lemma. To do so, we define the measures H, H' as

$$H = \left(\frac{1}{n} + \varepsilon U^{(1)}, \cdots, \frac{1}{n} + \varepsilon U^{(n)}\right) \quad H' = \left(\frac{1}{n} + \varepsilon U'^{(1)}, \cdots, \frac{1}{n} + \varepsilon U'^{(n)}\right) \tag{20}$$

where $U^{(1)}, \cdots, U^{(n)}$ and $U'^{(1)}, \cdots U'^{(n)}$ are n i.i.d. copies of the random variables U, U' defined in Lemma 6.We proceed to verify that each property in our goal is satisfied.

(i) We first observe that the mass of H is simply $1 + \varepsilon \sum_{i=1}^{n} U^{(i)}$. As stated in Lemma 6, we have $\mathbf{E}\left[U\right] = \frac{1}{n}\left(1 + O(\sqrt{k/n})\right)$. Since $U^{(i)}$s are just i.i.d. copies of U, this further implies that $\mathbf{E}\left[\sum_{i=1}^{n} U^{(i)}\right] = 1 + O(\sqrt{k/n}) = O(1)$. Hence, by Markov's inequality, we have $\sum_{i=1}^{n} U^i = \Theta(1)$ with probability at least $99/100$. Similar arguments hold for H'. This shows claim (i).

(ii) Turning to (ii), recall that by construction of U we have $\Pr\left[U \neq \frac{1}{n}\right] \ll k/n$. Hence, with probability $99/100$, there are at most $(k-1)/2$ entries in H with mass other than $1/n$, which makes H a k-histogram. To argue the second part, i.e., that H' is far away from any k-histogram, we first lower bound the number of adjacent pairs $(i, i+1)$ such that $H_i = 0, H_{i+1} = \frac{2}{n}$. We call such an adjacent pair a "right border pair." For U', we have $\Pr\left[U' = 0\right], \Pr\left[U' = \frac{2}{n}\right] > 1/3$. Hence, in expectation, there are at least $\frac{1}{9}(n-1)$ such right border pairs. On the other hand, the variance of the number of right border pairs is at most n. Therefore, by Chebyshev's inequality (and assuming n large enough), with probability $99/100$ there will be at least $n/100$ right border pairs. This implies that H' is at least $\left(\frac{n}{100} - k\right) \cdot \frac{\varepsilon}{n} = \Omega(\varepsilon)$ far from a k-histogram. This concludes claim (ii).

(iii) Let P_M and $P_{M'}'$ be the distributions of the tuple of m samples seen by the algorithm. By subadditivity of total variation distance, we have

$$\mathrm{TV}(P_M, P_M') \leq n \cdot \mathrm{TV}\left(\mathbf{E}_U\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U\right)\right], \mathbf{E}_{U'}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U'\right)\right]\right). \quad (21)$$

To handle the right-hand-side, we will use the following lemma about distance between mixtures of Poisson distributions.

Lemma 8 ([WY$^+$19, Lemma 4]). *Let V, V' be random variables taking values in $[0, \Lambda]$. If $\mathbf{E}\left[V^j\right] = \mathbf{E}\left[V'^j\right]$ for $1 \leq j \leq L$, then*

$$\mathrm{TV}(\mathbf{E}\left[\mathrm{Poi}(V)\right], \mathbf{E}\left[\mathrm{Poi}(V')\right]) \leq \left(\frac{e\Lambda}{2L}\right)^L.$$

Now, let $\Lambda \coloneqq m\left(\frac{1}{n} + c'\varepsilon \frac{\log^2 n}{\sqrt{kn}}\right)$ and $L \coloneqq c\log n$, where $c, c' > 0$ are as in Lemma 6. It is straightforward to check from Lemma 6 that the random variables $V \coloneqq \frac{m}{n} + \varepsilon m U$, $V' \coloneqq \frac{m}{n} + \varepsilon m U'$ satisfy the assumptions of Lemma 8. This implies the existence of a (small) absolute constant $c'' > 0$ such that, if $m \leq c'' \min\left(n\log n, \frac{\sqrt{kn}}{\varepsilon \log n}\right)$, then by (21)

$$\mathrm{TV}(P_M, P_M') \leq n \cdot \left(\frac{e}{2c} \cdot m\left(\frac{1}{n\log n} + c'\frac{\varepsilon \log n}{\sqrt{kn}}\right)\right)^{c\log n} \leq n \cdot \frac{1}{n^2} = \frac{1}{n} < \frac{1}{4}. \quad (22)$$

Combining Equations (22) and (21) then finishes the proof of claim (iii).

Since H and H' satisfy all the properties listed, by Le Cam's Lemma, no algorithm can distinguish between the two distributions with probability more than $3/4 + 2/100 \leq 9/10$ when

$$m = o\left(\min\left(n\log n, \frac{\sqrt{kn}}{\varepsilon \log n}\right)\right)$$

under the Poissonization model. Notice that $\min\left(n\log n, \frac{\sqrt{kn}}{\varepsilon \log n}\right)$ is exactly $\frac{\sqrt{kn}}{\varepsilon \log n}$ under the assumption $\varepsilon \cdot \log^2 n \geq \sqrt{k/n}$. This concludes the proof. $\qquad\square$

Proof of the second part of Proposition 2. We already showed above, while establishing the first part of Proposition 2, that, with 99% probability, the total masses of H, H' are $1 + \Theta(\varepsilon)$, and that H is a k-Histogram Approximate Probability Vector while H' is $\Omega(\varepsilon)$ far away any k-Histogram Approximate Probability Vector. It thus only remains to show the indistinguishability. To do so, let as before P_M and $P_{M'}$ denote the distributions of the tuple of m samples seen by the algorithm: we

then need to establish that P_M and $P_{M'}$ are within total variation distance $1/4$ when $m = \tilde{o}(k/\varepsilon^2)$, for which is suffices to bound by $1/4$ the RHS below:

$$\mathrm{TV}\left(P_M, P_M'\right) \leq n \cdot \mathrm{TV}\left(\mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U\right)\right], \mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U'\right)\right]\right). \qquad (23)$$

We will now use a result from [Han19], restated below:

Theorem 6 (Theorem 4 from [Han19]). *For any $\Lambda > 0$ and random variables X, X' supported on $[-\Lambda, \infty)$, we have*

$$\mathrm{TV}(\mathbf{E}[\mathrm{Poi}(\Lambda + X)], \mathbf{E}[\mathrm{Poi}(\Lambda + X')]) \leq \frac{1}{2}\left(\sum_{\ell=0}^{\infty} \frac{|\mathbf{E}[X^\ell] - \mathbf{E}[X'^\ell]|^2}{\ell!\Lambda^\ell}\right)^{1/2}.$$

Recall that the first few moments of U, U' are identical i.e., $\mathbf{E}[U^t] = \mathbf{E}[U'^t]$ for $1 \leq t \leq L := c \cdot \log n$. Let $X := \varepsilon m U$, $X' := \varepsilon m U'$, and $\Lambda := \frac{m}{n}$. Notice that we indeed have $|X|, |X'| \leq \Lambda$ under the assumption $\varepsilon \ll \sqrt{k/n}/\log^2 n$ since

$$\max(|X|, |X'|) = \varepsilon \cdot m \cdot \max(|U|, |U'|) \leq \varepsilon \cdot m \cdot O\left(\frac{\log^2 n}{\sqrt{kn}}\right) \leq m/n = \Lambda.$$

Applying Theorem 6 then gives

$$4 \cdot \mathrm{TV}\left(\mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U\right)\right], \mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U'\right)\right]\right)^2$$

$$\leq \sum_{\ell=0}^{\infty} (\varepsilon m)^{2\ell} \frac{|\mathbf{E}[U^\ell] - \mathbf{E}[U'^\ell]|^2}{\ell!(m/n)^\ell}$$

$$= \sum_{\ell=L+1}^{\infty} \left(\varepsilon^2 mn\right)^\ell \frac{|\mathbf{E}[U^\ell] - \mathbf{E}[U'^\ell]|^2}{\ell!} \qquad \text{(the first } L \text{ moments match)}$$

$$\leq \sum_{\ell=L+1}^{\infty} \left(\varepsilon^2 mn\right)^\ell \frac{1}{\ell!} \left(\frac{c'\log^2 n}{\sqrt{kn}}\right)^{2\ell} \qquad (|U|, |U'| \leq \frac{c'\log^2 n}{\sqrt{nk}} \text{ (Lemma 6))}$$

$$= \sum_{\ell=L+1}^{\infty} \left(c'^2 \cdot \log^4 n \cdot \varepsilon^2/k \cdot m\right)^\ell \frac{1}{\ell!}.$$

Set for convenience $\kappa := c'^2 \cdot \log^4 n \cdot \varepsilon^2/k \cdot m$. Notice that $\kappa \ll L$ when $m = o\left(\frac{k}{\varepsilon^2 \log^3 n}\right)$. Denoting by Y a $\mathrm{Poi}(\kappa)$ random variable, this leads to

$$4\mathrm{TV}\left(\mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U\right)\right], \mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U'\right)\right]\right)^2 \leq \sum_{\ell=L+1}^{\infty} \frac{\kappa^\ell}{\ell!} = e^\kappa \Pr[Y \geq L+1]$$

$$\leq e^\kappa e^{-\frac{(L+1-\kappa)^2}{2(L+1)}} = e^{-\frac{1}{2}\left(L+1+\frac{\kappa^2}{L+1}\right)} \leq e^{-\frac{L}{2}},$$

where the second inequality is by standard Poisson concentration (see, e.g., the note [Can]) and $\mathbf{E}[Y] = \kappa \ll L$. This immediately gives

$$\mathrm{TV}\left(\mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U\right)\right], \mathbf{E}\left[\mathrm{Poi}\left(\frac{m}{n} + \varepsilon m U'\right)\right]\right) \leq \frac{1}{2}e^{-L/4} \qquad (24)$$

Combining Equations (23) and (24) then yields

$$\mathrm{TV}\left(P_N, P_N'\right) \leq n \cdot e^{-L/4} = n \cdot e^{-c\log n/4} < 1/4.$$

for sufficiently large c. By Le Cam's Lemma, no algorithm can distinguish between the two distributions as constructed in Equation (20) with probability more than $3/4 + 2/100 < 9/10$ when $m = o\left(\frac{k}{\varepsilon^2 \log^3 n}\right)$ under the Poissonization model. $\qquad \square$

B.3 Proof of Lower Bound in Theorem 1

Proof of Lower Bound Part of Theorem 1. By Proposition 2, it holds that $m_{\mathrm{hist}}^{\mathrm{poi}} \geq \Omega(k/(\varepsilon^2 \cdot \log^3 k) + \sqrt{kn}/(\varepsilon \cdot \log k))$. We proceed to argue for sample complexity lower bound under standard sampling. Suppose we are given a tester for fixed sample size such that it succeeds with high probability as long as it is given more than m^* samples. Assume that we want to use it to test whether an unknown measure P is a k-histogram under Poisson sampling model with $\tilde{m} \sim \mathrm{Poi}(m_{\mathrm{hist}}^{\mathrm{poi}}(n, \ell, \varepsilon) \|P\|_1)$ samples. We can construct an estimator which invokes the fixed sample size tester whenever $\tilde{m} \geq m^*$ and outputs fail otherwise.

By our lower bound result for the Poisson sampling model, the estimator fails with probability at least $1/10$. On the other hand, the estimator based on the fixed-sample tester succeeds with high probability whenever $\tilde{m} > m^*$. Together this implies that

$$m^* \geq (1 - \nu) \cdot m \cdot \left(1 - O\left((1 - \nu)^{-1} \cdot m^{-\frac{1}{2}}\right)\right),$$

where $m := m_{\mathrm{hist}}^{\mathrm{poi}}(n, \ell, \varepsilon, \nu)$. Since $\nu < 1$, it then holds

$$m^* > \Omega(1) \cdot m_{\mathrm{hist}}^{\mathrm{poi}}(n, \ell, \varepsilon, \nu) \geq \Omega(k/(\varepsilon^2 \cdot \log^3 k) + \sqrt{kn}/(\varepsilon \cdot \log k)).$$

Finally, we remark that the standard lower bound construction and analysis for uniformity testing can be shown to still apply to testing k-histograms (see [Can16, Proposition 4.1]), making formal the intuitive statement that testing whether a distribution is a k-histogram is at least as hard as testing whether a distribution is uniform. This shows that we also have $m^* \geq \Omega(\sqrt{n}/\varepsilon^2)$, and concludes the proof. $\square$

C Application to Model Selection

In this section, we detail how our testing algorithm, by a standard reduction, can be used for *model selection*, i.e., to select a suitable parameter k in order to succinctly represent the data.

Theorem 7. *There exists an algorithm which, given samples from an unknown distribution $\mathbf{p}$ on $[n]$, an error parameter $\varepsilon \in (0, 1]$, and a failure probability $\delta \in (0, 1]$, outputs a parameter $1 \leq K \leq n$ such that the following holds. Denote by $1 \leq k \leq n$ the smallest integer such that $\mathbf{p} \in \mathcal{H}_k^n$. With probability at least $1 - \delta$, (1) the algorithm takes $\widetilde{O}((\sqrt{n} + k)/\varepsilon^2 \cdot \log(1/\delta))$ samples, (2) $1 \leq K \leq 2k$, and (3) $\mathbf{p}$ is at TV distance at most ε from some $H \in \mathcal{H}_K^n$. Moreover, the algorithm runs in time $\mathrm{poly}(m)$, where m is the number of samples taken.*

Before proving the theorem, we discuss the various aspects of its statement. (1) guarantees that, with high probability, the algorithm does not take more samples than what it would if it were *given k* and just needed to test whether it was the right value. Moreover, as most tasks on k-histograms require at least a number of samples growing linearly in k, this guarantes that whenever $k \gg \sqrt{n}$ the cost of the model selection step is negligible in front of all subsequent tasks. Item (2), ensures that the output of the algorithm is a good approximation of the true, *optimal* value k; that is, that the model selected is essentially as succinct as it gets. Finally, (3) guarantees that even when the output is such that $K \ll k$, the parameter K is still good: that is, approximating $\mathbf{p}$ by a K-histogram still leads to a sufficiently accurate representation (even though it is even more succinct that what the *true* parameter k would yield).

Proof. The model selection algorithm is quite simple, and works by maintaining a current "guess" K as part of a doubling search, and using the tester of to check if the current value is good. Specifically: for $0 \leq j \leq \lceil \log_2 n \rceil$, run the testing algorithm of Theorem 1 (**Divide-And-Learn-And-Sieve**) with parameters $n, \varepsilon, 2^j$, and $\delta_j := \frac{\delta}{2(j+1)^2}$. If the testing algorithm returns Accept, then return $K := 2^j$ and stop; otherwise, continue the loop.

Let $L \leq \lceil \log_2 n \rceil$ be the number of iterations before the algorithm stops. By a union bound, the probability that all L invocations of the testing algorithm behaved as they should is, by a union bound, at least

$$1 - \sum_{j=0}^{L} \delta_j = 1 - \sum_{j=0}^{L} \frac{\delta}{2(j+1)^2} \geq 1 - \sum_{j=0}^{\infty} \frac{\delta}{2(j+1)^2} \geq 1 - \delta.$$

Condition on this being the case. Since the algorithm, as soon as $j \geq \lceil \log_2 k \rceil$, returns Accept (since then $\mathbf{p} \in \mathcal{H}_{2^j}^n$), we get that $L := \lceil \log_2 k \rceil$, and that the output satisfies $K \leq 2k$, giving (2). Moreover, if $K < k$, then by the guarantee of the testing algorithm this means that $\mathbf{p}$ was *not* ε-far from $\mathcal{H}_K^n$, showing (3). Finally, the sample complexity is then

$$\sum_{j=0}^{L} \tilde{O}\left(\frac{\sqrt{n}}{\varepsilon^2} \log \frac{1}{\delta_j} + \frac{2^j}{\varepsilon^2} \log \frac{1}{\delta_j} \right) = \tilde{O}\left(\frac{\sqrt{n}}{\varepsilon^2} + \frac{k}{\varepsilon^2} \right) \log \frac{1}{\delta}$$

since $L = O(\log k)$. This shows (1), and concludes the proof. $\qquad\square$

As a final remark, we note that the guarantee provided in (2) can be improved to the optimal $1 \leq K \leq k$, by modifying slightly the above procedure. Namely, after finding some K such that $1 \leq K \leq 2k$ as before, one can run the testing algorithm for $K/2 \leq i \leq K$ (not a binary search anymore), each time with parameters n, ε, i, and $\delta_i = \delta/K$. By a union bound, this incurs an extra δ probability of failure, and an additional $\tilde{O}\left((\sqrt{n} + k)/\varepsilon^2 \cdot \log(1/\delta) \right)$ samples overall, but now the output after this second step will be guaranteed to be at most k (with high probability).