EXPONENTIAL GENERALIZATION BOUNDS WITH NEAR-OPTIMAL RATES FOR L_q-STABLE ALGORITHMS

ABSTRACT

The *stability* of learning algorithms to changes in the training sample has been actively studied as a powerful proxy for reasoning about generalization. Recently, exponential tail generalization and risk bounds with near-optimal rates have been obtained under the stringent and distribution-free notion of uniform stability (Bousquet et al., 2020; Klochkov & Zhivotovskiy, 2021). In the meanwhile, under the notion of L_q-stability, which is weaker and distribution dependent, exponential generalization bounds are also available yet so far only with sub-optimal rates. Therefore, a fundamental question we would like to address in this paper is whether it is possible to derive near-optimal exponential generalization bounds for L_q-stable learning algorithms. As the core contribution of the present work, we give an affirmative answer to this question by developing strict analogues of the near-optimal generalization and risk bounds of uniformly stable algorithms for L_q-stable algorithms. We demonstrate the power of our improved bounds by applying them to derive strong sparse excess risk bounds, under mild conditions, for computationally tractable sparsity estimation algorithms such as Iterative Hard Thresholding (IHT).

1 INTRODUCTION

A fundamental issue in statistical learning is to bound the generalization error of a learning algorithm for understanding its prediction performance on unseen data. It has long been recognized in literature that one of the key characteristics that permits learning algorithms to generalize is the *stability* of estimated model to perturbations in training data. The idea of using algorithmic stability as a proxy for generalization performance analysis dates back to the seventies (Rogers & Wagner, 1978; Devroye & Wagner, 1979). Since the seminal work of Bousquet & Elisseeff (2002), the search for generalization bounds under various notions of algorithmic stability has been a flourishing area of learning theory (Zhang, 2003; Mukherjee et al., 2006; Shalev-Shwartz et al., 2010; Kale et al., 2011; Hardt et al., 2016; Celisse & Guedj, 2016; Bousquet et al., 2020).

As one may expect, the stronger an algorithmic stability criterion is, the sharper the resulting generalization bound will be. On one end, exponential generalization bounds can be guaranteed by approaches under the most stringent notion of uniform stability (Bousquet & Elisseeff, 2002; Bousquet et al., 2020), which requires the change in the prediction loss to be uniformly small regardless data distribution. Despite the strength of generalization, the distribution-free nature makes uniform stability too restrictive to be fulfilled, e.g., by learning rules with unbounded losses (Celisse & Guedj, 2016). On the other end, based on weaker and distribution dependent notions of stability such as hypothesis stability and mean-square stability, only polynomial generalization bounds seem possible in general cases, although the corresponding stability criteria are more amenable to verification (Bousquet & Elisseeff, 2002). These observations have prompted the development of L_q-stability, as an in-between state, to achieve the best of two worlds (Celisse & Guedj, 2016; Abou-Moustafa & Szepesvári, 2019): it generalizes the notion of hypothesis stability from ℓ_1-norm criterion to L_q-norm criterion for $q \geq 2$ but remains distribution dependent and thus weaker than uniform stability; in the meanwhile it can still achieve the appealing exponential tail bounds of generalization matching those from Bousquet & Elisseeff (2002) for uniformly stable algorithms.

By far, the best known (and near-optimal) rates about exponential generalization bounds are offered by approaches based on uniform stability and certain fine-grained concentration inequalities for sum

of functions of independent random variables (Feldman & Vondrak, 2019; Bousquet et al., 2020). These rates are substantially sharper than those of Bousquet & Elisseeff (2002), which are implied by a naive use of McDiarmid's inequality, in terms of the overhead factors on stability coefficients. While for L_q-stable learning algorithms, the state-of-the-art exponential generalization bounds are derived based on the moments or exponential extensions of Efron-Stein inequality (Celisse & Guedj, 2016; Abou-Moustafa & Szepesvári, 2019), which yield similar rates of convergence to those of Bousquet & Elisseeff (2002) and thus are suspected to be *sub-optimal*.

Given the above observed gap in rates of convergence between the generalization bounds under uniform stability and L_q-stability, the following question is naturally raised:

> *Is it possible to derive sharper exponential generalization bounds for L_q-stable learning algorithms that match those recent breakthrough results for uniformly stable algorithms?*

As the core contribution of the present work, we give an affirmative answer to this open question by developing strict analogues of the near-optimal generalization bounds of uniformly stable algorithms for L_q-stable algorithms. To illustrate the power of our results, we have applied the improved bounds to derive sharper exponential risk bounds for computationally tractable sparsity recovery estimators, such as the Iterative Hard Thresholding (IHT) algorithms widely used in high dimensional sparse learning (Blumensath & Davies, 2009; Foucart, 2011; Jain et al., 2014).

Notation. Here we provide some notation that will be frequently used throughout the paper. Let $S = \{Z_1, Z_2, ..., Z_N\}$ be a set of independent random data samples valued in some measurable set $\mathcal{Z}$. For any indices set $I \subseteq [N] := \{1, ..., N\}$, we denote by $S_I = \{Z_i, i \in I\}$ and $S_{\bar{I}} = S \setminus S_I$. We denote by $S' = \{Z'_1, Z'_2, ..., Z'_N\}$ another i.i.d. sample from the same distribution as that of S and we write $S^{(i)} = \{Z_1, ..., Z_{i-1}, Z'_i, Z_{i+1}, ..., Z_N\}$. For a real-valued random variable Y, its L_q-norm for $q \geq 1$ is defined by $\|Y\|_q = (\mathbb{E}[|Y|^q])^{1/q}$. By definition it can be verified that $\forall q \geq 2$,

$$\|Y\|_q^2 = (\mathbb{E}[|Y|^q])^{2/q} = \left(\mathbb{E}[|Y^2|^{q/2}]\right)^{2/q} = \left\|Y^2\right\|_{q/2}. \tag{1}$$

Let $g : \mathcal{Z}^N \mapsto \mathbb{R}$ be some measurable function and consider the random variable $g(S) = g(Z_1, ...Z_N)$. For $g(S)$ and any index set $I \subseteq [N]$, we define the following abbreviations:

$$g(S_I) := \mathbb{E}[g(S) \mid S_I], \quad \|g\|_q(S_I) := (\mathbb{E}[|g(S)|^q \mid S_I])^{1/q}.$$

We say a real-valued function f is G-Lipschitz continuous over the domain $\mathcal{W}$ if

$$|f(w) - f(w')| \leq G\|w - w'\|, \quad \forall w, w' \in \mathcal{W}.$$

1.1 SETUP AND PRIOR RESULTS

Problem setup. We consider a statistical learning algorithm $A : \mathcal{Z}^N \mapsto \mathcal{W}$ that maps a training data set S to a model $A(S)$ in a closed subset $\mathcal{W}$ of an Euclidean space. The population risk and corresponding empirical risk evaluated at $A(S)$ are respectively given by

$$R(A(S)) := \mathbb{E}_Z[\ell(A(S); Z)] \text{ and } R_S(A(S)) := \frac{1}{N} \sum_{i=1}^{N} \ell(A(S); Z_i),$$

where $\ell : \mathcal{W} \times \mathcal{Z} \mapsto \mathbb{R}^+$ is a non-negative and potentially unbounded loss function whose value $\ell(w; z)$ measures the loss evaluated at z with parameter w. As a classic fundamental issue in statistical learning, we are interested in deriving the upper bounds on the difference between population and empirical risks, i.e., $|R(A(S)) - R_S(A(S))|$, which quantifies the generalization error of A.

We first introduce the following concept of uniform stability (Bousquet & Elisseeff, 2002) which requires the change in the prediction loss to be uniformly small regardless data distribution.

Definition 1 (Uniform stability). *A learning algorithm A is said to have uniform stability by $\gamma_u > 0$ if it satisfies the following uniform bound:*

$$\sup_{S, S^{(i)}, Z \in \mathcal{Z}} |\ell(A(S); Z) - \ell(A(S^{(i)}); Z)| \leq \gamma_u, \quad \forall i \in [N].$$

Bousquet & Elisseeff (2002) showed that a large class of regularized empirical risk minimization (ERM) algorithms has uniform stability, and using McDiarmids inequality yields the following exponential tail generalization bound that holds with probability at least $1-\delta$ over S for any $\delta \in (0, 1)$:

$$|R(A(S)) - R_S(A(S))| \lesssim \gamma_u \sqrt{N \log\left(\frac{1}{\delta}\right)} + \sqrt{\frac{\log(1/\delta)}{N}}. \tag{2}$$

Recently, equipped with a strong concentration inequality for sum of random functions, Bousquet et al. (2020) established the following moments bound of uniformly stable algorithms for all $q \geq 2$:

$$\|R(A(S)) - R_S(A(S))\|_q \lesssim q\gamma_u \log(N) + \sqrt{\frac{q}{N}}. \tag{3}$$

In view of the equivalence between tails and moments (see, e.g., Bousquet et al., 2020, Lemma 1), the above L_q-norm bound immediately implies the following tail bound that holds with probability at least $1 - \delta$ over S for any $\delta \in (0, 1)$:

$$|R(A(S)) - R_S(A(S))| \lesssim \gamma_u \log(N) \log\left(\frac{1}{\delta}\right) + \sqrt{\frac{\log(1/\delta)}{N}}. \tag{4}$$

This bound substantially improves the classic result in equation 2 by reducing the overhead factor on stability coefficient from $\mathcal{O}\left(\sqrt{N \log(\frac{1}{\delta})}\right)$ to $\mathcal{O}(\log(N) \log(\frac{1}{\delta}))$. For example, in regimes such as regularized ERM where $\gamma_u \lesssim \frac{1}{\sqrt{N}}$ is usually the case, the convergence rate in bound equation 2 becomes vacuous as it is not vanishing in sample size, while the bound in equation 4 still guarantees $\mathcal{O}\left(\frac{\log(N) \log(\frac{1}{\delta})}{\sqrt{N}}\right)$ rate of convergence. Indeed, up to logarithmic factors on sample size and tail bounds, the bound in equation 4 is nearly optimal in the sense of a lower bound on sum of random functions by Bousquet et al. (2020). Under the generalized Bernstein condition (Koltchinskii, 2006) and using the sharp concentration inequality for sum of random functions by Bousquet et al. (2020), Klochkov & Zhivotovskiy (2021) alternatively established the following deviation optimal excess risk bound that holds with probability at least $1 - \delta$ over S:

$$R(A(S)) - R^* \lesssim \Delta_{\text{opt}} + \mathbb{E}[\Delta_{\text{opt}}] + \gamma_u \log(N) \log\left(\frac{1}{\delta}\right) + \frac{\log(1/\delta)}{N}, \tag{5}$$

where R^* and Δ_{opt} are defined shortly by equation 8. This kind of bounds are of particular interest for understanding the population risk minimization performance of a learning algorithm.

While implying strong generalization guarantees, the uniform stability is also most stringent in the sense that it is distribution independent and hard to be fulfilled, e.g., by learning rules with unbounded losses. To address such an unpleasant restrictiveness, the notion of L_q-stability was alternatively introduced by Celisse & Guedj (2016) as a relaxation of uniform stability.

Definition 2 (L_q-Stability). *For $q \geq 1$, a learning algorithm A is said to have L_q-stability by coefficient $\gamma_q > 0$ if it satisfies the following moment bound:*

$$\left\|\ell(A(S); Z) - \ell(A(S^{(i)}); Z)\right\|_q \leq \gamma_q, \quad \forall i \in [N].$$

In the above definition, the expectation associated with L_q-norm is taken over $S, S^{(i)}, Z$, and the internal random bits of A, if any (such as in the case of stochastic optimization algorithms). Note that slightly different from that of Celisse & Guedj (2016), the random variable Z in the above definition is not necessarily required to be independent of S and $S^{(i)}$. By definition, L_q-stability is distribution dependent and thus is weaker than uniform stability which can be regarded as a special case of L_q-stability with $\gamma_q \equiv \gamma_u$ for all $q \geq 1$. Particularly for $q = 1$ and $q = 2$, the L_q-stability reduces to the notions of hypothesis stability (Bousquet & Elisseeff, 2002) and mean-square stability (Kale et al., 2011), respectively. For an instance, it has been shown that the classical ridge regression model with unbounded responses has L_q-stability for all $q \geq 1$ rather than uniform stability (Celisse & Guedj, 2016). As a novel and concrete example, we will see shortly in Section 3 that L_q-stability plays a crucial role for deriving strong sparse excess risk bounds for sparsity estimation algorithms such as IHT (Jain et al., 2014; Yuan et al., 2018). Alternatively, the definition of L_q-stability can be extended

to the L_q-argument-stability as $\left\| A(S) - A(S^{(i)}) \right\|_q \leq \gamma_q$, which generalizes the concept of uniform argument stability (Bassily et al., 2020) to the L_q-norm criterion. Obviously L_q-argument-stability is not at all relying on the random argument Z and it implies L_q-stability for Lipschitz losses.

The following is by far the best known moments generalization bound under L_q-stability that holds for all $q \geq 2$ (Celisse & Guedj, 2016; Abou-Moustafa & Szepesvári, 2019):

$$\|R(A(S)) - R_S(A(S))\|_q \lesssim \gamma_q \sqrt{Nq} + \sqrt{\frac{q}{N}}. \tag{6}$$

As one can see that the L_q-stability generalization bound equation 6 is significantly inferior to the near-optimal uniform stability generalization bound equation 3 in terms of the overhead on stability coefficient. Such a gap in rate of convergence is indeed unsurprising: the bound equation 6 was derived via more or less directly applying moments or exponential extensions of Efron-Stein inequality to generalization error (Celisse & Guedj, 2016; Abou-Moustafa & Szepesvári, 2019), and thus yields about the same overhead factor on stability coefficient as that of the sub-optimal exponential bound in equation 2 for uniform stable algorithms. In light of these observations, we are naturally motivated to derive sharper exponential generalization bounds for L_q-stable algorithms hopefully to match the near-optimal bound equation 3 achievable by uniformly stable algorithms.

1.2 OUR CONTRIBUTION

The core contribution of the present work is a set of substantially improved exponential generalization bounds for L_q-stable algorithms. The key ingredient of our analysis is a sharper concentration bound on sum of functions of independent random variables under the L_q-norm bounded difference conditions, which generalizes a previous counterpart under the uniform bounded difference conditions (Bousquet et al., 2020). With this generic concentration bound in hand, we are able to derive sharper generalization and excess risk bounds for L_q-stable learning algorithms that match those best known for uniform stable algorithms. The power of our results is demonstrated through deriving stronger exponential sparse excess risk bounds for computationally tractable sparsity estimation algorithms (such as IHT). Below we sketch the main results obtained in this work:

- In Section 2, we first establish in Theorem 1 an L_q-norm inequality for sum of functions of random variables with L_q-norm bounded difference. Then equipped with such a general-purpose concentration inequality, we prove in Theorem 2 the following L_q-norm generalization bound for L_q-stable learning algorithms for all $q \geq 2$:

$$\|R(A(S)) - R_S(A(S))\|_q \lesssim q\gamma_q \log N + \sqrt{\frac{q}{N}}.$$

  Compared to equation 6, the preceding bound improves the overhead factor on γ_q from $\sqrt{N}$ to $\log(N)$. As another consequence of our L_q-norm concentration inequality, we further derive in Theorem 3 the following excess risk bound for L_q-stable algorithms under Bernstein or quadratic-growth conditions:

$$\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \lesssim \mathbb{E}[\Delta_{\text{opt}}] + q\gamma_q \log(N) + \frac{q}{N}.$$

  Based on the equivalence between moments and tails, this above result implies the deviation optimal risk bound in equation 5 for uniformly stable algorithms.

- In Section 3, based on our L_q-stability generalization theory, we show in Theorem 4 a novel exponential sparse excess risk bound for inexact L_0-estimators. A key insight here is that L_0-estimators are in many cases "almost always" stable over any fixed supporting set, and thus can be shown to have L_q-stability over the same supporting set, which consequently makes the developed generalization theory applicable.

2 SHARPER EXPONENTIAL BOUNDS FOR L_q-STABLE ALGORITHMS

2.1 A MOMENT INEQUALITY FOR SUM OF RANDOM FUNCTIONS

We start by presenting in the following theorem a moment inequality for sum of random functions of N independent random variables that satisfy the L_q-norm bounded difference condition. See Appendix B.1 for a proof of this result.

Theorem 1. *Let $S = \{Z_1, Z_2, ..., Z_N\}$ be a set of independent random variables valued in $\mathcal{Z}$. Let $g_1, ..., g_N$ be a set of measurable functions $g_i : \mathcal{Z}^N \mapsto \mathbb{R}$ that satisfy the following conditions for any $i \in [N]$:*

- $\mathbb{E}[g_i(S) \mid S \setminus Z_i] = 0$ *almost surely;*

- $g_i(S)$ *has the following L_q-norm bounded difference property with respect to all variables in S except Z_i, i.e., $\forall j \neq i$, for all $q \geq 1$:*

$$\left\| g_i(S) - g_i(S^{(j)}) \right\|_q \leq \beta_q.$$

Then there exists a universal constant $\kappa < 1.271$ such that for all $q \geq 2$,

$$\left\| \sum_{i=1}^{N} g_i(S) - \mathbb{E}[g_i(S) \mid Z_i] \right\|_q \leq 4\kappa q N \lceil \log_2 N \rceil \beta_q.$$

Additionally, if $\|\mathbb{E}[g_i(S) \mid Z_i]\|_q \leq M_q$, then for all $q \geq 2$

$$\left\| \sum_{i=1}^{N} g_i(S) \right\|_q \leq 2\sqrt{2\kappa N q} M_q + 4\kappa q N \lceil \log_2 N \rceil \beta_q.$$

Remark 1. *Theorem 1 extends the moment inequality of Bousquet et al. (2020, Theorem 4) from under the distribution-free uniform bounded difference property to under the L_q-norm bounded difference property which is distribution dependent. Specially if $g_i(S)$ have uniform bounded difference property, then Theorem 1 reduces to the result of Bousquet et al. (2020, Theorem 4).*

Remark 2. *The L_q-norm boundedness condition $\|\mathbb{E}[g_i(S) \mid Z_i]\|_q \leq M_q$ in our theorem allows g_i to be potentially unbounded over domain of interest, which is weaker than the corresponding almost sure boundedness condition on $|\mathbb{E}[g_i(S) \mid Z_i]|$ as imposed by Bousquet et al. (2020, Theorem 4).*

2.2 GENERALIZATION BOUNDS FOR L_q-STABLE ALGORITHMS

As a consequence of Theorem 1, we can show the following main result on the generalization bound of L_q-stable learning algorithms. See Appendix B.2 for a proof of this result.

Theorem 2. *Let $A : \mathcal{Z}^N \mapsto \mathcal{W}$ be a learning algorithm that has L_q-stability by $\gamma_q > 0$ for $q \geq 1$. Suppose that $\|\ell(A(S); Z)\|_q \leq M_q$ for any $Z \in \mathcal{Z}$. Then for all $q \geq 2$,*

$$\|R(A(S)) - R_S(A(S))\|_q \lesssim q\gamma_q \log N + \sqrt{\frac{q}{N}} M_q.$$

Remark 3. *The L_q-norm boundedness condition $\|\ell(A(S); Z)\|_q \leq M_q$ allows for learning with unbounded losses over, e.g., data distribution with sub-Gaussian or sub-exponential tail bounds.*

To compare with the best known moments generalization bound equation 6 under the notion of L_q-stability, our bound in Theorem 2 substantially proves the overhead factor on γ_q from $\sqrt{N}$ to $\log(N)$. Specially when reduced to regime of uniform stability where $\gamma_q \equiv \gamma_u$ for all $q \geq 1$, our result revisits the moments generalization bound in equation 3 which is nearly tight, up to logarithmic factors on sample size, in the sense of a lower bound on sum of random functions from Bousquet et al. (2020). More broadly, suppose that the following exponential stability bound holds for any $\delta \in (0, 1)$ with a mixture of sub-Gaussian and sub-exponential tails [1] over $S, S^{(i)}, Z$:

$$\left| \ell(A(S); Z) - \ell(A(S^{(i)}); Z) \right| \leq a \log\left(\frac{e}{\delta}\right) + b\sqrt{\log\left(\frac{e}{\delta}\right)}. \tag{7}$$

Then according to the equivalence of tails and moments, as summarized in Lemma 4 (see Appendix A), we must have that A is L_q-stable by $\gamma_q = aq + b\sqrt{q}$. Assume that the loss is bounded

---

[1] In an exponential tail bound, the terms associated with $\sqrt{\log\left(\frac{1}{\delta}\right)}$ and $\log\left(\frac{1}{\delta}\right)$ are respectively referred to as sub-Gaussian and sub-exponential tails.

almost surely over data. Then the L_q-norm generalization bound in Theorem 2 combined with Lemma 4 immediately implies the following generalization bound:

$$|R(A(S)) - R_S(A(S))| \lesssim a \log(N) \log^2\left(\frac{1}{\delta}\right) + b \log(N) \log^{1.5}\left(\frac{1}{\delta}\right) + \sqrt{\frac{\log(1/\delta)}{N}}.$$

Compared with the uniform stability implied tail bound equation 4, the preceding L_q-stability bound is nearly identical up to slightly worse confidence tail terms which are caused by the uncertainty of L_q-stability with respect to data distribution. We conjecture that such a slight deterioration in tail bounds might possibly be remedied by using the exponential versions of Efron-Stein inequality (Boucheron et al., 2003; Abou-Moustafa & Szepesvári, 2019) instead of its currently used moments variant. We leave the improvement over poly-logarithmic terms for future investigation.

2.3 EXCESS RISK BOUNDS FOR L_q-STABLE ALGORITHMS

In addition to the generalization bounds, we further apply Theorem 1 to study the excess risk bounds of an L_q-stable learning algorithm which are of particular interest for understanding its population risk minimization performance. In the following analysis, the global minimum of the population risk and the empirical risk sub-optimality of the algorithm on training data are respectively denoted by

$$R^* := \min_{w \in \mathcal{W}} R(w), \quad \Delta_{\mathrm{opt}} := R_S(A(S)) - \min_{w \in \mathcal{W}} R_S(w). \tag{8}$$

Let us define $W^* := \mathrm{Argmin}_{w \in \mathcal{W}} R(w)$ as the optimal solution set of the population risk. In order to get sharper risk bounds, we need to impose some structural conditions on risk functions. Particularly, the following defined generalized Bernstein condition (Koltchinskii, 2006) is conventionally used with multiple global risk minimizers allowed.

Assumption 1 (Generalized Bernstein condition). *For some $B > 0$ and for any $w \in \mathcal{W}$, there exists $w^* \in W^*$ such that the following holds:*

$$\mathbb{E}\left[(\ell(w; Z) - \ell(w^*; Z))^2\right] \leq B(R(w) - R(w^*)).$$

We will also consider the quadratic growth condition which is widely used as an alternative condition for establishing fast rates of convergence in learning theory.

Assumption 2 (Quadratic growth condition). *For some $\mu > 0$ and for any $w \in \mathcal{W}$, there exists $w^* \in W^*$ such that the following holds:*

$$R(w) \geq R^* + \frac{\mu}{2}\|w - w^*\|^2.$$

Remark 4. *Clearly, when the loss is G-Lipschitz, the quadratic growth condition with parameter μ implies the Bernstein condition with parameter $B = \frac{2G^2}{\mu}$.*

The following is our main result on the excess risk bound of L_q-stable algorithms. A proof of this theorem can be found in Appendix B.3.

Theorem 3. *Let $A : \mathcal{Z}^N \mapsto \mathcal{W}$ be a learning algorithm that has L_q-stability by γ_q for $q \geq 1$.*

(a) If Assumption 1 holds and $\ell(\cdot; \cdot) \leq M$, then $\forall q \geq 2$,

$$\|R(A(S)) - R^* - \Delta_{opt}\|_q \lesssim \mathbb{E}[\Delta_{opt}] + q\gamma_q \log(N) + \frac{(M + B)q}{N}.$$

(b) If Assumption 2 holds and $\ell(\cdot; \cdot)$ is G-Lipschitz with respect to its first argument, then $\forall q \geq 2$,

$$\|R(A(S)) - R^* - \Delta_{opt}\|_q \lesssim \mathbb{E}[\Delta_{opt}] + q\gamma_q \log(N) + \frac{G^2 q}{\mu N}.$$

Remark 5. *Theorem 3 extends the near-optimal exponential risk bounds of Klochkov & Zhivotovskiy (2021) from uniform stable algorithms to L_q-stable algorithms. Suppose that A satisfies the exponential stability bound in equation 7, and thus A has L_q-stability by $\gamma_q = aq + b\sqrt{q}$. Then combined with Lemma 4, the L_q-norm risk bounds in Theorem 3 suggest that the following exponential tail bound holds:*

$$R(A(S)) - R^* \lesssim \Delta_{opt} + \mathbb{E}[\Delta_{opt}] + a \log(N) \log^2\left(\frac{1}{\delta}\right) + b \log(N) \log^{1.5}\left(\frac{1}{\delta}\right) + \frac{\log(1/\delta)}{N}.$$

Algorithm 1: `Inexact` L_0`-ERM Oracle`

Input : A training data set $S = \{Z_i\}_{i \in [N]}$ and the desired sparsity level k.
Output: $\tilde{w}_{S,k}$.
Compute an inexact k-sparse L_0-ERM estimation $\tilde{w}_{S,k}$ such that

- $\tilde{w}_{S,k} = w^*_{S|\tilde{J}} := \arg\min_{w \in \mathcal{W}, \mathrm{supp}(w) \subseteq \tilde{J}} R_S(w)$, where $\tilde{J} = \mathrm{supp}(\tilde{w}_{S,k})$;

- $\tilde{w}_{S,k}$ attains some $\bar{k}$-sparse sub-optimality $\Delta_{\bar{k},\mathrm{opt}} \geq 0$ for some $\bar{k} \leq k$, i.e.,
  $R_S(\tilde{w}_{S,k}) - R_S(w^*_{S,\bar{k}}) \leq \Delta_{\bar{k},\mathrm{opt}}$.

Remark 6. *We comment on the conditions required in the two parts of Theorem 3 for fast optimization. On one hand, the generalized Bernstein condition in part (a) is weaker than the quadratic growth condition in part (b). On the other hand, part (a) requires the loss to be uniformly bounded while part (b) only assumes the loss to be Lipschitz continuous.*

3 Application to Inexact L_0-ERM

In this section, we demonstrate an application of our L_q-stability and generalization theory to the following problem of high-dimensional stochastic risk minimization under hard sparsity constraint:

$$\min_{w \in \mathcal{W}} R(w) := \mathbb{E}_Z[\ell(w; Z)] \quad \text{subject to } \|w\|_0 \leq k,$$

where $\mathcal{W} \subseteq \mathbb{R}^d$ the cardinality constraint $\|w\|_0 \leq k$ for $k \ll d$ is imposed for enhancing the interpretability and learnability of model in cases where there are no clear favourite explanatory variables, or the model is over-parameterized. We consider the following L_0-ERM problem over training set $S = \{Z_i\}_{i \in [N]}$:

$$w^*_{S,k} = \arg\min_{\|w\|_0 \leq k} \left\{ R_S(w) := \frac{1}{N} \sum_{i=1}^{N} \ell(w; Z_i) \right\}. \tag{9}$$

Since the problem is known to be NP-hard (Natarajan, 1995), it is computationally intractable to solve it exactly in general cases. Alternatively, we consider the inexact L_0-ERM oracle as a meta-algorithm outlined in Algorithm 1. In order to avoid assuming unrealistic conditions like restricted isometry property (RIP), it is typically needed to allow sparsity relaxation for polynomial algorithms like IHT and HTP to achieve favorable converge behavior (Jain et al., 2014; Shen & Li, 2016; Yuan et al., 2018; Zhou et al., 2018). Therefore, we are particularly interested in the inexact L_0-ERM oracle with $\bar{k}$-sparse sub-optimality $\Delta_{\bar{k},\mathrm{opt}} \geq 0$ for some $\bar{k} \leq k$ such that

$$R_S(\tilde{w}_{S,k}) - R_S(w^*_{S,\bar{k}}) \leq \Delta_{\bar{k},\mathrm{opt}}.$$

It is typical that $\Delta_{\bar{k},\mathrm{opt}}$ is a random value over the training set S. For example, the sub-optimality guarantees of IHT for empirical risk usually hold with high probability over training data (Jain et al., 2014). Let $w^*_{\bar{k}} := \arg\min_{\|w\|_0 \leq \bar{k}} R(w)$ be the $\bar{k}$-sparse minimizer of population risk for some $\bar{k} \leq k$. We are interested in deriving exponential upper bounds for the $\bar{k}$-sparse excess risk given by $R(\tilde{w}_{S,k}) - R(w^*_{\bar{k}})$.

Our analysis also relies on the conditions of Restricted Strong Convexity (RSC) which extends the concept of strong convexity to the analysis of sparsity recovery methods (Bahmani et al., 2013; Blumensath & Davies, 2009; Jain et al., 2014; Yuan et al., 2020).

Definition 3 (Restricted Strong Convexity). *For any sparsity level $1 \leq s \leq d$, we say a function f is restricted μ_s-strongly convex if there exists $\mu_s > 0$ such that*

$$f(w) - f(w') - \langle \nabla f(w'), w - w' \rangle \geq \frac{\mu_s}{2} \|w - w'\|^2, \quad \forall \|w - w'\|_0 \leq s.$$

Particularly, we say f is μ-strongly convex if f is μ_d-strongly convex.

The following basic assumptions will be made in our theoretical analysis.

Assumption 3. *The loss function $\ell(\cdot; \cdot)$ is convex and G-Lipschitz with respect to its first argument.*

Assumption 4. *The population risk R is μ-strongly convex and the empirical risk R_S is μ_k-strongly convex with probability at least $1 - \delta_N$ over sample S for some $\delta_N \in (0, 1)$.*

Assumption 5. *The domain of interest is uniformly bounded such that $\|w\| \le D, \forall w \in \mathcal{W}$.*

Let $w^* := \arg\min_{w \in \mathcal{W}} R(w)$ be the global minimizer which is unique due to the strong convexity of R. For a given index set $J \subseteq [d]$, let us consider the following restrictive estimator over J:

$$w^*_{S|J} := \arg\min_{w \in \mathcal{W}, \text{supp}(w) \subseteq J} R_S(w). \tag{10}$$

We first present the following lemma that guarantees the L_q-stability of $w^*_{S|J}$ for any fixed J with $|J| = k$. See Appendix C.1 for its proof.

Lemma 1. *Asssume that Assumptions 3, 4 and 5 hold and $\frac{\log(1/\delta_N)}{\log(N)} \ge 2$. Let $J \subseteq [d], |J| = k$ be a set of indices of cardinality k. Then for any $q \ge 2$, the oracle estimator $w^*_{S|J}$ has L_q-stability by*

$$\gamma_q = \frac{1}{N}\left(\frac{4G^2}{\mu_k} + 2GD\right) + \frac{2GD\log(N)q}{\log(1/\delta_N)}.$$

Remark 7. *For sparse linear regression models, it can be verified based on the result by Agarwal et al. (2012, Lemma 6) that Assumptions 4 holds with $\delta_N = e^{-c_0 N}$ for some universal positive constant c_0. Then we have $\frac{\log(1/\delta_N)}{\log(N)} = \frac{c_0 N}{\log(N)} \ge 2$ for sufficiently large N, and Lemma 1 implies that for all $q \ge 2$,*

$$\gamma_q \lesssim \frac{1}{N}\left(\frac{G^2}{\mu_k} + \frac{GD\log(N)q}{c_0}\right).$$

The following theorem is our main result on the sparse excess risk of the inexact L_0-ERM oracle as defined in Algorithm 1. See Appendix C.2 for its proof.

Theorem 4. *Suppose that Assumptions 3, 4, 5 hold. Assume that $\frac{\log(1/\delta_N)}{\log(N)} \ge 2$. Then for any $\delta \in (0, e^{-1})$, the following $\bar{k}$-sparse excess risk bound holds with probability at least $1 - \delta$ over S:*

$$R(\tilde{w}_{S,k}) - R(w^*_{\bar{k}})$$

$$\lesssim \frac{GD\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)^2 \log^2(N)}{\log(1/\delta_N)} + \left(\log(N)\left(\frac{G^2}{\mu_k} + GD\right) + \frac{G^2}{\mu}\right)\frac{k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)}{N}$$

$$+ G\sqrt{\frac{\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N\mu}} + \Delta_{\bar{k},opt} + \mathbb{E}\left[\Delta_{\bar{k},opt}\right].$$

A few remarks are in order.

Remark 8. *For sparse linear regression models in which Assumptions 4 holds with $\delta_N = e^{-c_0 N}$ (Agarwal et al., 2012), we have that $\frac{\log(1/\delta_N)}{\log(N)} = \frac{c_0 N}{\log(N)} \ge 2$ can always be fulfilled for sufficiently large sample size N, and the sparse excess risk bound in Theorem 4 roughly scales as $R(\tilde{w}_{S,k}) - R(w^*_{\bar{k}}) \lesssim$*

$$\frac{(k\log(d) + \log(1/\delta))^2 \log^2(N)}{N} + \sqrt{\frac{(k\log(d) + \log(1/\delta))\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N}} + \Delta_{\bar{k},opt} + \mathbb{E}\left[\Delta_{\bar{k},opt}\right].$$

In the well-specified $\bar{k}$-sparse models such that $R(w^*_{\bar{k}}) = R(w^*)$, the above rate improves to

$$R(\tilde{w}_{S,k}) - R(w^*_{\bar{k}}) \lesssim \frac{(k\log(d) + \log(1/\delta))^2 \log^2(N)}{N} + \Delta_{\bar{k},opt} + \mathbb{E}\left[\Delta_{\bar{k},opt}\right].$$

Remark 9. *For the IHT-style algorithms, under proper sparsity level relaxation, the sparse optimization sub-optimality $\Delta_{\bar{k},opt}$ in the above sparse excess risk bounds can be arbitrarily small (with high probability) after sufficient iteration (Jain et al., 2014).*

Remark 10. *Compared to the $\mathcal{O}\left(\frac{1}{N}\right)$ sparse excess risk bound for IHT (Yuan & Li, 2022, Theorem 3) for misspecified sparsity models, our $\mathcal{O}\left(\frac{1}{\sqrt{N}}\right)$ bound in Theorem 4 is slower in rate but more broadly applicable without needing to assume any strong-signal conditions as imposed by that result.*

4 OTHER RELATED WORK

Uniform stability and exponential generalization. Stimulated by a recent landmark work of Hardt et al. (2016), there is a renewed interest in the use of uniform stability for deriving generalization bounds for various learning algorithms and paradigms including stochastic gradient descent (SGD) (Kuzborskij & Lampert, 2018; Charles & Papailiopoulos, 2018; Bassily et al., 2020; Lei & Ying, 2020), stochastic model based optimization (Wang et al., 2017; Deng & Gao, 2021), optimization based meta learning (Zhou et al., 2019), and differential privacy stochastic optimization (Bassily et al., 2019; Feldman et al., 2020). Compared to other stability arguments, uniform stability is notorious for implying high-probability generalization bounds in addition to the traditional in-expectation bounds. Until very recently, the basic result of Bousquet & Elisseeff (2002), as expressed in equation 2, remains the best known exponential generalization bound for uniformly stable algorithms. Using some elegant proof techniques from adaptive data analysis, Feldman & Vondrak (2018) managed to replace the first term in equation 2 by $\sqrt{\gamma_u \log(\frac{1}{\delta})}$, which leads to an improvement whenever $\gamma_u \gtrsim \frac{1}{N}$. Soon after, a series of breakthrough results were obtained (Feldman & Vondrak, 2019; Bousquet et al., 2020) using tighter concentration bounds for sum of random functions, which eventually improve the stability dependent rate to a near-optimal one $\gamma_u \log(N) \log\left(\frac{1}{\delta}\right)$ as shown in equation 4. Additionally under the generalized Bernstein condition, these state-of-the-art results lead to $\mathcal{O}(\frac{1}{N})$ excess risk bounds for uniformly stable algorithms (Klochkov & Zhivotovskiy, 2021).

Non-uniform stability and exponential generalization. More broadly for non-uniformly stable algorithms, exponential generalization bounds have also been shown to be possible under various weaker and distribution-dependent notions of stability. As an early work in this line, Kutin & Niyogi (2002) showed that under the so called "almost-everywhere" stability, which is a high-probability counterpart of uniform stability, generalization bounds that hold with overwhelming probability are still possible in view of certain modified McDiarmid's inequality (Kutin, 2002). Later, Rakhlin et al. (2005) revisited the bounded-difference results of Kutin & Niyogi (2002) in a more straightforward manner by using a powerful moment extension of Efron-Stein inequality (Boucheron et al., 2005). Recently for general L_q-stable algorithms, the exponential leave-one-out generalization bounds were derived using moment or exponential extensions of Efron-Stein inequality, with applications found in ridge regression, k-nearest neighbor classification and k-folds cross-validation (Celisse & Guedj, 2016; Celisse & Mary-Huard, 2018; Abou-Moustafa & Szepesvári, 2019). We refer the interested readers to Boucheron et al. (2013); Kontorovich (2014); Combes (2015); Warnke (2016) for more results on concentration inequalities under high-probability bounded-difference conditions, which are fundamental for deriving exponential bounds for non-uniformly stable algorithms.

Generalization analysis of ℓ_0-estimators. We further briefly review some prior results on the generalization guarantees for statistical learning with cardinality constraint, which is the theme of the application part of our work. For the ℓ_0-ERM estimator as expressed in equation 9, provided that the solution is exactly known, a series of uniform excess risk bounds were derived over binary prediction classes (Chen & Lee, 2018; 2020) and bounded liner prediction classes (Foster & Syrgkanis, 2019). The exact solutions of ℓ_0-ERM, however, is computationally intractable in general high-dimensional cases due to the NP-hardness of problem. Therefore, it is more realistic and desirable to establish generalization bounds for approximate ℓ_0-estimators such as the IHT-style algorithms (Yuan et al., 2018; Garg & Khandekar, 2009; Li et al., 2016). Particularly for misspecified sparsity models, a set of sparse excess risk bounds with slow and fast rates were established through the lens of uniform stability theory under proper regularity conditions (Yuan & Li, 2022).

5 CONCLUSIONS

There exits a clear discrepancy between the recently developed near-optimal generalization bounds for uniformly stable algorithms and the best known yet sub-optimal bounds for L_q-stable algorithms. Aiming at closing such a theoretical gap, we for the first time derive in this paper a set of near-optimal exponential generalization bounds for L_q-stable algorithms that match those of the uniformly stable algorithms. We have demonstrated the strength of our new bounds in deriving exponential risk bounds for computationally tractable sparsity estimation algorithms, under milder conditions than those of the previous results. To conclude, L_q-stable algorithms generalize almost as sharply as uniformly stable algorithms, though the former notion of stability is less stringent than the latter.

REFERENCES

Karim Abou-Moustafa and Csaba Szepesvári. An exponential efron-stein inequality for l_q stable learning rules. In *Algorithmic Learning Theory*, pp. 31–63. PMLR, 2019.

Alekh Agarwal, Sahand Negahban, Martin J Wainwright, et al. Fast global convergence of gradient methods for high-dimensional statistical recovery. *The Annals of Statistics*, 40(5):2452–2482, 2012.

Sohail Bahmani, Bhiksha Raj, and Petros T Boufounos. Greedy sparsity-constrained optimization. *Journal of Machine Learning Research*, 14(Mar):807–841, 2013.

Raef Bassily, Vitaly Feldman, Kunal Talwar, and Abhradeep Thakurta. Private stochastic convex optimization with optimal rates. *arXiv preprint arXiv:1908.09970*, 2019.

Raef Bassily, Vitaly Feldman, Cristobal Guzman, and Kunal Talwar. Stability of stochastic gradient descent on nonsmooth convex losses. In *Advances in Neural Information Processing Systems*, pp. 1–10, 2020.

Thomas Blumensath and Mike E Davies. Iterative hard thresholding for compressed sensing. *Applied and Computational Harmonic Analysis*, 27(3):265–274, 2009.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. Concentration inequalities using the entropy method. *The Annals of Probability*, 31(3):1583–1614, 2003.

Stéphane Boucheron, Olivier Bousquet, Gábor Lugosi, and Pascal Massart. Moment inequalities for functions of independent random variables. *The Annals of Probability*, 33(2):514–560, 2005.

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Olivier Bousquet and André Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2(Mar):499–526, 2002.

Olivier Bousquet, Yegor Klochkov, and Nikita Zhivotovskiy. Sharper bounds for uniformly stable algorithms. In *Conference on Learning Theory*, pp. 610–626, 2020.

Alain Celisse and Benjamin Guedj. Stability revisited: new generalisation bounds for the leave-one-out. *arXiv preprint arXiv:1608.06412*, 2016.

Alain Celisse and Tristan Mary-Huard. Theoretical analysis of cross-validation for estimating the risk of the k-nearest neighbor classifier. *The Journal of Machine Learning Research*, 19(1):2373–2426, 2018.

Zachary Charles and Dimitris Papailiopoulos. Stability and generalization of learning algorithms that converge to global optima. In *International Conference on Machine Learning*, pp. 744–753, 2018.

Le-Yu Chen and Sokbae Lee. Best subset binary prediction. *Journal of Econometrics*, 206(1):39–56, 2018.

Le-Yu Chen and Sokbae Lee. Binary classification with covariate selection through ℓ_0-penalized empirical risk minimization. *The Econometrics Journal*, pp. 1–16, 2020.

Yuan Shih Chow and Henry Teicher. *Probability theory: independence, interchangeability, martingales*. Springer Science & Business Media, 2003.

Richard Combes. An extension of mcdiarmid's inequality. *arXiv preprint arXiv:1511.05240*, 2015.

Qi Deng and Wenzhi Gao. Minibatch and momentum model-based methods for stochastic nonsmooth non-convex optimization. *arXiv preprint arXiv:2106.03034*, 2021.

Luc Devroye and Terry Wagner. Distribution-free inequalities for the deleted and holdout error estimates. *IEEE Transactions on Information Theory*, 25(2):202–207, 1979.

Vitaly Feldman and Jan Vondrak. Generalization bounds for uniformly stable algorithms. In *Advances in Neural Information Processing Systems*, pp. 9747–9757, 2018.

Vitaly Feldman and Jan Vondrak. High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. In *Conference on Learning Theory*, pp. 1270–1279, 2019.

Vitaly Feldman, Tomer Koren, and Kunal Talwar. Private stochastic convex optimization: optimal rates in linear time. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pp. 439–449, 2020.

Dylan J Foster and Vasilis Syrgkanis. Statistical learning with a nuisance component. In *Conference on Learning Theory*, pp. 1346–1348, 2019.

Simon Foucart. Hard thresholding pursuit: an algorithm for compressive sensing. *SIAM Journal on Numerical Analysis*, 49(6):2543–2563, 2011.

Rahul Garg and Rohit Khandekar. Gradient descent with sparsification: an iterative algorithm for sparse recovery with restricted isometry property. In *International Conference on Machine Learning*, volume 9, pp. 337–344, 2009.

Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, pp. 1225–1234, 2016.

Prateek Jain, Ambuj Tewari, and Purushottam Kar. On iterative hard thresholding methods for high-dimensional m-estimation. In *Advances in Neural Information Processing Systems*, pp. 685–693, 2014.

Satyen Kale, Ravi Kumar, and Sergei Vassilvitskii. Cross-validation and mean-square stability. In *In Proceedings of the Second Symposium on Innovations in Computer Science (ICS2011*. Citeseer, 2011.

Yegor Klochkov and Nikita Zhivotovskiy. Stability and deviation optimal risk bounds with convergence rate $o(1/n)$. In *Advances in Neural Information Processing Systems*, 2021.

Vladimir Koltchinskii. Local rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6):2593–2656, 2006.

Aryeh Kontorovich. Concentration in unbounded metric spaces and algorithmic stability. In *International Conference on Machine Learning*, pp. 28–36. PMLR, 2014.

Samuel Kutin. Extensions to mcdiarmid's inequality when differences are bounded with high probability. *Dept. Comput. Sci., Univ. Chicago, Chicago, IL, USA, Tech. Rep. TR-2002-04*, 2002.

Samuel Kutin and Partha Niyogi. Almost-everywhere algorithmic stability and generalization error. In *Proceedings of the Eighteenth conference on Uncertainty in artificial intelligence*, pp. 275–282, 2002.

Ilja Kuzborskij and Christoph Lampert. Data-dependent stability of stochastic gradient descent. In *International Conference on Machine Learning*, pp. 2820–2829, 2018.

Yunwen Lei and Yiming Ying. Fine-grained analysis of stability and generalization for sgd. In *International Conference on Machine Learning*, 2020.

Xingguo Li, Tuo Zhao, Raman Arora, Han Liu, and Jarvis Haupt. Stochastic variance reduced optimization for nonconvex sparse learning. *International Conference on Machine Learning*, 2016.

Sayan Mukherjee, Partha Niyogi, Tomaso Poggio, and Ryan Rifkin. Learning theory: stability is sufficient for generalization and necessary and sufficient for consistency of empirical risk minimization. *Advances in Computational Mathematics*, 25(1-3):161–193, 2006.

Balas Kausik Natarajan. Sparse approximate solutions to linear systems. *SIAM Journal on Computing*, 24(2):227–234, 1995.

Alexander Rakhlin, Sayan Mukherjee, and Tomaso Poggio. Stability results in learning theory. *Analysis and Applications*, 3(04):397–417, 2005.

William H Rogers and Terry J Wagner. A finite sample distribution-free performance bound for local discrimination rules. *The Annals of Statistics*, pp. 506–514, 1978.

Shai Shalev-Shwartz, Ohad Shamir, Nathan Srebro, and Karthik Sridharan. Learnability, stability and uniform convergence. *Journal of Machine Learning Research*, 11(Oct):2635–2670, 2010.

Jie Shen and Ping Li. A tight bound of hard thresholding. *arXiv preprint arXiv:1605.01656*, 2016.

Jialei Wang, Mladen Kolar, Nathan Srebro, and Tong Zhang. Efficient distributed learning with sparsity. In *International Conference on Machine Learning*, pp. 3636–3645, 2017.

Lutz Warnke. On the method of typical bounded differences. *Combinatorics, Probability and Computing*, 25(2):269–299, 2016.

Xiao-Tong Yuan and Ping Li. Stability and risk bounds of iterative hard thresholding. *IEEE Transactions on Information Theory*, 2022.

Xiao-Tong Yuan, Ping Li, and Tong Zhang. Gradient hard thresholding pursuit. *Journal of Machine Learning Research*, 18:1–43, 2018.

Xiao-Tong Yuan, Bo Liu, Lezi Wang, Qingshan Liu, and Dimitris N Metaxas. Dual iterative hard thresholding. *J. Mach. Learn. Res.*, 21(152):1–50, 2020.

Tong Zhang. Leave-one-out bounds for kernel methods. *Neural Computation*, 15(6):1397–1437, 2003.

Pan Zhou, Xiao-Tong Yuan, and Jiashi Feng. Efficient stochastic gradient hard thresholding. In *Advances in Neural Information Processing Systems*, pp. 1984–1993, 2018.

Pan Zhou, Xiaotong Yuan, Huan Xu, Shuicheng Yan, and Jiashi Feng. Efficient meta learning via minibatch proximal update. In *Advances in Neural Information Processing Systems*, pp. 1532–1542, 2019.

A PRELIMINARIES

In this section, we collect some preliminary results that will be used in our analysis. We start by introducing the following L_q-norm generalization of the celebrated Efron-Stein inequality (Boucheron et al., 2005; Celisse & Guedj, 2016).

Proposition 1 (Generalized Efron-Stein inequality). *Let $S = \{Z_1, ..., Z_N\}$ be a set of independent random variables valued in $\mathcal{Z}$ and $g : \mathcal{Z}^N \mapsto \mathbb{R}$ be some measurable function. Then there exists a universal constant $\kappa < 1.271$ such that for all $q \geq 2$,*

$$\|g(S) - \mathbb{E}[g(S)]\|_q \leq \sqrt{2\kappa q} \sqrt{\left\| \sum_{i=1}^{N} \left(g(S) - g(S^{(i)}) \right)^2 \right\|_{q/2}}.$$

The following result is an immediate consequence of Proposition 1 when applied to sum of independent random variables, which revisits a version of Marcinkiewicz-Zygmund inequality (Chow & Teicher, 2003).

Proposition 2. *Let $Z_1, ..., Z_N$ be a set of independent centered random variables. Then for all $q \geq 2$,*

$$\left\| \sum_{i=1}^{N} Z_i \right\|_q \leq 2\sqrt{2\kappa q} \sqrt{\left\| \sum_{i=1}^{N} Z_i^2 \right\|_{q/2}}.$$

The following lemma is simple yet useful in our analysis.

Lemma 2. *Let $S = \{Z_1, ..., Z_N\}$ be a set of independent random variables valued in some measure space $\mathcal{Z}$ and $g : \mathcal{Z}^N \mapsto \mathbb{R}$ be some measurable function. Then for all $I \subseteq [N]$ and $q \geq 1$, we have*

$$\|g(S_I)\|_q \leq \|g(S)\|_q = \|\|g\|_q(S_I)\|_q.$$

Proof. Recall $g(S_I) = \mathbb{E}[g(S) \mid S_I]$. Then using Jensen's inequality we can show that

$$\|g(S_I)\|_q = \left(\mathbb{E}\left[|\mathbb{E}[g(S) \mid S_I]|^q \right] \right)^{1/q} \leq \left(\mathbb{E}\left[\mathbb{E}[|g(S)|^q \mid S_I]] \right] \right)^{1/q} = \left(\mathbb{E}[|g(S)|^q] \right)^{1/q} = \|g(S)\|_q.$$

By definition we can also express $\|g(S)\|_q = \left(\mathbb{E}\left[\mathbb{E}[|g(S)|^q \mid S_I]] \right] \right)^{1/q} = \|\|g(S)\|_q(S_I)\|_q$. $\square$

As a direct consequence of Lemma 2, the following result indicates that conditional expectation does not expand the differences in L_q-norm.

Lemma 3. *Let $S = \{Z_1, ..., Z_N\}$ be a set of independent random variables valued in some measure space $\mathcal{Z}$ and $g : \mathcal{Z}^N \mapsto \mathbb{R}$ be some measurable function. Let $I \subseteq [N]$ be an index set. Then for all $i \in I$ and $q \geq 1$,*

$$\left\| g(S_I) - g(S_I^{(i)}) \right\|_q \leq \left\| g(S) - g(S^{(i)}) \right\|_q.$$

Proof. For each $i \in I$, by applying Lemma 2 to $g(S) - g(S^{(i)})$ we can show that $\|g(S_I) - g(S_I^{(i)})\|_q \leq \|g(S) - g(S^{(i)})\|_q$, which gives the desired result. $\square$

We also need the following lemma about the equivalence between tails and moments (see, e.g., Bousquet et al., 2020).

Lemma 4. *Let Y be a real-valued random variable.*

- *Suppose that Y satisfies the following inequality for some $a, b \geq 0$ with probability at least $1 - \delta$ for any $\delta \in (0, 1)$,*

$$|Y| \leq a \sqrt{\log\left(\frac{e}{\delta}\right)} + b \log\left(\frac{e}{\delta}\right).$$

  Then, for any $q \geq 1$ it holds that

$$\|Y\|_q \leq 3a\sqrt{q} + 9bq.$$

- *Suppose that Y satisfies $\|Y\|_q \leq f(q)$ for any $1 \leq q_l \leq q < q_u$ and some non-negative real function f. Then the following holds with probability at least $1 - \delta$ for any $\delta \in (e^{1-q_u}, e^{1-q_l}]$:*

$$|Y| \leq ef\left(\log\left(\frac{e}{\delta}\right)\right).$$

Proof. We only prove the second part which slightly generalizes the corresponding result of (Bousquet et al., 2020, Lemma 1). For any $\delta \in (e^{1-q_u}, e^{1-q_l}]$, we choose $q = \log(e/\delta) \in [q_l, q_u)$. Using the condiiton $\|Y\|_q \leq f(q)$ and Markov inequality yields

$$\mathbb{P}\left(|Y| > ef\left(\log\left(\frac{e}{\delta}\right)\right)\right) \leq \mathbb{P}\left(|Y| > e\|Y\|_q\right) \leq \frac{\mathbb{E}[|Y|^q]}{e^q\|Y\|_q^q} = \frac{\delta}{e} \leq \delta.$$

This proves the desired bound in the second part. $\qquad\square$

Remark 11. *Specially if $q_l = 1$ and $q_u = \infty$ is allowed, then the second bound in Lemma 4 holds with an arbitrary tail bound $\delta \in (0, 1)$. Suppose that it holds with probability at least $1 - \delta$ over $S, S^{(i)}, Z$ such that*

$$\left|\ell(A(S); Z) - \ell(A(S^{(i)}); Z)\right| \leq a\log\left(\frac{e}{\delta}\right) + b\sqrt{\log\left(\frac{e}{\delta}\right)},$$

then according to Lemma 4 we have that A is L_q-stable by $\gamma_q = 3aq + 9b\sqrt{q}$.

B PROOFS FOR SECTION 2

B.1 PROOF OF THEOREM 1

The proof is a generalization of that of Bousquet et al. (2020, Theorem 4) under the considered property of L_q-norm bounded difference. We reproduce the proof here for the sake of completeness.

Proof of Theorem 1. Consider k such that $2^{k-1} < N \leq 2^k$. If $N < 2^k$, we pad the set with extra zero-functions so that $N = 2^k$. Consider the partition $\mathcal{I}_0, ..., \mathcal{I}_k$ of $[N]$ given by

$$\mathcal{I}_0 = \{\{1\}, ..., \{2^k\}\}, \quad \mathcal{I}_1 = \{\{1, 2\}, \{3, 4\}..., \{2^k - 1, 2^k\}\}, \quad \mathcal{I}_k = \{\{1, ..., 2^k\}\}.$$

For any $i \in [N]$ and $l = 0, ..., k$, we denote by $I^l(i) \in \mathcal{I}_l$ the only set from $\mathcal{I}_l$ that contains i and consider the following random variables

$$g_i^l = \mathbb{E}\left[g_i \mid Z_i, S_{\overline{I^l(i)}}\right].$$

In particular, $g_i^0 = g_i$ and $g_i^k = \mathbb{E}[g_i \mid Z_i]$. Clearly we have the following telescope sum:

$$g_i = \sum_{l=0}^{k-1}(g_i^l - g_i^{l+1}) + \mathbb{E}[g_i \mid Z_i].$$

It follows that

$$\left\|\sum_{i=1}^N g_i - \mathbb{E}[g_i \mid Z_i]\right\|_q \leq \sum_{l=0}^{k-1}\left\|\sum_{i=1}^N g_i^l - g_i^{l+1}\right\|_q. \tag{11}$$

We need to upper bound the right hand side of the above inequality. To this end, it can be verified that

$$g_i^{l+1} = \mathbb{E}\left[g_i \mid Z_i, S_{\overline{I^{l+1}(i)}}\}\right] = \mathbb{E}\left[g_i^l \mid Z_i, S_{\overline{I^{l+1}(i)}}\right].$$

Since g_i has a bounded L_q-difference by β_q with respect to all variables except the i-th variable, it is known from Lemma 3 that so is g_i^l for each $l = 0, ..., k$. Conditioned on $Z_i, S_{\overline{I^{l+1}(i)}}$, invoking Proposition 1 to g_i^l yields

$$\|g_i^l - g_i^{l+1}\|_q\left(Z_i, S_{\overline{I^{l+1}(i)}}\right) \leq \sqrt{2\kappa q 2^l}\beta_q,$$

as there are 2^l indices in $I^{l+1}(i) \setminus I^l(i)$. It follows from Lemma 2 that

$$\|g_i^l - g_i^{l+1}\|_q = \left\| \|g_i^l - g_i^{l+1}\|_q (Z_i, S_{\overline{I^{l+1}(i)}}) \right\|_q \leq \sqrt{2\kappa q 2^l} \beta_q.$$

Now consider any $I^l \in \mathcal{I}_l$. Since for each $i \in I_l$, $g_i^l - g_i^{l+1}$ depends only on $Z_i, Z_{\overline{I^l}}$, these terms are independent and centered conditioned on $S_{\overline{I^l}}$. Therefore, applying Proposition 2 yields

$$\left\| \sum_{i \in I^l} g_i^l - g_i^{l+1} \right\|_q \left(S_{\overline{I^l}} \right) \leq 2\sqrt{2\kappa q 2^l} \times \sqrt{2\kappa q 2^l} \beta_q = 4\kappa q 2^l \beta_q,$$

which according to Lemma 2 implies that

$$\left\| \sum_{i \in I^l} g_i^l - g_i^{l+1} \right\|_q \leq 4\kappa q 2^l \beta_q.$$

Then based on triangle inequality we get

$$\left\| \sum_{i \in [N]} g_i^l - g_i^{l+1} \right\|_q \leq \sum_{I^l \in \mathcal{I}_l} \left\| \sum_{i \in I^l} g_i^l - g_i^{l+1} \right\|_q \leq 2^{k-l} \times 4\kappa q 2^l \beta_q = 4\kappa q 2^k \beta_q < 4\kappa q N \beta_q.$$

Finally, the right hand side of equation 11 can be bounded as

$$\left\| \sum_{i=1}^N g_i - \mathbb{E}[g_i \mid Z_i] \right\|_q \leq \sum_{l=0}^{k-1} \left\| \sum_{i=1}^N g_i^l - g_i^{l+1} \right\|_q \leq 4\kappa q N \lceil \log_2 N \rceil \beta_q, \tag{12}$$

which gives the first desired bound. In view of equation 12 and triangle inequality we have

$$\left\| \sum_{i=1}^N g_i \right\|_q \leq \left\| \sum_{i=1}^N \mathbb{E}[g_i \mid Z_i] \right\|_q + 4\kappa q N \lceil \log_2 N \rceil \beta_q. \tag{13}$$

Since $\|\mathbb{E}[g_i(S) \mid Z_i]\|_q \leq M_q$ and $\mathbb{E}[g_i(S) \mid S \setminus Z_i] = 0$, it follows from Proposition 2 that the first term at the right hand side of equation 13 can be bounded as

$$\left\| \sum_{i=1}^N \mathbb{E}[g_i \mid Z_i] \right\|_q \leq 2\sqrt{2\kappa N q} M_q. \tag{14}$$

The second desired bound is obtained by plugging equation 14 into equation 13. $\qquad \square$

B.2 Proof of Theorem 2

Proof. Let us consider

$$h_i(S) := R(A(S)) - \ell(A(S); Z_i), \quad g_i(S) = \mathbb{E}_{Z_i'} \left[R(A(S^{(i)})) - \ell(A(S^{(i)}); Z_i) \right].$$

Then the L_q-norm of the generalization gap can be bounded as

$$\|R(A(S)) - R_S(A(S))\|_q = \frac{1}{N} \left\| \sum_{i=1}^N h_i(S) \right\|_q \leq \frac{1}{N} \left(\underbrace{\left\| \sum_{i=1}^N g_i(S) \right\|_q}_{A} + \underbrace{\left\| \sum_{i=1}^N (h_i(S) - g_i(S)) \right\|_q}_{B} \right). \tag{15}$$

We next respectively upper bound the two terms A and B in equation 15. To bound the term A, by definition it holds that $\mathbb{E}[g_i(S) \mid S \setminus Z_i] = 0$. Based on triangle inequality we can show that

$$\|\mathbb{E}[g_i(S) \mid Z_i]\|_q \leq \|g_i(S)\|_q$$
$$= \left\| \mathbb{E}_{Z_i'}[\mathbb{E}_Z[\ell(A(S^{(i)}); Z)]] - \mathbb{E}_{Z_i'} \left[\ell(A(S^{(i)}); Z_i) \right] \right\|_q$$
$$\leq \|\ell(A(S^{(i)}); Z)\|_q + \|\ell(A(S^{(i)}); Z_i)\|_q \leq 2M_q,$$

where in the first and second inequalities we have twice used Lemma 2. Next we further show that g_i has a bounded L_q-norm difference by $2\gamma_q$ with respect to all variables in S except Z_i. Indeed, for each $j \neq i$ it can be verified that

$$\left\| g_i(S) - g_i(S^{(j)}) \right\|_q$$

$$\leq \left\| \mathbb{E}_{Z'_i} \left[R(A(S^{(i)})) - R(A((S^{(i)})^{(j)})) \right] \right\|_q + \left\| \mathbb{E}_{Z'_i} \left[\ell(A(S^{(i)}); Z_i) - \ell(A((S^{(i)})^{(j)}); Z_i) \right] \right\|_q$$

$$= \left\| \mathbb{E}_{Z'_i} \mathbb{E}_Z [\ell(A(S^{(i)}); Z) - \ell(A((S^{(i)})^{(j)}); Z)] \right\|_q + \left\| \mathbb{E}_{Z'_i} [\ell(A(S^{(i)}); Z_i) - \ell(A((S^{(i)})^{(j)}); Z_i)] \right\|_q$$

$$\leq \left\| \ell(A(S^{(i)}); Z) - \ell(A((S^{(i)})^{(j)}); Z) \right\|_q + \left\| \ell(A(S^{(i)}); Z_i) - \ell(A((S^{(i)})^{(j)}); Z_i) \right\|_q \leq 2\gamma_q,$$

where in the last but one inequality we have used Lemma 2, while in the last equality we have used the L_q-stability assumption on the algorithm A. Therefore, $\{g_i\}$ satisfy the conditions of Theorem 1 and thus

$$A = \left\| \sum_{i=1}^N g_i(S) \right\|_q \leq 2\sqrt{2\kappa N q} M_q + 8\kappa q N \lceil \log_2 N \rceil \gamma_q. \tag{16}$$

Now we proceed to bound the term B. It can be verified that

$$B \leq \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[R(A(S)) - R(A(S^{(i)})) \right] \right\|_q + \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[\ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right] \right\|_q$$

$$= \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \mathbb{E}_Z \left[\ell(A(S); Z) - \ell(A(S^{(i)}); Z) \right] \right\|_q + \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[\ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right] \right\|_q$$

$$\leq \sum_{i=1}^N \left\| \ell(A(S); Z) - \ell(A(S^{(i)}); Z) \right\|_q + \sum_{i=1}^N \left\| \ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right\|_q \leq 2N\gamma_q, \tag{17}$$

where in the last but one inequality we have used Lemma 2, and in the last equality we have used the L_q-stability assumption. Plugging bounds equation 16 and equation 17 into equation 15 and preserving leading terms yields the desired result. □

B.3 PROOF OF THEOREM 3

We need the following lemma which plays a fundamental role in proving the main result.

Lemma 5. *Let $A : \mathcal{Z}^N \mapsto \mathcal{W}$ be a learning algorithm that has L_q-stability by γ_q for $q \geq 1$. Suppose that $\|\ell(A(S); Z)\|_q \leq M_q$ for any $Z \in \mathcal{Z}$. Let S' be an independent copy of S. Then the following bound holds for all $q \geq 2$:*

$$\left\| R(A(S)) - R_S(A(S)) - \mathbb{E}[R(A(S))] + \frac{1}{N} \sum_{i=1}^N \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i] \right\|_q \lesssim q\gamma_q \log(N).$$

Proof. Let us again consider $g_i(S) = \mathbb{E}_{Z'_i} \left[R(A(S^{(i)})) - \ell(A(S^{(i)}); Z_i) \right]$. Then using similar proof arguments to those of Theorem 2 we can show that

$$\left\| N(R(A(S)) - R_S(A(S))) - \sum_{i=1}^N g_i(S) \right\|_q$$

$$\leq \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[R(A(S)) - R(A(S^{(i)})) \right] \right\|_q + \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[\ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right] \right\|_q$$

$$= \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \mathbb{E}_Z \left[\ell(A(S); Z) - \ell(A(S^{(i)}); Z) \right] \right\|_q + \left\| \sum_{i=1}^N \mathbb{E}_{Z'_i} \left[\ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right] \right\|_q$$

$$\leq \sum_{i=1}^N \left\| \ell(A(S); Z) - \ell(A(S^{(i)}); Z) \right\|_q + \sum_{i=1}^N \left\| \ell(A(S); Z_i) - \ell(A(S^{(i)}); Z_i) \right\|_q \leq 2N\gamma_q,$$

which implies

$$\left\| R(A(S)) - R_S(A(S)) - \frac{1}{N} \sum_{i=1}^{N} g_i(S) \right\|_q \le 2\gamma_q.$$

Also, $g_i(S)$ satisfies the conditions of Theorem 1 with $\beta_q = 2\gamma_q$ and it follows from the second bound of Theorem 1 that for all $q \ge 2$,

$$\left\| \frac{1}{N} \sum_{i=1}^{N} (g_i(S) - \mathbb{E}[g_i(S) \mid Z_i]) \right\|_q \le 8\kappa q \gamma_q \lceil \log_2 N \rceil.$$

Combining the above two yields

$$\left\| R(A(S)) - R_S(A(S)) - \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}[g_i(S) \mid Z_i] \right\|_q \lesssim q\gamma_q \log(N).$$

The desired result follows by noting that

$$\mathbb{E}[g_i(S) \mid Z_i] = \mathbb{E}[R(A(S'))] - \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i] = \mathbb{E}[R(A(S))] - \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i].$$

This completes the proof. $\qquad\square$

With Lemma 5 in place, we are ready to prove the main result of Theorem 3.

Proof of Theorem 3. Consider any $w^* \in W^*$. It is standard to decompose and bound the excess risk as

$$\begin{aligned}
&R(A(S)) - R^* \\
=&R(A(S)) - R_S(A(S)) + R_S(A(S)) - R_S(w^*) + R_S(w^*) - R^* \\
\le&\Delta_{\text{opt}} + R(A(S)) - R_S(A(S)) - (R^* - R_S(w^*)) \\
=&\Delta_{\text{opt}} + \Gamma(S) + \mathbb{E}[R(A(S))] - \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i] - (R^* - R_S(w^*)),
\end{aligned} \tag{18}$$

where

$$\Gamma(S) = R(A(S)) - R_S(A(S)) - \mathbb{E}[R(A(S))] + \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i].$$

Since we have the freedom to choose w^*, let us specify it in the above as $w^*(S') \in W^*$ which is the minimizer that satisfies the Bernstein condition in Assumption 1 associated with $A(S')$. Then, it follows from equation 18 that

$$\begin{aligned}
&R(A(S)) - R^* - \Delta_{\text{opt}} \\
\le&\Gamma(S) + \mathbb{E}[R(A(S))] - \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}[\ell(A(S'); Z_i) \mid Z_i] - (R^* - \mathbb{E}[R_S(w^*(S')) \mid S]).
\end{aligned}$$

Consequently,

$$\begin{aligned}
&\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \\
\le&\|\Gamma(S)\|_q + \left\| \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right] - (R^* - \mathbb{E}[R(A(S'))]) \right\|_q \\
\stackrel{\zeta_1}{\lesssim}&q\gamma_q \log(N) + \underbrace{\left\| \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right] - (R^* - \mathbb{E}[R(A(S'))]) \right\|_q}_{T},
\end{aligned} \tag{19}$$

where in "ζ_1" we have applied Lemma 5 to obtain $\|\Gamma(S)\|_q \le q\gamma_q \log(N)$, and the fact $\mathbb{E}[R(A(S))] = \mathbb{E}[R(A(S'))]$.

Part (a): To bound the term T, using Bernstein's inequality together with Bernstein condition we can show that (see the proof arguments of Klochkov & Zhivotovskiy (2021, Theorem 1.1) for more details):

$$T \lesssim \sqrt{\frac{qB\mathbb{E}[R(A(S)) - R^*]}{N}} + \frac{qM}{N} = \sqrt{\frac{qB(\mathbb{E}[R(A(S)) - R^* - \Delta_{\text{opt}}] + \mathbb{E}[\Delta_{\text{opt}}])}{N}} + \frac{qM}{N}$$
$$\leq \sqrt{\frac{qB(\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q + \mathbb{E}[\Delta_{\text{opt}}])}{N}} + \frac{qM}{N},$$

where the last inequality is due to Hölder's inequality. Therefore, combining the above and equation 19 yields that for some universal constant C:

$$\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \leq C\left(q\gamma_q \log(N) + \sqrt{\frac{qB(\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q + \mathbb{E}[\Delta_{\text{opt}}])}{N}} + \frac{qM}{N}\right),$$

which implies that

$$\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \lesssim \mathbb{E}[\Delta_{\text{opt}}] + q\gamma_q \log(N) + \frac{(M + B)q}{N}.$$

This gives the desired bound in part (a).

Part (b): Under the given conditions in part (b), we can bound the term T in equation 19 as follows for $q \geq 2$:

$$T = \left\| \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right] - (R^* - \mathbb{E}[R(A(S'))]) \right\|_q$$

$$\overset{\zeta_1}{\leq} \frac{2\sqrt{2\kappa q}}{N} \sqrt{\left\| \sum_{i=1}^{N} \left(\mathbb{E}\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right] - (R^* - \mathbb{E}[R(A(S'))])\right)^2 \right\|_{q/2}}$$

$$\leq \frac{4\sqrt{\kappa q}}{N} \sqrt{\sum_{i=1}^{N} \left\| \mathbb{E}^2\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right] + (R^* - \mathbb{E}[R(A(S'))])^2 \right\|_{q/2}}$$

$$\leq \frac{4\sqrt{\kappa q}}{N} \sqrt{\sum_{i=1}^{N} \left\| \mathbb{E}^2\left[\ell(w^*(S'); Z_i) - \ell(A(S'); Z_i) \mid Z_i\right]\right\|_{q/2} + N\mathbb{E}^2[R(w^*(S')) - R(A(S'))]}$$

$$\overset{\zeta_2}{\leq} \frac{4\sqrt{\kappa q}}{N} \sqrt{\sum_{i=1}^{N} \left\| \mathbb{E}^2\left[G\|w^*(S') - A(S')\| \mid Z_i\right]\right\|_{q/2} + N\mathbb{E}^2\left[G\|w^*(S') - A(S')\|\right]}$$

$$\overset{\zeta_3}{\leq} \frac{4G\sqrt{2\kappa q}}{\sqrt{N}} \sqrt{\mathbb{E}\left[\|w^*(S') - A(S')\|^2\right]}$$

$$\overset{\zeta_4}{\leq} \frac{8G\sqrt{\kappa q}}{\sqrt{N}} \sqrt{\frac{1}{\mu}\mathbb{E}\left[R(A(S')) - R^*\right]} = \frac{8G\sqrt{\kappa q}}{\sqrt{N\mu}} \sqrt{\mathbb{E}\left[R(A(S')) - R^* - \Delta_{\text{opt}}\right] + \mathbb{E}[\Delta_{\text{opt}}]}$$

$$\leq \frac{8G\sqrt{\kappa q}}{\sqrt{N\mu}} \sqrt{\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q + \mathbb{E}[\Delta_{\text{opt}}]}$$

where in "ζ_1" we have used Proposition 2, in "ζ_2" we have used the Lipschitz-loss condition, in "ζ_3" we have used $\left\|\mathbb{E}^2\left[G\|w^*(S') - A(S')\| \mid Z_i\right]\right\|_{q/2} = \mathbb{E}^2\left[G\|w^*(S') - A(S')\|\right] \leq G^2\mathbb{E}\left[\|w^*(S') - A(S')\|^2\right]$, in "$\zeta_4$" we have used Assumption 2, and the last inequality is due to Hölder's inequality. Then, plugging the above bound of term T into equation 19 yields that for some universal constant C:

$$\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \leq C\left(q\gamma_q \log(N) + G\sqrt{\frac{q}{N\mu}}\sqrt{\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q + \mathbb{E}[\Delta_{\text{opt}}]}\right),$$

which then implies

$$\|R(A(S)) - R^* - \Delta_{\text{opt}}\|_q \lesssim \mathbb{E}[\Delta_{\text{opt}}] + q\gamma_q \log(N) + \frac{qG^2}{\mu N}.$$

This gives the desired bound in part (b). The proof is completed. □

C PROOFS FOR SECTION 3

C.1 PROOF OF LEMMA 1

Proof. Let us consider the following event about the restricted strong convexity of R_S:

$$\mathcal{E} : R_S \text{ is } \mu_k\text{-strongly convex.}$$

Let $Y = 1_{\mathcal{E}}$ be the indication random variable associated with $\mathcal{E}$. Then by Assumption 4 we have $\mathbb{P}(Y = 1) \geq 1 - \delta_N$. Suppose that $\mathcal{E}$ occurs such that $Y = 1$. Then,

$$
\begin{aligned}
&R_S(w^*_{S^{(i)}|J}) - R_S(w^*_{S|J})\\
=&\frac{1}{N}\sum_{j\neq i}\left(\ell(w^*_{S^{(i)}|J}; Z_j) - \ell(w^* w_{S|J}; Z_j)\right) + \frac{1}{N}\left(\ell(w^*_{S^{(i)}|J}; Z_i) - \ell(w^*_{S|J}; Z_i)\right)\\
=&R_{S^{(i)}}(w^*_{S^{(i)}|J}) - R_{S^{(i)}}(w^*_{S|J}) + \frac{1}{N}\left(\ell(w^*_{S^{(i)}|J}; Z_i) - \ell(w^*_{S|J}; Z_i)\right)\\
&- \frac{1}{N}\left(\ell(w^*_{S^{(i)}|J}; Z'_i) - \ell(w^*_{S|J}; Z'_i)\right)\\
\leq&\frac{1}{N}\left|\ell(w^*_{S^{(i)}|J}; Z_i) - \ell(w^*_{S|J}; Z_i)\right| + \frac{1}{N}\left|\ell(w^*_{S^{(i)}|J}; Z'_i) - \ell(w^*_{S|J}; Z'_i)\right|\\
\leq&\frac{2G}{N}\left\|w^*_{S^{(i)}|J} - w^*_{S|J}\right\|,
\end{aligned}
$$

where we have used the optimality of $w^*_{S^{(i)}|J}$ with respect to $R_{S^{(i)}}(w)$ and the Lipschitz continuity of loss. Since $\mathcal{E}$ occurs by assumption, R_S is μ_k-strongly convex. Since $w^*_{S|J}$ is optimal for $R_S(w)$ over the supporting set J, we have

$$R_S(w^*_{S^{(i)}|J}) \geq R_S(w^*_{S|J}) + \frac{\mu_k}{2}\left\|w^*_{S^{(i)}|J} - w^*_{S|J}\right\|^2.$$

Combing the preceding two inequalities yields $\left\|w^*_{S^{(i)}|J} - w^*_{S|J}\right\| \leq \frac{4G}{\mu_k N}$. Consequently from the Lipschitz continuity of ℓ we have that for any $Z \in \mathcal{Z}$, the following holds conditioned on $Y = 1$:

$$\left|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)\right| \leq G\left\|w^*_{S^{(i)}|J} - w^*_{S|J}\right\| \leq \frac{4G^2}{\mu_k N}. \tag{20}$$

In the complementary case of $Y = 0$, in view of Assumption 5, it always holds that

$$|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)| \leq G\left\|w^*_{S^{(i)}|J} - w^*_{S|J}\right\| \leq 2GD. \tag{21}$$

Let us consider $q_u := \frac{\log\left(\frac{1}{\delta_N}\right)}{\log(N)}$. By assumption $q_u \geq 2$. Then for $2 \leq q \leq q_u$, it can be verified that

$$
\begin{aligned}
&\mathbb{E}\left[\left|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)\right|^q\right]\\
=&\mathbb{P}(Y = 1)\mathbb{E}\left[\left|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)\right|^q \mid Y = 1\right] + \mathbb{P}(Y = 0)\mathbb{E}\left[\left|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)\right|^q \mid Y = 0\right]\\
\leq&\left(\frac{4G^2}{\mu_k N}\right)^q + \delta_N(2GD)^q = \left(\frac{4G^2}{\mu_k N}\right)^q + \frac{1}{N^{q_u}}(2GD)^q \leq \left(\frac{4G^2}{\mu_k N}\right)^q + \frac{1}{N^q}(2GD)^q.
\end{aligned}
$$

It follows that for all $2 \leq q \leq q_u$

$$\left\|\ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z)\right\|_q \leq \left(\left(\frac{4G^2}{\mu_k N}\right)^q + \frac{1}{N^q}(2GD)^q\right)^{1/q} \leq \frac{1}{N}\left(\frac{4G^2}{\mu_k} + 2GD\right),$$

where we have used $a^q + b^q \leq (a+b)^q$ for $a, b > 0$ and $q \geq 2$. For the complementary case $q > q_u$, it is trivial to show that

$$\left\| \ell(w^*_{S^{(i)}|J}; Z) - \ell(w^*_{S|J}; Z) \right\|_q \leq 2GD \leq \frac{2GD}{q_u} q.$$

Assembling the preceding two bounds yields the desired L_q-stability bound. $\qquad \square$

C.2 PROOF OF THEOREM 4

Let us denote $a_+ = \max\{a, 0\}$. We need the following key result in our proof.

Lemma 6. *Suppose that Assumptions 3, 4, 5 hold. Assume that $\frac{\log(1/\delta_N)}{\log(N)} \geq 2$. Then for any $\delta \in (0, e^{-1})$, it holds with probability at least $1 - \delta$ that*

$$
\sup_{J \subseteq [d], |J| = k} R(w^*_{S|J}) - R(w^*_{\bar{k}})
$$

$$
\lesssim \frac{GD \left(k \log \left(\frac{ed}{k} \right) + \log \left(\frac{e}{\delta} \right) \right)^2 \log^2(N)}{\log(1/\delta_N)} + \left(\log(N) \left(\frac{G^2}{\mu_k} + GD \right) + \frac{G^2}{\mu} \right) \frac{k \log \left(\frac{ed}{k} \right) + \log \left(\frac{e}{\delta} \right)}{N}
$$

$$
+ G \sqrt{\frac{\left(k \log \left(\frac{ed}{k} \right) + \log \left(\frac{e}{\delta} \right) \right) \left(R(w^*_{\bar{k}}) - R(w^*) \right)}{N\mu}} + \left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}}) \right)_+
$$

$$
+ \mathbb{E} \left[\left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}}) \right)_+ \right].
$$

Proof. Given a fixed index set $J \subseteq [d]$ with $|J| = k$, we can show that the following holds for any $q \geq 1$:

$$
\left\| R(w^*_{S|J}) - R_S(w^*_{S|J}) + R_S(w^*) - R(w^*) \right\|_q
$$

$$
= \left\| \Gamma_{S|J} + \mathbb{E}[R(w^*_{S|J})] - \frac{1}{N} \sum_{i=1}^N \mathbb{E}[\ell(w^*_{S'|J}; Z_i) \mid Z_i] + R_S(w^*) - R(w^*) \right\|_q \tag{22}
$$

$$
\leq \left\| \Gamma_{S|J} \right\|_q + \underbrace{\left\| \mathbb{E}[R(w^*_{S|J})] - \frac{1}{N} \sum_{i=1}^N \mathbb{E}[\ell(w^*_{S'|J}; Z_i) \mid Z_i] - R(w^*) + R_S(w^*) \right\|_q}_{T},
$$

where

$$
\Gamma_{S|J} = R(w^*_{S|J}) - R_S(w^*_{S|J}) - \mathbb{E}[R(w^*_{S|J})] + \frac{1}{N} \sum_{i=1}^N \mathbb{E}[\ell(w^*_{S'|J}; Z_i) \mid Z_i].
$$

In view of Lemma 1 we have that $w^*_{S|J}$ has L_q-stability by

$$
\gamma_q = \frac{1}{N} \left(\frac{4G^2}{\mu_k} + 2GD \right) + \frac{2GD \log(N) q}{\log(1/\delta_N)}.
$$

Then invoking Lemma 5 over the supporting set J yields

$$
\left\| \Gamma_{S|J} \right\|_q \leq q \gamma_q \log(N) = \frac{q \log(N)}{N} \left(\frac{4G^2}{\mu_k} + 2GD \right) + \frac{2GD \log^2(N) q^2}{\log(1/\delta_N)}. \tag{23}
$$

We now bound the term T in equation 22 as follows:

$$T = \left\| \frac{1}{N} \sum_{i=1}^{N} \mathbb{E}\left[w^*; Z_i \right) - \ell(w^*_{S'|J}; Z_i) \mid Z_i \right] - (R(w^*) - \mathbb{E}[R(w^*_{S'|J})]) \right\|_q$$

$$\overset{\zeta_1}{\leq} \frac{2\sqrt{2\kappa q}}{N} \sqrt{ \left\| \sum_{i=1}^{N} \left(\mathbb{E}\left[\ell(w^*; Z_i) - \ell(w^*_{S'|J}; Z_i) \mid Z_i \right] - (R(w^*) - \mathbb{E}[R(w^*_{S'|J})]) \right)^2 \right\|_{q/2} }$$

$$\leq \frac{4\sqrt{\kappa q}}{N} \sqrt{ \sum_{i=1}^{N} \left\| \mathbb{E}^2 \left[\ell(w^*; Z_i) - \ell(w^*_{S'|J}; Z_i) \mid Z_i \right] + \left(R(w^*) - \mathbb{E}[R(w^*_{S'|J})] \right)^2 \right\|_{q/2} }$$

$$\leq \frac{4\sqrt{\kappa p}}{N} \sqrt{ \sum_{i=1}^{N} \left\| \mathbb{E}^2 \left[\ell(w^*; Z_i) - \ell(w^*_{S'|J}; Z_i) \mid Z_i \right] \right\|_{q/2} + N \mathbb{E}^2 \left[R(w^*) - R(w^*_{S'|J}) \right] } \quad (24)$$

$$\overset{\zeta_2}{\leq} \frac{4\sqrt{\kappa p}}{N} \sqrt{ \sum_{i=1}^{N} \left\| \mathbb{E}^2 \left[G \left\| w^* - w^*_{S'|J} \right\| \mid Z_i \right] \right\|_{q/2} + N \mathbb{E}^2 \left[G \left\| w^* - w^*_{S'|J} \right\| \right] }$$

$$\overset{\zeta_3}{\leq} \frac{4G\sqrt{2\kappa p}}{\sqrt{N}} \sqrt{ \mathbb{E}\left[\left\| w^* - w^*_{S'|J} \right\|^2 \right] }$$

$$\overset{\zeta_4}{\leq} \frac{8G\sqrt{\kappa q}}{\sqrt{N}} \sqrt{ \frac{1}{\mu} \mathbb{E}\left[R(w^*_{S'|J}) - R(w^*) \right] } = \frac{8G\sqrt{\kappa q}}{\sqrt{N\mu}} \sqrt{ \mathbb{E}\left[R(w^*_{S|J}) - R(w^*) \right] },$$

where in "ζ_1" we have used Proposition 2, in "ζ_2" we have used the Lipschitz-loss assumption, in "ζ_3" we have used $\left\| \mathbb{E}^2 \left[G\|w^* - w^*_{S'|J}\| \mid Z_i \right] \right\|_{q/2} = \mathbb{E}^2 \left[G\|w^* - w^*_{S'|J}\| \right] \leq G^2 \mathbb{E}\left[\|w^* - w^*_{S'|J}\|^2 \right]$, in "$\zeta_4$" we have used the strong convexity condition in Assumption 4. Plugging equation 23 and equation 24 into equation 22 yields that for any $q \geq 2$,

$$\left\| R(w^*_{S|J}) - R(w^*) - R_S(w^*_{S|J}) + R_S(w^*) \right\|_q$$

$$\leq \frac{q \log(N)}{N} \left(\frac{4G^2}{\mu_k} + 2GD \right) + \frac{2q^2 GD \log^2(N)}{\log(1/\delta_N)} + 8G\sqrt{ \frac{\kappa q \mathbb{E}\left[R(w^*_{S|J}) - R(w^*) \right]}{N\mu} }. \quad (25)$$

Next we need to upper bound the factor $\mathbb{E}\left[R(w^*_{S|J}) - R(w^*) \right]$ in the second term of the above bound. To do so, let us consider $q = 2$ in equation 25. It follows from the optimality of w^* and $w^*_{S|J}$ that

$$\mathbb{E}\left[R(w^*_{S|J}) - R(w^*) - R_S(w^*_{S|J}) + R_S(w^*) \right]$$

$$\leq \left\| R(w^*_{S|J}) - R(w^*) - R_S(w^*_{S|J}) + R_S(w^*) \right\|_2$$

$$\overset{equation\ 25}{\leq} \frac{\log(N)}{N} \left(\frac{8G^2}{\mu_k} + 4GD \right) + \frac{8GD \log^2(N)}{\log(1/\delta_N)} + 8G\sqrt{ \frac{2\kappa \mathbb{E}\left[R(w^*_{S|J}) - R(w^*) \right]}{N\mu} }$$

$$\leq \frac{\log(N)}{N} \left(\frac{8G^2}{\mu_k} + 4GD \right) + \frac{8GD \log^2(N)}{\log(1/\delta_N)} + \frac{\mathbb{E}\left[R(w^*_{S|J}) - R(w^*) \right]}{2} + \frac{64\kappa G^2}{N\mu},$$

where in the first inequality we have used Cauchy-Schwarz inequality, and in the last inequality we have used the fact $\sqrt{ab} \leq \frac{a}{2t} + \frac{bt}{2}$ for any $a, b, t > 0$. Rearranging both sides of the above inequality

with simple algebra leads to

$$
\begin{aligned}
&\mathbb{E}\left[R(w_{S|J}^*) - R(w^*)\right] \\
\leq& 2\mathbb{E}\left[R_S(w_{S|J}^*) - R_S(w^*)\right] + \frac{\log(N)}{N}\left(\frac{16G^2}{\mu_k} + 8GD\right) + \frac{16GD\log^2(N)}{\log(1/\delta_N)} + \frac{128\kappa G^2}{N\mu} \\
=& 2\mathbb{E}\left[R_S(w_{S|J}^*) - R_S(w_{\bar{k}}^*) + R_S(w_{\bar{k}}^*) - R_S(w^*)\right] \\
&+ \frac{\log(N)}{N}\left(\frac{16G^2}{\mu_k} + 8GD\right) + \frac{16GD\log^2(N)}{\log(1/\delta_N)} + \frac{128\kappa G^2}{N\mu} \\
=& 2\mathbb{E}\left[R_S(w_{S|J}^*) - R_S(w_{\bar{k}}^*)\right] + 2(R(w_{\bar{k}}^*) - R(w^*)) \\
&+ \frac{\log(N)}{N}\left(\frac{16G^2}{\mu_k} + 8GD\right) + \frac{16GD\log^2(N)}{\log(1/\delta_N)} + \frac{128\kappa G^2}{N\mu} \\
\leq& 2\mathbb{E}\left[R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*)\right] + 2(R(w_{\bar{k}}^*) - R(w^*)) \\
&+ \frac{\log(N)}{N}\left(\frac{16G^2}{\mu_k} + 8GD\right) + \frac{16GD\log^2(N)}{\log(1/\delta_N)} + \frac{128\kappa G^2}{N\mu} \\
\leq& 2\underbrace{\mathbb{E}\left[\left(R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*)\right)_+\right]}_{T_1} + \underbrace{2\left(R(w_{\bar{k}}^*) - R(w^*)\right)}_{T_2} \\
&+ \underbrace{\frac{\log(N)}{N}\left(\frac{16G^2}{\mu_k} + 8GD\right) + \frac{16GD\log^2(N)}{\log(1/\delta_N)} + \frac{128\kappa G^2}{N\mu}}_{T_3}.
\end{aligned}
\tag{26}
$$

Plugging equation 26 into equation 25 yields that for any $q \geq 2$

$$
\begin{aligned}
&\left\|R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*)\right\|_q \\
\leq& \frac{q\log(N)}{N}\left(\frac{4G^2}{\mu_k} + 2GD\right) + \frac{2GDq^2\log^2(N)}{\log(1/\delta_N)} + 8G\sqrt{\frac{\kappa q(T_1 + T_2 + T_3)}{N\mu}} \\
\leq& \frac{q\log(N)}{N}\left(\frac{4G^2}{\mu_k} + 2GD\right) + \frac{2GDq^2\log^2(N)}{\log(1/\delta_N)} + 8G\sqrt{\frac{\kappa q(T_1 + T_3)}{N\mu}} + 8G\sqrt{\frac{\kappa q T_2}{N\mu}} \\
\overset{\zeta_1}{\leq}& \frac{q\log(N)}{N}\left(\frac{4G^2}{\mu_k} + 2GD\right) + \frac{2GDq^2\log^2(N)}{\log(1/\delta_N)} + 8G\sqrt{\frac{\kappa q T_2}{N\mu}} + T_1 + T_3 + \frac{16\kappa q G^2}{N\mu} \\
\lesssim& \frac{q\log(N)}{N}\left(\frac{G^2}{\mu_k} + GD\right) + \frac{GDq^2\log^2(N)}{\log(1/\delta_N)} + \frac{qG^2}{N\mu} + \mathbb{E}\left[\left(R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*)\right)_+\right] \\
&+ G\sqrt{\frac{q\left(R(w_{\bar{k}}^*) - R(w^*)\right)}{N\mu}},
\end{aligned}
\tag{27}
$$

where in "ζ_1" we have again used the fact $\sqrt{ab} \leq \frac{a}{2t} + \frac{bt}{2}$ for $a, b, t > 0$. Now we are in the position to finally upper bound the desired sparse excess risk with respect to $w_{\bar{k}}^*$, which can be decomposed

in the following way:

$$
\begin{aligned}
R(w_{S|J}^*) - R(w_{\bar{k}}^*) =& R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*) + R_S(w_{S|J}^*) - R_S(w_{\bar{k}}^*) \\
& + R_S(w_{\bar{k}}^*) - R_S(w^*) + R(w^*) - R(w_{\bar{k}}^*) \\
\leq& R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*) + R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*) \\
& + R_S(w_{\bar{k}}^*) - R_S(w^*) + R(w^*) - R(w_{\bar{k}}^*) \\
\leq& \underbrace{R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*)}_{T_1'} + \left(R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*) \right)_+ \\
& + \underbrace{R_S(w_{\bar{k}}^*) - R_S(w^*) + R(w^*) - R(w_{\bar{k}}^*)}_{T_2'},
\end{aligned}
\tag{28}
$$

where in the first inequality we have used the fact $R_S(w_{S,\bar{k}}^*) \leq R_S(w_{\bar{k}}^*)$. We are going to bound the above two terms T_1' and T_2' respectively with high probability. Concerning T_1', since equation 27 holds for all $q \geq 2$, in view of the second part of Lemma 4 (with $q_l = 2$ and $q_u = \infty$) we can show that the following holds with probability at least $1 - \frac{\delta}{2}$ for any $\delta \in (0, e^{-1})$:

$$
\begin{aligned}
T_1' =& R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*) \\
\leq& \left| R(w_{S|J}^*) - R(w^*) - R_S(w_{S|J}^*) + R_S(w^*) \right| \\
\lesssim& \frac{GD \log^2\left(\frac{e}{\delta}\right) \log^2(N)}{\log(1/\delta_N)} + \left(\log(N) \left(\frac{G^2}{\mu_k} + GD \right) + \frac{G^2}{\mu} \right) \frac{\log\left(\frac{e}{\delta}\right)}{N} + G\sqrt{\frac{\log\left(\frac{e}{\delta}\right)\left(R(w_{\bar{k}}^*) - R(w^*)\right)}{N\mu}} \\
& + \mathbb{E}\left[\left(R_S(w_{S|J}^*) - R_S(w_{S,\bar{k}}^*) \right)_+ \right].
\end{aligned}
\tag{29}
$$

Regarding the term T_2', similar to the argument of equation 24, we can bound its L_q-norm for any $q \geq 2$ as follows:

$$
\begin{aligned}
\|T_2'\|_q =& \left\| \frac{1}{N} \sum_{i=1}^N \left(\ell(w_{\bar{k}}^*; Z_i) - \ell(w^*; Z_i) \right) - \left(R(w_{\bar{k}}^*) - R(w^*) \right) \right\|_q \\
\leq& \frac{2\sqrt{2\kappa q}}{N} \sqrt{\left\| \sum_{i=1}^N \left(\ell(w_{\bar{k}}^*; Z_i) - \ell(w^*; Z_i) - \left(R(w_{\bar{k}}^*) - R(w^*) \right) \right)^2 \right\|_{q/2}} \\
\leq& \frac{2\sqrt{2\kappa q}}{N} \sqrt{\sum_{i=1}^N \left\| \left(\ell(w_{\bar{k}}^*; Z_i) - \ell(w^*; Z_i) \right)^2 + \left(R(w_{\bar{k}}^*) - R(w^*) \right)^2 \right\|_{q/2}} \\
\leq& \frac{2\sqrt{2\kappa q}}{N} \sqrt{\sum_{i=1}^N \left\| \left(\ell(w_{\bar{k}}^*; Z_i) - \ell(w^*; Z_i) \right)^2 \right\|_{q/2} + N \left(R(w_{\bar{k}}^*) - R(w^*) \right)^2} \\
\leq& \frac{4\sqrt{\kappa q}}{N} \sqrt{N G^2 \left\| w_{\bar{k}}^* - w^* \right\|^2} \\
\leq& \frac{4G\sqrt{2\kappa q}}{\sqrt{N\mu}} \sqrt{R(w_{\bar{k}}^*) - R(w^*)}.
\end{aligned}
$$

By invoking the second part of Lemma 4 with $q_l = 2$ and $q_u = \infty$ we can translate the above moment bound into the following exponential tail bound that holds with probability at least $1 - \frac{\delta}{2}$ for any $\delta \in (0, e^{-1})$:

$$
T_2' \lesssim G\sqrt{\frac{\log\left(\frac{e}{\delta}\right)\left(R(w_{\bar{k}}^*) - R(w^*)\right)}{N\mu}}.
\tag{30}
$$

By plugging equation 29 and equation 30 into equation 28 and applying union probability argument we obtain that the following holds with probability at least $1 - \delta$ for any $\delta \in (0, e^{-1})$:

$$R(w^*_{S|J}) - R(w^*_{\bar{k}})$$

$$\lesssim \frac{GD \log^2\left(\frac{e}{\delta}\right) \log^2(N)}{\log(1/\delta_N)} + \left(\log(N)\left(\frac{G^2}{\mu_k} + GD\right) + \frac{G^2}{\mu}\right)\frac{\log\left(\frac{e}{\delta}\right)}{N} + G\sqrt{\frac{\log\left(\frac{e}{\delta}\right)\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N\mu}}$$

$$+ \left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}})\right)_+ + \mathbb{E}\left[\left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}})\right)_+\right].$$

This proves the desired sparse excess risk bound over the supporting set J.

As the final step, since there are at most $\binom{d}{k} \leq \left(\frac{ed}{k}\right)^k$ different J with $|J| = k$, by union probability we can show that the following holds with probability $1 - \delta$ for any $\delta \in (0, e^{-1})$:

$$\sup_{J \subseteq [d], |J| = k} R(w^*_{S|J}) - R(w^*_{\bar{k}})$$

$$\lesssim \frac{GD\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)^2 \log^2(N)}{\log(1/\delta_N)} + \left(\log(N)\left(\frac{G^2}{\mu_k} + GD\right) + \frac{G^2}{\mu}\right)\frac{k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)}{N}$$

$$+ G\sqrt{\frac{\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N\mu}} + \left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}})\right)_+$$

$$+ \mathbb{E}\left[\left(R_S(w^*_{S|J}) - R_S(w^*_{S,\bar{k}})\right)_+\right].$$

This completes the proof. $\qquad\square$

Now we are in the position to prove the main result in Theorem 4.

Proof of Theorem 4. Let us consider $\tilde{J} = \text{supp}(\tilde{w}_{S,k})$. By definition we have $\tilde{w}_{S,k} = w^*_{S|\tilde{J}}$. Then invoking Lemma 6 yields that for any $\delta \in (0, e^{-1})$, the following holds with probability at least $1 - \delta$:

$$R(\tilde{w}_{S,k}) - R(w^*_{\bar{k}}) = R(w^*_{S|\tilde{J}}) - R(w^*_{\bar{k}})$$

$$\lesssim \frac{GD\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)^2 \log^2(N)}{\log(1/\delta_N)} + \left(\log(N)\left(\frac{G^2}{\mu_k} + GD\right) + \frac{G^2}{\mu}\right)\frac{k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)}{N}$$

$$+ G\sqrt{\frac{\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N\mu}} + \left(R_S(w^*_{S|\tilde{J}}) - R_S(w^*_{S,\bar{k}})\right)_+$$

$$+ \mathbb{E}\left[\left(R_S(w^*_{S|\tilde{J}}) - R_S(w^*_{S,\bar{k}})\right)_+\right].$$

$$\lesssim \frac{GD\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)^2 \log^2(N)}{\log(1/\delta_N)} + \left(\log(N)\left(\frac{G^2}{\mu_k} + GD\right) + \frac{G^2}{\mu}\right)\frac{k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)}{N}$$

$$+ G\sqrt{\frac{\left(k\log\left(\frac{ed}{k}\right) + \log\left(\frac{e}{\delta}\right)\right)\left(R(w^*_{\bar{k}}) - R(w^*)\right)}{N\mu}} + \Delta_{\bar{k},\text{opt}} + \mathbb{E}\left[\Delta_{\bar{k},\text{opt}}\right].$$

This proves the desired sparse excess risk bound. $\qquad\square$