For high-dimensional hierarchical models, consider exchangeability of effects across covariates instead of across datasets

Abstract

When facing multiple related regression problems, it is standard practice to model effects as (1) exchangeable across datasets and (2) correlated to differing degrees across covariates. But we show that this classical approach can exhibit poor statistical performance when the number of covariates exceeds the number of datasets. For instance, in statistical genetics, we might regress dozens of traits (defining datasets) for thousands of individuals (responses) on up to millions of genetic variants (covariates). When an analyst has more covariates than datasets, we argue that it is often more natural to instead model effects as (1) exchangeable across *covariates* and (2) correlated to differing degrees across *datasets*. To this end, we propose a hierarchical model expressing our alternative perspective. We devise an empirical Bayes estimator for learning the degree of correlation between datasets. We show via theory and empirics that our method outperforms the classic approach when the number of covariates dominates the number of datasets.

1 Introduction

Hierarchical modeling is a mainstay of Bayesian inference. For instance, in (generalized) linear models, the unknown parameters are *effects*, each of which describes the association of a particular covariate with a response of interest. Often covariates are shared across multiple related datasets, but the effects are typically allowed to vary both by dataset and by covariate. A classic methodology, dating back to Lindley and Smith (1972) [43], models the effects as conditionally independent across datasets, with a latent (and learnable) degree of relatedness across covariates. From a practical standpoint, the model is motivated by the understanding that it "borrows strength" across different datasets [24, Chapter 5.6]. Mathematically, the model is motivated by assuming effects are exchangeable across datasets and applying a de Finetti theorem [43, 35]. The methodology of Lindley and Smith is ubiquitous when the number of datasets is larger than the number of covariates. It is a standard component of Bayesian pedagogy [[23, Chapter 13.3]; [24, Chapter 15.4]] and software; e.g. it is used by default in the mixed modeling package lme4 [5], which has over 13 million downloads at the time of writing.

Despite its resounding success when there are more datasets than covariates, we show in the present work that the approach of Lindley and Smith performs poorly when there are more covariates than datasets. To address the many-covariates case, we turn for inspiration to statistical genetics, where scientists commonly learn linear models relating genetic variants (covariates) to traits (corresponding to different datasets) across individuals (which each exhibit a response). These applications may exhibit millions of covariates, thousands of responses, and just a handful of datasets. In these cases, [39, 12, 54, 66, 45, 51] use a multivariate Gaussian prior akin to that of Lindley and Smith, but instead

assume conditional independence across *covariates* and prior parameters that encode correlations across *datasets*.

We can imagine using a similar model in applications beyond statistical genetics. Namely, when there are more covariates than datasets, we propose to model the effects as exchangeable across *covariates* (rather than datasets) and learn the degree of relatedness of effects across *datasets* (rather than covariates). Henceforth we refer to this framework as *ECov*, for *e*xchangeable effects across *cov*ariates, and distinguish it from *e*xchangeable effects across *data*sets, or *EData*. Though the ECov framework seems natural, to the best of our knowledge it is absent from existing literature on hierarchical Bayesian regression. Brown and Zidek (1980) [10] and Haitovsky (1987) [28] form two exceptions, but these two papers (1) consider only the situation in which a single covariate matrix is shared across all datasets (or equivalently, for each data point all responses are observed) and (2) include only theory and no empirics. We suspect that the historical origins of the statistical genetics models above have hindered their expansion to a wider audience. In particular, the statistical genetics literature traces back to mixed effects modeling for cattle breeding [55]; here, an even-earlier notion of the genetic contribution of trait correlation (i.e. "genetic correlation;" see Hazel (1943) [29]) informs the covariance structure of random effects. Although genetic correlation is now commonly understood to describe the correlation of effects of DNA sequence changes on different traits [12], its provenance predates even the first identification of DNA as the genetic material in 1944 [3]. See Appendix A for further discussion of related work.

In the present work, we propose ECov as a useful framework for hierarchical regression when the number of covariates exceeds the number of datasets. We show that the classic model structure from statistical genetics can be seen as an instance of this framework, much as Lindley and Smith give a (complementary) instance of an EData framework. To make the ECov approach generally practical, we devise an accurate and efficient algorithm for learning the dataset-correlation matrix. We demonstrate with theory and empirics that ECov is preferred when the number of covariates exceeds the number of datasets, while EData is preferred when the number of datasets exceeds the number of covariates. Our experiments analyze three real, non-genetic datasets in regression and classification, including an application to transfer learning with pre-trained neural network embeddings. We provide proofs of theoretical results in the appendix.

2 Exchangeability and its applications to hierarchical linear modeling

We start by establishing the data and model, motivating exchangeability among covariate effects (ECov), and motivating our Bayesian generative model.

Setup and notation. Consider Q datasets with D covariates. Let N^q be the number of data points in dataset q. For the qth dataset, the $N^q \times D$ real design matrix X^q collects the covariates, and Y^q is the N^q-vector of responses. The nth datapoint in dataset q consists of covariate D-vector X_n^q and scalar response Y_n^q. We let $\mathcal{D} := \{(X^q, Y^q)\}_{q=1}^Q$ denote the collection of all Q datasets. We take the generalized linear model $Y_n^q | X_n^q, \beta^q \overset{indep}{\sim} p(\cdot | X_n^{q\top} \beta^q)$ with unknown D-vector of real effects β^q. We collect all effects in a $D \times Q$ matrix β with (d, q) entry β_d^q. The linear form of the likelihood allows interpretation of β_d^q as the association between the dth covariate and the response in dataset q. In linear regression, the responses are real-valued and the conditional distribution is Gaussian. In logistic regression, the responses are binary, and we use the logit link. The independence assumption conflicts with some models that one might use, for example in some cases when the different datasets partially overlap.

Example. As a motivating non-genetics example, consider a study of the efficacy of microcredit. There are seven famous randomized controlled trials of microcredit, each in a different country [47]. We might be interested in the association between various aspects of small businesses (covariates), including whether or not they received microcredit, and their business profit (response). In this case, the dth element of X_n^q would be the dth characteristic of the nth small business in the qth country, and Y_n^q is the profit of this business. See the experiments for additional examples in rates of policing, web analytics, and transfer learning.

Exchangeable effects across datasets (EData). To fully specify a Bayesian model, we need to choose a prior over the parameters β. Lindley and Smith assume the effects are exchangeable across datasets. Namely, for every Q-permutation σ, $p(\beta^1, \beta^2, \ldots, \beta^Q) = p(\beta^{\sigma(1)}, \beta^{\sigma(2)}, \ldots, \beta^{\sigma(Q)})$.

88 Assuming exchangeability holds for an imagined growing Q and applying de Finetti's theorem

89 motivates a conditionally independent prior. Concretely, Lindley and Smith take $\beta^q \overset{i.i.d.}{\sim} \mathcal{N}(\xi, \Gamma)$, for

90 D-vector ξ and $D \times D$ covariance matrix Γ. The (d, d') entry of Γ captures the degree of relatedness

91 between the effects for covariates d and d'. Both ξ and Γ may be learned in an empirical Bayes

92 procedure.

93 **Exchangeable effects across covariates (ECov).** We here argue for a complementary approach

94 in settings where $D > Q$. In the microcredit example, notice that $D > Q$ will arise whenever

95 the experimenter records more characteristics of a small business than there are locations with

96 microcredit experiments; that is, $D > 7$ in this particular case. Concretely, let β_d be the Q-vector

97 of effects for covariate d across datasets. Then, in the ECov approach, we will assume that effects

98 are exchangeable across covariates instead of across datasets. Namely, for every D-permutation σ,

99 $p(\beta_1, \beta_2, \ldots, \beta_D) = p(\beta_{\sigma(1)}, \beta_{\sigma(2)}, \ldots, \beta_{\sigma(D)})$. We will see theoretical and empirical benefits to

100 ECov in later sections, but note that the ECov assumption is often a priori natural. For instance,

101 regarding microcredit, we may have no a priori beliefs about how effects differ for distinct small-

102 business characteristics. And we may a priori believe that different countries could exhibit more

103 similar effects – and wish to learn the degree of relatedness across those countries.

104 We may apply a similar rationale as Lindley and Smith to motivate a conditionally independent

105 model. Analogous to Lindley and Smith, we propose a Gaussian prior: $\beta_d \overset{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$. Σ is now

106 a $Q \times Q$ covariance matrix whose (q, q') entry captures the similarity between the effects in the q

107 and q' datasets. For simplicity, we restrict to $\mathbb{E}[\beta_d] = 0$; see Appendix E.3 for discussion. Another

108 potential benefit to ECov relative to EData is that we might expect a statistically easier problem, with

109 $O(Q^2)$ rather than $O(D^2)$ values to learn in the relatedness matrix. We provide a rigorous theoretical

110 analysis in Sections 4 and 5.

3 Our method

112 We next describe our inference method for specific instances of the exchangeable covariate effects

113 model of Section 2. We compute the β posterior and take an empirical Bayes approach to estimate Σ.

114 We find that an expectation maximization (EM) algorithm estimates Σ effectively; Appendix A.2

115 compares our approach to existing methods for the related problem of estimating Γ for EData.

116 **Notation.** We identify estimates of β and Σ with hats. For instance, $\hat{\beta}_{\text{LS}}$ is the least squares

117 estimate, with $\hat{\beta}_{\text{LS}}^q := (X^{q\top} X^q)^{-1} X^{q\top} Y^q$. We will sometimes find it useful to stack the columns

118 of β or its estimates into a length DQ vector; we denote such vectors with an arrow; for example,

119 $\vec{\beta} := [\beta^{1\top}, \beta^{2\top}, \ldots, \beta^{Q\top}]^\top$. For a natural number N, we use I_N, $\mathbf{1}_N$, and e_N to denote the $N \times N$

120 identity matrix, N-vector of ones, and Nth basis vector, respectively. We use $\otimes$ to denote the

121 Kronecker product.

3.1 Posterior inference with a Gaussian likelihood

123 We first consider a Gaussian likelihood: for each dataset q and observation n, we take $Y_n^q | X_n^q, \beta^q \overset{indep}{\sim}$

124 $\mathcal{N}(X_n^{q\top} \beta^q, \sigma_q^2)$ where σ_q^2 is a dataset-specific variance. When the relatedness matrix Σ is known,

125 a natural estimate of β is its posterior mean. We obtain the full posterior, including its mean, via a

126 standard conjugacy argument; see Appendix B.1:

Proposition 3.1. *For each covariate d, let $\beta_d \overset{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$ a priori. For each dataset q*

128 *and data point n, let $Y_n^q | X_n^q, \beta^q \overset{indep}{\sim} \mathcal{N}(X_n^{q\top} \beta^q, \sigma_q^2)$. Then $\vec{\beta} | \mathcal{D}, \Sigma \sim \mathcal{N}(\vec{\mu}, V)$ for $\vec{\mu} =$*

129 *$V[\sigma_1^{-2} Y^{1\top} X^1, \ldots, \sigma_Q^{-2} Y^{Q\top} X^Q]^\top$ and $V^{-1} = \Sigma^{-1} \otimes I_D + \text{diag}(\sigma_1^{-2} X^{1\top} X^1, \ldots, \sigma_Q^{-2} X^{Q\top} X^Q)$,*

130 *where $\text{diag}(\sigma_1^{-2} X^{1\top} X^1, \ldots, \sigma_Q^{-2} X^{Q\top} X^Q)$ denotes a $DQ \times DQ$ block-diagonal matrix.*

131 At first glance, the posterior mean $\vec{\mu}$ for this model might seem to introduce a computational challenge

132 because exact computation of V involves an $O(D^3 Q^3)$-time matrix inversion. Our experiments

133 (Section 6), however, involve on the order of $DQ \approx 1,000$ parameters, so direct inversion of V

134 demands less than a single second. Moreover, in much larger problems $\vec{\mu}$ may still be computed

135 very efficiently using the conjugate gradient algorithm [48, Chapter 5], with convergence in a small

136 number of $O(D^2 Q)$ time iterations; see Appendix B.2.

Algorithm 1 Expectation Maximization for Exchangeability Among Covariate Effects

1: // Initialize covariance
2: $\Sigma^{(0)} \leftarrow I_Q$
3: // Run EM algorithm
4: **for** $t = 0, 1, \ldots$ **do**
5:    // Expectation step
6:    $\mu_1, \ldots, \mu_D, V_1, \ldots, V_D \leftarrow \texttt{E\_Step}(\Sigma^{(t)})$
7:
8:    // Maximization step
9:    $\Sigma^{(t+1)} \leftarrow D^{-1} \sum_{d=1}^{D} (\mu_d \mu_d^\top + V_d)$
10:
11: Return $\Sigma^{(t+1)}$

Algorithm 2 E-Step: Linear Regression

1: $\vec{\mu}, V \leftarrow \mathbb{E}[\vec{\beta}|\mathcal{D}, \Sigma], \mathrm{Var}[\vec{\beta}|\mathcal{D}, \Sigma]$
2: **for** $d = 1, \ldots, D$ **do**
3:    $\mu_d \leftarrow (e_d \otimes I_Q)^\top \vec{\mu}$
4:    $V_d \leftarrow (e_d \otimes I_Q)^\top V (e_d \otimes I_Q)$
5: Return $\mu_1, \ldots, \mu_D, V_1, \ldots, V_D$

Algorithm 3 E-Step: Logistic Regression

1: $\vec{\mu}^* \leftarrow \arg\max_{\vec{\beta}} \log p(\vec{\beta}|\mathcal{D}, \Sigma)$
2: $V \leftarrow -[\nabla_{\vec{\beta}}^2 \log p(\vec{\beta}|\mathcal{D}, \Sigma)|_{\vec{\beta}=\vec{\mu}^*}]^{-1}$
3: **for** $d = 1, \ldots, D$ **do**
4:    $\mu_d \leftarrow (e_d \otimes I_Q)^\top \vec{\mu}^*$
5:    $V_d \leftarrow (e_d \otimes I_Q)^\top V (e_d \otimes I_Q)$
6: Return $\mu_1, \ldots, \mu_D, V_1, \ldots, V_D$

3.2 Empirical Bayes estimation of Σ by expectation maximization

The posterior mean of β in Proposition 3.1 requires Σ, which is typically unknown. Accordingly, we propose an empirical Bayes approach of estimating Σ by maximum marginal likelihood:

$$\hat{\beta}_{\mathrm{ECov}} := \mathbb{E}[\beta \mid \mathcal{D}, \hat{\Sigma}] \text{ where } \hat{\Sigma} := \arg\max_{\Sigma \succeq 0} p(\mathcal{D} \mid \Sigma). \tag{1}$$

Equation (1) defines a two step procedure. In the first step, we learn the similarity between datasets via estimation of Σ. In the second step, we use this similarity to compute an estimate, $\hat{\beta}_{\mathrm{ECov}}$, that correspondingly shares strength. Though we have been unable to identify a general analytic form for $\hat{\Sigma}$, we can compute it with an expectation maximization (EM) algorithm [46, Chapter 1.5]. Algorithm 1 summarizes this procedure; see Appendix B.3 for details.

3.3 Classification with logistic regression

We can extend the approach above to inference for multiple related classification problems. We assume a logistic likelihood; for each q and n, $Y_n^q|X_n^q, \beta^q \overset{indep}{\sim} \mathrm{Bern}[(1 + \exp\{-X_n^{q\top}\beta^q\})^{-1}]$. In the classification case, we cannot use Gaussian conjugacy directly, so we apply an approximation. Specifically, we adapt the original E-step in Algorithm 3 by using a Laplace approximation to the posterior [7, Chapter 4.4]. We approximate the posterior mean of β by the maximum a posteriori value. We leave extensions to other generalized linear models to future work.

4 Theoretical comparison of frequentist risk

In this section, we prove theory that suggests ECov has better frequentist risk than EData when D is large relative to Q. Analyzing $\hat{\beta}_{\mathrm{ECov}}$ directly is challenging due to its non-differentiability as a function of the data, so we take a multipart approach. First, in Theorem 4.3, we show that an ECov estimate based on moment-matching (MM), $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$, dominates least squares, $\hat{\beta}_{\mathrm{LS}}$, when D is large relative to Q; $\hat{\beta}_{\mathrm{LS}}$ in turn dominates $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ (a similar estimator for EData). Second, in Theorem 4.5, we show that $\hat{\beta}_{\mathrm{ECov}}$ uniformly improves on $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$.

Setup. Take a fixed value of β and an estimator $\hat{\beta}$. We use squared error risk, $\mathrm{R}(\beta, \hat{\beta}) := \mathbb{E}\left[\|\hat{\beta} - \beta\|_F^2 \mid \beta\right]$, as our measure of performance. $\| \cdot \|_F$ is the Frobenius norm of a matrix, and the expectation is over all observations $Y^1, \ldots, Y^Q$ jointly. We require the following orthogonal design condition.

Condition 4.1. *For each dataset q, $\sigma_q^{-2} X^{q\top} X^q = \sigma^{-2} I_D$ for some shared variance σ^2.*

164 Though restrictive, this condition is useful for theory, as other authors have found; see Appendix C.1.
165 We empirically demonstrate that our theoretical conclusions apply more broadly in Section 6.

166 **ECov vs. EData when using moment matching in high dimensions** For ECov, we show the
167 following estimate for Σ is unbiased under correct prior specification: $\hat{\Sigma}^{\mathrm{MM}} := D^{-1}\hat{\beta}_{\mathrm{LS}}^{\top}\hat{\beta}_{\mathrm{LS}} -$
168 $D^{-1}\mathrm{diag}(\sigma_1^2\|X^{1\dagger}\|_F^2, \ldots, \sigma_Q^2\|X^{Q\dagger}\|_F^2)$, where $\dagger$ denotes the Moore-Penrose pseudoinverse of a
169 matrix. We define $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}} := \mathbb{E}[\beta|\mathcal{D}, \hat{\Sigma}^{\mathrm{MM}}]$ to be the resulting parameter estimate. We define
170 $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ analogously for EData; see Appendix C.2 for details. While $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ are naturally
171 defined only when $D \geq Q$ and $D \leq Q$, respectively, we find it informative to compare how their
172 performances depend on D and Q nonetheless.

173 Before our theorem, a lemma provides concise expressions for the risks of $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$.

174 **Lemma 4.2.** *Under Condition 4.1 and when* $D \geq Q$, $\mathrm{R}(\beta, \hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}) = \sigma^2 DQ - \sigma^4 D(D -$
175 $2 - 2Q)\mathbb{E}[\|\hat{\beta}_{\mathrm{LS}}^{\dagger}\|_F^2 \mid \beta]$. *Additionally, when* $D \leq Q$, $\mathrm{R}(\beta, \hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}) = \sigma^2 DQ - \sigma^4 Q(Q - 2 -$
176 $2D)\mathbb{E}[\|\hat{\beta}_{\mathrm{LS}}^{\dagger}\|_F^2 \mid \beta]$.

177 Lemma 4.2 reveals forms for the risks of $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ that are surprisingly simple. The
178 symmetry between the forms and risks of these estimators, however, is intuitive; under Condition 4.1,
179 $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ can be seen as respectively arising from the same procedure applied to $\hat{\beta}_{\mathrm{LS}}$ and its
180 transpose.

181 With Lemma 4.2 in hand, we can now compare the risk of $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$, $\hat{\beta}_{\mathrm{LS}}$, and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$.

182 **Theorem 4.3.** *Let Condition 4.1 hold. Then (1) if* $D > 2Q + 2$, $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ *dominates* $\hat{\beta}_{\mathrm{LS}}$ *with respect*
183 *to squared error risk. In particular, for any* β, $\mathrm{R}(\beta, \hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}) < \mathrm{R}(\beta, \hat{\beta}_{\mathrm{LS}})$. *Additionally, (2) if*
184 $D > Q/2 - 1$, $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$ *is dominated by* $\hat{\beta}_{\mathrm{LS}}$.

185 Since $\hat{\beta}_{\mathrm{LS}}$ is minimax [40, Chapter 5], Theorem 4.3 implies that $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ has minimax risk in the
186 high-dimensional setting. It follows that, regardless of how well the ECov prior assumptions hold,
187 $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ will not perform very poorly.

188 **Further improvement with maximum marginal likelihood.** The moment based approach analyzed
189 above has a limitation: with positive probability, $\hat{\Sigma}^{\mathrm{MM}}$ is not positive semi-definite (PSD). Though
190 our expression for $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ remains well-defined in this case, this non-positive definiteness obscures
191 the interpretation of $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ as a Bayes estimate. We next show that performance further improves if
192 Σ is estimated by maximum marginal likelihood (Equation (1)) and is thereby constrained to be PSD.

193 Our next lemma characterizes the form of the resulting estimator, $\hat{\beta}_{\mathrm{ECov}}$, and establishes a connection
194 to the positive part James-Stein estimator [4].

195 **Lemma 4.4.** *Assume* $D > Q$ *and consider the singular value decomposition* $\hat{\beta}_{\mathrm{LS}} = V\mathrm{diag}(\lambda^{\frac{1}{2}})U^{\top}$
196 *where* V *and* U *satisfy* $V^{\top}V = U^{\top}U = I_Q$, *and* λ *is a* Q-vector *of non-negative reals. Un-*
197 *der Condition 4.1, Equation* (1) *reduces to* $\hat{\Sigma} = U\mathrm{diag}\left[(D^{-1}\lambda - \sigma^2\mathbf{1}_Q)_+\right]U^{\top}$ *and* $\hat{\beta}_{\mathrm{ECov}} =$
198 $V\mathrm{diag}\left[\lambda^{\frac{1}{2}} \odot (\mathbf{1}_Q - \sigma^2 D\lambda^{-1})_+\right]U^{\top}$, *where* $(\cdot)_+$ *is shorthand for* $\max(\cdot, 0)$ *element-wise,* $\odot$ *is the*
199 *Hadamard (i.e. element-wise) product, and the powers in* $\lambda^{\frac{1}{2}}$ *and* λ^{-1} *are applied element-wise.*

200 Lemma 4.4 allows us to see $\hat{\beta}_{\mathrm{ECov}}$ as shrinking $\hat{\beta}_{\mathrm{LS}}$ toward 0 in the direction of each singular vector
201 to an extent proportional to the inverse of the associated singular value. The transition from $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$
202 to $\hat{\beta}_{\mathrm{ECov}}$ is then analogous to the taking the "positive part" of the James-Stein estimator in vector
203 estimation, which provides a uniform improvement in risk [4]. Though $\mathrm{R}(\beta, \hat{\beta}_{\mathrm{ECov}})$ is not easily
204 available analytically, we nevertheless find that it dominates its moment-based counterpart.

205 **Theorem 4.5.** *Assume* $D > Q + 1$. *Under Condition 4.1* $\hat{\beta}_{\mathrm{ECov}}$ *dominates* $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ *with respect to*
206 *squared error loss, achieving strictly lower risk for every value of* β.

207 We establish Theorem 4.5 using a proof technique adapted from Baranchik [4]; see also Lehmann
208 and Casella [40][Thm. 5.5.4]. The standard approach we build upon is complicated by the fact that
209 the directions in which we apply shrinkage are themselves random.

Theorem 4.5 provides a strong line of support for using $\hat{\beta}_{\text{ECov}}$ over $\hat{\beta}_{\text{ECov}}^{\text{MM}}$ that does not rely on any assumption of "correct" prior specification; in particular the risk improves without any subjective assumptions on β. We discuss related earlier work in Appendix A.3.

5 Gains from ECov in the high-dimensional limit

The results of Section 4 give a promising endorsement of ECov but face two important limitations. First, the domination results relative to least squares do not directly demonstrate that $\hat{\beta}_{\text{ECov}}$ attain improvements by leveraging similarities across datasets in a meaningful way; indeed for a single dataset (i.e. $Q = 1$) $\hat{\beta}_{\text{ECov}}$ can be understood as a ridge regression estimate [31], and Theorems 4.3 and 4.5 provide that $\hat{\beta}_{\text{ECov}}$ dominates $\hat{\beta}_{\text{LS}}$ for $D > 3$. Second, domination results reveal nothing about the size of the improvement or how it depends on any structure of β; intuitively, we should expect better performance when β is in some way representative of the assumed prior. To address these limitations, we analyze the size of the gap between the risk of (1) $\hat{\beta}_{\text{ECov}}$ and (2) our method applied to each dataset independently, which we denote by $\hat{\beta}_{\text{ID}}$. We will characterize the dependence of this gap on β.

Reasoning quantitatively about the dependence of the risk on the unknown parameter poses significant analytical challenges. In particular, Lemma 4.2 shows that $\text{R}(\beta, \hat{\beta}_{\text{ECov}}^{\text{MM}})$ depends on β through $\mathbb{E}[\|\hat{\beta}_{\text{LS}}^{\dagger}\|_F^2|\beta]$; however, $\|\hat{\beta}_{\text{LS}}^{\dagger}\|_F^2$ is the sum of the eigenvalues of a non-central inverse Wishart matrix, a notoriously challenging quantity to work with; see e.g. [41, 30]. To regain tractability, we (1) develop an analysis asymptotic in the number of covariates D and (2) shift to a Bayesian analysis in order to sensibly consider a growing collection of covariate effects. In particular, we consider a sequence of regression problems, with parameters $\{\beta_d\}_{d=1}^{\infty}$ distributed as $\beta_d \overset{i.i.d.}{\sim} \pi$ for some distribution π. Accordingly, instead of using the frequentist risk as in Section 4, we now use the Bayes risk to measure performance. Specifically, for a dataset with D covariates and an estimator $\hat{\beta}$, the Bayes risk is $\text{R}_\pi^D(\hat{\beta}) := \mathbb{E}_\pi[\text{R}(\beta, \hat{\beta})]$ where $\text{R}(\beta, \hat{\beta})$ is the usual frequentist risk.

For a single metric characterizing the benefits of joint modeling, we will define the *asymptotic gain* as the relative performance between our two estimators of interest here, $\hat{\beta}_{\text{ECov}}$ and $\hat{\beta}_{\text{ID}}$.

Definition 5.1. *Consider a sequence of datasets of Q regression problems with an increasing number of covariates $D, \{\mathcal{D}_D\}_{D=1}^{\infty}$. Assume that for each dataset Condition 4.1 holds with variance σ^2 and that each $\beta_d \overset{i.i.d.}{\sim} \pi$. The asymptotic gain of joint modeling is $\text{Gain}(\pi, \sigma^2) := \lim_{D \to \infty} (\sigma^2 DQ)^{-1}[\text{R}_\pi^D(\hat{\beta}_{\text{ID}}) - \text{R}_\pi^D(\hat{\beta}_{\text{ECov}})]$.*

The factor of $\sigma^2 DQ$ in Definition 5.1 puts $\text{Gain}(\pi, \sigma^2)$ on a scale that is roughly invariant to the size and noise level of the problem; for example, $(\sigma^2 DQ)^{-1}\text{R}_\pi^D(\hat{\beta}_{\text{LS}}) = 1$ for any π.

Our next lemma gives an analytic expression for $\text{Gain}(\pi, \sigma^2)$ that provides a starting point for understanding its problem dependence.

Lemma 5.2. *Assume $\tilde{\Sigma} := \text{Var}_\pi[\beta_1]$ is finite and has eigenvalues $\lambda_1, \ldots, \lambda_Q$. Under Condition 4.1, $\text{Gain}(\pi, \sigma^2) = \sigma^2 Q^{-1}[\sum_{q=1}^{Q}(\lambda_q + \sigma^2)^{-1} - \sum_{q=1}^{Q}(\tilde{\Sigma}_{q,q} + \sigma^2)^{-1}]$.*

Lemma 5.2 reveals that the diagonals and eigenvalues and $\tilde{\Sigma}$ are key determinants of $\text{Gain}(\pi, \sigma^2)$, but does not directly provide an interpretation of when $\hat{\beta}_{\text{ECov}}$ offers benefits over $\hat{\beta}_{\text{ID}}$. Our next theorem demonstrates when an improvement can be achieved from joint modeling.

Theorem 5.3. *$\text{Gain}(\pi, \sigma^2) \geq 0$, with equality only when $\tilde{\Sigma} = \text{Var}_\pi[\beta_1]$ is diagonal.*

Proof. From Lemma 5.2 we see $\text{Gain}(\pi, \sigma^2)$ is the difference between a strictly Schur-convex function applied to the eigenvalues of $\tilde{\Sigma}$ and to its diagonals (since $(x + \sigma^2)^{-1}$ is convex on $\mathbb{R}_+$). By the Schur-Horn theorem, the eigenvalues of $\tilde{\Sigma}$ majorize its diagonals, providing the result. $\qquad \square$

Theorem 5.3 tells us that $\hat{\beta}_{\text{ECov}}$ succeeds at adaptively learning and leveraging similarities among datasets in the high-dimensional limit. In particular, $\text{Gain}(\pi, \sigma^2)$ reduces to zero only when the eigenvalues of $\tilde{\Sigma}$ are arbitrarily close to the entries of its diagonal, which occurs only when the

covariate effects are uncorrelated across datasets. However, when covariate effects are correlated, we obtain an improvement.

Our next theorem quantifies this relationship through upper and lower bounds.

Theorem 5.4. *Let $\lambda^{\downarrow}$ and $\ell^{\downarrow}$ denote the eigenvalues and diagonals of $\tilde{\Sigma}$, respectively, sorted in descending order. Then $\mathrm{Gain}(\pi, \sigma^2) \leq 2\sigma^2 Q^{-1} \|\lambda\|_2 \|\ell^{\downarrow} - \lambda^{\downarrow}\|_2 / (\lambda_{\min} + \sigma^2)^3$ and $\mathrm{Gain}(\pi, \sigma^2) \geq \sigma^2 Q^{-1} \|\ell^{\downarrow} - \lambda^{\downarrow}\|_2^2 / (\lambda_{\max} + \sigma^2)^3$, where $\lambda_{\max}$ and $\lambda_{\min}$ are the largest and smallest, respectively, eigenvalues of $\tilde{\Sigma}$.*

Theorem 5.4 allows us to see several aspects of when our method will and will not perform well. First, the presence of $\|\ell^{\downarrow} - \lambda^{\downarrow}\|_2^2$ in both the upper and lower bounds demonstrates that $\mathrm{Gain}(\pi, \sigma^2)$ will be small when the eigenvalues are close to the diagonal entries, with Euclidean distance as an informative metric.

As we find in our next corollary, Theorem 5.4 additionally allows us to see that nontrivial gains may be obtained only in an intermediate signal-to-noise regime, where signal is given by the size of the covariate effects and noise is the variance level σ^2. Notably, under Condition 4.1, σ^2 relates directly to the variance of $\hat{\beta}_{\mathrm{LS}}$, and is influenced by both the residual variances and the dataset sizes; see Appendix C.1. In particular we interpret $\lambda_{\min}$ as a proxy for signal strength since it captures the magnitude of typical β_d's along their direction of least variation.

Corollary 5.5. $\mathrm{Gain}(\pi, \sigma^2) \leq 4\kappa^2 \lambda_{\min}/\sigma^2$ *and* $\mathrm{Gain}(\pi, \sigma^2) \leq 4\kappa^2 (\lambda_{\min}/\sigma^2)^{-1}$, *where* $\kappa := \lambda_{\max}/\lambda_{\min}$ *is the condition number of* $\tilde{\Sigma}$.

Corollary 5.5 formalizes the intuitive result that with enough noise, the little recoverable signal is insufficient to effectively share strength. And furthermore, in the low-noise and high-signal regime $\hat{\beta}_{\mathrm{ID}}$ is very accurate on its own and there is little need for joint modeling. However, when there is a large gap between the largest and smallest eigenvalues of $\tilde{\Sigma}$, leading κ to be large, the gain could be larger. κ will be large, for example, when the covariate effects are very correlated across datasets.

6 Experiments

6.1 Simulated data

We first conduct simulations, where we can directly control the relatedness among datasets and where we know the ground truth values of the parameters. We show that $\hat{\beta}_{\mathrm{ECov}}$ is more accurate than $\hat{\beta}_{\mathrm{EData}}$ when covariates outnumber datasets, whether effects are correlated across datasets or not.

In particular, we simulated covariates, parameters and responses for $Q = 10$ datasets across a range of covariate dimensions. We generated covariate effects as $\beta_d \overset{i.i.d.}{\sim} \mathcal{N}(0, \Sigma)$. We chose Σ so that effects were either correlated (Figure 1 Left) or independent (Figure 1 Right) across datasets; see Appendix E for details.

Figure 1 reinforces our theoretical conclusions that (1) $\hat{\beta}_{\mathrm{ECov}}$ is more accurate when covariates outnumber datasets and (2) $\hat{\beta}_{\mathrm{EData}}$ is more accurate when datasets outnumber covariates. Our simulated X matrices are somewhat relaxed from a strict orthogonal design (Appendix E), so these experiments suggest that our conclusions may hold beyond Condition 4.1. Additionally, $\hat{\beta}_{\mathrm{ECov}}$ and $\hat{\beta}_{\mathrm{EData}}$ both outperform their moment based counterparts, $\hat{\beta}_{\mathrm{ECov}}^{\mathrm{MM}}$ and $\hat{\beta}_{\mathrm{EData}}^{\mathrm{MM}}$.

Even for the simulations with independent effects, Theorem 4.3 suggests $\hat{\beta}_{\mathrm{ECov}}$ should still outperform $\hat{\beta}_{\mathrm{LS}}$ and $\hat{\beta}_{\mathrm{EData}}$ in the higher dimensional regime, and we see this behavior in the right panel of Figure 1. Additionally, in agreement with Theorem 5.3, $\hat{\beta}_{\mathrm{ECov}}$ does not improve over $\hat{\beta}_{\mathrm{ID}}$ in the presence of independent effects, and the performances of these two estimators converge as D grows.

6.2 Real data

We find that ECov beats EData, as well as least squares and independent estimation, across three real datasets. We describe the datasets (with additional details in Appendix E.4) and then our results.

Figure 1: Dimension dependence of parameter estimation error in simulation. Covariate effects are either [Left] correlated or [Right] independent across the $Q = 10$ datasets. Each point is the mean $\pm 1\mathtt{SEM}$ across 20 replicates.

Community level law enforcement in the United States. Policing rates vary dramatically across different communities, mediating disparate impacts of criminal law enforcement across racial and socioeconomic groups [62, 52]. Understanding how demographic and socioeconomic attributes of communities relate to variation in rates of law enforcement is crucial to understanding these impacts. Linear models provide the desired interpretability. We use a dataset [49] consisting of $D = 117$ community characteristics and their rates of law enforcement (per capita) for different crimes. We consider $Q = 4$ data subsets corresponding to distinct (region, crime) pairs: (Midwest, Robbery), (South, Assault), (Northeast, Larceny), and (West, Auto-theft). This data setup illustrates a small Q and accords with the independent residuals assumption in the likelihood shared by ECov and EData (Section 2). Across q, N^q represents between 400 and 600 communities.

Blog post popularity. We regress reader engagement (responses) on $D = 279$ characteristics of blog posts (covariates) [13]. We divided the corpus based on an included length attribute into $Q = 3$ datasets, corresponding to (1) long posts, (2) short posts, and (3) posts from an earlier corpus with missing length attribute. We hypothesized that the relationships between the characteristics of posts and engagement would differ across these three datasets. We randomly downsampled to $N^q = 500$ posts in each dataset to mimic a low sample-size regime, in which sharing strength is crucial.

Figure 2: Prediction performance on held out data in three applications (mean $\pm 1\mathtt{SEM}$ across 5-fold cross-validation splits).

Multiple binary classifications using pre-trained neural network embeddings on CIFAR10. Modern machine learning methods have proved very successful on large datasets. Translating this success to smaller datasets is one of the most actively pursued algorithmic challenges in machine learning. It has spurred the development of frameworks from transfer learning [63] to one-shot learning [60] to meta-learning [21]. One common and simple strategy starts with a learned representation (or "embedding") from an expressive neural network fit to a large dataset. Then one can use this embedding as a covariate vector for classification tasks with few labeled data points.

We take a $D = 128$ dimensional embedding of the CIFAR10 image dataset [37, 58]. We create $Q = 8$ different binary classification tasks using the classes in CIFAR10 (Appendix E.4). We downsampled to N^q varying from 100 to 1000 to mimic a setting in which we hope to share strength from large datasets to improve performance on smaller datasets.

Discussion of evaluation and results. In previous sections we have focused on parameter estimation. Here we instead evaluate with prediction error on held-out data since the true parameters are not observed. Specifically we perform 5-fold cross-validation and report the mean squared errors and classification errors on test splits. To reduce variance of out-of-sample error estimates on the applications in which we downsampled, we also evaluate on the additional held-out data. Because the residual variances were unknown, we estimated these for each application and dataset as $\hat{\sigma}_q^2 :=$ $\|P_{X^q}^\perp Y^q\|^2/(N_q - D)$, where $P_{X^q}^\perp := I_{N_q} - X^q(X^{q\top}X^q)^{-1}X^{q\top}$ (see e.g. [23, Chapter 18.1]). All methods ran quickly on a 36 CPU machine; computation of $\hat{\beta}_{\mathrm{ECov}}$, including the EM algorithm, required 2.04 ± 0.64, 6.89 ± 3.19 and 37.14 ± 3.39 seconds (`mean` $\pm$ `st-dev` across splits) on the law enforcement, blog, and CIFAR10 tasks, respectively.

Our results further reinforce the main aspects of our theory. $\hat{\beta}_{\mathrm{ECov}}$ outperformed $\hat{\beta}_{\mathrm{EData}}$, independent Bayes estimates ($\hat{\beta}_{\mathrm{ID}}$), and least squares ($\hat{\beta}_{\mathrm{LS}}$) in all applications (at $> 95\%$ nominal confidence with a paired t-test).[1] Additionally, $\hat{\beta}_{\mathrm{ECov}}$ outperformed the baseline of ignoring heterogeneity, pooling datasets together, and using the same effect estimates for every dataset ("Least Sqrs./MLE Pool").

Appendix E includes additional results and comparisons. In particular, we provide the performance of the estimators on each component dataset for each application. Additionally, we report the performances of (1) stable and computationally efficient moment based alternatives to $\hat{\beta}_{\mathrm{ECov}}$ and $\hat{\beta}_{\mathrm{EData}}$ and (2) variants of $\hat{\beta}_{\mathrm{ECov}}$ and $\hat{\beta}_{\mathrm{EData}}$ that include a learned (rather than zero) prior mean. Appendix E.5 reports the licenses of software we used.

7 Discussion

The Bayesian community has long used hierarchical modeling with priors encoding exchangeability of effects across datasets (EData). In the present work, we have made a case for instead using priors that encode exchangeability across *covariates* (ECov) – in particular, when the number of covariates exceeds the number of datasets. We have presented a corresponding concrete model and inference method. We have shown that ECov outperforms EData in theory and practice when the number of covariates exceeds the number of datasets.

Our approach is, of course, not a panacea. In some settings, a priori exchangeability among covariate effects will be inconsistent with prior beliefs. For example, imagine in the CIFAR10 application if meta-data covariates (such as geo-location and date) were available, in addition to embeddings. Then we might achieve better performance by treating meta-data covariates as distinct from embedding covariates. Additionally, we focused on a Gaussian prior for convenience. In cases where practitioners have more specific prior beliefs about effects, alternative priors and likelihoods may be warranted, though they may be more computationally challenging. Moreover, while relatively interpretable, linear models have their downsides. The linear assumption can be overly simplistic in many applications. It is common to misinterpret effects as causal rather than associative. Both the linear model and squared error loss lend themselves naturally to reporting means, but in many applications a median or other summary is more appropriate; so using a mean for convenience can be misleading.

Nevertheless, we believe that, on net, our methodology is likely to produce positive societal impacts. Machine learning has recently trended toward black-box predictive models. Though linear models are already widely used in applied disciplines for their interpretability, we hope that our work will improve their accuracy and help further extend their reach.

And there remain many exciting directions for further investigation. For example, the covariance Σ may provide an informative measure of task similarity; this similarity measure can be useful in, e.g., meta learning [34] and statistical genetics [12]. It also remains to extend our methodology to other generalized linear models.

---

[1] We did not develop an extension akin to Algorithm 3 for EData, and so do not report $\hat{\beta}_{\mathrm{EData}}$ for CIFAR10. Additionally, we report a maximum likelihood estimate (MLE) instead of $\hat{\beta}_{\mathrm{LS}}$ for CIFAR10.

References

[1] Martín Abadi, Paul Barham, Jianmin Chen, Zhifeng Chen, Andy Davis, Matthieu Dean, Jeffrey an Devin, Sanjay Ghemawat, Geoffrey Irving, Michael Isard, Manjunath Kudlur, Josh Levenberg, Rajat Monga, Sherry Moore, Derek G. Murray, Benoit Steiner, Paul Tucker, Vijay Vasudevan, Pete Warden, Martin Wicke, Yuan Yu, Xiaoqiang Zheng, and Google Brain. Tensorflow: A system for large-scale machine learning. In *12th USENIX symposium on operating systems design and implementation ({OSDI} 16)*, pages 265–283, 2016.

[2] Tomohiro Ando and Arnold Zellner. Hierarchical Bayesian analysis of the seemingly unrelated regression and simultaneous equations models using a combination of direct Monte Carlo and importance sampling techniques. *Bayesian Analysis*, 5(1):65–95, 2010.

[3] Oswald T Avery, Colin M MacLeod, and Maclyn McCarty. Studies on the chemical nature of the substance inducing transformation of pneumococcal types: induction of transformation by a desoxyribonucleic acid fraction isolated from pneumococcus type III. *The Journal of Experimental Medicine*, 79(2):137–158, 1944.

[4] Alvin J Baranchik. Multiple regression and estimation of the mean of a multivariate normal distribution. Technical report, Stanford University, 1964.

[5] Douglas Bates, Martin Mächler, Ben Bolker, and Steve Walker. Fitting linear mixed-effects models using lme4. *Journal of Statistical Software*, 67(1):1–48, 2015.

[6] Anindya Bhadra and Bani K Mallick. Joint high-dimensional Bayesian variable and covariance selection with an application to eQTL analysis. *Biometrics*, 69(2):447–457, 2013.

[7] Christopher M. Bishop. *Pattern Recognition and Machine Learning*. Springer, 2006.

[8] Robert C Blattberg and Edward I George. Shrinkage estimation of price and promotional elasticities: Seemingly unrelated equations. *Journal of the American Statistical Association*, 86(414):304–315, 1991.

[9] Leo Breiman and Jerome H Friedman. Predicting multivariate responses in multiple linear regression. *Journal of the Royal Statistical Society: Series B*, 59(1):3–54, 1997.

[10] Philip J Brown and James V Zidek. Adaptive multivariate ridge regression. *The Annals of Statistics*, 8(1):64–74, 1980.

[11] Philip J Brown, Marina Vannucci, and Tom Fearn. Multivariate Bayesian variable selection and prediction. *Journal of the Royal Statistical Society: Series B*, 60(3):627–641, 1998.

[12] Brendan Bulik-Sullivan, Hilary K Finucane, Verneri Anttila, Alexander Gusev, Felix R Day, Po-Ru Loh, Laramie Duncan, John RB Perry, Nick Patterson, Elise B Robinson, Mark J Daly, Alkes L Price, and Benjamin M Neal. An atlas of genetic correlations across human diseases and traits. *Nature Genetics*, 47 (11):1236, 2015.

[13] Krisztian Buza. Feedback prediction for blogs. In *Data Analysis, Machine Learning and Knowledge Discovery*, pages 145–152. Springer, 2014.

[14] Diana Cai, Rishit Sheth, Lester Mackey, and Nicolo Fusi. Weighted meta-learning. *arXiv preprint arXiv:2003.09465*, 2020.

[15] Siddhartha Chib and Edward Greenberg. Hierarchical analysis of SUR models with extensions to correlated serial errors and time-varying parameter models. *Journal of Econometrics*, 68(2):339–360, 1995.

[16] A Philip Dawid. Some matrix-variate distribution theory: notational considerations and a Bayesian application. *Biometrika*, 68(1):265–274, 1981.

[17] Sameer K Deshpande, Veronika Ročková, and Edward I George. Simultaneous variable and covariance selection with the multivariate spike-and-slab lasso. *Journal of Computational and Graphical Statistics*, 2019.

[18] Bradley Efron and Carl Morris. Empirical Bayes on vector observations: An extension of Stein's method. *Biometrika*, 59(2):335–347, 1972.

[19] Bradley Efron and Carl Morris. Limiting the risk of Bayes and empirical Bayes estimators—Part II: The empirical Bayes case. *Journal of the American Statistical Association*, 67(337):130–139, 1972.

[20] Jianqing Fan and Runze Li. Variable selection via nonconcave penalized likelihood and its oracle properties. *Journal of the American Statistical Association*, 96(456):1348–1360, 2001.

[21] Chelsea Finn, Pieter Abbeel, and Sergey Levine. Model-agnostic meta-learning for fast adaptation of deep networks. In *International Conference on Machine Learning*, pages 1126–1135. PMLR, 2017.

[22] Alan E Gelfand, Susan E Hills, Amy Racine-Poon, and Adrian FM Smith. Illustration of Bayesian inference in normal data models using Gibbs sampling. *Journal of the American Statistical Association*, 85 (412):972–985, 1990.

[23] Andrew Gelman and Jennifer Hill. *Data Analysis using Regression and Multilevel/Hierarchical Models*. Cambridge University Press, 2006.

[24] Andrew Gelman, John B Carlin, Hal S Stern, David B Dunson, Aki Vehtari, and Donald B Rubin. *Bayesian Data Analysis*. Chapman and Hall/CRC, 2013.

[25] Amos Golan and Jeffrey M Perloff. Comparison of maximum entropy and higher-order entropy estimators. *Journal of Econometrics*, 107(1-2):195–211, 2002.

[26] Erin Grant, Chelsea Finn, Sergey Levine, Trevor Darrell, and Thomas Griffiths. Recasting gradient-based meta-learning as hierarchical Bayes. In *International Conference on Learning Representations*, 2018.

[27] William E Griffiths. Bayesian inference in the seemingly unrelated regressions model. In *Computer-Aided Econometrics*, pages 287–314. CRC Press, 2003.

[28] Yoel Haitovsky. On multivariate ridge regression. *Biometrika*, 74(3):563–570, 1987.

[29] Lanoy Nelson Hazel. The genetic basis for constructing selection indexes. *Genetics*, 28(6):476–490, 1943.

[30] Grant Hillier and Raymond Kan. Properties of the inverse of a noncentral Wishart matrix. *Available at SSRN 3370864*, 2019.

[31] Arthur E Hoerl and Robert W Kennard. Ridge regression: Biased estimation for nonorthogonal problems. *Technometrics*, 12(1):55–67, 1970.

[32] Alfred Horn. Doubly stochastic matrices and the diagonal of a rotation matrix. *American Journal of Mathematics*, 76(3):620–630, 1954.

[33] W. James and Charles Stein. Estimation with quadratic loss. *Proceedings of the Fourth Berkeley Symposium on Mathematical Statistics and Probability*, 1:361–379, 1961.

[34] Ghassen Jerfel, Erin Grant, Thomas L Griffiths, and Katherine Heller. Reconciling meta-learning and continual learning with online mixtures of tasks. *Advances in Neural Information Processing Systems*, 32, 2019.

[35] Michael I Jordan. Bayesian nonparametric learning: Expressive priors for intelligent systems. *Heuristics, Probability and Causality: A Tribute to Judea Pearl*, 11:167–185, 2010.

[36] Diederik P Kingma and Max Welling. Auto-encoding variational Bayes. *arXiv preprint arXiv:1312.6114*, 2013.

[37] Alex Krizhevsky. Learning multiple layers of features from tiny images. *Technical Paper, University of Toronto*, 2009.

[38] Nan M Laird and James H Ware. Random-effects models for longitudinal data. *Biometrics*, 38(4):963–974, 1982.

[39] Sang Hong Lee, Jian Yang, Michael E Goddard, Peter M Visscher, and Naomi R Wray. Estimation of pleiotropy between complex diseases using single-nucleotide polymorphism-derived genomic relationships and restricted maximum likelihood. *Bioinformatics*, 28(19):2540–2542, 2012.

[40] Erich L Lehmann and George Casella. *Theory of Point Estimation*. Springer Science & Business Media, 2006.

[41] Guy Letac and Hélène Massam. A tutorial on non central Wishart distributions. *Technical Paper, Toulouse University*, 2004.

[42] Alex Lewin, Habib Saadi, James E Peters, Aida Moreno-Moral, James C Lee, Kenneth GC Smith, Enrico Petretto, Leonardo Bottolo, and Sylvia Richardson. MT-HESS: an efficient Bayesian approach for simultaneous association detection in OMICS datasets, with application to eQTL mapping in multiple tissues. *Bioinformatics*, 32(4):523–532, 2015.

[43] Dennis V Lindley and Adrian FM Smith. Bayes estimates for the linear model. *Journal of the Royal Statistical Society: Series B*, 34(1):1–18, 1972.

[44] David G Luenberger. *Introduction to Linear and Nonlinear Programming*. Addison-Wesley Reading, MA, 1973.

[45] Robert Maier, Gerhard Moser, Guo-Bo Chen, Stephan Ripke, Cross-Disorder Working Group of the Psychiatric Genomics Consortium, William Coryell, James B Potash, William A Scheftner, Jianxin Shi, Myrna M Weissman, Christina M Hultman, Mikael Landén, Douglas F Levinson, Kenneth S Kendler, Jordan W Smoller, Naomi R Wray, and S Hong Lee. Joint analysis of psychiatric disorders increases accuracy of risk prediction for schizophrenia, bipolar disorder, and major depressive disorder. *The American Journal of Human Genetics*, 96(2):283–294, 2015.

[46] Geoffrey J McLachlan and Thriyambakam Krishnan. *The EM Algorithm and Extensions*, volume 382. John Wiley & Sons, 2007.

[47] Rachael Meager. Understanding the average impact of microcredit expansions: A Bayesian hierarchical analysis of seven randomized experiments. *American Economic Journal: Applied Economics*, 11(1):57–91, 2019.

[48] Jorge Nocedal and Stephen Wright. *Numerical Optimization*. Springer Science & Business Media, 2006.

[49] Michael Redmond and Alok Baveja. A data-driven software tool for enabling cooperative information sharing among police departments. *European Journal of Operational Research*, 141(3):660–678, 2002.

[50] Gregory C Reinsel. Mean squared error properties of empirical Bayes estimators in a multivariate random effects general linear model. *Journal of the American Statistical Association*, 80(391):642–650, 1985.

[51] Daniel E Runcie, Jiayi Qu, Hao Cheng, and Lorin Crawford. MegaLMM: Mega-scale linear mixed models for genomic predictions with thousands of traits. *BioRxiv*, 2020.

[52] Lee A Slocum, Beth M Huebner, Claire Greene, and Richard Rosenfeld. Enforcement trends in the city of St. Louis from 2007 to 2017: Exploring variability in arrests and criminal summonses over time and across communities. *Journal of Community Psychology*, 48(1):36–67, 2020.

[53] Michael Smith and Robert Kohn. Nonparametric seemingly unrelated regression. *Journal of Econometrics*, 98(2):257–281, 2000.

[54] Matthew Stephens. A unified framework for association analysis with multiple related phenotypes. *PloS One*, 8(7):e65245, 2013.

[55] Robin Thompson. The estimation of variance and covariance components with an application when records are subject to culling. *Biometrics*, pages 527–550, 1973.

[56] Hisayuki Tsukuma. Admissibility and minimaxity of Bayes estimators for a normal mean matrix. *Journal of Multivariate Analysis*, 99(10):2251–2264, 2008.

[57] A Van Der Merwe and JV Zidek. Multivariate regression analysis and canonical variates. *Canadian Journal of Statistics*, 8(1):27–39, 1980.

[58] Arnaud Van Looveren, Giovanni Vacanti, Janis Klaise, and Alexandru Coca. Alibi-Detect: Algorithms for outlier and adversarial instance detection, concept drift and metrics., 2019. URL https://github.com/SeldonIO/alibi-detect.

[59] Wessel N van Wieringen. Lecture notes on ridge regression. *arXiv preprint arXiv:1509.09169*, 2015.

[60] Oriol Vinyals, Charles Blundell, Timothy Lillicrap, Koray Kavukcuoglu, and Daan Wierstra. Matching networks for one shot learning. In *Proceedings of the 30th International Conference on Neural Information Processing Systems*, pages 3637–3645, 2016.

[61] Martin J Wainwright. *High-Dimensional Statistics: A Non-Asymptotic Viewpoint*. Cambridge University Press, 2019.

[62] David Weisburd, Malay K Majmundar, Hassan Aden, Anthony Braga, Jim Bueermann, Philip J Cook, Phillip Atiba Goff, Rachel A Harmon, Amelia Haviland, Cynthia Lum, Charles Manski, Stephen Mastrofski, Tracey Meares, Daniel Nagin, Emily Owens, Steven Raphael, Jerry Ratcliffe, and Tom Tyler. Proactive policing: A summary of the report of the national academies of sciences, engineering, and medicine. *Asian Journal of Criminology*, 14(2):145–177, 2019.

[63] Karl Weiss, Taghi M Khoshgoftaar, and DingDing Wang. A survey of transfer learning. *Journal of Big Data*, 3(1):1–40, 2016.

[64] Arnold Zellner. An efficient method of estimating seemingly unrelated regressions and tests for aggregation bias. *Journal of the American Statistical Association*, 57(298):348–368, 1962.

[65] Arnold Zellner and David S Huang. Further properties of efficient estimators for seemingly unrelated regression equations. *International Economic Review*, 3(3):300–313, 1962.

[66] Xiang Zhou and Matthew Stephens. Efficient multivariate linear mixed model algorithms for genome-wide association studies. *Nature Methods*, 11(4):407, 2014.

[67] Jim Zidek. Deriving unbiased risk estimators of multinormal mean and regression coefficient estimators using zonal polynomials. *The Annals of Statistics*, pages 769–782, 1978.