Distributed Inverse Constrained Reinforcement Learning for Multi-agent Systems

Shicheng Liu & Minghui Zhu
School of Electrical Engineering and Computer Science
Pennsylvania State University
University Park, PA 16802, USA
{sfl5539,muz16}@psu.edu

Abstract

This paper considers the problem of recovering the policies of multiple interacting experts by estimating their reward functions and constraints where the demonstration data of the experts is distributed to a group of learners. We formulate this problem as a distributed bi-level optimization problem and propose a novel bi-level "distributed inverse constrained reinforcement learning" (D-ICRL) algorithm that allows the learners to collaboratively estimate the constraints in the outer loop and learn the corresponding policies and reward functions in the inner loop from the distributed demonstrations through intermittent communications. We formally guarantee that the distributed learners asymptotically achieve consensus which belongs to the set of stationary points of the bi-level optimization problem. Simulations are done to validate the proposed algorithm.

1 Introduction

Multi-agent systems (MASs) have become an effective way to model large-scale networked systems, *e.g.*, multi-robot systems and the Internet-of-Things. Multi-agent reinforcement learning (MARL) is a systematic framework to coordinate MASs. It extends RL to address the decision-making problem of multiple interacting agents [1]. One challenge in (MA)RL is that manually crafting the reward functions or predesigning the policies can be infeasible for humans when the tasks are too complicated. Inverse reinforcement learning (IRL) presents a way to tackle this problem where a learner aims to recover a reward function (and a corresponding policy) that best explains the behavior in the demonstrations of an expert. Current state-of-the-arts on IRL [2, 3, 4, 5] can successfully solve the reward function inference problem but still face two other major challenges in MARL and MASs: (i) it is often the case that the expected behavior (experts' behavior) is more precisely explained by a reward function combined with a set of constraints instead of a single reward function; (ii) the demonstration data in MASs is usually distributed over multiple learners and unable to be shared due to some practical reasons including privacy concerns and communication burdens. This paper aims to bridge the gap and proposes the first algorithm to solve the three challenges (i.e., reward function inference, constraint inference, and distributed demonstration data) at once.

Related works. IRL is an ambiguous problem [6] as different reward functions could explain the demonstrations. To solve this problem, several fundamental approaches are proposed including feature expectation matching [7], maximum margin planning [8], maximum entropy (ME) [9], and maximum causal entropy (MCE) [10]. Recently, some advanced techniques baring more features are proposed. Paper [2] uses adversarial learning to learn reward functions robust to environmental changes, paper [4] applies strongly convex regularizers to the learner's policy to avoid the ambiguity problem, and paper [5] proposes a scalable Bayesian-based method working for large control tasks.

36th Conference on Neural Information Processing Systems (NeurIPS 2022).

In real-world settings, numerous constraints are present and a single reward function is not enough to explain complicated behavior. Paper [11] extends ME IRL to environments with constraints, formulates the constraint inference problem as a maximum likelihood estimation (MLE) problem, and leverages a greedy method to solve the problem. Paper [12] extends paper [11] to continuous environments. These two works both assume the access to the ground truth rewards.

While the discussions in the last two paragraphs are limited to single agents, there have been recent works which extend IRL techniques to multiple agents [13, 14, 15, 16, 17]. Notice that previous works on multi-agent IRL do not explicitly distinguish the notions of "experts" and "learners", their "multiple agents" means "multiple experts" and they only consider centralized learning (*i.e.*, a single learner who obtains all demonstrated behavior of the multiple experts). In this paper, we explicitly distinguish the notions of "experts" and "learners", and use "multiple experts" to represent the commonly used "multiple agents" in literature. Moreover, the aforementioned papers do not consider the more general situation where the demonstrations are distributed over multiple learners and unable to be shared. To handle this situation, the learners need to perform distributed learning.

As a basic topology in distributed learning, peer-to-peer (P2P) architecture has no central server and each node can communicate with a subset of all the nodes. It reduces the risk of single points of failures [18] in the system. This paper proposes D-ICRL based on the P2P structure where ICRL can learn policies by estimating the reward functions and constraints through demonstrations and distributed learners can perform ICRL without sharing the distributed demonstration data.

In specific, we formulate D-ICRL as a distributed bi-level optimization problem where the learners cooperatively learn the constraints through the outer problem and estimate the corresponding reward functions and policies through the inner problem. Paper [19] provides asymptotic convergence for centralized bi-level optimization problems by assuming that the inner problem can be completely solved and its optimal solution can be obtained. Papers [20, 21, 22] relax this assumption by partially solving the inner problem and provide convergence rate analysis for problems with strongly convex inner objective function. Paper [23] solves a special case of distributed bi-level optimization where the outer and inner problems use identical decision variables. To the best of our knowledge, there is no work on bi-level optimization which (i) assumes that the inner objective function is merely strictly convex and partially solves the inner problem; (ii) solves the general form of bi-level optimization in a distributed way. Notice that these two challenges arise in our D-ICRL problem.

Contribution statement. Our contributions are threefold. First, we extend IRL to a multi-expert-multi-learner (MEML) setting where a group of learners collaboratively learn policies by estimating the reward functions and constraints without sharing the distributed demonstrations performed by a set of cooperative experts. We formulate this D-ICRL problem as a distributed bi-level optimization problem. Second, we propose a distributed bi-level learning algorithm to solve this problem where the learners cooperatively estimate the constraints in the outer problem and learn the corresponding reward functions and policies in the inner problem. Third, we provide convergence rate analysis to the optimal solution of the inner problem and asymptotic convergence to the set of stationary points of the outer problem. Simulations are conducted to validate the proposed method.

2 Model

In this section, we present the models for cooperative experts and distributed learners.

Experts. There are N_E experts whose decision making is based on a constrained Markov game (CMG) [24]. A CMG $(\mathcal{S}, \mathcal{A}, \gamma, P_0, P, r_E, c_E, b)$ is defined via state set $\mathcal{S} \triangleq \prod_{i=1}^{N_E} \mathcal{S}^{[i]}$, action set $\mathcal{A} \triangleq \prod_{i=1}^{N_E} \mathcal{A}^{[i]}$, discount factor $\gamma \in (0, 1)$, and initial state distribution P_0. The state-action space can be either discrete or continuous. The system state transition function is given by P such that the probability (or probability density in continuous state-action space) of state transition to s' from s by taking action $a \triangleq (a^{[1]}, \cdots, a^{[N_E]})$ is $P(s'|s, a)$. The reward function of expert i is $r_E^{[i]} : \mathcal{S} \times \mathcal{A} \to \mathbb{R}$ and the experts are cooperative (*i.e.*, $r_E \triangleq \sum_{i=1}^{N_E} r_E^{[i]}$). The cost function of expert i is $c_E^{[i]} : \mathcal{S} \times \mathcal{A} \to \mathbb{R}_+$ and the cost function of the whole system is $c_E \triangleq \sum_{i=1}^{N_E} c_E^{[i]}$. Expert i's policy $\pi_E^{[i]}(a^{[i]}|s)$ represents the probability (or probability density) of expert i taking action $a^{[i]}$ at state s and the joint policy of all the experts is denoted by $\pi_E(a|s) \triangleq \prod_{i=1}^{N_E} \pi_E^{[i]}(a^{[i]}|s)$. Define $J_{r_E}(\pi) \triangleq E_{S,A}^\pi [\sum_{t=0}^\infty \gamma^t r_E(S_t, A_t)]$ as the expected cumulative reward where the initial

state is drawn from P_0 and $J_{c_E}(\pi) \triangleq E^{\pi}_{S,A}[\sum_{t=0}^{\infty} \gamma^t c_E(S_t, A_t)]$ as the expected cumulative cost. The experts' policy π_E wants to maximize $J_{r_E}(\pi)$ subject to $J_{c_E}(\pi) \leq b$. Define the constraint set indicated by $c_E^{[i]}$ as $\mathcal{C}_E^{[i]} \triangleq \{(s,a) \in \mathcal{S} \times \mathcal{A} : c_E^{[i]}(s,a) > 0\}$. Following [12], we study the case where $b = 0$, i.e., hard constraint. Thus, the constraint reduces to $J_{c_E}(\pi) = 0$, implying that the probability of reaching any $(s,a) \in \bigcup_i \mathcal{C}_E^{[i]}$ is zero under policy π. These experts use π_E to demonstrate a set $\mathcal{D} \triangleq \{\zeta^j\}_{j=1}^m$ of m trajectories, which is partitioned into N_L subsets. Each trajectory $\zeta^j \triangleq s_0^j, a_0^j, s_1^j, a_1^j \cdots$ is a state-action sequence of the experts.

Learners. There is a group of N_L learners where each learner v knows $(\mathcal{S}, \mathcal{A}, \gamma, P_0, P, b, \mathcal{D}^{[v]})$ and can communicate with other learners. The local demonstration subset $\mathcal{D}^{[v]}$ has $m^{[v]}$ trajectories. The learners choose an $l_r^{[i]}$-dimensional reward feature vector $\phi_r^{[i]} : \mathcal{S} \times \mathcal{A} \to [0, d_1]^{l_r^{[i]}}$ for expert i where d_1 is constant and it is assumed that $r_E^{[i]} = (\omega_{rE}^{[i]})^\top \phi_r^{[i]}$, where $\omega_{rE}^{[i]} \in \mathbb{R}^{l_r^{[i]}}$ is the (unknown) reward weight vector. The learners also choose an $l_c^{[i]}$-dimensional cost feature vector $\phi_c^{[i]} : \mathcal{S} \times \mathcal{A} \to [0, d_2]^{l_c^{[i]}}$ where d_2 is constant and it is assumed that $c_E^{[i]} = (\omega_{cE}^{[i]})^\top \phi_c^{[i]}$, where $\omega_{cE}^{[i]}$ is the (unknown) cost weight vector and $\omega_{cE,j}^{[i]}$ is the j-th component of $\omega_{cE}^{[i]}$. Let $\mathcal{C}_j^{[i]} \triangleq \{(s,a) \in \mathcal{S} \times \mathcal{A} : \phi_{c,j}^{[i]}(s,a) > 0\}$ be the constraint set indicated by the j-th cost feature $\phi_{c,j}^{[i]}$ of expert i. As the budget $b = 0$, the effect of $\omega_{cE,j}^{[i]} > 0$ is the same, i.e., $\mathcal{C}_j^{[i]}$ cannot be visited, thus the learners assume $\omega_{cE,j}^{[i]} \in [0,1]$.

Communication network. The learners execute a two-time-scale learning algorithm over a communication network. The fast-time-scale (inner) algorithm is executed at discrete time k over a time-varying directed graph $\mathcal{G}(k) \triangleq (\mathcal{V}, \mathcal{E}(k))$ where $\mathcal{V} \triangleq \{1, \cdots, N_L\}$ is the node (learner) set and $\mathcal{E}(k) \subseteq \mathcal{V} \times \mathcal{V}$ is the set of directed edges (communication links) at time k. The edge $(v, v') \in \mathcal{E}(k)$ means that learner v receives information from learner v' at time k and $(v,v) \in \mathcal{E}(k)$ for all $k \geq 0$. The adjacency matrix of the graph at time k is $W(k) \triangleq [W^{[vv']}(k)]_{v,v' \in \mathcal{V}} \in \mathbb{R}^{N_L \times N_L}$ where $W^{[vv']}(k) = 0$ if and only if $(v, v') \notin \mathcal{E}(k)$. We use $\mathcal{N}^{[v]}(k) \triangleq \{v' \in \mathcal{V} | (v, v') \in \mathcal{E}(k)\}$ to denote the set of neighbors of learner v at time k. The slow-time-scale (outer) algorithm is carried out at discrete time n over a graph $\bar{\mathcal{G}}(n) \triangleq (\mathcal{V}, \bar{\mathcal{E}}(n))$, where $\bar{\mathcal{E}}(n)$, $\bar{W}(n)$ and $\bar{\mathcal{N}}^{[v]}(n)$ are defined in an analogous way. The time unit for n is much larger than that for k.

Assumption 1. *There exists an integer $B \geq 1$ such that the graph $(\mathcal{V}, \mathcal{E}(k) \cup \cdots \cup \mathcal{E}(k+B-1))$ is strongly connected for all $k \geq 0$.*

Assumption 2. *The adjacency matrix $W(k)$ has the following properties:*
(i) (doubly stochastic) $\mathbf{1}^\top W(k) = \mathbf{1}^\top$ and $W(k)\mathbf{1} = \mathbf{1}$ where $\mathbf{1}$ is the column vector whose entries are all ones. (ii) (non-degenerate) There is an $\epsilon \in (0,1)$ such that $W^{[vv]}(k) \geq \epsilon$ for all $v \in \mathcal{V}$ and $W^{[vv']}(k) \geq \epsilon$ if $(v, v') \in \mathcal{E}(k)$.

Assumption 3. *The set $\bar{\mathcal{N}}^{[v]}(n) = \mathcal{V}$ for any $v \in \mathcal{V}$ and $n \geq 0$. The adjacency matrix $\bar{W}(n)$ is doubly stochastic and non-degenerate with constant $\bar{\epsilon}$.*

Figure 1: Relation between experts and learners

Figure 1 provides an illustration of the relation between the experts and the learners. The experts operate in a multi-agent system and demonstrate a set $\mathcal{D}$ of trajectories and the learners stand outside the system where each learner v can observe a subset $\mathcal{D}^{[v]}$ of the global demonstration.

3 Notions and notations

This section provides notions and notations which will be used in the remaining of the paper.

Reward feature expectation matching (RFEM). Given policy π, the reward feature expectation vector of expert i is defined as $\mu_r^{[i]}(\pi) \triangleq E_{S,A}^{\pi}[\sum_{t=0}^{\infty} \gamma^t \phi_r^{[i]}(S_t, A_t)]$ and the demonstration $\mathcal{D}$ is needed to estimate $\mu_r^{[i]}(\pi_E)$: $\hat{\mu}_r^{[i]} \triangleq \frac{1}{m}\sum_{j=1}^{m}\sum_{t=0}^{\infty} \gamma^t \phi_r^{[i]}(s_t^j, a_t^j)$, where $(s_t^j, a_t^j) \in \zeta^j$. The RFEM requires $\mu_r^{[i]}(\pi) = \hat{\mu}_r^{[i]}$ for every expert i. However, each learner v can only observe the subset $\mathcal{D}^{[v]}$ and its local estimate is $\hat{\mu}_r^{[i,v]} \triangleq \frac{1}{m^{[v]}}\sum_{\zeta^j \in \mathcal{D}^{[v]}}\sum_{t=0}^{\infty} \gamma^t \phi_r^{[i]}(s_t^j, a_t^j)$.

Cost expectation matching (CEM). Similar to RFEM, CEM requires $J_{c_E}(\pi) = J_{c_E}(\pi_E)$. As the true cost function c_E is unknown, given an estimate $c_{\omega_c} = \sum_{i=1}^{N_E}(\omega_c^{[i]})^{\top}\phi_c^{[i]}$, we define the empirical cost estimate of the experts as $\hat{b}_{\omega_c} \triangleq \frac{1}{m}\sum_{j=1}^{m}\sum_{t=0}^{\infty} \gamma^t c_{\omega_c}(s_t^j, a_t^j)$. The local estimate from learner v is $\hat{b}_{\omega_c}^{[v]} \triangleq \frac{1}{m^{[v]}}\sum_{\zeta^j \in \mathcal{D}^{[v]}}\sum_{t=0}^{\infty} \gamma^t c_{\omega_c}(s_t^j, a_t^j)$.

Remark 1. *As RFEM is a standard notion in IRL literature, it is "natural" to follow this and use "cost feature expectation matching" (CFEM). However, here we use CEM instead of CFEM because we want to leverage the budget information provided in CMG. We still use RFEM instead of "reward expectation matching" (REM) because REM does not embed any additional useful information as CMG does not have any "budget information" for cumulative reward.*

Notations. We use $\hat{\mu}_r \triangleq [(\hat{\mu}_r^{[1]})^{\top}, \cdots, (\hat{\mu}_r^{[N_E]})^{\top}]^{\top}$ to represent the concatenated empirical reward feature expectation vector. Similarly, we define learner v's concatenated empirical reward feature expectation vector estimate $\hat{\mu}_r^{[v]}$, concatenated reward feature expectation vector $\mu_r(\pi)$ under π, and concatenated cost feature vector ϕ_c. Let Π be the set of all valid stochastic policies such that every $\pi \in \Pi$ satisfies $\pi(a|s) \geq 0$ for any $s \in \mathcal{S}$ and $a \in \mathcal{A}$ and $\int_{a \in \mathcal{A}} \pi(a|s)da = 1$ for any $s \in \mathcal{S}$. We define the expected cumulative cost of policy π under cost function estimate c_{ω_c} as $J_{\omega_c}(\pi) \triangleq E_{S,A}^{\pi}[\sum_{t=0}^{\infty} \gamma^t c_{\omega_c}(S_t, A_t)]$ and use $\text{Project}_{\Omega_c}(\cdot)$ to denote the projection onto the set Ω_c.

4 Problem formulation

In an MEML D-ICRL problem, each learner v aims to use all the available information to collaboratively recover the policy π_E through communications by estimating the reward function r_E and the cost function c_E. In the learning process, learner v is unwilling to share its private data (*i.e.*, local demonstration and local estimate of reward feature expectation and cost expectation). Notice that estimating c_E is in essence estimating the constraint set $\bigcup_i \mathcal{C}_E^{[i]}$. We formulate the constraint learning problem as a distributed MLE problem:

$$\max_{\omega_c \in \Omega_c} \quad F(\omega_c) = \sum_{v=1}^{N_L} F^{[v]}(\omega_c), \tag{1}$$

where $\Omega_c \triangleq [0,1]^{\sum_{i=1}^{N_E} l_c^{[i]}}$, $F^{[v]}(\omega_c) \triangleq \sum_{j \in \mathcal{D}^{[v]}}\sum_{t=0}^{\infty} \gamma^t \ln \pi_{\omega_c}(a_t^j|s_t^j)$ is learner v's local likelihood function and π_{ω_c} is a policy parameterized by ω_c. To find a statistical distribution model for π_{ω_c}, we formulate the following problem based on MCE scheme:

$$\max_{\pi \in \Pi} \quad H(\pi), \qquad \text{s.t.} \quad \mu_r^{[i]}(\pi) = \hat{\mu}_r^{[i]} \quad \forall i \in \mathcal{I}, \quad J_{\omega_c}(\pi) = \hat{b}_{\omega_c}, \tag{2}$$

where $H(\pi) \triangleq \sum_{t=0}^{\infty} E_{S,A}[-\gamma^t \ln \pi(A_t|S_t)]$ is the (infinite-horizon) causal entropy [25].

Remark 2. *Directly matching the ground truth budget (i.e., $J_{\omega_c}(\pi) = b$) may cause infeasibility of (2), i.e., the feasible set of (2) is not guaranteed to be nonempty. Therefore, we match $\hat{b}_{\omega_c}$, which is an estimate of $J_{\omega_c}(\pi_E)$, to ensure the feasibility because now, π_E is a feasible solution of (2).*

The dual function of (2) is $G(\omega_r, \lambda; \omega_c) \triangleq \max_{\pi \in \Pi} H(\pi) + \sum_{i=1}^{N_E}(\omega_r^{[i]})^{\top}(\mu_r^{[i]}(\pi) - \hat{\mu}_r^{[i]}) + \lambda(J_{\omega_c}(\pi) - \hat{b}_{\omega_c})$ where the dual variable $\omega_r = [(\omega_r^{[1]})^{\top}, \cdots, (\omega_r^{[N_E]})^{\top}]^{\top}$ is to estimate ω_{r_E}. Let $\eta \triangleq [\omega_r^{\top}, \lambda]^{\top}$, it is well-known that $G(\eta; \omega_c)$ is convex in η since it is the pointwise maximum of a family of affine functions of η. As $G(\eta; \omega_c)$ is global, we introduce the local convex dual function $G^{[v]}(\eta; \omega_c) \triangleq \max_{\pi \in \Pi} H(\pi) + \sum_{i=1}^{N_E}(\omega_r^{[i]})^{\top}(\mu_r^{[i]}(\pi) - \hat{\mu}_r^{[i,v]}) + \lambda(J_{\omega_c}(\pi) - \hat{b}_{\omega_c}^{[v]})$.

Lemma 1. *(i) The optimal solution of problem (2) is the constrained soft Bellman policy $\pi_{\eta^*(\omega_c);\omega_c}$, where $\eta^*(\omega_c)$ is an optimal solution of $\min_\eta G(\eta;\omega_c)$. (ii) The local dual function $G^{[v]}(\eta;\omega_c)$ is differentiable and its gradient is $[(\mu_r(\pi_{\eta;\omega_c}) - \hat{\mu}_r^{[v]})^\top, J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c}^{[v]}]^\top$.*

The proof of Lemma 1 is similar to [10, 25, 26]. For the sake of completeness, we still include the proof and the expression of the constrained soft Bellman policy in the Appendix.

Lemma 1 indicates that $\pi_{\eta^*(\omega_c);\omega_c}$ is the statistical model we need in problem (1). Therefore, problem (1) can be reformulated as the following distributed bi-level optimization problem:

$$\max_{\omega_c \in \Omega_c} \quad F(\omega_c, \eta^*(\omega_c)) = \sum_{v=1}^{N_L} F^{[v]}(\omega_c, \eta^*(\omega_c)), \tag{3}$$

$$\text{s.t.} \quad \eta^*(\omega_c) = \arg\min_\eta \sum_{v=1}^{N_L} m^{[v]} G^{[v]}(\eta;\omega_c), \tag{4}$$

where $F^{[v]}$ and $m^{[v]} G^{[v]}$ are known only to learner v.

Remark 3. *Problem (4) and $\arg\min_\eta G(\eta;\omega_c)$ are equivalent because $\sum_{v=1}^{N_L} m^{[v]} G^{[v]}(\eta;\omega_c) = \sum_{v=1}^{N_L} m^{[v]} H(\pi_{\eta;\omega_c}) + \sum_{v=1}^{N_L} m^{[v]} \sum_{i=1}^{N_E} (\omega_r^{[i]})^\top (\mu_r^{[i]}(\pi_{\eta;\omega_c}) - \hat{\mu}_r^{[i,v]}) + \lambda \sum_{v=1}^{N_L} m^{[v]} (J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c}^{[v]}) = m H(\pi_{\eta;\omega_c}) + m \sum_{i=1}^{N_E} (\omega_r^{[i]})^\top (\mu_r^{[i]}(\pi_{\eta;\omega_c}) - \hat{\mu}_r^{[i]}) + \lambda m (J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c}) = m G(\eta;\omega_c)$.*

5 Algorithm and convergence guarantee

In this section, we develop a bi-level distributed learning algorithm to solve problem (3)-(4) and provide convergence rate to the optimal solution of the inner problem (4) and asymptotic convergence to the set of stationary points of the outer problem (3).

Algorithm 1 MEML D-ICRL

Input: $\{\omega_c^{[v]}(0)\}_{v=1}^{N_L}, \{\eta^{[v]}(0)\}_{v=1}^{N_L}, W(k), \bar{W}(n)$

Output: $\omega_c^{[v]}(n), \bar{\eta}^{[v]}(\omega_c^{[v]}(n)), \pi_{\bar{\eta}^{[v]}(\omega_c^{[v]}(n));\omega_c^{[v]}(n)} \quad \forall v \in \mathcal{V}$

1: **for** $n = 0, 1, \cdots$ **do**
2:   **for** $v \in \mathcal{V}$ **do**
3:     Receives $\omega_c^{[v']}(n)$ from $v' \in \bar{\mathcal{N}}^{[v]}(n)$
4:     $\bar{\eta}^{[v]}(\omega_c^{[v]}(n)), \pi_{\bar{\eta}^{[v]}(\omega_c^{[v]}(n));\omega_c^{[v]}(n)} = $ Inner process$(\omega_c^{[v]}(n))$
5:     **if** $n = 0$ **then**
6:       $\bar{\nabla}^{[v]}(0) = \bar{\nabla} F^{[v]}(\omega_c^{[v]}(0), \bar{\eta}^{[v]}(\omega_c^{[v]}(0)))$
7:     **else**
8:       Receives $\bar{\nabla}^{[v']}(n-1)$ from $v' \in \bar{\mathcal{N}}^{[v]}(n)$
9:       $\bar{\nabla}^{[v]}(n) = \sum_{v'=1}^{N_L} \bar{W}^{[vv']}(n) \bar{\nabla}^{[v']}(n-1) + \bar{\nabla} F^{[v]}(\omega_c^{[v]}(n), \bar{\eta}^{[v]}(\omega_c^{[v]}(n))) - \bar{\nabla} F^{[v]}(\omega_c^{[v]}(n-1), \bar{\eta}^{[v]}(\omega_c^{[v]}(n-1)))$
10:     **end if**
11:     $\tilde{\omega}_c^{[v]}(n) = \text{Project}_{\Omega_c}\left(\omega_c^{[v]}(n) + N_L \bar{\nabla}^{[v]}(n)\right)$
12:     $\omega_c^{[v]}(n+1/2) = \omega_c^{[v]}(n) + \beta(n)(\tilde{\omega}_c^{[v]}(n) - \omega_c^{[v]}(n))$
13:     Receives $\omega_c^{[v']}(n+1/2)$ from $v' \in \bar{\mathcal{N}}^{[v]}(n)$
14:     $\omega_c^{[v]}(n+1) = \sum_{v'=1}^{N_L} \bar{W}^{[vv']}(n) \omega_c^{[v']}(n+1/2)$
15:   **end for**
16: **end for**

At outer iteration n in Algorithm 1, the learners first cooperatively solve the inner problem for each learner through *Inner process* (lines 3-4) and then use the obtained result $\bar{\eta}^{[v]}(\omega_c^{[v]}(n))$ to

collaboratively solve the outer problem (3) through the outer process (lines 5-14). In what follows, we will elaborate each process.

Algorithm 2 Inner process($\omega_c^{[v]}$)

Input: $\omega_c^{[v]}$, $\{\eta^{[v]}(0)\}_{v=1}^{N_L}$, $W(k)$

Output: $\bar{\eta}^{[v]}(\omega_c^{[v]})$, $\pi_{\bar{\eta}^{[v]}(\omega_c^{[v]});\omega_c^{[v]}}$

1: **for** $k = 0, 1, \cdots, K - 2$ **do**
2:     **for** $\bar{v} \in \mathcal{V}$ **do**
3:         Receives $\eta^{[\bar{v}']}(\omega_c^{[v]}, k)$ from $\bar{v}' \in \mathcal{N}^{[\bar{v}]}(k)$
4:         $\eta^{[\bar{v}]}(\omega_c^{[v]}, k + 1) = \sum_{v'=1}^{N_L} W^{[\bar{v}\bar{v}']}(k)\eta^{[\bar{v}']}(\omega_c^{[v]}, k) - \alpha(k)\nabla_\eta G^{[\bar{v}]}(\eta^{[\bar{v}]}(\omega_c^{[v]}, k); \omega_c^{[v]})$
5:     **end for**
6: **end for**
7: $\bar{\eta}^{[v]}(\omega_c^{[v]}) \triangleq \frac{\sum_{k=0}^{K-1} \alpha(k)\eta^{[v]}(\omega_c^{[v]}, k)}{\sum_{k=0}^{K-1} \alpha(k)}$
8: $\pi_{\bar{\eta}^{[v]}(\omega_c^{[v]});\omega_c^{[v]}}$ is the corresponding constrained soft Bellman policy.

Inner process. As $\bar{\mathcal{N}}^{[v]}(n) = \mathcal{V}$, learner v knows the cost weight vector of all the learners. Given $\omega_c^{[v]}$, at inner iteration k, each learner $\bar{v}$ receives $\eta^{[\bar{v}']}(\omega_c^{[v]}, k)$ from neighbors and updates its estimate of the reward weight vector (embedded in $\eta^{[\bar{v}]}$). In specific, at line 4 in Algorithm 2, the first term (convex combination) on the right hand side encourages consensus and the second term (gradient) drives to the optimal solution. It is shown in Lemma 3 that each learner can reach $\eta^*(\omega_c^{[v]})$ at the rate of $O(\frac{1}{\sqrt{\log K}})$.

Outer process. Once obtaining $\bar{\eta}^{[v]}(\omega_c^{[v]})$, each learner v faces three challenges in solving the outer problem: (i) the gradient of $F^{[v]}(\omega_c^{[v]}, \eta^*(\omega_c^{[v]}))$ cannot be obtained because the inner problem is not fully solved, i.e., learner v does not find $\eta^*(\omega_c^{[v]})$ but its approximation $\bar{\eta}^{[v]}(\omega_c^{[v]})$, (ii) $\sum_{v' \neq v} F^{[v']}$ is unknown, and (iii) the local likelihood function $F^{[v]}$ is non-convex. To solve these challenges respectively, we propose three techniques: local gradient approximation (LGA), global gradient approximation (GGA), and local successive convex approximation (LSCA). By subtly designing these three approximations, our algorithm is guaranteed to converge to the set of stationary points. Notice that the second challenge also arises in the inner problem but we solve it in a simpler way because the inner problem is convex.

To approximate the local gradient $\nabla F^{[v]}(\omega_c, \eta^*(\omega_c))$, we propose the following LGA of learner v: $\bar{\nabla} F^{[v]}(\omega_c, \bar{\eta}(\omega_c)) = \sum_{\zeta^j \in \mathcal{D}^{[v]}} \sum_{t=0}^{\infty} \gamma^t \phi_c(s_t^j, a_t^j) - m^{[v]} E_{S,A}^{\pi_{\bar{\eta}(\omega_c);\omega_c}} [\sum_{t=0}^{\infty} \gamma^t \phi_c(S_t, A_t)]$, whose derivation is based on the strict convexity of $G(\eta; \omega_c)$ and $G^{[v]}(\eta; \omega_c)$. The derivation and the proof of strict convexity are both included in the Appendix. Moreover, it can be shown (in the Appendix) that the approximation error (defined in the Appendix) is upper bounded and reduces to zero as $K \to \infty$.

While learner v can use LGA to get a sound approximation of $\nabla F^{[v]}$, in order to solve problem (3), it still needs the knowledge of the gradients $\sum_{v' \neq v} \nabla F^{[v']}$ of other learners to get the global gradient. Therefore, we leverage a GGA $\bar{\nabla}^{[v]}$ (lines 5-10) who aims to track the average global gradient $\frac{1}{N_L} \nabla F(\omega_c^{[v]}, \eta^*(\omega_c^{[v]}))$. However, $\nabla F(\omega_c^{[v]}, \eta^*(\omega_c^{[v]}))$ is not available due to the inaccessibility of $\eta^*(\omega_c^{[v]})$. Therefore, $\bar{\nabla}^{[v]}$ is designed to track the approximation $\frac{1}{N_L} \sum_{v'=1}^{N_L} \bar{\nabla} F^{[v']}(\omega_c^{[v']}, \bar{\eta}^{[v']}(\omega_c^{[v']}))$ of the average global gradient.

To tackle the non-convexity issue, learner v solves a LSCA problem (line 11) and then follows two-step updates of the local cost weight vector $\omega_c^{[v]}$. LSCA proposes to solve a local convexification of problem (3). In specific, learner v solves the problem $\arg\max_{\omega_c \in \Omega_c} \tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ where $-\tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ is a local strongly-convex surrogate of $-F(\omega_c, \eta^*(\omega_c))$ at $\omega_c^{[v]}$ satisfying the gradient consistency condition $\nabla \tilde{F}^{[v]}(\omega_c; \omega_c^{[v]}) = N_L \bar{\nabla}^{[v]}$. An attractive property of LSCA [27] is

that its fixed point exists and is a stationary point of functions with gradient $N_L \bar{\nabla}^{[v]}$ under certain conditions (in the Appendix).

Once obtaining $\tilde{\omega}_c^{[v]}$, learner v executes two-step updates. The first update towards $\tilde{\omega}_c^{[v]}$ (line 12) drives learner v to a stationary point of problem (3) and the second update (lines 13-14) encourages the consensus among different learners.

Lemma 2. *The result $\tilde{\omega}_c^{[v]}(n)$ at line 11 in Algorithm 1 is the optimal solution of a LSCA problem at $\omega_c^{[v]}(n)$ under the GGA $N_L \bar{\nabla}^{[v]}(n)$.*

Lemma 3. *(Convergence rate of the inner problem) Suppose Assumptions 1 and 2 hold and let $\alpha(k) = \frac{\bar{\alpha}}{k+1}$ where $\bar{\alpha}$ is a positive constant. Then, for every learner $v \in \mathcal{V}$ in Algorithm 2,*

$$||\bar{\eta}^{[v]}(\omega_c) - \eta^*(\omega_c)|| \leq \frac{C_1^{[v]}}{\log K} + \frac{C_2^{[v]}}{\sqrt{\log K}},$$

where $C_1^{[v]}$ and $C_2^{[v]}$ are positive constants whose expression can be found in the Appendix.

We define the set of KKT points (also called stationary points) of problem (3)-(4) as [28] [29]: $\Omega_c^* \triangleq \{\omega_c^* \in \Omega_c : \max_{\omega_c \in \Omega_c}\{(\nabla F(\omega_c^*, \eta^*(\omega_c^*)))^\top(\omega_c - \omega_c^*)\} = 0\}$, and study the convergence property of the following metric [30]: $J(\omega_c(n)) \triangleq \max_{\omega_c \in \Omega_c}\{(\nabla F(\omega_c(n), \eta^*(\omega_c(n))))^\top(\omega_c - \omega_c(n))\}$.

Theorem 1. *(Asymptotic convergence of the outer problem) Suppose Assumptions 1, 2, and 3 hold. Let $\beta(n) \in (0,1)$, $\sum_{n=0}^{\infty} \beta(n) = +\infty$, and $\sum_{n=0}^{\infty}(\beta(n))^2 < +\infty$, then in Algorithm 1:*

$$(consensus): \quad \lim_{n\to\infty} \max_{v,v'\in\mathcal{V}} ||\omega_c^{[v]}(n) - \omega_c^{[v']}(n)|| = 0,$$

$$(convergence): \quad \limsup_{n\to\infty} J(\omega_c^{[v]}(n)) \leq \frac{\bar{M}}{\sqrt{\log K}},$$

where $\bar{M}$ is a positive constant whose existence is proved in the Appendix.

Lemma 3 and Theorem 1 show that the distributed learners can achieve common values of ω_c and η. As the constrained soft Bellman policy is continuous in (ω_c, η) (proved in the Appendix), they will also achieve a common policy.

6 Simulation

This section presents two simulation examples. In the first example, the experts are programmed to follow the optimal policy under discrete state and action spaces. In the second example, the experts are humans, i.e., may not be optimal, and the states and actions are both continuous.

6.1 Synthetic grid world

We consider the grid world introduced in [11]. Different from paper [11]'s MDP, we implement a CMG on this environment where three experts perform motion planning from $s_0/s_0'/s_0''$ to $s_G/s_G'/s_G''$ while avoiding collision with obstacles and each other. The environment consists of a 9-by-9 grid of states, and the actions of each expert are to stay still, move up, down, left, right, or diagonally by one cell. The dynamics is stochastic as each action of an expert has 20% probability of failure, resulting in staying still. While the grid world looks small, the CMG has more than $387,000,000$ state-action pairs due to multiple experts. Each state-action pair produces a distance-relevant reward feature and the ground truth reward of each expert increases if the distance to its goal decreases. The true CMG (Figure 2a), from which the experts generate demonstration, includes constraints (i.e., obstacles) and the nominal MG does not. We use heat maps to show the relevant visitation frequency of each state and action pair. Each expert will terminate its motion if violating a constraint, i.e, colliding with an obstacle.

In our algorithm, there are four distributed learners where each learner can only respectively get 10, 20, 30, 40 demonstrated trajectories of the experts. Notice that the learners and experts are not the same entities, i.e., the learners stand aside observing the experts. The four learners jointly

recover the experts' policies by learning the constraints and the reward functions over an underlying communication network. The detailed simulation setup is included in the Appendix.

Furthermore, we use the following three baselines for comparisons:

ME-greedy: This method is introduced in [11] where a greedy method is used to estimate the constraints based on the ME framework.

MCE-greedy: We construct this method by extending ME-greedy to the same greedy method based on the MCE framework which is more suitable for stochastic environments.

Centralized inverse constrained inverse reinforcement learning (C-ICRL): This method is a special case of D-ICRL where there is a centralized learner obtaining all the demonstration data.

Notice that ME-greedy and MCE-greedy assume the access to the ground truth rewards while C-ICRL does not have this assumption. All the three baselines are centralized learning where a single learner can get the total 100 demonstrated trajectories. Our algorithm is distributed learning and does not obtain the ground truth rewards, therefore, it solves a more difficult problem than the baselines do. However, the simulation shows that our algorithm can be on par with and even outperform the baselines in some aspects. In our bi-level algorithm, the outer loop is to estimate the constraints and the inner loop is to recover the corresponding policy by estimating the reward function. To reason about the performance of our algorithm, we use five metrics: cumulative rewards (CR), false positive rate (FPR), false negative rate (FNR), constraint violation rate (CVR), and reward function distance (RFD). The CR, commonly used in IRL and imitation learning literature [31, 32], illustrates the total reward collected by the three experts in an episode, FPR, introduced in [11], is the proportion of learned constraints that are not the ground truth constraints, FNR is the proportion of the ground truth constraints that are not learned, CVR [12] is the average percentage of the three experts' violating any constraint in an episode. Since the reward function is a linear combination of the reward features, we propose RFD to measure the distance between the learned reward weight vector ω_r and experts' reward weight vector ω_{r_E}: $||\omega_r - \omega_{r_E}||/||\omega_{r_E}||$.

Figure 2: Algorithm performance on a synthetic grid world CMG. In subfigures 2a 2b, each state and action cell is colored according to the relevant visitation frequency and the red crosses represent the constraints. In subfigures 2c-2g, D-learners 1-4 are distributed learners in our algorithm. Because ME-greedy and MCE-greedy are not bi-level, we only include their curves against outer iterations.

Figure 2b shows the constraints learned by our algorithm (each learner recovers the same constraints). Several action constraints are not recovered due to the reason that the experts will barely take these actions even if they are feasible. Figures 2c 2f show that our learning algorithm can successfully imitate the experts' behavior according to CR and CVR. Figures 2d 2e show that D-ICRL

and C-ICRL have better FPR performance compared to the greedy-based methods while MCE-greedy has the best FNR performance. From Figure 2e and the small window in Figure 2c, we can see that the distributed learners can quickly achieve consensus even if they start from different initial conditions in both outer and inner loops. Figures 2c and 2g show the performance of inner process in the last outer iteration. Figure 2g shows that D-ICRL and C-ICRL can approach the ground truth reward functions and stabilize.

Table 1: Performance Comparisons. Here, D means distributed and NATR means no access to the (ground truth) rewards.

		D	NATR	CR	FPR	FNR	CVR	RFD
Experts (ground truth)		N/A	N/A	2002.59 ± 125.71	N/A	N/A	0.000	0.000
D-ICRL (our method)	D-Learner 1	✓	✓	1890.67 ± 135.22	0.00	0.18	0.013	0.748
	D-Learner 2	✓	✓	1873.07 ± 162.14	0.00	0.18	0.020	0.750
	D-Learner 3	✓	✓	1878.51 ± 131.55	0.00	0.18	0.020	0.749
	D-Learner 4	✓	✓	1884.85 ± 167.16	0.00	0.18	0.010	0.749
C-ICRL		×	✓	1884.74 ± 158.90	0.00	0.18	0.013	0.731
ME-greedy		×	×	1162.24 ± 132.66	0.18	0.24	0.010	N/A
MCE-greedy		×	×	1776.84 ± 300.61	0.09	0.12	0.057	N/A
Experts (nominal)		N/A	N/A	18.64 ± 3.02	N/A	N/A	1.000	0.000

Table 1 shows the comparison result where each learner in our distributed algorithm can be on par with C-ICRL in all the five metrics. Our algorithm can outperform ME-greedy introduced in [11] in almost every metric except being on par with it in CVR. While MCE-greedy is centralized learning and assumes the access to the ground truth rewards, our method can still outperform it in three metrics, i.e., CR, FPR, and CVR.

6.2 Drones motion planning with obstacles

In the second example, we simulate in a physical setting with continuous state-action space. We build a simulator in Gazebo where each drone aims to reach the door in its diagonal direction while avoiding collisions. We first control the simulated drones to their target doors, record nine pairs of trajectories, and distribute four and five pairs to two learners respectively. The demonstration is shown in Figure 3b and each cell in the figure represents the constraint set indicated by a cost feature of each expert, i.e., $\mathcal{C}_j^{[i]}$. We define the reward feature of each state as its location multiplied with a coefficient. Figure 3c shows the learned constraints and trajectories.

(a) Simulator environment   (b) Demonstrated trajectories   (c) Learned trajectories

Figure 3: Drones motion planning with obstacles

We keep the three metrics, FPR, FNR, and CVR in last example. Because the ground truth reward is not accessible in this physical setting, we cannot use RFD and we use success rate (SR) instead

of CR. The SR is defined as the percentage of the drones' successfully reaching the doors without any collision. We use the reward function learned from D-ICRL as the input reward function of ME-greedy. Because this environment is deterministic, MCE-greedy reduces to ME-greedy.

Table 2: Performance Comparisons.

		D	NATR	FPR	FNR	CVR	SR
D-ICRL	D-Learner 1	✓	✓	0.028	0.00	0.03	0.97
	D-Learner 2	✓	✓	0.028	0.00	0.03	0.97
C-ICRL		×	✓	0.018	0.00	0.02	0.98
M(C)E-greedy		×	×	0.037	0.13	0.16	0.84

7 Discussion and future work

We propose D-ICRL, the first multi-expert-multi-learner MCE ICRL framework that is effective to CMG with both discrete and continuous environments. We employ MLE to estimate the constraints, use an MCE-based optimization problem to learn the corresponding reward function and policy, and derive our algorithm based on a novel distributed bi-level optimization theoretical framework. Experimental results show that D-ICRL can imitate the experts' behavior as well as recover the environmental constraints. Despite its benefits, the limitations are that (i) the learners assume that the reward/cost function is a linear combination of some features; (ii) the communication between learners is frequent. We will address the issues in the future.

8 Acknowledgements

This work is partially supported by National Science Foundation through grants CNS 1830390, ECCS 1846706, and ECCS 2140175. We would like to thank the reviewers for their insightful and constructive suggestions.

References

[1] K. Zhang, Z. Yang, and T. Başar, "Multi-agent reinforcement learning: A selective overview of theories and algorithms," *Handbook of Reinforcement Learning and Control*, pp. 321–384, 2021.

[2] J. Fu, K. Luo, and S. Levine, "Learning robust rewards with adverserial inverse reinforcement learning," in *International Conference on Learning Representations*, 2018.

[3] L. Yu, J. Song, and S. Ermon, "Multi-agent adversarial inverse reinforcement learning," in *International Conference on Machine Learning*, pp. 7194–7201, 2019.

[4] W. Jeon, C.-Y. Su, P. Barde, T. Doan, D. Nowrouzezahrai, and J. Pineau, "Regularized inverse reinforcement learning," in *International Conference on Learning Representations*, 2020.

[5] M. Imani and S. F. Ghoreishi, "Scalable inverse reinforcement learning through multifidelity Bayesian optimization," *IEEE Transactions on Neural Networks and Learning Systems*, pp. 1–8, 2021.

[6] A. Y. Ng and S. J. Russell, "Algorithms for inverse reinforcement learning," in *International Conference on Machine Learning*, pp. 663–670, 2000.

[7] P. Abbeel and A. Y. Ng, "Apprenticeship learning via inverse reinforcement learning," in *International Conference on Machine Learning*, pp. 1–8, 2004.

[8] N. D. Ratliff, J. A. Bagnell, and M. A. Zinkevich, "Maximum margin planning," in *International Conference on Machine Learning*, pp. 729–736, 2006.

[9] B. D. Ziebart, A. L. Maas, J. A. Bagnell, and A. K. Dey, "Maximum entropy inverse reinforcement learning," in *National Conference on Artificial intelligence*, vol. 8, pp. 1433–1438, 2008.

[10] B. D. Ziebart, J. A. Bagnell, and A. K. Dey, "Modeling interaction via the principle of maximum causal entropy," in *International Conference on Machine Learning*, pp. 1255–1262, 2010.

[11] D. R. Scobee and S. S. Sastry, "Maximum likelihood constraint inference for inverse reinforcement learning," in *International Conference on Learning Representations*, 2019.

[12] S. Malik, U. Anwar, A. Aghasi, and A. Ahmed, "Inverse constrained reinforcement learning," in *International Conference on Machine Learning*, pp. 7390–7399, 2021.

[13] S. Natarajan, G. Kunapuli, K. Judah, P. Tadepalli, K. Kersting, and J. Shavlik, "Multi-agent inverse reinforcement learning," in *International Conference on Machine Learning and Applications*, pp. 395–400, 2010.

[14] T. S. Reddy, V. Gopikrishna, G. Zaruba, and M. Huber, "Inverse reinforcement learning for decentralized non-cooperative multiagent systems," in *IEEE International Conference on Systems, Man, and Cybernetics*, pp. 1930–1935, 2012.

[15] J. Inga, E. Bischoff, F. Köpf, M. Flad, and S. Hohmann, "Inverse cooperative and non-cooperative dynamic games based on maximum entropy inverse reinforcement learning," *arXiv preprint arXiv:1911.07503*, 2019.

[16] W.-C. Ma, D.-A. Huang, N. Lee, and K. M. Kitani, "Forecasting interactive dynamics of pedestrians with fictitious play," in *IEEE Conference on Computer Vision and Pattern Recognition*, pp. 774–782, 2017.

[17] K. Bogert and P. Doshi, "Multi-robot inverse reinforcement learning under occlusion with interactions," in *International Conference on Autonomous Agents and Multi-agent Systems*, pp. 173–180, 2014.

[18] J. Verbraeken, M. Wolting, J. Katzy, J. Kloppenburg, T. Verbelen, and J. S. Rellermeyer, "A survey on distributed machine learning," *ACM Computing Surveys*, vol. 53, no. 2, pp. 1–33, 2020.

[19] N. Couellan and W. Wang, "On the convergence of stochastic bi-level gradient methods," *Optimization*, 2016.

[20] S. Ghadimi and M. Wang, "Approximation methods for bilevel programming," *arXiv preprint arXiv:1802.02246*, 2018.

[21] M. Hong, H.-T. Wai, Z. Wang, and Z. Yang, "A two-timescale framework for bilevel optimization: Complexity analysis and application to actor-critic," *arXiv preprint arXiv:2007.05170*, 2020.

[22] K. Ji, J. Yang, and Y. Liang, "Bilevel optimization: Convergence analysis and enhanced design," in *International Conference on Machine Learning*, pp. 4882–4892, 2021.

[23] F. Yousefian, "Bilevel distributed optimization in directed networks," in *American Control Conference*, pp. 2230–2235, 2021.

[24] E. Altman and A. Shwartz, "Constrained markov games: Nash equilibria," in *Advances in Dynamic Games and Applications*, pp. 213–221, 2000.

[25] Z. Zhou, M. Bloem, and N. Bambos, "Infinite time horizon maximum causal entropy inverse reinforcement learning," *IEEE Transactions on Automatic Control*, vol. 63, no. 9, pp. 2787–2802, 2017.

[26] T. Haarnoja, H. Tang, P. Abbeel, and S. Levine, "Reinforcement learning with deep energy-based policies," in *International Conference on Machine Learning*, pp. 1352–1361, 2017.

[27] F. Facchinei, G. Scutari, and S. Sagratella, "Parallel selective algorithms for nonconvex big data optimization," *IEEE Transactions on Signal Processing*, vol. 63, no. 7, pp. 1874–1889, 2015.

[28] P. Bianchi and J. Jakubowicz, "Convergence of a multi-agent projected stochastic gradient algorithm for non-convex optimization," *IEEE Transactions on Automatic Control*, vol. 58, no. 2, pp. 391–405, 2012.

[29] P. Di Lorenzo and G. Scutari, "Next: In-network nonconvex optimization," *IEEE Transactions on Signal and Information Processing over Networks*, vol. 2, no. 2, pp. 120–136, 2016.

[30] H.-T. Wai, J. Lafond, A. Scaglione, and E. Moulines, "Decentralized frank–wolfe algorithm for convex and nonconvex problems," *IEEE Transactions on Automatic Control*, vol. 62, no. 11, pp. 5522–5537, 2017.

[31] X. Wang and D. Klabjan, "Competitive multi-agent inverse reinforcement learning with sub-optimal demonstrations," in *International Conference on Machine Learning*, pp. 5143–5151, 2018.

[32] V. Tangkaratt, N. Charoenphakdee, and M. Sugiyama, "Robust imitation learning from noisy demonstrations," in *International Conference on Artificial Intelligence and Statistics*, pp. 298–306, 2021.

[33] V. I. Bogachev and O. G. Smolyanov, *Real and functional analysis*. Springer, 2020.

[34] W. Rudin, *Principles of mathematical analysis*. McGraw-hill New York, 1976.

[35] E. E. Tyrtyshnikov, *A brief introduction to numerical analysis*. Springer Science & Business Media, 1997.

[36] A. Nedic and A. Ozdaglar, "Distributed subgradient methods for multi-agent optimization," *IEEE Transactions on Automatic Control*, vol. 54, no. 1, pp. 48–61, 2009.

[37] G. Scutari and Y. Sun, "Parallel and distributed successive convex approximation methods for big-data optimization," in *Multi-agent Optimization*, pp. 141–308, 2018.

[38] G. Scutari and Y. Sun, "Distributed nonconvex constrained optimization over time-varying digraphs," *Mathematical Programming*, vol. 176, no. 1, pp. 497–544, 2019.

[39] J. Meyer, A. Sendobry, S. Kohlbrecher, U. Klingauf, and O. von Stryk, "Comprehensive simulation of quadrotor uavs using ros and gazebo," in *International Conference on Simulation, Modeling and Programming for Autonomous Robots*, p. to appear, 2012.

[40] N. Das, S. Bechtle, T. Davchev, D. Jayaraman, A. Rai, and F. Meier, "Model-based inverse reinforcement learning from visual demonstrations," in *Conference on Robot Learning*, pp. 1930–1942, 2021.

A   Appendix: notions and notations

We define the concatenated reward feature vector as $\phi_r \triangleq [(\phi_r^{[1]})^\top, \cdots, (\phi_r^{[N_E]})^\top]^\top$ and the augmented reward feature under ω_c as $\phi_{r,\omega_c} \triangleq [\phi_r^\top, c_{\omega_c}]^\top$. The augmented reward feature expectation starting from state-action pair (s,a) under policy π is $\mu_{r,\omega_c}^\pi(s,a) \triangleq \phi_{r,\omega_c}(s,a) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)\mu_{r,\omega_c}^\pi(s')ds'$ and the augmented reward feature expectation starting from state s is $\mu_{r,\omega_c}^\pi(s) \triangleq \int_{a\in\mathcal{A}} \pi(a|s)\mu_{r,\omega_c}^\pi(s,a)da$. Moreover, we denote $\mu_{r,\omega_c}(\pi) \triangleq [(\mu_r(\pi))^\top, J_{\omega_c}(\pi)]^\top$. The empirical augmented reward feature expectation vector is defined as $\hat{\mu}_{r,\omega_c} \triangleq [\hat{\mu}_r^\top, \hat{b}_{\omega_c}]^\top$ and its estimate from learner v is defined as $\hat{\mu}_{r,\omega_c}^{[v]} \triangleq [(\hat{\mu}_r^{[v]})^\top, \hat{b}_{\omega_c}^{[v]}]^\top$. For a given vector $\bar{\eta}$, we define the augmented Q-function as $Q_{\bar{\eta},\omega_c}^\pi(s,a) = \bar{\eta}^\top \mu_{r,\omega_c}^\pi(s,a)$ and the augmented value-function as $V_{\bar{\eta},\omega_c}^\pi(s) = \bar{\eta}^\top \mu_{r,\omega_c}^\pi(s)$. The concatenated cost feature expectation starting from state-action pair (s,a) under policy π is $\mu_c^\pi(s,a) \triangleq \phi_c(s,a) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)\mu_c^\pi(s')ds'$ and the concatenated cost feature expectation starting from state s is $\mu_c^\pi(s) \triangleq \int_{a\in\mathcal{A}} \pi(a|s)\mu_c^\pi(s,a)da$. We denote $\mu_c(\pi) \triangleq E_{S,A}^\pi[\sum_{t=0}^\infty \gamma^t \phi_c(S_t, A_t)]$. The empirical concatenated cost feature expectation is $\hat{\mu}_c \triangleq \frac{1}{m}\sum_{j=1}^m \sum_{t=0}^\infty \gamma^t \phi_c(s_t^j, a_t^j)$. The causal entropy staring from (s,a) under policy π is $H^\pi(s,a) = -\ln\pi(a|s) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)H^\pi(s')ds'$ and the casual entropy from state s is $H^\pi(s) = \int_{a\in\mathcal{A}} \pi(a|s)H^\pi(a|s)da$. We define the state-action visitation frequency as $\psi^\pi(s,a) \triangleq E^\pi[\sum_{t=0}^\infty \gamma^t \mathbb{1}\{S_t = s\}\mathbb{1}\{A_t = a\}]$ and state visitation frequency as $\psi^\pi(s) \triangleq E^\pi[\sum_{t=0}^\infty \gamma^t \mathbb{1}\{S_t = s\}]$, where $\mathbb{1}\{\cdot\}$ is the indicator function. It is obvious that $\psi^\pi(s) \leq \frac{1}{1-\gamma}$ for any $s \in \mathcal{S}$ and $\int_{s\in\mathcal{S}} \psi^\pi(s)ds = \frac{1}{1-\gamma}$. We use $\mathbf{0}_{m\times n}$ to denote an $m \times n$ matrix whose entries are all zero.

Lemma 4. *For any $(s,a) \in \mathcal{S} \times \mathcal{A}$, any $\omega_c \in \Omega_c$, and any π, $||\mu_{r,\omega_c}^\pi(s)||$, $||\mu_{r,\omega_c}^\pi(s,a)||$, $||\mu_c^\pi(s)||$, and $||\mu_c^\pi(s,a)||$ are bounded.*

Proof. We know that $\mu_{r,\omega_c}^\pi(s,a) = \phi_{r,\omega_c}(s,a) + E_{S,A}^\pi[\sum_{t=1}^\infty \gamma^t \phi_{r,\omega_c}(S_t, A_t)|S_0 = s, A_0 = a]$. Since $||\phi_{r,\omega_c}(s,a)|| \leq \sqrt{\sum_{i=1}^{N_E}(l_r^{[i]}d_1^2) + (\sum_{i=1}^{N_E} l_c^{[i]}d_2)^2)}$, we know that $||\mu_{r,\omega_c}^\pi(s,a)|| \leq \frac{1}{1-\gamma}\sqrt{\sum_{i=1}^{N_E}(l_r^{[i]}d_1^2) + (\sum_{i=1}^{N_E} l_c^{[i]}d_2)^2)}$. Because $\mu_{r,\omega_c}^\pi(s) = \int_{a\in\mathcal{A}} \pi(a|s)\mu_{r,\omega_c}^\pi(s,a)da$, we can see that $||\mu_{r,\omega_c}^\pi(s)||$ is bounded given that $||\mu_{r,\omega_c}^\pi(s,a)||$ is bounded. Analogously, $||\mu_c^\pi(s)||$ and $||\mu_c^\pi(s,a)||$ are also bounded. $\qquad\square$

Constrained soft Bellman policy. We provide the formula of the constrained soft Bellman policy under continuous state-action space and the policy can be approximated through soft Q learning [26]:

$$\pi_{\eta;\omega_c}(a|s) = \frac{\exp(Q_{\eta;\omega_c}^{\text{soft}}(s,a))}{\exp(V_{\eta;\omega_c}^{\text{soft}}(s))}, \qquad V_{\eta;\omega_c}^{\text{soft}}(s) = \ln\Big(\int_{a\in\mathcal{A}} \exp(Q_{\eta;\omega_c}^{\text{soft}}(s,a))\,da\Big),$$

$$Q_{\eta;\omega_c}^{\text{soft}}(s,a) = \sum_{i=1}^{N_E}(\omega_r^{[i]})^\top \phi_r^{[i]}(s,a) + \lambda c_{\omega_c}(s,a) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)V_{\eta;\omega_c}^{\text{soft}}(s')\,ds'.$$

It is obvious that the constrained soft Bellman policy is continuous in (ω_c, η) as it is a composition of continuous functions of (ω_c, η).

B   Appendix: proof

This section focuses on the continuous state-action space and has seven subsections. Subsection B.1 includes some preliminary results for the later subsections, subsection B.2 presents the proof of Lemma 1, subsection B.3 provides some intermediate results for the remaining subsections, subsection B.4 proves Lemma 3, subsection B.5 presents the derivation of LGA, subsection B.6 proves Lemma 2, and subsection B.7 presents the proof of the theorem.

B.1   Preliminary results

In this section, we prove two lemmas which serve as building blocks for the remaining subsections.

Lemma 5. *The gradient* $\nabla_\eta \ln \pi_{\eta;\omega_c}(a|s) = \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)$.

Proof. Define $Z_{s,\eta;\omega_c} \triangleq \exp(V_{\eta;\omega_c}^{\text{soft}}(s))$ and $Z_{a|s,\eta;\omega_c} \triangleq \exp(Q_{\eta;\omega_c}^{\text{soft}}(s,a))$, therefore $Z_{a|s,\eta;\omega_c}$ is smooth in η because it is a composition of logarithmic, exponential and linear functions of η. From Leibniz integral rule, we know that $\nabla_\eta \int_a Z_{a|s,\eta;\omega_c} da = \int_a \nabla_\eta Z_{a|s,\eta;\omega_c} da$. Therefore,

$$\nabla_\eta \ln Z_{s,\eta;\omega_c} = \frac{\int_{a\in\mathcal{A}} \nabla_\eta Z_{a|s,\eta;\omega_c} da}{Z_{s,\eta;\omega_c}} = \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\left[\phi_{r,\omega_c}(s,a) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)\cdot\right.$$

$$\left. \nabla_\eta \ln Z_{s',\eta;\omega_c} ds'\right] da,$$

$$= \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\left\{\phi_{r,\omega_c}(s,a) + \gamma \int_{s'\in\mathcal{S}} P(s'|s,a)\int_{a'\in\mathcal{A}} \pi_{\eta;\omega_c}(a'|s')\left[\phi_{r,\omega_c}(s',a')\right.\right.$$

$$\left.\left. + \gamma \int_{s''\in\mathcal{S}} P(s''|s',a')\nabla_\eta \ln Z_{s'',\eta;\omega_c} ds''\right] da'ds'\right\} da.$$

Continuing the expansion, we can get:

$$\nabla_\eta \ln Z_{s,\eta;\omega_c} = E_{S,A}^{\pi_{\eta;\omega_c}}\left[\sum_{t=0}^{\infty} \gamma^t \phi_{r,\omega_c}(S_t, A_t)|S_0 = s\right] = \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s),$$

$$\nabla_\eta \ln Z_{a|s,\eta;\omega_c} = \phi_{r,\omega_c}(s,a) + E_{S,A}^{\pi_{\eta;\omega_c}}\left[\sum_{t=1}^{\infty} \gamma^t \phi_{r,\omega_c}(S_t, A_t)|S_0 = s, A_0 = a\right] = \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a).$$

Therefore, $\nabla_\eta \ln \pi_{\eta;\omega_c}(a|s) = \nabla_\eta \ln Z_{a|s,\eta;\omega_c} - \nabla_\eta \ln Z_{s,\eta;\omega_c} = \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)$. □

Lemma 6. *The gradients* $\nabla_\eta \mu_{r,\omega_c}(\pi_{\eta;\omega_c}) = \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)\int_{a\in\mathcal{A}} \nabla_\eta \pi_{\eta;\omega_c}(a|s)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a))^\top da$ ds, $\nabla_\eta H(\pi_{\eta;\omega_c}) = \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)\int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\nabla_\eta \ln \pi_{\eta;\omega_c}(a|s)(H^{\pi_{\eta;\omega_c}}(s,a) - 1)dads$.

Proof.

$$\nabla_\eta \mu_{r,\omega_c}(\pi_{\eta;\omega_c}) = \int_{s_0\in\mathcal{S}} P_0(s_0)\int_{a_0\in\mathcal{A}}\left[\nabla_\eta \pi_{\eta;\omega_c}(a_0|s_0)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_0,a_0))^\top\right.$$

$$\left. + \pi_{\eta;\omega_c}(a_0|s_0)\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_0,a_0)\right] da_0 ds_0,$$

$$= \int_{s_0\in\mathcal{S}} P_0(s_0)\int_{a_0\in\mathcal{A}}\left[\nabla_\eta \pi_{\eta;\omega_c}(a_0|s_0)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_0,a_0))^\top + \pi_{\eta;\omega_c}(a_0|s_0)\nabla_\eta\left(\phi_{r,\omega_c}(s_0,a_0)\right.\right.$$

$$\left.\left. + \gamma \int_{s_1\in\mathcal{S}} P(s_1|s_0,a_0)\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_1)ds_1\right)\right] da_0 ds_0,$$

$$= \int_{s_0\in\mathcal{S}} P_0(s_0)\int_{a_0\in\mathcal{A}}\left[\nabla_\eta \pi_{\eta;\omega_c}(a_0|s_0)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_0,a_0))^\top + \pi_{\eta;\omega_c}(a_0|s_0)\gamma \int_{s_1\in\mathcal{S}} P(s_1|s_0,a_0)\cdot\right.$$

$$\left. \nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_1)ds_1\right] da_0 ds_0,$$

$$= \int_{s_0 \in \mathcal{S}} P_0(s_0) \int_{a_0 \in \mathcal{A}} \Big\{ \nabla_\eta \pi_{\eta;\omega_c}(a_0|s_0)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_0,a_0))^\top + \gamma \pi_{\eta;\omega_c}(a_0|s_0) \int_{s_1 \in \mathcal{S}} P(s_1|s_0,a_0) \cdot$$

$$\int_{a_1 \in \mathcal{A}} \Big[\nabla_\eta \pi_{\eta;\omega_c}(a_1|s_1)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_1,a_1))^\top + \pi_{\eta;\omega_c}(a_1|s_1) \int_{s_2 \in \mathcal{S}} P(s_2|s_1,a_1) \nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s_2) ds_2 \Big]$$

$$da_1 ds_1 \Big\} da_0 ds_0.$$

Keep the expansion and we can get:

$$\nabla_\eta \mu_{r,\omega_c}(\pi_{\eta;\omega_c}) = \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \nabla_\eta \pi_{\eta;\omega_c}(a|s)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a))^\top da ds,$$

$$= \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \pi_{\eta;\omega_c}(a|s) \nabla_\eta \ln \pi_{\eta;\omega_c}(a|s)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a))^\top da ds,$$

Analogously, we have that

$$\nabla_\eta H(\pi_{\eta;\omega_c}) = \int_{s_0 \in \mathcal{S}} P_0(s_0) \int_{a_0 \in \mathcal{A}} \Big[\nabla_\eta \pi_{\eta;\omega_c}(a_0|s_0)(H^{\pi_{\eta;\omega_c}}(s_0,a_0))$$

$$+ \pi_{\eta;\omega_c}(a_0|s_0) \nabla_\eta \Big(-\ln \pi_{\eta;\omega_c}(a_0|s_0) + \gamma \int_{s_1 \in \mathcal{S}} P(s_1|s_0,a_0) H^{\pi_{\eta;\omega_c}}(s_1) ds_1 \Big) \Big] da_0 ds_0.$$

Keep the expansion, we have:

$$\nabla_\eta H(\pi_{\eta;\omega_c}) = \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \pi_{\eta;\omega_c}(a|s) \nabla_\eta \ln \pi_{\eta;\omega_c}(a|s)(H^{\pi_{\eta;\omega_c}}(s,a) - 1) da ds.$$

$\square$

B.2 Proof of Lemma 1

The Lagrangian of problem (2) is $L(\pi, \eta; \omega_c) = H(\pi) + \sum_{i=1}^{N_E}(\omega_r^{[i]})^\top(\mu_r^{[i]}(\pi) - \hat{\mu}_r^{[i]}) + \lambda(J_{\omega_c}(\pi) - \hat{b}_{\omega_c})$. To find the maximizer of $\max_{\pi \in \Pi} L(\pi, \eta; \omega_c)$, we remove the constant terms and formulate the following problem:

$$\arg\max_{\pi \in \Pi} \sum_{t=0}^\infty \gamma^t E_{S,A}^\pi \Big[\sum_{i=1}^{N_E}(\omega_r^{[i]})^\top \phi_r^{[i]}(S_t, A_t) + \lambda c_{\omega_c}(S_t, A_t) - \ln \pi(A_t|S_t) \Big]. \tag{5}$$

From [26], we can see that the continuous constrained soft Bellman policy is the optimal solution of (5) by setting the hyperparameter $\alpha = 1$ in equation (2) in [26].

Let p^* be the optimal value of the primal problem (2) and d^* be the optimal value of the dual problem $\min_\eta G(\eta; \omega_c)$. We know that p^* exists (proof of Lemma 2 in [25]). For any η, $G(\eta; \omega_c)$ is an upper bound of p^* because any optimal solution of the primal problem (2) is a feasible solution of $\max_{\pi \in \Pi} L(\pi, \eta; \omega_c)$, therefore, d^* is finite.

We change the policy π to be time-dependent but force it to be stationary, then

$$\frac{\partial L(\pi, \eta; \omega_c)}{\partial \pi^t(a|s)} = -\gamma^t P(S_t = s)(\ln \pi^t(a|s) + 1)$$

$$+ P(S_t = s) E_{S,A}^\pi \Big[\sum_{\tau=t+1}^\infty -\gamma^\tau \ln \pi^\tau(A_\tau|S_\tau)|S_t = s, A_t = a \Big] + P(S_t = s) \cdot$$

$$\Big(\gamma^t \sum_{i=1}^{N_E}(\omega_r^{[i]})^\top \phi_r^{[i]}(s,a) + \lambda \gamma^t c_{\omega_c}(s,a) + E_{S,A}^\pi \Big[\sum_{\tau=t+1}^\infty \gamma^\tau \sum_{i=1}^{N_E}(\omega_r^{[i]})^\top \phi_r^{[i]}(S_\tau, A_\tau)$$

$$+ \lambda c_{\omega_c}(S_\tau, A_\tau)|S_t = s, A_t = a \Big] \Big),$$

$$= \gamma^t P(S_t = s)[H^\pi(s,a) - 1 + \eta^\top \mu_{r,\omega_c}^\pi(s,a)], \tag{6}$$

where $P(S_t = s)$ is the probability (density) of state s being reached at time t.

From the formula of the constrained soft Bellman policy, we know that $\pi_{\eta;\omega_c}(a|s) > 0$ for any $s \in \mathcal{S}$ and $a \in \mathcal{A}$, thus $\pi_{\eta;\omega_c}$ is an interior point of Π. Therefore $\frac{\partial L(\pi,\eta;\omega_c)}{\partial \pi(a|s)} = 0$ at $\pi_{\eta;\omega_c}$ and from (6) we have that for any $(s, a) \in \mathcal{S} \times \mathcal{A}$ such that $P(S_t = s) \neq 0$ for at least one time t:

$$H^{\pi_{\eta;\omega_c}}(s, a) - 1 + \eta^\top \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a) = 0. \tag{7}$$

Now, we derive the gradient of $G(\eta;\omega_c) = H(\pi_{\eta;\omega_c}) + \sum_{i=1}^{N_E}(\omega_r^{[i]})^\top(\mu_r^{[i]}(\pi_{\eta;\omega_c}) - \hat{\mu}_r^{[i]}) + \lambda(J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c})$:

$$\nabla_\eta G(\eta;\omega_c) = \nabla_\eta H(\pi_{\eta;\omega_c}) + \nabla_\eta \mu_{r,\omega_c}(\pi_{\eta;\omega_c})\eta + (\mu_{r,\omega_c}(\pi_{\eta;\omega_c}) - \hat{\mu}_{r,\omega_c}),$$

$$= \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\nabla_\eta \ln \pi_{\eta;\omega_c}(a|s)(H^{\pi_{\eta;\omega_c}}(s, a) - 1 + (\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a))^\top \eta)dads$$

$$+ (\mu_{r,\omega_c}(\pi_{\eta;\omega_c}) - \hat{\mu}_{r,\omega_c}),$$

$$= \mu_{r,\omega_c}(\pi_{\eta;\omega_c}) - \hat{\mu}_{r,\omega_c},$$

where the second equality follows from Lemma 6 and the third equality follows from (7) and the fact that $\psi^{\pi_{\eta;\omega_c}}(s) = 0$ if $P(S_t = s) = 0$ for all time t. Notice that $\mu_{r,\omega_c}(\pi_{\eta;\omega_c}) - \hat{\mu}_{r,\omega_c} = [(\mu_r(\pi_{\eta;\omega_c}) - \hat{\mu}_r)^\top, J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c}]^\top$. Analogously, we can see that $\nabla_\eta G^{[v]}(\eta;\omega_c) = [(\mu_r(\pi_{\eta;\omega_c}) - \hat{\mu}_r^{[v]})^\top, J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c}^{[v]}]^\top$.

As $G(\eta;\omega_c)$ attains its minimum at $\eta^*(\omega_c)$, given that $\min_\eta G(\eta;\omega_c)$ is unconstrained, the gradient of $G(\eta;\omega_c)$ with respect to η should be 0 at this point (i.e. $\mu_r(\pi_{\eta;\omega_c}) - \hat{\mu}_r = 0$ and $J_{\omega_c}(\pi_{\eta;\omega_c}) - \hat{b}_{\omega_c} = 0$). It implies that $\pi_{\eta;\omega_c}$ is a feasible solution of the primal problem (2). Thus, we have:

$$H(\pi_{\eta^*(\omega_c);\omega_c}) = G(\eta^*(\omega_c);\omega_c) = d^* \geq p^* \geq H(\pi_{\eta^*(\omega_c);\omega_c}),$$

Therefore, we know that $p^* = d^*$ and p^* is obtained at $\pi_{\eta^*(\omega_c);\omega_c}$.

B.3 Intermediate results

In this section, we prove three lemmas which will be used in the remaining subsections.

Lemma 7. *The dual functions $G(\eta;\omega_c)$ and $G^{[v]}(\eta;\omega_c)$ are both strictly convex in η for any nonzero cost weight vector ω_c.*

We know that

$$\nabla_{\eta\eta}^2 G(\eta;\omega_c) = \nabla_\eta \mu_{r,\omega_c}(\pi_{\eta;\omega_c}),$$

$$= \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\nabla_\eta \ln \pi_{\eta;\omega_c}(a|s)(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a))^\top dads,$$

$$= \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\left[\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)\right](\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a))^\top dads,$$

where the second equality follows from Lemma 6 and the last equality follows from Lemma 5.

For any nonzero vector $\bar{\eta}$, we have:

$$\bar{\eta}^\top \nabla_{\eta\eta}^2 G(\eta;\omega_c)\bar{\eta} = \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\omega_c,\eta}(a|s)\left[\eta^\top \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a) - \eta^\top \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)\right]$$

$$\cdot (\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s, a))\top\bar{\eta}dads,$$

$$= \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s)\left[Q_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}(s, a) - V_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}(s)\right](Q_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}(s, a))dads,$$

$$= \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)\mathrm{Var}(Q_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}(s, \cdot))ds,$$

where $\mathrm{Var}(Q_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}(s, \cdot))$ is the variance of the augmented Q-function $Q_{\bar{\eta},\omega_c}^{\pi_{\eta;\omega_c}}$ at state s.

We know that $Q_{\bar{\eta};\omega_c}^{\pi_{\eta;\omega_c}}(s,a) = \bar{\eta}^\top \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a)$ and $\mu_{r,\omega_c,j}^{\pi_{\eta;\omega_c}}(s,a)$ is in essence a discounted cumulative sum of $\phi_{r,\omega_c,j}$. Because each component of ϕ_{r,ω_c} can be chosen independently, the variance $\text{Var}(Q_{\bar{\eta};\omega_c}^{\pi_{\eta;\omega_c}}(s,\cdot)) = \sum_j \bar{\eta}_j^2 \text{Var}(\mu_{r,\omega_c,j}^{\pi_{\eta;\omega_c}}(s,\cdot))$. From the formula of the constrained soft Bellman policy, we know that whatever η and ω_c are, $\pi_{\eta;\omega_c}(a|s) > 0$ for any $a \in \mathcal{A}$ at state s. Therefore, the variance $\text{Var}(\mu_{r,\omega_c,j}^{\pi_{\eta;\omega_c}}(s,\cdot))$ is zero only under the situation that $\mu_{r,\omega_c,j}^{\pi_{\eta;\omega_c}}(s,a)$ is a constant almost everywhere over $\mathcal{A}$ and we can choose $\phi_{r,\omega_c,j}$ whose values always vary over a to avoid this. Therefore, $\text{Var}(\mu_{r,\omega_c,j}^{\pi_{\eta;\omega_c}}(s,\cdot)) > 0$ and thus $\text{Var}(Q_{\bar{\eta};\omega_c}^{\pi_{\eta;\omega_c}}(s,\cdot)) > 0$.

Because $\psi^{\pi_{\eta;\omega_c}}(s) \geq 0$ for all $s \in \mathcal{S}$ and $\int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)ds = \frac{1}{1-\gamma}$, the measure of the set where $\psi^{\pi_{\eta;\omega_c}}(s) > 0$ is strictly greater than 0, otherwise $\int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)ds = 0$. Thus, $\int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s)\text{Var}(Q_{\bar{\eta};\omega_c}^{\pi_{\eta;\omega_c}}(s,\cdot))ds > 0$ and $\nabla_{\eta\eta}^2 G(\eta;\omega_c)$ is positive definite. Analogously, $\nabla_{\eta\eta}^2 G^{[v]}(\eta;\omega_c)$ is also positive definite. Therefore, $G(\eta;\omega_c)$ and $G^{[v]}(\eta;\omega_c)$ are both strictly convex for any nonzero ω_c.

Lemma 8. *(i) There is a positive constant $C_{\nabla_\eta G^{[v]}}$ such that for any $\omega_c \in \Omega_c$ and $\eta \in \mathbb{R}^{\sum_{i=1}^{N_E} l_r^{[i]}+1}$, it holds that $||\nabla_\eta G^{[v]}(\eta;\omega_c)|| \leq C_{\nabla_\eta G^{[v]}}$ and $||\nabla_\eta G(\eta;\omega_c)|| \leq C_{\nabla_\eta G} \triangleq \sum_{v=1}^{N_L} C_{\nabla_\eta G^{[v]}}$.*

(ii) For any $\omega_c \in \Omega_c$ and $\eta \in \mathbb{R}^{\sum_{i=1}^{N_E}+1}$, $G^{[v]}(\eta;\omega_c)$ and $G(\eta;\omega_c)$ are continuously twice differentiable in (ω_c,η).

Proof. (i) From Lemma 1, we know that $||\nabla_\eta G^{[v]}(\eta;\omega_c)|| = ||\mu_{r,\omega_c}(\pi_{\eta;\omega_c}) - \hat{\mu}_{r,\omega_c}^{[v]}|| \leq ||\mu_{r,\omega_c}(\pi_{\eta;\omega_c})|| + ||\hat{\mu}_{r,\omega_c}^{[v]}||$, thus from Lemma 4, we know that $||\nabla_\eta G^{[v]}(\eta;\omega_c)|| \leq C_{\nabla_\eta G^{[v]}}$ for some positive constant $C_{\nabla_\eta G^{[v]}}$. Therefore, $||\nabla_\eta G(\eta;\omega_c)|| = ||\sum_{v=1}^{N_L} G^{[v]}(\eta;\omega_c)|| \leq \sum_{v=1}^{N_L} ||G^{[v]}(\eta;\omega_c)|| \leq \sum_{v=1}^{N_L} C_{\nabla_\eta G^{[v]}} \triangleq C_{\nabla_\eta G}$

(ii) From the proof in Lemma 7, we know that:

$$\nabla_{\eta\eta}^2 G(\eta;\omega_c) = \int_{s\in\mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a\in\mathcal{A}} \pi_{\eta;\omega_c}(a|s) \cdot \left[\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s) \right] (\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a))^\top dads,$$

$$= E_{S,A}^{\pi_{\eta;\omega_c}} \left[(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S))(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))^\top \right].$$

Thus,

$$\nabla_{\eta\eta\eta}^3 G(\eta;\omega) = E_{S,A}^{\pi_{\eta;\omega_c}} \left\{ \nabla_\eta \left[(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S))(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))^\top \right] \right\}$$

$$= E_{S,A}^{\pi_{\eta;\omega_c}} \left[2(\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))^\top - (\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S))(\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))^\top \right.$$

$$\left. - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S)(\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(S,A))^\top \right],$$

where the formula of $\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)$ can be found in Lemma 6 and the formula of $\nabla_\eta \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a)$ can be derived in a similar way to Lemma 6.

With similar process, we can also get the formulas of $\nabla_{\omega_c\omega_c\omega_c}^3 G(\eta;\omega_c)$, $\nabla_{\eta\omega_c\omega_c}^3 G(\eta;\omega_c)$, $\nabla_{\eta\eta\omega_c}^3 G(\eta;\omega_c)$, and $\nabla_{\eta\omega_c\eta}^3 G(\eta;\omega_c)$. Therefore, $G(\eta;\omega_c)$ is three-times differentiable. Analogously, $G^{[v]}(\eta;\omega_c)$ is also three-times differentiable. Thus, they are both continuously twice differentiable. $\square$

Lemma 9. *There are positive constants $\tau_{\nabla_{\eta\eta}^2 G}$ and $\tau_{\nabla_{\eta\eta}^2 G^{[v]}}$ such that $\nabla_{\eta\eta}^2 G(\omega_c, \eta^*(\omega_c')) \succeq \tau_{\nabla_{\eta\eta}^2 G} I$, $\nabla_{\eta\eta}^2 G(\omega_c, \bar{\eta}^{[v]}(\omega_c)) \succeq \tau_{\nabla_{\eta\eta}^2 G} I$, $\nabla_{\eta\eta}^2 G^{[v]}(\omega_c, \eta^*(\omega_c')) \succeq \tau_{\nabla_{\eta\eta}^2 G^{[v]}} I$ and $\nabla_{\eta\eta}^2 G^{[v]}(\omega_c, \bar{\eta}^{[v]}(\omega_c)) \succeq \tau_{\nabla_{\eta\eta}^2 G^{[v]}} I$ for all $\omega_c \in \Omega_c\backslash\{0\}$*

Proof. The proof of Lemma 9 is divided into two steps where the first step shows that $\nabla_{\eta\eta}^2 G(\omega_c, \eta^*(\omega_c')) \succeq \tau_{\nabla_{\eta\eta}^2 G} I$ and the second step shows that $\nabla_{\eta\eta}^2 G(\omega_c, \bar{\eta}^{[v]}(\omega_c)) \succeq \tau_{\nabla_{\eta\eta}^2 G} I$. The part for $\nabla_{\eta\eta}^2 G^{[v]}$ can be derived analogously.

Step (i). Lemma 7 shows that $G(\eta; \omega_c)$ is strictly convex in η, thus there is a unique $\eta^*(\omega_c)$. Since the problem $\min_\eta G(\eta; \omega_c)$ is unconstrained and $\nabla_\eta G(\eta^*(\omega_c); \omega_c) = 0$ holds for any $\omega_c \in \mathbb{R}^{\sum_{i=1}^{N_E} l_c^{[i]}}$, taking derivative with respect to ω_c on both sides renders

$$\nabla^2_{\eta\omega_c} G(\eta^*(\omega_c); \omega_c) + \nabla^2_{\eta\eta} G(\eta^*(\omega_c); \omega_c) \nabla\eta^*(\omega_c) = 0 \Rightarrow \nabla\eta^*(\omega_c) = -M(\omega_c, \eta^*(\omega_c))^\top \quad (8)$$

where $M(\omega_c, \eta) \triangleq \nabla^2_{\omega_c\eta} G(\eta; \omega_c) \big[\nabla^2_{\eta\eta} G(\eta; \omega_c)\big]^{-1}$.

Since $\eta^*(\omega_c')$ is differentiable (equation (8)) and $\nabla^2_{\eta\eta} G(\eta; \omega_c)$ is continuous in (ω_c, η) (Lemma 8 (ii)), $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ is continuous in (ω_c, ω_c'). Since Ω_c is compact, $\Omega_c \times \Omega_c$ is compact (Tychonoff's theorem, Theorem 1.9.7 in [33]), then the image of $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ is compact (Theorem 4.14 in [34]). We denote the eigenvalues of $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ by $\lambda_i(\omega_c, \omega_c')$ and we know that $\lambda_i(\omega_c, \omega_c') > 0$ for any ω_c and ω_c' because $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ is positive definite and symmetric.

Now, we prove that the image of every $\lambda_i(\omega_c, \omega_c')$ is compact. The characteristic polynomial of $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ is $(\lambda - \lambda_1(\omega_c, \omega_c')) \cdots (\lambda - \lambda_d(\omega_c, \omega_c')) = \lambda^d + p_{d-1}(\omega_c, \omega_c')\lambda^{d-1} + \cdots + p_1(\omega_c, \omega_c')\lambda + p_0(\omega_c, \omega_c')$, where $\lambda_i(\omega_c, \omega_c')$ is the root and $p_i(\omega_c, \omega_c')$ is the coefficient. Each $p_i(\omega_c, \omega_c')$ is in essence a polynomial function of the entries of $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ and the entries of $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ are continuous in (ω_c, ω_c') as $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c)$ is continuous in (ω_c, ω_c'), then $p_i(\omega_c, \omega_c')$ is continuous in (ω_c, ω_c') (Theorem 4.7 in [34]). Then $\lambda_i(\omega_c, \omega_c')$ is also continuous in (ω_c, ω_c') (Theorem 3.9.1 in [35]), thus the image of $\lambda_i(\omega_c, \omega_c')$ is compact.

According to Heine-Borel theorem (Theorem 2.41 in [34]), the image of $\lambda_i(\omega_c, \omega_c')$ is closed and bounded. Therefore, $\min \lambda_i(\omega_c, \omega_c')$ exists and belongs to the image of $\lambda_i(\omega_c, \omega_c')$, then $\min \lambda_i(\omega_c, \omega_c')$ is positive. Thus, we can choose a positive number $\tau_{\nabla^2_{\eta\eta} G^*} = \min\{\min \lambda_1(\omega_c, \omega_c'), \cdots, \min \lambda_d(\omega_c, \omega_c')\}$ and $\nabla^2_{\eta\eta} G(\eta^*(\omega_c'); \omega_c) \succeq \tau_{\nabla^2_{\eta\eta} G^*} I$.

Step (ii). For the distributed gradient descent in Algorithm 2, we know that (equation (5) in [36])

$$\eta^{[v]}(\omega_c, k) = \sum_{v'=1}^{N_L} [\Phi(k-1, 0)]_{v'}^v \eta^{[v']}(0) - \sum_{s=1}^{k-1} \alpha(s-1) \sum_{v'=1}^{N_L} [\Phi(k-1, s)]_{v'}^v \cdot$$
$$\nabla_\eta G^{[v']}(\eta^{[v']}(\omega_c, s-1); \omega_c) - \alpha(k-1) \nabla_\eta G^{[v]}(\eta^{[v]}(\omega_c, k-1); \omega_c),$$

where $\Phi(k, s) \triangleq W(s)W(s+1) \cdots W(k)$ is the state transition matrix and $[\Phi(k, s)]_{v'}^v$ is the entry at the v-th row and v'-th column.

We define $y(\omega_c, k) \triangleq \frac{1}{N_L} \sum_{v'=1}^{N_L} \eta^{[v']}(0) - \sum_{s=1}^{k} \alpha(s-1) \sum_{v'=1}^{N_L} \frac{1}{N_L} \nabla_\eta G^{[v']}(\eta^{[v']}(\omega_c, s-1); \omega_c)$, then $y(\omega_c, k+1) = y(\omega_c, k) - \frac{\alpha(k)}{N_L} \sum_{v'=1}^{N_L} \nabla_\eta G^{[v']}(\eta^{[v']}(\omega_c, k); \omega_c)$.

Following the proof of Lemma 5 (a) and proposition 3 in [36], we can get

$$||y(\omega_c, k+1) - \eta^*(\omega_c)||^2,$$

$$\leq ||y(\omega_c, k) - \eta^*(\omega_c)||^2 + \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)||$$

$$- \frac{2\alpha(k)}{N_L} \left[G(y(\omega_c, k); \omega_c) - G(\eta^*(\omega_c); \omega_c) \right] + \frac{(\alpha(k))^2}{N_L} C_{\nabla_\eta G}, \quad (9)$$

$$\leq ||y(\omega_c, k) - \eta^*(\omega_c)||^2 + \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| + \frac{(\alpha(k))^2}{N_L} C^2_{\nabla_\eta G}, \quad (10)$$

$$||y(\omega_c, k) - \eta^{[v]}(\omega_c, k)|| \leq 2 \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}} \sum_{v'=1}^{N_L} ||\eta^{[v']}(0)||$$

$$+ \sum_{s=1}^{k-1} 2\alpha(s-1) C_{\nabla_\eta G} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} + \frac{\alpha(k-1)}{N_L} (C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}}),$$

where $B_0 = (N_L - 1)B$.

We can find c_η such that $\sum_{v=1}^{N_L} ||\eta^{[v]}(0)|| \le c_\eta$. Therefore,

$$\frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| \le \frac{8\alpha(k)C_{\nabla_\eta G}c_\eta}{N_L} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}}$$

$$+ \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} \sum_{s=1}^{k-1} 2\alpha(s-1)C_{\nabla_\eta G} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}}$$

$$+ \frac{4\alpha(k)\alpha(k-1)}{N_L^2} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} (C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}}),$$

$$= \frac{8\alpha(k)C_{\nabla_\eta G}c_\eta}{N_L} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}} + \frac{8\alpha(k)C_{\nabla_\eta G}^2}{N_L} \sum_{s=1}^{k-1} \alpha(s-1) \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}}$$

$$+ \frac{4\alpha(k)\alpha(k-1)C_{\nabla_\eta G}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L^2}. \tag{11}$$

Notice that the second term in (11) does not exist if $k < 2$.

Claim 1. $\sum_{k=1}^{K} \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)||$ *is bounded for any K.*

Proof. Because the upper bound of $\frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)||$, i.e., formula (11), is positive, it suffices to show that $\sum_{k=1}^{\infty} \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)||$ is bounded.

Now, we prove that the summation of each term in (11) is finite one by one.

First, $\sum_{k=1}^{\infty} \frac{8\alpha(k)c_\eta C_{\nabla_\eta G}}{N_L} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}} \le \frac{8\bar{\alpha}c_\eta C_{\nabla_\eta G}}{N_L} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} \sum_{k=1}^{\infty} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}} = \frac{8\bar{\alpha}c_\eta C_{\nabla_\eta G}}{N_L} \frac{1 + \epsilon^{-B_0}}{(1 - \epsilon^{B_0})[1 - (1 - \epsilon^{B_0})^{\frac{1}{B_0}}]}$ is bounded.

Second,

$$\sum_{k=2}^{\infty} \frac{8\alpha(k)C_{\nabla_\eta G}^2}{N_L} \sum_{s=1}^{k-1} \alpha(s-1) \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} = \frac{8C_{\nabla_\eta G}^2(1 + \epsilon^{-B_0})}{N_L(1 - \epsilon^{B_0})} \cdot$$

$$\left[\sum_{k=2}^{\infty} \frac{\bar{\alpha}}{k+1} \sum_{s=1}^{k-1} \frac{\bar{\alpha}}{s} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} \right].$$

Let $S_k = \sum_{s=1}^{k-1} \frac{\bar{\alpha}}{s} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}}$, then $\frac{k}{\bar{\alpha}} S_{k-1} - \frac{k+1}{\bar{\alpha}} S_k = \frac{k}{\bar{\alpha}} \sum_{s=1}^{k-2} \frac{\bar{\alpha}}{s} (1 - \epsilon^{B_0})^{\frac{k-2-s}{B_0}} - \frac{k+1}{\bar{\alpha}} \sum_{s=1}^{k-1} \frac{\bar{\alpha}}{s} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} = \sum_{s=1}^{k-2} \frac{k}{s} (1 - \epsilon^{B_0})^{\frac{k-2-s}{B_0}} - \sum_{s=0}^{k-2} \frac{k+1}{s+1} (1 - \epsilon^{B_0})^{\frac{k-2-s}{B_0}} = \sum_{s=1}^{k-2} \frac{k-s}{s(s+1)} (1 - \epsilon^{B_0})^{\frac{k-2-s}{B_0}} - (k+1)(1 - \epsilon^{B_0})^{\frac{k-2}{B_0}} = \sum_{s=1}^{k-3} \frac{k-s}{s(s+1)} (1 - \epsilon^{B_0})^{\frac{k-2-s}{B_0}} + \frac{2}{(k-2)(k-1)} - (k+1)(1 - \epsilon^{B_0})^{\frac{k-2}{B_0}}$.

Because $(1 - \epsilon^{B_0})^{\frac{k-2}{B_0}}$ decays faster than $\frac{1}{(k-2)(k-1)(k+1)}$, there exists a positive integer $\bar{K}$ such that $\frac{2}{(k-2)(k-1)} - (k+1)(1 - \epsilon^{B_0})^{\frac{k-2}{B_0}} > 0$ if $k > \bar{K}$. Therefore, $\frac{k}{\bar{\alpha}} S_{k-1} - \frac{k+1}{\bar{\alpha}} S_k > 0$ if $k > \bar{K}$. We can find a positive number M such that $\frac{k+1}{\bar{\alpha}} S_k < M$ for any $k > 0$. Then, $\frac{\bar{\alpha}}{k+1} S_k < \frac{\bar{\alpha}^2 M}{(k+1)^2}$ and $\sum_{k=2}^{\infty} \frac{\bar{\alpha}}{k+1} S_k$ is finite.

Third, $\sum_{k=1}^{\infty} \frac{4\alpha(k)\alpha(k-1)C_{\nabla_\eta G}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L^2} = \frac{4C_{\nabla_\eta G}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L^2} \sum_{k=1}^{\infty} \alpha(k)\alpha(k - 1) \le \frac{4C_{\nabla_\eta G}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L^2} \sum_{k=1}^{\infty} \frac{\bar{\alpha}^2}{k^2}$ is bounded. $\square$

Equation (10) shows that $||y(\omega_c, k + 1) - \eta^*(\omega_c)||^2 \leq ||y(\omega_c, k) - \eta^*(\omega_c)||^2 + \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| + \frac{(\alpha(k))^2}{N_L} C_{\nabla_\eta G}^2$.

Telescoping from $k = 0$ to $K - 1$, we have:

$$||y(\omega_c, k) - \eta^*(\omega_c)||^2 \leq ||y(\omega_c, 0) - \eta^*(\omega_c)||^2 + \frac{4\alpha^{(0)}}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, 0) - \eta^{[v']}(0)||$$

$$+ \sum_{k=1}^{K-1} \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| + \sum_{k=0}^{K-1} \frac{(\alpha(k))^2 C_{\nabla_\eta G}^2}{N_L^2} \leq D_{\max},$$

where $D_{\max} = ||y(\omega_c, 0) - \eta^*(\omega_c)||^2 + \frac{4\alpha^{(0)}}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, 0) - \eta^{[v']}(0)|| + \sum_{k=1}^{\infty} \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| + \sum_{k=0}^{\infty} \frac{(\alpha(k))^2 C_{\nabla_\eta G}^2}{N_L^2}$ is finite.

Therefore,

$$||\eta^{[v]}(\omega_c, k) - \eta^*(\omega_c)|| \leq ||\eta^{[v]}(\omega_c, k) - y(\omega_c, k)|| + ||y(\omega_c, k) - \eta^*(\omega_c)||,$$

$$\leq 2c_\eta \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1}{B_0}} + \sum_{s=1}^{k-1} 2C_{\nabla_\eta G} \alpha(s-1) \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}}$$

$$+ \frac{\alpha(k-1)}{N_L} (C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}}) + \sqrt{D_{\max}},$$

$$\leq 2c_\eta \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} + 2C_{\nabla_\eta G} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} \sum_{s=1}^{k-1} \alpha(s-1)(1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} + \frac{\bar{\alpha}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L}$$

$$+ \sqrt{D_{\max}},$$

$$\leq 2c_\eta \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} + 2\bar{\alpha} C_{\nabla_\eta G} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} \sum_{s=1}^{k-1} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} + \frac{\bar{\alpha}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L},$$

$$+ \sqrt{D_{\max}} \leq \bar{D}_{\max},$$

where $\bar{D}_{\max} = 2c_\eta \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} + 2\bar{\alpha} C_{\nabla_\eta G} \frac{1 + \epsilon^{-B_0}}{1 - \epsilon^{B_0}} \sum_{s=1}^{\infty} (1 - \epsilon^{B_0})^{\frac{k-1-s}{B_0}} + \frac{\bar{\alpha}(C_{\nabla_\eta G} + N_L C_{\nabla_\eta G^{[v]}})}{N_L} + \sqrt{D_{\max}}$ is finite. Because $\eta^*(\omega_c)$ is bounded within a compact set and $||\eta^{[v]}(\omega_c, k)|| \leq ||\eta^*(\omega_c)|| + \bar{D}_{\max}$, we know that $\eta^{[v]}(\omega_c, k)$ is also bounded within a compact set. Then, $||\bar{\eta}^{[v]}(\omega_c)|| = ||\frac{\sum_{j=0}^{K-1} \alpha(k)\eta^{[v]}(\omega_c, k)}{\sum_{j=0}^{K-1} \alpha(k)}|| \leq \max ||\eta^{[v]}(\omega_c, j)|| \leq ||\eta^*(\omega_c)|| + \bar{D}_{\max}$ so that $\bar{\eta}^{[v]}(\omega_c)$ is also bounded within a compact set. Following the same idea of η^*, we can find a positive constant $\tau_{G_{\bar{\eta}\bar{\eta}}}$ such that $\nabla^2_{\eta\eta} G(\bar{\eta}^{[v]}(\omega_c); \omega_c) \succeq \tau_{G_{\bar{\eta}\bar{\eta}}} I$ for all $k > 0$.

Then, we can find the positive constant $\tau_{\nabla^2_{\eta\eta} G} = \min\{\tau_{\nabla^2_{\eta\eta} G^*}, \tau_{G_{\bar{\eta}\bar{\eta}}}\}$. With similar derivation, we can find the positive constant $\tau_{\nabla^2_{\eta\eta} G^{[v]}}$. $\qquad \square$

B.4 Proof of Lemma 3

From Claim 1, we know that there is a positive constant $c_{\max}$ such that $\sum_{k=0}^{K} 2\alpha(k) \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| \leq c_{\max}$ for all $K > 0$. From (9), we know that $||y(\omega_c, k+1) - \eta^*(\omega_c)||^2 \leq ||y(\omega_c, k) - \eta^*(\omega_c)||^2 + \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v]}} ||y(\omega_c, k) - \eta^{[v]}(\omega_c, k)|| - \frac{2\alpha(k)}{N_L} \left[G(y(\omega_c, k); \omega_c) - G(\eta^*(\omega_c); \omega_c) \right] + \frac{(\alpha(k))^2}{N_L} C_{\nabla_\eta G}^2$, then we have:

$$\sum_{k=0}^{K-1} \frac{2\alpha(k)}{N_L} \left[G(y(\omega_c, k); \omega_c) - G(\eta^*(\omega_c); \omega_c) \right] \leq ||y(\omega_c, 0) - \eta^*(\omega_c)||^2 - ||y(\omega_c, k) - \eta^*(\omega_c)||^2$$

$$+ \sum_{k=0}^{K-1} \frac{4\alpha(k)}{N_L} \sum_{v'=1}^{N_L} C_{\nabla_\eta G^{[v']}} ||y(\omega_c, k) - \eta^{[v']}(\omega_c, k)|| + \sum_{k=0}^{K-1} \frac{(\alpha(k))^2 C_{\nabla_\eta G}^2}{N_L^2},$$

$$\Rightarrow \frac{\sum_{k=0}^{K-1}\alpha(k)G(y(\omega_c,k);\omega_c)}{\sum_{k=0}^{K-1}\alpha(k)} - G(\eta^*(\omega_c);\omega_c) \le \frac{N_L}{2\sum_{k=0}^{K-1}\alpha(k)}||y(\omega_c,0)-\eta^*(\omega_c)||^2$$

$$+ \frac{1}{\sum_{k=0}^{K-1}\alpha(k)}\sum_{k=0}^{K-1}2\alpha(k)\sum_{v'=1}^{N_L}C_{\nabla_\eta G^{[v']}}||y(\omega_c,k)-\eta^{[v']}(\omega_c,k)||$$

$$+ \frac{1}{\sum_{k=0}^{K-1}\alpha(k)}\sum_{k=0}^{K-1}\frac{(\alpha(k))^2 C_{\nabla_\eta G}^2}{2N_L},$$

$$\le \frac{N_L}{2\sum_{k=0}^{K-1}\alpha(k)}||y(\omega_c,0)-\eta^*(\omega_c)||^2 + \frac{1}{\sum_{k=0}^{K-1}\alpha(k)}(c_{\max}+\frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha^2}{12N_L}),$$

$$\Rightarrow G(\frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)};x) - G(\eta^*(\omega_c);\omega_c) \le \frac{\sum_{k=0}^{K-1}\alpha(k)G(y(\omega_c,k);\omega_c)}{\sum_{k=0}^{K-1}\alpha(k)}$$

$$- G(\eta^*(\omega_c);\omega_c),$$

$$\le \frac{N_L}{2\sum_{k=0}^{K-1}\alpha(k)}||y(\omega_c,0)-\eta^*(\omega_c)||^2 + \frac{1}{\sum_{k=0}^{K-1}\alpha(k)}(c_{\max}+\frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha^2}{12N_L})$$

$$\le \frac{1}{\log K}\left[\frac{N_L||y(\omega_c,0)-\eta^*(\omega_c)||^2}{2\bar\alpha} + c_{\max} + \frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha}{12N_L}\right],$$

$$\Rightarrow ||\frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \eta^*(\omega_c)||^2 \le \frac{2}{\tau_{\nabla_{\eta\eta}^2 G}}\left[G(\frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)};\omega_c)\right.$$

$$\left.- G(\eta^*(\omega_c);\omega_c)\right],$$

$$\le \frac{2}{\tau_{\nabla_{\eta\eta}^2 G}\log K}\left[\frac{N_L||y(\omega_c,0)-\eta^*(\omega_c)||^2}{2\bar\alpha} + c_{\max} + \frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha}{12N_L}\right].$$

Then,

$$||\frac{\sum_{k=0}^{K-1}\alpha(k)\eta^{[v]}(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \eta^*(\omega_c)|| \le ||\frac{\sum_{k=0}^{K-1}\alpha(k)\eta^{[v]}(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)}||$$

$$+ ||\frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \eta^*(\omega_c)||,$$

$$\le \frac{\sum_{k=0}^{K-1}\alpha(k)||\eta^{[v]}(\omega_c,k)-y(\omega_c,k)||}{\sum_{k=0}^{K-1}\alpha(k)} + ||\frac{\sum_{k=0}^{K-1}\alpha(k)y(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \eta^*(\omega_c)||.$$

We know that $\frac{c_{\max}}{2} \ge \sum_{k=0}^{K-1}\alpha(k)\sum_{v'=1}^{N_L}C_{\nabla_\eta G^{[v']}}||y(\omega_c,k)-\eta^{[v']}(\omega_c,k)|| \ge C_{\nabla_\eta G^{[v]}}\sum_{k=0}^{K-1}\alpha(k)||y(\omega_c,k)-\eta^{[v]}(\omega_c,k)||$, then

$$||\frac{\sum_{k=0}^{K-1}\alpha(k)\eta^{[v]}(\omega_c,k)}{\sum_{k=0}^{K-1}\alpha(k)} - \eta^*(\omega_c)|| \le \frac{1}{\log K}\frac{c_{\max}}{2C_{\nabla_\eta G^{[v]}}}$$

$$+ \sqrt{\frac{2}{\tau_{\nabla_{\eta\eta}^2 G}\log K}\left[\frac{N_L||y(\omega_c,0)-\eta^*(\omega_c)||^2}{2\bar\alpha} + c_{\max} + \frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha}{12N_L}\right]},$$

$$= \frac{1}{\log K}\frac{c_{\max}}{2C_{\nabla_\eta G^{[v]}}} + \frac{1}{\sqrt{\log K}}\sqrt{\left[\frac{N_L||y(\omega_c,0)-\eta^*(\omega_c)||^2}{\tau_{\nabla_{\eta\eta}^2 G}\bar\alpha} + \frac{2c_{\max}}{\tau_{\nabla_{\eta\eta}^2 G}} + \frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha}{6\tau_{\nabla_{\eta\eta}^2 G}N_L}\right]},$$

$$= \frac{1}{\log K}\frac{c_{\max}}{2C_{\nabla_\eta G^{[v]}}} + \frac{1}{\sqrt{\log K}}\sqrt{\frac{N_L||\frac{1}{N_L}\sum_{v=0}^{N_L}y_v^{(0)}(x)-\eta^*(\omega_c)||^2}{\tau_{\nabla_{\eta\eta}^2 G}\bar\alpha} + \frac{2c_{\max}}{\tau_{\nabla_{\eta\eta}^2 G}} + \frac{C_{\nabla_\eta G}^2\pi^2\bar\alpha}{6\tau_{\nabla_{\eta\eta}^2 G}N_L}}.$$

We let $C_1^{[v]} = \frac{c_{\max}}{2C_{\nabla_\eta G^{[v]}}}$ and $C_2^{[v]} = \sqrt{\frac{N_L || \frac{1}{N_L} \sum_{v=0}^{N_L} y_v^{(0)}(x) - \eta^*(\omega_c) ||^2}{\tau_{\nabla_{\eta\eta}^2 G}} + \frac{2c_{\max}}{\tau_{\nabla_{\eta\eta}^2 G} \bar\alpha} + \frac{C_{\nabla_\eta G}^2 \pi^2 \bar\alpha}{6\tau_{\nabla_{\eta\eta}^2 G} N_L}}$.

B.5 The derivation of learner v's LGA

Using the chain rule, we know that:

$$\nabla F(\omega_c, \eta^*(\omega_c)) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c)) + [(\nabla_\eta F(\omega_c, \eta^*(\omega_c)))^\top \nabla \eta^*(\omega_c)]^\top,$$
$$= \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c)) - M(\omega_c, \eta^*(\omega_c)) \nabla_\eta F(\omega_c, \eta^*(\omega_c)),$$

where $M(\omega_c, \eta)$ is defined in (8).

Lemma 10. *The gradient* $\nabla F(\omega, \eta^*(\omega_c)) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c))$.

Proof. The global likelihood function is:

$$F(\omega_c, \eta) = \sum_{t=0}^\infty \gamma^t \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} m P_\mathcal{D}(S_t = s, A_t = a) \ln \pi_{\eta;\omega_c}(a|s) da ds,$$

where $P_\mathcal{D}(S_t = s, A_t = a)$ is the empirical probability of (s, a) occurring at time t in the demonstrations $\mathcal{D}$:

$$P_\mathcal{D}(S_t = s, A_t = a) \triangleq \frac{1}{m} \sum_{j=1}^m (\mathbb{1}\{s_t^j = s\} \mathbb{1}\{a_t^j = a\}),$$

From the proof of Lemma 5, we know that $\pi_{\eta;\omega_c}$ can be formulated as:

$$\pi_{\eta;\omega_c}(a|s) = \frac{Z_{a|s,\eta;\omega_c}}{Z_{s,\eta;\omega_c}},$$

$$\ln Z_{a|s,\eta;\omega_c} = \sum_{i=1}^{N_E} (\omega_r^{[i]})^\top \phi_r^{[i]}(s, a) + \lambda c_{\omega_c}(s, a) + \gamma \int_{s' \in \mathcal{S}} P(s'|s, a) \ln Z_{s',\eta;\omega_c} ds',$$

$$\ln Z_{s,\eta;\omega_c} = \ln \int_{a \in \mathcal{A}} Z_{a|s,\eta;\omega_c} da.$$

Thus,

$$\nabla_\eta F(\omega_c, \eta) = \sum_{t=0}^\infty \gamma^t \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} m P_\mathcal{D}(S_t = s, A_t = a) \nabla_\eta (\ln Z_{a|s,\eta;\omega_c} - \ln Z_{s,\eta;\omega_c}) da ds,$$

$$= \sum_{t=0}^\infty \gamma^t \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} m P_\mathcal{D}(S_t = s, A_t = a) \left\{ \phi_{r,\omega_c}(s, a) + E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t+1}^\infty \gamma^{\tau-t} \phi_{r,\omega_c}(S_\tau, A_\tau) | \right. \right.$$

$$\left. \left. S_t = s, A_t = a \right] - E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t}^\infty \gamma^{\tau-t} \phi_{r,\omega_c}(S_\tau, A_\tau) | S_t = s \right] \right\} da ds,$$

where the last inequality follows from Lemma 5. Here,

$$\gamma \int_{s' \in \mathcal{S}} \int_{a' \in \mathcal{A}} P_\mathcal{D}(S_{t+1} = s', A_{t+1} = a') E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t+1}^\infty \gamma^{\tau-t-1} \phi_{r,\omega_c}(S_\tau, A_\tau) | S_{t+1} = s' \right] da' ds',$$

$$= \gamma \int_{s' \in \mathcal{S}} P_\mathcal{D}(S_{t+1} = s') E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t+1}^\infty \gamma^{\tau-t-1} \phi_{r,\omega_c}(S_\tau, A_\tau) | S_{t+1} = s' \right] ds',$$

$$= \gamma \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} P_\mathcal{D}(S_t = s, A_t = a) \int_{s' \in \mathcal{S}} P(s'|s, a) \cdot$$

$$E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t+1}^\infty \gamma^{\tau-t-1} \phi_{r,\omega_c}(S_\tau, A_\tau) | S_{t+1} = s' \right] ds' da ds,$$

$$= \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} P_{\mathcal{D}}(S_t = s, A_t = a) E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{\tau=t+1}^{\infty} \gamma^{\tau-t} \phi_{r,\omega_c}(S_\tau, A_\tau) | S_t = s, A_t = a \right] dads.$$

Therefore,

$$\nabla_\eta F(\omega_c, \eta) = \sum_{t=0}^{\infty} \gamma^t \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} m P_{\mathcal{D}}(S_t = s, A_t = a) \phi_{r,\omega_c}(s, a) dads$$

$$- \int_{s \in \mathcal{S}} \int_{a \in \mathcal{A}} m P_{\mathcal{D}}(S_0 = s, A_0 = a) E_{S,A}^{\pi_{\eta;\omega_c}} \left[\sum_{t=0}^{\infty} \gamma^t \phi_{r,\omega_c}(S_t, A_t) | S_0 = s \right] dads,$$

$$= m(\hat{\mu}_{r,\omega_c} - \mu_{r,\omega_c}(\pi_{\eta;\omega_c})).$$

The above derivation shows that $\nabla_\eta F(\omega_c; \eta) = -m \nabla_\eta G(\eta; \omega_c)$. Therefore, we have that $\nabla F(\omega_c, \eta^*(\omega_c)) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c)) - M(\omega_c, \eta^*(\omega_c)) \nabla_\eta F(\omega_c, \eta^*(\omega_c)) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c)) + M(\omega_c, \eta^*(\omega_c)) \nabla_\eta m G(\eta^*(\omega_c); \omega_c) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c))$. $\qquad \square$

However, we cannot obtain the gradient $\nabla F(\omega_c, \eta^*(\omega_c))$ because each learner v does not find $\eta^*(\omega_c)$ but its approximation $\bar{\eta}^{[v]}(\omega_c)$. Therefore, we propose a surrogate gradient to approximate $\nabla F(\omega_c, \eta^*(\omega_c))$: $\bar{\nabla} F(\omega_c, \bar{\eta}^{[v]}(\omega_c)) \triangleq \nabla_{\omega_c} F(\omega_c, \bar{\eta}^{[v]}(\omega_c))$.

Notice that this surrogate gradient is global, to decompose it, we propose a local gradient approximation from learner v: $\bar{\nabla} F^{[v]}(\omega_c, \bar{\eta}^{[v]}(\omega_c)) \triangleq \nabla_{\omega_c} F^{[v]}(\omega_c, \bar{\eta}^{[v]}(\omega_c))$. This local approximation aims to approximate $\bar{\nabla} F^{[v]}(\omega_c, \eta^*(\omega_c)) = \nabla_{\omega_c} F^{[v]}(\omega_c, \eta^*(\omega_c))$. Notice that $\nabla F(\omega_c, \eta^*(\omega_c)) = \nabla_{\omega_c} F(\omega_c, \eta^*(\omega_c)) = \sum_{v=1}^{N_L} \nabla_{\omega_c} F^{[v]}(\omega_c, \eta^*(\omega_c)) = \sum_{v=1}^{N_L} \bar{\nabla} F^{[v]}(\omega_c, \eta^*(\omega_c))$.

Similar to the derivation of $\nabla_\eta F(\omega_c, \eta)$, we can get the gradient $\nabla_{\omega_c} F^{[v]}(\omega_c, \eta) = \sum_{\zeta^j \in \mathcal{D}^{[v]}} \sum_{t=0}^{\infty} \gamma^t \phi_c(s_t^j, a_t^j) - m^{[v]} E_{S,A}^{\pi_{\eta;\omega_c}} [\sum_{t=0}^{\infty} \gamma^t \phi_c(S_t, A_t)]$. We still use this approximation when $\omega_c = 0$.

Lemma 11. *The approximation error $||\bar{\nabla} F^{[v]}(\omega_c, \bar{\eta}(\omega_c)) - \bar{\nabla} F^{[v]}(\omega_c, \eta^*(\omega_c))||$ is upper bounded and decreases to zero at the rate of $O(\frac{1}{\sqrt{\log K}})$.*

Proof. At first, we first prove the following claim showing that bounded gradient can imply Lipschitz condition of a function if the domain of the function is convex.

Claim 2. *Suppose a function $f : E \subseteq \mathbb{R}^n \to \mathbb{R}^m$ is differentiable on the convex set E, and $||\nabla f(x)|| \leq M$ for any $x \in E$, where M is a finite positive constant. Then f satisfies the Lipschitz condition, i.e., $||f(b) - f(a)|| \leq M ||b - a||$ for any $a, b \in E$.*

Proof. Let $y(t) = (1 - t)a + tb$ where $t \in [0, 1]$ and $h(t) = f \circ y(t)$. Let $z = f(b) - f(a)$ and define $v(t) = z^\top h(t)$. Then v is real-valued and differentiable on $(0, 1)$. By the mean value theorem, there exists $\bar{t} \in (0, 1)$ such that $v(1) - v(0) = v'(t) \Rightarrow ||f(b) - f(a)||^2 = (f(b) - f(a))^\top (\nabla_y f(y(t)))^\top (b - a) \leq ||f(b) - f(a)|| \cdot ||\nabla_y f(y(t))|| \cdot ||b - a|| \Rightarrow ||f(b) - f(a)|| \leq M ||b - a||$. $\qquad \square$

Claim 3. *For any $\omega_c \in \Omega_c$, $\nabla_{\omega_c} F^{[v]}(\omega_c, \eta)$ is Lipschitz continuous (w.r.t. η) with positive constant $L_{\nabla_{\omega_c} F^{[v]}}$.*

Proof. From Claim 2, it suffices to show that $||\nabla_{\omega_c \eta} F^{[v]}(\omega_c, \eta)||$ is bounded. Following similar derivation in the proof of Lemma 10, we can get $\nabla_{\omega_c} F^{[v]}(\omega_c, \eta) = m^{[v]}(\hat{\mu}_c - \lambda \mu_c(\pi_{\eta;\omega_c}))$. Then following the similar idea of the proof in Lemma 7, we can see that

$$\nabla_{\omega_c \eta}^2 F^{[v]}(\omega_c, \eta) = -m^{[v]} \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \left[\nabla_\eta \pi_{\eta;\omega_c}(a|s)(\lambda \mu_c^{\pi_{\eta;\omega_c}}(s,a))^\top + \pi_{\eta;\omega_c}(a|s) \right.$$

$$\left. \cdot [\mathbf{0}_{\sum_{i=1}^{N_E} l_c^{[i]} \times \sum_{i=1}^{N_E} l_r^{[i]}}, \mu_c^{\pi_{\eta;\omega_c}}(s,a)]^\top \right] dads$$

$$= m^{[v]} \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \lambda \pi_{\eta;\omega_c}(a|s)[\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s) - \mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a)](\mu_c^{\pi_{\eta;\omega_c}}(s,a))^\top dads$$

$$- m^{[v]} \int_{s \in \mathcal{S}} \psi^{\pi_{\eta;\omega_c}}(s) \int_{a \in \mathcal{A}} \pi_{\eta;\omega_c}(a|s) [\mathbf{0}_{\sum_{i=1}^{N_E} l_c^{[i]} \times \sum_{i=1}^{N_E} l_r^{[i]}}, \mu_c^{\pi_{\eta;\omega_c}}(s,a)]^\top da ds.$$

Because $\psi^{\pi_{\eta;\omega_c}}(s)$, $\pi_{\eta;\omega_c}(a|s)$, $\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s)$, $\mu_{r,\omega_c}^{\pi_{\eta;\omega_c}}(s,a)$, $\mu_c^{\pi_{\eta;\omega_c}}(s)$, and $\mu_c^{\pi_{\eta;\omega_c}}(s,a)$ are finite (Lemma 4), and $\int_{s \in \mathcal{S}} ds = \mathcal{S}$ and $\int_{a \in \mathcal{A}} da = \mathcal{A}$ are finite, we know that $||\nabla_{\omega_c \eta} F^{[v]}(\omega_c, \eta)||$ is bounded and thus $L_{\nabla_{\omega_c} F^{[v]}}$ exists. $\qquad \square$

Therefore, the approximation error is

$$||\bar{\nabla} F^{[v]}(\omega_c, \bar{\eta}(\omega_c)) - \bar{\nabla} F^{[v]}(\omega_c, \eta^*(\omega_c))|| = ||\nabla_{\omega_c} F^{[v]}(\omega_c, \bar{\eta}(\omega_c)) - \nabla_{\omega_c} F^{[v]}(\omega_c, \eta^*(\omega_c))||,$$
$$\leq L_{\nabla_{\omega_c} F^{[v]}} ||\bar{\eta}(\omega_c) - \eta^*(\omega_c)||.$$

From Lemma 3, we know that $||\bar{\eta}(\omega_c) - \eta^*(\omega_c)||$ decreases to 0 at the rate of $O(\frac{1}{\sqrt{\log K}})$, thus $||\bar{\nabla} F^{[v]}(\omega_c, \bar{\eta}(\omega_c)) - \bar{\nabla} F^{[v]}(\omega_c, \eta^*(\omega_c))||$ decreases to 0 at the rate of $O(\frac{1}{\sqrt{\log K}})$. $\qquad \square$

B.6 Proof of Lemma 2

Learner v's LSCA problem at $\omega_c^{[v]}$ is $\arg\max_{\omega_c \Omega_c} \tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ where $\tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ needs to satisfy the following four conditions (Assumption 3.14 in [37]):
(i) $\nabla \tilde{F}^{[v]}(\omega_c; \omega_c^{[v]}) = N_L \bar{\nabla}^{[v]}$ at $\omega_c^{[v]}$.
(ii) $-\tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ is strongly convex in ω_c.
(iii) $\tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ is continuously differentiable in ω_c.
(iv) $\nabla \tilde{F}^{[v]}(\omega_c; \omega_c^{[v]})$ is Lipschitz continuous in $\omega_c^{[v]}$.

To satisfy these assumptions, we choose the function $\tilde{F}^{[v]}(\omega_c; \omega_c^{[v]}) = -\frac{1}{2}||\omega_c - \omega_c^{[v]}||^2 + (N_L \bar{\nabla}^{[v]})^\top (\omega_c - \omega_c^{[v]})$.

Then $\tilde{\omega}_c^{[v]} = \arg\max_{\omega_c \in \Omega_c} -\frac{1}{2}||\omega_c - \omega_c^{[v]}||^2 + (N_L \bar{\nabla}^{[v]})^\top (\omega_c - \omega_c^{[v]})$. Now, we prove that $\tilde{\omega}_c^{[v]} = \text{Project}_{\Omega_c}(\omega_c^{[v]} + N_L \bar{\nabla}^{[v]})$. Because $\omega_c \in \Omega_c$, by the property of the projection operator, we know that $\tilde{\omega}_c^{[v]} = \arg\min_{\omega_c \in \Omega_c} ||\omega_c - \omega_c^{[v]} - N_L \bar{\nabla}^{[v]}|| = \arg\min_{\omega_c \in \Omega_c} ||\omega_c - \omega_c^{[v]} - N_L \bar{\nabla}^{[v]}||^2 = \arg\min_{\omega_c \in \Omega_c} ||\omega_c - \omega_c^{[v]}||^2 + ||N_L \bar{\nabla}^{[v]}||^2 - 2(N_L \bar{\nabla}^{[v]})^\top (\omega_c - \omega_c^{[v]}) = \arg\max_{\omega_c \in \omega_c} -\frac{1}{2}||\omega_c - \omega_c^{[v]}||^2 + (N_L \bar{\nabla}^{[v]})^\top (\omega_c - \omega_c^{[v]})$.

B.7 Proof of Theorem 1

With the the result of LSCA in subsection B.6, SONATA (Theorem 4 in [38]) can prove that

$$\text{(consensus)} \qquad \lim_{n \to \infty} \max_{v, v' \in \mathcal{V}} ||\omega_c^{[v]}(n) - \omega_c^{[v']}(n)|| = 0,$$

$$\text{(convergence)} \qquad \lim_{n \to \infty} (\sum_{v=1}^{N_L} \bar{\nabla} F^{[v]}(\omega_c^{[v]}(n), \bar{\eta}^{[v]}(\omega_c^{[v]}(n))))^\top (\omega_c - \omega_c^{[v]}(n)) \leq 0$$

Then,

$$\limsup_{n \to \infty} (\nabla F(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))))^\top (\omega_c - \omega_c^{[v]}(n))$$

$$= \limsup_{n \to \infty} (\nabla F(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))) - \sum_{v=1}^{N_L} \bar{\nabla} F^{[v]}(\omega_c^{[v]}(n), \bar{\eta}^{[v]}(\omega_c^{[v]}(n))))^\top (\omega_c - \omega_c^{[v]}(n))$$

$$+ \limsup_{n \to \infty} (\sum_{v=1}^{N_L} \bar{\nabla} F^{[v]}(\omega_c^{[v]}(n), \bar{\eta}^{[v]}(\omega_c^{[v]}(n))))^\top (\omega_c - \omega_c^{[v]}(n)),$$

$$\leq \limsup_{n\to\infty}(\nabla F(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))) - \sum_{v=1}^{N_L}\bar\nabla F^{[v]}(\omega_c^{[v]}(n), \bar\eta^{[v]}(\omega_c^{[v]}(n))))^\top(\omega_c - \omega_c^{[v]}(n)),$$

$$\leq \limsup_{n\to\infty}||\nabla F(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))) - \sum_{v=1}^{N_L}\bar\nabla F^{[v]}(\omega_c^{[v]}(n), \bar\eta^{[v]}(\omega_c^{[v]}(n)))||\cdot||\omega_c - \omega_c^{[v]}(n)||,$$

$$\leq \limsup_{n\to\infty}2\sum_{i=1}^{N_E}l_c^{[i]}||\sum_{v=1}^{N_L}(\bar\nabla F^{[v]}(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))) - \bar\nabla F^{[v]}(\omega_c^{[v]}(n), \bar\eta^{[v]}(\omega_c^{[v]}(n))))||$$

$$\leq \limsup_{n\to\infty}2\sum_{i=1}^{N_E}l_c^{[i]}\sum_{v=1}^{N_L}||\bar\nabla F^{[v]}(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))) - \bar\nabla F^{[v]}(\omega_c^{[v]}(n), \bar\eta^{[v]}(\omega_c^{[v]}(n)))||$$

$$\leq \limsup_{n\to\infty}2\sum_{i=1}^{N_E}l_c^{[i]}\sum_{v=1}^{N_L}L_{\nabla_{\omega_c}F^{[v]}}||\eta^*(\omega_c^{[v]}(n)) - \bar\eta^{[v]}(\omega_c^{[v]}(n))||,$$

where the last inequality follows from Claim 3 and $||\eta^*(\omega_c^{[v]}(n)) - \bar\eta^{[v]}(\omega_c^{[v]}(n))|| = O(\frac{1}{\sqrt{\log K}})$ (Lemma 3). Then, we have

$$\limsup_{n\to\infty}(\nabla F(\omega_c^{[v]}(n), \eta^*(\omega_c^{[v]}(n))))^\top(\omega_c - \omega_c^{[v]}(n)) \leq \frac{\bar M}{\sqrt{\log K}},$$

where $\bar M$ is well-defined because $l_c^{[i]}$ and $L_{\nabla_{\omega_c}F^{[v]}}$ are both positive constants.

C   Appendix: simulation details

In this section, we provide the simulation details. The Python3 code was run on a laptop with one Intel Core i7-9750H 2.60GHz CPU and 16 GB of RAM under Ubuntu 18.04 operating system. Due to the well-known curse of dimensionality, the reinforcement learning (or dynamic programming) of multiple experts is hard to compute. To alleviate this issue, we model the experts as separate MDPs for most of the time and only model them as an MG when they are close to each other.

C.1   Synthetic grid world

The inner communication network for the four learners has two stages: in stage 1, learners 1 and 2 can communicate and learners 3 and 4 can communicate; in stage 2, learners 1 and 4 can communicate and learners 2 and 3 can communicate. Thus the adjacency matrix in stage 1 is

$$\begin{bmatrix} 0.5 & 0.5 & 0 & 0 \\ 0.5 & 0.5 & 0 & 0 \\ 0 & 0 & 0.5 & 0.5 \\ 0 & 0 & 0.5 & 0.5 \end{bmatrix} \text{ and in stage 2 is } \begin{bmatrix} 0.5 & 0 & 0 & 0.5 \\ 0 & 0.5 & 0.5 & 0 \\ 0 & 0.5 & 0.5 & 0 \\ 0.5 & 0 & 0 & 0.5 \end{bmatrix}.$$

The outer communication graph also has two stages, as we need to ensure $\bar N^{[v]}(n) = \mathcal N$, the adjacency matrix in two stages are respectively:

$$\begin{bmatrix} 0.49999 & 0.49999 & 0.00001 & 0.00001 \\ 0.49999 & 0.49999 & 0.00001 & 0.00001 \\ 0.00001 & 0.00001 & 0.49999 & 0.49999 \\ 0.00001 & 0.00001 & 0.49999 & 0.49999 \end{bmatrix} \text{ and } \begin{bmatrix} 0.49999 & 0.00001 & 0.00001 & 0.49999 \\ 0.00001 & 0.49999 & 0.49999 & 0.00001 \\ 0.00001 & 0.49999 & 0.49999 & 0.00001 \\ 0.49999 & 0.00001 & 0.00001 & 0.49999 \end{bmatrix}.$$

The discount factor is 0.9. For each $\omega_{c,j}^{[i]}$, we also need a threshold $\omega_{c,\text{th}}$ to judge whether $\mathcal C_j^{[i]}$ is identified as a part of the ground truth constraint, i.e., $\mathcal C_j^{[i]}$ is identified as a part of the ground truth constraint if $\omega_{c,j}^{[i]} \geq \omega_{c,\text{th}}$ and is not if otherwise. The principle of choosing the value of $\omega_{c,\text{th}}$ is that the probability of entering $\mathcal C_j^{[i]}$ when $\omega_{c,j}^{[i]} = \omega_{c,\text{th}}$ is less than 1% of the probability when $\omega_{c,j}^{[i]} = 0$. In this example, we choose $\omega_{c,\text{th}} = 0.1$.

C.2   Drones motion planning with obstacles

The simulator is built in Gazebo based on a package called hector_quadrotor [39].

In this experiment, there are only two learners who obtain, respectively, four and five pairs of demonstrated trajectories. The inner and outer communication graphs are the same and only have one stage. The adjacency matrix is $\begin{bmatrix} 0.5 & 0.5 \\ 0.5 & 0.5 \end{bmatrix}$. The discount factor is 1 and the drones fly at most 500 steps and we choose $\omega_{c,\text{th}} = 0.05$.

D Appendix: supplementary contexts

D.1 Intuition of the bi-level formulation

The intuition behind this bi-level formulation is inspired by hyperparameter learning [22] and an IRL literature [40]. Paper [40] interprets IRL as an bi-level problem where the outer level is to learn the reward function and the inner level is to learn the corresponding policy. In our case, we treat the constraint as a hyperparameter in our learned environment and we need to recover the corresponding reward function and policy given current constraint. Therefore, in our problem, the outer level is to learn the constraints and the inner level is to learn the corresponding reward function and policy.

D.2 Derivation of centralized bi-level problem and decomposition to the distributed bi-level problem

The global log likelihood of the global demonstration set $\mathcal{D} = \{\zeta^j\}_{j=1}^m$ is $F(\omega_c) = \sum_{j=1}^m \sum_{t=0}^\infty \gamma^t \ln \pi_{\omega_c}(a_t^j | s_t^j)$. To solve this MLE problem, we need to find a parametric policy model of π_{ω_c}. To find such a model, we use an optimization problem (2) based on MCE scheme. Notice that this optimization problem (2) is parameterized by ω_c, thus its optimal solution is also parameterized by ω_c. Its optimal solution is the policy model we want in the likelihood function $F(\omega_c)$. Following well-established MCE IRL literature [25], we can see that the optimal solution of this optimization problem is the constrained soft Bellman policy $\pi_{\eta^*(\omega_c);\omega_c}$, where $\eta^*(\omega_c)$ is the optimization solution of the problem $\min_\eta G(\eta;\omega_c)$. To fully define the MLE problem, we also need the optimal solution $\eta^*(\omega_c)$ of the problem $\min_\eta G(\eta;\omega_c)$. Therefore, we formulate the MLE problem as a bi-level optimization problem: $\max_{\omega_c} F(\omega_c, \eta^*(\omega_c)) = \sum_{j=1}^m \sum_{t=0}^\infty \gamma^t \ln \pi_{\eta^*(\omega_c);\omega_c}(a_t^j | s_t^j)$ s.t. $\eta^*(\omega_c) = \arg\min_\eta G(\eta;\omega_c)$. Now, we finish the derivation of the centralized bi-level problem.

As no learner knows the global demonstration data $\mathcal{D}$ and no learner can formulate F and G, the centralized bi-level problem cannot be directly solved. Therefore, we need to decompose the centralized problem into an equivalent distributed problem that the learners can solve even if each learner only knows its local demonstration. Therefore, we decompose F into multiple $F^{[v]}$ and G into multiple $G^{[v]}$. In remark 3, we show that the distributed problem (3)-(4) is equivalent to the centralized bi-level problem we derive here.

D.3 Explanation of the theoretical results

In our setting, each distributed learner only has a portion of the global demonstration data set and one goal of our distributed algorithm is that the distributed learners can learn as good as a centralized learner who obtains the global data set even if the distributed learners do not share their local demonstration set. In Lemma 3, it is shown that given a cost feature estimate ω_c, the distributed learners can converge to a consensus and the consensus is the optimal solution of the inner problem, i.e., their consensus is the best solution the centralized learner can achieve. In Theorem 1, it is shown that the distributed learners can achieve consensus on the cost weight ω_c which belongs to the stationary point set of the outer problem. Combining Lemma 3 and Theorem 1, we can see that the distributed learners will converge to the consensus on both the reward function and cost function where the learned cost weight belongs to the stationary point set of the outer problem and the learned reward weight is the optimal solution of the inner problem.