GRADIENT BOOSTING PERFORMS GAUSSIAN PROCESS INFERENCE

ABSTRACT

This paper shows that gradient boosting based on symmetric decision trees can be equivalently reformulated as a kernel method that converges to the solution of a certain Kernel Ridge Regression problem. Thus, we obtain the convergence to a Gaussian Process' posterior mean, which, in turn, allows us to easily transform gradient boosting into a sampler from the posterior to provide better knowledge uncertainty estimates through Monte-Carlo estimation of the posterior variance. We show that the proposed sampler allows for better knowledge uncertainty estimates leading to improved out-of-domain detection.

1 INTRODUCTION

Gradient boosting (Friedman, 2001) is a classic machine learning algorithm successfully used for web search, recommendation systems, weather forecasting, and other problems (Roe et al., 2005; Caruana & Niculescu-Mizil, 2006; Richardson et al., 2007; Wu et al., 2010; Burges, 2010; Zhang & Haghani, 2015). In a nutshell, gradient boosting methods iteratively combine simple models (usually decision trees), minimizing a given loss function. Despite the recent success of neural approaches in various areas, gradient-boosted decision trees (GBDT) are still state-of-the-art algorithms for *tabular* datasets containing heterogeneous features (Gorishniy et al., 2021; Katzir et al., 2021).

This paper aims at a better theoretical understanding of GBDT methods for regression problems assuming the widely used RMSE loss function. First, we show that the gradient boosting with regularization can be reformulated as an optimization problem in some Reproducing Kernel Hilbert Space (RKHS) with implicitly defined kernel structure. After obtaining that connection between GBDT and kernel methods, we introduce a technique for sampling from prior Gaussian process distribution with the same kernel that defines RKHS so that the final output would converge to a sample from the Gaussian process posterior. Without this technique, we can view the output of GBDT as the mean function of the Gaussian process.

Importantly, our theoretical analysis assumes the regularized gradient boosting procedure (Algorithm 2) without any simplifications — we only need decision trees to be symmetric (oblivious) and properly randomized (Algorithm 1). These assumptions are non-restrictive and are satisfied in some popular gradient boosting implementations, e.g., CatBoost (Prokhorenkova et al., 2018).

Our experiments confirm that the proposed sampler from the Gaussian process posterior outperforms the previous approaches (Malinin et al., 2021) and gives better knowledge uncertainty estimates and improved out-of-domain detection.

2 BACKGROUND

Assume that we are given a distribution ϱ over $X \times Y$, where $X \subset \mathbb{R}^d$ is called a feature space and $Y \subset \mathbb{R}$ — a target space. Further assume that we are given a dataset $\mathbf{z} = \{(x_i, y_i)\}_{i=1}^N \subset X \times Y$ of size $N \geq 1$ sampled i.i.d. from ϱ. Let us denote by $\rho(\mathrm{d}x) = \int_Y \mathrm{d}\varrho(\mathrm{d}x, \mathrm{d}y)$. W.l.o.g., we also assume that $X = \mathrm{supp}\,\rho = \overline{\{x \in \mathbb{R}^d : \forall \epsilon > 0, \rho(\{x' \in \mathbb{R}^d : \|x - x'\|_{\mathbb{R}^d} < \epsilon\}) > 0\}}$ which is a closed subset of $\mathbb{R}^d$. Moreover, for technical reasons, we assume that $\frac{1}{2N}\sum_{i=1}^N y_i^2 \leq R^2$ for some constant $R > 0$ almost surely, which can always be enforced by clipping. Throughout the paper, we also denote by $\mathbf{x}_N$ and $\mathbf{y}_N$ the matrix of all feature vectors and the vector of targets.

2.1 GRADIENT BOOSTED DECISION TREES

Given a loss function $L : \mathbb{R}^2 \to \mathbb{R}$, a classic gradient boosting algorithm (Friedman, 2001) iteratively combines weak learners (usually decision trees) to reduce the average loss over the train set $\mathbf{z}$: $\mathcal{L}(f) = \mathbb{E}_{\mathbf{z}}[L(f(x), y)]$. At each iteration τ, the model is updated as: $f_\tau(x) = f_{\tau-1}(x) + \epsilon w_\tau(x)$, where $w_\tau(\cdot) \in \mathcal{W}$ is a weak learner chosen from some family of functions $\mathcal{W}$, and ϵ is a learning rate. The weak learner w_τ is usually chosen to approximate the negative gradient of the loss function $-g_\tau(x, y) := -\frac{\partial L(s, y)}{\partial s}\big|_{s=f_{\tau-1}(x)}$:

$$w_\tau = \underset{w \in \mathcal{W}}{\arg\min}\, \mathbb{E}_{\mathbf{z}}\big[\big(-g_\tau(x, y) - w(x)\big)^2\big]. \tag{1}$$

The family $\mathcal{W}$ usually consists of decision trees. In this case, the algorithm is called GBDT (Gradient Boosted Decision Trees). A decision tree is a model that recursively partitions the feature space into disjoint regions called leaves. Each leaf R_j of the tree is assigned to a value, which is the estimated response y in the corresponding region. Thus, we can write $w(x) = \sum_{j=1}^d \theta_j \mathbf{1}_{\{x \in R_j\}}$, so the decision tree is a linear function of the leaf values θ_j.

A recent paper Ustimenko & Prokhorenkova (2021) proposes a modification of classic stochastic gradient boosting (SGB) called *Stochastic Gradient Langevin Boosting* (SGLB). SGLB combines gradient boosting with stochastic gradient Langevin dynamics to achieve global convergence even for non-convex loss functions. The algorithm has two modifications compared to the classic boosting. First, Gaussian noise (parametrized by a *temperature* parameter) is injected into the gradients, and the temperature regulates the level of exploration. Second, *shrinkage* is added to the boosting process: at each iteration, the currently built model is multiplied by a constant smaller than one. As a result, the obtained algorithm provably converges to some stationary distribution (invariant measure) concentrated near the global optimum of the loss function. We mention this method because it samples from similar distribution as our method but with a different kernel.

2.2 ESTIMATING UNCERTAINTY

In addition to the predictive quality, it is often important to detect when the system is uncertain and can be mistaken. For this, different measures of *uncertainty* can be used. There are two main sources of uncertainty: *data uncertainty* (a.k.a. aleatoric uncertainty) and *knowledge uncertainty* (a.k.a. epistemic uncertainty). Data uncertainty arises due to the inherent complexity of the data, such as additive noise or overlapping classes. For instance, if the target is distributed as $y|x \sim \mathcal{N}(f(x), \sigma^2(x))$, then $\sigma(x)$ reflects the level of data uncertainty. This uncertainty can be assessed if the model is probabilistic.

Knowledge uncertainty arises when the model gets input from a region either sparsely covered by or far from the training data. Since the model does not have enough data in this region, it will likely make a mistake. A standard approach to estimating knowledge uncertainty is based on *ensembles* (Gal, 2016; Malinin, 2019). Assume that we have trained an ensemble of several independent models. If all the models understand an input (low knowledge uncertainty), they will give similar predictions. However, for out-of-domain examples (high knowledge uncertainty), the models are likely to provide diverse predictions. For regression tasks, one can obtain knowledge uncertainty by measuring the variance of the predictions provided by multiple models (Malinin, 2019).

Such ensemble-based approaches are standard for neural networks (Lakshminarayanan et al., 2017). Recently, ensembles were also tested for GBDT models (Malinin et al., 2021). The authors consider two ways of generating ensembles: ensembles of independent SGB models and ensembles of independent SGLB models. While empirically methods are very similar, SGLB has better theoretical properties: the convergence of parameters to the stationary distribution allows one to sample models from a particular posterior distribution.

2.3 GAUSSIAN PROCESS INFERENCE

In this section, we briefly describe the basics of Bayesian inference with Gaussian processes that is closely related to our analysis. A random variable f with values in $L_2(\rho)$ is said to be a Gaussian process $f \sim \mathcal{GP}\big(f_0, \sigma^2\mathcal{K} + \delta^2\mathrm{Id}_{L_2}\big)$ with covariance defined via a kernel $\mathcal{K}(x, x') = \mathrm{cov}\big(f(x), f(x')\big)$

and mean value $f_0 \in L_2(\rho)$ iff $\forall g \in L_2(\rho)$ we have

$$\int_X f(x)g(x)\rho(\mathrm{d}x) \sim \mathcal{N}\Big(\int_X f_0(x)g(x)\rho(\mathrm{d}x), \sigma^2 \int_{X \times X} g(x)\mathcal{K}(x,x')g(x')\rho(\mathrm{d}x) \otimes \rho(\mathrm{d}x') + \delta^2 \|g\|_{L_2}^2\Big).$$

A typical Gaussian Process setup is to assume that the target $y|x$ is distributed as $\mathcal{GP}\big(\mathbb{0}_{L_2(\rho)}, \sigma^2 \mathcal{K} + \delta^2 \mathrm{Id}_{L_2}\big)$ for some kernel function[1] $\mathcal{K}(x,x')$ with scales $\sigma > 0$ and $\delta > 0$:

$$y|x = f(x), \ \ f \sim \mathcal{GP}(\mathbb{0}_{L_2(\rho)}, \sigma^2 \mathcal{K} + \delta^2 \mathrm{Id}_{L_2}).$$

The posterior distribution $f(x)|x, \mathbf{x}_N, \mathbf{y}_N$ is again a Gaussian Process $\mathcal{GP}(f_*, \sigma^2 \widetilde{\mathcal{K}} + \delta^2 \mathrm{Id}_{L_2})$ with mean and covariance given by (see Rasmussen & Williams (2006)):

$$f_*(x) = \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \frac{\delta^2}{\sigma^2} I_N\big)^{-1} \mathbf{y}_N,$$

$$\widetilde{\mathcal{K}}(x,x) = \mathcal{K}(x,x) - \mathcal{K}(x, \mathbf{x}_N) \cdot \big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \frac{\delta^2}{\sigma^2} I_N\big)^{-1} \mathcal{K}(\mathbf{x}_N, x)).$$

To estimate the posterior mean $f_*(x) = \int_{\mathbb{R}} f p(f|x, \mathbf{x}_N, \mathbf{y}_N) \, \mathrm{d}f$, we can use the maximum a posteriori probability estimate (MAP) — the minimal norm solution for the following Kernel Ridge Regression (KRR) problem:

$$L(f) = \frac{1}{2N} \sum_{i=1}^{N} \big(f(x_i) - y_i\big)^2 + \frac{\lambda}{2N} \|f\|_{\mathcal{H}}^2 \to \min_{f \in \mathcal{H}},$$

where $\mathcal{H} = \overline{\mathrm{span}\{\mathcal{K}(\cdot, x) | x \in X\}} \subset L_2(\rho)$ is the reproducing kernel Hilbert space (RKHS) for the kernel $\mathcal{K}(\cdot, \cdot)$ and $\lambda = \frac{\delta^2}{\sigma^2}$. We refer to Appendix D for the details on KRR and RKHS.

To solve the KRR problem, one can apply the gradient descent (GD) method in the functional space:

$$f_{\tau+1} = (1 - \frac{\lambda \epsilon}{N}) f_\tau - \epsilon \frac{1}{N} \sum_{i=1}^{N} (f_\tau(x_i) - y_i) \mathcal{K}_{x_i}, \ \ f_0 = \mathbb{0}_{L_2(\rho)}, \tag{2}$$

where $\epsilon > 0$ is a learning rate.

Since this objective is strongly convex due to the regularization, the gradient descent rapidly converges to the unique optimum:

$$f_*^\lambda = \lim_{\tau \to \infty} f_\tau = \mathcal{K}(\cdot, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1} \mathbf{y}_N,$$

see Appendices C and E for the details.

Gradient descent guides f_τ to the posterior mean f_* of the Gaussian Process with kernel $\sigma^2 \mathcal{K} + \delta^2 \mathrm{Id}_{L_2}$. To obtain the posterior variance estimate $\widetilde{\mathcal{K}}(x,x)$ for any x, one can use sampling and introduce a source of randomness in the above iterative scheme as follows:

1. sample $f^{init} \sim \mathcal{GP}(\mathbb{0}_{L_2(\rho)}, \sigma^2 \mathcal{K} + \delta^2 \mathrm{Id}_{L_2})$;
2. set new labels $\mathbf{y}_N^{new} = \mathbf{y}_N - f^{init}(\mathbf{x}_N)$;
3. fit GD $f_\tau(\cdot)$ on $\mathbf{y}_N^{new}$ assuming $F_0(\cdot) = \mathbb{0}_{L_2(\rho)}$;
4. output $f^{init}(\cdot) + f_\tau(\cdot)$ as final model.

This method is known as Sample-then-Optimize (Matthews et al., 2017) and is widely adopted for Bayesian inference. As $\tau \to \infty$, we get $f^{init} + f_\infty$ distributed as a Gaussian Process posterior with the desired mean and variance. Formally, the following result holds.

Lemma 2.1. $f^{init} + f_\infty$ follows the Gaussian Process posterior $\mathcal{GP}(f_*^\lambda, \sigma^2 \widetilde{\mathcal{K}} + \delta^2 \mathrm{Id}_{L_2})$ with:

$$f_*^\lambda(x) = \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1} \mathbf{y}_N,$$

$$\widetilde{\mathcal{K}}(x,x) = \mathcal{K}(x,x) - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1} \mathcal{K}(\mathbf{x}_N, x)).$$

---

[1] $\mathcal{K}(x,x')$ is a kernel function if $\big[\mathcal{K}(x_i, x_j)\big]_{i=1, j=1}^{N,N} \geq 0$ for any $N \geq 1$ and any $x_i \in X$ almost surely.

3 EVOLUTION OF GBDT IN RKHS

3.1 PRELIMINARIES

In our analysis, we assume that we are given a finite set $\mathcal{V}$ of weak learners used for the gradient boosting.[2] Each ν corresponds to a decision tree that defines a partition of the feature space into disjoint regions (leaves). For each $\nu \in \mathcal{V}$, we denote the number of leaves in the tree by $L_\nu \geq 1$. Also, let $\phi_\nu : X \to \{0, 1\}^{L_\nu}$ be a mapping that maps x to the vector indicating its leaf index in the tree ν. This mapping defines a decomposition of X into the disjoint union: $X = \cup_{j=1}^{L_\nu} \{x \in X | \phi_\nu^{(j)}(x) = 1\}$. Having ϕ_ν, we define a weak learner associated with it as $x \mapsto \langle \theta, \phi_\nu(x) \rangle_{\mathbb{R}^{L_\nu}}$ for any choice of $\theta \in \mathbb{R}^{L_\nu}$ which we refer to as 'leaf values'. In other words, θ corresponds to predictions assigned to each region of the space defined by ν.

Let us define a linear space $\mathcal{F} \subset L_2(\rho)$ of all possible ensembles of trees from $\mathcal{V}$:

$$\mathcal{F} = \mathrm{span}\left\{ \phi_\nu^{(j)}(\cdot) : X \to \{0, 1\} \big| \nu \in \mathcal{V}, j \in \{1, \dots, L_\nu\} \right\}.$$

We note that the space $\mathcal{F}$ can be data-dependent since $\mathcal{V}$ may depend on $\mathbf{z}$, but we omit this dependence in the notation for simplicity. Note that we do not take the closure w.r.t. the topology of $L_2(\rho)$ since we assume that $\mathcal{V}$ is finite and therefore $\mathcal{F}$ is finite-dimensional and thus topologically closed.

3.2 GBDT ALGORITHM UNDER CONSIDERATION

Our theoretical analysis holds for classic GBDT algorithms discussed in Section 2.1 equipped with regularization from Ustimenko & Prokhorenkova (2021). The only requirement we need is that the procedure of choosing each new tree has to be properly randomized. Let us discuss a tree selection algorithm that we assume in our analysis.

Each new tree approximates the gradients of the loss function with respect to the current predictions of the model. Since we consider the RMSE loss function, the gradients are proportional to the residuals $r_j = y_j - f(x_j)$, where f is the currently built model. The tree structure is defined by the features and the corresponding thresholds used to split the space. The analysis in this paper assumes the *SampleTree* procedure (see Algorithm 1), which is a classic approach equipped with proper randomization. *SampleTree* builds an *oblivious* decision tree (Prokhorenkova et al., 2018), i.e., all nodes at a given level share the same splitting criterion (feature and threshold).[3] To limit the number of candidate splits, each feature is quantized into $n + 1$ bins. In other words, for each feature, we have n thresholds that can be chosen arbitrarily.[4] The maximum tree depth is limited by m. Recall that we denote the set of all possible tree structures by $\mathcal{V}$.

We build the tree in a top-down greedy manner. At each step, we choose one split among all the remaining candidates based on the following *score* defined for $\nu \in \mathcal{V}$ and residuals r:

$$D(\nu, r) := \frac{1}{N} \sum_{j=1}^{L_v} \frac{\left(\sum_{i=1}^{N} \phi_\nu^{(j)}(x_i) \, r_i \right)^2}{\sum_{i=1}^{N} \phi_\nu^{(j)}(x_i)}. \tag{3}$$

In classic gradient boosting, one builds a tree recursively by choosing such split s that maximizes the score $D((\nu_i, s), r)$ (Ibragimov & Gusev, 2019).[5] Random noise is often added to the scores to improve generalization. In *SampleTree*, we choose a split that maximizes

$$D\big((\nu_i, s), r\big) + \varepsilon, \text{ where } \varepsilon \sim \mathrm{Gumbel}(0, \beta). \tag{4}$$

---

[2]The finiteness of $\mathcal{V}$ is important for our analysis, and it usually holds in practice, see Section 3.2.

[3]In fact, the procedure can be extended to arbitrary trees, but this would over-complicate formulation of the algorithm and would not change the space of tree ensembles as any non-symmetric tree can be represented as a sum of symmetric ones.

[4]A standard approach is to quantize the feature such that all $n + 1$ buckets have approximately the same number of training samples.

[5]Maximizing (3) is equivalent to minimizing the squared error between the residuals and the mean values in the leaves.

Algorithm 1 SampleTree$(r; m, n, \beta)$

> **input:** residuals $r = (r_i)_{i=1}^N$
> **output:** oblivious tree structure $\nu \in \mathcal{V}$
> **hyper-parameters:** number of feature splits n, max. tree depth m, random strength $\beta \in [0, \infty)$
> **definitions:**
> $\mathcal{S} = \{(j, k) | j \in \{1, \ldots, d\}, k \in \{1, \ldots, n\}\}$ — indices of all possible splits
> **instructions:**
> initialize $i = 0$, $\nu_0 = \emptyset$, $\mathcal{S}^{(0)} = \mathcal{S}$
> **repeat**
> > sample $(u_i(s))_{s \in \mathcal{S}^{(i)}} \sim \mathrm{U}([0, 1]^{nd-i})$
> > choose next split as $\{s_{i+1}\} = \underset{s \in \mathcal{S}^{(i)}}{\arg\max}\Big(D\big((\nu_i, s), r\big) - \beta \log(-\log u_i(s))\Big)$
> > update tree: $\nu_{i+1} = (\nu_i, s_{i+1})$
> > update candidate splits: $\mathcal{S}^{(i+1)} = \mathcal{S}^{(i)} \backslash \{s_{i+1}\}$
> >
> > $i = i + 1$
> **until** $i \geq m$ **or** $\mathcal{S}^{(i)} = \emptyset$
> **return:** ν_i

Algorithm 2 TrainGBDT$(\mathbf{z}; \epsilon, T, m, n, \beta, \lambda)$

> **input:** dataset $\mathbf{z} = (\mathbf{x}_N, \mathbf{y}_N)$
> **hyper-parameters:** learning rate $\epsilon > 0$, regularization $\lambda > 0$, iterations of boosting T, parameters of SampleTree m, n, β
>
> **instructions:**
> initialize $\tau = 0$, $f_0(\cdot) = \mathbb{0}_{L_2(\rho)}$
> **repeat**
> > $r_\tau = \mathbf{y}_N - f_\tau(\mathbf{x}_N)$ — compute residuals
> > $\nu_\tau = \text{SampleTree}(r_\tau; m, n, \beta)$ — construct a tree
> > $\theta_\tau = \left(\frac{\sum_{i=1}^N \phi_{\nu_\tau}^{(j)}(x_i) r_\tau^{(i)}}{\sum_{i=1}^N \phi_{\nu_\tau}^{(j)}(x_i)} \right)_{j=1}^{L_{\nu_\tau}}$ — set values in leaves
> > $f_{\tau+1}(\cdot) = (1 - \frac{\lambda \epsilon}{N}) f_\tau(\cdot) + \epsilon \Big\langle \phi_{\nu_\tau}(\cdot), \theta_\tau \Big\rangle_{\mathbb{R}^{L_{\nu_\tau}}}$ — update model
> > $\tau = \tau + 1$
> **until** $\tau \geq T$
> **return:** $f_T(\cdot)$

Here β is random strength: $\beta = 0$ gives the standard greedy approach, while $\beta \to \infty$ gives the uniform distribution among all possible split candidates.

To sum up, *SampleTree* is a classic oblivious tree construction but with added random noise. We do this to make the distribution of trees regular in a certain sense: roughly speaking, the distributions should stabilize with iterations by converging to some fixed distribution.

Given the algorithm *SampleTree*, we describe the gradient boosting procedure assumed in our analysis in Algorithm 2. It is a classic GBDT algorithm but with the update rule $f_{\tau+1}(\cdot) = (1 - \lambda\epsilon/N) f_\tau(\cdot) + \epsilon w_\tau(x)$. In other words, we shrink the model at each iteration, which serves as regularization (Ustimenko & Prokhorenkova, 2021). Such shrinkage is available, e.g., in the CatBoost library.

3.3 DISTRIBUTION OF TREES

The *SampleTree* algorithm induces a local family of distributions $p(\cdot | f, \beta)$ for each $f \in \mathcal{F}$:

$$p(\mathrm{d}\nu | f, \beta) = \mathbb{P}\Big(\Big\{\text{SampleTree}\big(\mathbf{y}_N - f(\mathbf{x}_N); m, n, \beta\big) \in \mathrm{d}\nu\Big\}\Big).$$

Remark 3.1. Lemma 3.7 ensure that such distribution coincides with the one where we use $f_*(\mathbf{x}_N)$ instead of $\mathbf{y}_N$. This is due to the fact that $D(\nu, \mathbf{y}_N - f(\mathbf{x}_N)) = D(\nu, f_* - f(\mathbf{x}_N)) \, \forall \nu \in \mathcal{V}, \, f \in \mathcal{F}$.

The following lemma describes the distribution $p(\mathrm{d}\nu | f, \beta)$, see Appendix F for the proof. Note that for oblivious decision trees, changing the order of splits does not affect the obtained partition. Hence, we assume that each tree is defined by an unordered set of splits.

Lemma 3.2. *(Probability of a tree)*

$$p(\nu | f, \beta) = \sum_{\varsigma \in \mathcal{P}_m} \prod_{i=1}^m \frac{e^{\frac{D(\nu_{\varsigma, i}, r)}{\beta}}}{\sum_{s \in \mathcal{S} \backslash \nu_{\varsigma, i-1}} e^{\frac{D((\nu_{\varsigma, i-1}, s), r)}{\beta}}},$$

where the sum is over all permutations $\varsigma \in \mathcal{P}_m$, $\nu_{\varsigma, i} = (s_{\varsigma(1)}, \ldots, s_{\varsigma(i)})$, and $\nu = (s_1, \ldots, s_m)$.

Let us define the stationary distribution of trees as $\pi(\cdot) = \lim_{\beta \to \infty} p(\cdot | f, \beta)$. It follows from Remark 3.1 that we also have $\pi(\cdot) = p(\cdot | f_*, \beta)$.

Corollary 3.3. *(Stationary distribution is the uniform distribution over tree structures)*
We have $\pi(\mathrm{d}\nu) = |\mathrm{d}\nu| / \binom{nd}{m}$, where $\binom{nd}{m} = \frac{(nd)!}{(nd-m)! m!}$.

3.4 RKHS Structure

In this section, we describe the evolution of GBDT in a certain Reproducing Kernel Hilbert Space (RKHS). Even though, the problem is finite dimensional, treating it as functional regression is more beneficial as dimension of the ensembles space growth rapidly and therefore we want to obtain dimension-free constants which is impossible if we treat it as finite dimensional optimization problem. Let us start with defining necessary kernels.

Definition 3.4. A weak learner's kernel $k_\nu(\cdot, \cdot)$ is a kernel function associated with a tree structure $\nu \in \mathcal{V}$ which can be defined as:

$$k_\nu(x, x') = \sum_{j=1}^{L_\nu} w_\nu^{(j)} \phi_\nu^{(j)}(x) \phi_\nu^{(j)}(x'), \text{ where } w_\nu^{(j)} = \frac{N}{\max\{N_\nu^{(j)}, 1\}}, N_\nu^{(j)} = \sum_{i=1}^{N} \phi_\nu^{(j)}(x_i).$$

This weak learner's kernel is a building block for any other possible kernel in boosting and is used to define the iterations of the boosting algorithm analytically.

Definition 3.5. We also define a *greedy* kernel of the gradient boosting algorithm as follows:

$$\mathcal{K}_f(x, x') = \int_{\mathcal{V}} k_\nu(x, x') p(\mathrm{d}\nu | f, \beta).$$

This greedy kernel is a kernel that guides the GBDT iterations, i.e., we can think of each iteration as SGD with a kernel from 3.5, and 3.4 is used as a stochastic gradient estimator of the Fréchet derivative in RKHS defined by the kernel from 3.5.

Definition 3.6. Finally, there is a *stationary* kernel $\mathcal{K}(x, x')$ that is independent from f:

$$\mathcal{K}(x, x') = \int_{\mathcal{V}} k_\nu(x, x') \pi(\mathrm{d}\nu),$$

which we call a *prior* kernel of the gradient boosting.

This kernel defines the limiting solution since the gradient projection on RKHS converges to zero, and thus 3.5 converges to 3.6.

Having the space of functions $\mathcal{F}$, we define a reproducing kernel Hilbert space (RKHS) structure $\mathcal{H} = (\mathcal{F}, \langle \cdot, \cdot \rangle_{\mathcal{H}})$ on it using a scalar product defined as

$$\langle f, f \rangle_{\mathcal{H}} = \min_{g \in L_2(\rho)} \int_X g^2(x) \rho(\mathrm{d}x) \text{ s.t. } \mathcal{K}^{\frac{1}{2}} g = f,$$

$$\langle f, g \rangle_{\mathcal{H}} = \frac{1}{4}\left(\langle f + g, f + g \rangle_{\mathcal{H}} - \langle f - g, f - g \rangle_{\mathcal{H}}\right),$$

where $\mathcal{K}^{\frac{1}{2}}$ is the unique symmetric positive definite (SPD) square root operator such that $\left(\mathcal{K}^{\frac{1}{2}}\right)^2 = \mathcal{K}$ assuming that it acts as $\mathcal{K}[g] = \int_X \mathcal{K}(\cdot, x) g(x) \rho(\mathrm{d}x) \in L_2(\rho)$ for any $g \in L_2(\rho)$.

Now, let us define the empirical error of a model f:

$$V(f, \lambda) = \frac{1}{2N}\left\|\mathbf{y}_N - f(\mathbf{x}_N)\right\|_{\mathbb{R}^N}^2 + \frac{\lambda}{2N}\|f\|_{\mathcal{H}}^2 - \inf_{f' \in \mathcal{F}}\left(\frac{1}{2N}\left\|\mathbf{y}_N - f'(\mathbf{x}_N)\right\|_{\mathbb{R}^N}^2 + \frac{\lambda}{2N}\|f'\|_{\mathcal{H}}^2\right).$$

Let us also define the following unique functions:

$$f_*^\lambda \in \arg\min_{f \in \mathcal{F}} V(f, \lambda).$$

And $f_* = \lim_{\lambda \to 0} f_*^\lambda \in \arg\min_{f \in \mathcal{F}} V(f)$ where $V(f) = V(f, 0)$. It is known that such f_* exists and is unique since the set of all solutions is convex, and therefore there is a unique minimizer of the norm $\|\cdot\|_{\mathcal{H}}$.

Finally, the following lemmas will be helpful for our analysis. In particular, we use them to analyze the distribution of trees in Section 3.3 and to derive the formula of the GBDT iteration in terms of kernels in Lemma 3.9. See Appendix D for the proofs.

Lemma 3.7. $\langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N} = 0$ *for any* $f \in \mathcal{F}$.

Lemma 3.8. $V(f, \lambda) = \frac{1}{2N}\|f_*(\mathbf{x}_N) - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 + \frac{\lambda}{2N}\|f\|_{\mathcal{H}}^2 + \text{const}.$

Lemma 3.9. *Iterations* f_τ *of Gradient Boosting (Algorithm 2) can be written in the form:*

$$f_{\tau+1} = (1 - \frac{\lambda\epsilon}{N})f_\tau + \frac{\epsilon}{N} k_{\nu_\tau}(\cdot, \mathbf{x}_N)\left[f_*(\mathbf{x}_N) - f_\tau(\mathbf{x}_N)\right], \nu_\tau \sim p(\nu | f_\tau, \beta).$$

4 KERNEL GRADIENT BOOSTING

4.1 KERNEL RIDGE REGRESSION

Consider the sequence $\{f_\tau\}_{\tau\in\mathbb{N}}$ generated by the gradient boosting algorithm. Its evolution is described by Lemma 3.9. The following theorem estimates the expected (w.r.t. the randomness of tree selection) empirical error of f_T relative to the best possible ensemble. The proof can be found in Appendix G.

Theorem 4.1. *Let* $C = \frac{8\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(8M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(8M_\beta(M_\beta^2-1)+3\right)\right)R_1^2$. *Assume that* $\frac{(1+\lambda)}{8N} \geq \epsilon$, $\frac{N+\lambda}{N}\frac{2mM_\beta C}{\beta} < \frac{1+\lambda}{2N}$ *and define* $T_1 = \left\lceil\frac{2M_\beta N}{(1+M_\beta\lambda))\epsilon}\log\frac{R^2}{C}\right\rceil + 1$ *then* $\forall T \geq T_1$ *it holds that*

$$V(f_T,\lambda) \leq Ce^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)N^2}R^2 + \frac{8}{1+\lambda}\left(\epsilon\left(1+\frac{\lambda}{N}\right) + 4M_\beta^2\frac{m\lambda C}{\beta N}\right)R_1^2$$

Corollary 4.2. *(Convergence to the solution of the KRR / Convergence to the Gaussian Process posterior mean function). Under assumptions of the previous theorem we have:*

$$\mathbb{E}\|f_T - f_*^\lambda\|_{L_2}^2 \leq \frac{NC}{1+\lambda}e^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)^2N}R^2 + \frac{8}{(1+\lambda)^2}\left(N\epsilon\left(1+\frac{\lambda}{N}\right) + 4M_\beta^2\frac{m\lambda C}{\beta}\right)R_1^2$$

We need this corollary to be sure that the Monte-Carlo sampling would approximately yield correct means and variances of the Gaussian process. We note that the estimate is pretty conservative. First, it relies on the estimate $\lambda_{\min}(\nabla_{\mathcal{H}}^2 V(f)) \geq \frac{1}{N}$ (when restricted on its image) which in fact in the $N \to \infty$ limit converges to $\lambda_{\min}(\lim_{N\to\infty}\mathcal{K}) > 0$ independent from N which makes the bound of order $\mathcal{O}(\frac{\lambda^2}{\beta N} + \epsilon)$ and requirement on $0 < \epsilon \leq \mathcal{O}(1)$ for any $T \geq T_1$ and $T_1 = \mathcal{O}\left(-\frac{1}{\epsilon}\log\mathcal{O}(\epsilon + \frac{\lambda^2}{\beta N})\right)$ which is linear convergence. Moreover, the condition $\frac{N+\lambda}{N}\frac{2mM_\beta C}{\beta} < \frac{1+\lambda}{2N}$ translates into $\beta \geq \mathcal{O}(1) > 0$. If we do not use liming argument and use the bound as is, then it can be made of order ε by setting $\beta \geq \frac{\lambda^2}{N\varepsilon}$ and $\epsilon \leq \frac{\varepsilon}{N}$ which imply $T_1 = \mathcal{O}\left(\frac{N^2}{\varepsilon}\log\mathcal{O}(\frac{1}{\varepsilon})\right)$ and $T - T_1 = \mathcal{O}\left(\frac{N^2}{\varepsilon}\log\mathcal{O}(\frac{\lambda}{\varepsilon})\right)$ which yields that $T = \mathcal{O}\left(\frac{N^2}{\varepsilon}\log\mathcal{O}(\frac{1}{\varepsilon})\right)$.

4.2 PRIOR SAMPLING

We introduce Algorithm 3 for sampling from the prior distribution. *SamplePrior* generates an ensemble of random trees (with random splits and random values in leaves). Note that while being random, the tree structure depends on the dataset features $\mathbf{x}_N$ since candidate splits are based on $\mathbf{x}_N$.

We first note that the process $h_T(\cdot)$ is centered with covariance operator $\mathcal{K}$:

$$\mathbb{E}h_T(x) = 0 \ \forall x \in X,$$
$$\mathbb{E}h_T(x)h_T(y) = \mathcal{K}(x,y) \ \forall x,y \in X. \quad (5)$$

Then, we show that $h_T(\cdot)$ converges to the Gaussian Process in the limit.

Lemma 4.3. *The following convergence holds almost surely in* $x \in X$:

$$h_T(\cdot) \xrightarrow[T\to\infty]{} \mathcal{GP}\left(\mathbb{0}_{L_2(\rho)},\mathcal{K}\right).$$

Algorithm 3 SamplePrior(T, m, n)

hyper-parameters: number of iterations T, parameters of SampleTree m, n

instructions:
initialize $\tau = 0$, $h_0(x) = 0$
repeat
 $\nu_\tau = \text{SampleTree}(\mathbb{0}_{\mathbb{R}^N}; m, n, 1)$ — sample random tree
 $\theta_\tau \sim \mathcal{N}\left(\mathbb{0}_{\mathbb{R}^{L_{\nu_\tau}}}, \text{diag}\left(\frac{N}{\max\{N_{\nu_\tau}^{(j)},1\}} : j \in \{1,\ldots,L_{\nu_\tau}\}\right)\right)$ — generate random values in leaves
 $h_{\tau+1}(\cdot) = h_\tau(\cdot) + \frac{1}{\sqrt{T}}\langle\phi_{\nu_\tau}(\cdot),\theta_\tau\rangle_{\mathbb{R}^{L_{\nu_\tau}}}$ — update model
 $\tau = \tau + 1$
until $\tau \geq T$
return: $h_T(\cdot)$

(a) D	(b) $\widetilde{\mu}$	(c) $\log\widetilde{\sigma}$

Figure 1: KGB on a synthetic dataset.

4.3 Posterior Sampling

Now we are ready to introduce Algorithm 4 for sampling from the posterior. The procedure is simple: we first perform T_0 iterations of SamplePrior to obtain a function $h_{T_0}(\cdot)$ and then we train a standard GBDT model $f_{T_1}(\cdot)$ approximating $\mathbf{y}_N - \sigma h_{T_0}(\mathbf{x}_N) + \mathcal{N}(\mathbb{0}_N, \delta^2 I_N)$. Our final model is $\sigma h_{T_0}(\cdot) + f_{T_1}(\cdot)$. We further refer to this procedure as *SamplePosterior* or *KGB* (Kernel Gradient Boosting) for brevity. Denote

$$h_\infty = \lim_{T_0 \to \infty} h_{T_0}, f_\infty = \lim f_{T_1},$$

Algorithm 4 SamplePosterior$(\mathbf{z}; \epsilon, T_1, T_0, m, n, \beta, \sigma, \delta)$

input: dataset $\mathbf{z} = (\mathbf{x}_N, \mathbf{y}_N)$
hyper-parameters: learning rate $\epsilon > 0$, boosting iteration T_1, SamplePrior iterations T_0, parameters of SampleTree m, n, β, kernel scale $\sigma > 0$ (default $\sigma = 1$), noise scale $\delta > 0$ (default: $\delta = 0.01$)
instructions:
$h_{T_0}(\cdot) = \text{SamplePrior}(T_0, m, n)$
$\mathbf{y}_N^{new} = \mathbf{y}_N - \sigma h_{T_0}(\mathbf{x}_N) + \mathcal{N}(\mathbb{0}_N, \delta^2 I_N)$
$f_{T_1}(\cdot) = \text{TrainGBDT}\big((\mathbf{x}_N, \mathbf{y}_N^{new}); \epsilon, T_1, m, n, \beta, \frac{\delta^2}{\sigma^2}\big)$

return: $\sigma h_{T_0}(\cdot) + f_{T_1}(\cdot)$

---

where the first limit is with respect to the point-wise convergence of stochastic processes and the second one with respect to $L_2(\rho)$ convergence.

The following theorem shows that KGB indeed samples from the desired posterior. The proof directly follows from Lemmas 4.3 and 2.1.

Theorem 4.4. *In the limit, the output of Algorithm 4 follows the Gaussian posterior:*

$$\sigma h_\infty(\cdot) + f_\infty(\cdot) + \mathcal{N}(0, \delta^2) \sim \mathcal{GP}\big(\widetilde{f}, \widetilde{\mathcal{K}}\big)$$

with mean $\widetilde{f}(x) = \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathbf{y}_N$ *and covariance* $\widetilde{\mathcal{K}}(x, x) = \delta^2 + \sigma^2\big(\mathcal{K}(x, x) - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x)\big)$.

5 Experiments

This section empirically evaluates the proposed KGB algorithm and shows that it indeed allows for better knowledge uncertainty estimates.

Synthetic experiment To illustrate the KGB algorithm in a controllable setting, we first conduct a synthetic experiment. For this, we defined the feature distribution as uniform over $D = \{(x, y) \in [0, 1]^2 : \frac{1}{10} \le (x - \frac{1}{2})^2 - (y - \frac{1}{2})^2 \le \frac{1}{4} \land (x \le \frac{2}{5} \lor x \ge \frac{3}{5}) \land (y \le \frac{2}{5} \lor y \ge \frac{3}{5})\}$. We sample 10K points from $U([0, 1]^2)$ and take into the train set only those that fall into D. The target is defined as $f(x, y) = x + y$. Figure 1(a) illustrates the training dataset colored with the target values. For evaluation, we take the same 10K points without restricting them to D.

For KGB, we fix $\epsilon = 0.3$, $T_0 = 100$, $T_1 = 900$, $\sigma = 10^{-2}$, $\delta = 10^{-4}$ $\beta = 0.1$, $m = 4$, $n = 64$, and sampled 100 KGB models. Figure 1(b) shows the estimated by Monte-Carlo posterior mean $\widetilde{\mu}$. On Figure 1(c), we show $\log\widetilde{\sigma}$, where $\widetilde{\sigma}^2$ is the posterior variance estimated by Monte-Carlo. One can see that the posterior variance is small in-domain and grows when we move outside the dataset D, as desired.

Table 1: Predictive performance, RMSE

Dataset	Single			Ensemble		
	SGB	SGLB	KGB	SGB	SGLB	KGB
Boston	3.06	3.12	**2.81**	3.04	3.10	**2.82**
Concrete	5.21	5.11	**4.36**	5.21	5.10	**4.30**
Energy	0.57	0.54	**0.33**	0.57	0.54	**0.33**
Kin8nm	0.14	0.14	**0.11**	0.14	0.14	**0.10**
Naval	0.00	**0.00**	0.00	0.00	**0.00**	0.00
Power	**3.55**	3.56	3.48	3.52	3.54	3.43
Protein	3.99	3.99	**3.79**	3.99	3.99	**3.76**
Wine	0.63	0.63	**0.61**	0.63	0.63	**0.60**
Yacht	0.82	0.84	**0.52**	0.83	0.84	**0.50**
Year	**8.99**	**8.96**	**8.97**	**8.97**	**8.93**	**8.94**

Table 2: Error and OOD detection

Dataset	PRR			AUC		
	SGB	SGLB	KGB	SGB	SGLB	KGB
Boston	36	37	**43**	80	80	**88**
Concrete	29	29	**37**	92	92	**93**
Energy	36	31	**60**	100	100	99
Kin8nm	18	**19**	20	45	45	41
Naval	**55**	56	35	100	100	**100**
Power	8	9	**31**	72	73	**76**
Protein	30	29	**35**	99	99	**100**
Wine	25	19	**37**	74	72	**87**
Yacht	74	78	**86**	62	60	**69**
Year	30	30	**32**	67	57	**71**

Experiment on real datasets Uncertainty estimates for GBDTs have been previously analyzed by Malinin et al. (2021). Our experiments on real datasets closely follow their setup, and we compare the proposed KGB with SGB, SGLB, and their ensembles. For the experiments, we use several standard regression datasets (Gal & Ghahramani, 2016). The implementation details can be found in Appendix H. The code of our experiments can be found on GitHub.[6]

We note that in our setup we *cannot* compute likelihoods as kernel $\mathcal{K}$ is defined implicitly and its evaluation requires summing up among all possible trees structures number of which grows as $(nd)^m$ which is unfeasible, not to mention requirement to inverse the kernel which requires $\mathcal{O}(N^{2+\omega})$ number of operations. Therefore, typical Bayesian setup is not applicable and that is why we resort to uncertainty estimation setup described in Malinin et al. (2021).

First, we compare KGB with SGLB since they both sample from similar posterior distributions. Thus, this comparison allows us to find out which of the algorithms does a better sampling from the posterior and thus provides us with more reliable estimates of knowledge uncertainty. Moreover, we consider the SGB approach as the most "straightforward" way to generate an ensemble of models.

In Table 1, we compare the predictive performance of the methods. Interestingly, we obtain improvements on almost all the datasets. Here we perform cross-validation to estimate statistical significance with paired t-test and highlight the approaches that are insignificantly different from the best one (p-value > 0.05). Then, we check whether uncertainty measured as the variance of the model's predictions can be used to detect errors and out-of-domain inputs. Detecting errors can be evaluated via the Prediction-Rejection Ratio (PRR) (Malinin, 2019; Malinin et al., 2020). PRR measures how well uncertainty estimates correlate with errors and rank-order them. Out-of-domain (OOD) detection is usually assessed via the area under the ROC curve (AUC-ROC) for the binary task of detecting whether a sample is OOD (Hendrycks & Gimpel, 2016). For this, one needs an OOD test set. We use the same OOD test sets (sampled from other datasets) as Malinin et al. (2021). The results of this experiment are given in Table 2. We can see that the proposed method significantly outperforms the baselines for out-of-domain detection. These improvements can be explained by the theoretical soundness of KGB: convergence properties are theoretically grounded and non-asymptotic. In contrast, for SGB, there are no general results applicable in our setting, while for SGLB the guarantees are asymptotic. In summary, these results show that our approach is superior to SGB and SGLB, achieving smaller values of RMSE and having better knowledge uncertainty estimates.

6 Conclusion

This paper theoretically analyses the classic gradient boosting algorithm. In particular, we show that GBDT converges to the solution of a certain Kernel Ridge Regression problem. We also introduce a simple modification of the classic algorithm allowing one to sample from the Gaussian posterior. The proposed method gives much better knowledge uncertainty estimates than the existing approaches.

We highlight the following important directions for future research. First, to explore how one can control the kernel and use it for better knowledge uncertainty estimates. Also, we do not analyze generalization in the current work, which is another important research topic. Finally, we need to establish universal approximation property which further justifies need for functional formalism.

---

[6]https://github.com/anon-author-123/Gradient-Boosting-Performs-Gaussian-Process-Inference

REFERENCES

Susan Athey, Julie Tibshirani, and Stefan Wager. Generalized random forests. *The Annals of Statistics*, 47(2):1148–1178, 2019.

Matej Balog, Balaji Lakshminarayanan, Zoubin Ghahramani, Daniel M Roy, and Yee Whye Teh. The mondrian kernel. *arXiv preprint arXiv:1606.05241*, 2016.

Christopher JC Burges. From RankNet to LambdaRank to LambdaMART: An overview. *Learning*, 11(23-581):81, 2010.

Rich Caruana and Alexandru Niculescu-Mizil. An empirical comparison of supervised learning algorithms. In *Proceedings of the 23rd international conference on Machine learning*, pp. 161–168. ACM, 2006.

Jerome H Friedman. Greedy function approximation: a gradient boosting machine. *Annals of statistics*, pp. 1189–1232, 2001.

Yarin Gal. *Uncertainty in Deep Learning*. PhD thesis, University of Cambridge, 2016.

Yarin Gal and Zoubin Ghahramani. Dropout as a Bayesian Approximation: Representing Model Uncertainty in Deep Learning. In *Proc. 33rd International Conference on Machine Learning (ICML-16)*, 2016.

Yury Gorishniy, Ivan Rubachev, Valentin Khrulkov, and Artem Babenko. Revisiting deep learning models for tabular data. In *Advances in Neural Information Processing Systems 34 (NeurIPS 2021)*, 2021.

Dan Hendrycks and Kevin Gimpel. A Baseline for Detecting Misclassified and Out-of-Distribution Examples in Neural Networks. http://arxiv.org/abs/1610.02136, 2016. arXiv:1610.02136.

Bulat Ibragimov and Gleb Gusev. Minimal variance sampling in stochastic gradient boosting. *Advances in Neural Information Processing Systems*, 32, 2019.

Liran Katzir, Gal Elidan, and Ran El-Yaniv. Net-dnf: Effective deep modeling of tabular data. In *International Conference on Learning Representations (ICLR)*, 2021.

B. Lakshminarayanan, A. Pritzel, and C. Blundell. Simple and Scalable Predictive Uncertainty Estimation using Deep Ensembles. In *Proc. Conference on Neural Information Processing Systems (NIPS)*, 2017.

David G. Luenberger. *Optimization by Vector Space Methods*. 1969.

Andrey Malinin. *Uncertainty Estimation in Deep Learning with application to Spoken Language Assessment*. PhD thesis, University of Cambridge, 2019.

Andrey Malinin, Bruno Mlodozeniec, and Mark JF Gales. Ensemble distribution distillation. In *International Conference on Learning Representations*, 2020. URL https://openreview.net/forum?id=BygSP6Vtvr.

Andrey Malinin, Liudmila Prokhorenkova, and Aleksei Ustimenko. Uncertainty in gradient boosting via ensembles. In *International Conference on Learning Representations*, 2021.

Matthews, Hron, Turner, and Ghahramani. Sample-then-optimize posterior sampling for bayesian linear models. In *NIPS 2017*, 2017.

Liudmila Prokhorenkova, Gleb Gusev, Aleksandr Vorobev, Anna Veronika Dorogush, and Andrey Gulin. Catboost: unbiased boosting with categorical features. In *Proceedings of the 32nd International Conference on Neural Information Processing Systems (NeurIPS)*, pp. 6638–6648, 2018.

C.E. Rasmussen and C.K.I. Williams. *Gaussian Processes for Machine Learning*. the MIT press, 2006.

Matthew Richardson, Ewa Dominowska, and Robert Ragno. Predicting clicks: estimating the click-through rate for new ads. In *Proceedings of the 16th international conference on World Wide Web*, pp. 521–530. ACM, 2007.

Byron P Roe, Hai-Jun Yang, Ji Zhu, Yong Liu, Ion Stancu, and Gordon McGregor. Boosted decision trees as an alternative to artificial neural networks for particle identification. *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 543(2):577–584, 2005.

Aleksei Ustimenko and Liudmila Prokhorenkova. Sglb: Stochastic gradient langevin boosting. In *International Conference on Machine Learning*, pp. 10487–10496. PMLR, 2021.

Qiang Wu, Christopher JC Burges, Krysta M Svore, and Jianfeng Gao. Adapting boosting for information retrieval measures. *Information Retrieval*, 13(3):254–270, 2010.

Yanru Zhang and Ali Haghani. A gradient boosting method to improve travel time prediction. *Transportation Research Part C: Emerging Technologies*, 58:308–324, 2015.

A   NOTATION USED IN THE PAPER

For convenience, let us list some frequently used notation:

- $X \subset \mathbb{R}^d$ — feature space;
- d — dimension of feature vectors;
- $Y \subset \mathbb{R}$ — target space;
- ρ — distribution of features;
- N — number of samples;
- $\mathbf{z} = (\mathbf{x}_N, \mathbf{y}_N)$ — dataset;
- $\mathcal{V}$ — set of all possible tree structures;
- $L_\nu : \mathcal{V} \to \mathbb{N}$ — number of leaves for $\nu \in \mathcal{V}$;
- $D(\nu, r)$ — score used to choose a split (3);
- $\mathcal{S}$ — indices of all possible splits;
- n — number of borders in our implementation of SampleTree;
- m — depth of the tree in our implementation of SampleTree;
- β — random strength;
- ϵ — learning rate;
- λ — regularization;
- $\mathcal{F}$ — space of all possible ensembles of trees from $\mathcal{V}$;
- $\phi_\nu : X \to \{0, 1\}^{L_\nu}$ — tree structure;
- $\phi_\nu^{(j)}$ — indicator of j-th leaf;
- $V(f)$ — empirical error of a model f relative to the best possible $f' \in \mathcal{F}$;
- $k_\nu(\cdot, \cdot)$ — single tree kernel;
- $\mathcal{K}(\cdot, \cdot)$ — stationary kernel of the gradient boosting;
- $p(\cdot | f, \beta)$ — distribution of trees, $f \in \mathcal{F}$;
- $\pi(\cdot) = \lim_{\beta \to \infty} p(\cdot | f, \beta) = p(\cdot | f_*, \beta)$ — stationary distribution of trees;
- σ — kernel scale.

B   ADDITIONAL RELATED WORK

Let us briefly discuss some additional related work. Mondrian forest method Balog et al. (2016) and Generalized Random Forests Athey et al. (2019) besides having links to the kernel methods, in fact, have the underlying limiting RKHS space that is much smaller than the space of all possible ensembles built on the same weak learners due to the independence of the trees that are added to the ensemble. Therefore, there is an issue of high bias when comparing plain gradient boosting with the plain random forest method. Also, these two methods are built from scratch to obtain an RKHS interpretation while we provide a link between the existing standard gradient boosting approaches to the kernel methods, i.e., we do not create a novel gradient boosting algorithm but rather show that the existing ones already have such a link to derive convergence rates and to exploit such linkage to obtain formal Gaussian process interpretation of the gradient boosting learning to get uncertainty estimates using well-established gradient boosting libraries.

C   CONVEX OPTIMIZATION IN FUNCTIONAL SPACES

In this section, we formulate basic definitions of differentiability in functional spaces and the theorem on the convergence of gradient descent in functional spaces. For the proof of the theorem and further details on convex optimization in functional space, the reader can consult Luenberger (1969).

We consider $\mathcal{H}$ to be a Hilbert space with some scalar product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$.

Definition C.1. We say that $F : \mathcal{H} \to \mathbb{R}$ is Fréchet differentiable if for any $f \in \mathcal{H}$ there exists a bounded linear functional $L_f : \mathcal{H} \to \mathbb{R}$ such that $\forall h \in \mathcal{H}$

$$F(f + h) = F(f) + L_f[h] + o(\|h\|).$$

The value of $L_f : \mathcal{H} \to \mathbb{R}$ is denoted by $\mathcal{D}_f F(f)$ and is called a Fréchet differential of F at point f. So, Fréchet differential is a functional $\mathcal{D}_f F : \mathcal{H} \to \mathcal{B}(\mathcal{H}, \mathbb{R})$, where $\mathcal{B}(X, Y)$ denotes a normed space of linear bounded functionals from X to Y.

Definition C.2. Let $F : \mathcal{H} \to \mathbb{R}$ be Fréchet differentiable with a Fréchet differential $\mathcal{D}_f F(f)$ that is a bounded linear functional. Then, by the Riesz theorem there exists a unique h_f such that $\big(\mathcal{D}_f F(f)\big)[h] = \langle h_f, h \rangle_{\mathcal{H}} \ \forall h \in \mathcal{H}$. We call such element a *gradient* of F in $\mathcal{H}$ at $f \in \mathcal{H}$ and denote it by $\nabla_{\mathcal{H}} F(f) = h_f \in \mathcal{H}$.

Definition C.3. $F : \mathcal{H} \to \mathbb{R}$ is said to be twice Fréchet differentiable if $\mathcal{D}_f F$ is Fréchet differentiable, where the definition of Fréchet differential is analogous to Definition C.1 with the only difference that $\mathcal{D}_f F$ takes values in $\mathcal{B}(\mathcal{H}, \mathbb{R})$. The second Fréchet differential is denoted by $\mathcal{D}_f^2 F : \mathcal{H} \to \mathcal{B}(\mathcal{H}, \mathcal{B}(\mathcal{H}, \mathbb{R}))$. As there is an isomorphism between $\mathcal{B}(\mathcal{H}, \mathcal{B}(\mathcal{H}, \mathbb{R}))$ and $\mathcal{B}(\mathcal{H} \times \mathcal{H}, \mathbb{R})$, we can consider the second Fréchet differential to take values in $\mathcal{B}(\mathcal{H} \times \mathcal{H}, \mathbb{R})$. Henceforth, we will not differentiate between $\mathcal{B}(\mathcal{H}, \mathcal{B}(\mathcal{H}, \mathbb{R}))$ and $\mathcal{B}(\mathcal{H} \times \mathcal{H}, \mathbb{R})$.

Definition C.4. A linear operator $P : \mathcal{H} \times \mathcal{H} \to \mathbb{R}$ is said to be semi-positive definite (denoted by $P \succeq 0$) if $\forall f \in \mathcal{H}$ we have $P(f, f) \geq 0$. P is said to be positive definite ($P \succ 0$) if $\forall f \in \mathcal{H} \setminus \{0\}$: $P(f, f) > 0$.

Definition C.5. Given two linear operators $P, G : \mathcal{H} \times \mathcal{H} \to \mathbb{R}$ we write $P \succeq G$ if $P - G \succeq 0$ and $P \succ G$ if $P - G \succ 0$.

Let $I \in B(\mathcal{H} \times \mathcal{H}, \mathbb{R})$ be a linear operator defined as $I(g, h) = (g, h)_{\mathcal{H}}$.

Theorem C.6. *Let F be bounded below and twice Fréchet differentiable functional on a Hilbert space $\mathcal{H}$. Assume that $\mathcal{D}_f^2 F(f)$ satisfies $0 \prec mI \preceq \mathcal{D}_f^2 F(f) \preceq \mu I$. Then the gradient descent scheme:*

$$f_{k+1} = f_k - \epsilon \nabla_{\mathcal{H}} F(f_k)$$

converges to f_* that minimizes F.

Proof. For the proof see Luenberger (1969). $\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad \square$

D KERNEL RIDGE REGRESSION AND RKHS

Definition D.1. $\mathcal{K} : X \times X \to \mathbb{R}$ is called a kernel function if it is positive semi-definite, i.e., $\forall N \in \mathbb{N}^+ \ \forall \mathbf{x}_N \in \mathbb{R}^N : \ \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) \succeq 0$.

Definition D.2. For any kernel function we can define a Reproducing Kernel Hilbert Space (RKHS)

$$\mathcal{H}(\mathcal{K}) = \overline{\mathrm{span}\left\{\mathcal{K}(\cdot, x) \big| x \in X\right\}} \subset L_2(\rho)$$

with a scalar product such that

$$\langle f, \mathcal{K}(\cdot, x) \rangle_{\mathcal{H}(\mathcal{K})} = f(x).$$

Alternatively, we can define the scalar product as

$$\langle f, f \rangle_{\mathcal{H}} = \min_{g \in L_2(\rho)} \int_X g^2(x)\rho(\mathrm{d}x) \ \text{s.t.} \ \mathcal{K}^{\frac{1}{2}} g = f,$$

$$\langle f, g \rangle_{\mathcal{H}} = \frac{1}{4}\left(\langle f + g, f + g \rangle_{\mathcal{H}} - \langle f - g, f - g \rangle_{\mathcal{H}}\right),$$

where $\mathcal{K}^{\frac{1}{2}}$ corresponds to the unique SPD square root operator such that $\left(\mathcal{K}^{\frac{1}{2}}\right)^2 = \mathcal{K}$ assuming that it acts as $\mathcal{K}[g] = \int_X \mathcal{K}(\cdot, x)g(x)\rho(\mathrm{d}x) \in L_2(\rho)$ for any $g \in L_2(\rho)$. Both define the same scalar product.

Consider the following Kernel Ridge Regression problem:

$$V(f, \lambda) = \frac{1}{2N}\|\mathbf{y}_N - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 + \frac{\lambda}{2N}\|f\|_{\mathcal{H}(K)}^2 - \min_{f \in \mathcal{F}} V(f, \lambda) \to \min_{f \in \mathcal{H}(\mathcal{K})}$$

and the following Kernel Ridgeless Regression problem:

$$V(f) = \frac{1}{2N}\|\mathbf{y}_N - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \min_{f \in \mathcal{F}} V(f) \to \min_{f \in \mathcal{H}(\mathcal{K})} \ .$$

Lemma D.3. $\min_{\mathcal{H}(\mathcal{K})} V(f, \lambda)$ *has the only solution*

$$f_*^\lambda = \mathcal{K}(\cdot, \mathbf{x}_N)(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I)^{-1}\mathbf{y}_N \ .$$

Proof. First, let us show that $f_*^\lambda \in \text{span}\{\mathcal{K}(\cdot, x_i)\}$. Let $\mathcal{H}(\mathcal{K}) = \text{span}\{\mathcal{K}(\cdot, x_i)\} \oplus \text{span}\{\mathcal{K}(\cdot, x_i)\}^\perp$ and consider the projector $P : \mathcal{H}(\mathcal{K}) \to \mathcal{H}(\mathcal{K})$ onto the space $\text{span}\{\mathcal{K}(\cdot, x_i)\}$.

It is easy to show that $P(f)(\mathbf{x}_N) = f(\mathbf{x}_N)$ for any $f \in \mathcal{H}(\mathcal{K})$. Indeed,

$$(f - P(f))[\mathbf{x}_N] = \langle f - P(f), \mathcal{K}(\cdot, \mathbf{x}_N)\rangle = 0 \ .$$

If f_*^λ does not lie in $\text{span}\{\mathcal{K}(\cdot, x_i)\}$, then $\|f_*^\lambda\|_{\mathcal{H}(\mathcal{K})} > \|P(f_*^\lambda)\|_{\mathcal{H}(\mathcal{K})}$ and $L(P(f_*^\lambda), \lambda) < L(f_*^\lambda, \lambda)$. We get a contradiction with the minimality of f_*^λ.

Now, let us prove the existence of f_*^λ. Consider $f = \mathcal{K}(\cdot, \mathbf{x}_N)c$, where $c \in \mathbb{R}^N$. Then, we find the optimal c by taking a derivative of $V(f, \lambda)$ with respect to c and equating it to zero:

$$\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)c - y_N) + \lambda\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)c = 0 \ .$$

Then,

$$c_v = (\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I)^{-1}(y_N + v) \ ,$$

where $v \in \ker \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$. Note that all $\mathcal{K}(\cdot, \mathbf{x}_N)c_v$ are equal. Then, we have the only solution of the KRR problem:

$$f_*^\lambda = \mathcal{K}(\cdot, \mathbf{x}_N)(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I)^{-1}\mathbf{y}_N \ .$$

$\square$

Lemma D.4. $\min_{\mathcal{H}(\mathcal{K})} V(f)$ *has the only solution in* $\text{span}\{\mathcal{K}(\cdot, \mathbf{x}_i)\}$ *and it is the solution of minimum RKHS norm:*

$$f_* = \mathcal{K}(\cdot, \mathbf{x}_N)\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)^\dagger\mathbf{y}_N \ .$$

Proof. Consider $f = \mathcal{K}(\cdot, \mathbf{x}_N)c$, where $c \in \mathbb{R}^N$. Now consider $L(f)$ and differentiate it with respect to c. If we equate the derivative to zero, we get:

$$\frac{1}{N}\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)c - \mathbf{y}_N) = 0 \ .$$

Then, $\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)c - (\mathbf{y}_N + v) = 0$ for some $v \in \ker \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$. Note that $\mathbf{y}_N + \ker \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) \cap \text{Im}\,\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) \neq \emptyset$. Then, for any v such that $\mathbf{y}_N + v \in \text{Im}\,\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$ there exist a solution $c_v = \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)^\dagger(\mathbf{y}_N + v) + \ker \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$. This follows from the fact that $\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)^\dagger$ is an orthoprojector onto $\text{Im}\,\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$. Note that

$$f_* = \mathcal{K}(\cdot, \mathbf{x}_N)(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)^\dagger(\mathbf{y}_N + v) + \ker \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)) = \mathcal{K}(\cdot, \mathbf{x}_N)\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)^\dagger\mathbf{y}_N.$$

Then, the existence and uniqueness of f_* follow. $\square$

Now, consider a linear space $\mathcal{F} \subset L_2(\rho)$ of all possible ensembles of trees from $\mathcal{V}$:

$$\mathcal{F} = \text{span}\left\{\phi_\nu^{(j)}(\cdot) : X \to \{0, 1\} \big| \nu \in \mathcal{V}, j \in \{1, \dots, L_\nu\}\right\} \ .$$

Consider such funtional on space F:

$$V(f) = \frac{1}{2N}\left\|\mathbf{y}_N - f(\mathbf{x}_N)\right\|_{\mathbb{R}^N}^2 - \inf_{f' \in \mathcal{F}}\left(\frac{1}{2N}\left\|\mathbf{y}_N - f'(\mathbf{x}_N)\right\|_{\mathbb{R}^N}^2\right) \ .$$

and define unique function:

$$f_* = \lim_{\lambda \to 0_+} \arg\min_{f \in \mathcal{F}} V(f, \lambda) \in \arg\min_{f \in \mathcal{F}} V(f).$$

Then following two lemmas hold.

Lemma D.5 (Lemma 3.7 in the main text). $\langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N} = 0$ *for any* $f \in \mathcal{F}$.

Proof. Assume that $\langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N} \neq 0$ for some $f \in \mathcal{F}$. Then, for some $f \in \mathcal{F}$, $\langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N} > 0$. We have:

$$\|\mathbf{y}_N - (f_* + \alpha f)(\mathbf{x}_N)\|_{\mathbb{R}^N}^2$$
$$= \|\mathbf{y}_N - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - 2\alpha \langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N} + \alpha^2 \|f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2$$
$$< \|\mathbf{y}_N - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2$$

for small enough $\alpha > 0$, which contradicts with the definition of f_*.

$\square$

Lemma D.6 (Lemma 3.8 in the main text). $V(f, \lambda) = \frac{1}{2N} \|f_*(\mathbf{x}_N) - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 + \frac{\lambda}{2N} \|f\|_{\mathcal{H}}^2 + C$.

Proof. We need to prove it only for $V(f)$ withour regularization as for regularized it follows immidiately (note that f_* is minimizer of not regularized objective). By definition,

$$V(f) = \frac{1}{2N} \|\mathbf{y}_N - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \frac{1}{2N} \|\mathbf{y}_N - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 .$$

Now, let us prove that

$$\|\mathbf{y}_N - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \|\mathbf{y}_N - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \|f_*(\mathbf{x}_N) - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 = 0 .$$

Indeed,

$$\|\mathbf{y}_N - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \|\mathbf{y}_N - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 - \|f_*(\mathbf{x}_N) - f(\mathbf{x}_N)\|_{\mathbb{R}^N}^2$$
$$= -2 \langle f_*(\mathbf{x}_N), f_*(\mathbf{x}_N) \rangle_{\mathbb{R}^N} - 2 \langle \mathbf{y}_N, f(\mathbf{x}_N) - f_*(\mathbf{x}_N) \rangle_{\mathbb{R}^N} + 2 \langle f_*(\mathbf{x}_N), f(\mathbf{x}_N) \rangle_{\mathbb{R}^N}$$
$$= -2 \langle \mathbf{y}_N - f_*(\mathbf{x}_N), f(\mathbf{x}_N) - f_*(\mathbf{x}_N) \rangle_{\mathbb{R}^N} = 0 ,$$

where the last equality follows from the previous lemma. $\square$

E GAUSSIAN PROCESS INFERENCE

In this section, we prove Lemma 2.1 from Section 2.3 of the main text.

Firstly, consider the Kernel Ridge Regression problem loss functional:

$$V(f, \lambda) = \frac{1}{2N} \sum_{i=1}^{N} \left(f(x_i) - y_i\right)^2 + \frac{\lambda}{2N} \|f\|_{\mathcal{H}}^2 - \min_{f \in \mathcal{F}} V(f, \lambda) \to \min_{f \in \mathcal{H}(\mathcal{K})} .$$

We will show that it satisfies the conditions needed for Theorem C.6. We will also deduce the formula of the gradient of L in order to show that gradient descent takes the form (2).

Lemma E.1. $L(f)$ *is Fréchet differentiable with the differential given by:*

$$\mathcal{D}_f V(f, \lambda) = \frac{\lambda}{N} \langle f, \cdot \rangle_{\mathcal{H}(\mathcal{K})} + \frac{1}{N} \sum_{i=1}^{N} \left(f(x_i) - y_i\right) ev_{x_i} ,$$

where $ev_{x_i} : \mathcal{H}(\mathcal{K}) \to \mathbb{R}$ is a bounded linear functional such that $ev_{x_i}(f) = f(x_i) = (f, \mathcal{K}(x_i, \cdot))_{\mathcal{H}(\mathcal{K})}$.[7]

Proof. As Fréchet differential is linear, we only need to find Fréchet differential for $(f(x_i) - y_i)^2$.

Note that $(f(x_i) - y_i)^2$ is a composition of two functions:

$$F : \mathcal{H}(\mathcal{K}) \to \mathbb{R}, \ F = ev_{x_i} - y_i ,$$

$$G : \mathbb{R} \to \mathbb{R}, \ G(x) = x^2 .$$

---

[7]We further use the notation $\mathcal{K}_{x_i} := \mathcal{K}(\cdot, x_i)$.

The differential of the composition can be found as:

$$\mathcal{D}_f G(F(f)) = \frac{\partial}{\partial x} G(F(f)) \mathcal{D}_f F(f) \, ,$$

$$\mathcal{D}_f G(F(f)) = 2(f(x_i) - y_i) ev_{x_i} \, ,$$

where $\mathcal{D}_f F(f) = ev_{x_i}$ because

$$ev_{x_i}(f + h) - y_i = ev_{x_i}(f) - y_i + ev_{x_i}(h) \, .$$

$\square$

Lemma E.2. *The gradient of $V(f)$, Riesz representative of the functional above, is given by:*

$$\nabla_f V(f, \lambda) = \frac{\lambda}{N} f + \frac{1}{N} \sum_{i=1}^{N} (f(x_i) - y_i) \mathcal{K}_{x_i} \, .$$

Proof. Trivially follows from the previous lemma. $\square$

Lemma E.3. *V is twice Fréchet differentiable with the differential given by:*

$$\mathcal{D}_f^2 V : \mathcal{H}(\mathcal{K}) \to B(\mathcal{H}(\mathcal{K}), B(\mathcal{H}(\mathcal{K}), \mathbb{R})) \, ,$$

$$\mathcal{D}_f^2 V(f, \lambda)[h] = \frac{\lambda}{N} h + \frac{1}{N} \sum_{i=1}^{N} h(x_i) ev_{x_i} \, .$$

Proof. Due to the linearity of Fréchet differential and lemma E.1 we need to find only Fréchet differential for $(f(x_i) - y_i) ev_{x_i}$.

Consider $S(f) = (f(x_i) - y_i) ev_{x_i}$. Then we need to find $V_f \in B(\mathcal{H}(\mathcal{K}), B(\mathcal{H}(\mathcal{K}), \mathbb{R}))$ such that $S(f + h) = S(f) + V_f[h] + o(\|h\|)$.

It is easy to show that $h \mapsto h(x_i) ev_{x_i} \in B(\mathcal{H}(\mathcal{K}), B(\mathcal{H}, \mathbb{R}))$ and $S(f + h) = S(f) + h(x_i) ev_{x_i}$. Thus, we get that $\mathcal{D}_f S(f)[h] = h(x_i) ev_{x_i}$. From this the statement of the lemma follows. $\square$

Given all the above lemmas, as a corollary of Theorem C.6, we have the following.

Corollary E.4. *Gradient descent, defined by the following iterative scheme:*

$$f_{\tau+1} = \left(1 - \frac{\lambda \epsilon}{N}\right) f_\tau - h \frac{1}{N} \sum_{i=1}^{N} (f_\tau(x_i) - y_i) \mathcal{K}_{x_i} \, ,$$

$$f_0 = \mathbb{0}_{L_2(\rho)}$$

converges to the optimum of $V(f, \lambda)$. Thus,

$$f_*^\lambda = \lim_{\tau \to \infty} f_\tau = \mathcal{K}(\cdot, \mathbf{x}_N) \left(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N \right)^{-1} \mathbf{y}_N \, . \tag{6}$$

Proof. As mentioned in Section 2.3, the solution of the Kernel Ridgeless Regression problem coincides with a posterior of a Gaussian. Since gradient descent converges to this solution and thus to the posterior of a Gaussian, equation (6) follows. $\square$

Lemma E.5. *Consider the gradient descent:*

$$f_{\tau+1} = \left(1 - \frac{\lambda \epsilon}{N}\right) f_\tau - \epsilon \frac{1}{N} \sum_{i=1}^{N} (f_\tau(x_i) - y_i) \mathcal{K}_{x_i} \, ,$$

$$f_0 = \mathbb{0}_{L_2(\rho)} \, ,$$

$$f_\infty = \lim_{\tau \to \infty} f_\tau$$

and the following randomization scheme:

> 1. *sample* $f^{init} \sim \mathcal{GP}(\mathbb{0}_{L_2(\rho)}, \sigma^2 \mathcal{K} + \delta^2 \mathrm{Id}_{L_2})$;
> 2. *set new labels* $\mathbf{y}_N^{new} = \mathbf{y}_N - f^{init}(\mathbf{x}_N)$;
> 3. *fit GD* $f_\tau(\cdot)$ *on* $\mathbf{y}_N^{new}$ *assuming* $f_0(\cdot) = \mathbb{0}_{L_2(\rho)}$;
> 4. *output* $\hat{f}(\cdot) = f^{init}(\cdot) + f_\infty(\cdot)$ *as final model.*

Then, $\hat{f}$ from the scheme above follows the Gaussian Process posterior with the following mean and covariance:

$$\mathbb{E}\hat{f}(x) = \mathcal{K}(x, \mathbf{x}_N)\Big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\Big)^{-1}\mathbf{y}_N$$

$$\mathrm{cov}(\hat{f}(x)) = \delta^2 + \sigma^2\Big(\mathcal{K}(x, x) - \mathcal{K}(x, \mathbf{x}_N)\Big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\Big)^{-1}\mathcal{K}(\mathbf{x}_N, x)\Big).$$

Proof.

$$f_\infty = \mathcal{K}(\cdot, \mathbf{x}_N)\Big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\Big)^{-1}\mathbf{y}_N^{new} = \mathcal{K}(\cdot, \mathbf{x}_N)\Big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\Big)^{-1}(\mathbf{y}_N - f^{init}(\mathbf{x}_N)).$$

Let us find the distribution of $\hat{f}$ at $x \in \mathbb{R}^n$. It can be easily seen that:

$$\mathbb{E}\hat{f}(x) = \mathcal{K}(x, \mathbf{x}_N)\Big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\Big)^{-1}\mathbf{y}_N.$$

Let us now calculate covariance:

$$\begin{aligned}
\mathrm{cov}\hat{f}(x) &= \mathbb{E}(\hat{f}(x) - \mathbb{E}\hat{f}(x))(\hat{f}(x) - \mathbb{E}\hat{f}(x))^T \\
&= \mathbb{E}\big(f^{init}(x) - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}f^{init}(\mathbf{x}_N)\big) \\
&\quad \cdot \big(f^{init}(x) - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}f^{init}(\mathbf{x}_N)\big)^T \\
&= \mathbb{E}f^{init}(x)f^{init}(x)^T - \mathbb{E}f^{init}(x)f^{init}(\mathbf{x}_N)^T\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x) \\
&\quad - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathbb{E}f^{init}(\mathbf{x}_N)f^{init}(x)^T \\
&\quad + \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathbb{E}f^{init}(\mathbf{x}_N)f^{init}(\mathbf{x}_N)^T\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x) \\
&= \delta^2 + \sigma^2\big(\mathcal{K}(x, x) - 2\mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x) \\
&\quad + \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x)\big) \\
&= \delta^2 + \sigma^2\big(\mathcal{K}(x, x) - \mathcal{K}(x, \mathbf{x}_N)\big(\mathcal{K}(\mathbf{x}_N, \mathbf{x}_N) + \lambda I_N\big)^{-1}\mathcal{K}(\mathbf{x}_N, x)\big),
\end{aligned}$$

which is exactly what we need. $\qquad\square$

F   DISTRIBUTION OF TREES

Lemma F.1 (Lemma 3.2 in the main text)**.**

$$p(\nu|f, \beta) = \sum_{\varsigma \in \mathcal{P}_m} \prod_{i=1}^{m} \frac{e^{\frac{D(\nu_{\varsigma,i}, r)}{\beta}}}{\sum_{s \in \mathcal{S}\backslash\nu_{\varsigma,i-1}} e^{\frac{D((\nu_{\varsigma,i-1}, s), r)}{\beta}}},$$

where the sum is over all permutations $\varsigma \in \mathcal{P}_m$, $\nu_{\varsigma,i} = (s_{\varsigma(1)}, \ldots, s_{\varsigma(i)})$, and $\nu = (s_1, \ldots, s_m)$.

Proof. Let us fix some permutation $\varsigma \in \mathcal{P}_m$. W.l.o.g., let $\varsigma = \mathrm{id}_{\mathcal{P}_m}$, i.e. $\varsigma(i) = i \,\forall i$. It remains to derive the formula for the fixed permutation. The probability of adding the next split given the previously build tree is:

$$P(\nu_{i-1} \cup s_i | \nu_{i-1}) = \frac{e^{\frac{1}{\beta} D(\nu_i, r)}}{\sum_{s \in \mathcal{S} \setminus \nu_{i-1}} e^{\frac{1}{\beta} D((\nu_{i-1}, s), r)}},$$

which comes from (4) and the Gumbel-SoftMax trick. Then, we decompose the probability $P(\nu)$ of a tree as:

$$P(\nu) = \prod_{i=1}^{m} P(\nu_{i-1} \cup s_i | \nu_{i-1}),$$

and so for the fixed permutation we have

$$P(\nu) = \prod_{i=1}^{m} \frac{e^{\frac{1}{\beta} D(\nu_i, r)}}{\sum_{s \in \mathcal{S} \setminus \nu_{i-1}} e^{\frac{1}{\beta} D((\nu_{i-1}, s), r)}}.$$

Then we sum over all permutations and the lemma follows.

$\square$

Now, let us define the following value indicating how different are the distribution of trees for f and f_*:

$$\Gamma_\beta(f) = \max \left\{ \left\| \frac{\mathrm{d}p(\cdot | f_*, \beta)}{\mathrm{d}p(\cdot | f, \beta)} \right\|_{L_\infty(\pi)}, \left\| \frac{\mathrm{d}p(\cdot | f, \beta)}{\mathrm{d}p(\cdot | f_*, \beta)} \right\|_{L_\infty(\pi)} \right\}.$$

Lemma F.2. *The following bound relates the distributions.*

$$\Gamma_\beta(f) \le e^{\frac{2mV(f)}{\beta}}.$$

Proof. Consider $\pi = p(\cdot | f_*, \beta)$ and the following expression $P(\nu, \varsigma)$:

$$P(\nu, \varsigma) := \prod_{i=1}^{m} \frac{e^{\frac{1}{\beta} D(\nu_{\varsigma, i}, r)}}{\sum_{s \in \mathcal{S} \setminus \nu_{\varsigma, i-1}} e^{\frac{1}{\beta} D((\nu_{\varsigma, i-1}, s), r)}}.$$

Then,

$$\sum_{\varsigma \in \mathcal{P}_m} P(\nu, \varsigma) \le \sum_{\varsigma \in \mathcal{P}_m} e^{\frac{m}{\beta} D(\nu, r)} \prod_{i=1}^{m} \frac{1}{\sum_{s \in \mathcal{S} \setminus \nu_{\varsigma, i-1}} 1} \le e^{\frac{2mV(f)}{\beta}} \pi(\nu).$$

where in second inequality we used $D(\nu, r) \le 2V(f)$ which straightly follows from the definition.

By noting that the probabilities remain the same if we shift $D(\cdot, r) \leftarrow D(\cdot, r) - 2V(f)$ which becomes everywhere non-positive and allows us to do the above trick once more but in reverse manner: if we formally replace the D with such modified function and repeat the steps with reversing the inequalities which is needed since the new function is everywhere negative then the lemma follows.

$$\sum_{\varsigma \in \mathcal{P}_m} P(\nu, \varsigma) \ge \sum_{\varsigma \in \mathcal{P}_m} e^{\sum_{i=1}^{m} \frac{1}{\beta} D(\nu_{\varsigma, i}, r) - \frac{m}{\beta} 2V(f)} \prod_{i=1}^{m} \frac{1}{\sum_{s \in \mathcal{S} \setminus \nu_{i-1}} 1} \ge e^{-\frac{2mV(f)}{\beta}} \pi(\nu).$$

$\square$

G PROOF OF THEOREM 4.1

G.1 RKHS STRUCTURE

For a weak learner ν, we define a covariance operator:

$$\Sigma_\nu[f] = \frac{1}{N} k_\nu(\cdot, \mathbf{x}_N) f(\mathbf{x}_N), \ \Sigma_\nu : \mathcal{H} \to \mathcal{H}.$$

Using this operator, we define $\Sigma_f = \int_\mathcal{V} \Sigma_\nu p(\mathrm{d}\nu | f, \beta)$ and $\Sigma = \int_\mathcal{V} \Sigma_\nu \pi(\mathrm{d}\nu)$. These operators are typically referred to as covariance operators.

Here we formulate and prove several lemmas about the RKHS structure and operators $\Sigma, \Sigma_f, \Sigma_\nu$.

Lemma G.1. *The following formula holds $\forall h, g \in \mathcal{H}$:*

$$\langle \Sigma[h], g \rangle_{\mathcal{H}} = \frac{1}{N} \sum_{i=1}^{N} h(x_i) g(x_i).$$

Lemma G.2. *(Covariation majorization) The following operator inequality holds $\forall \nu \in \mathcal{V}$ and $\forall f \in \mathcal{H}$:*

$$\lambda_{\max}(\Sigma_\nu) \leq 1$$

$$\lambda_{\min}(\Sigma_f) \geq \frac{1}{N}$$

Proof. We will show that $\lambda_{max}(\Sigma) \leq 1$ where π in the definition of Σ can be chosen arbitrary. This way the statement would follow for $\Sigma, \Sigma_f, \Sigma_\nu$ by taking π to be equal $p(\cdot \mid f_*, \beta), p(\cdot \mid f, \beta)$, or $\delta_\nu(\cdot)$ respectively. The operator Σ is dual to $K_N = \frac{1}{N} \mathcal{K}(\mathbf{x}_N, \mathbf{x}_N)$ since we can define the operator

$$S_N[f] = \left[\langle f, \mathcal{K}(\cdot, x_1) \rangle_{\mathcal{H}}, \ldots, \langle f, \mathcal{K}(\cdot, x_N) \rangle_{\mathcal{H}} \right] \in \mathbb{R}^N,$$

so that $\Sigma = \frac{1}{N} S_N^* S_N$ and $K_N = \frac{1}{N} S_N S_N^*$ and, thus, has the same spectra.

Therefore, instead of Σ, we can study the maximal eigenvalue of K_N but the latter admits another decomposition

$$K_N = \int_{\mathcal{V}} \Phi_\nu^T \left\{ \Phi_\nu \Phi_\nu^T \right\}^\dagger \Phi_\nu \pi(\mathrm{d}\nu),$$

where $\Phi_\nu = \phi_\nu(\mathbf{x}_N) \in \mathbb{R}^{L_\nu \times N}$. The key observation is that the sub-expectation expression $\Phi_\nu^T \left\{ \Phi_\nu \Phi_\nu^T \right\}^\dagger \Phi_\nu$ defines a matrix with maximal eigenvalue bounded by 1 since the former is an orthoprojector which is also shown in Ustimenko & Prokhorenkova (2021). Therefore, the expectation also inherits this bound on the maximal eigenvalue which with symmetry of K_N gives us the desired inequality.

Now we need to show that $\lambda_{\min}(\Sigma_f) \geq \frac{1}{N}$. Similarly to the previous proof, we need to show that $\lambda_{\min}(\Sigma_{f_*}) \geq \frac{1}{N}$ and then to replace $\pi \leftarrow p(\cdot | f, \beta)$. Consider the following formula:

$$\Sigma = \frac{1}{N} \sum_{i=1}^{N} \mathcal{K}(\cdot, x_i) \otimes_{\mathcal{H}} \mathcal{K}(\cdot, x_i),$$

$$\Sigma = \frac{1}{N} \sum_{i=1}^{N} \mathcal{K}(x_i, x_i) \left(\frac{\mathcal{K}(\cdot, x_i)}{\sqrt{\mathcal{K}(x_i, x_i)}} \otimes_{\mathcal{H}} \frac{\mathcal{K}(\cdot, x_i)}{\sqrt{\mathcal{K}(x_i, x_i)}} \right),$$

where $(a \otimes_{\mathcal{H}} b)[c] = \langle b, c \rangle_{\mathcal{H}} a$. If $a = b$ and $\|a\|_{\mathcal{H}} = 1$, then 1 and 0 are the only eigenvalues of $a \otimes_{\mathcal{H}} a$. Then by induction assuming that we already summed up some of such rank one operators one of two scenarios can happen if we add one more rank one operator: either the rank does not increase or the rank increases by one. In the first case we obtain the operator that is greater in the operator sense then the previously obtained one whilst both of them share the same linear space as image and coimage which due to positive definiteness of both of them on the space can happen only if all eigenvalues of the new one greater or equal compared to the ones of the previous one. In the second scenario the newly added rank one operator adds new eigenvalue 1 due to symmetry of the operators. Finally, by noting that $\inf_{x \in X} \mathcal{K}(x, x) \geq 1$, we obtain the bound.

$\square$

Corollary G.3. *(Expected squared norm majorization by RKHS norm) The following bound holds $\forall f \in \mathcal{H}$:*

$$\|f\|_{L_2(\rho)} \leq \|f\|_{\mathcal{H}}.$$

Proof. From $\lambda_{\max}(\Sigma) \leq 1$ it follows that $\lambda_{\max}(\mathcal{K}) = \lim_{N \to \infty} \lambda_{\max}(\Sigma) \leq 1$. Let $\mathcal{K}^{-1}$ be taken as the unique inverse $\mathcal{K}^{-1}$ on $\mathcal{F}$ and for arbitrary $f \in L_2(\rho)$ we consider $L_2(\rho) = \mathcal{F} \oplus \mathcal{F}^\perp$ which implies $f = f_1 + f_2, f_1 \in \mathcal{F}, f_2 \in \mathcal{F}^\perp \forall f \in L_2(\rho)$ and, thus, allows us to define $\mathcal{K}^{-1} f = \mathcal{K}^{-1} f_1 \in$

$\mathcal{F}$. Such definition of the inverse implies that $\lambda_{\min}(\mathcal{K}^{-1}) = \frac{1}{\lambda_{\max}(\mathcal{K})} \geq 1$ which when substituted back into the product

$$\langle f, f \rangle_{\mathcal{H}} = \langle \mathcal{K}^{-\frac{1}{2}} f, \mathcal{K}^{-\frac{1}{2}} f \rangle_{L_2(\rho)} = \langle \mathcal{K}^{-1} f, f \rangle_{L_2(\rho)}$$

implies the bound. $\qquad\square$

Lemma G.4. $\|k_\nu(\mathbf{x}_N, \mathbf{x}_N)\| \leq N$ for any $\nu \in \mathcal{V}$.

Proof. Consider $S_1, ..., S_{L_\nu}$ — unions such that $S_i = \{x_j | \phi_\nu^{(i)}(x_j) = 1\}$. Then, it is easy to see that

$$k_\nu = \sum_{i=1}^{L_\nu} w_\nu^{(i)} \mathbb{1}_{N_\nu^i \times N_\nu^i},$$

where $\mathbb{1}_{n \times n}$ is a matrix of size $n \times n$ consisting of ones. Then, we note that $\|\mathbb{1}_{n \times n}\| = n$ and now the statement of the lemma follows. $\qquad\square$

Lemma G.5. *For any* $\nu \in \mathcal{V}$, *we have* $k_\nu(\cdot, \mathbf{x}_N)[\mathbf{y}_N - f_*(\mathbf{x}_N)] = 0$.

Proof. Follows from Lemma 3.7. $\qquad\square$

Lemma G.6 (Lemma 3.9 in the main text). *Iterations* f_τ *of Gradient Boosting (Algorithm 2) can be written in the form:*

$$f_{\tau+1} = \left(1 - \frac{\lambda\epsilon}{N}\right) f_\tau + \frac{\epsilon}{N} k_{\nu_\tau}(\cdot, \mathbf{x}_N)[\mathbf{y}_N - f_\tau(\mathbf{x}_N)] = (1 - \frac{\lambda\epsilon}{N}) f_\tau + \frac{\epsilon}{N} k_{\nu_\tau}(\cdot, \mathbf{x}_N)[f_*(\mathbf{x}_N) - f_\tau(\mathbf{x}_N)],$$

$$\nu_\tau \sim p(\nu | f_\tau, \beta).$$

Proof. According to Algorithm 2:

$$f_{\tau+1}(\cdot) = \left(1 - \frac{\lambda\epsilon}{N}\right) f_\tau(\cdot) + \epsilon \langle \phi_{\nu_\tau}(\cdot), \theta_\tau \rangle_{\mathbb{R}^{L_{\nu_\tau}}} \text{ for } \theta_\tau = \left(\frac{\sum_{i=1}^N \phi_{\nu_\tau}^{(j)}(x_i) r_\tau^{(i)}}{\sum_{i=1}^N \phi_{\nu_\tau}^{(j)}(x_i)}\right)_{j=1}^{L_{\nu_\tau}}.$$

Thus,

$$f_{\tau+1} = \left(1 - \frac{\lambda\epsilon}{N}\right) f_\tau + \epsilon \frac{1}{N} \sum_{j=1}^{L_{\nu_\tau}} \omega_{\nu_\tau}^j \phi_{\nu_\tau}^j \sum_{i: \phi_{\nu_\tau}^j(x_i)=1} r_\tau^i.$$

Now note that $k_{\nu_\tau}(\cdot, x_i) = \omega_{\nu_\tau}^j \phi_{\nu_\tau}^j(\cdot)$, where j is such that $\phi_{\nu_\tau}^j(x_i) = 1$. From this the lemma follows. $\qquad\square$

Lemma G.7. *Let* $A, B \in \mathcal{B}(\mathcal{H}, \mathcal{H})$ *be two PSD operators such that* $\xi B - A$ *and* $\xi A - B$ *are PSD for some* $\xi \in (1, \infty)$. *Let* $g, h \in \mathcal{H}$ *two arbitrary vectors and* $\lambda \in \mathbb{R}_{++}$ *be a constant. Then:*

$$\langle A[g] + \lambda\xi h, B[g] + \lambda h \rangle_{\mathcal{H}} \geq \frac{1}{2}\left(\xi^{-1} \|A[g] + \lambda\xi h\|_{\mathcal{H}}^2 - \xi(\xi^2 - 1)\lambda^2 \|h\|_{\mathcal{H}}^2\right)$$

Proof. Consider the following equity:

$$\xi \langle \xi^{-1} A[g] + \lambda h, B[g] + \lambda h \rangle_{\mathcal{H}} = \frac{\xi}{2}\left(\|\xi^{-1} A[g] + \lambda h\|_{\mathcal{H}}^2 + \|B[g] + \lambda h\|_{\mathcal{H}}^2 - \|(B - \xi^{-1} A)[g]\|_{\mathcal{H}}^2\right)$$

which is basically classical decomposition of the dot product $\langle x, y \rangle = \frac{1}{2}\left(\|x\|^2 + \|y\|^2 - \|x - y\|^2\right)$. Then we note that $(1 - \xi^{-2})B - (B - \xi^{-1} A) = \xi^{-2}(\xi A - B)$ is PSD by assumption which implies:

$$\langle A[g] + \lambda\xi h, B[g] + \lambda h \rangle_{\mathcal{H}} \geq \frac{\xi}{2}\left(\|\xi^{-1} A[g] + \lambda h\|_{\mathcal{H}}^2 + \|B[g] + \lambda h\|_{\mathcal{H}}^2 - (1 - \xi^{-2})^2 \|B[g]\|_{\mathcal{H}}^2\right)$$

Finally, noting that $\frac{\xi^2}{2 - \xi^{-2}} - 1 = \frac{\xi^2(1 - \xi^{-2})^2}{2 - \xi^{-2}} \leq \xi^2 - 1$ the result directly follows from equity:

$$\|B[g] + \lambda h\|_{\mathcal{H}}^2 - (1 - \xi^{-2})^2 \|B[g]\|_{\mathcal{H}}^2 =$$

$$\left\|\xi^{-1} \sqrt{2 - \xi^{-2}} B[g] + \frac{\lambda}{\xi^{-1}\sqrt{2 - \xi^{-2}}} h\right\|_{\mathcal{H}}^2 - \lambda^2 \left(\frac{\xi^2}{2 - \xi^{-2}} - 1\right) \|h\|_{\mathcal{H}}^2$$

$\qquad\square$

Denote $\kappa(A, B) = \|\mathrm{Id}_{\mathcal{H}} - BA^{-1}\|_{\mathcal{B}(\mathcal{H},\mathcal{H})} = \|(B - A)A^{-1}\|_{\mathcal{B}(\mathcal{H},\mathcal{H})}$.

Lemma G.8. *If $\xi^{-1}\mathcal{K} \preceq \mathcal{K}_f \preceq \xi\mathcal{K}$ for $\xi > 1$ then $\kappa(\mathcal{K}, \mathcal{K}_f) \leq \xi - 1$.*

Proof. First we note that both operators are symmetric semi-positive definite in L_2. Now let us look at Rayleigh quotient:

$$\|(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-1}\|_{\mathcal{B}(\mathcal{H},\mathcal{H})} = \max_{f \in \mathcal{F}\backslash\{\mathbb{0}\}} \frac{\|(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-1}f\|_{\mathcal{H}}}{\|f\|_{\mathcal{H}}} = \max_{f \in \mathcal{F}\backslash\{\mathbb{0}\}} \frac{\|\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}}f\|_{L_2}}{\|f\|_{L_2}}.$$

In the last equality we used fact that $\mathcal{K}$ is symmetric positive definite and therefore $\mathcal{K}^{\frac{1}{2}}$ too and hence we can substitute $f \leftarrow \mathcal{K}^{\frac{1}{2}}f$ and use explicit formula for the dot product in $\mathcal{H}$ via product in L_2. Now we observe that

$$-(\xi - 1)\mathrm{Id}_{L_2} = \mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \xi\mathcal{K})\mathcal{K}^{-\frac{1}{2}} \preceq \mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}} \preceq$$
$$\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \xi^{-1}\mathcal{K})\mathcal{K}^{-\frac{1}{2}} = (1 - \xi^{-1})\mathrm{Id}_{L_2} \preceq (\xi - 1)\mathrm{Id}_{L_2}.$$

which implies that spectral radius $\rho(\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}})$ is bounded by $\xi - 1$. Therefore we obtain:

$$\max_{f \in \mathcal{F}\backslash\{\mathbb{0}\}} \frac{\|\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}}f\|_{L_2}}{\|f\|_{L_2}} = \max_{f \in \mathcal{F}\backslash\{\mathbb{0}\}} \frac{\|\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}}f\|_{L_2}}{\|f\|_{L_2}} \leq$$
$$\rho(\mathcal{K}^{-\frac{1}{2}}(\mathcal{K} - \mathcal{K}_f)\mathcal{K}^{-\frac{1}{2}}) \leq \xi - 1.$$
$\square$

Lemma G.9. *Iterations of gradient boosting can be represented as:*

$$\epsilon^{-1}\big(f_\tau - f_{\tau+1}\big) = \mathcal{K}_{f_\tau}D[f_\tau - f_*] + \frac{\lambda}{N}f_\tau.$$

where $D : \mathcal{F} \to \mathcal{F}$ is bounded and universal for f_τ, i.e. $\Sigma_{f_\tau} = \mathcal{K}_{f_\tau}D\forall\tau$. Similar decomposition holds for $\nabla_{\mathcal{H}}V(f, \lambda) = \mathcal{K}D[f - f_*] + \frac{\lambda}{N}f$.

Proof. First observe that iterations are basically $\nabla_{\mathcal{H}(\mathcal{K}_\tau)}V(f_\tau, \lambda)$ assuming that we replace $\mathcal{H}$ with $\mathcal{H}(\mathcal{K}_{f_\tau})$ in regularizer term. Therefore, we need to prove only $\nabla_{\mathcal{H}}V(f, \lambda) = \mathcal{K}D[f - f_*] + \frac{\lambda}{N}f$. To prove this observe that there is defined linear Frechet differential $\mathcal{D}_f V(f)[h] = \frac{1}{N}\sum_{i=1}^{N}\big(f(x_i) - f_*(x_i)\big)h(x_i) : \mathcal{F} \to \mathbb{R}$ which is bilinear functional by $f - f_*$ and h. Since $\mathcal{F}$ is finite dimensional then there is linear representation of the mapping in L_2 product restricted on $\mathcal{F}$, let us denote it $D : \mathcal{F} \to \mathcal{F}$. Now we have $\mathcal{D}_f V(f)[h] = \langle D[f - f_*], h\rangle_{L_2}$. Now we transform it to $\mathcal{D}_f V(f)[h] = \langle \mathcal{K}^{-1}\mathcal{K}D[f - f_*], h\rangle_{L_2} = \langle \mathcal{K}D[f - f_*], h\rangle_{\mathcal{H}}$ which implies that $\nabla_{\mathcal{H}}V(f) = \mathcal{K}D[f - f_*]$. $\square$

Lemma G.10. *Consider setting of the previous lemma with requirement $B - A$ being SPD being lifted. Then the inequality holds:*

$$\langle A[g] + \lambda h, B[g] + \lambda h\rangle_{\mathcal{H}} \geq \big(\frac{1}{2} - \kappa(A, B)\big)\big\|A[g] + \lambda h\big\|_{\mathcal{H}}^2 - \kappa^2(A, B)\lambda^2\|h\|_{\mathcal{H}}^2.$$

Proof. Consider rewriting the left part as

$$\langle A[g]+\lambda h, B[g]+\lambda h\rangle_{\mathcal{H}} = \langle A[g]+\lambda h, \big(\mathrm{Id}_{L_2}+(BA^{-1}-\mathrm{Id}_{L_2})\big)(A[g]+\lambda h)+\lambda(\mathrm{Id}_{L_2}-BA^{-1})h\rangle_{\mathcal{H}} \geq$$
$$\geq \|A[g]+\lambda h\|_{\mathcal{H}}^2 - \langle A[g]+\lambda h, (\mathrm{Id}_{L_2}-BA^{-1})(A[g]+\lambda h)\rangle_{\mathcal{H}} + \langle A[g]+\lambda h, \lambda(\mathrm{Id}_{L_2}-BA^{-1})h\rangle_{\mathcal{H}} \geq$$
$$\geq \big(\frac{3}{2}-\kappa(A, B)\big)\|A[g]+\lambda h\|_{\mathcal{H}}^2 + \frac{\lambda^2}{2}\|(\mathrm{Id}_{L_2}-BA^{-1})h\|_{\mathcal{H}}^2 - \frac{1}{2}\|(A[g]+\lambda h)-\lambda(\mathrm{Id}_{L_2}-BA^{-1})h\|_{\mathcal{H}}^2 \geq$$
$$\geq \big(\frac{3}{2} - \kappa(A, B)\big)\|A[g] + \lambda h\|_{\mathcal{H}}^2 - \|(A[g] + \lambda h)\|_{\mathcal{H}}^2 - \lambda^2\|(\mathrm{Id}_{L_2} - BA^{-1})h\|_{\mathcal{H}}^2 \geq$$
$$\geq \big(\frac{1}{2} - \kappa(A, B)\big)\big\|A[g] + \lambda h\big\|_{\mathcal{H}}^2 - \kappa^2(A, B)\lambda^2\|h\|_{\mathcal{H}}^2$$

$\square$

The following lemma holds.

Lemma G.11. *If $(\frac{\lambda}{N} + 1)\epsilon < 1$ and $f_0 = \mathbb{0}_{\mathcal{H}}$ then $\forall \tau$ the following holds almost surely*

$$\|f_\tau - f_*\| \leq \|f_*\|$$

for norms $\|\cdot\|_{L_2}$, $\|\cdot\|_{\mathcal{H}}$ and $\|\cdot\|_{\mathbb{R}^N}$.

Proof. Consider writing

$$f_{\tau+1} - f_* = \left(\mathrm{Id}_{L_2} - \frac{\lambda\epsilon}{N}\mathrm{Id}_{L_2} - \epsilon\Sigma_{\nu_\tau}\right)[f_\tau - f_*] - \frac{\lambda\epsilon}{N}f_* .$$

Now observe that $S := \left(\mathrm{Id}_{L_2} - \frac{\lambda\epsilon}{N}\mathrm{Id}_{L_2} - \epsilon\Sigma_{\nu_\tau}\right)$ is symmetric with eigenvalues $0 < \lambda_i(S) \leq 1 - \frac{\lambda\epsilon}{N}$ therefore its operator norm in $\mathcal{B}(L_2)$, $\mathcal{B}(\mathcal{H})$ and $\mathbb{R}^{N\times N}$ is lesser than $1 - \frac{\lambda\epsilon}{N}$. Taking norm from left and right side and using sub-additivity of norm we obtain:

$$\|f_{\tau+1} - f_*\| \leq (1 - \lambda\epsilon)\|f_\tau - f_*\| + \lambda\epsilon\|f_*\| .$$

Since $\|f_0 - f_*\| = \|f_*\|$ that recurrent relation inductively yields statement of the lemma. $\quad\square$

Corollary G.12. *Under the same conditions, $\|f_\tau\| \leq 2\|f_*\|$ and $V(f_\tau, 0) \leq R^2$ almost surely.*

G.2 MAIN THEOREM

Let us denote $R_1 = \|f_*\|_{\mathcal{H}}$. We argue that it is constant value since kernel $\mathcal{H}$ and f_* are convergent as $N \to \infty$ which makes it bounded by some constant with probability arbitrarily close to one. Moreover, denote $M_\beta = e^{\frac{2mR^2}{\beta}} > 1$.

Theorem G.13. *Consider an arbitrary ϵ, $0 < \epsilon(\frac{\lambda}{N} + 1) < 1$ and $\frac{(1+M_\beta\lambda)}{4M_\beta N} \geq \epsilon$. The following inequality holds:*

$$\mathbb{E}V(f_T) \leq R^2 e^{-\frac{1+M_\beta\lambda}{2M_\beta N}T\epsilon} + \frac{4\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(4M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(4M_\beta(M_\beta^2-1)+1\right)\right)R_1^2$$

Proof. First we deal with the term $V(f_{\tau+1})$:

$$\frac{1}{2N}\|f_{\tau+1}(\mathbf{x}_N) - f_*(\mathbf{x}_N)\|_{\mathbb{R}^N}^2 = V(f_\tau) + \mathcal{D}_f V(f_\tau)[f_{\tau+1} - f_\tau] + \frac{1}{2}\mathcal{D}_f^2 V(f_\tau)[f_{\tau+1} - f_\tau, f_{\tau+1} - f_\tau] \leq$$

$$\leq V(f_\tau) - \epsilon\langle\Sigma_{\nu_\tau}[f_\tau - f_*] + \frac{\lambda}{N}f_\tau, \mathcal{K}D[f_\tau - f_*]\rangle_{\mathcal{H}} + 2\epsilon^2 V(f_\tau) + 2\frac{\lambda^2\epsilon^2}{2N^3}\|f_\tau(\mathbf{x}_N)\|_{\mathbb{R}^N}^2.$$

Then with regularization:

$$\frac{\lambda}{2N}M_\beta\|f_{\tau+1}\|_{\mathcal{H}}^2$$

$$\leq \frac{\lambda}{2N}M_\beta\|f_\tau\|_{\mathcal{H}}^2 - \epsilon\langle\frac{\lambda}{N}M_\beta f_\tau, \Sigma_{\nu_\tau}[f_\tau - f_*] + \frac{\lambda}{N}f_\tau\rangle_{\mathcal{H}} + \frac{\epsilon^2\lambda}{N}M_\beta\|f_\tau - f_*\|_{\mathcal{H}}^2 + \frac{\epsilon^2\lambda^3}{N^3}M_\beta\|f_\tau\|_{\mathcal{H}}^2$$

$$\leq \frac{\lambda}{2N}M_\beta\|f_\tau\|_{\mathcal{H}}^2 - \epsilon\langle\frac{\lambda}{N}M_\beta f_\tau, \Sigma_{\nu_\tau}[f_\tau - f_*] + \frac{\lambda}{N}f_\tau\rangle_{\mathcal{H}} + \frac{\epsilon^2\lambda}{N}M_\beta\|f_\tau\|_{\mathcal{H}}^2 + \frac{\epsilon^2\lambda}{2N}M_\beta(3 + \frac{4\lambda^2}{N^2})\|f_*\|_{\mathcal{H}}^2.$$

Summing up those two expressions, applying Lemma G.7 and noting that Σ is non-degenerate if we restrict on its image (which we can since f_τ lies there) by using Polyak-Łojasiewicz inequality $\frac{1}{2}\|\nabla V\|_{\mathcal{H}} \geq \mu V$ for μ-strongly convex function V (restricted on image of Σ) with $\mu \geq \frac{1+M_\beta\lambda}{N} > 0$ we obtain the following bound:

$$\mathbb{E}V(f_{\tau+1}, M_\beta\lambda)$$

$$\leq \left(1 - \epsilon(\frac{(1+\lambda)}{M_\beta N} - 2\epsilon)\right)V(f_\tau) + \frac{2\lambda^2\epsilon^2}{N^3}R^2 + \frac{\epsilon\lambda M_\beta}{N}\left(\left(\epsilon(2 + \frac{2\lambda^2}{N^2}) + (M_\beta^2 - 1)\frac{\lambda}{N}\right)\right)R_1^2 \leq$$

$$\left(1 - \epsilon(\frac{(1+M_\beta\lambda)}{2M_\beta N})\right)V(f_\tau) + 2\frac{\lambda^2\epsilon^2}{N^3}R^2 + \frac{\epsilon\lambda M_\beta}{N}\left(\left(\epsilon(2 + \frac{2\lambda^2}{N^2}) + (M_\beta^2 - 1)\frac{\lambda}{N}\right)\right)R_1^2.$$

Which yields

$$\mathbb{E}V(f_T, M_\beta\lambda) \leq R^2 e^{-\frac{1+M_\beta\lambda}{2M_\beta N}T\epsilon} + \frac{4\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(2M_\beta^2\epsilon(2+\frac{2\lambda^2}{N^2}) + 2M_\beta^2(M_\beta^2 - 1)\frac{\lambda}{N}\right)R_1^2.$$

Which finally translates into bound

$$\mathbb{E}V(f_T) \leq R^2 e^{-\frac{1+M_\beta\lambda}{2M_\beta N}T\epsilon} + \frac{4\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(4M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(4M_\beta(M_\beta^2-1)+1\right)\right)R_1^2.$$

$\square$

Theorem G.14. *(Theorem 4.1 in the main text) Let* $C = \frac{8\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(8M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(8M_\beta(M_\beta^2 - 1) + 3\right)\right)R_1^2$. *Assume that* $0 < \epsilon(\frac{\lambda}{N} + 1) < 1$ *and* $\frac{(1+M_\beta\lambda)}{4M_\beta N} \geq \epsilon$, $\frac{(1+\lambda)}{8N} \geq \epsilon$, $\frac{N+\lambda}{N}\frac{2mM_\beta C}{\beta} < \frac{1+\lambda}{2N}$ *and define* $T_1 = \left[\frac{2M_\beta N}{(1+M_\beta\lambda))\epsilon}\log\frac{R^2}{C}\right] + 1$ *then* $\forall T \geq T_1$ *it holds that*

$$V(f_T, \lambda) \leq Ce^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)N^2}R^2 + \frac{8}{1+\lambda}\left(\epsilon(1+\frac{\lambda}{N}) + 4M_\beta^2\frac{m\lambda C}{\beta N}\right)R_1^2$$

Proof. If we take $T_1 = \left[\frac{2M_\beta N}{1+M_\beta\lambda}\log\left(\frac{4\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2} + \left(4M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(8M_\beta(M_\beta^2 - 1) + 3\right)\right)\frac{R_1^2}{R^2}\right)^{-1}\right] + 1$ then the following inequality holds $\forall\tau \geq T_1$:

$$\mathbb{E}V(f_\tau) + \frac{\lambda}{2N}\|f_\tau\|_\mathcal{H}^2 \leq \frac{8\lambda^2\epsilon M_\beta}{(1+M_\beta\lambda)N^2}R^2 + \left(8M_\beta^2\epsilon(1+\frac{\lambda^2}{N^2}) + \frac{M_\beta\lambda}{2N}\left(8M_\beta(M_\beta^2 - 1) + 3\right)\right)R_1^2.$$

Overall, the proof is similar to the previous theorem but with modifications, instead of Lemma G.7 we will use Lemma G.10. The only problem would arise in multiplier $\mathbb{E}\left(\frac{1}{2} - e^{\frac{2m(V(f_\tau)+\frac{\lambda}{2N}\|f_\tau\|_\mathcal{H}^2)}{\beta}}\right)\|\nabla_\mathcal{H}V(f_\tau, \lambda)\|_\mathcal{H}^2$. We can bound

$$\mathbb{E}\left(\frac{1}{2} - e^{\frac{2m(V(f_\tau)+\frac{\lambda}{2N}\|f_\tau\|_\mathcal{H}^2)}{\beta}}\right)\|\nabla_\mathcal{H}V(f_\tau, \lambda)\|_\mathcal{H}^2$$

$$\geq \left(\frac{1+\lambda}{N} - \frac{N+\lambda}{N}\frac{2mM_\beta C}{\beta}\right)\mathbb{E}V(f_\tau, \lambda) \geq \frac{1+\lambda}{2N}\mathbb{E}V(f_\tau, \lambda).$$

After using $(e^z - 1)^2 \leq e^{2z} - 1 \leq 2M_\beta^2 z$ for $z = \frac{2mV(f)}{\beta}$ satisfying $e^z \leq M_\beta$ we obtain the bound:

$$\mathbb{E}V(f_{\tau+1}, \lambda) \leq (1 - \frac{1+\lambda}{4N}\epsilon)\mathbb{E}V(f_\tau, \lambda) + 2\frac{\lambda^2\epsilon^2}{N^2}R^2 + \frac{\epsilon\lambda}{N}\left(2\epsilon(1+\frac{\lambda}{N}) + 8M_\beta^2\frac{m\lambda C}{\beta N}\right)R_1^2.$$

Iterating the bound yields

$$\mathbb{E}V(f_T, \lambda) = \mathbb{E}V(f_{T_1}, \lambda)e^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)N^2}R^2 + \frac{8}{1+\lambda}\left(\epsilon(1+\frac{\lambda}{N}) + 4M_\beta^2\frac{m\lambda C}{\beta N}\right)R_1^2$$

$$\leq Ce^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)N^2}R^2 + \frac{8}{1+\lambda}\left(\epsilon(1+\frac{\lambda}{N}) + 4M_\beta^2\frac{m\lambda C}{\beta N}\right)R_1^2.$$

$\square$

Corollary G.15. *(Convergence to the solution of the KRR / Convergence to the Gaussian Process posterior mean function). Under assumptions of both previous theorems we have that:*

$$\mathbb{E}\|f_T - f_*^\lambda\|_{L_2}^2 \leq \frac{NC}{1+\lambda}e^{-\frac{1+\lambda}{4N}\epsilon(T-T_1)} + 2\frac{\lambda^2\epsilon}{(1+\lambda)^2 N}R^2 + \frac{8}{(1+\lambda)^2}\left(N\epsilon(1+\frac{\lambda}{N}) + 4M_\beta^2\frac{m\lambda C}{\beta}\right)R_1^2.$$

Lemma G.16 (Lemma 4.3 in the main text). *The following convergence holds almost surely in* $x \in X$:

$$h_T(\cdot) \xrightarrow[T\to\infty]{} \mathcal{GP}\left(\mathbb{0}_{L_2(\rho)}, \mathcal{K}\right).$$

Proof. From (5), we have that the covariance of h_T is $\mathcal{K}$ independently from T. Thus, it remains to show that the limit is Gaussian almost surely which essentially holds due to the central limit theorem almost surely in $x \in X$:

$$h_T(x) = \frac{1}{\sqrt{T}} \sum_{i=1}^{T} h_{T,i}(x) \to \mathcal{N}\big(\mathbb{0}_{L_2}, \mathcal{K}(x,x)\big),$$

where each individual tree $h_{T,i}$ is centered i.i.d. (with the same distribution as h_1). $\qquad\square$

H IMPLEMENTATION DETAILS

In the experiments, we fix $\sigma = 10^{-2}$ (scale of the kernel) and $\delta = 10^{-4}$ (scale of noise), which theoretically can be taken arbitrarily. As a hyperparameter (that is estimated on the validation set), we consider $\beta \in \{10^{-2}, 10^{-1}, 1\}$. We use the standard CatBoost library and add the Gumbel noise term in selecting the trees for the "L2" scoring function, which is implemented in CatBoost out of the box but is not used by SGB and SGLB since it is not the default one for the library. Moreover, we do not consider subsampling of the data (as SGLB does also), and differently from SGB and SGLB, we disable the "boost-from-average" option. Finally, we set $l2-leaf-reg$ value to 0, as SGLB does.