The Lazy Online Subgradient Algorithm is Universal on Strongly Convex Domains

Abstract

We study the Lazy Subgradient algorithm for online linear optimisation on a strongly convex domain. It is known that Subgradient achieves $O(\sqrt{N})$ regret against adversarial opponents; here we show it is universal in the sense that it also achieves $O(\log N)$ regret against i.i.d opponents. This improves upon the more complex meta-algorithm of Huang et al [20] that only gets $O(\sqrt{N \log N})$ and $O(\log N)$ bounds. In addition we show that, unlike for the simplex, order bounds for pseudo-regret and expected regret are equivalent for strongly convex domains.

Introduction

Online linear optimisation is a repeated game with one opponent. On turn n we know the *cost vectors* $a_1, a_2, \ldots, a_{n-1} \in \mathbb{R}^d$ and select an *action* x_n from the *domain* $X \subset \mathbb{R}^d$. The opponent observes x_n and chooses the next cost vector $a_n \in \mathbb{R}^d$, and we pay cost $a_n \cdot x_n$. Our goal is to select actions to make the *regret* $\sum_{i=1}^N a_i \cdot (x_i - x^*)$ small for $x^* \in \operatorname{argmin}\{\sum_{i=1}^N a_i \cdot x : x \in X\}$.

If the cost vectors are selected by an adversarial opponent then $a_1, a_2, \ldots, a_{n-1}$ give no information about a_n and selecting a good action seems like an impossible task. Despite this, there are algorithms that get $O(\sqrt{N})$ regret against such opponents. That means the loss in performance against x^* is $O(1/\sqrt{N})$ per turn which vanishes as $N \to \infty$. The central algorithms in the field are Hedge [18, 34]; Prod [9]; and Subgradient, which has Greedy [41] and Lazy [34] variants. All three algorithms get $O(\sqrt{N})$ regret. The first two are specialised to X the simplex. The latter works for any compact convex domain. The bulk of online optimisation literature is refinements of the central algorithms.

In many cases however the cost vectors are non adversarial. For example they might be determined by fluctuations in traffic, the weather or the stock market over a short period of time. The standard model of *easy* opponents is cost vectors drawn independently from some fixed distribution. In the easy setting the history $a_1, \ldots, a_n$ provides information about the next cost vector. Indeed $\widetilde{a}_{n+1} = \frac{a_1 + \ldots + a_n}{n}$ is an increasingly reliable sequence of estimates for $a = \mathbb{E}[a_{n+1}]$ and it seems reasonable to play x_{n+1} to minimise $\widetilde{a}_{n+1} \cdot x$. This is called Follow-the-Leader (FTL). For example if X is the simplex FTL gives pseudo-regret $\mathbb{E}\left[\sum_{i=1}^N a \cdot (x_i - y^*)\right] \le O(1)$ for $y^* \in \operatorname{argmin}\{a \cdot x : x \in X\}$. Note this is weaker than bounding the expected regret $\mathbb{E}\left[\sum_{i=1}^N a_i \cdot (x_i - x^*)\right]$. On the other hand FTL gives no guarantee against adversarial opponents, and we might not know in advance whether the opponent is i.i.d or adversarial. Hence we would like an algorithm that is *universal* in that it always gets $O(\sqrt{N})$ regret but specialises to a much stronger bound against i.i.d opponents.

Recently Huang et al [20] solved this problem for smooth, strongly convex domains. For these domains they showed FTL gives $O(\log N)$ regret if $\|\sum_{i=1}^n a_i\|$ grows linearly.[1] The result was generalised to nonsmooth domains by [22, 26]. Huang et al then combine FTL with the (A,B)-Prod

---

[1] Without smoothness they show the growth condition occurs with high probability against i.i.d opponents.

Algorithm 1: Online Lazy Subgradient Algorithm

Data: Cost vectors $a_1, a_2, \ldots \in \mathbb{R}^d$. Parameter $\eta > 0$. Domain $X \subset \mathbb{R}^d$.

1 for $n = 0, 1, \ldots$ **do**

2 $\quad\Big|\quad x_{n+1} = P_X\left(-\frac{\eta}{\sqrt{n}} \sum_{i=1}^n a_i\right)$

3 $\quad\Big|\quad$ Receive a_{n+1} and pay cost $a_{n+1} \cdot x_{n+1}$

---

34 meta-algorithm of [31] to ensure $O(\sqrt{N \log N})$ regret against adversarial opponents. In this work
35 we improve and extend their results. In particular we show the meta-algorithm is unnecessary as the
36 much simpler Lazy Subgradient algorithm already achieves $O(\sqrt{N})$ and $O(\log N)$ regret bounds.

37 **Terminology and Notation**

38 Throughout $\|\cdot\|$ is the Euclidean norm. Given a vector subspace $V \subset \mathbb{R}^d$ and linear operator
39 $A : V \to V$ write the operator norm as $\|A\| = \max\{\|Av\| : v \in V, \|v\| = 1\}$. It is known (see [10]
40 Theorems 4.1 and 4.2) that $\|A\| = \max_j \sqrt{|\lambda_j|}$ for λ_j the eigenvalues of $A^T A$. Moreover if A is
41 symmetric we have $\|A\| = \max_j |\mu_j|$ for μ_j the eigenvalues of A. The set Z is called λ-strongly
42 convex to mean for each $x, y \in Z$ and $\alpha \in [0, 1]$ the ball $B(\alpha x + (1 - \alpha)y, r)$ is contained in Z for
43 $r = \frac{\lambda}{2}\alpha(1 - \alpha)\|x - y\|^2$. The pseudo-regret $\mathbb{P}[R_N]$ and expected regret $\mathbb{E}[R_N]$ are defined:

$$\mathbb{P}[R_N] = \mathbb{E}\left[\sum_{i=1}^N a \cdot x_i - \min_{x \in X} \sum_{i=1}^N a \cdot x\right] \qquad \mathbb{E}[R_N] = \mathbb{E}\left[\sum_{i=1}^N a_i \cdot x_i - \min_{x \in X} \sum_{i=1}^N a_i \cdot x\right] \qquad (1)$$

44 The Jensen Inequality implies $\mathbb{P}[R_N] \leq \mathbb{E}[R_N]$. We write P_X for the Euclidean projection onto
45 the domain X. The Lazy Subgradient algorithm is given as Algorithm 1. Note this is distinct
46 from the so-called Greedy Subgradient algorithm with starting point $x_1 = 0$ and recursive update
47 $x_{n+1} = P_X(x_n - \eta a_n/\sqrt{n})$. Section 5 Figure 2(b) suggests Greedy Subgradient is not universal.

48 **Summary of Contributions**

49 Throughout we make the following assumptions on the domain and cost vectors. Both assumptions
50 are nontrivial. For example Assumption 1 holds on the Euclidean and ℓ_p balls for $p \in (1, 2]$ but not
51 for the simplex or polytopes. Assumption 2 on the cost vectors is a typical model of an *easy opponent*
52 and fails if the opponent is intelligent and adapts to our past moves.

53 **Assumption 1.** The domain $X \subset \mathbb{R}^d$ is compact, m-strongly convex and contains the origin. Write
54 $D = \max\{\|x\| : x, y \in X\}$. The boundary $\mathcal{M} = \partial X$ is a $(d-1)$-dimensional C^2 manifold. Namely
55 each $z \in \mathcal{M}$ has a neighborhood U in $\mathbb{R}^d$ and C^2 function $F : U \to \mathbb{R}$ with nonzero gradient such that
56 $\mathcal{M} \cap U = \{x \in U : F(x) = 0\}$. Such a function is called a **coordinate patch** at z.

57 **Assumption 2.** The cost vectors $a_1, a_2, \ldots$ are i.i.d with expectation $\mathbb{E}[a_n] = a$. For all n we have
58 $\|a_n\| \leq L$ and $\|a_n - a\| \leq R$ and $\mathbb{E}\|a_n - a\| \leq \delta$. Let $x^* \in \operatorname{argmin}\{a \cdot x : x \in X\}$ be the expected
59 minimiser. Since we also assume X is strongly convex x^* is unique.

60 Our main result is Theorem 3 which says Lazy Subgradient achieves $O\left(\frac{L^2}{m\|a\|} \log N\right)$ expected-regret
61 under Assumptions 1 and 2. This bound is essentially tight by [20] Theorem 9. The improved
62 behaviour against i.i.d opponents is remarkable since the algorithm was designed with the adversarial
63 case in mind, and predates the recent interest in universal algorithms. Our second contribution is
64 Theorem 2 which states the gap between expected regret and pseudo-regret is $O\left(\frac{L^2}{m\|a\|}\right)$. Theorem 2
65 is used to prove Theorem 3 but is of independent interest, since it fails if the domain is the simplex.

66 From a technical standpoint, our Subgradient analysis goes beyond that of [20, 22, 26] for FTL. Their
67 central idea is that the normal vector to $\mathcal{M}$ at $x_n \in \operatorname{argmin}\{(a_1 + \ldots + a_{n-1}) \cdot x : x \in X\}$ points
68 along $A_{n-1} = -\frac{a_1 + \ldots + a_{n-1}}{n}$. Using $\|A_n - A_{n-1}\| = O(1/n)$ and how strong convexity says the
69 unit normal changes quickly as we vary the basepoint, they prove $\|\Theta_n - \Theta_{n+1}\| = O(1/n)$ for
70 the normal vectors Θ_n, Θ_{n+1} at x_n, x_{n+1}. Summing the resulting series gives a logarithmic regret
71 bound. Unfortunately this method is insufficient to analyse Subgradient instead of FTL, as it only
72 gives $\|\Theta_n - \Theta_{n+1}\| = O(1/\sqrt{n})$ which, when plugged into the second half of the existing proofs,
73 gives only an $O(\sqrt{N})$ regret bound. Thus we follow a new approach where our Theorem 1 first

establishes a high probability bound $\|x_n - x^*\| = O(1/\sqrt{n})$ for the distance from the expected minimiser. For comparison existing works do not mention the expected minimiser at all. Then we use strong convexity (Lemma 1) to see the surface locally looks like a quadratic with axis pointing along a. Hence only $O(1/n)$ of the displacement is in the a-direction; and the larger perpendicular component does not contribute to pseudo-regret. Summing the series gives an $O(\log N)$ bound for pseudo-regret. To convert this into a bound for expected regret we use our Theorem 2 which says the gap between expected regret and pseudo-regret is finite.

Our differential geometry methods are more nuanced than existing works. In particular we must consider both the magnitude and direction of $x_n - x^*$. For comparison existing works consider only the magnitude $\|x_n - x_{n+1}\|$. The relevant object is $N(z)v$ in Lemmas 1-5 and Proposition 1 which captures the rate of change of the unit normal as we move the basepoint in the v-direction.

Related Work

The first strand of related work is *universal algorithms*. That is to say online algorithms that get $\widetilde{O}(\sqrt{N})$ regret against adversarial opponents but specialise to get much stronger bounds against i.i.d opponents. The original research focus was on the simplex and bandit feedback [4, 8, 32, 33, 37, 40]. In the full information setting the FlipFlop algorithm [11] interpolates between FTL and Hedge on the simplex to get $O(1)$ and $O(\sqrt{N \log N})$ regret against i.i.d and adversarial opponents respectively. The (A,B)-Prod meta-algorithm [31] can be used for something similar in much greater generality. Another result comes from [16]. They design a variant of the Prod algorithm [9] with different learning rates for each arm. As a corollary they get $O(1)$ expected regret against i.i.d opponents with a unique best arm. Apart from [31] all the above algorithms apply only to the simplex.

However recent results suggest that such complicated algorithms are unnecessary. In the full information setting it was recently proved that two of the central algorithms do much better against i.i.d opponents. Hedge gets $O(\log(d)L_\infty^2/\Delta)$ pseudo-regret on the simplex [27]; and Lazy Subgradient gets $O(L_2^2/\Delta)$ pseudo-regret on polytopes[2] [1, 2].The latter is notable since it works for domains other than the simplex. Here d is the dimension; Δ the suboptimality gap; and we assume $\|a_n\| \leq L_2$ and $\|a_n\|_\infty \leq L_\infty$ where appropriate. The above bounds are in addition to the $O(\sqrt{N})$ regret bounds of Subgradient and Hedge (see [34] Chapter 2) against adversarial opponents.

The second strand of related work is the study of online algorithms on strongly convex domains. Typically these are unit balls with respect to some norm. For example [39] consider the ℓ_p unit ball with $p \in (1, 2)$ in the full-information setting. Another setting of interest [5, 12, 19] is when predictions are available for the direction of the next cost vector. In the bandit setting [7] consider ℓ^p balls; [6] consider the Euclidean ball; and [30] consider general strongly convex sets. They allow infinitely many arms but require the cost vectors be stochastically generated. In the offline setting [17] show the Frank-Wolfe method has $O(1/n^2)$ convergence rates − and hence finite regret − when the domain is strongly convex. [23] prove similar results for more general so-called uniformly convex sets. Under additional conditions the older works [13, 14, 25] show an exponential convergence rate.

Bound for Pseudo-Regret

Here we show Lazy Subgradient gets $O(\log N)$ pseudo-regret against i.i.d cost vectors on a C^2 strongly convex domain. This is in addition the $O(\sqrt{N})$ regret bound for adversarial cost vectors.

Theorem 1. Under Assumptions 1 and 2, Algorithm 1 with parameter $\eta > 0$ achieves pseudo-regret

$$\mathbb{P}[R_N] \leq \frac{144}{m\|a\|} \left(\frac{\sqrt{8\pi}DR}{\eta} + 4R^2 \right) \log N + O(1).$$

In particular for stepsize $\eta = \Theta(D/L)$ we have $\mathbb{P}[R_N] \leq O\left(\frac{L^2}{m\|a\|} \log N \right)$.

Observe by [34] Theorem 2.4 the stepsize $\eta = \Theta(D/L)$ is optimal for both the adversarial and i.i.d bound. The example to keep in mind through our analysis is $X = \{(\alpha, \beta) \in \mathbb{R}^2 : \beta \geq \alpha^2, y \leq 1\}$ the

---

[2]Huang et al [20] already proved an $O(1)$ bound for polytopes with unique optimal vertex. However dependence on dimension and suboptimality gap is not obvious.

epigraph of a quadradic function. Write the actions $x_n = (\alpha_n, \beta_n)$. For expected cost $a = (0,1)$ the minimiser is $x^* = (0,0)$. The proof strategy is to show $\left\| \sum_{i=1}^n a_i \right\| = \Omega(n)$ and $\left\| \sum_{i=1}^n (a_i - a) \right\| = O(\sqrt{n})$ with high probability. Hence the line from $y_n = -\frac{\eta}{\sqrt{n}} \sum_{i=1}^n a_i$ to any $x \in X$ has slope of absolute value $\Omega(\sqrt{n})$. In particular the line from y_n to $x_n = P_X(y_n)$ is normal to the graph. Since the normal at x_n has slope $-2/\alpha_n$ we get $2/\alpha_n = \Omega(\sqrt{n})$ and $\alpha_n = O(1/\sqrt{n})$. Finally write $a \cdot (x_n - x^*) = (0,1) \cdot x_n = \beta_n = \alpha_n^2 = O(1/n)$. Sum over n to get $O(\log N)$ pseudo-regret.

Lemma 1 generalises how in the example we convert an $\Omega(\sqrt{n})$ bound for the slope into an $O(1/\sqrt{n})$ bound for pseudo-regret. In higher dimension the result depends on the direction from x^* to x_n.

Definition 1. Given points $x, y \in \mathcal{M}$ a **geodesic** from x to y is a C^2 path in $\mathcal{M}$ with endpoints x, y with $\|\gamma'(t)\| = 1$ for each t, and such that γ has minimal length among such paths. That length is called the **geodesic distance** $d(x, y)$. The unit vector $v \in \mathbb{R}^d$ is called a **direction** from x to y to mean there is a geodesic $\gamma : [0, d(x,y)] \to \mathcal{M}$ from x to y with $\gamma'(0) = v$.

The generalisation of the height of the graph $y = x^2$ is the normal direction to the surface at the expected minimiser x^*. Henceforth write $N(x)$ for the normal outwards unit vector at $x \in \mathcal{M}$. To represent $N : \mathcal{M} \to \mathbb{R}^d$ locally we can choose a coordinate patch F and write $\nabla N = \frac{\nabla F}{\|\nabla F\|}$. Since F is C^2 the matrix ∇N of partial derivatives exists. In Lemma 1 the object $N(z)(x-z)$ is the component of $x - z$ in the normal direction.

Lemma 1. For each $z \in \mathcal{M}$ there exists a coordinate patch $F : U \to \mathbb{R}$ such that for each $x \in U \cap \mathcal{M}$ and direction v from z to x we have

$$|N(z)(x - z)| \leq \frac{|v^T \nabla^2 F(z) v|}{\|\nabla F(z)\|} d(x, z)^2.$$

Lemma 1 is proved using Lemmas 2-5 and Proposition 1 which are proved in Appendix II. The arguments there use routine differential geometry; but we provide self-contained proofs, since most of the literature focuses on intrinsically defined surfaces, rather than surfaces embedded in $\mathbb{R}^d$.

Lemma 2. For each $z \in \mathcal{M}$ and unit vector v tangent to $\mathcal{M}$ at z we have $\|\nabla N(z)v\| \geq m$.

Lemma 3. For each coordinate patch F at $z \in \mathcal{M}$ and each unit vector tangent v to $\mathcal{M}$ at z we have $\frac{v^T \nabla^2 F(z) v}{\|\nabla F(z)\|} \geq m$.

Lemma 4 states that the unit normal function has bounded second derivative, despite any complications caused by the function being only locally defined.

Lemma 4. There exist constants $M, r > 0$ such that each $z \in \mathcal{M}$ has a coordinate patch $F : B(z, r) \to \mathbb{R}$ with $\|\nabla N(y)\|, \frac{\|\nabla^2 F(y)\|}{\|\nabla F(y)\|} \leq M$ for all $y \in B(z, r)$.

Lemma 5. Let M be the constant from Lemma 4. For each geodesic $\gamma : [0, d(x,y)] \to \mathcal{M}$ from x to y we have for $d = d(x, y)$ and all $t \leq d$ the inequalities

$$(1)\ \|\gamma''(t)\| \leq M \qquad (2)\ \|\gamma(t) - \gamma(0) - t\gamma'(0)\| \leq \frac{M}{2} t^2 \qquad (3)\ \left| \|y - x\| - d \right| \leq \frac{M}{2} d^2$$

Proposition 1. Each $z \in \mathcal{M}$ has a neighborhood U in $\mathbb{R}^d$ such that for all $x \in U \cap \mathcal{M}$ and direction v from z to x we have

$$\|N(x) - N(z)\| \geq \frac{\|\nabla N(z)v\|}{2} d(x, z)$$

Proof of Lemma 1. Let F, M, r be from Lemma 4. Since F is C^2 Taylor's theorem says there is a function $\psi(t)$ defined on some neighborhood of the origin with $\lim_{t \to 0} \psi(t) = 0$ and

$$F(x) = F(z) + \nabla F(z)(x - z) + \frac{(x - z)^T \nabla^2 F(z)(x - z)}{2} + \psi(z - x)\|z - x\|^2$$

Hence there is $r > 0$ such that for all $x \in B(z, r)$ we have $|\psi(z - x)| \leq \frac{m}{4}\|\nabla F(z)\|$ and so

$$\left\| F(x) - F(z) - \nabla F(z)(x - z) - \frac{(x-z)^T \nabla^2 F(z)(x-z)}{2} \right\| \leq \frac{m}{4}\|\nabla F(z)\|\|x - z\|^2.$$

Shrink r if necessary to ensure $B(z, r)$ has geodesic diameter at most $\min\{m/2M^2, \sqrt{2m}/M^{3/2}\}$. By assumption $F(x) = F(z) = 0$. Hence for $d = d(x, z)$ the above gives

$$\left\| \nabla F(z)(x - z) - \frac{(x-z)^T \nabla^2 F(z)(x-z)}{2} \right\| \leq \frac{m}{4}\|\nabla F(z)\|\|x - z\|^2 \leq \frac{m}{4}\|\nabla F(z)\|d^2.$$

153   Let v be a direction from z to x. For $w = (x - z) - dv$. The reverse triangle inequality gives

$$\left\| |\nabla F(z)(x-z)| - \frac{|(dv)^T \nabla^2 F(z)(dv)|}{2} \right\| \le \frac{m}{4}\|\nabla F(z)\|d^2 + |w^T \nabla^2 F(z)(dv)| + \frac{|w^T \nabla^2 F(z)w|}{2}. \quad (2)$$

154   Lemma 5 says $\|w\| \le Md^2/2$ and Lemma 4 says $\frac{\|\nabla^2 F(z)\|}{\|\nabla F(z)\|} \le M$. Hence we get

$$|w^T \nabla^2 F(z)(dv)| \le \|w\|\|\nabla^2 F(z)\|\|dv\| \le \frac{M}{2}\frac{\|\nabla^2 F(z)\|}{\|\nabla F(z)\|}\|\nabla F(z)\|d^3$$
$$\le \frac{M^2}{2}\|\nabla F(z)\|d^3 \le \frac{\|\nabla F(z)\|}{4}md^2.$$

155   where the last inequality uses the assumption $d \le m/2M^2$. Similarly $d \le \sqrt{2m}/M^{3/2}$ gives
156    $|w^T \nabla^2 F(z)w| \le \frac{\|\nabla F(z)\|}{2}md^2$. Hence (2) becomes

$$\left\| |\nabla F(z)(x-z)| - \frac{|v^T \nabla^2 F(z)v|}{2}d^2 \right\| \le \frac{m}{4}\|\nabla F(z)\|d^2 + \frac{m}{4}\|\nabla F(z)\|d^2 = \frac{m}{2}\|\nabla F(z)\|d^2.$$

157   and so $|\nabla F(z)(x - z)| \le \left(\frac{|v^T \nabla^2 F(z)v|}{2} + \frac{m}{2}\|\nabla F(z)\| \right) d^2$. Since the unit normal is
158    $N(z) = \frac{\nabla F(z)}{\|\nabla F(z)\|}$ we can divide both sides to get $N(z)(x-z)| \le \frac{1}{2}\left(\frac{|v^T \nabla^2 F(z)v|}{\|\nabla F(z)\|} + m \right) d^2$. Lemma
159   3 says $m \le \frac{|v^T \nabla^2 F(z)v|}{\|\nabla F(z)\|}$. This completes the proof. $\qquad\square$
160

161   Lemma 6 generalises the statement $\beta_n = \alpha_n^2$ from the quadratic example. As mentioned earlier
162   the left-hand-side has the more general form $\beta_n = a \cdot (x_n - x^*)$ which is the pseudo-regret and
163   appears on the left-hand-side of the lemma. The general form of the right-hand-side is $\alpha_n^2 =$
164    $\frac{1}{2}(\nabla f(x_n) - \nabla f(x^*))^2$ for the function $f(z) = x^2 - y$ that defines the domain. The gradient
165   generalises to the normal vector to the surface and appears on the right-hand-side of the lemma.

166   **Lemma 6.** The minimiser x^* has a neighborhood U in $\mathbb{R}^d$ such that each $x \in U \cap \mathcal{M}$ has

$$a \cdot (x - x^*) \le 4\frac{\|a\|}{m}\|N(x) - N(x^*)\|^2.$$

Proof. Let $F : U \to \mathbb{R}$ be a coordinate patch at $z = x^*$ that meets the conditions in Lemma 1 and Proposition 1. Since $x^* = \operatorname{argmin}\{a \cdot x : x \in X\}$ the outwards unit normal is $N(x^*) = -\frac{a}{\|a\|}$ and the lemma gives

$$|a \cdot (x - x^*)| \le \|a\|\frac{|v^T \nabla^2 F(x^*)v|}{\|\nabla F(x^*)\|}d(x, x^*)^2.$$

167   For all $x \in U$ and direction v from x^* to x the proposition gives $d(x, x^*) \le 2\frac{\|N(x) - N(x^*)\|}{\|\nabla N(x^*)v\|}$. Combine
168   the two bounds to get

$$|a \cdot (x - x^*)| \le 4\|a\|\left(\frac{|v^T \nabla^2 F(x^*)v|}{\|\nabla F(x^*)\|}\frac{1}{\|\nabla N(x^*)v\|^2} \right)\|N(x) - N(x^*)\|^2.$$

169   To complete the proof Lemma 3 says $\frac{|v^T \nabla^2 F(x^*)v|}{\|\nabla F(x^*)\|} = |v^T \nabla N(x^*)v| \le \|v\|\|\nabla N(x^*)v\| \le \|N(x^*)v\|$.
170   Hence the middle factor above is at most $\frac{1}{\|N(x^*)v\|}$ which is at most $1/m$ by Lemma 2. $\qquad\square$

171   Lemmas 7 and 8 generalise the statement from the example that the normal at x_n has slope of absolute
172   value $\Omega(\sqrt{n})$ with high probability. The more general statement is the normal vector at x_n converges
173   to that at x^* with $O(1/\sqrt{n})$ rate. The proof is in Appendix III and follows easily from the vector
174   martingale theorems of Pinelis [28]. The use of these theorems $-$ stated only for bounded martingales
175    $-$ is the major obstruction to generalising our main results to unbounded but subgaussian cost vectors.

176   **Lemma 7.** Suppose $n \ge 16D^2/\eta^2\|a\|^2$. The Lazy Subgradient actions $x_1, x_2, ...$ give each

$$\mathbb{E}\|N(x^*) - N(x_{n+1})\|^2 \le \frac{1}{n}\frac{36}{\|a\|^2}\left(\frac{\sqrt{8\pi}DR}{\eta} + 4R^2 \right) + 8\exp\left(-\frac{\|a\|^2}{32R^2}n \right).$$

Lemma 8. For each neighborhood U of x^* in $\mathbb{R}^d$ the Lazy Subgradient actions $x_1, x_2, \ldots$ give $\sum_{i=1}^{\infty} P(x_i \notin U) < \infty$.

Proof of Theorem 1. It is enough to prove a tail bound $\mathbb{E}\left[\sum_{i>n_0}^{N} a \cdot (x_i - x^*) \right] \leq O(\log N)$ for some fixed n_0. To use Lemma 7 let $n_0 = \lfloor 16D^2/\eta^2 \|a\|^2 \rfloor$.[3] Let U be a neighborhood of x^* that meets the conditions of Lemma 6. Since each $a \cdot (x_{n+1} - x^*) \leq 2LD$ we see

$$\mathbb{E}|a \cdot (x_{n+1} - x^*)| \leq 2LD \cdot P(x_{n+1} \notin U) + \int_{x_{n+1} \in U} |a \cdot (x_{n+1} - x^*)| dP.$$

Lemma 6 says the integral is at most

$$\int_{x_{n+1} \in U} \frac{4\|a\|}{m} \|N(x_{n+1}) - N(x^*)\|^2 dP \leq \frac{4\|a\|}{m} \mathbb{E}\|N(x_{n+1}) - N(x^*)\|^2.$$

and we see $\mathbb{E}|a \cdot (x_{n+1} - x^*)|$ is at most

$$2LD \cdot P(x_{n+1} \notin U) + \frac{4\|a\|}{m} \mathbb{E}\|N(x_{n+1}) - N(x^*)\|^2.$$

Lemma 8 says the series $\sum_{i=1}^{\infty} P(x_i \notin U)$ converges. Lemma 7 gives constants $k \geq 0$ and $K = \frac{36}{\|a\|^2} \left(\frac{\sqrt{8\pi}DR}{\eta} + 4R^2 \right)$ such that $\mathbb{E}\|N(x_{n+1}) - N(x^*)\|^2 \leq K/n + 8e^{-kn}$. Hence we get

$$\mathbb{E}\left[\sum_{i>n_0}^{N} a \cdot (x_i - x^*) \right] = O(1) + \frac{4\|a\|}{m} \sum_{i=n_0}^{N} \left(K/i + 8e^{-ki} \right) = O(1) + \frac{4\|a\|}{m} \int_{n_0}^{N} \left(K/x + 8e^{-kx} \right) dx$$

$$= O(1) + \frac{4\|a\|}{m} K \log N = O(1) + \frac{4\|a\|}{m} K \log N = \frac{144}{m\|a\|} \left(\frac{\sqrt{8\pi}DR}{\eta} + 4R^2 \right) + O(1). \qquad \square$$

Equivalence of $\mathbb{E}[R_N]$ and $\mathbb{P}[R_N]$

So far Theorem 1 only gives an $O(\log N)$ bound on pseudo-regret. Hence it seems weaker than the expected regret bound of [20]. Here we show the two types of order bounds are in fact equivalent.

Theorem 2. Under Assumptions 1 and 2, Algorithm 1 with parameter $\eta > 0$ gives

$$\mathbb{E}[R_N] - \mathbb{P}[R_N] \leq \frac{31R^2}{m\|a\|} = O\left(\frac{L^2}{m\|a\|} \right).$$

Theorem 2 holds even for non smooth domains. We need only the following result of Vial [35].

Lemma 9. (Vial Theorem 1(v)) Suppose Z is m-strongly convex. For all $x, y \in \partial Z$ and outward unit normals p, q at x, y we have $\|x - y\| \leq \frac{1}{m} \|p - q\|$.

To prove Theorem 2 we first put the left-hand-side in a convenient form.

Lemma 10. Let $y^* \in \text{argmin} \left\{ \sum_{i=1}^{N} a_i \cdot x : x \in X \right\}$. Under Assumption 2 we have

$$\mathbb{E}[R_N] - \mathbb{P}[R_N] \leq \sqrt{\mathbb{E}\|x^* - y^*\|^2} \sqrt{\mathbb{E}\left\| \sum_{i=1}^{N} (a_i - a) \right\|^2}.$$

Proof. By definition $\mathbb{E}[R_N] - \mathbb{P}[R_N]$ equals $\mathbb{E}\left[\sum_{i=1}^{N} a_i \cdot (x_i - y^*) \right] - \mathbb{E}\left[\sum_{i=1}^{N} a \cdot (x_i - x^*) \right] = \mathbb{E}\left[\sum_{i=1}^{N} (a_i - a) \cdot x_i \right] + \mathbb{E}\left[\sum_{i=1}^{N} a_i \cdot (x^* - y^*) \right]$. To see the first term is zero recall each x_i is a function of $a_1, \ldots, a_{i-1}$. Since $a_1, a_2, \ldots$ are independent we see each x_i is independent of a_i and independent of $a_i - a$. Thus the expectation distributes over the product. Since $\mathbb{E}[a_i - a] = 0$ the first term vanishes. Write the second term as $\mathbb{E}\left[\sum_{i=1}^{N} (a_i - a) \cdot (x^* - y^*) \right] - \mathbb{E}\left[\sum_{i=1}^{N} a \cdot (y^* - x^*) \right]$. Since x^* minimises $a \cdot x$ the second term is positive and we get $\mathbb{E}[R_N] - \mathbb{P}[R_N] \leq \mathbb{E}\left[\sum_{i=1}^{N} (a_i - a) \cdot (x^* - y^*) \right]$. To complete the proof apply Cauchy-Schwarz to the L^2 norm. $\qquad \square$

---

[3] According to the adversarial bound the regret before turn n_0 is $O(\sqrt{n_0}) \leq O(D/\eta\|a\|)$.

To prove Theorem 2 we need the follwing three results from Appendix III. Like earlier, the first two results follow from the vector martingale theorems of Pinelis [28]. The third is well known but we were unable to find a sufficiently general proof in the literature.

Proposition 2. Suppose $X_1, X_2, \ldots \in \mathbb{R}^d$ are independent random variables with each $\mathbb{E}[X_n] = 0$ and $\|X_n\| \leq R$. For each $t > 0$ we have

$$P\left(\left\|\frac{1}{n}\sum_{i=1}^n X_i\right\| > t\right) \leq 2\exp\left(-\frac{t^2}{2R^2}n\right).$$

Lemma 11. For $y^* = \operatorname{argmin}\left\{\sum_{i=1}^N a_i \cdot x : x \in X\right\}$ The Lazy Subgradient actions give

$$\mathbb{E}\|N(x^*) - N(y^*)\|^2 \leq \frac{144R^2}{\|a\|^2 N} + 4\exp\left(-\frac{\|a\|^2}{8R^2}N\right).$$

Lemma 12. For X a nonnegative random variable we have $\mathbb{E}[X] = \int_0^\infty P(X > x)dx$.

Proof of Theorem 2. Proposition 2 gives $P\left(\|\sum_{i=1}^N (a_i - a)\| > t\right) \leq 2\exp\left(-\frac{t^2}{2NR^2}\right)$ for all $t \geq 0$. Replace t with t^2 to see $P\left(\|\sum_{i=1}^N (a_i - a)\|^2 > t\right) \leq 2\exp\left(-\frac{t}{2NR^2}\right)$. Lemma 12 says $\mathbb{E}\|\sum_{i=1}^N (a_i - a)\|^2 \leq \int_0^\infty P\left(\|\sum_{i=1}^N (a_i - a)\|^2 > t\right)dt \leq 4NR^2$. Hence Lemma 10 gives

$$\mathbb{E}[R_N] - \mathbb{P}[R_N] \leq 2R\sqrt{N}\sqrt{\mathbb{E}\|x^* - y^*\|^2}.$$

For the second factor Lemma 9 gives $\sqrt{\mathbb{E}\|x^* - y^*\|^2} \leq \frac{1}{m}\sqrt{\mathbb{E}\|N(x^*) - N(y^*)\|^2}$. Lemma 11 gives $\mathbb{E}\|N(x^*) - N(y^*)\|^2 \leq K/N + 4e^{-kN}$ for the constants $K = 144R^2/\|a\|^2$ and $k = \|a\|^2/8R^2$. Hence $\sqrt{\mathbb{E}\|x^* - y^*\|^2} \leq \frac{1}{m}\sqrt{K/N + 4e^{-kN}} \leq \frac{1}{m}\sqrt{\frac{K}{N}} + \frac{2}{m}e^{-(k/2)N}$ and so

$$\mathbb{E}[R_N] - \mathbb{P}[R_N] \leq \frac{2R\sqrt{K}}{m} + \frac{4R}{m}\sqrt{N}e^{-(k/2)N} = \frac{24R^2}{m\|a\|} + \frac{4R}{m}\sqrt{N}e^{-(k/2)N}. \tag{3}$$

To bound the last term consider the function $\sqrt{x}e^{-\beta x}$ for $\beta \geq 0$. It is straightforward to see the function is increasing and then decreasing. By differentiating we see the maximiser is $x = 1/2\beta$ and the function is at most $1/\sqrt{2e\beta}$. For $\beta = k/2$ we see $\sqrt{N}e^{-(k/2)N} \leq 1/\sqrt{ke} = 2\sqrt{2}R/\sqrt{e}\|a\|$. Thus the right-hand-side of (3) is at most $\left(24 + \frac{8\sqrt{2}}{\sqrt{e}}\right)\frac{R^2}{m\|a\|} \leq \frac{31R^2}{m\|a\|}$. $\square$

Finally we can combine Theorems 1 and 2:

Theorem 3. The Lazy Subgradient algorithm with parameter $\eta > 0$ gives

$$\mathbb{E}[R_N] \leq \frac{144}{m\|a\|}\left(\frac{\sqrt{8\pi}DR}{\eta} + 4R^2\right)\log N + O(1).$$

Numerical Simulations

Here we present numerical simulations for the Lazy Subgradient algorithm, using a range of strongly convex domains and i.i.d opponents. The higher dimensional simulations ran on the order of minutes, due to use of an all-purpose Python package to compute minimisers. Performance time can be improved by using the Franke Wolfe method of [17] to simplify the minimisation problems.

ℓ_p**-Norm Balls.** Here we examine performance for domains $X = \left\{x \in \mathbb{R}^d : \left(\sum_{i=1}^d |x(i)|^p\right)^{1/p} \leq 1\right\}$. For $p \in (1, 2]$ Corollary 1 of [17] says X is $(p-1)d^{\frac{1}{2}-\frac{1}{p}}$ strongly convex.

For simplicity we use stepsize $\eta = 1$. When searching for worst-case performance it is enough, by symmetry of the domain, to only consider $\mathbb{E}[a_n] = a$ with nonnegative nondecreasing components. We consider $a = b/\|b\|$ for $b(i) = (i/d)^r$ and a range of parameters $r \geq 0$. For example three degenerate cases are $a = \frac{1}{\sqrt{d}}(1, \ldots, 1)$ for $r = 0$; $a = \sqrt{\frac{2}{d(d+1)}}(1, 2, \ldots, d)$ for $r = 1$ and $a = (0, \ldots, 0, 1)$ for $r \to \infty$. We use cost vectors $a_n = a + \mu_n$ for two types of noise:

$$(1)\ \mu_n(j) = \frac{1}{\sqrt{d}}(B_1^n, \ldots, B_d^n) \qquad (2)\ \mu_n(j) = (0, \ldots, 0, B_d^n).$$

| (a) Noise type (1) | (b) Noise Type (2) |

Figure 1: Lazy Subgradient on the $\ell_{1.5}$ unit ball.

Here B_j^n are independent and take values ± 1 with equal probability. In both cases $\|\mu_n\| = 1$. Noise type (1) perturbs all components equally and individual perturbations are small for large d. Type (2) perturbs only the largest component, and the perturbation always has length 1. We run several instances (samples) of each online problem and plot the average pseudo-regret over instances.

Figure 1(a) shows that for noise type (1) performance degrades as r shrinks. This is consistent with the picture for $d = 2$ where the minimiser $(2^{-1/p}, 2^{-1/p})$ also minimises the curvature, and how Theorem 1 gives a worse bound for small curvature. For noise type (2) Figure 1(b) showslarger r gives worse performance. This can be attributed to how r is approximately $(0, \ldots, 0, 1)$ and the noise is focused on the final component rather than spreading over all components evenly.

For smaller of d, p the pattern is the same but takes longer to emerge. In Figure 3, Appendix I, we consider the worst cases in Figure 1 and vary the dimension. For $r = 0$ and noise type (1) the performance improves slightly with large d. This is perhaps due to the higher dimension having a regularising effect on the noise. For $r = 8$ and noise type (2) the performance deteriorates with large d. This is expected since the curvature $(p-1)d^{\frac{1}{2} - \frac{1}{p}}$ shrinks in high dimensions. In Figure 4, Appendix I, we vary the parameter p. In both worse cases the performance deteriorates as $p \to 1$ though it also takes longer for the worse performance to emerge.

For all cases the growth rates are orders of magnitude below those suggested by Theorem 1. For example in the setup of Figure 1 we have $p = 1.5$, $d = 32$ and the convexity parameter is $m = (p-1)d^{\frac{1}{2} - \frac{1}{p}} = 0.5 \cdot 32^{-\frac{1}{4}} \simeq 0.21$. This makes the coefficient in Theorem 1 approximately 6180.

Comparison with Greedy Subgradient. Theorem 3 says Lazy Subgradient gives logarithmic expected regret on a strongly convex domain against i.i.d cost vectors. One can ask does the Greedy Subgradient algorithm with update $x_{n+1} = P_X \left(x_n - \frac{\eta}{\sqrt{n}} a_n \right)$ give the same bound. Figure 2(b) suggests otherwise. The worst-case regret of Greedy Subgradient seems to be $\Omega(\sqrt{n})$. The coefficients D_1, C_1 in Figure 2(b) are chosen to match the regret on turns 1000 and 500, and the $\Theta(\sqrt{n})$ fit is very close over the entire range. The coefficient C_2 is chosen to match on turns 0 and 1000 and is a worse fit.

Schatten Matrix Norms. Schatten matrix norms are well-studied as regularisers in the context of machine learning. For example see [3, 21] and the citing works.

Definition 2. Given a real $n \times m$ matrix A there a unique decomposition $A = U\Sigma V$ such that V, U are real orthogonal, U is $n \times n$, V is $m \times m$, and Σ is $n \times m$ with real diagonal entries $\sigma_1 \geq \sigma_2 \geq \ldots \geq \sigma_{\min(n,m)}$ and all other entries zero. The values σ_i are called the singular values of A. For any $p \geq 0$ define the Schatten p-norm $\|A\|_{S(p)} = \|\sigma\|_p$ as the p-norm of the vector of singular values.

For simplicity we only consider square matrices. Corollary 2 of [17] says the Schatten unit balls $B_p(1) = \{X \in \mathbb{R}^{d \times d} : \|X\|_{S(p)} \leq 1\}$ are $(p-1)d^{\frac{1}{2} - \frac{1}{p}}$ strongly convex. When searching for worst case performance the following lemma says it is enough to only consider when $\mathbb{E}[a_n]$ is a diagonal matrix. See Appendix I for proof.

(a) Lazy Subgradient on the Schatten unit ball   (b) Greedy Subgradient on the Euclidean unit ball

Figure 2

Lemma 13. Suppose $a_1, a_2, \ldots \in \mathbb{R}^{d \times d}$ are i.i.d cost vectors with $\mathbb{E}[a_n] = a$. There is another set of i.i.d cost vectors $b_1, b_2, \ldots$ such that $\mathbb{E}[b_n] = b$ is diagonal and $\|a_n - a\| = \|b_n - b\|$; $\|a_n\| = \|b_n\|$; $\|a\| = \|b\|$ and the expected regret of playing Lazy Subgradient on $B_p(1)$ against $a_1, a_2, \ldots$ with $x_0 = 0$ is the same as playing against $b_1, b_2, \ldots$ with $x_0 = 0$.

Likewise by symmetry we can assume the diagonal entries are nonnegative and nondecreasing. For simplicity we use stepsize $\eta = 1$. We consider $a = b/\|b\|$ for $b = \mathrm{diag}\big((1/d)^r, (2/d)^r, \ldots, 1\big)$ and a range of parameters $r \geq 0$. We consider cost vectors $a_n = a + \mu^n$ and noise $\mu^n = \nu^n/\|\nu^n\|$ for three types of ν^n. Here B_{ij}^n are independent and take value ± 1 with equal probability.

    (1) $\nu_{ij}^n = B_{ij}^n$ for all i, j

    (2) $\nu_{ij}^n = B_{ij}^n$ for all i and some uniformly chosen j; and $\nu_{ik}^n = 0$ for all $k \neq j$.

    (3) $\nu_{ij}^n = B_{ij}^n$ for some uniformly chosen i, j; and $\nu_{lk}^n = 0$ for all $(l, k) \neq (i, j)$.

Compared to the examples for ℓ_p balls, fewer samples are needed to obtain smooth curves. For noise type (1) Figure 2(a) shows the worst performance occurs for $r = 0$. The different types of noise make minimal difference to performance. See Figure 5, Appendix I. Thus when varying p and d we only consider noise type (1). In Figure 6, Appendix I, we see large d leads to slightly better performance, and small p leads to worse performance. Though the infuence of p and d is minimal compared to that in Figures 5 and 2 for ℓ_p-balls.

For Schatten balls diagonal noise matrices seem to be the most difficult to deal with. In that case all a_n are also diagonal and the Schatten norm reduces to the p-norm of the diagonal entries.

Conclusion

The Lazy Subgradient algorithm on a strongly convex domain is universal in the sense that it achieves $O(\log N)$ regret against i.i.d opponents. This is in addition to the well-known $O(\sqrt{N})$ regret bound against adversarial opponents. Moreover for strongly convex domains, order bounds for pseudo-regret and expected regret are equivalent. For various types of noise on ℓ_p and Schatten unit balls, the growth coefficients are orders of magnitude smaller than those suggested by our analysis.

Disclosure of Funding. This work was supported by Science Foundation Ireland grant 16/IA/4610.

Broader Impact. This is theoretical work on special cases of online optimisation, with no particular application in mind. We do not forsee the work having any direct impact on society. The only ethical concerns are those related to the use of online optimisation in general.

References

[1] Daron Anderson and Douglas Leith. Optimality of the subgradient algorithm in the stochastic setting. *Arxiv E-prints*, 2020.

[2] Daron Anderson and Douglas Leith. Universal algorithms: Beyond the simplex. *Arxiv E-prints*, 2020.

[3] Andreas Argyriou, Charles A Micchelli, and Massimiliano Pontil. On spectral learning. *Journal of Machine Learning Research*, 11(2), 2010.

[4] Peter Auer and Chao-Kai Chiang. An algorithm with nearly optimal pseudo-regret for both stochastic and adversarial bandits. *CoRR*, abs/1605.08722, 2016.

[5] Aditya Bhaskara, Ashok Cutkosky, Ravi Kumar, and Manish Purohit. Online Learning with Imperfect Hints. *Proceedings of the 37th International Conference on Machine Learning, Vienna, Austria*, 119, 2020.

[6] Sébastien Bubeck, Nicolo Cesa-Bianchi, and Sham M Kakade. Towards minimax policies for online linear optimization with bandit feedback. In *Conference on Learning Theory*, pages 41–1, 2012.

[7] Sébastien Bubeck, Michael Cohen, and Yuanzhi Li. Sparsity, variance and curvature in multi-armed bandits. In *Algorithmic Learning Theory*, pages 111–127, 2018.

[8] Sébastien Bubeck and Aleksandrs Slivkins. The best of both worlds: stochastic and adversarial bandits. *CoRR*, abs/1202.4473, 2012.

[9] Nicolo Cesa-Bianchi, Yishay Mansour, and Gilles Stoltz. Improved second-order bounds for prediction with expert advice. *Machine Learning*, 66(2-3):321–352, 2007.

[10] Mohammed Dahleh, Munther A Dahleh, and George Verghese. Lectures on dynamic systems and control. 2011.

[11] Steven De Rooij, Tim Van Erven, Peter D Grünwald, and Wouter M Koolen. Follow the leader if you can, hedge if you must. *The Journal of Machine Learning Research*, 15(1):1281–1316, 2014.

[12] Ofer Dekel, Nika Haghtalab, Patrick Jaillet, et al. Online learning with a hint. In *Advances in Neural Information Processing Systems*, pages 5299–5308, 2017.

[13] V. F. Demyanov and A. M. Rubinov. Approximate Methods in Optimization Problems. (Modern Analytic and Computational Methods in Science and Mathematics). IX + 256 S. New York 1970. American Elsevier Publishing Company, Inc. Preis geb. Dfl. 77.50. *ZAMM - Journal of Applied Mathematics and Mechanics / Zeitschrift für Angewandte Mathematik und Mechanik*, 53(7):499–499, 1973.

[14] Joseph C Dunn. Rates of convergence for conditional gradient algorithms near singular and nonsingular extremals. *SIAM Journal on Control and Optimization*, 17(2):187–211, 1979.

[15] Ryszard Engelking. *General topology*. Sigma Series in Pure and Applied Mathematics, Volume 6. Heldermann Verlag Berlin, 1977.

[16] Pierre Gaillard, Gilles Stoltz, and Tim Van Erven. A second-order bound with excess losses. In *Conference on Learning Theory*, pages 176–196, 2014.

[17] Dan Garber and Elad Hazan. Faster rates for the frank-wolfe method over strongly-convex sets. In *32nd International Conference on Machine Learning, ICML 2015*, 2015.

[18] Geoffrey J Gordon. Regret bounds for prediction problems. In *Proceedings of the twelfth annual conference on Computational learning theory*, pages 29–40, 1999.

[19] Elad Hazan and Nimrod Megiddo. Online learning with prior knowledge. In *International Conference on Computational Learning Theory*, pages 499–513. Springer, 2007.

[20] Ruitong Huang, Tor Lattimore, András György, and Csaba Szepesvári. Following the leader and fast rates in online linear prediction: Curved constraint sets and other regularities. *J. Mach. Learn. Res.*, 18(1):5325–5355, January 2017.

[21] Sham M Kakade, Shai Shalev-Shwartz, and Ambuj Tewari. Regularization techniques for learning with matrices. *The Journal of Machine Learning Research*, 13(1):1865–1890, 2012.

[22] Ehsan Kazemi, Thomas Kerdreux, and Liqiang Wang. Trace-norm adversarial examples. *arXiv preprint arXiv:2007.01855*, 2020.

[23] Thomas Kerdreux, Alexandre d'Aspremont, and Sebastian Pokutta. Projection-Free Optimization on Uniformly Convex Sets. *arXiv e-prints*, page arXiv:2004.11053, April 2020.

[24] Peter M. Lee. The Probability Integral. *York College*.

[25] Evgeny S Levitin and Boris T Polyak. Constrained minimization methods. *USSR Computational mathematics and mathematical physics*, 6(5):1–50, 1966.

[26] Marco Molinaro. Curvature of feasible sets in offline and online optimization. *arXiv preprint arXiv:2002.03213*, 2020.

[27] Jaouad Mourtada and Stéphane Gaïffas. On the optimality of the Hedge algorithm in the stochastic regime. *Journal of Machine Learning Research*, 20:1–28, 2019.

[28] Iosif Pinelis. Optimum bounds for the distributions of martingales in Banach spaces. *The Annals of Probability*, 22(4):1679–1706, 10 1994.

[29] Pavel Pokorny. Geodesics revisited. *Chaotic Modeling and Simulation (CMSIM) 2012*, page 281–298, 2012.

[30] Paat Rusmevichientong and John N Tsitsiklis. Linearly parameterized bandits. *Mathematics of Operations Research*, 35(2):395–411, 2010.

[31] Amir Sani, Gergely Neu, and Alessandro Lazaric. Exploiting easy data in online optimization. *Proceedings of the 27th International Conference on Neural Information Processing Systems - Volume 1*, pages 810–818, 2014.

[32] Yevgeny Seldin and Gábor Lugosi. An improved parametrization and analysis of the EXP3++ algorithm for stochastic and adversarial bandits. *CoRR*, abs/1702.06103, 2017.

[33] Yevgeny Seldin and Aleksandrs Slivkins. One practical algorithm for both stochastic and adversarial bandits. In *Proceedings of the 31st International Conference on Machine Learning*, volume 32(2), 2014.

[34] Shai Shalev-Shwartz. Online learning and online convex optimization. *Found. Trends Mach. Learn.*, 4(2):107–194, February 2012.

[35] Jean-Philippe Vial. Strong convexity of sets and functions. *Journal of Mathematical Economics*, 9(1-2):187–205, 1982.

[36] Zuoqin Wang. Lecture 10: Tubular Neighborhood Theorem. *University of Science and Technology of China*.

[37] Chen-Yu Wei and Haipeng Luo. More adaptive algorithms for adversarial bandits. *CoRR*, abs/1801.03265, 2018.

[38] Dmitri Zaitsev. Differential geometry: Lecture notes. *Trinity College, Dublin, Ireland*, 2003.

[39] Julian Zimmert and Tor Lattimore. Connections between mirror descent, thompson sampling and the information ratio. In *Advances in Neural Information Processing Systems*, pages 11973–11982, 2019.

[40] Julian Zimmert and Yevgeny Seldin. An optimal algorithm for stochastic and adversarial bandits. *CoRR*, abs/1807.07623, 2018.

[41] Martin Zinkevich. Online Convex Programming and Generalized Infinitesimal Gradient Ascent. pages 928–935, 2003.