GENERALIZATION ERROR BOUNDS FOR NEURAL NETWORKS WITH ReLU ACTIVATION

ABSTRACT

We show rigorous bounds on the generalization error for Neural Networks with ReLU activation under the condition that the network size doesn't grow with the training set size. In order to prove these bounds we weaken the notion of uniform stability of a learning algorithm in a probabilistic way by positing the notion of almost sure (a.s.) support stability and proving that if an algorithm has low enough a.s. support stability its generalization error tends to 0 as the training set size increases. Further we show that for Stochastic Gradient Descent to be almost surely support stable we only need the loss function to be locally Lipschitz and locally smooth with probability 1, thereby showing low generalization error with weaker conditions than have been used in the literature. We then show that Neural Networks with ReLU activation and a doubly differentiable loss function possess these properties, thereby proving low generalization error. The caveat is that the size of NN must not grow with the size of the training set. Finally we present experimental evidence to validate our theoretical results.

1 INTRODUCTION

Given the importance of the generalization error in machine learning there have been numerous theoretical approaches to bounding it, one of the most significant being the concept of algorithmic stability whose origins can be traced back to Vapnik & Chervonenkis (1974). In this paradigm the definition of uniform stability by Bousquet & Elisseeff (2002) was a landmark because it allowed the authors to prove sharp concentration bounds showing that the generalization error for the algorithms that satisfy this property tends to 0 as the training set size increases. Thanks to the success of Bousquet & Elisseeff (2002) on a variety of non-neural methods, when Hardt et al. (2016) showed that a learning function computed by optimizing certain loss functions through Stochastic Gradient Descent is uniformly stable under mild conditions on the loss function, this clearly raised the expectation that the next step would be to show that Neural Network-based models are also uniformly stable and, hence, have a low generalization error. This analysis was then expected to corroborate the empirically observed fact that the generalization error is low for NN models (c.f., e.g., Krizhevsky et al. (2012), Hinton et al. (2012).) Case closed. However, this triumphal march of algorithmic stability was rudely interrupted by Zhang et al. (2017), who showed explicit examples of distributions that cannot be learned with low generalization error and yet can be learned with an NN of an appropriate size. Zhang et al. (2017) speculated that the problem arose from the fact that the concept of uniform stability was based *only* on properties of the algorithm and not on the properties of the unknown data distribution. While this was a reasonable speculation, it left open some questions: Is there a reasonable definition of stability that incorporates distribution properties and leads to small generalization error? And can we show that NNs with non-linear activation functions such as ReLU at the neurons have this property?

In this paper we give a strongly affirmative answer to the first question and a partially affirmative answer to the second question. We define a notion called *almost sure (a.s.) support stability* which is a probabilistic weakening of uniform stability. Unlike the data-dependent notions of stability defined in Kuzborskij & Lampert (2018), Lei & Ying (2020) that bound generalization error in expectation, a.s. support stability can be used to show high probability bounds on generalization error. This proof entails proving a mild generalization of McDiarmid's Inequality that could be of independent interest. The major contribution of our paper is that we show how to handle the non-linearity of ReLU in a mathematically rigorous way by modifying a result of Milne (2019) to show that ReLU

affects the smoothness of the gradients computed by Stochastic Gradient Descent with probability 0 for well-behaved distributions. To the best of our knowledge such a result has not been reported in the literature.

Clearly the question arises: What about the evidence presented by Zhang et al. (2017) that there exist unknown distributions that can be learned by an NN but will always have unbounded generalization error? To answer this we need to understand that our analysis of SGD presented requires the loss function to be smooth and Lipschitz in the parameter space (though with an appropriate probabilistic weakening of these properties.) Our analysis is able to demonstrate that these properties hold for NNs whose size doesn't grow with the size of the training set. So, our specific answer that we have for the second question listed above: *Almost support stability can be used to demonstrate that the generalization error for NNs with ReLU as long as the size of the NN doesn't grow too fast with the training set size, and if the unknown distribution places probability 0 on sets of Lebesgue measure 0.* This leaves open the matter of a theoretical explanation for why certain heavily overparametrized NNs generalize well in practice. However, we do provide some directions on how our analytical framework may be used to prove such results.

In particular our contributions are:

• In Section 3 we define a new notion of stability called *a.s. support stability* and show in Theorem 3 that algorithms a.s. support stability $o(1/\sqrt{m})$ have generalization error tending to 0 as $m \rightarrow \infty$ where m is the size of the training set. In order to prove this we state and prove Theorem 13 which is an extension of McDiarmid's Theorem.

• In Section 4 we show that if we run stochastic gradient descent on a parametrized optimization function that is only *locally* Lipschitz and *locally* smooth in the parameter space and that too only on input points selected with probability 1, then the replace-one error is bounded even if the training is conducted for $c \log m$ epochs. This implies (Corollary 8) that any learning algorithm trained this way has generalization error that goes to 0 as $m \rightarrow \infty$ for learning rate α_0/t at step t for an appropriately selected value of α_0 that does not depend on m.

• In Section 5 we show that the output of an NN with ReLU activations when used with a doubly differentable loss function is locally Lipschitz and locally smooth in the parameter space for all inputs except those from a set of Lebesgue measure 0 (Theorem 11). This allows us to show a.s. support stability and a generalization error bound for NNs with size that doesn't grow with m.

• We experimentally verify our theoretical results in Section 6, showing that our bounded Lipschitz and bounded smoothness conditions hold in practice.

2 RELATED WORK

Several authors have that although NNs are known to generalize well in practice, many different theoretical approaches have been tried without satifactorily explaining this phenomenon, c.f., Jin et al. (2020); Chatterjee & Zielinski (2022). We refer the reader to the work of Jin et al. (2020) which presents a concise taxonomy of these different theoretical approaches. There are also several works that seek to understand what a good theory of generalization should look like, c.f. Kawaguchi et al. (2017); Chatterjee & Zielinski (2022). Our own work falls within the paradigm that seeks to use notions of algorithmic stability to bound generalization error that began with Vapnik & Chervonenkis (1974) but gathered steam with the publication of the work by Bousquet & Elisseeff (2002).

The applicability of the algorithmic stability paradigm to the study of generalization error in NNs was brought to light by Hardt et al. (2016) who showed that functions optimized via Stochastic Gradient Descent have the property of uniform stability defined by Bousquet & Elisseeff (2002), implying that NNs should also have this property. Subsequently, there was renewed interest in uniform stability and a sequence of papers emerged using improved probabilistic tools to give better generalization bounds for uniformly stable algorithms, e.g., Feldman & Vondrak (2018; 2019) and Bousquet et al. (2020). Some other works, e.g. Klochkov & Zhivotovskiy (2021), took this line forward by focussing on the relationship of uniform stability with the excess risk. However the work of Zhang et al. (2017) complicated the picture by pointing out examples where the theory suggests the opposite of what happens in practice. This led to two different strands of research. In one thread an attempt was made to either discover those cases where uniform stability, (e.g. Charles & Papail-

iopoulos (2018)), or to show lower bounds on stability that ensure that uniform stability does not exists, (e.g. Zhang et al. (2022)). The other strand of research, our work falls in this category, focuses on weakening the notion of uniform stability, specifically by making it data-dependent, thereby following the suggestion made by Zhang et al. (2017). Kuzborskij & Lampert (2018) defined "on-average stability" which is weaker than our definition of a.s. support stability. Consequently, their definition leads to a weaker in-expectation bound on the generalization error where the expectation is over the training set as well as the random choices of the algorithm. Our Theorem 3, on the other hand, provides a sharp concentration bound on the choice of training set. Lei & Ying (2020) define an "on-average model stability" that requires the average replace-one error over all the training points to be bounded in expectation. While their smoothness requirements are less stringent, the problem is that their generalization results are all relative to the optimal choice of the weight vector, which implies a high generalization error in case of early stopping.

3 A.S. SUPPORT STABILITY AND GENERALIZATION

In this section we present a weakening of the notion of uniform stability defined by Bousquet & Elisseeff (2002) and show that exponential concentration bounds on the generalization error can be proved for learning algorithms that have this weaker form of stability.

3.1 TERMINOLOGY

Let $\mathcal{X}$ and $\mathcal{Y}$ be the input and output spaces respectively. We assume we have a training set $S \in \mathcal{Z}^m$ of size m where each point is chosen independently at random from an unknown distribution D over $\mathcal{Z} \subset \mathcal{X} \times \mathcal{Y}$. For $z = (x, y) \in \mathcal{Z}$ we will use the notation x_z to denote x and y_z to denote y. Let $\mathcal{R}$ be the set of all finite strings on some finite alphabet, and let us call the elements of $\mathcal{R}$ *decision strings* and let us assume that there is some probability distribution D_r according to which we will select r randomly from $\mathcal{R}$. Further, let $\mathcal{F}$ be the set of all functions from $\mathcal{X}$ to $\mathcal{Y}$. In machine learning settings we typically compute a map from $\mathcal{Z}^m \times \mathcal{R}$ to $\mathcal{F}$. We will denote the function computed by this map as $A_{S,r}$. Since the choice of S and r are both random, $A_{S,r}$ is effectively a random function and can also be thought of as a randomized algorithm.

Given a bound $M > 0$, we assume that we are given a bounded *loss function* $\ell : \mathcal{Y} \times \mathcal{Y} \to [0, M]$. We define the *risk* of A_S as
$$R(A, S, r) = \mathrm{E}_{z \sim D} \left[\ell(A_{S,r}(x_z), y_z) \right],$$
where the expectation is over the random choice of point z according to data distribution D. Note that the risk is a random variable since both S and r are randomly chosen. The *empirical risk* of A_S is defined as
$$R_e(A, S, r) = \frac{1}{|S|} \sum_{z \in S} \ell(A_{S,r}(x_z), y_z).$$

We are interested in bounding the *generalization error*
$$|R(A, S, r) - R_e(A, S, r)|.$$

3.2 A WEAKENING OF UNIFORM STABILITY

Given $S = \{Z_1, \ldots, Z_m\}$ chosen randomly according to D^m, we choose $\{Z_{1+m}, \ldots, Z_{2m}\}$ also from D^m independently from S and, for each $i \in [m]$, we define
$$S^i = \{Z_1, \ldots, Z_{i-1}, Z_{i+m}, Z_{i+1}, \ldots, Z_m\}.$$

Definition 1 (Almost (Sure) Support Stability). *We say an algorithm $A_{S,r}$ has η-almost support stability β with respect to the loss function $\ell(\cdot, \cdot)$ if for $Z_1, \ldots, Z_{2m}$ chosen i.i.d. according to an unknown distribution D defined over $\mathcal{Z}$,*
$$\forall i \in [m] : \forall z \in supp(D) : E_r \left[|\ell(A_{S,r}(x_z), y_z) - \ell(A_{S^i,r}(x_z), y_z)| \right] \leq \beta$$
with probability $1 - \eta$. If $\eta = 0$ then we say the algorithm has almost sure (a.s.) support stability β.

We note that this notion weakens the notion of uniform stability introduced by Bousquet & Elisseeff (2002) by requiring the bound on the difference in losses to hold with a certain probability. This probability is defined over the random choices of $Z_1, \ldots, Z_{2m}$. Besides the condition on the loss is required to hold only for those data points that lie in the support of D. These conditions make

almost support stability a *data-dependent quantity* on the lines of the suggestion made by Zhang et al. (2017). We also observe that almost support stability is comparable to but stronger than the hypothesis stability of Kearns & Ron (1999) as formulated by Bousquet & Elisseeff (2002).

3.3 EXPONENTIAL CONVERGENCE OF GENERALIZATION ERROR

Next, as promised, we prove that an η-almost β-support stable algorithm has low bias and converges exponentially in the size of S to its expected behavior. We first define a probabilistic weakening of Lipschitz functions.

Definition 2. *Given $2m$ i.i.d. random variables $X_1, \ldots, X_{2m}$ drawn from some domain $\mathcal{Z}$ according to some probability distribution D, for some $\beta > 0$ and $\eta \in [0, 1]$, a function $f : \mathcal{Z}^m \to \mathbb{R}$ is called η-almost β-Lipschitz w.r.t. D if*

$$\forall i \in \{1, \cdots, m\} : |f(X_1, \ldots, X_m) - f(X_1, \ldots, X_{i-1}, X_i', X_{i+1}, \ldots, X_m)| \le \beta,$$

with probability at least $1 - \eta$. In case $\eta = 0$ we say that f is almost surely β-Lipschitz w.r.t D. When D is understood we will omit it, and for the case $\eta = 1$ we will simply write that f is almost surely (or just a.s.) β-Lipschitz.

Bousquet & Elisseeff (2002) refer to a class of learning algorithms that they call "symmetric". This means that the value of the loss function is symmetric *in distribution* over the training points. We note though that if the training points are selected i.i.d. from a distribution then *any* loss function that depends on these training points will be symmetric in distribution (and therefore in expectation).

Theorem 3. *Let $A_{S,r}$ be a algorithm that is symmetric in distribution and is η-almost β-support stable with respect to the loss function $\ell(\cdot, \cdot)$ such that $0 \le \ell(A_{S,r}(x_z), y_z) \le M$ for all $S \in \mathcal{Z}^m$, for all $r \in \mathcal{R}$ and for all $z = (x_z, y_z) \in \mathcal{Z}$. Then, for any $m \ge 1$ and $\delta \in (0, 1)$ such that $\delta > \eta$, with probability $1 - \delta$*

$$E_r[R(S, r) - R_e(S, r)] \le (1 - \eta)\beta + \eta M + (2\beta m + M)\sqrt{\frac{\log(1/(\delta - \eta))}{m}}.$$

The bound given in this theorem should be compared with the bound given in Theorem 12 of Bousquet & Elisseeff (2002). When $\eta = 0$ the bound is exactly the same. But Theorem 3 is more useful in this case since it allows the stability condition to be violated on a training set that is picked with probability 0. This may seem like a triviality but we will see ahead that it is precisely this difference that allows us to obtain generalization bounds for Neural Networks with ReLU activation. The proof of the theorem is in Appendix B. The proof requires us to prove a mild generalization of McDiarmid's theorem that could be of independent interest and is presented in Appendix A.

4 A. S. SUPPORT STABILITY OF STOCHASTIC GRADIENT DESCENT

A large family of machine learning algorithms follow a paradigm in which the learned function is parametrized by a vector $\boldsymbol{w} \in \mathbb{R}^n$ for some $n \ge 1$, i.e., we have some fixed function $g : \mathbb{R}^n \times \mathcal{X} \to \mathcal{Y}$. The training set is used to learn a suitable parameter vector $\boldsymbol{w} \in \mathbb{R}^n$ such that the value $g(\boldsymbol{w}, x_z)$ is a good estimate of y_z for all $z \in \mathcal{Z}$. This value of $\boldsymbol{w}$ is learned by running Stochastic Gradient Descent (SGD) using a training set drawn from the unknown distribution. We will say that the size of the training set is m and the number of steps for which SGD is trained is T with the parameter vector at step t being denoted $\boldsymbol{w}_t$ for $0 \le t \le T$. Assuming that T is a multiple of m, the algorithm moves in *epochs* of m steps. To frame the learned function output by this algorithm in the terms defined in 3.1, the random decision string r consists of the pair $(\boldsymbol{w}_0, (\pi_0, \ldots, \pi_{\frac{T}{m}-1}))$, i.e., the random initial parameter vector $\boldsymbol{w}_0$ from which SGD beging and and the set of T/m random permutations used in the T/m epochs.

4.1 SOME PROPERTIES OF PARAMETERIZED FUNCTION

In Hardt et al. (2016), proving that the learning algorithm derived by SGD is stable requires smoothness and Lipschitz properties of f, but *only* for partial derivatives taken on $\mathbb{R}^n$, i.e., on the parameter space. The requirement there is that *every function* in the family of functions $\{f(\cdot, z) : z \in \mathcal{Z}\}$ is smooth and has a bounded Lipschitz constant. Our key insight is that this requirement is stronger

than required. All we need is that *the functions induced by the data points that we pick to train SGD have these properties.* We now present some definitions that encapsulate this idea.

Definition 4 (Almost Locally Parameter Lipschitz functions)**.** *Given a distribution D defined over $\mathcal{Z}$, a parametrized function $f : \mathbb{R}^n \times \mathcal{Z} \to \mathbb{R}$ is said to be β-almost Locally L_l Lipschitz w.r.t D for some $\beta \in (0,1]$ if for any $\boldsymbol{w} \in \mathbb{R}^n$ there exists constants $L_l > 0$ and $\epsilon > 0$ such that, with probability β (over the choice of z), for all $\boldsymbol{w}' \in \mathbb{R}^n$, $\|\boldsymbol{w}' - \boldsymbol{w}\| < \epsilon$ implies*
$$|f(\boldsymbol{w}', z) - f(\boldsymbol{w}, z)| \le L_l \|\boldsymbol{w}' - \boldsymbol{w}\|.$$
Note that L_l can depend on z. If $\beta = 1$ then we say that f is almost surely locally L_l-parameter Lipschitz *(*a.s. L_l-LPL *for short).*

Definition 5 (Almost Locally Parameter-Smooth functions)**.** *Given a distribution D defined over $\mathcal{Z}$, a parametrized function $f : \mathbb{R}^n \times \mathcal{Z} \to \mathbb{R}$ is said to be γ-almost K_l-Parameter Smooth w.r.t D for some $\gamma \in (0,1]$ if for any $\boldsymbol{w} \in \mathbb{R}^n$ there exist constants $K_l > 0$ and $\epsilon > 0$ such that, with probability γ (over the choice of z), for all $\boldsymbol{w}' \in \mathbb{R}^n$, $\|\boldsymbol{w}' - \boldsymbol{w}\| < \epsilon$ implies*
$$\|\nabla f(\boldsymbol{w}', z) - \nabla f(\boldsymbol{w}, z)\| \le K_l \|\boldsymbol{w}' - \boldsymbol{w}\|.$$
Note that K_l can depend on z. If $\gamma = 1$ then we say that f is almost surely locally K_l-parameter smooth *(*a.s. K_l-LPS *for short).*

If the function (or its gradient) is locally bounded, and, if we only look at this function at a finite number of points, we get a "global" property within this finite set of points:

Lemma 6. *Given $f : \mathbb{R}^n \to \mathbb{R}$ we have that (1) if f is bounded and is locally Lipschitz at a finite set of points $A \subseteq \mathbb{R}^n$, then there is a constant L such that for every pair $\boldsymbol{w}, \boldsymbol{w}' \in A$*
$$|f(\boldsymbol{w}) - f(\boldsymbol{w}')| \le L\|\boldsymbol{w} - \boldsymbol{w}'\|, and$$
2) if ∇f is bounded and is locally smooth at a finite set of points $A \subseteq \mathbb{R}^n$, then there is a constant K such that for every pair $\boldsymbol{w}, \boldsymbol{w}' \in A$
$$\|\nabla f(\boldsymbol{w}) - \nabla f(\boldsymbol{w}')\| \le K\|\boldsymbol{w} - \boldsymbol{w}'\|.$$

The proof is in Appendix C.

4.2   A.S. SUPPORT STABILITY OF SGD

We now work towards the a.s. support stability of SGD. First we state a theorem that bounds the replace-one error of SGD upto a certain number of epochs.

Theorem 7. *Suppose we are given a labelled data set $\mathcal{Z}$ and probability distribution D defined over it. Suppose we have a parametrized loss function $f : \mathbb{R}^n \times \mathcal{Z} \to \mathbb{R}$ that is a.s. L_l-LPL and a.s. K_l-LPS w.r.t D. Suppose for some $T > 0$ multiple of m, for each $i \in [m]$ we run Stochastic Gradient Descent on the f for T/m epochs using a training set S of size m chosen i.i.d. according to D with random choices r and, parallely, with the same set of random choices r, on a set S^i wherein the i-th data point z_i of S has been replaced by another data point z_i' chosen from D independent of all other random choices and with a decreasing learning rate of $\alpha_t \le \frac{\alpha_0}{t}$. Then, if for all $t > 0$, we denote by $\boldsymbol{w}_t$ and $\boldsymbol{w}_t'$ the parameter vectors obtained while training with S and S^i respectively, we have that with probability 1*
$$E_r\left[f(\boldsymbol{w}_T, z) - f(\boldsymbol{w}_T', z)\right] \le \frac{L_g^2}{K_g} F(T/m) \cdot \frac{(1 + \alpha_0 K_g)}{m^{1 - 2\alpha_0 K_g}},$$
where $F(T/m) := (T/m)^{2\alpha_0 K_g} \cdot 2^{T/m}$, and L_g and K_g are the "global" Lipschitz and "global" smoothness constants we get from Lemma 6.

The proof of Theorem 7 is in Appendix C. Let us look at its consequences for the generalization error of learning algorithms via this corollary written in plain English:

Corollary 8. *For a learning algorithm that is symmetric in distribution and trained as described in the statement of Theorem 7 with $\alpha_0 < 1/(4K_g)$, under the condition that L_g is a constant w.r.t. m, there is a constant $c > 0$ that depends on α_0 and K_g such that if it is trained for at most $c \log m$ epochs the expectation of the generalization error of the algorithm taken over the random choices of the algorithm tends to 0 as $m \to \infty$ with probability at least $1 - 1/m$ (over the choice of the training set.)*

Proof. It is easy to check that with the conditions given in the statement of Corollary 8 the learning algorithm has a.s. support stability β where β is $o(1/\sqrt{m})$. We can therefore apply Theorem 3 with $\delta = 1/m$ to get the result. □

If we compare this corollary with Theorem 4 of Kuzborskij & Lampert (2018), we note that their analysis requires an α_0 that decreases with m whereas our analysis allows for a value of α_0 that is constant. Importantly, their analysis bounds generalization error only up to the end of a single epoch whereas we can bound the error well beyond that. Kuzborskij & Lampert (2018) also require the Hessian to have a bounded Lipschitz constant, i.e., the third derivative of the loss function has to be bounded. We do not need this constraint.

5 NEURAL NETWORKS WITH ReLU ACTIVATION

The main result of this section presents the conditions required for low generalization error for Neural Networks with ReLU activation:

Theorem 9. *Suppose we have an unknown distribution D over $\mathbb{R}^d \times \mathcal{Y}$ such that $\mathcal{Y}$ is finite or countable and, for each $y \in \mathcal{Y}$, the set $\{x \in \mathbb{R}^d : Pr_D\{\mathsf{lab}(x) = y\} > 0\}$ is a countable union of products of intervals in some subset of $\mathbb{R}^d$. Suppose further that we have a learning algorithm $A_{S,r}$ that is a fully connected Neural Network with a single output and ReLU activations trained using Stochastic Gradient Descent (whose random choices are represented by the string r) with training set S drawn from D with learning rate α_0/t and a doubly differentiable loss function with bounded first and second derivatives. If the number of parameters of the Neural Network is a constant w.r.t. $|S| = m$ then there are constants (w.r.t. m) $c_1, c_2 > 0$ depending only on the size and structure of the Neural Network such that α_0 can be chosen to be at most c_1 and if the training is conducted for at most $c_2 \log m$ epochs, then*

$$\lim_{m \to \infty} E_r\left[|R(A, S, r) - R_e(A, S, r)|\right] \to 0,$$

with probability at least $1 - 1/m$, provided the data points of S and the parameter values explored by SGD are bounded.

We will now establish that the theory of a.s. support stability applies to NNs under the conditions specified.

5.1 PRELIMINARIES

5.1.1 NEURAL NETWORK TERMINOLOGY

Neural networks provide a family of parametrized functions of the form we have discussed in Section 4. The parameter vector $\boldsymbol{w} \in \mathbb{R}^n$ is applied over a network structure with layers. In this case we specify $\mathcal{Z}$ to be $\mathbb{R}^d \times \mathbb{R}$, i.e., the data points are from $\mathbb{R}^d$ and the label is from $\mathbb{R}$, i.e., the NN has a single output. We will denote the depth of the network by H. The layers will be numbered 0 to H with layer 0 being the *input layer*. The number of neurons in layer i will be k_i. For this discussion we assume a fully connected network. We will denote by $w_{j,k}^i$ the weight of the edge from the j neuron of the ith layer to the kth neuron of the $i + 1$st layer. For the NN with parameters $\boldsymbol{w}$ at a point $\boldsymbol{x} \in \mathbb{R}^d$ we will denote the input into the jth neuron of the ith layer by $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x})$ and its output by $\mathsf{out}_{i,j}(\boldsymbol{w}, \boldsymbol{x})$. These will be equal only if the former is non-negative. Further, we will assume that all neurons in all layers of the network except the input layer and the output layer have ReLU activation applied to them. In case the output of a node is 0 due to ReLU activation we will say the ReLU gate is *closed* otherwise we will say it is *open*. The label output by the network will be $\mathsf{out}_{H,1} = \mathsf{out}(\boldsymbol{w}, \boldsymbol{x})$. For each of exposition we will assume that $\mathsf{out}(\boldsymbol{w}, \boldsymbol{x}) = 1$ if $\mathsf{in}(\boldsymbol{w}, \boldsymbol{x}) > 0$ and 0 otherwise, i.e., there are only two labels in $\mathcal{Y}$.

5.1.2 MULTIVARIATE POLYNOMIALS ASSOCIATED WITH A NEURAL NETWORK

Given a set of indeterminates $\boldsymbol{x} = x_1, \ldots, x_l$, let $\mathcal{P}(\boldsymbol{x})$ be the set of multivariate polynomials on $x_1, \ldots, x_l$ with real coefficients. For any polynomial $p(\boldsymbol{x})$, $i_1, \ldots, i_q \in [l]$ and any $\alpha_1, \ldots, \alpha_q \in \mathbb{R}$ for some $q \le l$, we will denote by $p(\boldsymbol{x})\left\{x_{i_j} = \alpha_j : j \in [q]\right\}$ the polynomial in $\mathcal{P}(\boldsymbol{x} \setminus \{x_{i_1}, \ldots, x_{i_q}\})$ that is obtained by setting all occurences of x_{i_j} to α_j in $p(\boldsymbol{x})$. In particular, $p(\boldsymbol{x})\{x_i = 0\}$ is the polynomial

$p(\boldsymbol{x})$ with all monomials containing x_i removed, and $p(\boldsymbol{x})\{x_{i=1}\}$ retains all the monomials of $p(\boldsymbol{x})$ but those monomials that contain x_i appear without the term x_i.

Returning to NNs, let us consider two sets of indeterminates: $\boldsymbol{x} = \{x_i : i \in [d]\}$ and $\boldsymbol{w} = \{w_{j,k}^{(i)} : 0 \le i < H, 1 \le j \le k_i, 1 \le k \le k_{i+1}\}$ and $k_0 = d$. For a fully connected NN defined in Sec. 5.1.1 we will say that it has the following polynomial associated with it:

$$\phi(\boldsymbol{w}, \boldsymbol{x}) = \sum_{j_0=1}^{k_0} \sum_{j_1=1}^{k_1} \cdots \sum_{j_{H-1}=1}^{k_{H-1}} x_{j_0} w_{j_0,j_1}^{(0)} w_{j_1,j_2}^{(1)} \cdots w_{j_{H-1},1}^{(H-1)}.$$

Note that the output layer has only one neuron. We will refer to this as the *base polynomial* of the NN. The base polynomial associated with the jth neuron in layer i can be derived from the base polynomial of the network as follow

$$\phi_{i,j}(\boldsymbol{w}, \boldsymbol{x}) = \frac{\phi(\boldsymbol{w}, \boldsymbol{x})\left\{w_{l_1,l_2}^{(i)}=0, w_{l_4,l_5}^{(l_3)}=1:l_1\in[k_i]\setminus\{j\},l_2\in[k_{i+1}],l_3>i,l_4\in[k_{l_3}],l_5\in[k_{l_3+1}]\right\}}{\prod_{p=i+1}^{H} k_i}.$$

Due to ReLU activations varying at different points there is no single polynomial that captures the output of the NN everywhere in $\mathbb{R}^n \times \mathbb{R}^d$. However, the following observation shows a way of defining polynomials that describe the output over certain subsets of the space.

Observation 10. *Given $\boldsymbol{w} \in \mathbb{R}^n$ and $\boldsymbol{x} \ne (0, \ldots, 0) \in \mathbb{R}^d$, $i \in [H]$, and $j \in [k_i]$ such that* $\mathsf{in}_{l_1,l_2}(\boldsymbol{w}, \boldsymbol{x}) \ne 0$ *for all $1 \le l_1 \le i$ and all $1 \le l_2 \le k_{l_1}$, there is a $\varepsilon > 0$ depending on $\boldsymbol{w}, \boldsymbol{x}$ such that, for all $\boldsymbol{w}'$ with $\|\boldsymbol{w} - \boldsymbol{w}'\| < \varepsilon$,*
$$\mathsf{in}_{i,j}(\boldsymbol{w}', \boldsymbol{x}) = \mathsf{out}_{i,j}(\boldsymbol{w}', \boldsymbol{x}) = \phi_{i,j}(\boldsymbol{w}', \boldsymbol{x})\left\{w_{l_2,l_3}^{(l_1)}=0:l_1<i,l_2\in[k_{l_1}],l_3\in[k_{l_2+1}] \text{ and } \mathsf{in}_{l_1,l_2}(\boldsymbol{w},\boldsymbol{x})<0\right\}.$$

Proof. Since $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x})$ is strictly separated from 0 and there are only a finite number of neurons in the network, there must be a ε small enough for which all open ReLU gates remain open and all closed gates remain closed. It is easy to verify that the description of $\mathsf{out}(\boldsymbol{w}', \boldsymbol{x})$ given above is correct since it simply sets to 0 all the variables corresponding to weights originating in neurons whose ReLU gate is closed. □

5.2 Support Stability of Neural Networks with ReLU Activation

5.2.1 Smoothness Properties of Neural Networks with ReLU

We are now ready to state the main theorem of this section.

Theorem 11. *For every $\boldsymbol{w} \in \mathbb{R}^n$, a doubly differentiable loss function, $\ell : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$, applied to the output of a NN with ReLU activation is locally parameter-Lipschitz and locally parameter-smooth for all $\boldsymbol{x} \in \mathbb{R}^d$ except for a set of measure 0.*

The proof of this theorem is based on a idea of Milne (2019) and can be found in Appendix D.

Theorem 11 begs the question: How large are these Lipschitz and smoothness constants? We provide some general bounds that can be improved for specific architectures:

Proposition 12. *Suppose we have a fully connected NN with ReLU activation at the inner nodes that has depth $H + 1$ and at most K neurons in each layer. Then, if the coordinates of the parameter vectors have ℓ_∞-norm bounded by $B_{\boldsymbol{w}}$ and the data points have ℓ_∞-norm bounded by $B_{\boldsymbol{x}}$ then we have that*

1. *$L_g \le \sqrt{n} K^H |B_{\boldsymbol{x}}||B_{\boldsymbol{w}}|^{H-1}$, and*

2. *$K_g \le n K^H |B_{\boldsymbol{x}}||B_{\boldsymbol{w}}|^{H-2}$.*

The proof of the proposition is in Appendix D. We note that a result similar to Proposition 12(1) was presented by Szegedy et al. (2013) but in the context of the Lipschitz property over the data point not the parameter. We are now ready to prove our main theorem.

Proof of Theorem 9. Theorem 11 tells us that a NN with ReLU activations is locally parameter-Lipschitz and locally parameter-smooth. From Proposition 12 we see that the boundedness of the

first and second derivatives of the loss function and the boundedness of the parameters and data points ensures that the Lipschitz constants and smoothness constants associated with the NN's training are bounded w.r.t. m. With all these in place we have can apply Corollary 8 to get the result. The bound on the smoothness constant also tells us that the constraint on α_0 needed to apply Corollary 8 is not too stringent, i.e., α_0 can be chosen to be a value that is a constant w.r.t. m. □

5.3 DISCUSSION

Several of the constraints we have placed above are only for ease of exposition. We discuss which of these constraints can be removed.

• *Removing the fully connectedness constraint.* Although Theorem 9 is stated for a fully connected network, it can be applied to networks like CNNs which have partially connected convolutional layers with intermediate pooling and normalization layers (e.g., LeNet, AlexNet, etc.). In such cases the symmetry in distribution condition required for Theorem 3 holds as long as the training set is chosen i.i.d. from the unknown distribution.

• *Adding regularization terms to the loss function.* Several popular regularizers, the ℓ_2 regularizer being a prominent example, are doubly differentiable and therefore Theorem 9 can be applied when such regularizers are used along with a doubly differentiable loss function.

• *Activation functions apart from ReLU.* We present a comprehensive treatment of ReLU activation but that does not mean that our results are restricted to this kind of activation. Non-linearities like max-pool can also be handled in our framework by proving that, like with ReLU, the points of discontinuity of such a non-linearity also lie in a set of Lebesgue measure 0.

• *The case of multiple outputs* Although we state the Theorem 9 for the case of a NN with a single output, it is not difficult to extend the technique to cover the case of multiple outputs.

What about overparametrization? The question put by Zhang et al. (2017) as rearticulated by various authors, including Chatterjee & Zielinski (2022), can be put as follows: How do we speak theoretically about generalization in the overparametrized regime where NNs often generalize well even though they have enough parameters to overfit? Chatterjee & Zielinski (2022) make the point that some property of the training set is at play here, a claim more specific than the general claim that there is some property of the data distribution that has been missed. These are related, of course, but it is our feeling that a more fine-grained analysis of the growth in the replace-one error is required. We feel that it is possible look at the local Lipschitz and local smoothness constants in greater detail taking both network structure and input point values in mind. We hope that the polynomial characterization of the NN presented in Section 5.1.2 will help this process. We conjecture that it may be able to show that for certain distributions the constants actually improve (decrease) as the training proceeds, and this could lead to a proof of a.s. support stability in these cases.

6 EXPERIMENTS

In this section we will experimentally show that the Parameter Lipschitz and Parameter Smoothness constants that we reasoned with are indeed bounded, and that the theoretical upper bound that we derived for the generalization error of a neural network holds in practice.

Setup. For our experiments we use *MNIST* and *FashionMNIST* datasets. In both datasets, we randomly selected $20,000$ as training and $1,000$ as test points. All experiments were conducted using a fully connected feed forward neural network with single hidden layer and ReLU activation. We train the model using SGD (batch size = 1), with cross-entropy loss, starting with randomly initialized weights. As suggested in our analysis we use a decreasing learning rate $\alpha_t = \frac{\alpha_0}{t}$. In each epoch we consider a random permutation of the training set. L_g and K_g are computed by calculating the norm of gradients and Hessian across the training steps and taking their max.

Experiment 1. Our first experiment is aimed towards establishing that the "global" Lipschitz (L_g) and smoothness (K_g) values estimated using local values at each step are bounded. Figure 1 summarizes the results of these experiments over MNIST and FashionMNIST datasets ($\alpha_0 = 0.001$). The plots contain the maximum of the local parameter Lipschitz and smoothness values obtained after running each experiment 10 times with random weight initialization.

These results support our Theorem 11, since the upper bound values quickly stabilize and do not grow with the size of training set in both datasets. Similarly, the bounded smoothness constant supports our constraint on the learning rate, $\alpha_0 \leq \frac{1}{4K_g}$. We find L_g to be 8.1174 (MNIST) & 12.5737 (FashionMNIST), and K_g to be 58.185 (MNIST) and 102.7096 (FashionMNIST).

Experiment 2. We now turn our attention to the experiment to support our main result, i.e., the empirical generalization error estimated using validation set is upper bounded by our theoretical upper bound. We first split each dataset in 20:1 ratio into training and validation set, and train the model at varying size of training set. We empirically compute the generalization error at each training set size using the validation set. Figure 2 compares this empirical generalization error (in red) with the theoretical upper bound (in blue)From these results, we can see that our bound decreases along with the generalization error thus empirically validating our reasoning.

(a) MNIST

(b) FashionMNIST

7 CONCLUSION

We have devised a theoretical framework in this paper for using algorithmic stability to prove generalization bounds for NNs with ReLU nonlinearities. We feel that it is possible to prove stronger results in this framework than the ones we have presented here, and more widely applicable ones. Immediate lines of research that suggest themselves are to apply our methods for CNNs and to investigate what other architectures can be approached with our method. It would be particularly interesting to see if there is some analog of our polynomial characterisation for GNNs. Another clear challenge is the overparametrized case. We have not been able to tackle it but we feel a more fine-grained analysis in our framework may possibly lead to such a result.

Figure 1: Maximum of the Lipschitz and smoothness constants at every $p\ (=\ 20)$ interval of updates of SGD (we plot both the running average and the highest value found so far). Notice that these constants have a clear upper bound throughout the training process.

On another note, it is possible to define a version of a.s. support stability where we require a high probability bound on the replace-one error over *both* the random choices of the algorithm and the training set. It is possible to use this stronger definition to prove a version of Theorem 3 with high probability bounds on both the random choices of the algorithm and the training set. In this paper we did not present this result because we were not able to show that Neural Networks trained with SGD have this stronger property. However, it is our belief that such a result can be proved, perhaps if some information of the second moment of the Lipschitz and smoothness constants are known.

(a) MNIST   (b) FashionMNIST

Figure 2: Comparison of empirical generalization error (red) vs. theoretical upper bound (blue) with varying training set size for different datasets.

REFERENCES

Olivier Bousquet and André Elisseeff. Stability and Generalization. *J. Mach. Learn. Res.*, 2(Mar):
499–526, 2002.

Olivier Bousquet, Yegor Klochkov, and Nikita Zhivotovskiy. Sharper bounds for uniformly stable
algorithms. *Proc. Machine Learning Research*, 125:1–17, 2020.

Zachary Charles and Dimitris Papailiopoulos. Stability and generalization of learning algorithms
that converge to global optima. In Jennifer Dy and Andreas Krause (eds.), *Proceedings of the 35th
International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pp. 745–754. PMLR, 10–15 Jul 2018. URL https://proceedings.mlr.
press/v80/charles18a.html.

Satrajit Chatterjee and Piotr Zielinski. On the generalization mystery in Deep Learning.
arXiv:2203.10036, 2022.

Vitaly Feldman and Jan Vondrak. Generalization bounds for uniformly stable algorithms. In *Advances in Neural Information Processing Systems*, 2018.

Vitaly Feldman and Jan Vondrak. High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. In *Proceedings of the Thirty-Second Conference on Learning
Theory*, 2019.

Moritz Hardt, Ben Recht, and Yoram Singer. Train faster, generalize better: Stability of stochastic gradient descent. In Maria Florina Balcan and Kilian Q. Weinberger (eds.), *Proceedings of
The 33rd International Conference on Machine Learning*, volume 48 of *Proceedings of Machine
Learning Research*, pp. 1225–1234, New York, New York, USA, 20–22 Jun 2016. PMLR. URL
https://proceedings.mlr.press/v48/hardt16.html.

Geoffrey Hinton, Li Deng, Dong Yu, George E. Dahl, Abdel-rahman Mohamed, Navdeep Jaitly,
Andrew Senior, Vincent Vanhoucke, Patrick Nguyen, Tara N. Sainath, and Brian Kingsbury. Deep
neural networks for acoustic modeling in speech recognition: The shared views of four research
groups. *IEEE Signal Processing Magazine*, 29(6):82–97, 2012. doi: 10.1109/MSP.2012.2205597.

Pengzhan Jin, Lu Lu, Yifa Tang, and George Em Karniadakis. Quantifying the generalization error in
deep learning in terms of data distribution and neural network smoothness. *Neural Networks*, 130:
85–99, 2020. ISSN 0893-6080. doi: https://doi.org/10.1016/j.neunet.2020.06.024. URL https:
//www.sciencedirect.com/science/article/pii/S0893608020302392.

Kenji Kawaguchi, Leslie Pack Kaelbling, and Yoshua Bengio. Generalization in deep learning.
CoRR, abs/1710.05468, 2017.

M. Kearns and D. Ron. Algorithmic stability and sanity-check bounds for leave-one-out cross-
validation. *Neural Computation*, 11(6):1427–1453, 1999.

Yegor Klochkov and Nikita Zhivotovskiy. Stability and deviation optimal risk bounds with convergence rate o(1/n). In *Advances in Neural Information Processing Systems*, 2021.

Alex Krizhevsky, Ilya Sutskever, and Geoffrey E Hinton. Imagenet classification with deep
convolutional neural networks. In F. Pereira, C.J. Burges, L. Bottou, and K.Q. Weinberger (eds.), *Advances in Neural Information Processing Systems*, volume 25. Curran Associates, Inc., 2012. URL https://proceedings.neurips.cc/paper/2012/file/
c399862d3b9d6b76c8436e924a68c45b-Paper.pdf.

Ilja Kuzborskij and Christoph Lampert. Data-dependent stability of stochastic gradient descent.
In Jennifer Dy and Andreas Krause (eds.), *Proceedings of the 35th International Conference on Machine Learning*, volume 80 of *Proceedings of Machine Learning Research*, pp.
2815–2824. PMLR, 10–15 Jul 2018. URL https://proceedings.mlr.press/v80/
kuzborskij18a.html.

Yunwen Lei and Yiming Ying. Fine-grained analysis of stability and generalization for stochastic gradient descent. In Hal Daumé III and Aarti Singh (eds.), *Proceedings of the 37th International Conference on Machine Learning*, volume 119 of *Proceedings of Machine Learning Research*, pp. 5809–5819. PMLR, 13–18 Jul 2020. URL https://proceedings.mlr.press/v119/lei20c.html.

Tristan Milne. Piecewise strong convexity of neural networks. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett (eds.), *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019. URL https://proceedings.neurips.cc/paper/2019/file/b33128cb0089003ddfb5199e1b679652-Paper.pdf.

Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. arXiv:1312.6199 [cs.CV], 2013.

Vladimir Vapnik and Alexey Chervonenkis. *Theory of Pattern Recognition*. Nauka, Moscow, 1974.

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *International Conference on Learning Representations*, 2017. URL https://openreview.net/forum?id=Sy8gdB9xx.

Yikai Zhang, Wenjia Zhang, Sammy Bald, Vamsi Pritham Pingali, Chao Chen, and Mayank Goswami. Stability of SGD: Tightness analysis and improved bounds. In *The 38th Conference on Uncertainty in Artificial Intelligence*, 2022. URL https://openreview.net/forum?id=Sl-zmO8j5lq.

A MODIFICATION OF MCDIARMID'S THEOREM

We now state a modified version of McDiarmid's theorem that holds for η-almost β-Lipschitz functions.

Lemma 13. *Let $X_1, \ldots, X_m$ be i.i.d. random variables. If f is η-almost β-Lipschitz and takes values between 0 and M, then,*

$$Pr\{f(X_1, \ldots, X_m) - E[f(X_1, \ldots, X_m)] \geq \epsilon\} \leq \exp\left[\frac{-2\epsilon^2}{m\left((1-\eta)\beta + M\eta\right)^2}\right] + \eta.$$

For simplicity, we adopt the notation $A_{i,j} = A_i, \cdots, A_j$ (i.e. with subscript (i,j) we represent $j - i + 1$ variables) and A_i represents only one variable in the rest of the section. Before we prove Theorem 13, we state and prove the following lemma:

Lemma 14. *Let $f : \mathcal{Z}^m \to \mathbb{R}$ be a function such that $0 \leq f(\cdot) \leq M$. Define $V_i = E_{i+1,m}[f(X_{1,m})] - E_{i,m}[f(X_{1,m})]$, where $E_{i,j}[\cdot]$ denotes $E_{X_i, \cdots, X_j}[\cdot]$. Then, if f is η-almost β-Lipschitz, then with probability $1 - \eta$, $\forall i, V_i \leq (1 - \eta)\beta + \eta M$.*

Proof.
$$V_i = \mathrm{E}_{i+1,m}[f(X_{1,m})] - \mathrm{E}_{i,m}[f(X_{1,m})]$$
$$= \int \cdots \int_{y_{i+1}\cdots y_m} f(X_{1,i}, y_{i+1,m})d\mu(y_{i+1,m}) - \int \cdots \int_{y_i \cdots y_m} f(X_{1,i-1}, y_{i,m})d\mu(y_{i,m})$$
$$= \int \cdots \int_{y_{i+1}\cdots y_m} \left[f(X_{1,i}, y_{i+1,m})d\mu(y_{i+1,m}) - \int_{y_i} f(X_{1,i-1}, y_{i,m})d\mu(y_{i,m})d\mu(y_i)\right]d\mu(y_{i+1,m})$$

where μ is the measure associated with the random variables. Thus, V_i, a function of $X_{1,i}$, is an expectation over the random variables $y_{i+1,m}$. Thus, we can write $V_i = \mathrm{E}_{y_{i+1,m}}\left[\Phi(X_{1,i}, y_{i,m})|X_{1,i}\right]$.

Now, let's define two sets as follows:
$$S_i = \{x \in \mathcal{Z}^i : \exists y \in \mathcal{Z}^{m-i} : \Phi(x, y) \leq \beta\},$$
$$S'_x = \{y \in \mathcal{Z}^{m-i} : \Phi(x, y) \leq \beta\}.$$

Then, since f is η-almost β-Lipschitz, and renaming y_i as y_{i+m} in Φ, since they are *i.i.d.*, for any $x \in S_i$, we have

$$\mathrm{E}\left[\Phi(x,y)|x\right] \leq \sum_{y \in S'_x} \Pr\{y\}\beta + \left(1 - \sum_{y \in S'_x} \Pr\{y\}\right)M, \tag{1}$$

and for any $x \notin S_i$, we have

$$\mathrm{E}\left[\Phi(x,y)|x\right] \leq M.$$

We further note that $\sum_{x \in S_i} \Pr\{x\} \geq 1 - \eta$, and $\sum_{y \in S'_x} \Pr\{y\} \geq 1 - \eta$. Therefore, applying these bounds to Equation 1, we get, with probability at least $1 - \eta$,
$$V_i \leq (1 - \eta)\beta + \eta M.$$

$\square$

Proof of Theorem 13. We will use Azuma's inequality to prove the result. First we show that the sequence $V_1, \cdots, V_m$ is a martingale difference sequence with respect to $X_{1,m}$. To see this, we first notice that V_i is a function of $X_{1,i}$. Also, $\mathrm{E}\left[|V_i|\right]$ is finite since f is bounded. Finally,
$$\mathrm{E}\left[V_{i+1}|X_{1,i}\right] = \mathrm{E}_{i+1,m}\left[\mathrm{E}_{i+1,m}\left[f(X_{1,m})\right] - \mathrm{E}_{i,m}\left[f(X_{1,m})\right]|X_{1,i}\right]$$
$$= 0$$

Let $\boldsymbol{E}$ be the event that $|f(X_{1,m}) - \mathrm{E}\left[f(X_{1,m})\right]| \geq \epsilon$, and $\boldsymbol{F}$ the event that V_i is bounded by $\beta(1 - \eta) + \eta M$. We have,
$$\Pr\{\boldsymbol{E}\} \leq \Pr\{\boldsymbol{E}|\boldsymbol{F}\}\Pr\{\boldsymbol{F}\} + \Pr\{\overline{\boldsymbol{F}}\}. \tag{2}$$
Using Lemma 14, we have that with probability $1 - \eta$, for all i, V_i is bounded by $\beta(1 - \eta) + \eta M$.

Applying Azuma's inequality to $\Pr\{\boldsymbol{E}|\boldsymbol{F}\}$, we have the result. $\square$

B   PROOF OF CONVERGENCE OF GENERALIZATION ERROR FOR A.S. SUPPORT STABLE ALGORITHMS

Proof of Theorem 3. We will use Lemma 13 to prove the result. We will show that the η-almost β-support stability property implies that $\mathrm{E}_r\left[R(S,r) - R_e(S,r)\right]$, which is a function of random variable S, is η-almost β'-Lipschitz for a value of β' that depends on β, m and M. For the sake of brevity we will omit writing S in the following.

By taking expectations over r on both sides of the first claim of Lemma 7 from Bousquet & Elisseeff (2002) and flipping the order of expectations we have:
$$\mathrm{E}_S\left[\mathrm{E}_r\left[R(r) - R_e(r)\right]\right] \leq \mathrm{E}_{S,z'}\left[\mathrm{E}_r\left[|\ell(A_{S,r}(x_{z'}), y_{z'}) - \ell(A_{S^i,r}(x_{z'}), y_{z'})|\right]\right],$$
where the z' on the RHS is picked from D.

By the definition of η-almost β-support stability we can say that
$$\mathrm{E}_S\left[\mathrm{E}_r\left[|R(r) - R_e(r)|\right]\right] \leq (1 - \eta)\beta + \eta M. \tag{3}$$
We will now show that $\mathrm{E}_r\left[R(r) - R_e(r)\right]$ is η-almost β'-Lipschitz where $\beta' = 2\beta + M/m$. If we are able to show then, with (3) in hand, we can apply Lemma 13 to get

$$\Pr\left\{\mathrm{E}_r\left[R(r) - R_e(r)\right] - (1 - \eta)\beta - \eta M \geq \epsilon\right\} \leq \exp\left(\frac{-2\epsilon^2}{m(2\beta + \frac{M}{m})^2}\right) + \eta \tag{4}$$

Equating RHS of Equation 4 to δ and solving for ϵ we get
$$\epsilon = (2\beta m + M)\sqrt{\frac{\log(1/(\delta - \eta))}{2m}}.$$
Substituting this in Equation 4, we have the result.

So now we turn to showing the almost Lipschitz property for $\mathrm{E}_r\left[R(r) - R_e(r)\right]$. Denoting $R(S^i)$ as R^i and $R_e(S^i)$ as R_e^i we can say by linearity of expectation, the triangle inequality and the concavity of the absolute value function that
$$\left|\mathrm{E}_r\left[R(r) - R_e(r)\right] - \mathrm{E}_r\left[(R^i(r) - R_e^i(r))\right]\right| \leq \mathrm{E}_r\left[|R(r) - R^i(r)|\right]$$
$$+ \mathrm{E}_r\left[|R_e(r) - R_e^i(r)|\right] \tag{5}$$

If we bound the two terms on R.H.S with some probability then we can show $\mathrm{E}_r\left[R(S,r) - R_e(S,r)\right]$ is almost Lipschitz. Closely following the proof of Theorem 12 of Bousquet & Elisseeff (2002), we have that, with probability at least $1 - \eta$, the following two conditions hold $\forall i \in [m]$

$$\mathrm{E}_r\left[\left|R(r) - R^i(r)\right|\right] \le \beta \tag{6}$$

$$\mathrm{E}_r\left[\left|R_e(r) - R_e^i(r)\right|\right] \le \beta + \frac{M}{m} \tag{7}$$

For Eq. (6) since the risk $R(r)$ is an expectation of loss over $z \in D$ it is unchanged if we take the expectation over $z \in \mathrm{supp}\,(D)$ so using η-almost β-support stability we get the result. For (7), we have with probability at least $1 - \eta$

$$\mathrm{E}_r\left[\left|R_e(r) - R_e^i(r)\right|\right] = \frac{1}{m}\sum_{\substack{j=1 \\ j \neq i}}^{m}\mathrm{E}_r\left[\left|\ell(A_{S,r}(x_{z_j}), y_{z_j}) - \ell(A_{S^i,r}(x_{z_j}), y_{z_j})\right|\right]$$

$$+ \frac{1}{m}\mathrm{E}_r\left[\left|\ell(A_{S,r}(x_{z_i}), y_{z_i}) - \ell(A_{S^i,r}(x_{z'_i}), y_{z'_i})\right|\right]$$

$$\le \beta + \frac{M}{m},$$

In the second step we can use η-almost β-support stability because the points of S are picked from $\mathrm{supp}\,(D)$. Equations (6) and (7) along with (5) give us that $\mathrm{E}_r\left[R(r) - R_e(r)\right]$ is η-almost $2\beta + \frac{M}{m}$-Lipschitz. $\qquad\square$

C   A.S. SUPPORT STABILITY OF SGD PROVED

Proof of Lemma 6. If f is locally Lipschitz at $\boldsymbol{w} \in A$, there is an $\varepsilon_{\boldsymbol{w}} > 0$ and an $L_{\boldsymbol{w}} > 0$ such that for all $\boldsymbol{w}' \in \mathbb{R}^n$ with $\|\boldsymbol{w} - \boldsymbol{w}'\| \le \varepsilon_{\boldsymbol{w}}$, $|f(\boldsymbol{w}) - f(\boldsymbol{w}')| \le L_{\boldsymbol{w}}\|\boldsymbol{w} - \boldsymbol{w}'\|$. So, let us turn our attention to those $\boldsymbol{w}' \in A$ that lie outside the ball of radius $\varepsilon_{\boldsymbol{w}}$ around $\boldsymbol{w}$. Note that for such a $\boldsymbol{w}'$, if $B > 0$ is the bound on f, we have that

$$\frac{|f(\boldsymbol{w}) - f(\boldsymbol{w}')|}{\|\boldsymbol{w} - \boldsymbol{w}'\|} \le \frac{2B}{\varepsilon_{\boldsymbol{w}}}.$$

Therefore the "global" Lipschitz constant for f within A is $\max\{L_{\boldsymbol{w}}, 2B/\varepsilon_{\boldsymbol{w}} : \boldsymbol{w} \in A\}$ which is bounded since A is finite. This proves the first part of the lemma. The second part follows similarly. $\qquad\square$

Proof of Theorem 7. For some $i \in [m]$ we couple the trajectory of SGD on S and S^i where $z_i \in S$ has been replaced with z'_i. Our random string r in this case is a random choice of an initial parameter vector, $\boldsymbol{w}_0$, and a random set of T/m i.i.d permutations $\pi_0, \ldots, \pi_{T/m-1}$ of $[m]$ chosen uniformly at random. We use these random choices for training both the algorithms with S and S^i. For $0 \le j \le T/m - 1$, we denote $\pi_j^{-1}(i)$ by I_j, i.e., I_j is the (random) position where the ith training point is encountered in the jth training epoch. The key quantity we will track through the coupled training process will be

$$\delta_t = \|\mathrm{E}_r\left[\boldsymbol{w}_t - \boldsymbol{w}'_t\right]\|,$$

for $1 \le t \le T$. If we can show that δ_T is bounded by some quantity B almost surely, we can invoke the fact that f is a.s. L_l-LPL to say that $\|\mathrm{E}_r\left[f(\boldsymbol{w}_t, z) - f(\boldsymbol{w}'_t, z)\right]\| \le L_g B$ for all $z \in \mathrm{supp}\,(D)$, where L_g is the "global" Lipschitz constant we get from Lemma 6.

We argue differently for the first epoch and differently for later epochs. For the first epoch we note that for $t \le I_0$, $\delta_t = 0$ since SGD performs identical moves in both cases. At $t = I_0 + 1$
$$\delta_{I_0+1} = \|\boldsymbol{w}_{I_0} - \alpha_{I_0}\nabla f(\boldsymbol{w}_{I_0}, z_i) - (\boldsymbol{w}'_{I_0} - \alpha_{I_0}\nabla f(\boldsymbol{w}'_{I_0}, z'_i))\| = \alpha_{I_0}\|\nabla f(\boldsymbol{w}_{I_0}, z_i) - \nabla f(\boldsymbol{w}'_{I_0}, z'_i)\|,$$
where the second equality follows from the fact that $\boldsymbol{w}_{I_0} = \boldsymbol{w}'_{I_0}$ by the definition of I_0. Using Lemma 6 we can say that $\delta_{I_0+1} \le 2\alpha_{I_0}L_g$ almost surely. Now,
$$\delta_{I_0+2} \le \|\boldsymbol{w}_{I_0+1} - \boldsymbol{w}'_{I_0+1}\| + \alpha_{I_0+1}\|\nabla f(\boldsymbol{w}_{I_0+1}, z_i) - \nabla f(\boldsymbol{w}'_{I_0+1}, z'_i))\|.$$
Here although the parameter vectors $\boldsymbol{w}_{I_0+1}$ and $\boldsymbol{w}'_{I_0+1}$ are not the same, $z_{\pi_0(I_0+1)}$ and $z_{\pi_0(I_0+1)}$ are the same by the definition of I_0 (assuming that $I_0 \neq m$). Therefore we get that
$$\delta_{I_0+2} \le \delta_{I_0+1} + \alpha_{I_0+1}K_g\delta_{I_0+1}$$

with probability 1 since, from Lemma 6 we have that f has a "global" smoothness property for the entire set of at most $2T$ parameter vectors that will be encountered during the coupled training of S and S^i. Noting that a similar recursion can be applied all the way to the end of the first epoch, i.e. till $t = m$ we get

$$\delta_m \leq 2\alpha_{I_0} L_g \prod_{t=I_0+1}^{m} (1 + \alpha_t K_g) \leq 2\alpha_{I_0} L_g \exp\left\{ \sum_{t=I_0+1}^{m} \alpha_t K_g \right\}, \tag{8}$$

with probability 1. Moving on to the next epoch we note that we can make the argument above till the next point where the two training sequences differ, i.e., till the $m + I_1 + 1$st step. At this point we have,

$$\delta_{m+I_1+1} \leq \delta_{m+I_1} + \alpha_{m+I_1} \|\nabla f(\boldsymbol{w}_{m+I_1}, z_i) - \nabla f(\boldsymbol{w}'_{m+I_1}, z'_i))\|.$$

Since neither the parameter vector nor the training points are the same in the second term, we have no option but to use the almost "global" Lipschitz property to say that

$$\delta_{m+I_1+1} \leq \delta_{m+I_1} + \alpha_{m+I_1} 2L_g.$$

Since $\alpha_{m+I_1} < \alpha_{I_0}$, observing that our current bound for δ_{m+I_1} is larger than $\alpha_{m+I_1} 2L_g$. Therefore

$$\delta_{m+I_1+1} \leq 2\delta_{m+I_1}.$$

So, we see that in the second and subsequent epochs, for time step $jm + I_j + 1$, $1 \leq j < T/m$ we have the bound

$$\delta_{m+I_j+1} \leq 2\delta_{m+I_j},$$

and for all $t > m + I_1, t \neq I_1, \ldots, I_{T/m-1}$ we have, as before, by the smoothness property that

$$\delta_{t+1} \leq \delta_t (1 + \alpha_{t+1} K_g).$$

Therefore, we have that

$$\delta_T \leq 2\alpha_{I_0} L_g (2)^{\frac{T}{m}-1} \exp\left\{ \sum_{t=I_0+1}^{T} \alpha_t K_g \right\} \leq \alpha_0 L_g 2^{T/m} \left(\frac{T}{m} \right)^{2\alpha_0 K_g} \frac{m^{2\alpha_0 K_g}}{I_0^{1+\alpha_0 K_g}}. \tag{9}$$

where, in the first inequality for ease of calculation we have retained the terms of the form $(1 + \alpha_{I_j} K_g)$, $2 \leq j < T/m$ in the product on the right although we can ignore them. In the second inequality we have substituted $\alpha_t = \alpha_0/t$ and used the fact that $\log n \leq H_n \leq 2 \log n$ for the n-th Harmonic number H_n.

Finally, in order to compute $\mathrm{E}_r [\delta_T]$ we observe that, since π_0 is uniformly drawn from the set of permutations of $[m]$, I_0 is uniformly distributed on $[m]$. Summing up the last term of (9) over $I_0 \in [m]$ and dividing further by m we get the result. □

D   PROOFS FROM SECTION 5

Proof of Theorem 11. The idea behind this proof is due to Milne (2019) who used it for a different purpose. From Observation 10 it follows that if we have $\boldsymbol{x} \neq (0, \ldots, 0) \in \mathbb{R}^d$ such that $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x}) \neq 0$ for all $1 \leq i \leq H$ and all $1 \leq j \leq k_i$, then $\mathsf{out}(\boldsymbol{w}, \boldsymbol{x})$ is, in fact, just the polynomial $\phi(\boldsymbol{w}', \boldsymbol{x})$ within a small neighbourhood of $\boldsymbol{w}$. Therefore it is doubly differentiable. Since the loss function is also differentiable, we are done for all such values of $\boldsymbol{x}$.

So now let us consider the set of points $\boldsymbol{x}$ for which i is the smallest layer index such that $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x}) = 0$. In case there are two such indices, we break ties using the neuron index j. By Observation 10, in a neighbourhood of $\boldsymbol{w}$, $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x})$ is a polynomial in $\boldsymbol{w}$ and $\boldsymbol{x}$ for each $\boldsymbol{x}$.

Now, we consider two cases. In the first case, $\mathsf{out}_{i-1,j'}(\boldsymbol{w}, \boldsymbol{x}) = 0$ for all $j' \in [k_{i-1}]$, i.e., all the ReLU gates from the previous layers are closed because $\mathsf{in}_{i-1,j'}(\boldsymbol{w}, \boldsymbol{x}) < 0$ for all $j' \in [k_{i-1}]$. In this case $\mathsf{out}(\boldsymbol{w}', \boldsymbol{x}) = 0$ everywhere in the neighbourhood guaranteed by Observation 10 and therefore $\ell(\mathsf{out}(\boldsymbol{w}', \boldsymbol{x}), \mathsf{lab}(\boldsymbol{x}))$ is doubly differentiable in the parameter space at $\boldsymbol{w}$ for all such $\boldsymbol{x}$, where we assume that each data point has a label $\mathsf{lab}(\boldsymbol{x}) \in \{0, 1\}$ associated with it. We note that this argument is easily portable to the case of a more general label set $\mathcal{Y}$ with the property described in the statement of Theorem 9 since $\mathsf{in}_{H,1}$ will be 0 everywhere in a small neighbourhood.

In the second case we have some $j' \in [k_{i-1}]$ such that $\mathsf{out}_{i-1,j'}(\boldsymbol{w}, \boldsymbol{x}) > 0$. Let $C_{i,j} \subseteq \mathbb{R}^d$ be those $\boldsymbol{x}$ for which this case holds. $C_{i,j}$ contains the solutions to $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x}) = 0$. Since we are working with a specific value of $\boldsymbol{w}$, this simply becomes a polynomial in $\boldsymbol{x}$. In fact, inspecting the definition of base polynomials we note that when $\boldsymbol{w}$ is fixed $\mathsf{in}_{i,j}(\boldsymbol{w}, \boldsymbol{x})$ is simply a linear combination of

$x_1, \ldots, x_{\mathbb{R}}^d$. This implies that $C_{i,j}$ is a hyperplane in $\mathbb{R}^d$. We note that this argument can also be made of the output node under the condition on the labelset given in the statement of Theorem 9 because for $\text{in}_{H,1}(\boldsymbol{w}, \boldsymbol{x})$ to give a value that lies on the boundary between two sets with different labels for a given $\boldsymbol{w}$, $\boldsymbol{x}$ must be drawn from a set of Lebesgue measure 0.

Since the network size is finite the set of all possible values of $\boldsymbol{x}$ for which case 2 occurs, i.e., $\bigcup_{i \in [H], j \in [k_i]} C_{i,j}$ is a finite union of hyperplanes in $\mathbb{R}^d$ and therefore a set of Lebesgue measure 0. □

Proof of Proposition 12. Let us consider a coordinate $w_{\ell_1, \ell_2}^{(i)}$. Note that the number of monomials of $\phi(\boldsymbol{w}, \boldsymbol{x})$ it appears in is $k_0 \cdots k_{i-1} \cdot k_{i+1} \cdots k_H$. If we take the partial derivative of $\phi(\boldsymbol{w}, \boldsymbol{x})$ w.r.t $w_{\ell_1, \ell_2}^{(i)}$ each of these monomials has degree $H - 1$ in the indeterminates of $\boldsymbol{w}$ and degree 1 in the indeterminates of $\boldsymbol{x}$. Therefore we can say that

$$\left| \frac{\partial \phi(\boldsymbol{w}, \boldsymbol{x})}{\partial w_{\ell_1, \ell_2}^{(i)}} \right| \leq \frac{\prod_{j \in [H]} k_j}{k_i} \cdot |B_{\boldsymbol{x}}| |B_{\boldsymbol{w}}|^{H-1}.$$

Claim (1) of the proposition follows by bounding k_j by K for each j and taking the second norm.

For Claim (2) we use the fact that for a twice differentiable function the smoothness constant is bounded by the spectral norm of the Hessian. Since the degree of each $w_{\ell_1, \ell_2}^{(i)}$ is 1 in each monomial, the diagonal terms of the Hessian are 0. Additionally since each monomial of $\phi(\boldsymbol{w}, \boldsymbol{x})$ contains exactly one parameter from each layer we know that terms of the form $\partial \phi(\boldsymbol{w}, \boldsymbol{x}) / \partial w_{\ell_1, \ell_2}^{(i)} w_{\ell_3, \ell_4}^{(i)}$ are also 0. It is only when we take the partial derivatives with parameters from two different layers that we get a set of monomials with degree $H - 2$ in the indeterminates of $\boldsymbol{w}$ and degree 1 in the indeterminates of $\boldsymbol{x}$. Accordingly

$$\left| \frac{\partial \phi(\boldsymbol{w}, \boldsymbol{x})}{\partial w_{\ell_1, \ell_2}^{(i)} \partial w_{\ell_3, \ell_4}^{(j)}} \right| \leq \frac{\prod_{\ell_5 \in [H]} k_{\ell_5}}{k_i k_j} \cdot |B_{\boldsymbol{x}}| |B_{\boldsymbol{w}}|^{H-2}.$$

Since the diagonal terms are 0, by Gerschgorin's Circle Theorem the largest eigenvalue is bounded by the maximum of the sum of the absolute values of the entries of any row and the result follows. □