Nonlinear MCMC for Bayesian Machine Learning

Abstract

We explore the application of a nonlinear MCMC technique first introduced in [1] to problems in Bayesian machine learning. We provide a convergence guarantee in total variation that uses novel results for long-time convergence and large-particle ("propagation of chaos") convergence. We apply this nonlinear MCMC technique to sampling problems including a Bayesian neural network on CIFAR10.

1 Introduction

Characterizing uncertainty is a fundamental problem in machine learning. It is often desirable for a machine learning model to provide a prediction *and* a measure of how "certain" the model is about that prediction. Having access to a robust measure of uncertainty becomes particularly important in real-world, high risk scenarios such as self-driving cars [2–4], medical diagnosis [5, 6], and classifying harmful text [7].

However, despite the need for uncertainty in machine learning predictions, it is well known that traditional ML training, i.e. based on optimizing an objective function, frequently does not provide robust uncertainty measures [8], yielding overconfident predictions for popular neural networks such as ResNets [9]. [1] An appealing alternative to the traditional optimization paradigm for ML is the Bayesian probabilistic framework, due to its relatively simple formulation and extensive theoretical grounding; see for example [10].

In the probabilistic perspective of machine learning, one combines a prior $P(\theta)$ over the parameter space $\theta \in \Theta$ and a likelihood of the data given model parameters $P(\mathcal{D}|\theta)$ via Bayes' rule to obtain a posterior over the parameters $P(\theta|\mathcal{D}) \propto P(\mathcal{D}|\theta)P(\theta)$. The "traditional" approach in machine learning is to *optimize* the posterior (or the likelihood) to obtain $\theta^* \in \arg\max P(\theta|\mathcal{D})$ and generate predictions via $P(y|x, \mathcal{D}) = P(y|x, \theta^*)$. However, if we adopt the Bayesian approach, the posterior characterizes the uncertainty about the parameters of the model (i.e. epistemic uncertainty), which can propagate to uncertainty about a prediction by *integration*: $P(y|x, \mathcal{D}) = \int P(y|x, \theta)P(\theta|\mathcal{D})\mathrm{d}\theta$. This paper studies the problem of how to approximate this integration with samples from $P(\theta|\mathcal{D})$.

1.1 Contributions

- Our main contribution is the novel analysis of a modification of the general nonlinear Markov Chain Monte Carlo (MCMC) sampling method from [1] to obtain quantitative convergence guarantees in both the number of iterations and the number of samples.

- We apply the general results from above to determine the convergence of two specific nonlinear MCMC samplers.

- In experiments, we compare these nonlinear MCMC samplers to their linear counterparts, and find that nonlinear MCMC provides additional flexibility in designing sampling algorithms with as good, or better, performance than the linear variety.

---

[1] In Appendix C.2.3, we provide an experiment that demonstrates this effect.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

1.2 Background

Bayesian ML & MCMC. In Bayesian machine learning, the "computationally difficult" step is integration since the integral $\int P(y|x,\theta)P(\mathrm{d}\theta|\mathcal{D})$ is not analytically solvable except for rare cases. In practice, one typically uses a Monte Carlo approximation such as

$$\int P(y|x,\theta)P(\theta|\mathcal{D})\mathrm{d}\theta \approx \frac{1}{N}\sum_{i=1}^{N} P(y|x,\theta^i), \quad \text{where} \quad \theta^i \overset{iid}{\sim} P(\theta|\mathcal{D}).$$

where the expected error in this approximation is well known to converge to zero like $\mathcal{O}(1/\sqrt{N})$ by the Central Limit Theorem (CLT). There are various approaches to sampling from $P(\theta|\mathcal{D})$, but Markov chain Monte Carlo (MCMC) is perhaps the most widely used. The basic idea of MCMC is to use a Markov transition kernel $\mathcal{T}$ with stationary measure $P(\theta|\mathcal{D})$, and simulate a Markov chain $\theta_{n+1} \sim \mathcal{T}(\theta_n, \bullet)$ that converges rapidly to this stationary measure. Therefore we can estimate

$$\int P(y|x,\theta)P(\theta|\mathcal{D})\mathrm{d}\theta \approx \frac{1}{N}\sum_{i=1}^{N} P(y|x,\theta_\infty^i) \approx \frac{1}{N}\sum_{i=1}^{N} P(y|x,\theta_{n_{\mathrm{sim}}}^i)$$

where n_{sim} is some large number of simulation steps and $\{\theta_n^i\}_{n=0}^{\infty}$ are independent Markov chains governed by $\mathcal{T}$.

The basic problem is then to design an efficient transition kernel $\mathcal{T}$. There is a vast body of literature studying various choices of $\mathcal{T}$; some well known choices are the Metropolis-Hastings algorithm [11, 12], the Gibbs sampler [13], the Langevin algorithm [14–17], Metropolis-Adjusted Langevin [18, 19], and Hamiltonian Monte Carlo [20–24].

However there are various challenges in Bayesian ML that make applying these samplers difficult in practice. One challenge is that the posterior $P(\theta|\mathcal{D})$ can be highly multimodal [25, 26], which makes it difficult to ensure that the Markov chain θ_n explores all modes of the target distribution. One can combat this issue by employing auxiliary samplers that explore a more "tractable" variation of $P(\theta|\mathcal{D})$ [27, 28]. Other methods that empirically improve posterior sampling quality include tempering [29–32], RMSProp-style preconditioning [33] or adaptive MCMC algorithms [34, 35] such as the popular No U-Turn Sampler [36].

Nonlinear MCMC. Another class of powerful MCMC algorithms, which is less-studied in the context of Bayesian ML, arises from allowing the transition kernel $\mathcal{T}$ to depend on the distribution of the Markov chain as in $\theta_{n+1} \sim \mathcal{T}_{\mathrm{Distribution}(\theta_n)}(\theta_n, \bullet)$. This approach gives rise to so-called *nonlinear* MCMC since $\{\theta_n\}$ is no longer a true Markov chain. Nonlinear Markov theory is a rich area of research [37–42] and, in discrete-time, has strong connections to nonlinear filtering problems [43, 44], sequential Monte Carlo [45, 46], and nonlinear Feynman-Kac models [47]. One can replace $\mathrm{Distribution}(\theta_n)$, which is often intractable, with an empirical estimate $\mathrm{Distribution}(\theta_n) \approx \frac{1}{N}\sum_{i=1}^{N} \delta_{\theta_n^i}$ to obtain *interacting* particle MCMC (iMCMC, or iPMCMC) methods; see for example [48, 49, 1, 50, 51].

Our view is that nonlinear MCMC offers some appealing features that traditional linear MCMC lack. One such feature is the ability to leverage *global information* about the state space contained in $\mathrm{Distribution}(\theta_n^i)$ to improve exploration of the state space Θ — a central issue in Bayesian ML. Another feature is the increased flexibility of nonlinear MCMC algorithms, which can be leveraged to correct biases that are introduced by other design decisions in MCMC for Bayesian ML such as tempering. These features will be explored empirically in Section 4.

However, the theoretical analysis of nonlinear Markov MCMC presents an added difficulty in that the particles $\{\theta_n^1, \ldots, \theta_n^N\}$ of an interacting particle system are now *statistically dependent*. This means that, in addition to studying the large-time behaviour which is classical in MCMC [52, 53], one must study the large-particle behaviour separately to obtain Monte Carlo estimates since the CLT does not apply. One such large-particle behaviour is the propagation of chaos [41], which studies the tendency of interacting particles to become independent as the number of particles, N, increases; see [41]. We will need both of these elements — long-time and propagation of chaos behaviour — to properly characterize the convergence of nonlinear MCMC.

80 **Other Sampling Methods.** Finally, let us mention that MCMC is certainly not the only way to
81 obtain Monte Carlo sample estimates in Bayesian ML; some popular examples include MC dropout
82 [8], black-box variational inference [54], and normalizing flows [55, 56].

1.3 Common Notation

84 Let $\mathcal{P}(\mathbb{R}^d)$ be the set of probability measures on the meaurable space $(\mathbb{R}^d, \mathscr{B}(\mathbb{R}^d))$. If $\mu \in \mathcal{P}(\mathbb{R}^d)$
85 and $f \in \mathcal{B}_b(\mathbb{R}^d) := \{f : \mathbb{R}^d \to \mathbb{R} \mid f \text{ is bounded}\}$, we will denote $\mu(f) := \int f \mathrm{d}\mu$. If $K :$
86 $\mathbb{R}^d \times \mathscr{B}(\mathbb{R}^d) \to [0,1]$ is a Markov kernel[2] then we will denote $Kf(x) := \int f(y) K(x, \mathrm{d}y)$ and
87 $\mu K(\mathrm{d}y) := \int \mu(\mathrm{d}x) K(x, \mathrm{d}y)$. Finally, for $\overline{y} := \{y^1, \ldots, y^N\} \subset \mathbb{R}^d$, we will denote the empirical
88 measure of $\overline{y}$ as $m(\overline{y}) := \frac{1}{N} \sum_{i=1}^{N} \delta_{y^i}$.

2 Nonlinear MCMC

90 In this section, we outline the family of MCMC algorithms that will be used in rest of the work. We
91 will use general notation for simplicity but this difference is merely cosmetic; the "target distribution"
92 π in this section corresponds directly to $P(\theta|\mathcal{D})$ from the previous section.

2.1 Nonlinear Jump Interaction Markov Kernels

94 To specify a MCMC algorithm, we must specify the Markov transition kernel. The family of nonlinear
95 Markov kernels that we will be studying was introduced in [1] and is a mixture of a linear kernel,
96 denoted K, and a nonlinear jump-interaction kernel J_η indexed by a probability measure η, to obtain

$$K_\eta(x, \mathrm{d}y) := (1 - \varepsilon) K(x, \mathrm{d}y) + \varepsilon J_\eta(x, \mathrm{d}y) \tag{1}$$

97 where $\varepsilon \in]0, 1[$ is the mixture hyperparameter. The Markov kernel K_η will be the main object of
98 interest throughout this paper. We will give specific examples of J_η in Section 2.2. Furthermore, for
99 an expanded description of how our setup differs from [1], see Appendix A.1.

100 **Mean Field System.** Now we show how the kernel K_η can be used to construct a Markov chain.
101 Following [1], we use an auxiliary Markov chain $\{Y_n\}$ with transition kernel Q on the same state
102 space as K_η (i.e. $\mathbb{R}^d$) to obtain the nonlinear Markov chain $\{(Y_n, X_n)\}_{n=0}^{\infty}$ defined by

$$\begin{cases} Y_{n+1} & \sim Q(Y_n, \bullet) \\ \eta_{n+1} & := \text{Distribution}(Y_{n+1}) \quad Y_0 \sim \eta_0, \ X_0 \sim \mu_0 \\ X_{n+1} & \sim K_{\eta_{n+1}}(X_n, \bullet) \end{cases} \tag{2}$$

103 where μ_0, η_0 are the initial distributions and $\sim$ denotes "sample from". One should interpret this
104 as a sequence of steps where first we sample the auxiliary state Y_{n+1} from Q, then we obtain the
105 distribution of Y_{n+1} denoted η_{n+1}, and we use this distribution to index the primary kernel $K_{\eta_{n+1}}$
106 and obtain a sample X_{n+1}. We sample X_{n+1} with probability $(1 - \varepsilon)$ from the linear kernel K, and
107 with probability ε it will "jump" according to $J_{\eta_{n+1}}(X_n, \bullet)$. Because the Markov dynamics depend
108 on $\text{Distribution}(Y_n)$, we call this a "mean field system".

109 **Interacting Particle System.** One issue with the mean field system (2) is the fact that computing
110 $\text{Distribution}(Y_{n+1})$ is generally impossible except in special cases. Hence, to get a viable simulation
111 algorithm, we must approximate $\text{Distribution}(Y_n)$, and we do this by replacing $\text{Distribution}(Y_n)$
112 with its empirical measure estimated from a set of N samples $\overline{Y}_n := \{Y_n^1, \ldots, Y_n^N\}$ as follows:

$$\begin{cases} Y_{n+1}^i & \sim Q(Y_n^i, \bullet) \\ \eta_{n+1}^N & := m(\overline{Y}_{n+1}) \quad Y_0^i \overset{iid}{\sim} \eta_0, \ X_0^i \overset{iid}{\sim} \mu_0. \\ X_{n+1}^i & \sim K_{\eta_{n+1}^N}(X_n^i, \bullet) \end{cases} \tag{3}$$

2.2 Application to MCMC

114 Now we detail how to apply K_η and the Markov chains (2), and (3) to MCMC. In particular, we must
115 understand how to choose Q, K, J_η such that K_η will be invariant w.r.t. a target distribution π.

---

[2] i.e. $K(x, \bullet)$ is a probability measure $\forall x \in \mathbb{R}^d$ and $K(\bullet, A)$ is measurable $\forall A \in \mathscr{B}(\mathbb{R}^d)$

As is usually the case in probabilistic inference problems, we will assume that the target distribution π is known only up to a normalizing constant and that it has a density, also denoted π. We also make the simplifying assumption that Q has an invariant measure $\eta^\star$ (also with density denoted $\eta^\star$) i.e. $\eta^\star Q = \eta^\star$. This is not burdensome; in practice we can, and will, obtain Q from a *linear* MCMC algorithm for some choice of $\eta^\star$. In fact, being able to choose $\eta^\star$ is a powerful design parameter of our methods as we will see in Section 4. We will also assume that the linear kernel K is π-invariant, i.e. $\pi K = \pi$.

To see how we can ensure that π is K_η-invariant, consider the fact that we will design Q s.t. $\eta_n := \text{Distribution}(Y_n)$ converges to $\eta^\star$. This means we will eventually be sampling from the kernel $K_{\eta^\star}$ and we already have π-invariance of K. Therefore, if we arrange for $J_{\eta^\star}$ to be π-invariant, π will be invariant for $K_{\eta^\star}$ since

$$\pi K_{\eta^\star} = (1-\varepsilon)\pi K + \varepsilon \pi J_{\eta^\star} = (1-\varepsilon)\pi + \varepsilon\pi = \pi.$$

Intuitively, if the auxiliary chain converges to a steady state and the jumps in that steady state preserve π (and K preserves π), then so will $K_{\eta^\star}$. Now the remaining task is to design nonlinear interaction kernels J_η that will yield good performance. We detail two choices below.

Boltzmann-Gibbs Interaction.   The first choice of J_η we will investigate, from [1], relies on the Boltzmann-Gibbs transformation [47], which we now explain. Let $G : \mathbb{R}^d \to]0, \infty[$ be a potential function; then the Boltzmann-Gibbs (BG) transformation is a nonlinear mapping $\Psi_G : \mathcal{P}(\mathbb{R}^d) \to \mathcal{P}(\mathbb{R}^d)$ defined by

$$\Psi_G(\mu)(\mathrm{d}x) := \frac{G(x)}{\mu(G)}\mu(\mathrm{d}x) \quad \text{or equivalently} \quad \int f(x)\Psi_G(\mu)(\mathrm{d}x) := \int f(x)\frac{G(x)}{\mu(G)}\mu(\mathrm{d}x)$$

for any $f \in \mathcal{B}_b(\mathbb{R}^d)$ and whenever $\mu(G) \neq 0$. This transformation has many interesting properties and been extensively studied in [47] and related works.

To use the BG transformation in MCMC, we will assume that the densities π and $\eta^\star$ are positive[3] and make the choice that $G(x)$ will be the function

$$G(x) := \frac{\pi(x)}{\eta^\star(x)}. \tag{4}$$

With this choice, we get an interaction kernel $J_\eta^{BG}(x, \mathrm{d}y) := \Psi_G(\eta)(\mathrm{d}y)$. We can easily see that

$$\eta^\star(G) = \int \frac{\pi}{\eta^\star}\mathrm{d}\eta^\star = \int \mathrm{d}\pi = 1 \quad \text{and} \quad \Psi_G(\eta^\star)(\mathrm{d}x) = \frac{G(x)}{\eta^\star(G)}\eta^\star(\mathrm{d}x) = \frac{\pi(x)}{\eta^\star(x)}\eta^\star(\mathrm{d}x) = \pi(\mathrm{d}x),$$

i.e. Ψ_G is the multiplicative "change of measure" from $\eta^\star \to \pi$. Hence the first nonlinear Markov kernel we will investigate is

$$K_\eta^{BG}(x, \mathrm{d}y) := (1-\varepsilon)K(x, \mathrm{d}y) + \varepsilon\Psi_G(\eta)(\mathrm{d}y). \tag{5}$$

From the remarks above, is clear that π is $K_{\eta^\star}$-invariant.

Accept-Reject Interaction.   The second choice of jump interaction we will study, also introduced in [1], is a type of accept-reject interaction related to the Metropolis-Hastings algorithm. For the *same* choice of potential function G in (4), we can define the acceptance ratio

$$\alpha(x, y) := 1 \wedge \frac{G(y)}{G(x)} = 1 \wedge \frac{\pi(y)\eta^\star(x)}{\eta^\star(y)\pi(x)} \quad \text{and the quantity} \quad A_\eta(x) := \int \alpha(x, y)\eta(\mathrm{d}y)$$

for $\eta \in \mathcal{P}(\mathbb{R}^d)$. Hence we can define the accept-reject interaction kernel as[4]

$$J_\eta^{AR}(x, \mathrm{d}y) := \alpha(x, y)\eta(\mathrm{d}y) + (1 - A_\eta(x))\delta_x(\mathrm{d}y).$$

We can interpret this jump interaction as: starting in state x, we jump to a new state distributed according to $\eta(\mathrm{d}y)$ with probability $\alpha(x, y)$ (i.e. accept the proposed jump) and remain the in current state with probability $1 - A_\eta(x)$ (i.e. reject the proposed jump). This is a form of "adaptive Metropolis-Hastings" in which the proposal distribution evolves over time as the distribution of the auxiliary Markov chain. Hence we obtain the accept-reject nonlinear jump interaction kernel

$$K_\eta^{AR}(x, \mathrm{d}y) := (1-\varepsilon)K(x, \mathrm{d}y) + \varepsilon[\alpha(x, y)\eta(\mathrm{d}y) + (1 - A_\eta(x))\delta_x(\mathrm{d}y)]. \tag{6}$$

We note that π is also $J_{\eta^\star}^{AR}$-invariant; see Proposition 3 in Appendix G for a simple calculation.

---

[3] this can be relaxed to $\pi \ll \mu$ and $\mu \ll \pi$

[4] Given $f \in \mathcal{B}_b(\mathbb{R}^d)$, we can also write this as $J_\eta^{AR}f(x) = \int[f(y) - f(x)]\alpha(x, y)\eta(\mathrm{d}y) + f(x)$

152 **Simulation.** Let us note briefly that using both K_η^{BG} and K_η^{AR} in (3) produce interacting particle
153 systems that can, and will, be simulated. The simulation is relatively straightforward, see Appendix A
154 for pseudocode implementing the nonlinear MCMC algorithms we have now constructed.

3 Convergence Analysis

156 We will now study whether the nonlinear MCMC algorithms based on K_η from Section 2 — i.e.,
157 the interacting particle system (3) with the restrictions on K, Q, J_η from Section 2.2 — will actually
158 converge to the target distribution π. In other words, we would like to estimate $\|\mu_n^N - \pi\|$ for some
159 suitable notion of distance on $\mathcal{P}(\mathbb{R}^d)$, where $\mu_n^N := \text{Distribution}(X_n^1)$ is the distribution of a single
160 particle (it doesn't matter which particle, as the X_n^i are *exchangeable*).

161 The nonlinear nature of K_η makes the analysis more difficult than for a linear MCMC method. We
162 break the problem into two parts: one studying the convergence of the mean-field system (2) as the
163 number of steps $n \to \infty$, and one studying the convergence of the interacting particle system (3) to
164 the mean field system as the number of particles $N \to \infty$. This will allow us to apply the triangle
165 inequality as follows:

$$\|\mu_n^N - \pi\| \leq \underbrace{\|\mu_n^N - \mu_n\|}_{\text{large-particle convergence}} + \underbrace{\|\mu_n - \pi\|}_{\text{long-time convergence}}$$

166 where $\mu_n := \text{Distribution}(X_n)$ is the distribution of the mean-field system. Crucially, our analysis
167 of the large-particle limit is *uniform* in the number of steps n, which will allow us to establish bounds
168 above that hold as $n \to \infty$. The actual result is contained in Theorem 1.

169 While our analysis does not rely on heavy mathematical machinery, to state the full set of conditions
170 and results for long-time and large-particle convergence — each of which is a substantial result in its
171 own right — would occupy too much space in the main text. Instead, we will state the main result,
172 which is essentially a corollary of the long-time and large-particle analyses Theorems 2 and 3 in
173 Appendices E and F respectively, and below we will sketch the general arguments used in those
174 appendices. The proofs of the main results are in Appendix F.2. Note that our analysis, and the results
175 we obtain, are novel and not found in [1]; see Appendix D for an elaboration.

176 The following is stated in terms of the *total variation* metric, defined here for $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ as
177 $\|\mu - \nu\|_{tv} := \sup_{\|f\|_\infty \leq 1} |\mu(f) - \nu(f)|$ where $\| \bullet \|_\infty$ is the sup-norm on $\mathcal{B}_b(\mathbb{R}^d)$.

178 **Theorem 1.** *[Convergence of Nonlinear MCMC] Under suitable conditions on K_η and Q, there*
179 *exist fixed constants $C_1, C_2, C_3 > 0$, a function $\mathcal{R} : [0, \infty[\to [1, \infty[$, and $\rho > 0$ s.t.*

$$\|\mu_n^N - \pi\|_{tv} \leq C_1 \frac{1}{N} \mathcal{R}(1/N) + C_2 \rho^n + C_3 n \rho^n.$$

180 This result shows that, to control the approximation error $\|\mu_n^N - \pi\|_{tv}$, it does not necessarily suffice
181 to run the MCMC algorithm for a large number of steps n, since if $n \to \infty$ but $N < \infty$ then
182 $\|\mu_n^N - \pi\| \not\to 0$. However, this approximation cannot lead to arbitrarily bad results: Theorem 1
183 provides a quantitative upper bound on how much the MCMC algorithm can be biased. Note also that
184 this result uses total variation, which is a strong metric that represents a worst-case over *all* bounded
185 functions f (up to rescaling by $\|f\|_\infty$). It is entirely possible that, for many choices of practical f,
186 the approximation will be better as we will see empirically.

187 A corollary of Theorem 1 is that that we regain a Monte Carlo estimate for the interacting particle
188 system. This result is essentially due to [41] Theorem 2.2.

189 **Corollary 1.** *[Adapted from [41], Theorem 2.2] Suppose that Theorem 1 applies to K_η. Let*
190 $\overline{X}_n := \{X_n^1, \ldots, X_n^N\}$ *be from (3). Then for every $n \in \mathbb{N}$ and $f \in \mathcal{B}_b(\mathbb{R}^d)$ we have*

$$\lim_{N \to \infty} \mathbb{E}\left[\left\|\frac{1}{N} \sum_{i=1}^N f(X_n^i) - \mu_n(f)\right\|\right] = 0.$$

191 This corollary directly relates to the application of Bayesian ML we are interested in, where we would
192 have $f(\theta) = P(y|x, \theta)$.

3.1 Long-Time Bounds

There are two main ingredients in the general result on long-time convergence: ergodicity of K and Q, and Lipschitz regularity of the interaction kernel $\eta \mapsto J_\eta$. These, along with other technical conditions, produce an estimate of the form $\|\mu_n - \pi\| \leq C_2 \rho^n + C_3 n \rho^n$ where $\| \bullet \|$ is a weighted total variation norm. The full statement is in Theorem 2 of Appendix E.

Ergodicity of K and Q. A fundamental requirement of our results is that the *linear* building blocks of K_η must converge to their respective stationary measures in an appropriate metric. This type of result is now standard in the Markov chain literature, and we use a result from [57] for K and a result from [1] for Q. The former is actually able to ensure that K is a contraction on $\mathcal{P}(\mathbb{R}^d)$ w.r.t. a suitable weighted total variation; we use this feature repeatedly in our analysis.

Lipschitz Regularity of J_η. We also need that $\eta \mapsto J_\eta$ is *Lipschitz*-continuous w.r.t. a weighted total vartiation norm on Markov kernels (the Lipschitz constant does not have to be < 1). This regularity is used to translate the convergence of $\eta_n \to \eta^\star$ (as guaranteed by the ergodicity of Q) into convergence of $J_\eta \to J_{\eta^\star}$ with the Lipschitz estimate $\|J_{\eta_n} - J_{\eta^\star}\| \lesssim \|\eta_n - \eta^\star\|$. In Appendix G, we verify this analytically for J_η^{BG} and J_η^{AR}, see Lemma 5 and [1] Proposition 5.3.

3.2 Large-Particle Bounds

To study the large-particle behaviour, we would like to measure how close a group of $q \in \{1, \dots, N\}$ interacting particles $\{X_n^1, \dots, X_n^q\} \subset \{X_n^1, \dots, X_n^N\} =: \overline{X}_n$ from (3) is to being i.i.d. according to the mean-field measure μ_n. This analysis was pioneered in [41] under the name "propagation of chaos", and formalizes the intuition that, as $N \to \infty$, the influence of any individual particle $\to 0$.

To state this precisely, first note that if we had random variables $Z^i \overset{iid}{\sim} \eta \in \mathcal{P}(\mathbb{R}^d)$ then the joint distribution of $\overline{Z} := \{Z^1, \dots, Z^N\}$ would be $\eta^{\otimes N}$. Hence, for the interacting particles $\overline{X}_n$ at time n, we expect the distribution of $\{X_n^1, \dots, X_n^q\}$, denoted $\mu_n^{q,N}$, to get closer to the distribution of i.i.d. mean field particles from (2), denoted $\mu_n^{\otimes q}$, as $N \to \infty$. In other words, we expect $\|\mu_n^{q,N} - \mu_n^{\otimes q}\|_{tv} \to 0$ as $N \to \infty$. The full statement of this result is Theorem 3 of Appendix F.

The main condition in our propagation of chaos result is another type of regularity for $\eta \mapsto J_\eta$ which basically requires that, if one approximates a distribution $\eta \in \mathcal{P}(\mathbb{R}^d)$ by its empirical measure $m(\overline{Y})$ where $\overline{Y} := \{Y^1, \dots, Y^N\}$ and $Y^i \overset{iid}{\sim} \eta$, then $J_{m(\overline{Y})} \to J_\eta$ as $N \to \infty$. More precisely, there should be a nondecreasing function $\mathcal{R} : [0, \infty[\to [1, \infty[$ s.t.

$$|\mathbb{E}[J_{m(Y)}^{\otimes q} f(x)] - J_\eta^{\otimes q} f(x)| \lesssim \frac{q^2}{N} \mathcal{R}(q^2/N) \quad \forall x \in \mathbb{R}^d \text{ and } f \in \mathcal{B}_b(\mathbb{R}^d) \text{ with "oscillations"} \operatorname{osc}(f) \leq 1.$$

The expectation is taken over η, and "oscillations" are defined precisely in Appendix D. This inequality is analogous to the Lipschitz regularity from above since we can alternately write $\|\mu - \nu\|_{tv} = \sup\{|\mu(f) - \nu(f)| \mid f \in \mathcal{B}_b(\mathbb{R}^d), \operatorname{osc}(f) \leq 1\}$ [47].

3.3 Analysis of Specific Interaction Kernels

The main result Theorem 1 is in terms of conditions on a general K_η (i.e. general choices of K, Q, J_η). To apply this result to the samplers in Section 2, we must establish if these conditions hold for K_η^{BG} and K_η^{AR}. Fortunately, this can be done analytically; in Appendix G, we present conditions under which the Lipschitz regularity (see Lemma 5 and [1] Proposition 5.3) and large-particle regularity (see Corollaries 3 and 5) hold. These results, particularly for K_η^{BG}, are interesting and rely on novel techniques for controlling the various quantities, sometimes improving over previous methods. Due to space constraints, the results for K_η^{BG} and K_η^{AR} are in Corollaries 2 and 4 from Appendix G.

4 Experiments

In this section, we detail two experiments designed to explore how one might apply the nonlinear MCMC methods developed in the previous sections to Bayesian machine learning. Let us state

explicitly that our aim is *not* to achieve state-of-the-art with these experiments, nor do we claim that this method will necessarily lead to state-of-the-art results on a particular task. Rather, the aims of these experiments are: to demonstrate that nonlinear MCMC can be applied successfully to large-scale problems; compare linear vs nonlinear methods to understand what benefits and drawbacks nonlinear MCMC offers compared to linear MCMC; and to develop some recipes for choosing the various hyperparameters and samplers that determine a nonlinear MCMC method.

4.1 Two-Dimensional Toy Experiments

We use a toy setting of two-dimensional distributions to compare the relative benefits of linear and nonlinear MCMC. A benefit of this low-dimensional setting is that the multimodal toy distributions can be exactly sampled. This gives us the opportunity to quantitatively measure the quality of our samples via an unbiased estimator of the Maximum Mean Discrepancy (MMD) metric on $\mathcal{P}(\mathbb{R}^d)$ [58]. This approach stands in contrast to many previous works, which use very simple distributions (e.g. Gaussians) with analytically tractable statistics to measure quality. See Appendix C.1 for a brief overview of our methodology.

Setup. Our setup will compare the Metropolis Adjusted Langevin Algorithm (MALA) [18] (see Appendix B for an overview of MALA) with the nonlinear BG and AR samplers using MALA for the kernels K, Q in our nonlinear setup from Section 2. This will allow us to examine the effects of the nonlinearity in K_η^{AR} and K_η^{BG} while controlling for the type of sampler used and its hyperparameters.

The main difference between the linear and nonlinear algorithms, aside from the interaction itself, is the extra "design knob" to change in the form of the choice of the auxiliary density $\eta^\star$. We show how one can use $\eta^\star$ to incorporate additional insight to guide the sampling, such as regions of the state space to explore. In our experiments, we choose $\eta^\star$ to be a centered, 2-dimensional Gaussian with a large standard deviation ($\Sigma = 4I_2$ for the circular MoG and two-rings densities, and $\Sigma = 20I_2$ for the grid MoG). This conveys "coarse-grained" information of roughly where the support of the target density is located – in this case, a neighbourhood of $(0, 0)$. See Table 2 in Appendix C.1 for a full account of our experimental settings.

Results. From Figure 1, we see that having a simple auxiliary density with good coverage of the support of the target distribution is quite helpful. In all three examples, one or both of the nonlinear samplers outperformed the equivalent linear sampler in the empirical MMD metric. For the two most challenging densities, the "two rings" and "grid MoG" distributions, the improved exploration is evident, particularly for the BG interaction. However, the BG interaction also displays a substantially higher variance and can be less stable over time than the AR interaction. See Appendix C.1.3 for an analysis of the runtime of the algorithms.

4.2 CIFAR10

4.2.1 Setup

To examine the properties of the nonlinear sampler outside of a toy setting, we have also implemented a Bayesian neural network on the CIFAR10 dataset. We use a likelihood $P(y|x, \theta)$ parameterized by a ResNet-18 convolutional neural network [61] and a Gaussian prior $P(\theta)$ on the parameters of this neural network which are combined to form a posterior $P(\theta|\mathcal{D}_{\text{train}}) \propto P(\theta) \prod_{(x_i, y_i) \in \mathcal{D}_{\text{train}}} P(y_i|x_i, \theta)$. The goal is to sample θ^i, $i = 1, \ldots, N$ from this posterior. See Table 4 in Appendix C.2 for a full account of the experimental settings.

To deal with the fact that this sampling problem is very high-dimensional ($d \approx 11\text{M}$) and $|\mathcal{D}_{\text{train}}|$ is large ($|\mathcal{D}_{\text{train}}| = 60,000$) we use a variety of techniques:

1. We sample minibatches $\widehat{\mathcal{D}}$ of size 256 to obtain the surrogate target density $P(\theta|\widehat{\mathcal{D}})$ [17].
2. We use an RMSProp-like "preconditioned" Langevin algorithm, called RMS-Langevin or RMS-ULA, as in [33] for the auxiliary sampler Q; see Appendix B for details on this sampler. As shown in [33], this sampler is biased.
3. We use tempering, wherein we aim to sample from π or $\eta \propto P(\theta|\mathcal{D}_{\text{train}})^{1/\tau}$ where τ is a small number. This substantially improves mixing for hard-to-sample distributions such as $P(\theta|\mathcal{D}_{\text{train}})$ at the cost of bias since we are no longer sampling from the true posterior [32].

Target Density Ground Truth Linear Nonlinear (AR) Nonlinear (BG)

Figure 1: Visualizations of the 2d experiment. The first row is for the Circular Mixture of Gaussians (MoG) density [59] the second is for the Two Rings density [59], and the third is for the Grid Mixture of Gaussians density [60]. The right column shows the empirical MMD plotted over number of sampled steps, where the shaded region is ± 1 std with $N = 5$ trials. The leftmost four columns visualize the sampling problem for each distribution with $N = 2,000$ samples.

Using our nonlinear algorithm presents a novel opportunity to correct the bias introduced by tempering. For our experiments, we pick $\eta^\star \propto P(\theta|\mathcal{D}_{\text{train}})^{1/\tau}$ and $\pi = P(\theta|\mathcal{D}_{\text{train}})$. This means that the auxiliary chain Y_n explores a tempered version of the target, whereas the target chain X_n (in theory) explores the true target distribution. This is a novel strategy that is made possible by being able to select $\eta^\star$ independently of the target π. We study the case when π is tempered as well.

For our experiments, we use the RMS-Langevin sampler as the baseline, and we also use it for the auxiliary sampler Q. For the target sampler, it is not possible to use the RMS-Langevin algorithm because the smoothed square-gradient estimate is incompatible with the discontinuities (i.e. jumps) introduced by the nonlinear interaction. Instead, for the linear sampler K we use the unadjusted Langevin algorithm, ULA, [16] (see Appendix B). We also use the Boltzmann-Gibbs interaction since it is nonreversible, whereas the accept-reject interaction is reversible and this is not efficient for these high-dimensional settings (this is the same reason for using the ULA instead of MALA). We will investigate both test accuracy and calibration error [9] to assess performance.

Table 1: Results for CIFAR10 experiments. $\pm$ represents 1 standard deviation on $N = 5$ trials. The tempered results are using $\tau = 10^{-4}$. All results between linear and nonlinear are statistically significantly different ($p \ll 0.05$). See Appendix C.2 for an overview of expected calibration error. We also compute the maximum calibration error in Appendix C.2.

| | **Test Accuracy** ($\uparrow$) | | **Expected Calibration Error** ($\downarrow$) | |
Algorithm	Non-Tempered	Tempered	Non-Tempered	Tempered
Linear	$85.01_{\pm 0.10}$	$85.01_{\pm 0.19}$	$0.24_{\pm 0.017}$	$0.26_{\pm 0.014}$
Nonlinear (BG)	$84.28_{\pm 0.28}$	$84.74_{\pm 0.08}$	$0.14_{\pm 0.03}$	$0.16_{\pm 0.03}$

4.2.2 Results

Linear vs Nonlinear. From Table 1, we see that the linear (RMS-Langevin) sampler has slightly higher, but comparable test accuracy to the nonlinear Boltzmann-Gibbs sampler. This is likely because RMS-Langevin algorithm has better stability around the regions of high probability due to its adaptive stepsize scaling. However, from Figure 2, we see that for both the tempered and non-tempered cases, the nonlinear interaction appears to benefit during early exploration. This is an expected and desired

Non-Tempered   Tempered

Figure 2: Evaluation of test accuracy during sampling for CIFAR10. The shaded areas represent ± 1 standard deviation for $N = 5$ samples.

property of these nonlinear samplers, which incorporate global information about the sampler state (in this case, the relative potential G of each auxiliary chain's state) and are able to emphasize those states with higher probability. However, the linear method is able to eventually explore the relevant regions of the state space, and the difference disappears. See Appendix C.2.4 for a comparison linear vs nonlinear performance scaled by the number of gradient evaluations.

Tempering Vs Non-Tempering. The linear MCMC sampler is always tempered in our experiments so there should not be any statistically significant difference in the linear case. For the nonlinear sampler, the tempered version has slightly higher accuracy; this trend is also observed in [32]. On the other hand, the calibration errors are the same for both tempered and non-tempered variants. This is somewhat surprising, given the aggressive tempering used, and one would expect that this reduces the variance of the posterior estimate.[5] This observation can perhaps be explained by the fact that we are using $N = 10$ samples which may not be enough to accurately change the tempered auxiliary distribution $\eta^\star$ into the non-tempered primary distribution π in the jump interaction.

Calibration. Considering the expected calibration error (ECE) [9], in Table 1 we see that the nonlinear method has statistically significantly lower ECE ($p \ll 0.05$) compared to the linear method. We hypothesize that this is due to a tension between the RMS scaling of the gradient which improves the efficiency of each MCMC step but at the cost of bias, which may be measurable in the form of calibration error. The Langevin algorithm is also biased, but is generally known to have good convergence properties [16] and much less is known about the RMS-Langevin variant. By using our nonlinear setup, we are able to aggressively explore the auxiliary distribution without sacrificing calibration on the target distribution.

5 Conclusion

In this paper, we have studied the theoretical and empirical properties of nonlinear MCMC methods. We obtain powerful theoretical results for the convergence of our MCMC methods, and we apply these methods to Bayesian neural networks. The results on BNNs are comparable to, but not better than, the linear methods we studied. We hypothesize that this is because more investigation into choosing the best auxiliary density $\eta^\star$ is required; our choice is simplistic and may not be optimal. This hypothesis is supported by the toy experiments, which show significant improvement when $\eta^\star$ is able to incorporate some additional insight into the problem. How to do this in high dimensions is an exciting direction for future research.

Broader Impact. There are benefits and drawbacks to the nonlinear MCMC methods we describe. The benefits are mainly that properly accounting for uncertainty in machine learning will lead to better real-world outcomes for high-value scenarios such as self-driving cars or medical imaging. The drawbacks are that MCMC methods require $\mathcal{O}(N)$ storage and computations relative to the $\mathcal{O}(1)$ for deterministic methods; in fact, our nonlinear method would require $2N$ resources compared with N for a linear method (see also Appendices C.1.3 and C.2.4). If applied to a large swath of ML, this would mean a substantial increase in the energy consumption required for experimentation and deployment, worsening an already substantial issue in the field.

---

[5] As $\tau \to 0$, this $\theta \sim P(\theta|\mathcal{D}_{\text{train}})^{1/\tau}$ converges to the maximum *a posteriori* estimate with zero variance.

References

[1] Christophe Andrieu, Ajay Jasra, Arnaud Doucet, and Pierre Del Moral. On nonlinear Markov chain Monte Carlo. *Bernoulli*, 17(3):987 – 1014, 2011. doi: 10.3150/10-BEJ307. URL https://doi.org/10.3150/10-BEJ307.

[2] Rhiannon Michelmore, Marta Kwiatkowska, and Yarin Gal. Evaluating uncertainty quantification in end-to-end autonomous driving control. *arXiv preprint arXiv:1811.06817*, 2018.

[3] Sina Shafaei, Stefan Kugele, Mohd Hafeez Osman, and Alois Knoll. Uncertainty in machine learning: A safety perspective on autonomous driving. In *International Conference on Computer Safety, Reliability, and Security*, pages 458–464. Springer, 2018.

[4] Liuhui Ding, Dachuan Li, Bowen Liu, Wenxing Lan, Bing Bai, Qi Hao, Weipeng Cao, and Ke Pei. Capture uncertainties in deep neural networks for safe operation of autonomous driving vehicles. In *2021 IEEE Intl Conf on Parallel & Distributed Processing with Applications, Big Data & Cloud Computing, Sustainable Computing & Communications, Social Computing & Networking (ISPA/BDCloud/SocialCom/SustainCom)*, pages 826–835. IEEE, 2021.

[5] Benjamin Kompa, Jasper Snoek, and Andrew L Beam. Second opinion needed: communicating uncertainty in medical machine learning. *NPJ Digital Medicine*, 4(1):1–6, 2021.

[6] Roohallah Alizadehsani, Mohamad Roshanzamir, Sadiq Hussain, Abbas Khosravi, Afsaneh Koohestani, Mohammad Hossein Zangooei, Moloud Abdar, Adham Beykikhoshk, Afshin Shoeibi, Assef Zare, et al. Handling of uncertainty in medical data using machine learning and probability theory techniques: A review of 30 years (1991–2020). *Annals of Operations Research*, pages 1–42, 2021.

[7] Jianfeng He, Xuchao Zhang, Shuo Lei, Zhiqian Chen, Fanglan Chen, Abdulaziz Alhamadani, Bei Xiao, and ChangTien Lu. Towards more accurate uncertainty estimation in text classification. In *Proceedings of the 2020 Conference on Empirical Methods in Natural Language Processing (EMNLP)*, pages 8362–8372, 2020.

[8] Yarin Gal and Zoubin Ghahramani. Dropout as a bayesian approximation: Representing model uncertainty in deep learning. In *international conference on machine learning*, pages 1050–1059. PMLR, 2016.

[9] Chuan Guo, Geoff Pleiss, Yu Sun, and Kilian Q Weinberger. On calibration of modern neural networks. In *International Conference on Machine Learning*, pages 1321–1330. PMLR, 2017.

[10] K.P. Murphy. *Machine Learning: A Probabilistic Perspective*. Adaptive Computation and Machine Learning series. MIT Press, 2012. ISBN 9780262018029. URL https://books.google.ca/books?id=NZP6AQAAQBAJ.

[11] Nicholas Metropolis, Arianna W Rosenbluth, Marshall N Rosenbluth, Augusta H Teller, and Edward Teller. Equation of state calculations by fast computing machines. *The journal of chemical physics*, 21(6):1087–1092, 1953.

[12] W Keith Hastings. Monte Carlo sampling methods using Markov chains and their applications. 1970.

[13] Stuart Geman and Donald Geman. Stochastic relaxation, Gibbs distributions, and the Bayesian restoration of images. *IEEE Transactions on pattern analysis and machine intelligence*, (6): 721–741, 1984.

[14] Donald L. Ermak. A computer simulation of charged particles in solution. i. technique and equilibrium properties. *The Journal of Chemical Physics*, 62(10):4189–4196, 1975. doi: 10.1063/1.430300. URL https://doi.org/10.1063/1.430300.

[15] Gareth O Roberts and Richard L Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, pages 341–363, 1996.

[16] Alain Durmus and Eric Moulines. High-dimensional Bayesian inference via the unadjusted Langevin algorithm. *Bernoulli*, 25(4A):2854–2882, 2019.

[17] Max Welling and Yee W Teh. Bayesian learning via stochastic gradient Langevin dynamics. In *Proceedings of the 28th international conference on machine learning (ICML-11)*, pages 681–688. Citeseer, 2011.

[18] Gareth O Roberts and Jeffrey S Rosenthal. Optimal scaling of discrete approximations to Langevin diffusions. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 60(1):255–268, 1998.

[19] Gareth O. Roberts and Richard L. Tweedie. Exponential convergence of Langevin distributions and their discrete approximations. *Bernoulli*, 2(4):341 – 363, 1996. doi: bj/1178291835. URL https://doi.org/.

[20] Simon Duane, Anthony D Kennedy, Brian J Pendleton, and Duncan Roweth. Hybrid Monte Carlo. *Physics letters B*, 195(2):216–222, 1987.

[21] Michael Betancourt. A conceptual introduction to Hamiltonian Monte Carlo. *arXiv preprint arXiv:1701.02434*, 2017.

[22] Radford M. Neal. An improved acceptance procedure for the hybrid Monte Carlo algorithm. *Journal of Computational Physics*, 111:194–203, 1992.

[23] R.M. Neal. *Bayesian Learning for Neural Networks*. Lecture Notes in Statistics. Springer New York, 2012. ISBN 9781461207450. URL https://books.google.ca/books?id=LHHrBwAAQBAJ.

[24] Mark Girolami and Ben Calderhead. Riemann manifold Langevin and Hamiltonian Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 73 (2):123–214, 2011.

[25] Peter Müller and David Rios Insua. Issues in Bayesian analysis of neural network models. *Neural Computation*, 10(3):749–770, 1998.

[26] Arya A Pourzanjani, Richard M Jiang, and Linda R Petzold. Improving the identifiability of neural networks for Bayesian inference. In *NIPS Workshop on Bayesian Deep Learning*, volume 4, page 31, 2017.

[27] David M. Higdon. Auxiliary variable methods for Markov chain Monte Carlo with applications. *Journal of the American Statistical Association*, 93(442):585–595, 1998. ISSN 01621459. URL http://www.jstor.org/stable/2670110.

[28] Raza Habib and David Barber. Auxiliary variational MCMC. In *International Conference on Learning Representations*, 2018.

[29] Malcolm Sambridge. A Parallel Tempering algorithm for probabilistic sampling and multimodal optimization. *Geophysical Journal International*, 196(1):357–374, 10 2013. ISSN 0956-540X. doi: 10.1093/gji/ggt342. URL https://doi.org/10.1093/gji/ggt342.

[30] Robert Swendsen and Jian-Sheng Wang. Replica Monte Carlo simulation of spin-glasses. *Physical review letters*, 57:2607–2609, 12 1986. doi: 10.1103/PhysRevLett.57.2607.

[31] Rohitash Chandra, Konark Jain, Ratneel V Deo, and Sally Cripps. Langevin-gradient parallel tempering for Bayesian neural learning. *Neurocomputing*, 359:315–326, 2019.

[32] Florian Wenzel, Kevin Roth, Bastiaan Veeling, Jakub Swiatkowski, Linh Tran, Stephan Mandt, Jasper Snoek, Tim Salimans, Rodolphe Jenatton, and Sebastian Nowozin. How good is the Bayes posterior in deep neural networks really? In *International Conference on Machine Learning*, pages 10248–10259. PMLR, 2020.

[33] Chunyuan Li, Changyou Chen, David Carlson, and Lawrence Carin. Preconditioned stochastic gradient Langevin dynamics for deep neural networks. In *Thirtieth AAAI Conference on Artificial Intelligence*, 2016.

[34] Christophe Andrieu and Éric Moulines. On the ergodicity properties of some adaptive MCMC algorithms. *The Annals of Applied Probability*, 16(3):1462–1505, 2006.

[35] Christophe Andrieu and Johannes Thoms. A tutorial on adaptive MCMC. *Statistics and computing*, 18(4):343–373, 2008.

[36] Matthew D Hoffman, Andrew Gelman, et al. The no-u-turn sampler: adaptively setting path lengths in Hamiltonian Monte Carlo. *J. Mach. Learn. Res.*, 15(1):1593–1623, 2014.

[37] Henry P McKean Jr. A class of Markov processes associated with nonlinear parabolic equations. *Proceedings of the National Academy of Sciences of the United States of America*, 56(6):1907, 1966.

[38] Sylvie Méléard. Asymptotic behaviour of some interacting particle systems; mckean-vlasov and boltzmann models. In *Probabilistic models for nonlinear partial differential equations*, pages 42–95. Springer, 1996.

[39] José A Carrillo, Robert J McCann, and Cédric Villani. Kinetic equilibration rates for granular media and related equations: entropy dissipation and mass transportation estimates. *Revista Matematica Iberoamericana*, 19(3):971–1018, 2003.

[40] Arnaud Guillin and Pierre Monmarché. Uniform long-time and propagation of chaos estimates for mean field kinetic particles in non-convex landscapes. *Journal of Statistical Physics*, 185(2): 1–20, 2021.

[41] Alain-Sol Sznitman. Topics in propagation of chaos. In *Ecole d'été de probabilités de Saint-Flour XIX—1989*, pages 165–251. Springer, 1991.

[42] OA Butkovsky. On ergodic properties of nonlinear Markov chains and stochastic McKean–Vlasov equations. *Theory of Probability & Its Applications*, 58(4):661–674, 2014.

[43] Pierre Del Moral. Nonlinear filtering: Interacting particle resolution. *Comptes Rendus de l'Académie des Sciences-Series I-Mathematics*, 325(6):653–658, 1997.

[44] Pierre Del Moral and Laurent Miclo. Asymptotic results for genetic algorithms with applications to nonlinear estimation. In *Theoretical aspects of evolutionary computing*, pages 439–493. Springer, 2001.

[45] Arnaud Doucet, Nando De Freitas, and Neil Gordon. An introduction to sequential Monte Carlo methods. In *Sequential Monte Carlo methods in practice*, pages 3–14. Springer, 2001.

[46] Pierre Del Moral, Arnaud Doucet, and Ajay Jasra. Sequential Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(3):411–436, 2006.

[47] P.D. Moral. *Feynman-Kac Formulae: Genealogical and Interacting Particle Systems with Applications*. Probability and Its Applications. Springer New York, 2004. ISBN 9780387202686. URL https://books.google.ca/books?id=8LypfuG8ZLYC.

[48] Christophe Andrieu, Arnaud Doucet, and Roman Holenstein. Particle Markov chain Monte Carlo methods. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 72 (3):269–342, 2010.

[49] Tom Rainforth, Christian Naesseth, Fredrik Lindsten, Brooks Paige, Jan-Willem Vandemeent, Arnaud Doucet, and Frank Wood. Interacting particle Markov chain Monte Carlo. In *International Conference on Machine Learning*, pages 2616–2625. PMLR, 2016.

[50] Grégoire Clarté, Antoine Diez, and Jean Feydy. Collective proposal distributions for nonlinear MCMC samplers: Mean-field theory and fast implementation. *arXiv preprint arXiv:1909.08988*, 2019.

[51] Michael Arbel, Anna Korba, Adil Salim, and Arthur Gretton. Maximum mean discrepancy gradient flow. *Advances in Neural Information Processing Systems*, 32, 2019.

[52] Gareth O Roberts and Jeffrey S Rosenthal. General state space Markov chains and MCMC algorithms. *Probability surveys*, 1:20–71, 2004.

[53] Sean P Meyn and Richard L Tweedie. *Markov chains and stochastic stability*. Springer Science & Business Media, 2012.

[54] Rajesh Ranganath, Sean Gerrish, and David Blei. Black box variational inference. In *Artificial intelligence and statistics*, pages 814–822. PMLR, 2014.

[55] Danilo Rezende and Shakir Mohamed. Variational inference with normalizing flows. In *International conference on machine learning*, pages 1530–1538. PMLR, 2015.

[56] George Papamakarios, Eric Nalisnick, Danilo Jimenez Rezende, Shakir Mohamed, and Balaji Lakshminarayanan. Normalizing flows for probabilistic modeling and inference. *Journal of Machine Learning Research*, 22(57):1–64, 2021.

[57] Martin Hairer and Jonathan C Mattingly. Yet another look at Harris' ergodic theorem for Markov chains. In *Seminar on Stochastic Analysis, Random Fields and Applications VI*, pages 109–117. Springer, 2011.

[58] Arthur Gretton, Karsten M Borgwardt, Malte J Rasch, Bernhard Schölkopf, and Alexander Smola. A kernel two-sample test. *The Journal of Machine Learning Research*, 13(1):723–773, 2012.

[59] Vincent Stimper, Bernhard Schölkopf, and José Miguel Hernández-Lobato. Resampling base distributions of normalizing flows. In *International Conference on Artificial Intelligence and Statistics*, pages 4915–4936. PMLR, 2022.

[60] Ruqi Zhang, Chunyuan Li, Jianyi Zhang, Changyou Chen, and Andrew Gordon Wilson. Cyclical stochastic gradient MCMC for Bayesian deep learning. In *International Conference on Learning Representations*, 2019.

[61] Kaiming He, Xiangyu Zhang, Shaoqing Ren, and Jian Sun. Deep residual learning for image recognition. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 770–778, 2016.

[62] Michael K Pitt and Neil Shephard. Filtering via simulation: Auxiliary particle filters. *Journal of the American statistical association*, 94(446):590–599, 1999.

[63] James Bradbury, Roy Frostig, Peter Hawkins, Matthew James Johnson, Chris Leary, Dougal Maclaurin, George Necula, Adam Paszke, Jake VanderPlas, Skye Wanderman-Milne, and

Qiao Zhang. JAX: composable transformations of Python+NumPy programs, 2018. URL http://github.com/google/jax.

[64] Grigorios A Pavliotis. *Stochastic processes and applications: diffusion processes, the Fokker-Planck and Langevin equations*, volume 60. Springer, 2014.

[65] Denis Talay and Luciano Tubaro. Expansion of the global error for numerical schemes solving stochastic differential equations. *Stochastic analysis and applications*, 8(4):483–509, 1990.

[66] T. Tieleman and G. Hinton. Lecture 6.5—RmsProp: Divide the gradient by a running average of its recent magnitude. COURSERA: Neural Networks for Machine Learning, 2012.

[67] Chii-Ruey Hwang, Shu-Yin Hwang-Ma, and Shuenn-Jyi Sheu. Accelerating diffusions. *The Annals of Applied Probability*, 15(2):1433–1444, 2005.

[68] Andrew B Duncan, Tony Lelievre, and Grigorios A Pavliotis. Variance reduction using nonreversible Langevin samplers. *Journal of statistical physics*, 163(3):457–491, 2016.

[69] Alessandro Barp, Lancelot Da Costa, Guilherme França, Karl Friston, Mark Girolami, Michael I Jordan, and Grigorios A Pavliotis. Geometric methods for sampling, optimisation, inference and adaptive agents. *arXiv preprint arXiv:2203.10592*, 2022.

[70] Joris Bierkens, Paul Fearnhead, and Gareth Roberts. The zig-zag process and super-efficient sampling for Bayesian analysis of big data. *The Annals of Statistics*, 47(3):1288–1320, 2019.

[71] George Deligiannidis, Alexandre Bouchard-Côté, and Arnaud Doucet. Exponential ergodicity of the bouncy particle sampler. *The Annals of Statistics*, 47(3):1268–1287, 2019.

[72] Augustin Chevallier, Sam Power, Andi Q Wang, and Paul Fearnhead. PDMP Monte Carlo methods for piecewise-smooth densities. *arXiv preprint arXiv:2111.05859*, 2021.

[73] Alain Durmus, Arnaud Guillin, and Pierre Monmarché. Piecewise deterministic Markov processes and their invariant measures. In *Annales de l'Institut Henri Poincaré, Probabilités et Statistiques*, volume 57, pages 1442–1475. Institut Henri Poincaré, 2021.

[74] Pierre Monmarché. High-dimensional MCMC with a standard splitting scheme for the underdamped Langevin diffusion. *Electronic Journal of Statistics*, 15(2):4117–4166, 2021.

[75] Dimitris Bertsimas and John Tsitsiklis. Simulated annealing. *Statistical science*, 8(1):10–15, 1993.

[76] Alex Krizhevsky, Geoffrey Hinton, et al. Learning multiple layers of features from tiny images. 2009.

[77] Tom Hennigan, Trevor Cai, Tamara Norman, and Igor Babuschkin. Haiku: Sonnet for JAX, 2020. URL http://github.com/deepmind/dm-haiku.

[78] Yuval Netzer, Tao Wang, Adam Coates, Alessandro Bissacco, Bo Wu, and Andrew Y Ng. Reading digits in natural images with unsupervised feature learning. 2011.

A Pseudocode & Numerical Implementation Details

A.1 Simulation Algorithm

The method of simulating K_η is where this work and [1] diverge: we will provide an algorithm that uses a fixed number of samples $N \in \mathbb{N}$ to simulate the IPS (3), whereas [1] investigates an algorithm in which the empirical measure is formed from all the *past samples*.

While the algorithm in [1] yields a true estimate of the target measure in the limit as $n \to \infty$ (for $K_\bullet^{AR}$ at least, see [1]), their MCMC algorithm's storage complexity increases linearly with time. This behaviour is not well-suited to large-scale software implementations, which typically favour fixed-size "batches" of computations. Additionally, a practitioner cannot horizontally scale their algorithm to improve accuracy (e.g. by simply adding more samples) in a fixed time frame.

By contrast, Algorithm 1 below and the ensuing analysis use a fixed number of particles N throughout the lifetime of the algorithm. We will see that this produces a biased estimate of the target measure π, but that this bias can be controlled both in the number of particles, N, and the number of steps, n. In exchange, our algorithm can be efficiently implemented in vectorized computing frameworks, as we will demonstrate in Section 4.

Additionally, let us note that the decision to use an auxiliary Markov chain is highly pragmatic. It is possible to develop "autonomous" nonlinear MCMC algorithms of the form $\widetilde{\mu}_{n+1} = K_{\widetilde{\mu}_n}$ (see e.g. [47] Ch. 5) with strong theoretical guarantees but bad empirical performance. A primary reason for this is *sample degeneracy*: due to the properties of nonlinear interaction, in an autonomous nonlinear Markov kernel, quite often a single particle X_n^i will be given a large potential $G(X_n^i)$ which results in the jump interaction being concentrated on a single point. From this point onwards, the algorithm will be unable to generate enough diversity within its particles to correctly estimate the variance of the target measure. Due to the close relationship of nonlinear MCMC and nonlinear filtering, these issues have been studied in many settings such as [62, 48] and using auxiliary dynamics is indeed a common solution.

Algorithm 1: Sampling from a nonlinear Markov chain with transition K_η.

Input: Initial samples $X_0^i \overset{iid}{\sim} \mu_0$, $Y_0^i \overset{iid}{\sim} \eta_0$, $i = 1, \dots, N$
Input: Primary and auxiliary Markov kernels K, Q resp. and jump kernel J_η
Input: Number of iterations, n_{sim}, jump probability ε
Output: Collection of samples $\{X_{n_{\text{sim}}}^1, \dots, X_{n_{\text{sim}}}^N\}$.

1 **for** $n = 0, \dots, n_{sim-1}$ **do**
    // Sample auxiliary Markov chain
2   **for** $i = 1, \dots, N$ **do**
3        $Y_{n+1}^i \sim Q(Y_n^i, \bullet)$
    // Sample primary Markov chain
4   **for** $i = 1, \dots, N$ **do**
5        $Z^i \sim K(X_n^i, \bullet)$
    // Sample nonlinear Markov chain
6   **for** $i = 1, \dots, N$ **do**
7        $B^i \sim \text{Bern}(\varepsilon)$// Sample binary jump/no jump random variable
8   **if** $B^i == 0$ **then**
9   Set $X_{n+1}^i = Z^i$ // No jump; evolve according to K
10   **else**
11   Set $\overline{Y}_n = \{Y_n^1, \dots, Y_n^N\}$
12   Sample $X_{n+1}^i \sim J_{m(\overline{Y}_n)}(X_n^i, \bullet)$ // Jump; sample a new position.

A.2 Efficient Software Implementation

In Algorithm 1, we gave a high-level pseudocode implementation of the nonlinear sampler. In this implementation, for the sake of notational clarity, we used `for` loops to sample the individual

particles. However, with modern single instruction multiple data (SIMD) computing frameworks such as GPU accelerators, this is highly inefficient. Hence one can, and should, paralellize all for loops except the outer "time" loop in Algorithm 1.

We have done this using the variety of powerful tools provided by the Jax library [63] within Python, such as vmap which allows for automatic vectorization, and seamless targeting of GPU accelerators. This becomes particularly useful for automatic batch-wise and sample-wise vectorization in the case of stochastic gradient MCMC while writing functions in-terms of single inputs and outputs. We have also leveraged the library of linear MCMC algorithms provided by the *<REDACTED FOR ANONYMITY>* library *<REDACTED FOR ANONYMITY>*. All experiments were run on a single Titan X GPU; the code can be found at *<REDACTED FOR ANONYMITY>*.

B Linear MCMC Sampling Algorithms

Markov chain Monte Carlo algorithms form an integral part of Bayesian inference. In this section, we will review the basic MCMC algorithms used in this paper. In each case, we assume a C^1, strictly positive target density π known only up to a normalizing constant.

B.1 Unadjusted Langevin Algorithm

Consider the (overdamped) continuous Langevin diffusion [64]

$$\mathrm{d}X_t = \nabla \log \pi(X_t)\mathrm{d}t + \sqrt{2}\mathrm{d}B_t$$

where B_t is a standard Brownian motion. This is a fundamental stochastic process with far reaching consequences in many areas of math; in particular, the Langevin diffusion is a Markov process with π as a stationary measure [64]. This fact has long been exploited by the molecular dynamics literature [14]. We obtain the unadjusted Langevin algorithm, or ULA, by applying an Euler-Maruyama discretization to the SDE above yielding [64]

$$X_{n+1} = X_n + \delta \nabla \log \pi(X_n) + \sqrt{2\delta}Z_n, \quad Z_n \sim \mathcal{N}(0, I_d).$$

This is a popular and well-studied algorithm for MCMC, although it is known to be *biased* in the sense that the sationary measure of this Markov chain is not π [65]. However, this bias converges to 0 as $\delta \to 0$ [16].

B.2 Metropolis-Adjusted Langevin Algorithm

The Metropolis-Adjusted Langevin algorithm (MALA) [18, 19], as all basic Metropolis-Hastings MCMC methods, consists of two steps: a proposal step and an accept step. Starting from state X_n, in the proposal step, a candidate next state is sampled according to the Langevin dynamics above $\widetilde{X}_{n+1} = X_n + \delta \nabla \log \pi(X_n)\sqrt{2\delta}Z_n$. We will write $q(y|x)$ for the proposal distribution; in this case $q(y|x) = \mathcal{N}(x + \delta \nabla \log \pi(x); 2\delta)(y)$. However, unlike the ULA, MALA consists of a second step which will accept or reject the propsal $\widetilde{X}_{n+1}$ with probability

$$\alpha(\widetilde{X}_{n+1}, X_n) := 1 \wedge \frac{\pi(\widetilde{X}_{n+1})q(X_n|\widetilde{X}_{n+1})}{\pi(X_n)q(\widetilde{X}_{n+1}|X_n)}.$$

In other words

$$X_{n+1} = \left\{ \begin{array}{ll} \widetilde{X}_{n+1} & \text{if } U_n \leq \alpha(\widetilde{X}_{n+1}, X_n) \\ X_n & \text{otherwise} \end{array} \right. ; \quad U_n \sim [0, 1].$$

B.3 RMS-Unadjusted Langevin Algorithm

The ULA is a powerful technique for "black-box" MCMC in which one only has access to gradient information about the target density. However, for very high dimensional problems in which π is highly anisotropic, it can be inefficient to simulate an isotropic diffusion such as the Langevin algorithm. A simple and effective technique, borrowed from the optimization literature in which the

same phenomenon can cause problems, is to *precondition* the dynamics as follows. First we maintain an exponentially smoothed squared-gradient estimate r_n defined as

$$r_{n+1} = \beta r_n + (1 - \beta)\nabla \log \pi(X_n)^2; \quad \beta \in [0, 1[$$

where the gradient squared is meant elementwise. Then, as in RMSProp [66], we can construct an adaptive stepsize by dividing the square root of r_n as follows

$$\widehat{\delta}_{n+1} := \frac{\delta}{\sqrt{r_{n+1} + \epsilon}}$$

and then using this stepsize in the Langevin update

$$X_{n+1} = X_n + \widehat{\delta}_{n+1}\nabla \log \pi(X_n) + \sqrt{2\widehat{\delta}_{n+1}}Z_n$$

$$= X_n + \frac{\delta}{\sqrt{r_{n+1} + \epsilon}}\nabla \log \pi(X_n) + \sqrt{2\frac{\delta}{\sqrt{r_{n+1} + \epsilon}}}Z_n.$$

This has the effect of using gradient information to scale the stepsize of original Langevin diffusion independently along each dimension, which likely reduces the negative impacts of anisotropy and substantially increases mixing time. As studied in in [33], this algorithm, which we call RMS-ULA, is biased but this bias can be controlled with δ.

B.4   RMS-MALA.

Similar to the transition from ULA to MALA, we can "metropolize" the RMS-ULA algorithm to correct the bias. This follows the same structure as the MALA, except that the proposal depends on r[6]

$$r' = \beta r + (1 - \beta)\nabla \log \pi(x)^2$$

$$q(y|x) = \mathcal{N}\left(x + \frac{\delta}{\sqrt{r' + \epsilon}}\nabla \log \pi(X_n), 2\frac{\delta}{\sqrt{r' + \epsilon}}\right).$$

B.5   Practical Considerations for Bayesian Neural Networks: Reversibility & Tempering.

In practical settings, the high dimensional, anisotropic characteristics and limited computation budget of Bayesian neural networks (BNNs) necessitate some modifications to the above algorithms.

The first is that the Metropolis-Hastings (MH) step is rarely used except in simple settings. This is because, while ensuring that that π is the invariant measure, the reversibility of the MH step is too inefficient when we cannot affort to reject many samples. Because they cannot "backtrack", nonreversible dynamics are generally much more efficient than reversible dynamics [67, 68, 64, 69]. This has led to the recent interest in using piecewise deterministic Markov processes [70–73] and other nonreversible MCMC dynamics such as [74] for high dimensional Bayesian inference. See [69], Section 3.1 for an interesting dicussion on nonreversibility and efficiency.

The second modification is tempering. A tempered version of a distribution π is $\pi_\beta \propto \pi^\beta$. This is a pragmatic solution to speed up exploration of a target distribution when the noise component of Langevin-like algorithms causes slow or unstable dynamics. In the limit as $\beta \to \infty$, sampling from π_∞ is equivalent to the maximum *a posteriori* estimate (this is simulated annealing [75]). In practice, tempering can be achieved by simply scaling the noise by $\sqrt{\tau}$

$$X_{n+1} = X_n + \delta\nabla \log \pi(X_n) + \sqrt{2\delta}\sqrt{\tau}Z_n, \quad Z_n \sim \mathcal{N}(0, I_d).$$

which targets π^β where $\beta = 1/\tau$.[7] See also [32] for a discussion of tempering in BNNs.

---

[6] A more "proper" setup would expand the statespace to include be (x, r), which would restore the Markov property

[7] To see this, for $\pi_\beta \propto \pi^\beta$ we have $\nabla \log \pi_\beta = \beta\nabla \log \pi$ so by using a step size $\delta' = \delta/\beta$ we get $X_{n+1} = X_n + \beta\delta'\nabla \log \pi(X_n) + \sqrt{2\delta'}Z_n = X_n + \delta\nabla \log \pi(X_n) + \sqrt{2\delta/\beta}Z_n$ and $\tau = 1/\beta$.

C   Experimental Details

No hyperparameter sweeps were used in the any of these experiments. Hyperparameters were chosen based on reasonable guesses and minimal manual tuning.

C.1   2d Toy Experiments

C.1.1   Maximum Mean Discrepancy

We use the maximum mean discrepancy (MMD) metric [58] as a way to quantitatively evaluate our MCMC algorithms. The MMD metric is an integral probability metric of the form

$$\|\mu - \nu\|_{MMD} = \sup_{\|f\|_{\mathcal{H}} \leq 1} |\mu(f) - \nu(f)| = \sup_{\|f\|_{\mathcal{H}} \leq 1} |\mathbb{E}_{X \sim \mu}[f(X)] - \mathbb{E}_{Y \sim \nu}[f(Y)]|$$

where $\mathcal{H}$ is a reproducing kernel Hilbert space (RHKS) associated to a positive-definite kernel function $k : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$; see [58] for details. For our purposes, the important features of the MMD metric are that 1) it is similar to the total variation metric, which optimizes over the unit ball in $\mathcal{B}_b(\mathbb{R}^d)$ instead of $\mathcal{H}$; and 2) the MMD metric $\|\mu - \nu\|_{MMD}$ can be efficiently empirically estimated from samples of μ, ν. This is because, according to Lemma 6 in [58], we have

$$\|\mu - \nu\|_{MMD}^2 = \mathbb{E}_{X, X' \sim \mu \otimes \mu}[k(X, X')] - 2\mathbb{E}_{X, Y \sim \mu \otimes \nu}[k(X, Y)] + \mathbb{E}_{Y, Y' \sim \nu \otimes \nu}[k(Y, Y')]$$

which depends only on the kernel function k, and an *unbiased* estimator of $\|\mu - \nu\|_{MMD}^2$ is

$$\|\mu - \nu\|_{MMD}^2 \approx \frac{1}{N_\mu(N_\mu - 1)} \sum_{i,j=1,\ j \neq i}^{N_\mu} k(X^i, X^j) - 2\frac{1}{N_\mu N_\nu} \sum_{i=1}^{N_\mu} \sum_{j=1}^{N_\nu} k(X^i, Y^j) + \frac{1}{N_\nu(N_\nu - 1)} \sum_{i,j=1,\ j \neq i}^{N_\nu} k(Y^i, Y^j)$$

where $X^i \overset{iid}{\sim} \mu$ for $i = 1, \ldots, N_\mu$ and $Y^i \overset{iid}{\sim} \nu$ for $i = 1, \ldots, N_\nu$.

If we treat μ as the distribution μ_n of a (linear or nonlinear) MCMC algorithm at step n and ν as π, then we can apply this estimator provided we can sample from π exactly. This is not feasible for complex high-dimensional distributions, but it is possible for the distributions we have chosen, which are all mixtures or transformations of Gaussains.

C.1.2   Experimental Setup

Refer to Table 2 for the experimental settings. Additionally, we used $10,000$ samples from π and the kernel

$$k(x, y) = \exp(-\|x - y\|^2) + \exp(-2\|x - y\|^2)$$

to estimate the MMD as in the previous section.

Table 2: Experimental settings for the 2d toy experiments.

Setting	Symbol	Value
Auxiliary/Linear Markov kernel	Q	MALA
Auxiliary/Linear kernel stepsize	δ_{aux}	0.001
Auxiliary/Linear target density	$\eta^\star$	$\mathcal{N}(0, \sigma^2 I_2)$ with $\sigma = 4$ for circular MoG and two rings and $\sigma = 20$ for grid MoG
Primary Markov kernel	K	MALA
Primary kernel stepsize	δ	0.001
Number of samples	N	2000
Initial Auxiliary/Linear distribution	η_0	$\mathcal{U}([-7.5, 7.5] \times [-7.5, 7.5])$
Initial Primary distribution	η_0	$\mathcal{U}([-7.5, 7.5] \times [-7.5, 7.5])$
Jump probability	ε	0.1
Number of simulation steps	n_{sim}	$10,000$

C.1.3 Runtime Analysis

We also report on the performance vs runtime of the linear and nonlinear algorithms measured in terms of number of gradient executions and the wallclock time.

In general, the nonlinear methods will require two times the number of gradient executions as the linear methods due to the requirement that we sample from the auxiliary chain in addition to the primary chain. In Figure 3, we show the MMD-squared metric plotted against the number of gradient evaluations. Since we ran all algorithms for a fixed number of steps, this amounts to using $2\times$ the gradient evaluations for the nonlinear methods, although not $2\times$ the steps for a given particle. Despite this, the nonlinear methods offer better performance per gradient evaluation.

| Circular MoG | Two Rings | Grid MoG |

Figure 3: MMD^2 performance vs number of gradient executions (per sample) for the two-dimensional examples.

In Table 3, we also report the wallclock time for each of the methods and each of the distributions. We see that, despite using $2\times$ the gradient evaluations, the effective use of JAX and its vectorization and jit-compilation features allows us to achieve sublinear wallclock time, at least for this operating point of our GPU accelerator. We omit the first twenty iterations in our calculation since these can be slow while JAX compiles the program and we measure only the steady-state execution performance.

Tgt Density	Sampler	Wall time (s)	Slowdown vs Linear
Circular MoG	Linear	$2.57_{\pm 0.13}$	-
	AR	$3.00_{\pm 0.06}$	$1.17\times$
	BG	$3.29_{\pm 0.02}$	$1.10\times$
Two Rings	Linear	$2.58_{\pm 0.05}$	-
	AR	$3.00_{\pm 0.06}$	$1.16\times$
	BG	$3.13_{\pm 0.03}$	$1.21\times$
Grid MoG	Linear	$2.64_{\pm 0.05}$	-
	AR	$3.08_{\pm 0.04}$	$1.16\times$
	BG	$3.83_{\pm 0.01}$	$1.25\times$

Table 3: Wall Time analysis of the two-dimensional sampling problems (mean ± 1 standard deviation on 5 trials). We measure the steady-state wall time by omitting the first 20 iterations to account for the JAX jit-compilation time at the start of the program.

C.2 CIFAR10 Experiments

C.2.1 Experimental Setup

In this experiment, we implement a Bayesian neural network for image classification on the CIFAR10 dataset [76]. The neural network architecture we use is a standard ResNet-18 architecture [61] using the default settings implemented by the DeepMind `haiku` library [77]. The likelihood function $P(y|x, \theta)$ is the standard crossentropy on the logits y given an input image x and parameters θ. We also use standard data agumentation techniques including standardizing the images by the mean and variance, random crops and random flips. See the Table 4 for the experimental parameters.

Table 4: Experimental settings for the CIFAR10 experiments.

Setting	Symbol	Value
Auxiliary Markov kernel	Q	RMS-Langevin
Auxiliary kernel stepsize	$\delta_{aux}(n)$	$0.001 \times 0.1^{\lfloor n/2000 \rfloor}$
Auxiliary RMS β, ϵ parameters	N/A	$0.9, 1 \times 10^{-9}$
Auxiliary target density	$\eta^\star$	$\eta^\star \propto \pi^{0.9/\tau}$
Noise scaling (tempering)	$\sqrt{\tau}$	1×10^{-4}
Primary Markov kernel	K	Unadjusted Langevin
Primary kernel stepsize	δ	5×10^{-5}
Number of samples	N	10
Minibatch size	$|\widehat{\mathcal{D}}|$	256
Initial Auxiliary distribution	η_0	$\mathcal{N}(\theta_0, 0.001)$ where θ_0 is the initialized set of parameters from the `Haiku` implementation
Initial Primary distribution	μ_0	same as η_0
Jump probability	ε	0.05
Number of simulation steps	n_{sim}	50 passes through the dataset
Bayesian Prior	$P(\theta)$	$\mathcal{N}(0, 1 \times 10^{-4} I_d)$

C.2.2 Calibration Analysis

Calibration of a classifier is a measure of how well that classifier's logits represent probabilities [9]. We use this as a necessary but not sufficient test for correct sampling, since we know *a priori* that the Bayesian posterior will have much better calibration than an optimized version. We will use the methodology in [9], particularly "expected calibration error" and "maximum calibration error" to study the calibration of our Bayesian neural network, which we describe below.

To measure calibration, assume first that we have a test set of data $\mathcal{D}_{\text{test}} = \{(x_1, y_1), \ldots, (x_{N_{\text{test}}}, y_{N_{\text{test}}})\} \subset \mathcal{X} \times \{0, \ldots, C\}$ where $\mathcal{X}$ is the input space and there are $C \in \mathbb{Z}_{>1}$ labels. Suppose $\widehat{p} : \mathcal{X} \to \Delta^{C-1}$ is a supposed probability distribution representing $p(y|x)$. We would expect that the probabilities output by $\widehat{p}$ correlate with the accuracy of $\widehat{p}$, i.e. when $\widehat{p}$ is $\alpha\%$ certain, then $\alpha\%$ of the time $\widehat{p}$ is correct. More formally, perfect calibration is when

$$P(y = y_i | \widehat{p} = \alpha) = \alpha.$$

We can empirically estimate this relationship between predictied probability and true probability using histograms. I.e. given predicted probabilities $\widehat{p}_i = \widehat{p}(y_i | x_i)$, we can form bins B_k, $k = 1, \ldots, M$ on $[0, 1]$, we can estimate the accuracy in the k-th bin as A_k and the average probability in that bin as α_k. If $\widehat{p}$ represents a true probability distribution, then $A_k \approx \alpha_k$ for each $k = 1, \ldots, M$. The expected calibration error (ECE) is then

$$ECE(\widehat{p}) := \sum_{k=1}^{M} \frac{|B_k|}{N_{\text{test}}} |A_k - \alpha_k|.$$

We can also study the maximum calibrarion error

$$MCE(\widehat{p}) := \max_{k=1,\ldots,M} |A_k - \alpha_k|.$$

In Table 5, we report the ECE and MCE for the CIFAR10 experiment, where now

$$\widehat{p}(y|x, \mathcal{D}) = \sum_{i=1}^{N} p(y|x, \theta^i) \approx \int p(y|x, \theta) p(\mathrm{d}\theta | \mathcal{D}).$$

C.2.3 Distribution Shift

One useful outcome of using the probabilistic paradigm for ML is robustness to distribution. This is well-documented [8, 9]; traditional ML models tend to be overconfident even in settings where

Table 5: Calibration errors for the CIFAR10 experiment. $N = 5$ runs were used, and $\pm$ represents one standard deviation. Both ECEs and tempered MCE appear to be statistically significantly lower for the nonlinear than the linear algorithm, indicating better calibration in those cases.

Algorithm	Expected Calibration Error ($\downarrow$)		Max Calibration Error ($\downarrow$)	
	Non-Tempered	Tempered	Non-Tempered	Tempered
Linear	0.24 ± 0.017	0.26 ± 0.014	3.78 ± 0.43	4.88 ± 0.85
Nonlinear (BG)	0.14 ± 0.03	0.16 ± 0.03	3.21 ± 0.75	3.72 ± 0.40
p-values for Linear$-$Nonlinear	**0.00067**	**0.00017**	0.22084	**0.03925**

Figure 4: Demonstration of the in/out of domain performance of optimized, linear MCMC, and nonlinear MCMC. The linear and nonlinear methods perform essentially the same, which is *expected* since they both approximate the same distribution $P(y|x, \mathcal{D}_{\text{train}})$

they have no hope of good predictions. In short, traditional ML models lack the ability to say "I don't know" in a way that probabilistic ML models do not. This is particularly salient in real-world systems such as self-driving cars, where the distribution of images is highly nonstationary and the costs for overconfident predictions are high.

In Figure 4 below, we show how the predictive entropy on in-domain examples, i.e. the CIFAR10 test set, and out-of-domain examples differs. For the out-of-domain examples, we use the SVHN dataset [78] which also contains 32×32 RGB images, but this time of housing numbers (i.e. the digits 0-9) rather than objects such as airplanes or dogs. The entropy (a measure of uncertainty) is calculated as

$$H(\mathbf{p}) = -\sum_{i=1}^{10} p_i \log p_i$$

where $\mathbf{p} \in \Delta^9$ is the vector of probabilities for a prediction $p(y|x, \mathcal{D}_{\text{train}})$. In the traditional ML case, the probabilities are made with $p(y|x, \theta^*)$. We average $H(\mathbf{p})$ over the test data for the in-domain and out-of-domain examples. The optimized method was trained to the same test accuracy (85%) using essentially the same hyperparameters (learning rate, etc).

C.2.4 Runtime Analysis

In Figure C.2.4, we plot the performance of the tempered target vs the number of gradient evaluations. As described in Appendix C.1.3, the nonlinear algorithm uses $2\times$ the gradient computations. When plotting the eval performance vs the number of computations, we see that the linear algorithm is significantly more efficient. Further work into designing nonlinear samplers that achieve better performance (such as in the two-dimensional examples above) is required to make this tradeoff worthwhile in practice.

D Notation & Assumptions

The remaining sections will be devoted to proving the theoretical results contained in this paper. They should be read in sequence.

Note that, while some of the assumptions found below appear in [1], our analysis and the results we obtain are new. In particular, [1] studies a different MCMC algorithm based on K_η, see Appendix A

Figure 5: CIFAR10 test performance plotted against number of gradient evaluations (per sample) on the tempered target densit, for the linear and nonlinear (BG) MCMC algorithms.

for a discussion of algorithm differences. Moreover, they obtain an asymptotic "strong law of large numbers" result whereas we obtain nonasymptotic mean field long-time convergence and uniform propagation of chaos results, which are qualitatively different results using different proof techniques.

D.1 Notation and Definitions

D.1.1 Probability Spaces, Measures, and Kernels

We will be working on the measurable space $(\mathbb{R}^d, \mathscr{B}(\mathbb{R}^d))$ with $\mathscr{B}(\mathbb{R}^d)$ the Borel σ-algebra[8]; let $\mathcal{P}(\mathbb{R}^d)$ denote the space of probability measures on $\mathscr{B}(\mathbb{R}^d)$. Let $\mu \in \mathcal{P}(\mathbb{R}^d)$, $f : \mathbb{R}^d \to \mathbb{R}$ be a measurable function, and, $K : \mathbb{R}^d \times \mathscr{B}(\mathbb{R}^d) \to [0, 1]$ be a Markov kernel. Throughout the sequel, we will write

$$\mu(f) := \int f(x)\mu(\mathrm{d}x), \quad Kf(x) := \int f(y)K(x, \mathrm{d}y), \quad \mu K(\mathrm{d}y) := \int \mu(\mathrm{d}x)K(x, \mathrm{d}y).$$

An obvious consequence of this is the notation $\mu K(f) = \int f(y)\mu(\mathrm{d}x)K(x, \mathrm{d}y)$. We will also need the following subset of $\mathcal{P}(\mathbb{R}^d)$ for our results: let $G, U : \mathbb{R}^d \to [0, \infty[$ and fix constants $m_G, M_U > 0$; then we define

$$\mathcal{P}_{m_G, M_U}(\mathbb{R}^d) := \{\mu \in \mathcal{P}(\mathbb{R}^d) \mid \mu(G) > m_G, \ \mu(U) \le M_U\}.$$

We will abbreviate $\mathcal{P}_{m_G, M_U}(\mathbb{R}^d) = \mathcal{P}_{m, M}(\mathbb{R}^d)$ when there is no chance of confusion.

We will also frequently use the following notation for the empirical measure associated to $\overline{y} := \{y^1, \ldots, y^N\} \subset \mathbb{R}^d$:

$$m(\overline{y}) := \frac{1}{N} \sum_{i=1}^{N} \delta_{y^i}$$

with $\delta_x \in \mathcal{P}(\mathbb{R}^d)$ the Dirac measure. Additionally, we will need the notion of tensor products for functions and measures: let $q \in \mathbb{N}$, $f_i : \mathbb{R}^d \to \mathbb{R}$ and $\mu_i \in \mathcal{P}(\mathbb{R}^d)$ for $i = 1, \ldots, q$. Then for $x = (x^1, \ldots, x^q) \in (\mathbb{R}^d)^q$ and measurable $g : (\mathbb{R}^d)^q \to \mathbb{R}$

$$f_1 \otimes \cdots \otimes f_q(x) := f_1(x^1) \cdots f_q(x^q), \quad \mu_1 \otimes \cdots \otimes \mu_q(g) := \int g(x^1, \ldots, x^q)\mu_1(\mathrm{d}x^1) \cdots \mu_q(\mathrm{d}x^q).$$

When $f = f_1 = \cdots = f_q$ and $\mu = \mu_1 = \cdots = \mu_q$, we will write $f^{\otimes q}$ and $\mu^{\otimes q}$ respectively. Finally, we write $\mu \ll \nu$ to mean that ν dominates μ and we write $\frac{\mathrm{d}\mu}{\mathrm{d}\nu}$ for the Radon Nikodym derivative. The notation $\mu \sim \nu$ means that $\mu \ll \nu$ and $\nu \ll \mu$.

---

[8] This choice is merely for simplicity of exposition; most if not all our results will hold for any polish space and its Borel σ-algebra.

 ### D.1.2   Norms

787 Denote by $\mathcal{B}_b(\mathbb{R}^d)$ the set of bounded measurable functions $f : \mathbb{R}^d \to \mathbb{R}$. We will be working with a
788 family of norms on functions, and their dual norms on probability measures, which is parameterized
789 by functions of the form $U : \mathbb{R}^d \to [1, \infty[$. For such a U, we define

$$\|f\|_U := \sup_{x \in \mathbb{R}^d} \frac{|f(x)|}{U(x)}$$

790 for $f : \mathbb{R}^d \to \mathbb{R}$. The dual norm to $\|f\|_U$ corresponds to the weighted total variation distance; for
791 $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ we have

$$\|\mu - \nu\|_{tv,U} := \sup_{\|f\|_U \leq 1} |\mu(f) - \nu(f)|.$$

792 Note that we could replace the condition $\|f\|_U \leq 1$ by $|f| \leq U$. For $V : \mathbb{R}^d \to [0, \infty[$, we will
793 often work with $V_\beta(x) := 1 + \beta V(x)$ for $\beta > 0$; in this case, we will write $\|f\|_\beta := \|f\|_{V_\beta}$ and
794 $\| \bullet \|_{tv,\beta} := \| \bullet \|_{tv,V_\beta}$. We will also need the definition of the maximum oscillation of f, which we
795 denote and define as $\mathrm{osc}(f) := \sup\{|f(x) - f(y)| \mid x, y \in \mathbb{R}^d\}$.

796 For Markov kernels K, Q on $\mathbb{R}^d$, we obtain the weighted kernel distance

$$\|K - Q\|_{ker,U} := \sup_x \frac{\|K(x, \bullet) - Q(x, \bullet)\|_{tv,U}}{U(x)}.$$

797 It is worth noting the special case $U \equiv \mathbf{1}$ which corresponds to the usual total variation distance; in
798 this case we will write

$$\|f\|_\infty := \|f\|_{\mathbf{1}}, \quad \|\mu - \nu\|_{tv} := \|\mu - \nu\|_{tv,\mathbf{1}}.$$

799 Lastly, we will use the notation $\epsilon(K)$ to denote the contraction coefficient of a Markov kernel K
800 defined as $\epsilon(K) := \inf\{c \in [0, 1] \mid \|\mu K - \nu K\|_{tv} \leq c\|\mu - \nu\|_{tv} \ \forall \mu, \nu \in \mathcal{P}\}$ see, e.g. [47] Ch 4.

801 ### D.2   Assumptions

802 **Assumption 1** (Drift and Minorization)**.** *Let $K, Q : \mathbb{R}^d \times \mathscr{B}(\mathbb{R}^d) \to [0, 1]$ be Markov kernels.*

803 **K1** *K satisfies the drift criterion*

$$KV(x) \leq aV(x) + b$$

804 *for $a \in]0, 1[$, $b > 0$, and $V : \mathbb{R}^d \to [0, \infty[$ with $\lim_{\|x\| \to \infty} V(x) = \infty$.*

805 **K2** *K satisfies the following uniform minorization condition on the level sets of V: there exists*
806 *$\overline{\gamma} \in]0, 1[$ and $\nu \in \mathcal{P}(\mathbb{R}^d)$ and $R > 2b/(1 - a)$ s.t.*

$$\inf_{\{x \mid V(x) \leq R\}} K(x, A) \geq \overline{\gamma}\nu(A) \quad \forall A \in \mathscr{B}(\mathbb{R}^d).$$

807 **Q1** *Q satisfies the drift criterion*

$$QU(x) \leq \xi U(x) + c$$

808 *for $\xi \in]0, 1[$, $c > 0$, and $U : \mathbb{R}^d \to [1, \infty[$ with $\lim_{\|x\| \to \infty} U(x) = \infty$.*

809 **Q2** *Q satisfies the following uniform minorization condition on the level sets of U: there exists*
810 *$\zeta \in]0, 1[$ and $\nu' \in \mathcal{P}(\mathbb{R}^d)$ and $R' > 2c/(1 - \xi)$ s.t.*

$$\inf_{\{x \mid U(x) \leq R'\}} Q(x, A) \geq \zeta\nu'(A) \quad \forall A \in \mathscr{B}(\mathbb{R}^d).$$

811 The requirements in Assumption 1 are from [57] and are by now fairly standard in the Markov chain
812 literature. They will in particular imply the following properties for K and Q respectively:

813 **Proposition 1** (Basic Properties of K, Q; [57]&[1])**.**

814   *1. Suppose Assumptions 1-K1,K2 hold. Then there exists $\gamma \in]0, 1[$ and $\beta > 0$ s.t. $\forall \mu, \nu \in$*
815   *$\mathcal{P}(\mathbb{R}^d)$, we have*

$$\|\mu K - \nu K\|_{tv,\beta} \leq \gamma\|\mu - \nu\|_{tv,\beta}.$$

2. *Suppose that Assumptions 1-Q1,Q2 hold. Let $\eta_n = \eta_0 Q^n$ for $\eta_0 \in \mathcal{P}(\mathbb{R}^d)$, and $r \in]0,1]$. If $\eta_0(U^r) < \infty$, then for $r \in]0,1[$ there are constants $M(r) > 0$, $\delta \in]0,1[$ s.t.*

$$\|\eta_n - \eta^\star\|_{tv,U^r} \le M(r)\delta^n. \tag{7}$$

Proof.

1. This is directly from [57].

2. This uses Lemma 1 which shows Assumption 1 Q1,Q2 implies the assumptions in Lemma C.1 from [1]. Then we combine this with Lemma 2. Both lemmas can be found in Appendix E.

$\square$

Throughout the remainder of this paper, β and γ will be fixed and we will adopt the notation

$$V_\beta(x) := 1 + \beta V(x).$$

Additionally, references to $M(r)$ and δ are meant in the sense of the above proposition. Next, we introduce some criteria that ensure that the kernels K, Q are "compatible" in-terms of their drift criteria from Assumption 1.

Assumption 2 (Compatibility). **C1** *There is $r^* \in]0,1]$ s.t. $V_\beta(x) \le U(x)^{r^*} \; \forall x \in \mathbb{R}^d$ with V_β be as above.*

C2 *G satisfies the lower bound compatibility criterion with U: for every $R > 0$*

$$\theta(R) := \inf\{G(x) \mid x \in \mathbb{R}^d, \; U(x) \le R\} > 0.$$

Assumption 2-C1 is also present in [1], and ensures that V and U are sufficiently "comparable". Assumption 2-C2 is a novel assumption that we will use to obtain an *a priori* lower bound on $\eta_n(G)$. Finally, some straightforward boundedness assumptions on G.

Assumption 3 (Assumptions on G).

G1 *G is bounded, i.e. $\|G\|_\infty < \infty$.*

G2 *G is bounded in the weighted supremum norm for V_β; i.e. $\|G\|_\beta < \infty$.*

E Long-Time Convergence

This is the main long-time convergence result for the mean field system. We will revisit this theorem at the end of this section and provide a full proof. We will often use the equivalent "distribution flow" interpretation of system (2) defined as

$$\begin{cases} \eta_{n+1} &= \eta_n Q \\ \mu_{n+1} &= \mu_n K_{\eta_{n+1}}. \end{cases} \tag{8}$$

for our proofs.

Theorem 2. *Suppose that Assumption 1 and Assumption 2-C1 hold. Suppose also that J_η satisfies*

$$\|J_\eta - J_{\eta'}\|_{ker,\beta} \le C_J \|\eta - \eta'\|_{tv,\beta} \tag{9}$$

and that

$$K_\eta V(x) \le \widetilde{a} V(x) + \widetilde{b} \tag{10}$$

for $\widetilde{a} \in]0,1[$, $\widetilde{b} > 0$ and all $\eta \in \mathcal{P}_{m_G, M_U}$ for suitably chosen constants m_G, M_U. Suppose δ is from (7) and set

Case 1: *$\rho := (1 - \varepsilon)\gamma$ if $J_\eta(x, \mathrm{d}y)$ doesn't depend on x; or*

Case 2: *$\rho := (1 - \varepsilon)\gamma + \varepsilon\|J_{\eta^\star}V\|_V$ if $J_\eta(x, \mathrm{d}y)$ does depend on x.*

If $\mu_0(V), \eta_0(U) < \infty$ then there exists a constant $C > 0$ s.t.

$$\|\mu_n - \pi\|_{tv,\beta} \le \rho^n \|\mu_0 - \pi\|_{tv,\beta} + Cn \max(\rho, \delta)^n.$$

In particular, if $\rho < 1$, we have $\lim_{n \to \infty} \mu_n = \pi$ in V_β-total variation.

 ### E.1 Results about Weighted Total Variation

Lemma 1. *Suppose that a Markov kernel P satisfies a drift condition*

$$PV(x) \le aV(x) + b$$

for $V : \mathbb{R}^d \to [0, \infty[$, $V(x) \to \infty$ as $\|x\| \to \infty$, and the minorization condition with $\epsilon \in]0, 1[$, $\nu \in \mathcal{P}(\mathbb{R}^d)$, and $\mathcal{C} := \{x \mid V(x) \le R\}$ s.t.

$$\inf_{x \in \mathcal{C}} P(x, A) \ge \epsilon \nu(A).$$

holds for some $R > 2K/(1 - a)$. Then there exists $\bar{a} \in]a, 1[$, $\bar{b} > 0$, $\bar{\nu} \in \mathcal{P}(\mathbb{R}^d)$, $S \in \mathscr{B}(\mathbb{R}^d)$ s.t.

$$PV(x) \le \bar{a}V(x) + b\mathbb{1}_{\{x \in S\}}, \quad \inf_{x \in S} P(x, A) \ge \bar{\epsilon}\, \bar{\nu}(A).$$

Proof. This argument merely an elaboration on the remark from the end of [57]. Let $\bar{a} \in]a, 1[$ such that $R > b/(\bar{a} - a)$ and $S = \{x \mid V(x) \le R\}$. If $x \notin S$ then

$$PV(x) \le aV(x) + b = aV(x) + \frac{b}{V(x)}V(x) \le aV(x) + \frac{b}{R}V(x)$$

$$\le aV(x) + b\frac{\bar{a} - a}{b}V(x) = aV(x) + (\bar{a} - a)V(x)$$

$$= \bar{a}V(x).$$

If $x \in S$ then $PV(x) \le aV(x) + b \le \bar{a}V(x) + b$. Hence the choices are clear from fixing R as above. $\qquad\square$

Lemma 2. *Let P be a Markov kernel with invariant measure π and suppose there are constants $\rho \in]0, 1[$, $C > 0$, and $r \in]0, 1]$ s.t.*

$$\|P^n f - \pi(f)\|_{V^r} \le C\rho^n \|f\|_{V^r}$$

for any $f : \mathbb{R}^d \to \mathbb{R}$ with $\|f\|_{V^r} < \infty$. Then if $\mu_0(V^r) < \infty$ and μ_n is the flow of P then there is a constant $C' > 0$ s.t.

$$\|\mu_n - \pi\|_{tv, V^r} \le C'\rho^n.$$

Proof. Consider

$$\|\mu_n - \pi\|_{tv, V^r} = \sup_{\|f\|_{V^r} \le 1} |\mu_0 P^n(f) - \pi(f)|$$

$$= \sup_{\|f\|_{V^r} \le 1} |\mu_0(P^n f - \pi(f))|$$

$$\le \sup_{\|f\|_{V^r} \le 1} \mu_0(|P^n f(x) - \pi(f)|)$$

$$\le \sup_{\|f\|_{V^r} \le 1} \mu_0(V^r(x)\|P^n f - \pi(f)\|_{V^r})$$

$$\le C\rho^n \mu_0(V^r) \le C'\rho^n.$$

$\qquad\square$

Lemma 3. *Let P, Q be Markov kernels and suppose that P satisfies a drift condition*

$$PV(x) \le aV(x) + b$$

for $V : \mathbb{R}^d \to [1, \infty[$. Then

$$\|\mu P - \nu P\|_{tv, V} \le (a + b)\|\mu - \nu\|_{tv, V}$$

and

$$\|\mu P - \mu Q\|_{tv, V} \le \mu(V)\|P - Q\|_{ker, V}.$$

Proof. For the first part:

$$\|\mu P - \nu P\|_{tv,V} = \sup_{\|f\|_V \leq 1} |\mu P(f) - \nu P(f)| \leq \sup_{\|f\|_V, \|h\|_V \leq 1} \|Pf\|_V |\mu(h) - \nu(h)|$$

and

$$\|Pf\|_V = \sup_x \frac{|Pf(x)|}{V(x)} = \sup_x \frac{|\int P(x, \mathrm{d}y) f(y)|}{V(x)} = \sup_x \frac{|\int P(x, \mathrm{d}y) V(y) [f(y)/V(y)]}{V(x)}$$

$$\leq \|f\|_V \sup_x \frac{|PV(x)|}{V(x)} \leq \sup_x \frac{aV(x) + b}{V(x)} \leq a + b.$$

For the second part, let $\|f\|_V \leq 1$

$$|\mu P(f) - \mu Q(f)| \leq \mu(|Pf - Qf|) \leq \mu(V\frac{|Pf - Qf|}{V}) \leq \mu(V)\|Pf - Qf\|_V \leq \mu(V)\|P - Q\|_{ker,V}.$$

$\square$

E.2 Proof of Theorem 2

Theorem 2. *Suppose that Assumption 1 and Assumption 2-C1 hold. Suppose also that J_η satisfies*

$$\|J_\eta - J_{\eta'}\|_{ker,\beta} \leq C_J \|\eta - \eta'\|_{tv,\beta} \tag{9}$$

and that

$$K_\eta V(x) \leq \widetilde{a} V(x) + \widetilde{b} \tag{10}$$

for $\widetilde{a} \in]0, 1[$, $\widetilde{b} > 0$ and all $\eta \in \mathcal{P}_{m_G, M_U}$ for suitably chosen constants m_G, M_U. Suppose δ is from (7) *and set*

Case 1: $\rho := (1 - \varepsilon)\gamma$ *if $J_\eta(x, \mathrm{d}y)$ doesn't depend on x; or*

Case 2: $\rho := (1 - \varepsilon)\gamma + \varepsilon \|J_{\eta^\star} V\|_V$ *if $J_\eta(x, \mathrm{d}y)$ does depend on x.*

If $\mu_0(V), \eta_0(U) < \infty$ then there exists a constant $C > 0$ s.t.

$$\|\mu_n - \pi\|_{tv,\beta} \leq \rho^n \|\mu_0 - \pi\|_{tv,\beta} + Cn \max(\rho, \delta)^n.$$

In particular, if $\rho < 1$, we have $\lim_{n \to \infty} \mu_n = \pi$ in V_β-total variation.

Proof. Case 1: For $n = 1$, consider

$$\begin{aligned}
\|\mu_1 - \pi\|_{tv,\beta} &= \|\mu_0 K_{\eta_1} - \pi K_{\eta^\star}\|_{tv,\beta} \\
&\leq (1 - \varepsilon)\|\mu_0 K - \pi K\|_{tv,\beta} + \varepsilon \|\mu_0 J_{\eta_1} - \pi J_{\eta^\star}\|_{tv,\beta} \\
&= (1 - \varepsilon)\|\mu_0 K - \pi K\|_{tv,\beta} + \varepsilon \|J_{\eta_1} - J_{\eta^\star}\|_{tv,\beta} \\
&\leq (1 - \varepsilon)\gamma \|\mu_0 - \pi\|_{tv,\beta} + \varepsilon C_J \|\eta_1 - \eta^\star\|_{tv,\beta} \\
&\leq \rho \|\mu_0 - \pi\|_{tv,V} + \varepsilon C_J M(r^*)\delta \\
&\leq \rho \|\mu_0 - \pi\|_{tv,V} + \varepsilon C \max(\rho, \delta)
\end{aligned}$$

because[9]

$$\|\eta - \eta'\|_{tv,\beta} \leq \|\eta - \eta'\|_{tv,U^{r^*}}$$

due to the Assumption 2-C1. Hence assume true for $n - 1$. Then for n:

$$\begin{aligned}
\|\mu_n - \pi\|_{tv,\beta} &\leq (1 - \varepsilon)\|\mu_{n-1} K - \pi K\|_{tv,\beta} + \varepsilon \|\mu_{n-1} J_{\eta_n} - \pi J_{\eta^\star}\|_{tv,\beta} \\
&= (1 - \varepsilon)\|\mu_{n-1} K - \pi K\|_{tv,\beta} + \varepsilon \|J_{\eta_n} - J_{\eta^\star}\|_{tv,\beta} \\
&\leq (1 - \varepsilon)\gamma \|\mu_{n-1} - \pi\|_{tv,\beta} + \varepsilon C_J \|\eta_n - \eta^\star\|_{tv,\beta} \\
&\leq \rho(\rho^{n-1}\|\mu_0 - \pi\|_{tv,\beta} + C(n-1)\max(\rho, \delta)^{n-1}) + \varepsilon C_J M(r^*)\delta^n \\
&\leq \rho^n \|\mu_0 - \pi\|_{tv,\beta} + C[(n-1)\rho \max(\rho, \delta)^{n-1} + \max(\rho, \delta)^n] \\
&\leq \rho^n \|\mu_0 - \pi\|_{tv,\beta} + C[(n-1)\max(\rho, \delta)^n + \max(\rho, \delta)^n] \\
&= \rho^n \|\mu_0 - \pi\|_{tv,\beta} + Cn \max(\rho, \delta)^n.
\end{aligned}$$

---

[9] this is because $\|f\|_{U^{r^*}} \leq \|f\|_\beta \implies \|\mu - \nu\|_{tv,\beta} \leq \|\mu - \nu\|_{tv,U^{r^*}}$

884 hence Case 1 is done.

885 Case 2: We proceed the same way as case 1 except we need to use

$$\|\mu_{n-1}J_{\eta_n} - \pi J_{\eta^\star}\|_{tv,\beta} \le \|\mu_{n-1}J_{\eta_n} - \mu_{n-1}J_{\eta^\star}\|_{tv,\beta} + \|\mu_{n-1}J_{\eta^\star} - \pi J_{\eta^\star}\|_{tv,\beta}$$
$$\le \mu_{n-1}(V_\beta)\|J_{\eta_n} - J_{\eta^\star}\|_{tv,\beta} + \|J_{\eta^\star}V_\beta\|_\beta\|\mu_{n-1} - \pi\|_{tv,\beta}$$

886 using Lemma 3. Note that since $K_\bullet$ satisfies a uniform drift criterion for V, using linearity we have

$$\mu_n(V_\beta) \le \overline{a}^n\mu_0(V_\beta) + \frac{b}{1-\overline{a}} \le \mu_0(V_\beta) + \frac{b}{1-\overline{a}} =: C_0$$

887 Hence for $n = 1$:

$$\|\mu_1 - \pi\|_{tv,\beta} = \|\mu_0 K_{\eta_1} - \pi K_{\eta^\star}\|_{tv,\beta}$$
$$\le (1-\varepsilon)\|\mu_0 K - \pi K\|_{tv,\beta} + \varepsilon\|\mu_0 J_{\eta_1} - \pi J_{\eta^\star}\|_{tv,\beta}$$
$$\le (1-\varepsilon)\|\mu_0 K - \pi K\|_{tv,\beta} + \varepsilon\mu_0(V_\beta)\|J_{\eta_1} - J_{\eta^\star}\|_{tv,\beta} + \varepsilon\|J_{\eta^\star}V_\beta\|_\beta\|\mu_0 - \pi\|_{tv,\beta}$$
$$\le [(1-\varepsilon)\gamma + \varepsilon\|J_{\eta^\star}V_\beta\|_\beta]\|\mu_0 - \pi\|_{tv,\beta} + \varepsilon C_0 C_J\|\eta_1 - \eta^\star\|_{tv,\beta}$$
$$\le \rho\|\mu_0 - \pi\|_{tv,V} + \varepsilon C_0 C_J M(r^*)\max(\rho,\delta).$$

888 Hence assume true for $n-1$. Then for n:

$$\|\mu_n - \pi\|_{tv,\beta} \le (1-\varepsilon)\|\mu_{n-1}K - \pi K\|_{tv,\beta} + \varepsilon\|\mu_{n-1}J_{\eta_n} - \pi J_{\eta^\star}\|_{tv,\beta}$$
$$\le (1-\varepsilon)\|\mu_{n-1}K - \pi K\|_{tv,\beta} + \varepsilon\mu_{n-1}(V_\beta)\|J_{\eta_n} - J_{\eta^\star}\|_{tv,\beta} + \varepsilon\|J_{\eta^\star}V_\beta\|_\beta\|\mu_{n-1} - \pi\|_{tv,\beta}$$
$$\le (1-\varepsilon)\gamma\|\mu_{n-1} - \pi\|_{tv,\beta} + \varepsilon\mu_{n-1}(V_\beta)\|J_{\eta_n} - J_{\eta^\star}\|_{tv,\beta} + \varepsilon\|J_{\eta^\star}V_\beta\|_\beta\|\mu_{n-1} - \pi\|_{tv,\beta}$$
$$\le \rho(\rho^{n-1}\|\mu_0 - \pi\|_{tv,\beta} + C(n-1)\max(\rho,\delta)^{n-1}) + \varepsilon C_0 C_J M(r^*)\delta^n$$
$$\le \rho^n\|\mu_0 - \pi\|_{tv,\beta} + C[(n-1)\rho\max(\rho,\delta)^{n-1} + \max(\rho,\delta)^n]$$
$$\le \rho^n\|\mu_0 - \pi\|_{tv,\beta} + C[(n-1)\max(\rho,\delta)^n + \max(\rho,\delta)^n]$$
$$= \rho^n\|\mu_0 - \pi\|_{tv,\beta} + Cn\max(\rho,\delta)^n.$$

889 hence Case 2 is done. $\qquad\qquad\square$

F Large-Particle Convergence

891 We study the behaviour of the interacting particle system (3) as $N \to \infty$. The behaviour of this
892 system is probabilistically different from that of the mean field system (2) because the particles in (3)
893 are *coupled* whereas the particles in (2) are *independent*. The fact that we recover independence in
894 the limit of a collection of intercheangeable particles is a remarkable feature of interacting particle
895 systems [41].

F.1 Result

897 For a collection of N particles, under suitable assumptions on $K_\bullet$, we will show that any fixed-size
898 q-block of particles of $\{X_n^1, \ldots, X_n^q\}$ becomes independent as $N \to \infty$, and moreover this trend
899 towards independence happens *uniformly* in time. This phenomenon is called the uniform *propagation*
900 *of chaos* property[10].

901 Let us describe the dynamics of the distribution of the IPS (3). From (3), each X_n^i evolves according
902 to

$$X_n^i \sim K_{m(Y_n)}(X_{n-1}^i, \bullet)$$

903 which indicates that, given Y_n, X_n^i is sampled independently. Letting $X_n^{q,N} := (X_n^1, \ldots, X_n^q)$ and
904 $\mu_n^{q,N} := \text{Distribution}(X_n^{q,N}) \in \mathcal{P}((\mathbb{R}^d)^q)$, for measurable $f : (\mathbb{R}^d)^q \to \mathbb{R}$ one has

$$\mu_n^{q,N}(f) = \mathbb{E}[f(X_n^{q,N})] = \mathbb{E}[\mathbb{E}[f(X_n^{q,N})|Y_n]] = \mathbb{E}[\mu_{n-1}^{q,N}K_{m(Y_n)}^{\otimes q}(f)]$$

905 where the expectation is taken over the distribution of $Y_n = (Y_n^1, \ldots, Y_n^N)$, which is $\eta_n^{\otimes N}$. We will
906 use this decomposition to derive a uniform propagation of chaos result in the following theorem.

---

[10] here, "chaos" is synonymous with "statistical independence", coming from the statistical physics intuition
that a collection of independent particles are maximally disordered, or chaotic. This means that particles which
start chaotic will approximately "propagate their chaos" through time despite interactions between the particles

Theorem 3. *Let* $N > 0$, $q \in \{1, \ldots, N\}$, *and consider the interacting particle system* $X_n :=$ $(X_n^1, \ldots, X_n^N)$, $Y_n := (Y_n^1, \ldots, Y_n^N)$ *from* (3). *Let* $\mu_n^{q,N} := \text{Distribution}(X_n^1, \ldots, X_n^q)$, *and let* μ_n *be the distribution of the (independent) mean field system* (2), *with* $X_0^i \sim \mu_0$. *Suppose that* $\forall x \in (\mathbb{R}^d)^q$, $\eta \in \mathcal{P}(\mathbb{R}^d)$, *and* $f \in \mathcal{B}_b((\mathbb{R}^d)^q)$ *with* $osc(f) \leq 1$[11]

$$\left| \mathbb{E}[J_{m(Y)}^{\otimes q} f(x)] - J_\eta^{\otimes q} f(x) \right| \leq c \frac{q^2}{N} \mathcal{R}(q^2/N), \text{ where } Y = \{Y^1, \ldots, Y^N\}, Y^i \sim \eta.$$

Suppose finally that $\mu_0^{q,N} = \mu_0^{\otimes q}$ *for any* $1 \leq q \leq N$.

If **(Case 1)***:* $J_\eta(x, \bullet)$ *doesn't depend on* x, *or if* **(Case 2)***:* $J_\eta(x, \bullet)$ *does depend on* x *but additionally that* $\epsilon(K) < 1$, *then there exists a fixed constant* $C > 0$ *s.t.*

$$\sup_{n \geq 0} \|\mu_n^{q,N} - \mu_n^{\otimes q}\|_{tv} \leq C \frac{q^2}{N} \mathcal{R}(q^2/N).$$

Proof. **Case 1**: We first claim that

$$\|\mu_{n+1}^{q,N} - \mu_{n+1}^{\otimes q}\|_{tv} \leq c\varepsilon \sum_{j=0}^{n-1} (1-\varepsilon)^j \epsilon(K^{\otimes q})^j \cdot \frac{q^2}{N} \mathcal{R}(2q^2/N).$$

Let $f \in \mathcal{B}_b((\mathbb{R}^d)^q)$ s.t. $osc(f) \leq 1$.

Consider the following expression for general n:

$$\begin{aligned}
\left| \mu_{n+1}^{q,N}(f) - \mu_{n+1}^{\otimes q}(f) \right| &= \left| \mathbb{E}[\mu_n^{q,N} K_{m(Y_n)}^{\otimes q}(f)] - \mu_n^{\otimes q} K_{\eta_n}^{\otimes q}(f) \right| \\
&= \left| \mathbb{E}[\mu_n^{q,N} K_{m(Y_n)}^{\otimes q}(f)] - \mu_n^{q,N} K_{\eta_n}^{\otimes q}(f) + \mu_n^{q,N} K_{\eta_n}^{\otimes q}(f) - \mu_n^{\otimes q} K_{\eta_n}^{\otimes q}(f) \right| \\
&\leq \left| \mathbb{E}[\mu_n^{q,N} K_{m(Y_n)}^{\otimes q}(f)] - \mu_n^{q,N} K_{\eta_n}^{\otimes q}(f) \right| + \left| \mu_n^{q,N} K_{\eta_n}^{\otimes q}(f) - \mu_n^{\otimes q} K_{\eta_n}^{\otimes q}(f) \right| \\
&\leq \sup_x \varepsilon \left| \mathbb{E}[J_{m(Y_n)}^{\otimes q} f(x)] - J_{\eta_n}^{\otimes q} f(x) \right| + (1-\varepsilon) \left| \mu_n^{q,N} K^{\otimes q}(f) - \mu_n^{\otimes q} K^{\otimes q}(f) \right| \\
&\leq c\varepsilon \frac{q^2}{N} \mathcal{R}(q^2/N) + (1-\varepsilon)\epsilon(K^{\otimes q}) \left| \mu_n^{q,N}(f) - \mu_n^{\otimes q}(f) \right|.
\end{aligned}$$

where we have used the assumption on J.

Thus if $n = 1$ in the above expression, the base case holds since $\mu_0^{q,N} = \mu_0^{\otimes q}$. Now, if the claim holds for n, then for $n + 1$ we have

$$\begin{aligned}
|\mu_{n+1}^{q,N}(f) - \mu_{n+1}^{\otimes q}(f)| &\leq c\varepsilon \frac{q^2}{N} \mathcal{R}(q^2/N) + (1-\varepsilon)\epsilon(K^{\otimes q}) c\varepsilon \sum_{j=0}^{n-1} (1-\varepsilon)^j \epsilon(K^{\otimes q})^j \cdot \frac{q^2}{N} \mathcal{R}(q^2/N) \\
&= c\varepsilon \frac{q^2}{N} \mathcal{R}(q^2/N) \left[1 + (1-\varepsilon)\epsilon(K^{\otimes q}) \sum_{j=0}^{n-1} (1-\varepsilon)^j \epsilon(K^{\otimes q})^j \right] \\
&= c\varepsilon \frac{q^2}{N} \mathcal{R}(q^2/N) \sum_{j=0}^{n} (1-\varepsilon)^j \epsilon(K^{\otimes q})^j.
\end{aligned}$$

Thus the claim holds for $n + 1$. Since $(1-\varepsilon)\epsilon(K^{\otimes q}) < 1$, we have the result.

Case 2: We first claim that

$$\|\mu_{n+1}^{q,N} - \mu_{n+1}^{\otimes q}\|_{tv} \leq c\varepsilon \sum_{j=0}^{n-1} [\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q})]^j \cdot \frac{q^2}{N} \mathcal{R}(q^2/N).$$

---

[11] due to the characterization $\|\mu - \nu\|_{tv} = \sup\{|\mu(f) - \nu(f)| \mid f \in \mathcal{B}_b(\mathbb{R}^d), \ osc(f) \leq 1\}$ from [47] this regularity condition should be interpreted as a total variation Lipschitzness analogous to (9).

922   Let $f \in \mathcal{B}_b((\mathbb{R}^d)^q)$ with $osc(f) \leq 1$. We proceed by induction. Note that:

$$\left|\mu_{n+1}^{q,N}(f) - \mu_{n+1}^{\otimes q}(f)\right| = \left|\mathbb{E}[\mu_n^{q,N} K_{m(Y_n)}^{\otimes q}(f)] - \mu_n^{\otimes q} K_{\eta_n}^{\otimes q}(f)\right|$$

$$= \varepsilon \left|\mathbb{E}[\mu_n^{q,N} J_{m(Y_n)}^{\otimes q}(f)] - \mu_n^{\otimes q} J_{\eta_n}^{\otimes q}(f)\right| + (1 - \varepsilon)\left|\mu_n^{q,N} K(f) - \mu_n^{\otimes q} K(f)\right|$$

923   for the first term, we will use the decomposition

$$\left|\mathbb{E}[\mu_n^{q,N} J_{m(Y_n)}(f)] - \mu_n^{\otimes q} J_{\eta_n}(f)\right| \leq \left|\mathbb{E}[\mu_n^{q,N} J_{m(Y_n)}(f)] - \mu_n^{q,N} J_{\eta_n}(f)\right| + \left|\mu_n^{q,N} J_{\eta_n}(f) - \mu_n^{\otimes q} J_{\eta_n}(f)\right|$$

924   Then, using the assumption on J, we have

$$\left|\mathbb{E}[\mu_n^{q,N} J_{m(Y_n)}(f)] - \mu_n^{q,N} J_{\eta_n}(f)\right| \leq c\frac{q^2}{N}\mathcal{R}(q^2/N))$$

925   and also we know we can use

$$\left|\mu_n^{q,N} J_{\eta_n}(f) - \mu_n^{\otimes q} J_{\eta_n}(f)\right| \leq osc(J_{\eta_n} f)\left|\mu_n^{q,N}(h) - \mu_n^{\otimes q}(h)\right|$$

$$\leq \epsilon(J_{\eta_n})osc(f)\left|\mu_n^{q,N}(h) - \mu_n^{\otimes q}(h)\right|$$

$$\leq \left|\mu_n^{q,N}(h) - \mu_n^{\otimes q}(h)\right|$$

926   since $\epsilon(J_{\eta_n}) \leq 1$, with $osc(h) \leq 1$. Hence putting these together, we see

$$\left|\mu_{n+1}^{q,N}(f) - \mu_{n+1}^{\otimes q}(f)\right| \leq \varepsilon c\frac{q^2}{N}\mathcal{R}(q^2/2N) + \varepsilon\left|\mu_n^{q,N}(h) - \mu_n^{\otimes q}(h)\right| + (1-\varepsilon)\epsilon(K^{\otimes q})\left|\mu_n^{q,N}(f) - \mu_n^{\otimes q}(f)\right|$$

$$\leq \varepsilon c\frac{q^2}{N}\mathcal{R}(q^2/2N) + (\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q}))\|\mu_n^{q,N} - \mu_n^{\otimes q}\|_{tv}.$$

927   Now if $n = 0$ in the above expression, the base case holds since $\mu_0^{q,N} = \mu_0^{\otimes q}$. If it is true for n, then
928   for $n + 1$ we have

$$\left|\mu_{n+1}^{q,N}(f) - \mu_{n+1}^{\otimes q}(f)\right| = \left|\mathbb{E}[\mu_n^{q,N} K_{m(Y_n)}(f)] - \mu_n^{\otimes q} K_{\eta_n}(f)\right|$$

$$\leq c\varepsilon\frac{q^2}{N}\mathcal{R}(q^2/N) + (\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q}))\|\mu_n^{q,N}(f) - \mu_n^{\otimes q}(f)\|_{tv}$$

$$\leq c\varepsilon\frac{q^2}{N}\mathcal{R}(q^2/N) + (\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q}))\left[c\varepsilon\sum_{j=0}^{n-1}[\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q})]^j \cdot \frac{q^2}{N}\mathcal{R}(2q^2/N)\right]$$

$$= c\varepsilon\frac{q^2}{N}\mathcal{R}(q^2/N)\left[1 + \sum_{j=1}^{n}[\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q})]^j\right]$$

$$= c\varepsilon\frac{q^2}{N}\mathcal{R}(q^2/N)\sum_{j=0}^{n}[\varepsilon + (1-\varepsilon)\epsilon(K^{\otimes q})]^j.$$

929   Hence if $\epsilon(K^{\otimes q}) < 1$, i.e. $\epsilon(K) < 1$, the result holds. $\square$

F.2   Proof of Main Results (Theorem 1 & Corollary 1)

931   **Theorem 1.** *[Convergence of Nonlinear MCMC] Under suitable conditions on K_η and Q, there*
932   *exist fixed constants $C_1, C_2, C_3 > 0$, a function $\mathcal{R} : [0, \infty[\to [1, \infty[$, and $\rho > 0$ s.t.*

$$\|\mu_n^N - \pi\|_{tv} \leq C_1 \frac{1}{N}\mathcal{R}(1/N) + C_2\rho^n + C_3 n\rho^n.$$

933   *Proof.* This follows straightforwardly from the above discussion, the only technicality is converting
934   the results from Theorem 2 to the un-weighted total variation. But note that since $\{\|f\|_\infty \leq 1\} \subset$
935   $\{\|f\|_\beta \leq 1\}$ we have

$$\|\mu_n - \mu_\infty\|_{tv} = \sup_{\|f\|_\infty \leq 1} |\mu_n(f) - \mu_\infty(f)| \leq \sup_{\|f\|_\beta \leq 1} |\mu_n(f) - \mu_\infty(f)| = \|\mu_n - \mu_\infty\|_{tv,\beta}$$

936   so we're done. $\square$

Corollary 1. *[Adapted from [41], Theorem 2.2] Suppose that Theorem 1 applies to K_η. Let* $\overline{X}_n := \{X_n^1, \ldots, X_n^N\}$ *be from* (3). *Then for every $n \in \mathbb{N}$ and $f \in \mathcal{B}_b(\mathbb{R}^d)$ we have*

$$\lim_{N \to \infty} \mathbb{E}\left[\left\|\frac{1}{N}\sum_{i=1}^N f(X_n^i) - \mu_n(f)\right\|\right] = 0.$$

Proof. Let $f \in \mathcal{B}_b(\mathbb{R}^d)$ and consider

$$
\begin{aligned}
\mathbb{E}[(m(X_n)(f) - \mu_n(f))^2] &= \mathbb{E}\left[\left(\frac{1}{N}\sum_{i=1}^N f(X_n^i) - \mu_n(f)\right)^2\right] \\
&= \frac{1}{N^2}\sum_{i=1}^n \mathbb{E}[f(X_n^i)f(X_n^j)] - \frac{2}{N}\sum_{i=1}^N \mathbb{E}[f(X_n^i)]\mu_n(f) + \mu_n(f)^2 \\
&= \frac{1}{N}\mathbb{E}[f(X_n^1)^2] + \frac{N-1}{N}\mathbb{E}[f(X_n^1)f(X_n^2)] - 2\mathbb{E}[f(X_n^1)]\mu_n(f) + \mu_n(f)^2
\end{aligned}
$$

using interchangeability of the X_n^is. Since the strong propagation of chaos in Theorem 3 clearly implies weak propagation of chaos, i.e. for any bounded f $|\mu_n^{q,N}(f) - \mu_n^{\otimes q}(f)| \to 0$, we see that the expression above $\to 0$ and we have L^2-convergence which implies weak convergence. $\qquad\square$

G Analysis of the Kernels K_η^{BG} and K_η^{AR}

G.1 Analysis of K_η^{BG}

G.1.1 Statement of Results

A key difficulty that arises when working with $\Psi_G(\eta) = \frac{G\mathrm{d}\eta}{\eta(G)}$ is obtaining a uniform lower bound the denominator $\eta(G)$. This is important for deriving uniform regularity results and for generally establishing conditions under which the measures $\{\Psi_G(\eta_n)\}_{n=0}^\infty$ are well-defined. An effective approach used in [47] and related works is to assume the uniform lower bound $G(x) > \epsilon > 0 \ \forall x \in \mathbb{R}^d$ for some $\epsilon > 0$. While this assumption is self-contained in that it works for any choice of η, it eliminates ubiquitous families of measures such as Gaussians, e.g. $G(x) = \exp(-\|x\|^2/2)$ and $\mathrm{d}\eta = \mathrm{d}x$.

We will see below how to relax $G(x) > \varepsilon > 0$ using the structure of our problem. In particular, we can use the Lyapunov function U for Q to control the probability that the state $Y_n \sim \eta_n$ will venture "far away" from the center of the state space. This leads to the intuition that, if G is sufficiently "compatible" with U, i.e. the function G stays away from zero in the "center" of the state space as determined by U, then the expectation $\eta_n(G) = \mathbb{E}_{Y_n \sim \eta_n}[G(Y_n)]$ can be lower-bounded using the same Lyapunov function. This insight is encoded in the compatibility criterion Assumption 2-C2, and we can use it to prove the following *a priori* lower bound on $\eta_n(G)$.

Lemma 4. *Suppose that Assumption 1-Q1 and Assumption 2-C2 hold, and let $\eta_n = \eta_{n-1}Q$ with $\eta_0(U) < \infty$. Then we have the lower bound*

$$\eta_n(G) \geq \theta(R^*)\left(1 - \frac{\gamma^n \eta_0(U) + \frac{c}{1-\gamma}}{R^*}\right)$$

where R^* is fixed and doesn't depend on n.

This lemma essentially says that G should be bounded away from zero on the level sets of U. This is not a strong condition — if G is bounded away from zero on compact sets and U is continuous, the lemma applies. As an example, if $G = \exp(-\|x\|^2/2)$ and $U(x) = c\|x\|^2 + 1$ then this result applies. The constant R^* arises since $R \mapsto \theta(R)$ is nonincreasing and $R \mapsto 1 - \frac{1}{R}$ is increasing so we can optimize this bound as a function of R to get R^* (which may not be unique, but the value the bound attains will be). Together with the next lemma, we can obtain the desired convergence from Theorem 2 for the BG interaction.

Lemma 5. *Let $\eta, \eta' \in \mathcal{P}(\mathbb{R}^d)$, and suppose that Assumption 3 holds, i.e. $\|G\|_\infty, \|G\|_\beta < \infty$. Then*

$$\|\Psi_G(\eta) - \Psi_G(\eta')\|_{tv,\beta} \leq \left(\frac{\|G\|_\beta + \|G\|_\infty}{\eta(G) \vee \eta'(G)} + \eta(V) \wedge \eta'(V)\beta \frac{\|G\|_\beta \|G\|_\infty}{\eta(G)\eta'(G)} \right) \|\eta - \eta'\|_{tv,\beta}.$$

Clearly, we will use our knowledge that $\eta_n \in \mathcal{P}_{m_G, M_U}(\mathbb{R}^d) \; \forall n$ to make the Lipschitz constant in Lemma 5 uniform over $\mathcal{P}_{m_G, M_U}$.

Proposition 2. *Suppose that Assumption 1, 2, and Assumption 3-G1 hold. Then the drift criterion (10) holds uniformly over all $\eta \in \mathcal{P}_{m,M}(\mathbb{R}^d)$.*

Now we can apply Theorem 2 to obtain convergence.

Corollary 2. *Suppose that Assumptions 1, 2, 3 hold. If $\eta_0(G) > 0$, $\eta_0(U), \mu_0(V) < \infty$, then Theorem 2 holds for K^{BG}, i.e. the flow μ_n converges to π in V_β-total variation as long as $\rho = (1-\varepsilon)\gamma < 1$.*

Proof. Lemma 4 and Lemma 5 imply the regularity condition (9) for constants determined by those lemmas and by noting that, due to Asssumption 2-C1, $\eta(V_\beta) \leq \eta(U^{r^*}) \leq \eta(U)^{r^*} < \infty$ by Jensen's inequality. Additionally, Proposition 2 implies the condition (10). Hence, since we are in "Case 1" of Theorem 2, the result follows. $\qquad\square$

G.1.2 Proofs

Lemma 4. *Suppose that Assumption 1-Q1 and Assumption 2-C2 hold, and let $\eta_n = \eta_{n-1}Q$ with $\eta_0(U) < \infty$. Then we have the lower bound*

$$\eta_n(G) \geq \theta(R^*)\left(1 - \frac{\gamma^n \eta_0(U) + \frac{c}{1-\gamma}}{R^*}\right)$$

where R^* is fixed and doesn't depend on n.

Proof.

$$\begin{aligned}
QG(x) &= \mathbb{E}_{X \sim Q(x, \bullet)}[G(X)] \\
&= \mathbb{E}_{X \sim Q(x, \bullet)}[G(X)\mathbb{1}_{\{U(X) \leq R\}}] + \mathbb{E}_{X \sim Q(x, \bullet)}[G(X)\mathbb{1}_{\{U(X) > R\}}] \\
&\geq \theta(R) \cdot \mathbb{P}_x(U(X) \leq R) + \mathbb{E}_{X \sim Q(x, \bullet)}[G(X)\mathbb{1}_{\{U(X) > R\}}] \\
&\geq \theta(R) \cdot \mathbb{P}_x(U(X) \leq R).
\end{aligned}$$

But by Markov's inequality

$$\mathbb{P}_x(U(X) > R) \leq \frac{\mathbb{E}_x[U(X)]}{R} \leq \frac{\xi U(x) + c}{R}$$

so

$$\mathbb{P}_x(U(X) \leq R) = 1 - \mathbb{P}_x(U(X) > R) \geq 1 - \frac{\xi U(x) + c}{R}$$

and hence

$$QG(x) \geq \theta(R)\mathbb{P}_x(U(X) \leq R) \geq \theta(R)\left(1 - \frac{\xi U(x) + c}{R}\right).$$

Now

$$\begin{aligned}
Q^2 G(x) = Q(QG(x)) &\geq Q\left(\theta(R)\left(1 - \frac{\xi U(x) + c}{R}\right)\right) \\
&= \theta(R)\left(1 - \frac{\xi QU(x) + c}{R}\right) \geq \theta(R)\left(1 - \frac{\xi^2 U(x) + c(1 + \xi)}{R}\right)
\end{aligned}$$

and iterating this procedure gives

$$Q^n G(x) \geq \theta(R)\left(1 - \frac{\xi^n U(x) + \frac{c}{1-\xi}}{R}\right)$$

where we have used the sum of the geometric series to obtain $c/(1-\xi)$. Now, $\theta(R)$ is nonincreasing w.r.t. R and $1-1/R$ is increasing w.r.t. R, so optimizing to get R^* and integrating we obtain

$$\eta_n(G) = \eta_0(Q^n G) \geq \theta(R^*)\left(1 - \frac{\xi^n \eta_0(U) + \frac{c}{1-\xi}}{R^*}\right).$$

$\square$

Lemma 5. *Let $\eta, \eta' \in \mathcal{P}(\mathbb{R}^d)$, and suppose that Assumption 3 holds, i.e. $\|G\|_\infty, \|G\|_\beta < \infty$. Then*

$$\|\Psi_G(\eta) - \Psi_G(\eta')\|_{tv,\beta} \leq \left(\frac{\|G\|_\beta + \|G\|_\infty}{\eta(G) \vee \eta'(G)} + \eta(V) \wedge \eta'(V)\beta\frac{\|G\|_\beta\|G\|_\infty}{\eta(G)\eta'(G)}\right)\|\eta - \eta'\|_{tv,\beta}.$$

Proof. Let $\|f\|_\beta \leq 1$. then

$$|\Psi_G(\eta)(f) - \Psi_G(\eta')(f)| = \left|\int \frac{G(x)f(x)}{\eta(G)}\eta(\mathrm{d}x) - \int \frac{G(x)f(x)}{\eta'(G)}\eta'(\mathrm{d}x)\right|$$

$$\leq \left|\int \frac{G(x)f(x)}{\eta(G)}\eta(\mathrm{d}x) - \int \frac{G(x)f(x)}{\eta'(G)}\eta(\mathrm{d}x)\right| + \left|\int \frac{G(x)f(x)}{\eta'(G)}\eta(\mathrm{d}x) - \int \frac{G(x)f(x)}{\eta'(G)}\eta'(\mathrm{d}x)\right|$$

for the first term

$$\left|\int \left(\frac{G(x)f(x)}{\eta(G)} - \frac{G(x)f(x)}{\eta'(G)}\right)\eta(\mathrm{d}x)\right|$$

$$\leq \frac{|\eta(G) - \eta'(G)|}{\eta(G)\eta'(G)}\int G(x)|f(x)|\eta(\mathrm{d}x)$$

$$\leq \frac{\|G\|_\beta\|\eta - \eta'\|_{tv,\beta}}{\eta(G)\eta'(G)}\int G(x)|f(x)|\eta(\mathrm{d}x)$$

$$\leq \frac{\|G\|_\beta\|\eta - \eta'\|_{tv,\beta}}{\eta(G)\eta'(G)}\eta(G(1 + \beta V))\|f\|_\beta$$

$$= \left(\frac{\|G\|_\beta}{\eta'(G)} + \beta\eta(V)\frac{\|G\|_\beta\|G\|_\infty}{\eta(G)\eta'(G)}\right)\|\eta - \eta'\|_{tv,\beta}$$

and for the second

$$\left|\int \frac{G(x)f(x)}{\eta'(G)}\eta(\mathrm{d}x) - \int \frac{G(x)f(x)}{\eta'(G)}\eta'(\mathrm{d}x)\right|$$

$$= \frac{1}{\eta'(G)}\left|\int G(x)f(x)\eta(\mathrm{d}x) - \int G(x)f(x)\eta'(\mathrm{d}x)\right|$$

$$\leq \frac{\|fG\|_\beta}{\eta'(G)}\|\eta - \eta'\|_{tv,\beta}$$

$$\leq \frac{\|G\|_\infty\|f\|_\beta}{\eta'(G)}\|\eta - \eta'\|_{tv,\beta} = \frac{\|G\|_\infty}{\eta'(G)}\|\eta - \eta'\|_{tv,\beta}$$

since

$$\|fG\|_\beta = \sup_x \frac{|f(x)G(x)|}{1 + \beta V(x)} \leq \|G\|_\infty\|f\|_\beta$$

so putting these together

$$\|\Psi_G(\eta) - \Psi_G(\eta')\|_{tv,\beta} \leq \left(\frac{\|G\|_\beta + \|G\|_\infty}{\eta'(G)} + \eta(V)\beta\frac{\|G\|_\beta\|G\|_\infty}{\eta(G)\eta'(G)}\right)\|\eta - \eta'\|_{tv,\beta}$$

Using symmetry completes the proof. $\square$

Proposition 2. *Suppose that Assumption 1, 2, and Assumption 3-G1 hold. Then the drift criterion (10) holds uniformly over all $\eta \in \mathcal{P}_{m,M}(\mathbb{R}^d)$.*

Proof. Let $\eta \in \mathcal{P}_{m,M}(\mathbb{R}^d)$ and consider

$$K_\eta V(x) = (1 - \varepsilon)KV(x) + \varepsilon \Psi_G(\eta)(V)$$

$$\leq (1 - \varepsilon)aV(x) + (1 - \varepsilon)b + \varepsilon \frac{\eta(V)}{\eta(G)}$$

$$\leq (1 - \varepsilon)aV(x) + (1 - \varepsilon)b + \varepsilon \frac{M}{m}.$$

$\square$

Theorem 4 ([47] Thm 8.7.1 pp.283)**.** *Suppose that G has bounded oscillations, let $N \geq q \geq 1$, and $Y = (Y^1, \ldots, Y^N)$, $Y^i \sim \eta$. Then for any $f \in \mathcal{B}_b((\mathbb{R}^d)^q)$ with $\mathrm{osc}(f) \leq 1$, we have*

$$|\mathbb{E}[\Psi_{G^{\otimes q}}(m(Y)^{\otimes q})(f) - \Psi_{G^{\otimes q}}(\eta^{\otimes q})(f)]| \leq c \frac{q^2}{N} \mathcal{R}_{G,\eta}(2q^2/N)$$

where

$$\mathcal{R}_{G,\eta}(u) := 1 + \mathrm{osc}_\eta(G)^2(1 + \mathrm{osc}_\eta(G)\sqrt{u})\exp(\mathrm{osc}_\eta(G)^2 u), \quad \mathrm{osc}_\eta(G) := \mathrm{osc}(G/\eta(G)).$$

In particular, by picking $G \equiv 1$, we obtain

$$|\mathbb{E}[m(Y)^{\otimes q}(f) - \eta^{\otimes q}(f)]| \leq c \frac{q^2}{N}.$$

Corollary 3. *Suppose that Assumption 3 G1 holds (i.e. G is bounded) the compatibility criterion Assumption 2-C2 with $\eta_n(G) \geq m$. Then $K_\bullet^{BG}$ satisfies the uniform propagation of chaos in Case 1 with*

$$\mathcal{R}(u) = \mathcal{R}_G(u) := 1 + \frac{\mathrm{osc}(G)^2}{m^2}\left(1 + \frac{\mathrm{osc}(G)}{m}\sqrt{u}\right)\exp\left(\frac{\mathrm{osc}(G)^2}{m^2}u\right)$$

Proof. This follows from Theorem 3 and Theorem 4 above, noting 1) that $\|G\|_\infty$ implies $\mathrm{osc}(G) < \infty$, and 2) that we can obtain a bound on $\mathcal{R}_{G,\eta_n}$ independent of η_n by applying Lemma 4 to obtain $\eta_n(G) \geq m$ so

$$osc_{\eta_n}(G) = \mathrm{osc}(G/\eta_n(G)) \leq \mathrm{osc}(G)/m.$$

$\square$

G.2   Analysis of K_η^{AR}

G.2.1   Statement of Results

First, the result showing that π is $K_{\eta^\star}^{AR}$-invariant.

Proposition 3. $K_{\eta^\star}$ *is π-invariant.*

Proof. We have

$$\pi(J_{\eta^\star}f) = \iint [f(y) - f(x)]\alpha(x,y)\eta^\star(\mathrm{d}y)\pi(\mathrm{d}x)$$

$$= \iint [f(y) - f(x)]1 \wedge \frac{\pi(y)\eta^\star(x)}{\eta^\star(y)\pi(x)}\eta^\star(\mathrm{d}y)\pi(\mathrm{d}x) + \pi(f)$$

$$= \iint [f(y) - f(x)]\pi(x)\eta^\star(y) \wedge \pi(y)\eta^\star(x)\mathrm{d}y\mathrm{d}x + \pi(f)$$

$$= \pi(f)$$

since the integrals with $f(y)$ and $f(x)$ in the first term are equal. $\square$

We will establish equvalent results from Section G.1but for $K_\bullet^{AR}$. This will not require assumption 2-C2 since the interaction is well-defined for $\eta \in \mathcal{P}(\mathbb{R}^d)$ as $\alpha(x,y)$ is in fact *bounded*. K^{AR} is the main subject of study for [1], and in fact the regularity and uniform drift conditions were established there. Note that the statement $\eta \in \mathcal{P}_{0,\infty}$ is vaccuous since $G(x) > 0 \implies \eta(G) > 0$, and the second says that $\eta(V) < \infty$.

1028 **Lemma 6** ([1]). *Let $\eta, \eta' \in \mathcal{P}_{0,\infty}(\mathbb{R}^d)$. Then*

$$\|J_\eta^{AR} - J_{\eta'}^{AR}\|_{ker,\beta} \leq 2\|\eta - \eta'\|_{tv,\beta}.$$

1029 **Lemma 7.** *Suppose that Assumption 1, 2-C2, hold. Then the uniform drift criterion (10) holds for*
1030 $\eta \in \mathcal{P}_{0,M}(\mathbb{R}^d)$.

1031 **Proposition 4.** *Let $J = J^{AR}$, and $Y = \{Y^1, \ldots, Y^N\}$ where $Y^i \overset{iid}{\sim} \eta$. Then for any $f \in \mathcal{B}_b((\mathbb{R}^d)^q$*
1032 *with $\mathrm{osc}(f) \leq 1$ and $x \in (\mathbb{R}^d)^q$, we have*

$$|\mathbb{E}[J_{m(Y)}^{\otimes q} f(x)] - J_\eta^{\otimes q} f(x)| \leq 2c\frac{q^2}{N}.$$

1033 **Corollary 4.** *Suppose that Assumption 1, 2-C2 hold. If $\eta_0(U), \mu_0(V) < \infty$, then Theorem 2 holds for*
1034 K^{BG}, *i.e. the flow μ_n converges to π in V_β-total variation as long as $\rho = (1 - \varepsilon)\gamma + \varepsilon\|J_{\eta_0}V_\beta\| < 1$.*

G.2.2 Proofs

1036 **Lemma 7.** *Suppose that Assumption 1, 2-C2, hold. Then the uniform drift criterion (10) holds for*
1037 $\eta \in \mathcal{P}_{0,M}(\mathbb{R}^d)$.

1038 *Proof.* Let $\eta \in \mathcal{P}_{0,M}(\mathbb{R}^d)$ and consider

$$\begin{aligned}
K_\eta^{AR}V(x) &= (1 - \varepsilon)KV(x) + \varepsilon[\eta(\alpha(x, \bullet)V) + (1 - A_\eta(x))V(x)] \\
&\leq (1 - \varepsilon)aV(x) + (1 - \varepsilon)b + \varepsilon\eta(V) + \varepsilon V(x) \\
&\leq [(1 - \varepsilon)a + \varepsilon]V(x) + (1 - \varepsilon)b + \varepsilon M.
\end{aligned}$$

1039 $\qquad\square$

1040 **Proposition 4.** *Let $J = J^{AR}$, and $Y = \{Y^1, \ldots, Y^N\}$ where $Y^i \overset{iid}{\sim} \eta$. Then for any $f \in \mathcal{B}_b((\mathbb{R}^d)^q$*
1041 *with $\mathrm{osc}(f) \leq 1$ and $x \in (\mathbb{R}^d)^q$, we have*

$$|\mathbb{E}[J_{m(Y)}^{\otimes q} f(x)] - J_\eta^{\otimes q} f(x)| \leq 2c\frac{q^2}{N}.$$

1042 *Proof.* Starting with the first term, for each fixed x

$$\begin{aligned}
|\mathbb{E}[J_{m(Y_n)}^{\otimes q} f(x)] - J_{\eta_n}^{\otimes q} f(x)| &= \left| \mathbb{E}\left[\int [f(y) - f(x)]\alpha^{\otimes q}(x, y)m(Y_n)^{\otimes q}(\mathrm{d}y) \right] - \int [f(y) - f(x)]\alpha^{\otimes q}(x, y)\eta_n^{\otimes q}(\mathrm{d}y) \right| \\
&= |\mathbb{E}[m(Y_n)^{\otimes q}(\varphi_f(x, \bullet))] - \eta_n^{\otimes q}(\varphi_f(x, \bullet))|
\end{aligned}$$

1043 where

$$\varphi_f(x, y) := \alpha^{\otimes q}(x, y)[f(y) - f(x)].$$

1044 Now

$$\sup_y |\varphi_f(x, y)| = \sup_y |[f(y) - f(x)]\alpha^{\otimes q}(x, y)| \leq \sup_y |f(y) - f(x)|\|\alpha^{\otimes q}(x, \bullet)\|_\infty \leq \mathrm{osc}(f) \leq 1$$

1045 so automatically $\mathrm{osc}(\varphi_f(x, \bullet)) \leq 2$. Hence using Lemma 4

$$|\mathbb{E}[m(Y_n)^{\otimes q}(\varphi_f(x, \bullet))] - \eta_n^{\otimes q}(\varphi_f(x, \bullet))| \leq 2c\frac{q^2}{N}.$$

1046 $\qquad\square$

1047 **Corollary 5.** *If $\epsilon(K) < 1$, then $K_\bullet^{AR}$ satisfies the uniform propagation of chaos in Case 2 of*
1048 *Theorem 3 with $\mathcal{R} \equiv 1$.*