Can we globally optimize cross-validation loss? Quasiconvexity in ridge regression

Abstract

Models like LASSO and ridge regression are extensively used in practice due to their interpretability, ease of use, and strong theoretical guarantees. Cross-validation (CV) is widely used for hyperparameter tuning in these models, but do practical methods minimize the true out-of-sample loss? A recent line of research promises to show that the optimum of the CV loss matches the optimum of the out-of-sample loss (possibly after simple corrections). It remains to show how tractable it is to minimize the CV loss. In the present paper, we show that, in the case of ridge regression, the CV loss may fail to be quasiconvex and thus may have multiple local optima. We can guarantee that the CV loss is quasiconvex in at least one case: when the spectrum of the covariate matrix is nearly flat and the noise in the observed responses is not too high. More generally, we show that quasiconvexity status is independent of many properties of the observed data (response norm, covariate-matrix right singular vectors and singular-value scaling) and has a complex dependence on the few that remain. We empirically confirm our theory using simulated experiments.

1 Introduction

Linear models, including LASSO and ridge regression, are widely used for data analysis across diverse applied disciplines. Linear models are often preferred since they are straightforward to apply in various senses. In particular, (1) their parameters are readily interpretable. (2) They have strong theoretical guarantees on quality. And (3) standard optimization tools are often assumed to find useful parameter and hyperparameter values. Despite their seeming simplicity, though, mysteries remain about the quality of inference in linear models. Consider cross-validation (CV) [Stone, 1974, Allen, 1974], the de facto standard for hyperparameter selection across machine learning methods [Musgrave et al., 2020]. CV is an easy-to-evaluate proxy for the true out-of-sample loss. Is it a good proxy? [Homrighausen and McDonald, 2014, 2013, Chetverikov et al., 2020, Hastie et al., 2020] give conditions under which the optimum of the CV loss (possibly with some mild corrections) matches the optimum of the out-of-sample loss in LASSO and ridge regression. To complete the picture, we must understand whether standard methods for minimizing the CV loss find a global minimum.

It would be easy to find a unique minimum of the CV loss if the CV loss were convex. Alas (though perhaps unsurprisingly), we show below that in essentially every case of interest the CV loss is not convex. Indeed, the usual introductory cartoon of CV loss (left panel of Fig. 1; see also Fig. 5.9 of Hastie et al. [2017] or Fig. 1 of Rad and Maleki [2020]) is not convex. But the cartoon CV loss still exhibits a single global minimum and would be easy to globally minimize with popular approaches like gradient-based methods [Do et al., 2007, Maclaurin et al., 2015, Pedregosa, 2016, Lorraine et al., 2020] or grid search [Bergstra and Bengio, 2012, Pedregosa et al., 2011, Hsu et al., 2003]. Indeed, a

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

Figure 1: (*Left*): Idealized illustration of the leave-one-out CV loss, $\mathcal{L}$, (blue) and the true out-of-sample loss (black). The minimum of each curve is marked with a vertical line of the corresponding color. (*Center*): CV loss for a life-expectancy prediction problem after some standard data pre-processing (Condition 1 of Section 2). (*Right*): CV loss for wine-quality prediction problem on a subset of $N = 50$ data points after standard data pre-processing (Condition 1 of Section 2).

more plausible possibility (which holds for the typical cartoon CV loss) is that the CV loss might be *quasiconvex*. In one dimension, quasiconvexity implies a that any local optimum is a global optimum.

In what follows, we show that this cartoon need not hold in general. We consider using the leave-one-out CV (LOOCV) loss, $\mathcal{L}$, to select the regularization hyperparameter in ℓ_2-regularized linear regression (i.e. ridge regression). Our first contribution is to demonstrate that this loss can be non-quasiconvex; see the middle and right panel of Fig. 1 for some real-data examples, which we describe in detail in Section 3. Our second contribution is to characterize which aspects of the covariate matrix and observed responses affect quasiconvexity. We prove that the norm of the responses, the scale of the singular values of the covariate matrix, and the right singular vectors of the covariate matrix all have no effect on the quasiconvexity of $\mathcal{L}$. While this result places substantial constraints on what drives the quasiconvexity of $\mathcal{L}$, we show that the quasiconvexity of $\mathcal{L}$ is unfortunately still a complex function of the remaining quantities. Our third contribution is to prove conditions under which $\mathcal{L}$ is guaranteed to be quasiconvex. In particular, we show that if (1) the covariate matrix has a singular value spectrum sufficiently close to uniform, (2) the least-squares estimator fits the training data sufficiently well, and (3) the left singular vectors of the covariate matrix are sufficiently regular, then $\mathcal{L}$ is guaranteed to be quasiconvex. While the conditions of our theory are deterministic, we show that they have natural probabilistic interpretations; as a corollary to our theory, we demonstrate that many of our conditions are satisfied either empirically or theoretically by well-specified linear regression problems with i.i.d. sub-Gaussian covariates and moderate signal-to-noise ratios. Through empirical studies, we validate the conclusions of our theory and the necessity of our assumptions.

2 Setup and notation

For $n \in \{1, \ldots, N\}$, we observe covariates $x_n \in \mathbb{R}^D$ and responses $y_n \in \mathbb{R}$. In the present work, we consider the low to modest-dimensional case $D < N$. We are interested in learning a linear model between the covariates and responses, $\langle x_n, \theta \rangle \approx y_n$, for some parameter $\theta \in \mathbb{R}^D$. In ridge regression, i.e. ℓ_2-regularized linear regression, we take some $\lambda > 0$ and estimate:

$$\hat{\theta}(\lambda) := \arg\min_{\theta \in \mathbb{R}^D} \sum_{n=1}^{N} (\langle x_n, \theta \rangle - y_n)^2 + \frac{\lambda}{2} \|\theta\|_2^2 . \tag{1}$$

The regularization parameter λ is typically chosen by minimizing the cross-validation (CV) loss. Here we study the leave-one-out CV (LOOCV) loss:

$$\mathcal{L}(\lambda) := \sum_{n=1}^{N} \left(\langle x_n, \hat{\theta}^{\backslash n}(\lambda) \rangle - y_n \right)^2 , \tag{2}$$

where $\hat{\theta}^{\backslash n}(\lambda)$ is the solution to Eq. (1) with the nth datapoint left out. Let the covariate matrix $X \in \mathbb{R}^{N \times D}$ be the matrix with rows x_n, and let the vector $Y \in \mathbb{R}^N$ be the vector with entries y_n. We assume X and Y have undergone standard data pre-processing.

Condition 1. *Y is zero-mean, and X has zero-mean, unit variance columns. Equivalently, where* **1** *$\in \mathbb{R}^N$ is the vector of all ones,* $\mathbf{1}^T Y = 0$ *and* $X^T \mathbf{1} = \mathbf{0} \in \mathbb{R}^D$ *and for all* $d = 1, \ldots, D$, $\sum_{n=1}^{N} x_{nd}^2 = N$.

Preprocessing X and Y to satisfy Condition 1 represents standard best practice for ridge regression. First, using an unregularized bias parameter in Eq. (1) and setting Y to be zero-mean are equivalent; we choose to make Y zero-mean, as it simplifies our analysis below. The conditions on the covariate matrix X are important to ensure the use of ℓ_2-regularization is sensible. In particular, Eq. (1) penalizes all coordinates of θ equally. If e.g. some columns of X are measured in different scales or are centered differently, this uniform penalty will be inappropriate.

3   LOOCV loss is typically not convex and need not be quasiconvex

If the LOOCV loss $\mathcal{L}$ were convex or quasiconvex in λ, then any local minimum of $\mathcal{L}$ would be a global minimum, and we could trust gradient-based optimization methods or grid search methods to return a value near a global minimum. We next see that unfortunately $\mathcal{L}$ is typically not convex and is often not even quasiconvex. First we show that, in essentially all cases of interest, $\mathcal{L}$ is *not* convex.

Proposition 1. *If $\lambda = \infty$ is not a minimum of $\mathcal{L}$, then $\mathcal{L}$ is not a convex function.*

Proof. For the sake of contradiction, assume $\mathcal{L}$ is convex and $\lambda = \infty$ is not a minimum of $\mathcal{L}$. This implies that there is some maximal $\lambda^* < \infty$ such that $\mathcal{L}'(\lambda^*) = 0$. Let $\delta := \mathcal{L}'(\lambda^* + 1)$. By convexity, $\mathcal{L}'' \geq 0$, so we know that $\delta > 0$ and that for $\lambda \geq \lambda^* + 1$, we have $\mathcal{L}'(\lambda) \geq \delta$. Thus for $\lambda \geq \lambda^* + 1$, we have $\mathcal{L}(\lambda) \geq \lambda\delta$. So $\lim_{\lambda \to \infty} \mathcal{L}(\lambda) = \infty$. However, inspection of $\mathcal{L}$ shows $\lim_{\lambda \to \infty} \mathcal{L}(\lambda) = \sum_{n=1}^{N} y_n^2 < \infty$, which is a contradiction. $\qquad\square$

We say that the result covers essentially all cases of interest: if $\mathcal{L}$ continues to decrease as $\lambda \to \infty$, then there is so little signal in the data that the zero model $\theta = \mathbf{0} \in \mathbb{R}^D$ is the optimal predictor according to LOOCV.

Although $\mathcal{L}$ is generally not convex, $\mathcal{L}(\lambda)$ might still be easy to optimize if it satisfies an appropriate generalized notion of convexity. To that end, we recall the definition of quasiconvexity.

Definition 1. *A function $f : \mathbb{R}^p \to \mathbb{R}$ is* quasiconvex *if its level sets are convex.*

In one-dimension (i.e. $p = 1$ in Definition 1), quasiconvexity guarantees that any local optimum is a global optimum, just as convexity does. This property is often the key consideration in practical optimization algorithms. Moreover, it is not unreasonable to hope that the CV loss is quasiconvex: typical illustrations of the CV loss are not convex but are quasiconvex; see e.g. Hastie et al. [2015, Fig. 5.9], Rad and Maleki [2020, Fig. 1], or the left panel of Fig. 1. Illustrations of the out-of-sample loss are also typically quasiconvex; see e.g. Fig. 3.6 of Bishop [2006].

Unfortunately, we next demonstrate that the CV loss derived from real data analysis problems can be non-quasiconvex. Our first dataset contains $N = 2{,}938$ observations of life expectancy, along with $D = 20$ covariates such as country of origin or alcohol use; see Appendix A for a full description. In this case, $\mathcal{L}$ for the full dataset is quasiconvex. But now consider some standard data pre-processing. Practitioners often perform principal component regression (PCR) with the aim of reducing noise in the estimated θ. That is, they take the singular value decomposition of $X = USV$ and produce an $N \times R$ dimensional covariate matrix X' by retaining just the top R singular values of X: $X' = U_{.,:R}S_{:R}$. With this pre-processing, the resulting LOOCV curve $\mathcal{L}$ is non-quasiconvex for many values of R; in the center panel of Fig. 1 we show one example for $R = 15$.

Our second dataset consists of recorded wine quality of $N = 1{,}599$ red wines. The goal is to predict wine quality from $D = 11$ observed covariates relating to the chemical properties of each wine; see Appendix A for a full description. We find that subsets of this dataset often exhibit non-quasiconvex $\mathcal{L}$. In the right panel of Fig. 1, we show $\mathcal{L}$ for a subset of size $N = 50$. We see that this plot contains at least two local optima, with substantially different values of λ and substantially different values of the loss. A gradient-based algorithm initialized sufficiently far to the left would not find the global optimum, and grid search without sufficiently large values would not find the global optimum.

Now we know that $\mathcal{L}$ can be non-quasiconvex for real data. Given the difficulty of optimizing a function with several local minima, we next seek to understand *when* $\mathcal{L}$ is quasiconvex or not.

Figure 2: How does quasiconvexity depend on Y and U, for data with $N = 3$ and $D = 2$? The left, center, and right panels each correspond to a different setting of the singular values S. We divide the unit circle for each of U and Y into 100 equally spaced points. We check if $\mathcal{L}$ is quasiconvex over this 100×100 grid. If $\mathcal{L}$ is quasiconvex at a point, that point is colored dark purple; if $\mathcal{L}$ is not quasiconvex, the point is yellow.

4 What does the quasiconvexity of $\mathcal{L}$ depend on?

We have seen that $\mathcal{L}$ can be quasiconvex or non-quasiconvex, depending on the data at hand. If we could determine the quasiconvexity of $\mathcal{L}$ from the data, we might better understand how to efficiently tune hyperparameters from the CV loss. In what follows, we start by showing that the quasiconvexity of $\mathcal{L}$ is, in fact, independent of many aspects of the data (Proposition 2). We will see, however, that the dependence of quasiconvexity on the remaining aspects (though they are few) is complex.

A linear regression problem has a number of moving parts. The response Y may be an arbitrary vector in $\mathbb{R}^N$, and the covariate matrix X can be written in terms of its singular values and left and right singular vectors. More precisely, let $X = USV^T$ be the singular value decomposition of the covariate matrix X, where $U \in \mathbb{R}^{N \times D}$ is an $N \times D$ matrix with orthonormal columns, S is a diagonal matrix with non-negative diagonal, and $V \in \mathbb{R}^{D \times D}$ is an orthonormal matrix. With this notation in hand, we can identify aspects of the problem that do not contribute to quasiconvexity in the following result, which is proved in Appendix B.

Proposition 2. *The quasiconvexity of $\mathcal{L}$ is independent of*

   1. *the matrix of right singular vectors, V,*

   2. *the norm of the responses, $\|Y\|_2$, and*

   3. *the scaling of the singular values (i.e. changing S into S/c for $c \in \mathbb{R}_{>0}$),*

in the sense that altering any of these quantities does not change whether or not $\mathcal{L}$ is quasiconvex.

Remark 1. *Assume Condition 1 holds. Then by Proposition 2, without loss of generality we may (and do) assume that $V = I_D$ and that Y is a vector on the unit $(N-2)$-sphere.*

Recall X has zero-mean columns by preprocessing the data (Condition 1). By Proposition 2, we assume without loss of generality $V = I_D$. Thus, the columns of X have zero mean when $U^T \mathbf{1} = 0$, where $\mathbf{1} \in \mathbb{R}^N$ is the vector of all ones. Hence the quasiconvexity of $\mathcal{L}$ depends on three quantities: (1) the matrix of left singular vectors, U, an orthonormal matrix with $\mathbf{1} \in \mathbb{R}^N$ in its left null-space, (2) the (normalized) vector Y which sits on the unit $(N-2)$-sphere, and (3) the (normalized) singular values.

Now that we know the quasiconvexity of $\mathcal{L}$ depends on only three quantities, we might hope that quasiconvexity would be a simple function of the three. To investigate this dependence, we consider the case of $N = 3$ and $D = 2$, since this case is particularly easy to visualize. To see why it is easy to visualize, first note that Y is a three-dimensional vector. But due to the constraints $\|Y\|_2 = 1$ and $\mathbf{1}^T Y = 0$, Y can be taken to sit on the unit circle and hence can be parametrized by a scalar. Second, note that the matrix of left singular vectors U is parameterized by two orthonormal vectors, $U_{\cdot 1}$ and $U_{\cdot 2}$, on the unit 2-sphere. As both vectors must be orthogonal to $\mathbf{1} \in \mathbb{R}^3$, we can parameterize $U_{\cdot 1}$ and $U_{\cdot 2}$ by two orthonormal vectors on the unit circle. We parametrize $U_{\cdot 1}$ by a scalar that determines $U_{\cdot 2}$ up to a rotation of $U_{\cdot 2}$ by π. We fix a rotation for $U_{\cdot 2}$ relative to $U_{\cdot 1}$ and singular values S, so that the quasiconvexity of $\mathcal{L}$ is parameterized by two scalars.

Fig. 2 is the resulting visualization. Precisely, to make Fig. 2, we fix an orientation between $U_{\cdot 1}$ and $U_{\cdot 2}$: here, a rotation of $\pi/2$ on the unit circle. We create a uniform grid over the unit circles for $U_{\cdot 1}$ and Y. Fig. 2 visualizes the quasiconvexity of $\mathcal{L}$ as we move over this grid for three different settings of the singular values, corresponding to the left, center, and right panels. We see that, in general, U, Y, and S have a complicated interaction to determine the quasiconvexity of $\mathcal{L}$. The values of Y that produce quasiconvexity depend on U. No setting of Y or U guarantees quasiconvexity.

One trend does seem clear from Fig. 2: as the singular values become more similar (moving from left to right in the panels of Fig. 2), the fraction of Y values and U values that correspond to non-quasiconvexity (yellow regions) shrinks. Based on this behavior, one might conjecture that a sufficiently uniform spectrum of the covariate matrix could guarantee the quasiconvexity of $\mathcal{L}$.

5 Quasiconvexity of $\mathcal{L}$ with a nearly uniform spectrum S

We now build on the conjecture of the previous section to show that we can, in fact, guarantee quasiconvexity in certain cases. In particular, we will show conditions under which a sufficiently uniform spectrum S of the covariate matrix X guarantees that $\mathcal{L}$ is quasiconvex.

One might hope that for large N, eventually any Y or U would yield quasiconvexity. Even when S is exactly uniform, our experiments in Section 6 show that we cannot expect such a statement. Rather, we will devise conditions on Y and U to avoid "extreme" settings for either quantity. With these conditions, our main theorem will show that a sufficiently flat spectrum S does indeed guarantee quasiconvexity of $\mathcal{L}$. When our theorem applies, we can safely terminate an optimization procedure at the first local minimum of $\mathcal{L}$ that we discover.

We first establish notation and then state our assumptions.

Definition 2. *Let $\hat{\theta} := \left(X^T X \right)^{-1} X^T Y$ be the least-squares estimate. Define the least-squares residuals $\hat{E} := Y - X\hat{\theta}$. Let $\hat{\varepsilon}_n$ be the nth entry of $\hat{E}$.*

Note that $\hat{\theta}$ is well-defined since we have assumed that $D < N$ and that X is full-rank. For the tractability of our theory, all of our assumptions and conclusions will be asymptotic in N; in particular, our assumptions will use big-O and little-o statements, which are to be taken with respect to N growing large. Since LOOCV is useful precisely for finite N, we are careful to show in our experiments (Section 6) that these asymptotics take hold for small N.

Our first assumption concerns the magnitude of $\hat{E}$.

Assumption 1. $(1/N) \sum_{n=1}^{N} \hat{\varepsilon}_n^2$ *is $O(1)$ (i.e. it does not grow with N).*

This assumption is fairly lax. For example, suppose our linear model is well-specified. In particular, suppose there exists some $\theta^* \in \mathbb{R}^D$ such that $y_n = \langle x_n, \theta^* \rangle + \varepsilon_n$ where the ε_n are i.i.d. $N(0, \sigma^2)$ for some $\sigma > 0$. Stack the ε_n into a vector $E \in \mathbb{R}^N$. Then $\|\hat{E}\|^2 = \|(I_N - UU^T)E\|^2 < \|E\|^2$. Since $(1/N)\|E\|^2$ is $O(1)$ almost surely, it follows that Assumption 1 holds almost surely in this well-specified linear model. We emphasize that Assumption 1 depends on the residuals of the least squares estimate, not (directly) on the noise in the observations.

Our next assumption governs the size of the least squares estimate $\hat{\theta}$.

Assumption 2. $\|\hat{\theta}\|$ *is $O(1)$ (i.e. it does not grow with N).*

Again, this is a lax assumption. For example, given any statistical model for which $\hat{\theta}$ is a consistent estimator for some quantity, Assumption 2 holds.

Our next assumption constrains the uniformity of the left-singular value matrix U with rows $u_n \in \mathbb{R}^D$.

Assumption 3. *We have $\max_n \|u_n\|^2 := \|u_{\max}\|^2 = O(N^{-p})$ for some $p > 1/2$.*

Assumption 3 is an assumption about the coherence of the U matrix, a quantity of importance in compressed sensing and matrix completion [Candés and Recht, 2009]. In particular, Assumption 3 requires that the coherence of U decay sufficiently fast as a function of N. Suppose we remove the condition that U have zero-mean columns (see Condition 1 and the discussion after Remark 1) and assume a uniform distribution over valid U (i.e., matrices with orthonormal columns); then Assumption 3 is known to hold with high probability [Candés and Recht, 2009, Lemma 2.2].

There do exist matrices U with zero-mean columns that do not satisfy Assumption 3. For instance, take some small N_0 (say $N_0 = 5$) and a valid U' for this N_0. Then, for $N > N_0$, form U by appending $\mathbf{0} \in \mathbb{R}^{(N-N_0) \times D}$ to the bottom of U'. This construction yields an $N \times D$ matrix U with orthonormal and zero-mean columns for which the largest $\|u_n\|^2$ (across $n \in \{1, \ldots, N\}$) is constant as N grows. Still, in our experiments in Section 6 and Appendix C, we confirm that, for a uniform distribution over orthonormal U with zero-mean columns, Assumption 3 holds with high probability.

Our final assumption is a technical assumption relating $\|u_n\|^2$, $\hat{\varepsilon}_n$, and $\hat{\theta}$.

Assumption 4. *The following quantity is positive and $\Theta(1)$ (i.e. is bounded away from zero and does not grow with N):* $\|\hat{\theta}\|^2 - \sum_{n=1}^{N} \|u_n\|^2 \left(\langle u_n, \hat{\theta} \rangle^2 + 2\hat{\varepsilon}_n^2 \right)$

Roughly, this assumptions means that the largest $\|u_n\|^2$ and $\hat{\varepsilon}_n^2$ values do not occur for the same values of n. To see this relation, note that Assumption 3 implies $\|\hat{\theta}\|^2 - \sum_n \|u_n\|^2 \langle u_n, \hat{\theta} \rangle^2 \geq (1 - O(N^{-p}))\|\hat{\theta}\|^2$. If we assume that $\|\hat{\theta}\|^2 = \Theta(1)$ (i.e. Assumption 2 holds and $\hat{\theta}$ does not converge to $\mathbf{0} \in \mathbb{R}^D$), then we find $\|\hat{\theta}\|^2 - \sum_n \|u_n\|^2 \langle u_n, \hat{\theta} \rangle^2 = \Theta(1)$. So, we need only that $\sum_n \|u_n\|^2 \hat{\varepsilon}_n^2 = o(1)$ for Assumption 4 to hold; e.g. we need that the largest values of $\|u_n\|^2$ and the largest values of $\hat{\varepsilon}_n^2$ do not occur for the same values of n.

With our assumptions in hand, we can now state our main theorem. Our theorem relates the uniformity of the singular values of X to the quasiconvexity of $\mathcal{L}$. As we have shown in Proposition 2, the scaling of the singular values does not matter for the quasiconvexity of $\mathcal{L}$. We therefore take the singular values to be nearly uniform around $\mathbf{1} \in \mathbb{R}^D$.

Theorem 1. *Take Assumptions 1 to 4. For N sufficiently large, there is a neighborhood Δ of $\mathbf{1} \in \mathbb{R}^D$ such that if the spectrum $S \in \Delta$, $\mathcal{L}(\lambda)$ is quasiconvex.*

Proof sketch: For one-dimensional functions $\mathcal{L}$, a sufficient condition for quasiconvexity is that for all λ such that $\mathcal{L}'(\lambda) = 0$, we have $\mathcal{L}''(\lambda) > 0$ [Boyd and Vandenberghe, 2009, Chapter 3.4]. We first bound the region in which $\mathcal{L}'$ can be zero and then show $\mathcal{L}''$ is positive over this region. See Appendix D.1 for a full proof. □

In Section 4, we showed it can be difficult to guess when $\mathcal{L}$ is quasiconvex. But Theorem 1 yields one condition that guarantees $\mathcal{L}$ is quasiconvex: when X has a nearly uniform spectrum. A natural question then is: when is the spectrum of X nearly uniform? As it happens, a uniform spectrum occurs quite naturally, for example, when the x_{nd} are i.i.d. sub-Gaussian random variables.

Definition 3 (e.g. [Vershynin, 2018]). *A random variable V is sub-Gaussian if there exists a constant $c > 0$ such that $\mathbb{E}[\exp\left(V^2/c^2\right)] \leq 2$.*

Corollary 1. *Take Assumptions 3 and 4. Assume we have a well-specified linear model for some $\theta^* \in \mathbb{R}^D$; in particular, assume $y_n = \langle x_n, \theta^* \rangle + \varepsilon_n$, where $\varepsilon_n \overset{i.i.d.}{\sim} \mathcal{N}(0, \sigma^2)$. If σ is sufficiently small, $\hat{\theta}$ is consistent for θ^*, and the entries of the covariate matrix x_{nd} are i.i.d. sub-Gaussian random variables, then $\mathcal{L}$ is quasiconvex with probability tending to 1 as $N \to \infty$.*

Proof sketch: Assumptions 1 and 2 hold for a well-specified linear model. If the entries of X are i.i.d. sub-Gaussian random variables, standard concentration inequalities imply that its spectrum is nearly uniform with high probability; hence the result of Theorem 1 applies. See Appendix E for a full proof. □

6 Theorem 1 in practice

In Section 5, we established a number of assumptions that we then required in Theorem 1 to prove that $\mathcal{L}$ is quasiconvex. Moreover, Theorem 1 says that this quasiconvexity holds for some unspecified neighborhood Δ of the uniform spectrum $\mathbf{1} \in \mathbb{R}^D$ and asymptotically in N. In this section, we empirically investigate the necessity of our assumptions. And we check how large the neighborhood Δ in Theorem 1 might be and how quickly quasiconvexity appears as a function of N. To that end, we here focus on the small to modest-N regime of $N < 100$.

Figure 3: (*Upper left*): We generate many datasets and plot the fraction that are not quasiconvex, varying N and the distance of the spectrum from uniformity ($\|S - \mathbf{1}\|_1$). (*Upper right*): We generate two sets (orange, blue) of left-singular vector matrices U. In the blue case, we check that the maximum of $\log \|u_{\max}\|^2$ across all U for a particular N decreases roughly linearly on a log-log plot (i.e. the blue set satisfies Assumption 3). In the orange case, we check that the minimum of $\log \|u_{\max}\|^2$ across all U for a particular N is roughly constant (i.e. the orange set does not satisfy Assumption 3). (*Lower*): For all the U matrices from the upper right plot, we generate many datasets and plot the fraction that are not quasiconvex.

The only software dependency for our experiments is `NumPy` [Harris et al., 2020], which uses the BSD 3-Clause "New" or "Revised" License.

How large is the neighborhood Δ in Theorem 1? We first show that the neighborhood Δ is substantial, even for small to moderate N. We fix $D = 5$. To generate various spectra of X, we set $S_d = e^{\alpha d}/e^{\alpha D}$. For $\alpha \to 0$, we get $S \to \mathbf{1}$; we vary α from zero to one to generate spectra of varying distances from uniformity. For each α, we sample 100 left-singular-value matrices U from the uniform distribution over orthonormal U with column means equal to 0; see Appendix F for how to generate such matrices. We fix a unit-norm $\theta^* \in \mathbb{R}^D$ and for each U, we generate data from a well-specified linear model, $y_n = \langle x_n, \theta^* \rangle + \varepsilon_n$, where the ε_n are drawn i.i.d. from $\mathcal{N}(0, \sigma^2)$ with variance $\sigma^2 = 0.5$. In particular, for each setting of U, we generate 100 vectors Y. For each setting of U and Y, we compute $\mathcal{L}$ and check whether it is quasiconvex. In the top left panel of Fig. 3, we report the fraction of problems (out of the $100 * 100 = 10,000$ datasets for the corresponding α value) with a non-quasiconvex $\mathcal{L}$ versus the distance from uniformity, $\|S - \mathbf{1}\|_1$. We see that, even for $N = 20$, the fraction of non-quasiconvex problems quickly hits zero as $\|S - \mathbf{1}\|_1$ shrinks.

Importance of Assumption 3. We now establish the necessity of Assumption 3 on the decay of $\|u_{\max}\|^2$ with N. To do so, we generate two sets of matrices U as N grows. We generate the first set to satisfy Assumption 3, and we generate the second to violate Assumption 3. In both cases, we will take $D = 5$ and ten settings of N between $N = 10$ and $N = 300$.

To generate the assumption-satisfying matrices U, we proceed as follows. For each N, we draw 500 matrices U from the uniform distribution over orthonormal U matrices with column means equal to 0. For each N, we plot the *maximum* value of $\|u_{\max}\|^2$ across these 500 U matrices in Fig. 3 (top-right) as a blue dot. We fit a line to these values on a log-log plot, and find the slope is -0.74. This confirms that these matrices satisfy Assumption 3.

Figure 4: Checking Assumption 1. (*Left*): For each N and each $\nu = \|\hat{E}\|$, we generate many datasets and check if there is any non-quasiconvex $\mathcal{L}$. We plot the largest ν for which we find only quasiconvex $\mathcal{L}$. The growth is roughly linear, which suggests Assumption 1 cannot be loosened. (*Right*): For each N and each $\nu = \|\hat{E}\|$, we generate many data sets and plot the fraction of $\mathcal{L}$ that are non-quasiconvex. Vertical lines show ν_{max} for each N.

To generate assumption-violating matrices U, we proceed as follows. Recall that the smallest N is 10 and $D = 5$. 100 times, we randomly draw a $U_{\text{small}} \in \mathbb{R}^{8 \times 5}$. We then construct each U by appending $N - 8 \times D$ zeros to U_{small}. For each N, we plot the *minimum* value of $\|u_{\max}\|^2$ across these 100 U matrices in Fig. 3 (top-right) as an orange dot. Since the minimum of $\|u_{\max}\|^2$ is constant with N, Assumption 3 is violated.

Now we check quasiconvexity. To that end, we randomly select a fixed unit-norm vector $\theta^* \in \mathbb{R}^D$. For each N, we generate 100 noise vectors $E \in \mathbb{R}^N$, where the entries E_n are drawn i.i.d. from $\mathcal{N}(0, 0.5)$. For each U and E, we construct $Y = US\theta^* + E$, where $S = \mathbf{1} \in \mathbb{R}^D$. We then compute the fraction of these $(100 * 100 = 100{,}000)$ losses $\mathcal{L}$ that are non-quasiconvex. In the lower panel of Fig. 3, we plot this fraction against N for both for the assumption-satisfying case (blue) and the assumption-violating case (orange) in Assumption 3. When the assumption is satisfied (blue), we see that the conclusion of Theorem 1 holds: beyond a certain N, there are no settings of U or Y that generate a non-quasiconvex $\mathcal{L}$. We see that in practice, the boundary N is small or moderate (below 50). When the assumption is violated (orange dots), we see that the conclusion of Theorem 1 fails to hold: as N grows, there are still settings of U and Y for which quasiconvexity fails to hold. Finally, we call attention to the vertical axis. Even in the assumption-violating case, the fraction of non-quasiconvex losses is small. It follows that, even for our degenerate U's, nearly every combination of noise and U leads to a quasiconvex $\mathcal{L}$. An interesting challenge for future work is to provide a precise characterization of this effect.

Do the $\hat{\varepsilon}_n$ need to be small (Assumption 1)? Finally, we demonstrate the necessity of Assumption 1, which can be restated as requiring that the norm of $\hat{E}$ grows at most linearly in N. But we also find a suggestion that there may be even more permissive assumptions of interest. To this end, we vary N from 10 to 30. For each N, we generate 4,000 settings of U, each uniform over orthonormal U with column means equal to 0. For each U, we generate 250 unit vectors R such that $U^T R = 0$ (each generated uniformly over such R – see Appendix F). Separately, we consider 60 different norms ν for the vector $\hat{E}$ equally spaced between $\nu = 0$ and $\nu = 2$; these are the same across N. We generate a single unit-norm $\theta^* \in \mathbb{R}^D$.

For each setting of U, R, and ν, we consider the regression problem with covariate matrix $U\mathbf{1}$ and responses $Y = U\mathbf{1}\theta^* + \hat{E}$, where $\hat{E} := \nu R$. We record whether $\mathcal{L}$ is quasiconvex or not for this problem. For a particular N and particular error-norm ν, we check whether *any* of the $\mathcal{L}$ (across $4{,}000 * 250 = 1{,}000{,}000$ problems) were non-quasiconvex. Finally, for each N, we find the maximum error-norm $\nu_{\max,N}$ such that for all $\nu < \nu_{\max,N}$ every regression problem is quasiconvex. We plot a dot at $(N, \nu_{\max,N})$ in the left panel of Fig. 4. We see that in fact the boundary of allowable $\hat{E}$ norms does grow about linearly in N. Our next plot lends additional insight into how the boundary $\nu_{\max,N}$ varies with N and is also suggestive of other potential variations on Assumption 1 that might be of interest. In particular, in the right panel of Fig. 4, we consider two particular values of N: $N = 10$ (blue) and $N = 20$ (red). For each setting of ν on the horizontal axis, we compute the fraction of

non-quasiconvex losses $\mathcal{L}$ over all settings of U and R ($4{,}000 * 250 = 1{,}000{,}000$ problems for each ν).

First, we see that, as expected from the left panel of Fig. 4, the boundary $\nu_{\max,N}$ is higher for $N = 20$ than for $N = 10$. Second, we see that at high values of ν, the fraction of non-quasiconvex cases decreases again. We conjecture that this trend is more general (beyond these two particular N). Finally, we note that, as in the bottom panel of Fig. 3, the fraction of non-quasiconvex cases across all ν is low. Again, this small fraction suggests a direction for future work.

7   Discussion

We have shown that the LOOCV loss $\mathcal{L}$ for ridge regression can be non-quasiconvex in real-data problems. Local optima need not be global optima. These multiple local optima may pose a practical problem for common hyperparameter tuning methods like gradient-based optimizers, which may get stuck in a local optimum, and grid search, for which upper and lower bounds need to be set.

We proved that the quasiconvexity of $\mathcal{L}$ is determined by only a few aspects of a linear regression problem. But we also showed that the quasiconvexity of $\mathcal{L}$ is still a complicated function of the remaining quantities, and as of this writing the nature of this function is far from fully understood. Nonetheless, we have provided theory that guarantees at least some useful cases when $\mathcal{L}$ is quasiconvex: when the spectrum of the covariate matrix is sufficiently flat, the least-squares fit $\hat{\theta}$ fits the data reasonably well, and the left singular vectors of the covariate matrix are regular. In our experiments, we have confirmed that these assumptions are necessary to some extent: when they are not satisfied, $\mathcal{L}$ can be non-quasiconvex. Still, our empirical results make it clear there is more to be explored. We describe some of the directions we believe are most interesting for future work below.

Sharper characterization of when $\mathcal{L}$ is quasiconvex. Fig. 2 shows that non-quasiconvexity (yellow) disappears as the spectrum of X becomes uniform; however, it is clear that there is very regular behavior to the pattern of quasiconvexity even when the singular values of X are non-uniform. We are not able to characterize these patterns at this time but believe these patterns pose a fascinating challenge for future work. Relatedly, our experiments (Section 6) show that when our assumptions are violated, quasiconvexity of $\mathcal{L}$ is not guaranteed. However, we have observed that even when $\mathcal{L}$ is not guaranteed to be quasiconvex, many settings of U and Y still give quasiconvexity. In many of our experiments, the fraction of non-quasiconvex losses $\mathcal{L}$ was extremely small.

How many local optima and how bad are they? Without the guarantee of a single, global optimum, it is not clear that we can ever know that we have globally (near-)optimized $\mathcal{L}$. However, notice that our examples in Fig. 1 all have at most two local optima. In simulated experiments, we also typically encountered two local optima in non-quasiconvex losses, although we have not studied this behavior systematically. If $\mathcal{L}$ were guaranteed to have only two or some small number of optima, optimization might again be straightforward, even in the case of non-quasiconvexity; an algorithm could search until it finds the requisite number of optima and then report the one with the smallest value of $\mathcal{L}$. Alternatively, one might hope that all local optima have CV loss (and ideally out-of-sample error) close in value to that of the global optimum. Indeed, this closeness has been recently argued for certain losses in deep learning [Kawaguchi, 2016]. Presumably it is not universally the case that local optima exhibit similar loss since the right panel of Fig. 1 seems to give a counterexample. But it might be widely true, or true under mild conditions. Meanwhile, in the absence of such guarantees, optimization of $\mathcal{L}$ should proceed with caution.

Beyond ridge regression. We have shown – in our opinion – surprising non-quasiconvexity for the LOOCV loss for ridge regression. Do similar results hold for simple models outside ridge regression? The regularization parameter in other ℓ_2 or ℓ_1-regularized generalized linear models is often tuned by minimizing a cross-validation loss. In preliminary experiments, we have found non-quasiconvexity in ℓ_2-regularized logistic regression. To what extent do empirical results like those in Fig. 2 or theoretical results like those in Theorem 1 hold for other models and regularizers?

References

D. M. Allen. The relationship between variable selection and data augmentation and a method for prediction. *Technometrics*, 16(1):125–127, Feb 1974.

James Bergstra and Yoshua Bengio. Random search for hyper-parameter optimization. *Journal of Machine Learning Research*, 13(2), 2012.

C. M. Bishop. *Pattern Recognition and Machine Learning*. Springer, 2006.

S. Boyd and L. Vandenberghe. *Convex Optimization*. Cambridge University Press, 2009.

E. J. Candés and B. Recht. Exact matrix completion via convex optimization. *Foundations of Computational Mathematics*, 9, 2009.

D. Chetverikov, Z. Liao, and V. Chernozhukov. On cross-validated Lasso in high dimensions. *arXiv Preprint*, February 2020.

P. Cortez, A. Cerdeira, F. Almeida, T. Matos, and J. Reis. Modeling wine preferences by data mining from physicochemical properties. https://archive.ics.uci.edu/ml/datasets/wine+quality, 2009a. Accessed November 12, 2020.

P. Cortez, A. Cerdeira, F. Almeida, T. Matos, and J. Reis. Modeling wine preferences by data mining from physicochemical properties. *Decision Support Systems*, 47(4):547–553, 2009b.

C. B. Do, C. Foo, and A. Y. Ng. Efficient multiple hyperparameter learning for log-linear models. In *Neural Information Processing Systems (NIPS)*, 2007.

C. R. Harris, K. Jarrod Millman, S. J. van der Walt, R. Gommers, P. Virtanen, David C. Cournapeau, E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van Kerkwijk, M. Brett, A. Haldane, J. Fernández del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant, K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke, and T. E. Oliphant. Array programming with NumPy. *Nature*, 585(7825):357–362, September 2020. doi: 10.1038/s41586-020-2649-2. URL https://doi.org/10.1038/s41586-020-2649-2.

T. Hastie, R. Tibshirani, and M. Wainwright. *Statistical learning with sparsity: the Lasso and generalizations*. Chapman and Hall / CRC, 2015.

T. Hastie, R. Tibshirani, and J. Friedman. *The Elements of Statistical Learning*. Springer, Jan 2017.

T. Hastie, A. Montanari, S. Rosset, and R. J. Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *arXiv preprint*, Dec 2020.

D. Homrighausen and D. J. McDonald. The lasso, persistence, and cross-validation. In *International Conference in Machine Learning (ICML)*, 2013.

D. Homrighausen and D. J. McDonald. Leave-one-out cross-validation is risk consistent for Lasso. *Machine Learning*, 97(1-2):65–78, October 2014.

Chih-Wei Hsu, Chih-Chung Chang, Chih-Jen Lin, et al. A practical guide to support vector classification, 2003.

K. Kawaguchi. Deep learning without poor local minima. In *Neural Information Processing Systems (NIPS)*, 2016.

J. Lorraine, P. Vicol, and D. Duvenaud. Optimizing millions of hyperparameters by implicit differentiation. In *International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2020.

D. Maclaurin, D. Duvenaud, and R. P. Adams. Gradient-based hyperparameter optimization through reversible learning. In *International Conference on Machine Learning (ICML)*, 2015.

K. Musgrave, S. Belongie, and S. N. Lim. A machine learning reality check. *arXiv Preprint*, March 2020.

F. Pedregosa. Hyperparameter optimization with approximate gradient. In *International Conference in Machine Learning (ICML)*, 2016.

Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, et al. Scikit-learn: Machine learning in python. *The Journal of Machine Learning Research*, 12:2825–2830, 2011.

K. R. Rad and A. Maleki. A scalable estimate of the extra-sample prediction error via approximate leave-one-out. *arXiv Preprint*, January 2020.

K. Rajarshi. Life expectancy (WHO). `https://www.kaggle.com/kumarajarshi/life-expectancy-who`, 2018. Accessed May 02, 2021.

M. Stone. Cross-validatory choice and assessment of statistical predictions. *Journal of the American Statistical Association*, 36(2):111–147, 1974.

R. Vershynin. *High-dimensional probability: an introduction with applications in data science*. Cambridge University Press, August 2018.