LATENT CONVERGENT CROSS MAPPING

ABSTRACT

Discovering causal structures of temporal processes is a major tool of scientific inquiry because it helps us better understand and explain the mechanisms driving a phenomenon of interest, thereby facilitating analysis, reasoning, and synthesis for such systems. However, accurately inferring causal structures within a phenomenon based on observational data only is still an open problem. Indeed, this type of data usually consists in short time series with missing or noisy values for which causal inference is increasingly difficult. In this work, we propose a method to uncover causal relations between chaotic dynamical systems from short, noisy and *sporadic* time series (that is, incomplete observations at infrequent and irregular intervals) where the classical convergent cross mapping (CCM) fails. Our method works by learning a Neural ODE latent process modeling the state-space dynamics of the time series and by checking the existence of a continuous map between the resulting processes. We provide theoretical analysis and show empirically that Latent-CCM can reliably uncover the true causal pattern, unlike traditional methods.

1 INTRODUCTION

Inferring a right causal model of a physical phenomenon is at the heart of scientific inquiry. It is fundamental to how we understand the world around us and to predict the impact of future interventions (Pearl, 2009). Correctly inferring causal pathways helps us reason about a physical system, anticipate its behavior in previously unseen conditions, design changes to achieve some objective, or synthesize new systems with desirable behaviors. As an example, in medicine, causality inference could allow predicting whether a drug will be effective for a specific patient, or in climatology, to assess human activity as a causal factor in climate change. Causal mechanisms are best uncovered by making use of interventions because this framework leads to an intuitive and robust notion of causality. However, there is a significant need to identify causal dependencies when only observational data is available, because such data is more readily available as it is more practical and less costly to collect (*e.g.*, relying on observational studies when interventional clinical trials are not yet available).

However, real-world data arising from less controlled environment than, for instance, clinical trials poses many challenges for analysis. Confounding and selection bias come into play, which bias standard statistical estimators. If no intervention is possible, some causal configurations cannot be identified. Importantly, with real-world data comes the major issue of missing values. In particular, when collecting longitudinal data, the resulting time series are often *sporadic*: sampling is *irregular* in time and incomplete across dimensions leading to varying time intervals between observations of a given variable and typically multiple missing observations at any given time. This problem is ubiquitous in various fields, such as healthcare (De Brouwer et al., 2019), climate science (Thomson, 1990), or astronomy (Cuevas-Tello et al., 2010).

A key problem in causal inference is to assess whether one time series is causing another or is merely correlated with it. From assessing causal pathways for neural activity (Roebroeck et al., 2005) to ecology (Sugihara et al., 2012) or healthcare, it is a necessary step to unravel underlying generating mechanisms. A common way to infer causal direction between two time series is to use Granger causality (Granger, 1969), which defines "predictive causality" in terms of the predictability of one time series from the other. A key requirement of Granger causality is then separability (i.e., that information about causes are not contained in the caused variable itself). This assumption holds in purely stochastic linear systems, but fails in more general cases (such as weakly coupled

nonlinear dynamical systems) (Sugihara et al., 2012). To address this nonseparability issue, Sugihara *et al.* (Sugihara et al., 2012) introduced the Convergent Cross Mapping (CCM) method, which is based on the theory of chaotic dynamical systems, particularly on Takens' theorem. This method has been applied successfully in various fields such as ecology, climatology (Wang et al., 2018), and neuroscience (Schiecke et al., 2015). However, as the method relies on embedding the time series under study with time lags, it is highly sensitive to missing values and usually requires long uninterrupted time series. This method is thus not applicable in settings with repeated short sporadic time series, despite their occurrence in many practical situations.

To address this important limitation, we propose to learn the causal dependencies between time series by checking the existence of convergent cross mappings between latent processes of those time series. Using a joint model across all segments of partially observed time series and forcing the model to learn the inherent dynamic of the data, we show that our method can detect causal relationship on short and sporadic time series, without computing delay embeddings. To learn a continuous time latent representation of the system's state-space, we leverage GRU-ODE-Bayes (De Brouwer et al., 2019), a recently introduced filtering method that extends the Neural ODE model (Chen et al., 2018). Importantly for causal inference, the filtering nature of the model makes sure no future information can leak into the past. We then check the existence of continuous maps between the learnt latent representations and infer the causal direction accordingly.

In a series of increasingly challenging test cases, our method accurately detects the correct causal dependencies with high confidence, even when fed very few observations, and outperforms competing methods such as multi-spatial CCM or CCM with multivariate Gaussian process interpolation.

Figure 1: Schematic of the Latent-CCM rationale. If $X[t]$ causes $Y[t]$, there exists a continuous map (dotted line) from the latent process of Y ($\mathcal{H}_Y$) to the latent process of X ($\mathcal{H}_X$).

2 RELATED WORK

CCM to address failure of Granger causality. Granger causality (Granger, 1969) provided the first significant framework to infer causal dependencies from time series. Relying on predictability between dynamical systems, it was extended to account for different limitations, such as nonlinearity (Chen et al., 2004) or instantaneous relationships (Schiatti et al., 2015). However, the assumption of separability of information between causative and caused variables leads to the failure of the Granger paradigm for a significant number of time series coupling scenarios (Sugihara et al., 2012) (see Appendix D for a revealing worked out example). Convergent Cross Mapping, a technique based on nonlinear state space reconstruction was introduced to tackle this issue (Sugihara et al., 2012). Recently, several works have proposed extensions of CCM, such as the extended CCM, to address issues such as synchrony (Ye et al., 2015) or to improve the discrimination of the confounding case (Benkő et al., 2018). Huang et al. (2020) also proposed to predict directly the driving time series from the driven one with reservoir computing, bypassing the delay embedding step, making it more robust to noise. However, those methods still require long regularly sampled time series.

Causality for short or sporadic time series. Short time series are very common in practice and there has been some work proposing to learn causality from short time series relying on state space reconstruction. Ma et al. (2014) proposed a method for short, fully observed, unique time series. Multi-spatial CCM (Clark et al., 2015), considered the problem of inferring causality from several short fully observed snippets of the same dynamical system by computing delay embeddings compatible with the lengths of the time series and aggregating them. In comparison, on top of addressing irregular sampling, our approach computes more informative state-space representations by sharing a model across all segments. Techniques to infer causal direction from non-fully-observed time series have also been proposed, but all are relying on the Granger causality framework, which limits their applicability to separable dynamical systems. They use direct partial correlations on regularly sampled data (but with missing values) (Elsegai, 2019) or generalizations of similarity measures for irregular time series (Bahadori & Liu, 2012). To the best of our knowledge, this is the first work investigating the identification of causal dependencies from short sporadic time series using state-space reconstruction.

3 METHOD

We consider the problem of inferring a causal dependency between several segments of two multi-variate time series $X[t] \in \mathbb{R}^{d_X}$ and $Y[t] \in \mathbb{R}^{d_Y}$. To account for the more general and most frequent case, we consider those time series are only observed in segments of finite duration. $X[t]$ and $Y[t]$ then consist of collections of N short time series $(X^1[t],...,X^N[t]))$ and $(Y^1[t],...,Y^N[t]))$ respectively. Importantly, each segment of X and Y is observed concomitantly. To proceed with a lighter notation, we'll drop the superscript when referring to a segment of time series.

Each of those time series is *sporadic*, namely only observed at times $\mathbf{t}_X$ and $\mathbf{t}_Y$ respectively, which are typically not regularly spaced and for which not all dimensions are observed each time.

In this work, we define the notion of causality by considering the equations of the dynamical system as a structural causal model. In this framework, X causes Y if $p(Y|do(X)) \neq P(Y)$ where $do(X)$ is an intervention on X (Pearl, 2009).

3.1 CONVERGENT CROSS MAPPING AND TAKENS' THEOREM

CCM aims at discovering the causal direction between dynamical systems by checking if the state-space dynamics of one time series can be recovered from another. Intuitively, if X causes Y, information from X leaks into Y and it should be possible to recover X from Y, but not the other way round. The technique relies on Takens' embedding theorem (Takens, 1981). Let $X[t] \in \mathbb{R}^{d_x}$ be a chaotic dynamical system that has a strange attractor $\mathcal{M}$ with box-counting dimension d_M. The dynamics of this system are specified by a flow on $\mathcal{M}$, $\phi_{(\cdot)}(\cdot) : \mathbb{R} \times \mathcal{M} \to \mathcal{M}$, where $\phi_\tau(X[t]) = X[t+\tau]$. This flow is encoded in the ODE of the system. Takens' theorem then states that a delay embedding Φ with delay τ and embedding dimension k

$$\Phi_{\phi,\alpha}^{k,\tau}(X[t]) = (\alpha(\phi_0(X[t])), \alpha(\phi_{-\tau}(X[t])), \ldots, \alpha(\phi_{-k\tau}(X[t])))$$

is an embedding of the strange attractor $\mathcal{M}$ if $k > 2d_M$ and $\alpha : \mathbb{R}^{d_x} \to \mathbb{R}$ is a twice-differentiable observation function. More specifically, the embedding map Φ is a diffeomorphism between the original strange attractor manifold $\mathcal{M}$ and a shadow attractor manifold $\mathcal{M}'$. Under these assumptions, one can then theoretically reconstruct the original time series from the delay embedding.

The simplest observation function α consists in simply taking one of the dimensions of the dynamical system. In this case, writing $X_i[t]$ as the i-th dimension of $X[t]$, Takens' theorem ensures that there is a diffeomorphism between the original attractor manifold of the full dynamical system and the shadow manifold $\mathcal{M}'$ that would be generated by $X'[t] = (X_i[t], X_i[t-\tau], \ldots, X_i[t-k\tau])$. To see how this theorem can be used to infer the causal direction, let us consider the manifold $\mathcal{M}_Z$ of the joint dynamical system resulting of the concatenation of $X[t]$ and $Y[t]$. We then generate two shadow manifolds $\mathcal{M}'_X$ and $\mathcal{M}'_Y$ from the delay embeddings $X'[t] = (X_i[t], X_i[t-\tau], \ldots, X_i[t-k\tau])$ and $Y'[t] : (Y_j[t], Y_j[t-\tau], \ldots, Y_j[t-k\tau])$. Now, if X unidirectionally causes Y (*i.e.*, Y does not cause X), it means that X is part of an autonomous dynamical system and that Y is part of a larger one, containing X. The attractor of Y is then the same as the one of the joint dynamical system

Z. By contrast, the attractor of X is only a subset of it. From Taken's theorem, it is theoretically possible to recover the original $\mathcal{M}_Z$ from $\mathcal{M}'_Y$ and hence, by extension, recover $\mathcal{M}'_X$ from $\mathcal{M}'_Y$. However, the contrary is not true and it is in general not possible to recover $\mathcal{M}'_Y$ from $\mathcal{M}'_X$.

The CCM algorithm uses this property to infer causal dependency. It embeds both dynamical systems X and Y and use k-nearest neighbors to predict points on $\mathcal{M}'_X$ from $\mathcal{M}'_Y$ and inversely. The result then consists in the correlation of the predictions with the true values. We write $\mathbb{C}\mathrm{orr}(X, Y)$ the Pearson correlation for the task of reconstructing $\mathcal{M}'_X$ from $\mathcal{M}'_Y$. To infer that there is a causal link between the predictor dynamical system and the predicted one, this correlation should be high and, importantly, increase with the length of the observed time series, as the observed manifolds become denser.

The potential results are then interpreted in the following way (1) X causes Y if one can reconstruct with high accuracy $\mathcal{M}'_X$ from $\mathcal{M}'_Y$; (2) X and Y are not causally related (but not necessarily statistically independent) if nor $\mathcal{M}'_X$ nor $\mathcal{M}'_Y$ can be reconstructed from the other; (3) X and Y are in a circular causal relation if both $\mathcal{M}'_Y$ and $\mathcal{M}'_X$ can be reconstructed from the other. In the extreme case of strong coupling, the two systems are said to be in synchrony, and it becomes hard to distinguish between unidirectional or bidirectional coupling (Ye et al., 2015).

3.2 NEURAL ODEs

Many continuous-time deterministic dynamical systems are usefully described as ODEs. But in general, not all dimensions of the dynamical system will be observed so that the system is better described as an ODE on a continuous latent process $H(t)$, conditioned on which the observations $X[t]$ are generated. For instance, when observing only one dimension of a 2 dimensional dynamical system, we cannot find a flow $\phi_t(X)$ on that single dimension variable, but we can find one on the latent process $H(t)$. We then have the following description of the dynamics:

$$X[t] = g(H[t]) \ \text{ with } \ \frac{dH(t)}{dt} = f_\theta(H(t), t) \tag{1}$$

where θ represents the parameters of the ODE, $f_\theta(\cdot)$ is a uniformly Lipschitz continuous function and $g(\cdot)$ is a continuous function. Learning the dynamics of the system then consists in learning those parameters θ from a finite set of (potentially noisy) observations of the process X. Neural ODEs (Chen et al., 2018) parametrize this function by a neural network. Learning the weights of this network can be done using the adjoint method or by simply backpropagating through the numerical integrator. Note that one usually allows X[t] to be stochastic (*e.g.* observation noise). In that case, the mean of $X[t]$ (rather than $X[t]$ itself) follows Equation 1.

3.3 CAUSAL INFERENCE WITH LATENT CCM

A key step in the CCM methodology is to compute the delay embedding of both time series: $\Phi(X[t])$ and $\Phi(Y[t])$. However, when the data is only sporadically observed at irregular intervals, the probability of observing the delayed samples $X_i[t], X_i[t-\tau], \ldots, X_i[t-k\tau]$ is vanishing for any t. $X'[t]$ and $Y'[t]$ are then never fully observed (in fact, only one dimension is observed) and nearest neighbor prediction cannot be performed. What is more, short time series usually do not allow to compute a delay embedding of sufficient dimension (k) and lag (τ) (Clark et al., 2015).

Instead of computing delay embeddings, we learn the dynamics of the process with a continuous-time hidden process parametrized by a Neural ODE (as in Eq. 1) and use this hidden representation as a complete representation of the state-space, therefore eliminating the need for delay-embedding that was limiting the applicability of CCM to long, constant sampling time series. A graphical representation of the method is shown on Figure 1.

To infer causality between time series $X[t]$ and $Y[t]$, the first step is to train two GRU-ODE-Bayes models (De Brouwer et al., 2019), a filtering technique that extends Neural-ODEs. Being a filtering approach, GRU-ODE-Bayes ensures no leakage of future information backward in time, an important requirement for our notion of causality. The same model is used for all segments of each time series and is trained to minimize forecasting error. We learn the observation function g, the ODE

f_θ and the continuous-time latent process $H(t)$. We write the resulting space of latent vectors from time series X on all segments as $\mathcal{H}_X$.

Causality is then inferred by checking the existence of a continuous map between $\mathcal{H}_X$ and $\mathcal{H}_Y$. Analogously to CCM, we consider X causes Y if there exists a continuous map between $\mathcal{H}_Y$ and $\mathcal{H}_X$. This is consistent because $\mathcal{H}_X$, just as the delay embedding Φ_X, is a embedding of the strange attractor of the dynamical system as stated in Lemma 3.1 for which we give the proof in the Appendix E.

Lemma 3.1. *For a partially observed dynamical system $X[t] \in \mathcal{X}$ satisfying the following dynamics,*

$$X[t] = g(H(t)) \text{ with } \frac{dH}{dt} = f_\theta(H(t), t)$$

with $g(\cdot)$ and f_θ continuous functions. If there exists one observation function $\alpha_H \in \mathcal{C}^2 : \mathcal{X} \to \mathbb{R}$ along with a valid couple (k, τ) (in the Takens' embedding theorem sense) such that the map $\Phi^{k,\tau}_{g(\phi_H),\alpha_H}(H(t))$ is injective, the latent process $H(t)$ is an embedding of the strange attractor of the full dynamical system containing X.

The requirement of $\Phi^{k,\tau}_{g(\phi_H),\alpha_H}$ being injective is not enforced in our architecture. However, with sufficient regularization of the network, it is satisfied in practice as shown by our results in Section 4.5.

The same reasoning as in CCM then applies to the latent process and causal direction can be inferred. The existence of a continuous map between the latent spaces of both time series is quantitatively assessed with the correlation between the true latents of the driven time series and the reconstructions obtained with a k-nearest neighbors model on the latents of the driven time series. For instance, for a direction $X \to Y$, we report the correlation between predictions of $\mathcal{H}_X$ obtained from $\mathcal{H}_Y$ and the actual ones ($\mathcal{H}_X$). A strong positive correlation suggests an accurate reconstruction and thus a causal link in the studied direction between the variables (*e.g.*, $X \to Y$). By contrast, a weak correlation suggests no causal link in that direction.

4 EXPERIMENTS

We evaluate the performance of our approach on data sets from physical and neurophysiology models, namely a double pendulum and neurons activity data. We show that our method detects the right causal topology in all cases, outperforming multi-spatial CCM, as well as baselines designed to address the sporadicity of the time series. The code is available in the supplementary material.

4.1 BASELINE METHODS

To the best of our knowledge, this is the first time CCM is applied to short sporadic time series. Indeed, because of missing variables, many standard approaches are simply not applicable. The main baseline consists in multi-spatial CCM (Clark et al., 2015) applied to regularly sampled data with a sampling rate similar to the one of the sporadic data. We also compare our approach to variants where multi-spatial CCM is applied to an interpolation of the sporadic time series using (1) linear interpolation and (2) univariate and multivariate Gaussian Processes (GP and MVGP). For the Gaussian Process, we chose a mixture of RBF and identity kernel and learn the parameters from the data. To model multivariate GPs, we used the combination of a Matern and a periodic Matern kernel for the time dimension and used co-regionalization (Bonilla et al., 2008) with a full-rank interaction matrix. We then use the mean of the posterior process as the reconstruction subsequentially fed to the classical CCM method. Implementation was done with GPflow (Matthews et al., 2017).

4.2 PERFORMANCE METRICS

Our method assesses causality by detecting convergent reconstruction accuracy between the latent processes of different time series. To account for both aspects in a single score, we use the difference between the correlation of the reconstruction and the target latent vector using the whole

((a)) 3 Double Pendulums   ((b)) Neural Activity Data

Figure 2: Pearson correlation of the reconstructions in function of the proportion of data used for the Latent-CCM. Solid line represent the mean over the 5 folds and the shaded area the range of values. For both experiments, only the true causal directions show a convergent reconstruction. $\mathbb{C}\mathrm{orr}_{\mathrm{full}}$ is the score when the whole data is used. $\mathbb{C}\mathrm{orr}_0$ is the score when only 100 samples are used.

data ($\mathbb{C}\mathrm{orr}_{full}$) and the correlation using only 100 sample points ($\mathbb{C}\mathrm{orr}_0$), as shown on Figure 2 and suggested in Clark et al. (2015). The score for the causal coupling from $X[t]$ to $Y[t]$ is then defined as

$$\mathcal{S}c_{X \rightarrow Y} = \mathbb{C}\mathrm{orr}_{\mathrm{full}}(X, Y) - \mathbb{C}\mathrm{orr}_0(X, Y).$$

with a higher score implying more confidence in a causal relationship. Additionally, to quantify the certainty about the presence of a causal edge in the data generation graph, we compare the obtained scores with the ones that would be obtained with CCM on fully observed but independent time series. We compute the Mann-Whitney U-statistics (Ma et al., 2014) and provide the corresponding p-value.

Nevertheless, in practice, one might not have access to the score of independent time series, making it difficult to assess from the score only if a causal relationship is present. To address this issue, we visualize the results graphically as shown in Figure 2. Causal directions should then stand out clearly and have the characteristic convergent pattern (Sugihara et al., 2012).

4.3 DOUBLE PENDULUM

4.3.1 DESCRIPTION

The double pendulum is a simple physical system that is chaotic and exhibits rich dynamical behavior. It consists of two point masses m_1 and m_2 connected to a pivot point and to each other by weightless rods of length l_1 and l_2, as shown on Figure 4 in Appendix A. The trajectories of the double pendulum are described by the time series of the angles of the rods with respect to the vertical (θ_1 and θ_2), as well as the angular momenta p_1 and p_2 conjugate to these angles. Each trajectory is then a collection of 4-dimensional vector observations.

To introduce causal dependencies from pendulum X to Y, we include a non-physical asymmetrical coupling term in the update of the momentum conjugate to the first angle:

$$\dot{p}_1^Y = -\frac{\partial H^Y}{\partial \theta_1^Y} - 2 \cdot c_{X,Y}(\theta_1^Y - \theta_1^X),$$

where $c_{X,Y}$ is a coupling parameter. The term corresponding to a quadratic potential incorporated to the Hamiltonian of system Y results in an attraction on system Y by system X. Depending on

the values of $c_{X,Y}$ and $c_{Y,X}$, we have different causal relationships between X and Y. Namely, (1) X causes Y iff $c_{X,Y} \neq 0$, (2) Y causes X iff $c_{Y,X} \neq 0$ and (3) X is not causally related to Y if $c_{X,Y} = c_{Y,X} = 0$.

4.3.2 DATA GENERATION

We consider two cases of generating models. The first one consists of two double pendulums ($X[t]$ and $Y[t]$) with high observation noise with Y causing X. In this case, we set $c_{X,Y} = 0$ and $c_{Y,X} = 0.2$. The second consists of 3 double pendulums ($X[t]$, $Y[t]$ and $Z[t]$), with one of them causing the other two ($c_{Z,X} = 0.5$, $c_{Z,Y} = 1$). Remarkably, X and Y are here correlated but not causally related. Graphical representations of both considered cases are presented in Figure 3. Parameters of the pendulums (lengths and masses) are presented in Appendix A. We generate 5 trajectories with different initial conditions ($\theta_1 \sim \mathcal{N}(-1, 0.05)$ and $\theta_2 \sim \mathcal{N}(0.5, 0.05)$). We simulate observation noise by adding a random Gaussian noise n to the samples with $n \sim \mathcal{N}(\mu = 0, \sigma = 0.1)$ for the first case and $n \sim \mathcal{N}(\mu = 0, \sigma = 0.01)$ for the second. To account for the short length of time series usually encountered in the real world, we randomly split the trajectories in windows of 10 seconds. To simulate sporadicity, we sample observation uniformly at random with an average rate of 4 samples per second. Furthermore, for each of those samples, we apply an observation mask that keeps each individual dimension with probability 0.3. This whole procedure leads to a sporadic pattern with missing observed dimensions at each sample as shown in Figure 5 of Appendix A. We used 80% of available windows for training and used the remaining 20% for hyperparameter tuning.

Figure 3: Graphical model representation of both cases considered in the main body of the paper. Left: Case 1. Right: Case 2 (confounding).

4.4 NEURAL ACTIVITY DATA

We also evaluate our approach on neural activity data. We generate time series of the average membrane potential of two populations of leaky integrate-and-fire neurons with alpha-function shaped synaptic currents (*iaf_psc_alpha*) simulated by NEST-2.20.0 (Fardet et al., 2020). Each neuron population contains 100 units with sparse random excitatory synapses within the population. We consider two cases, one where population A unidirectionally excites population B and another case where both populations fire independently. To account for the short and sporadic nature of real-world data, we generate 5,000 windows of 20 seconds from which we sample 1 observation every second on average. This leads to 20 samples being available on average per time window.

4.5 RESULTS

Results over 5 folds for the double pendulums and the neural activity data are presented in Table 1 and in Figure 2.

Double Pendulum. Our approach is the only one to recover the right causal direction from the sporadic data. The other baselines do not detect any significant correlation and thus no causal link between double pendulums. Despite having access to constant sampling data, multi-spatial CCM is also not able to detect the right data structure. We argue this is caused by the short length of time series window, and thus the low number and quality of delay embeddings that can be computed. In contrast, as our method shares the same model across all time windows, it represents more reliably the (hidden) state-space at any point in time. Importantly, the perfect reconstruction for Case 2 shows that we can distinguish confounding from correlation between time series. Figure 2(a) graphically presents the results of our method for the second case, where it is obvious that the only two

convergent mappings are the ones corresponding to the true directions (solid blue and green lines), providing a strong signal for the right underlying causal mechanism.

Neural activity For the neuron activity data, we observe that our method delivers the largest effect size towards the true data generating model ($\mathcal{S}_c = 0.295$). Baselines methods relying on imputation do not provide any clear signal for a causal coupling (score 10 times lower). Multi-spatial CCM with the regularly sampled original data provides similar signal than our approach but dampened. Interestingly, we observe a small but significant correlation in the wrong direction ($X \leftarrow Y$) suggesting a small coupling in this direction. An inspection of Figure 2(b), however, will convince the reader that the main causal effect is indeed from X to Y. This small correlation in the direction $X \leftarrow Y$ is also observed in the fully observed data as shown in Figure 6 in Appendix B.

Table 1: Average reconstruction scores $\mathcal{S}_c$ (and their standard deviations) in all directions for the double pendulum and neural activity experiments. Standard deviations are computed using 5 repetitions. Significant correlations compared to noncoupled dynamical systems are in bold ($p < 0.01$). Significance is computed using the Mann-Whitney rank test. Our approach detects the correct causal structure. ✓and ✗ highlight correct and wrong direction detection respectively.

DATA	MULTI-SPATIAL CCM	LINEAR	MVGP	LATENT-CCM
CASE 1				
$X \rightarrow Y$	-0.017 ± 0.037	0.001 ± 0.006	0.009 ± 0.014	0.001 ± 0.013
$\mathbf{X \leftarrow Y}$	0.018 ± 0.0056	-0.011 ± 0.01	-0.001 ± 0.021	0.055 ± 0.001
$AUC_{1 \rightarrow 2}$	0.2 (P=0.98)✓	0.49 (P=0.52) ✓	0.66 (P=0.11) ✓	0.55 (P=0.35) ✓
$AUC_{2 \rightarrow 1}$	0.6 (P=0.23)✗	0.17 (P=0.99) ✗	0.44 (P=0.67) ✗	**1 (P<0.001)** ✓
CASE 2				
$X \rightarrow Y$	0.488 ± 0.074	-0.006 ± 0.005	-0.005 ± 0.009	0.001 ± 0.005
$X \leftarrow Y$	0.181 ± 0.119	-0.01 ± 0.01	-0.007 ± 0.012	0.009 ± 0.014
$X \rightarrow Z$	0.054 ± 0.021	-0.002 ± 0.007	-0.002 ± 0.014	0.035 ± 0.019
$\mathbf{Z \rightarrow X}$	0.324 ± 0.197	0.003 ± 0.004	0.012 ± 0.014	0.657 ± 0.105
$Y \rightarrow Z$	-0.071 ± 0.078	-0.005 ± 0.012	-0.003 ± 0.023	0.005 ± 0.011
$\mathbf{Z \rightarrow Y}$	0.101 ± 0.052	-0.002 ± 0.006	-0.003 ± 0.016	0.555 ± 0.109
$AUC_{1 \rightarrow 2}$	**1.00 (P<0.001)**✗	0.27 (P=0.95)✓	0.31 (P=0.91) ✓	0.78 (P=0.02) ✓
$AUC_{2 \rightarrow 1}$	**1.00 (P<0.001)**✗	0.2 (P=0.98)✓	0.31 (P=0.92) ✓	0.67 (P<0.09) ✓
$AUC_{1 \rightarrow 3}$	**0.98 (P<0.001)**✗	0.41 (P=0.74)✓	0.61 (P=0.19) ✓	0.79 (P=0.02) ✓
$AUC_{3 \rightarrow 1}$	**0.93 (P<0.001)**✓	0.25 (P=0.97) ✗	**0.81 (P=0.01)** ✗	**1.00 (P<0.001)** ✓
$AUC_{2 \rightarrow 3}$	0.26 (P=0.97)✓	0.4 (P=0.76)✓	0.45 (P=0.63) ✓	0.46 (P=0.62) ✓
$AUC_{3 \rightarrow 2}$	0.79 (P=0.02)✓	0.2(P=0.98) ✗	0.43 (P=0.69) ✗	**1.00 (P<0.001)** ✓
COUPLED				
$\mathbf{A \rightarrow B}$	0.267 ± 0.001	0.045 ± 0.014	0.028 ± 0.006	0.295 ± 0.012
$A \leftarrow B$	0.055 ± 0.003	0.014 ± 0.009	0.026 ± 0.010	0.033 ± 0.012
$AUC_{A \rightarrow B}$	**1.00 (P=0.006)**✓	**1.00(P=0.006)** ✓	**1.00 (P=0.006)** ✓	**1.00 (P=0.006)** ✓
$AUC_{B \rightarrow A}$	**1.00(P=0.006)** ✗	**1.00(P=0.006)** ✗	**1.00(P=0.006)** ✗	**1.00(P=0.006)** ✗
INDEPENDENT				
$X \rightarrow Y$	-0.012 ± 0.001	0.001 ± 0.003	-0.002 ± 0.008	-0.006 ± 0.007
$X \leftarrow Y$	-0.001 ± 0.001	-0.001 ± 0.003	-0.003 ± 0.005	-0.002 ± 0.008

5 CONCLUSION AND FUTURE WORK

In this work, we propose a novel way to detect causal structure linking chaotic dynamical systems that are sporadically observed using reconstruction of underlying latent processes learnt with Neural-ODE models. We show that our method correctly detects the causal directions between time series in a low and irregular sampling regime, when time series are observed in only short noncontiguous time windows and even in the case of hidden confounders, which are characteristics of real-world data. Despite the apparent limitation of our method to chaotic systems, it has been shown that CCM is broadly applicable in practice as many real dynamical systems are either chaotic or empirically allow Takens'-like embeddings. As our work builds upon CCM theoretically, we expect the range of application to be at least as large and leave the application to other real-world data for future work.

REFERENCES

Mohammad Taha Bahadori and Yan Liu. Granger causality analysis in irregular time series. In *Proceedings of the 2012 SIAM International Conference on Data Mining*, pp. 660–671. SIAM, 2012.

Zsigmond Benkő, Adám Zlatniczki, Dániel Fabó, András Sólyom, Loránd Erőss, András Telcs, and Zoltán Somogyvári. Complete inference of causal relations in dynamical systems. *arXiv preprint arXiv:1808.10806*, 2018.

Edwin V Bonilla, Kian M Chai, and Christopher Williams. Multi-task gaussian process prediction. In *Advances in neural information processing systems*, pp. 153–160, 2008.

Tian Qi Chen, Yulia Rubanova, Jesse Bettencourt, and David K Duvenaud. Neural ordinary differential equations. In *Advances in neural information processing systems*, pp. 6571–6583, 2018.

Yonghong Chen, Govindan Rangarajan, Jianfeng Feng, and Mingzhou Ding. Analyzing multiple nonlinear time series with extended granger causality. *Physics letters A*, 324(1):26–35, 2004.

Adam Thomas Clark, Hao Ye, Forest Isbell, Ethan R Deyle, Jane Cowles, G David Tilman, and George Sugihara. Spatial convergent cross mapping to detect causal relationships from short time series. *Ecology*, 96(5):1174–1181, 2015.

Juan C Cuevas-Tello, Peter Tiňo, Somak Raychaudhury, Xin Yao, and Markus Harva. Uncovering delayed patterns in noisy and irregularly sampled time series: an astronomy application. *Pattern Recognition*, 43(3):1165–1179, 2010.

Edward De Brouwer, Jaak Simm, Adam Arany, and Yves Moreau. Gru-ode-bayes: Continuous modeling of sporadically-observed time series. In *Advances in Neural Information Processing Systems*, pp. 7377–7388, 2019.

Heba Elsegai. Granger-causality inference in the presence of gaps: An equidistant missing-data problem for non-synchronous recorded time series data. *Physica A: Statistical Mechanics and its Applications*, 523:839–851, 2019.

Tanguy Fardet, Stine Brekke Vennemo, Jessica Mitchell, Håkon Mørk, Steffen Graber, Jan Hahne, Sebastian Spreizer, Rajalekshmi Deepu, Guido Trensch, Philipp Weidel, Jakob Jordan, Jochen Martin Eppler, Dennis Terhorst, Abigail Morrison, Charl Linssen, Alberto Antonietti, Kael Dai, Alexey Serenko, Binghuang Cai, Piotr Kubaj, Robin Gutzen, Hanjia Jiang, Itaru Kitayama, Björn Jürgens, and Hans Ekkehard Plesser. Nest 2.20.0, January 2020. URL https://doi.org/10.5281/zenodo.3605514.

Clive WJ Granger. Investigating causal relations by econometric models and cross-spectral methods. *Econometrica: journal of the Econometric Society*, pp. 424–438, 1969.

Yu Huang, Zuntao Fu, and Christian LE Franzke. Detecting causality from time series in a machine learning framework. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, 30(6):063116, 2020.

Huanfei Ma, Kazuyuki Aihara, and Luonan Chen. Detecting causality from nonlinear dynamics with short-term time series. *Scientific reports*, 4(1):1–10, 2014.

Alexander G. de G. Matthews, Mark van der Wilk, Tom Nickson, Keisuke. Fujii, Alexis Boukouvalas, Pablo León-Villagrá, Zoubin Ghahramani, and James Hensman. GPflow: A Gaussian process library using TensorFlow. *Journal of Machine Learning Research*, 18(40):1–6, apr 2017. URL http://jmlr.org/papers/v18/16-537.html.

Judea Pearl. *Causality*. Cambridge university press, 2009.

Alard Roebroeck, Elia Formisano, and Rainer Goebel. Mapping directed influence over the brain using granger causality and fmri. *Neuroimage*, 25(1):230–242, 2005.

L Schiatti, Giandomenico Nollo, G Rossato, and Luca Faes. Extended granger causality: a new tool to identify the structure of physiological networks. *Physiological measurement*, 36(4):827, 2015.

Karin Schiecke, Britta Pester, Martha Feucht, Lutz Leistritz, and Herbert Witte. Convergent cross mapping: Basic concept, influence of estimation parameters and practical application. In *2015 37th Annual International Conference of the IEEE Engineering in Medicine and Biology Society (EMBC)*, pp. 7418–7421. IEEE, 2015.

George Sugihara, Robert May, Hao Ye, Chih-hao Hsieh, Ethan Deyle, Michael Fogarty, and Stephan Munch. Detecting causality in complex ecosystems. *science*, 338(6106):496–500, 2012.

Floris Takens. Detecting strange attractors in turbulence. In *Dynamical systems and turbulence, Warwick 1980*, pp. 366–381. Springer, 1981.

David J Thomson. Time series analysis of holocene climate data. *Philosophical Transactions of the Royal Society of London. Series A, Mathematical and Physical Sciences*, 330(1615):601–616, 1990.

Yunqian Wang, Jing Yang, Yaning Chen, Philippe De Maeyer, Zhi Li, and Weili Duan. Detecting the causal effect of soil moisture on precipitation using convergent cross mapping. *Scientific reports*, 8(1):1–8, 2018.

Hao Ye, Ethan R Deyle, Luis J Gilarranz, and George Sugihara. Distinguishing time-delayed causal interactions using convergent cross mapping. *Scientific reports*, 5:14750, 2015.

A DOUBLE PENDULUM

Figure 4 presents a graphical representation of a double pendulum with its two masses and two weightless rods. Figure 5 shows examples of trajectories generated by a double pendulum.

Figure 4: Physical representation of the double pendulum[1]

The double pendulum is a simple physically system that is chaotic and exhibits rich dynamical behavior. The Lagrangian of the double pendulum is

$$\mathcal{L} = \frac{1}{2}(m_1 + m_2)l_1^2\dot{\theta}_1^2 + \frac{1}{2}m_2l_2^2\dot{\theta}_2^2 + m_2l_1l_2\dot{\theta}_1\dot{\theta}_2\cos(\theta_1 - \theta_2). \tag{2}$$

The corresponding Hamiltonian can be derived using Legendre transform $H = \sum_i \dot{\theta}_i p_i - \mathcal{L}$.

The system evolution can be simulated by integrating the Hamilton equations:

---

[1]Source: JabberWok / Wikimedia Commons, CC-by-3.0.

$$\dot{\theta}_i = \frac{\partial H}{\partial p_i}$$

$$\dot{p}_i = -\frac{\partial H}{\partial \theta_i}$$

The Jacobian of the right hand side is

$$
J = \begin{bmatrix}
\frac{\partial^2 H}{\partial \theta_1 \partial p_1} & \frac{\partial^2 H}{\partial^2 p_1} & \frac{\partial^2 H}{\partial p_1 \partial \theta_2} & \frac{\partial^2 H}{\partial p_1 \partial p_2} \\
-\frac{\partial^2 H}{\partial^2 \theta_1} & -\frac{\partial^2 H}{\partial \theta_1 \partial p_1} & -\frac{\partial^2 H}{\partial \theta_1 \partial \theta_2} & -\frac{\partial^2 H}{\partial \theta_1 \partial p_2} \\
\frac{\partial^2 H}{\partial \theta_1 \partial p_2} & \frac{\partial^2 H}{\partial p_1 \partial p_2} & \frac{\partial^2 H}{\partial \theta_2 \partial p_2} & \frac{\partial^2 H}{\partial^2 p_2} \\
-\frac{\partial^2 H}{\partial \theta_1 \partial \theta_2} & \frac{\partial^2 H}{\partial p_1 \partial \theta_2} & -\frac{\partial^2 H}{\partial^2 \theta_2} & -\frac{\partial^2 H}{\partial \theta_2 \partial p_2}
\end{bmatrix}.
$$

Note that the diagonal elements cancelling in pairs, which results in a trace of zero that indicates the volume-preserving property of the Hamiltonian flow according to Liouville's theorem. This property corresponds to information preservation in nondissipating physical systems. Consequently, a noncoupled double pendulum does not have a proper attractor. However, for a given initial condition, and thus given energy, the possible states still form a densely populated volume in state-space. Applying the nonphysical coupling term, the conservation rule do not hold anymore.

Figure 5: Example of trajectories generated by a double pendulum. The solid lines represent the true process and the dots the sampled measurements.

The real part of the eigenvalues of J are called the local Lyapunov exponents.

The direction of the largest expansion evolves as

$$\frac{d\mathbf{q}}{dt} = J\mathbf{q}$$
$$|\mathbf{q}(0)| = 1$$

The largest Lyapunov exponent is given by

$$\lambda_1 = \lim_{t \to \infty} \frac{1}{t} \log |\mathbf{q}(t)|.$$

Note that in stationary processes J is constant, and the differential equation have a closed form solution

$$\mathbf{q}(t) = \mathbf{q}(0)e^{Jt},$$

and the local and global Lyapunov exponents are equal.

The largest Lyapunov exponent can be described intuitively as

$$|\delta(t)| \approx |\delta(0)|e^{\lambda_1 t},$$

where $\delta(t)$ is defined as the difference between two phase-space trajectories, with initial condition infinitesimally close to each other:

$$\mathbf{x}'(t) = \mathbf{x}(t) + \delta(t), t \geq 0$$
$$|\delta(0)| \leq \epsilon.$$

We use numerical integration to compute the largest Lyapunov exponent of the double pendulum, and verify that it is in the chaotic regime.

Table 2: Parameters [m, kg] and the largest Lyapunov exponents of the uncoupled pendulums ($\lambda_1 > 0$ indicates chaotic behavior). We report means and their confidence interval over 10 repetitions with initial angles perturbed with $\sigma = 0.05$ normal distributed noise.

SYSTEM		l_1	l_2	m_1	m_2	θ_1	θ_2	λ_1 AND CI (80%)
$X \leftarrow Y$	X	1	0.5	2.0	1.0	1	-0.5	0.306 (0.149, 0.468)
	Y	0.5	1.0	0.5	4.0	1	-0.5	0.005 (0.001, 0.010)
	WHOLE $X \rightarrow Y$ SYSTEM							0.318 (0.183, 0.422)
$X \leftarrow Z \rightarrow Y$	X	0.5	1.0	2.0	1.0	1.0	-0.5	0.008 (0.006, 0.009)
	Y	0.5	1.0	2.0	1.0	1.0	-0.5	0.008 (0.006, 0.009)
	Z	1.0	1.0	1.0	3.0	1.0	-0.5	0.007 (0.005, 0.008)
	WHOLE $X \leftarrow Z \rightarrow Y$ SYSTEM							0.090 (0.027, 0.510)

B INTERACTING NEURON POPULATIONS

The time series is the average membrane potential of two populations of leaky integrate-and-fire neurons with alpha-function shaped synaptic currents (*iaf_psc_alpha*) simulated by NEST-2.20.0 (Fardet et al., 2020). Each population contains 100 units with sparse random excitatory synapses inside population, and unidirectionally from population A to population B. A Poisson generator with rate of 40kHz was used to excite the network.

Table 3: Neuron populations. Every non-specified model parameter is left at the default value.

Population	tau_m [ms]	I_e [μA]
A	$\mathcal{U}(15.0, 16.0)$	0.0
B	$\mathcal{N}(15.0, 1.0)$	60.0
C (not obs.)	10.0	0.0

In Figure 6, we plot the reconstruction correlations of the coupled neuron populations obtained with the fully observed time series (10 observations per second) and evaluated with standard CCM. We observe a small convergence in the reconstruction in the non-causal direction, indicating potential synchrony. As presented in Table 1, our approach also captures this small reconstruction signal.

Table 4: Synapses. Parameters have been tuned to achieve stable firing without depolarizing the neuron populations.

From	To	connection type	parameter
Poisson	A	fixed outdegree	outdegree = 10
Poisson	C	fixed outdegree	outdegree = 10
A	A	fixed indegree	indegree = 67
B	B	fixed indegree	indegree = 20
C	C	fixed indegree	indegree = 67
C	B	fixed outdegree	outdegree = 60

Figure 6: Result of CCM on fully observed neural activity data. Despite a clear signal of A driving B, we observe some positive correlation for the reconstruction in the noncausal direction.

C RESULTS WITH UNIVARIATE GAUSSIAN PROCESSES

In Table 5, we present the results of our experiments with univariate Gaussian Processes (GP). In this case, we only learn a GP on the dimension of interest to compute the delay embeddings. As we can see, results are slightly worse than when using multivariate Gaussian Processes (MVGP).

D FAILURE OF THE GRANGER CAUSALITY FRAMEWORK

To show how the Granger causality framework fails in the general nonlinear dynamical systems case, we consider the following coupled dynamical system:

$$X[t+1] = X[t](a - bX[t] - cY[t])$$
$$Y[t+1] = Y[t](d - eY[t])$$

Following Granger causality, including values of Y for predicting $X[t+1]$ should increase the prediction accuracy, and thus hint towards a causal effect of Y on X. However, dynamics of $X[t]$ can be rearranged such that all information about $Y[t]$ is contained in $X[t]$ already. Indeed,

Table 5: Average reconstruction scores $\mathcal{S}_c$ (and their standard deviations) in all directions for the double pendulum and neural activity experiments. Standard deviations are computed using 5 repetitions. Significant correlations compared to noncoupled dynamical systems are in bold ($p < 0.01$). Significance is computed using the Mann-Whitney rank test. Our approach detects the correct causal structure. ✓ and ✗ highlight correct and wrong direction detection respectively.

DATA	SPATIAL CCM	GP	MVGP	LATENT-CCM
CASE 1				
$X \to Y$	-0.017 ± 0.037	0.002 ± 0.004	0.009 ± 0.014	0.001 ± 0.013
$\mathbf{X \leftarrow Y}$	0.018 ± 0.0056	-0.001 ± 0.003	-0.001 ± 0.021	0.055 ± 0.001
$AUC_{1 \to 2}$	0.2 (P=0.98)✓	0.6 (P=0.22) ✓	0.66 (P=0.11) ✓	0.55 (P=0.35) ✓
$AUC_{2 \to 1}$	0.6 (P=0.23)✗	0.44 (P=0.67) ✗	0.44 (P=0.67) ✗	**1 (P<0.001)** ✓
CASE 2				
$X \to Y$	0.488 ± 0.074	-0.01 ± 0.008	-0.005 ± 0.009	0.001 ± 0.005
$X \leftarrow Y$	0.181 ± 0.119	-0.000 ± 0.003	-0.007 ± 0.012	0.009 ± 0.014
$X \to Z$	0.054 ± 0.021	-0.003 ± 0.003	-0.002 ± 0.014	0.035 ± 0.019
$\mathbf{Z \to X}$	0.324 ± 0.197	0.061 ± 0.004	0.012 ± 0.014	0.657 ± 0.105
$Y \to Z$	-0.071 ± 0.078	-0.003 ± 0.003	-0.003 ± 0.023	0.005 ± 0.011
$\mathbf{Z \to Y}$	0.101 ± 0.052	0.039 ± 0.008	-0.003 ± 0.016	0.555 ± 0.109
$AUC_{1 \to 2}$	**1.00 (P<0.001)**✗	0.21 (P=0.98) ✓	0.31 (P=0.91) ✓	0.78 (P=0.02) ✓
$AUC_{2 \to 1}$	**1.00 (P<0.001)**✗	0.49 (P=0.53)✓	0.31 (P=0.92) ✓	0.67 (P<0.09) ✓
$AUC_{1 \to 3}$	**0.98 (P<0.001)**✗	0.35 (P=0.87) ✓	0.61 (P=0.19) ✓	0.79 (P=0.02) ✓
$AUC_{3 \to 1}$	**0.93 (P<0.001)**✓	0.74 (P=0.03)✗	**0.81 (P=0.01)** ✓	**1.00 (P<0.001)** ✓
$AUC_{2 \to 3}$	0.26 (P=0.97)✓	0.36 (P=0.85) ✓	0.45 (P=0.63) ✓	0.46 (P=0.62) ✓
$AUC_{3 \to 2}$	0.79 (P=0.02)✓	0.58 (0.26) ✗	0.43 (P=0.69) ✗	**1.00 (P<0.001)** ✓
COUPLED				
$\mathbf{A \to B}$	0.267 ± 0.001	0.028 ± 0.006	0.028 ± 0.006	0.295 ± 0.012
$A \leftarrow B$	0.055 ± 0.003	0.026 ± 0.010	0.026 ± 0.010	0.033 ± 0.012
$AUC_{A \to B}$	**1.00 (P=0.006)**✓	**1.00 (P=0.006)** ✓	**1.00 (P=0.006)** ✓	**1.00 (P=0.006)** ✓
$AUC_{B \to A}$	**1.00(P=0.006)** ✗✗	**1(P=0.006)** ✗	**1.00(P=0.006)** ✗	**1.00(P=0.006)** ✗
INDEPENDENT				
$X \to Y$	-0.012 ± 0.001	-0.002 ± 0.008	-0.002 ± 0.008	-0.006 ± 0.007
$X \leftarrow Y$	-0.001 ± 0.001	-0.003 ± 0.005	-0.003 ± 0.005	-0.002 ± 0.008

$$Y[t] = \frac{-1}{c}(\frac{X[t]}{X[t-1]} - a + b)(d + \frac{e}{c}(\frac{X[t]}{X[t-1]} - a + b).$$

Conditioning on $Y[t]$ would not bring additional information and Granger causality would then fail to uncover the right causal structure.

E PROOF OF LEMMA 3.1

Proof. We first write the map $\Phi^k_{g(\phi_H),\alpha_H}(H(t))$ in its full form:

$$\Phi^k_{g(\phi_H),\alpha_H}(H(t)) : \mathcal{H} \to \mathbb{R}^k \text{ s.t.}$$
$$\Phi^k_{g(\phi_H),\alpha_H}(H(t)) = (\alpha_H(g(\phi_{H,0}(H(t)))), \alpha_H(g(\phi_{H,-\tau}(H(t)))), ..., \alpha_H(g(\phi_{H,-k\tau}(H(t)))))$$
$$= (\alpha_H(g(H(t))), \alpha_H(g(H(t-\tau))), ..., \alpha_H(g((H(t-k\tau)))))$$
$$= (\alpha_H(X[t]), \alpha_H(X[t-\tau]), ..., \alpha_H(X[t-k\tau])),$$

where the last line follows from the definition of the dynamical system. Φ then maps the latent process to the delay embedding of X obtained with observation function α_H.

As the observation function $\alpha_H \in \mathcal{C}^2$, the flow ϕ_H and the function $g(\cdot)$ are all continuous, this implies that the map Φ is also continuous in $H(t)$. It is also surjective as all delay embeddings (or points in the state-space) will have at least one latent value generating this delay embedding. Indeed, if we write $\mathcal{M}'_{\alpha_H}$ as the shadow manifold of the delay embeddings of X with observation function α_H, we have that

$$\forall m \in \mathcal{M}'_{\alpha_H}, \exists h \in \mathcal{H} \text{ s.t. } \Phi^k_{g(\phi_H),\alpha_H}(h) = m.$$

Let us now assume that there exists a specific observation function α^*_H such that Φ is injective. The map $\Phi_{\alpha^*_H}$ is then bijective. Furthermore, as both $\mathcal{H}$ and $\mathcal{M}'_{\alpha^*_H}$ are endowed with a metric, the map $\Phi_{\alpha^*_H}$ is a homeomorphism between $\mathcal{H}$ and $\mathcal{M}'_{\alpha^*_H}$.

We now show that Φ is a homeomorphism for any observation function. From Takens' theorem, any delay embedding with valid observation function α, dimension k, and delay τ is a valid embedding of the strange attractor of the dynamical system. There must then exists a homeomorphic map Ψ between any two valid delay embeddings with different observation functions:

$$\forall \alpha, \beta \in \mathcal{C}^2, \mathbb{R} \to \mathbb{R}, \exists \text{ homeomorphism } \Psi_{\alpha,\beta} : \mathcal{M}'_\alpha \to \mathcal{M}'_\beta \text{ s.t.}$$
$$\forall m_\alpha \in \mathcal{M}'_\alpha, m_\beta \in \mathcal{M}'_\beta, \Psi_{\alpha,\beta}(m_\alpha) = m_\beta.$$

By transitivity, there is now a homeorphism between $\mathcal{H}$ and any valid delay embedding defined with observation function α_H defined as $\Psi_{\alpha^*_H,\alpha_H} \circ \Phi^k_{g(\phi_H),\alpha^*_H}(H(t))$. By Takens' theorem, $\mathcal{H}$ is thus an embedding of the strange attractor of the dynamical system containing $X[t]$.

$\square$