Low-rank Optimal Transport: Approximation, Statistics and Debiasing

Abstract

The matching principles behind optimal transport (OT) play an increasingly important role in machine learning, a trend which can be observed when OT is used to disambiguate datasets in applications (e.g. single-cell genomics) or used to improve more complex methods (e.g. balanced attention in transformers or self-supervised learning). To scale to more challenging problems, there is a growing consensus that OT requires solvers that can operate on millions, not thousands, of points. The low-rank optimal transport (LOT) approach advocated in Scetbon et al. [2021] holds several promises in that regard, and was shown to complement more established entropic regularization approaches, being able to insert itself in more complex pipelines, such as quadratic OT. LOT restricts the search for low-cost couplings to those that have a low-nonnegative rank, yielding linear time algorithms in cases of interest. However, these promises can only be fulfilled if the LOT approach is seen as a legitimate contender to entropic regularization when compared on properties of interest, where the scorecard typically includes theoretical properties (statistical bounds, relation to other methods) or practical aspects (debiasing, hyperparameter tuning, initialization). We target each of these areas in this paper in order to cement the impact of low-rank approaches in computational OT.

1 Introduction

Optimal transport (OT) is used across data-science to put in correspondence different sets of observations. These observations may come directly from datasets, or, in more advanced applications, depict intermediate layered representations of data. OT theory provides a single grammar to describe and solve increasingly complex matching problems (linear, quadratic, regularized, unbalanced, etc...), making it gain a stake in various areas of science such as as single-cell biology Schiebinger et al. [2019], Yang et al. [2020], Demetci et al. [2020], imaging Schmitz et al. [2018], Heitz et al. [2020], Zheng et al. [2020] or neuroscience Janati et al. [2020], Koundal et al. [2020].

Regularized approaches to OT. Solving OT problems at scale poses, however, formidable challenges. The most obvious among them is computational: the Kantorovich [1942] problem on discrete measures of size n is a linear program that requires $O(n^3 \log n)$ operations to be solved. A second and equally important challenge lies in the estimation of OT in high-dimensional settings, since it suffers from the curse-of-dimensionality Fournier and Guillin [2015]. The advent of regularized approaches, such as entropic regularization [Cuturi, 2013], has pushed these boundaries thanks for faster algorithms [Chizat et al., 2020, Clason et al., 2021] and improved statistical aspects [Genevay et al., 2018a]. Despite these clear strengths, regularized OT solvers remain, however, costly as they typically scale quadratically in the number of observations.

Scaling up using low-rank couplings. While it is always intuitively possible to reduce the size of measures (e.g. using k-means) prior to solving an OT between them, a promising line of work

proposes to combine both [Forrow et al., 2019, Scetbon et al., 2021, 2022]. Conceptually, these low-rank approaches solve simultaneously both an optimal clustering/aggregation strategy with the computation of an effective transport. This intuition rests on an explicit factorization of couplings into two sub-couplings. This has several computational benefits, since its computational cost becomes linear in n if the ground cost matrix seeded to the OT problem has itself a low-rank. While these computational improvements, mostly demonstrated empirically, hold several promises, the theoretical properties of these methods are not yet well established. This stands in stark contrast to the Sinkhorn approach, which is comparatively much better understood.

Our Contributions. The goal of this paper is to advance our knowledge, understanding and practical ability to leverage low-rank factorizations in OT. This paper provides five contributions, targeting theoretical and practical properties of LOT: *(i)* We derive the rate of convergence of the low-rank OT to the true OT with respect to the non-nnegative rank parameter. *(ii)* We provide an upper-bound of the statistical error when estimating LOT using the plug-in estimator, and attain a parametric rate $\mathcal{O}(\sqrt{1/n})$ that is independent of the dimension. *(iii)* We introduce a debiased version of LOT: as the Sinkhorn divergence [Feydy et al., 2018], we show that it can interpolate between the maximum mean discrepancy [Gretton et al., 2012] and OT, it is nonnegative and it metrizes the weak convergence. *(iv)* We exhibit links between the bias induced by the low-rank factorization and clustering methods. *(v)* We propose solutions to practical issues such as the tuning the step-length or the choice of the initialization when applying the algorithm proposed in [Scetbon et al., 2021].

Notations. We consider $(\mathcal{X}, d_\mathcal{X})$ and $(\mathcal{Y}, d_\mathcal{Y})$ two nonempty compact Polish spaces and we denote $\mathcal{M}_1^+(\mathcal{X})$ (resp. $\mathcal{M}_1^+(\mathcal{Y})$) the space of positive Radon probability measures on $\mathcal{X}$ (resp. $\mathcal{Y}$). For all $n \geq 1$, we denote Δ_n the probability simplex of size n and Δ_n^* the subset of Δ_n of positive histograms. We write $\mathbf{1}_n \triangleq (1, \ldots, 1)^T \in \mathbb{R}^n$ and we denote similarly $\|\cdot\|_2$ the Euclidean norm and the Euclidean distance induced by this norm depending on the context.

2 Background on Low-rank Optimal Transport

Let $\mu \in \mathcal{M}_1^+(\mathcal{X})$, $\nu \in \mathcal{M}_1^+(\mathcal{Y})$ and $c : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}_+$ a nonnegative and continuous function. The Kantorovitch formulation of optimal transport between μ and ν is defined by

$$\mathrm{OT}_c(\mu, \nu) \triangleq \min_{\pi \in \Pi(\mu, \nu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi(x, y) \tag{1}$$

where the feasible set is the set of distributions over the product space $\mathcal{X} \times \mathcal{Y}$ with marginals μ and ν:

$$\Pi(\mu, \nu) \triangleq \left\{ \pi \in \mathcal{M}_1^+(\mathcal{X} \times \mathcal{Y}) \text{ s.t. } P_{1\#\pi} = \mu, P_{2\#\pi} = \nu \right\} ,$$

with $P_{1\#\pi}$ (resp. $P_{2\#\pi}$), the pushforward probability measure of π using the projection maps $P_1(x, y) = x$ (resp. $P_2(x, y) = y$). When there exists an optimal coupling solution of (1) supported on a graph of a function, we call such function a Monge map. In the discrete setting, one can reformulate the optimal transport problem as a linear program over the space of nonnegative matrices satisfying the marginal constraints. More precisely, let a and b be respectively elements of Δ_n^* and Δ_m^* and let also $\mathbf{X} \triangleq \{x_1, \ldots, x_n\}$ and $\mathbf{Y} \triangleq \{y_1, \ldots, y_m\}$ be respectively two subsets of $\mathcal{X}$ and $\mathcal{Y}$. By denoting $\mu_{a,\mathbf{X}} \triangleq \sum_{i=1}^n a_i \delta_{x_i}$ and $\nu_{b,\mathbf{Y}} \triangleq \sum_{j=1}^m b_j \delta_{y_j}$ the two discrete distributions associated and writing $C \triangleq [c(x_i, y_j)]_{i,j}$, the discrete optimal transport problem can be formulated as

$$\mathrm{OT}_c(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}}) = \min_{P \in \Pi_{a,b}} \langle C, P \rangle \quad \text{where} \quad \Pi_{a,b} \triangleq \{P \in \mathbb{R}_+^{n \times m} \text{ s.t. } P\mathbf{1}_m = a, P^T \mathbf{1}_n = b\}. \tag{2}$$

Scetbon et al. [2021] propose to constrain the discrete optimal transport problem to couplings that have a low-nonnegative rank:

Definition 1. *Given $M \in \mathbb{R}_+^{n \times m}$, the nonnegative rank of M is defined by:* $\mathrm{rk}_+(M) \triangleq \min\{q | M = \sum_{i=1}^q R_i, \forall i, \mathrm{rk}(R_i) = 1, R_i \geq 0\}$.

Note that for any $M \in \mathbb{R}_+^{n \times m}$, we always have that $\mathrm{rk}_+(M) \leq \min(n, m)$. For $r \geq 1$, we consider the set of couplings satisfying marginal constaints with nonnegative-rank of at most r as $\Pi_{a,b}(r) \triangleq \{P \in \Pi_{a,b}, \mathrm{rk}_+(P) \leq r\}$. The discrete Low-rank Optimal Transport (LOT) problem is defined by:

$$\mathrm{LOT}_{r,c}(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}}) \triangleq \min_{P \in \Pi_{a,b}(r)} \langle C, P \rangle . \tag{3}$$

To solve this problem, Scetbon et al. [2021] show that Problem (3) is equivalent to

$$\min_{(Q,R,g)\in\mathcal{C}_1(a,b,r)\cap\mathcal{C}_2(r)}\langle C, Q\operatorname{diag}(1/g)R^T\rangle \tag{4}$$

where $\mathcal{C}_1(a,b,r) \triangleq \left\{(Q,R,g)\in\mathbb{R}_+^{n\times r}\times\mathbb{R}_+^{m\times r}\times(\mathbb{R}_+^*)^r \text{ s.t. } Q\mathbf{1}_r = a, R\mathbf{1}_r = b\right\}$ and $\mathcal{C}_2(r) \triangleq \left\{(Q,R,g)\in\mathbb{R}_+^{n\times r}\times\mathbb{R}_+^{m\times r}\times\mathbb{R}_+^r \text{ s.t. } Q^T\mathbf{1}_n = R^T\mathbf{1}_m = g\right\}$. They propose to solve it using a mirror descent scheme and prove the non-asymptotic stationary convergence of their algorithm. While Scetbon et al. [2021] only focus on the discrete setting, we consider here its extension for arbitrary probability measures. Following [Forrow et al., 2019], we define the set of rank-r couplings satisfying marginal constraints by:

$$\Pi_r(\mu,\nu) \triangleq \left\{\pi\in\Pi(\mu,\nu) : \exists(\mu_i)_{i=1}^r\in\mathcal{M}_1^+(\mathcal{X})^r,\ (\nu_i)_{i=1}^r\in\mathcal{M}_1^+(\mathcal{Y})^r,\ \lambda\in\Delta_r^* \text{ s.t. } \pi = \sum_{i=1}^r \lambda_i\mu_i\otimes\nu_i\right\} .$$

This more general definition of LOT between $\mu\in\mathcal{M}_1^+(\mathcal{X})$ and $\nu\in\mathcal{M}_1^+(\mathcal{Y})$ reads:

$$\operatorname{LOT}_{r,c}(\mu,\nu) \triangleq \inf_{\pi\in\Pi_r(\mu,\nu)}\int_{\mathcal{X}\times\mathcal{Y}}c(x,y)d\pi(x,y) . \tag{5}$$

Note that this definition of $\operatorname{LOT}_{r,c}$ is consistent as it coincides with the one defined in (3) on discrete probability measures. Observe also that $\Pi_r(\mu,\nu)$ is compact for the weak topology and therefore the infimum in (5) is attained. See Appendix A for more details.

3 Approximation Error of LOT to original OT as a function of rank

Our goal in this section is to obtain a control of the error induced by the low-rank constraint when trying to approximate the true OT cost. We provide first a control of the approximation error in the discrete setting. The proof is given in Appendix B.1.

Proposition 1. *Let $n,m\geq 2$, $\mathbf{X}\triangleq\{x_1,\dots,x_n\}\subset\mathcal{X}$, $\mathbf{Y}\triangleq\{y_1,\dots,y_m\}\subset\mathcal{Y}$ and $a\in\Delta_n^*$ and $b\in\Delta_m^*$. Then for $2\leq r\leq\min(n,m)$, we have that*

$$|\operatorname{LOT}_{r,c}(\mu_{a,\mathbf{X}},\nu_{b,\mathbf{Y}}) - \operatorname{OT}_c(\mu_{a,\mathbf{X}},\nu_{b,\mathbf{Y}})| \leq \|C\|_\infty\, ln(\min(n,m)/(r-1))$$

Remark 1. *Note that this result improves the control obtained in [Liu et al., 2021], where they obtain that $|\operatorname{LOT}_{r,c}(\mu_{a,\mathbf{X}},\nu_{b,\mathbf{Y}}) - \operatorname{OT}_c(\mu_{a,\mathbf{X}},\nu_{b,\mathbf{Y}})| \lesssim \|C\|_\infty\sqrt{nm}(\min(n,m) - r)$ as we have for any $z,z'\geq 1$, $|\ln(z) - \ln(z')|\leq|z - z'|$.*

It is in fact possible to obtain another control of the approximation error by partitioning the space where the measures are supported. For that purpose let us introduce the notion of entropy numbers.

Definition 2. *Let $(\mathcal{Z},d)$ a metric space, $\mathcal{W}\subset\mathcal{Z}$ and $k\geq 1$ an integer. Then by denoting $B_\mathcal{Z}(z,\varepsilon)\triangleq\{y\in\mathcal{Z}:\ d(z,y)\leq\varepsilon\}$, we define the k-th (dyadic) entropy number of $\mathcal{W}$ as*

$$\mathcal{N}_k(\mathcal{W},d) \triangleq \inf\{\varepsilon \text{ s.t. } \exists\, z_1,\dots,z_{2^k}\in\mathcal{Z}:\ \mathcal{W}\subset\cup_{i=1}^{2^k}B_\mathcal{Z}(z_i,\varepsilon)\}$$

For example, any compact set $\mathcal{W}$ of $\mathbb{R}^d$ admits finite entropy numbers, and by denoting $R\triangleq\sup_{w\in\mathcal{W}}\|w\|_2$, we have $\mathcal{N}_k(\mathcal{W},\|\cdot\|_2)\leq 4R/2^{k/d}$. We obtain next a control of the approximation error of $\operatorname{LOT}_{r,c}$ to the true OT cost using entropy numbers (see proof in Appendix B.2).

Proposition 2. *Let $\mu\in\mathcal{M}_1^+(\mathcal{X})$, $\nu\in\mathcal{M}_1^+(\mathcal{Y})$ and assume that c is L-Lipschitz w.r.t. x and y. Then for any $r\geq 1$, we have*

$$|\operatorname{LOT}_{r,c}(\mu,\nu) - \operatorname{OT}_c(\mu,\nu)| \leq 2L\max(\mathcal{N}_{\lfloor\log_2(\lfloor\sqrt{r}\rfloor)\rfloor}(\mathcal{X},d_\mathcal{X}),\mathcal{N}_{\lfloor\log_2(\lfloor\sqrt{r}\rfloor)\rfloor}(\mathcal{Y},d_\mathcal{Y}))$$

This results in the following bound for the p-Wasserstein distance for any $p\geq 1$ on $\mathbb{R}^d$.

Corollary 1. *Let $d\geq 1$, $p\geq 1$, $\mathcal{X}$ a compact subspace of $\mathbb{R}^d$ and $\mu,\nu\in\mathcal{M}_1^+(\mathcal{X})$. By denoting $R\triangleq\sup_{x\in\mathcal{X}}\|x\|_2$, we obtain that for any $r\geq 1$,*

$$|\operatorname{LOT}_{r,\|\cdot\|_2^p}(\mu,\nu) - \operatorname{OT}_{\|\cdot\|_2^p}(\mu,\nu)| \leq 4dp\frac{(8R^2)^p}{r^{p/2d}} .$$

106 As per the Proof of Proposition 2 we can provide a tighter control, assuming a Monge map exists.

107 **Corollary 2.** *Under the same assumptions of Proposition 2 and by assuming in addition that there*
108 *exists a Monge map solving* $OT_c(\mu, \nu)$*, we obtain that for any* $r \geq 1$*,*

$$|\text{LOT}_{r,c}(\mu, \nu) - \text{OT}_c(\mu, \nu)| \leq L\mathcal{N}_{\lfloor \log_2(r) \rfloor}(\mathcal{Y}, d_{\mathcal{Y}})$$

109 When $\mathcal{X} = \mathcal{Y}$ are a subspaces of $\mathbb{R}^d$, a sufficient condition for a Monge map to exists is that either μ
110 or ν is absolutely continuous with respect to the Lebesgue measure and that c is of the form $h(x - y)$
111 where $h : \mathcal{X} \to \mathbb{R}_+$ is a strictly convex function [Santambrogio, 2015, Theorem 1.17]. Therefore if
112 μ is absolutely continuous with respect to the Lebesgue measure, we obtain for any $r \geq 1$ and $p > 1$

$$|\text{LOT}_{r,\|\cdot\|_2^p}(\mu, \nu) - \text{OT}_{\|\cdot\|_2^p}(\mu, \nu)| \leq 2dp\frac{(8R^2)^p}{r^{p/d}} .$$

4 Sample Complexity of LOT

114 We now focus on the statistical performance of the plug-in estimator for LOT. In the following
115 we assume that $\mathcal{X} = \mathcal{Y}$ for simplicity. Given $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, we denote the empirical measures
116 associated $\hat{\mu}_n \triangleq \frac{1}{n}\sum_{i=1}^n \delta_{X_i}$ and $\hat{\nu}_n \triangleq \frac{1}{n}\sum_{i=1}^n \delta_{Y_i}$, where $(X_i, Y_i)_{i=1}^n$ are sampled independently
117 from $\mu \otimes \nu$. We consider the plug-in estimator defined as $\text{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n)$, and we aim at quantifying
118 the rate at which it converges towards the true low-rank optimal transport cost $\text{LOT}_{r,c}(\mu, \nu)$. Before
119 doing so, in the next Proposition we show that this estimator is consistent on compact spaces. The
120 proof is given in Appendix B.3.

121 **Proposition 3.** *Let* $r \geq 1$ *and* $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$*, then* $\text{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) \xrightarrow[n \to +\infty]{} \text{LOT}_{r,c}(\mu, \nu)$ *a.s.*

122 Next we aim at obtaining the convergence rates of our plug-in estimator. In the following Proposition,
123 we obtain a non-asymptotic upper-bound of the statistical error. See Appendix B.4 for the proof.

124 **Proposition 4.** *Let* $r \geq 1$ *and* $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$*. Then, there exists a constant* K_r *such that for any*
125 $\delta > 0$ *and* $n \geq 1$*, we have, with a probability of at least* $1 - 2\delta$*, that*

$$\text{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) \leq \text{LOT}_{r,c}(\mu, \nu) + 11\|c\|_\infty\sqrt{\frac{r}{n}} + K_r\|c\|_\infty\left[\sqrt{\frac{\log(40/\delta)}{n}} + \frac{\sqrt{r}\log(40/\delta)}{n}\right]$$

126 This result is, to the best of our knowledge, the first attempt at providing a statistical control of
127 low-rank optimal transport. We provide an upper-bound of the plug-in estimator which converges
128 towards $\text{LOT}_{r,c}$ at a parametric rate and which is independent of the dimension on general compact
129 metric spaces. While we fall short of providing a lower bound that could match that upper bound,
130 and therefore provide a complete statistical complexity result, we believe this result might provide a
131 first explanation on why, in practice, $\text{LOT}_{r,c}$ displays better statistical properties than unregularized
132 OT and its curse of dimensionality [Dudley, 1969]. In addition, that upper bound compares favorably
133 to known results on entropic optimal transport. The rate of entropy regularized OT does not depend
134 on the ambient dimension with respect to n, but carries an exponential dependence in dimension
135 with respect to the regularization parameter ε [Mena and Niles-Weed, 2019]. By contrast, the term
136 associated with the nonnegative rank r in our bound has no direct dependence on dimension.

137 Our next aim is to obtain an explicit rate with respect to r and n. In Proposition 4, we cannot control
138 explicitly K_r in the general setting. Indeed, in our proof, we obtain that $K_r \triangleq 14/\min_i \lambda_i^*$ where
139 $(\lambda_i^*)_{i=1}^r \in \Delta_r^*$ are the weights involved in the decomposition of one optimal solution of the true
140 $\text{LOT}_{r,c}(\mu, \nu)$. Therefore the control of K_r requires additional assumptions on the optimal solutions
141 of $\text{LOT}_{r,c}(\mu, \nu)$. In the following Proposition, we obtain an explicit upper-bound of the plug-in
142 estimator with respect to r and n in the asymptotic regime.

143 **Proposition 5.** *Let* $r \geq 1$*,* $\delta > 0$ *and* $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$*. Then there exists a constant* $N_{r,\delta}$ *such that if*
144 $n \geq N_{r,\delta}$ *then with a probability of at least* $1 - 2\delta$*, we have*

$$\text{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) \leq \text{LOT}_{r,c}(\mu, \nu) + 11\|c\|_\infty\sqrt{\frac{r}{n}} + 77\|c\|_\infty\sqrt{\frac{\log(40/\delta)}{n}} .$$

Note that one cannot recover the result obtained in Proposition 5 from the one obtained in Proposition 4 as we have that $K_r \geq 14r \xrightarrow[r \to +\infty]{} +\infty$. In order to prove the above result, we use an extension of the McDiarmid's inequality when differences are bounded with high probability [Kutin, 2002]. See proof in Appendix B.5 for more details.

5 Debiased Formulation of LOT

We introduce here the debiased formulation of $\mathrm{LOT}_{r,c}$ and show that it is able to distinguish two distributions, metrize the convergence in law and can be used as a new objective in order to learn distributions. We focus next on the debiasing terms involving measures with themselves $\mathrm{LOT}_{r,c}(\mu, \mu)$ in this new divergence, and show that they can be interpreted as defining a new clustering method generalizing k-means for any geometry.

5.1 On the Proprieties of the Debiased Low-rank Optimal Transport

When it comes to learn (or generate) a distribution in ML applications given samples, it is crucial to consider a divergence that is able to distinguish between two distributions and metrize the convergence in law. In general, $\mathrm{LOT}_{r,c}(\mu, \mu) \neq 0$ and the minimum of $\mathrm{LOT}_{r,c}(\nu, \mu)$ with respect to ν will not necessarily recover μ. In order to alleviate this issue we propose a debiased version of $\mathrm{LOT}_{r,c}$ defined for any $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$ as

$$\mathrm{DLOT}_{r,c}(\mu, \nu) \triangleq \mathrm{LOT}_{r,c}(\mu, \nu) - \frac{1}{2}[\mathrm{LOT}_{r,c}(\mu, \mu) + \mathrm{LOT}_{r,c}(\nu, \nu)] .$$

Note that $\mathrm{DLOT}_{r,c}(\nu, \nu) = 0$. In the next Proposition, we show that, as the Sinkhorn divergence [Genevay et al., 2018b, Feydy et al., 2018], $\mathrm{DLOT}_{r,c}$ interpolates between the Maximum Mean Discrepancy (MMD) and OT. See proof in Appendix B.6.

Proposition 6. *Let $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$. Let us assume that c is symmetric, then we have*

$$\mathrm{DLOT}_{1,c}(\mu, \nu) = \frac{1}{2} \int_{\mathcal{X}^2} -c(x,y)d[\mu - \nu] \otimes d[\mu - \nu](x,y) .$$

If in addition we assume the c is Lipschitz w.r.t to x and y, then we have

$$\mathrm{DLOT}_{r,c}(\mu, \nu) \xrightarrow[r \to +\infty]{} \mathrm{OT}_c(\mu, \nu) .$$

Next, we aim at showing some useful properties of the debiased low-rank OT for machine learning applications. For that purpose, let us first recall some definitions.

Definition 3. *We say that the cost $c : \mathcal{X} \times \mathcal{X} \to \mathbb{R}_+$ is a semimetric on $\mathcal{X}$ if for all $x, x' \in \mathcal{X}$, $c(x, x') = c(x', x)$ and $c(x, x') = 0$ if and only if $x = x'$. In addition we say that c has a negative type if $\forall n \geq 2$, $x_1, \ldots, x_n \in \mathcal{X}$ and $\alpha_1, \ldots, \alpha_n \in \mathbb{R}$ such that $\sum_{i=1}^n \alpha_i = 0$, $\sum_{i,j=1}^n \alpha_i \alpha_j c(x_i, x_j) \leq 0$. We say also that c has a strong negative type if for all $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, $\mu \neq \nu \implies \int_{\mathcal{X}^2} c(x, y)d[\mu - \nu] \otimes [\mu - \nu] < 0$.*

Note that if c has a strong negative type, then c has a negative type too. For example, all Euclidean spaces and even separable Hilbert spaces endowed with the metric induced by their inner products have strong negative type. Also, on $\mathbb{R}^d$, the squared Euclidean distance has a negative type [Sejdinovic et al., 2013].

We can now provide stronger geometric guarantees for $\mathrm{DLOT}_{r,c}$. In the next Proposition, we show that for a large class of cost functions, $\mathrm{DLOT}_{r,c}$ is nonnegative, able to distinguish two distributions, and metrizes the convergence in law. The proof is given in Appendix B.8.

Proposition 7. *Let $r \geq 1$, and let us assume that c is a semimetric of negative type. Then for all $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, we have that*

$$\mathrm{DLOT}_r(\mu, \nu) \geq 0 .$$

In addition, if c has strong negative type then we have also that

$$\mathrm{DLOT}_{r,c}(\mu, \nu) = 0 \iff \mu = \nu \ \text{and}$$
$$\mu_n \to \mu \iff \mathrm{DLOT}_{r,c}(\mu_n, \mu) \to 0 .$$

where the convergence of the sequence of probability measures considered is the convergence in law.

Observe that when c has strong negative type, $\nu \to \mathrm{DLOT}_{r,c}(\nu, \mu) \geq 0$ and it admits a unique global minimizer at $\nu = \mu$. Therefore, $\mathrm{DLOT}_{r,c}$ has desirable properties to be used as a loss. It is also worth noting that, in order to obtain the metrization of the convergence in law, we show the following Proposition. See proof in Appendix B.7.

Proposition 8. *Let $r \geq 1$ and $(\mu_n)_{n \geq 0}$ and $(\nu_n)_{n \geq 0}$ two sequences of probability measures such that $\mu_n \to \mu$ and $\nu_n \to \nu$ with respect to the convergence in law. Then we have that*

$$\mathrm{LOT}_{r,c}(\mu_n, \nu_n) \to \mathrm{LOT}_{r,c}(\mu, \nu) \ .$$

5.2 Low-Rank Transport Bias and Clustering

We turn next to the debiasing terms appearing in DLOT and exhibit links between LOT and clustering methods. Indeed, in the discrete setting, the low-rank bias of a probability measure μ defined as $\mathrm{LOT}_{k,c}(\mu, \mu)$ can be seen as a generalized version of the k-means method for any geometry. In the next Proposition we obtain a new formulation of $\mathrm{LOT}_{k,c}(\mu, \mu)$ viewed as a general clustering method on arbitrary metric space. See proof in Appendix B.9.

Proposition 9. *Let $n \geq k \geq 1$, $\mathbf{X} \triangleq \{x_1, \ldots, x_n\} \subset \mathcal{X}$ and $a \in \Delta_n^*$. If c is a semimetric of negative type, then by denoting $C = (c(x_i, x_j))_{i,j}$, we have that*

$$\mathrm{LOT}_{k,c}(\mu_{a,\mathbf{X}}, \mu_{a,\mathbf{X}}) = \min_Q \langle C, Q diag(1/Q^T \mathbf{1}_n) Q^T \rangle \ \ s.t. \ \ Q \in \mathbb{R}_+^{n \times k} \ , \ \ Q\mathbf{1}_k = a \ . \tag{6}$$

Let us now explain in more details the link between (6) and k-means. When $\mathcal{X}$ is a subspace of $\mathbb{R}^d$, c is the squared Euclidean distance and $a = \mathbf{1}_n$, we recover exactly the k-means algorithm.

Corollary 3. *Let $n \geq k \geq 1$ and $\mathbf{X} \triangleq \{x_1, \ldots, x_n\} \subset \mathbb{R}^d$. We have that*

$$\mathrm{LOT}_{k,\|\cdot\|_2^2}(\mu_{\mathbf{1}_n,\mathbf{X}}, \mu_{a,\mathbf{X}}) = 2 \min_{Q, z_1, \ldots, z_k} \sum_{i=1}^n \sum_{q=1}^k Q_{i,q} \|x_i - z_q\|_2^2 \ \ s.t. \ \ Q \in \{0,1\}^{n \times k}, \ \ Q\mathbf{1}_k = \mathbf{1}_n \ .$$

In the general setting, solving $\mathrm{LOT}_{k,c}(\mu_{a,\mathbf{X}}, \mu_{a,\mathbf{X}})$ for a given geometry c, and a prescribed histogram a offers a new clustering method where the assignment of the points to the clusters is determined by the matrix Q^* solution of (6).

6 Computing LOT: Adaptive Stepsizes and Better Initializations

We target in this section practical issues that arises when using [Scetbon et al., 2021, Algo.3] to solve (4). Scetbon et al. [2021] propose to apply a mirror descent scheme with respect to the Kullback-Leibler divergence which boils down to solve at each iteration $k \geq 0$ the following convex problem using the Dykstra's Algorithm [Dykstra, 1983]:

$$(Q_{k+1}, R_{k+1}, g_{k+1}) \triangleq \operatorname*{argmin}_{\boldsymbol{\zeta} \in \mathcal{C}_1(a,b,r) \cap \mathcal{C}_2(r)} \mathrm{KL}(\boldsymbol{\zeta}, \boldsymbol{\xi}_k) \tag{7}$$

where $(Q_0, R_0, g_0) \in \mathcal{C}_1(a,b,r) \cap \mathcal{C}_2(r)$, $\boldsymbol{\xi}_k \triangleq (\xi_k^{(1)}, \xi_k^{(2)}, \xi_k^{(3)})$, $\xi_k^{(1)} \triangleq Q_k \odot \exp(-\gamma_k C R_k \operatorname{diag}(1/g_k))$, $\xi_k^{(2)} \triangleq R_k \odot \exp(-\gamma_k C^T Q_k \operatorname{diag}(1/g_k))$, $\xi_k^{(3)} \triangleq g_k \odot \exp(\gamma_k \omega_k / g_k^2)$ with $[\omega_k]_i \triangleq [Q_k^T C R_k]_{i,i}$ for all $i \in \{1, \ldots, r\}$, $\mathrm{KL}(\mathbf{w}, \mathbf{r}) \triangleq \sum_i w_i \log(w_i / r_i)$ and $(\gamma_k)_{k \geq 0}$ is a sequence of positive step sizes. In the general setting, each iteration of their algorithm requires $\mathcal{O}(nmr)$ operations and when the ground cost matrix C admits a low-rank factorization of the form $C = AB^T$ where $A \in \mathbb{R}^{n \times q}$ and $B \in \mathbb{R}^{m \times q}$ with $q \ll \min(n, m)$, then the total complexity per iteration becomes linear $\mathcal{O}((n + m)rq)$. Note that for the squared Euclidean cost on $\mathbb{R}^d$, we have that $q = d + 2$. In the following we investigate two practical aspects of the algorithm: the choice of the step sizes and the initialization.

Adaptive choice of γ_k. Scetbon et al. [2021] show experimentally that the choice of $(\gamma_k)_{k \geq 0}$ does not impact the solution obtained upon convergence, but rather the speed at which it is attained. Indeed the larger γ_k is, the faster the algorithm will converge. As a result, their algorithm simply relies on a fixed γ schedule. However, the range of admissible γ depends on the problem considered and it may

vary from one problem to another. Indeed, the algorithm might fail to converge as one needs to ensure at each iteration k of the mirror descent scheme that the kernels ξ_k do not admit 0 entries in order to solve (7) using the Dykstra's Algorithm. Such a situation can occur when the terms involved in the exponentials become too large which may depend on the problem considered. Therefore, it may be of particular interest for practitioners to have a generic range of admissible values for γ independently of the considered problem, in order to alleviate parameter tuning issues. We propose to consider instead an adaptive choice of $(\gamma_k)_{k \geq 0}$ along iterations. D'Orazio et al. [2021], Bayandina et al. [2018] have proposed adaptive mirror descent schemes where, at each iteration, the step-size is normalized by the squared dual-norm of the gradient. Applying such a strategy in our case amounts to consider at each iteration

$$\gamma_k = \frac{\gamma}{\| (CR\operatorname{diag}(1/g), C^T Q\operatorname{diag}(1/g), -\mathcal{D}(Q^T RC)/g^2) \|_\infty^2}, \tag{8}$$

where the initial $\gamma > 0$ is fixed. By doing so, we are able to guarantee a lower-bound of the exponential terms involved in the expression of the kernels ξ_k at each iteration and prevent them from having 0 entries. We recommend to set such as global $\gamma \in [1, 10]$, and observe that this range works whatever the problem considered.

On the choice of the initialization. As $\text{LOT}_{r,c}$ (4) is a non-convex optimization problem, the question of choosing an efficient initialization arises in practice. Scetbon et al. [2021] show experimentally that the convergence of the algorithm does not depend on the initalization chosen if no stopping criterion is used. Indeed, their experimental findings support that only well behaved local minimas are attractive. However, in practice one needs to use a stopping criterion in order to terminate the algorithm. We do observe in many instances that using trivial initializers may result in spurious local minima, which trigger the stopping criterion early on and prevent the algorithm to reach a good solution. Based on various experimentations, we propose to consider a novel initialization of the algorithm. Our initialization aims at being close to a well-behaved local minimum by clustering the input measures. When the measures are supported on Euclidean space, we propose to find r centroids $(z_i)_{i=1}^r$ of one of the two input discrete probability measures using k-means and to solve the following convex barycenter problem:

$$\min_{Q,R} \langle C_{X,Z}, Q \rangle + \langle C_{Y,Z}, R \rangle - \varepsilon H(Q) - \varepsilon H(R) \ \text{ s.t. } \ Q\mathbf{1}_n = a, \ R\mathbf{1}_n = b, \ Q^T\mathbf{1}_r = R^T\mathbf{1}_r \tag{9}$$

where $C_{X,Z} = (c(x_i, z_j))_{i,j}$, $C_{Y,Z} = (c(y_i, z_j))_{i,j}$, and $H(P) = -\sum_{i,j} P_{i,j}(\log(P_{i,j} - 1)$. In practice we fix $\varepsilon = 1/10$ and we then initialize $\text{LOT}_{r,c}$ using (Q, R) solution of (9) and $g \triangleq Q^T\mathbf{1}_r (= R^T\mathbf{1}_r)$. Note that (Q, R, g) is an admissible initialization and finding the centroids as well as solving (9) requires $\mathcal{O}((n + m)r)$ algebraic operations. Therefore such initialization does not change the total complexity of the algorithm.

In the general (non-Euclidean) case, we propose to initialize the algorithm by applying our generalized k-means approach defined in (6) on each input measure where we fix the common marginal to be $g = \mathbf{1}_r/r$. More precisely, by denoting $C_{X,X} = (c(x_i, x_j))_{i,j}$ and $C_{Y,Y} = (c(y_i, y_j))_{i,j}$, we initialize the algorithm by solving:

$$\begin{aligned} Q &\in \operatorname*{argmin}_{Q} \langle C_{X,X}, Q\operatorname{diag}(1/Q^T\mathbf{1}_n)Q^T \rangle \ \text{ s.t. } \ Q \in \mathbb{R}_+^{n \times k}, \ Q\mathbf{1}_k = a, \ Q^T\mathbf{1}_n = \mathbf{1}_r/r \\ R &\in \operatorname*{argmin}_{R} \langle C_{Y,Y}, R\operatorname{diag}(1/R^T\mathbf{1}_m)R^T \rangle \ \text{ s.t. } \ R \in \mathbb{R}_+^{m \times k}, \ R\mathbf{1}_k = b, \ R^T\mathbf{1}_n = \mathbf{1}_r/r. \end{aligned} \tag{10}$$

Note that again the (Q, R, g) obtained is an admissible initialization and the complexity of solving (10) is of the same order as solving (4), thus the total complexity of the algorithm remains the same.

7 Experiments

In this section, we illustrate experimentally our theoretical findings and show how our initialization provide practical improvements. For that purpose we consider 3 synthetic problems and one real world dataset to: *(i)* provide illustrations on the statistical rates of $\text{LOT}_{r,c}$, *(ii)* exhibit the gradient flow of the debiased formulation $\text{DLOT}_{r,c}$, *(iii)* use the clustering method induced by $\text{LOT}_{r,c}$, and *(iv)* show the effect of the initialization. All experiments were run on a MacBook Pro 2019 laptop.

Figure 1: In this experiment, we consider a mixture of 10 anisotropic Gaussians supported on $\mathbb{R}^d$ and we plot the value of $\mathrm{DLOT}_{r,c}$ between two independent empirical measures associated to this mixture when varying the number of samples n and the dimension d for multiple ranks r. The ground cost considered is the squared Euclidean distance. Note that $\mathrm{LOT}_r(\mu,\mu) \neq 0$ and therefore we use $\mathrm{DLOT}_{r,c}(\mu,\mu)$ instead to evaluate the rates. Each point has been obtained by repeating 10 times the experiment. We compare the empirical rates obtained with the theoretical one derived in Proposition 4 for $r = 1$. We observe that our theoretical results match the empirical ones and, as expected, the rates do not depend on d.

Figure 2: We compare the gradient flows $(\mu_t)_{t \geq 0}$ (in red) starting from a Gaussian distribution, μ_0, to a moon shape distribution (in blue), ν, in 2D when minimizing either $L(\mu) \triangleq \mathrm{DLOT}_{r,c}(\mu,\nu)$ or $L(\mu) \triangleq \mathrm{LOT}_{r,c}(\mu,\nu)$. The ground cost is the squared Euclidean distance and we fix $r = 100$. We consider 1000 samples from each distribution and and we plot the evolution of the probability measure obtained along the iterations of a gradient descent scheme. We also display in green the vector field in the descent direction. We show that the debiased version allows to recover the target distribution while $\mathrm{LOT}_{r,c}$ is learning a biased version with a low-rank structure.

Statistical rates. We aim at showing the statistical rates of the plug-in estimator of $\mathrm{LOT}_{r,c}$. As $\mathrm{LOT}_{r,c}(\mu,\mu) \neq 0$ and as we do not have access to this value given samples from μ, we consider instead the debiased version of the low-rank optimal transport, $\mathrm{DLOT}_{r,c}$. In figure 1, we show that the empiricial rates match the theoretical bound obtained in Proposition 4. In particular, we show that that these rates does not depend on the dimension of the ground space. Note also that we recover our theoretical dependence with respect to the rank r: the higher the rank, the slower the convergence.

Gradient Flows using DLOT. We illustrate here a practical use of DLOT for ML application. In figure 6, we consider $Y_1, \ldots, Y_n$ independent samples from a moon shape distribution in 2D, and by denoting $\hat{\nu}_n$ the empirical measure associated, we show the iterations obtained by a gradient descent scheme on the following optimization problem:

$$\min_{\mathbf{X} \in \mathbb{R}^{n \times 2}} \mathrm{DLOT}_{r,c}(\mu_{\mathbf{1}_n/n, \mathbf{X}}, \hat{\nu}_n) \ .$$

We initialize the algorithm using $n = 1000$ samples drawn from a Gaussian distribution. We show that the gradient flow of our debiased version is able to recover the target distribution. We also compare it with the gradient flow of the biased version (LOT) and show that it fails to reproduce the target distribution as it is learning a biased one with a low-rank structure.

Application to Clustering. In this experiment we show some applications of the clustering method induced by $\mathrm{LOT}_{r,c}$. In figure 3, we consider 6 datasets with different structure and we aim at recovering the clusters using (6) for some well chosen costs. We compare the clusters obtained when considering either the squared Euclidean cost (which amounts at applying the k-means) and the

Figure 3: In this experiment, we draw 1000 samples from multiple distributions from the python package scikit-learn [Pedregosa et al., 2011] and we apply the method proposed in (6) for two different costs: in the top row we consider the squared Euclidean distance while in the bottom row, we consider the shortest path distance on the graph associated with the ground cost $c(x, y) = 1 - k(x, y)$ where k is a Gaussian kernel. In the two first problem (starting from the left), we fix $r = 2$, in the next three problem we fix $r = 3$ and in the last one we fix $r = 4$. We observe that the flexibility of our method allows to recover the clustering for a well chosen ground cost.

Figure 4: In this experiment, we consider the Newsgroup20 dataset [Pedregosa et al., 2011] constituted of texts and we embed them into distributions in 50D using the same pre-processing steps as in [Cuturi et al., 2022]. We compare different initialization when applying the algorithm of [Scetbon et al., 2021] to compare random texts viewed as distributions for multiple choices of rank r. The ground cost considered in the squared Euclidean distance. We repeat the experiments 50 times by sampling randomly multiple problems of similar size ($\simeq 250$ samples). We normalize the cost matrix by its maximum value in order to have comparable LOT cost. We consider 4 different initialization: the one using k-means algorithm (9), the one using the generalized k-means (10), the rank-2 initialization [Scetbon et al., 2021] and a random initialization where Q, R and g are drawn from Gaussians. We compare both the cost value and the criterion value (Δ_k) along the iterations of the MD scheme. Note that the curves obtained do not start at the same point in time as we start plotting the curves after obtaining the initial point which in some case requires more algebraic operations (e.g. kmeans methods). First we observe that whatever the initialization considered, the algorithm converges toward the same value. In addition, we observe that both k-means and general k-means are able to initialize well the algorithm by avoiding bad local minima at initialization while the two other initialization are close to spurious local minima at initialization.

278 shortest-path distance on the data viewed as a graph. We show that our method is able to recover the
279 clusters on these settings for well chosen costs and therefore the proposed algorithm in Scetbon et al.
280 [2021] can be seen as a new alternative in order to clusterize data.

281 **Effect of the Initialization.** Our goal here is to show the effect of the initialization. In figure 4, we
282 display the evolution of the cost as well as the value of the stopping criterion along the iterations of
283 the MD scheme solving (4) when considering different initialization. The x-axis corresponds to the
284 total number of algebraic operations. This number is computed at each iteration of the outer loop
285 of the algorithm proposed in Scetbon et al. [2021] and is obtained by computing the complexity of
286 all the operations involved in their algorithm to reach it. We consider this notion of time instead of

CPU/GPU time as we do not want to be architecture/machine dependent. Recall also that the stopping criterion introduced in [Scetbon et al., 2021] is defined for all $k \geq 1$ by

$$\Delta_k \triangleq \frac{1}{\gamma_k^2}(\mathrm{KL}((Q_k, R_k, g_k), (Q_{k-1}, R_{k-1}, g_{k-1})) + \mathrm{KL}((Q_{k-1}, R_{k-1}, g_{k-1}), (Q_k, R_k, g_k))),$$

where $((Q_k, R_k, g_k))_{k \geq 0}$ is the sequence solution of (7). First, we show that whatever the initialization chosen, the algorithm manages to converge to an efficient solution if no stopping criterion is used. However, the choice of the initialization may impact the termination of the algorithm as some initialization might be too close to some spurious local minima. Indeed, the initial points obtained using a "rank 2" or random initialization can be close to spurious and non-attractive local minima, which may trigger the stopping criterion too early and prevent the algorithm from continuing to run in order to converge towards an attractive and well behaved local minimum. We show also that the initialization we propose in (9) and (10) are sufficiently far away from bad local minima and allow the algorithm to converge directly toward the desired solution.

The right figure of Fig.4 shows two main observations: (i) that the initial point obtained using a "rank 2" or random initialization can be close to spurious and non-attractive local minima, which may trigger the stopping criterion too early and prevent the algorithm from continuing to run in order to converge towards an attractive and well behaved local minimum. (ii) When initialiazing the algorithm using kmeans methods, we show that our stopping criterion is a decreasing function of time meaning that the algorithm converges directly towards the desired solution.

Conclusion. We assembled in this work theoretical and practical arguments to support low-rank factorizations for OT. We have presented two controls: one concerning the approximation error to the true optimal transport and another concerning the statistical rates of the plug-in estimator. The latter is showed to be independent of the dimension, which is of particular interest when studying OT in ML settings. We have motivated further the use of LOT as a loss by introducing its debiased version and showed that it possesses desirable properties: positivity and metrization of the convergence in law. We have also presented the links between the bias induced by such regularization and clustering methods, and studied empirically the effects of hyperparameters involved in the practical estimation of LOT. The strong theoretical foundations provided in this paper motivate further studies of the empirical behaviour of LOT estimator, notably on finding suitable local minima and on improvements on the convergence of the MD scheme using other adaptive choices for step sizes.

References

Anastasia Bayandina, Pavel Dvurechensky, Alexander Gasnikov, Fedor Stonyakin, and Alexander Titov. Mirror descent and convex optimization problems with non-smooth inequality constraints. In *Large-scale and distributed optimization*, pages 181–213. Springer, 2018.

Lenaic Chizat, Pierre Roussillon, Flavien Léger, François-Xavier Vialard, and Gabriel Peyré. Faster wasserstein distance estimation with the sinkhorn divergence. *Advances in Neural Information Processing Systems*, 33, 2020.

Christian Clason, Dirk A. Lorenz, Hinrich Mahler, and Benedikt Wirth. Entropic regularization of continuous optimal transport problems. *Journal of Mathematical Analysis and Applications*, 494 (1):124432, 2021. ISSN 0022-247X. doi: https://doi.org/10.1016/j.jmaa.2020.124432.

Marco Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. In *Advances in neural information processing systems*, pages 2292–2300, 2013.

Marco Cuturi, Laetitia Meng-Papaxanthos, Yingtao Tian, Charlotte Bunne, Geoff Davis, and Olivier Teboul. Optimal transport tools (ott): A jax toolbox for all things wasserstein. *arXiv preprint arXiv:2201.12324*, 2022.

Pinar Demetci, Rebecca Santorella, Björn Sandstede, William Stafford Noble, and Ritambhara Singh. Gromov-wasserstein optimal transport to align single-cell multi-omics data. *BioRxiv*, 2020.

Ryan D'Orazio, Nicolas Loizou, Issam Laradji, and Ioannis Mitliagkas. Stochastic mirror descent: Convergence analysis and adaptive variants via the mirror stochastic polyak stepsize. *arXiv preprint arXiv:2110.15412*, 2021.

Richard Mansfield Dudley. The speed of mean glivenko-cantelli convergence. *The Annals of Mathematical Statistics*, 40(1):40–50, 1969.

Richard L Dykstra. An algorithm for restricted least squares regression. *Journal of the American Statistical Association*, 78(384):837–842, 1983.

Jean Feydy, Thibault Séjourné, François-Xavier Vialard, Shun-Ichi Amari, Alain Trouvé, and Gabriel Peyré. Interpolating between optimal transport and mmd using sinkhorn divergences. *arXiv preprint arXiv:1810.08278*, 2018.

Aden Forrow, Jan-Christian Hütter, Mor Nitzan, Philippe Rigollet, Geoffrey Schiebinger, and Jonathan Weed. Statistical optimal transport via factored couplings. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 2454–2465. PMLR, 2019.

Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3-4):707–738, 2015.

Aude Genevay, Lénaic Chizat, Francis Bach, Marco Cuturi, and Gabriel Peyré. Sample complexity of sinkhorn divergences. *arXiv preprint arXiv:1810.02733*, 2018a.

Aude Genevay, Gabriel Peyre, and Marco Cuturi. Learning generative models with sinkhorn divergences. In *Proceedings of the Twenty-First International Conference on Artificial Intelligence and Statistics*, Proceedings of Machine Learning Research. PMLR, 09–11 Apr 2018b.

Arthur Gretton, Karsten M Borgwardt, Malte J Rasch, Bernhard Schölkopf, and Alexander Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(Mar):723–773, 2012.

Matthieu Heitz, Nicolas Bonneel, David Coeurjolly, Marco Cuturi, and Gabriel Peyré. Ground metric learning on graphs. *Journal of Mathematical Imaging and Vision*, pages 1–19, 2020.

Hicham Janati, Thomas Bazeille, Bertrand Thirion, Marco Cuturi, and Alexandre Gramfort. Multi-subject meg/eeg source imaging with sparse multi-task regression. *NeuroImage*, page 116847, 2020.

Leonid Kantorovich. On the transfer of masses (in russian). *Doklady Akademii Nauk*, 37(2):227–229, 1942.

Sunil Koundal, Rena Elkin, Saad Nadeem, Yuechuan Xue, Stefan Constantinou, Simon Sanggaard, Xiaodan Liu, Brittany Monte, Feng Xu, William Van Nostrand, et al. Optimal mass transport with lagrangian workflow reveals advective and diffusion driven solute transport in the glymphatic system. *Scientific reports*, 10(1):1–18, 2020.

Samuel Kutin. Extensions to mcdiarmid's inequality when differences are bounded with high probability. *Dept. Comput. Sci., Univ. Chicago, Chicago, IL, USA, Tech. Rep. TR-2002-04*, 2002.

Weijie Liu, Chao Zhang, Nenggan Zheng, and Hui Qian. Approximating optimal transport via low-rank and sparse factorization. *arXiv preprint arXiv:2111.06546*, 2021.

Gonzalo Mena and Jonathan Niles-Weed. Statistical bounds for entropic optimal transport: sample complexity and the central limit theorem. *Advances in Neural Information Processing Systems*, 32, 2019.

Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, et al. Scikit-learn: Machine learning in python. *the Journal of machine Learning research*, 12:2825–2830, 2011.

Filippo Santambrogio. Optimal transport for applied mathematicians. *Birkäuser, NY*, 55(58-63):94, 2015.

Meyer Scetbon, Marco Cuturi, and Gabriel Peyré. Low-rank sinkhorn factorization, 2021.

Meyer Scetbon, Gabriel Peyré, and Marco Cuturi. Linear-time gromov wasserstein distances using low rank couplings and costs. *ICML*, 2022.

Geoffrey Schiebinger, Jian Shu, Marcin Tabaka, Brian Cleary, Vidya Subramanian, Aryeh Solomon, Joshua Gould, Siyan Liu, Stacie Lin, Peter Berube, et al. Optimal-transport analysis of single-cell gene expression identifies developmental trajectories in reprogramming. *Cell*, 176(4):928–943, 2019.

Morgan A Schmitz, Matthieu Heitz, Nicolas Bonneel, Fred Ngole, David Coeurjolly, Marco Cuturi, Gabriel Peyré, and Jean-Luc Starck. Wasserstein dictionary learning: Optimal transport-based unsupervised nonlinear dictionary learning. *SIAM Journal on Imaging Sciences*, 11(1):643–678, 2018.

Dino Sejdinovic, Bharath Sriperumbudur, Arthur Gretton, and Kenji Fukumizu. Equivalence of distance-based and rkhs-based statistics in hypothesis testing. *The Annals of Statistics*, pages 2263–2291, 2013.

Karren Dai Yang, Karthik Damodaran, Saradha Venkatachalapathy, Ali C Soylemezoglu, GV Shivashankar, and Caroline Uhler. Predicting cell lineages using autoencoders and optimal transport. *PLoS computational biology*, 16(4):e1007828, 2020.

Xinye Zheng, Jianbo Ye, James Z Wang, and Jia Li. Scott: Shape-location combined tracking with optimal transport. *SIAM Journal on Mathematics of Data Science*, 2(2):284–308, 2020.

Supplementary materials

A   On the Definition of $\text{LOT}_{r,c}$

Let $(\mathcal{X}, d_{\mathcal{X})}$ and $(\mathcal{Y}, d_{\mathcal{Y}})$ two nonempty compact Polish spaces, $\mu \in \mathcal{M}_1^+(\mathcal{X})$, $\nu \in \mathcal{M}_1^+(\mathcal{Y})$ two probability measures on these spaces and $c : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}_+$ a nonnegative and continuous function. We define the generalized low-rank optimal transport between μ and ν as

$$\text{LOT}_{r,c}(\mu, \nu) \triangleq \inf_{\pi \in \Pi_r(\mu,\nu)} \int_{\mathcal{X} \times \mathcal{Y}} c(x,y) d\pi(x,y) \ .$$

where

$$\Pi_r(\mu, \nu) \triangleq \{\pi \in \Pi(\mu, \nu) : \exists (\mu_i)_{i=1}^r \in \mathcal{M}_1^+(\mathcal{X})^r, \ (\nu_i)_{i=1}^r \in \mathcal{M}_1^+(\mathcal{Y})^r, \ \lambda \in \Delta_r^* \ \text{s.t.} \ \pi = \sum_{i=1}^r \lambda_i \mu_i \otimes \nu_i \} \ .$$

As $\mathcal{X}$ and $\mathcal{Y}$ are compact, $\Pi_r(\mu, \nu)$ is tight, then Prokhorov's theorem applies and the closure of $\Pi_r(\mu, \nu)$ is sequentially compact. Let us now show that $\Pi_r(\mu, \nu)$ is closed. Indeed, Let $(\pi_n)_{n \geq 0}$ a sequence of $\Pi_r(\mu, \nu)$ converging towards π_*. Then by definition there exists for all $k \in [[1, r]]$, $(\mu_n^{(k)})_{n \geq 0}$, $(\nu_n^{(k)})_{n \geq 0}$ and $(\lambda_n^{(k)})_{n \geq 0}$ such that for all $n \geq 0$

$$\pi_n = \sum_{i=1}^r \lambda_n^{(k)} \mu_n^{(k)} \otimes \nu_n^{(k)} \ .$$

However, $(\mu_n^{(k)})_{n \geq 0}$ and $(\nu_n^{(k)})_{n \geq 0}$ are also tight, and Prokhorov's theorem applies, therefore we can extract a common subsequence such that for all k,

$$\mu_n^{(k)} \to \mu_*^{(k)} \ \text{and} \ \nu_n^{(k)} \to \nu_*^{(k)}$$

In addition as $(\lambda_n)_{n \geq 0}$ live in the simplex Δ_r, we can also extract a sub-sequence, such that $\lambda_n \to \lambda_* \in \Delta_r$. Finally by unicity of the limit we obtain that

$$\pi_* = \sum_{k=1}^r \lambda_*^{(k)} \mu_*^{(k)} \otimes \nu_*^{(k)} \ .$$

Finally, by denoting $I \triangleq \{k : \lambda_*^{(k)} > 0\}$, and by considering $i^* \in I$, we obtain that

$$\pi_* = \sum_{i \in I \setminus \{i^*\}} \lambda_*^{(i)} \mu_*^{(i)} \otimes \nu_*^{(i)} + \sum_{j=1}^{r-|I|+1} \frac{\lambda_*^{(i^*)}}{r-|I|+1} \mu_*^{(i^*)} \otimes \nu_*^{(i^*)} \ .$$

from which follows that $\pi_* \in \Pi_r(\mu, \nu)$.

B   Proofs

B.1   Proof of Proposition 1

Proposition. *Let $n, m \geq 2$, $\mathbf{X} \triangleq \{x_1, \ldots, x_n\} \subset \mathcal{X}$, $\mathbf{Y} \triangleq \{y_1, \ldots, y_m\} \subset \mathcal{Y}$ and $a \in \Delta_n^*$ and $b \in \Delta_m^*$. Then for $2 \leq r \leq \min(n, m)$, we have that*

$$|\text{LOT}_{r,c}(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}}) - \text{OT}_c(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}})| \leq \|C\|_\infty \ ln(\min(n,m)/(r-1))$$

Proof. Let $P \in \text{argmin}_{P \in \Pi_{a,b}} \langle C, P \rangle$. As P is a nonnegative matrix, its nonnegative rank cannot exceed $\min(n, m)$. Assume for simplicity, that $n = m$, then there exists $(R_i)_{i=1}^n$ nonnegative matrices of rank 1 such that

$$P = \sum_{i=1}^n R_i \ .$$

As for all $i \in [|1, n|]$, R_i is a rank 1 matrix, there exist $\tilde{q}_i, \tilde{r}_i \in \mathbb{R}^n_+$ such that $R_i = \tilde{q}_i \tilde{r}_i^T$. Then by denoting $q_i = \tilde{q}_i/|\tilde{q}_i|$, $r_i = \tilde{r}_i/|\tilde{r}_i|$ and $\lambda_i = |\tilde{q}_i||\tilde{r}_i|$ where for any $h \in \mathbb{R}^n$ $|h| \triangleq \sum_{i=1}^n h_i$, we obtain that

$$P = \sum_{i=1}^n \lambda_i q_i r_i^T .$$

454   Without loss of generality, we can consider the case where $\lambda_1 \geq \cdots \geq \lambda_n$. Let us now denote
455   $\lambda := (\lambda_1, \ldots, \lambda_n)$, and by using the fact the P is a coupling we obtain that $\lambda \in \Delta_n$. Also, by
456   definition of λ, we have that for all $k \in [|1, n|]$, $\lambda_k \leq 1/k$. Let us now define

$$\tilde{P} \triangleq \sum_{i=1}^{r-1} \lambda_i q_i r_i^T + \left(\sum_{i=r}^n \lambda_i \right) \alpha_r \beta_r^T$$

457   where

$$\alpha_r \triangleq \frac{\sum_{i=r}^n \lambda_i q_i}{\sum_{i=r}^n \lambda_i}$$
$$\beta_r \triangleq \frac{\sum_{i=r}^n \lambda_i r_i}{\sum_{i=r}^n \lambda_i}$$

458   Remark that $\tilde{P} \in \Pi_{a,b}(r)$, therefore we obtain that

$$|\text{LOT}_{r,c}(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}}) - \text{OT}_c(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}})| = \text{LOT}_{r,c}(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}}) - \text{OT}_x(\mu_{a,\mathbf{X}}, \nu_{b,\mathbf{Y}})$$
$$\leq \langle C, \tilde{P} \rangle - \langle C, P \rangle$$
$$\leq \langle C, \left(\sum_{i=r}^n \lambda_i \right) \alpha_r \beta_r^T \rangle - \langle C, \sum_{i=r}^n \lambda_i q_i r_i^T \rangle$$
$$\leq \langle C, \left(\sum_{i=r}^n \lambda_i \right) \alpha_r \beta_r^T \rangle$$
$$\leq \|C\|_\infty \sum_{i=r}^n \lambda_i \leq \|C\|_\infty \sum_{i=r}^n \frac{1}{i} \leq \|C\|_\infty \ln(n/(r-1))$$

459   $\square$

B.2   Proof of Proposition 2

461   **Proposition 10.** *Let $\mu \in \mathcal{M}_1^+(\mathcal{X})$, $\nu \in \mathcal{M}_1^+(\mathcal{Y})$ and let us assume that c is L-Lipschitz w.r.t. x and*
462   *y. Then for any $r \geq 1$, we have*

$$|LOT_{r,c}(\mu, \nu) - OT_c(\mu, \nu)| \leq 2L \max(\mathcal{N}_{\lfloor \log_2(\lfloor \sqrt{r} \rfloor) \rfloor}(\mathcal{X}, d_\mathcal{X}), \mathcal{N}_{\lfloor \log_2(\lfloor \sqrt{r} \rfloor) \rfloor}(\mathcal{Y}, d_\mathcal{Y}))$$

463   *Proof.* As $\mathcal{X}$ and $\mathcal{Y}$ are compact, $\mathcal{N}_{\lfloor \log_2(\lfloor \sqrt{r} \rfloor) \rfloor}(\mathcal{X}, d), \mathcal{N}_{\lfloor \log_2(\lfloor \sqrt{r} \rfloor) \rfloor}(\mathcal{Y}, d) < +\infty$ and then by de-
464   noting $\varepsilon_\mathcal{X} \triangleq \mathcal{N}_{\lfloor \log_2(\lfloor \sqrt{r} \rfloor) \rfloor}(\mathcal{X}, d_\mathcal{X})$, there exists $x_1, \ldots, x_{\lfloor \sqrt{r} \rfloor} \in \mathcal{X}$, such that $\mathcal{X} \subset \bigcup_{i=1}^r \mathcal{B}_\mathcal{X}(x_i, \varepsilon)$
465   from which we can extract a partition $(S_{i,\mathcal{X}})_{i=1}^{\lfloor \sqrt{r} \rfloor}$ of $\mathcal{X}$ such that for all $i \in [|1, \lfloor \sqrt{r} \rfloor|]$, and
466   $x, y \in S_{i,\mathcal{X}}$, $d_\mathcal{X}(x, y) \leq \varepsilon_\mathcal{X}$. Similarly we can build a partition $(S_{i,\mathcal{Y}})_{i=1}^{\lfloor \sqrt{r} \rfloor}$ of $\mathcal{Y}$. Let us now
467   define for all $k \in [|1, \lfloor \sqrt{r} \rfloor|]$,

$$\mu_k \triangleq \frac{\mu|_{S_{k,\mathcal{X}}}}{\mu(S_{k,\mathcal{X}})} \text{ and } \nu_k \triangleq \frac{\nu|_{S_{k,\mathcal{Y}}}}{\nu(S_{k,\mathcal{Y}})}$$

468   with the convention that $\frac{0}{0} = 0$, we can define

$$\pi_r \triangleq \sum_{i,j=1}^{\lfloor \sqrt{r} \rfloor} \pi^*(S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}) \nu_j \otimes \mu_i .$$

First remarks that $\pi_r \in \Pi_r(\mu, \nu)$. Indeed we have for any measurable set B

$$
\begin{aligned}
\pi_r(\mathcal{X} \times B) &= \sum_{j=1}^{\lfloor \sqrt{r} \rfloor^2} \nu_j(B) \sum_{i=1}^{r} \pi^*(S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}) \\
&= \sum_{j=1}^{\lfloor \sqrt{r} \rfloor} \nu_j(B) \nu(S_{j,\mathcal{Y}}) \\
&= \sum_{j=1}^{\lfloor \sqrt{r} \rfloor} \nu|_{S_{j,\mathcal{X}}}(B) \\
&= \nu(B) ,
\end{aligned}
$$

similarly $\pi_r(A \times \mathcal{Y}) = \mu(A)$ and we have that $\lfloor \sqrt{r} \rfloor^2 \leq r$. Therefore we obtain that

$$
\begin{aligned}
|\mathrm{LOT}_{r,c}(\mu, \nu) - \mathrm{OT}_c(\mu, \nu)| &= \mathrm{LOT}_{r,c}(\mu, \nu) - \mathrm{OT}_c(\mu, \nu) \\
&\leq \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi_r(x, y) - \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi^*(x, y) \\
&\leq \sum_{i,j=1}^{\lfloor \sqrt{r} \rfloor} \int_{S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}} c(x, y) d[\pi_r(x, y) - \pi^*(x, y)] \\
&\leq \sum_{i,j=1}^{\lfloor \sqrt{r} \rfloor} \pi^*(S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}) \\
&\quad \times [\sup_{(x,y) \in S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}} c(x, y) - \inf_{(x,y) \in S_{i,\mathcal{X}} \times S_{j,\mathcal{Y}}} c(x, y)] \\
&\leq L[\varepsilon_{\mathcal{X}} + \varepsilon_{\mathcal{Y}}]
\end{aligned}
$$

from which the result follows. $\qquad \square$

Corollary. *Under the same assumptions of Proposition 2 and by assuming in addition that there exists a Monge map solving $OT_c(\mu, \nu)$, we obtain that for any $r \geq 1$,*

$$
|\mathrm{LOT}_{r,c}(\mu, \nu) - \mathrm{OT}_c(\mu, \nu)| \leq L\mathcal{N}_{\lfloor \log_2(r) \rfloor}(\mathcal{Y}, d_{\mathcal{Y}})
$$

Proof. Let us denote T a Monge map solution of $\mathrm{OT}_c(\mu, \nu)$ and as in the proof above, let us consider a partition of $(S_{i,\mathcal{Y}})_{i=1}^r$ of $\mathcal{Y}$ such that for all $i \in [\![1, r]\!]$, and $x, y \in S_{i,\mathcal{Y}}$, $d_{\mathcal{Y}}(x, y) \leq \varepsilon_{\mathcal{Y}}$ with $\varepsilon_{\mathcal{Y}} \triangleq \mathcal{N}_{\lfloor \log_2(r) \rfloor}(\mathcal{Y}, d_{\mathcal{Y}})$. Let us now define for all $k \in [\![1, \lfloor \sqrt{r} \rfloor]\!]$,

$$
\mu_k \triangleq \frac{\mu|_{T^{-1}(S_{k,\mathcal{Y}})}}{\mu(T^{-1}(S_{k,\mathcal{Y}}))} \quad \text{and} \quad \nu_k \triangleq \frac{\nu|_{S_{k,\mathcal{Y}}}}{\nu(S_{k,\mathcal{Y}})}
$$

with the convention that $\frac{0}{0} = 0$, we can define

$$
\pi_r \triangleq \sum_{k=1}^{r} \pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) \nu_k \otimes \mu_k .
$$

478   Again we have that $\pi_r \in \Pi_r(\mu, \nu)$, and we obtain that

$$|\text{LOT}_{r,c}(\mu, \nu) - \text{OT}_c(\mu, \nu)| = \text{LOT}_{r,c}(\mu, \nu) - \text{OT}_c(\mu, \nu)$$

$$\leq \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi_r(x, y) - \int_{\mathcal{X} \times \mathcal{Y}} c(x, y) d\pi^*(x, y)$$

$$\leq \sum_{k=1}^{r} \pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) \int_{T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}} c(x, y) d\mu_k(y) \otimes \nu_k(y)$$

$$- \sum_{k=1}^{r} \int_{T^{-1}(S_{k,\mathcal{Y}})} c(x, T(x)) d\mu(x)$$

$$\leq \sum_{k=1}^{r} \pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) \int_{T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}} c(x, y) d\mu_k(y) \otimes \nu_k(y)$$

$$- \sum_{k=1}^{r} \pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) \int_{T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}} c(x, T(x)) d\mu_k(x) \otimes \nu_k(y)$$

$$\leq \sum_{k=1}^{r} \pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) \int_{T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}} [c(x, y) - c(x, T(x))] d\mu_k(x) \otimes \nu_k(x)$$

$$\leq L\varepsilon_{\mathcal{Y}}$$

479   from which the result follows. Note that to obtain the above inequalities, we use the fact that π^* is
480   supported on the graph of T, and therefore we have have for all $k \in [\![1, r]\!]$,

$$\pi^*(T^{-1}(S_{k,\mathcal{Y}}) \times S_{k,\mathcal{Y}}) = \mu(T^{-1}(S_{k,\mathcal{Y}})) = \nu(S_{k,\mathcal{Y}}).$$

481   $\qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \qquad \square$

B.3   Proof of Proposition 3

483   **Proposition.** *Let $r \geq 1$ and $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, then $\text{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) \xrightarrow[n \to +\infty]{} \text{LOT}_{r,c}(\mu, \nu)$ a.s.*

Proof. Let π^* solution of $\text{LOT}_{r,c}(\mu, \nu)$. Then there exists $\lambda^* \in \Delta_r^*$, $(\mu_i^*)_{i=1}^r$, $(\nu_i^*)_{i=1}^r \in \mathcal{M}_1^+(\mathcal{X})^r$ such that

$$\pi^* = \sum_{i=1}^{r} \lambda_i^* \mu_i^* \otimes \nu_i^*.$$

484   Note that by definition, we have that

$$\mu = \sum_{i=1}^{r} \lambda_i^* \mu_i^* \quad \text{and} \quad \nu = \sum_{i=1}^{r} \lambda_i^* \nu_i^* .$$

485   Let us now define π_μ and π_μ both elements of $\mathcal{M}_1^+(\mathcal{X} \times [\![1, r]\!])$ as follows:

$$\pi_\mu(A \times \{k\}) \triangleq \lambda_k \mu_k(A) \quad \text{and} \quad \pi_\nu(A \times \{k\}) \triangleq \lambda_k \nu_k(A) \quad \text{for any measurable set } A \text{ and } k \in [\![1, r]\!] .$$

486   Observe that the right marginals of π_μ and π_ν is the same and we will denote it ρ. We can now
487   define for all $x, y \in \mathcal{X}$ the family of kernels $(k_\mu(\cdot, x))_{x \in \mathcal{X}} \in \mathcal{M}_1^+([\![1, r]\!])^{\mathcal{X}}$ and $(k_\nu(\cdot, y))_{y \in \mathcal{X}} \in$
488   $\mathcal{M}_1^+([\![1, r]\!])^{\mathcal{X}}$ corresponding to the disintegration with respect to the projection of respectively μ
489   and ν. Let us now consider n independent samples $(Z_i^\mu)_{i=1}^n$ and $(Z_i^\nu)_{i=1}^n$ such that for all $i \in [\![1, n]\!]$,
490   $Z_i^\mu \sim k_\mu(\cdot, X_i)$ and $Z_i^\nu \sim k_\nu(\cdot, Y_i)$ and let us define for all $k \in [\![1, r]\!]$

$$\tilde{\mu}_k \triangleq \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}_{Z_i^\mu = k} \delta_{X_i} \quad \text{and} \quad \tilde{\nu}_k \triangleq \frac{1}{n} \sum_{i=1}^{n} \mathbf{1}_{Z_i^\nu = k} \delta_{Y_i} .$$

491   Let us now define

$$\tilde{\pi} \triangleq \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k| |\tilde{\nu}_k|} \tilde{\mu}_k \otimes \tilde{\nu}_k$$

$$+ \frac{1}{1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)} \left[\hat{\mu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k \right] \otimes \left[\hat{\nu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \tilde{\nu}_k \right]$$

with the convention that $\frac{0}{0} = 0$. Now it is easy to check that $\tilde{\pi} \in \Pi_r(\hat{\mu}, \hat{\nu})$, indeed we have that

$$
\begin{aligned}
\tilde{\pi}(A \times \mathcal{X}) &= \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k(A) \\
&\quad + \frac{1}{1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)} \left[\hat{\mu}(A) - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k(A) \right] \left[1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|) \right] \\
&= \hat{\mu}(A)
\end{aligned}
$$

in addition by construction we have that

$$
\left| \hat{\mu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k \right| = \left| \hat{\nu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \tilde{\nu}_k \right| = 1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)
$$

and both $\hat{\mu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k$ and $\hat{\nu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \tilde{\nu}_k$ are positive measures. Therefore we obtain that

$$
\text{LOT}_{r,c}(\hat{\mu}, \hat{\nu}) \leq \int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}(x, y)
$$

Now we aim at showing at $\int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}(x, y) \to \text{LOT}_{r,c}(\mu, \nu)$ $a.s.$. Indeed first observe that from the law of large numbers we have that for all $k \in [|1, r|]$, $|\tilde{\mu}_k| \to \lambda_k^*$ and similarly $|\tilde{\nu}_k| \to \lambda_k^*$. In addition, for all k, q we have that almost surely, $\tilde{\mu}_k \otimes \tilde{\nu}_q$ converges weakly towards $\lambda_k^* \lambda_q^* \mu_k \otimes \nu_q$. Indeed one can consider the following algebra $\mathcal{F} \triangleq \{(x, y) \in \mathcal{X}^2 \to f(x)g(y) \ f, g \in \mathcal{C}(\mathcal{X})\}$, and then by Stone-Weierstrass, one obtains by density the desired result. Now remark that

$$
\begin{aligned}
\int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}(x, y) &= \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \int_{\mathcal{X}^2} c(x, y) d\tilde{\mu}_k \otimes \tilde{\nu}_k \\
&\quad + \frac{1}{\tilde{\lambda}_r} \int_{\mathcal{Z}^2} c(x, y) d\tilde{\mu}_r \otimes \tilde{\nu}_r \\
&\quad + \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) \int_{\mathcal{X}^2} c(x, y) d\tilde{\mu}_r \otimes \tilde{\nu}_k \\
&\quad + \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \int_{\mathcal{X}^2} c(x, y) d\tilde{\mu}_k \otimes \tilde{\nu}_r \\
&\quad + \frac{1}{\tilde{\lambda}_r} \sum_{k,q=1}^{r-1} \int_{\mathcal{X}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \left(1 - \frac{\min(|\tilde{\mu}_q|, |\tilde{\nu}_q|)}{|\tilde{\nu}_q|} \right) c(x, y) d\tilde{\mu}_k(x) d\tilde{\nu}_q(y)
\end{aligned}
$$

from which follows directly that $\int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}(x, y) \to \text{LOT}_{r,c}(\mu, \nu)$ $a.s.$ Let us now denote for all $n \geq 1$, π_n a solution of $\text{LOT}_{r,c}(\hat{\mu}, \hat{\nu})$. Let $\omega \in \Omega$ an element of the probability space where live the random variables $(X_i)_{i \geq 0}$ and $(Y_i)_{i \geq 0}$ such that $\int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}^{(\omega)}(x, y) \to \text{LOT}_{r,c}(\mu, \nu)$. As $\mathcal{X}$ is compact Thanks to Prokhorov's Theorem, we can extract a sequence such that $(\pi_n^{(\omega)})_{n \geq 0}$ converge weakly towards $\pi^{(\omega)} \in \Pi_r(\mu, \nu)$. In addition we have that for all $n \geq 1$

$$
\int_{\mathcal{X}^2} c(x, y) d\pi_n^{(\omega)}(x, y) \leq \int_{\mathcal{X}^2} c(x, y) d\tilde{\pi}^{(\omega)}(x, y)
$$

And by considering the limit we obtain that

$$
\int c(x, y) d\pi^{(\omega)}(x, y) \leq \text{LOT}_{r,c}(\mu, \nu)
$$

However $\pi^{(\omega)} \in \Pi_r(\mu, \nu)$ and by optimality we obtain that

$$
\int c(x, y) d\pi^{(\omega)}(x, y) = \text{LOT}_{r,c}(\mu, \nu)
$$

This holds for an arbitrary subsequence of $(\pi_n^{(\omega)})_{n \geq 0}$, from which follows that $\int c(x, y) d\pi_n^{(\omega)}(x, y) \to \text{LOT}_{r,c}(\mu, \nu)$. Finally this holds almost surely and the result follows. $\quad \square$

 B.4 Proof of Proposition 4

 Proposition. *Let $r \geq 1$ and $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$. Then, there exists a constant K_r such that for any*
 $\delta > 0$ and $n \geq 1$, we have, with a probability of at least $1 - 2\delta$, that

$$\mathrm{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) - \mathrm{LOT}_{r,c}(\mu, \nu) \leq 11\|c\|_\infty \sqrt{\frac{r}{n}} + K_r\|c\|_\infty \left[\sqrt{\frac{\log(40/\delta)}{n}} + \frac{\sqrt{r}\log(40/\delta)}{n} \right]$$

Proof. We reintroduce the same notation as in the proof of Proposition 3. Let π^* solution of $\mathrm{LOT}_{r,c}(\mu, \nu)$. Then there exists $\lambda^* \in \Delta_r^*$, $(\mu_i^*)_{i=1}^r, (\nu_i^*)_{i=1}^r \in \mathcal{M}_1^+(\mathcal{Z})^r$ such that

$$\pi^* = \sum_{i=1}^r \lambda_i^* \mu_i^* \otimes \nu_i^*.$$

 As before let us also consider π_μ and π_μ defined as $\pi_\mu(A \times \{k\}) \triangleq \lambda_k \mu_k(A)$ and $\pi_\nu(A \times \{k\}) \triangleq$
 $\lambda_k \nu_k(A)$ for any measurable set A and $k \in [\![1, r]\!]$ and denote ρ their common right marginal. We
 also consider n independent samples $(Z_i^\mu)_{i=1}^n$ and $(Z_i^\nu)_{i=1}^n$ such that for all $i \in [\![1, n]\!]$, $Z_i^\mu \sim$
 $k_\mu(\cdot, X_i)$ and $Z_i^\nu \sim k_\nu(\cdot, Y_i)$ and we denote for all $k \in [\![1, r]\!]$

$$\tilde{\mu}_k \triangleq \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{Z_i^\mu = k} \delta_{X_i} \text{ and } \tilde{\nu}_k \triangleq \frac{1}{n} \sum_{i=1}^n \mathbf{1}_{Z_i^\nu = k} \delta_{Y_i}$$

Let us now define

$$\hat{\pi} \triangleq \sum_{i=1}^r \frac{1}{\lambda_k^*} \tilde{\mu}_k \otimes \tilde{\nu}_k .$$

 Our goal is to control the following quantity:

$$\left| \mathrm{LOT}_{r,c}(\mu, \nu) - \int_{\mathcal{Z}^2} c(x, y) d\hat{\pi}(x, y) \right|,$$

 First observe that

$$\mathbb{E}\left[\int_{\mathcal{Z}^2} c(x, y) d\hat{\pi}(x, y) \right] = \sum_{i=1}^r \frac{1}{\lambda_k^*} \mathbb{E}\left[\int_{\mathcal{Z}^2} c(x, y) d\tilde{\mu}_k(x) d\tilde{\nu}_k(y) \right]$$

$$= \sum_{i=1}^r \frac{1}{\lambda_k^* n^2} \times \sum_{i,j} \mathbb{E}\left[c(X_i, Y_j) \mathbf{1}_{Z_i^\mu = k} \mathbf{1}_{Z_j^\nu = k} \right]$$

 Moreover, we have that

$$\mathbb{E}\left[c(X_i, Y_j) \mathbf{1}_{Z_i^\mu = k} \mathbf{1}_{Z_j^\nu = k} \right] = \int_{(\mathcal{Z} \times [\![1,r]\!])^2} c(x, y) \mathbf{1}_{z=k} \mathbf{1}_{z'=k} d\pi_\mu(x, z) d\pi_\nu(y, z')$$

$$= \int_{(\mathcal{Z} \times [\![1,r]\!])^2} c(x, y) \mathbf{1}_{z=k} \mathbf{1}_{z'=k} d\mu_z(x) d\nu_{z'}(y) d\rho(z) d\rho(z')$$

$$= \lambda_k^2 \int_{\mathcal{Z}^2} c(x, y) d\mu_k(x) d\nu_k(y)$$

 from which follows that

$$\mathbb{E}\left[\int_{\mathcal{Z}^2} c(x, y) d\hat{\pi}(x, y) \right] = \sum_{i=1}^r \lambda_k^* \int_{\mathcal{Z}^2} c(x, y) d\mu_k(x) d\nu_k(y) = \mathrm{LOT}_{r,c}(\mu, \nu)$$

 Now let us define for all $(x_i, z_i)_{i=1}^n, (y_i, z_i') \in (\mathcal{Z} \times [\![1, r]\!])^n$,

$$g((x_1, z_1), \ldots, (x_n, z_n), (y_1, z_1'), \ldots, (y_n, z_n')) \triangleq \sum_{q=1}^r \frac{1}{\lambda_q^* n^2} \sum_{i,j} c(x_i, y_j) \mathbf{1}_{z_i = q} \mathbf{1}_{z_j' = q} ,$$

since $\mathcal{Z}$ is compact and c is continuous, we have that

$$
\begin{aligned}
|g(\dots, (x_k, z_k), \dots) - g(\dots, (\tilde{x}_k, \tilde{z}_k), \dots)| &= \left| \sum_{q=1}^{r} \frac{1}{\lambda_q^* n^2} \sum_j [c(x_k, y_j)\mathbf{1}_{z_k=q} - c(\tilde{x}_k, y_j)\mathbf{1}_{\tilde{z}_k=q}]\mathbf{1}_{z_j'=q} \right| \\
&= \left| \frac{1}{\lambda_{z_k}^* n^2} \sum_{j=1}^{n} c(x_k, y_j)\mathbf{1}_{z_j'=z_k} - \frac{1}{\lambda_{\tilde{z}_k}^* n^2} \sum_{j=1}^{n} c(\tilde{x}_k, y_j)\mathbf{1}_{z_j'=\tilde{z}_k} \right| \\
&\leq \frac{\|c\|_\infty}{n^2} \left[\frac{\sum_{j=1}^{n} \mathbf{1}_{z_j'=z_k}}{\lambda_{z_k}^*} + \frac{\sum_{j=1}^{n} \mathbf{1}_{z_j'=\tilde{z}_k}}{\lambda_{z_k}^*} \right] \\
&\leq \frac{2\|c\|_\infty}{\min\limits_{1\leq q\leq r} \lambda_q^*} \frac{1}{n}
\end{aligned}
$$

Then by applying the McDiarmid's inequality we obtain that for $\delta > 0$, with a probability at least of $1 - \delta$, we have

$$
\left| \mathrm{LOT}_{r,c}(\mu, \nu) - \int_{\mathcal{Z}^2} c(x, y) d\hat{\pi}(x, y) \right| \leq \frac{2\|c\|_\infty}{\min\limits_{1\leq q\leq r} \lambda_q^*} \sqrt{\frac{\log(2/\delta)}{n}}
$$

Now we aim at building a coupling $\tilde{\pi} \in \Pi_r(\hat{\mu}, \hat{\nu})$ from $\hat{\pi}$. Let us consider the same as the one introduce in the proof of Proposition B.3, that is

$$
\begin{aligned}
\tilde{\pi} &\triangleq \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \tilde{\mu}_k \otimes \tilde{\nu}_k \\
&+ \frac{1}{1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)} \left[\hat{\mu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \tilde{\mu}_k \right] \otimes \left[\hat{\nu} - \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \tilde{\nu}_k \right]
\end{aligned}
$$

with the convention that $\frac{0}{0} = 0$. Let us now expand the above expression, and by denoting $\tilde{\lambda}_r = 1 - \sum_{k=1}^{r-1} \min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)$ we obtain that

$$
\begin{aligned}
\tilde{\pi} &= \sum_{k=1}^{r-1} \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \tilde{\mu}_k \otimes \tilde{\nu}_k \\
&+ \frac{1}{\tilde{\lambda}_r} \tilde{\mu}_r \otimes \tilde{\nu}_r \\
&+ \frac{1}{\tilde{\lambda}_r} \tilde{\mu}_r \otimes \left[\sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) \tilde{\nu}_k \right] \\
&+ \frac{1}{\tilde{\lambda}_r} \left[\sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \tilde{\mu}_k \right] \otimes \tilde{\nu}_r \\
&+ \frac{1}{\tilde{\lambda}_r} \left[\sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \tilde{\mu}_k \right] \otimes \left[\sum_{k=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) \tilde{\nu}_k \right]
\end{aligned}
$$

528 Now we aim at controlling the following quantity $\left| \int_{\mathcal{Z}^2} c(x,y) d\hat{\pi}(x,y) - \int_{\mathcal{Z}^2} c(x,y) d\tilde{\pi}(x,y) \right|$ and
529 we observe that

$$\int_{\mathcal{Z}^2} c(x,y) d[\hat{\pi}(x,y) - \tilde{\pi}(x,y)] = \sum_{k=1}^{r-1} \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_k^*} - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \right] d\tilde{\mu}_k(x)\tilde{\nu}_k(y) \quad (11)$$

$$+ \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_r^*} - \frac{1}{\tilde{\lambda}_r} \right] d\tilde{\mu}_r(x)\tilde{\nu}_r(y) \quad (12)$$

$$+ \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) c(x,y) d\tilde{\mu}_r(x) d\tilde{\nu}_k(y) \quad (13)$$

$$+ \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) c(x,y) d\tilde{\mu}_k(x) d\tilde{\nu}_r(y) \quad (14)$$

$$+ \frac{1}{\tilde{\lambda}_r} \sum_{k,q=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \left(1 - \frac{\min(|\tilde{\mu}_q|,|\tilde{\nu}_q|)}{|\tilde{\nu}_q|} \right) c(x,y) d\tilde{\mu}_k(x) d\tilde{\nu}_q(y) \quad (15)$$

530 Let us now control each term of the RHS of the above equality. Let us first consider the term in
531 Eq. 11, remark that we have

$$\left| \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_k^*} - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \right] d\tilde{\mu}_k(x)\tilde{\nu}_k(y) \right|$$

$$\leq \left| \left[\frac{1}{\lambda_k^*} - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \right] \right| \|c\|_\infty |\tilde{\mu}_k||\tilde{\nu}_k|$$

$$\leq \left| \left[\frac{|\tilde{\mu}_k||\tilde{\nu}_k|}{\lambda_k^*} - \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|) \right] \right| \|c\|_\infty$$

$$\leq \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|) \left| \frac{\max(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{\lambda_k^*} - 1 \right| \|c\|_\infty$$

$$\leq \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{\lambda_k^*} |\max(|\tilde{\mu}_k|,|\tilde{\nu}_k|) - \lambda_k^*| \|c\|_\infty$$

$$\leq \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{\lambda_k^*} \max(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty, \|\tilde{\lambda}_\nu - \lambda^*\|_\infty) \|c\|_\infty$$

$$\leq \|c\|_\infty \max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right) \max(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty, \|\tilde{\lambda}_\nu - \lambda^*\|_\infty)$$

532 where we have denoted $\tilde{\lambda}_\mu \triangleq (|\tilde{\mu}_k|)_{k=1}^r$ and $\tilde{\lambda}_\nu \triangleq (|\tilde{\nu}_k|)_{k=1}^r$. Now observe that

$$\mathbb{P}\left(\max(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty, \|\tilde{\lambda}_\nu - \lambda^*\|_\infty) \geq t \right) \leq 2\mathbb{P}\left(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty \geq t \right)$$

$$\leq \mathbb{P}\left(d_K(\lambda^*, \tilde{\lambda}_\mu) \geq \frac{t}{2} \right)$$

$$\leq 4\exp(-nt^2/2)$$

533 where d_K is the Kolmogorov distance. In addition we have

$$\max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right) \leq 1 + \frac{1}{\min_{1\leq i \leq r} \lambda_i^*} \max\left(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty, \|\tilde{\lambda}_\nu - \lambda^*\|_\infty \right)$$

534 Combining the two above controls, we obtain that for all $\delta > 0$, with a probability of at least $1 - \delta$,

$$\left| \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_k^*} - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k||\tilde{\nu}_k|} \right] d\tilde{\mu}_k(x)\tilde{\nu}_k(y) \right| \leq \|c\|_\infty \sqrt{\frac{2\ln 8/\delta}{n}} + \frac{\|c\|_\infty}{n} \frac{2\ln 8/\delta}{\min_{1\leq i \leq r} \lambda_i^*}$$

Let us now consider the term in Eq. 12, we have that

$$\left| \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_r^*} - \frac{1}{\tilde{\lambda}_r} \right] d\tilde{\mu}_r(x)\tilde{\nu}_r(y) \right| \leq \frac{|\tilde{\mu}_r||\tilde{\nu}_r|}{\lambda_r^*\tilde{\lambda}_r} \left| 1 - \sum_{i=1}^{r} \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|) - \lambda_r \right| \|c\|_\infty$$

$$\leq \max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right) \sum_{k=1}^{r-1} |\lambda_k^* - \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)| \, \|c\|_\infty$$

$$\leq \max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right) \|c\|_\infty (\|\lambda^* - \tilde{\lambda}_\mu\|_1 + \|\lambda^* - \tilde{\lambda}_\nu\|_1)$$

$$\leq 2\|c\|_\infty \max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right) \max(\|\lambda^* - \tilde{\lambda}_\mu\|_1, \|\lambda^* - \tilde{\lambda}_\nu\|_1)$$

However we have that

$$\mathbb{P}\left(\max(\|\lambda^* - \tilde{\lambda}_\mu\|_1, \|\lambda^* - \tilde{\lambda}_\nu\|_1) \geq t \right) \leq 2\mathbb{P}\left(\|\lambda^* - \tilde{\lambda}_\mu\|_1 \geq t \right)$$

In addition we have that $\mathbb{E}(\|\lambda^* - \tilde{\lambda}_\mu\|_1) \leq \sqrt{\frac{r}{n}}$ and by applying the McDiarmid's Inequality, we obtain that for all $\delta > 0$, with a probability of $1 - \delta$

$$\|\lambda^* - \tilde{\lambda}_\mu\|_1 \leq \sqrt{\frac{r}{n}} + \sqrt{\frac{2\ln(2/\delta)}{n}}$$

Therefore we obtain that with a probability of at least $1 - \delta$,

$$\left| \int_{\mathcal{Z}^2} c(x,y) \left[\frac{1}{\lambda_r^*} - \frac{1}{\tilde{\lambda}_r} \right] d\tilde{\mu}_r(x)\tilde{\nu}_r(y) \right| \leq 2\|c\|_\infty \left[\sqrt{\frac{r}{n}} + \sqrt{\frac{2\ln(8/\delta)}{n}} + \frac{2\ln(8/\delta) + \sqrt{2r\ln(8/\delta)}}{n \times \min_{1 \leq i \leq r} \lambda_i^*} \right]$$

For the term in Eq. 13 and 14, we obtain that

$$\left| \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) c(x,y) d\tilde{\mu}_r(x) d\tilde{\nu}_k(y) \right|$$

$$\leq \frac{|\tilde{\mu}_r|}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} (|\tilde{\nu}_k| - \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)) \|c\|_\infty$$

$$\leq \frac{|\tilde{\mu}_r|}{\tilde{\lambda}_r} [\tilde{\lambda}_r - |\tilde{\nu}_r|] \|c\|_\infty$$

$$\leq [|\tilde{\lambda}_r - \lambda_r^*| + |\lambda_r^* - \tilde{\nu}_r|] \|c\|_\infty$$

$$\leq 3\|c\|_\infty \max(\|\lambda^* - \tilde{\lambda}_\mu\|_1, \|\lambda^* - \tilde{\lambda}_\nu\|_1)$$

Therefore we obtain that with a probability of at least $1 - \delta$,

$$\left| \frac{1}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\nu}_k|} \right) c(x,y) d\tilde{\mu}_r(x) d\tilde{\nu}_k(y) \right| \leq 3\|c\|_\infty \left[\sqrt{\frac{r}{n}} + \sqrt{\frac{2\ln(2/\delta)}{n}} \right]$$

Finally the last term in Eq. 15 can be controlled as the following:

$$\left| \frac{1}{\tilde{\lambda}_r} \sum_{k,q=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \left(1 - \frac{\min(|\tilde{\mu}_q|,|\tilde{\nu}_q|)}{|\tilde{\nu}_q|} \right) c(x,y) d\tilde{\mu}_k(x) d\tilde{\nu}_q(y) \right|$$

$$\leq \frac{\|c\|_\infty}{\tilde{\lambda}_r} \sum_{k,q=1}^{r-1} \left(1 - \frac{\min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \left(1 - \frac{\min(|\tilde{\mu}_q|,|\tilde{\nu}_q|)}{|\tilde{\nu}_q|} \right) |\tilde{\mu}_k||\tilde{\nu}_q|$$

$$\leq \frac{\|c\|_\infty}{\tilde{\lambda}_r} \sum_{k=1}^{r-1} (|\tilde{\mu}_k| - \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|)) \sum_{k=1}^{r-1} (|\tilde{\nu}_k| - \min(|\tilde{\mu}_k|,|\tilde{\nu}_k|))$$

$$\leq 3\|c\|_\infty \max(\|\lambda^* - \tilde{\lambda}_\mu\|_1, \|\lambda^* - \tilde{\lambda}_\nu\|_1)$$

543 and we obtain that with a probability of at least $1 - \delta$,

$$
\left| \frac{1}{\tilde{\lambda}_r} \sum_{k,q=1}^{r-1} \int_{\mathcal{Z}^2} \left(1 - \frac{\min(|\tilde{\mu}_k|, |\tilde{\nu}_k|)}{|\tilde{\mu}_k|} \right) \left(1 - \frac{\min(|\tilde{\mu}_q|, |\tilde{\nu}_q|)}{|\tilde{\nu}_q|} \right) c(x,y) d\tilde{\mu}_k(x) d\tilde{\nu}_q(y) \right|
$$

$$
\leq 3\|c\|_\infty \left[\sqrt{\frac{r}{n}} + \sqrt{\frac{2\ln(2/\delta)}{n}} \right]
$$

544 Then by applying a union bound we obtain that with a probability of at least $1 - \delta$

$$
\left| \int_{\mathcal{Z}^2} c(x,y) d[\hat{\pi}(x,y) - \tilde{\pi}(x,y)] \right| \leq \|c\|_\infty \left[11\sqrt{\frac{r}{n}} + 12\sqrt{\frac{2\ln 40/\delta}{n}} + \frac{6\ln(40/\delta) + 2\sqrt{2r\ln(40/\delta)}}{n \times \min\limits_{1 \leq i \leq r} \lambda_i^*} \right]
$$

545 Now observe that

$$
\mathrm{LOT}_{r,c}(\hat{\mu}, \hat{\nu}) - \mathrm{LOT}_{r,c}(\mu, \nu) \leq \int_{\mathcal{Z}^2} c(x,y) d\tilde{\pi}(x,y) - \int_{\mathcal{Z}^2} c(x,y) d\pi^*(x,y)
$$

$$
\leq \int_{\mathcal{Z}^2} c(x,y) d[\tilde{\pi} - \hat{\pi}](x,y) + \int_{\mathcal{Z}^2} c(x,y) d[\hat{\pi} - \pi^*](x,y)
$$

546 and by combining the two control we obtain that with a probability of at least $1 - 2\delta$,

$$
\mathrm{LOT}_{r,c}(\hat{\mu}, \hat{\nu}) - \mathrm{LOT}_{r,c}(\mu, \nu) \leq \|c\|_\infty \left[11\sqrt{\frac{r}{n}} + 12\sqrt{\frac{2\ln 40/\delta}{n}} + \frac{1}{\alpha}\left(2\sqrt{\frac{\log(2/\delta)}{n}} + \frac{6\ln(40/\delta) + 2\sqrt{2r\ln(40/\delta)}}{n} \right) \right]
$$

$$
\leq 11\|c\|_\infty \sqrt{\frac{r}{n}} + \frac{14\|c\|_\infty}{\alpha}\sqrt{\frac{\log(40/\delta)}{n}} + \frac{2\|c\|_\infty \max(6, \sqrt{2r})\log(40/\delta)}{n\alpha}
$$

547 where $\alpha \triangleq \min\limits_{1 \leq i \leq r} \lambda_i^*$ and the result follows. $\qquad\square$

B.5 Proof Proposition 5

549 **Proposition.** *Let $r \geq 1$, $\delta > 0$ and $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$. Then there exists a constant $N_{r,\delta}$ such that if*
550 $n \geq N_{r,\delta}$ *then with a probability of at least $1 - 2\delta$, we have*

$$
\mathrm{LOT}_{r,c}(\hat{\mu}_n, \hat{\nu}_n) - \mathrm{LOT}_{r,c}(\mu, \nu) \leq 11\|c\|_\infty \sqrt{\frac{r}{n}} + 77\|c\|_\infty \sqrt{\frac{\log(40/\delta)}{n}} \ .
$$

551 *Proof.* We consider the same notations as in the proof of Proposition 4. In particular let us define for
552 all $(x_i, z_i)_{i=1}^n, (y_i, z_i') \in (\mathcal{Z} \times [\![1, r]\!])^n$,

$$
g((x_1, z_1), \dots, (x_n, z_n), (y_1, z_1'), \dots, (y_n, z_n')) \triangleq \sum_{q=1}^r \frac{1}{\lambda_q^* n^2} \sum_{i,j} c(x_i, y_j) \mathbf{1}_{z_i=q} \mathbf{1}_{z_j'=q} \ ,
$$

553 Recall that we have

$$
|g(\dots, (x_k, z_k), \dots) - g(\dots, (\tilde{x}_k, \tilde{z}_k), \dots)| \leq \frac{\|c\|_\infty}{n^2} \left[\frac{\sum_{j=1}^n \mathbf{1}_{z_j'=z_k}}{\lambda_{z_k}^*} + \frac{\sum_{j=1}^n \mathbf{1}_{z_j'=\tilde{z}_k}}{\lambda_{\tilde{z}_k}^*} \right]
$$

$$
\leq \frac{2\|c\|_\infty}{n} \max\left(\left\| \frac{\tilde{\lambda}_\mu}{\lambda^*} \right\|_\infty, \left\| \frac{\tilde{\lambda}_\nu}{\lambda^*} \right\|_\infty \right)
$$

$$
\leq \frac{2\|c\|_\infty}{n} + \frac{2\|c\|_\infty}{n \times \min\limits_{1 \leq i \leq r} \lambda_i^*} \max\left(\|\tilde{\lambda}_\mu - \lambda^*\|_\infty, \|\tilde{\lambda}_\nu - \lambda^*\|_\infty \right)
$$

554 In fact if we have a control in probability of the bounded difference we can use an extension of the
555 McDiarmid's Inequality. For that purpose let us first introduce the following definition.

Definition 4. *Let $(X_i)_{i=1}^m$, m independent random variables and g a measurable function. We say that g is weakly difference-bounded with respect to $(X_i)_{i=1}^m$ by (b, β, δ) if*

$$\mathbb{P}\left(|g(X_1, \ldots, X_m) - g(X_1', \ldots, X_m')| \leq \beta\right) \geq 1 - \delta$$

with $X_i' = X_i$ except for one coordinate k where X_k' is an independent copy of X_k. Furthermore for any $(x_i)_{i=1}^m$ and $(x_i')_{i=1}^m$ where for all coordinate except on $x_j = x_j'$

$$|g(x_1, \ldots, x_m) - g(x_1', \ldots, x_m')| \leq b \, .$$

Let us now introduce an extension of McDiarmid's Inequality [Kutin, 2002].

Theorem 1. *Let $(X_i)_{i=1}^m$, m independent random variables and g a measurable function which is weakly difference-bounded with respect to $(X_i)_{i=1}^m$ by $(b, \beta/m, \exp(-Km))$, then if $0 < \tau \leq T(b, \beta, K)$ and $m \geq M(b, \beta, K, \tau)$, then*

$$\mathbb{P}(|g(X_1, \ldots, X_m) - \mathbb{E}(g(X_1, \ldots, X_m))| \geq \tau) \leq 4\exp\left(\frac{-\tau^2 m}{8\beta^2}\right)$$

where

$$T(b, \beta, K) \triangleq \min\left(\frac{14c}{2}, 4\beta\sqrt{K}, \frac{\beta^2 K}{b}\right)$$

$$M(b, \beta, K, \tau) \triangleq \max\left(\frac{b}{\beta}, \beta\sqrt{40}, 3\left(\frac{24}{K} + 3\right)\log\left(\frac{24}{K} + 3\right), \frac{1}{\tau}\right)$$

Given the above Theroem we can obtain an asymptotic control of the deviation of g from its mean. Let $\delta' > 0$ and let us denote

$$m \triangleq 2n$$

$$b \triangleq \frac{2\|c\|_\infty}{n \times \min\limits_{1 \leq i \leq r} \lambda_i^*}$$

$$K \triangleq \frac{\log(1/\delta')}{2n}$$

$$\beta \triangleq 4\|c\|_\infty \left[1 + \frac{1}{\min\limits_{1 \leq i \leq r} \lambda_i^*}\sqrt{\frac{2\log(4/\delta')}{n}}\right]$$

Observe now that with a probability of at least $1 - \exp(-Km)$

$$|g(\ldots, (x_k, z_k), \ldots) - g(\ldots, (\tilde{x}_k, \tilde{z}_k), \ldots)| \leq \frac{2\|c\|_\infty}{n}\left[1 + \frac{1}{\min\limits_{1 \leq i \leq r} \lambda_i^*}\sqrt{\frac{2\log(4/\delta')}{n}}\right]$$

Let us now fix $\delta > 0$ and let us choose δ' such that $\delta' \triangleq 4/n$ and $\tau \triangleq \beta\sqrt{\frac{4\log(4/\delta)}{n}}$, then we obtain that for n sufficiently large (such that $n \geq M(b, \beta, K, \tau)/2$ and $\tau \leq T(b, \beta, K)$), we have that with a probability of at least $1 - \delta$

$$\left|\text{LOT}_{r,c}(\mu, \nu) - \int_{\mathcal{Z}^2} c(x, y)d\hat{\pi}(x, y)\right| \leq 4\|c\|_\infty\left[1 + \frac{1}{\min\limits_{1 \leq i \leq r} \lambda_i^*}\sqrt{\frac{2\log(n)}{n}}\right]\sqrt{\frac{4\log(4/\delta)}{n}}$$

$$\leq 4\|c\|_\infty\sqrt{\frac{4\log(4/\delta)}{n}} + \frac{16\sqrt{5}\|c\|_\infty\sqrt{\log(n)\log(4/\delta)}}{n \times \min\limits_{1 \leq i \leq r} \lambda_i^*}$$

Recall also from the proof of Proposition 4, that we have with a probability of at least $1 - \delta$

$$\left|\int_{\mathcal{Z}^2} c(x, y)d[\hat{\pi}(x, y) - \tilde{\pi}(x, y)]\right| \leq \|c\|_\infty\left[11\sqrt{\frac{r}{n}} + 12\sqrt{\frac{2\ln 40/\delta}{n}} + \frac{6\ln(40/\delta) + 2\sqrt{2r\ln(40/\delta)}}{n \times \min\limits_{1 \leq i \leq r} \lambda_i^*}\right]$$

572 Finally by imposing in addition that

$$\sqrt{\frac{n}{\log(n)}} \geq \frac{1}{\min\limits_{1 \leq i \leq r} \lambda_i^*} \ , \ \sqrt{n} \geq \frac{\sqrt{\log(40/\delta)}}{\min\limits_{1 \leq i \leq r} \lambda_i^*} \ \text{ and } \ \sqrt{n} \geq \frac{\sqrt{r}}{\min\limits_{1 \leq i \leq r}}$$

573 we obtain that for n is large enough (such that (such that $n \geq M(b, \beta, K, \tau)/2$ and $\tau \leq T(b, \beta, K)$)
574 and satysfing the above inequalities, we have with a probability of at least $1 - 2\delta$ that

$$\text{LOT}_{r,c}(\hat{\mu}, \hat{\nu}) - \text{LOT}_{r,c}(\mu, \nu) \leq 11\|c\|_\infty \sqrt{\frac{r}{n}} + 77\|c\|_\infty \sqrt{\frac{\log(40/\delta)}{n}}$$

575 □

B.6 Proof Proposition 6

576

Proposition. *Let $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$. Let us assume that c is symmetric, then we have*

$$\text{DLOT}_{1,c}(\mu, \nu) = \frac{1}{2} \int_{\mathcal{X}^2} -c(x, y) d[\mu - \nu] \otimes d[\mu - \nu](x, y) \ .$$

577 *If in addition we assume the c is Lipschitz w.r.t to x and y, then we have*

$$\text{DLOT}_{r,c}(\mu, \nu) \xrightarrow[r \to +\infty]{} \text{OT}_c(\mu, \nu) \ .$$

578 *Proof.* When $r = 1$, it is clear that for any $\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, $\Pi_r(\mu, \nu) = \{\mu \otimes \nu\}$ and thanks to the
579 symmetry of c, we have directly that

$$\text{DLOT}_{1,c}(\mu, \nu) = \frac{1}{2} \int_{\mathcal{X}^2} -c(x, y) d[\mu - \nu] \otimes d[\mu - \nu](x, y) = \frac{1}{2}\text{MMD}_{-c}(\mu, \nu) \ .$$

580 The limit is a direct consequence of Proposition 2. □

B.7 Proof of Proposition 8

581

Proposition. *Let $r \geq 1$ and $(\mu_n)_{n \geq 0}$ and $(\nu_n)_{n \geq 0}$ two sequences of probability measures such that
$\mu_n \to \mu$ and $\nu_n \to \nu$ with respect to the convergence in law. Then we have that*

$$\text{LOT}_{r,c}(\mu_n, \nu_n) \to \text{LOT}_{r,c}(\mu, \nu) \ .$$

582 *Proof.* Let us denote π an optimal solution of $\text{LOT}_{r,c}(\mu, \nu)$ and let us denote $(\mu^{(i)})_{i=1}^r$, $(\nu^{(i)})_{i=1}^r$
583 and $(\lambda^{(i)})_{i=1}^r$ the decomposition associated. In the following Lemma, we aim at building specific
584 decompositions of the sequences $(\mu_n)_{n \geq 0}$ and $(\nu_n)_{n \geq 0}$.

585 **Lemma 1.** *Let $r \geq 1$, $\mu \in \mathcal{M}_1^+(\mathcal{X})$ and $(\mu^{(i)})_{i=1}^r \in \mathcal{M}_1^+(\mathcal{X})$ and $(\lambda^{(i)})_{i=1}^r \in \Delta_r^*$ such that*
586 *$\mu = \sum_{i=1}^r \lambda_i \mu^{(i)}$. Then for any sequence of probability measures $(\mu_n)_{\geq 0}$ such that $\mu_n \to \mu$, there*
587 *exist for all $i \in [|1, r|]$ a sequence of nonnegative measures $(\mu_n^{(i)})_{n \geq 0}$ such that*

$$\mu_n^{(i)} \to \lambda_i \mu^{(i)} \ \text{for all } i \in [|1, r|] \text{ and}$$

$$\sum_{i=1}^r \mu_n^{(i)} = \mu_n \ \text{for all } n \geq 0$$

588 *Proof.* For $r = 1$ the result is clear. Let us now show the result for $r = 2$. Let us denote $(\tilde{\mu}_n^{(1)})$ a
589 sequence converging weakly towards $\lambda_1 \mu^{(1)}$. Then by denoting $\mu_n^{(1)} \triangleq \mu_n - (\mu_n - \tilde{\mu}_n^{(1)})_+$ where
590 $(\cdot)_+$ correspond to the non-negative part of the measure, we have that

$$\mu_n^{(1)} \geq 0, \ \mu_n^{(1)} \to \lambda_1 \mu^{(1)},$$

$$\mu_n^{(2)} \triangleq \mu_n - \mu_n^{(1)} \geq 0, \ \mu_n^{(2)} \to \lambda_2 \mu^{(2)} \text{ and}$$

$$\mu_n = \mu_n^{(1)} + \mu_n^{(2)} \ \text{for all } n \geq 0$$

591 which is the result. Let $r \geq 2$ and let us assume that the result holds for all $1 \leq k \leq r$. Let us now
592 consider a decomposition of μ such that $\mu = \sum_{i=1}^{r+1} \lambda_i \mu^{(i)}$. By denoting $\tilde{\mu}^{(1)} \triangleq \frac{\sum_{i=1}^{r} \lambda_i \mu^{(i)}}{\sum_{i=1}^{r} \lambda_i}$, we
593 obtain that

$$\mu = \left(\sum_{i=1}^{r} \lambda_i \right) \tilde{\mu}^{(1)} + \lambda_{r+1} \mu^{(r+1)} .$$

594 Then by recursion we have that there exists sequences of nonnegative measures $(\tilde{\mu}_n^{(1)})$ and $(\mu_n^{(r+1)})$
595 such that

$$\tilde{\mu}_n^{(1)} \rightarrow \left(\sum_{i=1}^{r} \lambda_i \right) \tilde{\mu}^{(1)}, \ \mu_n^{(r+1)} \rightarrow \lambda_{r+1} \mu^{(r+1)} \ \text{and} \ \mu_n = \tilde{\mu}_n^{(1)} + \mu_n^{(r+1)} \ \text{for all} \ n \geq 0$$

596 Now observe that $\frac{\tilde{\mu}_n^{(1)}}{|\tilde{\mu}_n^{(1)}|} \rightarrow \tilde{\mu}^{(1)} = \sum_{i=1}^{r} \frac{\lambda_i}{\sum_{i=1}^{r} \lambda_i} \mu^{(i)}$. Therefore applying the recursion on this
597 problem allows us to obtain a decomposition of $\tilde{\mu}_n^{(1)}$ of the form

$$\frac{\tilde{\mu}_n^{(1)}}{|\tilde{\mu}_n^{(1)}|} = \sum_{i=1}^{r} \mu_n^{(i)} \ \text{where}$$

$$\mu_n^{(i)} \geq 0 \ \text{and} \ \mu_n^{(i)} \rightarrow \frac{\lambda_i}{\sum_{i=1}^{r} \lambda_i} \mu^{(i)} .$$

598 Therefore we obtain that

$$\mu_n = \sum_{i=1}^{r} |\tilde{\mu}_n^{(1)}| \mu_n^{(i)} + \mu_n^{(r+1)} \ \text{where}$$

$$\mu_n^{(i)} \geq 0, \ |\tilde{\mu}_n^{(1)}| \mu_n^{(i)} \rightarrow \lambda_i \mu^{(i)} \ \text{for all} \ i \in [|1, r|] \ \text{and}$$

$$\mu_n^{(r+1)} \geq 0, \ \mu_n^{(r+1)} \rightarrow \lambda_{r+1} \mu^{(r+1)}$$

599 from which follows the result. $\qquad \square$

600 Let us now consider such decompositions of $(\mu_n)_{n \geq 0}$ and $(\nu_n)_{n \geq 0}$ such that each factor converges
601 toward the target decomposition of μ. Now let us build the following coupling:

$$\tilde{\pi}_n \triangleq \sum_{k=1}^{r-1} \frac{\min(|\mu_n^{(k)}|, |\nu_n^{(k)}|)}{|\mu_n^{(k)}||\nu_n^{(k)}|} \mu_n^{(k)} \otimes \mu_n^{(k)}$$

$$+ \frac{1}{1 - \sum_{k=1}^{r-1} \min(|\mu_n^{(k)}|, |\nu_n^{(k)}|)} \left[|\mu_n| - \sum_{k=1}^{r-1} \frac{\min(|\mu_n^{(k)}|, |\nu_n^{(k)}|)}{|\mu_n^{(k)}|} \mu_n^{(k)} \right] \otimes \left[\nu_n - \sum_{k=1}^{r-1} \frac{\min(|\mu_n^{(k)}|, |\nu_n^{(k)}|)}{|\nu_n^{(k)}|} \nu_n^{(k)} \right]$$

602 with the convention that $\frac{0}{0} = 0$. Now it is easy to check that $\tilde{\pi}_n \in \Pi_r(\mu_n, \nu_n)$, and we have that

$$\text{LOT}_{r,c}(\mu_n, \nu_n) \leq \int_{\mathcal{X}^2} d(x, y) d\tilde{\pi}_n(x, y) \rightarrow \text{LOT}_{r,c}(\mu, \nu)$$

603 and by Prokhorov's theorem and the optimality of the limit of $(\tilde{\pi}_n)_{n \geq 0}$ (up to an extraction) we
604 obtain that $\text{LOT}_{r,c}(\mu_n, \nu_n) \rightarrow \text{LOT}_{r,c}(\mu, \nu)$. $\qquad \square$

605 **B.8 Proof Proposition 7**

Proposition. *Let $r \geq 1$, and let us assume that c is a semimetric of negative type. Then for all
$\mu, \nu \in \mathcal{M}_1^+(\mathcal{X})$, we have that*

$$DLOT_r(\mu, \nu) \geq 0 .$$

606 *In addition, if c has strong negative type then we have also that*

$$\text{DLOT}_{r,c}(\mu, \nu) = 0 \iff \mu = \nu \ and$$

$$\mu_n \rightarrow \mu \iff \text{DLOT}_{r,c}(\mu_n, \mu) \rightarrow 0 .$$

607 *where the convergence of the sequence of probability measures considered is the convergence in law.*

Proof. Let π^* solution of $\mathrm{LOT}_{r,c}(\mu, \nu)$. Then there exists $\lambda^* \in \Delta_r^*$, $(\mu_i^*)_{i=1}^r, (\nu_i^*)_{i=1}^r \in \mathcal{M}_1^+(\mathcal{X})^r$ such that

$$\pi^* = \sum_{i=1}^r \lambda_i^* \mu_i^* \otimes \nu_i^*.$$

608   Note that by definition, we have that

$$\mu = \sum_{i=1}^r \lambda_i^* \mu_i^* \ \text{ and } \ \nu = \sum_{i=1}^r \lambda_i^* \nu_i^*,$$

609   By definition we have also that

$$\mathrm{LOT}_{r,c}(\mu, \mu) \leq \sum_{k=1}^r \lambda_k^* \int_{\mathcal{X}^2} c(x,y) d\mu_k^* \otimes \mu_k^*$$

610   similarly for $\mathrm{LOT}_{r,c}(\nu, \nu)$ we have

$$\mathrm{LOT}_{r,c}(\nu, \nu) \leq \sum_{k=1}^r \lambda_k^* \int_{\mathcal{X}^2} c(x,y) d\nu_k^* \otimes \nu_k^*$$

611   Therefore we have

$$\mathrm{DLOT}_{r,c}(\mu, \nu) \geq \sum_{k=1}^r \lambda_k^* \left(\int_{\mathcal{X}^2} c(x,y) d\mu_k^* \otimes \nu_k^* - \frac{1}{2} \left[\int_{\mathcal{X}^2} c(x,y) d\mu_k^* \otimes \mu_k^* + \int_{\mathcal{X}^2} c(x,y) d\nu_k^* \otimes \nu_k^* \right] \right)$$

$$\geq \sum_{k=1}^r \lambda_k^* \int_{\mathcal{X}^2} -c(x,y) d[\mu_k^* - \nu_k^*] \otimes [\mu_k^* - \nu_k^*]$$

$$\geq \sum_{k=1}^r \frac{\lambda_k^*}{2} D_c(\mu_k^*, \nu_k^*)$$

612   where for any any probability measures α, β on $\mathcal{X}$ we define

$$D_c(\alpha, \beta) \triangleq 2 \int_{\mathcal{X}^2} c(x,y) d\alpha \otimes \beta - \int_{\mathcal{X}^2} c(x,y) d\alpha \otimes \alpha - \int_{\mathcal{X}^2} c(x,y) d\beta \otimes \beta$$

613   However, as c is assumed to have a negative type, we have that

$$D_c(\mu_k^*, \nu_k^*) \geq 0 \ \forall k$$

614   In addition if we assume that c has a strong negative type, then we obtain directly that

$$\mathrm{DLOT}_{r,c}(\mu, \nu) = 0 \implies \mu_k^* = \nu_k^* \ \forall k \ .$$

615   Let us now show that $\mathrm{DLOT}_{r,c}$ metrize the convergence in law. The direct implication is a direct
616   consequence of the Proposition 8. Conversely, if $\mathrm{DLOT}_{r,c}(\mu_n, \mu) \to 0$, then by compacity of $\mathcal{X}$
617   and thanks to the Prokhorov's theorem we can extract a subsequence of $\mu_n \to \mu^*$, and thanks to
618   Proposition 8, we also obtain that $\mathrm{DLOT}_{r,c}(\mu_n, \mu) \to \mathrm{DLOT}_{r,c}(\mu^*, \mu)$. Finally we deduce that
619    $\mathrm{DLOT}_{r,c}(\mu^*, \mu) = 0$ and $\mu^* = \mu$.

620

$\square$

621   ## B.9   Proof Proposition 9

622   **Proposition.** *Let $n \geq k \geq 1$, $\mathbf{X} \triangleq \{x_1, \dots, x_n\} \subset \mathcal{X}$ and $a \in \Delta_n^*$. If c is a semimetric of negative*
623   *type, then by denoting $C = (c(x_i, x_j))_{i,j}$, we have that*

$$\mathrm{LOT}_{k,c}(\mu_{a,\mathbf{X}}, \mu_{a,\mathbf{X}}) = \min_Q \langle C, Q diag(1/Q^T \mathbf{1}_n) Q^T \rangle \ s.t. \ Q \in \mathbb{R}_+^{n \times k} \ , \ Q\mathbf{1}_k = a \ . \tag{16}$$

624   *Proof.* First remarks that one can reformulate the $\mathrm{LOT}_{k,c}$ problem as

$$\mathrm{LOT}_{k,c}(\mu, \mu) \triangleq \min_{g \in \Delta_k^*} \min_{(\mathbf{x}, \mathbf{y}) \in K_{a,g}^2} \sum_{i=1}^k \frac{\mathbf{x}_i^T C \mathbf{y}_i}{g_i}$$

where

$$K_{a,g} \triangleq \{\mathbf{x} \in \mathbb{R}^{nk} \text{ s.t. } A\mathbf{x} = [a,g]^T, \ \mathbf{x} \geq 0\}$$

$$A \triangleq \begin{pmatrix} \mathbf{1}_n^T \otimes \mathbb{I}_k \\ \mathbb{I}_n^T \otimes \mathbf{1}_k \end{pmatrix} \text{ and}$$

$$\mathbf{x}_i \triangleq [x_{(i-1)\times n+1}, \ldots, x_{i\times n}]^T, \ \mathbf{y}_i \triangleq [y_{(i-1)\times n+1}, \ldots, y_{i\times n}]^T \text{ for all } i \in [|1,k|]$$

Indeed the above optimization problem is just a reformulation of $\mathrm{LOT}_{k,c}(\mu,\mu)$ where we have vectorized the couplings in a column-wise order. Let us now show the following lemma from which the result will follow.

Lemma 2. *Under the same assumption of Proposition 9 we have that for all $g \in \Delta_k^*$*

$$\min_{(\mathbf{x},\mathbf{y}) \in K_{a,g}^2} \sum_{i=1}^k \frac{\mathbf{x}_i^T C \mathbf{y}_i}{g_i} = \min_{\mathbf{x} \in K_{a,g}} \sum_{i=1}^k \frac{\mathbf{x}_i^T C \mathbf{x}_i}{g_i}$$

Proof. Let $(\mathbf{x}^*, \mathbf{y}^*)$ solution of the LHS optimization problem. Then we have that

$$\sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{x}_i^*}{g_i} \geq \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i}$$

$$\sum_{i=1}^k \frac{(\mathbf{y}_i^*)^T C \mathbf{y}_i^*}{g_i} \geq \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i}$$

Therefore we obtain that

$$0 \leq \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{x}_i^*}{g_i} - \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i} = \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C(\mathbf{x}_i^* - \mathbf{y}_i^*)}{g_i}$$

$$0 \leq \sum_{i=1}^k \frac{(\mathbf{y}_i^*)^T C \mathbf{y}_i^*}{g_i} - \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i} = \sum_{i=1}^k \frac{(\mathbf{y}_i^* - \mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i}$$

Then by symmetry of C, we obtain by adding the two terms that

$$\sum_{i=1}^k \frac{(\mathbf{x}_i^* - \mathbf{y}_i^*)^T C(\mathbf{x}_i^* - \mathbf{y}_i^*)}{g_i} \geq 0$$

However, thanks to the linear constraints, we have that for all $i \in [|1,k|]$,

$$\sum_{q=0}^{n-1} x_{(i-1)\times n+1+q}^* = \sum_{q=0}^{n-1} y_{(i-1)\times n+1+q}^* = g_i$$

Therefore $(\mathbf{x}_i^* - \mathbf{y}_i^*)^T \mathbf{1}_n = 0$ and thanks to the negativity of the cost function c we obtain that

$$(\mathbf{x}_i^* - \mathbf{y}_i^*)^T C(\mathbf{x}_i^* - \mathbf{y}_i^*) \leq 0$$

Therefore we have that

$$(\mathbf{x}_i - \mathbf{y}_i)^T C(\mathbf{x}_i - \mathbf{y}_i) = 0$$

from which follows that

$$\sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{x}_i^*}{g_i} = \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{y}_i^*}{g_i} = \sum_{i=1}^k \frac{(\mathbf{y}_i^*)^T C \mathbf{y}_i^*}{g_i}$$

and the result follows. □

As the above result holds for any $g \in \Delta_k^*$, we obtain that

$$\mathrm{LOT}_{k,c}(\mu,\mu) = \min_{g \in \Delta_k^*} \min_{\mathbf{x} \in K_{a,g}} \sum_{i=1}^k \frac{(\mathbf{x}_i^*)^T C \mathbf{x}_i^*}{g_i}$$

Then by formulating back this problem in term of matrices, we obtain that

$$\mathrm{LOT}_{k,c}(\mu,\mu) = \min_{g \in \Delta_k^*} \min_{Q \in \Pi_{a,g}} \langle C, Q\mathrm{diag}(1/g)Q^T \rangle$$

from which the result follows. □

C Additional Experiments

C.1 Comparison of the γ schedules

Figure 5: In this experiment, we compare two strategies for the choice of the step-size in the MD scheme proposed by Scetbon et al. [2021] on two different problems. More precisely, we compare the constant γ schedule with the proposed adaptive one and compare them when the distributions are sampled from either uniform distributions (*left*) or mixtures of anisotropic Gaussians (*right*). We show that the range of admissible γ when considering a constant schedule varies from one problem to another. Indeed, in the right plot, we observe that the algorithm converges only when $\gamma \leq 1$, while in the left plot, the algorithm manages to converge for $\gamma \leq 100$. We also observe that our adaptive strategy allows to have a consistent choice of admissible values for γ whatever the problem considered. It is worth noticing that whatever the γ chosen, the algorithm converges towards the same value, however the larger γ is chosen in its admissible range, the faster the algorithm converges.

C.2 Gradient Flows between two Moons

Figure 6: We compare the gradient flows $(\mu_t)_{t \geq 0}$ (in red) starting from a moon shape distribution, μ_0, to another moon shape distribution (in blue), ν, in 2D when minimizing either $L(\mu) \triangleq \text{DLOT}_{r,c}(\mu, \nu)$ or $L(\mu) \triangleq \text{LOT}_{r,c}(\mu, \nu)$. The ground cost is the squared Euclidean distance and we fix $r = 100$. We consider 1000 samples from each distribution and and we plot the evolution of the probability measure obtained along the iterations of a gradient descent scheme. We also display in green the vector field in the descent direction. We show that the debiased version allows to recover the target distribution while $\text{LOT}_{r,c}$ is learning a biased version with a low-rank structure.