Risk-Seeking Reinforcement Learning via Multi-Timescale EVaR Optimization

Abstract

Risk sensitivity is pivotal in shaping agents' behavior when navigating uncertainty and diverging from risk-neutral scenarios. Risk measures such as Value at Risk (`VaR`) and Conditional Value at Risk (`CVaR`) have shown promising results in risk-sensitive reinforcement learning. In this paper, we study the incorporation of a relatively new coherent risk measure, Entropic Value at Risk (`EVaR`), as the objective the agent seeks to optimize. We propose a multi-timescale stochastic approximation algorithm to seek the optimal parameterized `EVaR` policy. Our algorithm facilitates effective exploration of the policy space and robust approximation of the gradient, leading to the optimization of the `EVaR` objective. We analyze the asymptotic behavior of our proposed algorithm and rigorously evaluate it across various discrete and continuous benchmark environments. The results highlight that the `EVaR` policy achieves higher cumulative returns and corroborates that `EVaR` is indeed a competitive risk seeking objective for RL.

1 Introduction

Sequential decision making modelled as a Markov decision process (MDP) (Puterman, 2014) is the underlying formulation of Reinforcement Learning (RL) (Sutton & Barto, 1998). The primary objective in RL is typically to maximize the expected cumulative return, which may be discounted, undiscounted, or averaged (Puterman, 2014), depending on whether the horizon is finite or infinite. Optimizing for expected return has demonstrated remarkable success in structured and less volatile applications like atari games, board games, scientific experiments (Chen et al., 2022) and other regularized simulated environments (Shao et al., 2019; Sethy et al., 2015; Silver et al., 2016) due to stable and predictable environment dynamics. This approach often struggles in environments characterized by high uncertainty and variability in returns, such as safety-critical systems (Zhang et al., 2020; Martín H. & de Lope, 2009), finance (Filos, 2019; Cao et al., 2021), navigation (Wu et al., 2024), industrial automation, healthcare (Wang et al., 2023) and robotics (Kober et al., 2013; Gu et al., 2023), where robustness and adaptability are essential. In such high-stakes scenarios, accounting for risk becomes crucial.

To handle these challenges, reinforcement learning must go beyond simply maximizing the expected return. Instead, it should incorporate risk-awareness by modifying the objective to account for return variability, giving rise to risk-sensitive RL (García & Fernández, 2015; Howard & Matheson, 1972b). The standard expected cumulative reward criterion does not inherently avoid rare but severe negative outcomes, nor does it consider the impact of large positive rewards. Consider two policies π_1 and π_2, where $\mathbb{P}_{\pi_1}(\texttt{Reward} = -100) = 0.5$ and $\mathbb{P}_{\pi_1}(\texttt{Reward} = +100) = 0.5$. Also, $\mathbb{P}_{\pi_2}(\texttt{Reward} = -1) = 1.0$. Although π_1 has a higher expected return, it exhibits extreme volatility, leading to highly variable outcomes which can be categorized as risky. In contrast, π_2 provides a stable and predictable outcome. The expected cumulative reward criterion would favor π_1, despite its high risk, highlighting the need for alternative evaluation metrics that account for risk. Classical RL objectives based on expected cumulative reward can be classified as risk-neutral, meaning they do not explicitly consider the uncertainties associated with actions. In contrast, risk-sensitive decision-making can be divided into two approaches: *risk-averse RL* – the agent prioritizes stability, favoring policies with low reward variability, which is crucial in safety-critical applications, and *risk-seeking RL* – the agent pursues higher mean returns, even at the cost of greater potential losses, resembling human decision-making patterns

in portfolio management (Gollier, 2001) and super-human racing AI (Wurman et al., 2022; Kaufmann et al., 2023), as explained by cumulative prospect theory (Tversky & Kahneman, 1992). By integrating risk-sensitive objectives into RL, agents can achieve a balance between maximizing expected rewards and managing uncertainty. This leads to more robust and adaptive behaviors in dynamic and high-risk environments.

Risk Measures: In this paper, we assume the existence of a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, where Ω represents the sample space, $\mathcal{F}$ a σ-field over Ω, and $\mathbb{P}$ signifies a probability measure over $\mathcal{F}$. In this paper, we consider random variables defined over this probability space. Several risk measures are commonly used in decision-making, each with different assumptions and applications. These include the Markowitz Mean-Variance risk measure (Markowitz & Todd, 2000), which assumes that returns follow a normal distribution and balances expected return against variance. Another approach is the Wang transform function (Wang, 1996), which distorts the cumulative distribution function to model risk aversion or risk-seeking behavior. More widely used in risk-sensitive optimization are Value at Risk (VaR) (Rockafellar et al., 2000) and Conditional Value at Risk (CVaR) (Rockafellar et al., 2000; Rockafellar & Uryasev, 2002). $\mathtt{CVaR}_\alpha$ and $\mathtt{VaR}_\alpha$ of a random variable $\mathbf{X}$ at confidence level $\alpha \in [0, 1]$ are defined as follows:

$$\mathtt{CVaR}_\alpha(\mathbf{X}) = \mathbb{E}[\mathbf{X} \mid \mathbf{X} \geqslant \mathtt{VaR}_\alpha(\mathbf{X})], \text{ where} \tag{1}$$

$$\mathtt{VaR}_\alpha(\mathbf{X}) = \sup\{\beta \in \mathbb{R} \mid \mathbb{P}(\mathbf{X} \geqslant \beta) \geqslant \alpha\}. \tag{2}$$

While $\mathtt{VaR}_\alpha$ identifies a gain threshold, $\mathtt{CVaR}_\alpha$ provides a more comprehensive assessment by evaluating the expected gains in the best-case scenarios. This makes it particularly useful in risk-sensitive decision-making, financial risk management, and safety-critical applications. Acceptance of the risk measures depend on the stability of their estimation procedures and the simplicity of optimization. The incorporation of these risk measures in policy optimization can be achieved either by modifying the objective function—where cumulative rewards are transformed non-linearly using a risk measure, commonly, $\mathtt{CVaR}_\alpha$ (Chow et al., 2015; Tamar et al., 2015; Kashima, 2007; Keramati et al., 2020; Singh et al., 2020)—or by considering the risk measure as a constraint in the optimization setting, also commonly $\mathtt{CVaR}_\alpha$ (Prashanth, 2014; Chow & Ghavamzadeh, 2014; Zhang et al., 2024; Ahmadi et al., 2021).$\mathtt{CVaR}_\alpha$ is more widely accepted due to its coherent nature.

Entropic value at risk ($\mathtt{EVaR}$) (Ahmadi-Javid, 2012) is a fairly new risk measure based on the exponential moment of gains, derived from the Chernoff bound, that provides a convex and coherent lower bound on gains, making it particularly useful for managing tail risk. It is defined as follows: For a random return $\mathbf{X} \in \mathbb{R}$, Entropic Value-at-Risk ($\mathtt{EVaR}$ - right tail) is defined with a confidence parameter $\alpha \in [0, 1)$, as

$$\mathtt{EVaR}_\alpha[\mathbf{X}] = \inf_{\beta > 0} \left(\frac{1}{\beta} \log \frac{\mathbb{E}[e^{\beta \mathbf{X}}]}{\alpha} \right), \tag{3}$$

For the validity of the above definition, we assume that the moment-generating function $M_{\mathbf{X}}(\beta) = \mathbb{E}[e^{\beta \mathbf{X}}]$ exists for all $\beta \geqslant 0$. $\mathtt{EVaR}$ minimizes the worst-case bound on the right tail by optimizing over exponential moment bounds. It acts as a dual to the Legendre-Fenchel transform of the cumulant function $\log \mathbb{E}[e^{\beta \mathbf{X}}]$, ensuring a convex upper bound on extreme right-tail values. From large deviation theory, the cumulant function has a dual formulation: $\log \mathbb{E}[e^{\beta \mathbf{X}}] = \sup_{\xi \ll \mathbb{P}} \{\mathbb{E}_\xi[\mathbf{X}] - \mathtt{KL}(\xi \| \mathbb{P})\}$. Hence, we obtain the dual representation of $\mathtt{EVaR}_\alpha$ as follows:

$$\mathtt{EVaR}_\alpha[\mathbf{X}] = \inf_{\beta > 0} \sup_{\xi \ll P} \left(\frac{1}{\beta} \left(\mathbb{E}_\xi[\mathbf{X}] - \mathtt{KL}(\xi \| \mathbb{P}) - \log \alpha \right) \right)$$

$$= \sup_{\xi \ll P} \inf_{\beta > 0} \left(\mathbb{E}_\xi[\mathbf{X}] - \frac{1}{\beta}\mathtt{KL}(\xi \| \mathbb{P}) - \frac{1}{\beta} \log \alpha \right)$$

$$= \sup_{\xi \ll \mathbb{P}} \mathbb{E}_\xi[\mathbf{X}] \text{ with } \mathtt{KL}(\xi \| \mathbb{P}) \leqslant \log(1/\alpha).$$

From the above characterization, it easily follows that, $\mathbb{E}(\mathbf{X}) \leqslant \mathtt{EVaR}_\alpha(\mathbf{X}) \leqslant \mathrm{esssup}(\mathbf{X})$, where $\mathrm{esssup}(\mathbf{X}) = \inf\{x \in \mathbb{R} : \mathbb{P}(\mathbf{X} \leqslant x) = 1\}$ is the essential sup of $\mathbf{X}$. Note that $\mathtt{CVaR}_\alpha(\mathbf{X}) = \int \mathbb{X} d\mathbb{P}_{\mathtt{CVaR}}$, where the probability measure $\mathbb{P}_{\mathtt{CVaR}}(A) = \frac{1}{\alpha}\mathbb{P}(\{\mathbf{X} \geqslant \mathtt{VaR}_\alpha(\mathbf{X})\} \cap A))$, for Borel set A. Furthermore, $\mathtt{KL}(\mathbb{P}_{\mathtt{CVaR}}, \mathbb{P}) = \int \log \frac{1}{\alpha} d\mathbb{P}_{\mathtt{CVaR}} = \int \frac{\alpha}{\alpha} \log \frac{1}{\alpha} d\mathbb{P} = \log \frac{1}{\alpha}$. Also, $\mathbb{P}_{\mathtt{CVaR}} \ll \mathbb{P}$. This leads to the well-known ordering of risk measures:

$$\mathtt{VaR}_\alpha(\mathbf{X}) \leqslant \mathtt{CVaR}_\alpha(\mathbf{X}) \leqslant \mathtt{EVaR}_\alpha(\mathbf{X}).$$

This ordering indicates that `VaR` provides the least conservative risk assessment, while `EVaR` offers the most robust and conservative measure, with `CVaR` serving as an intermediate risk measure.

For continuous $r.v.s.$, `VaR`$_\alpha$ estimate follows the asymptotic normality $\mathcal{N}\left(\mathtt{VaR}_\alpha, \frac{\alpha(1-\alpha)}{nf(\mathtt{VaR}_\alpha)^2}\right)$ (Serfling, 2009), where f is the PDF. This implies that as one approaches extreme tails (α small), the variance of the estimator becomes prohibitively large, making both `VaR`$_\alpha$ and `CVaR`$_\alpha$ difficult to estimate accurately. `EVaR` is a coherent risk measure as it satisfies subadditivity, positive homogeneity, monotonicity, and translation invariance. Its convex formulation ensures both robustness and tractability, making `EVaR` more suitable for optimal decision-making in risk-sensitive settings. `EVaR`$_\alpha$ avoids explicit tail estimation by leveraging the moment-generating function for its computation.

Figure 1: Comparison of `VaR`$_\alpha$, `CVaR`$_\alpha$, and `EVaR`$_\alpha$ for heavy-tailed (Pareto) and light-tailed (Normal) distributions, showing `EVaR`'s stronger tail-risk sensitivity.

Problem Statement: In this paper, our objective is to seek optimal decision making under uncertainity which is modeled as a Markov Decision Process (MDP). An MDP is defined by the tuple $(\mathcal{S}, \mathcal{A}, R, P, \gamma)$, where $\mathcal{S} = [n]$ and $\mathcal{A} = [m]$ are the state and action spaces, respectively. $R : \mathcal{S} \times \mathcal{A} \times \mathcal{S} \to \mathbb{R}$ is the reward function, where $R(s, a, s')$ represents the reward received for each state transition from $s \xrightarrow{a} s'$ taking action a. The transition probabilities are $\mathbb{P} : \mathcal{S} \times \mathcal{A} \to \Delta^S$, where Δ^S is the probability simplex in $\mathbb{R}^S$ and for a particular state-action pair, $\mathbb{P}(\cdot|s, a)$ is the transition probability, $\mathbb{P}_0(\cdot)$ is the initial state distribution, and $\gamma \in [0, 1)$ is the discount factor. For each state s, the set $\mathcal{A}(s)$ gives all available actions. A stationary policy $\pi(\cdot|s)$ is a probability distribution over actions that depends on the current state s. Here we consider parameterized stochastic policies, which are parameterized by a p-dimensional vector θ, which means that the policy space can be written as $\Pi_\Theta = \{\pi_\theta(\cdot|s), s \in \mathcal{S}, \theta \in \Theta \subseteq \mathbb{R}^p\}$. In this paper, we consider the following risk-sensitive control problem:

$$\theta^* = \arg\max_{\theta \in \Theta} J_{\mathtt{EVaR}}(\theta) = \mathtt{EVaR}_\alpha[R(\tau)]$$

$$= \arg\max_{\theta \in \Theta} \inf_{\beta > 0} \frac{1}{\beta} \log \frac{\mathbb{E}_{\tau \sim \pi_\theta}\left[e^{\beta R(\tau)}\right]}{\alpha}, \tag{4}$$

where $R(\tau) = \sum_{t=0}^T \gamma^t R(\mathbf{s}_t, \mathbf{a}_t, \mathbf{s}_{t+1})$, with $\mathbf{s}_0 \sim \mathbb{P}_0, \mathbf{a}_t \sim \pi_\theta(\cdot|\mathbf{s}_t), \mathbf{s}_{t+1} \sim \mathbb{P}(\cdot|\mathbf{s}_t, \mathbf{a}_t)$, and $T \in \mathbb{N} \cup \{\infty\}$. Note that $J_{\mathtt{EVaR}}(\theta)$ exists for $\forall\theta$, since $R(\cdot)$ is bounded.

The entropic value-at-risk is inherently connected to *exponential tilting*, a technique that modifies the probability distribution to emphasize higher-risk, high-reward outcomes. To illustrate this connection,

(a)   (b)

Figure 2: **(a)** Plot shows exponentially tilted distributions for two RL policies with moderate risk sensitivity. Policy A remains stable, while Policy B shifts significantly, emphasizing its long tail. Tilting reweights probability mass toward high-reward outcomes, enhancing risk-sensitive learning. **(b)** Low β encourages exploration with stable learning, moderate β balances exploration and risk sensitivity, while high β prioritizes high-reward strategies but introduces higher variability.

consider the exponentially tilted probability measure $\mathbb{Q}_\beta$, defined as:

$$\frac{d\mathbb{Q}_\beta}{d\mathbb{P}_{\pi_\theta}} = \frac{e^{\beta r}}{\mathbb{E}_{\tau \sim \pi_\theta}[e^{\beta R(\tau)}]}, \tag{5}$$

This transformation effectively reweights the original probability distribution $\mathbb{P}_{\pi_\theta}$, increasing the probability of trajectories with higher cumulative rewards $R(\tau)$. As a result, the modified measure $\mathbb{Q}_\beta$ gives greater importance to *risk-sensitive outcomes*, ensuring that extreme rewards are more heavily considered. Using this tilted distribution, we obtain the following relationship between the KL divergence and the EVaR objective:

$$\text{KL}(\mathbb{Q}_\beta \| \mathbb{P}_{\pi_\theta}) = \beta \mathbb{E}_{\mathbb{Q}_\beta}[\mathbf{R}] - \log \mathbb{E}_{\tau \sim \pi_\theta}[e^{\beta R(\tau)}] \text{ and}$$

$$J_{\text{EVaR}}(\theta) = \inf_{\beta > 0} \left(\mathbb{E}_{\mathbb{Q}_\beta}[\mathbf{R}] - \frac{1}{\beta}\text{KL}(\mathbb{Q}_\beta \| \mathbb{P}_{\pi_\theta}) - \frac{1}{\beta}\log \alpha \right)$$

$$= \sup_{\mathbb{Q} \in \mathcal{M}_\theta} \left\{ \mathbb{E}_{\mathbb{Q}}[\mathbf{R}] \mid \text{KL}(\mathbb{Q} \| \mathbb{P}_{\pi_\theta}) \leqslant \log \frac{1}{\alpha} \right\},$$

$$\text{where } \mathcal{M}_\theta = \left\{ \mathbb{Q}_\beta \mid \exists \beta > 0, \ \frac{d\mathbb{Q}_\beta}{d\mathbb{P}_{\pi_\theta}} = \frac{e^{\beta r}}{\mathbb{E}_{\tau \sim \pi_\theta}[e^{\beta R(\tau)}]} \right\}.$$

This formulation highlights that EVaR selects the extreme expectation over a set of tilted distributions, each constrained by a KL-divergence bound. In other words, it finds the most extreme risk-sensitive expectation while ensuring the alternative probability distribution remains within a reasonable divergence from the original measure, thereby maintaining robustness in decision-making under uncertainty. The EVaR objective does not merely maximize expected returns but incorporates a risk-seeking adjustment that prioritizes high-reward yet riskier trajectories. Further, the KL-divergence constraint prevents excessive deviation from the original probability distribution, ensuring a balanced trade-off between exploration and risk seeking.

Related Literature: The field of risk-sensitive control and reinforcement learning has been well studied, beginning with the seminal work of (Howard & Matheson, 1972a) which introduced the application of an exponential utility function to rewards. A substantial body of research has concentrated on risk-sensitive control in continuous-time, finite-horizon settings, particularly for problems with known transition kernels (Fleming & McEneaney, 1995; Whittle, 1990; Coraluppi & Marcus, 1999; Koenig & Simmons, 1994). Early developments extended optimal control techniques to reinforcement learning (Littman & Szepesvári, 1996; Borkar, 2001; 2002; 2010; Mihatsch & Neuneier, 2002; Heger, 1994). The use of exponential ergodic

performance in discrete time control (Di Masi & Stettner, 1999; 2007) was further generalized to address risk-sensitive average cost criteria (Cavazos-Cadena & Hernandez-Hernandez, 2011). In reinforcement learning, risk-sensitive methods have been adapted for model-free settings with unknown transition dynamics, employing techniques such as relative entropy stochastic search and Q-learning (Borkar & Meyn, 2002; Osogami, 2012). Recent advances in risk-sensitive reinforcement learning integrate conditional value-at-risk (CVaR) objectives for robust policy optimization (Chow et al., 2015; Chow & Ghavamzadeh, 2014), quantile temporal-difference learning to capture return distributions more accurately (Rowland et al., 2024), leverage sample-based dynamic programming techniques that augment Bayes-adaptive MDPs with CVaR constraints to derive risk-averse policies (Rigter et al., 2021), and incorporate coherent risk measures alongside non-linear function approximation (Lam et al., 2022). EVaR policy optimization in RL is a relatively new and less explored area. (Ni & Lai, 2022) proposed a trajectory-based policy gradient method to optimize EVaR-induced risk-sensitive criteria, (Dixit et al., 2021) developed nested EVaR-constrained models, and (Hau et al., 2023) introduced a dynamic programming approach for EVaR objectives by formulating EVaR-based Bellman equations under known transition dynamics.

Our Contribution: In this paper, we provide an online multi time scale stochastic approximation algorithm to estimate the `EVaR` of the reward distribution and also seek the optimal `EVaR` policy in the context of model-free risk seeking reinforcement learning setting.

2 Proposed Method

In this section, we recast the control objective (4) into two tightly coupled sub-problems. *Prediction* demands, for any fixed policy parameter θ, an online, sample-efficient estimate of the entropic value-at-risk $J_{\texttt{EVaR}}(\theta)$ and *optimisation* then needs a dependable ascent direction built from those estimates to steer θ towards an `EVaR`-optimal policy. We tackle both challenges simultaneously within a multi-timescale stochastic-approximation framework: a fast inner loop continually callibrates the `EVaR` estimate, while a slower outer loop performs simultaneous-perturbation-based gradient ascent, which seeks the risk-seeking solution.

2.1 `EVaR` Estimation

In this section, we propose an online, multi-timescale approach to estimate `EVaR`. From Eq.(4) we have

$$J_{\texttt{EVaR}}(\theta) = \inf_{\beta > 0} \frac{1}{\beta} \big(\log \mathbb{E}_{\tau \sim \pi_\theta} \left[e^{\beta \cdot R(\tau)} \right] - \log \alpha \big) \tag{6}$$

Let $H(\beta) = \beta^{-1} \big(\log \mathbb{E}_{\tau \sim \pi_\theta} \left[e^{\beta \cdot R(\tau)} \right] - \log \alpha \big)$, where $\beta > 0$. We establish the following result for H.

Proposition 1. *The function $G(\beta) = H\left(\frac{1}{\beta}\right)$ is convex in β over $\beta > 0$.*

Proof. Given, $H\left(\frac{1}{\beta}\right) = \beta \left(\log \mathbb{E} \left[e^{\frac{R(\tau)}{\beta}} \right] - \log \alpha \right)$. To prove convexity, for any $\lambda \in [0, 1]$ and $\beta_1, \beta_2 > 0$, let $\beta = \lambda \beta_1 + (1 - \lambda)\beta_2$. Define normalized weights:

$$\mu = \frac{\lambda \beta_1}{\beta}, \quad \nu = \frac{(1 - \lambda)\beta_2}{\beta}, \quad \mu + \nu = 1.$$

By Hölder's inequality:

$$\mathbb{E}\left[e^{\frac{R(\tau)}{\beta}} \right] \leqslant \mathbb{E}\left[e^{\frac{R(\tau)}{\beta_1}} \right]^\mu \mathbb{E}\left[e^{\frac{R(\tau)}{\beta_2}} \right]^\nu.$$

Taking logarithms:

$$\log \mathbb{E}\left[e^{\frac{R(\tau)}{\beta}} \right] \leqslant \mu \log \mathbb{E}\left[e^{\frac{R(\tau)}{\beta_1}} \right] + \nu \log \mathbb{E}\left[e^{\frac{R(\tau)}{\beta_2}} \right].$$

Multiply by β and subtract $\beta \log \alpha$:

$$H\left(\frac{1}{\beta}\right) \leqslant \lambda H\left(\frac{1}{\beta_1}\right) + (1 - \lambda)H\left(\frac{1}{\beta_2}\right).$$

Thus, $H\left(\frac{1}{\beta}\right)$ is convex in β. $\qquad\square$

(a) $H(\beta)$ *vs.* β for different confidence levels α. (b) $G(x) = H(1/x)$ *vs.* x for different confidence levels α.

Figure 3: Dependence of the `EVaR` curves on the confidence level α for a truncated Pareto reward distribution (`shape`= 3, `scale`= 1). **(a)** The function $H(\beta)$ diverges more steeply as $\beta \to 0$ for lower α, indicating stronger tail-risk sensitivity. **(b)** The convex transformation $G(x) = H(1/x)$ attains its minimizer at larger x as α decreases, reflecting a more conservative risk-seeking nature.

The original function $H(\beta)$ exhibits mixed curvature due to the β-scaling in the denominator and the exponential term. However, the transformation $\beta \to \frac{1}{\beta}$, reparameterizes the function, flipping the curvature to enforce convexity. This property is critical because it simplifies the optimization landscape, making it much easier and more reliable to find the unique global optimum.

Corollary 1.

$$\min_{\beta > 0} H(\beta) = \min_{x > 0} G(x),$$

Moreover, the minimizers satisfy $\beta^* = 1/x^*$.

Proof. Define the change of variable $x = \frac{1}{\beta}$. Then $G(x) = H(\frac{1}{x})$. Let β^* be a global minimizer of $H(\beta)$ and $x^* = 1/\beta^*$. Then $G(x^*) = H(\frac{1}{x^*}) = H(\beta^*)$. Since, β^* minimizes H, we have for any $x > 0$, $H(\beta^*) \leqslant H(\frac{1}{x}) = G(x)$. Hence, $G(x^*) = H(\beta^*) \leqslant G(x), \quad \forall\, x > 0$. Thus x^* is a global minimizer of G, and

$$\min_{x > 0} G(x) \leqslant G(x^*) = H(\beta^*) = \min_{\beta > 0} H(\beta).$$

Conversely, suppose x^* minimizes $G(x)$. Define $\beta^* = 1/x^*$. By an identical argument,

$$\min_{\beta > 0} H(\beta) \leqslant H(\beta^*) = G(x^*) = \min_{x > 0} G(x).$$

Putting these two inequalities together,

$$\min_{\beta > 0} H(\beta) = \min_{x > 0} G(x).$$

Furthermore, the minimizers match by $\beta^* = 1/x^*$. □

To establish the convergence and stability of our approach, we require certain regularity conditions on the risk-sensitive objective and the policy-induced reward distribution. These conditions ensure that the EVaR optimization problem remains well-posed and that the gradient estimates concentrate sufficiently around their expected values.

Assumption 1. *The variable $x = 1/\beta$ is restricted to a compact set $I = [x_{\min}, x_{\max}]$ with $0 < x_{\min} < x_{\max} < \infty$.*

Assumption 2. *There exists $\sigma > 0$ such that for all $x \in I$,*

$$Var_{p_x}(R) = \mathbb{E}_{p_x}[R^2] - (\mathbb{E}_{p_x}[R])^2 \geqslant \sigma.$$

This assumption ensures that the reward distribution under the exponentially tilted measure p_x always retains a minimum level of variability. Specifically, it guarantees that the variance of rewards does not collapse to zero for any risk sensitivity parameter x. This is critical because the policy search requires sufficient variability in rewards to effectively explore the policy space and avoid degenerate solutions.

Lemma 1. *Under Assumptions 1 and 2:*

1. *The variance $\mathbf{Var}_{p_x}(R(\tau))$ is continuous in x over $I = [x_{\min}, x_{\max}]$.*

2. *$\exists \bar{\sigma} > 0$ such that $\mathbf{Var}_{p_x}(R(\tau)) \geqslant \bar{\sigma}$, $\forall x \in I$.*

Proof. Let $Z(x) = \mathbb{E}_\tau[e^{R(\tau)/x}]$. The tilted expectation $\mathbb{E}_{p_x}[R]$ and $\mathbb{E}_{p_x}[R^2]$ are given by:

$$\mathbb{E}_{p_x}[R] = \frac{\mathbb{E}_\tau[R(\tau)e^{\frac{R(\tau)}{x}}]}{Z(x)}, \ \mathbb{E}_{p_x}[R^2] = \frac{\mathbb{E}_\tau[R(\tau)^2 e^{\frac{R(\tau)}{x}}]}{Z(x)}.$$

By ($|R(\tau)| \leqslant \frac{R_\infty}{1-\gamma}$,) and Assumption 1 ($x \in I$), the terms $Re^{R/x}$ and $R(\tau)^2 e^{R(\tau)/x}$ are bounded. By the Dominated Convergence Theorem (DCT), $\mathbb{E}_\tau[R(\tau)e^{R(\tau)/x}]$, $\mathbb{E}_\tau[R(\tau)^2 e^{R(\tau)/x}]$, and $Z(x)$ are continuous in x. Since $Z(x) \geqslant e^{\frac{-R_\infty}{(1-\gamma)x_{\max}}} > 0$, the ratios $\mathbb{E}_{p_x}[R]$ and $\mathbb{E}_{p_x}[R^2]$ are continuous. Thus, $\mathbf{Var}_{p_x}(R) = \mathbb{E}_{p_x}[R^2] - (\mathbb{E}_{p_x}[R])^2$ is continuous. By Assumption 2, $\mathbf{Var}_{p_x}(R) \geqslant \sigma > 0$ for all $x \in I$. Continuity of $\mathbf{Var}_{p_x}(R)$ and compactness of I imply $\mathbf{Var}_{p_x}(R)$ attains its minimum on I. Let $\bar{\sigma} = \min_{x \in I} \mathbf{Var}_{p_x}(R)$. By Assumption 2, $\bar{\sigma} > 0$. $\qquad \square$

Theorem 1. *Under Assumptions 1 and 2, the function*

$$G(x) = x \left[\log \mathbb{E}\left(e^{R(\tau)/x} \right) - \log \alpha \right]$$

is m-strongly convex on $I = [x_{\min}, x_{\max}]$ with modulus $m = \frac{\bar{\sigma}}{x_{\max}^3}$.

Proof. One can easily find the second-order derivative of $G(x)$ as follows:

$$G''(x) = \frac{\mathbb{E}_{p_x}[R^2] - (\mathbb{E}_{p_x}[R])^2}{x^3} = \frac{\mathrm{Var}_{p_x}(R)}{x^3}.$$

By Lemma 1, $\mathbf{Var}_{p_x}(R) \geqslant \bar{\sigma} > 0$. Since $x \leqslant x_{\max}$,

$$G''(x) \geqslant \frac{\bar{\sigma}}{x_{\max}^3} = m > 0.$$

Thus, $G(x)$ is strongly convex on I. $\qquad \square$

We now propose a multi-timescale framework to estimate $J_{\texttt{EVaR}}$ by solving the above optimization problem via finding the roots of the derivate G'. For this purpose, we derive a closed-form expression for G with respect to x as follows:

$$\begin{aligned} G'(x) &= \frac{d}{dx} \left[x \left(\log \mathbb{E}_\tau \left[e^{R(\tau)/x} \right] - \log \alpha \right) \right] \qquad (7) \\ &= \log \frac{\mathbb{E}_\tau \left(e^{R(\tau)/x} \right)}{\alpha} - \frac{1}{x} \frac{\mathbb{E}_\tau[R(\tau)\, e^{R(\tau)/x}]}{\mathbb{E}_\tau[e^{R(\tau)/x}]}. \end{aligned}$$

The interchange of $\mathbb{E}_\tau[\cdot]$ and $\frac{d}{dx}$ in the above equality is possible through the bounded convergence theorem. In practice, the exact expectations $\mathbb{E}_\tau[e^{R(\tau)/x}]$ and $\mathbb{E}_\tau[R(\tau)e^{R(\tau)/x}]$ are often intractable to compute directly. To address this, we replace these expectations with online average estimates, computed from observed samples over time. Consequently, we arrive at a two-timescale stochastic approximation algorithm for finding $x^* = \arg\min_{x>0} G(x)$. In this framework, one timescale is used to update the estimate of G', while the other

handles the solution update in the direction of the estimate. This separation enables more efficient and stable convergence of the gradient estimation process.

$$\vartheta_{t+1} = \vartheta_t + \delta_t \left(e^{R(\tau_{t+1})/x_t} - \vartheta_t \right), \qquad \text{where } \tau_{t+1} \sim \pi_\theta \text{ and the step-size } \delta_t \in (0,1) \tag{8}$$
$$\omega_{t+1} = \omega_t + \delta_t e^{R(\tau_{t+1})/x_t} \left(R(\tau_{t+1}) - x_t\omega_t \right),.$$

The above single timescale stochastic recursions estimate the expectations $\mathbb{E}_\tau[e^{R(\tau)/x}]$ and $\mathbb{E}_\tau[R(\tau)e^{R(\tau)/x}]/x\mathbb{E}_\tau[e^{R(\tau)/x}]$ which amount to estimating G'. Now we can apply these estimates to seek x^* the by calibrating the iterates in the negative direction of the derivative estimate as follows:

$$x_{t+1} = x_t - \xi_t \left(\log \frac{\vartheta_t}{\alpha} - \omega_t \right), \text{ where } \xi_t \in (0,1) \text{ is the step-size.} \tag{9}$$

Note that the above recursion is maintained at a slower time scale relative to the recursions Eq.(8). This is required because one needs a good estimate of G' to efficiently calibrate x_t. This develops a bidirectional coupling where the recursion Eq.(9) can be considered quasistatic *w.r.t.* recursion Eq.(8). This can be illustrated as follows:

$$x_{t+1} = x_t - \delta_t \frac{\xi_t}{\delta_t} \left(\log \frac{\vartheta_t}{\alpha} - \omega_t \right) \tag{10}$$

By stacking Eqs.(8), and (10) in vector notation, we obtain the following

$$\begin{bmatrix} \vartheta_{t+1} \\ \omega_{t+1} \\ x_{t+1} \end{bmatrix} = \begin{bmatrix} \vartheta_t \\ \omega_t \\ x_t \end{bmatrix} + \delta_t \begin{bmatrix} e^{R(\tau_{t+1})/x_t} - \vartheta_t \\ e^{R(\tau_{t+1})/x_t} \left(R(\tau_{t+1}) - x_t\omega_t \right) \\ \frac{-\xi_t}{\delta_t} \left(\log \frac{\vartheta_t}{\alpha} - \omega_t \right) \end{bmatrix}$$

If we let $\lim_{t\to\infty} \frac{\xi_t}{\delta_t} \to 0$, then from the above equation, one can find that while x_t is quasi-static, the estimates ϑ_t and ω_t get nearly equilibrated to their mean-field limits $\vartheta^*(x_t)$ and $\omega^*(x_t)$ respectively. This quasi-static equilibrium eliminates bias in gradient estimates.

Assumption 3. *We assume that the step-size schedules $\{\delta_t\}_{t\in\mathbb{N}}$ and $\{\xi_t\}_{t\in\mathbb{N}}$ are real-valued, positive, deterministic and pre-determined sequences and they satisfy*

$$\sum_{t\in\mathbb{N}} \left(\delta_t^2 + \xi_t^2 \right) < \infty, \quad \sum_{t\in\mathbb{N}} \delta_t = \sum_{t\in\mathbb{N}} \xi_t = \infty, \quad \lim_{t\to\infty} \frac{\xi_t}{\delta_t} = 0.$$

Examples of such step sizes can be $\xi_t = \frac{1}{t}, \delta_t = \frac{1}{1+t\log t}$ or $\xi_t = \frac{1}{t^{2/3}}, \delta_i = 1/t$. The above assumptions are required as they are critical technical requirements for ensuring almost sure convergence. The first condition ensures that they decay fast enough. The second condition ensures that updates happen through the entire time continuum. The final condition (time scale separation) ensures that the ratio of the step sizes must approach zero, ensuring the updates for x_t occur on a slower timescale compared to ϑ_t and ω_t.

Theorem 2 (Convergence of J_{EVaR} under static θ). *Given policy π_θ, under Assumptions 1–3, the coupled stochastic recursions equation 8 & equation 9 constitute a stochastic Euler discretization of the ODE system:*

$$\dot{\vartheta} = \mathbb{E}_{\tau\sim\pi_\theta} \left[e^{R(\tau)/x} \right] - \vartheta,$$

$$\dot{\omega} = \frac{\mathbb{E}_{\tau\sim\pi_\theta} \left[R(\tau)e^{R(\tau)/x} \right]}{x\mathbb{E}_{\tau\sim\pi_\theta} \left[e^{R(\tau)/x} \right]} - \omega,$$

$$\dot{x} = -\underbrace{\left(\log \frac{\mathbb{E}_{\tau\sim\pi_\theta} \left[e^{R(\tau)/x} \right]}{\alpha} - \frac{\mathbb{E}_{\tau\sim\pi_\theta} \left[R(\tau)e^{R(\tau)/x} \right]}{x\mathbb{E}_{\tau\sim\pi_\theta} \left[e^{R(\tau)/x} \right]} \right)}_{-G'(x)},$$

Furthermore, the iterates satisfy:

$$(\vartheta_t, \omega_t, x_t) \xrightarrow{\text{a.s.}} (\vartheta^*(\theta), \omega^*(\theta), x^*(\theta)),$$

with equilibrium $(\vartheta^*(\theta), \omega^*(\theta), x^*(\theta))$ characterized by:

$$\vartheta^*(\theta) = \mathbb{E}_{\tau \sim \pi_\theta}\left[e^{R(\tau)/x^*(\theta)}\right], \quad \omega^*(\theta) = \frac{\mathbb{E}_{\tau \sim \pi_\theta}\left[R(\tau)e^{R(\tau)/x^*(\theta)}\right]}{x^*(\theta)\vartheta^*(\theta)} \quad and,$$

$$x^*(\theta) \in \left\{x \in \mathbb{R}^+ \,|\, G'(x) = 0\right\}.$$

Corollary 2. *For a given policy π_θ, as $t \to \infty$, the iterates $\{x_t\}$ satisfy the following*

$$\lim_{t \to \infty} G(x_t) = J_{EVaR}(\theta). \tag{11}$$

Proof. Follows from the continuity of G and Theorem 2. $\qquad\square$

Theorem 3. *Let $\delta_t = \frac{c}{t}, \xi_t = \frac{d}{t^b}, b \in (0.5, 1)$. Then:*

1. ***Convergence of ϑ_t:** For $c > \frac{1}{2}$,*

$$\mathbb{E}\left[|\vartheta_t - \vartheta^*(x)|^2\right] \leqslant \frac{K_1}{t}, \quad where \quad K_1 = C\left(1 + c^2 e^{\frac{2R_\infty}{(1-\gamma)x_{\min}}}\frac{\pi^2}{6}\right).$$

2. ***Convergence of ω_t:** For $c > \frac{1}{2x_{\min}\mathbb{E}\left[e^{R(\tau)/x}\right]}$,*

$$\mathbb{E}\left[|\omega_t - \omega^*(x)|^2\right] \leqslant \frac{K_2}{t}, \quad where \quad K_2 = C\left(1 + c^2 \frac{R_\infty^2 e^{\frac{2R_\infty}{x_{\min}(1-\gamma)}}}{x_{\min}^2(1-\gamma)^2}\frac{\pi^2}{6}\right).$$

3. ***Convergence of x_t:** With $d = \frac{x_{\max}^3(1-b)}{2\bar{\sigma}} \cdot b$,*

$$\mathbb{E}\left[(x_t - x^*)^2\right] \leqslant \frac{K}{t^b}, \quad where \quad x^* = \arg\min_x G(x) \quad and$$

$$K = \frac{x_{\max}^6(1-b)^2}{4\bar{\sigma}^2}\left(2e^{\frac{2R_\infty}{(1-\gamma)x_{\min}}}K_1 + 2K_2 + \left(\frac{2R_\infty}{x_{\min}(1-\gamma)} + |\log\alpha|\right)^2\right).$$

2.2 EVaR Optimization

The gradient estimation of the EVaR objective with respect to policy parameters θ employs the simultaneous perturbation stochastic approximation method (Spall, 1992), a computationally efficient technique for high-dimensional optimization. This approach perturbs all parameters simultaneously using a randomized direction vector $\Delta_t \in \mathbb{R}^p$ circumventing the $O(p)$ computational complexity of finite-difference methods. The gradient estimate is constructed as follows:

$$\widehat{\nabla_\theta J}_{\text{EVaR}}(\theta) = \frac{J_{\text{EVaR}}(\theta + c_t\Delta_t) - J_{\text{EVaR}}(\theta - c_t\Delta_t)}{2c_t\Delta_t}. \tag{12}$$

where $c_t > 0$ with $\lim_{t \to \infty} c_t \downarrow 0$ and $\Delta_t \in \mathbb{R}^p$ with each of the components are $\Delta_{t_i} \overset{\text{iid}}{\sim} \texttt{Bernoulli}(\pm 1)$ *w.p.* 0.5. Also, $\Delta_t^{-1} = [\Delta_{t_1}^{-1}, \Delta_{t_2}^{-1} \dots \Delta_{t_p}^{-1}]^\top$. Since the true J_{EVaR} is not available, we estimate it as follows:

$$\widehat{\nabla_\theta J}_{\text{EVaR}}(\theta) = \frac{J_{\text{EVaR}}(\theta + c_t\Delta_t) - J_{\text{EVaR}}(\theta - c_t\Delta_t)}{2c_t\Delta_t} \approx G(x_t^+) - G(x_t^-)$$

$$= x_t^+\left(\log\frac{\vartheta_t^+}{\alpha}\right) - x_t^-\left(\log\frac{\vartheta_t^-}{\alpha}\right) \tag{13}$$

The algorithm maintains parallel estimators $\{\vartheta_t^+, \omega_t^+, x_t^+\}$ and $\{\vartheta_t^-, \omega_t^-, x_t^-\}$ for the perturbed policies $\theta + c_t\Delta_t$ and $\theta - c_t\Delta_t$, respectively, to prevent cross–contamination of gradient signals. Each estimator executes N_t

inner iterations of the recursions equation 8–equation 9 per outer policy update, so that $\vartheta_t^\pm$ and $x_t^\pm$ approach quasi–stationary values before the gradient approximation 13 is computed.

The random perturbations (with zero mean) can introduce variance in gradient estimates, which gets asymptotically averaged over multiple random perturbations during the stochastic gradient recursion. As the number of perturbations increases, the average gradient estimate approaches the true gradient asymptotically, meaning that with enough samples, the variance in the gradient estimate becomes negligible and the estimate becomes increasingly accurate. Also, this approach is advantageous as it does not require explicit computation of individual partial derivatives. Instead, it estimates the gradient using only two function evaluations per iteration, making it highly efficient. Unlike finite difference methods that perturb each parameter separately, this method perturbs randomnly chosen parameters simultaneously, reducing computational complexity while maintaining a robust gradient estimate. Furthermore, it benefits from asymptotic unbiasedness - as more iterations accumulate, the stochastic noise from perturbations cancels out, ensuring convergence to the true gradient. Finally, the updation of the policy parameter using the above gradient estimate is:

$$\hat{g}_t \leftarrow \frac{G_t^+ - G_t^-}{2\,c_t}\,\Delta_t^{-1} \quad \text{where } G_t^+ \leftarrow x_t^+\,\ln\!\big(\tfrac{\vartheta_t^+}{\alpha}\big), \quad G_t^- \leftarrow x_t^-\,\ln\!\big(\tfrac{\vartheta_t^-}{\alpha}\big)$$

$$\theta_{t+1} \leftarrow \theta_t + a_t\,\hat{g}_t, \tag{14}$$

where $a_t, c_t \in (0,1)$ are learning rate and perturbation parameter respectively. Now this procedure is illustrated in Algorithm 1.

Note that the above updation rule can be rewritten as follows:

$$\theta_{t+1} = \theta_t + a_t\left(b_t + e_t + \varphi_t + \nabla J_{\texttt{EVaR}}(\theta_t)\right), \text{ where } e_t = \widehat{\nabla J}_{\texttt{EVaR}}(\theta_t) - \mathbb{E}\left[\widehat{\nabla J}_{\texttt{EVaR}}(\theta_t) \mid \mathcal{F}_t\right]$$

$$b_t = \mathbb{E}\left[\widehat{\nabla J}_{\texttt{EVaR}}(\theta_t) - \nabla J_{\texttt{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] \tag{15}$$

$$\text{and } \varphi_t = \frac{z(\theta_t^+) - z(\theta_t^-)}{2c_t\Delta_t} \text{ with } z(\theta_t^+) = G(x_t^+) - J_{\texttt{EVaR}}(\theta_t^+) \text{ and}$$

$$z(\theta_t^-) = G(x_t^-) - J_{\texttt{EVaR}}(\theta_t^-).$$

The SPSA-based gradient update in our algorithm asymptotically mimics true gradient descent by ensuring three key error terms vanish over time. Bias (b_t), arising from finite-difference gradient approximations, scales with the perturbation size c_t^2. Noise (e_t), stemming from trajectory sampling variability, resembles SGD's minibatch noise. If step sizes a_t decay sufficiently, these zero-mean fluctuations average out due to the martingale structure, preventing erratic updates. Drift (φ_t), caused by finite inner-loop EVaR estimation, fades by increasing inner-loop iterations N_t ensuring inner approximations align with outer updates. Collectively, these decays—bias, noise, and drift –ensure the update trajectory converges to the true gradient flow of $J_{\texttt{EVaR}}$, guaranteeing eventual convergence to an $\texttt{EVaR}$-optimal policy.

Assumption 4. *For the stepsize sequences, $a_t > 0, c_t > 0, a_t \to 0, c_t \to 0, \sum_{t=0}^{\infty} a_t = \infty$, and $\sum_{t=0}^{\infty} a_t^2 c_t^2 < \infty$.*

The following lemma analyzes the behaviour of the iterates ϑ_t and ω_t, during the quasi-stagnant phase of β_t and θ_t.

We define the filtration $\{\mathcal{F}_t\}_{t\in\mathbb{N}}$, where the σ-field $\mathcal{F}_t := \sigma\left(\theta_i, \Delta_i, \vartheta_i^\pm, \omega_i^\pm, x_i^\pm, 1 \leqslant i \leqslant t\right)$.

Lemma 2. *Let $J_{EVaR}^{(3)}(\theta) \equiv \partial^3 J_{EVaR}/\partial\theta^T\partial\theta^T\partial\theta^T$ exists and $\max_{i_1,i_2,i_3} \sup_\theta \|J_{EVaR_{i_1 i_2 i_3}}^{(3)}(\theta)\|_\infty \leqslant \epsilon$. Then $\forall \theta \in \Theta$*

$$b_t(\theta_t) = \mathbb{E}\left[\widehat{\nabla_\theta J}_{EVaR}(\theta_t) - \nabla J_{EVaR}(\theta_t) \mid \mathcal{F}_t\right] = \mathcal{O}(c_t^2).$$

Theorem 4. *Assume the conditions mentioned in Theorem 3. Further, assume that $\frac{a_t\sqrt{K}}{c_t N_t^{\frac{b}{2}}} < \infty$. Then the iterates $\{\theta_t\}$ generated by Algorithm satisfy the following:*

$$\theta_t \to \mathcal{E} = \{\theta | \nabla J_{EVaR}(\theta) = 0\} \text{ on the event } \{\sup_t \|\theta_t\| < \infty\} \text{ as } tss \to \infty.$$

Algorithm 1 Multi-timescale EVaR optimization

Require: risk level $\alpha \in (0,1)$, initial $\theta_0 \in \mathbb{R}^p$, step-sizes $\{a_t, c_t, \delta_t, \xi_t\}$, inner lengths N_t

1: $\theta_0 \leftarrow \theta_0$, $\quad \vartheta_0^+ \leftarrow 0$, $\omega_0^+ \leftarrow 0$, $x_0^+ \leftarrow 1$, $\quad \vartheta_0^- \leftarrow 0$, $\omega_0^- \leftarrow 0$, $x_0^- \leftarrow 1$

2: **for** $t = 0, \ldots, T-1$ **do**

3: $\quad$ Draw $\Delta_t \in \{\pm 1\}^p$ IID.

4: $\quad \theta_t^+ \leftarrow \theta_t + c_t \Delta_t$, $\quad \theta_t^- \leftarrow \theta_t - c_t \Delta_t$

5: $\quad$ **for** $k = 1, \ldots, N_t$ **do** $\qquad\qquad\qquad\qquad\qquad\qquad$ ▷ EVaR estimation for "+"

6: $\qquad$ Sample trajectory $\tau_{t,k}^+ \sim \pi_{\theta_t^+}$, compute $R_{t,k}^+ = \sum_{u=0}^H \gamma^u r_u$

7: $\qquad \vartheta_t^+ \leftarrow \vartheta_t^+ + \delta_t \left(e^{R_{t,k}^+/x_t^+} - \vartheta_t^+ \right)$

8: $\qquad \omega_t^+ \leftarrow \omega_t^+ + \delta_t \left(R_{t,k}^+ e^{R_{t,k}^+/x_t^+} - x_t^+ \omega_t^+ \right)$

9: $\qquad x_t^+ \leftarrow x_t^+ - \xi_t \left[\ln\left(\frac{\vartheta_t^+}{\alpha} \right) - \omega_t^+ \right]$

10: $\quad$ **end for**

11: $\quad$ **for** $k = 1, \ldots, N_t$ **do** $\qquad\qquad\qquad\qquad\qquad\qquad$ ▷ EVaR estimation for "-"

12: $\qquad$ Sample trajectory $\tau_{t,k}^- \sim \pi_{\theta_t^-}$, compute $R_{t,k}^- = \sum_{u=0}^H \gamma^u r_u$

13: $\qquad \vartheta_t^- \leftarrow \vartheta_t^- + \delta_t \left(e^{R_{t,k}^-/x_t^-} - \vartheta_t^- \right)$

14: $\qquad \omega_t^- \leftarrow \omega_t^- + \delta_t \left(R_{t,k}^- e^{R_{t,k}^-/x_t^-} - x_t^- \omega_t^- \right)$

15: $\qquad x_t^- \leftarrow x_t^- - \xi_t \left[\ln\left(\frac{\vartheta_t^-}{\alpha} \right) - \omega_t^- \right]$

16: $\quad$ **end for**

17: $\quad G_t^+ \leftarrow x_t^+ \ln\left(\frac{\vartheta_t^+}{\alpha} \right)$, $\quad G_t^- \leftarrow x_t^- \ln\left(\frac{\vartheta_t^-}{\alpha} \right)$

18: $\quad \hat{g}_t \leftarrow \dfrac{G_t^+ - G_t^-}{2\,c_t} \Delta_t^{-1}$

19: $\quad \theta_{t+1} \leftarrow \theta_t + a_t \hat{g}_t$

20: **end for**

21: **return** θ_T

Further, if $\mathcal{E}$ is a discrete set, then we have the following.

$$\theta_t \to \{\theta | \nabla J_{EVaR}(\theta) = 0 \text{ and } \nabla^2 J_{EVaR}(\theta) \prec 0\} \text{ on the event } \{\sup_t \|\theta_t\| < \infty\} \text{ as } t \to \infty.$$

In our multi-timescale algorithm, φ_t acts as a bias drift superimposed on the true gradient $\nabla J_{\text{EVaR}}(\theta_t)$. If φ_t does not vanish fast enough—-for example, if c_t decays too quickly or N_t grows too slowly—then this residual bias can dominate the small gradient signal and prevent the iterates from homing in on a stationary point. In Theorem 4 , the condition $a_t \left(c_t N_t^{b/2} \right)^{-1} \longrightarrow 0$ ensures that the product $a_t \mathbb{E}[|\varphi_t|]$ vanishes, so that φ_t only perturbs the trajectory transiently but does not alter the limiting ODE $\dot{\theta} = \nabla J_{\text{EVaR}}(\theta)$, and hence does not affect almost-sure convergence to the EVaR-critical set.

The aforementioned result indicates that the distribution parameters θ_t converge to the local maxima of the objective EVaR$_\alpha$, provided that the iterates θ_t remains bounded which is denoted by the condition $\sup_t \|\theta_t\| < \infty$. This condition is necessary because noise can cause the iterates to gradually drift outward, potentially leading to divergence. This can be achieved by constraining the iterates to remain within a convex compact set, and if they drift beyond its boundary, they can be projected back onto the set. The projected version of the recursion is as follows:

$$\theta_{t+1} = \Pi^\Theta \left[\theta_t + \frac{a_t}{(2c_t \Delta_t)} \left(x_t^+ \left(\log \frac{\vartheta_t^+}{\alpha} \right) - x_t^- \left(\log \frac{\vartheta_t^-}{\alpha} \right) \right) \right], \tag{16}$$

where $\Pi^\Theta(v) = \arg\min_{\theta \in \Theta} \|v - \theta\|_2^2$ and Θ is convex and compact.

Figure 4: Dependency of the policy-parameter error $|\theta_t - \theta*|$ on the three key hyperparameter schedules in a 50-state heavy-tailed MDP (Pareto($\alpha = 2$) rewards, truncated at 10, $\gamma = 0.95$). Curves show the mean over 10 runs for: (i) inner-loop length $N_t = t$ (yellow), which reduces finite-difference bias and accelerates convergence; (ii) step-size $a_t = t^{-0.6}$ (orange), which maintains larger updates early at the cost of greater long-term variability; and (iii) perturbation size $c_t = t^{-0.5}$ (red), which aggressively shrinks gradient bias $O(c_t^2)$ and yields the fastest, most stable descent.

Figure 5: A minimal MDP with state space $S = \{s_1, s_2\}$, uniform transition probabilities $P(s' \mid s, a) = 0.5$, and a quadratic reward $R(s, a, s') = -\|\theta\|^2$, yielding an EVaR objective $J_{\text{EVaR}}(\theta) \approx -\frac{1}{2}\|\theta\|^2$. Gray arrows depict the true gradient field $\nabla J_{\text{EVaR}}(\theta)$. Starting from $\theta_0 = (1.5, 1.5)$, the *orange dotted* curve shows 20 exact gradient-ascent updates $\theta_{t+1} = \theta_t + 0.1\,\nabla J_{\text{EVaR}}(\theta_t)$, while the *red dotted* curve shows 20 SPSA-estimated updates using $\hat{g}_t$. This comparison illustrates how $\hat{g}_t$ tracks—but noisily perturbs—the true ascent path toward the EVaR-optimal parameter.

The above recursion can be rearranged as follows:

$$
\theta_{t+1} = \Pi^\Theta \Bigg[\theta_t + a_t \bigg(\underbrace{\mathbb{E}\Big[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\Big]}_{b_t} - \mathbb{E}\Big[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\Big] +
$$

$$
\widehat{\nabla J}_{\text{EVaR}}(\theta_t) + \underbrace{\frac{z(\theta_t^+) - z(\theta_t^-)}{2c_t\Delta_t}}_{\varphi_t} \bigg) \Bigg]
$$

Let

$$\bar{\theta}_t \;=\; \theta_t + a_t\Big(b_t + \varphi_t + \nabla J_{\texttt{EVaR}}(\theta_t) + e_t\Big) \qquad \text{and } u_t \;=\; \frac{\Pi^{\Theta}[\bar{\theta}_t] - \bar{\theta}_t}{a_t}.$$

Then,

$$\theta_{t+1} \;=\; \theta_t + a_t\big[\nabla J_{\texttt{EVaR}}(\theta_t) + e_t + b_t + \varphi_t + u_t\big]. \tag{18}$$

Because θ_{t+1} minimises $\|\theta - \bar{\theta}_{t+1}\|^2$ over Θ, the first–order optimality condition gives

$$\langle\, u_t,\; \theta - \theta_{t+1}\,\rangle \;\geqslant 0 \quad \forall \theta \in \Theta \quad \Leftrightarrow \quad -u_t \in N_{\Theta}(\theta_{t+1}) \quad \Leftrightarrow \quad u_t \in -\,N_{\Theta}(\theta_{t+1}), \tag{19}$$

where $N_{\Theta}(x) := \{\nu : \langle \nu, \theta - x \rangle \leqslant 0,\ \forall \theta \in \Theta\}$ is the normal cone at x. Thus the correction produced by the projection lies in the negative normal cone at the projected point. Therefore, Eq. (18) can be considered as

$$\theta_{t+1} - \theta_t \;\in\; a_t\Big[\,\nabla J_{\texttt{EVaR}}(\theta_t) \;-\; N_{\Theta}(\theta_{t+1}) \;+\; e_{t+1} + \varphi_t + b_t\Big]. \tag{20}$$

Therefore, θ_t will asymptotically follow the set-valued ODE (Benaïm, 2006; Borkar, 2009)

$$\frac{d\theta}{dt} \;\in\; J_{\texttt{EVaR}}(\theta) \;-\; N_{\Theta}(\theta). \tag{21}$$

Applying Corollary 4, Chapter 5 of (Borkar, 2009), to the stochastic inclusion equation 20, we obtain

$$\theta_t \;\xrightarrow{\text{a.s.}}\; \theta^{\star} \in \mathcal{E} = \{\theta \in \Theta| -\nabla J(\theta) \;+\; N_{\Theta}(\theta) = 0\} = \{\theta \in \Theta | \nabla J(\theta) \;=\; N_{\Theta}(\theta)\},$$

i.e. every sample path converges almost surely to the set $\mathcal{E}$. If the attractor set $\mathcal{E}$ is finite, then the iterates $\{\theta_t\}$ converge *a.s.* to a single locally EVaR-optimal policy $\theta^{\star} \in \mathcal{E}$.

Projected ascent iterates (Eq. 15) up to θ₃ — descent flow backdrop

Figure 6: Projected ascent on the toy two–state MDP. The return, $R(\tau) = -\|\boldsymbol{\theta}\|^2$, so the risk–seeking objective is $J_{\text{EVaR}}(\boldsymbol{\theta}) = -\frac{1}{2}\|\boldsymbol{\theta}\|^2$. Parameters are constrained to the shaded square $\Theta = [-0.6, 0.6]^2$. We follow $\boldsymbol{\theta}_{t+1} = \Pi^{\Theta}\big(\boldsymbol{\theta}_t + \alpha\,\hat{\mathbf{g}}\big)$ to produce the shown iterates $\boldsymbol{\theta}_0 \to \boldsymbol{\theta}_1 \to \boldsymbol{\theta}_2 \to \boldsymbol{\theta}_3$ (blue). Red ×'s mark the raw updates $\bar{\boldsymbol{\theta}}_t = \boldsymbol{\theta}_t + \alpha\hat{\mathbf{g}}$; grey arrows project them back to Θ, and orange wedges depict the outward normal cone $N_{\Theta}(\boldsymbol{\theta}_{t+1})$ at each boundary hit. The faint vector field in the background is the flow $\dot{\boldsymbol{\theta}} = -\boldsymbol{\theta}$ of J_{EVaR}.

Remark. *To improve the quality of the solution, one can inject a decaying Gaussian noise (Maryak & Chin, 2008) into the iterates θ_t as follows:*

$$\theta_{t+1} = \theta_t + \frac{a_t}{2c_t\Delta_t}\left(x_t^{+}\left(\log\frac{\vartheta_t^{+}}{\alpha}\right) - x_t^{-}\left(\log\frac{\vartheta_t^{-}}{\alpha}\right)\right) + q_t\varepsilon_t \tag{22}$$

where $q_t > 0$ is the step schedule and $\varepsilon_t \overset{\text{iid}}{\sim} \mathcal{N}(0, \mathcal{I})$. The noise term $q_t\epsilon_t$ introduces randomness into the update process, but this randomness is controlled by q_t which typically decreases over time to ensure that the influence of noise diminishes. The finite-difference gradient estimator has controlled bias $O(c_t^2)$ and the injected noise $q_t\varepsilon_t$ adds exploration without affecting asymptotic convergence. The perturbation leaves the limiting ODE unchanged – so the interpolated trajectory remains attracted to $\mathcal{E}$–but it alters the transient behaviour of the iterates, supplying occasional random deviations that let the iterate escape any strict, non-global local optimum. When the noise is suitably behaved and certain other conditions are satisfied, the iterates θ_t so generated converge to the global maxima of J_{EVaR}:

$$\lim_{t \to 0}[J_{EVaR}(\theta_t)] = J_{EVaR}(\theta^*) \tag{23}$$

Please refer to (Maryak & Chin, 2008) for the conditions required to ensure convergence.

Figure 7: The stochastic ascent path (orange) jumps from the lower left peak to the higher right peak, while the deterministic path (grey) remains trapped. Gaussian kicks $q_t\varepsilon_t$ enable this valley-crossing, and once the iterate enters the right basin, ascent resumes toward the global maximum.

3 Experimental Results

We conduct our experiments over various values of risk sensitivity, in each environment, we learn policies at different `EVaR` values α and evaluate the learned policies. For more details about the details and reproducibility of the experiments, please refer Appendix section(s) D and C.

3.1 GridWorld

We evaluate our algorithm in a discrete $m \times m$ grid environment, where agents must navigate from a predefined start state to a goal state avoiding obstacles. Each movement incurs a small cost, while reaching the goal provides a significant reward, encouraging efficient path selection. To promote generalization and robust policy learning, obstacles are randomly placed in each training batch, preventing the agent from overfitting to a fixed obstacle configuration.

Figure 8: The start state is fixed at $(0,0)$ [green], and the goal is at $(4,4)$[red]. [*Left*] Plot shows that mean rewards obtained for various levels of the threshold α for $m = 5$. [*Right*] Optimal path chosen to reach goal.

The agent exhibits risk-aware efficiency, consistently choosing the shortest path while successfully navigating random obstacle placements (Figure 8). We aslo evaluate our algorithm for various levels of $\alpha = [0.2, 0.15, 0.1, 0.05, 0.01]$. Interestingly, we observe that as α decreases, the agent converges to optimality more slowly. This behavior can be attributed to the effect of α on the function $H(\beta)$: for smaller values of α, $H(\beta)$ exhibits a flatter region near its base, forming a saddle point basin. As a result, the optimization process experiences slower convergence to the true `EVaR`, requiring more training iterations to refine the policy.

3.2 MuJoCo

Figure 9: For various $\alpha \in \{0.1, 0.01, 0.05\}$ we have the policy return values (a) and the respective `EVaR` objective value (b) using the finite difference optimization.

We consider the OpenAI Gym environments *InvertedDoublePendulum-v4, Swimmer-v4, Hopper-v4, Walker2d-v2* from the MuJoCo framework (Tassa et al., 2018) and *MountainCarContinuous-v0* from the Box2D Gym framework (Towers et al., 2023). These environments incorporate penalty terms in their reward functions, discouraging unsuccessful task completion. As a result, they exhibit higher reward variance, increasing the likelihood of worst-case scenarios, regardless of expected rewards. In Figure 9, we see the increase in rewards as well as the `EVaR` objective function value for *InvertedDoublePendulum*. More environments are in Appendix C.

3.3 Glycemic Control

Figure 10: We compare our proposed method with SAC (Haarnoja et al., 2018), demonstrating reduced variability in blood glucose levels and more robust insulin administration. We observe that when blood glucose reaches risky levels (red), our method effectively course-corrects insulin administration without prolonged exposure to risk, ensuring better patient stability.

We demonstrate our algorithm's ability to manage high-risk insulin administration for T1DM using the Simglucose simulator (Xie, 2018). Simglucose mimics real-world scenarios, providing a controlled environment to test control algorithms before clinical deployment. In our experiments, a PID controller regulates insulin based on blood glucose levels, aiming to keep them within a safe range. We evaluate performance on both adult and adolescent patient profiles to minimize the risk of hyper- and hypoglycemia, illustrated in Figure 10. Glycemic control is a high-stakes problem where inaction or overly conservative actions can lead to severe health risks such as prolonged hyperglycemia. A risk-averse policy (SAC) favors safe, incremental insulin adjustments, but this could result in suboptimal glucose regulation. This is illustrated in Figure 10. Our risk-seeking policy seems to be more focused in optimizing rewards, actively pushing for tight glycemic control, and taking bold insulin actions when necessary to rapidly correct glycemic levels.

3.4 Portfolio Optimization

The portfolio optimization problem seeks an optimal portfolio allocation among N assets by maximizing the `EVaR` of the portfolio returns $\mathcal{R}$, which captures the upside tail of the return distribution. Here policy represents the action chosen which includes sell, buy or hold. Constraints are kept to ensure that the portfolio weights w_i are nonnegative and sum to one, representing a fully invested portfolio. For our portfolio (top 10 stocks of DJIA) weights $\mathbf{w} \in \mathbb{R}^{10}$ are constrained such that $\sum_{i=1}^{10} w_i = 1$, $w_i \geqslant 0 \forall i$. A constant transaction cost of 0.1% is applied, computed as - Cost $= 0.001 \times \sum_{i=1}^{10} \left| w_i^{\text{new}} - w_i^{\text{old}} \right| \times$ Portfolio Value. In Table 1 we provide the backtesting (Wong, 2010) results of our `EVaR` strategy compared against other risk sensitive strategies as follows:

Portfolio	Cum Ret	Exp Ret	Ann Vol	Sharpe
EVaR	265.7%	13.3%	15.0%	0.75
VaR	250.0%	12.6%	14.1%	0.75
CVaR	257.5%	12.8%	10.2%	0.99

Table 1: Portfolio Backtest Summary

In Figure 11, we observe the upward trend of the portfolio, demonstrating resilience despite adverse market conditions. This is evident in the drawdown, which represents the maximum decline from peak to trough during the given time period. The portfolio's strong performance can be attributed to the risk-seeking nature of the algorithm, which strategically places aggressive bets to navigate and withstand market turbulence.

Further implementation details are provided in Appendix.

Figure 11: Drawdown analysis of our portfolio

3.5 Sensitivity to Learning rate

Figure 12: **Parameter sensitivity.** The figure shows the evolution of the policy parameter θ_t, the `EVaR` shape parameter x_t, the finite–difference bias b_t (`log axis`), and the drift term φ_t on a 50-state heavy-tailed MDP. The *baseline* schedule $\left[a_t = 0.10(t+1)^{-0.6}, c_t = 0.10(t+1)^{-0.10}, \delta_t = \xi_t = 0.05(t+1)^{-0.6}, N_t = 8\right]$ is compared with three single–factor variants: (i) a larger outer step–size $a_t = 0.20(t+1)^{-0.4}$, (ii) a faster perturbation decay $c_t \propto (t+1)^{-0.20}$, and (iii) a linearly growing inner batch $N_t = \min\{t+1, 15\}$. Solid lines show the mean over six i.i.d. runs; shaded envelopes indicate ± 1 standard deviation.

Figure 13: Sensitivity of the optimizer to the learning rate placed at $[0.01, 0.1, 0.2, 0.5]$ with ADAM for the top row for J_{EVaR} perturbation (a) and the expected returns (b) and similarly, the bottom row shows the similar setup without ADAM in (c) and (d).

To analyze hyper-parameter sensitivity, we consider an episodic, two–action MDP with $|S|=50$ states arranged on a ring. Action 0 (*safe*) keeps the agent in place and yields a deterministic reward 0.5; action 1 (*risky*) moves one step clockwise and delivers a heavy–tailed reward distributed as Pareto($k=1$, $b=2$). Episodes last $H = 5$ time steps with discount $\gamma = 0.95$. The policy is a one–parameter Bernoulli, $\Pr(a_t=1) = \sigma(\theta)$, so the return distribution is regularly varying, which stresses the `EVaR` objective.

With the baseline hyper–parameters, the bias decays like $t^{-0.20}$, the drift remains at $1/8$, and the variance of the stochastic gradient contracts as $t^{-0.60}$, bringing (θ_t, x_t) close to their limits by iteration 50. Increasing the outer step–size accelerates early progress but leaves a wider asymptotic band because the noise term $a_t^2 \text{Var}$ dominates more slowly. A steeper perturbation decay eliminates bias faster yet stalls convergence once the finite–difference signal falls below the simulation noise floor. Growing the inner batch length suppresses the drift from 10^{-1} to 10^{-2} in fewer than twenty iterations, yielding the tightest bands at the expense of a quadratic increase in sample complexity. These trends corroborate theoretical predictions: a_t sets the speed–variance trade-off, c_t governs asymptotic bias, and N_t controls late–stage variance once the iterate enters the EVaR basin of attraction.

We further analyze the impact of learning rates on algorithm performance using *InvertedPendulumDouble-v4*. We evaluate learning rates $0.01, 0.1, 0.2,$ and 0.5 by averaging results over 5 independent batches, each with 50 episodes of 500 steps. The optimizer runs for 500 iterations with the risk parameter $\alpha = 0.1$. Using ADAM(Kingma, 2014) improves iterate stability, with 10^{-2} yielding the best performance. In this context of our gradient estimator with ADAM, our learning rate parameter update rule is:

$$a_{t+1} = a_t - \psi \cdot \frac{\hat{m}_t}{\sqrt{\hat{v}_t} + \epsilon}$$

where: $\hat{m}_t$ is the bias-corrected first moment estimate and $\hat{v}_t$ is the bias-corrected second moment estimate. To show that under the ADAM condition the convergence is guaranteed, we need to show that $\mathbb{E}[\|a_t - a^*\|^2] \to 0$ as $t \to \infty$, where a^* is the optimal learning rate parameter. We give a proof sketch as follows

Proof. We define

$$\mathcal{L}_t = \mathbb{E}[\|a_t - a^*\|^2]$$

Now, the expected change $\mathbb{E}[\mathcal{L}_{t+1} - \mathcal{L}_t] = \mathbb{E}[\|a_{t+1} - a^*\|^2 - \|a_t - a^*\|^2]$, which is upper bounded as

$$\mathbb{E}[L_{t+1} - L_t] = \mathbb{E}[\|a_{t+1} - a^*\|^2 - \|a_t - a^*\|^2]$$

$$= \mathbb{E}[\|a_t - \psi\frac{\hat{m}_t}{\sqrt{\hat{v}_t} + \epsilon} - a^*\|^2 - \|a_t - a^*\|^2]$$

$$\leqslant -2\psi\mathbb{E}[(a_t - a^*)\frac{\hat{m}_t}{\sqrt{\hat{v}_t} + \epsilon}] + \psi^2\mathbb{E}[\|\frac{\hat{m}_t}{\sqrt{\hat{v}_t} + \epsilon}\|^2]$$

Choosing a decreasing learning rate schedule $\psi_t = \frac{\psi_0}{\sqrt{t}}$ and under conditions $\sum_{t=1}^{\infty}\psi_t = \infty, \sum_{t=1}^{\infty}\psi_t^2 < \infty$, we can conclude that under appropriate choice of constants and applying the above conditions, we can show that $\mathbb{E}[\mathcal{L}_{t+1} - \mathcal{L}_t] \leqslant 0$ for sufficiently large t, which implies convergence. $\qquad\square$

For bounding the variance $\mathbb{V}[a_t]$, we proceed as follows

$$\mathbb{V}[a_{t+1}] = \mathbb{V}[a_t - \psi_t\frac{\hat{m}_t}{\sqrt{\hat{v}_t} + \epsilon}] \leqslant (1 - \psi_t\lambda)^2\mathbb{V}[a_t] + \psi_t^2\sigma^2$$

where λ is related to the smallest eigenvalue of the Hessian of the objective function, and σ^2 bounds the variance of the gradient estimate. Now unrolling the recursion, we get

$$\mathbb{V}[a_t] \leqslant \prod_{i=1}^{t}(1 - \psi_i\lambda)^2\mathbb{V}[a_0] + \sum_{i=1}^{t}\psi_i^2\sigma^2\prod_{j=i+1}^{t}(1 - \psi_j\lambda)^2$$

Using the decreasing learning rate schedule with $\psi_t = \frac{\psi_0}{\sqrt{t}}$, we show that $\mathbb{V}[a_t] = \mathcal{O}\left(\frac{1}{\sqrt{t}}\right)$. As learning rate sensitivity affects both convergence and variance a higher sensitivity can lead to faster initial convergence but may impact long-term stability. Very high sensitivity can increase the upper bound on variance, potentially leading to less stable convergence. The adaptive nature of ADAM helps mitigate these effects by adjusting the effective learning rate based on the moments of the gradients. From Fig. 13(a) which depicts the movement of the iterates and Fig.13(b) the expected, it is evident that the introduction of an adaptive learning schedule for gradient estimator of `EVaR` controls rapid movement of the iterates and is resilient against environment dynamics. When compared against the non-adaptive case Fig. 13(c) and (d), we clearly see increased movement as the initial learning rate decreases depicting high susceptibility to the initial choice of the learning rate.

4   Conclusion

We optimize risk-seeking RL policies using `EVaR` of discounted rewards, leveraging its monotonicity and interpretability. By employing randomly perturbed finite difference approximation, we optimize the `EVaR` policy. Experiments across discrete and continuous tasks show that `EVaR` achieves higher rewards and demonstrates its effectiveness for risk-sensitive RL.

References

Mohamadreza Ahmadi, Ugo Rosolia, Michel D Ingham, Richard M Murray, and Aaron D Ames. Constrained risk-averse markov decision processes. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 35, pp. 11718–11725, 2021.

Amir Ahmadi-Javid. Entropic value-at-risk: A new coherent risk measure. *Journal of Optimization Theory and Applications*, 155:1105–1123, 2012.

Michel Benaïm. Dynamics of stochastic approximation algorithms. In *Seminaire de probabilites XXXIII*, pp. 1–68. Springer, 2006.

Vivek S. Borkar. Convex analytic methods in markov decision processes. *Handbook of Markov Decision Processes*, pp. 347–375, 2001.

Vivek S. Borkar. Q-learning for risk-sensitive control. *Mathematics of Operations Research*, 27(2):294–311, 2002.

Vivek S Borkar. *Stochastic approximation: a dynamical systems viewpoint*, volume 9. Springer, 2008.

Vivek S Borkar. *Stochastic approximation: a dynamical systems viewpoint*, volume 48. Springer, 2009.

Vivek S. Borkar. Risk-constrained markov decision processes. *Stochastics*, 82(3):337–361, 2010.

Vivek S. Borkar and Sean P. Meyn. Risk-sensitive control with discounted cost in discrete time. *SIAM Journal on Control and Optimization*, 40(3):681–707, 2002.

Greg Brockman, Vicki Cheung, Ludwig Pettersson, Jonas Schneider, John Schulman, Jie Tang, and Wojciech Zaremba. Openai gym. *arXiv preprint arXiv:1606.01540*, 2016.

Dany Cajas. Entropic portfolio optimization: a disciplined convex programming framework. *Available at SSRN 3792520*, 2021.

Dany Cajas. Riskfolio-lib (6.1.0), 2024. URL `https://github.com/dcajasn/Riskfolio-Lib`.

Jay Cao, Jacky Chen, John Hull, and Zissis Poulos. Deep hedging of derivatives using reinforcement learning. *arXiv preprint arXiv:2103.16409*, 2021.

Rolando Cavazos-Cadena and Daniel Hernandez-Hernandez. Risk-sensitive control of discrete-time markov systems with feedback information. *SIAM Journal on Control and Optimization*, 49(4):1593–1612, 2011.

Venkat Chandrasekaran and Parikshit Shah. Relative entropy optimization and its applications. *Mathematical Programming*, 161:1–32, 2017.

Robert Chares. Cones and interior-point algorithms for structured convex optimization involving powers andexponentials, 2009.

I-Ju Chen, Markus Aapro, Abraham Kipnis, Alexander Ilin, Peter Liljeroth, and Adam S Foster. Precise atom manipulation through deep reinforcement learning. *Nature Communications*, 13(1):7499, 2022.

Yinlam Chow and Mohammad Ghavamzadeh. Algorithms for cvar optimization in mdps. *Advances in neural information processing systems*, 27, 2014.

Yinlam Chow, Aviv Tamar, Shie Mannor, and Marco Pavone. Risk-sensitive and robust decision-making: a cvar optimization approach. *Advances in neural information processing systems*, 28, 2015.

Stefano P Coraluppi and Steven I Marcus. Risk-sensitive and minimax control of discrete-time, finite-state markov decision processes. *Automatica*, 35(2):301–309, 1999.

Giovanni B. Di Masi and Lukasz Stettner. Risk-sensitive control of discrete-time markov processes. *Mathematics of Operations Research*, 24(3):552–568, 1999.

Giovanni B. Di Masi and Lukasz Stettner. Risk-sensitive control of continuous-time markov processes with unbounded cost rates. *SIAM Journal on Control and Optimization*, 45(3):1061–1079, 2007.

Anushri Dixit, Mohamadreza Ahmadi, and Joel W Burdick. Risk-sensitive motion planning using entropic value-at-risk. In *2021 European Control Conference (ECC)*, pp. 1726–1732. IEEE, 2021.

Angelos Filos. Reinforcement learning for portfolio management. *arXiv preprint arXiv:1909.09571*, 2019.

Wendell H. Fleming and William M. McEneaney. Risk-sensitive control and differential games. *Applied Mathematics and Optimization*, 31(3):235–261, 1995.

Javier Garcıa and Fernando Fernández. A comprehensive survey on safe reinforcement learning. *Journal of Machine Learning Research*, 16(1):1437–1480, 2015.

Christian Gollier. *The economics of risk and time*. MIT press, 2001.

Shangding Gu, Alap Kshirsagar, Yali Du, Guang Chen, Jan Peters, and Alois Knoll. A human-centered safe robot reinforcement learning framework with interactive behaviors. *Frontiers in Neurorobotics*, 17:1280341, 2023.

Tuomas Haarnoja, Aurick Zhou, Pieter Abbeel, and Sergey Levine. Soft actor-critic: Off-policy maximum entropy deep reinforcement learning with a stochastic actor. In *International conference on machine learning*, pp. 1861–1870. PMLR, 2018.

Jia Lin Hau, Marek Petrik, and Mohammad Ghavamzadeh. Entropic risk optimization in discounted mdps. In *International Conference on Artificial Intelligence and Statistics*, pp. 47–76. PMLR, 2023.

Matthias Heger. Consideration of risk in reinforcement learning. In *Machine Learning Proceedings 1994*, pp. 105–111. Elsevier, 1994.

Ashley Hill, Antonin Raffin, Maximilian Ernestus, Adam Gleave, Anssi Kanervisto, Rene Traore, Prafulla Dhariwal, Christopher Hesse, Oleg Klimov, Alex Nichol, Matthias Plappert, Alec Radford, John Schulman, Szymon Sidor, and Yuhuai Wu. Stable baselines. `https://github.com/hill-a/stable-baselines`, 2018.

Ronald A. Howard and James E. Matheson. Risk-sensitive markov decision processes. *Management Science*, 18(7):356–369, 1972a.

Ronald A Howard and James E Matheson. Risk-sensitive markov decision processes. *Management science*, 18 (7):356–369, 1972b.

Hisashi Kashima. Risk-sensitive learning via minimization of empirical conditional value-at-risk. *IEICE TRANSACTIONS on Information and Systems*, 90(12):2043–2052, 2007.

Elia Kaufmann, Leonard Bauersfeld, Antonio Loquercio, Matthias Müller, Vladlen Koltun, and Davide Scaramuzza. Champion-level drone racing using deep reinforcement learning. *Nature*, 620(7976):982–987, 2023.

Ramtin Keramati, Christoph Dann, Alex Tamkin, and Emma Brunskill. Being optimistic to be conservative: Quickly learning a cvar policy. In *Proceedings of the AAAI conference on artificial intelligence*, volume 34, pp. 4436–4443, 2020.

Diederik P Kingma. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.

Jens Kober, J. Andrew Bagnell, and Jan Peters. Reinforcement learning in robotics: A survey. *Int. J. Rob. Res.*, 32(11):1238–1274, September 2013. ISSN 0278-3649. doi: 10.1177/0278364913495721. URL `https://doi.org/10.1177/0278364913495721`.

Sven Koenig and Reid G Simmons. Risk-sensitive planning with probabilistic decision graphs. In *Principles of Knowledge Representation and Reasoning*, pp. 363–373. Elsevier, 1994.

Thanh Lam, Arun Verma, Bryan Kian Hsiang Low, and Patrick Jaillet. Risk-aware reinforcement learning with coherent risk measures and non-linear function approximation. In *The Eleventh International Conference on Learning Representations*, 2022.

Michael L Littman and Csaba Szepesvári. A generalized reinforcement-learning model: Convergence and applications. In *ICML*, volume 96, pp. 310–318, 1996.

Harry M Markowitz and G Peter Todd. *Mean-variance analysis in portfolio choice and capital markets*, volume 66. John Wiley & Sons, 2000.

José Antonio Martín H. and Javier de Lope. Learning autonomous helicopter flight with evolutionary reinforcement learning. In Roberto Moreno-Díaz, Franz Pichler, and Alexis Quesada-Arencibia (eds.), *Computer Aided Systems Theory - EUROCAST 2009*, pp. 75–82, Berlin, Heidelberg, 2009. Springer Berlin Heidelberg. ISBN 978-3-642-04772-5.

John L Maryak and Daniel C Chin. Global random optimization by simultaneous perturbation stochastic approximation. *IEEE TRANSACTIONS ON AUTOMATIC CONTROL*, 53(3):781, 2008.

Oliver Mihatsch and Ralph Neuneier. Risk-sensitive reinforcement learning. In *Advances in Neural Information Processing Systems*, pp. 172–178, 2002.

Xinyi Ni and Lifeng Lai. Policy gradient based entropic-var optimization in risk-sensitive reinforcement learning. In *2022 58th Annual Allerton Conference on Communication, Control, and Computing (Allerton)*, pp. 1–6. IEEE, 2022.

Takayuki Osogami. Risk-sensitive reinforcement learning by relative entropy stochastic search. In *Advances in Neural Information Processing Systems*, pp. 1647–1655, 2012.

LA Prashanth. Policy gradients for cvar-constrained mdps. In *International Conference on Algorithmic Learning Theory*, pp. 155–169. Springer, 2014.

Martin L Puterman. *Markov decision processes: discrete stochastic dynamic programming*. John Wiley & Sons, 2014.

Marc Rigter, Bruno Lacerda, and Nick Hawes. Risk-averse bayes-adaptive reinforcement learning. *Advances in Neural Information Processing Systems*, 34:1142–1154, 2021.

R Tyrrell Rockafellar and Stanislav Uryasev. Conditional value-at-risk for general loss distributions. *Journal of banking & finance*, 26(7):1443–1471, 2002.

R Tyrrell Rockafellar, Stanislav Uryasev, et al. Optimization of conditional value-at-risk. *Journal of risk*, 2: 21–42, 2000.

Mark Rowland, Rémi Munos, Mohammad Gheshlaghi Azar, Yunhao Tang, Georg Ostrovski, Anna Harutyunyan, Karl Tuyls, Marc G Bellemare, and Will Dabney. An analysis of quantile temporal-difference learning. *Journal of Machine Learning Research*, 25:1–47, 2024.

Robert J Serfling. *Approximation theorems of mathematical statistics*. John Wiley & Sons, 2009.

Harshit Sethy, Amit Patel, and Vineet Padmanabhan. Real time strategy games: a reinforcement learning approach. *Procedia Computer Science*, 54:257–264, 2015.

Kun Shao, Zhentao Tang, Yuanheng Zhu, Nannan Li, and Dongbin Zhao. A survey of deep reinforcement learning in video games. *arXiv preprint arXiv:1912.10944*, 2019.

David Silver, Aja Huang, Chris J Maddison, Arthur Guez, Laurent Sifre, George Van Den Driessche, Julian Schrittwieser, Ioannis Antonoglou, Veda Panneershelvam, Marc Lanctot, et al. Mastering the game of go with deep neural networks and tree search. *nature*, 529(7587):484–489, 2016.

Rahul Singh, Qinsheng Zhang, and Yongxin Chen. Improving robustness via risk averse distributional reinforcement learning. In *Learning for Dynamics and Control*, pp. 958–968. PMLR, 2020.

James C Spall. Multivariate stochastic approximation using a simultaneous perturbation gradient approximation. *IEEE transactions on automatic control*, 37(3):332–341, 1992.

Richard S Sutton and Andrew G Barto. Reinforcement learning: an introduction mit press. *Cambridge, MA*, 22447, 1998.

Aviv Tamar, Yonatan Glassner, and Shie Mannor. Optimizing the cvar via sampling. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 29, 2015.

Yuval Tassa, Yotam Doron, Alistair Muldal, Tom Erez, Yazhe Li, Diego de Las Casas, David Budden, Abbas Abdolmaleki, Josh Merel, Andrew Lefrancq, et al. Deepmind control suite. *arXiv preprint arXiv:1801.00690*, 2018.

Mark Towers, Jordan K. Terry, Ariel Kwiatkowski, John U. Balis, Gianluca de Cola, Tristan Deleu, Manuel Goulão, Andreas Kallinteris, Arjun KG, Markus Krimmel, Rodrigo Perez-Vicente, Andrea Pierré, Sander Schulhoff, Jun Jet Tai, Andrew Tan Jin Shen, and Omar G. Younis. Gymnasium, March 2023. URL https://zenodo.org/record/8127025.

Amos Tversky and Daniel Kahneman. Advances in prospect theory: Cumulative representation of uncertainty. *Journal of Risk and uncertainty*, 5:297–323, 1992.

Guangyu Wang, Xiaohong Liu, Zhen Ying, Guoxing Yang, Zhiwei Chen, Zhiwen Liu, Min Zhang, Hongmei Yan, Yuxing Lu, Yuanxu Gao, et al. Optimized glycemic control of type 2 diabetes with reinforcement learning: a proof-of-concept trial. *Nature Medicine*, 29(10):2633–2642, 2023.

Shaun Wang. Premium calculation by transforming the layer premium density. *ASTIN Bulletin: The Journal of the IAA*, 26(1):71–92, 1996.

Peter Whittle. Risk-sensitive linear/quadratic/gaussian control. *Advances in Applied Probability*, 22(4): 764–774, 1990.

Woon K. Wong. Backtesting value-at-risk based on tail losses. *Journal of Empirical Finance*, 17(3):526–538, 2010. ISSN 0927-5398. doi: https://doi.org/10.1016/j.jempfin.2009.11.004. URL https://www.sciencedirect.com/science/article/pii/S0927539809000929.

Jingda Wu, Chao Huang, Hailong Huang, Chen Lv, Yuntong Wang, and Fei-Yue Wang. Recent advances in reinforcement learning-based autonomous driving behavior planning: A survey. *Transportation Research Part C: Emerging Technologies*, 164:104654, 2024.

Peter R Wurman, Samuel Barrett, Kenta Kawamoto, James MacGlashan, Kaushik Subramanian, Thomas J Walsh, Roberto Capobianco, Alisa Devlic, Franziska Eckert, Florian Fuchs, et al. Outracing champion gran turismo drivers with deep reinforcement learning. *Nature*, 602(7896):223–228, 2022.

J. Xie. Jxx123/simglucose: A type-1 diabetes simulator implemented in python for reinforcement learning purpose. https://github.com/jxx123/simglucose, 2018. Accessed: 2024-05-22.

Jesse Zhang, Brian Cheung, Chelsea Finn, Sergey Levine, and Dinesh Jayaraman. Cautious adaptation for reinforcement learning in safety-critical settings. In *International Conference on Machine Learning*, pp. 11055–11065. PMLR, 2020.

Qiyuan Zhang, Shu Leng, Xiaoteng Ma, Qihan Liu, Xueqian Wang, Bin Liang, Yu Liu, and Jun Yang. Cvar-constrained policy optimization for safe reinforcement learning. *IEEE Transactions on Neural Networks and Learning Systems*, 2024.

A   Proofs

Consider the filtration $\mathcal{F}_t = \sigma\{\vartheta_t, J, \omega_t, \beta_t, \theta_t\}$. We prove Theorem 4 in two parts.

A.1   Proof of Theorem 2

Proof of Theorem 2: Part I

Proof. Consider the recursion

$$
\begin{aligned}
\vartheta_{t+1} &= \vartheta_t + \delta_t \left(e^{R(\tau)/x} - \vartheta_t \right) \\
&= \vartheta_t + \delta_t \left(\mathbb{E}\left[e^{R(\tau)/x} \right] + e^{R(\tau)/x} - \mathbb{E}\left[e^{R(\tau)/x} \right] - \vartheta_t \right) \\
&= \vartheta_t + \delta_t \left(h_1(\vartheta_t) + \mathbb{M}_{t+1} \right), \\
&\text{where } h_1(\vartheta) = \mathbb{E}\left[e^{R(\tau)/x} \right] - \vartheta, \text{ and } \mathbb{M}_{t+1} = e^{R(\tau)/x} - \mathbb{E}\left[e^{R(\tau)/x} \right]
\end{aligned}
\tag{24}
$$

It is easy to verify the $\{\mathbb{M}_t\}$ is a Martingale difference with respect to the filtration $\{\mathcal{F}_t\}$ *i.e.*, $\mathbb{M}_t$ is $\mathcal{F}_t$-measurable and $\mathbb{E}\left[\mathbb{M}_{t+1}|\mathbb{F}_t\right] = 0$ *a.s.* Also, note that h_1 is Lipschitz continuous. Further, by Borkar-Meyn Theorem (Theorem 7, Chapter 3 of Borkar (2009)), one can show that $\sup_t |\vartheta_t| < \infty$ *a.s.* Hence, by Theorem 1 of Chapter 2 of Borkar (2009), the iterates $\{\vartheta_t\}$ asymptotically tracks the following ODE:

$$
\dot{\vartheta} = \mathbb{E}_{\tau \sim \pi_\theta}\left[e^{R(\tau)/x} \right] - \vartheta.
\tag{25}
$$

Since h_1 is linear, we conclude that

$$
\lim_{t \to \infty} \vartheta_t = \{\vartheta^* | h(\vartheta^*) = 0\} \Rightarrow \lim_{t \to \infty} \vartheta_t = \mathbb{E}_{\tau \sim \pi_\theta}\left[e^{\beta R(\tau)} \right].
\tag{26}
$$

Now consider the stochastic recursion

$$
\begin{aligned}
\omega_{t+1} &= \omega_t + \delta_t \left(R(\tau) e^{\beta R(\tau)} - \omega_t e^{R(\tau)/x} \right) \\
&= \omega_t + \delta_t \left(R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} - \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} \right] + \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} \right] \right)
\end{aligned}
$$

Rewriting the above equation, we get

$$
\omega_{t+1} = \omega_t + \delta_t \left(h_\omega(\omega_t) + \mathbb{M}^\omega_{t+1} \right), \text{ where}
\tag{27}
$$

$$
h^\omega(\omega) = \mathbb{E}_\tau \left[R(\tau) e^{R(\tau)/x} - \omega e^{R(\tau)/x} \right] \text{ and}
\tag{28}
$$

$$
\mathbb{M}^\omega_{t+1} = R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} - \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} \Big| \mathcal{F}_t \right].
\tag{29}
$$

Now,

$$
\begin{aligned}
\mathbb{E}\left[\mathbb{M}^\omega_{t+1}|\mathcal{F}_t\right] &= \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} - \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} \Big| \mathcal{F}_t \right] \Big| \mathcal{F}_t \right] \\
&= \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{R(\tau)/x} \Big| \mathcal{F}_t \right] - \mathbb{E}\left[R(\tau) e^{R(\tau)/x} - \omega_t e^{\beta R(\tau)} \Big| \mathcal{F}_t \right] \Big| \mathcal{F}_t \right] = 0
\end{aligned}
$$

Hence $\{\mathbb{M}^\omega_t\}$ is a Martingale-difference noise adapted to the filtration $\{\mathcal{F}_t\}$. Also since $\gamma \in [0, 1)$ and we have $|R(\tau)| < \frac{R_\infty}{1-\gamma}$. Hence,

$$
\exists K_w > 0 \ s.t. \ \mathbb{E}\left[|\mathbb{M}^\omega_t|^2|\mathcal{F}_t\right] < (1 + K_w)(|\omega_t|)^2.
\tag{30}
$$

Also, for $\omega_1, \omega_2 \in \mathbb{R}$, we have

$$
\begin{aligned}
|h^\omega(\omega_1) - h^\omega(\omega_2)| &= \mathbb{E}_\tau \left[e^{R(\tau)/x} \right] |\omega_1 - \omega_2| \\
&\leqslant \exp\left\{ \left(\frac{R_\infty}{x(1-\gamma)} \right) \right\} |\omega_1 - \omega_2|.
\end{aligned}
$$

Hence h^ω is Lipschitz continuous.

Now, we will show that the iterates θ_t are stable, *i.e.*, $\sup_t |\omega_t| < \infty$ *a.s.* Hence, we consider the following scaled functions

$$
h_c(\omega) = \frac{h^\omega(c\omega)}{c}, c > 0. \tag{31}
$$

Now consider the ∞-ODE given by

$$
\dot{\omega} = h^\omega_\infty(\omega). \tag{32}
$$

where

$$
\begin{aligned}
h^\omega_\infty(\omega) = \lim_{c \to \infty} h^\omega_c(\omega) = \lim_{c \to \infty} \frac{h(c\omega)}{c} &= \lim_{c \to \infty} \frac{1}{c} \left(\mathbb{E}_\tau \left[R(\tau) e^{R(\tau)/x} - c\omega e^{R(\tau)/x} \right] \right) \\
&= -\omega \mathbb{E}_\tau \left[e^{R(\tau)/x} \right]
\end{aligned} \tag{33}
$$

Hence ODE (32) becomes

$$
\dot{\omega} = -U\omega, \text{ where } U = \mathbb{E}_\tau \left[e^{R(\tau)/x} \right]. \tag{34}
$$

Note that since $\beta > 0$, we have $U > 0$. Hence the ∞-ODE given above has a unique globally asymptotically stable equilibrium point. Hence by Borkar-Meyn Theorem (Theorem 7 Chapter 3 of Borkar (2009)), we haves

$$
\sup_t |\omega_t| < \infty \text{ a.s.} \tag{35}
$$

Now by Theorem 1 of Chapter 2 of (Borkar, 2009), the sequence (ω_t^θ) converges to a compact connected internally chain transitive invariant set of the ODE given by

$$
\dot{\omega} = h^\omega(\omega). \tag{36}
$$

Since h^ω is a linear function, the only compact connected internally chain transitive invariant set is $\{\omega | h^\omega(\omega) = 0\}$. Hence

$$
\begin{aligned}
h^\omega(\omega) = 0 &\Rightarrow \mathbb{E}_\tau \left[R(\tau) e^{R(\tau)/x} \right] - \omega \mathbb{E}_\tau \left[e^{R(\tau)/x} \right] = 0 \\
&\Rightarrow \omega = \frac{\mathbb{E}_\tau \left[R(\tau) e^{R(\tau)/x} \right]}{\mathbb{E}_\tau \left[e^{R(\tau)/x} \right]}
\end{aligned}
$$

Therefore

$$
\lim_{t \to \infty} \omega_t = \frac{\mathbb{E}_\tau \left[R(\tau) e^{R(\tau)/x} \right]}{\mathbb{E}_\tau \left[e^{R(\tau)/x} \right]} \quad a.s. \tag{37}
$$

$\square$

Lemma 3. *The function* $\vartheta^*(x, \theta) = \mathbb{E}_{\tau \sim \pi_\theta} \left[e^{R(\tau)/x} \right]$ *is Lipschitz continuous on* $x \in [x_{min}, x_{max}]$, *i.e.,* $\exists L_\vartheta > 0$ *such that:*

$$
|\vartheta^*(x_1, \theta) - \vartheta^*(x_2, \theta)| \leqslant L_\vartheta |x_1 - x_2|, \quad \forall x_1, x_2 \in [x_{min}, x_{max}].
$$

Proof. We compute the derivative of $\vartheta^*(x, \theta)$ with respect tot x as follows:

$$\frac{d}{dx}\vartheta^*(x,,\theta) = \mathbb{E}_\tau\left[-\frac{R(\tau)}{x^2}e^{R(\tau)/x}\right].$$

Since $|R| \leqslant \frac{R_\infty}{1-\gamma}$ and $x \geqslant x_{min}$, we have:

$$\left|\frac{d}{dx}\vartheta^*(x,\theta)\right| \leqslant \frac{R_\infty}{(1-\gamma)x_{min}^2}\mathbb{E}_\tau\left[e^{R(\tau)/x}\right] \leqslant \frac{R_\infty}{(1-\gamma)x_{min}^2}e^{\frac{R_\infty}{(1-\gamma)x_{min}}}.$$

Let $L_\vartheta = \frac{R_\infty}{(1-\gamma)x_{min}^2}e^{\frac{R_\infty}{(1-\gamma)x_{min}}}$. By the Mean Value Theorem, $\forall x_1, x_2 \in [x_{min}, x_{\max}]$:

$$|\vartheta^*(x_1,\theta) - \vartheta^*(x_2,\theta)| \leqslant L_\vartheta|x_1 - x_2|.$$

Hence, $\vartheta^*(x,\theta)$ is Lipschitz continuous in x. $\qquad\square$

Lemma 4. *The function* $\omega^*(x,\theta) = \frac{\mathbb{E}_{\tau\sim\pi_\theta}\left[R(\tau)e^{R(\tau)/x}\right]}{x\mathbb{E}_\tau\left[e^{R(\tau)/x}\right]}$ *is Lipschitz continuous on* $x \in [x_{min}, x_{max}]$, *i.e., there exists* $L_\omega > 0$ *such that:*

$$|\omega^*(x_1,\theta) - \omega^*(x_2,\theta)| \leqslant L_\omega|x_1 - x_2|, \quad \forall x_1, x_2 \in [x_{min}, x_{max}].$$

Proof. Let $N(x) = \mathbb{E}_\tau[R(\tau)e^{R(\tau)/x}]$ and $D(x) = x\mathbb{E}_\tau[e^{R(\tau)/x}]$. Then: $\omega^*(x,\theta) = \frac{N(x)}{D(x)}$. Their derivatives are:

$$N'(x) = -\frac{1}{x^2}\mathbb{E}_\tau[R(\tau)^2 e^{R(\tau)/x}], \quad D'(x) = \mathbb{E}_\tau[e^{R(\tau)/x}] - \frac{1}{x}\mathbb{E}_\tau[R(\tau)e^{R(\tau)/x}].$$

Using the quotient rule, we obtain:

$$\frac{d}{dx}\omega^*(x,\theta) = \frac{N'(x)D(x) - N(x)D'(x)}{D(x)^2}.$$

Bounding each term ($|R(\tau)| \leqslant \frac{R_\infty}{(1-\gamma)}$, $x \geqslant x_{min}$) from above as follows:

$$|N'(x)| \leqslant \frac{R_\infty^2}{(1-\gamma)^2 x_{min}^2}e^{\frac{R_\infty}{(1-\gamma)x_{min}}}$$
$$|D(x)| \geqslant x_{min}e^{-C/a},$$
$$|D'(x)| \leqslant e^{\frac{R_\infty}{(1-\gamma)x_{min}}} + \frac{R_\infty}{(1-\gamma)x_{min}}e^{\frac{R_\infty}{(1-\gamma)x_{min}}},$$
$$|N(x)| \leqslant \frac{R_\infty}{(1-\gamma)}e^{\frac{R_\infty}{(1-\gamma)x_{min}}}.$$

Substituting these bounds, we get:

$$\left|\frac{d}{dx}\omega^*(x,\theta)\right| \leqslant \left(\frac{R_\infty^2}{(1-\gamma)^2 x_{min}^2}e^{\frac{2R_\infty}{(1-\gamma)x_{min}}} + \frac{R_\infty}{(1-\gamma)}e^{\frac{2R_\infty}{(1-\gamma)x_{min}}}(1 + \frac{R_\infty}{(1-\gamma)x_{min}})\right)x_{min}^{-2}e^{\frac{2R_\infty}{(1-\gamma)x_{min}}} \triangleq L_\omega.$$

Thus $|\omega^*(x_1,\theta) - \omega^*(x_2,\theta)| \leqslant L_\omega|x_1 - x_2|$. $\qquad\square$

Lemma 5. *The first order derivative* $G'(x) = \log\frac{\vartheta^*(x,\theta)}{\alpha} - \omega^*(x,\theta)$ *is Lipschitz continuous on* $x \in [x_{min}, x_{max}]$, *i.e.,* $\exists L_{G'} > 0$ *such that:*

$$\left|G'(x_1) - G'(x_2)\right| \leqslant L_{G'}|x_1 - x_2|, \quad \forall x_1, x_2 \in [x_{min}, x_{max}].$$

Proof. Consider the derivative:

$$\frac{d}{dx}\left(\log\frac{\vartheta^*(x,\theta)}{\alpha}\right) = \frac{\vartheta^*(x,\theta)'}{\vartheta^*(x,\theta)}.$$

Using Lemma 3, we obtain:

$$\left| \frac{d}{dx} \log \frac{\vartheta^*(x,\theta)}{\alpha} \right| \leqslant L_\vartheta e^{\frac{R_\infty}{x_{min}(1-\gamma)}}.$$

From Lemma 4, $\omega^*(x,\theta)$ has Lipschitz constant L_ω. Therefore:

$$|G'(x_1) - G'(x_2)| \leqslant \left(L_\vartheta e^{\frac{R_\infty}{x_{min}(1-\gamma)}} + L_\omega \right) |x_1 - x_2| \triangleq L_G |x_1 - x_2|.$$

$\square$

Proof of Theorem 2 Part II:

Proof. On the slower timescale, the x–update approximates the ODE:

$$\dot{x} = -G'(x) = -\left(\log \frac{\mathbb{E}_{\tau \sim \pi_\theta}\left[e^{R(\tau)/x} \right]}{\alpha} - \frac{\mathbb{E}_{\tau \sim \pi_\theta}\left[R(\tau) e^{R(\tau)/x} \right]}{x \, \mathbb{E}_{\tau \sim \pi_\theta}\left[e^{R(\tau)/x} \right]} \right).$$

By Theorem 1, $G(x)$ is m–strongly convex. Using $G(x)$ as a Lyapunov function:

$$\frac{dG}{dt} = G'(x) \, \dot{x} = -\left(G'(x) \right)^2 \leqslant 0,$$

with equality only at $x = x^*(\theta)$. By LaSalle's invariance principle,

$$x_t \to x^*(\theta) \quad \text{almost surely.}$$

Now regarding the joint convergence, note that the time–scale separation ensures

$$\vartheta_t \to \vartheta^*(x_t,\theta) \quad \text{and} \quad \omega_t \to \omega^*(x_t,\theta)$$

before x_t updates significantly. Thus the system converges to

$$(\vartheta^*(x^*(\theta),\theta), \, \omega^*(x^*(\theta),\theta), \, x^*(\theta)) \quad a.s.$$

$\square$

A.2 Proof of Theorem 3

Lemma 6. *Let $\delta_t = \frac{c}{t}$ with $c > \frac{1}{2}$. Then for any fixed $x > 0$,*

$$\mathbb{E}\left[|\vartheta_t - \vartheta^*(x,\theta)|^2 \right] \leqslant \frac{K_1}{t},$$

where $K_1 = C\left(1 + c^2 \nu^2 \frac{\pi^2}{6} \right)$, $\nu^2 = e^{\frac{2R_\infty}{(1-\gamma)x_{min}}}$, and $C > 0$ is a constant.

Proof. Define the error process $\epsilon_t^\vartheta := \vartheta_t - \vartheta^*(x,\theta)$, where $\vartheta^*(x,\theta) = \mathbb{E}_{\tau \sim \pi_\theta}[e^{R(\tau)/x}]$. The update rule becomes:

$$\epsilon_{t+1}^\vartheta = (1 - \delta_t)\epsilon_t^\vartheta + \delta_t \eta_{t+1}, \tag{38}$$

where $\eta_{t+1} = e^{R(\tau_{t+1})/x} - \vartheta^*(x,\theta)$ is a martingale difference sequence.

Square both sides and take conditional expectations:

$$\mathbb{E}\left[(\epsilon_{t+1}^\vartheta)^2 \mid \mathcal{F}_t \right] = (1 - \delta_t)^2 (\epsilon_t^\vartheta)^2 + \delta_t^2 \mathbb{E}[\eta_{t+1}^2]$$
$$\leqslant \left(1 - 2\delta_t + \delta_t^2 \right) (\epsilon_t^\vartheta)^2 + \delta_t^2 \nu^2 \tag{39}$$

where $\nu^2 = \mathbb{E}[\eta_{t+1}^2] \leqslant e^{\frac{2R_\infty}{(1-\gamma)x_{min}}}.$

Taking total expectations and using $\delta_t = \frac{c}{t}$:

$$\mathbb{E}[(\epsilon_{t+1}^\vartheta)^2] \leq \left(1 - \frac{2c}{t}\right)\mathbb{E}[(\epsilon_t^\vartheta)^2] + \frac{c^2\nu^2}{t^2} \tag{40}$$

Unroll the recursion from $t = 1$ to T:

$$\mathbb{E}[(\epsilon_T^\vartheta)^2] \leq \mathbb{E}[(\epsilon_1^\vartheta)^2]\prod_{t=1}^{T-1}\left(1 - \frac{2c}{t}\right) + c^2\nu^2\sum_{t=1}^{T-1}\frac{1}{t^2}\prod_{k=t+1}^{T-1}\left(1 - \frac{2c}{k}\right) \tag{41}$$

For $x = \frac{2c}{k}$ (assuming $2c < k$ to ensure $1 - \frac{2c}{k} > 0$). Now apply $1 - x \leq e^{-x}$, to obtain

$$\prod_{k=1}^{T}\left(1 - \frac{2c}{k}\right) \leq \prod_{k=1}^{T}e^{-2c/k} = e^{-2c\sum_{k=1}^{T}\frac{1}{k}}.$$

For large T:

$$\sum_{k=1}^{T}\frac{1}{k} \approx \ln T + a,$$

where $a \approx 0.577$ is the Euler-Mascheroni constant. Substitute into the exponential, to obtain:

$$e^{-2c\sum_{k=1}^{T}\frac{1}{k}} \approx e^{-2c(\ln T + a)} = e^{-2ca} \cdot T^{-2c}.$$

Let $C = e^{-2ca}$. Then:

$$\prod_{k=1}^{T}\left(1 - \frac{2c}{k}\right) \leq \frac{C}{T^{2c}}, \text{ and } \prod_{k=t+1}^{T-1}\left(1 - \frac{2c}{k}\right) \leq \frac{C}{(T/t)^{2c}} \tag{42}$$

Combining these results, we get:

$$\mathbb{E}[(\epsilon_T^\vartheta)^2] \leq \frac{C}{T^{2c}} + c^2\nu^2\sum_{t=1}^{T-1}\frac{1}{t^2}\frac{C}{(T/t)^{2c}}$$

$$\leq \frac{C}{T^{2c}} + \frac{Cc^2\nu^2}{T^{2c}}\sum_{t=1}^{\infty}t^{2c-2} \leq \frac{K_1}{T} \tag{43}$$

where $K_1 = C(1 + c^2\nu^2\zeta(2 - 2c))$ and ζ is the Riemann zeta function. $\qquad\square$

Lemma 7. *For a given policy π_θ, let the step size $\delta_t = \frac{c}{t}$ with $c > \frac{1}{2x_{\min}\mathbb{E}_{\tau\sim\pi_\theta}\left[e^{R(\tau)/x}\right]}$. Then, the stochastic variable ω_t satisfies:*

$$\mathbb{E}\left[|\omega_t - \omega^*(x,\theta)|^2\right] \leq \frac{K_2}{t},$$

where $\omega^*(x,\theta) = \frac{\mathbb{E}_{\tau\sim\pi_\theta}[R(\tau)e^{R(\tau)/x}]}{x\mathbb{E}_{\tau\sim\pi_\theta}[e^{R(\tau)/x}]}$, and $K_2 = C\left(1 + c^2\sigma_\omega^2\frac{\pi^2}{6}\right)$ with $\sigma_\omega^2 = \frac{R_\infty^2 e^{2R\infty/(x_{\min}(1-\gamma))}}{x_{\min}^2(1-\gamma)^2}$, and $C > 0$ is some constant.

Proof. Define the error $\epsilon_t^\omega := \omega_t - \omega^*(x,\theta)$. The update rule of ω_t becomes:

$$\epsilon_{t+1}^\omega = \epsilon_t^\omega\left(1 - \delta_t x e^{R(\tau_{t+1})/x}\right) + \delta_t e^{R(\tau_{t+1})/x}\left(R(\tau_{t+1}) - x\omega^*(x)\right).$$

Square both sides and take expectations, we obtain:

$$\mathbb{E}\left[(\epsilon_{t+1}^\omega)^2\right] = \mathbb{E}_\tau\left[\left(1 - \delta_t x e^{R(\tau)/x}\right)^2\right]\mathbb{E}\left[(\epsilon_t^\omega)^2\right] + \delta_t^2\mathbb{E}_\tau\left[e^{2R(\tau)/x}\left(R(\tau) - x\omega^*(x)\right)^2\right].$$

Using $|R| \leqslant \frac{R_\infty}{1-\gamma}$ and $x \geqslant x_{\min}$, we bound:

$$\mathbb{E}_\tau \left[e^{2R(\tau)/x} \left(R(\tau) - x\omega^*(x)\right)^2 \right] \leqslant \frac{R_\infty^2 e^{2R_\infty/(x_{\min}(1-\gamma))}}{(1-\gamma)^2} = \sigma_\omega^2.$$

Substitute $\delta_t = \frac{c}{t}$ and approximate $\mathbb{E}_\tau \left[\left(1 - \delta_t x e^{R(\tau)/x}\right)^2 \right] \approx 1 - \frac{2cx\mathbb{E}_\tau\left[e^{R(\tau)/x}\right]}{t}$:

$$\mathbb{E}\left[(\epsilon_{t+1}^\omega)^2\right] \leqslant \left(1 - \frac{\lambda}{t}\right) \mathbb{E}\left[(\epsilon_t^\omega)^2\right] + \frac{c^2 \sigma_\omega^2}{t^2},$$

where $\lambda = 2cx\mathbb{E}_\tau[e^{R(\tau)/x}] \geqslant 2cx_{\min}e^{-R_\infty/(x_{\max}(1-\gamma))}$.

Unroll the recursion from $t = 1$ to T:

$$\mathbb{E}\left[(\epsilon_T^\omega)^2\right] \leqslant \mathbb{E}\left[(\epsilon_1^\omega)^2\right] \prod_{t=1}^{T-1} \left(1 - \frac{\lambda}{t}\right) + c^2\sigma_\omega^2 \sum_{t=1}^{T-1} \frac{1}{t^2} \prod_{k=t+1}^{T-1} \left(1 - \frac{\lambda}{k}\right).$$

Using the inequality $\prod_{k=1}^{T} \left(1 - \frac{\lambda}{k}\right) \leqslant \frac{C}{T^\lambda}$, and for $\lambda \geqslant 1$ we get:

$$\mathbb{E}\left[(\epsilon_T^\omega)^2\right] \leqslant \frac{C}{T^\lambda} \left(\mathbb{E}\left[(\epsilon_1^\omega)^2\right] + c^2\sigma_\omega^2 \frac{\pi^2}{6} \right).$$

Hence,

$$\mathbb{E}\left[|\omega_t - \omega^*(x)|^2\right] \leqslant \frac{K}{t}.$$

$\qquad$ □

Theorem 5. *Let step sizes $\delta_t = \frac{c}{t}$, $\xi_t = \frac{d}{t^b}$ with $b \in (0.5, 1)$. Under the Assumption 1, the iterates $x_t \in \mathbb{R}$ satisfy:*

$$\mathbb{E}\left[(x_t - x^*)^2\right] \leqslant \frac{K}{t^b},$$

where $x^* = \arg\min_x G(x)$, and the constant $K > 0$ is explicitly given by:

$$K = \frac{x_{\max}^6(1-b)^2 \left(2e^{\frac{2R_\infty}{((1-\gamma)x_{\min})}} K_1 + 2K_2 + \left(\frac{2R_\infty}{x_{\min}(1-\gamma)} + |\log\alpha| \right)^2 \right)}{4\bar{\sigma}^2}.$$

Proof. Define $V_t = |x_t - x^*|^2$. The update rule is:

$$x_{t+1} = x_t - \xi_t \left(G'(x_t) + \eta_t \right),$$

where the gradient estimation error $\eta_t = \log \frac{\vartheta_t}{\vartheta^*(x)} - (\omega_t - \omega^*(x))$

Using the Lipschitz continuity of $\log(\cdot)$ near $\vartheta^*(x)$:

$$\left| \log \frac{\vartheta_t}{\vartheta^*(x)} \right| \leqslant L_1 |\epsilon_t^\vartheta|,$$

where $L_1 = \frac{1}{\inf_x \vartheta^*(x)} \leqslant e^{R_\infty/((1-\gamma)x_{\min})}$. Then:

$$\|\eta_t\| \leqslant L_1 \|\epsilon_t^\vartheta\| + \|\epsilon_t^\omega\|.$$

Squaring and taking expectations:

$$\mathbb{E}[\|\eta_t\|^2] \leqslant 2L_1^2 \mathbb{E}[\|\epsilon_t^\vartheta\|^2] + 2\mathbb{E}[\|\epsilon_t^\omega\|^2] \leqslant \frac{2L_1^2 K_1 + 2K_2}{t}.$$

Let $C = 2L_1^2 K_1 + 2K_2$. Expand $V_{t+1} = (x_t - \xi_t(G'(x_t) + \eta_t) - x^*)^2$. Taking conditional expectations:

$$\mathbb{E}[V_{t+1} \mid \mathcal{F}_t] \leqslant V_t - 2\xi_t G'(x_t)(x_t - x^*) + \xi_t^2 \mathbb{E}\left[(G'(x_t) + \eta_t)^2 \mid \mathcal{F}_t\right].$$

Using strong convexity $G'(x_t)(x_t - x^*) \geqslant \mu V_t$ with $\mu = \frac{\bar{\sigma}}{x_{\max}^3}$ and $|G'(x)| \leqslant \frac{2R_\infty}{x_{\min}(1-\gamma)} + |\log \alpha| \triangleq L$:

$$\mathbb{E}[V_{t+1}] \leqslant (1 - 2\mu\xi_t)\,\mathbb{E}[V_t] + \xi_t^2\left(L^2 + \frac{C}{t}\right),$$

Substitute $\xi_t = \frac{d}{t^b}$:

$$\mathbb{E}[V_{t+1}] \leqslant \left(1 - \frac{2\bar{\sigma}d}{x_{\max}^3 t^b}\right)\mathbb{E}[V_t] + \frac{d^2}{t^{2b}}\left(L^2 + \frac{C}{t}\right).$$

Unroll recursively for $T \geqslant 1$:

$$\mathbb{E}[V_T] \leqslant \mathbb{E}[V_1] \prod_{t=1}^{T-1}\left(1 - \frac{2\bar{\sigma}d}{x_{\max}^3 t^b}\right) + d^2 \sum_{t=1}^{T-1} \frac{L^2 + C/t}{t^{2b}} \prod_{k=t+1}^{T-1}\left(1 - \frac{2\bar{\sigma}d}{x_{\max}^3 k^b}\right).$$

Using $1 - x \leqslant e^{-x}$:

$$\prod_{t=1}^{T-1}\left(1 - \frac{2\bar{\sigma}d}{x_{\max}^3 t^b}\right) \leqslant \exp\left(-\frac{2\bar{\sigma}d}{x_{\max}^3}\sum_{t=1}^{T-1}\frac{1}{t^b}\right).$$

Approximate $\sum_{t=1}^{T-1}\frac{1}{t^b} \geqslant \frac{T^{1-b}}{1-b}$:

$$\exp\left(-\frac{2\bar{\sigma}d}{x_{\max}^3} \cdot \frac{T^{1-b}}{1-b}\right) \leqslant \frac{1}{T^b}.$$

Choose $d = \frac{x_{\max}^3(1-b)}{2\bar{\sigma}} \cdot b$. The noise term becomes:

$$d^2 \sum_{t=1}^{\infty} \frac{L^2 + C}{t^b} \leqslant \frac{d^2(L^2 + C)}{b}.$$

Substituting d:

$$K = \frac{x_{\max}^6(1-b)^2\left(2L_1^2 K_1 + 2K_2 + \left(\frac{2R_\infty}{x_{\min}(1-\gamma)} + |\log \alpha|\right)^2\right)}{4\bar{\sigma}^2}.$$

$\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad\qquad$ $\square$

A.3 Proof of Theorem 4

Recall the definition,

$$\widehat{\nabla J}_{\text{EVaR}}(\theta_t) = \frac{J_{\text{EVaR}}(\theta_t + c_t\Delta_t) - J_{\text{EVaR}}(\theta_t - c_t\Delta_t)}{2c_t\Delta_t}, \tag{44}$$

where $z(\theta_t^+) = G(x_t^+) - J_{\text{EVaR}}(\theta_t^+)$ and $z(\theta_t^-) = G(x_t^-) - J_{\text{EVaR}}(\theta_t^-)$.

Lemma 8. *Let* $J_{\text{EVaR}}^{(3)}(\theta) \equiv \partial^3 J_{\text{EVaR}}/\partial\theta^T\partial\theta^T\partial\theta^T$ *exist and* $\max_{i_1,i_2,i_3} \sup_\theta \|EVaR_{\alpha_{i_1 i_2 i_3}}^{(3)}(\theta)\|_\infty \leqslant \epsilon$. *Then* $\forall \theta \in$ *interior*(Θ)

$$b_t(\theta_t) = \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] = \mathcal{O}(c_t^2).$$

Proof. By the continuity of $J_{\text{EVaR}}^{(3)}$ and Δ_t being a Bernoulli random variable, we have, by Taylor's theorem,

$$J_{\text{EVaR}}(\theta_t + c_t\Delta_t) \approx J_{\text{EVaR}}(\theta_t) + c_t\Delta_t^\top \nabla J_{\text{EVaR}}(\theta_t) + \frac{c_t J_{\text{EVaR}}^2}{2!}\Delta_t^\top \nabla^2 \text{EVaR}_\alpha(\theta_t)\Delta_t + \frac{c_t^3}{3!}\nabla^3 J_{\text{EVaR}}(\theta_t)\Delta_t \otimes \Delta_t \otimes \Delta_t,$$

where $\bar{\theta}_t$ lies on the line segment between θ_t and $\theta_t + c_t\Delta_t$. Hence,

$$\frac{J_{\text{EVaR}}(\theta_t + c_t\Delta_t) - J_{\text{EVaR}}(\theta_t - c_t\Delta_t)}{2c_t\Delta_t} = \frac{\Delta_t^\top}{\Delta_t}\nabla J_{\text{EVaR}}(\theta_t) + \frac{c_t^2}{12\Delta_t}\nabla^3\left(J_{\text{EVaR}}(\bar{\theta}_t) + J_{\text{EVaR}}(\bar{\theta}_t')\right)\Delta_t \otimes \Delta_t \otimes \Delta_t$$

Now,

$$b_t(\theta_t) = \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right]$$

$$= \mathbb{E}\left[\frac{J_{\text{EVaR}}(\theta_t + c_t\Delta_t) - J_{\text{EVaR}}(\theta_t - c_t\Delta_t)}{2c_t\Delta_t} - \nabla J_{\text{EVaR}}(\theta_t)\Big|\mathcal{F}_t\right].$$

Let b_{t_l} denote the l^{th} term of the bias vector b_t. Then

$$b_{t_l} = \mathbb{E}\left[\frac{\Delta_{t_l}}{\Delta_{t_l}}\nabla J_{\text{EVaR}}(\theta_t) + \frac{c_t^2}{12\Delta_{t_l}}\nabla^3\left(J_{\text{EVaR}}(\bar{\theta}_t) + J_{\text{EVaR}}(\bar{\theta}_t')\right)\Delta_t \otimes \Delta_t \otimes \Delta_t - \nabla_\ell J_{\text{EVaR}}(\theta_t)\Big|\mathcal{F}_t\right] \quad (45)$$

where $\bar{\theta}_t'$ lies on the line segment between θ_t and $\theta_t - c_t\Delta_t$. Now note that,

$$\mathbb{E}\left[\Delta_t^{-1}\Delta_t^\top \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] = (\nabla J_{\text{EVaR}}(\theta_t))_1 \mathbb{E}\left[\Delta_t^{-1}\Delta_{t_1} \mid \mathcal{F}_t\right] + \cdots + (\nabla J_{\text{EVaR}}(\theta_t))_p \mathbb{E}\left[\Delta_t^{-1}\Delta_{t_p} \mid \mathcal{F}_t\right]$$

$$\text{(Since, } \nabla J_{\text{EVaR}}(\theta_t) \text{ is measurable } w.r.t. \ \mathcal{F}_t\text{)}$$

$$= (\nabla J_{\text{EVaR}}(\theta_t))_1 \mathbb{E}\left[\begin{pmatrix} 1 \\ \Delta_{t_2}^{-1}\Delta_{t_1} \\ \vdots \\ \Delta_{t_p}^{-1}\Delta_{t_1} \end{pmatrix} \mid \mathcal{F}_t\right] + \cdots + (\nabla J_{\text{EVaR}}(\theta_t))_p \mathbb{E}\left[\begin{pmatrix} \Delta_{t_1}^{-1}\Delta_{t_p} \\ \Delta_{t_2}^{-1}\Delta_{t_p} \\ \vdots \\ 1 \end{pmatrix} \mid \mathcal{F}_t\right]$$

$$= (\nabla J_{\text{EVaR}}(\theta_t))_1 \begin{pmatrix} 1 \\ \mathbb{E}\Delta_{t_2}^{-1}\mathbb{E}\Delta_{t_1} \\ \vdots \\ \mathbb{E}\Delta_{t_p}^{-1}\mathbb{E}\Delta_{t_1} \end{pmatrix} + \cdots + (\nabla J_{\text{EVaR}}(\theta_t))_p \begin{pmatrix} \mathbb{E}\Delta_{t_1}^{-1}\mathbb{E}\Delta_{t_p} \\ \mathbb{E}\Delta_{t_2}^{-1}\mathbb{E}\Delta_{t_p} \\ \vdots \\ 1 \end{pmatrix}$$

$$\text{(Since, } \mathbb{E}\Delta_{t_i} = 0, \forall i \in [1 \ldots p] \text{ and } \Delta_{t_i} \text{ is independent of } \Delta_{t_j} \forall i \neq j\text{)}$$

$$= (\nabla J_{\text{EVaR}}(\theta_t))_1 \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \cdots + (\nabla J_{\text{EVaR}}(\theta_t))_p \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{pmatrix} = \nabla J_{\text{EVaR}}(\theta_t) \quad (46)$$

Therefore, from Eq. 45 and Eq. 46, we get,

$$b_{t_l} = \frac{1}{12}\mathbb{E}\left[\frac{1}{\Delta_{t_l}}\left(\nabla^3 J_{\text{EVaR}}(\bar{\theta}_t) + \nabla^3 J_{\text{EVaR}}(\bar{\theta}_t')\right)\bar{\Delta}_t \otimes \bar{\Delta}_t \otimes \bar{\Delta}_t \mid \mathcal{F}_t\right] \quad (47)$$

We can bound the term on the right-hand side of eq. 47 in magnitude as follows:

$$b_t(\theta_t) = \frac{1}{12}\mathbb{E}\left[\frac{1}{\Delta_{t_l}}\left(\nabla^3 J_{\text{EVaR}}(\bar{\theta}_t) + \nabla^3 J_{\text{EVaR}}(\bar{\theta}_t')\right)\bar{\Delta}_t \otimes \bar{\Delta}_t \otimes \bar{\Delta}_t \,\Big|\, \mathcal{F}_t\right]$$

$$\leqslant \frac{\epsilon c_t^2}{6}\sum_{i_1}\sum_{i_2}\sum_{i_3}\mathbb{E}\left[\frac{\Delta_{t_{i_1}}\Delta_{t_{i_2}}\Delta_{t_{i_3}}}{\Delta_{k_l}}\right] \leqslant \frac{p^3\epsilon c_t^2}{6} = \mathcal{O}(c_t^2). \quad (48)$$

The first inequality follows as $\nabla^3 J_{\text{EVaR}}(\bar{\theta}) \leqslant \epsilon, \forall \theta$ and the latter inequality follows since $\frac{\Delta_{t_{i_1}}\Delta_{t_{i_2}}\Delta_{t_{i_3}}}{\Delta_{t_l}} \leqslant 1$. $\quad\square$

Proof of Theorem 4

Proof. Consider the recursion from Step 12 of the algorithm:

$$\theta_{t+1} = \theta_t + a_t \left(\underbrace{\mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right]}_{b_t} - \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \nabla J_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] + \right.$$

$$\left. \widehat{\nabla J}_{\text{EVaR}}(\theta_t) + \frac{z(\theta_t^+) - z(\theta_t^-)}{2c_t \Delta_t} \right)$$

$$= \theta_t + a_t \left(b_t + \varphi_t + \nabla J_{\text{EVaR}}(\theta_t) + \widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] \right)$$

$$= \theta_t + a_t \left(b_t + e_t + \varphi_t + \nabla J_{\text{EVaR}}(\theta_t) \right), \text{ where } e_t = \widehat{\nabla J}_{\text{EVaR}}(\theta_t) - \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right]$$

$$\text{and } \varphi_t = \frac{z(\theta_t^+) - z(\theta_t^-)}{2c_t \Delta_t}. \tag{49}$$

Then

$$\mathbb{E}\left[|\varphi_t||\mathcal{F}_t\right] = \mathbb{E}\left[\left|\frac{z(\theta_t^+) - z(\theta_t^-)}{2c_t \Delta_t}\right| \Big| \mathcal{F}_t\right] \leqslant \frac{1}{2c_t}\mathbb{E}\left[\Delta_t^{-1}(z(\theta_t^+) - z(\theta_t^-))|\mathcal{F}_t\right]$$

$$\leqslant \frac{1}{2c_t}\mathbb{E}\left[|\Delta_t^{-1}|(|z(\theta_t^+)| + |z(\theta_t^-)|)|\mathcal{F}_t\right]$$

$$= \frac{1}{2c_t}\mathbb{E}\left[|\Delta_t^{-1}||z(\theta_t^+)||\mathcal{F}_t\right] + \mathbb{E}\left[|\Delta_t^{-1}||z(\theta_t^-)||\mathcal{F}_t\right]$$

$$\leqslant \frac{L_G}{2c_t}\mathbb{E}\left[|\Delta_t^{-1}||x_t^+ - x^*(\theta_t^+)||\mathcal{F}_t\right] + \frac{L_G}{2c_t}\mathbb{E}\left[|\Delta_t^{-1}||x_t^- - x^*(\theta_t^-)||\mathcal{F}_t\right] \quad \text{(Since } G \text{ is Lipschitz}$$

$$\text{with constant } L_G \text{ which follows from Theorem 1)}$$

$$\leqslant \frac{1}{2c_t}\left(\underbrace{\mathbb{E}\left[\Delta_t^{-2}\right]^{\frac{1}{2}}}_{=1}\mathbb{E}\left[|x_t^+ - x^*(\theta_t^+)|^2|\mathcal{F}_t\right]^{\frac{1}{2}} + \underbrace{\mathbb{E}\left[\Delta_t^{-2}\right]^{\frac{1}{2}}}_{=1}\mathbb{E}\left[|x_t^- - x^*(\theta_t^-)|^2|\mathcal{F}_t\right]^{\frac{1}{2}} \right)$$

$$\text{(by Cauchy-Schwartz Inequality)}$$

$$\leqslant \frac{\sqrt{K}}{c_t N_t^{\frac{b}{2}}} \quad \text{(by Theorem 5)}$$

Now by the Monotone Convergence Theorem, we have

$$\mathbb{E}\left[\sum_{t \geqslant 1} a_t|\varphi_t|\right] = \sum_{t \geqslant 1} a_t \mathbb{E}\left[|\varphi_t|\right] \leqslant \sum_{t \geqslant 1} \frac{a_t\sqrt{K}}{c_t N_t^{\frac{b}{2}}} < \infty.$$

Therefore

$$\mathbb{P}\left(\sum_{t \geqslant 1} a_t|\varphi_t| < \infty\right) = 1 \quad \Rightarrow \sum_{t \geqslant 1} a_t\varphi_t < \infty \quad a.s. \quad \Rightarrow \sum_{t \geqslant k} a_t\varphi_t \xrightarrow[k \to 0]{} 0. \tag{50}$$

Define

$$\xi_{t+1} = \sum_{i=0}^{t} a_t e_t, t \geqslant 0. \tag{51}$$

Then

$$\mathbb{E}\left[\xi_{t+1}|\mathcal{F}_t\right] = \mathbb{E}\left[\sum_{i=0}^{t-1} a_t e_t \Big| \mathcal{F}_t\right] = \sum_{i=0}^{t} a_t \mathbb{E}\left[e_t|\mathcal{F}_t\right] + a_t \left(\mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t)|\mathcal{F}_t\right] - \mathbb{E}\left[\widehat{\nabla J}_{\text{EVaR}}(\theta_t) \mid \mathcal{F}_t\right] \right) = \xi_t. \tag{52}$$

This implies that $\{\xi_t\}$ is a martingale with respect to filtration $\{\mathcal{F}_t\}$. Also, since $J_{\texttt{EVaR}}$ is continuously differentiable, we have ξ_t is square-integrable, $\forall t$, *i.e*, $\mathbb{E}\left[\|\xi_t\|^2\right] < \infty$, $\forall t$. Again, by the continuous differentiability of $J_{\texttt{EVaR}}$ and Assumption 4, we obtain

$$\sum_t \mathbb{E}\left[\|\xi_{t+1} - \xi_t\|^2 | \mathcal{F}_t\right] = \sum_t a_t^2 \mathbb{E}\left[\|e_t\|^2\right] < \infty \text{ on the set } \{\sup_t \|\theta_t\| < \infty\}.$$

Therefore, by the Martingale convergence theorem, we get

$$\lim_{t \to \infty} \xi_t \text{ exists on the event } \{\sup_t \|\theta_t\| < \infty\}. \tag{53}$$

Hence, by Theorem 2, Chapter 2 of Borkar (2008), the asymptotic behavior of the sample paths belonging to the event $\{\sup_t \|\theta_t\| < \infty\}$ is equivalent to the long-term behavior of the dynamical system induced by the ODE

$$\frac{d\theta(t)}{dt} = \nabla J_{\texttt{EVaR}}(\theta(t)), t \geqslant 0. \tag{54}$$

This further implies that the iterates θ_t corresponding to the sample paths belonging to the event $\{\sup_t \|\theta_t\| < \infty\}$ converge to any of the compact transitive invariant sets connected internally in chains of (54). Invariant sets are subsets of the state space that remain unchanged under the flow of the dynamical system. The dynamical system (54) driven by the gradient of the $J_{\texttt{EVaR}}$ is a gradient flow where the only possible invariant sets are the subsets of $H = \{\theta | \nabla \texttt{EVaR}_\alpha(\theta) = 0\}$ (Lemma 1, Section 10.2 of Borkar (2008)). Further, by invoking the LaSalle invariance principle and the Lyapunov theorem, one can obtain that the asymptotically stable points inside H are given by $\{\theta \in H | \nabla^2 J_{\texttt{EVaR}}(\theta) \preccurlyeq 0\}$. □

B Comparison of EVaR estimation methods

One can also estimate `EVaR` of discounted cumulative rewards for sample trajectory using a disciplined convex programming characterization as stated by Cajas (2021) as follows

$$J_{\texttt{EVaR}}(\theta) = \min_{\beta, t, u}\ t - \beta \ln\left(\alpha N\right), \tag{55}$$

$$\text{subject to } \beta \geqslant \sum_{j=1}^N u_j \text{ and } (J_{\texttt{EVaR}}(\theta)_j - t, \beta, u_j) \in K_{\exp}, \forall j \in [1, N]$$

where $\{J_{\texttt{EVaR}}(\theta)_j\}_{j=1}^N$ are N realizations of $J_{\texttt{EVaR}}(\theta)$, β, t and u are variables and $K_{\exp}$ is an exponential cone (Chares, 2009) which is defined as follows

$$K_{\exp} = \left\{(a, b, c) \mid b > 0, c \geqslant b \exp\left(\frac{a}{b}\right)\right\} \cup \{(a, b, c) \mid a \leqslant 0, b = 0, c \geqslant 0\}. \tag{56}$$

The above optimization can be solved efficiently to any desired level of accuracy via interior-point methods due to the existence of computationally tractable barrier functions which enables the efficient exploration of the solution space (Chandrasekaran & Shah, 2017). In Algorithm 2 we give an `EVaR` optimization algorithm where the `EVaR` estimate computation is done using the disciplined convex cone method by using solving the optimization problem. This requires access to N trajectories for the computation and which decides the desired accuracy of the estimate. This process is not an online process with respect to the estimation and the movement of the samples have little influence over the `EVaR` estimate. When this process is compared with Stochastic Approximation (SA) version of `EVaR` estimation we see in Figure 14 is more resilient to changing environment dynamics and a higher error band justifies the online nature and adaptability to the samples.

Remark. *The rewards for the environment is negated to comply with structure of the solver in Algorithm 2.*

Figure 14: `EVaR` estimation using two-time scale schotastic approximation where the blue represents the estimate using the convex optimization and the orange band represents the estimate using SA method. Run for 10 batches with CartPole as the environment with episode length of 100.

Algorithm 2 `EVaR` Optimization using Disciplied Convex Cone

Require: risk level $\alpha \in (0,1)$, initial $\theta_0 \in \mathbb{R}^p$, step-sizes $\{a_t, c_t, \delta_t, \xi_t\}$, inner lengths N_t
1: Initialize policy network (π_θ) parameters $\theta \in \Theta$
2: **for** $t = 0, \ldots, T-1$ **do**
3:     Draw $\Delta_t \in \{\pm 1\}^p$ IID.
4:     $\theta_t^+ \leftarrow \theta_t + c_t \Delta_t, \quad \theta_t^- \leftarrow \theta_t - c_t \Delta_t$
5:     **for** $k = 1, \ldots, N_t$ **do**   ▷ `EVaR estimation for "+"`
6:         `Sample trajectory` $\tau_{t,k}^+ \sim \pi_{\theta_t^+}$, `compute` $R_{t,k}^+ = \sum_{u=0}^H \gamma^u r_u$
7:     **end for**
8:     Solve using Interior point method

$$G_{t,j}^+ \leftarrow \texttt{Solve}\Big(\min_{\beta,t,u} \ t - \beta \ln(\alpha N), \ \texttt{subject to} \ \beta \geqslant \sum_{j=1}^N u_j$$

$$\texttt{and} \ \Big(R_{t,k}^+ - t, \beta, u_j \Big) \in K_{\exp}, \forall j = 1, \ldots, N \Big)$$

9:     **for** $k = 1, \ldots, N_t$ **do**   ▷ `EVaR estimation for "-"`
10:         `Sample trajectory` $\tau_{t,k}^- \sim \pi_{\theta_t^-}$, `compute` $R_{t,k}^- = \sum_{u=0}^H \gamma^u r_u$
11:     **end for**
12:     Solve using Interior point method

$$G_{t,j}^- \leftarrow \texttt{Solve}\Big(\min_{\beta,t,u} \ t - \beta \ln(\alpha N), \ \texttt{subject to} \ \beta \geqslant \sum_{j=1}^N u_j$$

$$\texttt{and} \ \Big(R_{t,k}^- - t, \beta, u_j \Big) \in K_{\exp}, \forall j = 1, \ldots, N \Big)$$

13:     $\hat{g}_t \leftarrow \dfrac{G_t^+ - G_t^-}{2\,c_t} \Delta_t^{-1}$
14:     $\theta_{t+1} \leftarrow \theta_t + a_t \hat{g}_t$
15: **end for**
16: **return** θ_T

C Experiments Detail

C.1 Finite Difference Gradient Estimation

The hyperparameters used in our finite difference-based gradient estimation have a significant impact on the algorithm's performance. The timeout parameter controls the time allotted for each function evaluation, ensuring efficient computation. Iterations govern the number of optimization steps, balancing convergence speed and accuracy. Learning rate decay and power adjust the step size dynamically, allowing the learning process to slow down as the model converges. Perturbation size affects the extent of exploration in the gradient estimation, while its decay and power ensure that perturbations shrink over time, refining the gradient's precision. Momentum helps maintain stable updates by incorporating past gradients, and the Adam parameters (Beta and Epsilon) enhance robustness, particularly in the face of noisy gradient estimates, providing smoother and more reliable updates. Together, these hyperparameters create a dynamic and adaptable optimization process, crucial for navigating uncertain and noisy environments.

Hyperparameter	Value
Timeout	1×10^{-4}
Iterations	10,000
Learning Rate Decay	1×10^{-3}
Learning Rate Power	0.5
Perturbation Size (px)	2.0
Perturbation Decay	1×10^{-2}
Perturbation Power	0.161
Momentum	0.9
Beta (Adam Parameter)	0.999
Epsilon (Adam Parameter)	1×10^{-7}

Table 2: Hyperparameters for Finite Difference Gradient Estimation

C.2 MuJoCo and Gridworld

We evaluate our proposed algorithm on various continuous control tasks from the OpenAI Gym suite Brockman et al. (2016), where we augment the environment with random normal noise to introduce uncertainty. For the discrete setting, we use a custom GridWorld environment with randomly placed obstacles covering approximately 30% of the grid. Importantly, we ensure that a path always exists from the start to the goal state, even with obstacles. The hyperparameters used across all experiments are summarized below :

Hyperparameter	Value
Learning rate	$a_k = \frac{a}{(k+1+A)^{0.602}}, c_k = \frac{c}{(k+1)^{0.101}}$
Constant A	10^{-7}
Constant c	0.999
Random noise parameter δ	$1 - \alpha$
Action sampling	$a_t \sim \pi_\theta(\cdot|s) + \delta\mathcal{N}(0,1)$
Step size δ	0.12
Step size ξ	3×10^{-10}

Table 3: Hyperparameters used in experiments

Remark. *The confidence value α that is used for `EVaR` is flipped i.e $1 - \alpha$ is used for the same experiment for `CVaR` and `VaR` estimates to align the measures in the direction of the upward risk which is used for most reward distributions.*

Figure 15: Comparison of EVaR, CVaR and VaR at $\alpha \in \{0.9, 0.95\}$ for *InvertedDoublePendulum-v4*, the left (a) and (c) represents the rewards received following the EVaR policy and (b) and (d) shows the maximization of EVaR policy.

Figure 16: On the left column (a) and (b) shows the maximization of the objective values and the cumulative returns received following EVaR policy at $\alpha \in \{0.9, 0.95, 0.99\}$ respectively and on the right column (c) and (d) visualizes comparision of EVar, CVaR and VaR for objective and return.

Figure 17: Comparison of EVaR, CVaR and VaR at $\alpha = 0.9$ for Swimmer, the left (a) and (c) represents the rewards received following the EVaR policy and (b) and (d) shows the maximization of EVaR policy.

C.3 Glycemic Control

The environment is modelled as a MDP is such that the state space consisting multiple noisy glucose measurements at various time points in the past, the carbohydrate intakes and other relevant information pertaining to the patient. The action is the amount of insulin to be delivered to the patient also a scalar which indicates the insulin to be administered. We observe from our experiments our algorithm is able to keep the patient in admissible levels of risk. There are instances where the glucose breaches into the hypo or hyper region the policy course corrects to maintain the stable condition.

Figure 18: In plots (a),(b),(c) and (d) represents the patient profile of *adoloscent001* and the rest of the plots represent the patient *adult002*. Both patients show up to be stable and alive under the influence of the administration of the insulin via the controller where the signals are optimized by the EVaR optimizer.

We show that our algorithm is equipped to handle high risk scenarios like administrating insulin to Type-1 diabetes patients of type *adult#002, adolescent#001* registered on the Gym environment using the simulator developed by Xie (2018) for RL control tasks. The simulator emulates a PID controller that provides the insulin to maintain the Blood Glucose levels which is the action for the agent and we try to optimize by increasing the longevity of the patient using the finite difference `EVaR` optimization. The patient profiles are given below where, CF: Carbohydrate Factor, CR: Carb Ratio (often referred to as Carb-to-Insulin Ratio) and TDI: Total Daily Insulin.

Name	CR	CF	Age	TDI
adolescent#001	12	15.0360	18	36.7339
adult#002	8	9.2128	65	57.8688

Table 4: Patient profiles

C.4 Portfolio Optimization

Problem Definition : We focus on the top 10 US DJIA stocks—[`KO, AAPL, MSFT, JPM, WMT, UNH, V, PG, JNJ, HD`]—using market data from January 1, 2014, to January 1, 2024, which provides a decade of market information encompassing various economic cycles. Our custom `PortfolioEnv` class, designed to adhere to the OpenAI Gym interface (Brockman et al., 2016) and compatible with Stable Baselines 3 (Hill et al., 2018), represents a 50-day window of historical returns as a matrix $\mathbf{O} \in \mathbb{R}^{50 \times 10}$. To minimize the `EVaR` value of the negative returns, which in our case are the portfolio values represented as $\mathcal{R}$ under the constraints that the weights of the portfolio always sum up to 1 after each allocation step. We find the optimal EVaR portfolio with N-assets represented by the optimization problem:

$$\max_{\beta > 0} \ \beta^{-1} \log\left(\alpha^{-1} \mathbb{E}[e^{\beta \mathcal{R}}]\right) \tag{57}$$

$$\text{s.t.} \ \sum_{i=1}^{N} w_i = 1 \tag{58}$$

$$\beta > 0, w \geqslant 0 \tag{59}$$

here the weights of the N assets in the portfolio is represented by the weight vector $\boldsymbol{w} = [w_1, \dots, w_n]^\top$ such that $\sum_{i=1}^{N} w_i = 1$, where $0 \leqslant w_i \leqslant 1$. An asset weight of 0 indicates zero holdings of a particular asset in a portfolio, whereas a weight of 1 means that the entire portfolio is concentrated in the said asset.

Action Space : For each action $A \in \mathcal{A}$, the action A represents portfolio weights for portfolio allocation on N assets. The weight constraint makes each action represented as $A = [a_1, \dots, a_N]^\top$ so that $\sum_{i=1}^{N} a_i = 1$ where $0 \leqslant a_i \leqslant 1$. In alternative implementations of this framework, $a_i < 0$ would permit short-selling an asset, while $a_i > 1$ would allow for leveraged positions. However, in our scenario, we limit the actions to non-leveraged, long-only positions. To enforce these constraints, we apply the softmax function to the agent's continuous actions.

State Space : The state space $\mathcal{S}$ is represented by a matrix $S_t \in \mathbb{R}^{w \times n}$, where w is a predefined window size and n is the number of assets. Each element $s_{i,j}$ of the matrix represents the return of the asset j at the time step $t - w + i$. Formally:

$$S_t = \{r_{i,j} | i \in [t - w + 1, t], j \in [1, n]\} \tag{60}$$

where $r_{i,j}$ is the return of asset j at time i. The observation space is bounded, with $S_t \in [-1, 1]^{w \times n}$.

At each time step t, the state S_t is represented by a matrix $S_t \in \mathbb{R}^{w \times n}$, where:

- w is the window size (number of historical time steps considered)

- n is the number of assets in the portfolio

Formally, we define the state matrix as follows:

$$S_t = \begin{bmatrix} r_{t-w+1,1} & r_{t-w+1,2} & \cdots & r_{t-w+1,n} \\ r_{t-w+2,1} & r_{t-w+2,2} & \cdots & r_{t-w+2,n} \\ \vdots & \vdots & \ddots & \vdots \\ r_{t,1} & r_{t,2} & \cdots & r_{t,n} \end{bmatrix} \tag{61}$$

where $r_{i,j}$ represents the return of asset j at time step i. Each element $s_{i,j}$ of the matrix S_t corresponds to the return of a specific asset at a specific time:

$$s_{i,j} = r_{t-w+i,j} = \frac{P_{t-w+i,j} - P_{t-w+i-1,j}}{P_{t-w+i-1,j}} \tag{62}$$

where $P_{t,j}$ is the price of asset j at time t. To ensure numerical stability and consistent scale across different assets, we bound the elements of the state matrix $S_t \in [-1,1]^{w \times n}$. The rows of the matrix S_t represent different time steps, with the most recent returns in the bottom row and the oldest returns in the top row. The rows of the matrix S_t represent different time steps, with the most recent returns in the bottom row and the oldest returns in the top row. This structure allows the agent to identify temporal patterns or trends in asset returns potentially.

Reward Function : The reward function is designed to balance portfolio return with risk. We use the Entropic Value at Risk (`EVaR`) as a risk measure. The reward R_t at time t is defined as:

$$R_t = -\texttt{EVaR}_\alpha(\mathcal{R}_t) \tag{63}$$

where the negative portfolio returns are given by $\mathcal{R}_t = (V_0 - V_t)/V_0$ is the portfolio return up to time t, and α is the confidence level for `EVaR` calculation. The negative sign before `EVaR` ensures that minimizing risk corresponds to maximizing reward.

An episode terminates when either:

- The end of the available market data is reached ($t = T$), or

- The portfolio value drops to zero ($V_t \leqslant 0$).

Transistion Dynamics : The environment functions as a comprehensive interface to the market, utilizing a technique known as market replay to traverse historical data. Additionally, it operates as both a broker and an exchange; at each timestep, it processes the agent's actions to rebalance the portfolio according to the latest prices and specified allocations. As the trading day progresses and new price data is acquired, the environment provides these updates to the agent as observations, accompanied by the Entropic Value at Risk (`EVaR`) reward. For the scope of this study, we permit instantaneous rebalancing of the portfolio. Given an action a_t, the environment updates the portfolio value V_t according to:

$$V_{t+1} = V_t \cdot \left(1 + \sum_{i=1}^{n} a_{t,i} \cdot r_{t+1,i} - c \cdot \sum_{i=1}^{n} |a_{t,i} - a_{t-1,i}|\right) \tag{64}$$

where $r_{t+1,i}$ is the return of asset i at time $t+1$, and c is the transaction cost rate which is kept constant in our case. After rebalancing, the environment creates the next state S_{t+1} and proceeds to the next timestep $t+1$. It calculates the new portfolio value based on V_{t+1} and computes the reward R_t which it returns to the agent.

C.4.1 Algorithm

We use Soft Actor Critic Haarnoja et al. (2018) for policy optimization and the incorporate the stochastic recursive updates for the convergence of EVaR sample estimate by augmenting the critic function. We employ

(a) (b)

Figure 19: **(a)** Optimal portfolio allocation obtained by the `EVaR` agent. **(b)** Efficient Frontier of `EVaR` at $\alpha = 0.01$.

the SAC algorithm with its default MLP policy as implemented in Stable Baselines 3. The actor (policy) and critic (value) networks consist of two hidden layers with 256 units each, using ReLU activations. The model is trained for 1000 timesteps, which we found sufficient for convergence in our environment.

Table 5: Hyperparameters used for training

Hyperparameter	Value
Learning rate	0.0003
Buffer size	1,000,000
Batch size	256
Gamma	0.99
Train frequency	TrainFreq(frequency=1, unit=STEP)
Gradient steps	1
Entropy coefficient (ent coef)	auto
Target entropy	-10.0
Tau	0.005
Policy kwargs	{use_sde: False}

We see that our `EVaR` estimation is more robust as it adapts to the samples and error band is consistent which shows a contained variance and provides a more conservative estimate of the sample averages which is desirable in minimal risk portfolio setting. In Figs.(20a,19a,19b) we visualise our portfolio and how it fairs against other risk measures like `VaR` and `CVaR`, where we employ RiskFolioCajas (2024) to generate the plots. The importance of these plots are discussed below.

Date	AAPL	HD	JNJ	JPM	KO	MSFT	PG	UNH	V	WMT
2014-01-03	-2.20%	-0.16%	0.90%	0.77%	-0.49%	-0.67%	-0.11%	0.71%	0.07%	-0.33%
2014-01-06	0.55%	-0.96%	0.52%	0.58%	-0.47%	-2.11%	0.24%	-1.15%	-0.60%	-0.56%
2014-01-07	-0.72%	0.49%	2.12%	-1.15%	0.30%	0.78%	0.97%	3.06%	0.76%	0.31%
⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮	⋮
2024-01-26	-0.90%	1.23%	-0.04%	-0.38%	0.36%	-0.23%	0.33%	1.99%	-1.71%	0.88%
2024-01-29	-0.36%	0.11%	-0.09%	0.26%	0.61%	1.43%	0.01%	0.27%	2.13%	0.47%

Table 6: Snapshot of portfolio assets and their adjusted returns over a decade.

C.4.2 Our Portfolio and Results

Returns histogram : provides a visual representation of the distribution of portfolio returns, allowing for a comprehensive analysis of the performance of our `EVaR` agent compared to other risk measures such as `VaR` and `CVaR`. This comparison is crucial as it reveals how our `EVaR`-based portfolio exhibits a more conservative allocation, which is characterized by a lower frequency of extreme negative returns. By evaluating the shape and spread of the distribution, we can gauge the robustness of our portfolio against adverse market conditions, highlighting the potential benefits of utilizing `EVaR` as a risk measure.

| | (a) | | (b) |

Figure 20: **(a)** Returns histogram with comparison of `VaR`, `CVaR` and `EVaR` which shows `EVaR` portfolio being the most conservative allocation. **(b)** Comparison of various risk measures using a risk adjusted portfolio.

The comparison of various risk measures is essential for understanding the effectiveness of our portfolio strategy in managing risk. By analyzing the performance of the `EVaR` portfolio against traditional measures like `VaR` and `CVaR`, we can illustrate the advantages of adopting an entropic approach to risk assessment. In Figure.(20b) illustrating these comparisons indicate that our `EVaR` strategy not only limits potential losses but also adapts more dynamically to changing market conditions. This adaptability is reflected in the reduced variance of estimated returns, which is desirable for investors seeking minimal risk.

Efficient Frontier : is a key concept in modern portfolio theory, representing the set of optimal portfolios that offer the highest expected return for a given level of risk or the lowest risk for a given level of expected return. By analyzing the Efficient Frontier in the context of our portfolio optimization with `EVaR` measure, we can visually assess the trade-offs between risk and return for various portfolio allocations. In Figure.(19b) showcasing the Efficient Frontier for our `EVaR`-based portfolio indicates that it achieves a favorable balance between risk and return, demonstrating that our optimization strategy effectively identifies portfolios that maximize returns while minimizing downside risk.

D Reproducibility Details

The experiments were conducted on NVIDIA DGX A100 having an AMD EPYC 7742 64-core processor operating at 1.5 GHz $\sim$ 3.39 GHz, with GDDR5 32 GB RAM, NVIDIA A100-SXM4-40gb GPU at 1.41 GHz and memory clocked at 1.21 GHz.The operating CUDA version for PyTorch 1.13.1 is 11.6 for Python version 3.10.13. The supplementary material provided has all experiments with their obtained values that have been reported here in visual format.