A SPARSE, FAST, AND STABLE REPRESENTATION FOR MULTIPARAMETER TOPOLOGICAL DATA ANALYSIS

ABSTRACT

Topological data analysis (TDA) is a new area of geometric data analysis that focuses on using invariants from algebraic topology to provide multiscale shape descriptors for point clouds. One of the most important shape descriptors is *persistent homology*, which studies the topological variations as a filtration parameter changes; a typical parameter is the feature scale.

For many data sets, it is useful to consider varying multiple filtration parameters at once, for example scale and density. While the theoretical properties of single parameter persistent homology are well understood, less is known about the multiparameter case. Of particular interest is the problem of representing multiparameter persistent homology by elements of a vector space for integration with traditional machine learning.

Existing approaches to this problem either ignore most of the multiparameter information to reduce to the one-parameter case or are heuristic and potentially unstable in the face of noise. In this article, we introduce a general representation framework for multiparameter persistent homology that encompasses previous approaches. We establish theoretical stability guarantees under this framework as well as efficient algorithms for practical computation, making this framework an applicable and versatile tool for TDA practitioners. We validate our stability results and algorithms with numerical experiments that demonstrate statistical convergence, prediction accuracy, and fast running times on several real data sets.

1 INTRODUCTION

Topological Data Analysis (TDA) Carlsson (2009) is a methodology for analyzing data sets using multiscale shape descriptors coming from algebraic topology. There has been intense interest in the field in the last decade, since topological features have allowed practitioners to compute and encode information that classical approaches do not capture. Moreover, TDA rests on solid theoretical grounds, with guarantees accompanying many of its methods and descriptors. TDA has proved useful in a wide variety of application areas, including computer graphics Carrière et al. (2015a); Poulenard et al. (2018), computational biology Rabadán & Blumberg (2019), and material science Buchet et al. (2018); Saadatfar et al. (2017).

The main tool of TDA is *persistent homology*. In its most standard form, one is given a finite metric space (e.g., a finite set of points and their pairwise distances) and a continuous function $f : X \to \mathbb{R}$. This function usually represents a parameter of interest (such as, e.g., scale or density for point clouds, marker genes for single-cell data, etc), and the goal of persistent homology is to characterize the topological variations of this function on the data. More formally, this is achieved by computing the so-called *persistence diagram* of f, which is obtained by looking at all sublevel sets of the form $\{f^{-1}((-\infty, \alpha])\}_\alpha$ and by computing a *decomposition* of these sets, that is, by recording the appearances and disappearances of topological features (connected components, loops, enclosed spheres, etc) in these sets. When such a feature appears (resp. disappears), e.g., in a sublevel set $f^{-1}((-\infty, \alpha_b])$, we call the corresponding threshold α_b (resp. α_d) the *birth time* (resp. *death time*) of the topological feature, and we summarize this information in the persistence diagram $D(f) := \{(\alpha_b, \alpha_d)\}_{\alpha \in A} \subset \mathbb{R}^2$. It is usual to consider the distance to the diagonal $\alpha_d - \alpha_b$ as a proxy for the statistical significance of the corresponding feature. The most important foundational

Figure 1: **(left)** Example of point cloud X filtered by both feature scale (computed from unions of balls centered on X with increasing radii) and (co)density. A point will be in the bifiltration if its density is high enough, and if it is close enough to X. Note that the circle is present in the red zone, and can be detected as a large summand in the decomposition of the corresponding module **(upper right)**, or in a representation of it **(bottom right)**.

result in the subject establishes that persistence diagrams are *stable* — they will not change much when computed on a small perturbation $\tilde{f}$ of the function f Cohen-Steiner et al. (2007).

However, an inherent limitation of the formulation of persistent homology is that it can handle only a single filtration parameter f. However, in practice it is common that one has to deal with multiple filtration functions: a standard example is given in Figure 1, where both feature scale and density functions are necessary to obtain meaningful topological representation of a noisy point cloud. An extension of persistent homology to this more general setting is called *multiparameter* persistent homology Botnan & Lesnick (2022); Carlsson & Zomorodian (2009), since roughly speaking it amounts to studying the topological variation of a continuous *multiparameter* function $f : X \to \mathbb{R}^n$ with $n \in \mathbb{N}^*$. This setting is notoriously difficult to analyze theoretically, as there is no general decomposition theorem and hence no analogue of persistence diagrams.

To make multiparameter persistent homology tractable, a natural approach is to study *representations*, i.e., embeddings into Hilbert spaces, of multiparameter persistent homology. A number of such representations have now been defined Corbet et al. (2019); Coskunuzer et al. (2021); Vipond (2020). Most of them reduce to the single-parameter case by *slicing*, that is, by transforming multiparameter functions into scalar ones so that the usual decomposition result and the stability theorem of persistence diagrams still apply. Another recent line of work Carrière & Blumberg (2020) presented a heuristic that was able to encode more than just the slicing information, but has no accompanying robustness guarantee and incurred substantial computational overhead. The problem of defining a representation of multiparameter persistent homology that is fast, rich and robust thus has remained open and pressing.

Contributions. Our contributions in this work are listed below:

- We provide a general framework that parametrizes representations of multiparameter persistent homology (Definition 3.1), and which encompasses most of the previous approaches in the literature.

- We identify candidate parameters in this framework that result in representations that have stability guarantees while still encoding more information about multiparameter persistent homology than slicing (see Theorem 3.4).

- We propose a sparse implementation for our framework which allows dramatically improved memory usage and running times compared to previous implementations (see Section 4.3).

- We illustrate the performance of our framework with numerical experiments: (1) We demonstrate the practical consequences of the stability theorem by measuring the statistical convergence of our representations. (2) We achieve the best performance with the lowest runtime on several classification tasks on public data sets (see Sections 4.1 and 4.2).

Outline. Our work is organized as follows. In Section 2, we recall the basics of multiparameter persistent homology. Next, in Section 3 we present our general framework and state our associated stability result. Finally, we showcase the numerical performances of our representations in Section 4, and we conclude in Section 5.

2 BACKGROUND

In this section, we recall the basics of single and multiparameter persistent homology. We only provide a brief introduction, and we refer the reader to Chazal et al. (2016a); Oudot (2015); Rabadán & Blumberg (2019) for a thorough treatment.

Persistent homology. A standard procedure for studying a data set provided as a finite metric space, i.e., a set X with a distance $d_X \colon X \times X \to \mathbb{R}$, is to form a graph with vertex set X and an each edge connecting x_i and x_j when $d_X(x_i, x_j) < \epsilon$. Persistent homology is a simultaneous generalization of this procedure, replacing the graph with a *simplicial complex* (a combinatorial model of a topological space) and aggregating information across different scale parameters ϵ.

More generally, given a filtered topological space X, by which we mean a topological space X together with a function $f \colon X \to \mathbb{R}$, the system of inverse images $F(\alpha) := f^{-1}((-\infty, \alpha])$ fit together to produce a system $F(\alpha_1) \subseteq F(\alpha_2) \subseteq \ldots \subseteq F(\alpha_i) \subseteq \ldots$, where $\alpha_i \leq \alpha_{i+1}$. This system is an example of *filtration* of X, where a filtration is generally defined as a sequence of nested subspaces $X_1 \subseteq \ldots \subseteq X_i \subseteq \ldots \subseteq X$. Then, one can apply the kth *homology functor* H_k on each $F(\alpha_i)$. We do not define the homology functor explicitly here, but simply recall that each $H_k(F(\alpha_i))$ is a vector space, whose basis elements represent the kth dimensional topological features of $F(\alpha_i)$ (connected components for $k = 0$, loops for $k = 1$, spheres for $k = 2$, etc). Moreover the inclusions $F(\alpha_i) \subseteq F(\alpha_{i+1})$ translate into linear maps $H_k(F(\alpha_i)) \to H_k(F(\alpha_{i+1}))$, which connect the features of $F(\alpha_i)$ and $F(\alpha_{i+1})$ together. Such a sequence of vector spaces connected with linear maps is a *persistence module*, which is generally defined as a functor from $(\mathbb{R}, \leq)$ (regarded as a category) to **Vec**, the category of vector spaces.

Multiparameter persistent homology. The persistence modules defined above extend straightforwardly when there are multiple filtration directions. An n-filtration, or multifiltration, is a family of subspaces $\mathcal{F} = \{X_\alpha \subseteq X\}_{\alpha \in \mathbb{R}^n}$ such that $X_\alpha \subseteq X_{\alpha'}$ whenever $\alpha \leq \alpha'$ (in the partial order of $\mathbb{R}^n$). Moreover, we say that a multifiltration is *1-critical* if there exists a function $f : X \to \mathbb{R}^n$ such that X_α is homeomorphic to the sublevel set $\{x \in X : f(x) \leq \alpha\}$ (note that single parameter filtrations are always 1-critical). Again, applying the homology functor H_k on a multifiltration $\mathcal{F}$ induces a *multiparameter persistence module* $\mathbb{M} = \mathbb{M}(\mathcal{F})$. The *support* of $\mathbb{M}$ is then defined as the set $\operatorname{supp}(\mathbb{M}) := \{\alpha \in \mathbb{R}^n : \dim(\mathbb{M}(\alpha)) > 0\}$. Any module which is of dimension 1 everywhere on its support is called an *indicator module*, and, in addition, when its support is an interval of $\mathbb{R}^n$ (Botnan & Lesnick, 2022, Definition 2.1), it is called an *interval module*. We also use $\mathbb{M}\big|_K$ to denote the module obtained by restricting $\mathbb{M}$ on a subset K of $\mathbb{R}^n$. Notably, when K is a line, $\mathbb{M}\big|_K$ is a single parameter module: this is called *slicing* the module along K.

A multiparameter persistence module $\mathbb{M}$ can always be decomposed in a unique way: $\mathbb{M} = \oplus_{i=1}^m M_i$ (Botnan & Lesnick, 2022, Theorem 4.2), where the M_i are indecomposable modules that are called the *summands* of $\mathbb{M}$. See Figure 1 (upper right) for an example of decomposition. For single parameter filtrations, the summands of the decomposition are known to be interval modules $\mathbb{M} = \oplus_{i=1}^m \mathbb{I}[\alpha_{b_i}, \alpha_{d_i}]$, and can be summarized in a *persistence diagram* $D(\mathbb{M}) = \{(\alpha_{b_i}, \alpha_{d_i})\}_{1 \leq i \leq m}$. This result does not extend to the multiparameter setting however, which motivates the use of slicing in order to reduce to the single parameter case and its decomposition guarantees.

Finally, multiparameter persistence modules can be compared with two standard distances: the *interleaving* and *bottleneck* (or ℓ^∞) distances. Their explicit definitions are technical and not necessary for our main exposition, so we refer the reader to, e.g., [Sections 6.1, 6.4]Botnan & Lesnick (2022) for more details. The *stability* theorem [Theorem 5.3]Lesnick (2015) states that 1-critical multiparameter persistence modules are stable:

$$d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}') \leq \|f - f'\|_\infty, \tag{1}$$

where f and f' are functions associated to the 1-critical modules $\mathbb{M}$ and $\mathbb{M}'$ respectively.

3 A FRAMEWORK FOR PERSISTENCE MODULE REPRESENTATION

Even though multiparameter persistence modules are known to encode useful data information, their algebraic definitions make them not suitable for subsequent data science and machine learning purposes. Hence, in this section, we propose a general framework for studying *representations* of multiparameter persistence modules, i.e., maps defined on the space of multiparameter persistence modules and taking values in an (implicit or explicit) Hilbert space. Defining such maps is crucial for subsequent data science tasks, as most of the standard methods in statistical machine learning expect the input to be Euclidean vectors, or at least to be equipped with a scalar product.

3.1 PERSISTENCE MODULE REPRESENTATION

Framework. In this article, we study the following general representation framework:

Definition 3.1. Let $\mathbb{M}$ be a multiparameter persistence module. Let $\mathbb{M} = \oplus_{i=1}^m M_i$ be its decomposition (note that the M_i need not be intervals), and let $\mathcal{M}$ be the space of indecomposable persistence modules. A *multiparameter representation* of $\mathbb{M}$ is:

$$V_{\mathrm{op},w,\phi}(\mathbb{M}) = \mathrm{op}(\{w(M_i) \cdot \phi(M_i)\}_{i=1}^m), \tag{2}$$

where op is a permutation invariant operation (sum, max, min, mean, etc), $w : \mathcal{M} \to \mathbb{R}$ is a weight function, and $\phi : \mathcal{M} \to \mathcal{H}$ sends any indecomposable module to a vector in a Hilbert space $\mathcal{H}$.

This general definition is inspired from a similar framework that was introduced for single-parameter persistence with the automatic representation method *Perslay* Carrière et al. (2020).

Remark 3.2. Definition 3.1 requires being able to decompose multiparameter persistence modules. Several approaches have been proposed for this problem in the literature, that provide either exact Dey & Xin (2022); Lesnick & Wright (2015) or approximate Loiseaux et al. (2022) decompositions, or decompositions that are restricted to specific homology dimensions Cai et al. (2020).

Relation to previous work. Interestingly, specific choices of op, w and ϕ can reproduce the previous work from the literature. For sake of simplicity, assume $\mathbb{M} = \oplus_{i=1}^m M_i$ can be decomposed into indicator modules (we can thus, by a slight abuse of notation, identify any summand M_i with its support), then:

- Using $w : M_i \mapsto 1$, $\phi : M_i \mapsto \left\{ \begin{array}{ll} \mathbb{R}^n & \to \mathbb{R} \\ x & \mapsto \Lambda(x, M_i \cap \ell_{\mathbf{e}_\Delta,x}) \end{array} \right.$ and op $= k$th maximum, where $\Lambda(\cdot, \ell)$ denotes the tent function associated to any segment $\ell \subset \mathbb{R}^n$, induces the kth multiparameter persistence landscape Vipond (2020).

- Using $w : M_i \mapsto 1$, $\phi : M_i \mapsto \left\{ \begin{array}{ll} \mathbb{R}^n \times \mathbb{R}^n & \to \mathbb{R}^d \\ p, q & \mapsto w'(M_i \cap [p,q]) \cdot \phi'(M_i \cap [p,q]) \end{array} \right.$ and op $=$ op$'$, where op$'$, w' and ϕ' are the parameters of any persistence diagram representation from Perslay, induces the multiparameter persistence kernel Corbet et al. (2019).

- Using $w : M_i \mapsto \mathrm{vol}(M_i)$, $\phi : M_i \mapsto \left\{ \begin{array}{ll} \mathbb{R}^n & \to \mathbb{R} \\ x & \mapsto \exp(-\min_{\ell \in L} \|x, M_i \cap \ell\|_2^2 / \sigma^2) \end{array} \right.$ and op $= \sum$, where L is a set of (pre-defined) diagonal lines, induces the multiparameter persistence image Carrière & Blumberg (2020).

Concerning the first two approaches, it is worth noting that they do not depend on the decompositions and the global geometric shapes of the support of the summands since they are based on slicing modules, which is stable but not very informative about the summand shapes (see (Loiseaux et al., 2022, Section 7.1) for examples of different modules with similar slicing). This means in particular that a lot of information about the module is lost during the representation step. On the other hand, the third approach does use the module decomposition but is known to be unstable (Figure 11 in Botnan & Lesnick (2018) provides two modules that are close in the interleaving distance but that also have multiparameter persistence images that are far away from each other). In the next section, we study representations that allow to encode the summand shapes while also enjoying stability guarantees.

3.2 METRIC PROPERTIES

In this section, we study specific parameters within our framework in Definition 3.1, and prove associated theoretical properties about their associated representations.

Definition. We begin by defining parameters that we call *geometric and stable* (GS) in the sense that they capture shape information about the summands, and that they also lead to robust representations.
Definition 3.3. The *geometric and stable* (GS) representation parameters are:

1. the weight function $w : M \mapsto d_{\mathrm{I}}(M, 0)$,

2. the individual summand representations $\phi_\delta(M) : \mathbb{R}^n \to \mathbb{R}$:

   (a) $\phi_\delta(M)(x) = \frac{1}{\delta} d_{\mathrm{I}}(M\big|_{R_{x,\boldsymbol{\delta}}}, 0)$,   (b) $\phi_\delta(M)(x) = \frac{1}{(2\delta)^n} \mathrm{vol}\left(\mathrm{supp}\left(M\right) \cap R_{x,\boldsymbol{\delta}}\right)$,

   (c) $\phi_\delta(M)(x) = \frac{1}{(2\delta)^n} \sup_{x',\boldsymbol{\delta}'} \left\{\mathrm{vol}(R_{x',\boldsymbol{\delta}'}) : R_{x',\boldsymbol{\delta}'} \subseteq \mathrm{supp}\left(M\right) \cap R_{x,\boldsymbol{\delta}}\right\}$,

   where $R_{x,\boldsymbol{\delta}} := \left\{z \in \mathbb{R}^n : x - \boldsymbol{\delta} \le z \le x + \boldsymbol{\delta}\right\} \subseteq \mathbb{R}^n$, $\boldsymbol{\delta} := \delta \cdot [1, \ldots, 1] \in \mathbb{R}^n$ for any $\delta > 0$, and vol denotes the volume of a set in $\mathbb{R}^n$.

3. the permutation invariant operators $\mathrm{op} = \sum$ and $\mathrm{op} = \sup$.

Equipped with these GS parameters, we can now define the two following GS representations, that can be applied on any multiparameter persistence module $\mathbb{M} = \oplus_{i=1}^m M_i$:

$$V_{p,\delta}(\mathbb{M}) := \sum_{i=1}^m \frac{w(M_i)^p}{\sum_{j=1}^m w(M_j)^p} \phi_\delta(M_i), \text{ for some } p \in \mathbb{R}_+, \tag{3}$$

$$V_{\infty,\delta}(\mathbb{M}) := \sup_{1 \le i \le m} \phi_\delta(M_i). \tag{4}$$

These GS parameters allow for some trade-off between computational cost and the amount of information that is kept: (a) and (c) are very easy to compute on interval modules, but (b) encodes more information about summand shapes. See Figures 1 (upper right) and 2b for visualizations.

Stability. The main motivation for introducing our GS representations is that one can show that they are stable in the interleaving and bottleneck distances, as stated in the following theorem. We assume that persistence modules have support included in a compact set $K \subseteq \mathbb{R}^n$ for simplicity (but our results extend straightforwardly to modules restricted to K if their supports are infinite).
Theorem 3.4. *Let $\mathbb{M} = \oplus_{i=1}^m M_i$ and $\mathbb{M}' = \oplus_{j=1}^{m'} M_j'$ be two multiparameter persistence modules that can be decomposed with indicator modules. Assume that we have $\frac{1}{m}\sum_i w(M_i), \frac{1}{m'}\sum_j w(M_j') \ge C$, for some $C > 0$. Then for any parameter $\delta > 0$, one has*

$$\|V_{0,\delta}(\mathbb{M}) - V_{0,\delta}(\mathbb{M}')\|_\infty \le 2(d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta, \tag{5}$$

$$\|V_{1,\delta}(\mathbb{M}) - V_{1,\delta}(\mathbb{M}')\|_\infty \le \left[4 + \frac{2}{C}\right](d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta, \tag{6}$$

$$\|V_{\infty,\delta}(\mathbb{M}) - V_{\infty,\delta}(\mathbb{M}')\|_\infty \le (d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta. \tag{7}$$

Again, we emphasize that our GS representations encode strictly more information than slicing (whereas the multiparameter persistence landscape Vipond (2020) and kernel Corbet et al. (2019) do not). The multiparameter persistence image Carrière & Blumberg (2020) is also richer than slicing, but does not satisfy any robustness properties. Theorem 3.4 shows that our GS representations bring the best of both worlds: they are richer than slicing and stable at the same time.

A proof of Theorem 3.4 can be found in Appendix B. One might worry that the minimum with δ encountered in the upper bounds in Theorem 3.4 show that our GS representations are too stable and not discriminative enough for practical purposes; while we did not observe such behaviour in our numerical experiments, we also provide an additional stability result with a similar, yet slightly different representation in Appendix C, with an associated upper bound without minimum.
Remark 3.5. It can easily be seen that the GS representations are injective: if the support of the modules are different, then the GS representations (evaluated on a point that belongs to the support of one summand but not on the support of the other) will differ, provided that δ is sufficiently small.

4 NUMERICAL EXPERIMENTS

In this section, we illustrate the various properties of our GS representations with numerical experiments. First, we explore the stability theorem in Section 4.1 by studying the convergence rates of our GS representations on various data sets. Then, we showcase the efficiency of our GS representations on classification tasks in Section 4.2. Finally, we discuss running times in Section 4.3.

Decomposition. The representations presented in this article require us to know the persistence module decomposition. Many different approaches can be used for this task Loiseaux et al. (2022); Dey & Xin (2022); Cai et al. (2020); Lesnick & Wright (2015). In our experiments, we always use the MMA approximate decomposition Loiseaux et al. (2022). Our code is available in the supplementary material and will be released publicly upon acceptance.

4.1 CONVERGENCE RATE

In this section, we study the convergence rate of our representation with respect to the number of sampled points. Similar to the single parameter persistence setting Chazal et al. (2015), these rates are derived for free from our stability theorem. Indeed, since concentration inequalities for multiparameter persistence modules have already been showed in the literature, these concentration inequalities apply directly to our representations. Note that while Equations (8) and (9), which provide such rates, are stated for GS representation (4), they also hold for GS representation (3).

Measure bifiltration. Let μ be a compactly supported probability measure of $\mathbb{R}^D$, and let μ_n be the discrete measure associated to a sampling of n points from μ. The *measure bifiltration* associated to μ and μ_n is defined as $\mathcal{F}^\mu_{r,t} := \{x \in \mathbb{R}^D : \mu(B(x,r)) \leq t\}$, where $B(x,r)$ denotes the Euclidean ball centered on x with radius r. Now, let $\mathbb{M}$ and $\mathbb{M}_n$ be the multiparameter persistence modules obtained from applying the homology functor on top of the measure bifiltrations $\mathcal{F}^\mu$ and $\mathcal{F}^{\mu_n}$. These modules are known to enjoy the following stability result (Blumberg & Lesnick, 2021, Theorem 3.1, Proposition 2.23 (i)):

$$d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}_n) \leq d_{\mathrm{Pr}}(\mu, \mu_n) \leq \min(d_W^p(\mu, \mu_n)^{\frac{1}{2}}, d_W^p(\mu, \mu_n)^{\frac{p}{p+1}}),$$

where d_W^p and d_{Pr} stand for the p-Wasserstein and Prokhorov distances between probability measures. Combining these inequalities with Theorem 3.4, then taking expectations and applying the concentration inequalities of the Wasserstein distance (see (Lei, 2020, Theorem 3.1) and (Fournier & Guillin, 2015, Theorem 1)) lead to:

$$\delta \mathbb{E}\left[\|V_{\infty,\delta}(\mathbb{M}) - V_{\infty,\delta}(\mathbb{M}_n)\|_\infty\right] \leq \left(c_{p,q} \|x\|_{L_\mu^q} n^{-\left(\frac{1}{2p \vee d}\right) \wedge \frac{1}{p} - \frac{1}{q}} \log^{\alpha/q} n\right)^{\frac{p}{p+1}}, \qquad (8)$$

where $\alpha = 2$ if $2p = q = d$, $\alpha = 1$ if $d \neq 2p$ and $q = dp/(d-p) \wedge 2p$ or $q > d = 2p$ and $\alpha = 0$ otherwise, $c_{p,q}$ is a constant that depends on p and q, and $\|x\|_{L_\mu^q}$ denotes the first moment of μ.

Čech complex and density. A limitation of the measure bifiltration is that it can be difficult to compute. Hence, we now focus on another, easier to compute bifiltration. Let X be a compact d-submanifold of $\mathbb{R}^D$ ($d \leq D$), and μ be a measure on X with density f with respect to the uniform measure on X. We now define the 1-critical bifiltration $\mathcal{F}^{C,f}$ swith:

$$\mathcal{F}^{C,f}_{u,v} := \check{\mathrm{Cech}}(u) \cap f^{-1}([v,\infty)) = \left\{x \in \mathbb{R}^D : d(x,X) \leq u, f(x) \geq v\right\}.$$

Moreover, given a set X_n of n points sampled from μ, we also consider the approximate bifiltration $\mathcal{F}^{C,f_n}$, where $f_n \colon X \to \mathbb{R}$ is an estimation of f (such as, e.g., a kernel density estimator). Let $\mathbb{M}$ and $\mathbb{M}_n$ be the multiparameter persistence modules associated to $\mathcal{F}^{C,f}$ and $\mathcal{F}^{C,f_n}$. Then, the stability of the interleaving distance (Lesnick, 2015, Theorem 5.3) ensures:

$$d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}_n) \leq \|f - f_n\|_\infty \vee d_H(X, X_n),$$

where d_H stands for the Hausdorff distance. Moreover, concentration inequalities for the Hausdorff distance and kernel density estimators are also available in the literature under some conditions (see (Chazal et al., 2015, Theorem 4) and (Kim et al., 2019, Corollary 15)). More precisely, when the

density f is L-Lipschitz and bounded from above and from below, i.e., when $0 < f_{\min} \leq f \leq f_{\max} < \infty$, and when f_n is a kernel density estimator of f with associated kernel k, one has:

$$\mathbb{E}(d_H(X, X_n)) \lesssim \left(\frac{\log n}{n}\right)^{\frac{1}{d}} \text{ and } \mathbb{E}(\|f - f_n\|_\infty) \lesssim Lh_n \|x\|_{L_k \, d\lambda_d} + \sqrt{\frac{\log(1/h_n)}{nh_n^d}}$$

where h_n is the (adaptive) bandwidth of the kernel k. In particular, if μ is a measure comparable to the uniform measure of a $d = 2$-manifold, then for any stationary sequence $h_n := h > 0$, one has:

$$\delta\mathbb{E}\left[\|V_{\infty,\delta}(\mathbb{M}) - V_{\infty,\delta}(\mathbb{M}_n)\|_\infty\right] \lesssim \sqrt{\frac{\log n}{n}} + L\sqrt{2}h. \tag{9}$$

Empirical convergence rates. Now that we have established the theoretical convergence rates of our GS representations, we estimate and validate them on data sets. We will study a synthetic data set and a real data set of point clouds obtained with immunohistochemistry.

Annulus with non-uniform density. In this synthetic example, we generate an annulus in $\mathbb{R}^2$ with a non-uniform density, displayed in Figure 2a. Then, we compute the bifiltration $\mathcal{F}^{C,f_n}$ corresponding to the Alpha filtration and the sublevel set filtration of a kernel density estimator, with bandwidth parameter $h = 0.1$, on the complete Alpha simplicial complex. Finally, we compute the representations (see Figure 2b) of the associated multiparameter module (in homology dimension 1), and their normalized distances to the target representation, using either $\|\cdot\|_2^2$ or $\|\cdot\|_\infty$.

The corresponding distances for various number of sample points are displayed in log-log plots in Figure 2c. One can see that the empirical rate is roughly consistent with the theoretical one ($-1/2$ for $\|\cdot\|_\infty$ and -1 for $\|\cdot\|_2$), even when $p \neq \infty$ (in which case our GS representations are stable for the bottleneck distance but not for the interleaving distance).

(a) Scatter plot of the synthetic data set, comprised of 25,000 points, and colored by a kernel density estimator with bandwidth $h = 0.5$.

(b) GS representations with parameters, $p = 0$ (upper left), $p = 1$ (upper right), $p = \infty$ (bottom).

(c) **(left)** $\|\cdot\|_2^2$ and **(right)** $\|\cdot\|_\infty$ (right) between the target representation and the empirical one w.r.t. n.

Figure 2: Convergence rate of synthetic data set.

Immunohistochemistry data. In our second experiment, we consider a point cloud representing cells, taken from Vipond et al. (2021), see Figure 3a. These cells are given with biological markers, which are typically used to assess, e.g., cell types and functions. In this experiment, we first triangulate the point cloud by placing a 100×100 grid on top of it. Then, we filter this grid using the sublevel set filtrations of kernel density estimators (with Gaussian kernel and bandwidth $h = 1$) associated to the CD8 and CD68 biological markers for immune cells. Finally, we compute the associated multiparameter modules in homology dimensions 0 and 1, and we compute and concatenate their corresponding GS representations.

The convergence rates are displayed in Figure 3b. Similar to the previous experiment, the theoretical convergence rate of our representations is upper bounded by the one for kernel density estimators with the ∞-norm, which is of order $\frac{1}{\sqrt{n}}$ with respect to the number n of sampled points. Again, the observed convergence rate are consistent with the theoretical ones.

(a) Point cloud representing cells colored by CD8 (in red) and CD68 (in black).

(b) **(left)** $\|\cdot\|_2^2$ and **(right)** $\|\cdot\|_\infty$ distances between the target representation and the empirical one w.r.t. n.

Figure 3: Convergence rate of immunofluorescence data set.

4.2 CLASSIFICATION

In this section, we illustrate the efficiency of our GS representations by using them for classification purposes, and showing that they perform at least as good as already existing representations (we also provide non-topological baselines for context, even though our primary goal is not to compete against those). We focus on a data set of point clouds representing cells from Vipond et al. (2021). Every point cloud has a label corresponding to the type of its cells, and our goal is to check whether we can predict these labels using the bifiltration defined in the previous section. We also compare the performances of our GS representations (evaluated on a 50×50 grid) to the one of the multiparameter persistence landscape (MPL)[1] and the one of the Ripley function, which is a classical statistical tool for analyzing point processes. Our accuracy scores were cross-validated with a 10-fold cross-validation over the following parameters; $p \in \{0, 1\}$, op $\in \{\text{sum}, \text{mean}\}$, $\delta \in \{0.01, 0.1, 0.5\}$, homological dimensions $\{\{0, 1\}, \{0, 1, 2\}\}$. Results obtained after training a standard random forest classifier can be found in Table 1. One can see that our GS representations outperform both MPL and Ripley function for almost all GS parameters. This indicates in particular that our representation contains indeed more information than slicing. We also provide accuracy scores for several UCR data sets in Appendix D, in which our GS representations also compare favorably to competitors.

---

[1]Note that the sizes of the point clouds were too large for practical computations of the multiparameter persistence kernel Corbet et al. (2019) and the multiparameter persistence image Carrière & Blumberg (2020).

Ours	p	δ	Accuracy (%)
		10^{-2}	74.1 ± 2.9
-	1	10^{-3}	75.6 ± 3.4
		10^{-4}	68.7 ± 3.0
		10^{-2}	77.81 ± 2.0
-	0	10^{-3}	76.0 ± 3.3
		10^{-4}	70.5 ± 3.0
MPL	-	-	60.5 ± 3.5
Ripley	-	-	67.2 ± 2.2

Table 1: Classification results for immuno-histochemistry data.

	Annulus	Immuno
Ours (GS)	$250ms \pm 2ms$	$275ms \pm 9.8ms$
Ours (MPL)	$36.9ms \pm 0.8ms$	$65.9ms \pm 0.9ms$
MPI (50)	$6.43s \pm 25ms$	$5.67s \pm 23.3ms$
MPL (50)	$17s \pm 39ms$	$15.6s \pm 14ms$
MPI (100)	$13.1s \pm 125ms$	$11.65s \pm 7.9ms$
MPL (100)	$35s \pm 193ms$	$31.3s \pm 23.3ms$

Table 2: Running times for our GS representations and competitors.

4.3 RUNNING TIMES

Finally, we end this section with running time comparisons. It is worth noting that our implementation (with pseudo-code available in Appendix E) relies on an implicit description of the summands in the module decomposition, which are characterized with their *corners* Loiseaux et al. (2022). This is in contrast with the implementations of the other representations that are available in the `multipers`[2] library, which require slicing the modules with many lines first. This shows from Table 2, where it can be seen that our representations (computed on the pinched annulus and immunohistochemistry data sets defined in the previous sections) can be computed much faster than the others, by a factor of at least 25 (all representations are evaluated on a grid of size 100×100, and we provide the maximum running time over $p \in \{0, 1, \infty\}$). All computations were done using a Ryzen 4800 laptop CPU, with 16GB of RAM. Interestingly, this sparse and fast implementation based on corners can also be used to improve on the running time of the multiparameter landscape (see second line of Table 2).

5 CONCLUSION

In this article, we study the general question of representing multiparameter persistence modules in Topological Data Analysis. We first introduce a general theoretical framework that allows to recover existing approaches in the literature, as well as to define new ones. Then, we identify and prove the stability of some specific representations in that framework, that we call *geometric and stable* (GS) representations since they are able to encode the shapes of the summands comprised in the modules, making these representations more powerful than already existing ones. We finally showcase the stability and efficiency of these GS representations in various theoretical and numerical applications about statistical convergence and classification accuracies.

Several questions remain open for future works:

- The framework that we introduced in Definition 3.1 is similar to the Perslay framework of single parameter persistence Carrière et al. (2020). In this work, each of the framework parameter was optimized by a neural network. It is thus natural to investigate whether one can optimize the module representation within our framework in a data-driven way as well.

- In our numerical applications, we focused on representations computed off of MMA decompositions Loiseaux et al. (2022). In the future, we plan to investigate whether working with other decomposition methods Cai et al. (2020); Dey & Xin (2022) lead to better numerical performances when combined with our representation framework.

- Even though our experiments were done on data sets equipped with bifiltrations, our code and framework is well-defined for any number of filtrations. Identifying and investivating the dependences and performances of our framework on more than two filtrations is thus of particular interest.

---

[2]https://github.com/MathieuCarriere/multipers

REFERENCES

Andrew J. Blumberg and Michael Lesnick. Stability of 2-parameter persistent homology. *arXiv:2010.09628 [cs, math]*, February 2021. URL http://arxiv.org/abs/2010.09628.

Magnus Botnan and Michael Lesnick. An introduction to multiparameter persistence. In *CoRR*. arXiv:2203.14289, 2022.

Magnus Bakke Botnan and Michael Lesnick. Algebraic stability of zigzag persistence modules. *Algebraic & Geometric Topology*, 18(6):3133–3204, October 2018. ISSN 1472-2739, 1472-2747. doi: 10/gkw5xh.

Mickaël Buchet, Yasuaki Hiraoka, and Ippei Obayashi. Persistent homology and materials informatics. In *Nanoinformatics*, pp. 75–95. Springer-Verlag, 2018.

Chen Cai, Woojin Kim, Facundo Mémoli, and Yusu Wang. Elder-rule-staircodes for augmented metric spaces. In *36th International Symposium on Computational Geometry (SoCG 2020)*, pp. 26:1–26:17. Schloss Dagstuhl–Leibniz-Zentrum fuer Informatik, 2020.

Gunnar Carlsson. Topology and data. *Bulletin of the American Mathematical Society*, 46(2):255–308, 2009.

Gunnar Carlsson and Afra Zomorodian. The theory of multidimensional persistence. *Discrete & Computational Geometry*, 42(1):71–93, 2009.

Mathieu Carrière and Andrew Blumberg. Multiparameter persistence image for topological machine learning. In *Advances in Neural Information Processing Systems 33 (NeurIPS 2020)*, pp. 22432–22444. Curran Associates, Inc., 2020.

Mathieu Carrière, Steve Oudot, and Maks Ovsjanikov. Stable topological signatures for points on 3D shapes. *Computer Graphics Forum*, 34(5):1–12, 2015a.

Mathieu Carrière, Steve Oudot, and Maks Ovsjanikov. Local signatures using persistence diagrams. In *HAL*. HAL:01159297, 2015b.

Mathieu Carrière, Frédéric Chazal, Yuichi Ike, Théo Lacombe, Martin Royer, and Yuhei Umeda. PersLay: a neural network layer for persistence diagrams and new graph topological signatures. In *23rd International Conference on Artificial Intelligence and Statistics (AISTATS 2020)*, pp. 2786–2796. PMLR, 2020.

Frédéric Chazal, Marc Glisse, Catherine Labruère, and Bertrand Michel. Convergence rates for persistence diagram estimation in topological data analysis. *Journal of Machine Learning Research*, 16(110):3603–3635, 2015.

Frédéric Chazal, Vin de Silva, Marc Glisse, and Steve Oudot. *The structure and stability of persistence modules*. SpringerBriefs in Mathematics. Springer-Verlag, 2016a.

Frédéric Chazal, Vin de Silva, Marc Glisse, and Steve Oudot. *The Structure and Stability of Persistence Modules*. SpringerBriefs in Mathematics. Springer International Publishing, Cham, 2016b. ISBN 978-3-319-42543-6 978-3-319-42545-0. doi: 10.1007/978-3-319-42545-0.

David Cohen-Steiner, Herbert Edelsbrunner, and John Harer. Stability of persistence diagrams. *Discrete & Computational Geometry*, 37(1):103–120, 2007.

René Corbet, Ulderico Fugacci, Michael Kerber, Claudia Landi, and Bei Wang. A kernel for multi-parameter persistent homology. *Computers & Graphics: X*, 2:100005, 2019.

Baris Coskunuzer, Ignacio Akcora, Cuneyt Dominguez, Yuzhou Chen, Zhiwei Zhen, Murat Kantarcioglu, and Yulia Gel. Smart Vectorizations for Single and Multiparameter Persistence. In *CoRR*. arXiv:2104.04787, 2021.

Tamal Dey and Cheng Xin. Generalized persistence algorithm for decomposing multiparameter persistence modules. *Journal of Applied and Computational Topology*, 2022.

Nicolas Fournier and Arnaud Guillin. On the rate of convergence in Wasserstein distance of the empirical measure. *Probability Theory and Related Fields*, 162(3):707–738, August 2015. ISSN 1432-2064. doi: 10.1007/s00440-014-0583-7.

Jisu Kim, Jaehyeok Shin, Alessandro Rinaldo, and Larry Wasserman. Uniform convergence rate of the kernel density estimator adaptive to intrinsic volume dimension. In *Proceedings of the 36th International Conference on Machine Learning (ICML 2019)*, volume 97, pp. 3398–3407. PMLR, 2019.

Claudia Landi. The rank invariant stability via interleavings. In *Research in Computational Topology*, pp. 1–10. Springer, 2018.

Jing Lei. Convergence and Concentration of Empirical Measures under Wasserstein Distance in Unbounded Functional Spaces. *Bernoulli*, 26(1), February 2020. ISSN 1350-7265. doi: 10.3150/19-BEJ1151.

Michael Lesnick. The theory of the interleaving distance on multidimensional persistence modules. *Foundations of Computational Mathematics*, 15(3):613–650, 2015.

Michael Lesnick and Matthew Wright. Interactive visualization of 2D persistence modules. In *CoRR*. arXiv:1512.00180, 2015.

David Loiseaux, Mathieu Carrière, and Andrew Blumberg. Efficient Approximation of Multiparameter Persistence Modules. In *CoRR*. arXiv:2206.02026, 2022.

Steve Oudot. *Persistence theory: from quiver representations to data analysis*, volume 209 of *Mathematical Surveys and Monographs*. American Mathematical Society, 2015.

Adrien Poulenard, Primoz Skraba, and Maks Ovsjanikov. Topological function optimization for continuous shape matching. *Computer Graphics Forum*, 37(5):13–25, 2018.

Raúl Rabadán and Andrew Blumberg. *Topological data analysis for genomics and evolution*. Cambridge University Press, 2019.

Mohammad Saadatfar, Hiroshi Takeuchi, Vanessa Robins, Nicolas François, and Yasuaki Hiraoka. Pore configuration landscape of granular crystallization. *Nature Communications*, 8:15082, 2017.

Oliver Vipond. Multiparameter persistence landscapes. *Journal of Machine Learning Research*, 21 (61):1–38, 2020.

Oliver Vipond, Joshua Bull, Philip Macklin, Ulrike Tillmann, Christopher Pugh, Helen Byrne, and Heather Harrington. Multiparameter persistent homology landscapes identify immune cell spatial patterns in tumors. *Proceedings of the National Academy of Sciences of the United States of America*, 118(41), 2021.

A   ADDITIONAL BACKGROUND

Definition A.1. Let $n \in \mathbb{N}^*$ and $\mathcal{F} = \{\mathcal{F}^{(1)}, \ldots, \mathcal{F}^{(n)}\}$ be a multi-filtration on a topological space X. Let $\mathbf{e}, b \in \mathbb{R}^n$, and $\ell_{\mathbf{e},b} : \mathbb{R} \to \mathbb{R}^n$ be the line in $\mathbb{R}^n$ defined with $\ell_{\mathbf{e},b}(t) = t \cdot \mathbf{e} + b$, that is, $\ell_{\mathbf{e},b}$ is the line of direction $\mathbf{e}$ passing through b. Let $\mathcal{F}_{\mathbf{e},b} : \mathbb{R} \to \mathcal{P}(X)$ defined with $\mathcal{F}_{\mathbf{e},b}(t) = \bigcap_{i=1}^n \mathcal{F}^{(i)}([\ell_{\mathbf{e},b}(t)]_i)$, where $[\cdot]_i$ denotes the i-th coordinate. Then, each $\mathcal{F}_{\mathbf{e},b}$ is a single-parameter filtration and has a corresponding persistence diagram, or barcode $B_{\mathbf{e},b}$. The set $\mathcal{B}(\mathcal{F}) = \{B_{\mathbf{e},b} : \mathbf{e}, b \in \mathbb{R}^n\}$ is called the *fibered barcode* of $\mathcal{F}$.

The two following lemmas from Landi (2018) describe two useful properties of the fibered barcode.

Lemma A.2 (Lemma 1 in Landi (2018))**.** *Let $\mathbf{e}, b \in \mathbb{R}^n$ and $\ell_{\mathbf{e},b}$ be the corresponding line. Let $\hat{e} = \min_i[\mathbf{e}]_i$. Let $\mathcal{F}, \mathcal{F}'$ be two multi-filtrations, $\mathbb{M}, \mathbb{M}'$ be the corresponding persistence modules and $B_{\mathbf{e},b} \in \mathcal{B}(\mathcal{F})$ and $B'_{\mathbf{e},b} \in \mathcal{B}(\mathcal{F}')$ be the corresponding barcodes in the fibered barcodes of $\mathcal{F}$ and $\mathcal{F}'$. Then, the following stability property holds:*

$$d_{\mathrm{B}}(B_{\mathbf{e},b}, B'_{\mathbf{e},b}) \leq \frac{d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}')}{\hat{e}}. \tag{10}$$

Lemma A.3 (Lemma 2 in Landi (2018)). *Let $\mathbf{e}, \mathbf{e}', b, b' \in \mathbb{R}^n$ and $\ell_{\mathbf{e},b}$, $\ell_{\mathbf{e}',b'}$ be the corresponding lines. Let $\hat{e} = \min_i[\mathbf{e}]_i$ and $\hat{e}' = \min_i[\mathbf{e}']_i$. Let $\mathcal{F}$ be a multi-filtration, $\mathbb{M}$ be the corresponding persistence module and $B_{\mathbf{e},b}, B_{\mathbf{e}',b'} \in \mathcal{B}(\mathcal{F})$ be the corresponding barcodes in the fibered barcode of $\mathcal{F}$. Assume $\mathbb{M}$ is decomposable $\mathbb{M} = \oplus_{i=1}^m M_i$, and let $K > 0$ such that $M_i \subseteq B_\infty(0, K) := \{x \in \mathbb{R}^n : \|x\|_\infty \le K\}$ for all $i \in [\![1, m]\!]$. Then, the following stability property holds:*

$$d_{\mathrm{B}}(B_{\mathbf{e},b}, B_{\mathbf{e}',b'}) \le \frac{(K + \max\{\|b\|_\infty, \|b'\|_\infty\}) \cdot \|\mathbf{e} - \mathbf{e}'\|_\infty + \|b - b'\|_\infty}{\hat{e} \cdot \hat{e}'}. \tag{11}$$

B PROOF OF THEOREM 3.4

Our proof is based on several lemmas. In the first one, we focus on the GS weight function $w : M \mapsto d_{\mathrm{I}}(M, 0)$.

Lemma B.1. *Let M and M' be two indicator modules with compact support. Then, one has*

$$|w(M) - w(M')| \le d_{\mathrm{I}}(M, M'). \tag{12}$$

Furthermore, one has the equality

$$w(M) = \frac{1}{2} \sup_{b,d \in \mathrm{supp}(M)} \min_j (d_j - b_j)_+. \tag{13}$$

Proof. Inequality (12) follows directly from the triangle inequality applied on d_{I}.

We now show Equation (13) with two inequalities.

First inequality: $\le$ Let M be an indicator module. If $w(M) := d_{\mathrm{I}}(M, 0) = 0$, then the inequality is trivial, so we now assume that $w(M) > 0$. Let $\delta > 0$ such that $\delta < w(M)$. By definition of d_{I}, the identity morphism $M \to M[2\boldsymbol{\delta}]$ cannot be factorized by 0. This implies the existence of some $b \in \mathbb{R}^n$ such that $\mathrm{Rk}(M(b \le b + 2\boldsymbol{\delta})) > 0$; in particular, $b, b + 2\boldsymbol{\delta} \in \mathrm{supp}(M)$. Making δ converge to $w(M)$ yields the desired inequality.

Second inequality: $\ge$ Let $(K_n)_{n \in \mathbb{N}}$ be a convex compact exhaustion of $\mathrm{supp}(M)$, and $b_n, d_n \in K_n$ be two points that achieve the maximum in

$$\frac{1}{2} \sup_{b,d \in K_n} \min_j (d_j - b_j)_+.$$

Now, by functoriality of persistence modules, we can assume without loss of generality that b_n and d_n are on the same diagonal line (indeed, if they are not, it is possible to transform d_n into $\tilde{d}_n$ such that b_n and $\tilde{d}_n$ are on the same diagonal line and also achieve the maximum). Thus, $\mathrm{rank}(M(b_n \le d_n)) > 0$, and $d_{\mathrm{I}}(M, 0) \ge \frac{1}{2} \|d_n - b_n\|_\infty$. Taking the limit over $n \in \mathbb{N}$ leads to the desired inequality. $\square$

In the following lemma, we rewrite volumes of indicator module supports using interleaving distances.

Lemma B.2. *Let M be an indicator module, and $R \subseteq \mathbb{R}^n$ be a compact rectangle, with $n \ge 2$. Then, one has:*

$$\mathrm{vol}(\mathrm{supp}(M) \cap R) = 2 \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}\left(M\big|_{l_y \cap R}, 0\right) \mathrm{d}\lambda^{n-1}(y)$$

where l_y is the diagonal line crossing y, and λ^{n-1} denotes the Lebesgue measure in $\mathbb{R}^{n-1}$.

Proof. Using the change of variables $y_i = x_i - x_n$ and $t = x_n$ (which has a trivial Jacobian) yields the following inequalities:

$$\mathrm{vol}(\mathrm{supp}(M) \cap R) = \int_{\mathrm{supp}(M) \cap R} \mathrm{d}\lambda^n(x)$$

$$= \int_{\{y \in \mathbb{R}^n : y_n = 0\}} \int_{t \in \mathbb{R}} \mathbb{1}_{\mathrm{supp}(M) \cap R}(y + \boldsymbol{t}) \, \mathrm{d}t \, \mathrm{d}\lambda^{n-1}(y)$$

$$= \int_{\{y \in \mathbb{R}^n : y_n = 0\}} \|\mathrm{supp}(M) \cap l_y \cap R\| \, \mathrm{d}\lambda^{n-1}(y)$$

where l_y is the diagonal line passing through y. Now, since M is an indicator module, one has $\|\mathrm{supp}(M) \cap l_y \cap R\| = 2 d_{\mathrm{I}}(M\big|_{l_y \cap R}, 0)$, which concludes the proof. $\square$

In the following proposition, we provide stability bounds for single indicator modules.

Proposition B.3. *If M and M' are two indicator modules, then for any $\delta > 0$ and GS parameter ϕ_δ in Definition 3.3, one has:*

1. $0 \leq \phi_\delta(M)(x) \leq ((w(M)/\delta) \wedge 1)$, *for any $x \in \mathbb{R}^n$,*

2. $\|\phi_\delta(M) - \phi_\delta(M')\|_\infty \leq 2(d_{\mathrm{I}}(M, M') \wedge \delta)/\delta$.

Proof. Claim 1. is a simple consequence of Equation (13).

Claim 2. for GS parameter (a) is a simple consequence of the triangle inequality.

Let us prove Claim 2. for (b). Let $x \in \mathbb{R}^n$ and $\delta > 0$. One has:

$$
\begin{aligned}
|\phi_\delta(M)(x) - \phi_\delta(M')(x)| &\leq \frac{2}{(2\delta)^n} \int_{\{y : y_n = 0\}} |d_{\mathrm{I}}\left(M\big|_{l_y \cap R_{x,\delta}}, 0\right) - d_{\mathrm{I}}\left(M'\big|_{l_y \cap R_{x,\delta}}, 0\right)| \, \mathrm{d}\lambda^{n-1}(y) \\
&\leq \frac{2}{(2\delta)^n} \int_{\{y : y_n = 0\}} d_{\mathrm{I}}\left(M\big|_{l_y \cap R_{x,\delta}}, M'\big|_{l_y \cap R_{x,\delta}}\right) \, \mathrm{d}\lambda^{n-1}(y) \\
&\leq 2(d_{\mathrm{I}}(M, M') \wedge \delta)/\delta,
\end{aligned}
$$

where the first inequality comes from Lemma B.2, the second inequality is an application of the triangle inequality, and the third inequality comes from Lemma A.2.

Finally, let us prove Claim 2. for (c). Let $x \in \mathbb{R}^n$ and $\delta > 0$. Let $b \leq d \in \mathrm{supp}\,(M) \cap R_{x,\delta}$. Let also $\gamma > 0$. Then, using Lemma B.2, one has:

$$
\begin{aligned}
\frac{1}{(2\delta)^n} \mathrm{vol}(\mathrm{supp}\,(M) \cap R_{b,d}) &= \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}(M\big|_{R_{b,d} \cap l_y}, 0) \, \mathrm{d}\lambda^{n-1}(y) \\
&\leq \frac{2}{\delta}\gamma + \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}(M\big|_{R_{b+\gamma, d-\gamma} \cap l_y}, 0) \, \mathrm{d}\lambda^{n-1}(y),
\end{aligned}
$$

using the convention $R_{a,b} = \varnothing$ if $a \not\leq b$. Now, set $\gamma := d_{\mathrm{I}}(M\big|_{R_{x,\delta}}, M'\big|_{R_{x,\delta}})$. If $b + \gamma$ or $d - \gamma \notin \mathrm{supp}\,(M')$ then $d_{\mathrm{I}}(M\big|_{R_{x,\delta}}, M'\big|_{R_{x,\delta}}) = \gamma > d_{\mathrm{I}}(M, M')$ which is impossible. Thus,

$$
\begin{aligned}
\frac{1}{(2\delta)^n} \mathrm{vol}(R_{b,d}) &\leq 2 d_{\mathrm{I}}(M\big|_{R_{x,\delta}}, M'\big|_{R_{x,\delta}})/\delta \\
&\quad + \sup_{a,c \in R_{x,\delta} \cap \mathrm{supp}(M')} \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}(M\big|_{R_{a,c} \cap l_y}, 0) \, \mathrm{d}\lambda^{n-1}(y) \\
&= 2 d_{\mathrm{I}}(M\big|_{R_{x,\delta}}, M'\big|_{R_{x,\delta}})/\delta + \phi_\delta(M')(x)
\end{aligned}
$$

Finally, taking the supremum on $b \leq d \in \mathrm{supp}\,(M) \cap R_{x,\delta}$ yields

$$
\phi_\delta(M)(x) - \phi_\delta(M')(x) \leq 2 d_{\mathrm{I}}(M\big|_{R_{x,\delta}}, M'\big|_{R_{x,\delta}})/\delta \leq 2\left(d_{\mathrm{I}}(M, M') \wedge \delta\right)/\delta.
$$

The desired inequality follows by symmetry on M and M'.

$\square$

Equipped with these results, we can finally prove Theorem 3.4.

Proof. Theorem 3.4.

Let $\mathbb{M} = \oplus_{i=1}^m M_i$ and $\mathbb{M}' = \oplus_{j=1}^{m'} M_j'$ be two modules that are decomposable into indicator modules and $x \in \mathbb{R}^n$.

Inequality 6. To simplify notations, we define the following: $w_i := w(M_i)$, $\phi_{i,x} := \phi_\delta(M_i)(x)$ and $w_j' := w(M_j')$, $\phi_{j,x}' := \phi_\delta(x, M_j')$. Let us also assume without loss of generality that the indices are consistent with a matching achieving the bottleneck distance. In other words, the bottleneck distance is achieved for a matching that matches M_i with M_i' for every i (up to adding 0 modules in

the decompositions of $\mathbb{M}$ and $\mathbb{M}'$ so that $m = m'$). Finally, assume without loss of generality that $\sum_i w_i' \geq \sum_i w_i$. Then, one has:

$$
|V_{1,\delta}(\mathbb{M})(x) - V_{1,\delta}(\mathbb{M}')(x)| = \left| \frac{1}{\sum_i w_i} \sum_i w_i \phi_{i,x} - \frac{1}{\sum_i w_i'} \sum_i w_i' \phi_{i,x}' \right|
$$
$$
\leq \frac{1}{\sum_i w_i'} \left| \sum_i w_i \phi_{i,x} - \sum_i w_i' \phi_{i,x}' \right| + \left| \frac{1}{\sum_i w_i} - \frac{1}{\sum_i w_i'} \right| \left| \sum_i w_i \phi_{i,x} \right|.
$$

Now, for any index i, since $d_{\mathrm{I}}(M_i, M_i') \leq d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}')$ and $|w_i - w_i'| \leq d_{\mathrm{I}}(M_i, M_i') \leq d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}')$ by Lemma B.1, Proposition B.3 ensures that:

$$
|w_i \phi_{i,x} - w_i' \phi_{i,x}'| \leq |w_i - w_i'|\phi_{i,x} + w_i'|\phi_{i,x} - \phi_{i,x}'| \leq 2(w_i + w_i')(d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta
$$

and

$$
\left| \frac{1}{\sum_i w_i} - \frac{1}{\sum_i w_i'} \right| \leq \frac{1}{\sum_i w_i'} \left| \frac{\sum_i w_i' - w_i}{\sum_i w_i} \right| \leq \frac{m d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}')}{(\sum_i w_i')(\sum_i w_i)}.
$$

Finally,

$$
|V_{1,\delta}(\mathbb{M})(x) - V_{1,\delta}(\mathbb{M}')(x)| \leq \left[\frac{\sum_i w_i + w_i'}{\sum_i w_i'} + \frac{\sum_i w_i}{\frac{1}{m}(\sum_i w_i)(\sum_i w_i')} \right] 2(d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta
$$
$$
\leq \left[4 + \frac{2}{C} \right](d_{\mathrm{B}}(\mathbb{M}, \mathbb{M}') \wedge \delta)/\delta.
$$

Inequality 5 can be proved using the proof of Inequality 6 by replacing every w_i by 1.

Inequality 7. Let us prove the inequality for (b). Let $R := R_{x-\boldsymbol{\delta}, x+\boldsymbol{\delta}}$. One has:

$$
V_{\infty,\delta}(\mathbb{M})(x) - V_{\infty,\delta}(\mathbb{M}')(x) =
$$
$$
\sup_i \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}(M_i|_{l_y \cap R}, 0) \, \mathrm{d}\lambda^{n-1}(y)
$$
$$
- \sup_j \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_I(M_j'|_{l_y \cap R}, 0) \, \mathrm{d}\lambda^{n-1}(y)
$$
$$
\textit{(for any index } j) \leq \sup_i \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} d_{\mathrm{I}}(M_i|_{l_y \cap R}, 0) - d_{\mathrm{I}}(M_j'|_{l_y \cap R}, 0) \, \mathrm{d}\lambda^{n-1}(y)
$$
$$
\leq \frac{2}{(2\delta)^n} \int_{\{y \in \mathbb{R}^n : y_n = 0\}} \sup_i \inf_j d_{\mathrm{I}}(M_i|_{l_y \cap R}, M_j'|_{l_y \cap R}) \, \mathrm{d}\lambda^{n-1}(y)
$$

Now, as the interleaving distance is equal to the bottleneck distance for single parameter persistence (Chazal et al., 2016b, Theorem 5.14), one has:

$$
\sup_i \inf_j d_{\mathrm{I}}(M_i|_{l_y \cap R}, M_j'|_{l_y \cap R}) \leq d_{\mathrm{B}}(\mathbb{M}|_{l_y \cap R}, \mathbb{M}'|_{l_y \cap R}) = d_{\mathrm{I}}(\mathbb{M}|_{l_y \cap R}, \mathbb{M}'|_{l_y \cap R}) \leq d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}') \wedge \delta
$$

which leads to the desired inequality. The proofs for (a) and (c) follow the same lines (upper bound the suprema in the right hand term with either infima or appropriate choices in order to reduce to the single parameter case). $\qquad\square$

C AN ADDITIONAL STABILITY THEOREM

In this section, we define a new GS representation, with a slightly different type of upper bound. It relies on the notations introduced in Appendix A. We also slightly abuse notations and use M_i to denote both an indicator module and its support.

Proposition C.1. *Let $\sigma > 0$, and let $0 \leq \delta \leq \delta(\mathbb{M})$, where*

$$
\delta(\mathbb{M}) := \inf\{\delta \geq 0 : \Gamma_{\mathbb{M}} \text{ achieves } d_{\mathrm{B}}(B_{\mathbf{e}_\Delta, x}, B_{\mathbf{e}_\Delta, x+\delta\mathbf{u}}) \text{ for all } x, \mathbf{u} \text{ s.t. } \|\mathbf{u}\|_\infty = 1, \langle \mathbf{e}_\Delta, \mathbf{u} \rangle = 0\},
$$

where $\Gamma_{\mathbb{M}}$ is the partial matching induced by the decomposition of $\mathbb{M}$. Let $\mathcal{N}(x, \sigma)$ denote the function $\mathcal{N}(x, \sigma) : \begin{cases} \mathbb{R}^n & \to \mathbb{R} \\ p & \mapsto \exp\left(-\frac{\|p-x\|^2}{2\sigma^2}\right) \end{cases}$ and let

$$V_{\delta,\sigma}(\mathbb{M}) : \begin{cases} \mathbb{R}^n & \to \mathbb{R} \\ x & \mapsto \max_{1 \le i \le m} \ \max_{f \in \mathcal{C}(x,\delta,M_i)} \ \|\mathcal{N}(x,\sigma) \cdot f\|_1 \end{cases} \tag{14}$$

where $\mathcal{C}(x, \delta, M_i)$ stands for the set of indicator functions from $\mathbb{R}^n$ to $\{0, 1\}$ whose support is $T_\delta(\ell) \cap M_i$, where ℓ is a connecting component of $\mathrm{im}(\ell_{\mathbf{e}_\Delta, x}) \cap M_i$ and $\mathbf{e}_\Delta = [1, \dots, 1] \in \mathbb{R}^n$, and where $T_\delta(\ell)$ is the δ-thickening of the line $L(\ell)$ induced by ℓ: $T_\delta(\ell) = \{x \in \mathbb{R}^k : \|x, L(\ell)\|_\infty \le \delta\}$.

Then, $V_{\delta,\sigma}$ satisfies the following stability property:

$$\|V_{\delta,\sigma}(\mathbb{M}) - V_{\delta,\sigma}(\mathbb{M}')\|_\infty \le (\sqrt{\pi}\sigma)^n \cdot \sqrt{2^{n+1}\delta^{n-1} d_{\mathrm{I}}(\mathbb{M}, \mathbb{M}') + C_n(\delta)}, \tag{15}$$

where $C_n(\cdot)$ is a continuous function such that $C_n(\delta) \to 0$ when $\delta \to 0$.

Proof. Let $\mathbb{M} = \oplus_{i=1}^m M_i$ and $\mathbb{M}' = \oplus_{j=1}^{m'} M_j'$ be two persistence modules that are decomposable into indicators, let $x \in \mathbb{R}^k$ and let $0 \le \delta \le \min\{\delta(\mathbb{M}), \delta(\mathbb{M}')\}$.

Notations. We first introduce some notations. Let N (resp. N') be the number of bars in $B_{\mathbf{e}_\Delta, x}$ (resp. $B'_{\mathbf{e}_\Delta, x}$), and assume without loss of generality that $N \le N'$. Let Γ be the partial matching achieving $d_{\mathrm{B}}(B_{\mathbf{e}_\Delta, x}, B'_{\mathbf{e}_\Delta, x})$. Let N_1 (resp. N_2) be the number of bars in $B_{\mathbf{e}_\Delta, x}$ that are matched (resp. not matched) to a bar in $B'_{\mathbf{e}_\Delta, x}$ under Γ, so that $N = N_1 + N_2$. Finally, note that $B_{\mathbf{e}_\Delta, x} = \{\ell : \exists i \text{ such that } \ell \in \mathcal{C}(\mathrm{im}(\ell_{\mathbf{e}_\Delta, x}) \cap M_i) \text{ and } \mathrm{im}(\ell_{\mathbf{e}_\Delta, x}) \cap M_i \ne \emptyset\}$, where $\mathcal{C}$ stands for the set of connected components (and similarly for $B'_{\mathbf{e}_\Delta, x}$), and let $F_\Gamma : B_{\mathbf{e}_\Delta, x} \to B'_{\mathbf{e}_\Delta, x}$ be a function defined on all bars of $B_{\mathbf{e}_\Delta, x}$ that coincides with Γ on the N_1 bars of $B_{\mathbf{e}_\Delta, x}$ that have an associated bar in $B'_{\mathbf{e}_\Delta, x}$, and that maps the N_2 remaining bars of $B_{\mathbf{e}_\Delta, x}$ to some arbitrary bars in the $(N' - N_1)$ remaining bars of $B'_{\mathbf{e}_\Delta, x}$.

A reformulation of the problem with vectors. We now derive vectors that allow to reformulate the problem in a useful way. Let $\hat{V}$ be the sorted vector of dimension N containing all weights $\|\mathcal{N}(x, \sigma) \cdot f\|_1$, where f is the indicator function whose support is $T_\delta(\ell) \cap M_i$ for some M_i, where $\ell \in B_{\mathbf{e}_\Delta, x}$ is a connected component of $\mathrm{im}(\ell_{\mathbf{e}_\Delta, x}) \cap M_i$. Now, let $\hat{V}'$ be the vector of dimension N' obtained by concatenating the two following vectors:

- the vector $\hat{V}_1'$ of dimension N whose ith coordinate is $\|\mathcal{N}(x, \sigma) \cdot f'\|_1$, where f' is the indicator function whose support is $T_\delta(\ell') \cap M_j'$ for some M_j', and $\ell' \in B'_{\mathbf{e}_\Delta, x}$ is the image under Γ of the bar $\ell \in B_{\mathbf{e}_\Delta, x}$ corresponding to the ith coordinate of $\hat{V}$, i.e., $\ell' = F_\Gamma(\ell)$ where $[\hat{V}]_i = \|\mathcal{N}(x, \sigma) \cdot f\|_1$ and f is the indicator function whose support is $T_\delta(\ell) \cap M_{i_0}$ for some M_{i_0}. In other words, $\hat{V}_1'$ is the (not necessarily sorted) vector of weights computed on the bars of $B'_{\mathbf{e}_\Delta, x}$ that are images (under the partial matching Γ achieving the bottleneck distance) of the bars of $B_{\mathbf{e}_\Delta, x}$ that were used to generate the (sorted) vector $\hat{V}$.

- the vector $\hat{V}_2'$ of dimension $(N' - N)$ whose jth coordinate is $\|\mathcal{N}(x, \sigma) \cdot f'\|_1$, where f' is an indicator function whose support is $T_\delta(\ell') \cap M_j'$ for some M_j', and $\ell' \in B'_{\mathbf{e}_\Delta, x}$ satisfies $\ell' \notin \mathrm{im}(F_\Gamma)$. In other words, $\hat{V}_2'$ is the vector of weights computed on the bars of $B'_{\mathbf{e}_\Delta, x}$ (in an arbitrary order) that are not images of bars of $B_{\mathbf{e}_\Delta, x}$ under Γ.

Finally, we let V be the vector of dimension N' obtained by filling $\hat{V}$ (whose dimension is $N \le N'$) with null values until its dimension becomes N', and we let $V' = \mathrm{sort}(\hat{V}')$ be the vector obtained after sorting the coordinates of $\hat{V}'$. Observe that:

$$|V_{\delta,\sigma}(\mathbb{M})(x) - V_{\delta,\sigma}(\mathbb{M}')(x)| = [V - V']_1 = [V - \mathrm{sort}(\hat{V}')]_1 \tag{16}$$

An upper bound. We now upper bound $\left\|V - \hat{V}'\right\|_\infty$. Let $q \in [\![1, N']\!]$. Then, one has $[V]_q = \|\mathcal{N}(x, \sigma) \cdot f\|_1$, where f is an indicator function whose support is $T_\delta(\ell) \cap M_i$ for some M_i with $\ell \in B_{\mathbf{e}_\Delta, x}$ if $q \leq N$ and $\ell = \emptyset$ otherwise; and similarly $[\hat{V}']_q = \|\mathcal{N}(x, \sigma) \cdot f'\|_1$, where f' is an indicator function whose support is $T_\delta(\ell') \cap M_j'$ for some M_j' with $\ell' \in B'_{\mathbf{e}_\Delta, x}$. Thus, one has:

$$
\begin{aligned}
[V - \hat{V}']_q &= |\, \|\mathcal{N}(x, \sigma) \cdot f\|_1 - \|\mathcal{N}(x, \sigma) \cdot f'\|_1 \,| \\
&\leq \|\mathcal{N}(x, \sigma) \cdot f - \mathcal{N}(x, \sigma) \cdot f'\|_1 \text{ by the reverse triangle inequality} \\
&= \|\mathcal{N}(x, \sigma) \cdot (f - f')\|_1 \text{ by linearity} \\
&\leq \|\mathcal{N}(x, \sigma)\|_2 \cdot \|f - f'\|_2 \text{ by Hölder's inequality} \\
&= (\sqrt{\pi}\sigma)^k \cdot \|f - f'\|_2
\end{aligned}
$$

Since $(f - f')$ is an indicator function whose support is $(T_\delta(\ell) \cap M_i) \triangle (T_\delta(\ell') \cap M_j')$, one has $\|f - f'\|_2 = \sqrt{|(T_\delta(\ell) \cap M_i) \triangle (T_\delta(\ell') \cap M_j')|}$. Given a segment ℓ and a vector $\mathbf{u}$, we let $\ell_\mathbf{u}$ denote the segment $\mathbf{u} + \ell$, and we let $\ell^\mathbf{u}$ denote the (infinite) line induced by $\ell_\mathbf{u}$. More precisely:

$$
\begin{aligned}
\|f - f'\|_2^2 &= |(T_\delta(\ell) \cap M_i) \triangle (T_\delta(\ell') \cap M_j')| \\
&= |\bigcup_\mathbf{u} (\ell^\mathbf{u} \cap M_i) \triangle \bigcup_\mathbf{u} ((\ell')^\mathbf{u} \cap M_j')| \\
&\qquad \text{where } \mathbf{u} \text{ ranges over the vectors such that } \|\mathbf{u}\|_\infty \leq \delta, \langle \mathbf{u}, \mathbf{e}_\Delta \rangle = 0 \\
&\leq \int_\mathbf{u} |(\ell^\mathbf{u} \cap M_i) \triangle ((\ell')^\mathbf{u} \cap M_j')| \mathrm{d}\mathbf{u} \\
&\leq \int_\mathbf{u} |(\ell^\mathbf{u} \cap M_i) \triangle (\ell \cap M_i)_\mathbf{u}| + |(\ell \cap M_i)_\mathbf{u} \triangle (\ell' \cap M_j')_\mathbf{u}| + |(\ell' \cap M_j')_\mathbf{u} \triangle ((\ell')^\mathbf{u} \cap M_j')| \mathrm{d}\mathbf{u} \\
&\leq \int_\mathbf{u} 4 d_\mathrm{B}(B_{\mathbf{e}_\Delta, x}, B_{\mathbf{e}_\Delta, x+\mathbf{u}}) + 4 d_\mathrm{B}(B_{\mathbf{e}_\Delta, x}, B'_{\mathbf{e}_\Delta, x}) + 4 d_\mathrm{B}(B'_{\mathbf{e}_\Delta, x}, B'_{\mathbf{e}_\Delta, x+\mathbf{u}}) \mathrm{d}\mathbf{u} \qquad (17) \\
&\leq 4 \int_\mathbf{u} \|\mathbf{u}\|_\infty + d_\mathrm{I}(\mathbb{M}, \mathbb{M}') + \|\mathbf{u}\|_\infty \, \mathrm{d}\mathbf{u} \text{ by Lemma A.2 and A.3}
\end{aligned}
$$

Inequality (17) comes from the fact that the symmetric difference between two bars (in two different barcodes) that are both matched (or unmatched) by the optimal partial matching is upper bounded by four times the bottleneck distance between the barcodes, and that (by assumption) the partial matchings achieving $d_\mathrm{B}(B_{\mathbf{e}_\Delta, x}, B_{\mathbf{e}_\Delta, x+\mathbf{u}})$ and $d_\mathrm{B}(B'_{\mathbf{e}_\Delta, x}, B'_{\mathbf{e}_\Delta, x+\mathbf{u}})$ are induced by $\mathbb{M}$ and $\mathbb{M}'$.

Conclusion. Finally, one has

$$
\begin{aligned}
|V_{\delta, \sigma}(\mathbb{M})(x) - V_{\delta, \sigma}(\mathbb{M}')(x)| = [V - V']_1 &= [V - \mathrm{sort}(\hat{V}')]_1 \text{ from Equation (16)} \\
&\leq \|V - V'\|_\infty \\
&\leq (\sqrt{\pi}\sigma)^k \cdot \sqrt{2^{n+1}\delta^{n-1} d_\mathrm{I}(\mathbb{M}, \mathbb{M}') + C_n(\delta)}, \qquad (18)
\end{aligned}
$$

with $C_n(\delta) = 8 \int_\mathbf{u} \|\mathbf{u}\|_\infty \mathrm{d}\mathbf{u} \to 0$ when $\delta \to 0$. Inequality (18) comes from the fact that any upper bound for the norm of the difference between a given vector $\hat{V}'$ and a sorted vector V, is also an upper bound for the norm of the difference between the sorted version V' of $\hat{V}'$ and the same vector V (see Lemma 3.9 in Carrière et al. (2015b)). $\qquad \square$

While the stability constant is not upper bounded by δ, $V_{\delta, \sigma}$ is more difficult to compute than the GS representations presented in Definition 3.3.

D ADDITIONAL RESULTS FOR UCR DATA SETS

In this section, we provide accuracy scores (obtained with the same parameters than Section 4.2) for several UCR data sets. The procedures for computing bifiltrations and the baseline scores were obtained from Carrière & Blumberg (2020). Baselines include random forest on the multiparameter

persistence kernel (MPK), landscape (MPL) and image (MPI), k-nearest neighbor classifier with Euclidean distance (B1) and dynamic time warping with optimized (B2) and constant width (B3).

Dataset	B1	B2	B3	MPK	MPL	MPI	Ours
DistalPhalanxOutlineAgeGroup	62.6	62.6	**77.0**	67.6	70.5	**71.9**	**71.9**
DistalPhalanxOutlineCorrect	71.7	**72.5**	71.7	74.6	69.6	71.7	**76.8**
DistalPhalanxTW	**63.3**	**63.3**	59.0	61.2	56.1	61.9	**65.4**
ProximalPhalanxOutlineAgeGroup	78.5	78.5	**80.5**	78.0	78.5	81.0	**85.9**
ProximalPhalanxOutlineCorrect	**80.8**	79.0	78.4	78.7	78.7	**81.8**	**81.8**
ProximalPhalanxTW	70.7	**75.6**	**75.6**	**79.5**	73.2	76.1	73.2
ECG200	**88.0**	**88.0**	77.0	77.0	74.0	**83.0**	80.0
ItalyPowerDemand	**95.5**	**95.5**	95.0	**80.7**	78.6	79.8	79.1
MedicalImages	68.4	**74.7**	73.7	55.4	55.7	60.0	**60.9**
Plane	96.2	**100.0**	**100.0**	92.4	84.8	**97.1**	**97.1**
SwedishLeaf	78.9	**84.6**	79.2	78.2	64.6	**83.8**	81.3
GunPoint	**91.3**	**91.3**	90.7	88.7	**94.0**	90.7	90.7
GunPointAgeSpan	89.9	**96.5**	91.8	93.0	85.1	90.5	**94.3**
GunPointMaleVersusFemale	97.5	97.5	**99.7**	**96.8**	88.3	95.9	94.0
GunPointOldVersusYoung	95.2	**96.5**	83.8	99.0	97.1	**100.0**	**100.0**
PowerCons	**93.3**	92.2	87.8	85.6	84.4	86.7	**90.0**
SyntheticControl	88.0	98.3	**99.3**	50.7	**60.3**	60.0	58.0

Table 3: Classification results for time series.

E   PSEUDO-CODE FOR OUR REPRESENTATIONS

In this section, we briefly present the pseudo-code that we use to compute our GS representations. In order to compute our representations, we implement the following procedures:

1. Given an indicator module M and a rectangle $R \subset \mathbb{R}^n$, compute the restriction $M\big|_R$,

2. Given a module M (or $M\big|_R$), compute $d_{\mathrm{I}}(M, 0)$,

3. Given a module restricted to a rectangle, compute the volume of the biggest rectangle in the support of this module,

4. Given a module M and a line l, compute the persistence diagram associated to $M\big|_l$.

For the first point, this correspond to "pushing" the corners in the rectangle.

Algorithm 1: Restriction of a module to a rectangle

Data: birth/death corners of a module M, and $R = \{z \in \mathbb{R}^n : m \le z \le M\}$
Result: The birth and death corners of $M\big|_R$.
for *Summand* **in** M **do**
    new_birth_list $\leftarrow$ [ ];
    for b **in** Summand_birth_corners **do**
        if $b \le M$ **then**
            $b' = \{\max(b_i, m_i) \text{ for } i \in [\![1, n]\!]\}$;
            Append b' to new_birth_list.
        end
    end
    new_death_list $\leftarrow$ [ ];
    for d **in** Summand_death_corners **do**
        if $d \ge m$ **then**
            $d' = \{\min(d_i, M_i) \text{ for } i \in [\![1, n]\!]\}$;
            Append d' to new_death_list.
        end
    end
    new_summands_corners $\leftarrow$ [new_birth_list, new_death_list];
end

For the second point, we proved in Lemma B.1 that $d_{\mathrm{I}}(M, 0)$ has a closed-form formula with corners.

Algorithm 2: Interleaving to 0

Data: birth/death corners of a module M
Result: The interleaving distance to 0 $d_I(M, 0)$.
result $\leftarrow 0$;
for *Summand* **in** M **do**
    for b **in** birthcorners **do**
        for d **in** deathcorners **do**
            result $\leftarrow \max\left(\text{result}, \frac{1}{2}\min_i(d_i - b_i)_+\right)$.
        end
    end
end

The 3rd point also has a closed-form formula with corners, leading to the following algorithm.

Algorithm 3: Volume of the biggest rectangle in $\operatorname{supp}(M)$

Data: birth/death corners of a module M
Result:
result $\leftarrow 0$;
for b **in** birthcorners **do**
    for d **in** deathcorners **do**
        result $\leftarrow \max\left(\text{result}, \Pi_i(d_i - b_i)_+\right)$.
    end
end

Finally, the following algorithm retrieves persistence diagrams of a multiparameter persistence module along any line, which is enough to compute the multiparameter persistence landscapes.

Algorithm 4: Restriction of a module to a line

Data: birth/death corners of a module M, and a diagonal line l
Result: $M\big|_l$
barcode $\leftarrow [\,]$;
$y \leftarrow$ a point in l;
for *Summand* **in** M **do**
    birth $\leftarrow y + \mathbf{1} \times \min_{b \in \text{summand\_birth\_corners}} \max_i b_i - y_i$.
    death $\leftarrow y + \mathbf{1} \times \max_{d \in \text{summand\_death\_corners}} \min_i d_i - y_i$.
    bar $\leftarrow [\text{birth}, \text{death}]$.
    Append bar to barcode.
end