HOW DOES SHARPNESS-AWARE MINIMIZATION MINIMIZE SHARPNESS?

ABSTRACT

Sharpness-Aware Minimization (SAM) is a highly effective regularization technique for improving the generalization of deep neural networks for various settings. However, the underlying working of SAM remains elusive because of various intriguing approximations in the theoretical characterizations. SAM intends to penalize a notion of sharpness of the model but implements a computationally efficient variant; moreover, a third notion of sharpness was used for proving generalization guarantees. The subtle differences in these notions of sharpness can indeed lead to significantly different empirical results. This paper rigorously nails down the exact sharpness notion that SAM regularizes and clarifies the underlying mechanism. We also show that the two steps of approximations in the original motivation of SAM individually lead to inaccurate local conclusions, but their combination accidentally reveals the correct effect, when full-batch gradients are applied. Furthermore, we also prove that the stochastic version of SAM in fact regularizes the third notion of sharpness mentioned above, which is most likely to be the preferred notion for practical performance. The key mechanism behind this intriguing phenomenon is the alignment between the gradient and the top eigenvector of Hessian when SAM is applied.

1 INTRODUCTION

Modern deep nets are often overparametrized and have the capacity to fit even randomly labeled data (Zhang et al., 2016). Thus, a small training loss does not necessarily imply good generalization. Yet, standard gradient-based training algorithms such as SGD are able to find generalizable models. Recent empirical and theoretical studies suggest that generalization is well-correlated with the sharpness of the loss landscape at the learned parameter (Keskar et al., 2016; Dinh et al., 2017; Dziugaite & Roy, 2017; Neyshabur et al., 2017; Jiang et al., 2019). Partly motivated by these studies, Foret et al. (2021); Wu et al. (2020); Zheng et al. (2021) propose to penalize the sharpness of the landscape to improve the generalization. We refer this method to *Sharpness-Aware Minimization* (SAM) and focus on the version of Foret et al. (2021) in this paper.

Despite its empirical success, the underlying working of SAM remains elusive because of the various intriguing approximations made in its derivation and analysis. There are three different notions of sharpness involved – SAM intends to optimize the first notion, the sharpness along the worst direction but actually implements a computationally efficient notion, the sharpness along the direction of the gradient. But in the analysis, a third notion of sharpness is actually used to prove generalization guarantees, which admits the first notion as an upper bound. The subtle difference between the three notions can lead to very different explicit biases. (see Figure 1 for demonstraion)

More concretely, let L be the training loss, x be the parameter and ρ be the *perturbation radius*, a hyperparameter requiring tuning. The first notion corresponds to the following optimization problem (1), where we call $R_\rho^{\text{Max}}(x) = L_\rho^{\text{Max}}(x) - L(x)$ the *worst-direction sharpness* at x and thus SAM is intended to minimize the original training loss plus the worst-direction sharpness at x.

$$\min_x L_\rho^{\text{Max}}(x), \quad \text{where} \quad L_\rho^{\text{Max}}(x) = \max_{\|v\|_2 \leq 1} L(x + \rho v), \tag{1}$$

However, even evaluation of $L_\rho^{\text{Max}}(x)$ is computationally expensive, not to mention optimization. Thus Foret et al. (2021); Zheng et al. (2021) propose to approximate the worst perturbation direction by the direction of the gradient and implement the second notion of sharpness, which corresponds

Type of Sharpness	Symbol	Definition	Explicit Bias Among Minimizers
Worst-direction	L_ρ^{Max}	$\max_{\|v\|_2 \leq 1} L(x + \rho v)$	$\min_x \lambda_1(\nabla^2 L(x))$
Ascent-direction	L_ρ^{Asc}	$L\left(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right)$	$\min_x \lambda_{\min}(\nabla^2 L(x))$
Average-direction	L_ρ^{Avg}	$\mathbb{E}_{v \sim N(0,I)} L(x + \rho \frac{v}{\|v\|_2})$	$\min_x \text{Tr}(\nabla^2 L(x))$

Table 1: Definitions and explicit biases of different notions of sharpness. Here $\lambda_{\min}$ denotes to the smallest *non-zero* eigenvalue.

to (2). We call $R_\rho^{\text{Asc}}(x) = L_\rho^{\text{Asc}}(x) - L(x)$ the *ascent-direction sharpness* at x.

$$\min_x L_\rho^{\text{Asc}}(x), \quad \text{where} \quad L_\rho^{\text{Asc}}(x) = L\left(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right). \tag{2}$$

Intriguingly, in the generalization analysis of SAM by Foret et al. (2021); Wu et al. (2020), the first notion of sharpness, *i.e.*, the worst-direction sharpness, is only used for upper bounding the third notion of sharpness via the PAC Bayesian theory (McAllester, 2003). We call the third notion $R_\rho^{\text{Avg}}(x) = L_\rho^{\text{Avg}}(x) - L(x)$ the *average-direction sharpness* at x, where $L_\rho^{\text{Avg}}(x) = \mathbb{E}_{g \sim N(0,I)} L(x + \rho g/\|g\|)$.

For further acceleration, Foret et al. (2021); Zheng et al. (2021) omit the gradient through other occurrence of x and approximate the gradient of ascent-direction sharpness by gradient taken after one-step ascent, *i.e.*, $\nabla L_\rho^{\text{Asc}}(x) \approx \nabla L\left(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right)$ and derive the update rule of SAM, where η is the learning rate.

$$\text{Sharpness-Aware Minimization (SAM):} \quad x(t+1) = x(t) - \eta \nabla L\left(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right) \tag{3}$$

In this paper, we analyze the explicit bias of various notions of sharpness and the optimization trajectory of SAM. Our analysis is performed for small perturbation radius ρ and learning rate η under the setting where the minimizers of loss form a manifold following Fehrman et al. (2020); Li et al. (2021) In particular, we make the following theoretical contributions.

1. We prove that full-batch SAM does minimize worst-direction sharpness. (Theorem 4.9)
2. Surprisingly, when batch size is 1, SAM minimizes average-direction sharpness. (Theorem 4.9)
3. We characterize the explicit biases of three notions of sharpness among minimizers when perturbation radius ρ goes to zero. (Theorems 4.5 to 4.7, also see Table 1) Surprisingly, both heuristic approximations made for the update rule of SAM lead to inaccurate solutions, that is, (1) minimizing worst-direction sharpness and ascent-direction sharpness induce different biases among minimizers, and (2) SAM doesn't minimize ascent-direction sharpness.

The key mechanism behind this implicit bias of SAM is the implicit alignment between the gradient and the top eigenvector of Hessian when running SAM.

2 RELATED WORKS

Sharpness and Generalization. The study on the connection between sharpness and generalization can be traced back to Hochreiter & Schmidhuber (1997). Keskar et al. (2016) observed a positive correlation between the batch size, the generalization error, and the sharpness of the loss landscape when changing the batch size. Jastrzebski et al. (2017) extends this by finding a correlation of the sharpness and the ratio between learning rate to batch size. Dinh et al. (2017) show that one can easily construct networks with good generalization but with arbitrary large sharpness by reparametrization. Dziugaite & Roy (2017); Neyshabur et al. (2017); Wei & Ma (2019) give theoretical guarantees on the generalization error using sharpness-related measures. Jiang et al. (2019) perform a large-scale empirical study on various generalization measures and showed that sharpness-based measures have the highest correlation with generalization.

Background on Sharpness Aware-Minimization. Foret et al. (2021); Zheng et al. (2021) concurrently proposed to minimize the loss at the perturbed from current parameter towards the worst direction to improve generalization. Wu et al. (2020) propose the almost identical method for a different purpose, robust generalization of adversarial training. Kwon et al. (2021) proposed a different metric for SAM to fix the rescaling problem pointed out by Dinh et al. (2017). Liu et al. (2022) propose an more computationally efficient version of SAM. Zhao et al. (2022) proposed to improve generalization by penalizing gradient norm. Their proposed algorithm can be viewed as a

Figure 1: Visualization of the different biases of different sharpness notions on a 4D-toy example. For $x, y \in \mathbb{R}^2$, consider loss $L(x, y) = F_0(x)y_0^2 + F_1(x)y_1^2$ with $F_0(x) = x_0^2 + 6x_1^2 + 8$ and $F_1(x) = 4(1 - x_0)^2 + (1 - x_1)^2 + 1$. L has a zero loss manifold $\{y = 0\}$ and the eigenvalues of its Hessian on manifold are $F_0(x)$ and $F_1(x)$ with $F_0(x) \geq 8 > 6 \geq F_1(x)$ on $[0, 1]^2$. As our theory predicts, (1). full-batch SAM (3) finds the minimizer with the smallest top eigenvalue, $F_0(x)$; (2). GD on Ascent-direction Loss L_ρ^{Asc} (2) finds the minimizer with the smallest bottom eigenvalue, $F_0(x)$. (3). Stochastic SAM (16) (with $L_0(x, y) = F_0(x)x_0^2, L_1(x, y) = F_1(x)y_1^2$) finds the minimizer with smallest trace of Hessian. Loss landscape $L(x, \cdot)$ are visualized as 3D plots at converged x to illustrate the different biases. (*cf.*Table 1)

generalization of SAM. Andriushchenko & Flammarion (2022) studied a variant of SAM where the step size of ascent step is ρ instead of $\frac{\rho}{\|\nabla L(x)\|_2}$. They showed that for a simple model this variant of SAM has a stronger regularization effect when batch size is 1 compared to the full-batch case and argued that this might be the explanation that SAM generalizes better with small batch sizes.

3 NOTATIONS AND ASSUMPTIONS

For any integer k, we define $\mathcal{C}^k$ as the set of k times continuously differentiable functions. For any mapping F, we define $\partial F(x)[u]$ and $\partial^2 F(x)[u, v]$ as the first and second order directional derivative of $F(x)$ along the direction u (and v). Given a differential submanifold Γ of $\mathbb{R}^D$ and a point $x \in \Gamma$, define $P_{x,\Gamma}$ as the projection operator onto the manifold of the normal space of Γ at x and $P_{x,\Gamma}^\perp = I_D - P_{x,\Gamma}$. We fix our initialization as x_{init} and our loss function as $L : \mathbb{R}^D \rightarrow \mathbb{R}$. Given the loss function, the gradient flow can be defined as mapping $\phi : \mathbb{R}^D \times [0, \infty) \rightarrow \mathbb{R}^D$ satisfying $\phi(x, \tau) = x - \int_0^\tau \nabla L(\phi(x, t))dt$. We further define the limiting map $\Phi : \mathbb{R}^D \rightarrow \mathbb{R}^D$ as $\Phi(x) = \lim_{\tau \rightarrow \infty} \phi(x, \tau)$. For any positive definite symmetry matrix $A \in \mathbb{R}^{D \times D}$, define $\{\lambda_i(A), v_i(A)\}_{i \in [D]}$ as all its eigenvalues and eigenvectors satisfying $\lambda_1(A) \geq \lambda_2(A)... \geq \lambda_3(A)$ and $\|v_i(A)\|_2 = 1$. Our analysis assumes sufficiently small η and ρ and uses $O(\cdot)$ to hide constant.

Following Fehrman et al. (2020); Li et al. (2021); Arora et al. (2022), we make the below assumption.

Assumption 3.1. *Assume loss $L : \mathbb{R}^D \rightarrow \mathbb{R}$ belongs to $\mathcal{C}^4$, and there exists a manifold Γ that is $D - M$ dimensional $\mathcal{C}^2-$submanifold of $\mathbb{R}^D$ for some integer $1 \leq M \leq D$, where for all $x \in \Gamma$, x is a global minimizer of L, $L(x) = 0$ and $rank(\nabla^2 L(x)) = M$.*

The smoothness assumption is meet with networks with smooth activation functions and the existence of the manifold is due to the vast overparameterization of the modern neural network. The full rank assumption is necessary for the analysis to guarantee the differentiability of Φ.

Definition 3.2 (Attraction Set). *Let U be the attraction set of Γ under gradient flow, that is, a neighborhood of Γ containing all points starting from which gradient flow w.r.t. loss L converges to some point in Γ, or mathematically, $U \triangleq \{x \in \mathbb{R}^D | \Phi(x) \text{ exists and } \Phi(x) \in \Gamma\}$.*

Assumption 3.1 implies that U is open and Φ is in $\mathcal{C}^3$ on U (from Lemma B.15 Arora et al. (2022)).

4 EXPLICIT AND IMPLICIT BIAS IN THE FULL-BATCH SETTING

Section 4.1 provides a general theorem to properly analyze the explicit bias of various notions of sharpness among different minimizers. We then apply our machinery on *ascent-direction sharpness* and *worst-direction sharpness* and show that they have different explicit biases. In Section 4.2 we provide our main theorem in the full-batch setting, that SAM implicitly minimizes the worst-direction sharpness, via characterizing its limiting dynamics as learning rate ρ and η goes to 0 with

a Riemmanian gradient flow with respect to the top eigenvalue of the Hessian of the loss on the manifold of global minimizers. In Section 4.3 we sketch the proof of the implicit bias of SAM and identified a key property behind the implicit bias, which is the implicit alignment between the gradient and the top eigenvector of the Hessian throughout the training.

4.1 WORST- AND ASCENT-DIRECTION SHARPNESS HAVE DIFFERENT EXPLICIT BIAS

The intuition of approximating R_ρ^{Max} by R_ρ^{Asc} comes from the following Taylor expansions (Foret et al., 2021; Wu et al., 2020), which works for small ρ:

$$R_\rho^{\text{Max}}(x) = \sup_{\|v\|_2 \leq 1} L(x + \rho v) - L(x) = \sup_{\|v\|_2 \leq 1} \left(\rho v^\top \nabla L(x) + \frac{\rho^2}{2} v^\top \nabla^2 L(x) v + O(\rho^3) \right) \quad (4)$$

$$R_\rho^{\text{Asc}}(x) = L\left(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2} \right) - L(x) = \rho \|\nabla L(x)\|_2 + \frac{\rho^2}{2} \frac{\nabla L(x)^\top \nabla^2 L(x) \nabla L(x)}{\|\nabla L(x)\|_2^2} + O(\rho^3) \quad (5)$$

Here, the preference among the global minima is what we are mainly concerned with. Since $\sup_{\|v\|_2 \leq 1} v^\top \nabla L(x) = \|\nabla L(x)\|_2$ when $\|\nabla L(x)\|_2 > 0$, the leading terms in Equations (4) and (5) are both the first order term, $\rho \|\nabla L(x)\|_2$, and are the same. However, it is erroneous to think that the first order term decides the explicit bias, as the first order term $\|\nabla L(x)\|_2$ vanishes at the global minimizers of the loss L and thus the second order term becomes the leading term. Any global minimizer x of the original loss L is an $O(\rho^2)$-approximate minimizer of the sharpness-aware loss because $\nabla L(x) = 0$. Therefore, the sharpness-aware loss needs to be of order ρ^2 so that we can guarantee the second-order terms in Equation (4) and/or Equation (5) to be non-trivially small.

In this section, we aim to understand under Assumption 3.1 what the explicit biases of various notions of sharpness among different minimizers are. Theorem 4.3 will be our main theoretic tool to analyze the explicit bias for small perturbation radius ρ. Below we first introduce some notations and assumption.

Notation for Regularizers. Let $R_\rho : \mathbb{R}^D \to \mathbb{R} \cup \{\infty\}$ be a family of regularizers parameterized by ρ. If R_ρ is not well-defined at some x, then we let $R_\rho(x) = \infty$. This convention will be useful when analyzing ascent-direction sharpness $R_\rho^{\text{Asc}} = L_\rho^{\text{Asc}} - L$ which is not defined when $\nabla L(x) = 0$. This convention will not change the minimizers of the regularized loss. Intuitively, a regularizer should always be non-negative, but however, when far away from manifold, regularizer $R_\rho(x)$ can actually be negative, *e.g.*, $R_\rho^{\text{Avg}}(x) \approx \frac{\rho^2}{2D} \text{Tr}[\nabla^2 L(x)]$. Therefore we make the following assumption to allow the regularizer to be mildly negative.

Assumption 4.1. *Suppose for any bounded closed set $B \subset U$, there exists $C > 0$, such that for sufficiently small ρ, $\forall x \in B, R_\rho(x) \geq -C\rho^2$.*

The following concept of limiting regularizer is of crucial role in our analysis.

Definition 4.2 (Limiting Regularizer). *We define the* limiting regularizer *of $\{R_\rho\}$ as the function*

$$S : \Gamma \to \mathbb{R}, \quad S(x) = \lim_{\rho \to 0} \lim_{r \to 0} \inf_{\|x' - x\|_2 \leq r} R_\rho(x')/\rho^2.$$

We say the limiting regularizer S of $\{R_\rho\}$ is a good *around some $x^* \in \Gamma$, if there is an open set V containing x^*, such that S is a non-negative continuous function in V and for any $\epsilon > 0$, there is some $\rho_{x^*} > 0$, it holds that $\forall x \in \Gamma \cap V, \rho \leq \rho_{x^*}, \left| S(x) - \inf_{\|x' - x\|_2 \leq \epsilon \cdot \rho} R_\rho(x')/\rho^2 \right| < \epsilon$. We say the limiting regularizer S is* good *on Γ, if S is good around every point $x \in \Gamma$.*

The high-level intuition behind the definition of limiting regularizer is that we want to use it to capture the second order term in the Taylor expansion of regularizer R_ρ when $\rho \to 0$. When the second order term is continuous in x, the definition of $S(x)$ can also be simplified as $R_\rho(x)/\rho^2$. The intuition of the concept of a good limiting regularizer is that, the regularizer should not change very fast, especially in an $O(\rho)$ neighborhood of the minimizer. If so, the minimizer of the regularized loss may be $\Omega(\rho)$ away from any minimizer to reduce the regularizer at the cost of increasing the original loss, which makes the limiting regularizer unable to capture the explicit bias of the regularizer.

Theorem 4.3. *Let U' be any bounded open set such that its closure $\overline{U'}$ is contained in U and that $\overline{U'} \cap \Gamma = \overline{U'} \cap \overline{\Gamma}$. Then for any family of parametrized regularizers $\{R_\rho\}$ admitting a good limiting*

regularizer on Γ and satisfying Assumption 4.1 and any $\epsilon \geq 0$, there is a $\rho_0 > 0$, such that for all $u \in U'$ and $\rho < \rho_0$, it holds that

$$L(u) + R_\rho(u) \leq \inf_{x \in U'} \left(L(x) + R_\rho(x) \right) + \epsilon\rho^2 + o(\rho^2)$$

$$\iff \left(L(u) - \inf_{x \in U'} L(x) \right) + \left| R_\rho(u) - \rho^2 \inf_{x \in U' \cap \Gamma} S(x) \right| \leq \epsilon\rho^2 + o(\rho^2)$$

For the applications we are interested in in this paper, the good limiting regularizer S can be continuously extended to the entire space $\mathbb{R}^D$. In such a case, the implication of "$\implies$" of Theorem 4.3 also admits the following alternative form which doesn't involve R_ρ. Corollary 4.4 implies minimizing regularized loss $L(x) + R_\rho(x)$ is equivalent to minimizing the limiting regularizer of $\{R_\rho\}_\rho$, $S(x)$ on the global minimizer manifold Γ.

Corollary 4.4. *Under the setting of Theorem 4.3, let $\overline{S}$ be an continuous extension of S to $\mathbb{R}^d$, if $L(u) + R_\rho(u) \leq \inf_{x \in U'} \left(L(x) + R_\rho(x) \right) + \epsilon\rho^2 + o(\rho^2)$, then we have that $L(u) - \inf_{x \in U'} L(x) = O(\rho^2)$ and that $\left| \overline{S}(u) - \inf_{x \in U' \cap \Gamma} \overline{S}(x) \right| = \epsilon + o(1)$.*

Corollary 4.4 suggests a sharp phase transition of the property of the solution of $\min_x L(x) + R_\rho(x)$ when the optimization error drops from $\omega(\rho^2)$ to $O(\rho^2)$. when the optimization error is larger than $\omega(\rho^2)$, no regularization effect happens and any minimizer satisfies the requirement. When the error becomes $O(\rho^2)$, there is a non-trivial restriction on the (extended) limiting regularizer.

Below we show all of worst-, ascent- and average-direction sharpness admit good limiting regularizers. Thus we can apply Corollary 4.4 to characterize their explicit biases, which are all different.

Theorem 4.5. *Worst-direction sharpness R_ρ^{Max} satisfies Assumption 4.1 and admits $\lambda_1(\nabla^2 L(\cdot))/2$ as a good limiting regularizer on Γ.*

Theorem 4.6. *Ascent-direction sharpness R_ρ^{Asc} satisfies Assumption 4.1 and admits $\lambda_M(\nabla^2 L(\cdot))/2$ as a good limiting regularizer on Γ.*

Theorem 4.7. *Average-direction sharpness R_ρ^{Avg} satisfies Assumption 4.1 and admits $\text{Tr}(\nabla^2 L(\cdot))/(2D)$ as a good limiting regularizer on Γ.*

When R_ρ is continuous at some $x \in \Gamma$, the definition of $S(x)$ can be simplified as $\lim_{\rho \to 0} R_\rho(x)/\rho^2$. Worst- and average- direction sharpness fall into this type and the limiting regularizer can be solved straightforwardly by Taylor expansion, which is well-known in literature (Keskar et al., 2016; Kaur et al., 2022; Zhuang et al., 2022; Orvieto et al., 2022). Theorems 4.5 and 4.7 can be viewed as a restatement of the existing results in our manifold setting.

Our major contribution in this subsection is the analysis for ascent-direction sharpness, which is more tricky as $R_\rho^{\text{Asc}}(x) = \infty$ and thus is not continuous for any $x \in \Gamma$. To minimize R_ρ^{Asc} around x, we can pick $x' \to x$ to make $\|\nabla L(x)\|_2 \to 0$ but not equal to 0. By (5), we have $R_\rho^{\text{Asc}}(x') \approx \rho^2/2 \cdot \nabla L(x')^\top \nabla^2 L(x) \nabla L(x')/\|\nabla L(x')\|^2$. Here the crucial step of the proof is that because of Assumption 3.1, $\nabla L(x)/\|\nabla L(x)\|_2$ must almost lie in the column span of $\nabla^2 L(x)$, and thus implies $\inf_{x'} \nabla L(x')^\top \nabla^2 L(x) \nabla L(x')/\|\nabla L(x')\|^2 \overset{\rho \to 0}{\to} \lambda_M(\nabla^2 L(x))$. The above alignment property between the gradient and the column space of Hessian can be checked directly for any non-negative quadratic function and the maximal Hessian rank assumption in Assumption 3.1 ensures this property extends to general losses.

4.2 SAM Provably Decreases Worst-direction Sharpness Locally

Though ascent-direction sharpness has different explicit bias from worst-direction sharpness, in this subsection we will show that surprisingly, SAM, an heuristic method designed to minimize ascent-direction sharpness, provably decreases worst-direction sharpness. The main result here is an exact characterization of the trajectory of SAM (3) via the following ordinary differential equation (ODE) (6), when learning rate η and perturbation radius ρ are small and the initialization $x(0) = x_{\text{init}}$ is in U. We call the solution of (6) the *limiting flow* of SAM, which is exactly the Riemannian Gradient Flow on the manifold Γ with respect to $\lambda_1(\nabla^2 L(\cdot))$. In other words, the ODE (6) is essentially

a projected gradient descent algorithm with loss $\lambda_1(\nabla^2 L(\cdot))$ on the constraint set Γ and an infinitesimal learning rate.

$$X(\tau) = X(0) - \frac{1}{2}\int_{s=0}^{\tau} P_{X(s),\Gamma}^{\perp} \nabla\lambda_1(\nabla^2 L(X(s)))ds, X(0) = \Phi(x_{\text{init}}). \tag{6}$$

Note $\lambda_1(\nabla^2 L(x))$ may not be differentiable at x if $\lambda_1(\nabla^2 L(x)) = \lambda_2(\nabla^2 L(x))$, thus to ensure the (6) is well-defined, we assume there is a positive eigengap for L on Γ. Assuming ODE (6) has a solution till time T_3, we have Theorem 4.9, which is the main theorem of this section.

Assumption 4.8. *For $x \in \Gamma$, there exists a positive eigengap,i.e., $\lambda_1(\nabla^2 L(x)) > \lambda_2(\nabla^2 L(x))$.*

Theorem 4.9 (Main, Theorems C.35 and C.36 stated informally). *Let $\{x(t)\}$ be the iterates of SAM (3) with $x(0) = x_{init} \in U$, then under Assumptions 3.1 and 4.8, for all η, ρ such that $\eta \ln(1/\rho)$ and ρ/η are sufficiently small, the dynamics of SAM can be split into two phases:*

- **Phase I:** *SAM follows Gradient Flow with respect to L until entering an $O(\eta\rho)$ neighborhood of the manifold Γ in $\tilde{O}(\frac{1}{\eta})$ steps;*
- **Phase II:** *SAM tracks the solution X of (6), the Riemannian Gradient Flow with respect to $\lambda_1(\nabla^2 L(\cdot))$ in the $O(\eta\rho)$ neighborhood in the sense that $\max_{0 \le T \le T_3}\|\Phi(x(\lceil T/(\eta\rho^2)\rceil)) - X(T)\| = O((\eta + \rho)\log(1/\eta\rho))$. Moreover, the angle between $\nabla L(x(t))$ and $v_1(\nabla^2 L(x(t)))$ is at most $O(\rho)$.*

Theorem 4.9 shows that SAM decreases the largest eigenvalue of Hessian of loss locally around the manifold of global minimizers. As a corollary of Theorem 4.9, we can also show that the largest eigenvalue of the limiting flow closely tracks the regularized training loss.

Corollary 4.10. *For all $T_3' > 0$, for all ρ, η such that $\eta \ln(1/\rho)$ and ρ/η are sufficiently small we have $\forall T_3' < \eta\rho^2 t \le T_3$, $\|R_\rho^{\text{Max}}(x(t)) - \rho^2\lambda_1(\nabla^2 L(X(\eta\rho^2 t))/2\| = \tilde{O}(\eta\rho^2)$*

Recall Theorem 4.6 shows that the largest eigenvalue of Hessian is the limiting regularizer of the *worst-direction sharpness*, leveraging the equivalence relationship in Theorem 4.3, below we show that full-batch SAM provably minimizes *worst-direction sharpness* if we additionally assume the limiting flow converges to a local minimizer of the top eigenvalue of Hessian.

Corollary 4.11. *Define U' as in Theorem 4.3, suppose $X(\infty) = \lim_{t\to\infty} X(t)$ exists and is a minimizer of $\lambda_1(\nabla^2 L(x))$ in $U' \cap \Gamma$, for all $\epsilon > 0$, there exists a constant $T > 0$, then for all ρ, η such that $\eta \ln(1/\rho)$ and ρ/η are sufficiently small we have $L_\rho^{\text{Max}}(x(\lceil T/(\eta\rho^2)\rceil)) \le \epsilon\rho^2 + \inf_{x\in U'} L_\rho^{\text{Max}}(x)$*

4.3 PROOF SKETCH FOR PHASE II IN THEOREM 4.9

Now we sketch the proof for the ODE-based characterization of the trajectory of SAM in Phase II. The framework of the analysis is similar to Arora et al. (2022); Lyu et al. (2022), where the high-level idea is to use $\Phi(x(t))$ as a proxy for $x(t)$ and study the dynamics of $\Phi(x(t))$ via Taylor expansion, which turns out to be dependent on the alignment between the gradient and the eigenvectors of the Hessian. In particular, like Arora et al. (2022); Lyu et al. (2022), we show that the gradient aligns to the top eigenvector of Hessian, and thus encourages the SAM dynamics to reduce the top eigenvalue of the Hessian.

Taylor Expansion on Φ. In Phase II, it can be shown that $\|x(t) - \Phi(x(t))\| = O(\eta\rho)$ holds for every step, this implies $\|x(t+1) - x(t)\|_2 = O(\rho\eta)$. (See Lemma C.12) Therefore we have that

$$\Phi(x(t+1)) - \Phi(x(t)) = -\eta\partial\Phi(x(t))(x(t+1) - x(t)) + O(\eta\|x(t+1) - x(t)\|^2)$$

$$= -\eta\partial\Phi(x(t))\nabla L\left(x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right) + O(\eta^3\rho^2) \tag{7}$$

Now we apply Taylor expansion on $\nabla L\left(x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right)$ around x and get

$$\nabla L\left(x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right)$$
$$= \nabla L(x) - \rho\nabla^2 L(x)\frac{\nabla L(x)}{\|\nabla L(x)\|_2} + \frac{\rho^2}{2}\partial^2(\nabla L)(x)\left[\frac{\nabla L(x)}{\|\nabla L(x)\|_2}, \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right] + O(\rho^3). \tag{8}$$

Lemma 4.12 (Lemma B.16 Arora et al. (2022)). *For $x \in U$, $\partial\Phi(x)\nabla L(x) = 0$, $\partial\Phi(x)\nabla^2 L(x)\nabla L(x) = -\partial^2\Phi(x)[\nabla L(x), \nabla L(x)]$.*

Now we plug (8) into (7) and simplify the expression using Lemma 4.12. We have that

$$\Phi(x(t+1)) - \Phi(x(t)) = 0 + O(\rho \left\|\partial\Phi(x(t))\nabla^2 L(x(t))\right\| \left\|\nabla L(x(t))\right\|_2) \tag{9}$$
$$- \frac{\eta\rho^2}{2}\partial\Phi(x)\partial^2(\nabla L)(x)\left[\frac{\nabla L(x)}{\|\nabla L(x)\|_2}, \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right] + O(\eta^3\rho^2 + \eta\rho^3)$$

where in the last step we use the property that $\|x(t) - \Phi(x(t))\| = O(\eta\rho)$, which further implies $\|\nabla L(x(t))\|_2 = O(\eta\rho)$ in Phase II (See Lemma C.7).

Lemma 4.13. *For $x \in \Gamma$, $\partial\Phi(x) = P_{x,\Gamma}^\perp$, $\partial\Phi(x)\nabla^2 L(x) = 0$, where $P_{x,\Gamma}^\perp$ is the orthogonal projection matrix of the tangent space of Γ at x.*

Next we use Lemma 4.13 to further simplify (9). Since the entire phase II happens in an $O(\eta\rho)$-neighborhood of manifold Γ, we have that $\|x(t) - \Phi(x(t))\|_2 = O(\eta\rho)$, thus both $\left\|\partial\Phi(x(t))\nabla^2 L(x(t))\right\|$ and $\|\nabla L(x(t))\|_2$ are $O(\eta\rho)$. So far, the proof is almost completed, with the implicit alignment between gradient and top eigenvector of Hessian being the last missing piece. Suppose we have $\left\|\frac{\nabla L(x)}{\|\nabla L(x)\|_2} - v_1(\nabla^2 L(x))\right\| = O(\rho)$, then it holds that

$$\Phi(x(t+1)) - \Phi(x(t)) = -\frac{\eta\rho^2}{2}\partial\Phi(x)\partial^2(\nabla L)(x)\left[\frac{\nabla L(x)}{\|\nabla L(x)\|_2}, \frac{\nabla L(x)}{\|\nabla L(x)\|_2}\right] + O(\eta^3\rho^2 + \eta\rho^3)$$
$$= -\frac{\eta\rho^2}{2}\partial\Phi(x)\partial^2(\nabla L)(x)\left[v_1(\nabla^2 L(x)), v_1(\nabla^2 L(x))\right] + O(\eta^3\rho^2 + \eta\rho^3)$$
$$= -\frac{\eta\rho^2}{2}\partial\Phi(x(t))\nabla\lambda_1(\nabla^2 L(x(t))) + O(\eta^3\rho^2 + \eta\rho^3)$$
$$= -\frac{\eta\rho^2}{2}\partial\Phi(\Phi(x(t)))\cdot\nabla\lambda_1(\nabla^2 L(x))|_{x=\Phi(x(t))} + O(\eta^3\rho^2 + \eta\rho^3), \tag{10}$$

where the second to last step we use the following property about the derivative of eigenvalue (Lemma C.60) and the last step is due to Taylor expansion.

Implicit Hessian-gradient Alignment. It remains to explain why the gradient implicitly aligns to the top eigenvector of Hessian, which is the key component of the analysis in Phase II. The proof strategy here is to first show alignment for a quadratic loss function, and then generalize its proof to general loss functions satisfying Assumption 3.1. Below we first give the formal statement of the implicit alignment on quadratic loss Theorem 4.14. Note this alignment property is an implicit property of the SAM algorithm – it is not due to the fact that SAM is an approximation of GD on L_ρ^{Asc}, because optimizing L_ρ^{Asc} would rather make the gradient align to the smallest eigenvector!

Theorem 4.14. *Suppose A is a positive definite symmetric matrix with unique top eigenvalue. Consider running SAM on loss $L(x) := \frac{1}{2}x^T Ax$ as (11), then for almost every $x(0)$, we have $x(t)$ converges in direction to $v_1(A)$ and $\lim_{t\to\infty}\|x(t)\| = \frac{\eta\rho\lambda_1(A)}{2-\eta\lambda_1(A)}$ with $\eta\lambda_1(A) < 1$.*

$$x(t+1) = x(t) - \eta A\left(x(t) + \rho\frac{Ax(t)}{\|Ax(t)\|}\right), \tag{11}$$

Equivalently, we can reformulate the update rule through the lens of the gradient $\nabla L(x) = Ax$:

$$\nabla L(x(t+1)) = \nabla L(x(t)) - \eta(\nabla^2 L(x(t)))^2\left(\nabla L(x(t)) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}\right), \tag{12}$$

and the result of Theorem 4.14 becomes the alignment between gradient and the top eigenvector of Hessian. This property can be generalized to the analysis of general loss as the dynamic of SAM near a fixed point on Γ is similar to the quadratic case with small perturbation. To see this, we apply Taylor expansion on the update rule of SAM (3):

$$x(t+1) = x(t) - \eta\nabla L(x(t)) - \eta\rho\nabla^2 L(x(t))\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|_2} + O(\eta\rho^2). \tag{13}$$

Since phase II happens in an $O(\eta\rho)$-neighborhood of manifold Γ, we have $\|x(t+1) - x(t)\|_2 = O(\eta\rho)$. Then by (13) and Taylor expansion on $\nabla L(x(t+1))$ at $x(t)$, we have that

$$\nabla L[x(t+1)] = \nabla L(x(t)) - \nabla^2 L(x(t)) \left(x(t+1) - x(t) \right) + O(\eta^2\rho^2) \tag{14}$$

$$= \nabla L(x(t)) - \eta(\nabla^2 L(x(t)))^2 \left(\nabla L(x(t)) + \rho \frac{\nabla L(x(t))}{\|\nabla L(x(t)))\|} \right) + O(\eta\rho^2) \tag{15}$$

Equation (14) is a $O(\eta\rho^2)$-perturbed version of the update rule in the quadratic case. Note this is a higher order term comparing to the other two terms, which have orders $\Theta(\eta^2\rho)$ and $\Theta(\eta\rho)$ respectively, the implicit alignment between Hessian and gradient happens for the same reason as in the quadratic case. We further show once this alignment happens, it will be kept until the end of our analysis, which is $\Theta(\eta^{-1}\rho^{-2})$ steps.

5 EXPLICIT AND IMPLICIT BIAS IN THE STOCHASTIC SETTING

In practice, people usually use SAM in the stochastic mini-batch setting, and the test accuracy improves as the batch size decreases (Foret et al., 2021). Towards explaining this phenomenon, Foret et al. (2021) argue intuitively that stochastic SAM minimizes stochastic worst-direction sharpness. Given our results in Section 4, it's natural to ask if we can justify the above intuition by showing the implicit Hessian-gradient alignment in the stochastic setting. Unfortunately, such alignment is not possible in the most general setting. For example, take a simple quadratic loss $L(x) = \frac{L_1(x) + L_2(x)}{2}$, where $L_k(x) = 0.5 x^\top A_k x$ and A_k is a positive definite matrix for $k = 1, 2$. If A_1 and A_2 have different top eigendirection, then no x can simultaneously satisfy that $\nabla L_k(x) = A_k x$ aligns to the top eigenvector of $\nabla^2 L_k(x) = A_k$. Yet when the batch size is 1, we can prove rigorously that stochastic SAM minimizes stochastic worst-direction sharpness (Section 5.2), which is the main result in this section.

Below we start by introducing our setting for SGD with batch size 1. We still need Assumption 3.1 in this section. We first by analyzing the explicit bias of the stochastic ascent- and worst-direction sharpness in Section 5.1 via the tools developed in Section 4.1. It turns out they are all proportional to the trace of hessian as $\rho \to 0$. In Section 5.2, we show stochastic SAM locally decreases trace of Hessian.

Setting Let $f_k(x)$ be the model output on kth data where f_k is a $\mathcal{C}^4$-smooth function and y_k be the ith label for $l = 1, \ldots, M$. We define the loss on kth data as $L_k(x) = \ell(f_k(x), y_k)$ and the total loss $L = \sum_{k=1}^{M} L_k/M$, where $\ell(y', y)$ is a $\mathcal{C}^4$-smooth function satisfying the following properties[1]: (1). $\arg\min_{y' \in \mathbb{R}} \ell(y', y) = y$, for any $y \in \mathbb{R}$; (2). $\frac{d^2\ell(y',y)}{d^2 y'}|_{y'=y} > 0$, for any $y \in R$; (3). There exists x_0, such that for any $k \in [M]$, it holds that $f_k(x_0) = y_k$.

5.1 STOCHASTIC WORST- AND ASCENT-DIRECTION SHARPNESS HAVE SAME EXPLICIT BIAS

Below we specify *stochastic worst-direction sharpness* and *stochastic ascent-direction sharpness* as $\mathbb{E}_k[R_{k,\rho}^{\text{Max}}] = \mathbb{E}_k[L_{k,\rho}^{\text{Max}}] - L$ and $\mathbb{E}_k[R_{k,\rho}^{\text{Asc}}] = \mathbb{E}_k[L_{k,\rho}^{\text{Asc}}] - L$. Here $L_{k,\rho}^{\max}$ and $L_{k,\rho}^{\text{asc}}$ is defined as in Equations (1) and (2) with L replaced by L_k. Unlike the full-batch setting, these two sharpness have same explicit bias, or more precisely, they have the same limiting regularizers. We omit the result on the stochastic average-direction sharpness as it is the same as its counterpart in the full-batch case.

Theorem 5.1. *Stochastic worst-direction sharpness $\mathbb{E}_k[R_{k,\rho}^{\text{Max}}]$ satisfies Assumption 4.1 and admits* $\text{Tr}(\nabla^2 L(\cdot))/2$ *as a good limiting regularizer on Γ.*

Theorem 5.2. *Stochastic ascent-direction sharpness $\mathbb{E}_k[R_{k,\rho}^{\text{Asc}}]$ satisfies Assumption 4.1 and admits* $\text{Tr}(\nabla^2 L(\cdot))/2$ *as a good limiting regularizer on Γ.*

Unlike in the full-batch setting where the implicit regularizer of ascent-direction sharpness and worst-direction sharpness have different explicit bias, here they are the same because there is no difference between the maximum and minimum of its non-zero eigenvalue for rank-1 Hessian of each individual loss, and that the average of limiting regularizers is equal to the limiting regualrizer of the average regularizers by definition.

---

[1]Examples include ℓ_2 loss: $\ell(y, y') = 0.5(y - y')^2$.

5.2 STOCHASTIC SAM MINIMIZES STOCHASTIC WORST-DIRECTION SHARPNESS

Stochastic SAM: Recall L_k is the loss on kth data, we use *stochastic SAM* to denote the following update rule, where k_t is sampled i.i.d from uniform distribution on $[M]$.

$$x(t+1) = x(t) - \eta \nabla L_{k_t} \left(x + \rho \frac{\nabla L_{k_t}(x)}{\|\nabla L_{k_t}(x)\|_2} \right), \tag{16}$$

The goal of this section is to show stochastic SAM tracks the following Riemannian gradient flow with respect to $\text{Tr}(\nabla^2 L(\cdot))$ on the manifold for sufficiently small η and ρ,

$$X(\tau) = X(0) - \frac{1}{2} \int_{s=0}^{\tau} P_{X(s),\Gamma}^{\perp} \nabla \text{Tr}(X(s))ds, X(0) = \Phi(x_{\text{init}}). \tag{17}$$

The fundamental difference between stochastic SAM (16) with batch size 1 and that in the general setting is that the rank of the Hessian of each stochastic loss at minimizers is only rank-1, which enforces the gradient $\nabla L_k(x) \approx \nabla^2 L_k(x)(x - \Phi(x))$ to (almost) lie in the direction of the top eigenvalue of the Hessian. Lemma 5.3 formally states this property. For convenience, we denote $\frac{d^2 \ell(y',y_k)}{d^2 y'}|_{y'=f_k(x)}\|\nabla f_k(x)\|^2, \nabla f_k(x)/\|\nabla f_k(x)\|$ as $\Lambda_k(x), w_k(x)$.

Lemma 5.3. *Under Assumption 3.1, for any* $x \in U$ *and* $p \in \Gamma, \nabla^2 L_k(p) = \Lambda_k(p)w_k(p)w_k(p)^T$ *and* $\exists s \in \{1, -1\} \frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = sw_k(p) + O(\|x - p\|)$

With Lemma 5.3, we can show $\Phi(x(t+1)) - \Phi(x(t)) = -\eta \rho^2 P_{t,\Gamma}^{\perp} \nabla(\Lambda_k(x))/2 + \tilde{O}(\eta^2 \rho^2)$ when $\|x(t+1) - x(t)\| = \tilde{O}(\eta \rho)$, via a similar but slightly more complicated analysis on $\Phi(x(t+1)) - \Phi(x(t))$ as in Section 4.2. As we have $\mathbb{E}[\Lambda_k(x)] = (\sum_k \Lambda_k(x))/M = \text{Tr}(\nabla^2 L(x)), x \in \Gamma$, we have that locally SAM is essentially performing a stochastic projected gradient descent with respect to $\text{Tr}(\nabla^2 L(\cdot))$.

It still remains to prove stochastic SAM gets $\tilde{O}(\eta \rho)$ close of the manifold, which is addressed by Theorem C.54. Put everything together, we have the following main theorem for this section.

Theorem 5.4. *Let* $\{x(t)\}$ *be the iterates defined by SAM (16) and* $x(0) = x_{init} \in U$, *then under Assumption 3.1, for all* η *and* ρ *such that* $(\eta + \rho) \log(1/\eta\rho)$ *is sufficiently small, the dynamics of SAM can be split into two phases:*

- **Phase I:** *Stochastic SAM follows Gradient Flow with respect to* L *until entering an* $O(\eta\rho)$ *neighborhood of the manifold* Γ *in* $\tilde{O}(\frac{1}{\eta})$ *steps, with probability at least* $1 - O(\sqrt{\rho})$;
- **Phase II:** *Stochastic SAM tracks the solution* X *of (17), the Riemannian Gradient Flow with respect to* $\text{Tr}(\nabla^2 L(\cdot))$, *with time scaling as* $\|\Phi(x(\lceil T_3/(\eta\rho^2)\rceil)) - X(T_3)\| = O((\eta + \rho)\log(1/\eta\rho))$.

Corollaries 5.5 and 5.6 below are stochastic counterparts of Corollaries 4.10 and 4.11, which says that the trace of Hessian of the limiting flow of well-tracks the stochastic worst-direction sharpness, and therefore when the limiting flow converges to a local minimizer of trace of Hessian, stochastic SAM minimizes the stochastic worst-direction sharpness.

Corollary 5.5. *For all* $T_3' > 0$, *for all* ρ, η *such that* $(\eta + \rho) \log(1/\eta\rho)$ *are sufficiently small we have* $\forall T_3' \leq \eta\rho^2 t \leq T_3, \|\mathbb{E}_k[R_{k,\rho}^{\text{Max}}](x(t)) - \rho^2 \text{Tr}(\nabla^2 L(X(\eta\rho^2 t)))/2\| = \tilde{O}((\eta + \rho)\rho^2)$.

Corollary 5.6. *Define* U' *as in Theorem 4.3, suppose* $X(\infty) = \lim_{t \to \infty} X(t)$ *exists and is a minimizer of* $\text{Tr}(\nabla^2 L(x))$ *in* $U' \cap \Gamma$, *for all* $\epsilon > 0$, *there exists a constant* $T > 0$, *then for all* ρ, η *such that* $(\eta + \rho) \log(1/\eta\rho)$ *are sufficiently small we have* $\mathbb{E}_k[L_{k,\rho}^{\text{Max}}](\lceil T/(\eta\rho^2)\rceil)) \leq \epsilon\rho^2 + \inf_{x \in U'} \mathbb{E}_k[L_{k,\rho}^{\text{Max}}](x)$

6 CONCLUSION

In this work, we have performed a rigorous mathematical analysis of the explicit bias of various notions of sharpness when used as regularizers and the implicit bias of the SAM algorithm. In particular, we show the explicit biases of worst-, ascent- and average-direction sharpness around the manifold of minimizers are minimizing the largest eigenvalue, the smallest nonzero eigenvalue, and the trace of Hessian of the loss function. We show that in the full-batch setting, SAM provably decreases the largest eigenvalue of Hessian, while in the stochastic setting when batch size is 1, SAM provably decreases the trace of Hessian.

REFERENCES

Maksym Andriushchenko and Nicolas Flammarion. Towards understanding sharpness-aware minimization. In *International Conference on Machine Learning*, pp. 639–668. PMLR, 2022.

Sanjeev Arora, Zhiyuan Li, and Abhishek Panigrahi. Understanding gradient descent on edge of stability in deep learning. *arXiv preprint arXiv:2205.09745*, 2022.

Peter L Bartlett, Philip M Long, and Olivier Bousquet. The dynamics of sharpness-aware minimization: Bouncing across ravines and drifting towards wide minima. *arXiv preprint arXiv:2210.01513*, 2022.

Guy Blanc, Neha Gupta, Gregory Valiant, and Paul Valiant. Implicit regularization for deep neural networks driven by an ornstein-uhlenbeck like process. *arXiv preprint arXiv:1904.09080*, 2019.

Yaim Cooper. The loss landscape of overparameterized neural networks. *arXiv preprint arXiv:1804.10200*, 2018.

Alex Damian, Tengyu Ma, and Jason Lee. Label noise sgd provably prefers flat global minimizers, 2021.

Chandler Davis and William Morton Kahan. The rotation of eigenvectors by a perturbation. iii. *SIAM Journal on Numerical Analysis*, 7(1):1–46, 1970.

Laurent Dinh, Razvan Pascanu, Samy Bengio, and Yoshua Bengio. Sharp minima can generalize for deep nets. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pp. 1019–1028. JMLR. org, 2017.

Felix Draxler, Kambis Veschgini, Manfred Salmhofer, and Fred Hamprecht. Essentially no barriers in neural network energy landscape. In *International conference on machine learning*, pp. 1309–1318. PMLR, 2018.

Gintare Karolina Dziugaite and Daniel M Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. *arXiv preprint arXiv:1703.11008*, 2017.

Benjamin Fehrman, Benjamin Gess, and Arnulf Jentzen. Convergence rates for the stochastic gradient descent method for non-convex objective functions. *Journal of Machine Learning Research*, 21:136, 2020.

Pierre Foret, Ariel Kleiner, Hossein Mobahi, and Behnam Neyshabur. Sharpness-aware minimization for efficiently improving generalization. *arXiv preprint arXiv:2010.01412*, 2020.

Pierre Foret, Ariel Kleiner, Hossein Mobahi, and Behnam Neyshabur. Sharpness-aware minimization for efficiently improving generalization. In *International Conference on Learning Representations*, 2021.

Timur Garipov, Pavel Izmailov, Dmitrii Podoprikhin, Dmitry P Vetrov, and Andrew G Wilson. Loss surfaces, mode connectivity, and fast ensembling of dnns. *Advances in neural information processing systems*, 31, 2018.

E. Hairer, S.P. Nørsett, and G. Wanner. *Solving Ordinary Differential Equations I: Nonstiff Problems*. Springer Series in Computational Mathematics. Springer Berlin Heidelberg, 2008. ISBN 9783540566700. URL https://books.google.com/books?id=F93u7VcSRyYC.

Sepp Hochreiter and Jürgen Schmidhuber. Flat minima. *Neural Computation*, 9(1):1–42, 1997.

Roger A. Horn and Charles R. Johnson. *Matrix analysis*. Cambridge university press, 2012.

Stanisław Jastrzebski, Zachary Kenton, Devansh Arpit, Nicolas Ballas, Asja Fischer, Yoshua Bengio, and Amos Storkey. Three factors influencing minima in sgd. *arXiv preprint arXiv:1711.04623*, 2017.

Yiding Jiang, Behnam Neyshabur, Hossein Mobahi, Dilip Krishnan, and Samy Bengio. Fantastic generalization measures and where to find them. *arXiv preprint arXiv:1912.02178*, 2019.

Simran Kaur, Jeremy Cohen, and Zachary C Lipton. On the maximum hessian eigenvalue and generalization. *arXiv preprint arXiv:2206.10654*, 2022.

Nitish Shirish Keskar, Dheevatsa Mudigere, Jorge Nocedal, Mikhail Smelyanskiy, and Ping Tak Peter Tang. On large-batch training for deep learning: Generalization gap and sharp minima. *arXiv preprint arXiv:1609.04836*, 2016.

Jungmin Kwon, Jeongseop Kim, Hyunseo Park, and In Kwon Choi. Asam: Adaptive sharpness-aware minimization for scale-invariant learning of deep neural networks. In *International Conference on Machine Learning*, pp. 5905–5914. PMLR, 2021.

Zhiyuan Li, Tianhao Wang, and Sanjeev Arora. What happens after sgd reaches zero loss?–a mathematical framework. In *International Conference on Learning Representations*, 2021.

Yong Liu, Siqi Mai, Xiangning Chen, Cho-Jui Hsieh, and Yang You. Towards efficient and scalable sharpness-aware minimization. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 12360–12370, 2022.

Kaifeng Lyu, Zhiyuan Li, and Sanjeev Arora. Understanding the generalization benefit of normalization layers: Sharpness reduction. *arXiv preprint arXiv:2206.07085*, 2022.

Jan R Magnus. On differentiating eigenvalues and eigenvectors. *Econometric theory*, 1(2):179–191, 1985.

David McAllester. Simplified pac-bayesian margin bounds. In *Learning theory and Kernel machines*, pp. 203–215. Springer, 2003.

Behnam Neyshabur, Srinadh Bhojanapalli, David McAllester, and Nati Srebro. Exploring generalization in deep learning. In *Advances in Neural Information Processing Systems*, pp. 5947–5956, 2017.

Antonio Orvieto, Anant Raj, Hans Kersting, and Francis Bach. Explicit regularization in overparametrized models via noise injection. *arXiv preprint arXiv:2206.04613*, 2022.

Colin Wei and Tengyu Ma. Improved sample complexities for deep networks and robust classification via an all-layer margin. *arXiv preprint arXiv:1910.04284*, 2019.

Dongxian Wu, Shu-Tao Xia, and Yisen Wang. Adversarial weight perturbation helps robust generalization. *Advances in Neural Information Processing Systems*, 33:2958–2969, 2020.

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. *arXiv preprint arXiv:1611.03530*, 2016.

Yang Zhao, Hao Zhang, and Xiuyuan Hu. Penalizing gradient norm for efficiently improving generalization in deep learning. *arXiv preprint arXiv:2202.03599*, 2022.

Yaowei Zheng, Richong Zhang, and Yongyi Mao. Regularizing neural networks via adversarial model perturbation. In *Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition*, pp. 8156–8165, 2021.

Juntang Zhuang, Boqing Gong, Liangzhe Yuan, Yin Cui, Hartwig Adam, Nicha Dvornek, Sekhar Tatikonda, James Duncan, and Ting Liu. Surrogate gap minimization improves sharpness-aware training. *arXiv preprint arXiv:2203.08065*, 2022.

CONTENTS

A ADDITIONAL RELATED WORKS

Sharpness Minimization as Implicit Bias. Recent theoretical works Blanc et al. (2019); Damian et al. (2021); Li et al. (2021) showed that SGD with label noise is implicitly biased to local minimizers with a smaller trace of Hessian. Arora et al. (2022) showed that normalized GD implicitly decreases the largest eigenvalue of the Hessian. Lyu et al. (2022) showed that GD with weight decay on a scale invariant loss function implicitly decreases the spherical sharpness, *i.e.*, the largest eigenvalue of the Hessian evaluated at the normalized parameter.

In a concurrent work, Bartlett et al. (2022) prove that on quadratic loss, the iterate of SAM (Equation (11)) and its gradient converges to the top eigenvector of Hessian, which is almost the same as our Theorem 4.14. Assuming such alignment for a general loss, the work of Bartlett et al. (2022) shows that the largest eigenvalue of Hessian decreases in the next step. This paper also proves such a Hessian-gradient alignment for general loss functions and an end-to-end theorem showing that the largest eigenvalue of Hessian and worst-direction sharpness decrease along the trajectory of SAM (Theorem 4.9), which are not shown in Bartlett et al. (2022). Moreover, this paper also characterizes implicit bias of stochastic SAM with batch size 1, which is minimizing the average-direction sharpness, while Bartlett et al. (2022) only considers the deterministic case.

Comparison with Arora et al. (2022). Our proof uses similar framework as Arora et al. (2022). However our analysis has its own difficulty for the following reasons. First, Arora et al. (2022) only deal with deterministic case, while our analysis extends to stochastic SAM as well (Section 5). Second, our analysis for the deterministic case is different from that of Arora et al. (2022) in the following two aspects:

1. The alignment analysis is more complicated because we have two hyperparameters, learning rate η and perturbation radius ρ, while Arora et al. (2022) only needs to deal with one hyperparameter, learning rate η.
2. The mechanism of penalizing worst-direction sharpness is different, which can be seen from the dependency of the sharpness-reduction rate over learning rate η. In Arora et al. (2022), normalized GD reduces the sharpness via a second-order effect of GD and thus the sharpness is reduced by $O(\eta^2)$ per step. In our analysis, for fixed small perturbation radius ρ, the sharpness is reduced by $O(\rho^2\eta)$ per step, which is linear in η.

B MORE DISCUSSIONS OF MANIFOLD ASSUMPTION

Validity. Recent empirical studies have shown that there are essentially no barriers in loss landscape between different minimizers, that is, the set of minimizers are path-connected (Draxler et al., 2018; Garipov et al., 2018). Motivated by this empirical discovery, we make the assumption below following Fehrman et al. (2020); Damian et al. (2021); Li et al. (2021), which is theoretically justified by Cooper (2018) under a generic setting.

Importance. The connectivity of the set of local minimizers implied by the manifold assumption above allows us to take limits of perturbation radius $\rho \to 0$ while still yield interesting and insightful implicit bias results. So far almost all analysis of implicit bias for general model parameterizations relies on Taylor expansion, e.g. Blanc et al. (2019); Damian et al. (2021); Li et al. (2021); Arora et al. (2022), so does the derivation of the SAM algorithm (Foret et al., 2020; Wu et al., 2020). Thus it's crucial to consider small perturbation size ρ. On the contrary, if the set of global minimizers are a set of discrete points, then with small perturbation radius ρ, implicit bias of optimizers is not sufficient to drive the iterate from global minimum to the other one.

Limitation. As proved by Cooper (2018), the manifold setting holds generically for regression problems with smooth loss. The manifold assumption doesn't hold for networks with ReLU (which is non-smooth) and classification problems with softmax loss (where an overparametrized network may achieve arbitrarily small but non-zero loss). In the latter case, there may not even exist a global minimizer, not to mention a manifold of minimizers.

C   PROOF

C.1   PROOF SETUPS

In this section we provide details of our proof setups, including notations and assumptions/settings.

We first introduce some additional notations that will be used in the proofs. For any subset $S \in \mathbb{R}^D$, we define $\mathrm{Dist}(x, S) \triangleq \inf_{y \in S} \|x - y\|_2$. For any $d > 0$ and any subset $S \in \mathbb{R}^D$, we define $S^d \triangleq \{x \in \mathbb{R}^D \mid \mathrm{Dist}(x, S) \le d\}$. Our convention is to use K to denote a compact set and U to denote an open set.

Below we restate our main assumption in the full-batch case and related notations in Section 3. Throughout the analysis, we fix our initialization as x_{init}, our loss function as $L : \mathbb{R}^D \to \mathbb{R}$.

Assumption 3.1. *Assume loss $L : \mathbb{R}^D \to \mathbb{R}$ belongs to $\mathcal{C}^4$, and there exists a manifold Γ that is $D - M$ dimensional $\mathcal{C}^2-$submanifold of $\mathbb{R}^D$ for some integer $1 \le M \le D$, where for all $x \in \Gamma$, x is a global minimizer of L, $L(x) = 0$ and $rank(\nabla^2 L(x)) = M$.*

Notations for Full-Batch Setting: Given any point $x \in \Gamma$, define $P_{x,\Gamma}$ as the projection operator onto the manifold of the normal space of Γ at x and $P^{\perp}_{x,\Gamma} = I_D - P_{x,\Gamma}$. Given the loss function L, its gradient flow is denoted by mapping $\phi : \mathbb{R}^D \times [0, \infty) \to \mathbb{R}^D$. Here, $\phi(x, \tau)$ denotes the iterate at time τ of a gradient flow starting at x and is defined as the unique solution of $\phi(x, \tau) = x - \int_0^\tau \nabla L(\phi(x, t)) dt, \forall x \in \mathbb{R}^D$. We further define the limiting map of $\phi(x, \cdot)$ as $\Phi(x) = \lim_{\tau \to \infty} \phi(x, \tau)$, that is, $\Phi(x)$ denotes the convergent point of the gradient flow starting from x. For convenience, we define $\lambda_i(x), v_i(x)$ as $\lambda_i(\nabla^2 L(\Phi(x))), v_i(\nabla^2 L(\Phi(x)))$ whenever the latter is well defined. When $x(t)$ and Γ is clear from context, we also use $\lambda_i(t) := \lambda_i(x(t)), v_i(t) := v_i(x(t)), P^{\perp}_{t,\Gamma} := P^{\perp}_{\Phi(x(t)),\Gamma}, P_{t,\Gamma} := P_{\Phi(x(t)),\Gamma}$.

Definition 3.2 (Attraction Set). *Let U be the attraction set of Γ under gradient flow, that is, a neighborhood of Γ containing all points starting from which gradient flow w.r.t. loss L converges to some point in Γ, or mathematically, $U \triangleq \{x \in \mathbb{R}^D | \Phi(x) \text{ exists and } \Phi(x) \in \Gamma\}$.*

Below we restate the setting for stochastic loss of batch size one in Section 5.

Setting C.1. *Let $f_k(x)$ be the model output on kth data where f_k is a $\mathcal{C}^4$-smooth function and y_k be the ith label for $l = 1, \ldots, M$. We define the loss on kth data as $L_k(x) = \ell(f_k(x), y_k)$ and the total loss $L = \sum_{k=1}^M L_k / M$, where $\ell(y', y)$ is a $\mathcal{C}^4$-smooth function satisfying the following properties[2]: (1). $\arg\min_{y' \in \mathbb{R}} \ell(y', y) = y$, for any $y \in \mathbb{R}$; (2). $\frac{d^2 \ell(y', y)}{d^2 y'}|_{y'=y} > 0$, for any $y \in R$; (3). There exists x_0, such that for any $k \in [M]$, it holds that $f_k(x_0) = y_k$.*

In our analysis, we prove our main theorems in the stochastic setting under a more general condition than Setting C.1, which is Assumption C.2 (on top of Assumption 3.1). The only usage of Setting C.1 in the proof is Lemma 5.3 and theorem C.53.

Assumption C.2. *Total loss $L = \frac{1}{M} \sum_{k=1}^M L_k$. For each $k \in [M]$, L_k is $\mathcal{C}^4$, and there exists a $(D - 1)$-dimensional $\mathcal{C}^2$-submanifold of $\mathbb{R}^D$, Γ_k, where for all $x \in \Gamma_k$, x is a global minimizer of L_k, $L_k(x) = 0$ and $rank(\nabla^2 L_k(x)) = 1$. Moreover, $\Gamma = \cap_{k=1}^M \Gamma_k$ for Γ defined in Assumption 3.1.*

Notations for Stochastic Setting: Since L_k is rank-1 on Γ_k for each $k \in [M]$, we can write it as $L_k(x) = \Lambda_k(x) w_k(x) w_k^\top(x)$ for any $x \in \Gamma$, where w_k is a continuous function on Γ with pointwise unit norm. Given the loss function L_k, its gradient flow is denoted by mapping $\phi_k : \mathbb{R}^D \times [0, \infty) \to \mathbb{R}^D$. Here, $\phi_k(x, \tau)$ denotes the iterate at time τ of a gradient flow starting at x and is defined as the unique solution of $\phi_k(x, \tau) = x - \int_0^\tau \nabla L_k(\phi_k(x, t)) dt, \forall x \in \mathbb{R}^D$. We further define the limiting map Φ_k as $\Phi_k(x) = \lim_{\tau \to \infty} \phi_k(x, \tau)$, that is, $\Phi_k(x)$ denotes the convergent point of the gradient flow starting from x. Similar to Definition 3.2, we define $U_k = \{x \in \mathbb{R}^D | \Phi(x) \text{ exists and } \Phi_k(x) \in \Gamma_k\}$ be the attraction set of Γ_i. We have that each U_k is open and Φ_k is $\mathcal{C}^2$ on U_k by Lemma B.15 in Arora et al. (2022).

Definition C.3. *A function L is μ-PL in a set U iff $\forall x \in U$, $\|\nabla L(x)\|_2^2 \ge 2\mu(L(x) - \inf_{x \in U} L(x))$.*

---

[2]Examples include ℓ_2 loss: $\ell(y, y') = 0.5(y - y')^2$.

Definition C.4. *The spectral 2-norm of a k-order tensor $X_{i_1,...,i_k} \in R^{d_1 \times ... \times d_k}$ is defined as*

$$\|X\|_2 = \max_{x_i \in R^{d_i}, \|x_i\|_2 = 1} X[x_1, ..., x_k].$$

Lemma C.5 (Arora et al. (2022) Lemma B.2). *Given any compact set $K \subseteq \Gamma$, there exist $r(K), \mu(K), \Delta(K) \in \mathbb{R}^+$ such that*

1. *$K^{r(K)} \cap \Gamma$ is compact.*
2. *$K^{r(K)} \subset U \cap (\cap_k U_k)$.*
3. *L is $\mu(K)$-PL on $K^{r(K)}$.*
4. *$\inf_{x \in K^{r(K)}}(\lambda_1(\nabla^2 L(x)) - \lambda_2(\nabla^2 L(x))) \geq \Delta(K) > 0$.*
5. *$\inf_{x \in K^{r(K)}} \lambda_M(\nabla^2 L(x)) \geq \mu(K) > 0$.*
6. *$\inf_{x \in K^{r(K)}} \lambda_1(\nabla^2 L_k(x)) \geq \mu(K) > 0$.*

Given compact set $K \subset \Gamma$, we further define

$$\zeta(K) = \sup_{x \in K^{r(K)}} \|\nabla^2 L(x)\|_2, \ \nu(K) = \sup_{x \in K^{r(K)}} \|\nabla^3 L(x)\|_2, \ \Upsilon(K) = \sup_{x \in K^{r(K)}} \|\nabla^4 L(x)\|_2,$$

$$\xi(K) = \sup_{x \in K^{r(K)}} \|\nabla^2 \Phi(x)\|_2, \ \chi(K) = \sup_{x \in K^{r(K)}} \|\nabla^3 \Phi(x)\|_2.$$

Similarly, we use notations like $\zeta_k(K), \nu_k(K), \Upsilon_k(K), \xi_k(K), \chi_k(K)$ to denote the counterpart of the above quantities defined for stochastic loss L_k and its limiting map Φ_k for $k \in [M]$.

Lemma C.6 (Arora et al. (2022), Lemma B.5 and B.7). *Given any compact subset $K \subset \Gamma$, let $r(K)$ be defined in Lemma C.5, there exist $0 < h(K) < r(K)$ such that*

1. *$\sup_{x \in K^{h(K)}} L(x) - \inf_{x \in K^{h(K)}} L(x) \leq \frac{\mu(K)\rho^2(K)}{8}$.*
2. *$\forall x \in K^{h(K)}, \Phi(x) \in K^{r(K)/2}$.*
3. *$\forall x \in K^{h(K)}, \|x - \Phi(x)\|_2 \leq \frac{8\mu(K)^2}{\zeta(K)\nu(K)}$.*
4. *The whole segment $\overline{x\Phi(x)}$ lies in $K^{r(K)}$, so does $\overline{x\Phi_k(x)}$, for any $k \in [D]$.*

The proof of the lemmas above can be found in Arora et al. (2022). Readers should note that although Arora et al. (2022) only prove these lemmas when K is a special compact set (the trajectory of an ODE), all the proof does not use any property of K other than it is a compact subset of Γ, and thus our Lemmas C.5 and C.6 hold for general compact subsets of Γ.

In the rest part of the appendix, for convenience we will drop the dependency on K in various constants when there is no ambiguity.

C.2 PROPERTIES OF LIMITING MAP OF GRADIENT FLOW, Φ

In our analysis, the property of Φ will be heavily used. In this section, we will recap some related lemmas from Arora et al. (2022), and then introduce some new lemmas for the stochastic setting with batch size one.

Lemma C.7 (Arora et al. (2022) Lemma B.6). *Given any compact set $K \subset \Gamma$, for any $x \in K^h$,*

$$\|x - \Phi(x)\|_2 \leq \int_0^\infty \|\frac{d\phi(x,t)}{dt}\|_2 \leq \sqrt{\frac{2(L(x) - L(\Phi(x)))}{\mu}} \leq \frac{\|\nabla L(x)\|_2}{\mu}.$$

Lemma C.8. *Given any compact set $K \subset \Gamma$, for any $x \in K^h$,*

$$\|\nabla L(x)\|_2 \leq \zeta\|x - \Phi(x)\|_2 \leq \zeta\sqrt{\frac{2(L(x) - L(\Phi(x)))}{\mu}}.$$

Proof. The first inequality is by Lemma C.5 and Taylor Expansion. The second inequality is by Lemma C.7. $\square$

Lemma C.9 (Arora et al. (2022) Lemmas B.16 and B.22). *Given any compact set $K \subset \Gamma$, for any $x \in K^h$,*

$$\partial\Phi(x)\nabla L(x) = 0, x \in U;$$
$$\partial\Phi(x)\nabla^2 L(x)\nabla L(x) = -\partial^2\Phi(x)[\nabla L(x), \nabla L(x)], x \in U;$$
$$\partial\Phi(x)\partial^2(\nabla L)(x)[v_1, v_1] = P^{\perp}_{X,\Gamma}\nabla(\lambda_1(\nabla^2(L(x)))), x \in \Gamma.$$

Lemma C.10 (Arora et al. (2022) Lemmas B.8 and B.9). *Given any compact set $K \subset \Gamma$, for any $x \in K^h$,*

$$\|P^{\perp}_{\Phi(x),\Gamma}(x - \Phi(x))\|_2 \leq \frac{\zeta\nu}{4\mu^2}\|x - \Phi(x)\|_2^2;$$

$$\|\nabla L(x) - \nabla^2 L(\Phi(x))(x - \Phi(x))\|_2 \leq \frac{\nu}{2}\|x - \Phi(x)\|_2^2;$$

$$\left|\frac{\|\nabla L(x)\|_2}{\|\nabla^2 L(\Phi(x))(x - \Phi(x))\|_2} - 1\right| \leq \frac{2\nu}{\mu}\|x - \Phi(x)\|_2;$$

$$\frac{\nabla L(x)}{\|\nabla L(x)\|} = \frac{\nabla^2 L(\Phi(x))(x - \Phi(x))}{\|\nabla^2 L(\Phi(x))(x - \Phi(x))\|_2} + O(\frac{\nu}{\mu}\|x - \Phi(x)\|_2).$$

The proof of above lemmas can be found in Arora et al. (2022).

Lemma C.11. *Given any compact set $K \subset \Gamma$, for any $x \in K^h$,*

$$\|\partial\Phi(x)\nabla L_k(x)\|_2 \leq (\nu_k + \zeta_k\xi)\|x - \Phi(x)\|_2^2$$
$$\|\partial\Phi(x)\nabla^2 L_k(x)\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 \leq (\nu_k + \zeta_k\xi)\|x - \Phi(x)\|_2$$

Proof. By Lemma C.6 and Taylor Expansion,

$$\|\partial\Phi(x)\nabla L_k(x)\|_2 \leq \|\partial\Phi(x)\nabla^2 L_k(\Phi(x))(x - \Phi(x))\|_2 + \nu_k\|x - \Phi(x)\|_2^2$$
$$\leq \|\partial\Phi(\Phi(x))\nabla^2 L_k(\Phi(x))(x - \Phi(x))\|_2 + \nu_k\|x - \Phi(x)\|_2^2 + \zeta_k\xi\|x - \Phi(x)\|_2^2$$
$$= \|P^{\perp}_{x,\Gamma}\partial\Phi(\Phi(x))\nabla^2 L_k(\Phi(x))(x - \Phi(x))\|_2 + \nu_k\|x - \Phi(x)\|_2^2 + \zeta_k\xi\|x - \Phi(x)\|_2^2$$
$$= (\nu_k + \zeta_k\xi)\|x - \Phi(x)\|_2^2,$$

this proves the first claim.

Again by Lemma C.5 and Taylor Expansion,

$$\|\partial\Phi(x)\nabla^2 L_k(x)\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 \leq \|\partial\Phi(x)\nabla^2 L_k(\Phi(x))\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 + \nu_k\|x - \Phi(x)\|_2$$
$$\leq \|\partial\Phi(\Phi(x))\nabla^2 L_k(\Phi(x))\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 + (\nu_k + \zeta_k\xi)\|x - \Phi(x)\|_2$$
$$= (\nu_k + \zeta_k\xi)\|x - \Phi(x)\|_2,$$

this proves the second claim. $\qquad\square$

Lemma C.12. *Given any compact set $K \subset \Gamma$, for any $x \in K^h$ and $y = x - \eta\nabla L\left(x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|}\right)$,*

$$\|y - x\|_2 \leq \eta\|\nabla L(x)\|_2 + \eta\zeta\rho$$
$$\|\Phi(x) - \Phi(y)\|_2 \leq \xi\eta\rho\|\nabla L(x)\|_2 + \nu\eta\rho^2 + \xi\eta^2\|\nabla L(x)\|_2^2 + \xi\zeta^2\eta^2\rho^2$$
$$\leq \zeta\xi\eta\rho\|x - \Phi(x)\|_2 + \zeta^2\xi\eta^2\|x - \Phi(x)\|_2^2 + \nu\eta\rho^2 + \xi\zeta^2\eta^2\rho^2$$

Proof. For sufficient small ρ, $x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|} \in K^r$. By Taylor Expansion,

$$\|y - x\|_2 = \eta\|\nabla L\left(x + \rho\frac{\nabla L(x)}{\|\nabla L(x)\|}\right)\|_2 \leq \eta\|\nabla L(x)\|_2 + \eta\zeta\rho$$

This further implies that for sufficiently small η and ρ, $\overline{xy} \in K^r$.

Again by Taylor Expansion,

$$\|\partial\Phi(x)(y-x)\|_2 \leq \eta\|\partial\Phi(x)\nabla L(x) + \rho\partial\Phi(x)\nabla^2 L(x)\frac{\nabla L(x)}{\|\nabla L(x)\|}\|_2 + \eta\rho^2\nu/2.$$

By Lemma C.9, $\partial\Phi(x)\nabla L(x) = 0$ and $\partial\Phi(x)\nabla^2 L(x)\nabla L(x) = -\partial^2\Phi(x)[\nabla L(x),\nabla L(x)]$. Hence,

$$\|\partial\Phi(x)(y-x)\|_2 \leq \eta\rho\|\nabla L(x)\|_2\|\partial^2\Phi(x)\left[\frac{\nabla L(x)}{\|\nabla L(x)\|},\frac{\nabla L(x)}{\|\nabla L(x)\|}\right]\|_2 + \eta\rho^2\nu/2$$

$$\leq \xi\eta\rho\|\nabla L(x)\|_2 + \eta\rho^2\nu/2.$$

As $\overline{xy} \in K^r$, by Taylor Expansion,

$$\|\Phi(y) - \Phi(x)\|_2 \leq \|\partial\Phi(x)(y-x)\|_2 + \xi\|y-x\|_2^2/2$$

Putting together we have

$$\|\Phi(x) - \Phi(y)\|_2 \leq \xi\eta\rho\|\nabla L(x)\|_2 + \eta\rho^2\nu + \xi\eta^2\|\nabla L(x)\|_2^2 + \xi\zeta^2\eta^2\rho^2.$$

Finally, by Lemma C.8, we have

$$\|\Phi(x) - \Phi(y)\|_2 \leq \xi\eta\rho\|\nabla L(x)\|_2 + \nu\eta\rho^2 + \xi\eta^2\|\nabla L(x)\|_2^2 + \xi\zeta^2\eta^2\rho^2$$

$$\leq \zeta\xi\eta\rho\|x - \Phi(x)\|_2 + \zeta^2\xi\eta^2\|x - \Phi(x)\|_2^2 + \nu\eta\rho^2 + \xi\zeta^2\eta^2\rho^2.$$

$\square$

Lemma C.13. *Given any compact set $K \subset \Gamma$, for any $x \in K^h$ and $y = x - \eta\nabla L_k\left(x + \rho\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\right)$,*

$$\|y - x\|_2 \leq \eta\|\nabla L_k(x)\|_2 + \eta\zeta\rho$$

$$\|\Phi(x) - \Phi(y)\|_2 \leq O(\eta\|\nabla L(x)\|_2^2 + \eta\rho\|\nabla L(x)\|_2 + \eta\rho^2)$$

Proof. For sufficient small ρ, $x + \rho\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} \in K^r$. By Taylor Expansion,

$$\|y - x\|_2 = \eta\|\nabla L_k\left(x + \rho\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\right)\|_2 \leq \eta\|\nabla L_k(x)\|_2 + \eta\zeta\rho.$$

This further implies that for sufficiently small η and ρ, $\overline{xy} \in K^r$.

Again by Taylor Expansion,

$$\|\partial\Phi(x)(y-x)\|_2 \leq \eta\|\partial\Phi(x)\nabla L_k(x) + \rho\partial\Phi(x)\nabla^2 L_k(x)\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 + \eta\rho^2\nu/2.$$

We further have by Lemma C.7,

$$\|\partial\Phi(x)\nabla L_k(x)\|$$

$$\leq\|\partial\Phi(\Phi(x))\nabla L_k(x)\| + \xi\|\nabla L_k(x)\|_2\|x - \Phi(x)\|$$

$$\leq\|\partial\Phi(\Phi(x))\nabla^2 L_k(\Phi(x))(x - \Phi(x))\| + \nu\|x - \Phi(x)\|_2^2 + \zeta\xi\|x - \Phi(x)\|_2^2$$

$$\leq\frac{\nu}{\mu}\|\nabla L(x)\|_2^2 + \frac{\zeta\xi}{\mu^2}\|\nabla L(x)\|_2^2.$$

Similarly

$$\|\rho\partial\Phi(x)\nabla^2 L_k(x)\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2$$

$$\leq\|\rho\partial\Phi(\Phi(x))\nabla^2 L_k(x)\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 + \rho\zeta\xi\|x - \Phi(x)\|_2$$

$$\leq\|\rho\partial\Phi(\Phi(x))\nabla^2 L_k(\Phi(x))\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}\|_2 + \rho\zeta\xi\|x - \Phi(x)\|_2 + \rho\nu\|x - \Phi(x)\|_2$$

$$\leq\rho\frac{\zeta\xi}{\mu^2}\|\nabla L(x)\|_2 + \rho\frac{\nu}{\mu}\|\nabla L(x)\|_2\,.$$

$\square$

Lemma C.14 (A general version of Lemma 5.3). *Under Assumption 3.1 and Assumption C.2, for* $x \in K^h$ *and* $p \in C$, $\nabla^2 L_k(p) = \Lambda_k(p)w_k(p)w_k(p)^\top$, *there exists* $s \in \{1, -1\}$,

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = sw_k(p) + O(\|x - p\|_2)\,.$$

Further if $|w_k^\top(x - p)| \geq \|x - p\|_2^{3/2}$, *then* $s = \text{sign}(w_k^\top(x - p))$. *This implies*

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}^\top (x - p) \geq sw_k^\top(x - p) - O(\|x - p\|_2^2)$$

$$\geq \|w_k^\top(x - p)\|_2 - O(\|x - p\|_2^{3/2})\,.$$

Proof. We will calculate the direction of $\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|}$ using two different approximations and compare them to get our result.

1. According to Lemma C.10,

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = \frac{\nabla^2 L_k(\Phi_k(x))(x - \Phi_k(x))}{\|\nabla^2 L_k(\Phi_k(x))(x - \Phi_k(x))\|_2} + O(\|x - \Phi_k(x)\|_2).$$

Suppose $\nabla^2 L_k(\Phi_k(x)) = \Lambda_k(\Phi_k(x))w_k(\Phi_k(x))w_k(\Phi_k(x))^\top$, then

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = w_k(\Phi_k(x)) + O(\|x - \Phi_k(x)\|_2)$$

As $\nabla^2 L_k(p) = \Lambda_k(p)w_k(p)w_k(p)^\top$, using Davis-Kahan Theorem C.58, we would have $\exists s \in \{-1, 1\}$, such that $\|w_k(\Phi_k(x)) - sw_k(p)\|_2 \leq \zeta\|\Phi_k(x) - p\|_2$.

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = sw_k(p) + O(\|\Phi_k(x) - p\|_2 + \|x - p\|_2).$$

According to Lemma C.7, we have $\|x - \Phi_k(x)\|_2 \leq \frac{\|\nabla L_k(x)\|_2}{\mu} \leq \frac{\zeta\|x-p\|_2}{\mu}$. This implies,

$$\frac{\nabla L_k(x)}{\|\nabla L_k(x)\|} = sw_k(p) + O(\|x - p\|_2). \tag{18}$$

Equation (18) is our first statement.
2. By Taylor expansion at p,

$$\nabla L_k(x) = \Lambda_k(x)w_k(p)w_k(p)^\top(x - p) + O(\nu\|x - p\|_2^2).$$

That being said, when $|w_k^\top(x - p)| \geq \|x - p\|_2^{3/2}$, we have

$$\|\nabla L_k(x) - \Lambda_k w_k w_k^\top(x - p)\|_2 \leq O(\|x - p\|_2^2)\,.$$
$$\|\nabla L_k(x)\| \geq \|\Lambda_k w_k w_k^\top(x - p)\|_2 - O(\|x - p\|_2^2) \geq \Omega(\|x - p\|_2^{3/2}).$$

Concluding,

$$\|\frac{\nabla L_k\left(x\right)}{\|\nabla L_k\left(x\right)\|} - \frac{\Lambda_k w_k w_k^\top \left(x-p\right)}{\|\Lambda_k w_k w_k^\top \left(x-p\right)\|}\|_2 \leq O(\|x-p\|_2^{1/2})$$

Hence we have

$$\frac{\nabla L_k\left(x\right)}{\|\nabla L_k\left(x\right)\|} = \operatorname{sign}(w_k^\top\left(x-p\right))w_k + O(\|x-p\|_2^{1/2})\,. \tag{19}$$

Comparing (18) and (19), we have $s = \operatorname{sign}(w_k(p)^\top\left(x-p\right))$ when $|w_k^\top\left(x-p\right)| \geq \|x-p\|_2^{3/2}$.

□

C.3   WELL-DEFINEDNESS OF SAM

In this section, we discuss the well-definedness of SAM. When $\nabla L(x) = 0$, SAM (Equation (3)) is not well-defined, as the normalized gradient $\frac{\nabla L(x)}{\|\nabla L(x)\|_2}$ is not well-defined. The usual intuition here is that, for a continuously differentiable update rule, as long as the points where the update rule is ill-defined (here it means the points with zero gradient) has zero measure, then almost surely for all initialization, gradient-based optimization algorithms like SAM will not reach exactly at any stationary point. However the following example shows this intuition is not completely correct.

Theorem C.15. *For any $\eta, \rho > 0$, there is a $\mathcal{C}^2$ loss function $L : \mathbb{R} \to \mathbb{R}$ satisfying that (1) L has a unique stationary point and (2) the set of initialization that makes SAM with learning rate η and perturbation radius ρ to reach the unique stationary point has positive Lebesgue measure.*

Proof of Theorem C.15. We first consider the case with $\rho = \eta = 1$ with

$$L(x) = \begin{cases} x^2/2 + x + 1/2, & \text{for } x \in (-\infty, -2); \\ x^4/64 + x^2/8, & \text{for } x \in [-2, 2]; \\ x^2/2 - x + 1/2, & \text{for } x \in (2, \infty). \end{cases} \tag{20}$$

We first check L is indeed $\mathcal{C}^1$: $L(2) = L(-2) = 1/2$, $L'(2) = -L'(-2) = 1$ and $L''(2) = L''(-2) = 1$. Now we claim that for all $|x(0)| > 2$, $x(1) = 0$, which is a stationary point. Note that L is even and monotone increasing on $[0, \infty)$, we have $\nabla L(x)/|\nabla L(x)| = \operatorname{sign}(x)$. Thus for $|x(t)| > 1$, it holds that $|x(t) + \operatorname{sign}(x(t)| > 2$ and therefore

$$\begin{aligned} x(t+1) =& x(t) - \eta L'(x(t) + \rho \nabla L(x)/|\nabla L(x)|) \\ =& x(t) - L'(x(t) + \operatorname{sign}(x(t))) \\ =& x(t) - (x(t) + \operatorname{sign}(x(t)) - \operatorname{sign}(x(t) + \operatorname{sign}(x(t)))) \\ =& x(t) - x(t) = 0. \end{aligned} \tag{21}$$

Now we turn to the case with arbitrary positive η, ρ. It suffices to consider $L_{\eta,\rho}(x) \triangleq \frac{\rho}{\eta}L(\frac{x}{\rho})$. We can use the calculation for $\rho = \eta = 1$ to verify for any $|x| > 2\rho$,

$$L'_{\eta,\rho}(x + \rho\operatorname{sign}(L'_{\eta,\rho}(x))) = L'_{\eta,\rho}(x + \rho\operatorname{sign}(x)) = \frac{1}{\eta}L(x/\rho + \operatorname{sign}(x)) = \frac{x}{\eta},$$

namely $x - L'_{\eta,\rho}(x + \rho\operatorname{sign}(L'_{\eta,\rho}(x))) = 0$. This completes the proof.   □

The issue here is that though differentiable map of SAM $x \mapsto x - \eta\nabla L(x - \rho\frac{\nabla L(x)}{\|\nabla L(x)\|_2})$ always maps the zero-measure set to zero-measure set, the preimage of zero-measure set is not necessarily zero-measure, as the map $x \mapsto x - \eta\nabla L(x - \rho\frac{\nabla L(x)}{\|\nabla L(x)\|_2})$ is not necessarily invertible. The update rule of SAM is not invertible at 0 is exactly the reason of why preimage of 0 has a positive measure.

Now we are going to present the main theorems in this section, which says that the above issue only happen for countably many learning rates for any general $\mathcal{C}^1$ update rule, including full-batch SAM (Theorem C.16) and stochastic SAM (Theorem C.17). Note results in this section does not rely on the manifold assumption, *i.e.*, Assumption 3.1.

Theorem C.16. *Consider any $\mathcal{C}^2$ loss L with zero-measure stationary set $\{x \mid \nabla L(x) = 0\}$. For every $\rho > 0$, except countably many learning rates, for almost all initialization and all t, the iterate of full-batch SAM (Equation (3)) $x(t)$ has non-zero gradient and is thus well-defined.*

Theorem C.17. *Consider any $\mathcal{C}^2$ losses $\{L_k\}_{k=1}^M$ with zero-measure stationary set $\{x \mid \nabla L_i(x) = 0\}$ for each $k \in [M]$. For every $\rho > 0$, except countably many learning rates, for almost all initialization and all t, with probability one of the randomness of the algorithm, the iterate of stochastic SAM (Equation (16)) $x(t)$ has non-zero gradient and is thus well-defined.* [3]

Before present our main theorem (Theorem C.18), we need to introduce some notations first. For a map F mapping from $\mathbb{R}^D \setminus Z \to \mathbb{R}^D$, we define that $F_\eta : \mathbb{R}^D \setminus Z \to \mathbb{R}^D$ as $F_\eta(x) \triangleq x - \eta F(x)$ for any $\eta \in \mathbb{R}^+$. Given a sequence of functions $\{F^n\}_{n=1}^\infty$, we define $F_\eta^n(x) \triangleq x - \eta F^n(x)$, for any $x \in \mathbb{R}^D$. We further define that $\overline{F}_\eta^n(x) \triangleq F_\eta^n(\overline{F}_\eta^{n-1}(x))$ for any $n \geq 1$ and that $\overline{F}_\eta^0(x) = x$.

Theorem C.18. *Let Z be a closed subset of $\mathbb{R}^D$ with zero Lebesgue measure and μ be any probability measure on $\mathbb{R}^D$ that is absolutely continuous to the Lesbegue measure. For any sequence of continuously differentiable functions $F^n : \mathbb{R}^D \setminus Z \to \mathbb{R}^D, n \in \mathbb{N}^+$, the following claim holds for at most countably many $\eta \in \mathbb{R}^+$:*

$$\mu\left(\{x \in \mathbb{R}^D \mid \exists n \in \mathbb{N}, \overline{F}_\eta^n(x) \in Z \text{ and } \forall 0 \leq i \leq n-1, \overline{F}_\eta^i(x) \notin Z\}\right) = 0.$$

In other words, for almost all η (except countably many positive numbers), iteration $x(t+1) = x(t) - \eta F(x(t)) = \overline{F}_\eta^t(x(0))$ will not enter Z almost surely, provided that $x(0)$ is sampled from μ.

Theorem C.16 and Theorem C.17 follows immediately from Theorem C.18.

Proof sketch of Theorem C.16. Let $F(x) = \nabla L(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|_2})$ and $Z = \{x \in \mathbb{R}^D \mid \nabla L(x) = 0\}$. We can easily check F is $\mathcal{C}^1$ on $\mathbb{R}^D \setminus Z$ and by assumption Z is a zero-measure set. Applying Theorem C.18 with $F^n \equiv F$ for all $n \in \mathbb{N}^+$, we get the desired results. $\qquad\square$

Proof sketch of Theorem C.17. Let $G^k(x) = \nabla L(x + \rho \frac{\nabla L_k(x)}{\|\nabla L_k(x)\|_2})$ and $Z = \cup_{k=1}^M \{x \in \mathbb{R}^D \mid \nabla L_k(x) = 0\}$. We can easily check F_k is $\mathcal{C}^1$ on $\mathbb{R}^D \setminus Z$ and by assumption Z is a zero-measure set. Applying Theorem C.18 with $F^n = G^{k_n}$ for all $n \in \mathbb{N}^+$ where k_n is the nth data/batch sampled by the algorithm, we get the desired results. $\qquad\square$

Now we will turn to the proof of Theorem C.18, which is based on the following two lemmas.

Lemma C.19. *Let Z be a closed subset of $\mathbb{R}^D$ with zero Lebesgue measure and $F : \mathbb{R}^D \setminus Z \to \mathbb{R}^D$ be a continuously differentiable function. Then except countably many $\eta \in \mathbb{R}^+$, $\{x \in \mathbb{R}^d \setminus Z \mid \det(\partial F_\eta(x)) = 0\}$ is a zero-measure set under Lebesgue measure.*

Lemma C.20. *Let Z be a closed subset of $\mathbb{R}^D$ with zero Lebesgue measure and $H : \mathbb{R}^D \setminus Z \to \mathbb{R}^D$ be a continuously differentiable function. If $\{x \in \mathbb{R}^d \setminus Z \mid \det(\partial H(x)) = 0\}$ is a zero-measure set, then for any zero-measure set Z', $H^{-1}(Z')$ is a zero-measure set.*

Proof of Theorem C.18. It suffices to prove that for every $N \in \mathbb{N}^+$, at most for countably many η:

$$\mu\left(\{x \in \mathbb{R}^D \mid \overline{F}_\eta^N(x) \in Z \text{ and } \forall 0 \leq i \leq N-1, \overline{F}_\eta^i(x) \notin Z\}\right) = 0. \tag{22}$$

The desired results is immediately implied by the above claim because the countable union of countable set is still countable, and countable union of zero-measure set is still zero measure.

To prove *Equation* (22), we first introduce some notations. For any $\eta > 0$, $0 \leq n \leq N-1$, and $x \in \mathbb{R}^D$, we define $\overline{F}_\eta^{-(n+1)}(x) \triangleq (F_\eta^{N-n})^{-1}(\overline{F}_\eta^{-n}(x))$, where $\overline{F}_\eta^0(x) = x$. We extend the

---

[3]Though we call Equation (16) 1-SAM, but our result here applies to any batch size where L_k can be regarded as the loss for k-th possible batch and M is the number of the total number of batches.

definition to set in a natural way, namely $\overline{F}_\eta^{-n}(S) \triangleq \cup_{x \in S} \overline{F}_\eta^{-n}(x)$ for any $S \subseteq \mathbb{R}^D$. Under this notation, we have that

$$F_\eta^{-N}(Z) = \mu \left(\{ x \in \mathbb{R}^D \mid \overline{F}_\eta^N(x) \in Z \text{ and } \forall 0 \leq i \leq N-1, \overline{F}_\eta^i(x) \notin Z \} \right)$$

We will prove by induction. We claim that for each $0 \leq n \leq N$ except for countably many $\eta \in \mathbb{R}^+$, $\overline{F}_\eta^{-n}(Z)$ has zero Lebesgue measure. The base case $n = 0$ is by trivial as Z is assumed to be zero-measure. Suppose this holds for n. By Lemma C.19, except countably many $\eta \in \mathbb{R}^+$, $\{ x \in \mathbb{R}^d \setminus \overline{F}_\eta^{-n}(Z) \mid \det(\partial F_\eta^{N-n-1}(x)) = 0 \}$ is a zero-measure set. Next by Lemma C.20 if for some $\eta \in \mathbb{R}^+$, $\{ x \in \mathbb{R}^d \setminus \overline{F}_\eta^{-n}(Z) \mid \det(\partial F_\eta^{N-n-1}(x)) = 0 \}$ is a zero-measure set, then $\overline{F}_\eta^{-n-1}(Z) = (F_\eta^{N-n-1})^{-1}(\overline{F}_\eta^{-n}(Z))$ is a zero-measure set. Then by induction, we know that except countably many $\eta \in \mathbb{R}^+$, for all integer $0 \leq n \leq N$, $\overline{F}_\eta^{-n}(Z)$ is zero-measure. Since μ is absolutely continuous to Lebesgue measure, $\mu(F_\eta^{-N}(Z)) = 0$. $\qquad \square$

We end this section with the proofs of Lemmas C.19 and C.20.

Proof of Lemma C.19. We use $\lambda_i(x)$ to denote that the real part of the ith eigenvalue of the matrix $\partial F(x)$ in the descending order. Since $\partial F(x)$ is continuous in x, $\lambda_i(x)$ is continuous in x as well, for any $i \in [D]$, and thus $\{ x \in \mathbb{R}^D \setminus Z \mid \lambda_i(x) = 1/\eta \}$ is a measurable set. Note that for a fixed $i \in [D]$, for each positive integer n, let I_n be the set of η where $\mu(\{ x \in \mathbb{R}^D \setminus Z \mid \lambda_i(x) = 1/\eta \}) > 1/n$, then $|I_n| \leq n$, because

$$\frac{|I_n|}{n} \leq \sum_{\eta \in I_n} \mu((\{ x \in \mathbb{R}^D \setminus Z \mid \lambda_i(x) = 1/\eta \}) \leq \mu((\{ x \in \mathbb{R}^D \setminus Z \mid 1/\lambda_i(x) \in I_n \}) \leq 1.$$

Therefore, there are at most countably many $\eta \in \mathbb{R}^+$, such that $\mu(\{ x \in \mathbb{R}^D \setminus Z \mid \lambda_i(x) = 1/\eta \}) > 0$. Further note that $\det(\partial F_\eta(x)) = 0 \iff \exists i \in [D], \lambda_i(x) = 1/\eta$, we know that there are at most countably many $\eta \in \mathbb{R}^+$, such that $\mu(\{ x \in \mathbb{R}^D \setminus Z \mid \det(\partial F_\eta(x)) = 0 \}) = 0$. This completes the proof. $\qquad \square$

Proof of Lemma C.20. Denote $\{ x \in \mathbb{R}^D \setminus Z \mid \det(\partial H(x)) = 0 \}$ by Z'', since $\det(\partial H(x))$ is continuous in x as F is $\mathcal{C}^1$, Z'' is relatively closed in $\mathbb{R}^D \setminus Z$. Since Z' is a closed set, $\mathbb{R}^D \setminus (Z' \cup Z'')$ is open. Thus for all $x \in \mathbb{R}^D \setminus (Z' \cup Z'')$ with $\det(\partial H(x)) \neq 0$, there exists a open neighborhood of x, U, where for all $x' \in U$, $\det(\partial H(x')) \neq 0$, since thus $\det(\partial H(x))$ is continuous. This further implies H is invertible on U and its inverse $(H|_U)^{-1}$ is differentiable on $F(U)$. Therefore, $(H|_U)^{-1}$ maps any zero-measure set to a zero-measure set. In particular, $(H|_U)^{-1}(Z' \cap H(U))$ is zero measure, so is $(H)^{-1}(Z') \cap U \subset (H|_U)^{-1}(Z' \cap H(U))$. Now for every $x \in \mathbb{R}^D \setminus Z$ we take an open neighborhood $U_x \subseteq \mathbb{R}^D \setminus (Z' \cup Z'')$. Since $\mathbb{R}^D$ is a separable metric space, the open cover of $\mathbb{R}^D$, $\{ U_x \}_{x \in \mathbb{R}^D \setminus (Z' \cup Z'')}$ has a countable subcover, $\{ U_x \}_{x \in I}$, where I is a countable set of $\mathbb{R}^D \setminus (Z' \cup Z'')$. Therefore we have that $H^{-1}(Z') \setminus (Z' \cup Z'') = H^{-1}(Z') \cap (\mathbb{R}^D \setminus (Z' \cup Z'')) = \cup_{x \in I} H^{-1}(Z') \cap U_x$ is a zero-measure set. Thus $H^{-1}(Z')$ is also zero-measure since Z', Z'' are both zero-measure. This completes the proof. $\qquad \square$

C.4 EXPLICIT BIAS

We will first prove the generic Theorem 4.3 and then apply this theorems to characterize a variety of regularizers.

Lemma C.21. *Let U' be any bounded open set such that its closure $\overline{U'} \subseteq U$. Further assume $\overline{U'} \cap \Gamma = \overline{U'} \cap \overline{\Gamma}$, then for all $h_2 > 0, \exists \rho_0 > 0$ if $x \in U', \text{dist}(x, \Gamma) \leq \rho_0 \Rightarrow \text{dist}(x, \overline{U'} \cap \Gamma) \leq h_2$*

Proof. We will prove by contradiction. Suppose there exists $h_2 > 0$ and $\{x_k\}_{k=1}^\infty \in U'$, such that $\lim_{k \to \infty} \text{dist}(x_k, \Gamma) = 0$ but $\forall k > 0, \text{dist}(x_k, \overline{U'} \cap \Gamma) \geq h_2$. Since U' is bounded, $\overline{U'}$ is compact and thus $\{x_k\}_{k=1}^\infty$ has at least one accumulate point x^* in $\overline{U'} \subseteq U$. Since U is the attraction set of Γ under gradient flow, we know that $\Phi(x^*) \in \Gamma$. Now we claim $x^* \in \Gamma$. This is because $\lim_{k \to \infty} \text{dist}(x_k, \Gamma) = 0$ and thus there exists a sequence of points on Γ, $\{y_k\}_{k=1}^\infty$, where

$\lim_{k\to\infty} \|x_k - y_k\| = 0$. Thus we have that $x^* = \lim_{k\to\infty} y_k = \lim_{k\to\infty} \Phi(y_k) = \Phi(x^*)$, where the last step we used that $x^* \in U$ and Φ is continuous on U. By the definition of U, $x^* \in U \iff \Phi(x^*) \in \Gamma$, thus $x^* \in \Gamma$. Then we would have $x^* \in \overline{U'} \cap \Gamma$, which is contradictory to $\mathrm{dist}(x_k, \overline{U'} \cap \Gamma) \geq \mathrm{dist}(x_k, \overline{U' \cap \Gamma}) \geq h_2, \forall k > 0$. This completes the proof. $\quad\square$

Lemma C.22. *Let U' be any bounded open set such that its closure $\overline{U'} \subseteq U$. then for all $h_2 > 0, \exists \rho_1 > 0$ if $x \in U', L(x) \leq \rho_1 \Rightarrow \mathrm{dist}(x, \overline{U' \cap \Gamma}) \leq h_2$*

Proof. We will prove by contradiction. If there exists a list of $\rho_1, ..., \rho_k, ...$, such that $\rho_k \to 0$ and there exists $x_k \in U'$, such that $L(x_k) \leq \inf_{x\in U'} L(x) + \rho_k$ and $\mathrm{dist}(x_k, \overline{U' \cap \Gamma}) \geq h_2$. Since U' is bounded, $\overline{U'}$ is compact and thus $\{x_k\}_{k=1}^{\infty}$ has at least one accumulate point x^* in $\overline{U'} \subseteq U$. Since L is continuous in U, $L(x^*) = \lim_{k\to\infty} L(x_k) = \inf_{x\in U'} L(x)$. Thus x^* is a local minimizer of L and thus has zero gradient, which further implies that $x^* = \Phi(x^*)$. Thus $x^* \in \overline{U'} \cap \Gamma$, which is contradictory to $\mathrm{dist}(x_k, \overline{U'} \cap \Gamma) \geq \mathrm{dist}(x_k, \overline{U' \cap \Gamma}) \geq h_2, \forall k > 0$. This completes the proof. $\quad\square$

Proof of Theorem 4.3 . We will first prove $\Rightarrow$ side. The proof consists of four steps. Define h as the constant in Lemma C.5 with $K = \overline{U' \cap H}$.

Step 1 Consider $x_0 \in U' \cap \Gamma$, satisfying

$$S(x_0) \leq \inf_{x\in U'\cap\Gamma} S(x) + O(\rho).$$

Then using definition 4.2, we have there exists sufficiently small $\epsilon_\rho < \rho^2$, such that exists $\|x_1 - x_0\| \leq \epsilon_\rho < \rho^2$ and $\|S(x_0) - R_\rho(x_1)/\rho^2\| = O(\rho^2)$. Now that as $x_0 \in U'$, we have for sufficiently small ρ, $x_1 \in U'$. By Lemma C.21, we have for sufficiently small ρ, $\overline{x_0 x_1} \in \overline{U' \cap H}^h$, then consider doing Taylor Expansion, we would have $L(x_1) + R_\rho(x_1) \leq \rho^2 S(x_0) + o(\rho^2)$.

Step 2 Now consider $L(u) + R_\rho(u) \leq \rho^2 \inf_{x\in U'\cap\Gamma} S(x) + \epsilon\rho^2 + o(\rho^2)$. We easily have $L(u) \leq O(\rho^2)$ by Assumption 4.1. By Lemma C.22, we have $\overline{u\Phi(u)} \in K^h$ and that $\|u - \Phi(u)\| = O(\rho)$ by Lemma C.7. This implies $\mathrm{dist}(\overline{U' \cap H}, \Phi(u)) = o(1)$. Notice now, we have shown $R_\rho(u) \leq \rho^2 \inf S(x) + \rho^2 \epsilon + o(\rho^2)$

Step 3 Now applying definition of a good limiting regularizer and Finite covering theorem, we have this further implies $S(\Phi(u))\rho^2 \leq R_\rho(u) + o(\rho^2)$. We also have $S(\Phi(u)) \geq \inf_{x\in U'\cap\Gamma} S(x) - o(1)$ as $S \in \mathcal{C}^0$. Finally $R_\rho(u) \geq S(\Phi(u))\rho^2 - o(\rho^2) \geq \rho^2 \inf S(x) - o(\rho^2)$. We also have $L(u) \leq o(\rho^2)$, proving our claim.

For the $\Leftarrow$ side, we need to provide a lower bound for $L(x) + R_\rho(x)$. Define constant C_1 such that for all $x, \|x - \Phi(x)\| \geq C_1\rho$, we have $L(x) \geq (C_{U'} + \inf_{x\in U'\cap\Gamma} S(x) + 1)\rho^2$ where $C_{U'}$ is the constant in Assumption 4.1. Then we have for x such that $\|x - \Phi(x)\| \geq C_1\rho$, we have $L(x) + R_\rho(x) \geq (\inf_{x\in U'\cap\Gamma} S(x) + 1)\rho^2$. For $\|x - \Phi(x)\| \leq C_1\rho$, we have by definition of good limiting regularizer and Finite covering theorem, $R_\rho(x) \geq \rho^2 S(\Phi(x)) - o(\rho^2)$. As $\mathrm{dist}\Phi(x), \overline{U' \cap \Gamma} = o(1)$, we have $R_\rho(x) \geq \inf_{x\in U'\cap\Gamma} S(x) - o(\rho^2)$, hence $L(x) + R_\rho(x) \geq \inf_{x\in U'\cap\Gamma} S(x)\rho^2$. $\quad\square$

Theorem 4.3 implies Corollary C.23 saying that the minimum of the regularized loss is approximately the sum of the minimum of loss and the minimum of limiting regularizer.

Corollary C.23. *Under the setting of Theorem 4.3,*

$$\left| \inf_{x\in U'} (L(x) + R_\rho(x)) - \inf_{x\in U'} L(x) - \rho^2 \inf_{x\in U'\cap\Gamma} S(x) \right| \leq o(\rho^2)$$

Proof of Corollary C.23. In Theorem 4.3, choose $\epsilon = 0$, and choose u such that $L(u) + R_\rho(u) \leq \inf_{x\in U'}(L(x) + R_\rho(x)) + o(\rho^2)$, the result is then clear. $\quad\square$

We will then prove the Corollary 4.4.

Proof of Corollary 4.4. By Theorem 4.3, we have $\|L(u) - \inf_{x\in U'} L(x)\| = O(\rho^2)$. This implies $\|u - \Phi(u)\| = O(\rho)$ and also $\mathrm{dist}(\overline{U' \cap H}, \Phi(u)) = o(1)$.

We also have $\rho^2 S(\Phi(x)) - o(\rho^2) \leq R_\rho(u) \leq \rho^2 \inf_{x \in U' \cap H} S(x)$. We have $\|S(\Phi(x)) - \inf_{x \in U' \cap H} S(x)\| = o(1)$. This further implies $\|\overline{S}(\Phi(x)) - \inf_{x \in U' \cap H} S(x)\| = o(1)$

$\square$

Proof of Theorem 4.5.

1. We will first verify Assumption 4.1. For fixed compact set $B \subset U$, as $\|\nabla^3 L(x)\|_2$ is continuous, there exists constant ν, such that $\forall x \in B^1$, $\|\nabla^3 L(x)\|_2 \leq \nu$. Then by Taylor Expansion,

$$R_\rho^{\mathrm{Max}}(x) = \max_{\|v\|_2 \leq 1} L(x + \rho v) - L(x)$$
$$\geq \max_{\|v\|_2 \leq 1} \left(\rho \langle \nabla L(x), v \rangle + \rho^2 v^T \nabla^2 L(x) v / 2 \right) - \nu \rho^3 / 6$$
$$\geq -\nu \rho^3 / 6 .$$

2. Now we verify $S^{\mathrm{Max}}(x) = \lambda_1(\nabla^2 L(\cdot))/2$ is the limiting regularizer of R_ρ^{Max}. Let x be any point in Γ, by continuity of R_ρ^{Max},

$$\lim_{\rho \to 0} \lim_{r \to 0} \inf_{\|x' - x\|_2 \leq r} \frac{R_\rho^{\mathrm{Max}}(x')}{\rho^2} = \lim_{\rho \to 0} \frac{R_\rho^{\mathrm{Max}}(x)}{\rho^2} = \lambda_1(\nabla^2 L(x))/2 .$$

3. Finally we verify definition of good limiting regularizer, by Assumption 3.1, $S^{\mathrm{Max}}(x) = \lambda_1(x)/2$ is non-negative and continuous on Γ. For any $x^* \in \Gamma$, choose a sufficiently small open convex set V containing x^* such that $\forall x \in V^1$, $\|\nabla^3 L(x)\|_2 \leq \nu$. For any $x \in V \cap \Gamma$, for any x' satisfying that $\|x' - x\|_2 \leq C\rho$, by Theorem C.58,

$$R_\rho^{\mathrm{Max}}(x') = \max_{\|v\|_2 \leq 1} L(x' + \rho v) - L(x') \geq \max_{\|v\|_2 \leq 1} \left(\rho \langle \nabla L(x'), v \rangle + \rho^2 v^T \nabla^2 L(x') v / 2 \right) - \nu \rho^3 / 6$$
$$\geq \rho^2 \lambda_1(\nabla^2 L(x'))/2 - \nu \rho^3 / 6 \geq \rho^2 \lambda_1(\nabla^2 L(x))/2 - O(\rho^3) .$$

This implies $\inf_{\|x' - x\|_2 \leq C\rho} R_\rho^{\mathrm{Max}}(x') \geq \rho^2 \lambda_1(\nabla^2 L(x))/2 - O(\rho^3)$.
On the other hand, for any $x \in V \cap \Gamma$,

$$R_\rho^{\mathrm{Max}}(x) = \max_{\|v\|_2 \leq 1} L(x + \rho v) - L(x) \leq \max_{\|v\|_2 \leq 1} \left(\rho \langle \nabla L(x), v \rangle + \rho^2 v^T \nabla^2 L(x) v / 2 \right) + \nu \rho^3$$
$$= \max_{\|v\|_2 \leq 1} \rho^2 v^T \nabla^2 L(x) v / 2 + \nu \rho^3 = \rho^2 \lambda_1(\nabla^2 L(x'))/2 + O(\rho^3) .$$

This implies $\inf_{\|x' - x\|_2 \leq C\rho} R_\rho^{\mathrm{Max}}(x') \leq \rho^2 \lambda_1(\nabla^2 L(x))/2 + O(\rho^3)$.

Thus, we conclude that $\left| \inf_{\|x' - x\|_2 \leq C\rho} R_\rho^{\mathrm{Max}}(x')/\rho^2 - \lambda_1(\nabla^2 L(x))/2 \right| = O(\rho), \forall x \in V \cap \Gamma$, indicating S^{Max} is a good limiting regularizer of R_ρ^{Max} on Γ.

$\square$

Proof of Theorem 4.6.

1. We will first prove Assumption 4.1 holds. For any fixed compact set $B \subset U$, as $\lambda_1(\nabla^2 L)$ and $\|\nabla^3 L\|$ is continuous, there exists constant C, such that $\forall x \in B^2$, $\lambda_1(\nabla^2 L) > -\zeta$ and $\|\nabla^3 L(x)\| < \nu$. Then by Taylor Expansion,

$$R_\rho^{\mathrm{Asc}}(x) = L(x + \rho \frac{\nabla L(x)}{\|\nabla L(x)\|}) - L(x)$$
$$\geq \left(\rho \|\nabla L(x)\|_2 + \rho^2 (\frac{\nabla L(x)}{\|\nabla L(x)\|})^T \nabla^2 L(x) \frac{\nabla L(x)}{\|\nabla L(x)\|} / 2 \right) - \nu \rho^3 / 6$$
$$\geq -(\zeta + \nu/6)\rho^2 .$$

2. Now we verify $S^{\text{Asc}}(x) = \text{Tr}(\nabla^2 L(\cdot))/2$ is the limiting regularizer of R_ρ^{Asc}. Let x be any point in Γ. Let $K = \{x\}$ and choose $h = h(K)$ as in Lemma C.5. For any $x' \in K^h \cap U'$,

$$R_\rho^{\text{Asc}}(x') = L(x' + \rho\frac{\nabla L(x')}{\|\nabla L(x')\|}) - L(x')$$

$$\geq \rho\|\nabla L(x')\|_2 + \rho^2(\frac{\nabla L(x')}{\|\nabla L(x')\|})^T \nabla^2 L(x')\frac{\nabla L(x')}{\|\nabla L(x')\|}/2 - \nu\rho^3/6$$

$$\geq \rho^2(\frac{\nabla L(x')}{\|\nabla L(x')\|})^T \nabla^2 L(\Phi(x'))\frac{\nabla L(x')}{\|\nabla L(x')\|}/2 - \nu\rho^3/6\,.$$

By Lemma C.10, we have $\frac{\nabla L(x')}{\|\nabla L(x')\|} = \frac{\nabla^2 L(\Phi(x'))(x' - \Phi(x'))}{\|\nabla^2 L(\Phi(x'))(x' - \Phi(x'))\|_2} + O(\frac{\nu}{\mu}\|x' - \Phi(x')\|_2)$. Hence

$$R_\rho^{\text{Asc}}(x') \geq \rho^2\lambda_M(\nabla^2 L(\Phi(x')))/2 - \zeta\rho^2 O(\|x' - \Phi(x')\|_2) - \nu\rho^3/6\,.$$

This implies $\lim_{\rho\to 0}\lim_{r\to 0}\inf_{\|x'-x\|_2\leq r}\frac{R_\rho^{\text{Asc}}(x')}{\rho^2} \geq \lambda_M(\nabla^2 L(\Phi(x')))/2$.
We now show the above inequality is in fact equality. If we choose $x_r'' = x + rv_M$, then by Taylor Expansion

$$\nabla L(x_r'') = \nabla L(x) + \nabla^2 L(x)(x_r'' - x) + O(\|x_r'' - x\|^2)$$
$$= rv_M + O(r^2)$$

This implies

$$\lim_{r\to 0}\frac{\nabla L(x_r'')}{\|\nabla L(x_r'')\|} = v_M.$$

We also have $\lim_{r\to 0}\nabla^2 L(x_r'') = \nabla^2 L(x)$ and $\lim_{r\to 0}\nabla^L(x_r'') = 0$. Putting together,

$$\lim_{r\to 0} R_\rho^{\text{Asc}}(x_r'') = \lim_{r\to 0} L(x_r'' + \rho\frac{\nabla L(x_r'')}{\|\nabla L(x_r'')\|}) - L(x_r'')$$

$$= \lim_{r\to 0}\left(\rho\|\nabla L(x_r'')\|_2 + \rho^2(\frac{\nabla L(x_r'')}{\|\nabla L(x_r'')\|})^T \nabla^2 L(x_r'')\frac{\nabla L(x_r'')}{\|\nabla L(x_r'')\|}/2 + O(\nu\rho^3)\right)$$

$$= \rho^2\lambda_M(\nabla^2 L(x))/2 + O(\rho^3).$$

This implies $\lim_{\rho\to 0}\lim_{r\to 0}\inf_{\|x'-x\|_2\leq r}\frac{R_\rho^{\text{Asc}}(x')}{\rho^2} \leq \lim_{\rho\to 0}\lim_{r\to 0}\frac{R_\rho^{\text{Asc}}(x_r'')}{\rho^2} = \lambda_M(\nabla^2 L(x))/2$.
Hence the limiting regularizer S is exactly $\lambda_M(\nabla^2 L(\cdot))/2$.

3. Finally we verify definition of good limiting regularizer, by Assumption 3.1, $S^{\text{Max}}(x) = \lambda_M(x)/2$ is non-negative and continuous on Γ. For any $x^* \in \Gamma$, choose a sufficiently small open convex set V containing x^* such that $\forall x \in V^1, \|\nabla^3 L(x)\|_2 \leq \nu$. For any $x \in V \cap \Gamma$, for any x' satisfying that $\|x' - x\|_2 \leq C\rho$,

$$R_\rho^{\text{Asc}}(x') = L(x' + \rho\frac{\nabla L(x')}{\|\nabla L(x')\|}) - L(x')$$

$$\geq \rho\|\nabla L(x')\|_2 + \rho^2(\frac{\nabla L(x')}{\|\nabla L(x')\|})^T \nabla^2 L(x')\frac{\nabla L(x')}{\|\nabla L(x')\|}/2 - \nu\rho^3/6$$

$$\geq \rho^2(\frac{\nabla L(x')}{\|\nabla L(x')\|})^T \nabla^2 L(\Phi(x'))\frac{\nabla L(x')}{\|\nabla L(x')\|}/2 - \nu\rho^3/6\,.$$

By Lemma C.10, we have $\frac{\nabla L(x')}{\|\nabla L(x')\|} = \frac{\nabla^2 L(\Phi(x'))(x' - \Phi(x'))}{\|\nabla^2 L(\Phi(x'))(x' - \Phi(x'))\|_2} + O(\frac{\nu}{\mu}\|x' - \Phi(x')\|_2)$. This implies $\inf_{\|x'-x\|_2\leq C\rho} R_\rho^{\text{Asc}}(x') \geq \rho^2\lambda_M(\nabla^2 L(x))/2 - O(\rho^3)$.

On the other hand, similar to the proof in the second part, we have $\inf_{\|x'-x\|_2\leq C\rho} R_\rho^{\text{Asc}}(x') \leq \rho^2\lambda_M(\nabla^2 L(x))/2 + O(\rho^3)$.

Thus, we conclude that $\left|\inf_{\|x'-x\|_2\leq C\rho} R_\rho^{\text{Max}}(x')/\rho^2 - \lambda_1(\nabla^2 L(x))/2\right| = O(\rho), \forall x \in V \cap \Gamma$, indicating S^{Max} is a good limiting regularizer of R_ρ^{Max} on Γ.

$\square$

Proof of Theorem 4.7.

1. We will first verify Assumption 4.1. For fixed compact set $B \subset U$, as $\|\nabla^3 L(x)\|_2$ is continuous, there exists constant ν, such that $\forall x \in B^1$, $\|\nabla^3 L(x)\|_2 \le \nu$. Then by Taylor Expansion,

$$R_\rho^{\text{Avg}}(x) = \mathbb{E}_{g \sim N(0,I)} L(x + \rho \frac{g}{\|g\|}) - L(x)$$

$$\ge \mathbb{E}_{g \sim N(0,I)} \left(\rho \langle \nabla L(x), \frac{g}{\|g\|} \rangle + \rho^2 (\frac{g}{\|g\|})^T \nabla^2 L(x) \frac{g}{2\|g\|} \right) - \nu \rho^3 / 6$$

$$\ge -\nu \rho^3 / 6 \,.$$

2. Now we verify $S^{\text{Max}}(x) = \text{Tr}(\nabla^2 L(\cdot))/2D$ is the limiting regularizer of R_ρ^{Avg}. Let x be any point in Γ, by continuity of R_ρ^{Avg},

$$\lim_{\rho \to 0} \lim_{r \to 0} \inf_{\|x'-x\|_2 \le r} \frac{R_\rho^{\text{Avg}}(x')}{\rho^2} = \lim_{\rho \to 0} \frac{R_\rho^{\text{Avg}}(x)}{\rho^2} = \text{Tr}(\nabla^2 L(x))/2D \,.$$

3. Finally we verify definition of good limiting regularizer, by Assumption 3.1, $S^{\text{Avg}}(x) = \text{Tr}(x)/2D$ is non-negative and continuous on Γ. For any $x^* \in \Gamma$, choose a sufficiently small open convex set V containing x^* such that $\forall x \in V^1, \|\nabla^3 L(x)\|_2 \le \nu$. For any $x \in V \cap \Gamma$, for any x' satisfying that $\|x'-x\|_2 \le C\rho$, by Theorem C.58,

$$R_\rho^{\text{Avg}}(x') = \mathbb{E}_{g \sim N(0,I)} L(x' + \rho \frac{g}{\|g\|}) - L(x')$$

$$\ge \mathbb{E}_{g \sim N(0,I)} \left(\rho \langle \nabla L(x'), \frac{g}{\|g\|} \rangle + \rho^2 \frac{g}{\|g\|}^T \nabla^2 L(x') \frac{g}{\|2g\|} \right) - \nu \rho^3 / 6$$

$$\ge \rho^2 \text{Tr}(\nabla^2 L(x'))/2D - \nu \rho^3 / 6 \ge \rho^2 \text{Tr}(\nabla^2 L(x))/2D - O(\rho^3) \,.$$

This implies $\inf_{\|x'-x\|_2 \le C\rho} R_\rho^{\text{Avg}}(x') \ge \rho^2 \text{Tr}(\nabla^2 L(x))/2D - O(\rho^3)$.
On the other hand, for any $x \in V \cap \Gamma$,

$$R_\rho^{\text{Avg}}(x) = \mathbb{E}_{g \sim N(0,I)} L(x + \rho \frac{g}{\|g\|}) - L(x)$$

$$\le \mathbb{E}_{g \sim N(0,I)} \left(\rho \langle \nabla L(x), \frac{g}{\|g\|} \rangle + \rho^2 \frac{g}{\|g\|}^T \nabla^2 L(x) \frac{g}{2\|g\|} \right) + \nu \rho^3$$

$$= \mathbb{E}_{g \sim N(0,I)} \rho^2 \frac{g}{\|g\|}^T \nabla^2 L(x) \frac{g}{2\|g\|} + \nu \rho^3 = \rho^2 \text{Tr}(\nabla^2 L(x'))/2D + O(\rho^3) \,.$$

This implies $\inf_{\|x'-x\|_2 \le C\rho} R_\rho^{\text{Avg}}(x') \le \rho^2 \text{Tr}(\nabla^2 L(x))/2D + O(\rho^3)$.

Thus, we conclude that $\left| \inf_{\|x'-x\|_2 \le C\rho} R_\rho^{\text{Avg}}(x')/\rho^2 - \text{Tr}(\nabla^2 L(x))/2D \right| = O(\rho), \forall x \in V \cap \Gamma$, indicating S^{Avg} is a good limiting regularizer of R_ρ^{Avg} on Γ.

$\square$

Proof of Theorem 5.1. By Theorem C.53, Assumption C.2 holds.

Easily deducted from Theorem 4.5 $\Lambda_k(x)$ is a good limiting regularizer for $R_{k,\rho}^{\max}$ on Γ_k. Then as $\Gamma \subset \Gamma_k$, $\Lambda_k(x)$ is a good limiting regularizer for $R_{k,\rho}^{\max}$ on Γ. Hence $S(x) = \sum_k \Lambda_k(x)/2M = \text{Tr}(\nabla^2 \text{L}(x))/2$ is a good limiting regularizer of $\mathbb{E}_k[R_{k,\rho}^{\text{Max}}](x)$ on Γ. $\square$

Proof of Theorem 5.2. By Theorem C.53, Assumption C.2 holds.

Easily deducted from Theorem 4.6 $\Lambda_k(x)$ is a good limiting regularizer for $R_{k,\rho}^{\text{asc}}$ on Γ_k as the codimension of Γ_k is 1. Then as $\Gamma \subset \Gamma_k$, $\Lambda_k(x)$ is a good limiting regularizer for $R_{k,\rho}^{\max}$ on Γ. Hence $S(x) = \sum_k \Lambda_k(x)/2M = \text{Tr}(\nabla^2 L(x))/2$ is a good limiting regularizer of $\mathbb{E}_k[R_{k,\rho}^{\text{Asc}}](x)$ on Γ. $\square$

C.5 FULL-BATCH SAM ON QUADRATIC LOSS (PROOF OF THEOREM 4.14)

The goal of this section is to prove Theorem 4.14. In this section, we use $A \prec B$ to indicate $B - A$ is positive semi-definite.

Theorem 4.14. *Suppose A is a positive definite symmetric matrix with unique top eigenvalue. Consider running SAM on loss $L(x) := \frac{1}{2} x^T A x$ as (11), then for almost every $x(0)$, we have $x(t)$ converges in direction to $v_1(A)$ and $\lim_{t \to \infty} \|x(t)\| = \frac{\eta \rho \lambda_1(A)}{2 - \eta \lambda_1(A)}$ with $\eta \lambda_1(A) < 1$.*

$$x(t+1) = x(t) - \eta A \left(x(t) + \rho \frac{A x(t)}{\|A x(t)\|} \right), \tag{11}$$

Proof of Theorem 4.14. We first rewrite the iterate as

$$x(t+1) = x(t) - \eta A x(t) - \eta \rho \frac{A^2 x(t)}{\|A x(t)\|_2}.$$

Define $\tilde{x}(t) \triangleq \frac{\nabla L(x(t))}{\rho} = \frac{A x(t)}{\rho}$, and we have

$$\tilde{x}(t+1) = \tilde{x}(t) - \eta A \tilde{x}(t) - \eta \frac{A^2 \tilde{x}(t)}{\|\tilde{x}(t)\|_2}. \tag{23}$$

We suppose $A \in R^{D \times D}$ and use λ_i, v_i to denote $\lambda_i(A), v_i(A)$.

Further, we define that

$$P^{(j:D)} \triangleq \sum_{i=j}^{D} v_i(A) v_i(A)^T,$$

$$\mathbb{I}_j \triangleq \{\tilde{x} \mid \|P^{(j:D)} \tilde{x}\|_2 \le \eta \lambda_j^2\},$$

$$\tilde{x}_i(t) \triangleq \langle \tilde{x}(t), v_i \rangle,$$

$$S \triangleq \{t \mid \|\tilde{x}(t)\|_2 \le \frac{\eta \lambda_1^2}{2 - \eta \lambda_1}, t > T_1\}.$$

By Lemma C.24, $\mathbb{I}_j$ is an invariant set for update rule Equation (23).

Our proof consists of two steps.

(1) *Entering Invariant Set.* Lemma C.25 implies that there exists constant $T_1 > 0$, such that $\forall t > T_1, \|P^{(j:D)} \tilde{x}(t)\|_2 \le \eta \lambda_j^2$

(2) *Alignment to Top Eigenvector.* Lemmas C.33 and C.34 show that $\|\tilde{x}(t)\|_2$ and $|\tilde{x}_1(t)|$ converge to $\frac{\eta \lambda_1^2}{2 - \eta \lambda_1}$, which implies our final results.

$\square$

C.5.1 ENTERING INVARIANT SET

In this subsection, we will prove the following three lemmas.

1. Lemma C.24 shows $\mathbb{I}_j$ is an invariant set for update rule (Equation (23)).
2. Lemma C.25 shows that under the update rule (Equation (23)), all iterates not in $\mathbb{I}_j$ will shrink exponentially in ℓ_2 norm.
3. Lemma C.26 combines Lemmas C.24 and C.25 to show that for sufficiently large t, $x(t) \in \cap_j \mathbb{I}_j$.

Lemma C.24. *For $t \ge 0$, if $\eta \lambda_1(A) < 1$ and $\tilde{x}(t) \in \mathbb{I}_j$, then $\tilde{x}(t+1) \in \mathbb{I}_j$.*

Proof of Lemma C.24. By (Equation (23)), we have that

$$P^{(j:D)} \tilde{x}(t+1) = (I - P^{(j:D)} \eta A - \eta \frac{P^{(j:D)} A^2}{\|\tilde{x}(t)\|_2}) P^{(j:D)} \tilde{x}(t).$$

Hence we have that

$$\|P^{(j:D)}\tilde{x}(t+1)\|_2 = \|(I - P^{(j:D)}\eta A - \eta \frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2})P^{(j:D)}\tilde{x}(t)\|_2$$

$$\leq \|I - P^{(j:D)}\eta A - \eta \frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2}\|_2\|P^{(j:D)}\tilde{x}(t)\|_2.$$

Because $\tilde{x}(t) \in \mathbb{I}_j$, $\|\tilde{x}(t)\|_2 \leq \frac{\eta\lambda_j^2}{1-\eta\lambda_j}$. This implies,

$$I(1 - \eta\lambda_j - \eta\frac{\lambda_j^2}{\|P^{(j:D)}\tilde{x}(t)\|_2}) \prec I(1 - \eta\lambda_j - \eta\frac{\lambda_j^2}{\|\tilde{x}(t)\|_2}) \prec I - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2} \prec I.$$

Furthermore, $\|I - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|_2\tilde{x}(t)\|}\|_2 \leq \max(1, \eta\lambda_j + \eta\frac{\lambda_j^2}{\|P^{(j:D)}\tilde{x}(t)\|_2} - 1).$

Hence it holds that

$$\|P^{(j:D)}\tilde{x}(t+1)\|_2 \leq \max(\|P^{(j:D)}\tilde{x}(t)\|_2, \eta\lambda_j^2 - (1 - \eta\lambda_j)\|P^{(j:D)}\tilde{x}(t)\|_2) \leq \eta\lambda_j^2,$$

where the last equality is because $1 - \eta\lambda_j \geq 0$. This above inequality is exactly the definition of $\tilde{x}(t+1) \in \mathbb{I}_j$ and thus is proof is completed. □

Lemma C.25. *For $t \geq 0$, if $\eta\lambda_1(A) < 1$ and $\tilde{x}(t) \notin \mathbb{I}_j$, then*

$$\|P^{(j:D)}\tilde{x}(t+1)\|_2 \leq \max\left(1 - \eta\lambda_D - \eta\frac{\lambda_D^2}{\|\tilde{x}(t)\|_2}, \eta\lambda_j\right)\|P^{(j:D)}\tilde{x}(t)\|_2 \qquad (24)$$

$$\leq \max\left(1 - \eta\lambda_D, \eta\lambda_j\right)\|P^{(j:D)}\tilde{x}(t)\|_2.$$

Proof of Lemma C.25. Note that

$$\|P^{(j:D)}\tilde{x}(t+1)\|_2 = \|(I - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2})P^{(j:D)}\tilde{x}(t)\|_2$$

$$\leq \|P^{(j:D)} - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2}\|_2\|P^{(j:D)}\tilde{x}(t)\|_2.$$

As $\tilde{x}(t) \notin \mathbb{I}_j$, We have $\|\tilde{x}(t)\|_2 \geq \|P^{(j:D)}\tilde{x}(t)\|_2 > \eta\lambda_j^2$, hence $\eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2} \prec \eta\frac{P^{(j:D)}A^2}{\eta\lambda_j^2} \prec P^{(j:D)}$.

This implies that

$$-\eta\lambda_j P^{(j:D)} \prec -P^{(j:D)}\eta A \prec P^{(j:D)} - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2},$$

and

$$P^{(j:D)} - P^{(j:D)}\eta A - \eta\frac{P^{(j:D)}A^2}{\|\tilde{x}(t)\|_2} \prec P^{(j:D)}(1 - \eta\lambda_D) - \eta\frac{\lambda_D^2}{\|\tilde{x}(t)\|_2}.$$

Hence we have that

$$\|P^{(j:D)}\tilde{x}(t+1)\|_2 \leq \max\left(1 - \eta\lambda_D - \eta\frac{\lambda_D^2}{\|\tilde{x}(t)\|_2}, \eta\lambda_j\right)\|P^{(j:D)}\tilde{x}(t)\|_2.$$

$$\leq \max\left(1 - \eta\lambda_D, \eta\lambda_j\right)\|P^{(j:D)}\tilde{x}(t)\|_2$$

This completes the proof. □

Lemma C.26. *Choosing $T_1 = \max_j\left(-\log_{\max(1-\eta\lambda_D,\eta\lambda_j)}\max(\frac{\|\tilde{x}(0)\|_2}{\eta\lambda_j^2}, 1)\right)$, then $\forall t \geq T_1, D > j \geq 1, \tilde{x}(t) \in \mathbb{I}_j$*

Proof of Lemma C.26. We will prove by contradiction. Suppose $\exists j \in [D]$ and $T > T_1$, such that $\tilde{x}(T) \notin \mathbb{I}_j$. By Lemma C.24, it holds that $\forall t < T, \tilde{x}(t) \notin \mathbb{I}_j$. Then by Lemma C.25,

$$\|P^{(j:D)}\tilde{x}(T)\|_2 \leq \max\left(1 - \eta\lambda_D, \eta\lambda_j\right)^T \|P^{(j:D)}\tilde{x}(0)\|_2 \leq \eta\lambda_j^2,$$

which leads to a contradiction. □

C.5.2   ALIGNMENT TO TOP EIGENVECTOR

In this subsection, we prove the following lemmas towards showing that $\tilde{x}(t)$ converges in direction to $v_1(A)$ up to a proper sign flip.

1. Corollary C.27 show that for almost every learning rate η and initialization x_{init}, $\tilde{x}_1(t) \neq 0$, for every $t \geq 0$. This condition is important because if $\tilde{x}_1(t) = 0$ at some step t, then for any $t' \geq t$, $\tilde{x}_1(t')$ will also be 0 and thus alignment is impossible.
2. Lemma C.28 shows that under update rule (Equation (23)), $t \notin S \Rightarrow t + 1 \in S$ for sufficiently large t, where the definition of S is $\{t | \|\tilde{x}(t)\|_2 \leq \frac{\eta \lambda_1^2}{2 - \eta \lambda_1}, t > T_1\}$.
3. Lemma C.32, a combination of Lemmas C.29 and C.30, shows that following update rule (Equation (23)), $\tilde{x}_1(t)$ increases for $t \in S$.
4. Lemma C.33 shows that $\|\tilde{x}(t)\|$ converges to $\frac{\eta \lambda_1^2}{2 - \eta \lambda_1}$ under Equation (23).
5. Lemma C.34 shows that $\|\tilde{x}_1(t)\|_2$ converges to $\frac{\eta \lambda_1^2}{2 - \eta \lambda_1}$ under Equation (23).

We will first prove that $\forall t, \tilde{x}_1(t) \neq 0$ happens for almost every learning rate η and initialization x_{init} (Corollary C.27), using a much more general result (Theorem C.18).

Corollary C.27. *Except for countably many $\eta \in \mathbb{R}^+$, for almost all initialization $x_{init} = x(0)$, it holds that for all natural number t, $\tilde{x}_1(t) \neq 0$.*

Proof of Corollary C.27. Let $F_n(x) \equiv F(x) \triangleq A(x + \rho \frac{Ax}{\|Ax\|_2})$, $\forall n \in \mathbb{N}^+, x \in \mathbb{R}^D$ and $Z = \{x \in \mathbb{R}^D \mid \langle x, v_1 \rangle = 0\}$. We can easily check F is $\mathcal{C}^1$ on $\mathbb{R}^D \setminus Z$ and Z is a zero-measure set. Applying Theorem C.18, we have the following corollary. $\square$

Lemma C.28. *For $t \geq 0$, if $\|\tilde{x}(t)\|_2 > \frac{\eta \lambda_1^2}{2 - \eta \lambda_1}, \tilde{x}(t) \in \cap \mathbb{I}_j$, then*

$$\|\tilde{x}(t+1)\|_2 \leq \max(\frac{\eta \lambda_1^2}{2 - \eta \lambda_1} - \eta \frac{\lambda_D^4}{2\lambda_1^2}, \eta \lambda_1^2 - (1 - \eta \lambda_1)\|\tilde{x}(t)\|_2)$$

Proof of Lemma C.28. Note that

$$\tilde{x}(t+1) = (I - \eta A - \eta \frac{A^2}{\|\tilde{x}(t)\|_2})\tilde{x}(t)$$

$$= \frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=1}^{D} \left((1 - \eta \lambda_j)\|\tilde{x}(t)\|_2 - \eta \lambda_j^2\right) \tilde{x}_j(t) v_j$$

Consider the following two cases.

1 If for any i, such that $|(1 - \eta \lambda_1)\|\tilde{x}(t)\|_2 - \eta \lambda_1^2| \geq |(1 - \eta \lambda_i)\|\tilde{x}(t)\|_2 - \eta \lambda_i^2|$, then we have
$$\|\tilde{x}(t+1)\|_2 \leq |(1 - \eta \lambda_1)\|\tilde{x}(t)\|_2 - \eta \lambda_1^2| = \eta \lambda_1^2 - (1 - \eta \lambda_1)\|\tilde{x}(t)\|_2.$$

2 If there exists i, such that $|(1 - \eta \lambda_1)\|\tilde{x}(t)\|_2 - \eta \lambda_1^2| < |(1 - \eta \lambda_i)\|\tilde{x}(t)\|_2 - \eta \lambda_i^2|$, then suppose WLOG, i is the smallest among such index.
As
$$\eta \lambda_i^2 - (1 - \eta \lambda_i)\|\tilde{x}(t)\|_2 < \eta \lambda_1^2 - (1 - \eta \lambda_1)\|\tilde{x}(t)\|_2 = |(1 - \eta \lambda_1)\|\tilde{x}(t)\|_2 - \eta \lambda_1^2|$$
We have $-\eta \lambda_i^2 + (1 - \eta \lambda_i)\|\tilde{x}(t)\|_2 > \eta \lambda_1^2 - (1 - \eta \lambda_1)\|\tilde{x}(t)\|_2$. Equivalently,
$$\|\tilde{x}(t)\|_2 > \frac{\eta \lambda_1^2 + \eta \lambda_i^2}{2 - \eta \lambda_1 - \eta \lambda_i} \tag{25}$$
Combining with $\tilde{x}(t) \in \mathbb{I}_1 \Rightarrow \|\tilde{x}(t)\|_2 \leq \eta \lambda_1^2$, we have $\eta < \frac{\lambda_1 - \lambda_i}{\lambda_1^2}$.
Now consider the following vertors,
$$v^{(1)}(t) \triangleq (\eta \lambda_1^2 - (1 - \eta \lambda_1)\|\tilde{x}(t)\|_2)\tilde{x}(t),$$
$$v^{(2)}(t) \triangleq ((2 - \eta \lambda_1 - \eta \lambda_i)\|\tilde{x}(t)\|_2 - \eta \lambda_i^2 - \eta \lambda_1^2)P^{(i:D)}\tilde{x}(t),$$
$$v^{(2+j)}(t) \triangleq ((\eta \lambda_{i+j-1} - \eta \lambda_{i+j})\|\tilde{x}(t)\|_2 - \eta \lambda_{i+j}^2 + \eta \lambda_{i+j-1}^2)P^{(i+j:D)}\tilde{x}(t), 1 \leq j \leq D - i.$$

Then we have

$$\|\tilde{x}(t+1)\|_2 = \|\frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=1}^{D} \left((1-\eta\lambda_j)\|\tilde{x}(t)\|_2 - \eta\lambda_j^2\right)\tilde{x}_j(t)v_j\|_2$$

$$\leq \|\frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=1}^{i-1} \left(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2\right)\tilde{x}_j(t)v_j\| +$$

$$\|\frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=i}^{D} \left((1-\eta\lambda_j)\|\tilde{x}(t)\|_2 - \eta\lambda_j^2\right)\tilde{x}_j(t)v_j\|_2$$

$$\leq \frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=1}^{D+1-i} \|v^{(j)}\|_2$$

By assumption, we have $\tilde{x}(t) \in \cap \mathbb{I}_j$, hence we have

$$\|v^{(1)}(t)\|_2 = (\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)\|\tilde{x}(t)\|_2\,,$$
$$\|v^{(2)}(t)\|_2 \leq \eta((2-\eta\lambda_1-\eta\lambda_i)\|\tilde{x}(t)\|_2 - \eta\lambda_i^2 - \eta\lambda_1^2)\lambda_i^2\,,$$
$$\|v^{(2+j)}(t)\|_2 \leq \eta((\eta\lambda_{i+j-1}-\eta\lambda_{i+j})\|\tilde{x}(t)\|_2 - \eta\lambda_{i+j}^2 + \eta\lambda_{i+j-1}^2)\lambda_{i+j}^2, 1 \leq j \leq D-i\,.$$

Using AM-GM inequality, we have

$$\lambda_{i+j-1}\lambda_{i+j}^2 \leq \frac{\lambda_{i+j-1}^3 + 2\lambda_{i+j}^3}{3}\,,$$
$$\lambda_{i+j-1}^2\lambda_{i+j}^2 \leq \frac{\lambda_{i+j-1}^4 + \lambda_{i+j}^4}{2}\,.$$

Hence

$$\|v^{(2+j)}(t)\|_2 \leq \eta((\eta\lambda_{i+j-1}-\eta\lambda_{i+j})\|\tilde{x}(t)\|_2 - \eta\lambda_{i+j}^2 + \eta\lambda_{i+j-1}^2)\lambda_{i+j}^2$$

$$\leq \eta^2\|\tilde{x}(t)\|_2 \frac{\lambda_{i+j-1}^3 - \lambda_{i+j}^3}{3} + \eta^2 \frac{\lambda_{i+j-1}^4 - \lambda_{i+j}^4}{2}, 1 \leq j \leq D-i$$

$$\sum_{j=1}^{D-i} \|v^{(2+j)}(t)\|_2 \leq \eta^2\|\tilde{x}(t)\|_2 \frac{\lambda_i^3 - \lambda_D^3}{3} + \eta^2 \frac{\lambda_i^4 - \lambda_D^4}{2}\,.$$

Putting together,

$$\|\tilde{x}(t+1)\|_2 \leq \frac{1}{\|\tilde{x}(t)\|_2} \sum_{j=1}^{D+1-i} \|v^{(i)}\|_2$$

$$\leq \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\eta\lambda_i) + \eta^2 \frac{\lambda_i^3 - \lambda_D^3}{3} - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2$$

$$- \eta^2\lambda_i^2(\lambda_i^2 + \lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} + \eta^2 \frac{\lambda_i^4 - \lambda_D^4}{2}\frac{1}{\|\tilde{x}(t)\|_2}$$

$$\leq \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2$$

$$- \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2 + \lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} - \eta^2 \frac{\lambda_D^4}{2\|\tilde{x}(t)\|_2}$$

$$\leq \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2$$

$$- \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2 + \lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} - \eta\frac{\lambda_D^4}{2\lambda_1^2}\,.$$

We further discuss three cases

1. If $\eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}} < \frac{\eta\lambda_1^2+\eta\lambda_i^2}{2-\eta\lambda_1-\eta\lambda_i}$, we have $\|\tilde{x}(t)\|_2 > \frac{\eta\lambda_1^2+\eta\lambda_i^2}{2-\eta\lambda_1-\eta\lambda_i} > \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}}$,then

$$
\begin{aligned}
\|\tilde{x}(t+1)\|_2 \leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2 \\
& - \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\frac{\eta\lambda_1^2+\eta\lambda_i^2}{2-\eta\lambda_1-\eta\lambda_i} \\
& - \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{2-\eta\lambda_1-\eta\lambda_i}{\eta\lambda_1^2+\eta\lambda_i^2} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \frac{\eta\lambda_1^2}{2-\eta\lambda_1} - \eta\frac{\lambda_D^4}{2\lambda_1^2}.
\end{aligned}
$$

The second line is because $(1-\eta\lambda_1)\|\tilde{x}(t)\|_2 + \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2}$ monotonously increase w.r.t $\|\tilde{x}(t)\|_2$ when $\|\tilde{x}(t)\|_2 > \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}}$. The last line is due to technical lemma Lemma C.62.

2. If $\eta\lambda_1^2 \geq \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}} \geq \frac{\eta\lambda_1^2+\eta\lambda_i^2}{2-\eta\lambda_1-\eta\lambda_i}$, then

$$
\begin{aligned}
\|\tilde{x}(t+1)\|_2 \leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2 \\
& - \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - 2\eta\lambda_i\sqrt{(\lambda_1^2+\frac{1}{2}\lambda_i^2)(1-\eta\lambda_1)} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \frac{\eta\lambda_1^2}{2-\eta\lambda_1} - \eta\frac{\lambda_D^4}{2\lambda_1^2}.
\end{aligned}
$$

The second line is because of AM-GM inequality. The last line is due to technical lemma Lemma C.64.

3. If $\eta\lambda_1^2 < \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}}$, we have $\|\tilde{x}(t)\|_2 < \eta\lambda_1^2 < \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}}$, then

$$
\begin{aligned}
\|\tilde{x}(t+1)\|_2 \leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2 \\
& - \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \eta\lambda_1^2 + \eta\lambda_i^2(2-\eta\lambda_1-\frac{2}{3}\eta\lambda_i) - (1-\eta\lambda_1)\eta\lambda_1^2 \\
& - \eta\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\lambda_1^2} - \eta\frac{\lambda_D^4}{2\lambda_1^2} \\
\leq & \frac{\eta\lambda_1^2}{2-\eta\lambda_1} - \eta\frac{\lambda_D^4}{2\lambda_1^2}.
\end{aligned}
$$

The second line is because $(1-\eta\lambda_1)\|\tilde{x}(t)\|_2 + \eta^2\lambda_i^2(\frac{1}{2}\lambda_i^2+\lambda_1^2)\frac{1}{\|\tilde{x}(t)\|_2}$ monotonously decrease w.r.t $\|\tilde{x}(t)\|_2$ when $\|\tilde{x}(t)\|_2 < \eta\lambda_i\sqrt{\frac{\frac{1}{2}\lambda_i^2+\lambda_1^2}{1-\eta\lambda_1}}$. The last line is due to technical lemma Lemma C.63.

$\square$

Lemma C.29. *if* $\|\tilde{x}(t)\|_2 \leq \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$*, it holds that*

$$
|\tilde{x}_1(t+1)| \geq |\tilde{x}_1(t)|.
$$

Proof. We have $|\tilde{x}_1(t+1)| = |1-\eta\lambda_1 - \eta\frac{\lambda_1^2}{\|\tilde{x}(t)\|_2}||\tilde{x}_1(t)|$.

We also have $\eta \frac{\lambda_1^2}{\|\tilde{x}(t)\|_2} > 2 - \eta\lambda_1^2$, hence $1 - \eta\lambda_1 - \eta\frac{\lambda_1^2}{\|\tilde{x}(t)\|_2} < -1$.

Hence we have that $|\tilde{x}_1(t+1)| > |\tilde{x}_1(t)|$. $\qquad\square$

Lemma C.30. *For any $t \geq 0$, if $\|\tilde{x}(t)\|_2 \leq \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$, $\tilde{x}(t) \in \cap\mathbb{I}_j$, it holds that*

$$\|\tilde{x}(t+1)\|_2 \leq \eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2.$$

Proof. The result holds as

$$\|I - \eta A - \eta\frac{A^2}{\|\tilde{x}(t)\|_2}\|_2 \leq \max_{1 \leq j \leq D}\{|1 - \eta\lambda_j - \eta\frac{\lambda_j^2}{\|\tilde{x}(t)\|}|\} = \eta\frac{\lambda_1^2}{\|\tilde{x}(t)\|} - (1 - \eta\lambda_j).$$

$\qquad\square$

Lemma C.31. *For any $t \geq 0$, if $\|\tilde{x}(t)\|_2 \leq \frac{\eta\lambda_1^2}{1-\eta\lambda_1}$, it holds that*

$\|\tilde{x}(t+1)\|_2$
$\leq(\lambda_1^2 - (1+\eta\lambda_1)\|\tilde{x}(t)\|_2)\times$
$$\sqrt{\frac{|\tilde{x}_1(t)|^2}{\|\tilde{x}(t)\|^2} + \left(\max\{\frac{|(1-\eta\lambda_2)\|\tilde{x}(t)\|_2 - \eta\lambda_2^2|}{\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2}, \frac{|(1-\eta\lambda_D)\|\tilde{x}(t)\|_2 - \eta\lambda_D^2|}{\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2}\}\right)^2(1 - \frac{|\tilde{x}_1(t)|^2}{\|\tilde{x}(t)\|^2})}.$$

Proof. We will discuss the movement along v_1 and orthogonal to v_1. First,

$$\|P^{(2:D)}\tilde{x}(t+1)\|_2 = \|(I - P^{(2:D)}\eta A - \eta\frac{P^{(2:D)}A^2}{\|\tilde{x}(t)\|_2})P^{(2:D)}\tilde{x}(t)\|_2$$

$$\leq \|P^{(2:D)} - P^{(2:D)}\eta A - \eta\frac{P^{(2:D)}A^2}{\|\tilde{x}(t)\|_2}\|_2\|P^{(2:D)}\tilde{x}(t)\|_2$$

$$\leq \max\{|1 - \eta\lambda_2 - \frac{\eta\lambda_2^2}{\|\tilde{x}(t)\|_2}|, |1 - \eta\lambda_D - \frac{\eta\lambda_D^2}{\|\tilde{x}(t)\|_2}|\}\|P^{(2:D)}\tilde{x}(t)\|_2.$$

Second

$$|\tilde{x}_1(t+1)| = (\frac{\lambda_1^2}{\|\tilde{x}(t)\|_2} - 1 + \eta\lambda_1)|\tilde{x}_1(t)|.$$

Hence we have that

$\|\tilde{x}(t+1)\|_2$
$\leq(\lambda_1^2 - (1+\eta\lambda_1)\|\tilde{x}(t)\|_2)\times$
$$\sqrt{\frac{|\tilde{x}_1(t)|^2}{\|\tilde{x}(t)\|^2} + \left(\max\{\frac{|(1-\eta\lambda_2)\|\tilde{x}(t)\|_2 - \eta\lambda_2^2|}{\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2}, \frac{|(1-\eta\lambda_D)\|\tilde{x}(t)\|_2 - \eta\lambda_D^2|}{\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2}\}\right)^2(1 - \frac{|\tilde{x}_1(t)|^2}{\|\tilde{x}(t)\|^2})}.$$

$\qquad\square$

Lemma C.32. *For $t, t' \in S, 0 \leq t \leq t'$, then $|\tilde{x}_1(t)| \leq |\tilde{x}_1(t')|$.*

Proof. For $t \in S$, by Lemma C.28, $t+1 \in S$ or $t+1 \notin S, t+2 \in S$. We will discuss by case.

1. If $t+1 \in S$, we can use Lemma C.29 to show $|\tilde{x}_1(t)| \leq |\tilde{x}_1(t+1)|$.
2. If $t+1 \notin S, t+2 \in S$, then

$$|\tilde{x}_1(t+2)| = \frac{(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t+1)\|_2)}{\|\tilde{x}(t)\|_2\|\tilde{x}(t+1)\|_2}|\tilde{x}_1(t)|.$$

As

$$(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t+1)\|_2) \geq \|\tilde{x}(t)\|_2\|\tilde{x}(t+1)\|_2$$
$$\Longleftrightarrow \eta^2\lambda_1^4 - \eta\lambda_1^2(1-\eta\lambda_1)(\|\tilde{x}(t)\|_2 + \|\tilde{x}(t+1)\|_2)$$
$$\geq (2\eta\lambda_1 - \eta^2\lambda_1^2)\|\tilde{x}(t)\|_2\|\tilde{x}(t+1)\|_2$$
$$\Longleftrightarrow \eta^2\lambda_1^4 - \eta\lambda_1^2(1-\eta\lambda_1)\|\tilde{x}(t)\|_2$$
$$\geq \left((2\eta\lambda_1 - \eta^2\lambda_1^2)\|\tilde{x}(t)\|_2 + \eta\lambda_1^2(1-\eta\lambda_1)\right)\|\tilde{x}(t+1)\|_2,$$

combining with Lemma C.30, we only need to prove,

$$\eta^2\lambda_1^4 - \eta\lambda_1^2(1-\eta\lambda_1)\|\tilde{x}(t)\|_2$$
$$\geq \left((2\eta\lambda_1 - \eta^2\lambda_1^2)\|\tilde{x}(t)\|_2 + \eta\lambda_1^2(1-\eta\lambda_1)\right)\left(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2\right).$$

Through some calculation, this is equivalent to

$$((2-\eta\lambda_1)\|\tilde{x}(t)\|_2 - \eta\lambda_1^2)((1-\eta\lambda_1)\|\tilde{x}(t)\|_2 - \eta\lambda_1^2) \geq 0.$$

which holds for $\|\tilde{x}(t)\|_2 \leq \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$.

Combining the two cases and using induction, we can get the desired result.

$\square$

Lemma C.33. $\|\tilde{x}(t)\|$ *converges to* $\frac{\eta\lambda_1^2}{2-\eta\lambda_1}$ *when* $t \to \infty$.

Proof. By Lemma C.32, $|\tilde{x}_1(t)|$ increases monotonously for $t \in S$. By Lemma C.28, S is infinite. By Lemma C.25, for sufficiently large t, $|\tilde{x}_1(t)|$ is bounded. Combining the three facts, we know $\tilde{x}_1(t)$ for $t \in S$ converges.

Formally $\forall \epsilon > 0$, there exists $T_\epsilon > 0$ such that $\forall t, t' \in S, t' > t > T_\epsilon, \frac{\|\tilde{x}_1(t')\|_2}{\|\tilde{x}_1(t)\|_2} < 1 + \epsilon$.

Then by Lemma C.28, $\forall t \in S, t+1 \in S$ or $t+2 \in S$, we will discuss by case. For $t \geq T_\epsilon$,

1. If $t+1 \in S$, then

$$1 + \epsilon \geq \frac{\|\tilde{x}_1(t+1)\|_2}{\|\tilde{x}_1(t)\|_2} = \frac{\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2}{\|\tilde{x}(t)\|_2}.$$

2. If $t+1 \notin S$ and $t+2 \in S$, then

$$1 + \epsilon \geq \frac{\|\tilde{x}_1(t+2)\|_2}{\|\tilde{x}_1(t)\|_2}$$
$$= \frac{(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t+1)\|_2)}{\|\tilde{x}(t)\|_2\|\tilde{x}(t+1)\|_2}$$
$$\geq \frac{(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)\left(\eta\lambda_1^2 - (1-\eta\lambda_1)\left(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2\right)\right)}{\|\tilde{x}(t)\|_2\left(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2\right)}$$
$$= \frac{\eta\lambda_1^2 - (1-\eta\lambda_1)\left(\eta\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2\right)}{\|\tilde{x}(t)\|_2}.$$

Here in the last inequality, we apply Lemma C.30.

Concluding, $\|\tilde{x}(t)\|_2 \geq \min\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1^2+\epsilon}, \frac{\eta^2\lambda_1^3}{(2-\lambda_1\eta)\lambda_1\eta+\epsilon}\right), \forall t > T_\epsilon, t \in S$. As $\forall t \notin S, t > T_\epsilon$, we have $\|\tilde{x}(t)\|_2 \geq \frac{\eta\lambda_1^2}{2-\eta\lambda_1^2}$. Hence we have $\forall t > T_\epsilon, \|\tilde{x}(t)\|_2 \geq \min\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1^2+\epsilon}, \frac{\eta^2\lambda_1^3}{(2-\lambda_1\eta)\lambda_1\eta+\epsilon}\right)$.

Further by Lemma C.30, $\forall t > T_\epsilon + 1, \|\tilde{x}(t)\|_2 \leq \eta\lambda_1^2 - (1-\eta\lambda_1)\min\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1^2+\epsilon}, \frac{\eta^2\lambda_1^3}{(2-\lambda_1\eta)\lambda_1\eta+\epsilon}\right)$.

Combining both bound, we have $\lim_{t\to\infty}\|\tilde{x}(t)\|_2 = \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$. $\square$

Lemma C.34. $\|\tilde{x}_1(t)\|_2$ *converges to* $\frac{\eta\lambda_1^2}{2-\eta\lambda_1}$, *when* $t \to \infty$.

Proof. Notice that

$$\|P^{(2:D)}\tilde{x}(t+1)\|_2 \le \max\left(|1 - \eta\lambda_2 - \eta\frac{\lambda_2^2}{\|\tilde{x}(t)\|_2}|, |1 - \eta\lambda_D - \eta\frac{\lambda_D^2}{\|\tilde{x}(t)\|_2}|\right)\|P^{(2:D)}\tilde{x}(t)\|_2.$$

When $\|\tilde{x}(t)\|_2 > \frac{\eta\lambda_2^2}{2-\eta\lambda_2-\delta}$,

$$-1 + \delta \le 1 - \eta\lambda_2 - \eta\frac{\lambda_2^2}{\|\tilde{x}(t)\|_2} \le 1 - \eta\lambda_D - \eta\frac{\lambda_D^2}{\|\tilde{x}(t)\|_2} \le 1 - \eta\lambda_D$$

$$\|P^{(2:D)}\tilde{x}(t+1)\|_2 \le \max(1 - \eta\lambda_D, 1 - \delta)\|P^{(2:D)}\tilde{x}(t)\|_2$$

Hence for sufficiently large t, $\|P^{(2:D)}\tilde{x}(t)\|_2$ shrinks exponentially, showing that $\lim_{t\to\infty}\|\tilde{x}_1(t)\|_2 = \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$. □

C.6 FULL-BATCH SAM ON GENERAL LOSS (PROOF OF THEOREM 4.9)

The goal of this section is to prove the following theorem.

Theorem 4.9 (Main, Theorems C.35 and C.36 stated informally). *Let $\{x(t)\}$ be the iterates of SAM (3) with $x(0) = x_{init} \in U$, then under Assumptions 3.1 and 4.8, for all η, ρ such that $\eta\ln(1/\rho)$ and ρ/η are sufficiently small, the dynamics of SAM can be split into two phases:*

- *Phase I: SAM follows Gradient Flow with respect to L until entering an $O(\eta\rho)$ neighborhood of the manifold Γ in $\tilde{O}(\frac{1}{\eta})$ steps;*
- *Phase II: SAM tracks the solution X of (6), the Riemannian Gradient Flow with respect to $\lambda_1(\nabla^2 L(\cdot))$ in the $O(\eta\rho)$ neighborhood in the sense that $\max_{0\le T\le T_3}\|\Phi(x(\lceil T/(\eta\rho^2)\rceil)) - X(T)\| = O((\eta + \rho)\log(1/\eta\rho))$. Moreover, the angle between $\nabla L(x(t))$ and $v_1(\nabla^2 L(x(t)))$ is at most $O(\rho)$.*

To prove the theorem, we will separate the dynamic of SAM on general loss L to two phases.

Define

$$R_j(x) = \sqrt{\sum_{i=j}^{M}\lambda_i^2(x)\langle v_i(x), x - \Phi(x)\rangle^2} - \eta\rho\lambda_j(x), j \in [M], x \in U,$$

which is the length projection of $x - \Phi(x)$ on button$-k$ non-zero eigenspace of $\nabla^2 L(\Phi(x))$. We will provide a fine-grained convergence bound on $R_j(x)$.

Theorem C.35 (Phase I). *Let $\{x(t)\}$ be the iterates defined by SAM (Equation (3)) and $x(t) = x_{init} \in U$, then under Assumption 3.1 there exists a constant T_1, such that for any $T_1' > T_1$, it holds for all η, ρ such that $(\eta + \rho)\ln(1/\eta\rho)$ is sufficiently small, we have*

$$\max_{T_1\ln(1/\eta\rho)\le\eta t\le T_1'\ln(1/\eta\rho)}\max_{j\in[M]}R_j(x) = O(\eta\rho^2)$$

$$\max_{T_1\ln(1/\eta\rho)\le\eta t\le T_1'\ln(1/\eta\rho)}\|\Phi(x(t)) - \Phi(x_{init})\| = O((\eta + \rho)\ln(1/\eta\rho))$$

Theorem C.35 implies SAM will converge to an $O(\eta\rho)$ neighbor of Γ. Notice in the time frame defined by Theorem C.35, $x(t)$ effectively operates at a local regime around $\Phi(\lceil T_1\ln(1/\eta\rho)/\eta\rceil)$, this allows us to approximate L with the quadratic Taylor expansion of L at $\Phi(\lceil T_1\ln(1/\eta\rho)/\eta\rceil)$ and give us the following theorem.

Theorem C.36 (Phase II). *Let $\{x(t)\}$ be the iterates defined by SAM (Equation (3)) under Assumptions 3.1 and 4.8, for all η, ρ such that $\eta\ln(1/\rho)$ and ρ/η is sufficiently small, further assuming that (1) $\max_j R_j(x(0)) = O(\eta\rho^2)$, (2) $\|\Phi(x(0)) - \Phi(x_{init})\| = O((\eta + \rho)\ln(1/\eta\rho))$ and (3) $|\langle x(0) - \Phi(x(0)), v_1(x(0))\rangle| \ge \Omega(\rho^2)$, the iterates $x(t)$ tracks the solution X of Equation (6). Quantitatively for $t = \lceil T_3/\eta\rho^2\rceil$, we have that*

$$\|\Phi(x(t)) - X(\eta\rho^2 t)\| = O(\eta\ln(1/\rho)).$$

Moreover the angle between $\nabla L(x(t))$ and the top eigenspace of $\nabla^2 L(\Phi(x(t)))$ is at most $O(\rho)$. Quantitatively,

$$|\langle x(t) - \Phi(x(t)), v_1(x(t))\rangle| = \Theta(\eta\rho)\,.$$
$$\max_{j\in[2:M]} |\langle x(t) - \Phi(x(t)), v_j(x(t))\rangle| = O(\eta\rho^2)\,.$$

In this section we will define K as $\{X(t) \mid 0 \leq t \leq T_3\}$ where X is the solution of Equation (6). To simplify our proof, we assume WLOG $L(x) = 0$ for $x \in \Gamma$.

C.6.1   PHASE I (PROOF OF THEOREM C.35)

The goal of this section is to prove Theorem C.35.

Proof. The proof consists of three major steps.

1. *Tracking Gradient Flow* Lemma C.37 shows the existence of step $t_{\mathrm{GF}} = O(1/\eta)$ such that $x(t_{\mathrm{GF}})$ is in a subset of K^h and $\Phi(x(t_{\mathrm{GF}}))$ is $O(\eta + \rho)$ close to $\Phi(x_{\mathrm{init}})$.
2. *Decreasing Loss* Lemma C.39 shows the existence of step $t_{\mathrm{DEC}} = O(\ln(1/\rho)/\eta)$ such that $x(t_{\mathrm{DEC}})$ is in $O(\rho)$ neighbor of Γ and $\Phi(x(t_{\mathrm{DEC}}))$ is $O((\eta + \rho)\ln(1/\rho))$ close to $\Phi(x_{\mathrm{init}})$.
3. *Entering Invariant Set* Lemmas C.44 and C.46 shows the existence of step $t_{\mathrm{INV}} = O(\ln(1/\rho\eta)/\eta)$ such that for any t satisfying $t_{\mathrm{INV}} \leq t \leq t_{\mathrm{INV}} + \Theta(\ln(1/\eta)/\eta)$, we have that $x(t) \in \cap_{k\in[M]}\mathbb{I}_k$ and $\Phi(x(t))$ is $O((\eta + \rho)\ln(1/\eta\rho))$ close to $\Phi(x_{\mathrm{init}})$.

$\square$

Tracking Gradient Flow   Lemma C.37 shows that the iterates $x(t)$ tracks gradient flow to an $O(1)$ neighbor of Γ.

Lemma C.37. *Under condition of Theorem C.35, there exists $t_{\mathrm{GF}} = O(1/\eta)$, such that the iterate $x(t_{\mathrm{GF}})$ is $O(1)$ close to the manifold Γ and $\Phi(x(t_{\mathrm{GF}}))$ is $O(\eta+\rho)$ is close to $\Phi(x_{init})$. Quantitatively,*

$$L(x(t_{\mathrm{GF}})) \leq \frac{\mu h^2}{32}$$
$$\|x(t_{\mathrm{GF}}) - \Phi(x(t_{\mathrm{GF}}))\| \leq h/4\,,$$
$$\|\Phi(x(t_{\mathrm{GF}})) - \Phi(x_{init})\| = O(\eta + \rho)\,.$$

Proof. Choose $C = \frac{1}{4}\sqrt{\frac{\mu}{\zeta}}$.

There exists $T > 0$, such that

$$\|\phi(x_{\mathrm{init}}, T) - \Phi(x_{\mathrm{init}})\|_2 \leq Ch/2\,.$$

Consider

$$x(t+1) = x(t) - \eta\nabla L(x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|})$$
$$= x(t) - \eta\nabla L(x(t)) + O(\eta\rho)\,.$$

By standard approximation theory (e.g., Section I.10 in Hairer et al. (2008)), for sufficiently small η and ρ, the iterates $x(t)$ tracks gradient flow $\phi(x_{\mathrm{init}}, T)$ in $O(1/\eta)$ steps. Quantitatively for $t_{\mathrm{GF}} = \lceil\frac{T}{\eta}\rceil$, we have that

$$\|x(t_{\mathrm{GF}}) - \phi(x_{\mathrm{init}}, T)\|_2 \leq O(\eta + \rho)\,.$$

This implies $x(t_{\mathrm{GF}}) \in K^h$, hence by Taylor Expansion on Φ,

$$\|\Phi(x(t_{\mathrm{GF}})) - \Phi(x_{\mathrm{init}})\|_2 = \|\Phi(x(t_{\mathrm{GF}})) - \Phi(\phi(x_{\mathrm{init}}, T))\|_2$$
$$\leq O(\|x(t_{\mathrm{GF}}) - \phi(x_{\mathrm{init}}, T)\|_2)$$
$$\leq O(\eta + \rho)\,.$$

This implies

$$\begin{aligned} \|x(t_{\text{GF}}) - \Phi(x(t_{\text{GF}}))\|_2 \leq & \|x(t_{\text{GF}}) - \phi(x_{\text{init}}, T_0)\|_2 + \|\phi(x_{\text{init}}, T_0) - \Phi(x_{\text{init}})\|_2 \\ & + \|\Phi(x_{\text{init}}) - \Phi(x(t_{\text{GF}}))\|_2 \\ \leq & Ch/2 + O(\eta + \rho) \leq Ch \leq h/4 \,. \end{aligned}$$

By Taylor Expansion,

$$L(x(t_{\text{GF}})) \leq \zeta \|x(t_{\text{GF}}) - \Phi(x(t_{\text{GF}}))\|_2^2/2 \leq \frac{\mu h^2}{32}$$

$\square$

Decreasing Loss Lemma C.39 shows that the iterates $x(t)$ converges to an $O(\rho)$ neighbor of Γ in $O(\ln(1/\rho)/\eta)$ steps.

Lemma C.38. *Under condition of Theorem C.35, if $x(t) \in K^h$ and $\|\nabla L(x(t))\| \geq 4\zeta\rho$, then we have that $L(x(t+1))$ decreases with respect to $L(x(t))$, quantitatively, we have that*

$$L(x(t+1)) \leq L(x(t))(1 - \eta\mu/8) \,.$$

Moreover the movement of the projection of the iterates on the manifold is bounded, quantitatively, we have that

$$\|\Phi(x(t+1)) - \Phi(x(t))\| \leq O(\eta^2) \,.$$

Proof. As $x(t) \in K^h$ and L is μ-PL in K^h, we have $L(x(t)) \geq 0$.

As $x(t) \in K^h$, by Lemma C.12 and Taylor Expansion, we have $\|\overline{x(t)x(t+1)}\| = O(\eta)$. hence for sufficiently small η, $\overline{x(t)x(t+1)} \subset K^r$. Using similar argument, the segment from $x(t)$ to $x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}$ is in K^r.

Then by Taylor Expansion on L,

$$\begin{aligned} L(x(t+1)) = & L\left(x(t) - \eta\nabla L\left(x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}\right)\right) \\ \leq & L(x(t)) - \eta\left\langle\nabla L(x(t)), \nabla L\left(x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}\right)\right\rangle \\ & + \frac{\zeta\eta^2\|\nabla L\left(x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}\right)\|^2}{2} \,. \end{aligned} \tag{26}$$

By Taylor Expansion on ∇L, we have that

$$\|\nabla L\left(x(t) + \rho\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}\right) - \nabla L(x(t))\| \leq \zeta\rho \,.$$

After plugging in Equation (26), we have that

$$L(x(t+1)) \leq L(x(t)) - \eta\|\nabla L(x(t))\|^2 + \eta\zeta\rho\|\nabla L(x(t))\| + \zeta\eta^2\|\nabla L(x(t))\|^2 + \zeta^3\eta^2\rho^2 \,. \tag{27}$$

As $\|\nabla L(x(t))\| \geq 4\zeta\rho$, we have that the following term is bounded.

$$\zeta\eta^2\|\nabla L(x(t))\|^2 \leq \frac{1}{2}\eta\|\nabla L(x(t))\|^2 \,,$$

$$\eta\zeta\rho\|\nabla L(x(t))\| \leq \frac{1}{4}\eta\|\nabla L(x(t))\|^2 \,,$$

$$\zeta^3\eta^2\rho \leq \zeta^2\eta\rho^2 \leq \frac{1}{16}\eta\|\nabla L(x(t))\|^2 \,.$$

After plugging in Equation (27), by Lemma C.8,

$$L(x(t+1)) \leq L(x(t)) - \frac{1}{16}\eta\|\nabla L\left(x(t)\right)\|^2$$
$$\leq L(x(t))(1 - \eta\mu/8)\,.$$

As $x(t) \in K^h$, by Taylor Expansion, we have

$$\|\nabla L\left(x(t)\right)\| \leq \zeta h\,.$$

Hence by Lemma C.12 and Taylor Expansion,

$$\|\Phi(x(t+1)) - \Phi(x(t))\| \leq \xi\eta\rho\|\nabla L\left(x\right)\|_2 + \nu\eta\rho^2 + \xi\eta^2\|\nabla L\left(x\right)\|_2^2 + \xi\zeta^2\eta^2\rho^2$$
$$= O(\eta^2)\,.$$

$\square$

Lemma C.39. *Under condition of Theorem C.35, assuming there exists t_{GF} such that $L(x(t_{\mathrm{GF}})) \leq \frac{\mu h^2}{32}$ and $x(t_{\mathrm{GF}}) \in K^{h/4}$, then there exists $t_{\mathrm{DEC}} = t_{\mathrm{GF}} + O(\ln(1/\rho)/\eta)$, such that $x(t_{\mathrm{DEC}})$ is in $O(\rho)$ neighbor of Γ, quantitatively, we have that*

$$\|\nabla L(x(t_{\mathrm{DEC}}))\|_2 \leq 4\zeta\rho\,.$$

Moreover the movement of the projection of $\Phi(x(\cdot))$ on the manifold is bounded,

$$\|\Phi(x(t_{\mathrm{GF}})) - \Phi(x(t_{\mathrm{DEC}}))\|_2 = O(\eta\ln(1/\rho))\,.$$

Proof. Choose t_{DEC} as the minimal $t \geq t_{\mathrm{GF}}$ such that $\|\nabla L(x(t_{\mathrm{DEC}}))\|_2 \leq 4\zeta\rho$. Define $C = \lceil\ln_{1-\frac{\eta\mu}{8}}(64\rho^2/h^2)\rceil = O(\ln(1/\rho)/\eta)$.

We will first perform an induction on $t \leq \min\{t_{\mathrm{DEC}}, t_{\mathrm{GF}} + C\} = t_{\mathrm{GF}} + O(\ln(1/\rho)/\eta)$ to show that

$$L(x(t)) \leq (1 - \eta\mu/8)^{t-t_{\mathrm{GF}}}L(x(t_{\mathrm{GF}}))$$
$$\|\Phi(x(t)) - \Phi(x(t_{\mathrm{GF}}))\| = O(\eta^2(t - t_{\mathrm{GF}}))$$

For $t = t_{\mathrm{GF}}$, the result holds trivially. Suppose the induction hypothesis holds for t. Then by C.7 and Taylor Expansion,

$$\|\Phi(x(t)) - x(t)\| \leq \sqrt{\frac{2L(x(t_{\mathrm{GF}}))}{\mu}}$$
$$\leq h/4\,.$$

Then we have that

$$\mathrm{dist}(K, x(t)) \leq \mathrm{dist}(K, x(t_{\mathrm{GF}})) + \|x(t_{\mathrm{GF}}) - \Phi(x(t_{\mathrm{GF}}))\|_2$$
$$+ \|\Phi(x(t_{\mathrm{GF}})) - \Phi(x(t))\| + \|\Phi(x(t)) - x(t)\|$$
$$\leq 3h/4 + O(\eta^2(t - t_{\mathrm{GF}})) = 3h/4 + O(\eta\ln(1/\rho)) \leq h\,.$$

That is $x(t) \in K^h$. Then as $t \leq t_{\mathrm{DEC}}$, $\|\nabla L(x(t))\|_2 \geq 4\zeta\rho$. Then by Lemma C.38, we have that

$$L(x(t+1)) \leq (1 - \eta\mu/8)L(x(t)) \leq (1 - \eta\mu/8)^{t+1-t_{\mathrm{GF}}}L(x(t_{\mathrm{GF}}))\,,$$
$$\|\Phi(x(t+1)) - \Phi(x(t_{\mathrm{GF}}))\| \leq \|\Phi(x(t+1)) - \Phi(x(t))\| + \|\Phi(x(t)) - \Phi(x(t_{\mathrm{GF}}))\|$$
$$\leq O(\eta^2(t - t_{\mathrm{GF}}))\,.$$

The induction is complete.

Now if $t_{\mathrm{DEC}} \geq t_{\mathrm{GF}} + C = t_{\mathrm{GF}} + O(\ln(1/\rho)/\eta)$, As the result of the induction, we have that

$$L(x(t_{\mathrm{GF}} + C)) \leq (1 - \frac{\eta\mu}{8})^C L(t_{\mathrm{GF}})$$
$$\leq \frac{64\rho^2}{h^2}L(t_{\mathrm{GF}})$$
$$\leq 8\rho^2\mu\,.$$

By Lemma C.8, we have that

$$\|\nabla L(x(t_{\mathrm{GF}} + C))\|_2 \le \zeta \sqrt{\frac{2L(x(t_{\mathrm{GF}} + C))}{\mu}} = 4\zeta\rho \,.$$

This is a contradiction.

Hence we have that $t_{\mathrm{DEC}} \le t_{\mathrm{GF}} + C = t_{\mathrm{GF}} + O(\ln(1/\rho)/\eta)$. By induction, we have that

$$\|\Phi(x(t_{\mathrm{DEC}})) - \Phi(x(t_{\mathrm{GF}}))\| = O(\eta^2(t_{\mathrm{DEC}} - t_{\mathrm{GF}})) = O(\eta \ln(1/\rho)) \,.$$

$\square$

Entering Invariant Set   We will introduce some notations that is required for the proof in this and following subsection.

Define

$$\hat{x} = x - \Phi(x) \,,$$
$$A(x) = \nabla^2 L\left(\Phi(x)\right) \,,$$
$$\tilde{x} = A(x)\hat{x} \,,$$
$$\tilde{x}_j = \langle \tilde{x}, v_j(x) \rangle \,,$$
$$P^{(j:D)}(x) = \sum_{i=j}^{M} v_i(x)v_i^T(x) \,.$$

Note $\tilde{x} \approx \nabla L(x)$ for x near the manifold Γ. We also use $\tilde{x}(t)$, $A(t)$ and $\hat{x}(t)$ to denote $\tilde{x(t)}$, $A(x(t))$ and $\hat{x(t)}$.

Recall the original definition of $R_j(x)$ is

$$R_j(x) = \sqrt{\sum_{i=j}^{M} \lambda_i^2(x)\langle v_i(x), x - \Phi(x)\rangle^2 - \eta\rho\lambda_j^2(x)} \,,$$

Based on the above notions, we can rephrase the notion R as

$$R_j(x) = \|P^{(j:D)}(x)\tilde{x}\|^2 - \eta\rho\lambda_j^2(x) \,.$$

We additionally define the approximate invariant set $\mathbb{I}_j$ as

$$\mathbb{I}_j = \{R_j(x) \le O(\eta\rho^2)\} \,.$$

Lemma C.40. *Under condition of Theorem C.35, assuming t satisfy that $x(t) \in K^h$, then we have that*

$$\frac{\mu}{2}\|x(t) - \Phi(x(t))\| \le \|\tilde{x}(t)\| \le \zeta\|x(t) - \Phi(x(t))\|$$

Proof. First by Lemma C.10, $\Phi(x(t)) \in K^r$, hence

$$\|\tilde{x}(t)\| = \|\nabla^2 L(\Phi(x(t)))(x(t) - \Phi(x(t)))\|$$
$$\le \zeta\|x(t) - \Phi(x(t))\| \,.$$

Also

$$\|\tilde{x}(t)\| = \|\nabla^2 L(\Phi(x(t)))(x(t) - \Phi(x(t)))\|$$
$$\ge \mu\|P^\perp_{\Phi(x(t)),\Gamma}(x(t) - \Phi(x(t)))\| \,.$$

By Lemma C.10 and Lemma C.6, we have

$$\|x(t) - \Phi(x(t))\| \le \|P^\perp_{\Phi(x(t)),\Gamma}(x(t) - \Phi(x(t)))\| + \|P_{\Phi(x(t)),\Gamma}(x(t) - \Phi(x(t)))\|$$
$$\le \frac{\zeta\nu}{4\mu^2}\|x(t) - \Phi(x(t))\|^2 + \frac{1}{\mu}\|\tilde{x}(t)\|$$
$$\le \frac{1}{2}\|x(t) - \Phi(x(t))\| + \frac{1}{\mu}\|\tilde{x}(t)\|.$$

Hence $\|x(t) - \Phi(x(t))\| \le \frac{2}{\mu}\|\tilde{x}(t)\|$. $\square$

Lemma C.41. *Under condition of Theorem C.35, assuming t satisfy that $x(t) \in K^h$ and $\|\tilde{x}(t)\|_2 = O(\rho)$, then we have that*

$$\|\Phi(x(t+1)) - \Phi(x(t))\| = O(\eta\rho^2).$$

Proof. By Lemma C.40, we have $\|x(t) - \Phi(x(t))\| = O(\rho)$. By Lemma C.12, we have that

$$\|\Phi(x(t+1)) - \Phi(x(t))\| \leq \zeta\xi\eta\rho\|x - \Phi(x)\|_2 + \zeta^2\xi\eta^2\|x - \Phi(x)\|_2^2 + \nu\eta\rho^2 + \xi\zeta^2\eta^2\rho^2$$

$$\leq O(\eta\rho^2)$$

$\square$

Lemma C.42. *Under condition of Theorem C.35, assuming t satisfy $x(t) \in K^{h/2}$ and $\|x(t) - \Phi(x(t))\|_2 = O(\rho)$, define x' as $x'(t) = x(t)$ and*

$$x'(t+1) = x'(t) - \eta\nabla^2 L(\Phi(x(t)))(x'(t) - \Phi(x))$$

$$-\eta\rho\nabla^2 L(\Phi(x(t)))\frac{\nabla^2 L(\Phi(x(t)))(x'(t) - \Phi(x))}{\|\nabla^2 L(\Phi(x(t)))(x'(t) - \Phi(x))\|_2}.$$

Then

$$\|x'(t+1) - x(t+1)\|_2 = O(\eta\rho^2)$$

and further if $\|x(t+1) - \Phi(x(t+1))\|_2 = \Omega(\eta\rho)$, then

$$\|x'(t+2) - x(t+2)\|_2 = O(\eta\rho^2).$$

Proof. For simplicity, we will sometime omit the t in $x(t)$.

By $\|x(t) - \Phi(x(t))\| = O(\rho)$, $x(t) \in K^{h/2}$, and Lemma C.12, we have that $\|x(t+1) - x(t)\| = O(\eta\rho)$ and hence $x(t+1) \in K^{3h/4}$. This also implies $\|x(t+1) - \Phi(x(t+1))\|_2 = O(\rho)$. Similarly we have $x(t+2) \in K^{3h/4}$.

For $k \in \{1, 2\}$, by Taylor Expansion,

$$x(t+k+1) = x(t+k) - \eta\nabla L(x(t+k)) - \eta\rho\nabla^2 L(x(t+k))\frac{\nabla L(x(t+k))}{\|\nabla L(x(t+k))\|} + O(\eta\rho^2)$$

$$= x(t+k) - \eta\nabla^2 L(\Phi(x(t+k)))(x(t+k) - \Phi(x(t+k))) + O(\eta\rho^2)$$

$$- \eta\rho\nabla^2 L(\Phi(x(t+k)))\frac{\nabla L(x(t+k))}{\|\nabla L(x(t+k))\|} + O(\eta\rho^2)$$

$$= x(t+k) - \eta\nabla^2 L(\Phi(x(t+k)))(x(t+k) - \Phi(x(t+k)))$$

$$- \eta\rho\nabla^2 L(\Phi(x(t+k)))\frac{\nabla L(x(t+k))}{\|\nabla L(x(t+k))\|} + O(\eta\rho^2).$$

Now by Lemmas C.40 and C.41, $\|\Phi(x(t+k)) - \Phi(x(t))\|_2 = O(\eta\rho^2)$,

$$x(t+k+1) = x(t+k) - \eta\nabla^2 L(\Phi(x))(x(t+k) - \Phi(x(t)))$$

$$- \eta\rho\nabla^2 L(\Phi(x))\frac{\nabla L(x(t+k))}{\|\nabla L(x(t+k))\|} + O(\eta\rho^2). \tag{28}$$

Now we first prove the first claim, we have for $k = 0$, $\|x(t+k) - x'(t+k)\|_2 = 0$, by Lemma C.10 and eq. (28),

$$x(t+1) = x(t) - \eta\nabla^2 L(\Phi(x))(x(t) - \Phi(x)) - \eta\rho\frac{\nabla^2 L(\Phi(x))(x(t) - \Phi(x))}{\|\nabla^2 L(\Phi(x))(x(t) - \Phi(x))\|_2} + O(\eta\rho^2)$$

$$= x'(t+1) + O(\eta\rho^2).$$

The second claim is slightly more complex. By the first claim and Lemma C.10, we have that

$$\frac{\nabla L\left(x(t+1)\right)}{\|\nabla L\left(x(t+1)\right)\|} = \frac{\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))}{\|\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))\|_2} + O(\|x(t+1) - \Phi(x(t+1))\|_2). \tag{29}$$

We first show $\|\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))\|_2$ is of order $\|x(t+1) - \Phi(x(t+1))\|_2 = \Omega(\rho^2)$ to show that the normalized gradient term is stable with respect to small perturbation,

$$\|\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))\|_2$$
$$\geq \|P_{\Phi(x(t+1)),\Gamma} \nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))\|_2$$
$$\geq \|\nabla^2 L(\Phi(x(t+1))) P_{\Phi(x(t+1)),\Gamma}(x(t+1) - \Phi(x(t+1)))\|_2$$
$$\geq \mu \|P_{\Phi(x(t+1)),\Gamma}(x(t+1) - \Phi(x(t+1)))\|_2$$
$$\geq \mu(\|(x(t+1) - \Phi(x(t+1)))\|_2 - \|P^{\perp}_{\Phi(x(t+1)),\Gamma}(x(t+1) - \Phi(x(t+1)))\|_2)$$
$$\geq \mu(\|(x(t+1) - \Phi(x(t+1)))\|_2 - \frac{\nu\zeta}{4\mu^2}\|x(t+1) - \Phi(x(t+1))\|_2^2)$$
$$\geq \frac{\mu}{2}\|(x(t+1) - \Phi(x(t+1)))\|_2 = \Omega(\eta\rho).$$

Based on Lemma C.12, we have

$$\Phi(x(t+1)) - \Phi(x(t)) = O(\eta\rho^2).$$

We further have by the first claim and Lemma C.41,

$$\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1))) - \nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x(t)))$$
$$= \nabla^2 L(\Phi(x))(x(t+1) - \Phi(x(t+1))) - \nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x(t)))$$
$$\quad + O(\|x(t+1) - \Phi(x(t+1))\|_2 \|\Phi(x(t+1)) - \Phi(x)\|_2)$$
$$= \nabla^2 L(\Phi(x))(x(t+1) - \Phi(x(t+1))) - \nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x(t))) + O(\eta\rho^3)$$
$$= \nabla^2 L(\Phi(x))(x(t+1) - x'(t+1)) + \nabla^2 L(\Phi(x))(\Phi(x(t+1)) - \Phi(x(t))) + O(\eta\rho^3)$$
$$= O(\eta\rho^2)$$

This implies

$$\frac{\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))}{\|\nabla^2 L(\Phi(x(t+1)))(x(t+1) - \Phi(x(t+1)))\|_2} = \frac{\nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x))}{\|\nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x))\|_2} + O(\rho)$$

Combining with Equation (29), we have

$$\frac{\nabla L\left(x(t+1)\right)}{\|\nabla L\left(x(t+1)\right)\|} = \frac{\nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x))}{\|\nabla^2 L(\Phi(x))(x'(t+1) - \Phi(x))\|_2} + O(\rho)$$

By the above approximation and Equation (28),

$$x(t+2) = x'(t+2) + O(\eta\rho^2).$$

$\square$

Lemma C.43. *Under condition of Theorem C.35, assuming assuming t satisfy that $x(t) \in K^{3h/4}$ and $\|\tilde{x}(t)\|_2 = O(\rho)$, then we have that*

$$\|\tilde{x}(t+1) - \tilde{x}(t) + \eta A(t)\tilde{x}(t) + \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|_2 = O(\eta\rho^2).$$

Proof. By Lemma C.42, we know

$$\|x(t+1) - x(t) + \eta\tilde{x}(t) + \eta\rho A(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\| \leq O(\eta\rho^2).$$

This implies

$$\|A(t)(x(t+1) - \Phi(x(t))) - \tilde{x}(t) + \eta A(t)\tilde{x}(t) + \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|$$

$$=\|A(t)(x(t+1) - x(t) + \eta\tilde{x}(t) + \eta\rho A(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|})\|$$

$$\leq\zeta\|x(t+1) - x(t) + \eta\tilde{x}(t) + \eta\rho A(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\| = O(\eta\rho^2). \tag{30}$$

We also have

$$\tilde{x}(t+1) - A(t)(x(t+1) - \Phi(x(t)))$$
$$=(A(t+1) - A(t))(x(t+1) - \Phi(x(t+1))) - A(t)(\Phi(x(t)) - \Phi(x(t+1)))$$
$$=O(\eta\rho^2).$$

Plugging in Equation (30), we have that

$$\|\tilde{x}(t+1) - \tilde{x}(t) + \eta A(t)\tilde{x}(t) + \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|_2 = O(\eta\rho^2).$$

$\square$

Lemma C.44. *Under condition of Theorem C.35, assuming there exists t_{DEC} such that $x(t_{\mathrm{DEC}}) \in K^{h/2}$ and $\|\nabla L(x(t_{\mathrm{DEC}}))\| \leq 4\zeta\rho$, then there exists $t_{\mathrm{DEC2}} = t_{\mathrm{DEC}} + O(\ln(1/\eta)/\eta)$, such that $x(t_{\mathrm{DEC2}})$ is in $\mathbb{I}_1 \cap K^{3h/4}$.*

Furthermore, for any t satisfying $t_{\mathrm{DEC2}} \leq t \leq t_{\mathrm{DEC2}} + \Theta(\ln(1/\eta)/\eta)$, we have that $x(t) \in \mathbb{I}_1 \cap K^{3h/4}$ and $\|\Phi(x(t)) - \Phi(x(t_{\mathrm{DEC}}))\| = O(\rho^2 \ln(1/\eta))$.

Proof. For simplicity, denote $C = \lceil \ln_{1-\eta\mu} \frac{\eta\mu^3}{4\zeta^2} \rceil + \Theta(\ln(1/\rho)/\eta) = O(\ln(1/\eta)/\eta)$. Here the quantity $\Theta(\ln(1/\rho)/\eta)$ is the same quantity in the statement of the lemma.

We will prove the induction hypothesis for $t_{\mathrm{DEC}} \leq t \leq t_{\mathrm{DEC}} + 2C$,

$$\begin{cases} \|\tilde{x}(t-1)\| \geq \eta\rho\lambda_1^2(t), t > t_{\mathrm{DEC}} \Rightarrow \|\tilde{x}(t)\| \leq (1-\eta\mu)\|\tilde{x}(t-1)\|, \\ \|\tilde{x}(t-1)\| \leq \eta\rho\lambda_1^2(t-1), t > t_{\mathrm{DEC}} \Rightarrow \|\tilde{x}(t)\| \leq \eta\rho\lambda_1^2(t) + O(\eta\rho^2), \\ \qquad\qquad\qquad \|\Phi(x(t)) - \Phi(x(t_{\mathrm{DEC}}))\| \leq O(\eta\rho^2(t - t_{\mathrm{DEC}})), \\ \qquad\qquad\qquad\qquad\qquad x(t) \in K^{3h/4}. \end{cases}$$

The induction hypothesis holds trivially for $t = t_{\mathrm{DEC}}$.

Assume the induction hypothesis holds for $t' \leq t$. By Lemmas C.7 and C.40, $\|\tilde{x}(t_{\mathrm{DEC}})\|_2 \leq \zeta\|x(t_{\mathrm{DEC}}) - \Phi(x(t_{\mathrm{DEC}}))\| \leq \frac{\zeta}{\mu}\|\nabla L(x(t_{\mathrm{DEC}}))\| \leq \frac{4\zeta^2}{\mu}\rho$. Combining with the induction hypothesis, we have $\|\tilde{x}(t)\| \leq \frac{4\zeta^2}{\mu}\rho$.

By $x(t) \in K^{3h/4}$ and Lemma C.41, we have that

$$\|\Phi(x(t+1)) - \Phi(x(t))\| \leq O(\eta\rho^2).$$

Hence we have that

$$\|\Phi(x(t+1)) - \Phi(x(t_{\mathrm{DEC}}))\| \leq \|\Phi(x(t+1)) - \Phi(x(t))\| + \|\Phi(x(t)) - \Phi(x(t_{\mathrm{DEC}}))\|$$
$$\leq O(\eta\rho^2(t+1 - t_{\mathrm{DEC}})). \tag{31}$$

This proves the third statement of the induction hypothesis.

By $\|\tilde{x}(t)\| = O(\rho)$ and Lemma C.43, we have that

$$\|\tilde{x}(t+1) - \tilde{x}(t) + \eta A(t)\tilde{x}(t) + \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|_2 = O(\eta\rho^2).$$

Analogous to the proof of Lemmas C.24 and C.25, we have

1. If $\|\tilde{x}(t)\| > \eta\rho\lambda_1^2(t)$, we would have

$$\|\tilde{x}(t) - \eta A(t)\tilde{x}(t) - \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|$$

$$\leq \|\tilde{x}(t)\|\|I - \eta A(t) - \eta\rho A^2(t)\frac{1}{\|\tilde{x}(t)\|}\|$$

$$\leq \|\tilde{x}(t)\| \max\{\eta\lambda_1, 1 - \eta\lambda_D - \eta\rho\lambda_D^2\frac{1}{\|x(t)\|}\}$$

$$\leq \max\{(1 - \eta\lambda_D)\|\tilde{x}(t)\| - \eta\rho\lambda_D^2, \eta\lambda_1\|\tilde{x}(t)\|\}$$

$$\leq \max\{(1 - \eta\mu)\|\tilde{x}(t)\| - \eta\rho\mu^2, \eta\zeta\|\tilde{x}(t)\|\}$$

   Hence we have

$$\|\tilde{x}(t+1)\| \leq \max\{(1 - \eta\mu)\|\tilde{x}(t)\| - \eta\rho\mu^2, \eta\zeta\|\tilde{x}(t)\|\} + O(\eta\rho^2) \leq (1 - \eta\mu)\|\tilde{x}(t)\|.$$

2. If $\|\tilde{x}(t)\|_2 \leq \eta\rho\lambda_1^2(t)$, then by Lemma C.24, we have that

$$\|\tilde{x}(t) - \eta A(t)\tilde{x}(t) - \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\|_2 \leq \eta\rho\lambda_1^2(t).$$

   Hence by Lemma C.56

$$\|\tilde{x}(t+1)\| \leq \eta\rho\lambda_1^2(t) + O(\eta\rho^2)$$
$$\leq \eta\rho\lambda_1^2(t+1) + O(\eta\rho^2).$$

Concluding the two cases, we have shown the first and second claim of the induction hypothesis holds. Hence we can show that $\|\tilde{x}(t+1)\| \leq \frac{4\zeta^2}{\mu}\rho$. Then by Lemma C.40, we have that $\|x(t+1) - \Phi(x(t+1))\| \leq \frac{8\zeta^2}{\mu^2}\rho$.

As $t \leq t_{\text{DEC}} + 2C = t_{\text{DEC}} + O(\ln(1/\eta)/\eta)$, by Equation (31),

$$\|\Phi(x(t+1)) - \Phi(x(t_{\text{DEC}}))\| \leq O(-\rho^2\ln\eta).$$

This implies

$$\text{dist}(x(t+1), K) \leq \text{dist}(x(t_{\text{DEC}}), K) + \|x(t_{\text{DEC}}) - \Phi(x(t_{\text{DEC}}))\|$$
$$+ \|\Phi(x(t_{\text{DEC}})) - \Phi(x(t+1))\| + \|x(t+1) - \Phi(x(t+1))\|$$
$$= h/2 + O(\rho^2\ln(1/\eta)) + O(\rho) \leq 3h/4.$$

This proves the fourth claim of the inductive hypothesis.

The induction is complete.

Now define t_{DEC2} the minimal $t \geq t_{\text{DEC}}$, such that $\|\tilde{x}(t)\| \leq \eta\rho\lambda_1^2(t)$.

If $t_{\text{DEC2}} > t_{\text{DEC}} + C$, then by the induction, Lemmas C.7 and C.40,

$$\|\tilde{x}(t_{\text{DEC}} + C)\| \leq (1 - \eta\mu)^C\|\tilde{x}(t_{\text{DEC}})\|$$

$$\leq \frac{\eta\mu^3}{4\zeta^2}\|\tilde{x}(t_{\text{DEC}})\|$$

$$\leq \frac{\eta\mu^3}{4\zeta^2}\zeta\|x(t_{\text{DEC}}) - \Phi(x(t_{\text{DEC}}))\|$$

$$\leq \frac{\eta\mu^2}{4\zeta}\|\nabla L(t_{\text{DEC}})\|$$

$$\leq \mu^2\eta\rho$$

$$\leq \lambda_1^2(t_{\text{DEC}} + C)\eta\rho.$$

This is a contradiction. Hence we have $t_{\text{DEC2}} \leq t_{\text{DEC}} + C$. By the induction hypothesis $x(t_{\text{DEC2}}) \in \mathbb{I}_1 \cap K^{3h/4}$.

Furthermore by induction, for any t satisfying $t_{\text{DEC2}} \leq t \leq t_{\text{DEC}} + 2C$, we have that
$$\|\tilde{x}(t)\| \leq \eta\rho\lambda_1^2(t) + O(\eta\rho^2).$$
By the induction hypothesis $x(t) \in \mathbb{I}_1 \cap K^{3h/4}$ and $\|\Phi(x(t)) - \Phi(x(t_{\text{DEC}}))\| = O(\rho^2 \ln(1/\rho))$. $\square$

Lemma C.45. *Under condition of Theorem C.35, assuming assuming t satisfy that $x(t) \in \mathbb{I}_1 \cap K^{3h/4}$, then we have that*
$$\begin{cases} R_k(x(t)) \geq 0 \Rightarrow R_k(x(t+1)) + \lambda_k^2(t+1)\eta\rho \leq (1 - \eta\mu)(R_k(x(t)) + \lambda_k^2(t)\eta\rho), \\ R_k(x(t)) \leq 0 \Rightarrow R_k(x(t+1)) \leq O(\eta\rho^2). \end{cases}$$

Proof. As $x(t) \in \mathbb{I}_1$, $\|\tilde{x}(t)\|_2 \leq \zeta\eta\rho + O(\eta\rho^2)$.

As $\|\tilde{x}(t)\|_2 = O(\rho)$, we have $\overline{x(t)x(t+1)} \subset K^h$ and $\overline{\Phi(x(t))\Phi(x(t+1))} \subset K^r$.

We will begin with a quantization technique separating $[M]$ into disjoint continuous subset $S_1, ..., S_p$ such that $\forall i \neq j$,
$$\min_{k \in S_i, l \in S_j} |\lambda_k(t) - \lambda_l(t)| \geq \rho.$$

By Lemmas C.41 and C.56, we have that for any $n \in [M]$,
$$\begin{aligned} |\lambda_k(t) - \lambda_k(t+1)| &= O(\|\nabla^2 L(\Phi(x(t))) - \nabla^2 L(\Phi(x(t+1)))\|) \\ &= O(\|\Phi(x(t)) - \Phi(x(t+1))\|) \\ &= O(\eta\rho^2). \end{aligned}$$

This implies
$$\min_{k \in S_i, l \in S_j} |\lambda_k(t+1) - \lambda_l(t+1)| \geq \rho - O(\eta\rho^2) \geq 0.99\rho.$$

Define
$$P_{S^{(i)}}^{(t)} \triangleq \sum_{k \in S_i} v_n(t)v_n(t)^T.$$

By Theorem C.58, for any k,
$$\|P_{S_k}^{(t)} - P_{S_k}^{(t+1)}\| \leq O\left(\frac{\|\nabla^2 L(\Phi(x(t))) - \nabla^2 L(\Phi(x(t+1)))\|}{\rho}\right) = O(\eta\rho).$$

By Lemma C.43, we have that
$$\left\|\tilde{x}(t+1) - \tilde{x}(t) + \eta A(t)\tilde{x}(t) + \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\right\|_2 = O(\eta\rho^2).$$

We will write $x'(t+1)$ as shorthand of $\tilde{x}(t) - \eta A(t)\tilde{x}(t) - \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}$.

Now we discuss by cases,

1. If $\sqrt{\sum_{i=j}^p \|P_{S^{(i)}}^{(t)}\tilde{x}(t)\|^2} > \max_{k \in S_j} \lambda_k^2(t)\eta\rho > \mu^2\eta\rho$, by Lemma C.26,

$$\sqrt{\sum_{i=j}^p \|P_{S^{(i)}}^{(t)}\tilde{x}(t+1)\|^2} \leq \sqrt{\sum_{i=j}^p \|P_{S^{(i)}}^{(t)}x'(t+1)\|^2} + O(\eta\rho^2)$$

$$\leq \max\left\{\left(1 - \eta\lambda_D(t+1)\right)\|\sum_{i=j}^p P_{S^{(i)}}^{(t)}\tilde{x}(t)\| - \eta\rho\lambda_D(t+1)^2\frac{\|\sum_{i=j}^p P_{S^{(i)}}^{(t)}\tilde{x}(t)\|}{\|\tilde{x}(t)\|},\right.$$

$$\eta\max_{k \in S_j}\lambda_k(t+1)\|\sum_{i=j}^p P_{S^{(i)}}^{(t)}\tilde{x}(t)\|\right\} + O(\eta\rho^2)$$

$$\leq \max\left\{\left(1 - \eta\mu\right)\|\sum_{i=j}^p P_{S^{(i)}}^{(t)}\tilde{x}(t)\| - \eta\rho\frac{\mu^3}{2\zeta}, \eta\zeta\|\sum_{i=j}^p P_{S^{(i)}}^{(t)}\tilde{x}(t)\|\right\} + O(\eta\rho^2).$$

This further implies

$$\sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t+1)}\tilde{x}(t+1)\|^2} \leq \sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t)}\tilde{x}(t+1)\|^2} + O(\eta\rho\|\tilde{x}(t+1)\|)$$

$$\leq \max\{\big(1-\eta\mu\big)\|\sum_{i=j}^{p} P_{S^{(i)}}^{(t)}\tilde{x}(t)\| - \eta\rho\frac{\mu^3}{2\zeta}, \eta\zeta\|\sum_{i=j}^{p} P_{S^{(i)}}^{(t)}\tilde{x}(t)\|\} + O(\eta\rho^2)$$

$$\leq \big(1-\eta\mu\big)\|\sum_{i=j}^{p} P_{S^{(i)}}^{(t)}\tilde{x}(t)\|.$$

2. If $\sqrt{\sum_{i=j}^{p}\|P_{S^{(i)}}^{(t)}\tilde{x}(t)\|^2} \leq \max_{k\in S_j}\lambda_k^2(t)\eta\rho$, then by Lemma C.24, we have that

$$\|\sum_{i=j}^{p} P_{S^{(i)}}^{(t)}x'(t+1)\|_2 \leq \eta\rho\max_{k\in S_j}\lambda_k^2(t).$$

Hence we have that

$$\sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t)}\tilde{x}(t+1)\|^2} \leq \sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t)}x'(t+1)\|^2} + O(\eta\rho^2)$$

$$\leq \max_{k\in S_j}\lambda_k^2(t)\eta\rho + O(\eta\rho^2)$$

$$\leq \max_{k\in S_j}\lambda_k^2(t+1)\eta\rho + O(\eta\rho^2).$$

This further implies

$$\sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t+1)}\tilde{x}(t+1)\|^2} \leq \sqrt{\sum_{i=j}^{p} \|P_{S^{(i)}}^{(t)}\tilde{x}(t+1)\|^2} + O(\eta\rho\|\tilde{x}(t+1)\|)$$

$$\leq \max_{k\in S_j}\lambda_k^2(t)\eta\rho + O(\eta\rho^2)$$

$$\leq \max_{k\in S_j}\lambda_k^2(t+1)\eta\rho + O(\eta\rho^2).$$

Finally taking into quantization error, as all the eigenvalue in the same group at most differ $D\rho$, for any $i \in S_j$, we have that $-\lambda_i^2(t+1) + \max_{k\in S_j}\lambda_k^2(t+1) \leq 2D\zeta\rho + D^2\rho^2$.

Hence the previous discussion concludes as

1. If $R_k(x(t)) \geq 0$

$$R_k(x(t+1)) + \lambda_k^2(t+1)\eta\rho \leq (1-\eta\mu)(R_k(x(t)) + \lambda_k^2(t).\eta\rho)$$

2. If $R_k(x(t)) < 0$

$$R_k(x(t+1)) \leq O(\eta\rho^2).$$

$\square$

Lemma C.46. *Under condition of Theorem C.35, assuming there exists t_{DEC2} such that for any t satisfying $t_{\mathrm{DEC2}} \leq t \leq t_{\mathrm{DEC2}} + \Theta(\ln(1/\eta)/\eta)$, we have that $x(t) \in \mathbb{I}_1 \cap K^{3h/4}$. Then there exists $t_{\mathrm{INV}} = t_{\mathrm{DEC2}} + O(\ln(1/\eta)/\eta))$ such that for any t satisfying $t_{\mathrm{INV}} \leq t \leq t_{\mathrm{INV}} + \Theta(\ln(1/\eta)/\eta)$, we have that*

$$x(t) \in \cap_{k\in[M]}\mathbb{I}_k \cap K^{7h/8}.$$

$$\|\Phi(x(t)) - \Phi(x(t_{\mathrm{DEC2}}))\| = O(\rho^2\ln(1/\eta)).$$

Proof. The proof is almost identical with Lemma C.44 replacing the first two iterative hypothesis to Lemma C.45 and is omitted here. $\square$

C.6.2 PHASE II (PROOF OF THEOREM C.36)

The goal of this section is to prove Theorem C.36.

Proof. First by Lemma C.51, there exists t_{ALIGN} such that $\nabla L(x(t_{\text{ALIGN}}))$ aligns with $v_1(t_{\text{ALIGN}})$ with $O(\rho)$ angle. Then we can inductively use Lemma C.52 to show that $x(t)$ maintains in K^h and Lemma C.51 to show that $x(t)$ keeps alignment with $v_1(t)$ for $T_3/\eta\rho^2$ step. $\qquad\square$

Alignment to Top Eigenvector   We will continue to use the notations introduced in Appendix C.6.1.

We further define

$$S = \left\{ t \Big| \|\tilde{x}(t)\| \leq \frac{\eta\lambda_1^2}{2 - \eta\lambda_1}\rho + O(\eta\rho^2) \right\},$$

$$T = \left\{ t \Big| \|\tilde{x}(t)\| \leq \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2 - \eta\lambda_1} + \frac{\eta\lambda_2^2}{2 - \eta\lambda_2}\right)\rho \right\},$$

$$U = \left\{ t \Big| \Omega(\rho^2) \leq \|\tilde{x}_1(t)\| \leq \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2 - \eta\lambda_1} + \frac{\eta\lambda_2^2}{2 - \eta\lambda_2}\right)\rho \right\}.$$

Here the constant in O depends on the constant in $\mathbb{I}_j$ and will be made clear in Lemma C.48.

For $s \in S$, define $\text{next}(t)$ as the smallest integer greater than s in S.

Lemma C.47. *Under the condition of Theorem C.36, there exist constants $C_1, C_2 < 1$ independent of η and ρ, if $\|\tilde{x}_1(t)\|_2 \leq \frac{1}{2}\left(\frac{\eta\lambda_1(t)^2}{2 - \eta\lambda_1(t)} + \frac{\eta\lambda_2(t)^2}{2 - \eta\lambda_2(t)}\right)\rho$ and $x(t) \in \cap_j\mathbb{I}_j \cap K^{7h/8}$, then*

$$\|\tilde{x}(t)\|_2 \geq C_1\frac{\eta\lambda_1^2}{2 - \eta\lambda_1}\rho \Rightarrow \|\tilde{x}(t+1)\|_2 \leq C_2\frac{\eta\lambda_1^2}{2 - \eta\lambda_1}\rho$$

Proof. By Lemma C.43, if we write $x'(t+1)$ as shorthand of $\tilde{x}(t) - \eta A(t)\tilde{x}(t) - \eta\rho A^2(t)\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}$, then $\|\tilde{x}(t+1) - x'(t+1)\| = O(\eta\rho^2)$.

Define $\mathbb{I}_j^{\text{quad}}$ as $\{x | R_j(x) \leq 0\}$. Then we can find a surrogate $x_{\text{sur}}(t)$ such that $x_{\text{sur}}(t) \in \cap_j\mathbb{I}_j^{\text{quad}} \cap K^h$ and $\|x_{\text{sur}}(t) - \tilde{x}(t)\|_2 = O(\eta\rho^2)$. We will write $x_{\text{sur}}'(t+1)$ as shorthand of $x_{\text{sur}}(t) - \eta A(t)x_{\text{sur}}(t) - \eta\rho A^2(t)\frac{x_{\text{sur}}(t)}{\|x_{\text{sur}}(t)\|}$.

Let

$$h(t) \triangleq (2 - t)\sqrt{\frac{1}{2t}\left(\frac{(\zeta - \Delta)^2}{\zeta^2} + 1\right) + \left(1 - \frac{1}{2t}\left(\frac{(\zeta - \Delta)^2}{\zeta^2} + 1\right)\right)\max\{\frac{\zeta^2 - \mu^2}{\zeta^2}, \frac{(\zeta - \Delta)^2}{\zeta^2}\}}.$$

As $h(1) < 1$, we can choose $C_1 < 1$, such that $h(C_1) < 1$.

We can further choose $C_2 = \max\{(h(C_1) + 1)/2, 1 - \frac{\mu^2}{3\zeta^2}\} < 1$.

We will discuss by cases

1. If

$$\|\tilde{x}(t)\|_2 \geq \frac{\eta\lambda_1^4}{\lambda_1^2(1 - \eta\lambda_D) + (\lambda_1^2 - \lambda_D^2)(1 - \eta\lambda_1)}\rho$$

   Then

$$\frac{\|\tilde{x}(t)\|_2}{\frac{\eta\lambda_1^2}{2 - \eta\lambda_1}\rho} = \frac{\lambda_1^2(2 - \eta\lambda_1)}{\lambda_1^2(1 - \eta\lambda_D) + (\lambda_1^2 - \lambda_D^2)(1 - \eta\lambda_1)}$$

$$= \frac{\lambda_1^2(2 - \eta\lambda_1)}{\lambda_1^2(2 - \eta\lambda_1 - \eta\lambda_D) - \lambda_D^2(1 - \eta\lambda_1)}$$

$$\geq \frac{1}{1 - \frac{\lambda_D^2}{\lambda_1^2}\frac{1 - \eta\lambda_1}{2 - \eta\lambda_1}} \geq 1 + \frac{\lambda_D^2}{\lambda_1^2}\frac{1 - \eta\lambda_1}{2 - \eta\lambda_1} \geq 1 + \frac{\mu^2}{3\zeta^2}.$$

In such case we have

$$\|\frac{\tilde{x}(t)}{\|\tilde{x}(t)\|} - \frac{x_{\text{sur}}(t)}{\|x_{\text{sur}}(t)\|}\| = O(\rho).$$

Then we have $\|x'_{\text{sur}}(t+1) - x'(t+1)\| = O(\eta\rho^2)$. By Lemma C.28, we have that

$$\|\tilde{x}(t+1)\|_2 \le \|\tilde{x}(t+1) - x'(t+1)\|_2 + \|\tilde{x}(t+1) - x'_{\text{sur}}(t+1)\| + \|x'_{\text{sur}}(t+1)\|$$

$$\le \max(\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho - \eta\rho\frac{\lambda_D^4}{2\lambda_1^2}, \eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2) + O(\eta\rho^2)$$

$$\le \max(1 - \frac{\lambda_D^4(2-\eta\lambda_1)}{2\lambda_1^4}, (2-\eta\lambda_1) - (1-\eta\lambda_1)(1+\frac{\mu^2}{3\zeta^2}))\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho$$

$$\le (1 - \frac{\mu^2}{3\zeta^2})\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho \le C_2\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho$$

2. If

$$\|\tilde{x}(t)\|_2 \le \frac{\eta\lambda_1^4}{\lambda_1^2(1-\eta\lambda_D) + (\lambda_1^2 - \lambda_D^2)(1-\eta\lambda_1)}\rho \le \frac{\eta\lambda_1^2}{1-\eta\lambda_1}\rho.$$

Then we have

$$\frac{|-\eta\rho\lambda_D^2 + (1-\eta\lambda_D)\|\tilde{x}(t)\|_2|}{\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2} \le \frac{\lambda_1^2 - \lambda_D^2}{\lambda_1^2}.$$

$$\frac{|\eta\rho\lambda_2^2 - (1-\eta\lambda_2)\|\tilde{x}(t)\|_2|}{\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2} \le \frac{\lambda_2^2}{\lambda_1^2}.$$

By Lemma C.31,

$$\|x'(t+1)\|_2$$

$$\le(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)\sqrt{\frac{\|\tilde{x}_1^2(t)\|_2}{\|\tilde{x}(t)\|_2^2} + (1 - \frac{\|\tilde{x}_1^2(t)\|_2}{\|\tilde{x}(t)\|_2^2})\max\{\frac{\lambda_1^2 - \lambda_D^2}{\lambda_1^2}, \frac{\lambda_2^2}{\lambda_1^2}\}}$$

$$\le(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|_2)\sqrt{\frac{\|\tilde{x}_1^2(t)\|_2}{\|\tilde{x}(t)\|_2^2} + (1 - \frac{\|\tilde{x}_1^2(t)\|_2}{\|\tilde{x}(t)\|_2^2})\max\{\frac{\zeta^2 - \mu^2}{\zeta^2}, \frac{(\zeta-\Delta)^2}{\zeta^2}\}}.$$

.

As

$$\|\tilde{x}_1(t)\|_2 \le \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2}\right)\rho$$

For $\|\tilde{x}(t)\|_2 \ge \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho C_1$,

$$\frac{\|\tilde{x}_1(t)\|_2}{\|\tilde{x}(t)\|_2} \le \frac{1}{2}\left(\frac{\lambda_2^2(2-\eta\lambda_1)}{\lambda_1^2(2-\eta\lambda_2)} + 1\right)/C_1 \le \frac{1}{2}\left(\frac{\lambda_2^2}{\lambda_1^2} + 1\right)/C_1 \le \frac{1}{2C_1}\left(\frac{(\zeta-\Delta)^2}{\zeta^2} + 1\right).$$

After plugging in, we have that

$$\|\tilde{x}(t+1)\|_2 \le \|x'(t+1)\|_2 + O(\eta\rho^2) \le h(C_1)\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho + O(\eta\rho^2) \le C_2\frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho.$$

This concludes the proof. $\square$

Lemma C.48. *Under the condition of Theorem C.36, assuming that (1) $x(t) \in \cap_j \mathbb{I}_j$, (2) $t \in S \cap K^{15h/16}$ then* $\text{next}(t)$ *is always well defined and* $\text{next}(t) \le t + 2$.

Proof. If $t + 1 \notin S$, then

$$\|\tilde{x}(t+1)\| \ge \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho + \Theta(\eta\rho^2).$$

Following similar argument in Lemma C.24, we have that $x(t+1) \in \cap_j \mathbb{I}_j$.

By Lemma C.43, if we write $x'(t+2)$ as shorthand of $\tilde{x}(t+1) - \eta A(t+1)\tilde{x}(t+1) - \eta \rho A^2(t+1)\frac{\tilde{x}(t+1)}{\|\tilde{x}(t+1)\|}$, then $\|\tilde{x}(t+2) - x'(t+2)\| = O(\eta \rho^2)$.

Define $\mathbb{I}_j^{\text{quad}}$ as $\{x | R_j(x) \leq 0\}$. Then we can find a surrogate $x_{\text{sur}}(t+1)$ such that $x_{\text{sur}}(t+1) \in \cap_j \mathbb{I}_j^{\text{quad}} \cap K^h$, and $\|x_{\text{sur}}(t+1) - \tilde{x}(t+1)\|_2 = O(\eta \rho^2)$. We will write $x'_{\text{sur}}(t+2)$ as shorthand of $x_{\text{sur}}(t+1) - \eta A(t+1)x_{\text{sur}}(t+1) - \eta \rho A^2(t+1)\frac{x_{\text{sur}}(t+1)}{\|x_{\text{sur}}(t+1)\|}$.

As $\|\tilde{x}(t+1)\| = \Omega(\eta \rho)$, we have

$$\|\frac{x_{\text{sur}}(t+1)}{\|x_{\text{sur}}(t+1)\|} - \frac{\tilde{x}(t+1)}{\|\tilde{x}(t+1)\|}\|_2 = O(\rho).$$

Hence we have that $\|\tilde{x}(t+2) - x'_{\text{sur}}(t+2)\| = \|\tilde{x}(t+2) - x'(t+2)\| + \|x'(t+2) - x'_{\text{sur}}(t+2)\| = O(\eta \rho^2)$

Notice we have $\|x_{\text{sur}}(t)\|_2 \geq \frac{\eta \lambda_1^2}{2 - \eta \lambda_1} \rho$ for properly chosen function in the definition S, hence, by Lemma C.28

$$\|x'_{\text{sur}}(t+1)\|_2 \leq \frac{\eta \lambda_1^2}{2 - \eta \lambda_1} \rho.$$

This further implies $t+2 \in S$. $\qquad \square$

Lemma C.49. *Under the condition of Theorem C.36, there exists constant $C > 0$ independent of η and ρ, assuming that (1) $x(t) \in \cap_j \mathbb{I}_j$, (2) $t \in S \cap K^{7h/8}$ (3) $\Omega(\rho^2) \leq \|\tilde{x}_1(t)\|$, then*

$$\|\tilde{x}_1(\text{next}(t))\| \geq \|\tilde{x}_1(t)\| - O(\eta \rho^2).$$

Proof. This is by standard approximation as in previous proof and Lemma C.32. $\qquad \square$

Lemma C.50. *Under the condition of Theorem C.36, there exists constant $C > 0$ independent of η and ρ, assuming that (1) $x(t) \in \cap_j \mathbb{I}_j$, (2) $t \in S \cap K^{7h/8}$ (3) $\Omega(\rho^2) \leq \|\tilde{x}_1(t)\| \leq \frac{1}{2}\left(\frac{\eta \lambda_1^2}{2 - \eta \lambda_1} + \frac{\eta \lambda_2^2}{2 - \eta \lambda_2}\right)\rho$, then*

$$\|\tilde{x}_1(\text{next}(t))\| \geq \min\{(1 + C\eta)\|\tilde{x}_1(t)\|, \frac{1}{2}\left(\frac{\eta \lambda_1^2}{2 - \eta \lambda_1} + \frac{\eta \lambda_2^2}{2 - \eta \lambda_2}\right)\rho\},$$

$$\text{or } \|\tilde{x}_1(\text{next}(\text{next}(t)))\| \geq \min\{(1 + C\eta)\|\tilde{x}_1(t)\|, \frac{1}{2}\left(\frac{\eta \lambda_1^2}{2 - \eta \lambda_1} + \frac{\eta \lambda_2^2}{2 - \eta \lambda_2}\right)\rho\}.$$

Proof. In this proof, we will sometime drop the t in $\lambda_k(t)$ or $A(t)$. Applying Lemma C.48, we have $\text{next}(t)$ and $\text{next}(\text{next}(t))$ are well-defined. We can suppose $\|\tilde{x}_1(\text{next}(t))\|_2 \leq \frac{1}{2}\left(\frac{\eta \lambda_1^2}{2 - \eta \lambda_1} + \frac{\eta \lambda_2^2}{2 - \eta \lambda_2}\right)$, else the result holds already.

By assumption, we have $\|\tilde{x}_1(t)\| \geq \Omega(\rho^2)$.

Using Lemma C.43,

$$\|\tilde{x}(t+1) - \tilde{x}(t) + \eta A\tilde{x}(t) + \eta \rho A^2 \frac{\tilde{x}(t)}{\|\tilde{x}(t)\|}\| \leq O(\eta \rho^2).$$

Denote

$$x'(t+1) = \tilde{x}(t) + \eta A\tilde{x}(t) + \eta \rho A^2 \frac{\tilde{x}(t)}{\|\tilde{x}(t)\|},$$

as the one step update of SAM on the quadratic approximation of the general loss.

Now using Lemma C.47 and the induction hypothesis, we have for some C_1 and C_2 smaller than 1, $\|\tilde{x}(t)\| \geq C_1 \frac{\eta \lambda_1^2}{2 - \eta \lambda_1} \rho \Rightarrow \|x'(t+1)\| \leq C_2 \frac{\eta \lambda_1^2}{2 - \eta \lambda_1} \rho$.

We will discuss by cases,

1 If $\|\tilde{x}(t)\| \leq C_1 \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho$

   If $\text{next}(t) = t+1$, then

$$\frac{\|x_1'(t+1)\|}{\|\tilde{x}_1(t)\|} = \frac{\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|}{\|\tilde{x}(t)\|} \geq \frac{(2-C_1) - \eta\lambda_1 + C_1\eta\lambda_1}{C_1} \geq \frac{1}{C_1}$$

As we have $\tilde{x}_1(t) = \Omega(\rho^2)$, we have $x_1'(t+1) = \Omega(\rho^2)$, then as $\|\tilde{x}_1(t+1) - x_1'(t+1)\| = O(\eta\rho^2)$, this implies

$$\|\tilde{x}_1(t+1)\| \geq \|x_1'(t+1)\| - O(\eta\rho^2) \geq \frac{1}{C_1}\|\tilde{x}_1(t+1)\| - O(\eta\rho^2) \geq \frac{1}{2}(\frac{1}{C_1}+1)\|\tilde{x}_1(t)\|.$$

If $\text{next}(t) = t+2$, define $x'(t+2) = x'(t+1) - \eta A x'(t+1) - \eta\rho A^2 \frac{x'(t+1)}{\|x'(t+1)\|}$, as $\|\tilde{x}_1(t+1)\| = \Omega(\eta\rho)$, by Lemma C.42, we have $\|x'(t+2) - \tilde{x}(t+2)\| = O(\eta\rho^2)$.

$$\begin{aligned}
\frac{\|x'(t+2)\|}{\|\tilde{x}_1(t)\|} &= \frac{(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|)(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|x'(t+1)\|)}{\|\tilde{x}(t)\|\|x'(t+1)\|} \\
&\geq \frac{(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|)\left(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\left(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|\right)\right)}{\|\tilde{x}(t)\|\left(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|\right)} \\
&= \frac{\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\left(\eta\rho\lambda_1^2 - (1-\eta\lambda_1)\|\tilde{x}(t)\|\right)}{\|\tilde{x}(t)\|} \\
&\geq (1-\eta\lambda_1)^2 + \frac{\eta\lambda_1}{C_1}(2-\eta\lambda_1) \geq 1 + 4C\eta.
\end{aligned}$$

   Combining with $|\tilde{x}_1(t)| \geq \Omega(\rho^2)$, we have that

$$\|\tilde{x}_1(\text{next}(t))\| \geq (1+C\eta)\|\tilde{x}_1(t)\|$$

2 *Case 2* $\|\tilde{x}(t)\| > C_1 \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho$, then $\|\tilde{x}(t+1)\| \leq C_2 \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho$, $\text{next}(t) = t+1$

   By Lemma C.49, $\|\tilde{x}_1(t+1)\| \geq (1-C\eta)\|\tilde{x}_1(t)\|$.

   As $\|\tilde{x}(\text{next}(t))\| \leq C_2 \frac{\eta\lambda_1^2}{2-\eta\lambda_1}$, similar to the first case,

$$\|\tilde{x}_1(\text{next}(\text{next}(t)))\| \geq (1+4C\eta)\|\tilde{x}_1(\text{next}(t))\| \geq (1+C\eta)\|\tilde{x}_1(t)\|.$$

In conclusion, if $\|\tilde{x}_1(t)\| \leq \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2}\right)\rho$, we would have there exists $C > 0$

$$\|\tilde{x}_1(\text{next}(t))\| \geq (1+C\eta)\|\tilde{x}_1(t)\| \text{ or } \|\tilde{x}_1(\text{next}(\text{next}(t)))\| \geq (1+C\eta)\|\tilde{x}_1(t)\|.$$

$\square$

Lemma C.51. *Under the condition of Theorem C.36, there exists constant $T_2 > 0$ independent of η and ρ, we would have that when $t = t_{\text{ALIGN}} = \lceil T_2 \ln(1/\rho)/\eta \rceil$,*

$$|\langle x(t) - \Phi(x(t)), v_1(x(t))\rangle| = \Theta(\eta\rho),$$
$$\max_{j\in[2:M]}|\langle x(t) - \Phi(x(t)), v_j(x(t))\rangle| = O(\eta\rho^2).$$

Further if $x(t') \in K^h$ holds for $t' = 0,1,...,t_{\text{LOCAL}}$, then for t satisfying $t_{\text{ALIGN}} \leq t \leq t_{\text{LOCAL}}$

$$|\langle x(t) - \Phi(x(t)), v_1(x(t))\rangle| = \Theta(\eta\rho),$$
$$\max_{j\in[2:M]}|\langle x(t) - \Phi(x(t)), v_j(x(t))\rangle| = O(\eta\rho^2).$$

Proof. Let C be the constant defined in Lemma C.50.

Suppose WLOG $\|\tilde{x}_1(0)\| \geq \rho^2$. Define

$$C_1 \triangleq \lceil \log_{1+C\eta}(\frac{\eta\lambda_1^2}{2-\eta\lambda_1}/\rho)\rceil$$

$$C_2 \triangleq C_1 + \lceil \ln_{\max\{1-\frac{\mu^2}{2\zeta^2}, 1-\frac{\Delta^2}{4\zeta^2}\}}\frac{\rho^2}{\zeta^2}\rceil = O(\log(1/\rho)/\eta).$$

We will choose t_{ALIGNMID} as the minimal $t \in S$, such that $\|\tilde{x}_1(t)\| \geq \frac{1}{2} \left(\frac{\eta \lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2} \right) \rho$.

Then by induction and Lemmas C.49 and C.50, we easily have that for $t \leq \min\{C_2+1, t_{\text{ALIGNMID}}\}$ and $t \in S$, we have that

$$x(t) \in K^{7h/8} \cap \cap_j \mathbb{I}_j \,,$$

$$\|\tilde{x}_1(t)\| \geq \min\{(1+C\eta)^{t/4}\|\tilde{x}_1(0)\|, \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2} \right)\rho\}$$

$$\text{or} \quad \|\tilde{x}_1(\text{next}(t))\| \geq \min\{(1+C\eta)^{t/4}\|\tilde{x}_1(0)\|, \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2} \right)\rho\}\,.$$

The detailed induction is analogous to previous inductive argument and is omitted. If $t_{\text{ALIGNMID}} \geq C_1$, then we have for the minimal $t \geq C_1$ and $t \in S$

$$\|\tilde{x}_1(t)\| \geq \frac{\eta\lambda_1^2}{2-\eta\lambda_1}\rho\,.$$

This is a contradiction and we have that $t_{\text{ALIGNMID}} \leq C_1$.

As we have $\|\tilde{x}_1(\text{next}(t))\| \geq \|\tilde{x}_1(t)\| - O(\eta\rho^2)$ for $\|\tilde{x}_1(t)\| \geq \frac{1}{2}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + \frac{\eta\lambda_2^2}{2-\eta\lambda_2} \right)\rho$,

$$\|\tilde{x}(t)\| \geq \|\tilde{x}_1(t))\| \geq \frac{1}{4}\left(\frac{\eta\lambda_1^2}{2-\eta\lambda_1} + 3\frac{\eta\lambda_2^2}{2-\eta\lambda_2} \right)\rho.$$

for $C_2 \geq t \geq t_{\text{ALIGNMID}}$.

We will then show that for $t \geq t_{\text{ALIGNMID}} + C_1$ iteration, $\|P^{(2:D)}\bar{x}(t+1)\| \leq O(\eta\rho^2)$.

For $C_1 \geq t \geq t_{\text{ALIGNMID}}$,

$$1 - \eta\lambda_2 - \eta\rho\frac{\lambda_2^2}{\|\tilde{x}(t)\|} \leq 1 - \eta\lambda_D - \eta\rho\frac{\lambda_D^2}{\|\tilde{x}(t)\|} \leq 1 - \frac{\lambda_D^2}{2\lambda_1^2} \leq 1 - \frac{\mu^2}{\zeta^2}$$

Notice that,

$$1 - \eta\lambda_2 - \eta\rho\frac{\lambda_2^2}{\|\tilde{x}(t)\|}$$

$$\geq 1 - \eta\lambda_2 - \eta\rho\frac{\lambda_2^2}{\|\tilde{x}(t)\|}$$

$$\geq 1 - \eta\lambda_2 - \frac{4\lambda_2^2}{\lambda_1^2 + 3\lambda_2^2}(2 - \eta\lambda_2)$$

$$\geq -1 + \frac{2(\lambda_1^2 - \lambda_2^2)}{\lambda_1^2 + 3\lambda_2^2} \geq -1 + \frac{\Delta^2}{2\zeta^2}$$

Hence,

$$\|P^{(2:D)}(t)x'(t+1)\|_2 \leq \max\{1 - \frac{\mu^2}{\zeta^2}, 1 - \frac{\Delta^2}{2\zeta^2}\}\|P^{(2:D)}(t)\tilde{x}(t)\|_2$$

Now by Lemma C.56 and Theorem C.58,

$$\|P^{(2:D)}(t) - P^{(2:D)}(t+1)\| \leq O(\eta\rho^2)$$
$$\|v_1(t) - v_1(t+1)\| \leq O(\eta\rho^2)$$
$$\|\lambda_1(t) - \lambda_1(t+1)\| \leq O(\eta\rho^2)$$

By Lemma C.43,

$$\|x'(t+1) - \tilde{x}(t+1)\| = O(\eta\rho^2)$$

Combining,

$$\|P^{(2:D)}(t+1)\tilde{x}(t+1)\| \le \max\{1 - \frac{\mu^2}{2\zeta^2}, 1 - \frac{\Delta^2}{4\zeta^2}\}\|P^{(2:D)}(t)\tilde{x}(t)\| + O(\eta\rho^2)$$

Hence when $t = t_{\text{ALIGN}} = t_{\text{ALIGNMID}} + C_2$,

$$\|\tilde{x}(t)\| \ge \|\tilde{x}_1(t)\| \ge \Omega(\eta\rho)$$
$$\|P^{(2:D)}(t)\tilde{x}(t)\| \le O(\eta\rho^2)$$

By $x(t) \in \mathbb{I}_1$, we easily have $\|\tilde{x}_1(t)\| = O(\eta\rho)$. Hence we conclude that

$$\|\tilde{x}_1(t)\| = \Theta(\eta\rho)$$
$$\|P^{(2:D)}(t)\tilde{x}(t)\| = O(\eta\rho^2) \,.$$

The second claim is just another induction similar to previous steps and is omitted as well.

$\square$

Tracking Riemannian Gradient Flow   We are now ready to show that $\Phi(x(t))$ will track the solution of Equation (6). The main principal of this proof has been introduced in Section 4.3.

Lemma C.52. *Under the condition of Theorem C.36, for any t satisfying that*

$$x(t) \in K^h,$$
$$\|\tilde{x}_1(t)\| = \Theta(\eta\rho),$$
$$\|P^{(2:D)}(t)\tilde{x}(t)\| = O(\eta\rho^2) \,,$$

it holds that

$$\|\Phi(x(t+1)) - \Phi(x(t)) + \eta\rho^2 P_{X,\Gamma}^{\perp}\nabla\lambda_1(t)/2\| \le O(\eta\rho^3 + \eta^2\rho^2) \,.$$

Proof. To begin with, we can approximate $\Phi(x(t+1)) - \Phi(x(t))$ by its first order Taylor Expansion, by Lemma C.12,

$$\|\Phi(x(t+1)) - \Phi(x(t)) - \partial\Phi(x(t))(x(t+1) - x(t))\| = O(\|x(t+1) - x(t)\|^2) = O(\eta^2\rho^2) \,.$$

Then by plugging in the update rule and another Taylor Expansion,

$$\|\partial\Phi(x(t))(x(t+1) - x(t)) - \eta\rho\partial\Phi(x(t))\nabla^2 L(x)\frac{\nabla L(x)}{\|\nabla L(x)\|}$$
$$- \eta\rho^2\partial\Phi(x(t))\partial\nabla^2 L(x)[\frac{\nabla L(x)}{\|\nabla L(x)\|}, \frac{\nabla L(x)}{\|\nabla L(x)\|}]/2\|_2 = O(\eta\rho^3).$$

Using Lemma C.9, we have

$$\|\eta\rho\partial\Phi(x(t))\nabla^2 L(x)\frac{\nabla L(x)}{\|\nabla L(x)\|}\| = \eta\rho\|\nabla L(x)\|\|\partial^2\Phi(x(t))\left[\frac{\nabla L(x)}{\|\nabla L(x)\|}, \frac{\nabla L(x)}{\|\nabla L(x)\|}\right]\| = O(\eta\rho\|\nabla L(x)\|) \,.$$

Putting together, we have that

$$\|\Phi(x(t+1)) - \Phi(x(t)) - \eta\rho^2\partial\Phi(x(t))\partial\nabla^2 L(\Phi(x(t)))[\frac{\nabla L(x)}{\|\nabla L(x)\|}, \frac{\nabla L(x)}{\|\nabla L(x)\|}]/2\|$$
$$\le O(\eta^2\rho^2 + \eta\rho^3) + O(\eta\rho\|\nabla L(x)\|) \,.$$

As we have $\|\tilde{x}(t)\| = \Theta(\eta\rho)$, hence by Lemmas C.8 and C.40,

$$\|\Phi(x(t+1)) - \Phi(x(t)) - \eta\rho^2\partial\Phi(x(t))\partial\nabla^2 L(\Phi(x(t)))[\frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}, \frac{\nabla L(x(t))}{\|\nabla L(x(t))\|}]/2\|$$
$$\le O(\eta\rho^3 + \eta^2\rho^2)$$

Finally, we have that

$$\|\eta\rho^2 \partial\Phi(x(t))\partial\nabla^2 L\left(\Phi(x(t))\right)[\frac{\nabla L\left(x(t)\right)}{\|\nabla L\left(x(t)\right)\|}, \frac{\nabla L\left(x(t)\right)}{\|\nabla L\left(x(t)\right)\|}]/2$$
$$- \eta\rho^2 \partial\Phi(x(t))\partial\nabla^2 L\left(\Phi(x(t))\right)[v_1(t), v_1(t)]/2\| \leq O(\eta\rho^3)$$

as the angle between $\frac{\nabla L(x)}{\|\nabla L(x)\|}$ and $v_1(t)$ is $O(\rho)$.

By Lemma C.9, it holds that

$$\partial\Phi(x(t))\partial\nabla^2 L\left(\Phi(x(t))\right)[v_1(t), v_1(t)] = P_{X,\Gamma}^\perp \nabla(\lambda_1(t))$$

Putting together we have that,

$$\|\Phi(x(t+1)) - \Phi(x(t)) + \eta\rho^2 P_{X,\Gamma}^\perp \nabla\lambda_1(t)/2\| \leq O(\eta\rho^3 + \eta^2\rho^2).$$

It completes the proof. $\qquad\square$

C.6.3 PROOFS OF COROLLARIES 4.10 AND 4.11

Proof of Corollary 4.10. We will do a Taylor expansion on L_ρ^{Max}. By Theorem C.35 and C.36, we have $\|x(\lceil T_3/\eta\rho^2\rceil)) - X(T_3)\| = \tilde{O}(\eta + \rho)$ and $\|x(\lceil T_3/\eta\rho^2\rceil)) - \Phi(x(\lceil T_3/\eta\rho^2\rceil)))\|_2 = O(\eta\rho)$. For convenience, we denote $x(\lceil T_3/\eta\rho^2\rceil)$ by x.

$$R_\rho^{\text{Max}}(x) = \max_{\|v\|_2 \leq 1} \rho v^T \nabla L(x) + \rho^2 v^T \nabla^2 L(x)v/2 + O(\rho^3)$$

Since $\max_{\|v\|_2 \leq 1} \|v^T \nabla L(x)\|_2 = O(\|x - \Phi(x)\|_2) = O(\eta\rho)$, it holds that

$$R_\rho^{\text{Max}}(x) = \rho^2 \max_{\|v\|_2 \leq 1} v^T \nabla^2 L(x)v/2 + O(\eta^2\rho^2 + \rho^3)$$
$$= \rho^2 \lambda_1(\nabla^2 L(x)) + O(\eta^2\rho^2 + \rho^3)$$
$$= \rho^2 \lambda_1(\nabla^2 L(X(T_3))) + \tilde{O}(\eta\rho^2),$$

which completes the proof. $\qquad\square$

Proof of Corollary 4.11. We choose T such that $X(T_\epsilon)$ is sufficiently close to $X(\infty)$, such that $\lambda_1(X(T_\epsilon)) \leq \lambda_1(X(\infty)) + 2\epsilon$. By Corollary 4.10 (let $T_3 = T_\epsilon$), we have that for all ρ, η such that $\eta\ln(1/\rho)$ and ρ/η are sufficiently small, $\|R_\rho^{\text{Max}}(x(\lceil T_\epsilon/(\eta\rho^2)\rceil)) - \rho^2\lambda_1(X(T_\epsilon))/2\| \leq \tilde{O}(\eta\rho^2)$. This further implies $\|R_\rho^{\text{Max}}(x(\lceil T_\epsilon/(\eta\rho^2)\rceil)) - \rho^2\lambda_1(X(\infty))/2\| \leq \epsilon\rho^2 + \tilde{O}(\eta\rho^2)$. We also have $L(x(\lceil T_\epsilon/(\eta\rho^2)\rceil)) - \inf_{x \in U'} L(x) = O(\eta^2\rho^2)$. Then we can leverage Theorem 4.3 and Theorem 4.5 to get the desired bound. $\qquad\square$

C.7 STOCHASTIC SAM: PROOF OF THEOREM 5.4

We will first prove Lemma 5.3,

Proof of Lemma 5.3. Note that $\frac{d\ell(y',y_k)}{dy'}|_{y'=f_k(p)} = 0$, we have $\nabla^2 L_k(p) = \Lambda_k(p)w_k(p)w_k(p)^T$. Also note $\nabla L_k(x)/\|\nabla L_k(x)\| = \text{sign}(\frac{d\ell(y',y_k)}{dy'}|_{y'=f_k(x)})\nabla f_k(x)/\|\nabla f_k(x)\|$. By Assumption 3.1, $\nabla^2 L(p) = \sum_k \nabla^2 L_k(p)/M = \sum_k \Lambda_k(p)w_k(p)w_k(p)^T/M$ has rank M, this implies $\forall k, \nabla f_k(p) \neq 0$, hence $\nabla f_k(x)/\|\nabla f_k(x)\|$ is in C^1 near p and we have proved our claim. $\qquad\square$

We will prove under a more general assumption.

Assumption C.2. *Total loss $L = \frac{1}{M}\sum_{k=1}^M L_k$. For each $k \in [M]$, L_k is C^4, and there exists a $(D-1)$-dimensional C^2-submanifold of $\mathbb{R}^D$, Γ_k, where for all $x \in \Gamma_k$, x is a global minimizer of L_k, $L_k(x) = 0$ and $rank(\nabla^2 L_k(x)) = 1$. Moreover, $\Gamma = \cap_{k=1}^M \Gamma_k$ for Γ defined in Assumption 3.1.*

We have U_k is open and Φ_k is in $\mathcal{C}^3$ on U_k.(from Lemma B.15 Arora et al. (2022))

Theorem C.53 shows that Setting in Theorem 5.4 satisfies assumption C.2.

Theorem C.53. *Suppose $L(x) = \sum_{k=1}^M L_k(x)/M$ and manifold Γ satisfy Assumption 3.1, and that $f_k(x) = y_k$ for every $x \in \Gamma, k \in [M]$. Then there exists $(D-1)$-dimensional $\mathcal{C}^2$ submanifolds of $\mathbb{R}^D$, such that $\cap_{k=1}^M \Gamma_k = \Gamma$ and for every $x \in \Gamma_k$ and $k \in [M]$, $f_k(x) = y_k$, that is, Γ_k is a manifold of global minimizers of L_k.*

Proof of Theorem C.53. By standard calculus, for $x \in \Gamma$, we have $\nabla^2 L(x) = \sum_{k=1}^M w_k w_k^T/M$. By Assumption 3.1,$\nabla^2 L(x)$ is full rank, this implies $w_k \neq 0$. Then we have in an open set $V(x)$ containing x, $\nabla f_k(x) \neq 0$. Then consider $V = \cup_{x \in \Gamma} V(x)$, which is an open set and in which $\nabla f_k \neq 0$. Now apply preimage theorem, we would have $\Gamma_k = \{x \in V | f_k(x) \neq y_k\}$ forms $\mathcal{C}^2$ dimensional sub-manifolds. We also easily have $\cap \Gamma_k = \Gamma$ from definitions. $\square$

The following theorems shows that stochastic SAM (16) essentially minimize trace of Hessian of loss. Analogous to the full-batch setting, we will split the trajectory into two phase.

Theorem C.54 (Phase I). *Let $\{x(t)\}$ be the iterates defined by SAM (3) and $x(0) = x_{init} \in U$, then under Assumption 3.1 and C.2 there exists a constant T_1, it holds for sufficiently small $-(\eta + \rho) \log(\eta\rho)$, we have with probability $1 - O(\sqrt{\rho})$, $\min_{-T_1 \log \rho/\eta \geq t} \|x(t) - \Phi(x(t))\| = O(-(\eta\rho + \rho^2) \log \eta\rho)$ and $\max_{-T_1 \log \rho/\eta \geq t} \|\Phi(x_{init}) - \Phi(x(t))\| = O(-(\eta + \rho) \log(\eta\rho))$.*

Theorem C.54 shows that SAM will converges to an $\tilde{O}(\eta\rho)$ neighborhood of the manifold without getting far away from $\Phi(x(0))$, where we can perform a local analysis on the trajectory of $\Phi(x(t))$.

Under Assumptions 3.1 and C.2, we have $\text{Tr}(\nabla^2 L_k(x)) = \lambda_1(\nabla^2 L_k(x))$ is differentiable for $x \in \Gamma_i$. Hence $\text{Tr}(\nabla^2 L(x)) = \sum_i \text{Tr}(\nabla^2 L_k(x))$ is also differentiable and we have (17) is well defined for some finite time T_2.

Theorem C.55 (Phase II). *Let $\{x(t)\}$ be the iterates defined by SAM (16) under Assumptions 3.1 and C.2, assuming (1) $\forall t, k, L_k(x(t)) \neq 0$, (2) $\|x(0) - \Phi(x(0))\| = O(-(\eta\rho + \rho^2) \log \eta\rho)$ and (3) $\|\Phi(x_{init}) - \Phi(x(0))\| = O(-(\eta + \rho) \log(\eta\rho))$, then for any $T_2 > 0$ till which solution of (17) exists, for sufficiently small $-(\eta + \rho) \log(\eta\rho)$, we have with probability $1 - O(\eta\rho)$, for all $\eta\rho^2 t < T_2$, $\|\Phi(x(t)) - X(\eta\rho^2 t)\| = O(-(\eta + \rho) \log \eta\rho)$ and $\|x(t) - \Phi(x(t))\| = O(-(\eta\rho + \rho^2) \log \eta\rho)$.*

In this section we will define K as $\{X(t)\}$ where X is the solution of (17).

We will prove these theorems respectively in the following sections.

C.7.1   PHASE I: PROOF OF THEOREM C.54

We will now discuss the convergence of 1-SAM to the manifold of minimizer. We will separate the dynamics into the following phase. Define ϕ as the gradient flow projection as in deterministic case.

Subphase A Gradient flow approximation, using standard approximation, we can show that $x(t)$ with high probability falls into a region K^h where the loss satisfies PL condition.

Subphase B After reaching the region, a detailed analysis will show that loss continue to decrease until $\|x(t) - \Phi(x(t))\| = \tilde{O}(\rho)$.

Subphase C Consider a quadratic approximation and we will get with high probability $x(t)$ will falls into $O(\eta\rho + \rho^2)$ neighbor of $\Phi(x(t))$.

Subphase A   This is analogous to Subphase A in Section C.6.1 and we can suppose $\exists t_1, x(t_1) \in K^h$

Subphase B   Define event $A(t)$ as $\{\nabla L(x(\tau)) \geq 4\zeta\rho, \forall \tau \leq t\}$.

We have if $\|\nabla L\left(x(t)\right)\| \geq 4\zeta\rho$

$$
\begin{aligned}
\mathbb{E}[L(x(t+1))|x(t)] &= \mathbb{E}\left[L\left(x(t) - M\eta\nabla L_k[x(t) + \rho\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|}]\right)|x(t)\right] \\
&\leq \mathbb{E}\left[L(x(t)) - M\eta\left\langle\nabla L\left(x(t)\right), \nabla L_k[x(t) + \rho\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|}]\right\rangle\right] \\
&\quad + \mathbb{E}\left[\frac{\zeta M\eta^2}{2}\|\nabla L_k[x(t) + \rho\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|}]\|^2\right] \\
&\leq L(x(t)) - M\eta\|\nabla L\left(x(t)\right)\|^2 + M\eta\rho\zeta\|\nabla L\left(x(t)\right)\| + \zeta M\eta^2\mathbb{E}[\|\nabla L_k(x(t))\|^2] + \zeta^3 M\eta^2\rho^2 \\
&\leq L(x(t)) - \frac{M\eta}{2}\|\nabla L\left(x(t)\right)\|^2 \\
&\leq L(x(t)) - \frac{M\eta\mu}{2}L(x(t))
\end{aligned}
$$

We have

$$
\mathbb{E}[L(x(t+1))\mathbf{1}A(t+1)] \leq \mathbb{E}[L(x(t+1))\mathbf{1}A(t)] \leq (1 - \frac{M\eta\mu}{2})\mathbb{E}[L(x(t))\mathbf{1}A(t)].
$$

We can then conclude that with $t_2 = \frac{2\log\frac{h^2}{16\rho^2\mu M\eta}}{M\eta\mu} + t_1$

$$
16\zeta^2\rho^2\mu\mathbb{P}\left(A(t_2+1)\right) \leq \mathbb{E}[L(x(t_2+1))\mathbf{1}A(t_2+1)] \leq (1 - \frac{M\eta\mu}{2})^{t_2-t_1}L(x(t_1)) \leq \zeta^2 h^2(1 - \frac{M\eta\mu}{2})^{t_2-t_1}
$$

We have

$$
\mathbb{P}\left(A(t_2+1)\right) \leq M\eta
$$

With an abuse of notation, suppose $\nabla L\left(x(t_2)\right) \leq 4\zeta\rho$, which implies $\|x(t_2) - \Phi(x(t_2))\| = O(\rho)$.

Subphase C   After $\|\nabla L\left(x(t_2)\right)\| \leq 4\zeta\rho$, it becomes difficult to prove the loss continue to decrease. We proceed by consider a quadratic approximation. Now consider

$$
\begin{aligned}
x(t+1) &= x(t) - M\eta\nabla L_k\left(x(t) + \rho\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|}\right) \\
&= x(t) - M\eta\nabla L_k\left(x(t)\right) - M\eta\rho\nabla^2 L_k(x(t))\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|} + O(M\eta\rho^2) \\
&= x(t) - M\eta\nabla L_k\left(x(t)\right) - M\eta\rho\Lambda_k w_k w_k^T\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|} + O(M\eta\rho^2)
\end{aligned}
$$

Iteratively define $t_{2,j}$, for $1 \leq j \leq 3$, $t_{2,1} = t_2$.

For $j \leq 2$, inductively suppose $\|x(t_{2,j}) - \Phi(x(t_{2,j}))\| \leq O(\rho^{(j+1)/2} + M\eta\rho)$, let $p_j = \Phi(x(t_{2,j}))$, further assume $\nabla L_k(\Phi(x(t_{2,j}))) = v_i v_i^T$. Further suppose N as the normal space of Γ at p_j and T as the tangent space. Define P_N and P_T as projection to the space.

We have $\|P_T(x(t_{2,j}) - p_j)\| = O((x(t_{2,j}) - p_j)^2)$.

Consider $t_{2,j} \leq t \leq t_{2,j} + \frac{\rho^{\frac{j-2}{2}}}{M\eta}$

Define event $A_j(t) = \{\|x(\tau) - p_j\| \geq \rho^{(j+2)/2}|\forall\tau \leq t\}$

We have $\|P_N(x(t) - p_j)\| = O(\rho^{k+1} + M\eta\rho^2(t - t_{2,j})) \leq O(\rho^{(j+2)/2})$

Under $A_j(t)$, we would have $\|P_T(x(t) - p_j)\| = O(\|x(t) - p_j\|)$

By Lemma C.14, we have

$$
\frac{\nabla L_k\left(x(t)\right)}{\|\nabla L_k\left(x(t)\right)\|} = s_k(t)w_k + O(\|x(t) - p_j\|)
$$

We also have

$$s_k(t) \neq sign(w_k^T(x(t) - p_j)) \Rightarrow \|w_k^T(x(t) - p_j)\| \leq \|x(t) - p_j\|^{3/2}$$

Now by Taylor Expansion,

$$
\begin{aligned}
x(t+1) - p_j &= (x(t) - p_j) - M\eta\Lambda_k w_k w_k^T(x(t) - p_j) + O(M\eta\|x(t) - p_j\|^2) \\
&\quad - M\eta\rho\Lambda_k s_k(t) w_k w_k^T w_k + O(M\eta\rho\|x(t) - p_j\|) + O(M\eta\rho^2) \\
&= (x(t) - p_j) - M\eta\Lambda_k w_k w_k^T(x(t) - p_j) - M\eta\rho\Lambda_k s_k(t) w_k w_k^T w_k + O(M\eta\rho^2)
\end{aligned}
$$

We then have

$$
\begin{aligned}
\mathbb{E}[\|x(t+1) - p_j\|^2 | x(t)] &= \|x(t) - p_j\|^2 + 2\eta^2 \sum_k \Lambda_k^2 |w_k^T(x(t) - p_j)|^2 + \eta^2\rho^2 \sum_k \Lambda_k^2 + O(M\eta\rho^2\|x(t) - p_j\|) \\
&\quad - \eta \sum_k \Lambda_k |w_k^T(x(t) - p_j)|^2 - \eta\rho \sum_k \Lambda_k s_k(t) w_k^T(x(t) - p_j)
\end{aligned}
$$

We lower bound $\sum_k \Lambda_k s_k(t) v_i^T(x(t) - p_j)$ again by Lemma C.14, there exists C such that

$$
\begin{aligned}
\sum_k \|v_i\| s_k(t) v_i^T(x(t) - p_j) &\geq \sum_k \|v_i^T(x(t) - p_j)\| - 2\kappa \sum_k \|v_i\| \|x(t) - p_j\|^{3/2} \\
&\geq C\|x(t) - p_j\|.
\end{aligned}
$$

and

$$
\begin{aligned}
\mathbb{E}[\|x(t+1) - p_j\|^2 | x(t)] &\leq \|x(t) - p_j\|^2 + 2\eta^2 C_1 \|x(t) - p_j\|^2 + \eta^2\rho^2 C_1 + \eta\rho^2 C_1 \|x(t) - p_j\| \\
&\quad - \eta C\|x(t) - p_j\|^2 - 3\eta\rho C\|x(t) - p_j\| \\
&\leq \|x(t) - p_j\|^2 - 2\eta\rho C\|x(t) - p_j\| + \eta^2\rho^2 C_1
\end{aligned}
$$

Hence we have if $\|x(t) - p_j\| \geq O(\eta\rho)$

$$\mathbb{E}[\|x(t+1) - p_j\| | x(t)] \leq \|x(t) - p_j\| - \alpha\eta\rho.$$

which implies,

$$\mathbb{E}[\|x(t+1) - p_j\| \mathbf{1} A_j(t+1)] \leq \mathbb{E}[\|x(t+1) - p_j\| \mathbf{1} A_j(t)] \leq \mathbb{E}[\|x(t) - p_j\| \mathbf{1} A_j(t)] - \alpha\eta\rho P(A_j(t))$$

Hence

$$\alpha\rho^{j/2} P(A_j(t_{2,j} + \frac{\rho^{(j-2)/2}}{\eta})) \leq \alpha\eta\rho \sum_{t=t_{2,j}}^{t=t_{2,j}+\frac{\rho^{(j-2)/2}}{\eta}} P(A_j(t)) \leq \|x(t_{2,j}) - \Phi(x(t_{2,j}))\| = O(\rho^{(j+1)/2})$$

Hence with probability $O(\sqrt{\rho})$, there exists $t_{2,j+1}$, such that $\|x(t_{2,j+1}) - p_j\| \leq O(\rho^{(j+2)/2})$, which further implies $\|x(t_{2,j+1}) - \Phi(x(t_{2,j+1}))\| \leq O(\rho^{(j+2)/2})$ using Lemma C.7.

Define $t_3 = t_{2,3}$

C.7.2 PHASE II: PROOF OF THEOREM C.55

We will inductively prove the following claim holds with probability $1 - O(\eta\rho)$,

$$
\begin{aligned}
\|\Phi(x(t)) - X(\eta\rho^2 t)\| &\leq O((\eta + \rho)\log(1/\rho)) \\
\|x(t) - \Phi(x(t))\| &= O(\rho(\eta + \rho)\log(1/\rho))
\end{aligned}
$$

To be more precise, the induction mainly consists of two parts. The first part shows that $x(t)$ will stay close to the manifold with large probability and the second part shows the direction $\Phi(x(t))$ moves.

To be more succinct with previous section, we abuse notation and suppose the iteration starts at t_3.

Part I: Convergence Near Manifold   We have $\|x(t_3) - \Phi(x(t_3))\| = O(\eta\rho + \rho^2)$. By induction hypothesis, we have $x(t) \in K^h$.

According to Lemma C.11

$$\|\Phi(x(t+1)) - \Phi(x(t))\| = O(\eta\rho^2)$$

Using the same argument in previous section, we have for sufficienly large constant A, if $A(\eta\rho + \rho^2)\log(1/\eta\rho) \geq \|x(t) - \Phi(x(t))\| \geq A(\eta\rho + \rho^2)$, then there exists constant α, B independent of A,

$$\mathbb{E}[\|x(t+1) - \Phi(x(t+1))\| | x(t)] \leq \|x(t) - \Phi(x(t))\| - \alpha\eta\rho$$
$$\|(x(t+1) - \Phi(x(t+1))) - (x(t) - \Phi(x(t)))\| \leq B\eta\rho$$

.

We then have

$$Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho)) = \sum_{\tau=t_3}^{t} Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho)$$

$$\text{and } y(\tau) < A(\eta\rho + \rho^2) \text{ and}$$
$$\forall t + 1 \geq \tau' \geq \tau + 1, y(\tau') > A(\eta\rho + \rho^2)$$

We then consider each term,

$$Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho) \text{ and } y(\tau) < A(\eta\rho + \rho^2) \text{ and } \forall t+1 \geq \tau' \geq \tau+1, y(\tau') > A(\eta\rho + \rho^2))$$
$$\leq Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho) \text{ and } \forall t+1 \geq \tau' \geq \tau+1, y(\tau') > A(\eta\rho + \rho^2)$$
$$|A(\eta\rho + \rho^2) < y(\tau+1) < (A+B)(\eta\rho + \rho^2))$$

Define a coupled process $\tilde{y}(\tau+1) = y(\tau+1)$ and

$$\tilde{y}(\tau') = \begin{cases} \|x(\tau') - \Phi(x(\tau'))\|, & \text{if } \tilde{y}(\tau'-1) = \|x(\tau'-1) - \Phi(x(\tau'-1))\| > A(\eta\rho + \rho^2) \\ \tilde{y}(\tau'-1) - \alpha\eta\rho, & \text{if otherwise} \end{cases}$$

Then clearly

$$Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho) \text{ and } \forall t+1 \geq \tau' \geq \tau+1, y(\tau') > A(\eta\rho + \rho^2)$$
$$|A(\eta\rho + \rho^2) < y(\tau+1) < (A+B)(\eta\rho + \rho^2)) \leq Pr(\tilde{y}(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho))$$

We have

$$|\tilde{y}(t+1) - \tilde{y}(t)| \leq B\eta\rho$$
$$\mathbb{E}[\tilde{y}(t+1)] - \mathbb{E}[\tilde{y}(t)] \leq -\alpha\eta\rho$$

Now applying Azuma-Hoeffding bound(Lemma C.59), we have

$$P(\tilde{y}(t+1) \geq \tilde{y}(\tau+1) - \alpha\eta\rho(t-\tau) + h) \leq P(\tilde{y}(t+1) \geq \mathbb{E}[\tilde{y}(t+1)] + h)$$
$$\leq exp(-\frac{2h^2}{(t-\tau)B^2\eta^2\rho^2})$$

Choosing $h = \alpha\eta\rho(t-\tau) - y(\tau+1) + A(\eta\rho + \rho^2)\log(1/\eta\rho)$

$$P(\tilde{y}(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho)) \leq exp(-2\frac{(\alpha\eta\rho(t-\tau) - y(\tau+1) + A(\eta\rho + \rho^2)\log(1/\eta\rho))^2}{(t-\tau)B^2\eta^2\rho^2})$$

$$\leq exp(-2\frac{(\alpha\eta\rho(t-\tau) - A(\eta\rho + \rho^2) + A(\eta\rho + \rho^2)\log(1/\eta\rho))^2}{(t-\tau)B^2\eta^2\rho^2})$$

$$\leq exp(-2\frac{(\alpha\eta\rho(t-\tau) - A(\eta\rho + \rho^2)\log(\eta\rho)/2)^2}{(t-\tau)B^2\eta^2\rho^2})$$

$$= exp(-2\frac{(\alpha(t-\tau) - A\log(\eta\rho)/2)^2}{(t-\tau)B^2})$$

$$\leq exp(-2\frac{\sqrt{A\alpha}}{B^2}\log(\eta\rho)) = \eta^{10}\rho^{10}$$

We then have

$$Pr(y(t+1) \geq A(\eta\rho + \rho^2)\log(1/\eta\rho)) \leq \eta^{10}\rho^{10}(t-t_3) \leq \eta^8\rho^8$$

Part II: Direction of $\Phi(x(t+1)) - \Phi(x(t))$ We shall do a Taylor expansion and show that

$$x(t+1) = x(t) - \eta\nabla L_k\left(x(t) + \rho\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}\right)$$

$$= x(t) - \eta\nabla L_k(x(t)) - \eta\rho\nabla^2 L_k(x(t))\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}$$

$$- \eta\rho^2\partial^2(\nabla L_k)[\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}, \frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}]/2 + O(\eta\rho^3)$$

Now by induction we have,

$$\|x(t) - \Phi(x(t))\| = \tilde{O}(\eta\rho + \rho^2)$$

, then by Lemma C.12, it implies

$$\|x(t+1) - x(t)\| = O(\eta\rho)$$

Then we have

$$\|\Phi(x(t+1)) - \Phi(x(t)) - \partial\Phi(x(t))(x(t+1) - x(t))\| \leq \xi\|x(t+1) - x(t)\|^2 = O(\eta^2\rho^2)$$

Using Lemma C.11, we have

$$\|\eta\partial\Phi(x(t))\nabla L_k(x(t))\| = O(\eta\|x(t) - \Phi(x(t))\|^2) = O(\eta^3\rho^2 + \eta\rho^4)$$

$$\|\eta\rho\partial\Phi(x(t))\nabla^2 L_k(x(t))\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}\| = O(\eta\rho\|x(t) - \Phi(x(t))\|) = \tilde{O}(\eta^2\rho^2 + \eta\rho^3)$$

Hence

$$\|\Phi(x(t+1)) - \Phi(x(t)) + \eta\rho^2\partial\Phi(x(t))\partial^2(\nabla L_k)[\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}, \frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}]/2\| = \tilde{O}(\eta^2\rho^2 + \eta\rho^3)$$

Notice finally that by Lemma C.14

$$\partial\Phi(x(t))\partial^2(\nabla L_k)[\frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}, \frac{\nabla L_k(x(t))}{\|\nabla L_k(x(t))\|}] = \partial\Phi(\Phi(x(t)))\partial^2(\nabla L_k)[w_k, w_k] + O(\|x(t) - \Phi(x(t))\|)$$

$$= P_{x,\Gamma}^\perp\Phi(x(t))\nabla(\lambda_1(\nabla^2 L_k(\Phi(x(t))))) + O(\|x(t) - \Phi(x(t))\|)$$

Hence we have

$$\Phi(x(t+1)) - \Phi(x(t)) = -\eta\rho^2 P_{x,\Gamma}^\perp\Phi(x(t))\nabla(\lambda_1(\nabla^2 L_k(\Phi(x(t)))))/2 + \tilde{O}(\eta^2\rho^2 + \eta\rho^3)$$

Notice finally,

$$\mathbb{E}_k[M P_{x,\Gamma}^\perp\Phi(x(t))\nabla(\lambda_1(\nabla^2 L_k(\Phi(x(t)))))] = P_{x,\Gamma}^\perp\Phi(x(t))\nabla(\text{Tr}(\nabla^2 L(\Phi(x(t)))))$$

Together with standard concentration bound, we would have $x(t)$ follows the Riemannian gradient flow on Γ for loss

$$\mathbb{E}_i\lambda_1\left(\nabla^2 L_k(\Phi(x(t)))\right) = \mathbb{E}_i\text{Tr}\left(\nabla^2 L_k(\Phi(x(t)))\right) = \text{Tr}\nabla^2 L(\Phi(x(t))).$$

C.7.3   PROOF OF COROLLARIES

Proof of Corollary 5.5. We will do a Taylor expansion on $\mathbb{E}_k[L_{k,\rho}^{\text{Max}}](x)$. By Theorem C.54 and C.55, for $t > \eta\rho^2 T_3'$, we have $\|X(\eta\rho^2 t) - x(t)\| = \tilde{O}(\eta + \rho)$ and $\|x(t) - \Phi(x(t))\| = \tilde{O}(\eta\rho + \rho^2)$

$$\mathbb{E}_k[R_{k,\rho}^{\text{Max}}](x) = \max_v \mathbb{E}_k[\rho v^T\nabla L_k(x) + \rho^2 v^T\nabla^2 L_k(x)v/2] + O(\rho^3)$$

Then as $L_k(x) = \tilde{O}(\eta^2\rho^2 + \rho^4)$ and $\|v^T\nabla L_k(x)\| = \tilde{O}(\eta\rho + \rho^2)$, this implies

$$\mathbb{E}_k[R^{\text{Max}}_{k,\rho}](x) = \rho^2\mathbb{E}_k \max_v v^T\nabla^2 L(x)v/2 + \tilde{O}(\eta^2\rho^2 + \rho^3)$$
$$= \rho^2\mathbb{E}_k \max_v v^T\nabla^2 L(X(\eta\rho^2 t))v/2 + \tilde{O}(\eta\rho^2 + \rho^3)$$
$$= \rho^2\text{Tr}(X(\eta\rho^2 t))/2 + \tilde{O}(\eta\rho^2)$$

$\square$

Proof of Corollary 5.6. Choose T such that $X(T)$ is sufficiently close to $X(\infty)$, such that $\text{Tr}(X(T)) \leq \text{Tr}(X(\infty)) + 2\epsilon$

By corollary 5.5, we have $\|\mathbb{E}_k[R^{\text{Max}}_{k,\rho}](x(\lceil T/(\eta\rho^2)\rceil)) - \rho^2\text{Tr}(X(T))/2\| \leq \tilde{O}(\eta\rho^2)$. This further implies $\|\mathbb{E}_k[R^{\text{Max}}_{k,\rho}](x(\lceil T/(\eta\rho^2)\rceil)) - \rho^2\text{Tr}(X(\infty))/2\| \leq \epsilon\rho^2 + \tilde{O}(\eta\rho^2)$. We also have $\|L(x(\lceil T/(\eta\rho^2)\rceil))\| = O(\eta^2\rho^2)$. Then we can leverage Theorem 4.3 and Theorem 5.1 to get the desired bound. $\square$

C.8 TECHNICAL LEMMAS

Lemma C.56 (Cor. 4.3.15, Horn & Johnson (2012)). *Let $\Sigma, \hat{\Sigma} \in R^{D\times D}$ be symmetric and non-negative with eigenvalues $\lambda_1 \geq ... \geq \lambda_D$ and $\hat{\lambda}_1 \geq ... \geq \hat{\lambda}_D$, then for any i,*

$$|\hat{\lambda}_i - \lambda_i| \leq \|\Sigma - \hat{\Sigma}\|_2$$

Definition C.57 (Unitary invariant norms). *A matrix norm $\|\cdot\|_*$ on the space of matrices in $\mathbb{R}^{p\times d}$ is unitary invariant if for any matrix $K \in \mathbb{R}^{p\times d}$, $\|UKW\|_* = \|K\|_*$ for any unitary matrices $U \in \mathbb{R}^{p\times p}, W \in \mathbb{R}^{d\times d}$.*

Theorem C.58. *[Davis-Kahan $\sin(\theta)$ theorem, Davis & Kahan (1970)] Let $\Sigma, \hat{\Sigma} \in \mathbb{R}^{p\times p}$ be symmetric, with eigenvalues $\lambda_1 \geq \ldots \geq \lambda_p$ and $\hat{\lambda}_1 \geq \ldots \geq \hat{\lambda}_p$ respectively. Fix $1 \leq r \leq s \leq p$, let $d := s - r + 1$ and let $V = (v_r, v_{r+1}, \ldots, v_s) \in \mathbb{R}^{p\times d}$ and $\hat{V} = (\hat{v}_r, \hat{v}_{r+1}, \ldots, \hat{v}_s) \in \mathbb{R}^{p\times d}$ have orthonormal columns satisfying $\Sigma v_j = \lambda_j v_j$ and $\hat{\Sigma}\hat{v}_j = \hat{\lambda}_j\hat{v}_j$ for $j = r, r+1, \ldots, s$. Define $\Delta := \min\left\{\max\{0, \lambda_s - \hat{\lambda}_{s+1}\}, \max\{0, \hat{\lambda}_{r-1} - \lambda_r\}\right\}$, where $\hat{\lambda}_0 := \infty$ and $\hat{\lambda}_{p+1} := -\infty$, we have for any unitary invariant norm $\|\cdot\|_*$,*

$$\Delta \cdot \|\sin\Theta(\hat{V}, V)\|_* \leq \|\hat{\Sigma} - \Sigma\|_*.$$

Here $\Theta(\hat{V}, V) \in \mathbb{R}^{d\times d}$, with $\Theta(\hat{V}, V)_{j,j} = \arccos\sigma_j$ for any $j \in [d]$ and $\Theta(\hat{V}, V)_{i,j} = 0$ for all $i \neq j \in [d]$. $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_d$ denotes the singular values of $\hat{V}^\top V$. $[\sin\Theta]_{ij}$ is defined as $\sin(\Theta_{ij})$.

Lemma C.59 (Azuma-Hoeffding Bound). *Suppose Z_n is a super-martingale, suppose $-\alpha \leq Z_{i+1} - Z_i \leq \beta$, then for all $n > 0, a > 0$, we have*

$$P(|Z_n - Z_0| \geq a) \leq 2exp(-a^2/(2N(\alpha + \beta)^2))$$

Lemma C.60 (Magnus (1985)). *Let $A : \mathbb{R}^D \to \mathbb{R}^{D\times D}$ be any $\mathcal{C}^1$ symmetric matrix function and $x^* \in \mathbb{R}^D$ satisfying $\lambda_1(A(x^*)) > \lambda_2(A(x^*))$ and v_1 be the top eigenvector of $A(x^*)$. It holds that $\nabla\lambda_1(A(x))|_{x=x^*} = \nabla(v_1^\top A(x)v_1)|_{x=x^*}$.*

We then present some of the technical lemmas we required to prove Lemma C.28.

Lemma C.61. *If $0 < c < \frac{b-a}{b^2}, a\sqrt{\frac{a^2+2b^2}{2(1-cb)}} \geq \frac{a^2+b^2}{2-ca-cb}$, then $a > \frac{1}{2}b, cb \leq \frac{1}{2}$*

Proof. Notice that

$$ca\sqrt{\frac{a^2+b^2}{1-cb}} \geq ca\sqrt{\frac{a^2+2b^2}{2(1-cb)}} \geq \frac{cb^2+ca^2}{2-cb-ca} \geq \frac{cb^2+ca^2}{2-cb}$$

.

So

$$\sqrt{1 - cb} + \frac{1}{\sqrt{1 - cb}} \geq \sqrt{1 + \frac{b^2}{a^2}}$$

As $c < \frac{b-a}{b^2}$, we have $1 > 1 - cb > \frac{a}{b}$.

So

$$\sqrt{\frac{a}{b}} + \sqrt{\frac{b}{a}} \geq \sqrt{1 + \frac{b^2}{a^2}}$$

The above inequality implies $a \geq \frac{1}{2}b$. As $c < \frac{b-a}{b^2}$, $cb \leq \frac{1}{2}$

$\square$

Lemma C.62. *When* $0 < a < b, 0 < c < \frac{b-a}{b^2}$, *we have*

$$cb^2 + ca^2(2 - cb - \frac{2}{3}ca) - (1 - cb)\frac{c(a^2 + b^2)}{2 - ca - cb} - ca^2(\frac{1}{2}a^2 + b^2)\frac{2 - ca - cb}{(a^2 + b^2)} \leq \frac{cb^2}{2 - cb}$$

Proof. Equivalently, we are going to prove

$$(1 - cb)b^2 \left(\frac{1}{2 - ca - cb} - \frac{1}{2 - cb} \right) + a^2\frac{1 - cb}{2 - ca - cb} + a^2(\frac{1}{2}a^2 + b^2)\frac{2 - ca - cb}{(a^2 + b^2)} \geq a^2(2 - cb - \frac{2}{3}ca)$$

Further simplifying, we only need to prove

$$\frac{(1 - cb)cab^2}{(2 - cb)(2 - ca - cb)} + a^2\frac{1 - cb}{2 - ca - cb} \geq \frac{1}{3}ca^3 + \frac{a^4}{2(a^2 + b^2)}(2 - ca - cb)$$

We have the following auxiliary inequalities,

$$(1 - cb)b > a$$

$$\frac{1 - cb}{2 - ca - cb} = \frac{1}{\frac{a+b}{b} + \frac{b-a}{1-cb}} \geq \frac{1}{\frac{a+b}{b} + \frac{b^2-ab}{a}} = \frac{ab}{a^2 + b^2} \geq \frac{a^2}{a^2 + b^2}$$

Using the above anxiliary inequalities we have

$$\frac{(1 - cb)cab^2}{(2 - cb)(2 - ca - cb)} + a^2\frac{1 - cb}{2 - ca - cb} \geq \frac{1}{3}ca^3 + \frac{a^4}{2(a^2 + b^2)}(2 - ca - cb)$$

$$\Leftarrow \frac{ca^2b}{(2 - cb)(2 - ca - cb)} + \left(1 - \frac{1}{2}(2 - ca - cb)\right)\frac{a^2(1 - cb)}{2 - ca - cb} \geq \frac{1}{3}ca^3$$

$$\Leftarrow \frac{ca^2b}{(2 - cb)(2 - ca - cb)} + \frac{ca^2(a + b)(1 - cb)}{2(2 - ca - cb)} \geq \frac{1}{3}ca^3$$

$$\Leftarrow \frac{ca^2b}{(2 - cb)(2 - ca - cb)} + \frac{ca^2b(1 - cb)}{2(2 - ca - cb)} \geq \frac{1}{3}ca^2b$$

$$\Leftarrow \frac{1}{(2 - cb)^2} + \frac{1 - cb}{2(2 - cb)} \geq \frac{1}{3}$$

$$\Leftarrow 3(1 - cb)(2 - cb) + 6 \geq 2(2 - cb)^2$$

$$\Leftarrow (cb)^2 - cb + 4 \geq 0$$

$\square$

Lemma C.63. *When* $0 < a < b, 0 < c < \frac{b-a}{b^2}, a\sqrt{\frac{a^2+2b^2}{2(1-cb)}} \geq \frac{a^2+b^2}{2-ca-cb}$, *we have*

$$cb^2 + ca^2(2 - cb - \frac{2}{3}ca) - (1 - cb)cb^2 - ca^2(\frac{1}{2}a^2 + b^2)\frac{1}{b^2} \leq \frac{cb^2}{2 - cb}$$

Proof. Equivalently, we are going to prove,

$$cb^3 + a^2(2 - cb - \frac{2}{3}ca) \le \frac{b^2}{2 - cb} + \frac{a^2(\frac{1}{2}a^2 + b^2)}{b^2}$$

$$\iff cb^3 + a^2(1 - cb - \frac{2}{3}ca) \le \frac{b^2}{2 - cb} + \frac{a^4}{2b^2}$$

We have the auxiliary inequality $\frac{1}{2 - cb} > \frac{1}{2} + \frac{cb}{4}$.

Hence

$$cb^3 + a^2(1 - cb - \frac{2}{3}ca) \le \frac{b^2}{2 - cb} + \frac{a^4}{2b^2}$$

$$\Leftarrow cb^3 + a^2(1 - cb - \frac{2}{3}ca) \le \frac{b^2}{2} + \frac{a^4}{2b^2} + \frac{cb^3}{4}$$

$$\Leftarrow c(\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3) \le \frac{b^2}{2} + \frac{a^4}{2b^2} - a^2$$

Case 1 If $\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3 \le 0$, then

$$c(\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3) \le 0 \le \frac{b^2}{2} + \frac{a^4}{2b^2} - a^2$$

Case 2 If $\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3 > 0$, then

$$c(\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3) \le \frac{b^2}{2} + \frac{a^4}{2b^2} - a^2$$

$$\Leftarrow \frac{b - a}{b^2}(\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3) \le \frac{(b^2 - a^2)^2}{2b^2}$$

$$\Leftarrow 2(\frac{3b^3}{4} - ba^2 - \frac{2}{3}a^3) \le (b - a)(b + a)^2$$

$$\Leftarrow 2(b^3 - ba^2) - (b - a)(b + a)^2 \le \frac{b^3}{2} + \frac{4a^3}{3}$$

$$\Leftarrow (b - a)(2b(a + b) - (a + b)^2) \le \frac{b^3}{2} + \frac{4a^3}{3}$$

$$\Leftarrow (b - a)^2(b + a) \le \frac{b^3}{2} + \frac{4a^3}{3}$$

Using Lemma C.61,$a > \frac{b}{2}$,$(b - a)^2(b + a) = (b^2 - a^2)(b - a) \le b^2(b - a) \le \frac{b^3}{2}$

$\square$

Lemma C.64. *When* $0 \le a \le b, 0 \le c \le \frac{b - a}{b^2}, b^2 \ge a\sqrt{\frac{a^2 + 2b^2}{2(1 - cb)}} \ge \frac{a^2 + b^2}{2 - ca - cb}$, *we have*

$$cb^2 + ca^2(2 - cb - \frac{2}{3}ca) - 2ca\sqrt{(b^2 + \frac{1}{2}a^2)(1 - cb)} \le \frac{cb^2}{2 - cb}$$

Proof. Define

$$F(a) := a^2(2 - cb - \frac{2}{3}ca) - 2a\sqrt{(b^2 + \frac{1}{2}a^2)(1 - cb)}$$

$$S_a(c, b) := \{a | 0 \le a \le b, 0 < c \le \frac{b - a}{b^2}, b^2 \ge a\sqrt{\frac{a^2 + 2b^2}{2(1 - cb)}} \ge \frac{a^2 + b^2}{2 - ca - cb}\}$$

$$a_{min}(c, b) := \inf S_a(c, b)$$

$$a_{max}(c, b) := \sup S_a(c, b) \le b - cb^2$$

Here we suppose WLOG $S_a(c, b) \ne \phi$.

Consider

$$\frac{dF(a)}{da} = 2a(2 - cb - \frac{2}{3}ca) - \frac{2}{3}ca^2 - 2\sqrt{(b^2 + \frac{1}{2}a^2)(1 - cb)} - a^2\sqrt{\frac{1 - cb}{b^2 + \frac{1}{2}a^2}}$$

$$\frac{d^2F(a)}{da^2} = 2(2 - cb - \frac{2}{3}ca) - \frac{4}{3}ca - \frac{4}{3}ca - a\sqrt{\frac{1 - cb}{b^2 + \frac{1}{2}a^2}} - 2a\sqrt{\frac{1 - cb}{b^2 + \frac{1}{2}a^2}} + \frac{a^3}{2(b^2 + \frac{1}{2}a^2)^{\frac{3}{2}}}\sqrt{1 - cb}$$

$$\geq 4 - 2cb - 4ca - 3a\sqrt{\frac{1 - cb}{b^2 + \frac{1}{2}a^2}}$$

Define $u := cb, v := \frac{a}{b}$, then $u + v \leq 1$.

$$\frac{d^2F(a)}{da^2} \geq 4 - 2u - 4uv - 3\sqrt{1 - u}\frac{1}{\sqrt{\frac{1}{2} + \frac{1}{v^2}}}$$

$$\geq 4 - 2u - 4u(1 - u) - 3\sqrt{1 - u}\frac{1}{\sqrt{\frac{1}{2} + \frac{1}{(1-u)^2}}}$$

$$\geq 4u^2 - 6u + 4 - 3\sqrt{1 - u}\frac{(1 - u)}{\sqrt{\frac{(1-u)^2}{2} + 1}}$$

As $\sqrt{\frac{(1-u)^2}{2} + 1} \geq \sqrt{\frac{(1-u)^2+1}{2}} \geq (1 - u)$, we have

$$\frac{d^2F(a)}{da^2} \geq 4u^2 - 6u + 4 - 3(1 - u) = 4u^2 + 1 - 3u > 0$$

The above inequality shows that $F(a)$ is convex w.r.t to a for $a_{min}(c, b) \leq a \leq a_{max}(c, b)$.

Hence $F(a) \leq \max\left(F(a_{min}(c, b)), F(a_{max}(c, b))\right)$

Part 1 We abuse the notation and use a_{min} a shorthand for $a_{min}(c, b)$.

We have $a_{min}\sqrt{\frac{a_{min}^2 + 2b^2}{2(1 - cb)}} = \frac{a_{min}^2 + b^2}{2 - ca_{min} - cb}$. This implies

$$2a_{min}\sqrt{(b^2 + \frac{1}{2}a_{min}^2)(1 - cb)} = (1 - cb)\frac{(a_{min}^2 + b^2)}{2 - ca_{min} - cb} + a_{min}^2(\frac{1}{2}a_{min}^2 + b^2)\frac{2 - ca_{min} - cb}{(a_{min}^2 + b^2)}$$

Hence using Lemma C.28,

$$F(a_{min}) = a_{min}^2(2 - cb - \frac{2}{3}ca_{min}) - (1 - cb)\frac{c(a_{min}^2 + b^2)}{2 - ca_{min} - cb} - ca_{min}^2(\frac{1}{2}a_{min}^2 + b^2)\frac{2 - ca_{min} - cb}{(a_{min}^2 + b^2)}$$

$$\leq \frac{1}{c}(\frac{cb^2}{2 - cb} - cb^2)$$

Part 2 We abuse the notation and use a_{max} a shorthand for $a_{max}(c, b)$.

It's not easy to see which boundary condition a_{max} satisfy, hence we will discuss by cases.

Case 1 $a_{max}\sqrt{\frac{a_{max}^2 + 2b^2}{2(1 - cb)}} = \frac{a_{max}^2 + b^2}{2 - ca_{max} - cb}$, in this case we simply redo the calculation in Part 1.

Case 2 $b^2 = a_{max}\sqrt{\frac{a_{max}^2 + 2b^2}{2(1 - cb)}}$. This implies

$$2a_{max}\sqrt{(b^2 + \frac{1}{2}a_{max}^2)(1 - cb)} = (1 - cb)b^2 + a_{max}^2(\frac{1}{2}a_{max}^2 + b^2)\frac{1}{b^2}$$

Hence using Lemma C.63,

$$F(a_{max}) = a_{max}^2(2 - cb - \frac{2}{3}ca_{max}) - (1 - cb)cb^2 - ca_{max}^2(\frac{1}{2}a_{max}^2 + b^2)\frac{1}{b^2}$$

$$\leq \frac{1}{c}(\frac{cb^2}{2 - cb} - cb^2)$$

Case 3 $cb^2 = b - a_{max}$ As $1 - cb = \frac{a_{max}}{b}$ and $b^2 \geq a_{max}\sqrt{\frac{b(a_{max}^2 + 2b^2)}{2a_{max}}}$.

This implies $a_{max}^3 + 2a_{max}b^2 - 2b^3 \leq 0 \Rightarrow a_{max} < \frac{9}{10}b$.

Define $v := \frac{a_{max}}{b}, cb = 1 - v$

Then by Lemma C.61, $\frac{1}{2} \leq v \leq \frac{9}{10}$

$$F(a_{max}) = a_{max}^2\left(2 - cb - \frac{2}{3}ca_{max}\right) - 2a_{max}\sqrt{\left(b^2 + \frac{1}{2}a_{max}^2\right)(1 - cb)}$$

$$= b^2\left(v^2(2 - (1 - v) - \frac{2}{3}(1 - v)v) - 2v\sqrt{(1 + \frac{v^2}{2})v}\right)$$

We will prove the following inequality,

$$v^2(2 - (1 - v) - \frac{2}{3}(1 - v)v) - 2v\sqrt{(1 + \frac{v^2}{2})v} \leq \frac{1}{2 - cb} - 1 = \frac{-v}{1 + v}$$

In fact we can directly show

$$v^2(1 + v) + \frac{v}{1 + v} \leq 2v\sqrt{(1 + \frac{v^2}{2})v}$$

$$\Longleftrightarrow v(1 + v) + \frac{1}{1 + v} \leq 2\sqrt{(1 + \frac{v^2}{2})v}$$

for $v \in [0.5, 0.9]$.

Hence we have

$$F(a) \leq \max\left(F(a_{min}(c, b)), F(a_{max}(c, b))\right)$$

$$\leq \frac{1}{c}\left(\frac{cb^2}{2 - cb} - cb^2\right)$$

$\square$