Federated Learning on Adaptively Weighted Nodes by Bilevel Optimization

Abstract

We propose a federated learning method with weighted nodes in which the weights can be modified to optimize the model's performance on a separate validation set. The problem is formulated as a bilevel optimization problem where the inner problem is a federated learning problem with weighted nodes and the outer problem focuses on optimizing the weights based on the validation performance of the model returned from the inner problem. A communication-efficient federated optimization algorithm is designed to solve this bilevel optimization problem. We analyze the generalization performance of the output model and identify the scenarios when our method is in theory superior to training a model locally and superior to federated learning with static and evenly distributed weights.

1 Introduction

Federated learning (FL) is an emerging technique for training a model using data distributed over a network of nodes without sharing data between nodes (Konečný et al., 2016; McMahan et al., 2017). In this paper, we focus on the case where data distributions across nodes are heterogeneous and each node aims at a model with an optimal local generalization performance. In the classical setting of FL, a globally shared model is learned by minimizing a weighted average loss across all nodes. However, given the heterogeneity of data distributions, a global model is likely to be sub-optimal for some node (Fallah et al., 2020). Alternatively, each node can train a model only using its local data, but such a local model may not generalize well neither when the volume of local data is small.

To achieve a good local generalization performance, each node can still exploit global training data through FL but, at the same time, identify and collaborate only with the nodes whose data distributions are similar or identical to its local distribution. One way to implement this strategy is to allow each node to solve its own weighted average loss minimization problem with weights designed based on the performance on a separate set of local (validation) data. Ideally, each node can learn a better model by allocating more weights on its peers whose data distribution is similar to its local distribution. In this paper, we formulate the choice of the weights as a bilevel optimization (BO) problem (Colson et al., 2005; Vicente & Calamai, 1994), which can be solved by a federated bilevel optimization algorithm, and analyze the generalization performances of the resulting model.

We consider a standard learning problem where the goal is to learn a vector of model parameters θ from a set Θ that minimizes a generalization loss. This problem can be formulated as

$$\theta^* \in \underset{\theta \in \Theta}{\arg \min} \left\{ L_0(\theta) := \mathbb{E}_{z \sim p_0} \left[l(\theta; z) \right] \right\}, \tag{P}$$

where $l(\theta; z)$ is the loss of θ on a data point z from a space $\mathcal{Z}$, and $\mathbb{E}_{z \sim p_0}$ represents the expectation taken over z when z follows an unknown ground truth distribution p_0.

Directly solving (P) is challenging as p_0 is unknown, and, typically, training data sampled from p_0 is needed for learning an approximation of θ^*. In this paper, we consider the scenario where the amount of data sampled directly from p_0 may not be sufficient to learn a good approximation of θ^*, but there exist external data distributed on K nodes that can potentially help the learning on θ^*. In particular, we denote the set of nodes by $\mathcal{K} := \{1, \dots, K\}$ and assume a training set D_k^{train} is stored in node k. We also define $D^{\text{train}} := \left\{ D_k^{\text{train}} \right\}_{k=1}^K$ and assume $|D_k^{\text{train}}| = n_k$ and $D_k^{\text{train}} = \{z_k^{(i)}\}_{i=1}^{n_k}$, where $z_k^{(i)} \in \mathcal{Z}$ is an i.i.d. sample from an unknown distribution p_k for $k \in \mathcal{K}$.

We assume node k is weighted by w_k and the vector of weights $w = (w_1, \ldots, w_K) \in [0,1]^K$ is located on the capped simplex Δ_K^b defined as

$$\Delta_K^b = \left\{ w = (w_1, \ldots, w_K) \Big| \sum_{k=1}^K w_k = 1, 0 \le w_k \le b, k \in \mathcal{K} \right\},$$

where $b \in [\frac{1}{K}, 1]$ is a user-defined parameter. The FL on weighted nodes can be formulated as

$$\widehat{\theta}(w) \in \underset{\theta \in \Theta}{\arg\min} \sum_{k=1}^K w_k \widehat{L}_k(\theta), \tag{1}$$

where $\widehat{L}_k(\theta)$ is the empirical loss of θ on D_k^{train}, namely,

$$\widehat{L}_k(\theta) := \frac{1}{n_k} \sum_{i=1}^{n_k} l(\theta; z_k^{(i)}), \quad k \in \mathcal{K}. \tag{2}$$

When some p_k's are different from p_0, w in (1) must be chosen adaptively to ensure $\widehat{\theta}(w)$ is a good approximation of θ^* in (P). To do so, we assume that there is a validation dataset D^{valid} with $|D^{\text{valid}}| = n_0 = n_{\text{valid}}$ and $D^{\text{valid}} = \{z^{(i)}\}_{i=1}^{n_0}$, where $z^{(i)} \in \mathcal{Z}$ is an i.i.d. sample from p_0. We assume D^{valid} is stored in a node called node 0 or *center*, which may or may not be a node in $\mathcal{K}$. Set D^{valid} alone may not be sufficient for learning θ^* precisely but can be used to assist the selection of w. We then propose to estimate the generalization loss of $\widehat{\theta}(w)$ using the loss on D^{valid}, i.e.,

$$\widehat{L}_0(\theta) := \frac{1}{n_0} \sum_{i=1}^{n_0} l(\theta; z^{(i)}) \tag{3}$$

and use this validation loss to guide the procedure for updating w. Presumably, when both the training and validation sets are large enough, the weights in w will be shifted towards the nodes where the data is helpful for learning θ^*. Following this idea, we formulate the *federated learning problem on adaptively weighted nodes* as the following *bilevel optimization* (BO) problem:

$$\widehat{w} \in \underset{w \in \Delta_K^b}{\arg\min} \left\{ \widehat{F}(w) := \widehat{L}_0(\widehat{\theta}(w)) \text{ s.t. } \widehat{\theta}(w) \text{ is defined as in (1)} \right\}. \tag{P}$$

In Section 4, we will present a federated optimization algorithm for solving (P). Suppose an algorithm can find the optimal solution $\widehat{w}$ of (P) and the corresponding model parameter $\widehat{\theta}(w)$. We are interested in the optimality gap of the generalization loss of $\widehat{\theta}(\widehat{w})$, namely, $L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*)$, where L_0 is defined as in (P). The *main contribution of this paper* is to establish a high-probability bound of this gap as a function of the sizes of D^{train} and D^{valid} as well as a statistical distance between p_0 and p_k's. Moreover, we compare our generalization bound with the bound achieved by learning only locally from D^{valid} and the bound achieved by solving (1) with evenly distributed weights, and identify the parameter regimes where our method is preferred in theory.

2 RELATED WORK

The work most related to ours is Chen et al. (2021a) in which the authors proposed a target-aware weighted training algorithm for cross-task learning. Although their problem is completely different from FL, the bilevel optimization model they studied contains (P) as a special case. In fact, some steps in the proofs of the generalization bounds in the current work are borrowed from Chen et al. (2021a) with some modifications. However, our work extends their results in several valuable directions. First, the generalization bound in Chen et al. (2021a) is shown for any weight w without any small or zero components, which is not necessarily the case for the optimal solution $\widehat{w}$ of (P). Second, their generalization bound contains a term of task distance whose convergence rate is not characterized. On the contrary, we show the convergence of the entire generalization bound for $\widehat{w}$ without any conditions on its components. Third, the generalization bound in Chen et al. (2021a) has a dominating term $O(1/\sqrt{n_{\text{valid}}})$, which is the same as the generalization bound obtained by directly training with the local data D^{valid}. However, we show that, when there exist identical neighbors and an error bound condition holds (Assumptions 2′ and 3), the model learned by (P) can be superior to a model trained locally when the p_k's are similar enough to (but still different from) p_0, providing an insight on when a node with insufficient data should actively seek collaboration with others.

FL has become a prominent machine learning paradigm for training models with distributed data Konečný et al. (2016); McMahan et al. (2017). Many federated optimization algorithms have been

developed for solving (1) or its expectation form (with $\widehat{L}_k$ replaced by L_k). A well-known method is the federated averaging (FedAvg) method (McMahan et al., 2017), which applies a local optimization method (e.g., stochastic gradient descend (Robbins & Monro, 1951)) to $\widehat{L}_k$ or L_k in each node and periodically aggregates the solutions from all nodes by averaging. Many variants of FedAvg and other federated learning methods have been proposed to reduce the computation and communication complexity. A partial list includes Gorbunov et al. (2021); Lee et al. (2017); Karimireddy et al. (2020); Liang et al. (2019); Li et al. (2020); Yuan & Ma (2020); Wu & Wang (2021); Zhao et al. (2021). In our setting, (1) is a sub-problem we need to solve multiple times with different w's. We then apply the Local-SVRG method by Gorbunov et al. (2021) to (1) because it has the lowest communication complexity for finite-sum problems like (1).

Most FL methods produce a globally-shared model which may not perform well on each node when data is heterogeneous across nodes. To address this challenge, many personalized FL methods, including but not limited to Smith et al. (2017); Tan et al. (2022); Fallah et al. (2020); Li & Wang (2019); Deng et al. (2020); Li et al. (2021), have been developed, where a global model is tailored using local data for a good local performance. However, many personalized FL methods use a fixed weight in (1) to obtain the global model. Such a global model may be dominated by the majority of the data distributions in the network and is hard to personalize for a minority group with unique data patterns. On the contrary, our method can produce a personalized weight so a node from the minority group can still find and collaborate with its peers.

BO has a long history of study in operations research and mathematical programming (see Colson et al. (2005); Vicente & Calamai (1994) and references therein). It recently has also been studied actively by the machine learning community because of its application in hyper-parameter optimization (Franceschi et al., 2018), model-agnostic meta learning (Finn et al., 2017; Rajeswaran et al., 2019) and data hypercleaning (Shaban et al., 2019). Many efficient optimization algorithms have been developed recently for BO, including but not limited to Chen et al. (2021c;b); Ghadimi & Wang (2018); Hong et al. (2020); Guo et al. (2021). However, these algorithms are designed for a single-machine setting and may not be communication efficient if implemented directly in a distributed environment. There are much fewer studies on BO in a distributed setting. The recent works (Li et al., 2022; Tarzanagh et al., 2022) consider a BO where both the outer and inner problems are defined with the expectations over data distributed across nodes. They analyze the communication complexity of their methods in a non-convex setting. We propose a different FL algorithm based on Local-SVRG because our problem ($\widehat{P}$) has a finite-sum structure that allows periodically going through all the data points in each node to obtain exact gradient information and achieving lower communication complexity than Li et al. (2022); Tarzanagh et al. (2022).

3 GENERALIZATION PERFORMANCE

The following assumption on ($\widehat{P}$) is made for analyzing the generalization performance of $\widehat{\theta}(\widehat{w})$ in ($\widehat{P}$) and the convergence property of the optimization algorithm for solving ($\widehat{P}$) in Section 4.

Assumption 1 (Well-behaved function). *The following statements hold. (1) $l(\theta; z) \in [0, 1]$ and $\nabla l(\theta; z)$ is ℓ_1-Lipschitz continuous in θ for any $z \in \mathcal{Z}$. (2) $\widehat{L}_k(\theta)$ and $\nabla^2 \widehat{L}_k(\theta)$ are ℓ_0 and ℓ_2-Lipschitz continuous, respectively, for $k \in \mathcal{K}$. (3) $\widehat{L}_k(\theta)$ is μ-strongly convex for $k \in \mathcal{K}$.*

These are standard regularity assumptions in recent literature on bilevel optimization (e.g. Ghadimi & Wang (2018)). Assuming the strong convexity in the lower-level problem, (1) has a unique solution so that the inclusion there can be replaced by equality. Similar to L_0 in (P), we define

$$L_k(\theta) := \mathbb{E}_{z \sim p_k}[l(\theta; z)] \text{ for } k \in \mathcal{K},$$

and we consider the following auxiliary problem

$$\mathcal{W}^* = \underset{w \in \Delta_K^b}{\arg\min} \left\{ F(w) := L_0(\theta(w)) \text{ s.t. } \theta(w) \in \underset{\theta \in \Theta}{\arg\min} \sum_{k=1}^K w_k L_k(\theta) \right\}. \tag{P_*}$$

Problem ($\widehat{P}$) can be viewed as an empirical approximation of (P_*) in both inner and outer problems.

Even if all p_k's are different from p_0, it is still possible to learn θ^* correctly by solving (P_*). A simple example on mean estimation is $\min_{w \in \Delta_2^1} \mathbb{E}(\theta(w) - z_0)^2$ s.t. $\theta(w) \in \arg\min_\theta \sum_{k=1}^2 w_k \mathbb{E}(\theta - z_k)^2$,

where z_0, z_1 and z_2 follow normal distributions $\mathcal{N}(0,1)$, $\mathcal{N}(a,1)$ and $\mathcal{N}(-a,1)$, respectively, for any $a \neq 0$. Obviously, $w^* = (0.5, 0.5)$ is the optimal weight and $\theta(w^*) = 0 = \theta^*$. Throughout the paper, we assume θ^* can be learned by solving (P$_*$), which is stated formally below.

Assumption 2 (Learnability of θ^* by (P$_*$)). *$\theta(w) = \theta^*$ for any $w \in \mathcal{W}^*$, where θ^* satsifes (P).*

Besides the situation like the aforementioned simple example, Assumption 2 holds obviously when $p_k = p_0$ for at least one $k \in \mathcal{K}$. In fact, the latter case happens when node 0 is a node in $\mathcal{K}$, so w equal to one on that node and zero on others is optimal. Moreover, we will later on provide a refined generalization performance analysis for the latter case, so we state the latter case as a separate assumption below.

Assumption 2$'$ (Existance of identical neighbors). *There exists a strict subset $\mathcal{J} \subset \mathcal{K}$ with $|\mathcal{J}| = J$ such that $p_k = p_0$ for $k \in \mathcal{J}$. Moreover,*

$$\mathcal{W}^* = \left\{ w \in \Delta_K^b \big| w_k = 0 \, for \, k \in \mathcal{K} \backslash \mathcal{J} \right\}. \tag{4}$$

The first statement in Assumption 2$'$ implies that the right-hand side of (4) is contained by the left-hand side. The second statement further assumes that they are equal. Assumption 2$'$ implies Assumption 2 because $\sum_{k=1}^{K} w_k L_k(\theta) = L_0(\theta)$ for any $\theta \in \Theta$ and any $w \in \mathcal{W}^*$ satisfying (4).

Assumption 3 (Error bound condition). *There exist $C_r > 0$ and $r \geq 1$ such that*

$$\mathrm{Dist}(w, \mathcal{W}^*) := \min_{w' \in \mathcal{W}^*} \|w - w'\| \leq C_r \left[F(w) - \min_{w \in \Delta_K^b} F(w) \right]^{1/r}. \tag{5}$$

Inequality (5) means problem (P$_*$) satisfies the *error bound condition*, which has impact on the convergence property of many optimization algorithms (Johnstone & Moulin, 2020; Yang & Lin, 2018; Lewis & Pang, 1998; Pang, 1997; Lin et al., 2020). Due to the limit of space, we refer readers to Appendix A for a practical example satisfying Assumption 3.

We are interested in the generalization performance of $\widehat{\theta}(\widehat{w})$, represented by the gap $L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*)$, as both D^{valid} and D^{train} grow. For simplicity of notation, we assume $n_k = n_{\mathrm{train}}$ for any $k \in \mathcal{K}$ for some integer $n_{\mathrm{train}} \gg n_{\mathrm{valid}}$. To facilitate the analysis, we need to introduce a few notations. Given a probability measure $\mathbb{Q}$ on $\mathcal{Z}$, let $\mathcal{H} = \{l(\theta; \cdot) : \theta \in \Theta\}$ be a pseudometric metric space equipped with the pseudometric metric $\rho_{\mathbb{Q}}$, which is the L_2 distance metric with respect $\mathbb{Q}$, i.e., $\rho_{\mathbb{Q}}(l, l') := \sqrt{\int_{\mathcal{Z}} (l(z) - l'(z))^2 d\mathbb{Q}(z)}$ for $l, l' \in \mathcal{H}$. The ball with radius $\epsilon > 0$ centered at $l \in \mathcal{H}$ is defined as $B_\epsilon(l) := \{l' \in \mathcal{H} | \rho_{\mathbb{Q}}(l, l') \leq \epsilon\}$. Let $\mathcal{N}(\mathcal{H}; \rho_{\mathbb{Q}}, \epsilon)$ be the ϵ-covering number of $\mathcal{H}$ with respect to $\rho_{\mathbb{Q}}$, i.e., $\mathcal{N}(\mathcal{H}; \rho_{\mathbb{Q}}, \epsilon) := \min\{m | \exists l_1, \ldots, l_m \in \mathcal{H}, \mathcal{H} \subset \cup_{i=1}^{m} B_\epsilon(l_i)\}$.

Following Chen et al. (2021a), we make the following assumption on $\mathcal{N}(\mathcal{H}; \rho_{\mathbb{Q}}, \epsilon)$, which is important for analyzing the generalization performance (Koltchinskii, 2006; Kakade et al., 2008).

Assumption 4. *There exist $C_{\mathcal{H}} > 0$ and $\nu_{\mathcal{H}} > 0$ such that, for any probability measure $\mathbb{Q}$ on $\mathcal{Z}$,*

$$\mathcal{N}(\mathcal{H}; \rho_{\mathbb{Q}}, \epsilon) \leq (C_{\mathcal{H}}/\epsilon)^{\nu_{\mathcal{H}}}, \quad \forall \epsilon > 0. \tag{6}$$

With the these assumptions, we obtain the following theorems whose proofs are in Appendix C.

Theorem 1 (Bound independent of statistical distance). *Suppose Assumptions 1, 2 and 4 hold. There exists a universal constant[1] $C_g > 0$ such that, with a probability of at least $1 - \delta$,*

$$L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \leq C_g \left(\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{valid}} \right)^{\frac{1}{2}} + C_g \frac{\ell_0}{\sqrt{\mu}} \left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{train}/(Kb^2)} \right)^{\frac{1}{4}}. \tag{7}$$

Under the same assumptions,[2] the generalization bound by Chen et al. (2021a) becomes

$$\begin{aligned} L_0(\widehat{\theta}(w)) - L_0(\theta^*) \leq & C_g' \left(\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\mathrm{valid}}} \right)^{\frac{1}{2}} + C_g' \frac{\sqrt{\beta}}{\sqrt{\mu}} \left(\frac{\nu_{\mathcal{H}} + K \log(K) + \log(1/\delta)}{K n_{\mathrm{train}}} \right)^{\frac{1}{4}} \\ & + L_0(\theta(w)) - L_0(\theta^*) \end{aligned} \tag{8}$$

---

[1]We define a universal constant as a constant that does not depend on any parameter of the problem except $C_{\mathcal{H}}$ and C_r. This definition is made only to simply the constant factors in our bounds.

[2]A (ρ, C_ρ)-transferable assumption is needed in Chen et al. (2021a), which also holds in our case with $\rho = 2$ because of the Lipschitz continuity and strong convexity assumed in Assumption 1.

for a universal constant C'_g and any w satisfying $\beta^{-1} \leq w_k/w_j \leq \beta$ with $k \neq j$ for some $\beta > 0$. However, it is likely that the optimal solution w^* has zero components (e.g., when $p_k = p_0$ for some k). If so, β on the right-hand side of (8) needs to be arbitrarily large for $\widehat{w}(\approx w^*)$ in (P̂) to satisfy the aforementioned condition. Moreover, when $w = \widehat{w}$, the convergence of the last term $L_0(\theta(w)) - L_0(\theta^*)$ in (8) is not characterized in Chen et al. (2021a). On the contrary, Theorem 1 holds without any assumption on $\widehat{w}$ (zero components are allowed), does not depend on β and provides a generalization bound converging in every term.[3] When $b = c/K$ with a constant $c \geq 1$, the right-hand side of (7) improves the first two terms on the right-hand side of (8) by a $\log(K)$ term.

The bounds (7) and (8) may both be dominated by the $O(1/\sqrt{n_{\text{valid}}})$ term, which is the same as the generalization bound achieved simply by learning locally with D^{valid} (see Proposition 1 in Appendix D). However, if Assumption 3 holds and Assumption 2 is strengthened to Assumption 2', we can establish a generalization bound different from Theorem 1 which suggests that (P̂) can still outperform local training when the following statistical distance between p_0 and p_k's is small:

$$G := \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K}} \left(L_0(\theta) - L_k(\theta) \right)^2}. \tag{9}$$

Theorem 2 (Bound dependent on statistical distance). *Suppose Assumptions 1, 2', 3 and 4 hold. There exists universal constants $C_e > 0$ and $C_w > 0$ such that, with a probability of at least $1 - 3\delta$,*

$$\text{Dist}(\widehat{w}, \mathcal{W}^*) \leq \varepsilon(n_{\text{valid}}, n_{\text{train}}) := C_w \left(\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\text{valid}}} \right)^{\frac{1}{2r}} + C_w \frac{\ell_0}{\sqrt{\mu}} \left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{\text{train}}/(Kb^2)} \right)^{\frac{1}{4r}} \tag{10}$$

and

$$L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \leq C_e \sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{N_\varepsilon}} + C_e \frac{\varepsilon(n_{\text{valid}}, n_{\text{train}})(K - J)}{b\sqrt{N_\varepsilon}} + 2\varepsilon(n_{\text{valid}}, n_{\text{train}})G, \tag{11}$$

where $N_\varepsilon = \frac{n_{\text{train}}}{b^2 J + \varepsilon^2(n_{\text{valid}}, n_{\text{train}})(K-J)}$ and G is defined in (9).

Note that $N_\varepsilon = \Theta(n_{\text{train}})$. Based on the decreasing rate of $\varepsilon(n_{\text{valid}}, n_{\text{train}})$ in (10), we simplify (11) by only showing the bounds in terms of n_{valid}, n_{train} and G for a clear comparison with local training.

Corollary 1. *Suppose the assumptions of Theorem 2 hold and n_{valid} and n_{train} are large enough such that $\varepsilon(n_{valid}, n_{train})$ defined in (10) satisfies $\varepsilon(n_{valid}, n_{train}) \leq \frac{b\sqrt{J}}{K-J}$. With a probability of at least $1 - 3\delta$, we have $L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \leq O\left(1/n_{train}^{\frac{1}{2}} + G \cdot \left(1/n_{valid}^{\frac{1}{2r}} + 1/n_{train}^{\frac{1}{4r}} \right) \right).$*

When $G = o(1/n_{\text{valid}}^{\frac{1}{2} - \frac{1}{2r}})$ and $G = o(n_{\text{train}}^{\frac{1}{4r}}/n_{\text{valid}}^{\frac{1}{2}})$, the bound in Corollary 1 becomes $o(1/n_{\text{valid}}^{\frac{1}{2}})$, meaning that method (P̂) has a better generalization guarantee than training locally. Since G is small in this case, a natural question is whether optimizing the weight in (P̂) is still needed because the FL with equally weighted nodes may already have a good performance with respect to p_0. However, we show in Proposition 2 in Appendix D that (P̂) is still preferred to FL with equally weighted nodes for any G. We show the impacts of b, J and K through Corollary 2 in Appendix C.

4 FEDERATED BILEVEL OPTIMIZATION ALGORITHM

Although our main focus is the generalization performance of (P̂), we present a federated optimization algorithm for (P̂) based on the existing techniques by Gorbunov et al. (2021) and Ghadimi & Wang (2018). Different from a single-level optimization problem, the outer objective $\widehat{F}(w)$ in (P̂) depends implicitly on w through the inner optimal solution $\widehat{\theta}(w)$, which makes the exact gradient $\nabla \widehat{F}(w)$ difficult to compute. A commonly used solution is to exploit implicit function as shown in the following lemma, which is from Lemma 2.1 and 2.2 in Ghadimi & Wang (2018).

Lemma 1. *Under Assumption 1, $\nabla \widehat{F}(w)$ is ℓ_F-Lipschitz continuous with*

$$\ell_F := \left(\frac{2\ell_0 \ell_1}{\mu} + \frac{\ell_2 \ell_0^2}{\mu^2} \right) \frac{\sqrt{K} \ell_0}{\mu} + \frac{K \ell_1 \ell_0^2}{\mu^2}, \tag{12}$$

---

[3]As a by-product of our analysis, we show in (38) that $L_0(\theta(\widehat{w})) - L_0(\theta^*)$ also satisfies (7).

Algorithm 1: Local-SVRG method for (16): `Local-SVRG`$(\{f_{k,i}\}, w, \gamma, \tau, q, T)$

---

1 **Input**: functions $\{f_{k,i}\}$, weight w, initial vector $x^{(0)} \in \mathbb{R}^d$, learning rate γ, communication period $\tau \geq 1$, probability q of updating reference point, and the total number of iterations T
2 $x_k^{(0)} = x^{(0)}, y_k^{(0)} = x^{(0)}, k = 1, \ldots, K$
3 **for** $t = 0, 1, \ldots, T-1$ **do**
4   **for** $k = 1, \ldots, K$ in parallel **do**
5   Choose i_k from $\{1, \ldots, n_k\}$ uniformly at random
6          $g_k^{(t)} = \nabla f_{k,i_k}(x_k^{(t)}) - \nabla f_{k,i_k}(y_k^{(t)}) + \nabla f_k(y_k^{(t)})$
7          $y_k^{(t+1)} = x_k^{(t)}$ with probability q and $y_k^{(t+1)} = y_k^{(t)}$ with probability $1 - q$
8   **if** $t + 1 \mod \tau = 0$ **then**
9              $x_k^{(t+1)} = x^{(t+1)} := \sum_{k=1}^K w_k \left(x_k^{(t)} - \gamma g_k^{(t)} \right)$
10   **else**
11              $x_k^{(t+1)} = x_k^{(t)} - \gamma g_k^{(t)}$
12   **end**
13   **end**
14 **end**
15 **Return:** $\bar{x}^{(T)} = U_T^{-1} \sum_{t=0}^T u_t x^{(t)}$ with $u_t = (1 - \min\{\gamma\mu, q/4\})^{-(t+1)}$ and $U_T = \sum_{t=0}^T u_t$.

---

and $\nabla \widehat{F}(w) = (\nabla_k \widehat{F}(w))_{k=1,\ldots,K}$, where $\nabla_k \widehat{F}(w)$ is the partial derivative of $\widehat{F}$ w.r.t. w_k and

$$\nabla_k \widehat{F}(w) = -\nabla \widehat{L}_k(\widehat{\theta}(w))^\top \left(\sum_{k=1}^K w_k \nabla^2 \widehat{L}_k(\widehat{\theta}(w)) \right)^{-1} \nabla \widehat{L}_0(\widehat{\theta}(w)). \tag{13}$$

By Lemma 1, computing $\nabla_k \widehat{F}(w)$ requires solving (1) exactly and taking the inverse of the Hessian matrix in (13), both of which are challenging. Hence, for a given w, we will find an approximate solution of (1), denoted by $\bar{\theta}(w)(\approx \widehat{\theta}(w))$, and approximate the matrix inversion in (13) by solving a strongly convex quadratic program. In particular, we will approximate $\nabla \widehat{F}(w)$ by

$$\bar{\nabla} \widehat{F}(w) := (\bar{\nabla}_k \widehat{F}(w))_{k=1,\ldots,K} \quad \text{with} \quad \bar{\nabla}_k \widehat{F}(w) := -\nabla \widehat{L}_k(\bar{\theta}(w))^\top \bar{h}, \tag{14}$$

$$\text{where} \qquad \bar{h} \approx \arg\min_h \tfrac{1}{2} h^\top \left(\sum_{k=1}^K w_k \nabla^2 \widehat{L}_k(\bar{\theta}(w)) \right) h - h^\top \nabla \widehat{L}_0(\bar{\theta}(w)). \tag{15}$$

Both (1) and (15) can be written as a *distributed finite-sum* minimization on K weighted nodes:

$$\min_{x \in \mathbb{R}^d} f(x) := \sum_{k=1}^K w_k f_k(x), \quad \text{where} \quad f_k(x) = \tfrac{1}{n_k} \sum_{i=1}^{n_k} f_{k,i}(x), \quad k = 1, \ldots, K. \tag{16}$$

When $f_{k,i}(\theta) = l(\theta; z_k^{(i)})$, (16) becomes (1). When $f_{k,i}(h) = \tfrac{1}{2} h^\top \nabla^2 l(\bar{\theta}(w); z_k^{(i)}) h - h^\top \nabla \widehat{L}_0(\bar{\theta}(w))$, (16) becomes (15).

With this observation, we apply Local-SVRG by Gorbunov et al. (2021) to the aforementioned two instances (16) to obtain $\bar{\theta}(w)$ and $\bar{h}$, which are used to construct the approximate gradient $\bar{\nabla} \widehat{F}(w)$ in (14). Then we update w using $\bar{\nabla} \widehat{F}(w)$ based on the accelerated bilevel approximation method (ABA) by Ghadimi & Wang (2018). We choose the combination of Local-SVRG and the ABA methods because it leads to the lowest communication complexity in literature for solving $(\widehat{P})$. We formally present this approach in Algorithms 1 and 2. Recall that we have assumed D^{valid} is stored in node 0, which is called center in Algorithm 2.

In each iteration of Algorithm 2, in addition to the communication within Local-SVRG, constantly many rounds of communication are needed to exchange $\theta^{(s)}, h^{(s)} \nabla \widehat{L}_0(\theta^{(s)})$ and $\nabla \widehat{L}_k(\theta^{(s)})$ between the center and node k. We present the communication complexity of Algorithm 2 which can be proved by adapting the analysis in Gorbunov et al. (2021) and Ghadimi & Wang (2018) to our setting. The proofs are deferred to Sections E.1 and E.2.

Theorem 3. *Suppose Assumption 1 holds and $\widehat{F}(w)$. Let $R := \max_{w \in \Delta_K^b} \|\widehat{\theta}(w)\|$ and*

$$\gamma_0 := \min \left\{ \frac{3}{80\ell_1}, \frac{1}{\ell_1 \sqrt{5e(\tau-1)[6(\tau-1) + 8 + 16/(1-q)]}}, \frac{q}{4\mu} \right\}. \tag{17}$$

Algorithm 2: Federated Learning Method for Bilevel Optimization ($\widehat{P}$)

---

1 **Input**: initial weight $w^{(0)}$, learning rate η, training data D_k^{train} for $k \in \mathcal{K}$, validation data D^{valid}, the number of outer iterations S, parameters (γ, τ, q) for `Local-SVRG`, and the number of inner iterations T_s for $s = 0, \ldots, S-1$

2 Set $w_{\text{ag}}^{(0)} = w^{(0)}$

3 **for** $s = 0, 1, \ldots, S-1$ **do**

4 $\quad$ $w_{\text{md}}^{(s)} = \frac{2}{s+2} w^{(s)} + \frac{s}{s+2} w_{\text{ag}}^{(s)}$

5 $\quad$ Compute $\bar{\nabla} \widehat{F}(w_{\text{md}}^{(s)})$ as follows:

6 $\quad\quad$ Set $f_{k,i}(\theta) = l(\theta; z_k^{(i)}), \quad i = 1, \ldots, n_k, \quad k = 1, \ldots, K$

7 $\quad\quad$ Compute $\theta^{(s)} = \text{Local-SVRG}(\{f_{k,i}\}, w_{\text{md}}^{(s)}, \gamma, \tau, q, T_s)$ and send it to each node.

8 $\quad\quad$ Compute $\nabla \widehat{L}_0(\theta^{(s)})$ at center and send it to each node.

9 $\quad\quad$ Set $f_{k,i}(h) = \frac{1}{2} h^\top \nabla^2 l(\theta^{(s)}; z_k^{(i)}) h - h^\top \nabla \widehat{L}_0(\theta^{(s)}), \quad i = 1, \ldots, n_k, \quad k = 1, \ldots, K$

10 $\quad\quad$ Compute $h^{(s)} = \text{Local-SVRG}(\{f_{k,i}\}, w_{\text{md}}^{(s)}, \gamma, \tau, q, T_s)$ and send it to each node.

11 $\quad\quad$ Each node computes $\nabla \widehat{L}_k(\theta^{(s)})$ in parallel and send it to the center.

12 $\quad\quad$ Set $\bar{\nabla}_k \widehat{F}(w_{\text{md}}^{(s)}) = -\nabla \widehat{L}_k(\theta^{(s)})^\top h^{(s)}$ for $k = 1, \ldots, K$

13 $\quad$ $w^{(s+1)} = \arg\min_{w \in \Delta_K^b} \left\langle \bar{\nabla} \widehat{F}(w_{\text{md}}^{(s)}), w \right\rangle + \frac{2}{\eta(s+1)} \| w - w^{(s)} \|^2$

14 $\quad$ $w_{\text{ag}}^{(s+1)} = \arg\min_{w \in \Delta_K^b} \left\langle \bar{\nabla} \widehat{F}(w_{\text{md}}^{(s)}), w \right\rangle + \frac{1}{2\eta} \| w - w_{\text{md}}^{(s)} \|^2$

15 **end**

16 **Return:** $w_{\text{ag}}^{(S)}$

Suppose $\eta = \frac{1}{3\ell_F}$ in Algorithm 2 with ℓ_F defined as in (12). There exist constants A_1, A_2 and A_3 that only depend on ℓ_0, ℓ_1, ℓ_2, μ, R, q and K but not on τ such that the following statements hold.

- *Suppose $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0 \mu} \ln\left(\frac{A_1(s+1)^4}{\gamma_0}\right)$. Algorithm 2 finds an ϵ-optimal solution of ($\widehat{P}$) with $\tilde{O}\left(\epsilon^{-0.5}\right)$ rounds of communication.*

- *Suppose $\tau > 1$, $\gamma = \frac{1}{M_s}$ and $T_s = \mu^{-1} M_s \ln\left(M_s^3\right)$, where*

$$M_s = \max\left\{ 1/\gamma_0, (s+1)^2 \sqrt{[A_1 + A_2(\tau-1) + A_3(\tau-1)^2]} \right\}, s = 0, 1, \ldots. \quad (18)$$

Algorithm 2 finds an ϵ-optimal solution of ($\widehat{P}$) with $\tilde{O}\left(\epsilon^{-1.5}\right)$ rounds of communication.

When $\widehat{F}$ in ($\widehat{P}$) is non-convex, we aim at finding an ϵ-stationary point of ($\widehat{P}$). Following Ghadimi & Wang (2018), we apply a standard proximal gradient method to ($\widehat{P}$) based on the approximate gradient $\bar{\nabla} \widehat{F}(w)$ in (14). This method and its analysis are standard and we include them in Section E.3 due to the limit of space. In Remark 1 in Section E.3, we also show that the complexity of our method is lower than those of Li et al. (2022) and Tarzanagh et al. (2022).

5 NUMERICAL EXPERIMENT

In this section, we demonstrate the performance of our methods on image classification tasks. We compare our method, denoted by **Bi-level**, against four baselines, including (1) **Local-train**, which solves $\min_{\theta \in \Theta} \widehat{L}_0(\theta)$ locally; (2) **FedAvg** (McMahan et al., 2017), which solves (1) with $w_k = 1/K$; (3) **Ditto** (Li et al., 2021); and (4) **pFedMe** (T Dinh et al., 2020). Ditto and pFedMe are two personalized FL methods. We apply all methods to train a convolutional neural network (CNN) on multiple image datasets: Fashion-MNIST (Xiao et al., 2017), MNIST (Deng, 2012), CIFAR-10 (Krizhevsky et al., 2009) and downsampled 32×32 ImageNet (Chrabaszcz et al., 2017). We denote Fashion-MNIST and downsampled ImageNet by F-MNist and DS-ImageNet, respectively. See Appendix F for more details on the CNN and the computing environment we use.

Figure 1: Comparison in test accuracy for the minority group vs number of synchronizations.

We use a mini batch of size 50 to construct the stochastic gradients in all methods. In the Bi-level method, Algorithm 3 is applied to $(\widehat{P})$ with $b = 1/3$ and five epochs are performed within each call of Local-SVRG (i.e., $T_s = 5n_{\text{train}}/50$). We set $q = 1/50$, choose γ from $\{0.05, 0.02, 0.01\}$ when solving (1) and from $\{0.0005, 0.0002, 0.0001\}$ when solving (15), choose τ from $\{10, 20\}$, and η from $\{0.025, 0.02, 0.015\}$. We choose the combination that produces the highest validation accuracy after five outer iterations. SVRG is applied to $\min_{\theta \in \Theta} \widehat{L}_0(\theta)$ in Local-train and Local-SVRG is applied to (1) with $w_k = 1/K$ in FedAvg. Parameters τ, q, and η in FedAvg and Local-train are set the same as in our method. Ditto is implemented by setting $S_t = \mathcal{K}$, $r = \tau$, $s = 25$ and $\eta_g = \eta_l = \gamma$ in Algorithm 2 in Li et al. (2021), where τ and γ are set the same as in our method. Similar to Li et al. (2021), we choose λ in Ditto from $\{0.05, 0.1, 0.2\}$ to maximize the validation accuracy after five outer iterations. pFedMe is implemented by setting $\beta = 1$, $\delta = 0.005$, $R = \tau$ and $\eta = \gamma$ in Algorithm 1 in T Dinh et al. (2020) with τ and γ set the same as in our method. Each subproblem in pFedMe is solved by gradient descend with a maximum iterations of 20. Like Li et al. (2021), we choose λ in pFedMe from $\{5, 10, 15\}$ to maximize the validation accuracy after five outer iterations.

We set $\mathcal{K} = \{1, \ldots, 15\}$ (i.e., $K = 15$) and partition it into two groups, a minority group $\mathcal{J}_m = \{1, \ldots, 5\}$ and a majority group $\mathcal{J}_M = \{6, \ldots, 15\}$. We then generate D_k^{train} for $k \in \mathcal{K}$ by randomly sampling data from the training sets with some artificial distributions, such that the data distributions (i.e., p_k's) are the same within each group but different between groups. In particular, we create the data distributions of $\mathcal{J}_m$ and $\mathcal{J}_M$ under four different settings. In **Setting 1**, we create two different distributions over the classes and use them to sample D_k^{train} with $k \in \mathcal{J}_m$ and $k \in \mathcal{J}_M$, respectively. In **Setting 2**, **Setting 3** and **Setting 4**, we first sample data in the same way as Setting 1 and, additionally, we permute the class labels among a few classes in D_k^{train} with $k \in \mathcal{J}_M$ under Setting 2, rotate each image in D_k^{train} with $k \in \mathcal{J}_M$ by 90 degrees in the same but random direction under Setting 3, and do both under Setting 4. This creates nodes with different levels of heterogeneity.

To compare the performances of the methods on both groups, we conduct two sets of experiments under each setting, one with p_0 being the distribution of $\mathcal{J}_m$ (i.e., $\mathcal{J} = \mathcal{J}_m$) and the other with p_0 being the distribution of $\mathcal{J}_M$ (i.e., $\mathcal{J} = \mathcal{J}_M$). D^{valid} is then sampled from p_0. For out-of-sample evaluation, we generate testing data by sampling from the testing set of each dataset using

Figure 2: How w evolves during the Bi-level method under Setting 1.

distribution p_0 described above under each setting. We denote the testing set by D^{test} and let $n_{\text{test}} = |D^{\text{test}}|$. We repeat all experiments five times using different random seeds. The values of n_{valid}, n_{train}, n_{test} and the details of data generation are presented in Sections F.1, F.2 and F.3.

We plot the test (top-1) accuracy each method obtains during iterations for the minority group in Figure 1, where the horizontal axis represents the number of synchronizations, i.e., the rounds of communications the method performs. Since Local-train does not require any communication, we just plot a horizontal line positioned at its final accuracy. Due to space limit, we present the accuracy for the majority group in Figure 6 in Section F.4. We also report the same results in Figure 7 and Figure 8 but the horizontal axis there represents the cumulative number of data points each method processes in parallel. In each figure, we show the confidence intervals of the curves as shaded areas.

According to Figure 1, our Bi-level method performs better than the four benchmarks on the minority group on all datasets under all settings. Local-train does not perform well because it only gets access to a small amount of data. The poor performance of FedAvg is because of the heterogeneity we created across nodes. In fact, FedAvg is even worse than Local-train in many cases, especially in Settings 2, 3 and 4 where the heterogeneity is high. This is consistent with the findings in literature. Although Ditto and pFedMe are designed for heterogeneous nodes, they still use a fixed weight on each node to train a global model, which may not provide a good starting point for personalization due to the high heterogeneity. In fact, their performances drop more or less as the data heterogeneity increases from Setting 1 to Settings 2, 3 and 4. On the contrary, by updating the weights, our method filters the information in the network and help the node in the minority group to find its similar peers and produce a good model through intra-group collaboration. Comparing Figure 1 with Figure 6, we find that the performances of FedAvg, Ditto and pFedMe are improved on the majority group. This is again because they utilize the information aggregated from all nodes, which is in favor of the majority. However, our method perform similarly on both groups and is still overall the best for the majority group. Similar phenomena are found in Figure 7 and Figure 8.

In addition, we also plot in Figure 2 how the weight w_k for each node evolves during the Bi-level method under Setting 1. We show the results when p_0 is the distribution of the majority and the minority groups separately. In each case, we call the nodes in $\mathcal{J}$ similar nodes (to node 0) and call the others dissimilar nodes. According to Figure 2, our method successfully detects similar nodes in both cases and increases their weights but decreases the weights of dissimilar nodes. We present the weights under Settings 2, 3 and 4 in Section F.4. Similar phenomenons are observed.

6 CONCLUSION

We propose a FL approach on a network with weighted nodes and develop a federated bilevel optimization algorithm to optimize the weights based on the model's performance on a validation set. We analyze the generalization performance of the resulting model and identify the scenarios where our method theoretically outperforms training with local data and FL with even weights.

REFERENCES

Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities: A nonasymptotic theory of independence*. Oxford university press, 2013.

Shuxiao Chen, Koby Crammer, Hangfeng He, Dan Roth, and Weijie J Su. Weighted training for cross-task learning. *arXiv preprint arXiv:2105.14095*, 2021a.

Tianyi Chen, Yuejiao Sun, and Wotao Yin. Closing the gap: Tighter analysis of alternating stochastic gradient methods for bilevel problems. *Advances in Neural Information Processing Systems*, 34, 2021b.

Tianyi Chen, Yuejiao Sun, and Wotao Yin. A single-timescale stochastic bilevel optimization method. *arXiv preprint arXiv:2102.04671*, 2021c.

Patryk Chrabaszcz, Ilya Loshchilov, and Frank Hutter. A downsampled variant of imagenet as an alternative to the cifar datasets. *arXiv preprint arXiv:1707.08819*, 2017.

Benoît Colson, Patrice Marcotte, and Gilles Savard. Bilevel programming: A survey. *4or*, 3(2): 87–107, 2005.

Jia Deng, Wei Dong, Richard Socher, Li-Jia Li, Kai Li, and Li Fei-Fei. Imagenet: A large-scale hierarchical image database. In *2009 IEEE conference on computer vision and pattern recognition*, pp. 248–255. Ieee, 2009.

Li Deng. The mnist database of handwritten digit images for machine learning research. *IEEE Signal Processing Magazine*, 29(6):141–142, 2012.

Yuyang Deng, Mohammad Mahdi Kamani, and Mehrdad Mahdavi. Adaptive personalized federated learning. *arXiv preprint arXiv:2003.13461*, 2020.

Alireza Fallah, Aryan Mokhtari, and Asuman Ozdaglar. Personalized federated learning: A meta-learning approach. *arXiv preprint arXiv:2002.07948*, 2020.

Chelsea Finn, Pieter Abbeel, and Sergey Levine. Model-agnostic meta-learning for fast adaptation of deep networks. In *International conference on machine learning*, pp. 1126–1135. PMLR, 2017.

Luca Franceschi, Paolo Frasconi, Saverio Salzo, Riccardo Grazzi, and Massimiliano Pontil. Bilevel programming for hyperparameter optimization and meta-learning. In *International Conference on Machine Learning*, pp. 1568–1577. PMLR, 2018.

Saeed Ghadimi and Mengdi Wang. Approximation methods for bilevel programming. *arXiv preprint arXiv:1802.02246*, 2018.

Pinghua Gong and Jieping Ye. Linear convergence of variance-reduced stochastic gradient without strong convexity. *arXiv preprint arXiv:1406.1102*, 2014.

Eduard Gorbunov, Filip Hanzely, and Peter Richtárik. Local sgd: Unified theory and new efficient methods. In *International Conference on Artificial Intelligence and Statistics*, pp. 3556–3564. PMLR, 2021.

Bin Gu, De Wang, Zhouyuan Huo, and Heng Huang. Inexact proximal gradient methods for non-convex and non-smooth optimization. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 32, 2018.

Zhishuai Guo, Yi Xu, Wotao Yin, Rong Jin, and Tianbao Yang. On stochastic moving-average estimators for non-convex optimization. *arXiv preprint arXiv:2104.14840*, 2021.

Mingyi Hong, Hoi-To Wai, Zhaoran Wang, and Zhuoran Yang. A two-timescale framework for bilevel optimization: Complexity analysis and application to actor-critic. *arXiv preprint arXiv:2007.05170*, 2020.

Patrick R Johnstone and Pierre Moulin. Faster subgradient methods for functions with hölderian growth. *Mathematical Programming*, 180(1):417–450, 2020.

Sham M Kakade, Karthik Sridharan, and Ambuj Tewari. On the complexity of linear prediction: Risk bounds, margin bounds, and regularization. *Advances in neural information processing systems*, 21, 2008.

Sai Praneeth Karimireddy, Satyen Kale, Mehryar Mohri, Sashank Reddi, Sebastian Stich, and Ananda Theertha Suresh. Scaffold: Stochastic controlled averaging for federated learning. In *International Conference on Machine Learning*, pp. 5132–5143. PMLR, 2020.

Vladimir Koltchinskii. Local rademacher complexities and oracle inequalities in risk minimization. *The Annals of Statistics*, 34(6):2593–2656, 2006.

Jakub Konečný, H Brendan McMahan, Felix X Yu, Peter Richtárik, Ananda Theertha Suresh, and Dave Bacon. Federated learning: Strategies for improving communication efficiency. *arXiv preprint arXiv:1610.05492*, 2016.

Alex Krizhevsky, Geoffrey Hinton, et al. Learning multiple layers of features from tiny images. 2009.

Jason D Lee, Qihang Lin, Tengyu Ma, and Tianbao Yang. Distributed stochastic variance reduced gradient methods by sampling extra data with replacement. *The Journal of Machine Learning Research*, 18(1):4404–4446, 2017.

Adrian S Lewis and Jong-Shi Pang. Error bounds for convex inequality systems. In *Generalized convexity, generalized monotonicity: recent results*, pp. 75–110. Springer, 1998.

Daliang Li and Junpu Wang. Fedmd: Heterogenous federated learning via model distillation. *arXiv preprint arXiv:1910.03581*, 2019.

Junyi Li, Feihu Huang, and Heng Huang. Local stochastic bilevel optimization with momentum-based variance reduction. *arXiv preprint arXiv:2205.01608*, 2022.

Tian Li, Anit Kumar Sahu, Manzil Zaheer, Maziar Sanjabi, Ameet Talwalkar, and Virginia Smith. Federated optimization in heterogeneous networks. *Proceedings of Machine Learning and Systems*, 2:429–450, 2020.

Tian Li, Shengyuan Hu, Ahmad Beirami, and Virginia Smith. Ditto: Fair and robust federated learning through personalization. In *International Conference on Machine Learning*, pp. 6357–6368. PMLR, 2021.

Xianfeng Liang, Shuheng Shen, Jingchang Liu, Zhen Pan, Enhong Chen, and Yifei Cheng. Variance reduced local sgd with lower communication complexity. *arXiv preprint arXiv:1912.12844*, 2019.

Qihang Lin, Runchao Ma, Selvaprabu Nadarajah, and Negar Soheili. First-order methods for convex constrained optimization under error bound conditions with unknown growth parameters. *arXiv preprint arXiv:2010.15267*, 2020.

Brendan McMahan, Eider Moore, Daniel Ramage, Seth Hampson, and Blaise Aguera y Arcas. Communication-efficient learning of deep networks from decentralized data. In *Artificial intelligence and statistics*, pp. 1273–1282. PMLR, 2017.

Jong-Shi Pang. Error bounds in mathematical programming. *Mathematical Programming*, 79(1): 299–332, 1997.

Aravind Rajeswaran, Chelsea Finn, Sham M Kakade, and Sergey Levine. Meta-learning with implicit gradients. *Advances in neural information processing systems*, 32, 2019.

Herbert Robbins and Sutton Monro. A stochastic approximation method. *The annals of mathematical statistics*, pp. 400–407, 1951.

Amirreza Shaban, Ching-An Cheng, Nathan Hatch, and Byron Boots. Truncated back-propagation for bilevel optimization. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pp. 1723–1732. PMLR, 2019.

Virginia Smith, Chao-Kai Chiang, Maziar Sanjabi, and Ameet S Talwalkar. Federated multi-task learning. *Advances in neural information processing systems*, 30, 2017.

Canh T Dinh, Nguyen Tran, and Josh Nguyen. Personalized federated learning with moreau envelopes. *Advances in Neural Information Processing Systems*, 33:21394–21405, 2020.

Alysa Ziying Tan, Han Yu, Lizhen Cui, and Qiang Yang. Towards personalized federated learning. *IEEE Transactions on Neural Networks and Learning Systems*, 2022.

Davoud Ataee Tarzanagh, Mingchen Li, Christos Thrampoulidis, and Samet Oymak. Fednest: Federated bilevel, minimax, and compositional optimization. *arXiv preprint arXiv:2205.02215*, 2022.

Luis N Vicente and Paul H Calamai. Bilevel and multilevel programming: A bibliography review. *Journal of Global optimization*, 5(3):291–306, 1994.

Hongda Wu and Ping Wang. Fast-convergent federated learning with adaptive weighting. *IEEE Transactions on Cognitive Communications and Networking*, 7(4):1078–1088, 2021.

Han Xiao, Kashif Rasul, and Roland Vollgraf. Fashion-mnist: a novel image dataset for benchmarking machine learning algorithms. *arXiv preprint arXiv:1708.07747*, 2017.

Tianbao Yang and Qihang Lin. Rsg: Beating subgradient method without smoothness and strong convexity. *The Journal of Machine Learning Research*, 19(1):236–268, 2018.

Honglin Yuan and Tengyu Ma. Federated accelerated stochastic gradient descent. *Advances in Neural Information Processing Systems*, 33:5332–5344, 2020.

Haoyu Zhao, Zhize Li, and Peter Richtárik. Fedpage: A fast local stochastic gradient method for communication-efficient federated learning. *arXiv preprint arXiv:2108.04755*, 2021.

A   EXAMPLES SATISFYING ASSUMPTION 3

We consider (P_*) in the setting of linear regression. Consider data $z = (x, y)$, where $x \in \mathbb{R}^d$ is a feature vector and $y \in \mathbb{R}$ is a continuous target variable, and consider the quadratic loss $l(\theta; z) = \frac{1}{2}(x^\top \theta - y)^2$. We assume x in all nodes, including node 0 (center) and the nodes in $\mathcal{K}$, follows the same distribution, and matrix $\mathbb{E}[xx^\top]$ is non-singular. Moreover, we assume that there is a vector $\theta_k^* \in \mathbb{R}^d$ associated to node k, and y in node k is generated as $y = x^\top \theta_k^* + \epsilon_k$ for $k = 0, 1, \ldots, K$, where ϵ_k is a zero-mean random noise indepedent of x. In this problem, we have

$$L_k(\theta) = \tfrac{1}{2}\mathbb{E}\big[(x^\top \theta - y)^2\big] = \tfrac{1}{2}\mathbb{E}\big[(x^\top \theta - x^\top \theta_k^* - \epsilon_k)^2\big] = \tfrac{1}{2}(\theta - \theta_k^*)^\top \mathbb{E}\big[xx^\top\big](\theta - \theta_k^*) + \tfrac{1}{2}\mathbb{E}\big[\epsilon_k^2\big].$$

We can easily show that $\theta(w)$ in (P_*) has the closed form $\theta(w) = \sum_{k=1}^K w_k \theta_k^*$, which means

$$F(w) = L_0(\theta(w)) = \tfrac{1}{2}(\sum_{k=1}^K w_k \theta_k^* - \theta_0^*)^\top \mathbb{E}\big[xx^\top\big](\sum_{k=1}^K w_k \theta_k^* - \theta_0^*) + \tfrac{1}{2}\mathbb{E}\big[\epsilon_0^2\big].$$

This is a quadratic function of w over the polyhedral set Δ_K^b and thus satisfies the error bound condition (5) with $r = 2$ according to Lemma 1 in Gong & Ye (2014).

B   TECHNICAL LEMMAS

In this section, we provide some technical lemmas with proofs which are necessary for establishing the main theorems. The main steps in the proofs of Lemma 2 and 3 are borrowed from Chen et al. (2021a). However, the generalization bound in Theorem 3.1 in Chen et al. (2021a) is for a w satisfying $\beta^{-1} \le w_k/w_j \le \beta$ with $k \ne j$ for some $\beta > 0$, and their bound increases with β. When applied to $w = \hat{w}$ with zero or nearly zero components (that happens when $p_k = p_0$ for some k), such a β is very large or equals infinity. Therefore, we make necessary changes in the proofs to extend the results for a generic w and a w that is ε-away from $\mathcal{W}^*$ (see Lemma 3) where the components can be nearly zero. These extensions are important for proofing our main theorems.

Lemma 2. *Suppose Assumptions 1 and 4 hold. There exists a universal constant $C_0 > 0$ such that, with a probability of at least $1 - \delta$,*

$$\sup_{\theta \in \Theta} \left| L_0(\theta) - \widehat{L}_0(\theta) \right| \le C_0 \sqrt{\frac{\nu_\mathcal{H} + \log(1/\delta)}{n_{valid}}}.$$

Proof. For simplicity of notation, we write n_{valid} as n in this proof. Let

$$G_{\text{valid}}(D^{\text{valid}}) := \sup_{\theta \in \Theta} \left[L_0(\theta) - \widehat{L}_0(\theta) \right] \quad \text{and} \quad G'_{\text{valid}}(D^{\text{valid}}) := \sup_{\theta \in \Theta} \left[\widehat{L}_0(\theta) - L_0(\theta) \right].$$

Consider any $i \in \{1, 2, \ldots, n\}$. Let D_i^{valid} be the same as D^{valid} except that $z^{(i)}$ is replaced by another data point $z'^{(i)}$ sampled from p_0. Recall (3). We have

$$\left| G_{\text{valid}}(D^{\text{valid}}) - G_{\text{valid}}(D_i^{\text{valid}}) \right|$$

$$= \left| \sup_{\theta \in \Theta} \left[L_0(\theta) - \widehat{L}_0(\theta) \right] - \sup_{\theta \in \Theta} \left[L_0(\theta) - \widehat{L}_0(\theta) + \frac{1}{n}\left(l(\theta; z^{(i)}) - l(\theta; z'^{(i)}) \right) \right] \right| \le \frac{1}{n},$$

where the inequality is because the loss is in $[0, 1]$ (Assumption 1). This inequality means we can apply the McDiarmid's inequality to obtain that, for any $\epsilon > 0$,

$$\mathbb{P}\left(G_{\text{valid}}(D^{\text{valid}}) \ge \mathbb{E}[G_{\text{valid}}(D^{\text{valid}})] + \epsilon \right) \le \exp(-2\epsilon^2 n),$$

or equivalently, with a probability of at least $1 - \delta$,

$$G_{\text{valid}}(D^{\text{valid}}) \le \mathbb{E}[G_{\text{valid}}(D^{\text{valid}})] + \sqrt{\frac{\log(1/\delta)}{2n}}.$$

Next, we apply the standard symmetrization argument by introducing a ghost dataset

$$D_{\text{ghost}}^{\text{valid}} := \left\{ z'^{(i)} \right\}_{i=1}^n,$$

which is independent of D^{valid} and sampled from p_0. Let $\{\sigma_i\}_{i=1}^n$ be Rademacher random variables. We have

$$
\begin{aligned}
\mathbb{E}[G_{\text{valid}}(D^{\text{valid}})] =& \mathbb{E}\left[\sup_{\theta \in \Theta}\left[L_0(\theta) - \widehat{L}_0(\theta)\right]\right] = \mathbb{E}\left[\sup_{\theta \in \Theta}\left[\mathbb{E}\left[L(\theta; D_{\text{ghost}}^{\text{valid}})\right] - L(\theta; D^{\text{valid}})\right]\right] \\
\leq& \mathbb{E}\left[\sup_{\theta \in \Theta}\left[L(\theta; D_{\text{ghost}}^{\text{valid}}) - L(\theta; D^{\text{valid}})\right]\right] = \mathbb{E}\left[\sup_{\theta \in \Theta}\frac{1}{n}\sum_{i=1}^n \sigma_i\left(l(\theta; z'^{(i)}) - l(\theta; z^{(i)})\right)\right] \\
\leq& 2\mathbb{E}\left[\sup_{\theta \in \Theta}\frac{1}{n}\sum_{i=1}^n \sigma_i l(\theta; z^{(i)})\right] = 2\mathbb{E}\widehat{R}_n(\mathcal{H}),
\end{aligned}
$$

where $\widehat{R}_n(\mathcal{H}) = \mathbb{E}\left[\sup_{\theta \in \Theta}\frac{1}{n}\sum_{i=1}^n \sigma_i l(\theta; z^{(i)})\Big|D^{\text{valid}}\right]$ and $\mathcal{H} = \{l(\theta; \cdot) : \theta \in \Theta\}$.

Let $M_\theta := \frac{1}{\sqrt{n}}\sum_{i=1}^n \sigma_i l(\theta; z^{(i)})$ for $\theta \in \Theta$. By Hoeffding's Lemma, we have for any $\theta, \theta' \in \Theta$,

$$
\begin{aligned}
\mathbb{E}\left[\exp\left(\lambda(M_\theta - M_{\theta'})\right)\Big|D^{\text{valid}}\right] &= \prod_{i=1}^n \mathbb{E}\left[\exp\left(\frac{\lambda}{\sqrt{n}}\sigma_i\left(l(\theta; z^{(i)}) - l(\theta'; z^{(i)})\right)\right)\Big|D^{\text{valid}}\right] \\
&\leq \prod_{i=1}^n \exp\left(\frac{\lambda^2}{2n}\left(l(\theta; z^{(i)}) - l(\theta'; z^{(i)})\right)^2\right) = \exp\left(\frac{\lambda^2}{2}\mathrm{d}^2(\theta, \theta')\right),
\end{aligned}
$$

where

$$
\mathrm{d}(\theta, \theta') = \sqrt{\sum_{i=1}^n \frac{1}{n}\left(l(\theta; z^{(i)}) - l(\theta'; z^{(i)})\right)^2} \leq 1
$$

is the L_2-distance between mappings $l(\theta; \cdot)$ and $l(\theta'; \cdot)$ with respect to the empirical distribution over D^{valid} and is a pseudometric in $\mathcal{H}$. Hence, by Dudley's entropy integral inequality (see Corollary 13.2 in Boucheron et al. (2013)), there exists a universal constant C_d such that

$$
\widehat{R}_n(\mathcal{H}) = \frac{1}{\sqrt{n}}\mathbb{E}\left[\sup_{\theta \in \Theta}M_\theta\Big|D^{\text{valid}}\right] \leq \frac{C_d}{\sqrt{n}}\int_0^1 \sqrt{\log\left(\mathcal{N}(\mathcal{H}; \mathrm{d}; \epsilon)\right)}d\epsilon
$$

According to Assumption 4, we have

$$
\mathbb{E}[G_{\text{train}}] \leq 2\widehat{R}_n(\mathcal{H}) \leq \frac{2C_d}{\sqrt{n}}\int_0^1 \sqrt{\nu_{\mathcal{H}}\log\left(\frac{C_{\mathcal{H}}}{\epsilon}\right)}d\epsilon
$$

Hence, with a probability of at least $1 - \delta$,

$$
G_{\text{valid}}(D^{\text{valid}}) \leq \frac{2C_d}{\sqrt{n}}\int_0^1 \sqrt{\nu_{\mathcal{H}}\log\left(\frac{C_{\mathcal{H}}}{\epsilon}\right)}d\epsilon + \sqrt{\frac{\log(1/\delta)}{2n}}.
$$

Applying the same argument to $G'_{\text{valid}}(D^{\text{valid}})$, we can show that, with a probability of at least $1 - \delta$

$$
G'_{\text{valid}}(D^{\text{valid}}) \leq \frac{2C_d}{\sqrt{n}}\int_0^1 \sqrt{\nu_{\mathcal{H}}\log\left(\frac{C_{\mathcal{H}}}{\epsilon}\right)}d\epsilon + \sqrt{\frac{\log(1/\delta)}{2n}}.
$$

By a union bound, we have, with a probability of at least $1 - \delta$

$$
\sup_{\theta \in \Theta}\left|L_0(\theta) - \widehat{L}_0(\theta)\right| \leq \frac{2C_d}{\sqrt{n}}\int_0^1 \sqrt{\nu_{\mathcal{H}}\log\left(\frac{C_{\mathcal{H}}}{\epsilon}\right)}d\epsilon + \sqrt{\frac{\log(2/\delta)}{2n}},
$$

which completes the proof.

$\square$

Given $\varepsilon > 0$, we define

$$\mathcal{W}_\varepsilon^* := \left\{ w \in \Delta_K^b \,\middle|\, \text{Dist}(w, \mathcal{W}^*) \leq \varepsilon \right\} \tag{19}$$

$$N_\varepsilon := \frac{n_{\text{train}}}{b^2 J + \varepsilon^2 (K - J)}. \tag{20}$$

Lemma 3. *Suppose Assumptions 1,2 and 4 hold. There exists a universal constant $C_a > 0$ such that, with a probability of at least $1 - \delta$,*

$$\sup_{\theta \in \Theta, w \in \Delta_K^b} \left| \sum_{k=1}^K w_k \left[\widehat{L}_k(\theta) - L_k(\theta) \right] \right| \leq C_a \sqrt{\frac{\nu_\mathcal{H} + K + \log(1/\delta)}{n_{train}/(Kb^2)}}. \tag{21}$$

Suppose Assumptions 1, 2′, 3 and 4 hold. There exists a universal constant $C_a' > 0$ such that, with a probability of at least $1 - \delta$,

$$\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left| \sum_{k=1}^K w_k \left[\widehat{L}_k(\theta) - L_k(\theta) \right] \right| \leq C_a' \left(\sqrt{\frac{\nu_\mathcal{H} + J + \log(1/\delta)}{N_\varepsilon}} + \frac{\varepsilon(K - J)}{b\sqrt{N_\varepsilon}} \right). \tag{22}$$

Proof. We prove (22) first. Suppose Assumptions 1, 2′, 3 and 4 hold. By Assumptions 2′, we have $w_j^* = 0$ for $w \in \mathcal{W}^*$ and $j \in \mathcal{K} \backslash \mathcal{J}$, which means $w_j \leq \sqrt{\sum_{k \in \mathcal{K} \backslash \mathcal{J}} w_k^2} \leq \text{Dist}(w, \mathcal{W}^*) \leq \varepsilon$ for any $w \in \mathcal{W}_\varepsilon^*$ and $j \in \mathcal{K} \backslash \mathcal{J}$.

Let

$$G_{\text{train}}(D^{\text{train}}) := \sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left[L_k(\theta) - \widehat{L}_k(\theta) \right] \right\} \tag{23}$$

$$G'_{\text{train}}(D^{\text{train}}) := \sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left[\widehat{L}_k(\theta) - L_k(\theta) \right] \right\}. \tag{24}$$

Consider an index $j \in \{1, 2, \ldots, K\}$ and $i \in \{1, 2, \ldots, n_j\}$. Let $D_{j,i}^{\text{train}}$ be the same as D^{train} except that $z_j^{(i)}$ is replaced by another data point $z_j'^{(i)}$ sampled from p_j. Recall (2). We have

$$\left| G_{\text{train}}(D^{\text{train}}) - G_{\text{train}}(D_{j,i}^{\text{train}}) \right|$$

$$= \left| \sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left[L_k(\theta) - \widehat{L}_k(\theta) \right] \right\} \right.$$

$$\left. - \sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left[L_k(\theta) - \widehat{L}_k(\theta) \right] + \frac{w_j}{n_j} \left(l\left(\theta; z_j^{(i)}\right) - l\left(\theta; z_j'^{(i)}\right) \right) \right\} \right|$$

$$\leq \frac{w_j}{n_j} \leq \begin{cases} \frac{b}{n_j} & j \in \mathcal{J} \\ \frac{\varepsilon}{n_j} & j \in \mathcal{K} \backslash \mathcal{J}. \end{cases} \tag{25}$$

With this inequality, we can apply the McDiarmid's inequality to show that, for any $\epsilon > 0$,

$$\mathbb{P}\left(G_{\text{train}}(D^{\text{train}}) \geq \mathbb{E}[G_{\text{train}}(D^{\text{train}})] + \epsilon \right) \leq \exp\left(\frac{-2\epsilon^2 n_{\text{train}}}{b^2 J + \varepsilon^2 (K - J)} \right),$$

which implies that, with a probability of at least $1 - \delta$,

$$G_{\text{train}}(D^{\text{train}}) \leq \mathbb{E}[G_{\text{train}}(D^{\text{train}})] + \sqrt{\frac{\log(1/\delta)}{N_\varepsilon}}. \tag{26}$$

Next, we apply the standard symmetrization strategy by introducing a ghost dataset $D_{\text{ghost}}^{\text{train}} := \left\{ D_{k,\text{ghost}}^{\text{train}} \right\}_{k=1}^K$ where

$$D_{k,\text{ghost}}^{\text{train}} := \left\{ z_k'^{(i)} \right\}_{i=1}^{n_k} \quad \text{for } k = 1, \ldots, K$$

is a dataset independent of D_k^{train} sampled from p_k. Let $\{\sigma_{k,i}\}_{i=1}^{n_k}$ for $k = 1, \ldots, K$ be Rademacher random variables. We have

$$
\begin{aligned}
\mathbb{E}[G_{\text{train}}(D^{\text{train}})] &= \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left[L_k(\theta) - \widehat{L}_k(\theta) \right] \right\}\right] \\
&= \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left(\mathbb{E}\left[L_k(\theta; D_{k,\text{ghost}}^{\text{train}}) \right] - \widehat{L}_k(\theta) \right) \right\}\right] \\
&\leq \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left\{ \sum_{k=1}^K w_k \left(L_k(\theta; D_{k,\text{ghost}}^{\text{train}}) - \widehat{L}_k(\theta) \right) \right\}\right] \\
&= \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \sum_{k=1}^K \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} \left(l(\theta; z_k'^{(i)}) - l(\theta; z_k^{(i)}) \right)\right] \\
&\leq 2\mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \sum_{k=1}^K \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} l(\theta; z_k^{(i)})\right] = 2\mathbb{E}\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \mathcal{W}_\varepsilon^*), \quad (27)
\end{aligned}
$$

where $\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \mathcal{W}_\varepsilon^*) := \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \sum_{k=1}^K \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} l(\theta; z_k^{(i)}) \Big| D^{\text{train}}\right]$ and $\mathcal{H} = \left\{ l(\theta; \cdot) : \theta \in \Theta \right\}$.

Let $M_{\theta,w} := \sqrt{N_\varepsilon} \sum_{k=1}^K \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} l(\theta; z_k^{(i)})$ for $\theta \in \Theta$ and $w \in \mathcal{W}_\varepsilon^*$. Hence, by Hoeffding's Lemma, we have

$$
\begin{aligned}
&\mathbb{E}\left[\exp\left(\lambda(M_{\theta,w} - M_{\theta',w'}) \right) \Big| D^{\text{train}}\right] \\
&= \prod_{k=1}^K \prod_{i=1}^{n_k} \mathbb{E}\left[\exp\left(\lambda \sqrt{N_\varepsilon} \frac{\sigma_{k,i}}{n_k} \left(w_k l(\theta; z_k^{(i)}) - w_k' l(\theta'; z_k^{(i)}) \right) \right) \Big| D^{\text{train}}\right] \\
&\leq \prod_{k=1}^K \prod_{i=1}^{n_k} \exp\left(\frac{\lambda^2 N_\varepsilon}{2 n_k^2} \left(w_k l(\theta; z_k^{(i)}) - w_k' l(\theta'; z_k^{(i)}) \right)^2 \right) \\
&= \exp\left(\frac{\lambda^2}{2} \mathrm{d}^2(\theta, w, \theta', w') \right), \quad (28)
\end{aligned}
$$

where

$$
\mathrm{d}(\theta, w, \theta', w') = \sqrt{\sum_{k=1}^K \sum_{i=1}^{n_k} \frac{N_\varepsilon}{n_k^2} \left(w_k l(\theta; z_k^{(i)}) - w_k' l(\theta'; z_k^{(i)}) \right)^2} \leq \sqrt{\sum_{k \in \mathcal{J}} \frac{N_\varepsilon b^2}{n_{\text{train}}} + \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \frac{N_\varepsilon \varepsilon^2}{n_{\text{train}}}} = 1
$$

is a pseudo distance metric between $(l(\theta; \cdot), w)$ and $(l(\theta'; \cdot), w')$ in $\mathcal{H} \times \mathcal{W}_\varepsilon^*$. Hence, by Dudley's entropy integral inequality (see Corollary 13.2 in Boucheron et al. (2013)), there exists a universal constant C_d such that

$$
\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \mathcal{W}_\varepsilon^*) = \frac{1}{\sqrt{N_\varepsilon}} \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} M_{\theta,w} \Big| D^{\text{train}}\right] \leq \frac{C_d}{\sqrt{N_\varepsilon}} \int_0^1 \sqrt{\log\left(\mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon) \right)} d\epsilon,
$$

where $\mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon)$ is the ϵ-covering number of $\mathcal{H} \times \mathcal{W}_\varepsilon^*$ w.r.t. d.

We next need to bound $\mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon)$. Note that

$$\mathrm{d}^2(\theta, w, \theta', w')$$

$$= \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{N_\varepsilon}{n_k^2} \left(w_k l(\theta; z_k^{(i)}) - w_k' l(\theta'; z_k^{(i)}) \right)^2$$

$$\leq \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{2N_\varepsilon}{n_k^2} w_k^2 \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2 + \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{2N_\varepsilon}{n_k^2} (w_k - w_k')^2 l^2(\theta'; z_k^{(i)})$$

$$\leq \sum_{k \in \mathcal{J}} \sum_{i=1}^{n_k} \frac{2N_\varepsilon}{n_k^2} b^2 \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2 + \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \sum_{i=1}^{n_k} \frac{2N_\varepsilon}{n_k^2} \varepsilon^2 \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2$$

$$+ \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{2N_\varepsilon}{n_k^2} (w_k - w_k')^2$$

$$\leq \frac{2N_\varepsilon}{n_{\text{train}}} \left(\sum_{k \in \mathcal{J}} \sum_{i=1}^{n_{\text{train}}} \frac{b^2}{n_{\text{train}}} \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2 + \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \sum_{i=1}^{n_{\text{train}}} \frac{\varepsilon^2}{n_{\text{train}}} \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2 \right)$$

$$+ \frac{2N_\varepsilon}{n_{\text{train}}} \|w - w'\|^2. \tag{29}$$

We then define a probability measure

$$\mathbb{Q} = \frac{N_\varepsilon}{n_{\text{train}}} \left(\sum_{k \in \mathcal{J}} \sum_{i=1}^{n_{\text{train}}} \frac{b^2}{n_{\text{train}}} \delta_{z_k^{(i)}} + \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \sum_{i=1}^{n_{\text{train}}} \frac{\varepsilon^2}{n_{\text{train}}} \delta_{z_k^{(i)}} \right) \tag{30}$$

on $\mathcal{Z}$, where $\delta_{z_k^{(i)}}$ is a point mass at $z_k^{(i)}$. Then, we can construct an ϵ-cover for $\mathcal{H} \times \mathcal{W}_\varepsilon^*$ w.r.t. d by taking the Cartesian product of an $\frac{\epsilon}{2}$-cover for $\mathcal{H}$ w.r.t. distance metric $\rho_{\mathbb{Q}}(l, l') = \sqrt{\int_{\mathcal{Z}} (l(z) - l'(z))^2 d\mathbb{Q}(z)}$ for $l, l' \in \mathcal{H}$ and a $\sqrt{\frac{n_{\text{train}}}{N_\varepsilon}} \frac{\epsilon}{2}$-cover for $\mathcal{W}_\varepsilon^*$ w.r.t. the Euclidean distance. According to Assumption 4, the former has a cardinality of $(2C_{\mathcal{H}}/\epsilon)^{\nu_{\mathcal{H}}}$. To construct the latter, we create a $\sqrt{\frac{n_{\text{train}}}{(J+1)N_\varepsilon}} \frac{\epsilon}{2}$-cover for $[0, b]$ corresponding to a coordinate in $\mathcal{J}$ and create a $\sqrt{\frac{n_{\text{train}}}{(K-J)(J+1)N_\varepsilon}} \frac{\epsilon}{2}$-cover for $[0, \varepsilon]$ corresponding to a coordinate in $\mathcal{K} \setminus \mathcal{J}$. (Recall that $w_j \leq \varepsilon$ for $j \in \mathcal{K} \setminus \mathcal{J}$.) Then we take the Cartesian product of these K one-dimensional covers and project it to $\mathcal{W}_\varepsilon^*$. This provides a $\sqrt{\frac{n_{\text{train}}}{N_\varepsilon}} \frac{\epsilon}{2}$-cover for $\mathcal{W}_\varepsilon^*$ with a cardinality of

$$\left(\left\lceil \frac{b\sqrt{(J+1)N_\varepsilon}}{\sqrt{n_{\text{train}}}\epsilon} \right\rceil \right)^J \left(\left\lceil \frac{\varepsilon\sqrt{(K-J)(J+1)N_\varepsilon}}{\sqrt{n_{\text{train}}}\epsilon} \right\rceil \right)^{K-J} \leq \left(\frac{2}{\epsilon} \right)^J \left(\left\lceil \frac{2\varepsilon\sqrt{K-J}}{b\epsilon} \right\rceil \right)^{K-J}, \tag{31}$$

where the inequality is because $N_\varepsilon \leq \frac{n_{\text{train}}}{b^2 J}$ by the definition of N_ε. This implies

$$\mathbb{E}[G_{\text{train}}(D^{\text{train}})] \leq 2\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \mathcal{W}_\varepsilon^*) \leq \frac{2C_d}{\sqrt{N_\varepsilon}} \int_0^1 \sqrt{\log \mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon)} d\epsilon$$

$$= O\left(\frac{1}{\sqrt{N_\varepsilon}} \int_0^1 \sqrt{(\nu_{\mathcal{H}} + J) \log\left(\frac{1}{\epsilon}\right) + (K-J) \log\left(\left\lceil \frac{2\varepsilon\sqrt{K-J}}{b\epsilon} \right\rceil\right)} d\epsilon \right)$$

$$= O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J}{N_\varepsilon}} \right) + O\left(\sqrt{\frac{K-J}{N_\varepsilon}} \frac{2\varepsilon\sqrt{K-J}}{b} \right) = O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J}{N_\varepsilon}} \right) + O\left(\frac{\varepsilon(K-J)}{b\sqrt{N_\varepsilon}} \right), \tag{32}$$

where the first equality is because

$$\mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon) \leq \left(\frac{2C_{\mathcal{H}}}{\epsilon} \right)^{\nu_{\mathcal{H}}} \times \left(\frac{2}{\epsilon} \right)^J \left(\left\lceil \frac{2\varepsilon\sqrt{K-J}}{b\epsilon} \right\rceil \right)^{K-J}$$

according to Assumption 4 and (31) and the second equality is by changing variable ϵ to $\frac{b\epsilon}{2\varepsilon\sqrt{K-J}}$ in the integral and the fact that $\left\lceil \frac{1}{\epsilon} \right\rceil = 0$ when $\epsilon > 1$.

Combining (32) with (26), we have that, with a probability of at least $1 - \delta$,

$$G_{\text{train}}(D^{\text{train}}) \leq O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{N_\varepsilon}} \right) + O\left(\frac{\varepsilon(K - J)}{b\sqrt{N_\varepsilon}} \right).$$

Applying the same argument to $G'_{\text{valid}}(D^{\text{valid}})$, we can show that the same inequality as above holds for $G'_{\text{train}}(D^{\text{train}})$ with a probability of at least $1 - \delta$. By a union bound, we have, with a probability of at least $1 - \delta$

$$\sup_{\theta \in \Theta, w \in \mathcal{W}_\varepsilon^*} \left| \sum_{k=1}^{K} w_k \left[\widehat{L}_k(\theta) - L_k(\theta) \right] \right| \leq O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{N_\varepsilon}} \right) + O\left(\frac{\varepsilon(K - J)}{b\sqrt{N_\varepsilon}} \right),$$

which completes the proof (22).

Next we prove (21). Since the proof is similar to (22), we will mainly elaborate the parts that are different. Suppose Assumptions 1,2 and 4 hold. We define $G_{\text{train}}(D^{\text{train}})$ and $G'_{\text{train}}(D^{\text{train}})$ the same as in (23) and (24) except that $\mathcal{W}_\varepsilon^*$ is replaced by the entire domain Δ_K^b. Following the same proof of (25), we have

$$\left| G_{\text{train}}(D^{\text{train}}) - G_{\text{train}}(D_{j,i}^{\text{train}}) \right| \leq \frac{w_j}{n_j} \leq \frac{b}{n_j} \quad \text{for all } i \in \{1, 2, \ldots, n_j\} \text{ and } j \in \mathcal{K}.$$

Then the McDiarmid's inequality implies that, with a probability of at least $1 - \delta$,

$$G_{\text{train}}(D^{\text{train}}) \leq \mathbb{E}[G_{\text{train}}(D^{\text{train}})] + \sqrt{\frac{Kb^2 \log(1/\delta)}{2n_{\text{train}}}}. \tag{33}$$

By replacing $\mathcal{W}_\varepsilon^*$ with Δ_K^b in the proof of (27), we can show that

$$\mathbb{E}[G_{\text{train}}(D^{\text{train}})] \leq 2\mathbb{E}\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \Delta_K^b)$$

where $\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \Delta_K^b) := \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \Delta_K^b} \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} l(\theta; z_k^{(i)}) \middle| D^{\text{train}} \right].$

Let N_b defined as (20) with ε replaced by b, i.e., $N_b = n_{\text{train}}/(Kb^2)$. Let $M_{\theta,w} := \sqrt{N_b} \sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{\sigma_{k,i} w_k}{n_k} l(\theta; z_k^{(i)})$ for $\theta \in \Theta$ and $w \in \Delta_K^b$. With $\mathcal{J}$ replaced by $\emptyset$ and N_b replaced by N_ε in the proof of (28), we have

$$\mathbb{E}\left[\exp\left(\lambda(M_{\theta,w} - M_{\theta',w'}) \right) \middle| D^{\text{train}} \right] = \exp\left(\frac{\lambda^2}{2} \mathrm{d}^2(\theta, w, \theta', w') \right),$$

where

$$\mathrm{d}(\theta, w, \theta', w') = \sqrt{\sum_{k=1}^{K} \sum_{i=1}^{n_k} \frac{N_b}{n_k^2} \left(w_k l(\theta; z_k^{(i)}) - w_k' l(\theta'; z_k^{(i)}) \right)^2} \leq \sqrt{\sum_{k \in \mathcal{K}} \frac{N_b b^2}{n_{\text{train}}}} = 1$$

is a pseudo distance metric between $(l(\theta; \cdot), w)$ and $(l(\theta'; \cdot), w')$ in $\mathcal{H} \times \Delta_K^b$. Hence, by Dudley's entropy integral inequality (see Corollary 13.2 in Boucheron et al. (2013)), there exists a universal constant C_d such that

$$\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \Delta_K^b) = \frac{1}{\sqrt{N_b}} \mathbb{E}\left[\sup_{\theta \in \Theta, w \in \Delta_K^b} M_{\theta,w} \middle| D^{\text{train}} \right] \leq \frac{C_d}{\sqrt{N_b}} \int_0^1 \sqrt{\log\left(\mathcal{N}(\mathcal{H} \times \Delta_K^b; \mathrm{d}; \epsilon) \right)} d\epsilon,$$

where $\mathcal{N}(\mathcal{H} \times \Delta_K^b; \mathrm{d}; \epsilon)$ is the ϵ-covering number of $\mathcal{H} \times \Delta_K^b$ w.r.t. d.

Next, we just need to bound $\mathcal{N}(\mathcal{H} \times \mathcal{W}_\varepsilon^*; \mathrm{d}; \epsilon)$. Similar to (29), we can show that

$$\mathrm{d}^2(\theta, w, \theta', w') \leq \frac{2N_b}{n_{\text{train}}} \left(\sum_{k \in \mathcal{K}} \sum_{i=1}^{n_{\text{train}}} \frac{b^2}{n_{\text{train}}} \left(l(\theta; z_k^{(i)}) - l(\theta'; z_k^{(i)}) \right)^2 \right) + \frac{2N_b}{n_{\text{train}}} \|w - w'\|^2.$$

Similar to (30), we define a probability measure $\mathbb{Q} = \frac{N_b}{n_{\text{train}}} \sum_{k \in \mathcal{K}} \sum_{i=1}^{n_{\text{train}}} \frac{b^2}{n_{\text{train}}} \delta_{z_k^{(i)}}$ on $\mathcal{Z}$, where $\delta_{z_k^{(i)}}$ is a point mass at $z_k^{(i)}$. Then, we only need to construct an ϵ-cover for $\mathcal{H} \times \Delta_K^b$ by taking the Cartesian product of an $\frac{\epsilon}{2}$-cover for $\mathcal{H}$ w.r.t. distance metric $\rho_\mathbb{Q}$ and a $\sqrt{\frac{n_{\text{train}}}{N_b}} \frac{\epsilon}{2}$-cover for Δ_K^b w.r.t. the Euclidean distance. According to Assumption 4, the former has a cardinality of $(2C_\mathcal{H}/\epsilon)^{\nu_\mathcal{H}}$. To construct a $\sqrt{\frac{n_{\text{train}}}{N_b}} \frac{\epsilon}{2}$-cover for Δ_K^b, we first construct a $\sqrt{\frac{n_{\text{train}}}{KN_b}} \frac{\epsilon}{2}$-cover for $[0, b]$, take its K-fold Cartesian product, and project it to Δ_K^b. This provides a $\sqrt{\frac{n_{\text{train}}}{N_b}} \frac{\epsilon}{2}$-cover for Δ_K^b with a cardinality of $\left\lceil \frac{b\sqrt{KN_b}}{\sqrt{n_{\text{train}}}\epsilon} \right\rceil^K = \left\lceil \frac{1}{\epsilon} \right\rceil^K$. This implies $\mathcal{N}(\mathcal{H} \times \Delta_K^b; \mathrm{d}; \epsilon) \leq \left(\frac{2C_\mathcal{H}}{\epsilon} \right)^{\nu_\mathcal{H}} \times \left\lceil \frac{1}{\epsilon} \right\rceil^K$ and thus

$$\mathbb{E}[G_{\text{train}}(D^{\text{train}})] \leq 2\widehat{R}_{n_{\text{train}}}(\mathcal{H}, \Delta_K^b) \leq \frac{2C_d}{\sqrt{N_b}} \int_0^1 \sqrt{\log \mathcal{N}(\mathcal{H} \times \Delta_K^b; \mathrm{d}; \epsilon)} d\epsilon$$

$$= O\left(\frac{1}{\sqrt{N_b}} \int_0^1 \sqrt{\nu_\mathcal{H} \log\left(\frac{1}{\epsilon}\right) + K \log\left(\left\lceil \frac{1}{\epsilon} \right\rceil\right)} d\epsilon \right) = O\left(\sqrt{\frac{\nu_\mathcal{H} + K}{N_b}} \right). \tag{34}$$

Combining (34) with (33), we have that, with a probability of at least $1 - \delta$,

$$G_{\text{train}}(D^{\text{train}}) \leq O\left(\sqrt{\frac{\nu_\mathcal{H} + K + \log(1/\delta)}{N_b}} \right).$$

Applying the same argument to $G'_{\text{valid}}(D^{\text{valid}})$, we can show that the same inequality as above holds for $G'_{\text{train}}(D^{\text{train}})$ with a probability of at least $1 - \delta$. By taking a union bound, we have that, with a probability of at least $1 - \delta$,

$$\sup_{\theta \in \Theta, w \in \Delta_K^b} \left| \sum_{k=1}^K w_k \left[\widehat{L}_k(\theta) - L_k(\theta) \right] \right| \leq O\left(\sqrt{\frac{\nu_\mathcal{H} + K + \log(1/\delta)}{N_b}} \right),$$

which completes the proof of (21) as $N_b = n_{\text{train}}/(Kb^2)$. $\qquad\square$

C PROOFS OF MAIN THEOREMS AND COROLLARIES

In this section, we provide the proofs of Theorem 1, Theorem 2 and Corollary 1.

Proof of Theorem 1. By the strong convexity of the loss function and the optimality of $\theta(w)$ and $\widehat{\theta}(w)$ in the inner problems in (P$_*$) and ($\widehat{\text{P}}$), we have, for any $w \in \Delta_K^b$,

$$\frac{\mu}{2} \left\| \theta(w) - \widehat{\theta}(w) \right\|^2 \leq \sum_{k=1}^K w_k L_k(\widehat{\theta}(w)) - \sum_{k=1}^K w_k L_k(\theta(w)) \tag{35}$$

$$\frac{\mu}{2} \left\| \theta(w) - \widehat{\theta}(w) \right\|^2 \leq \sum_{k=1}^K w_k \widehat{L}_k(\theta(w)) - \sum_{k=1}^K w_k \widehat{L}_k(\widehat{\theta}(w)). \tag{36}$$

Adding (35) and (36) on both sides leads to, with a probability of at least $1 - \delta$,

$$\mu \left\| \theta(w) - \widehat{\theta}(w) \right\|^2$$

$$\leq \sum_{k=1}^K w_k L_k(\widehat{\theta}(w)) - \sum_{k=1}^K w_k \widehat{L}_k(\widehat{\theta}(w)) + \sum_{k=1}^K w_k \widehat{L}_k(\theta(w)) - \sum_{k=1}^K w_k L_k(\theta(w))$$

$$\leq 2C_a \sqrt{\frac{\nu_\mathcal{H} + K + \log(1/\delta)}{n_{\text{train}}/(Kb^2)}}, \quad \forall w \in \Delta_K^b, \tag{37}$$

where the second inequality is because the first conclusion in Lemma 3.

Let $w^* = \text{Proj}_{\mathcal{W}^*}(\widehat{w})$. Then we have, with a probability of $1 - 2\delta$, that

$$
\begin{aligned}
F(\widehat{w}) - \min_{w \in \Delta_K^b} F(w) =& L_0(\theta(\widehat{w})) - L_0(\theta(w^*)) \\
\leq& L_0(\widehat{\theta}(\widehat{w})) - L_0(\widehat{\theta}(w^*)) + \ell_0 \|\widehat{\theta}(\widehat{w}) - \theta(\widehat{w})\| + \ell_0 \|\widehat{\theta}(w^*) - \theta(w^*)\| \\
\leq& \widehat{L}_0(\widehat{\theta}(\widehat{w})) - \widehat{L}_0(\widehat{\theta}(w^*)) + 2C_0 \sqrt{\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\text{valid}}}} \\
& + \ell_0 \|\widehat{\theta}(\widehat{w}) - \theta(\widehat{w})\| + \ell_0 \|\widehat{\theta}(w^*) - \theta(w^*)\| \\
\leq& 2C_0 \sqrt{\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\text{valid}}}} + 2\ell_0 \sqrt{\frac{2C_a}{\mu}} \left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{\text{train}}/(Kb^2)} \right)^{\frac{1}{4}},
\end{aligned}
\tag{38}
$$

where the first inequality is because of Assumption 1, the second is due to Lemma 2, and the last is due to (37) and the optimality of $\widehat{w}$ for problem $(\widehat{P})$. Therefore, we can show that, with a probability of $1 - 2\delta$,

$$
\begin{aligned}
L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) =& L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta(\widehat{w})) + L_0(\theta(\widehat{w})) - L_0(\theta(w^*)) \\
\leq& \ell_0 \|\widehat{\theta}(\widehat{w}) - \theta(\widehat{w})\| + L_0(\theta(\widehat{w})) - L_0(\theta(w^*)) \\
\leq& 2C_0 \sqrt{\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\text{valid}}}} + 3\ell_0 \sqrt{\frac{2C_a}{\mu}} \left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{\text{train}}/(Kb^2)} \right)^{\frac{1}{4}}
\end{aligned}
$$

where the equality is because of Assumption 2, the first inequality is by Assumption 1 and the second by (37) and (38). This completes the proof. $\qquad\square$

Proof of Theorem 2. Since Assumption $2'$ implies Assumption 2, the proof and the conclusion of Theorem 1 also hold under the assumptions of Theorem 2. In particular, inequality (38) holds with a probability of $1 - 2\delta$. According to Assumption 3 and (38), we have with a probability of $1 - 2\delta$ that

$$
\left[C_r^{-1} \text{Dist}(\widehat{w}, \mathcal{W}^*) \right]^r \leq 2C_0 \sqrt{\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{\text{valid}}}} + 2\ell_0 \sqrt{\frac{2C_a}{\mu}} \left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{\text{train}}/(Kb^2)} \right)^{\frac{1}{4}}.
$$

Applying the fact that $s + t \leq (s^{\frac{1}{r}} + t^{\frac{1}{r}})^r$ for any $s > 0$ and $t > 0$ to the right-hand side of the inequality above, we obtain (10) with an appropriately defined C_w.

Suppose $\text{Dist}(\widehat{w}, \mathcal{W}^*) \leq \varepsilon(n_{\text{valid}}, n_{\text{train}})$, which happens with a probability of $1 - 2\delta$ according to the proof above. We have $\widehat{w} \in \mathcal{W}_\varepsilon^*$ with $\varepsilon = \varepsilon(n_{\text{valid}}, n_{\text{train}})$ according to the definition in (19). We then decompose the optimality gap of the generalization loss as follows

$$
\begin{aligned}
& L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \\
=& \underbrace{L_0(\widehat{\theta}(\widehat{w})) - \sum_{k=1}^K \widehat{w}_k L_k(\widehat{\theta}(\widehat{w}))}_{T_1} + \underbrace{\sum_{k=1}^K \widehat{w}_k L_k(\widehat{\theta}(\widehat{w})) - \sum_{k=1}^K \widehat{w}_k \widehat{L}_k(\widehat{\theta}(\widehat{w}))}_{T_2} \\
& + \underbrace{\sum_{k=1}^K \widehat{w}_k \widehat{L}_k(\widehat{\theta}(\widehat{w})) - \sum_{k=1}^K \widehat{w}_k \widehat{L}_k(\theta(\widehat{w}))}_{T_3} + \underbrace{\sum_{k=1}^K \widehat{w}_k \widehat{L}_k(\theta(\widehat{w})) - \sum_{k=1}^K \widehat{w}_k L_k(\theta(\widehat{w}))}_{T_4} \\
& + \underbrace{\sum_{k=1}^K \widehat{w}_k L_k(\theta(\widehat{w})) - \sum_{k=1}^K \widehat{w}_k L_k(\theta(w^*))}_{T_5} + \underbrace{\sum_{k=1}^K \widehat{w}_k L_k(\theta^*) - L_0(\theta^*)}_{T_6}.
\end{aligned}
\tag{39}
$$

It is clear that $T_3 \leq 0$ and $T_5 \leq 0$ by the optimality of $\widehat{\theta}(\widehat{w})$ and $\theta(\widehat{w})$ in $(\widehat{P})$ and (P_*), respectively. Moreover, by Assumption $2'$, we have $w_j^* = 0$ for $w \in \mathcal{W}^*$ and $j \in \mathcal{K} \backslash \mathcal{J}$. Using the fact that

$\widehat{w} \in \mathcal{W}_\varepsilon^*$, we have $\sum_{k \in \mathcal{K} \setminus \mathcal{J}} \widehat{w}_k^2 \leq \text{Dist}^2(\widehat{w}, \mathcal{W}^*) \leq \varepsilon^2$, which implies

$$T_1 = \sum_{k=1}^K \widehat{w}_k \left(L_0(\widehat{\theta}(\widehat{w})) - L_k(\widehat{\theta}(\widehat{w})) \right) = \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \widehat{w}_k \left(L_0(\widehat{\theta}(\widehat{w})) - L_k(\widehat{\theta}(\widehat{w})) \right)$$

$$\leq \sqrt{\sum_{k \in \mathcal{K} \setminus \mathcal{J}} \widehat{w}_k^2} \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2} \leq \varepsilon \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2}.$$

Using a similar argument, we can also show

$$T_6 \leq \varepsilon \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2}.$$

According to Lemma 3 and the fact that $\widehat{w} \in \mathcal{W}_\varepsilon^*$, we have that, with a probability of at least $1 - \delta$,

$$T_2, T_4 \leq C_a' \left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{N_\varepsilon}} + \frac{\varepsilon(K - J)}{b\sqrt{N_\varepsilon}} \right).$$

Note that $G = \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2}$ under Assumption $2'$. Applying the upper bounds of the six terms to (39) and taking a union bound, we can show that

$$L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \leq 2C_a' \left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{N_\varepsilon}} + \frac{\varepsilon(K - J)}{b\sqrt{N_\varepsilon}} \right) + 2\varepsilon(n_{\text{valid}}, n_{\text{train}})G$$

with a probability of at least $1 - 3\delta$, which completes the proof. $\qquad\square$

Before we prove Corollary 1, we first present another corollary of Theorem 2 where we can see the impact of b more clearly.

Corollary 2. *Suppose the assumptions of Theorem 2 hold and n_{valid} and n_{train} are large enough such that $\varepsilon(n_{valid}, n_{train})$ defined in (10) satisfies $\varepsilon(n_{valid}, n_{train}) \leq \frac{b\sqrt{J}}{K - J}$. With a probability of at least $1 - 3\delta$, we have*

$$L_0(\widehat{\theta}(\widehat{w})) - L_0(\theta^*) \leq O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{n_{train}/(Jb^2)}} \right)$$

$$+ G \cdot O\left(\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{valid}} \right)^{\frac{1}{2r}} + G \cdot O\left(\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{n_{train}/(Kb^2)} \right)^{\frac{1}{4r}}, \qquad (40)$$

Proof. Theorem 2 guarantees that (11) holds with a high probability. When $\varepsilon(n_{\text{valid}}, n_{\text{train}}) \leq \frac{b\sqrt{J}}{K - J}$, the second term on the right-hand side of (11) can be merged with the first term. Also, we have $N_\varepsilon = \frac{n_{\text{train}}}{b^2 J + \varepsilon^2 (n_{\text{valid}}, n_{\text{train}})(K - J)} \geq \frac{n_{\text{train}}}{b^2 J + b^2 J/(K - J)} \geq \frac{n_{\text{train}}}{2b^2 J}$, where the last inequality is because $K - J \geq 1$. As a result, the first two terms in (11) together has the order of

$$O\left(\sqrt{\frac{\nu_{\mathcal{H}} + J + \log(1/\delta)}{n_{\text{train}}/(Jb^2)}} \right).$$

Then (40) is proved by applying the definition of $\varepsilon(n_{\text{valid}}, n_{\text{train}})$ to the third term on the right-hand side of (11). $\qquad\square$

Suppose $n_{\text{train}} \gg n_{\text{valid}}$. The first term on the right-hand side of (40) is smaller than the entire right-hand side of (7). If, in addition, $G = o(1/n_{\text{valid}}^{\frac{1}{2} - \frac{1}{2r}})$ and $G = o(1/n_{\text{train}}^{\frac{1}{4} - \frac{1}{4r}})$, the other two terms in (40) are also smaller than the two terms in (7), respectively, so the bound in (40) is tighter than (7).

Proof of Corollary 1. Corollary 1 is directly from Corollary (2) by only keeping G, n_{train} and n_{test} in the order of magnitude given in (40). $\qquad\square$

D GENERALIZATION PERFORMANCE BY TRAINING LOCALLY AND TRAINING WITH EQUALLY WEIGHTED NODES

In this section, we first consider a model locally trained only with data D^{valid} in node 0, namely,

$$\widehat{\theta}_{\text{valid}} \in \arg\min_{\theta\in\Theta} \widehat{L}_0(\theta) \tag{41}$$

where $\widehat{L}_0$ is given in (3). The generalization bound of $\widehat{\theta}_{\text{valid}}$ is well-known, so we omit the proof but directly give the result.

Proposition 1. *Suppose Assumptions 1 and 4 hold. There exists a universal constant $C_0 > 0$ such that, with a probability of at least $1 - \delta$,*

$$L_0(\widehat{\theta}_{\text{valid}}) - L_0(\theta^*) \leq 2C_0\sqrt{\frac{\nu_{\mathcal{H}} + \log(1/\delta)}{n_{valid}}}.$$

For the purpose of theoretical comparison, we also consider a model trained only with data D^{train} distributed over equally weighted nodes, namely,

$$\widehat{\theta}_{\text{equal}} \in \arg\min_{\theta\in\Theta} \frac{1}{K}\sum_{k=1}^{K}\widehat{L}_k(\theta), \tag{42}$$

where $\widehat{L}_k$ is defined as in (2). It is easy to construct an example where each p_k with $k \in \mathcal{K}\backslash\mathcal{J}$ is significantly different from p_0 so that $L_0(\widehat{\theta}_{\text{equal}})$ does not convergence to $L_0(\theta^*)$ as n_{train} goes to infinity. Motivated by Corollary 1 and the discussion afterwards, it will be interesting to show the generalization bound of $\widehat{\theta}_{\text{equal}}$ when each p_k with $k \in \mathcal{K}\backslash\mathcal{J}$ is similar to p_0 with a small G defined in (9).

Proposition 2. *Suppose Assumptions 1, 2′ and 4 hold. There exists a universal constant $C_a > 0$ such that, with a probability of at least $1 - 3\delta$,*

$$L_0(\widehat{\theta}_{\text{equal}}) - L_0(\theta^*) \leq 2C_a\sqrt{\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{Kn_{train}}} + \frac{2\sqrt{K-J}}{K}G,$$

where G is defined in (9).

Proof. We first define

$$\theta_{\text{equal}} \in \arg\min_{\theta\in\Theta} \frac{1}{K}\sum_{k=1}^{K}L_k(\theta). \tag{43}$$

We first decompose the optimality gap of the generalization loss as follows

$$
\begin{aligned}
& L_0(\widehat{\theta}_{\text{equal}}) - L_0(\theta^*) \\
= \; & \underbrace{L_0(\widehat{\theta}_{\text{equal}}) - \frac{1}{K}\sum_{k=1}^{K}L_k(\widehat{\theta}_{\text{equal}})}_{T_1} + \underbrace{\frac{1}{K}\sum_{k=1}^{K}L_k(\widehat{\theta}_{\text{equal}}) - \frac{1}{K}\sum_{k=1}^{K}\widehat{L}_k(\widehat{\theta}_{\text{equal}})}_{T_2} \\
& + \underbrace{\frac{1}{K}\sum_{k=1}^{K}\widehat{L}_k(\widehat{\theta}_{\text{equal}}) - \frac{1}{K}\sum_{k=1}^{K}\widehat{L}_k(\theta_{\text{equal}})}_{T_3} + \underbrace{\frac{1}{K}\sum_{k=1}^{K}\widehat{L}_k(\theta_{\text{equal}}) - \frac{1}{K}\sum_{k=1}^{K}L_k(\theta_{\text{equal}})}_{T_4} \\
& + \underbrace{\frac{1}{K}\sum_{k=1}^{K}L_k(\theta_{\text{equal}}) - \frac{1}{K}\sum_{k=1}^{K}L_k(\theta^*)}_{T_5} + \underbrace{\frac{1}{K}\sum_{k=1}^{K}L_k(\theta^*) - L_0(\theta^*)}_{T_6}
\end{aligned}
\tag{44}
$$

It is clear that $T_3 \leq 0$ and $T_5 \leq 0$ by the optimality of $\widehat{\theta}_{\text{equal}}$ and θ_{equal} in (42) and (43), respectively. Moreover, we have

$$
\begin{aligned}
T_1 =& \frac{1}{K} \sum_{k=1}^{K} \left(L_0(\widehat{\theta}_{\text{equal}}) - L_k(\widehat{\theta}_{\text{equal}}) \right) = \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \frac{1}{K} \left(L_0(\widehat{\theta}_{\text{equal}}) - L_k(\widehat{\theta}_{\text{equal}}) \right) \\
\leq& \sqrt{\frac{K-J}{K^2}} \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2}.
\end{aligned}
$$

Using a similar argument, we can also show

$$
T_6 \leq \sqrt{\frac{K-J}{K^2}} \sqrt{\max_{\theta \in \Theta} \sum_{k \in \mathcal{K} \setminus \mathcal{J}} \left(L_0(\theta) - L_k(\theta) \right)^2}.
$$

Since Assumption $2'$ implies Assumption 2, by the first statement of Lemma 3 with $b = \frac{1}{K}$, we have, with a probability of at least $1 - \delta$, that

$$
T_2, \, T_4 \leq C_a \sqrt{\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{K n_{\text{train}}}}.
$$

Applying the upper bounds of the six terms to (44) and a taking union bound, we have

$$
L_0(\widehat{\theta}_{\text{equal}}) - L_0(\theta^*) \leq 2 C_a \sqrt{\frac{\nu_{\mathcal{H}} + K + \log(1/\delta)}{K n_{\text{train}}}} + \frac{2\sqrt{K-J}}{K} G
$$

with a probability of at least $1 - 3\delta$, which completes the proof. $\qquad\square$

Note that the bound in Proposition 2 is strictly worse than the one we showed in Corollary 1 for any value of G. In fact, the former is $O(1/\sqrt{n_{\text{train}}} + G)$ and the latter is $O(1/\sqrt{n_{\text{train}}}) + o(G)$

E   COMMUNICATION COMPLEXITY OF ALGORITHM 2 AND EXTENSION TO NON-CONVE CASE

In this section, we present the communication complexity of Algorithm 2 for convex problems as well as the corresponding algorithm and complexity for non-convex problems. To do so, we first present the convergence property of Algorithm 1, which is originally established by Gorbunov et al. (2021). Then, we combine the analysis by Gorbunov et al. (2021) and Ghadimi & Wang (2018) with some minor but necessary modifications, for example, to allow for a generic weight w instead of the uniform weight in Gorbunov et al. (2021), and to handle the approximation error between $\nabla \widehat{F}(w)$ and $\bar{\nabla} \widehat{F}(w)$, which is a little different from the one considered in Ghadimi & Wang (2018).

E.1   CONVERGENCE PROPERTY OF ALGORITHM 1

As mentioned in Section 4, we need to solve subproblems (1) and (15) in Algorithm 2, both of which are instances of (16). Because of Assumption 1, problem (16) in these two cases satisfies the following assumption with ℓ_1 and μ exactly the same as the ℓ_1 and μ in Assumption 1.

Assumption 5. *$f_{k,i}(x)$ is convex, $\nabla f_{k,i}(x)$ is ℓ_1-Lipschitz continuous and $f_k(x)$ is μ-strongly convex for $i = 1, \ldots, n_k$ and $k = 1, 2, \ldots, K$.*

A unified analysis is provided in Gorbunov et al. (2021) for a large class of FL methods including Local-SVRG given in Algorithm 1. The following proposition is obtained by applying Theorem 2.1 in Gorbunov et al. (2021) to Local-SVRG under our setting after minor modifications. It characterizes the convergence property of Algorithm 1. We omit its proof because it is the almost the same as the proof of Theorem G.7 in Gorbunov et al. (2021).

Proposition 3. *Suppose Assumption 5 holds for (16) and $\gamma \leq \gamma_0$ with γ_0 defined in (17). Algorithm 1 guarantees*

$$\mathbb{E}\left[f(\overline{x}^{(T)}) - f(x^*)\right] \leq \frac{1}{\gamma}\left(1 - \gamma\mu\right)^{T+1}\left(4 + \frac{32\gamma^2\ell_1^2}{3q} + 30e\gamma^3\ell_1^3(\tau-1)\frac{2+q}{q}\right)\left\|x^{(0)} - x^*\right\|^2$$

$$+ \frac{45e}{2}\ell_1\gamma^2(\tau-1)^2\sum_{k=1}^{K}w_k\left\|\nabla f_k(x^*)\right\|^2, \tag{45}$$

where x^* be the optimal solution of (16).

E.2 COMMUNICATION COMPLEXITY OF ALGORITHM 2

To analyze the complexity of Algorithm 2, we needs to bound the error of the approximate gradient of $\widehat{F}$, namely, the quantity

$$\left\|\bar{\nabla}\widehat{F}(w_{\mathrm{md}}^{(s)}) - \nabla\widehat{F}(w_{\mathrm{md}}^{(s)})\right\|, \quad s = 0, 1, \ldots,$$

and then the convergence analysis in Ghadimi & Wang (2018) can be directly applied. This error depends the suboptimality of $\theta^{(s)}$ and $h^{(s)}$ in iteration s of Algorithm 2, which can be characterized using Proposition 3. To do so, we first bound $\|\nabla f_k(x^*)\|$ and $\|x^{(0)} - x^*\|$ appearing in Proposition 3 for these two instances. For simplicity, we assume Local-SVRG is initialized at $x^{(0)} = 0$ when it is applied to any instance of (16).

Suppose $f_{k,i}(\theta) = l(\theta; z_k^{(i)}), i = 1, \ldots, n_k, k = 1, \ldots, K$ and x^* is the optimal solution of (16), namely, $x^* = \widehat{\theta}(w)$. Because of Assumption 1, we have that $\left\|\nabla f_k(x^*)\right\| \leq \ell_0$ for any k in any iteration s of Algorithm 2 and $x^* = \widehat{\theta}(w)$ is a continuous of w on Δ_K^b, which means $\left\|x^{(0)} - x^*\right\|^2 = \left\|x^*\right\|^2 \leq \max_{w\in\Delta_K^b}\|\widehat{\theta}(w)\|^2$ in any iteration s of Algorithm 2.

Suppose $f_{k,i}(h) = \frac{1}{2}h^\top\nabla^2 l(\theta^{(s)}; z_k^{(i)})h - h^\top\nabla\widehat{L}_0(\theta^{(s)}), i = 1, \ldots, n_k, k = 1, \ldots, K$. The optimal solution of (16) in this case is

$$x^* = \left(\sum_{k=1}^{K}w_k^{(s)}\nabla^2\widehat{L}_k(\theta^{(s)})\right)^{-1}\nabla\widehat{L}_0(\theta^{(s)})$$

which means $\left\|x^{(0)} - x^*\right\|^2 = \left\|x^*\right\|^2 \leq \frac{1}{\mu^2}\|\nabla\widehat{L}_0(\theta^{(s)})\|^2 \leq \frac{\ell_0^2}{\mu^2}$ because of Assumption 1. Moreover,

$$\nabla f_k(x^*) = \nabla^2\widehat{L}_k(\theta^{(s)})\left(\sum_{k=1}^{K}w_k^{(s)}\nabla^2\widehat{L}_k(\theta^{(s)})\right)^{-1}\nabla\widehat{L}_0(\theta^{(s)}) - \nabla\widehat{L}_0(\theta^{(s)}),$$

so $\|\nabla f_k(x^*)\| \leq \frac{\ell_1\ell_0}{\mu} + \ell_0$ for any k and s by Assumption 1.

Since f is μ-strongly convex, we have $\frac{\mu}{2}\|\overline{x}^{(T)} - x^*\|^2 \leq f(\overline{x}^{(T)}) - f(x^*)$. With this inequality and the discussion observations, we can derive from (45) that

$$\mathbb{E}\left[\left\|\overline{x}^{(T)} - x^*\right\|^2\right] \leq \frac{2}{\gamma\mu}\left(1 - \gamma\mu\right)^{T+1}\left(4 + \frac{32\gamma^2\ell_1^2}{3q} + 30e\gamma^3\ell_1^3(\tau-1)\frac{2+q}{q}\right)\cdot\max\left\{R^2, \frac{\ell_0^2}{\mu^2}\right\}$$

$$+ \frac{45e}{\mu}\ell_1\gamma^2(\tau-1)^2(\frac{\ell_1}{\mu}+1)^2\ell_0^2 \tag{46}$$

with $R := \max_{w\in\Delta_K^b}\|\widehat{\theta}(w)\|$ when Local-SVRG is applied to either (1) or (15) in any iteration of Algorithm 2.

The following lemma bounds the error of the approximate gradient for $\widehat{F}$.

Lemma 4. *Suppose Assumption 5 holds and $\gamma \leq \gamma_0$ with γ_0 defined in (17). We have*

$$\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{md}^{(s)}) - \bar{\nabla}\widehat{F}(w_{md}^{(s)})\big\|^2\big] \leq \frac{C_1}{\gamma}\left(1 - \gamma\mu\right)^{T_s+1}\left(C_2 + C_3(\tau - 1)\right) + C_4\gamma^2(\tau - 1)^2, \quad (47)$$

where C_1, C_2 and C_3 are constants that depend on ℓ_0, ℓ_1, ℓ_2, μ, R, q and K but not τ, T_s and γ. Consequently,

- *when $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0\mu}\ln\left(\frac{C_1C_2(s+1)^4}{\gamma_0}\right)$ OR*

- *when $\tau > 1$, $\gamma = \frac{1}{M_s}$ and $T_s = \mu^{-1}M_s\ln\left(M_s^3\right)$,*

where

$$M_s = \max\left\{\frac{1}{\gamma_0}, (s+1)^2\sqrt{[C_1(C_2 + C_3(\tau - 1)) + C_4(\tau - 1)^2]}\right\},$$

we have $\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{md}^{(s)}) - \bar{\nabla}\widehat{F}(w_{md}^{(s)})\big\|^2\big] \leq \frac{1}{(s+1)^4}$.

Moreover,

- *when $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0\mu}\ln\left(\frac{C_1C_2(s+1)^2}{\gamma_0}\right)$ OR*

- *when $\tau > 1$, $\gamma = \frac{1}{M_s'}$ and $T_s = \mu^{-1}M_s'\ln\left(M_s'^3\right)$,*

where

$$M_s' = \max\left\{\frac{1}{\gamma_0}, (s+1)\sqrt{[C_1(C_2 + C_3(\tau - 1)) + C_4(\tau - 1)^2]}\right\},$$

we have $\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{md}^{(s)}) - \bar{\nabla}\widehat{F}(w_{md}^{(s)})\big\|^2\big] \leq \frac{1}{(s+1)^2}$.

Proof. Let $\widetilde{\nabla}\widehat{F}(w_{md}^{(s)}) = (\widetilde{\nabla}_k\widehat{F}(w_{md}^{(s)}))_{k=1,\ldots,K}$, where

$$\widetilde{\nabla}_k\widehat{F}(w_{md}^{(s)}) = -\nabla\widehat{L}_k(\theta^{(s)})^\top\left(\sum_{k=1}^K w_{md,k}^{(s)}\nabla^2\widehat{L}_k(\theta^{(s)})\right)^{-1}\nabla\widehat{L}_0(\theta^{(s)}). \quad (48)$$

Recall that

$$\bar{\nabla}_k\widehat{F}(w_{md}^{(s)}) = -\nabla\widehat{L}_k(\theta^{(s)})^\top h^{(s)}.$$

We then obtained from (46) that

$$\mathbb{E}\big[\big\|\widetilde{\nabla}\widehat{F}(w_{md}^{(s)}) - \bar{\nabla}\widehat{F}(w_{md}^{(s)})\big\|^2\big]$$

$$\leq K\ell_0^2\mathbb{E}\left[\left\|h^{(s)} - \left(\sum_{k=1}^K w_{md,k}^{(s)}\nabla^2\widehat{L}_k(\theta^{(s)})\right)^{-1}\nabla\widehat{L}_0(\theta^{(s)})\right\|^2\right]$$

$$\leq \frac{2K\ell_0^2}{\gamma\mu}\left(1 - \gamma\mu\right)^{T_s+1}\left(4 + \frac{32\gamma_0^2\ell_1^2}{3q} + 30e\gamma_0^3\ell_1^3(\tau - 1)\frac{2+q}{q}\right)\cdot\max\left\{R^2, \frac{\ell_0^2}{\mu^2}\right\}$$

$$+ \frac{45eK\ell_0^2}{\mu}\ell_1\gamma^2(\tau - 1)^2(\frac{\ell_1}{\mu} + 1)^2\ell_0^2 \quad (49)$$

According to Lemma 2.2 in Ghadimi & Wang (2018) and (46), we have

$$\mathbb{E}\big[\big\|\widetilde{\nabla}\widehat{F}(w_{md}^{(s)}) - \nabla\widehat{F}(w_{md}^{(s)})\big\|^2\big]$$

$$\leq K\left(\frac{2\ell_0\ell_1}{\mu} + \frac{\ell_2\ell_0^2}{\mu^2}\right)^2\mathbb{E}\big[\|\theta^{(s)} - \widehat{\theta}(w_{md}^{(s)})\|^2\big]$$

$$\leq K\left(\frac{2\ell_0\ell_1}{\mu} + \frac{\ell_2\ell_0^2}{\mu^2}\right)^2\frac{2}{\gamma\mu}\left(1 - \gamma\mu\right)^{T_s+1}\left(4 + \frac{32\gamma_0^2\ell_1^2}{3q} + 30e\gamma_0^3\ell_1^3(\tau - 1)\frac{2+q}{q}\right)\cdot\max\left\{R^2, \frac{\ell_0^2}{\mu^2}\right\}$$

$$+ K\left(\frac{2\ell_0\ell_1}{\mu} + \frac{\ell_2\ell_0^2}{\mu^2}\right)^2\frac{45e}{\mu}\ell_1\gamma^2(\tau - 1)^2(\frac{\ell_1}{\mu} + 1)^2\ell_0^2 \quad (50)$$

Combining (49) and (50) by the triangle inequity leads to (47).

When $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0 \mu} \ln\left(\frac{C_1 C_2 (s+1)^4}{\gamma_0 \mu}\right)$, it is easy to show that $\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{\mathrm{md}}^{(s)}) - \bar{\nabla}\widehat{F}(w_{\mathrm{md}}^{(s)})\big\|^2\big] \leq \frac{1}{(s+1)^4}$.

Suppose $\tau > 1$, $\gamma = \frac{1}{M_s}$ and $T_s = \mu^{-1} M_s \ln\left(M_s^3\right)$. We have

$$
\begin{aligned}
\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{\mathrm{md}}^{(s)}) - \bar{\nabla}\widehat{F}(w_{\mathrm{md}}^{(s)})\big\|^2\big] \leq & \frac{C_1}{\gamma}\left(1 - \gamma\mu\right)^{T_s+1}(C_2 + C_3(\tau-1)) + C_4\gamma^2(\tau-1)^2 \\
\leq & \exp\left(-\frac{\mu T_s}{M_s}\right)M_s C_1\left(C_2 + C_3(\tau-1)\right) + \frac{C_4(\tau-1)^2}{M_s^2} \\
\leq & \exp\left(-\ln(M_s^3)\right)M_s C_1\left(C_2 + C_3(\tau-1)\right) + \frac{C_4(\tau-1)^2}{M_s^2} \\
\leq & \frac{C_1\left(C_2 + C_3(\tau-1)\right)}{M_s^2} + \frac{C_4(\tau-1)^2}{M_s^2} \leq \frac{1}{(s+1)^4}.
\end{aligned}
$$

The conclusion with $\mathbb{E}\big[\big\|\nabla\widehat{F}(w_{\mathrm{md}}^{(s)}) - \bar{\nabla}\widehat{F}(w_{\mathrm{md}}^{(s)})\big\|^2\big] \leq \frac{1}{(s+1)^2}$ can be proved in the same way except that $(s+1)^4$ must be changed to $(s+1)^2$, and thus we omit the proof. □

The complexity of Algorithm 2 when $\widehat{F}(w)$ is convex can be showed using the proof in Ghadimi & Wang (2018) with their gradient approximation error replaced by the one in Lemma 4.

Proof of Theorem 3. According to Lemma 2.2 in Ghadimi & Wang (2018), $\widehat{F}(w)$ is ℓ_F-smooth with ℓ_F defined in (12). Let $E_s := \big\|\nabla\widehat{F}(w_{\mathrm{md}}^{(s)}) - \bar{\nabla}\widehat{F}(w_{\mathrm{md}}^{(s)})\big\|$ According to (2.51) in Ghadimi & Wang (2018), we have

$$
\begin{aligned}
\widehat{F}(w_{\mathrm{ag}}^{(s+1)}) \leq & \frac{s}{s+2}\widehat{F}(w_{\mathrm{ag}}^{(s)}) + \frac{2}{s+2}\widehat{F}(\widehat{w}) + \frac{16}{\eta(s+1)(s+2)}\left(\|\widehat{w} - w^{(s)}\|^2 - \|\widehat{w} - w^{(s+1)}\|^2\right) \\
& + \frac{2}{s+2}\|\widehat{w} - w^{(s+1)}\|E_s + \frac{\eta}{2}E_s^2 \\
\leq & \frac{s}{s+2}\widehat{F}(w_{\mathrm{ag}}^{(s)}) + \frac{2}{s+2}\widehat{F}(\widehat{w}) + \frac{16}{\eta(s+1)(s+2)}\left(\|\widehat{w} - w^{(s)}\|^2 - \|\widehat{w} - w^{(s+1)}\|^2\right) \\
& + \frac{4}{s+2}E_s + \frac{\eta}{2}E_s^2,
\end{aligned}
$$

where the second inequality is because $\|\widehat{w} - w^{(s+1)}\| \leq 2$ as both $\widehat{w}$ and $w^{(s+1)}$ are on a simplex. Subtracting $\widehat{F}(\widehat{w})$ from both sides of the inequality above and dividing both sides by $\frac{2}{(s+1)(s+2)}$, we have

$$
\begin{aligned}
& \frac{(s+1)(s+2)}{2}\left[\widehat{F}(w_{\mathrm{ag}}^{(s+1)}) - \widehat{F}(\widehat{w})\right] \\
\leq & \frac{s(s+1)}{2}\left[\widehat{F}(w_{\mathrm{ag}}^{(s)}) - \widehat{F}(\widehat{w})\right] + \frac{8}{\eta}\left(\|\widehat{w} - w^{(s)}\|^2 - \|\widehat{w} - w^{(s+1)}\|^2\right) \\
& + 2(s+1)E_s + \frac{\eta(s+1)(s+2)}{4}E_s^2.
\end{aligned}
$$

Summing up this inequality for $s = 0, 1, \ldots, S-1$ gives

$$
\frac{S(S+1)}{2}\mathbb{E}\left[\widehat{F}(w_{\mathrm{ag}}^{(S)}) - \widehat{F}(\widehat{w})\right] \leq \frac{16}{\eta} + \sum_{s=0}^{S-1}2(s+1)\sqrt{\mathbb{E}\big[E_s^2\big]} + \sum_{s=0}^{S-1}\frac{\eta(s+1)(s+2)}{4}\mathbb{E}\big[E_s^2\big],
$$

which, when $\mathbb{E}\big[E_s^2\big] \leq \frac{1}{(s+2)^4}$, implies

$$
\mathbb{E}\left[\widehat{F}(w_{\mathrm{ag}}^{(S)}) - \widehat{F}(\widehat{w})\right] \leq \frac{32}{\eta S(S+1)} + \frac{4\log(S)}{S(S+1)} + \frac{\pi^2\eta}{12 S(S+1)}.
$$

This means, as long as $\mathbb{E}\left[E_s^2\right] \leq \frac{1}{(s+2)^4}$, Algorithm 2 finds an ϵ-optimal solution of $(\widehat{P})$ in $\tilde{O}(\epsilon^{-0.5})$ iterations.

Let $A_1 = C_1 C_2$, $A_2 = C_1 C_3$ and $A_3 = C_1 C_4$ with C_1, C_2, C_3 and C_4 defined in Lemma 4.

Suppose $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0 \mu} \ln \left(\frac{A_1 (s+1)^4}{\gamma_0} \right)$. By Lemma 4, we have $\mathbb{E}\left[E_s^2\right] \leq \frac{1}{(s+2)^4}$. In iteration s of Algorithm 2, the total number of rounds of communication needed in Local-SVRG is $O(T_s) = O(\ln(s))$ so that means the total number of rounds is $\tilde{O}(\epsilon^{-0.5})$.

Suppose $\tau > 1$, $\gamma = \frac{1}{M_s}$ and $T_s = \mu^{-1} M_s \ln \left(M_s^3 \right)$. By Lemma 4, we have $\mathbb{E}\left[E_s^2\right] \leq \frac{1}{(s+2)^4}$. In iteration s of Algorithm 2, the total number of rounds of communication needed in Local-SVRG is $O(T_s/\tau) = O(s^2)$ so that means the total number of rounds is $\tilde{O}(\epsilon^{-1.5})$. $\qquad\square$

E.3 Algorithm and Communication Complexity for Non-Convex $\widehat{F}$

When $\widehat{F}$ in $(\widehat{P})$ is non-convex, we no long expect any algorithm to find an ϵ-optimal solution and change our goal to finding an ϵ-*stationary* point of $(\widehat{P})$, which is defined as a solution $\bar{w} \in \Delta_K^b$ satisfying

$$\eta^{-1} \left\| \bar{w} - \text{Proj}_{\Delta_K^b}(\bar{w} - \eta \nabla \widehat{F}(\bar{w})) \right\| \leq \epsilon.$$

for some $\eta > 0$. There exist multiple numerical techniques for finding an ϵ-stationary, among which the proximal gradient method is the simplest one. When the gradient can only be computed inexactly, there exist studies on the iteration complexity of the proximal gradient method for finding an ϵ-stationary point, including Ghadimi & Wang (2018) for bilevel optimization and Gu et al. (2018) for a general problem. We will simply apply the proximal gradient method to $(\widehat{P})$ using the approximate gradient $\bar{\nabla} \widehat{F}(w)$ in (14). We formally present this approach in Algorithm 3. Again, the center is node 0, i.e., the node where D^{valid} is stored.

Algorithm 3: Federated Learning Method for Bilevel Optimization $(\widehat{P})$ (Non-Convex Case)

---

1 **Input**: initial weight $w^{(0)}$, learning rate η, training data D_k^{train} for $k \in \mathcal{K}$, validation data D^{valid}, the number of outer iterations S, parameters (γ, τ, q) for Local-SVRG, and the number of inner iterations T_s for $s = 0, \ldots, S-1$

2 **for** $s = 0, 1, \ldots, S-1$ **do**

3   Compute $\bar{\nabla} \widehat{F}(w^{(s)})$ as follows:

4   Set $f_{k,i}(\theta) = l(\theta; z_k^{(i)})$, $\quad i = 1, \ldots, n_k$, $\quad k = 1, \ldots, K$

5   Compute $\theta^{(s)} = $ Local-SVRG$(\{f_{k,i}\}, w^{(s)}, \gamma, \tau, q, T_s)$ and send it to each node.

6   Compute $\nabla \widehat{L}_0(\theta^{(s)})$ at center and send it to each node.

7   Set $f_{k,i}(h) = \frac{1}{2} h^\top \nabla^2 l(\theta^{(s)}; z_k^{(i)}) h - h^\top \nabla \widehat{L}_0(\theta^{(s)})$, $\quad i = 1, \ldots, n_k$, $\quad k = 1, \ldots, K$

8   Compute $h^{(s)} = $ Local-SVRG$(\{f_{k,i}\}, w^{(s)}, \gamma, \tau, q, T_s)$ and send it to each node.

9   Each node computes $\nabla \widehat{L}_k(\theta^{(s)})$ in parallel and send it to the center.

10   Set $\bar{\nabla}_k \widehat{F}(w^{(s)}) = -\nabla \widehat{L}_k(\theta^{(s)})^\top h^{(s)}$ for $k = 1, \ldots, K$

11      $w^{(s+1)} = \arg\min_{w \in \Delta_K^b} \left\langle \bar{\nabla} \widehat{F}(w^{(s)}), w \right\rangle + \frac{1}{2\eta} \|w - w^{(s)}\|^2$

12 **end**

13 **Return**: $w^{(\bar{s})}$ with $\bar{s}$ sampled randomly from $\{0, 1, \ldots, S-1\}$.

The convergence result of Algorithm 3 can be proved in a standard way (e.g., see Theorem 2.1 in Ghadimi & Wang (2018)). We present it below only for the sake of completeness.

Theorem 4. *Suppose Assumption 1 holds. Let $R := \max_{w \in \Delta_K^b} \|\widehat{\theta}(w)\|$ and ℓ_F and γ_0 defined as in (12) and (17). Suppose $\eta = \frac{1}{3\ell_F}$ in Algorithm 3. There exist constants A_1, A_2 and A_3 that only depend on ℓ_0, ℓ_1, ℓ_2, μ, R, q and K but not on τ such that the following statements hold.*

- *Suppose $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0 \mu} \ln \left(\frac{A_1 (s+1)^2}{\gamma_0} \right)$. Algorithm 3 finds an ϵ-stationary solution of $(\widehat{P})$ with $\tilde{O}\left(\epsilon^{-2}\right)$ rounds of communication.*

- *Suppose $\tau > 1$, $\gamma = \frac{1}{M'_s}$ and $T_s = \mu^{-1} M'_s \ln\left(M'^3_s\right)$, where*

$$M'_s = \max\left\{\frac{1}{\gamma_0}, (s+1)\sqrt{[A_1 + A_2(\tau - 1) + A_3(\tau - 1)^2]}\right\}, s = 0, 1, \ldots. \quad (51)$$

Algorithm 3 finds an ϵ-stationary solution of ($\widehat{P}$) with $\tilde{O}\left(\epsilon^{-4}\right)$ rounds of communication.

Proof. Let $E_s := \left\|\nabla\widehat{F}(w^{(s)}) - \bar\nabla\widehat{F}(w^{(s)})\right\|$. Since $w^{(s+1)} = \text{Proj}_{\Delta^b_K}(w^{(s)} - \eta\bar\nabla\widehat{F}(w^{(s)}))$, by the property of projection mapping, we have

$$\left\|w^{(s)} - \text{Proj}_{\Delta^b_K}(w^{(s)} - \eta\nabla\widehat{F}(w^{(s)}))\right\|^2$$

$$\leq \quad 2\left\|w^{(s)} - w^{(s+1)}\right\|^2 + 2\left\|\text{Proj}_{\Delta^b_K}(w^{(s)} - \eta\bar\nabla\widehat{F}(w^{(s)})) - \text{Proj}_{\Delta^b_K}(w^{(s)} - \eta\nabla\widehat{F}(w^{(s)}))\right\|^2$$

$$\leq \quad 2\left\|w^{(s)} - w^{(s+1)}\right\|^2 + 2\eta^2 E_s^2. \quad (52)$$

By the definition of $w^{(s+1)}$ and the $\frac{1}{\eta}$-strong convexity of function $\left\langle\bar\nabla\widehat{F}(w^{(s)}), w\right\rangle + \frac{1}{2\eta}\|w - w^{(s)}\|^2$, we have, for any $w \in \Delta^b_K$

$$\left\langle\bar\nabla\widehat{F}(w^{(s)}), w^{(s+1)} - w^{(s)}\right\rangle + \frac{1}{2\eta}\|w^{(s+1)} - w^{(s)}\|^2 + \frac{1}{2\eta}\|w - w^{(s+1)}\|^2$$

$$\leq \quad \left\langle\bar\nabla\widehat{F}(w^{(s)}), w - w^{(s)}\right\rangle + \frac{1}{2\eta}\|w - w^{(s)}\|^2. \quad (53)$$

Taking $w = w^{(s)}$ in (53) gives

$$\left\langle\bar\nabla\widehat{F}(w^{(s)}), w^{(s+1)} - w^{(s)}\right\rangle + \frac{1}{\eta}\|w^{(s+1)} - w^{(s)}\|^2 \leq 0. \quad (54)$$

Since $\widehat{F}$ is ℓ_F-Lipschitz continuous and $\eta \leq \frac{1}{\ell_F}$, we have

$$\widehat{F}(w^{(s+1)}) - \widehat{F}(w^{(s)}) \quad \leq \quad \left\langle\nabla\widehat{F}(w^{(s)}), w^{(s+1)} - w^{(s)}\right\rangle + \frac{1}{2\eta}\|w^{(s+1)} - w^{(s)}\|^2. \quad (55)$$

Adding (54) and (55) gives us

$$\widehat{F}(w^{(s+1)}) - \widehat{F}(w^{(s)}) + \frac{1}{2\eta}\|w^{(s+1)} - w^{(s)}\|^2 \quad \leq \quad \left\langle\nabla\widehat{F}(w^{(s)}) - \bar\nabla\widehat{F}(w^{(s)}), w^{(s+1)} - w^{(s)}\right\rangle$$

$$\leq \quad E_s\|w^{(s+1)} - w^{(s)}\| \leq 2E_s,$$

which, together with (52), implies

$$\frac{1}{4\eta}\left\|w^{(s)} - \text{Proj}_{\Delta^b_K}(w^{(s)})\right\|^2 \leq \widehat{F}(w^{(s)}) - \widehat{F}(w^{(s+1)}) + 2E_s + \frac{\eta E_s^2}{2}.$$

Summing this inequality and taking expectation give us

$$\eta^{-2}\mathbb{E}\left[\left\|w^{(\bar s)} - \text{Proj}_{\Delta^b_K}(w^{(\bar s)})\right\|^2\right] \leq \frac{4}{\eta S}\left[\widehat{F}(w^{(0)}) - \widehat{F}(\widehat{w})\right] + \frac{8}{\eta S}\sum_{s=0}^{S-1}\mathbb{E}[E_s] + \frac{2}{S}\sum_{s=0}^{S-1}\mathbb{E}[E_s^2].$$

When $\mathbb{E}[E_s^2] \leq \frac{1}{(s+1)^2}$, the inequality above implies

$$\eta^{-2}\mathbb{E}\left\|w^{(\bar s)} - \text{Proj}_{\Delta^b_K}(w^{(\bar s)})\right\|^2 \leq \frac{4}{\eta S}\left[\widehat{F}(w^{(0)}) - \widehat{F}(\widehat{w})\right] + \frac{8\log(S)}{\eta S} + \frac{\pi^2}{3S}.$$

This means, as long as $\mathbb{E}[E_s^2] \leq \frac{1}{(s+1)^2}$, Algorithm 3 finds an ϵ-stationary solution of ($\widehat{P}$) in $\tilde{O}(\epsilon^{-2})$ iterations.

Note that Lemma 4 still holds with $w^{(s)}_{\text{md}}$ replaced by $w^{(s)}$ in Algorithm 3. Let $A_1 = C_1 C_2$, $A_2 = C_1 C_3$ and $A_3 = C_1 C_4$ with C_1, C_2, C_3 and C_4 defined in Lemma 4.

Suppose $\tau = 1$, $\gamma = \gamma_0$ and $T_s = \frac{1}{\gamma_0 \mu} \ln\left(\frac{A_1(s+1)^2}{\gamma_0}\right)$. By Lemma 4, we have $\mathbb{E}\left[E_s^2\right] \leq \frac{1}{(s+2)^2}$. In iteration s of Algorithm 3, the total number of rounds of communication needed in Local-SVRG is $O(T_s) = O(\ln(s))$ so that means the total number of rounds is $\tilde{O}(\epsilon^{-2})$.

Suppose $\tau > 1$, $\gamma = \frac{1}{M_s'}$ and $T_s = \mu^{-1} M_s' \ln\left(M_s'^3\right)$. By Lemma 4, we have $\mathbb{E}\left[E_s^2\right] \leq \frac{1}{(s+2)^4}$. In iteration s of Algorithm 3, the total number of rounds of communication needed in Local-SVRG is $O(T_s/\tau) = O(s)$ so that means the total number of rounds is $\tilde{O}(\epsilon^{-4})$. $\qquad\square$

Remark 1. *The federated bilevel optimization methods by Li et al. (2022) and Tarzanagh et al. (2022) can find an ϵ-stationary point within $\tilde{O}(\epsilon^{-3})$ and $\tilde{O}(\epsilon^{-4})$ rounds of communication, respectively. In the first setting of Theorem 4 ($\tau = 1$), Algorithm 3 has complexity of $\tilde{O}(\epsilon^{-2})$, which is better than Li et al. (2022) and Tarzanagh et al. (2022). We want to point out that the lower complexity of Algorithm 3 is because it utilizes the finite-sum structure in $(\widehat{P})$, which allows computing a deterministic gradient infrequently to accelerate the convergence. However, Li et al. (2022) and Tarzanagh et al. (2022) both consider objective functions given in expectation, which does not allow computing a deterministic gradient in general.*

F   ADDITIONAL MATERIALS FOR NUMERICAL EXPERIMENTS

In this section, we present additional details and results of our numerical experiments in Section 5. The CNN we train in the experiments consists of two layers of 2D convolution, each equipped with 2D batch normalization and ReLU activation, and followed by a fully connected layer to generate predictions. The first convolution layer is set to output the same number of channels as the input, and uses kernels with a size of 4, a stride of 4 and one padding. The second convolution layer returns two output channels for Fashion-MNIST and MNIST and five for CIFAR-10 and ImageNet, and uses kernels with a size of 2, a stride of 2 and one padding. All experiments are conducted with PyTorch 1.9.0 and CUDA 11.1 computing platform on a computer with the CPU Intel Xeon Gold 6330@2.0GHz (Turbo up to 3.1GHz) and the GPU NVIDIA GeForce RTX 2080 Ti.

F.1   DATA GENERATION WITH DIFFERENT CLASS DISTRIBUTIONS (SETTING 1)

In this section, we describe in details how we generate D^{valid}, D^{train} and D^{test} from each original dataset for our experiments under Setting 1.

F.1.1   FASHION-MNIST

Fashion-MNIST (Xiao et al., 2017) contains a training set of 60,000 images and a testing set of 10,000 images. Each image is in grayscale, has a size of 28×28, and is associated with a label from ten classes: 0: T-shirt/top, 1: Trouser, 2: Pullover, 3: Dress, 4: Coat, 5: Sandal, 6: Shirt, 7: Sneaker, 8: Bag and 9: Ankle boot. We merge the ten classes into four classes as follows:

C1: Classes 2, 4 and 6 which include long-sleeve upper-body clothes;

C2: Classes 0 and 3 which include short-sleeve upper-body clothes;

C3: Classes 1 and 8 which include pants and bags;

C4: Classes 5, 7 and 9 which include only shoes.

Note that these four merged classes are only used for generating data. In the classification task, we still have ten classes. This is the same for the other three datasets. We set $n_{\text{train}} = 4000$, $n_{\text{valid}} = 500$ and $n_{\text{test}} = 5000$. Each image is sampled from one of the four merged classes with a probability distribution (P_1, P_2, P_3, P_4). Once a merged class is chosen, each image in that merged class has an equal chance to be sampled. For D_k^{train} with $k \in \mathcal{J}$, we sample data from the training set with $P_1 = 0.42$, $P_2 = 0.08$, $P_3 = 0.38$ and $P_4 = 0.12$. For D_k^{train} for $k \in \mathcal{J}_M$, we sample data with $P_1 = 0.12$, $P_2 = 0.38$, $P_3 = 0.08$ and $P_4 = 0.42$. Depending on p_0 is the distribution of $\mathcal{J}_m$ or $\mathcal{J}_M$, D^{valid} and D^{test} are sampled from p_0 the corresponding distribution. Note that D^{valid} and D^{test} are sampled from the training set and testing set of the original data, respectively, although they have the same probability distribution over the four merged classes. Since D^{valid} and D^{test} are generated

in the similar way for the other three datasets, we will only discuss the generation of D_k^{train}'s in the subsequent sections.

F.1.2 MNIST

MNIST (Deng, 2012) contains a training set of 60,000 images and a testing set of 10,000 images. Each image is a handwritten digit in grayscale and has a size of 28×28. Since MNIST has the same number of classes, same class distribution, and same data size as Fashion-MNIST, we directly apply the same procedure in Section F.1.1 to sample data. In particular, we merge the ten digits into four classes as follows:

> C1: Digits 2, 4 and 6;
>
> C2: Digits 0 and 3;
>
> C3: Digits 1 and 8;
>
> C4: Digits 5, 7 and 9.

We set $n_{\text{train}} = 4000$, $n_{\text{valid}} = 500$ and $n_{\text{test}} = 5000$. Following Section F.1.1, for D_k^{train} with $k \in \mathcal{J}_m$, we sample data from the four merged classes with $P_1 = 0.42$, $P_2 = 0.08$, $P_3 = 0.38$ and $P_4 = 0.12$. For D_k^{train} with $k \in \mathcal{J}_M$, we sample data with $P_1 = 0.12$, $P_2 = 0.38$, $P_3 = 0.08$ and $P_4 = 0.42$.

F.1.3 CIFAR-10

CIFAR-10 (Krizhevsky et al., 2009) contains a training set of 50,000 images and a testing set of 10,000 images. Each image is in color, has a size of 32×32, and is associated with a label from ten classes: 0: airplane, 1: automobile, 2: bird, 3: cat, 4: deer, 5: dog, 6: frog, 7: horse, 8: ship and 9: truck. We merge the ten classes into four classes as follows:

> C1: Classes 1 and 9 which are related to ground transportation;
>
> C2: Classes 0 and 8 which are related to non-ground transportation;
>
> C3: Classes 2, 3 and 4 which form a set of animals;
>
> C4: Classes 5, 6 and 7 which form another set of animals.

We set $n_{\text{train}} = 4000$, $n_{\text{valid}} = 500$ and $n_{\text{test}} = 5000$. Similar to the procedure with Fashion-MNIST, for D_k^{train} with $k \in \mathcal{J}_m$, we sample data from the four merged classes with $P_1 = 0.36$, $P_2 = 0.04$, $P_3 = 0.54$ and $P_4 = 0.06$. For D_k^{train} with $k \in \mathcal{J}_M$, we sample data with $P_1 = 0.04$, $P_2 = 0.36$, $P_3 = 0.06$ and $P_4 = 0.54$.

F.1.4 DOWNSAMPLED IMAGENET

Downsampled ImageNet (Chrabaszcz et al., 2017) is created by downsampling each image in ImageNet (Deng et al., 2009) to 32×32 pixels without changing the class labels. Just as ImageNet, downsampled ImageNet has 1000 classes and we choose ten classes and merge them into four classes as follows. (The class labels listed below are consistent with ImageNet.)

> C1: Classes $7, 9, 10, 29, 54, 75, 84$ and 189, which are cats or animals similar to cat;
>
> C2: Classes $61, 66, 68, 101, 114, 124, 131$ and 148, which are dogs or animals similar to cat;
>
> C2: Classes $383, 397, 403, 404, 405, 406, 412, 414, 420, 426, 433$ and 434, which are all birds;
>
> C3: Classes $224, 441, 442, 443, 444, 445, 449, 453, 454, 498, 499$ and 500, which are either fishes or frogs.

We set $n_{\text{train}} = 4000$, $n_{\text{valid}} = 1500$, $n_{\text{test}} = 1000$. For D_k^{train} for $k \in \mathcal{J}_m$, we sample data from the four merged classes with $P_1 = 0.36$, $P_2 = 0.04$, $P_3 = 0.54$ and $P_4 = 0.06$. For D_k^{train} with $k \in \mathcal{J}_M$, we sample data with $P_1 = 0.04$, $P_2 = 0.36$, $P_3 = 0.06$ and $P_4 = 0.54$.

F.2 DATA GENERATION WITH DIFFERENT CLASS DISTRIBUTIONS AND LABEL PERMUTATION (SETTING 2)

In this section, we discuss in details how D^{train}, D^{valid} and D^{test} are generated from each original dataset for our experiments under Setting 2. For each dataset, we first sample data D^{train}, D^{valid} and D^{test} in the same way as Setting 1 described in Section F.1. Then we permute the class labels in D_k^{train} with $k \in \mathcal{J}_M$. For each dataset, the class labels in D_k^{train} with $k \in \mathcal{J}_m$ are unchanged. If p_0 is the distribution of $\mathcal{J}_M$, the labels of D^{valid} and D^{test} are permuted in the same way as D_k^{train} with $k \in \mathcal{J}_M$. If p_0 is the distribution of $\mathcal{J}_m$, the labels of D^{valid} and D^{test} are unchanged.

We then describe how the class labels in D_k^{train} for $k \in \mathcal{J}_M$ are permuted for each dataset. For Fashion-MNIST and MNIST, we permute the class labels in D_k^{train} for $k \in \mathcal{J}_M$ by changing label 2 to 0, 0 to 1, 1 to 5, and 5 to 2. For CIFAR-10, we permute the class labels in D_k^{train} for $k \in \mathcal{J}_M$ by changing label 1 to 0, 0 to 2, 2 to 5, and 5 to 1. For downsampled ImageNet, we permute the class labels in D_k^{train} for $k \in \mathcal{J}_M$ by changing label 7 to 61, 61 to 383, 383 to 224, 224 to 9, 9 to 66, 66 to 397, 397 to 441, 441 to 10, 10 to 68, 68 to 403, 403 to 442, and 442 to 7.

F.3 DATA GENERATION WITH DIFFERENT CLASS DISTRIBUTIONS, LABEL PERMUTATION AND/OR RANDOM ROTATION (SETTINGS 3 AND 4)

In this section, we discuss in details how we generate D^{valid}, D^{train} and D^{test} from each original dataset for our experiments under Settings 3 and 4.

Under Setting 3, we first generate data in the same way as in Setting 1 described in Section F.1. Then, we randomly choose a rotation direction, clockwise or anti-clockwise, and rotate each image in D_k^{train} with $k \in \mathcal{J}_M$ toward that direction for 90 degrees. Under Setting 4, we first generate data in the same way as in Setting 2 described in Section F.2. Then, we apply the same rotation procedure as we do in Setting 3.

F.4 ADDITIONAL NUMERICAL RESULTS

In this section, we first plot how the weight w_k for each node evolves during the Bi-level method under Setting 2, 3 and 4 respectively in Figure 3, Figure 4 and Figure 5. We show the results when p_0 is the distribution of the majority and the minority groups separately.

Since the performance of an algorithm may fluctuate during training, we are interested in comparing the methods in the best performance they achieved during training. To do so, we save the model generated by each method at the iteration where the highest accuracy is achieved on the validation data. Then, we report the performance of the saved model's by each method on the testing set in Table 1. Again, we show the results when p_0 is the distribution of the majority and the minority groups separately. Our method outperforms the baselines in most of the cases.

At last, we plot the test (top-1) accuracy each method obtains during iterations for the minority group and the majority group in Figure 7 and Figure 8, respectively, where the horizontal axis represents the cumulative number of data points each method processes in parallel.

Figure 3: How w evolves during the Bi-level method under Setting 2.

Figure 4: How w evolves during the Bi-level method under Setting 3.

Figure 5: How w evolves during the Bi-level method under Setting 4.

Table 1: Test accuracy when reaching the highest validation accuracy.

Setting 1	Minority Model				Majority Model			
Method\Data	F-MNIST	MNIST	CIFAR-10	DS-ImageNet	F-MNIST	MNIST	CIFAR-10	DS-ImageNet
Bi-level	**0.7758±0.0059**	**0.8824±0.0198**	**0.5175±0.0065**	**0.2668±0.0079**	**0.8364±0.0121**	0.8443±0.0096	**0.5928±0.0068**	**0.2630±0.0091**
Local-train	0.6926±0.0175	0.7850±0.0269	0.3036±0.0102	0.1108±0.0127	0.7427±0.0110	0.7550±0.0095	0.3568±0.0144	0.1150±0.0072
FedAvg	0.7507±0.0097	0.8297±0.0171	0.2965±0.0075	0.2008±0.0117	0.8327±0.0119	**0.8484±0.0123**	0.5705±0.0052	0.2612±0.0116
Ditto	0.7297±0.0146	0.8358±0.0201	0.4361±0.0166	0.1582±0.0037	0.8086±0.0141	0.8059±0.0053	0.5021±0.0204	0.1874±0.0126
pFedMe	0.7418±0.0235	0.8568±0.0204	0.4860±0.0067	0.2084±0.0068	0.8297±0.0070	0.8320±0.0162	0.5588±0.0110	0.2324±0.0087
Setting 2	Minority Model				Majority Model			
Method\Data	F-MNIST	MNIST	CIFAR-10	DS-ImageNet	F-MNIST	MNIST	CIFAR-10	DS-ImageNet
Bi-level	**0.7754±0.0082**	**0.8580±0.0283**	**0.5148±0.0051**	**0.2734±0.0052**	**0.8332±0.0063**	**0.8384±0.0196**	**0.5979±0.0044**	**0.2694±0.0104**
Local-train	0.6763±0.0116	0.7874±0.0129	0.3159±0.0110	0.1260±0.0033	0.7343±0.0104	0.7650±0.0146	0.3594±0.0086	0.1276±0.0110
FedAvg	0.5686±0.0211	0.5809±0.0405	0.2870±0.0106	0.1676±0.0090	0.7824±0.0062	0.7746±0.0121	0.5431±0.0087	0.2542±0.0089
Ditto	0.7204±0.0059	0.7841±0.0217	0.3951±0.0121	0.1536±0.0178	0.7889±0.0093	0.7841±0.0089	0.4946±0.0131	0.1724±0.0068
pFedMe	0.7207±0.0130	0.7506±0.0224	0.4422±0.0131	0.1936±0.0113	0.8091±0.0044	0.7911±0.0283	0.5574±0.0104	0.2146±0.0081
Setting 3	Minority Model				Majority Model			
Method\Data	F-MNIST	MNIST	CIFAR-10	DS-ImageNet	F-MNIST	MNIST	CIFAR-10	DS-ImageNet
Bi-level	**0.7726±0.0170**	**0.8736±0.0055**	**0.5235±0.0060**	**0.2680±0.0072**	**0.8234±0.0119**	**0.8342±0.0135**	**0.5861±0.0064**	**0.2788±0.0142**
Local-train	0.6762±0.0126	0.7803±0.0204	0.3061±0.0107	0.1162±0.0054	0.7262±0.0074	0.7698±0.0100	0.3632±0.0083	0.1224±0.0071
FedAvg	0.5940±0.0211	0.6225±0.0195	0.3084±0.0099	0.1756±0.0088	0.7702±0.0143	0.7632±0.0171	0.5705±0.0123	0.2676±0.0074
Ditto	0.6856±0.0247	0.7452±0.0160	0.4184±0.0149	0.1626±0.0093	0.7628±0.0115	0.7742±0.0167	0.4986±0.0175	0.1792±0.0133
pFedMe	0.7123±0.0239	0.7203±0.0201	0.4535±0.0068	0.1938±0.0144	0.7871±0.0104	0.7650±0.0341	0.5563±0.0038	0.2188±0.0122
Setting 4	Minority Model				Majority Model			
Method\Data	F-MNIST	MNIST	CIFAR-10	DS-ImageNet	F-MNIST	MNIST	CIFAR-10	DS-ImageNet
Bi-level	**0.7658±0.0117**	**0.8588±0.0133**	**0.5172±0.0107**	**0.2698±0.0115**	**0.8160±0.0051**	**0.8455±0.0064**	**0.5935±0.0074**	**0.2678±0.0062**
Local-train	0.6809±0.0079	0.7645±0.0270	0.3086±0.0052	0.1148±0.0059	0.7012±0.0066	0.7476±0.0282	0.3709±0.0079	0.1172±0.0087
FedAvg	0.5648±0.0481	0.5475±0.0205	0.3612±0.0181	0.1722±0.0090	0.7644±0.0258	0.7577±0.0127	0.5529±0.0077	0.2600±0.0068
Ditto	0.6974±0.0130	0.7176±0.0185	0.3970±0.0105	0.1538±0.0077	0.7646±0.0085	0.7608±0.0063	0.4986±0.0102	0.1744±0.0159
pFedMe	0.7035±0.0187	0.6932±0.0140	0.4548±0.0069	0.1794±0.0114	0.7910±0.0110	0.7688±0.0171	0.5489±0.0060	0.2232±0.0063

Figure 6: Comparison in test accuracy for the majority group vs number of synchronizations.

Figure 7: Comparison in test accuracy for the minority group vs number of points processed.

Figure 8: Comparison in test accuracy for the majority group vs number of points processed.