A PAC-Bayesian Generalization Bound for Equivariant Networks

Arash Behboodi[*]
Qualcomm AI Research, Amsterdam[†]
behboodi@qti.qualcomm.com

Gabriele Cesa[*]
Qualcomm AI Research, Amsterdam[†]
AMLab, University of Amsterdam
gcesa@qti.qualcomm.com

Taco Cohen
Qualcomm AI Research, Amsterdam[†]
tacos@qti.qualcomm.com

Abstract

Equivariant networks capture the inductive bias about the symmetry of the learning task by building those symmetries into the model. In this paper, we study how equivariance relates to generalization error utilizing PAC Bayesian analysis for equivariant networks, where the transformation laws of feature spaces are determined by group representations. By using perturbation analysis of equivariant networks in Fourier domain for each layer, we derive norm-based PAC-Bayesian generalization bounds. The bound characterizes the impact of group size, and multiplicity and degree of irreducible representations on the generalization error and thereby provide a guideline for selecting them. In general, the bound indicates that using larger group size in the model improves the generalization error substantiated by extensive numerical experiments.

1 Introduction

Equivariant networks are widely believed to enjoy better generalization properties than their fully-connected counterparts by including an inductive bias about the invariance of the task. A canonical example is a convolutional layer for which the translation of the input image (over $\mathbb{R}^2$) leads to translation of convolutional features. This construction can be generalized to actions of other groups including rotation and permutation. Intuitively, the inductive bias about invariance and equivariance helps to focus on features that matter for the task, and thereby help the generalization. Recently, many works provide theoretical support for this claim mainly for general equivariant and invariant models. In this work, however, we focus on a class of equivariant neural networks built using representation theoretic tools. As in Shawe-Taylor and Wood [1996], Cohen and Welling [2016b], Weiler et al. [2018a], this approach can build general equivariant networks using irreducible representations of a group. An open question is about how to design feature spaces and combine representations. Besides, the question of impact of these choices on generalization remain. On the other hand, while it is rather well understood how to build a G equivariant model when G is finite, the choice of the group G to consider is not always obvious. Indeed, using a subgroup $H < G$ smaller than G is usually sufficient to achieve significant improvements over a non-equivariant baseline. Moreover, the symmetries G of the data are often continuous, which means G-equivariance can not be built into the model using a group convolution design as in Cohen and Welling [2016a]. In such cases, the most successful

---

[*]equal contribution

[†]Qualcomm AI Research is an initiative of Qualcomm Technologies, Inc.

approaches approximate it with a discrete subgroup $H < G$, e.g. see Weiler and Cesa [2019] for the $G = O(2)$ case. In addition, the choice of the equivariance group H can affect either the complexity or the expressiveness of the model and, therefore, its generalization capability, depending on how the architecture is adapted. For instance, if one can not freely increase the model size, using a larger group H only increases the parameters sharing but reduces the expressiveness of the model. For this reason, using the whole group G might not be beneficial even when feasible. In that line, we consider how the choice of group size affects generalization.

Contributions In this paper, we utilize PAC Bayes framework to derive generalization bounds on equivariant networks. Our focus is on equivariant networks built following Weiler et al. [2018a], Cohen and Welling [2016b]. We combine the representation theoretic framework of Weiler and Cesa [2019] with PAC-Bayes framework of Neyshabur et al. [2018] to get generalization bounds on equivariant networks as a function of irreducible representations (irreps) and their multiplicities. Different from previous PAC-Bayes analysis, we derive the perturbation analysis in Fourier domain. Without this approach, as we will show, the naive application of Neyshabur et al. [2018] lead to degenerate bounds. As part of the proof, we derive a tail bound on the spectral norm of random matrices characterized as direct sum of irreps. Note that for building equivariant networks, we need to work with real representations and not complex ones, which are conventionally studied in representation theory. The obtained generalization bound provide new insights about the effect of design choices on the generalization error verified via numerical results. To the best of our knowledge, this is the first generalization bound for equivariant networks with an explicit equivariant structure. We conduct extensive experiments to verify our theoretical insights, as well as some of the previous observations about neural networks.

2 Related Works

Equivariant networks. In machine learning, there has been many works on incorporating symmetries and invariances in neural network design. This class of models has been extensively studied from different perspectives Shawe-Taylor [1993, 1989], Kondor and Trivedi [2018], Cohen et al. [2018], Mallat [2012], Dieleman et al. [2016], Cohen and Welling [2016a,b], Worrall et al. [2017], Weiler et al. [2018b,a], Thomas et al. [2018], Bekkers et al. [2018], Weiler and Cesa [2019], Bekkers [2020], Defferrard et al. [2019], Finzi et al. [2020], Weiler et al. [2021], Cesa et al. [2022] to mention only some. Indeed, the most recent developments in the field of equivariant networks suggest a design which enforces equivariance not only at a global level Laptev et al. [2016] but also at each layer of the model. An interesting question is how a particular choice for designing equivariant networks affect its performance and in particular generalization. A related question is whether including this inductive bias helps the generalization. The authors in Sokolić et al. [2017a] consider general invariant classifiers and obtain robustness based generalization bounds, as in Sokolić et al. [2017b], assuming that data transformation changes the inputs drastically. For finite groups, they reported a scaling of generalization error with $1/\sqrt{|H|}$ where $|H|$ is the cardinality of underlying group. In Elesedy [2022], the gain of invariant or equivariant hypotheses was studied in PAC learning framework, where the gain in generalization is attributed to shrinking the hypothesis space to the space of orbit representatives. A similar argument can be found in Sannai et al. [2021], where it is shown that the invariant and equivariant models effectively operate on a shrunk space called Quotient Feature Space (QFS), and the generalization error is proportional to its volume. They generalize the result of Sokolić et al. [2017a] and relax the robustness assumption, although their bound shows suboptimal exponent for the sample size. Similar to Sokolić et al. [2017a], they focus on scaling improvement of the generalization error with invariance and do not consider computable architecture dependent bounds. The authors in Lyle et al. [2020] use PAC-Bayesian framework to study the effect of invariance on generalization, although do not obtain any explicit bound. Bietti and Mairal [2019] studies the stability of models equivariant to compact groups from a RKHS point of view, relating this stability to a Rademacher complexity of the model and a generalization bound. The authors in Elesedy and Zaidi [2021] provided the generalization gain of equivariant models more concretely and reported VC-dimension analysis. Subsequent works also considered the connection of invariance and generalization Zhu et al. [2021]. In contrast with these models, we consider the equivariant networks with representation theoretic construction. Beyond generalization, we are interested in getting design insights from the bound.

Generalization error for neural networks. A more general study of generalization error in neural networks, however, has been ongoing for some time with huge body of works on the topic. We refer to some highlights here. A challenge for this problem was brought up in Zhang et al. [2017], where it was shown using the image-net dataset with random labels, that the generalization error can be arbitrarily large for neural networks, as they achieve small training error but naturally cannot generalize because of random labels of images. Therefore, any complexity measure for neural networks should be consistent with the above observation. As a result, uniform complexity measures like VC-dimension do not satisfy this requirement as they provide a uniform measure over the whole hypothesis space.

In this light, recent works related the generalization errors to quantities like margin and different norms of weights for example in, among others, Wei and Ma [2019], Sokolić et al. [2017b], Neyshabur et al. [2018], Arora et al. [2018], Bartlett et al. [2017], Golowich et al. [2018], Dziugaite and Roy [2018a], Long and Sedghi [2019], Vardi et al. [2022], Ledent et al. [2021], Valle-Pérez and Louis [2020]. Some of these bounds are still vacuous or dimension-dependent (see Jiang et al. [2020] for detailed experimental investigation and Nagarajan and Kolter [2019], Koehler et al. [2021], Negrea et al. [2021] for follow-up discussions on uniform complexity measures). Class of convolutional neural networks are particularly relevant as they can be considered as a special case of equivariant models. These models are discussed in Pitas et al. [2019], Long and Sedghi [2019], Vardi et al. [2022], Ledent et al. [2021]. We choose PAC Bayesian framework for deriving generalization bounds. There are many works on PAC Bayesian bounds for neural networks Neyshabur et al. [2018], Biggs and Guedj [2022], Dziugaite and Roy [2017, 2018b,a], Dziugaite et al. [2020] ranging from randomized and deterministic bounds to choosing priors for non-vacuous bounds. Our method combines the machinery of Neyshabur et al. [2018] with representation theoretic tools.

3 Background

3.1 Preliminaries and notation

We fix some notations first. We consider a classification task with input space $\mathcal{X}$ and output space $\mathcal{Y}$. The input is assumed to be bounded in ℓ_2-norm by B. The data distribution, over $\mathcal{X} \times \mathcal{Y}$, is denoted by $\mathcal{D}$. The hypothesis space, $\mathcal{H}$, consists of all functions realized by a L-layer general neural network with 1-Lipschitz homogeneous activation functions[3]. The network function, $f_{\mathbb{W}}(\cdot)$ with fixed architecture, is specified by its parameters $\mathbb{W} := (\boldsymbol{W}_1, \dots, \boldsymbol{W}_L)$. The operations on $\mathbb{W}$ are extended from the underlying vector spaces of weights. For any function $f(\cdot)$, the k'th component of the function is denoted by $f(\boldsymbol{x})[k]$ throughout the text. For a loss function $\mathrm{L} : \mathcal{H} \times \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$, the empirical loss is defined by $\hat{\mathcal{L}}(f) := \frac{1}{m} \sum_{i=1}^{m} \mathrm{L}(f, \boldsymbol{x}_i, y_i))$ and the true loss is defined by $\mathcal{L}(f) = \mathbb{E}_{(\boldsymbol{x},y) \sim \mathcal{D}}[\mathrm{L}(f, \boldsymbol{x}, y))]$. For classification, the margin loss is given by

$$\mathcal{L}_\gamma(f_{\mathbb{W}}) = \mathbb{P}_{(\boldsymbol{x},y) \sim \mathcal{D}} \left(f_{\mathbb{W}}(\boldsymbol{x})[y] \leq \gamma + \max_{j \neq y} f_{\mathbb{W}}(\boldsymbol{x})[j] \right). \tag{1}$$

The true loss of a given classifier is then given by $\mathcal{L}(f_{\mathbb{W}}) = \mathcal{L}_0(f_{\mathbb{W}})$. The empirical margin loss is similarly defined and denoted by $\hat{\mathcal{L}}_\gamma$. In this work, we consider invariance with respect to transformations modeled as the action of a compact group G. See Supplementary 7 for a brief introduction to group theory and the concepts that we will use throughout this paper.

3.2 Equivariant Neural networks

We provide a concise introduction to equivariant networks here with more details given in Supplementary 8. We adopt a representation theoretic framework for building equivariant models.

Example. Consider a square integrable complex valued function f with bandwidth B defined on 2D-rotation group $SO(2)$ and represented using its Fourier series as $f(\theta) = \sum_{n=0}^{B} a_n e^{in\theta}$. Consider an equivariant linear functional of f mapping it to another function on $SO(2)$. In Fourier space, the linear transformation is given by $\boldsymbol{W}\boldsymbol{a}$ with $\boldsymbol{a}$ as the Fourier coefficient vector $\boldsymbol{a} = (a_0, \dots, a_B)$. The action of rotation group on the input and output space is simply given by $f(\theta) \to f(\theta - \theta_0)$ with θ_0 as the rotation angle, or equivalently in Fourier space as a linear transformation $\boldsymbol{a} \to \rho(\theta_0)\boldsymbol{a}$

---

[3]The function $\sigma(\cdot)$ is homogeneous if and only if for all $\boldsymbol{x}$ and $\lambda \in \mathbb{R}$, we have $\sigma(\lambda \boldsymbol{x}) = \lambda \sigma(\boldsymbol{x})$.

Figure 1: An equivariant layer maps the coefficients corresponding to each frequency (irrep) ψ to a new set of coefficients based on the matrix $\widehat{W}_l(\psi)$, which include blocks of $\widehat{W}_l(\psi, i, j)$ with $i \in [m_{l-1,\psi}]$ and $j \in [m_{l,\psi}]$.

with $\rho(\theta_0) = \mathrm{diag}\left(1, e^{-i\theta_0} \ldots e^{-iB\theta_0}\right)$. Since it holds for all a, Equivariance condition means $W\rho(\theta_0) = \rho(\theta_0)W$ for all θ_0, which implies that W should be diagonal. A general feature space can be built by stacking multiple linear transformations W, and more flexibly, by letting the transformations acting only on some of the frequencies (for example, a transformation that acts only on a_B for frequency B). This machinery can be generalized to represent equivariant networks with respect to any compact group, by using irreducible representations (irreps) of the group in place of complex exponential $e^{in\theta}$ of this example. The key is to notice that $e^{in\theta}$'s are irreps of $SO(2)$.

General Model. Consider an Multi-Layer Perceptron (MLP) with L layers with the layer $l \in [L]$ given by the matrix $W_l \in \mathbb{R}^{c_l \times c_{l-1}}$. The action of the group H on layer l is determined by a linear transformation ρ_l, which is called the group representation on $\mathbb{R}^{c_l}$ (see section 8 for more details). A group element $h \in H$ linearly transforms the layer l by the matrix $\rho_l(h)$. The matrix W_l is equivariant w.r.t. the representations ρ_l and ρ_{l-1} acting on its input and output if and only if for all $h \in H$, we have:

$$W_l \rho_{l-1}(h) = \rho_l(h) W_l \quad .$$

Representation theory provides a way of characterizing equivariant layers using Maschke's theorem and Schur's lemma. The key step is to start from characterizing the irreducible representations (irreps) $\{\psi\}$ of the group H. As we will see, irreps are closely related to generalization of Fourier analysis for functions defined on H. Maschke's theorem implies that the representation ρ_l decomposes into direct sum of irreps as $\rho_l = Q_l \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right) Q_l^{-1}$, where $m_{l,\psi}$ is the number of times the irrep ψ is present in the representation ρ_l, that is, its multiplicity. Each ψ is a $\dim_\psi \times \dim_\psi$ matrix, and the direct sum is effectively a block diagonal matrix with matrix irreps ψ on the diagonal. Through this decomposition, we can parameterize equivariant networks in terms of irreps, that is in Fourier space. Defining $\widehat{W}_l = Q_l^{-1} W_l Q_{l-1}$, the equivariance condition writes as

$$\widehat{W}_l \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l-1,\psi}} \psi \right) = \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right) \widehat{W}_l.$$

The block diagonal structure of $\left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right)$ induces a similar structure on $\widehat{W}_l$ (see Figure 6). By $\widehat{W}_l(\psi_2, j, \psi_1, i)$ denote the block in $\widehat{W}$ that relates i'th instance of ψ_1 to j'th instance of ψ_2, with $i \in [m_{l-1,\psi_1}]$ and $j \in [m_{l,\psi_2}]$, as:

$$\widehat{W}_l(\psi_2, j, \psi_1, i)\psi_1 = \psi_2 \widehat{W}_l(\psi_2, j, \psi_1, i).$$

Schur's lemma helps us to characterize the equivariant kernels. Note for neural network implementation, we need to work with a version of Schur's lemma for real-valued representations given in Supplementary 8. Schur's lemma implies that if $\psi_1 \neq \psi_2$, the block needs to be zero to guarantee equivariance. Otherwise, it needs to have one of the three forms in Supplementary 8, depending on

the *type* of ψ_1. To simplify the notation, we write $\widehat{W}_l(\psi, j, \psi, i)$ as $\widehat{W}_l(\psi, j, i)$. As it is shown in Supplementary 8, the matrix $\widehat{W}_l(\psi, j, i)$ can be written as the linear sum

$$\widehat{W}_l(\psi, j, i) = \sum_{k=1}^{c_\psi} \widehat{w_{l,i,j}}(\psi)_k B_{l,\psi,i,j,k} \tag{2}$$

where $\{\widehat{w_{l,i,j}}(\psi)_k\}_{k=1}^{c_\psi}$ are learnable parameters with fixed matrices $B_{l,\psi,i,j,k}$. The value of c_ψ is either 1,2 or 4. The structure of $B_{l,\psi,i,j,k}$ changes according to each type c_ψ. Each layer is parameterized by the matrices $\widehat{W}_l(\psi)$, which include c_ψ parameters $\widehat{w_{l,i,j}}(\psi) \in \mathbb{R}^{c_\psi}$ with $i \in [m_{l-1,\psi}]$ and $j \in [m_{l,\psi}]$. Therefore, for each layer, there will be $\sum_\psi m_{l,\psi} m_{l-1,\psi} c_\psi$ parameters with the width of the layer l given as $c_l = \sum_\psi m_{l,\psi} \dim_\psi$. Note that, if $x_l = W_l x_{l-1}$, then $\widehat{x}_l = \widehat{W}_l \widehat{x}_{l-1}$, where $\widehat{x}_l = Q_l^{-1} x_l$. The non-linearities should satisfy equivariance condition to have full end-to-end equivariance. An important question is which representation ρ_l of H should be used for each layer. For the input, the representation is fixed by the input space and its structure. However, the intermediate layers can choose ρ_l by selecting irreps and their multiplicity. We explain how this choice impacts generalization bounds. We consider a general equivariant network denoted by $f_\mathbb{W}$ with the parameters of l'th layer given by $\widehat{w_{l,i,j}}(\psi)$ for all irreps ψ, $i \in [m_{l-1,\psi}]$ and $j \in [m_{l,\psi}]$.

4 PAC-Bayesian bound

We derive PAC-Bayes bounds for equivariant networks $f_\mathbb{W}$ presented in the previous section. Although the proof follows a similar strategy as in Neyshabur et al. [2018], it differs in two important aspects. First, the weights of equivariant MLPs have specific structure requiring reworking proof steps. Second, we carry out the analysis in Fourier domain. In Supplementary 11, we also show that naively applying norm-bounds of Neyshabur et al. [2018] cannot explain the generalization behaviour of equivariant networks. A simple version of our result is given below.

Theorem 4.1 (Homogeneous Bounds for Equivariant Networks). *For any equivariant network, with high probability we have:*

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \le \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) + \tilde{\mathcal{O}}\left(\sqrt{ \frac{\prod_l \|W_l\|_2^2}{\gamma^2 m \eta} \left(\sum_{l=1}^L \sqrt{M(l,\eta)} \right)^2 \left(\sum_l \frac{\sum_{\psi,i,j} \left\| \widehat{W}_l(\psi, i, j) \right\|_F^2 / \dim_\psi}{\|W_l\|_2^2} \right) } \right)$$

where $\eta \in (0, 1)$ and

$$M(l, \eta) := \log\left(\frac{\sum_{l=1}^L \sum_\psi m_{l,\psi}}{1 - \eta} \right) \max_\psi \left(5 m_{l-1,\psi} m_{l,\psi} c_\psi \right). \tag{3}$$

The complete version of the theorem is given in Theorem 9.1.

We comment briefly on the proof steps. PAC-Bayesian generalization bounds start with defining a prior P and posterior Q over the network parameters. The network is drawn randomly using Q, which is also used to evaluate the average loss and generalization error. The PAC-Bayesian bound provides a bound on the average generalization error of networks randomly drawn from Q (Section 4.1). The next step is to de-randomize the bound and to obtain the generalization error for a specific instance $f_\mathbb{W} \in \mathcal{H}$. Among different strategies, we follow the perturbation based method of Neyshabur et al. [2018]. The main idea is to carefully define a posterior Q, such that the networks randomly drawn from Q have small *distance* with the specific $f_\mathbb{W}$, and therefore the loss averaged over Q can be used to bound the loss of $f_\mathbb{W}$. One way to choose Q is to define a Gaussian distribution around the parameters $\mathbb{W}$ and choose the variance to control the output perturbation (Section 4.2).

In what follows we provide more details about these steps.

4.1 PAC-Bayesian Bounds for Randomized Networks

The starting point is the PAC-Bayes theorem Langford and Shawe-taylor [2002], McAllester [1998], which provides a generalization bound on randomized classifiers. If the function is chosen randomly

from a distribution Q, then define the average losses as $\mathcal{L}(Q) = \mathbb{E}_{f \sim Q} \mathcal{L}(f)$ and $\hat{\mathcal{L}}(Q) = \mathbb{E}_{f \sim Q} \hat{\mathcal{L}}(f)$. We use the following version from Germain et al. [2009].

Lemma 4.2. *For any hypothesis space $\mathcal{H}$, let P be a probability distribution on $\mathcal{H}$. Then for all $\delta \in (0, 1]$, with probability $1 - \delta$, for all Q on $\mathcal{H}$, we have:*

$$\mathcal{L}(Q) \leq \hat{\mathcal{L}}(Q) + \sqrt{\frac{D(Q \| P) + \log(\xi(m)/\delta)}{2m}}. \tag{4}$$

where $\xi(m) := \sum_{k=0}^{m} \binom{m}{k} (k/m)^k (1 - k/m)^{m-k}$.

In PAC-Bayes vernacular, the distributions P and Q are called *a prior* and *a posterior* distribution on $\mathcal{H}$. This indicates that the distribution P and Q are chosen independently with Q typically chosen based on the training data and the obtained model, and P chosen independently based on any information available before training. Choosing an appropriate prior plays a central role in getting non-vacuous bounds. For instance in Dziugaite and Roy [2017], the prior P is chosen using a separate dataset not used during training. Lemma 4.2 provides immediately a generalization bound for randomized equivariant classifiers. The hypothesis space is parameterized by the network parameters $\mathbb{W}$, which contains kernel parameters $\widehat{\boldsymbol{w}_{l,i,j}}(\psi)$. We choose a prior distribution P as zero-mean normal with covariance matrix $\sigma^2 \boldsymbol{I}$ over the parameters. The posterior Q is chosen as normal distribution with the final parameters $\widehat{\boldsymbol{w}_{l,i,j}}(\psi)$ as mean value and the same covariance matrix $\sigma^2 \boldsymbol{I}$. The KL-divergence in the PAC-Bayesian theorem is then given by:

$$D(Q \| P) = \frac{\sum_{l,\psi,i,j} \left\| \widehat{\boldsymbol{w}_{l,i,j}}(\psi) \right\|_2^2}{2\sigma^2}. \tag{5}$$

The generalization bound depends depends on the sum of the norm of kernels. In the next step, we de-randomize the bound to get a bound on the generalization error of $f_{\mathbb{W}}$.

4.2 De-Randomization and Perturbation Bounds

Our de-randomization technique follows closely that of Neyshabur et al. [2018]. We provide the sketch of derivations here, and the full proof is relegated to the supplementary materials. To de-randomize the previous bound, a common step is to choose Q such that the probability of margin violation is controlled. Let $\mathcal{S}_{\mathbb{W}}$ be defined as:

$$\mathcal{S}_{\mathbb{W}} := \{ h \in \mathcal{H} : \| h - f_{\mathbb{W}} \|_\infty \leq \gamma/4 \}. \tag{6}$$

Any function on this set can change the margin of $f_{\mathbb{W}}$ at most by $\gamma/2$. Therefore, for any h, $\mathcal{L}(f_{\mathbb{W}}) \leq \mathcal{L}_{\gamma/2}(h)$, which implies $\mathcal{L}(f_{\mathbb{W}}) \leq \mathcal{L}_{\gamma/2}(Q)$ if Q is supported only on this set. Similarly $\hat{\mathcal{L}}_{\gamma/2}(Q) \leq \hat{\mathcal{L}}_\gamma(f_{\mathbb{W}})$. To choose Q, first, we characterize the output perturbation of equivariant networks for a given input perturbation. Next, we determine the variance σ such that the output perturbation is bounded by $\gamma/4$ with probability[4] $1/2$. The first step is the perturbation analysis.

Lemma 4.3 (Perturbation Bound). *For a neural network $f_{\mathbb{W}}(\cdot)$ with input space of ℓ_2-norm bounded by B, and for any weight perturbations $\mathbb{U} = (\boldsymbol{U}_1, \ldots, \boldsymbol{U}_L)$, we have:*

$$\| f_{\mathbb{W}+\mathbb{U}}(\boldsymbol{x}) - f_{\mathbb{W}}(\boldsymbol{x}) \|_2 \leq eB \left(\prod_{i=1}^{L} \| \boldsymbol{W}_i \|_2 \right) \sum_{i=1}^{L} \frac{\| \boldsymbol{U}_i \|_2}{\| \boldsymbol{W}_i \|_2}. \tag{7}$$

For equivariant networks, where the weights are parameterized as in eq. 2, the spectral norm of the perturbation $\boldsymbol{U}_l$ is bounded as

$$\| \boldsymbol{U}_l \|_2 \leq \sqrt{\max_\psi \max_{1 \leq i \leq m_{l-1,\psi}} m_{l-1,\psi} \frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{U}_l}(\psi, j, i) \right\|_F^2} \tag{8}$$

The first inequality is already given in Neyshabur et al. [2018]. The inequality 8 will be shown in supplementary materials.

---

[4]This value can be changed to any probability with proper adjustments. See the supplementary materials.

The perturbation bound already suggests the benefit of equivariant Kernels. As it will be shown in Supplementary materials, the equivariant kernels w.r.t. ψ satisfy

$$\|\boldsymbol{W}\|_2 = \frac{1}{\sqrt{\dim_\psi}} \|\boldsymbol{W}\|_F, \tag{9}$$

which is the smallest possible norm (compare with general relation for $\dim_\psi \times \dim_\psi$ matrices: $\|\boldsymbol{W}\|_2 \geq \frac{1}{\sqrt{\dim_\psi}} \|\boldsymbol{W}\|_F$). Therefore, the term $\frac{1}{\dim_\psi} \left\|\widehat{\boldsymbol{U}_l}(\psi, j, i)\right\|_F^2$ in the perturbation bound is already tight.

The perturbation bound helps defining a posterior Q by adding zero-mean Gaussian perturbations $\widehat{u_{l,i,j}}(\psi)_k$ of variance σ^2 to the learnable parameters $\widehat{w_{l,i,j}}(\psi)_k$. The output perturbation is given by the above theorem in terms of norm of $\boldsymbol{U}_i$. The following lemma controls the norm of this random matrix, and thereby the output perturbation.

Lemma 4.4. *Consider a random equivariant matrix $\boldsymbol{U}_l$ defined by i.i.d. Gaussian $N(0, \sigma^2)$ choice of $\widehat{u_{l,i,j}}(\psi)_k$ in eq. 2. We have:*

$$\mathbb{P}\left(\|\boldsymbol{U}_l\|_2 \geq \sigma\sqrt{\max_\psi\left(5m_{l-1,\psi}m_{l,\psi}c_\psi t\right)}\right) \leq \left(\sum_\psi m_{l,\psi}\right)e^{-t} \tag{10}$$

The proof is given in supplementary materials. With Lemma 4.4, we can choose the variance σ^2 such that the output perturbation does not violate the intended margin. Using an union bound over all layers, one finds that by choosing $t = \log\left(2\sum_l\sum_\psi m_{l,\psi}\right)$, with probability at least $\frac{1}{2}$ the following inequality holds for every layer $l \in [L]$:

$$\|\boldsymbol{U}_l\|_2 < \sigma\sqrt{\max_\psi 5m_{l-1,\psi}m_{l,\psi}c_\psi}\sqrt{\log\left(2\sum_l\sum_\psi m_{l,\psi}\right)} \tag{11}$$

We can use this bound jointly with the perturbation bound eq. 7 to choose σ such that the margin is below $\gamma/4$ with high probability. Note that the σ chosen in this way is a function of learned weights. This cannot be the case because of the prior P should be fixed prior to learning. A trick for circumventing this issue is to select many priors that can adequately cover the space of possible weights, and take the union bound over it. The complete proof of the theorem is given in Supplementary 9, where we also consider special cases of this generalization bound for group convolutional networks Cohen and Welling [2016a].

Remarks on the generalization error from the theory. The generalization error of Theorem 4.1 contains multiple terms. First, the term η is a hyperparameter, which will be fixed to $1/2$ for the rest. The term $M(l, \eta)$ encodes the multiplicity and type of used irreps and their impact on the generalization error. Let the hidden layer dimension $c_l = \sum_\psi \dim_\psi m_{l,\psi}$ be fixed to a constant. For Abelian groups, all real irreps are at most 2-dimensional, therefore, restricting the network to 2-dimensional irreps, we have:

$$M(l, \eta) = \log\left(\frac{\sum_{l=1}^L c_l}{2(1-\eta)}\right)\max_\psi\left(5m_{l-1,\psi}m_{l,\psi}c_\psi\right).$$

The bound can be controlled further by appropriate choice of $m_{l-1,\psi}, m_{l,\psi}, c_\psi$, which favors smaller multiplicity and type c_ψ. Therefore, the bound favors using different irreps instead of repeating them. For non-abelian cases, irreps can have larger $\dim_\psi$, which keeping the layer dimension c_l fixed, amounts to smaller multiplicity. This can improve the sum $\sum_{l=1}^L\sum_\psi m_{l,\psi}$ in $M(l,\eta)$ and indicates a potential improved generalization for non-abelian groups. For dihedral groups, which is in general non-abelian, the real irreps are also at most two dimensional, however, of type $c_\psi = 1$. Note that for finite groups, there is a connection between irrep degrees and group order $|H| = \sum_\psi \dim_\psi^2/c_\psi$. The impact of group size manifests itself in the whole generalization error term. We numerically verify this in the experimental results. The inverse dependence on margin γ is expected and similar to other

Figure 2: Comparison between the bound in Theorem 4.1 (1st row) and the one in Supplementary 11 (2nd row) on the synthetic $O(2)$ datasets with different frequencies. The bound from Supplementary 11 clearly does not capture the effect of group equivariance.

works for instance Neyshabur et al. [2018]. The rest of the terms contain different norms of neural network kernels. We discuss them more in connection with Neyshabur et al. [2018].

Comparison to Neyshabur et al. [2018]. There are two main differences with the bound in Neyshabur et al. [2018]. First, their bound has a norm dependence of $\tilde{\mathcal{O}}\left((\prod_l \|\boldsymbol{W}\|_2^2)(\sum_l \|\boldsymbol{W}\|_F^2 / \|\boldsymbol{W}\|_2^2)\right)$, while in our result, the Frobenius norm includes an additional scaling of $1/\dim_\psi$ in $\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2 / \dim_\psi$. This term is strictly smaller in our bound for $\dim_\psi > 1$. The second difference is the term $\left(\sum_l M(l,\eta)\right)^2$, which is replaced with $L^2 h \ln(Lh)$ where h is the layer width. In our case, the layer width is $c_l = \sum_\psi \dim_\psi m_{l,\psi}$. Choosing all layer widths c_l equal to h, we recover the $\ln(Lh)$ term from Neyshabur et al. [2018] in our bound. The remaining term is $O\left(\left(\sum_l \sqrt{m_{l-1,\psi} m_{l,\psi} c_\psi}\right)^2\right)$. As an example, assume the same multiplicity $m_{l,\psi} = m_l$ and same dimension $\dim_\psi = \dim$ for all irreps, that is, $h = N \dim m_l$ with N being the number of used irreps. Then, we get a term as $O\left(h^2 L^2 / (N \dim)^2\right)$. As long as the multiplicity m_l is smaller than $N \dim$, i.e. number of used irreps times their dimension, our bound is strictly tighter. Intuitively, this means that using more diverse irreps leads to better generalization. See Section 11 for more detailed discussions.

5 Experiments

In this section, we also validate the theoretical insights derived in the previous section. More experiments and discussions are included in Supplementary 12.

We have used datasets based on natural images and synthetic data. In all experiments, we consider data symmetries G which are subgroups of the group $O(2)$ of planar rotations and reflections. This includes a number of finite/continuous and commutative/non-commutative groups, in particular C_N (N rotations), D_N (N rotations and reflections), $SO(2)$ (continuous planar rotations) and $O(2)$ itself (see Supplementary 7). Moreover, all models are based on group convolution over a finite subgroup $H < G$ Cohen and Welling [2016a]. This means that the features of the layer l have shape $|H| \times c_l^H$, that is, c_l^H channels of dimension $|H|$. For each layer l, we keep the total number of features (approximatively) constant when changing the group H. Models are trained until 99% of the training set is correctly classified with at least a margin γ. We used $\gamma = 10$ in the synthetic datasets and $\gamma = 2$ in the image ones.

First, we compare our result with the bound derived in Supplementary 11 using naive application of Neyshabur et al. [2018]. In Fig. 2 and Fig. 3, we compare these two bounds on 6 synthetic datasets with $O(2)$ and $SO(2)$ symmetries (one per column). Each dataset lives on high dimensional tori and $O(2)$ (or $SO(2)$) rotates the circles composing them, with different frequencies. We consider

Figure 3: Comparison between Theorem 4.1 (1st row) and the one in Supplementary 11 (2nd row) on the synthetic SO(2) datasets.

different maximum frequencies $F \in \{1, 2, \ldots, 6\}$ in the experiments. For example, in Fig. 2, the first plot on the left corresponds to a synthetic dataset with O(2) symmetry generating only using frequency 1. See Supplementary 12.1 for more details and visualization of the synthetic data.

The results in Fig. 2 and 3 indicate that the bound built using the strategy from Neyshabur et al. [2018] cannot explain the effect of equivariance on generalization. Conversely, the bound in Theorem 4.1, which uses the parametrization in the Fourier domain, correlates with the measured generalization error, suggesting that it can account for the effect of different groups on generalization. In addition, note that it also correlates with the saturation effect observed when using larger discrete groups H to approximate the continuous group $G = O(2)$. Indeed, on low frequency datasets, the bound tends to saturate faster as we increase the group size $|H|$. Conversely, on high frequency datasets, choosing a larger group H results in larger improvements in the estimated bound.

Remark 5.1. As it can be seen in our experiments in Fig 2 and 3, our bound is larger that the one in Neyshabur et al. [2018]. This is partially due to coarse approximation of Frobenius norm and the presence of the term h^2 in our bound. This dependency could be alleviated if we increase the number of used irreps in the group convolutional network, namely the larger group size. It is an interesting research direction to explore the optimal dependency on the width for equivariant networks.

In Fig. 4, we perform a larger study on the transformed MNIST datasets to investigate the simultaneous effect of the group size $|H|$, data augmentation and training set size m. For each training data size, the generalization error improves with larger equivariance group H.

We study the relation between the term $\sum_l \sqrt{M(l, \eta)}$ and the generalization error observed. Note that this term depends only on the architecture design, i.e. it is data and training agnostic. In Fig. 5, we show the correlation between these two quantities for two synthetic datasets. Both datasets have $G = O(2)$ symmetry, but the data rotates with frequencies up to $F = 1$ in the first case and $F = 6$ in the second. We observe that the term $\sum_l \sqrt{M(l, \eta)}$ correlates strongly with the generalization in the high-frequency dataset ($F = 6$), where the generalization benefits from increasing the size of the equivariance group. Conversely, in the low-frequency case ($F = 1$), we observe a saturation of the performance, as increasing the group size does not improve generalization. In this setting, the term $\sum_l \sqrt{M(l, \eta)}$ is only a weak predictor of generalization. From these observations, we interpret this term as an indicator of the expressiveness of a model. Particularly, when the irreps are chosen compatible with the dataset symmetries, the term $\sum_l \sqrt{M(l, \eta)}$ provides a good guideline for the architecture choice. However, similar to general neural networks, the generalization error is dependent on other factors as well, which can be for example norms of network weights.

6 Conclusion

We provided learning theoretic frameworks to study the effect of invariance and equivariance on the generalization error and provide guidelines for design choices. The obtained bounds provide

(a) $H = C_N, G = \mathrm{SO}(2)$ (b) $H = C_N, G = \mathrm{O}(2)$ (c) $H = D_N, G = \mathrm{O}(2)$

(d) $H = C_N, G = \mathrm{SO}(2)$(augment) (e) $H = C_N, G = \mathrm{O}(2)$ (augment) (f) $H = D_N, G = \mathrm{O}(2)$ (augment)

Figure 4: Generalization Error (GE) of different H-equivariant models ($H = C_N$ or $H = D_N$) on different G-MNIST datasets ($G = \mathrm{SO}(2)$ or $G = \mathrm{O}(2)$) for different training set sizes m.

(a) $G = \mathrm{O}(2)$, Frequency $F = 1$ (b) $G = \mathrm{O}(2)$, Frequency $F = 6$

Figure 5: Correlation between the term $\sum_l \sqrt{M(l, \eta)}$ and the Generalization Error (GE) of different H-equivariant models ($H = C_N$ or $H = D_N$) on the synthetic $\mathrm{O}(2)$ dataset with frequency $F = 1$ and $F = 6$, with a training set size $m = 3200$.

useful insights and guidelines for understanding these architectures. As limitation of our work, we show experimentally in supplementary materials that the obtained bounds do not decrease with larger training sizes m as expected. Nevertheless, for a fixed training set, the trend of the generalization bound correlates with the generalization error.

References

Sanjeev Arora, Rong Ge, Behnam Neyshabur, and Yi Zhang. Stronger Generalization Bounds for Deep Nets via a Compression Approach. In *International Conference on Machine Learning*, pages 254–263, 2018.

Peter L Bartlett, Dylan J Foster, and Matus J Telgarsky. Spectrally-normalized margin bounds for neural networks. In I. Guyon, U. V. Luxburg, S. Bengio, H. Wallach, R. Fergus, S. Vishwanathan, and R. Garnett, editors, *Advances in Neural Information Processing Systems 30*, pages 6240–6249. Curran Associates, Inc., 2017.

Erik J Bekkers. B-spline CNNs on lie groups. In *International Conference on Learning Representations*, 2020.

Erik J. Bekkers, Maxime W Lafarge, Mitko Veta, Koen A.J. Eppenhof, Josien P.W. Pluim, and Remco Duits. Roto-translation covariant convolutional networks for medical image analysis. In *International Conference on Medical Image Computing and Computer-Assisted Intervention (MICCAI)*, 2018.

Alberto Bietti and Julien Mairal. Group invariance, stability to deformations, and complexity of deep convolutional representations. *J. Mach. Learn. Res.*, 20(1):876–924, January 2019. ISSN 1532-4435.

Felix Biggs and Benjamin Guedj. Non-Vacuous Generalisation Bounds for Shallow Neural Networks. *arXiv:2202.01627 [cs, stat]*, February 2022. URL http://arxiv.org/abs/2202.01627. arXiv: 2202.01627.

Gabriele Cesa, Leon Lang, and Maurice Weiler. A program to build e(n)-equivariant steerable CNNs. In *International Conference on Learning Representations*, 2022. URL https://openreview.net/forum?id=WE4qe9xlnQw.

Taco S. Cohen and Max Welling. Group Equivariant Convolutional Networks. *arXiv:1602.07576 [cs, stat]*, February 2016a. arXiv: 1602.07576.

Taco S. Cohen and Max Welling. Steerable CNNs. In *ICLR 2017*, November 2016b.

Taco S. Cohen, Mario Geiger, and Maurice Weiler. A general theory of equivariant CNNs on homogeneous spaces. *arXiv preprint arXiv:1811.02017*, 2018.

Michaël Defferrard, Martino Milani, Frédérick Gusset, and Nathanaël Perraudin. DeepSphere: a graph-based spherical CNN. In *International Conference on Learning Representations*, 2019.

Sander Dieleman, Jeffrey De Fauw, and Koray Kavukcuoglu. Exploiting cyclic symmetry in convolutional neural networks. In *International Conference on Machine Learning (ICML)*, 2016.

Gintare Karolina Dziugaite and Daniel Roy. Entropy-SGD optimizes the prior of a PAC-Bayes bound: Generalization properties of Entropy-SGD and data-dependent priors. In *International Conference on Machine Learning*, pages 1377–1386, July 2018a.

Gintare Karolina Dziugaite and Daniel M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. *arXiv preprint arXiv:1703.11008*, 2017.

Gintare Karolina Dziugaite and Daniel M. Roy. Data-dependent PAC-Bayes priors via differential privacy. In *Advances in Neural Information Processing Systems*, pages 8430–8441, 2018b.

Gintare Karolina Dziugaite, Alexandre Drouin, Brady Neal, Nitarshan Rajkumar, Ethan Caballero, Linbo Wang, Ioannis Mitliagkas, and Daniel M. Roy. In search of robust measures of generalization. *Advances in Neural Information Processing Systems*, 33:11723–11733, 2020.

Bryn Elesedy. Group symmetry in PAC learning. In *ICLR 2022 Workshop on Geometrical and Topological Representation Learning*, page 9, 2022.

Bryn Elesedy and Sheheryar Zaidi. Provably strict generalisation benefit for equivariant models. In *International Conference on Machine Learning*, pages 2959–2969. PMLR, 2021.

Marc Finzi, Samuel Stanton, Pavel Izmailov, and Andrew Gordon Wilson. Generalizing convolutional neural networks for equivariance to lie groups on arbitrary continuous data. In *International Conference on Machine Learning*, pages 3165–3176. PMLR, 2020.

Pascal Germain, Alexandre Lacasse, François Laviolette, and Mario Marchand. PAC-Bayesian learning of linear classifiers. In *Proceedings of the 26th Annual International Conference on Machine Learning*, pages 353–360, 2009.

Noah Golowich, Alexander Rakhlin, and Ohad Shamir. Size-Independent Sample Complexity of Neural Networks. In *Conference On Learning Theory*, pages 297–299, July 2018.

Yiding Jiang, Behnam Neyshabur, Dilip Krishnan, Hossein Mobahi, and Samy Bengio. Fantastic Generalization Measures and Where to Find Them. In *International Conference on Learning Representations*, 2020.

Frederic Koehler, Lijia Zhou, Danica J. Sutherland, and Nathan Srebro. Uniform Convergence of Interpolators: Gaussian Width, Norm Bounds and Benign Overfitting. In *Advances in Neural Information Processing Systems*, May 2021. URL `https://openreview.net/forum?id=FyOhThdDBM`.

Risi Kondor and Shubhendu Trivedi. On the generalization of equivariance and convolution in neural networks to the action of compact groups. In *International Conference on Machine Learning (ICML)*, 2018.

John Langford and John Shawe-taylor. PAC-Bayes & Margins. In *Advances in Neural Information Processing Systems 15*. Citeseer, 2002.

Dmitry Laptev, Nikolay Savinov, Joachim M Buhmann, and Marc Pollefeys. Ti-pooling: transformation-invariant pooling for feature learning in convolutional neural networks. In *Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition*, pages 289–297, 2016.

B. Laurent and P. Massart. Adaptive estimation of a quadratic functional by model selection. *Annals of Statistics*, 28(5):1302–1338, October 2000. ISSN 0090-5364, 2168-8966. doi: 10.1214/aos/1015957395. Publisher: Institute of Mathematical Statistics.

Antoine Ledent, Waleed Mustafa, Yunwen Lei, and Marius Kloft. Norm-based generalisation bounds for deep multi-class convolutional neural networks. In *35th AAAI Conference on Artificial Intelligence*, pages 8279–8287. AAAI Press, 2021.

Philip M. Long and Hanie Sedghi. Size-free generalization bounds for convolutional neural networks. *arXiv:1905.12600 [cs, math, stat]*, May 2019. arXiv: 1905.12600.

Clare Lyle, Mark van der Wilk, Marta Kwiatkowska, Yarin Gal, and Benjamin Bloem-Reddy. On the benefits of invariance in neural networks. *arXiv preprint arXiv:2005.00178*, 2020.

Stéphane Mallat. Group invariant scattering. *Communications on Pure and Applied Mathematics*, 65 (10):1331–1398, 2012.

David A. McAllester. Some PAC-Bayesian Theorems. In *Machine Learning*, pages 230–234. ACM Press, 1998.

Vaishnavh Nagarajan and J. Zico Kolter. Uniform convergence may be unable to explain generalization in deep learning. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d\textquotesingle Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems 32*, pages 11615–11626. Curran Associates, Inc., 2019.

Jeffrey Negrea, Gintare Karolina Dziugaite, and Daniel M. Roy. In Defense of Uniform Convergence: Generalization via derandomization with an application to interpolating predictors. Technical Report arXiv:1912.04265, arXiv, September 2021. URL `http://arxiv.org/abs/1912.04265`.

Behnam Neyshabur, Srinadh Bhojanapalli, and Nathan Srebro. A PAC-Bayesian Approach to Spectrally-Normalized Margin Bounds for Neural Networks. In *International Conference on Learning Representations*, 2018.

Konstantinos Pitas, Andreas Loukas, Mike Davies, and Pierre Vandergheynst. Some limitations of norm based generalization bounds in deep neural networks. *arXiv:1905.09677 [cs, stat]*, May 2019. arXiv: 1905.09677.

Siamak Ravanbakhsh. Universal equivariant multilayer perceptrons. *arXiv preprint arXiv:2002.02912*, 2020.

Akiyoshi Sannai, Masaaki Imaizumi, and Makoto Kawano. Improved generalization bounds of group invariant/equivariant deep networks via quotient feature spaces. In *Uncertainty in Artificial Intelligence*, pages 771–780. PMLR, 2021.

Jean-Pierre Serre. *Linear representations of finite groups*. Springer, 1977.

J. Shawe-Taylor. Building symmetries into feedforward networks. In *1989 First IEE International Conference on Artificial Neural Networks, (Conf. Publ. No. 313)*, page 158–162, Oct 1989.

J Shawe-Taylor. Symmetries and discriminability in feedforward network architectures. *IEEE Trans. Neural Netw.*, page 1–25, 1993. ISSN 1045-9227.

John Shawe-Taylor and Jeffrey Wood. Representation theory and invariant neural networks. *Discrete Applied Mathematics*, 69(1-2):33–60, August 1996. Publisher: North-Holland.

Jure Sokolić, Raja Giryes, Guillermo Sapiro, and Miguel Rodrigues. Generalization Error of Invariant Classifiers. In *Artificial Intelligence and Statistics*, pages 1094–1103, 2017a.

Jure Sokolić, Raja Giryes, Guillermo Sapiro, and Miguel Rodrigues. Robust Large Margin Deep Neural Networks. *IEEE Transactions on Signal Processing*, 65(16):4265–4280, August 2017b. ISSN 1053-587X. doi: 10.1109/TSP.2017.2708039.

Nathaniel Thomas, Tess Smidt, Steven M. Kearnes, Lusann Yang, Li Li, Kai Kohlhoff, and Patrick Riley. Tensor field networks: Rotation- and translation-equivariant neural networks for 3d point clouds. *arXiv preprint arXiv:1802.08219*, 2018.

Joel A Tropp. User-friendly tail bounds for sums of random matrices. *Foundations of computational mathematics*, 12(4):389–434, 2012.

Guillermo Valle-Pérez and Ard A. Louis. Generalization bounds for deep learning, December 2020. URL http://arxiv.org/abs/2012.04115. arXiv:2012.04115 [cs, stat].

Gal Vardi, Ohad Shamir, and Nathan Srebro. The Sample Complexity of One-Hidden-Layer Neural Networks. *arXiv:2202.06233 [cs, stat]*, February 2022. URL http://arxiv.org/abs/2202.06233.

Colin Wei and Tengyu Ma. Improved Sample Complexities for Deep Networks and Robust Classification via an All-Layer Margin. *arXiv:1910.04284 [cs, stat]*, October 2019. arXiv: 1910.04284.

Maurice Weiler and Gabriele Cesa. General $E(2)$-Equivariant Steerable CNNs. In *Conference on Neural Information Processing Systems (NeurIPS)*, 2019.

Maurice Weiler, Mario Geiger, Max Welling, Wouter Boomsma, and Taco S. Cohen. 3D steerable CNNs: Learning rotationally equivariant features in volumetric data. In *Conference on Neural Information Processing Systems (NeurIPS)*, 2018a.

Maurice Weiler, Fred A. Hamprecht, and Martin Storath. Learning steerable filters for rotation equivariant CNNs. In *Conference on Computer Vision and Pattern Recognition (CVPR)*, 2018b.

Maurice Weiler, Patrick Forré, Erik Verlinde, and Max Welling. Coordinate Independent Convolutional Networks–Isometry and Gauge Equivariant Convolutions on Riemannian Manifolds. *arXiv preprint arXiv:2106.06020*, 2021.

Jeffrey Wood and John Shawe-Taylor. Representation theory and invariant neural networks. *Discrete Applied Mathematics*, 69(1):33 – 60, 1996. ISSN 0166-218X. doi: https://doi.org/10.1016/0166-218X(95)00075-3.

Daniel E. Worrall, Stephan J. Garbin, Daniyar Turmukhambetov, and Gabriel J. Brostow. Harmonic networks: Deep translation and rotation equivariance. In *Conference on Computer Vision and Pattern Recognition (CVPR)*, 2017.

Chiyuan Zhang, Samy Bengio, Moritz Hardt, Benjamin Recht, and Oriol Vinyals. Understanding deep learning requires rethinking generalization. In *ICLR 2017*, 2017.

Sicheng Zhu, Bang An, and Furong Huang. Understanding the Generalization Benefit of Model Invariance from a Data Perspective. In *Advances in Neural Information Processing Systems*, 2021.

Supplementary Material

7   Elements of Group and Representation Theory

In this section, we provide a brief introduction to the concepts from Group Theory which we need in our derivations.

Definition 7.1 (Group). A *group* is a pair $(G, \cdot)$ containing a set G and a binary operation $\cdot : G \times G \to G, (h, g) \mapsto h \cdot g$ which satisfies the group axioms:

- Associativity: $\forall a, b, c \in G \quad a \cdot (b \cdot c) = (a \cdot b) \cdot c$

- Identity: $\exists e \in G : \forall g \in G \quad g \cdot e = e \cdot g = g$

- Inverse: $\forall g \in G \ \exists g^{-1} \in G : \quad g \cdot g^{-1} = g^{-1} \cdot g = e$

The operation $\cdot$ is the *group law* of G.

The inverse elements g^{-1} of an element g, and the identity element e are *unique*. In addition, if the group law is also *commutative*, the group G is an *abelian group*. To simplify the notation, we commonly write ab instead of $a \cdot b$. It is also common to denote the group $(G, \cdot)$ just with the name of its underlying set G.

The **order** of a group G is the *cardinality* of its set and is indicated by $|G|$. A group G is **finite** when $|G| \in \mathbb{N}$, i.e., when it has a finite number of elements. A **compact** group is a group that is also a compact topological space with continuous group operation.

Definition 7.2 (Group Action). Given a group G, its *action* on a set $\mathcal{X}$ is a map $. : G \times \mathcal{X} \to \mathcal{X}, (g, x) \mapsto g.x$ which satisfies the axioms:

- identity: $\forall x \in \mathcal{X} \quad e.x = x$

- compatibility: $\forall a, b \in G \ \forall x \in \mathcal{X} \quad a.(b.x) = (ab).x$

A simple example of group action is the group law itself $\cdot : G \times G \to G$ which defines an action of G on its own elements ($\mathcal{X} = G$). Another important action is the one defined on signals overs the group G. Given a signal $x : G \to \mathbb{R}$, the action of an element $g \in G$ maps $x \mapsto g.x, [g.x](h) := x(g^{-1}h)$.

The **orbit** of $x \in \mathcal{X}$ through G is the set $G.x := \{g.x | g \in G\}$. The orbits of the elements in $\mathcal{X}$ form a *partition* of $\mathcal{X}$. By considering the equivalence relation $\forall x, y \in \mathcal{X} \quad x \sim_G y \iff x \in G.y$ (or, equivalently, $y \in G.x$), one can define the **quotient space** $\mathcal{X}/G := \{G.x | x \in \mathcal{X}\}$, i.e. the set of all different orbits.

Definition 7.3 (Subgroup). Given a group $(G, \cdot)$, a non-empty subset $H \subseteq G$ is a **subgroup** of G if it forms a group $(H, \cdot)$ under the same group law, restricted to the elements of H. This is usually denoted as $H \leq G$.

H is a subgroup of G if and only if the subset H is *closed* under the group law and the inverse operations, i.e.:

- $\forall a, b \in H \quad a \cdot b \in H$
- $\forall h \in H \quad h^{-1} \in H$

Note that a subgroup H needs to contain the identity element $e \in G$.

Linear representations

Definition 7.4 (Linear representation). Given a group G and a vector space V, a linear representation of G is a homomorphism $\rho : G \to \mathrm{GL}(V)$ which associates to each element of $g \in G$ an element of general linear group $\mathrm{GL}(V)$, i.e. an invertible matrix acting on V, such that the condition below is satisfied:

$$\forall g, h \in G, \quad \rho(gh) = \rho(g)\rho(h).$$

The most simple representation is the trivial representation $\psi : G \to \mathbb{R}, g \mapsto 1$, mapping all element to the multiplicative identity $1 \in \mathbb{R}$. The common 2-dimensional rotation matrices are an example of representation on $\mathbb{R}^2$ of the group $\mathrm{SO}(2)$.

For a finite group G, an important representation is the *regular representation* ρ_{reg}. It acts on the space $V = \mathbb{R}^{|G|}$ of vectors representing signals over the group G. The regular representation $\rho_{\mathrm{reg}}(g)$ of an element $g \in G$ is a $|G| \times |G|$ matrix which permutes the entries of vectors $\boldsymbol{x} \in \mathbb{R}^{|G|}$. Each vector in $\mathbb{R}^{|G|}$ represents a function over the group, $x : G \to \mathbb{R}$, with $x(g_i)$ being i-th entries of $\boldsymbol{x}$. Then, the group action $\rho_{\mathrm{reg}}(g)\boldsymbol{x}$ represents the function $g.x$, i.e. the signal x shifted by g. In other words, $\rho_{\mathrm{reg}}(g)$ is a permutation matrix moving the i-th entry of $\boldsymbol{x}$ to the j-th entry such that $g_j = gg_i$. Regular representations are of high importance, because they describe the features of group convolution networks.

Given two representations $\rho_1 : G \to \mathrm{GL}(\mathbb{R}^{n_1})$ and $\rho_2 : G \to \mathrm{GL}(\mathbb{R}^{n_2})$, their *direct sum* $\rho_1 \oplus \rho_2 : G \to \mathrm{GL}(\mathbb{R}^{n_1+n_2})$ is a representation obtained by stacking the two representations as follow:

$$(\rho_1 \oplus \rho_2)(g) = \begin{bmatrix} \rho_1(g) & 0 \\ 0 & \rho_2(g) \end{bmatrix} .$$

Note that this representation acts on $\mathbb{R}^{n_1+n_2}$ which contains the concatenation of the vectors in $\mathbb{R}^{n_1}$ and $\mathbb{R}^{n_2}$.

Fourier Transform Fourier analysis can be generalized for square integrable complex functions $f : G \to \mathbb{C}$ defined over a compact group G. Fourier components in this case are a particular set of complex representations of G called *irreducible representations* (or *irreps*) of G. Irreps are defined as representation with no $G-$invariant subspaces (see Serre [1977] for rigorous details). The key role of irreps is clear from the following theorem.

Theorem 7.5 (Maschke's Theorem). *Every representation of a finite group G on a nonzero, finite dimensional complex vector space is a direct sum of irreducible representations.*

Maschke's theorem can be generalized to compact groups. Irreps are, therefore, simple blocks for decomposing representations. Irreps are denoted by ψ with dimension $\dim_\psi$, namely, $\psi : G \to \mathrm{GL}(\mathbb{C}^{\dim_\psi})\}_\psi$. The set of irreps is denoted by $\hat{G} = \{\psi : G \to \mathrm{GL}(\mathbb{C}^{\dim_\psi})\}$. The matrix coefficients of the irreps (i.e. each entry $\psi_{ij} : G \to \mathbb{C}$ of each irrep ψ, interpreted as a function over G) form an orthogonal basis for square integrable functions over G. Therefore, we can project a function $f : G \to \mathbb{C}$ on this basis and get the Fourier components as:

$$\hat{f}(\psi) := \int_{g \in G} f(g)\psi(g)\, d\mu(g),$$

where μ is the Haar measure over G. Note that $\hat{f}(\psi)$ is a $\dim_\psi \times \dim_\psi$ matrix containing a coefficient for each entry of ψ. The inverse Fourier transform is defined as:

$$f(g) = \sum_{\psi \in \hat{G}} \dim_\psi \mathrm{Tr}(\hat{f}(\psi)\psi(g^{-1}))$$

Note that the trace is equivalent to computing the elementwise product between the matrices $\hat{f}(\psi)$ and $\psi(g^{-1})^T = \overline{\psi(g)}$ and then summing over all entries.

If G is the cyclic group, $G = \mathrm{C}_N$, one recovers the usual *discrete Fourier transform*, because the complex irrep of C_N are 1-dimensional and correspond to the the complex exponentials up to order $N - 1$. When G is finite, the number of Fourier coefficients $\sum_{\psi \in \hat{G}} \dim_\psi^2$ is equal to $|G|$. Indeed, the Fourier transform of a signal $x : G \to \mathbb{C}$ can be interpreted as a $|G| \times |G|$ change of basis which maps the vector $\boldsymbol{x} \in \mathbb{C}^{|G|}$ representing x in the group domain to $\hat{\boldsymbol{x}} \in \mathbb{C}^{|G|}$ containing all the Fourier coefficients.

Given two signals $w, x : G \to \mathbb{C}$, the following property (similar to the common Fourier transform) holds:

$$\widehat{g.x}(\psi) = \psi(g)\hat{x}(\psi) \tag{12}$$

Eq. 12 guarantees that a transformation by g of x does not mix the coefficients associated to different irreps. Moreover, the coefficients associated with the irrep ψ are mixed precisely by the matrix $\psi(g)$, i.e. they transform according to ψ. Indeed, if all the coefficients associated to the same irrep are grouped together, one can decompose the regular representation ρ_{reg} of G in a *direct sum* of the irreps of G, up to the change of basis B as above. More precisely, the irreps decomposition of ρ_{reg} contains $\dim_\psi$ copies of ψ, one for each column[5] of $\hat{f}(\psi)$, i.e.:

$$\rho_{\text{reg}}(g) = B^{-1} \bigoplus_{\psi \in \hat{G}} \left(\underbrace{\psi(g) \oplus \psi(g) \oplus \dots}_{\dim_\psi \text{ times}} \right) B \; .$$

This is the key insight used in the analysis of group convolution in this work.

Group Convolution Given a space $\mathcal{X}$ associated with the action of a group G and an invariant inner product $\langle \cdot, \cdot \rangle : \mathcal{X} \times \mathcal{X} \to \mathbb{R}$, we define the *group convolution* $w \circledast_G x : G \to \mathbb{R}$ of two elements $w, x \in \mathcal{X}$ as

$$\forall g \in G \quad (w \circledast_G x)(g) := \langle g.w, x \rangle \; . \tag{13}$$

Group convolution satisfies the following equality:

$$\widehat{w \circledast_G x}(\psi) = \hat{x}(\psi)\hat{w}(\psi)^T \tag{14}$$

The classical definition of *group convolution* between two signals $w, x : G \to \mathbb{R}$ is a special case of the one above in the case $\mathcal{X}$ is the space of square integrable functions over G and the invariant inner product is defined as:

$$\langle w, x \rangle := \int_{g \in G} w(g)x(g) \, d\mu(g)$$

where $\mu : G \to \mathbb{R}$ is a Haar measure over G. The group convolution, then, becomes

$$(w \circledast_G x)(g) := \int_{h \in G} w(g^{-1}h)x(h) \, d\mu(h) = \int_{h \in G} w(h)x(g.h) \, d\mu(h) \; .$$

Note that what we defined is technically a **group cross-correlation**, and so it differs from the usual definitions of convolution over groups. We still refer to is as group convolution to follow the common terminology in the deep learning literature.

When G is finite, the Haar measure is the counting measure and the integral becomes a sum. Fix an ordering of group elements as $(g_0, g_1, \dots, g_i, \dots, g_{|G|-1})$ with $g_0 = e$ the identity element. The signals $x, w : G \to \mathbb{R}$ can be stored as $|G|$ dimensional vectors $\boldsymbol{x}, \boldsymbol{w} \in \mathbb{R}^{|G|}$, where the i-th entry contains the value of the function evaluated on g_i:

$$\boldsymbol{x} = \begin{pmatrix} x(g_0) \\ \vdots \\ x(g_{|G|-1}) \end{pmatrix}, \quad \boldsymbol{w} = \begin{pmatrix} w(g_0) \\ \vdots \\ w(g_{|G|-1}) \end{pmatrix} . \tag{15}$$

The same holds for the output signal $w \circledast_G x$. Define matrix $\boldsymbol{W}$ as follows:

$$\boldsymbol{W} = \begin{pmatrix} w(g_0) & \cdots & w(g_{|G|-1}) \\ w(g_1^{-1}g_0) & \cdots & w(g_1^{-1}g_{|G|-1}) \\ \vdots & \ddots & \vdots \\ w(g_{|G|-1}^{-1}g_0) & \cdots & w(g_{|G|-1}^{-1}g_{|G|-1}) \end{pmatrix} \tag{16}$$

Then, the group convolution can then be expressed as a matrix multiplication between the vector $\boldsymbol{x}$ and a matrix $\boldsymbol{W}$ containing at position (i, j) the entry of $w_{g_i^{-1}g_j}$:

$$(w \circledast_G h.x)(g_i) = (\boldsymbol{W}\boldsymbol{x})[i] \tag{17}$$

The matrix $\boldsymbol{W}$ contains permuted copies of its first row, where the respective permutation matrix is determined by group actions. Assuming $g_0 = e$ is the identity, the first row of $\boldsymbol{W}$ contains $\boldsymbol{w}^T$ while the i-th row contains $(g_i.\boldsymbol{w})^T$, i.e. the vector representing the signal $g_i.w$. The matrix $\boldsymbol{W}$ is a G-**circulant matrix**.

---

[5] In Eq. 12, $\psi(g)$ acts on each column of the matrix $\hat{f}(\psi)$ independently.

Equivariance. For two vector spaces V and V' with group representations respectively ρ and ρ', the linear transformation $T : V' \to V$ is equivariant if

$$\forall g \in G : T \circ \rho'(g) = \rho(g) \circ T.$$

Now, since an invariant inner product satisfies $\langle g.w, g.x \rangle = \langle w, x \rangle$ for any element $g \in G$, one can show that the group convolution above is **equivariant**:

$$
\begin{aligned}
(w \circledast_G h.x)(g) &= \langle g.w, h.x \rangle \\
&= \langle (h^{-1}g).w, (h^{-1}h).x \rangle \\
&= \langle (h^{-1}g).w, x \rangle \\
&= (w \circledast_G x)(h^{-1}g) \ .
\end{aligned}
$$

By definition of action of G on signals over G, it follows that $(w \circledast_G h.x) = h.(w \circledast_G x)$.

We finish the part by presenting Schur's lemma, [Serre, 1977, Section 2.2] which helps characterizing equivariant maps.

Theorem 7.6 (Schur's lemma). *Let ρ^1 and ρ^2 be irreducible representations of G respectively on vector spaces V_1 and V_2. Suppose that the linear transformation $T : V_1 \to V_2$ is equivariant. Then:*

1. *If ρ^1 and ρ^2 are non-isomorphic, then $T = 0$.*

2. *If $V_1 = V_2$ and $\rho^1 = \rho^2$, then T is a homothety, i.e, a scalar multiple of identity.*

Equivariant neural networks. To build equivariant networks, without loss of generality, we assume the feature $x \in \mathbb{R}^{c_l}$ of the neural network at layer l transforms according to a generic representation c_l-dimensional representation ρ_l and that it is decomposed[6] in a number of subvectors $\hat{x}_i \in \mathbb{R}^{c_{l,i}}$, with $\sum_i c_{l,i} = c_l$, each transforming according to an irrep $\psi_i : G \to \mathrm{GL}(\mathbb{C}^{c_{l,i}})$. To see this more precisely, consider the layer l given by the matrix $W_l \in \mathbb{R}^{c_l \times c_{l-1}}$. The group representation is ρ_l acting on $\mathbb{R}^{c_l}$. The equivariance relation is given by:

$$W_l \rho_{l-1}(h) = \rho_l(h) W_l \quad .$$

We first decompose representations using Maschke's theorem. The representation ρ_l can be written direct sum of irreps as

$$\rho_l = Q_l \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right) Q_l^{-1}$$

$$= Q_l \times \text{block-diagonal} \left(\psi \in \hat{G} : \underbrace{\begin{pmatrix} \psi & \dots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \dots & \psi \end{pmatrix}}_{m_{l,\psi} \text{ blocks}} \right) \times Q_l^{-1}$$

The multiplicity of the irrep ψ in ρ_l is $m_{l,\psi}$. Using $\widehat{W}_l = Q_l^{-1} W_l Q_{l-1}$, we have

$$\widehat{W}_l \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l-1,\psi}} \psi \right) = \left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right) \widehat{W}_l.$$

Since $\left(\bigoplus_\psi \bigoplus_{i=1}^{m_{l,\psi}} \psi \right)$ is block diagonal, we can partition $\widehat{W}_l$ similarly with the sub-block (j, i) of blocks relating ψ_1 to ψ_2, denoted by $\widehat{W}_l(\psi_2, j, \psi_1, i)$. The block relates ψ_1 to ψ_2 as:

$$\widehat{W}_l(\psi_2, j, \psi_1, i)\psi_1 = \psi_2 \widehat{W}_l(\psi_2, j, \psi_1, i).$$

---

[6]The decomposition can include an orthogonal change of basis Q.

Figure 6: The block diagonal structure in $\boldsymbol{W}_l \rho_{l-1}(h) = \rho_l(h)\boldsymbol{W}_l$

This setup follows the premise of Schur's lemma, exept that in practice, a neural network uses real valued features and weights. Hence, we need to consider *real* irreps, which we denote with $\{\psi_i\}$. This requires some adaptation to the Fourier analysis and Schur's lemma. This is explained more precisely in Supplementary 8. If $\psi_1 \neq \psi_2$, the block needs to be zero. Otherwise, each block $\boldsymbol{W}$ equivariant to irrep ψ can be expressed as

$$\boldsymbol{W} = \sum_{k=1}^{c_\psi} w_k \boldsymbol{B}_{\psi,k}$$

where matrices $\boldsymbol{B}_{\psi,k}$ are given in Supplementary 8, c_ψ is either 1,2 or 4, and $\{w_k\}_{k=1}^{c_\psi}$ are the c_ψ free parameters. Therefore, each block $\widehat{\boldsymbol{W}}_l(\psi, j, \psi, i)$ is characterized by c_ψ free parameters.

Note that, the main difference of real irreps is that only the matrix coefficients in a fraction $\frac{1}{c_\psi}$ of the columns of ψ are part of the orthogonal basis for real functions on G, while the other coefficients are redundant. It follows that, when computing the Fourier transform of a function $x : G \to \mathbb{R}$, only the first $\frac{\dim_\psi}{c_\psi}$ columns of $\hat{x}(\psi)$ needs to be considered and, therefore, the regular representation of H only contains $\frac{\dim_\psi}{c_\psi}$ copies of ψ.

Group convolutional networks. Group convolutional networks is an important special case of the setup presented above. Group convolution based architectures of Cohen and Welling [2016a] can be obtained by choosing multiple copies of regular representations at each layer.

Consider an input vector space $\mathcal{X}$ associated with an action $H \times \mathcal{X} \to \mathcal{X}$ of H. We use group convolution as defined in eq. 13. Each convolutional kernel can be parametrized by a kernel $\boldsymbol{w}$, and its action is represented by a group circulant matrix $\boldsymbol{W}$ as in eq. 16.

This is the building blocks of the linear layers of group equivariant networks. Each intermediate convolutional layer maps a feature space with c_{l-1} channels of dimension equal to group size $|H|$ to a new feature space with c_l channels of dimension $|H|$. Any such layer can be visualized as follows:

$$\begin{pmatrix} \mathbf{y}_1 \in \mathbb{R}^{|H|} \\ \mathbf{y}_2 \in \mathbb{R}^{|H|} \\ \vdots \\ \mathbf{y}_{c_l} \in \mathbb{R}^{|H|} \end{pmatrix} = \underbrace{\begin{pmatrix} \boldsymbol{W}_{l,(1,1)} & \boldsymbol{W}_{l,(2,1)} & \cdots & \boldsymbol{W}_{l,(c_{l-1},1)} \\ \boldsymbol{W}_{l,(1,2)} & \boldsymbol{W}_{l,(2,2)} & \cdots & \boldsymbol{W}_{l,(c_{l-1},2)} \\ \vdots & \ddots & & \vdots \\ \boldsymbol{W}_{l,(1,c_l)} & \boldsymbol{W}_{l,(2,c_l)} & \cdots & \boldsymbol{W}_{l,(c_{l-1},c_l)} \end{pmatrix}}_{\boldsymbol{W}_l} \cdot \underbrace{\begin{pmatrix} \mathbf{x}_1 \in \mathbb{R}^{|H|} \\ \mathbf{x}_2 \in \mathbb{R}^{|H|} \\ \vdots \\ \mathbf{x}_{c_{l-1}} \in \mathbb{R}^{|H|} \end{pmatrix}}_{\boldsymbol{x}} \tag{18}$$

Here, any $\boldsymbol{W}_{l,(i,j)} \in \mathbb{R}^{|H| \times |H|}$ block an group circulant matrix corresponding to H, which encodes a H-convolution parametrised by a filter $\boldsymbol{w}_{l,i,j} \in \mathbb{R}^{|H|}$. Note that the output of the l-th layer is seen as a $|H| \times c_l$ signal, i.e. it stores a different value $(\boldsymbol{w}_1 \circledast_H \boldsymbol{x})(h, i)$ for each group element h and channel i.

Usually, the group action on the input space $\mathcal{X}$ is given by the task. The first linear layer transforms the input space to multiple signals (channels) over H. From that moment, we can use the convolution in Eq. 13 in the following layers.

Finally, the last layer produces an invariant output for each class. This can be interpreted as a group convolution with constant filters or, equivalently, as an average pooling within the $|H|$ channels in each block to produce a c_{L-1} dimensional invariant vector followed by a linear classifier.

The resulting architecture can be written as a MLP with structured weights similar to Wood and Shawe-Taylor [1996], Ravanbakhsh [2020]. Such network, denoted by $f(\boldsymbol{x}; \mathbb{W})$, is an MLP with L layers, input $\boldsymbol{x}$ and the set of effective parameters $\mathbb{W}$. The l-th layer consists of a linear map $\boldsymbol{W}_l$ from a vector space $\mathcal{V}_{l-1}$ to another vector space $\mathcal{V}_l$ followed by an activation function $\sigma_l(\cdot)$. The network output at the layer l, denoted by $f_l(\boldsymbol{x}; \mathbb{W})$, is given by $f_l(\boldsymbol{x}; \mathbb{W}) = \sigma_l(\boldsymbol{W}_l f_{l-1}(\boldsymbol{x}; \mathbb{W}))$.

This design guarantees that each linear layer commutes with the group action. If also all non-linearities commute with it, i.e.
$$\sigma_l(\boldsymbol{W}_l h.\boldsymbol{x}) = \sigma_l(h.(\boldsymbol{W}_l \boldsymbol{x})) = h.\sigma_l(\boldsymbol{W}_l \boldsymbol{x}) \;,$$
it follows that the output of the model is invariant to H.

Note that the number of *effective channels* after the l-th hidden layer is $|H| c_l$ but its number of trainable parameters is $|H| c_l c_{l-1}$. In total, if number of channels across layers are all equal to c, then the number trainable parameters is equal to $\mathcal{O}\left(L|H|c^2\right)$. In our experiments, we always consider fixed architectures where the number of *effective channels* is kept constant when changing the group, i.e. different group choices only affect the number of learnable parameters and the structure of the weight matrices but not their size and, therefore, not the computational cost of the model.

Groups considered in this work. In the experiments, we consider 4 different families of compact groups. The special orthogonal group $\mathrm{SO}(2)$ is the group of all planar rotations. It is an abelian (commutative) group and contains any rotation by an angle $\theta \in [0, 2\pi)$. $\mathrm{SO}(2)$ contains infinite elements. The orthogonal group $\mathrm{O}(2)$ is the group of all planar rotations and reflections. Indeed, $\mathrm{SO}(2)$ is a subgroup of $\mathrm{O}(2)$. The elements of $\mathrm{O}(2)$ are planar rotations, or planar rotations combined with a reflection along a fixed axis. It is not commutative and contains infinite elements. The cyclic group C_N of order N is a finite group containing the N rotations by angles multiple of $\frac{2\pi}{N}$. It is a subgroup of both $\mathrm{SO}(2)$ and $\mathrm{O}(2)$ and it is abelian. Finally, the dihedral group D_N is a finite group containing $2N$ elements. It contains the N rotations in C_N and but also their composition with a reflection along a fixed axis. It is a subgroup of $\mathrm{O}(2)$ and it is not commutative. If $G = \mathrm{C}_N$, the matrix $\boldsymbol{W}$ is a circulant matrix in the classical sense, where each row is a cyclic shift by one step of the previous one.

8 Schur's Lemma for intertwiners between real representations

Given two *complex* irreps $\sigma_i : G \to \mathrm{GL}(\mathbb{C}^{\dim \psi_i})$ and $\sigma_j : G \to \mathrm{GL}(\mathbb{C}^{\dim \sigma_j})$, Schur's Lemma guarantees that the set of equivariant matrices (**intertwiners**)
$$\mathrm{Hom}_{G,\mathbb{C}}(\psi_i, \psi_j) := \{M \mid M\psi_i(g) = \psi_j(g)M \quad \forall g \in G\}$$
is either zero dimensional containing only zero function (if $\psi_i \not\cong \psi_j$), or 1-dimensional and contains only scalar multiples of the identity (if $\psi_i = \psi_j$), i.e.
$$\forall g \in G \quad M\psi(g) = \psi(g)M \iff \exists \lambda \in \mathbb{C} \text{ s.t. } M = \lambda I,$$
where $\cong$ denotes isomorphism. In other words, there is a change of basis matrix such that the two representations are equal. The space $\mathrm{Hom}_{G,\mathbb{C}}(\psi_i, \psi_j)$ is called the space of homomorphisms. The subscript $\mathbb{C}$ indicates the complex-valued intertwiners are intended. This lemma is the core of equivariant (complex valued) linear layers and provides a way to parametrize the space of all such operations.

In this section, we discuss Schur's lemma for real irreps and characterize the space of homomorphisms $\mathrm{Hom}_{G,\mathbb{R}}(\psi_i, \psi_j)$ for real valued representations. We still use σ to denote *complex irreps* and will refer to *real irreps* with ψ. First we note that if $\psi_i \not\cong \psi_j$ are different *real* irreps, by using the complex version of Schur's Lemma, one can show that $\mathrm{Hom}_{G,\mathbb{R}}(\psi_i, \psi_j)$ still contains only the zero matrix.

When we have $\psi_i \cong \psi_j$, the derivations are more tricky. For this cases, $\psi_i = \psi_j = \psi$, we have to consider $\mathrm{Hom}_{G,\mathbb{R}}(\psi, \psi)$. For complex-valued representations, as seen above from Schur's lemma, this space is one dimensional spanned by the identity matrix $\boldsymbol{I}$. However, for real representations, this space can be either 1, 2 or 4 dimensional, corresponding to three types: real $\mathbb{R}$, complex $\mathbb{C}$ and quanternion $\mathbb{H}$ (see [Cesa et al., 2022, Section C] for accessible explanations). Based on this, any real irrep ψ can be classified in three categories:

- **real-type**: $\exists \psi$ real $: \psi \cong \sigma \cong \overline{\sigma}$

- **complex-type**: $\exists \psi$ real $: \psi = \sigma_{\mathbb{R}} \cong \sigma \oplus \overline{\sigma}$, with $\sigma \not\cong \overline{\sigma}$

- **quaternionic-type**: $\exists \psi$ real $: \psi = \sigma_{\mathbb{R}} \cong \sigma \oplus \overline{\sigma} \cong \sigma \oplus \sigma$, as $\sigma \cong \overline{\sigma}$

where $\overline{\sigma}$ is a representation such that $\overline{\sigma} : g \mapsto \overline{\sigma(g)}$ and $\overline{\sigma(g)}$ is the complex-conjugation of the entries of the matrix $\sigma(g)$. We also have defined

$$\sigma_{\mathbb{R}}(g) := \begin{bmatrix} Re\left(\sigma(g)\right) & -Im\left(\sigma(g)\right) \\ Im\left(\sigma(g)\right) & Re\left(\sigma(g)\right) \end{bmatrix} \in \mathbb{R}^{2d \times 2d}$$

where $Re\left(X\right)$ (respectively, $Im\left(X\right)$) is a matrix containing the *real* (respectively, *imaginary*) part of the complex entries of the matrix X, i.e.:

$$Re\left(X\right) = \frac{1}{2}\left(X + lineX\right)$$

$$Im\left(X\right) = -i\frac{1}{2}\left(X - \overline{X}\right)$$

$$X = Re\left(X\right) + iIm\left(X\right)$$

Using these definitions, one can also verify that for any d-dimensional complex irrep σ:

$$\sigma_{\mathbb{R}}(g) = D_d \begin{bmatrix} \sigma(g) & 0 \\ 0 & \overline{\sigma}(g) \end{bmatrix} D_d^* = D_d(\sigma(g) \oplus \overline{\sigma}(g))D_d^*$$

where

$$D_d = \frac{1}{\sqrt{2}} \begin{bmatrix} iI_d & -iI_d \\ I_d & I_d \end{bmatrix}$$

where I_d is the $d \times d$ identity matrix and $*$ indicates the conjugate transpose. Note that D_d is a unitary matrix.

We can then distinguish three cases.

Real type   If $\psi \cong \sigma \cong \overline{\sigma}$, we have $\psi = A\sigma A^*$, with A unitary matrix ($A^{-1} = A^*$). An intertwiner matrix M satisfies $M\psi(g) = \psi(g)M$. It can be seen that the matrix $\tilde{M} := A^*MA$, $\tilde{M}$ is an intertwiner of σ:

$$\tilde{M}\sigma(g) = A^*MA\sigma(g) = A^*M\psi(g)A = A^*\psi(g)MA = \sigma(g)A^*MA = \sigma(g)\tilde{M},$$

and, therefore, has form λI for $\lambda \in \mathbb{C}$ from complex-valued Schur's lemma. It follows that $M = \lambda I$, allowing only real intertwiners,

$$M_{\mathbb{R}} = \lambda I$$

with $\lambda \in \mathbb{R}$.

Complex type   If $\psi \cong \sigma \oplus \overline{\sigma}$, we have $\psi = D_d(\sigma \oplus \overline{\sigma})D_d^*$ with d being the dimension of irrep σ. If M satisfies $M\psi(g) = \psi(g)M$, then the matrix $\tilde{M} = D_d^*MD_d$ is an intertwiner of $\sigma \oplus \overline{\sigma}$:

$$\begin{aligned} \tilde{M}(\sigma(g) \oplus \overline{\sigma}(g)) &= D_d^*MD_d(\sigma(g) \oplus \overline{\sigma}(g)) \\ &= D_d^*M\psi D_d \\ &= D_d^*MD_d(\sigma(g) \oplus \overline{\sigma}(g)) = (\sigma(g) \oplus \overline{\sigma}(g))\tilde{M}. \end{aligned}$$

We can apply complex-valued Schur's lemma again. The matrix $\tilde{M}$ needs to have form $\alpha I \oplus \beta I$ for $\alpha, \beta \in \mathbb{C}$. Re-applying the change of basis D_d and requiring only real entries, one finds that M needs to have the following form:

$$M_{\mathbb{C}} = \begin{bmatrix} aI_d & -bI_d \\ bI_d & aI_d \end{bmatrix}$$

with $a, b \in \mathbb{R}$.

Quaternionic type If $\psi \cong \sigma \oplus \sigma$, there is a basis where $\psi = D_d(\sigma \oplus \overline{\sigma})D_d^*$ and $\overline{\sigma} = T_d^T \sigma T_d$ with $T_d = \begin{bmatrix} 0 & -I_{d/2} \\ I_{d/2} & 0 \end{bmatrix}$, that is $\psi = D_d(I_d \oplus T_d^T)(\sigma \oplus \sigma)(I_d \oplus T_d)D_d^*$. If M satisfies $M\psi(g) = \psi(g)M$, we define $\tilde{M} = D_d^*(I_d \oplus T_d)M(I_d \oplus T_d^T)D_d$, intertwiner of $\sigma \oplus \sigma$. $\tilde{M}$ needs to have form $\tilde{M} = \begin{bmatrix} \alpha I_d & \gamma I_d \\ \beta I_d & \delta I_d \end{bmatrix}$ for $\alpha, \beta, \gamma, \delta \in \mathbb{C}$. Re-applying the change of basis $D_d(I_d \oplus T_d^T)$ and requiring only real entries, one can show that M needs to have the following form:

$$M_{\mathbb{H}} = \begin{bmatrix} aI_d + cT_d & -bI_d + dT_d \\ bI_d + dT_d & aI_d - cT_d \end{bmatrix} = \begin{bmatrix} aI_d & -cI_d & -bI_d & -dI_d \\ cI_d & aI_d & dI_d & -bI_d \\ bI_d & -dI_d & aI_d & cI_d \\ dI_d & bI_d & -cI_d & aI_d \end{bmatrix} = \begin{bmatrix} a & -c & -b & -d \\ c & a & d & -b \\ b & -d & a & c \\ d & b & -c & a \end{bmatrix} \otimes I_d$$

with $a, b, c, d \in \mathbb{R}$.

Application to equivariant networks. Real-valued version of Schur's lemma characterizes equivariant kernel $\boldsymbol{W}$ parametrized by $M_{\mathbb{R}}, M_{\mathbb{C}}, M_{\mathbb{H}}$. The columns of $\boldsymbol{W}$ are all orthogonal to each other and each contains only one copy of each of its c_ψ free parameters, where

$$c_\psi = \begin{cases} 1 & \text{real type} \\ 2 & \text{complex type} \\ 4 & \text{quaternionic type} \end{cases}$$

It follows that all columns have the same norm, equal to the sum of the c_ψ free parameters squared, and therefore that $\boldsymbol{W}$ is a scalar multiple of an orthonormal matrix (with scale equal to the norm of any of its columns). The matrix $\boldsymbol{W}$ satisfies the following property:

$$\|\boldsymbol{W}\boldsymbol{x}\|_2^2 = \frac{1}{\dim_\psi} \|\boldsymbol{W}\|_F^2 \|\boldsymbol{x}\|_2^2 .$$

From the above identity, it follows that the spectral norm of $\boldsymbol{W}$ is

$$\|\boldsymbol{W}\|_2 = \frac{1}{\sqrt{\dim_\psi}} \|\boldsymbol{W}\|_F \tag{19}$$

For any irrep ψ, we can express any matrix $\boldsymbol{W}$ equivariant to ψ as

$$\boldsymbol{W} = \sum_{k=1}^{c_\psi} w_k \boldsymbol{B}_{\psi,k}$$

where $\{w_k\}_{k=1}^{c_\psi}$ are the c_ψ free parameters. When ψ has real type, $c_\psi = 1$ and $\boldsymbol{B}_{\psi,1} = \boldsymbol{I}$ is the identity matrix. When ψ has complex type, $c_\psi = 2$ and $\boldsymbol{B}_{\psi,1} = \boldsymbol{I}_{2d}$ is the identity matrix while $\boldsymbol{B}_{\psi,2} = \begin{bmatrix} 0 & -\boldsymbol{I}_d \\ \boldsymbol{I}_d & 0 \end{bmatrix}$, where $d = \dim_\psi / 2$. Finally, if ψ has quaternionic type, $c_\psi = 4$

$$\boldsymbol{B}_{\psi,1} = \boldsymbol{I}_{4d}, \boldsymbol{B}_{\psi,2} = \begin{bmatrix} 0 & 0 & -\boldsymbol{I}_d & 0 \\ 0 & 0 & 0 & -\boldsymbol{I}_d \\ \boldsymbol{I}_d & 0 & 0 & 0 \\ 0 & \boldsymbol{I}_d & 0 & 0 \end{bmatrix}, \boldsymbol{B}_{\psi,3} = \begin{bmatrix} 0 & -\boldsymbol{I}_d & 0 & 0 \\ \boldsymbol{I}_d & 0 & 0 & 0 \\ 0 & 0 & 0 & \boldsymbol{I}_d \\ 0 & 0 & -\boldsymbol{I}_d & 0 \end{bmatrix}, \boldsymbol{B}_{\psi,4} = \begin{bmatrix} 0 & 0 & 0 & -\boldsymbol{I}_d \\ 0 & 0 & \boldsymbol{I}_d & 0 \\ 0 & -\boldsymbol{I}_d & 0 & 0 \\ \boldsymbol{I}_d & 0 & 0 & 0 \end{bmatrix}$$

where $d = \dim_\psi / 4$. Note that, for any ψ and $1 \leq k \leq c_\psi$, $\boldsymbol{B}_{\psi,k}$ is an orthonormal matrix.

9 Proof of Theorem 4.1

In this section, we provide the detailed proof of Theorem 4.1. The complete version of the theorem, including all consants and dependences, is given below.

Theorem 9.1 (Homogeneous Bounds for Equivariant Networks). *For any equivariant network, with probability $1 - \delta$, we have:*

$$\mathcal{L}_{\mathcal{D}}(f_{\mathbb{W}}) \leq \hat{\mathcal{L}}_{\gamma}(f_{\mathbb{W}})+ \tag{20}$$

$$\sqrt{\frac{32e^4 B^2 \beta^2}{\gamma^2 m \eta} \left(\sum_{l=1}^{L} \sqrt{M(l,\eta)} \right)^2 \left(\sum_{l} \frac{\sum_{\psi,i,j} \left\| \widehat{\boldsymbol{W}}_l(\psi,i,j) \right\|_F^2 / \dim_\psi}{\|\boldsymbol{W}_l\|_2^2} \right) + \frac{\log(\xi(m)\frac{Lm^{1+1/2L}}{\delta})}{2\gamma^2 m}} \tag{21}$$

where $\beta = \left(\prod_l \|\boldsymbol{W}_l\|_2 \right)$, and

$$M(l,\eta) := \log \left(\frac{\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}}{1 - \eta} \right) \max_{\psi} \left(5 m_{l-1,\psi} m_{l,\psi} c_\psi \right). \tag{22}$$

9.1   Proof of Lemma 4.3: Perturbation Analysis

The proof of perturbation bound uses standard inequalities and given in Neyshabur et al. [2018]. We derive a bound on the spectral norm of an equivariant matrix $\boldsymbol{W}_l$.

$$
\begin{aligned}
\|\boldsymbol{W}_l \boldsymbol{x}\|_2^2 = \left\| \widehat{\boldsymbol{W}}_l \widehat{\boldsymbol{x}} \right\|_2^2 &= \sum_{\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{W}_l \boldsymbol{x}}(\psi,j) \right\|_2^2 \\
&= \sum_{\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \sum_{i=1}^{m_{l-1,\psi}} \widehat{\boldsymbol{W}}_l(\psi,j,i) \widehat{\boldsymbol{x}}(\psi,i) \right\|_2^2 \\
&\overset{(a)}{\leq} \sum_{\psi} \sum_{j=1}^{m_{l,\psi}} m_{l-1,\psi} \sum_{i=1}^{m_{l-1,\psi}} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \widehat{\boldsymbol{x}}(\psi,i) \right\|_2^2 \\
&\leq \sum_{\psi} \sum_{j=1}^{m_{l,\psi}} m_{l-1,\psi} \sum_{i=1}^{m_{l-1,\psi}} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \right\|_2^2 \|\widehat{\boldsymbol{x}}(\psi,i)\|_2^2 \\
&\overset{(b)}{=} \sum_{\psi} \sum_{j=1}^{m_{l,\psi}} m_{l-1,\psi} \sum_{i=1}^{m_{l-1,\psi}} \frac{1}{\dim_\psi} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \right\|_F^2 \|\widehat{\boldsymbol{x}}(\psi,i)\|_2^2 \\
&= \sum_{\psi} \sum_{i=1}^{m_{l-1,\psi}} \left(m_{l-1,\psi} \frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \right\|_F^2 \right) \|\widehat{\boldsymbol{x}}(\psi,i)\|_2^2 \\
&\leq \left(\max_{\psi} \max_{1 \leq i \leq m_{l-1,\psi}} m_{l-1,\psi} \frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \right\|_F^2 \right) \|\boldsymbol{x}\|_2^2
\end{aligned}
$$

where (a) follows from Cauchy-Schwarz inequality, and (b) from eq. 19, namely that for equivariant matrices, we have:

$$\|\boldsymbol{W}\|_2 = \frac{1}{\sqrt{\dim_\psi}} \|\boldsymbol{W}\|_F \tag{23}$$

Therefore, we get:

$$\|\boldsymbol{W}_l\|_2 \leq \sqrt{\max_{\psi} \max_{1 \leq i \leq m_{l-1,\psi}} m_{l-1,\psi} \frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{W}}_l(\psi,j,i) \right\|_F^2} \tag{24}$$

9.2 Proofs of Lemma 4.4 and Tail Bound for Spectral Norms of Equivariant Kernels

Proof. Recall that the matrix $\widehat{\boldsymbol{U}}_l(\psi, j, i)$ has only c_ψ learnable entries and that $\frac{1}{\dim_\psi} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2$ is equal to the sum of the squares (see Supplementary 8):

$$\frac{1}{\dim_\psi} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2 = \sum_{k=1}^{c_\psi} \left| \widehat{u_{l,i,j}}(\psi)_k \right|^2 .$$

To find tail bounds on the spectral norm, note that $\sum_{j=1}^{m_{l,\psi}} \sum_k \left| \widehat{u_{l,i,j}}(\psi)_k \right|^2$ is a χ^2-random variable with $m_{l,\psi} \times c_\psi$ degrees of freedom. We will use the following inequality for bounding the tail.

Lemma 9.2 ([Laurent and Massart, 2000, Lemma 1.]). *Let $X_1, \ldots, X_n$ be i.i.d. Gaussian random variable with zero mean and variance 1. For any vector $\boldsymbol{a} \in \mathbb{R}^n$, we have*

$$\mathbb{P}(\sum_{i=1}^n a_i (X_i^2 - 1) \geq 2 \|\boldsymbol{a}\|_2 \sqrt{x} + 2 \|\boldsymbol{a}\|_\infty x) \leq \exp(-x). \tag{25}$$

Then, using this theorem, we have:

$$\mathbb{P}\left(\frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2 \geq m_{l,\psi} c_\psi \sigma^2 + 2 m_{l,\psi} c_\psi \sigma^2 \sqrt{t} + 2\sigma^2 t \right) \leq e^{-t}$$

Then:

$$\mathbb{P}\left(m_{l-1,\psi} \frac{1}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2 \geq m_{l-1,\psi} \left(m_{l,\psi} c_\psi \sigma^2 + 2 m_{l,\psi} c_\psi \sigma^2 \sqrt{t} + 2\sigma^2 t \right) \right) \leq e^{-t}$$

Using the union bound:

$$\left(\sum_\psi m_{l,\psi} \right) e^{-t} \geq \mathbb{P}\left(\bigvee_\psi^{m_{l,\psi}} \bigvee_i \left[\frac{m_{l-1,\psi}}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2 \geq m_{l-1,\psi} \left(m_{l,\psi} c_\psi \sigma^2 + 2 m_{l,\psi} c_\psi \sigma^2 \sqrt{t} + 2\sigma^2 t \right) \right] \right)$$

$$\geq \mathbb{P}\left(\max_\psi \max_i^{m_{l,\psi}} \frac{m_{l-1,\psi}}{\dim_\psi} \sum_{j=1}^{m_{l,\psi}} \left\| \widehat{\boldsymbol{U}}_l(\psi, j, i) \right\|_F^2 \geq \max_\psi m_{l-1,\psi} \left(m_{l,\psi} c_\psi \sigma^2 + 2 m_{l,\psi} c_\psi \sigma^2 \sqrt{t} + 2\sigma^2 t \right) \right)$$

We can combine this bound with the bound on the spectral norm in Supplementary 9.1 to obtain a bound similar to the one in Lemma 4.4:

$$\mathbb{P}\left(\|\boldsymbol{U}_l\|_2 \geq \sigma \sqrt{\max_\psi m_{l-1,\psi} \left(m_{l,\psi} c_\psi + 2 m_{l,\psi} c_\psi \sqrt{t} + 2t \right)} \right) \leq \left(\sum_\psi m_{l,\psi} \right) e^{-t} \tag{26}$$

Using $5 m_{l-1,\psi} m_{l,\psi} c_\psi t \geq m_{l-1,\psi} (m_{l,\psi} c_\psi + 2 m_{l,\psi} c_\psi \sqrt{t} + 2t)$:

$$\mathbb{P}\left(\|\boldsymbol{U}_l\|_2 \geq \sigma \sqrt{\max_\psi \left(5 m_{l-1,\psi} m_{l,\psi} c_\psi t \right)} \right) \leq \left(\sum_\psi m_{l,\psi} \right) e^{-t} \tag{27}$$

Using an union bound over all layers, one finds that by choosing $t = \log\left(2 \sum_l \sum_\psi m_{l,\psi} \right)$, with probability at least $\frac{1}{2}$ the following inequality holds for every layer $l \in [L]$:

$$\|\boldsymbol{U}_l\|_2 < \sigma \sqrt{\max_\psi 5 m_{l-1,\psi} m_{l,\psi} c_\psi} \sqrt{\log\left(2 \sum_l \sum_\psi m_{l,\psi} \right)} \tag{28}$$

$\square$

9.3 A KL Divergence Identity

We use the following lemma later in context of PAC-Bayes bounds.

Lemma 9.3. *For any two probability measures μ, ρ defined on the probability space $(\Omega, \mathcal{F})$, assume that $\mu_{|A}$ is defined as the normalized probability measure restricted to the set $A \in \mathcal{F}$. Then we have:*

$$D(\mu\|\rho) = \mu(A)D(\mu_{|A}\|\rho) + \mu(A^c)D\left(\mu_{|A^c}\|\rho\right) + h_b(\mu(A)), \tag{29}$$

where $h_b(\cdot)$ is the binary entropy defined as $h_b(p) = -p\log p - (1-p)\log(1-p)$.

The proof of this lemma follows from standard integral manipulations.

9.4 PAC-Bayesian Generalization Bound - Derandomization Technique

The PAC-Bayesian generalization lemma of 4.2 provides a bound on the generalization error of randomized classifiers drawn from the posterior distribution Q. To obtain a bound that holds for individual hypothesis, we need to de-randomize the bound. The following lemma, a reformulation of Neyshabur et al. [2018], provides a way of achieving this goal.

Lemma 9.4. *Let $\mathcal{P}$ be a finite set of priors P over the hypothesis space. Let the set $\mathcal{S}_{h_0}$ be defined as:*

$$\mathcal{S}_{h_0} = \{h \in \mathcal{H} : \|h - h_0\|_\infty \leq \gamma/4\}.$$

If the hypothesis after training is chosen using the posterior Q, set $\eta := Q(\mathcal{S}_{h_0})$. Then with probability $1 - \delta$, we have for all Q:

$$\mathcal{L}(h_0) \leq \hat{\mathcal{L}}_\gamma(h_0) + \sqrt{\frac{\min_{P \in \mathcal{P}} D(Q\|P) + \eta \log(\xi(m)|\mathcal{P}|/\delta)}{2m\eta}}, \tag{30}$$

where $\xi(m) := \sum_{k=0}^m \binom{m}{k}(k/m)^k(1-k/m)^{m-k}$.

Proof. PAC-Bayesian bound in Lemma 4.2 holds for a single P. Taking the union bound over all $P \in \mathcal{P}$, we can see that with probability at least $1 - |\mathcal{P}|\delta$ for all $P \in \mathcal{P}$ and for all Q over $\mathcal{H}$, the bound in Lemma 4.2 holds. Choose the margin loss eq. 1 with margin $\gamma/2$ as the loss function. After replacing δ with $\delta/|\mathcal{P}|$, the following inequality holds with probability $1 - \delta$:

$$\mathcal{L}_{\gamma/2}(Q) \leq \hat{\mathcal{L}}_{\gamma/2}(Q) + \sqrt{\frac{\min_{P \in \mathcal{P}} D(Q\|P) + \log(\xi(m)|\mathcal{P}|/\delta)}{2m}}. \tag{31}$$

For each Q, define $\hat{Q}$ as:

$$\hat{Q}(h) := \begin{cases} \frac{1}{\eta}Q(h) & h \in \mathcal{S}_{h_0} \\ 0 & \text{otherwise.} \end{cases} \tag{32}$$

Since eq. 31 holds for all Q, we can use $\hat{Q}$ in the bound. Let h be drawn from $\hat{Q}$. Then h belongs to $\mathcal{S}_{h_0}$. Knowing that, we follow the argument we sketched above to obtain bound on the risks of h_0. Since $h \in \mathcal{S}_{h_0}$, $\|h - h_0\|_\infty \leq \gamma/4$. We have:

$$\left(h(\boldsymbol{x})[y] - \max_{j \neq y} h(\boldsymbol{x})[j]\right) \leq \left(h_0(\boldsymbol{x})[y] + \gamma/4 - \max_{j \neq y}(h_0(\boldsymbol{x})[j] - \gamma/4)\right) \tag{33}$$

$$\leq \left(h_0(\boldsymbol{x})[y] - \max_{j \neq y} h_0(\boldsymbol{x})[j]\right) + \gamma/2, \tag{34}$$

and therefore:

$$\mathcal{L}(h_0) = \mathbb{P}_{(\boldsymbol{x},y)\sim\mathcal{D}}\left(h_0(\boldsymbol{x})[y] \leq \max_{j \neq y} h_0[j]\right) \leq \mathcal{L}_{\gamma/2}(h) = \mathbb{P}_{(\boldsymbol{x},y)\sim\mathcal{D}}\left(h(\boldsymbol{x})[y] \leq \gamma/2 + \max_{j \neq y} h[j]\right). \tag{35}$$

Similarly we have:

$$\left(h_0(\boldsymbol{x})[y] - \max_{j \neq y} h_0(\boldsymbol{x})[j]\right) \leq \left(h(\boldsymbol{x})[y] - \max_{j \neq y} h(\boldsymbol{x})[j]\right) + \gamma/2, \tag{36}$$

which provides a bound on the empirical risk as follows:

$$\hat{\mathcal{L}}_{\gamma/2}(h) \leq \hat{\mathcal{L}}_{\gamma}(h_0). \tag{37}$$

Since inequalities 35 and 37 hold for all $h \in \mathcal{S}_{h_0}$, we get:

$$\mathcal{L}(h_0) \leq \mathcal{L}_{\gamma/2}(\hat{Q}) \quad \text{and} \quad \hat{\mathcal{L}}_{\gamma/2}(\hat{Q}) \leq \hat{\mathcal{L}}_{\gamma}(h_0), \tag{38}$$

from which we immediately obtain the following de-randomized bound:

$$\mathcal{L}(h_0) \leq \hat{\mathcal{L}}_{\gamma}(h_0) + \sqrt{\frac{\min_{P \in \mathcal{P}} D(\hat{Q}\|P) + \log(\xi(m)|\mathcal{P}|/\delta)}{2m}}. \tag{39}$$

Applying the lemma 9.3, we get $\eta D\left(\hat{Q}\|P\right) \leq D\left(Q\|P\right)$. Using this inequality, the lemma is obtained from eq. 39.

$\square$

9.5 Proof of Theorem 9.1 and Generalization Bounds for Equivariant Networks

Proof. We consider the perturbation bound in Eq. 7:

$$\|f_{\mathbb{W}+\mathbb{U}} - f_{\mathbb{W}}\|_{\infty} \leq \|f_{\mathbb{W}+\mathbb{U}} - f_{\mathbb{W}}\|_2 \leq eB \left(\prod_{i=1}^{L} \|\boldsymbol{W}_i\|_2\right) \sum_{i=1}^{L} \frac{\|\boldsymbol{U}_i\|_2}{\|\boldsymbol{W}_i\|_2}. \tag{40}$$

By applying the results found in Supplementary 9.2 together with a union bound argument, with probability $1 - \left(\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}\right) e^{-t}$, for all l, it holds (for $t > 1$):

$$\|\boldsymbol{U}_l\|_2 \leq \sigma \sqrt{\max_{\psi} \left(5m_{l-1,\psi} m_{l,\psi} c_{\psi} t\right)}. \tag{41}$$

Therefore, with probability $1 - \left(\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}\right) e^{-t}$,

$$\|f_{\mathbb{W}+\mathbb{U}} - f_{\mathbb{W}}\|_{\infty} \leq eB \left(\prod_{i=1}^{L} \|\boldsymbol{W}_i\|_2\right) \sum_{l=1}^{L} \frac{\sigma \sqrt{\max_{\psi} \left(5m_{l-1,\psi} m_{l,\psi} c_{\psi} t\right)}}{\|\boldsymbol{W}_l\|_2}. \tag{42}$$

Choosing $t = \log\left(\frac{\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}}{1-\eta}\right)$, with probability η:

$$\|f_{\mathbb{W}+\mathbb{U}} - f_{\mathbb{W}}\|_{\infty} \leq eB \left(\prod_{i=1}^{L} \|\boldsymbol{W}_i\|_2\right) \sum_{l=1}^{L} \frac{\sigma \sqrt{\max_{\psi} \left(5m_{l-1,\psi} m_{l,\psi} c_{\psi} \log\left(\frac{\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}}{1-\eta}\right)\right)}}{\|\boldsymbol{W}_l\|_2}. \tag{43}$$

To reduce clutter in the next equations, we define

$$M(l,\eta) := \log\left(\frac{\sum_{l=1}^{L} \sum_{\psi} m_{l,\psi}}{1-\eta}\right) \max_{\psi} \left(5m_{l-1,\psi} m_{l,\psi} c_{\psi}\right) \tag{44}$$

such that the previous bound becomes:

$$\|f_{\mathbb{W}+\mathbb{U}} - f_{\mathbb{W}}\|_{\infty} \leq eB \left(\prod_{i=1}^{L} \|\boldsymbol{W}_i\|_2\right) \sum_{l=1}^{L} \frac{\sigma \sqrt{M(l,\eta)}}{\|\boldsymbol{W}_l\|_2}. \tag{45}$$

We can then consider a set $\mathcal{P}$ of possible values for σ, such that for any σ_0 there exists a $\sigma \in \mathcal{P}$ satisfying

$$|\sigma - \sigma_0| \leq \kappa \sigma_0 \tag{46}$$

with

$$\sigma_0 := \frac{\gamma}{4eB(1+\kappa)\left(\prod_{i=1}^{L}\|\boldsymbol{W}_i\|_2\right)\sum_{i=1}^{L}\frac{\sqrt{M(l,\eta)}}{\|\boldsymbol{W}_i\|_2}}. \tag{47}$$

By choosing the covariance of the distribution Q from this set $\mathcal{P}$ of values for σ, we get

$$Q(\mathcal{S}_{f_{\mathbb{W}}}) \geq \eta. \tag{48}$$

Again, suppose that such a set $\mathcal{P}$ of σ values can be chosen and that it has cardinality $|\mathcal{P}|$. We can then use the PAC-Bayesian KL-bound in Eq. 5 and choose a σ satisfying the above inequality to obtain:

$$D(Q\|P) \leq \frac{(1+\kappa)^2 32e^2 B^2\left(\prod_{i=1}^{L}\|\boldsymbol{W}_i\|_2^2\right)}{(1-\kappa)^2\gamma^2}\left(\sum_{l=1}^{L}\frac{\sqrt{M(l,\eta)}}{\|\boldsymbol{W}_l\|_2}\right)^2\left(\sum_{l,\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2/\dim_\psi\right) \tag{49}$$

$$\leq \frac{(1+\kappa)^2 32Le^2 B^2\left(\prod_{i=1}^{L}\|\boldsymbol{W}_i\|_2^2\right)}{(1-\kappa)^2\gamma^2}\left(\sum_{l=1}^{L}\frac{M(l,\eta)}{\|\boldsymbol{W}_l\|_2^2}\right)\left(\sum_{l,\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2/\dim_\psi\right). \tag{50}$$

We need to construct a set $\mathcal{P}$ such that for any σ_0 there exists a $\sigma \in \mathcal{P}$ satisfying inequality 46. Using a similar argument to Neyshabur et al. [2018], it can be seen that the bound is non-vacuous if $\prod_{l=1}^{L}\|\boldsymbol{W}_l\|_2 \in [\gamma/2B, \gamma\sqrt{m}/2B]$. If the product of norms after training is outside the interval, the bound holds trivially, and any (P, Q) can be chosen. Therefore, we cover only this interval. Details follow below.

In homogeneous networks, we can always rescale the weights such that the margin is not touched, and thereby change all the norms. The above generalization bound is not invariant to rescaling of weights. Here, we derive a scaling-invariant version of it. Define $\beta^L = \prod_{i=1}^{L}\|\boldsymbol{W}_i\|_2$ and normalize every layer to have norm β. Then, σ_0 can be written as:

$$\sigma_0 = \frac{\gamma}{4eB(1+\kappa)\beta^{L-1}\sum_{l=1}^{L}\sqrt{M(l,\eta)}}. \tag{51}$$

To cover the set σ_0, we require only to cover β effectively. Again, we consider only $\beta^L > \frac{\gamma}{2B}$ and $\beta^{2L} < \left(\frac{\gamma^2 m}{4B^2}\right)$, such that the bound is not vacuous. Note that the generalization bound in Eq. 49 is simplified to (removing κ as we cover our set differently):

$$\frac{32e^2 B^2\beta^{2L}}{\gamma^2}\left(\sum_{l=1}^{L}\frac{\sqrt{M(l,\eta)}}{\beta}\right)^2\left(\sum_{l}\frac{\beta^2}{\|\boldsymbol{W}_l\|_2^2}\sum_{\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2/\dim_\psi\right) \tag{52}$$

$$= \frac{32e^2 B^2\beta^{2L}}{\gamma^2}\left(\sum_{l=1}^{L}\sqrt{M(l,\eta)}\right)^2\left(\sum_{l}\frac{\sum_{\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2/\dim_\psi}{\|\boldsymbol{W}_l\|_2^2}\right). \tag{53}$$

Therefore, if $\beta > (\frac{\gamma\sqrt{m}}{2B})^{1/L}$ all the remaining terms are bigger than one. Specifically note that:

$$\|\boldsymbol{W}_l\|_2^2 \leq \sum_{\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_2^2 = \sum_{\psi,i,j}\left\|\widehat{\boldsymbol{W}}_l(\psi,i,j)\right\|_F^2/\dim_\psi \tag{54}$$

where we first used the fact that the spectral norm of a block matrix is upperbounded by the sum of the spectral norm of each block and then we used Eq. 19 to express the spectral norm of each block.

Therefore we need only to cover β within the interval $(\frac{\gamma}{2B})^{1/L} \leq \beta \leq (\frac{\gamma\sqrt{m}}{2B})^{1/L}$ with radius $1/d(\frac{\gamma}{2B})^{1/L}$.

This choice guarantees that there is a $\tilde{\beta}$ such that $\left\|\beta - \tilde{\beta}\right\| \leq \beta/d$. From which, it can be concluded that $1/e\beta^{L-1} \leq \hat{\beta}^{L-1} \leq e\beta^{L-1}$. Using a cover of size $Lm^{1/2L}$ provides the desired cover. $\qquad\square$

10 Derivations for Group Convolutional Networks

In this section, we provide a special case of Theorem 9.1 for group convolutional networks.

In a group convolution architecture, we assume the features at the l-th layer are $C_l = c_l|H|$-dimensional vectors representing a c_l-channels signal over H. This means ρ_l contains c_l copies of the regular representation. The linear layer consists of a matrix $\boldsymbol{W}_l$ with a block structure as in Eq. 18, where the block $\boldsymbol{W}_{l,(i,j)}$ is an H-circulant matrix mapping the i-th input channel to the j-th output channel via group convolution with a filter $\boldsymbol{w}_{l,i,j} \in \mathbb{R}^{|H|}$. The parametrization of this block is in term of the Fourier transform of $\boldsymbol{w}_{l,i,j}$, i.e. the coefficient in the matrices $\widehat{\boldsymbol{w}_{l,i,j}}(\psi)$, indexed by the irreps $\psi \in \hat{H}$. Since we use real irreps, we only need the first $\dim_\psi /c_\psi$ columns of $\widehat{\boldsymbol{w}_{l,i,j}}(\psi)$, so we assume it is a $\dim_\psi \times \frac{\dim_\psi}{c_\psi}$ matrix. Similarly, the Fourier transform of a single channel $\boldsymbol{x}_{l,i} \in \mathbb{R}^{|H|}$ only contains the first $\dim_\psi /c_\psi$ columns of $\widehat{x}_{l,i}(\psi)$. In the notation used before, this means that ψ is contained in each channel $\dim_\psi /c_\psi$ times and, therefore, for each (intermediate) layer l and irrep ψ, it holds that:

$$m_{l,\psi} = c_l \dim_\psi /c_\psi.$$

Note also that, for each block (i,j), the matrix $\widehat{\boldsymbol{w}_{l,i,j}}(\psi) \in \mathbb{R}^{\dim_\psi \times \frac{\dim_\psi}{c_\psi}}$ contains the c_ψ coefficients in $\widehat{\boldsymbol{w}_{l,\cdot,\cdot}}(\psi)$ associated to each of the $\frac{\dim_\psi}{c_\psi} \times \frac{\dim_\psi}{c_\psi}$ pairs of input and output occurrences of ψ in $\boldsymbol{x}_{l-1,i}$ and $\boldsymbol{x}_{l,j}$.

Assuming a group-convolutional architecture, we can use the identity $m_{l,\psi} = c_l \frac{\dim_\psi}{c_\psi}$. Define:

$$D_H := \max_\psi \frac{\dim_\psi^2}{c_\psi} \tag{55}$$

$$E_H := \sum_\psi \frac{\dim_\psi}{c_\psi}. \tag{56}$$

In case of a group convolution architecture, we get:

$$M(l,\eta) = \log\left(\frac{E_H \sum_{l=1}^{L} c_l}{1-\eta}\right) 5c_l c_{l-1} D_H \tag{57}$$

and the bound in Theorem 9.1 becomes:

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \leq \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) +$$

$$\sqrt{\frac{32e^4 B^2 \beta^2}{\gamma^2 m\eta}\left(\sum_{l=1}^{L}\sqrt{5c_l c_{l-1}}\right)^2 D_H \log\left(\frac{E_H \sum_{l=1}^{L} c_l}{1-\eta}\right)\left(\sum_l \frac{\sum_{\psi,i,j}\left\|\widehat{\boldsymbol{w}_{l,j,i}}(\psi)\right\|_F^2}{\|\boldsymbol{W}_l\|_2^2}\right) + \frac{\log(\xi(m)^{\frac{Lm^{1+1/2L}}{\delta}})}{2\gamma^2 m}}$$

By choosing $\eta = 1/2$ and by defining $Q_H = \left(\sum_{l=1}^{L}\sqrt{c_{l-1}c_l}\right)^2 D_H \log 2E_H \sum_{l=1}^{L} c_{l-1}$, this bound corresponds to Theorem 10.1.

Theorem 10.1 (Generalization Bounds for Group Convolutional Networks). *For any group convolutional network, with high probability, we have:*

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \leq \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) + \tilde{\mathcal{O}}\left(\sqrt{\frac{\left(\prod_{l=1}^{L}\|\boldsymbol{W}_l\|_2^2\right)Q_H\left(\sum_l^L \frac{\sum_{\psi,i,j}\left\|\widehat{\boldsymbol{w}_{l,i,j}}(\psi)\right\|_F^2}{\|\boldsymbol{W}_l\|_2^2}\right)}{\gamma^2 m}}\right)$$

with $Q_H = \left(\sum_{l=1}^{L}\sqrt{c_{l-1}c_l}\right)^2 D_H \log 2E_H \sum_{l=1}^{L} c_{l-1}$.

First, note that the constant factor D_H is at most 2 for commutative groups (all subgroups of $\mathrm{SO}(2)$) and at most 4 for other subgroups of $\mathrm{O}(2)$. In practical networks, the number channels are inversely scaled with the group size. This means that the $c_l|H|$ are kept constant for different values of $|H|$.

In terms of generalization error, this implies an improvement of order $1/|H|$ not withstanding the impact of group size on the ratio of Frobenius and Spectral norms.

It is worth mentioning an intermediate result for proving the above theorem. We can adapt the tail bound of eq. 11 to perturbations for group convolutional networks:

$$\|\boldsymbol{U}_l\|_2 < \sigma \sqrt{5c_{l-1}c_l \left(\max_\psi \frac{\dim_\psi^2}{c_\psi} \right) \log \left(2 \sum_l c_l \sum_\psi \frac{\dim_\psi}{c_\psi} \right)}.$$

which is equal to Eq. 11 when defining $D_H = \max_\psi \frac{\dim_\psi^2}{c_\psi}$ and $E_H = \sum_\psi \frac{\dim_\psi}{c_\psi}$. The bound can be further simplified by noting that $|H| = \sum_\psi \dim_\psi^2 / c_\psi \geq \sum_\psi \dim_\psi / c_\psi = E_H$:

$$\|\boldsymbol{U}_l\|_2 < \sigma \sqrt{5c_{l-1}c_l D_H \log \left(2|H| \sum_l c_l \right)}.$$

This tail bound can be of independent interest.

11 Comparison to Neyshabur et al. [2018]

In this section, we try to directly adapt the proof of Neyshabur et al. [2018] to show the benefit of working in space of irreps.

First, note that the Forbenius norm of kernels is an upper-bound on the norm of parameters. To see this, similar to Neyshabur et al. [2018], define $h = \max_l C_l$, where $C_l = c_l|H|$ is the effective number of channels, as the upper bound on the number of channels, and similarly $h_{|H|} = \max_l c_l = h/|H|$. In the hidden layers and the input layer,

$$\sum_\psi \left\| \widehat{w_{l,i,j}}(\psi) \right\|_F^2 \leq \sum_\psi \dim_\psi \left\| \widehat{w_{l,i,j}}(\psi) \right\|_F^2 = |H| \left\| \boldsymbol{w}_{l,i,j} \right\|_2^2 = \left\| \boldsymbol{W}_{l,(j,i)} \right\|_F^2$$

and, therefore, $\sum_{i,j,\psi} \left\| \widehat{w_{l,i,j}}(\psi) \right\|_F^2 \leq \|\boldsymbol{W}_l\|_F^2$. By also upperbounding $c_l = C_l/|H| \leq h/|H|$ and ignoring the constant factors, the result in Theorem 10.1 becomes

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \leq \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) + \tilde{\mathcal{O}} \left(\frac{1}{|H|} \sqrt{\frac{\left(\prod_{l=1}^L \|\boldsymbol{W}_l\|_2^2 \right) L^2 h^2 D_H \log(2Lh \frac{E_H}{|H|}) \left(\sum_l^L \frac{\|\boldsymbol{W}_l\|_F^2}{\|\boldsymbol{W}_l\|_2^2} \right)}{\gamma^2 m}} \right)$$

which can be further simplified to

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \leq \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) + \tilde{\mathcal{O}} \left(\frac{1}{|H|} \sqrt{\frac{\left(\prod_{l=1}^L \|\boldsymbol{W}_l\|_2^2 \right) L^2 h^2 \log(2Lh) \left(\sum_l^L \frac{\|\boldsymbol{W}_l\|_F^2}{\|\boldsymbol{W}_l\|_2^2} \right)}{\gamma^2 m}} \right)$$

by assuming D_H is approximately constant and by noting $E_H \leq |H|$.

This bound looks very similar to the one proposed in Neyshabur et al. [2018], with the difference that it contains 1) h^2 instead of h and 2) an additional term $\frac{1}{|H|}$. The additional h factor is constant in our experiments as we always consider fixed architectures when changing the equivariance group H, so we ignore it in this discussion. Instead, we want to drive the attention to the $\frac{1}{|H|}$ term.

The presence of this $\frac{1}{|H|}$ in our bound implies that, when changing the equivariance group H for a fixed architecture, the bound from Neyshabur et al. [2018] scales at least $|H|$ times worse than ours. In the experiments, we empirically observe that our bound is approximately scaling like $\frac{1}{\sqrt{|H|}}$. This can be explained by the fact that with larger group size, the ratio of Frobenious norm and spectral norm increases similar to $O(|H|)$, thereby increasing the generalization error by $O(\sqrt{|H|})$. This

leads to an overall scaling of $O(1/\sqrt{|H|})$ observed by our experiments too. On the other hand, it follows that the bound from Neyshabur et al. [2018] scales like $\sqrt{H}$, which is undesirable.

However, it is still possible to adapt the method in Neyshabur et al. [2018] by considering, as done in this work, perturbations only in the equivariant subspace of the weights. The main difference with respect to our method is how the spectral norm of the perturbation matrices $\|U_l\|_2$ is bounded. In the rest of this section, we follow this strategy to derive an alternative generalization bound.

The strategy used in Neyshabur et al. [2018] can be adapted to any parametrization of the linear layers. In particular, assume c_{l-1} and c_l input and output channels and $W_l = \sum_k w_{l,k} B_{l,k}$ for some set of matrices $\{B_{l,k} \in \mathbb{R}^{c_l \times c_{l-1}}\}$ forming a basis for the linear layer W_l.

Theorem 11.1 (Tropp [2012]). *Assume $\{u_{l,k}\}_k$ are i.i.d. random variables with a standard gaussian distribution $\mathcal{N}(0,1)$. Define the variance parameter*

$$v_l^2 = \max\left\{\left\|\sum_k B_{l,k}^* B_{l,k}\right\|_2 , \left\|\sum_k B_{l,k} B_{l,k}^*\right\|_2\right\} .$$

Then, for any $t \geq 0$:

$$\mathbb{P}\left(\left\|\sum_k u_{l,k} B_{l,k}\right\|_2 \geq t\right) \leq (c_l + c_{l-1})e^{-\frac{t^2}{2v^2}}$$

If perturbation is done on the weights $\{w_{l,k}\}_k$ with i.i.d. $\{u_{l,k} \sim \mathcal{N}(0, \sigma^2)\}_k$ and $h = \max c_l, c_{l-1}$, this result can be used to bound the spectral norm of the perturbation matrix as

$$\|U_l\|_2 \leq \sigma\sqrt{2v_l^2\sigma^2 \log(4h)}$$

Using an union bound over all layers $l \in [L]$, one gets:

$$\|U_l\|_2 \leq \sqrt{2v^2\sigma^2 \log(4hL)}$$

where $v^2 = \max_l v_l^2$.

In particular, in Neyshabur et al. [2018], the parametrization $W_l = \sum_{i,j} w_{l,i,j} E_{i,j}$ was considered, where $w_{l,i,j}$ is the entry of W_l at row j and column i while $E_{i,j}$ is a matrix of the same shape containing 0 everywhere but 1 at position (i,j). In that case, $v^2 = h$.

We can express the basis considered in this work as

$$W_l = \sum_{\psi,i,j,k} \widehat{w_{l,i,j}}(\psi)_k B_{l,\psi,i,j,k} ,$$

where $\widehat{w_{l,i,j}}(\psi)_k$ is the k-th entry of the vector $\widehat{w_{l,i,j}}(\psi)$ containing the c_ψ learnable coefficients which parametrize the block $\widehat{W}_l(\psi,i,j)$. Indeed, recall that a block $\widehat{W}_l(\psi,i,j)$ always has one of the three forms described in Supplementary 8 and can be written as $\widehat{W}_l(\psi,i,j) = \sum_{k=1}^{c_\psi} \widehat{w}_l(\psi,i,j)_k B_{\psi,k}$. The matrix $B_{l,\psi,i,j,k}$ is simply a zero matrix containing the matrix $B_{\psi,k}$ at the same position of the corresponding block $\widehat{W_l(\psi,i,j)}$. By using the structure of the matrices $B_{l,\psi,i,j,k}$ and the orthonormality of all matrices $B_{\psi,k}$, one can show that in this case $v_l^2 = \max_\psi c_\psi \max_l m_{l,\psi}$. When using a group convolution architecture, because $m_{l,\psi} = c_l \frac{\dim_\psi}{c_\psi}$, this becomes $v_l^2 = \max_\psi \dim_\psi h_{|H|} = \max_\psi \dim_\psi h/|H|$. We can use this result in a similar manner to obtain a bound on the generalization error: Assuming again that $\max_\psi \dim_\psi$ is a constant factor, it follows that the bound on the spectral norm in our parametrization is $|H|$ times tighter than the one obtained in Neyshabur et al. [2018].

$$\mathcal{L}_\mathcal{D}(f_\mathbb{W}) \leq \hat{\mathcal{L}}_\gamma(f_\mathbb{W}) + \tilde{\mathcal{O}}\left(\frac{1}{\sqrt{|H|}}\sqrt{\frac{(\max_\psi \dim_\psi) L^2 h \log(2Lh)\left(\prod_{l=1}^L \|W_l\|_2^2\right)\left(\sum_l^L \frac{\|W_l\|_F^2}{\|W_l\|_2^2}\right)}{\gamma^2 m}}\right)$$

$$(58)$$

This bound explicitly shows the scaling of the generalization error as $\frac{1}{\sqrt{|H|}}$ which we also verified experimentally. A similar scaling was reported in Sokolić et al. [2017a], however it was mainly based on the assumption that the transformations change the input space in a considerable way. However, in our experiments we observe the bound in Theorem 9.1 empirically scales like $\frac{1}{\sqrt{|H|}}$. Because the bound in Eq. 58 is at least $\frac{1}{\sqrt{|H|}}$ times worst, we do not expect it to correlate significantly with the equivariance group H. We also verify this hypothesis empirically in our experiments. See Fig. 2 and 3 for a comparison of the two bounds on different continuous synthetic datasets.

A similar strategy could also be applied by considering bases for group convolution layers directly in terms of the filters parametrizing each H-circulant matrix (i.e. by performing the perturbation analysis in the group domain). More precisely, one could consider a basis

$$W_l = \sum_{i,j,k} \boldsymbol{w}_{l,i,j,k} \boldsymbol{C}_{l,i,j,k} \ ,$$

where $\boldsymbol{w}_{l,i,j} \in \mathbb{R}^{|H|}$ and $\boldsymbol{w}_{l,i,j,k}$ is its k-th entry. $\boldsymbol{C}_{l,i,j,k}$ is then a zero matrix containing the a base circulant matrix $C_{H,k}$ in the block (i,j). The matrix $C_{H,k}$ is an H-circulant matrix generated by the vector $\boldsymbol{e}_k = (0, \ldots, 0, 1, 0, \ldots 0)^T \in \mathbb{R}^{|H|}$ (with 1 in the k-th entry). For instance, for $H = \mathrm{C}_3$

$$\{C_{\mathrm{C}_3,k}\}_{k=1}^{3} = \left\{ \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \right\} \ .$$

In this parametrization, $v_l^2 = |H| \max\{\mathrm{c}_l, \mathrm{c}_{l-1}\}$ and, therefore, $v^2 = h$ as in Neyshabur et al. [2018]. Indeed, this leads exactly to their same bound, which we have already discussed at the beginning of this section.

12   Experiments

We experiment on synthetic datasets characterized by discrete or continuous symmetries as well as with transformed MNIST 12K and CIFAR10 datasets.

We choose a *discrete* subgroup $H < G$ and build a shallow MLP classifier equivariant to H. In particular, denoting as ρ_0 the representation of G acting on the samples in a G symmetric dataset, the input representation of the MLP is chosen to be $\mathrm{Res}_H^G \rho_0$ while in the hidden layers we use multiple copies of the regular representation ρ_{reg}^H of H, making the linear layers effectively group convolutions over H. For different choices of H, we preserve the size of each layer, which means that the number of parameters of the model is proportional to $\frac{1}{|H|}$. Note that a larger group H results in a higher level of symmetry for the network but also in a loss in capacity, as less channels are allocated per each group element.

We train each model on a fixed training set until the model correctly classifies 99% of it with a margin greater than γ. We use $\gamma = 10$ in the synthetic datasets and $\gamma = 2$ on the image datasets. In the synthetic datasets, we test the models on a testset consisting of $10K$ fixed samples from the full distribution $\mathbb{P}_G$ in order to measure the generalization on the whole symmetry group G.

The MLP used in the experiments on the synthetic datasets consists of 3 linear layers, alternated with ReLU non-linearities, and has 2048 and 512 channels in the intermediate features. In the MNIST experiments, we use 4 linear layers with 1024, 512 and 512 channels in the intermediate features. Finally, in the CIFAR experiments we use 4 layers with 2048, 1536 and 512 channels. The first linear layer of the last two models uses steerable filters from Weiler and Cesa [2019] to process the input images. The models are trained with *Adam* without any weight decay or additional regularization. Neither batch normalization nor dropout are used in the models.

To handle input images, we use an $E(2)$-convolution layer as described in Weiler and Cesa [2019]: with a filter as large as the input image, the output produced is a single vector which is only H equivariant. The implementation of the kernel constraint from Weiler and Cesa [2019] automatically performs the necessary band-limiting and implicitly decomposes the image as a signal over the group H considered.

12.1 Synthetic Datasets

Continuous Symmetries For $G = \mathrm{SO}(2)$, the data distribution is defined over a D-dimensional torus $\mathcal{T}^D$ embedded in a $2D$ dimensional Euclidean space. The action of an element $r_\theta \in \mathrm{SO}(2)$ simultaneously rotates the D circles in $\mathcal{T}^D$, potentially at different integer frequencies in each of them. Note that each circle is isomorphic to $\mathrm{SO}(2)$. We generate one point in the first circle and two points in all the others. Then, we generate 2^{D-1} representative points X by tacking every combination of points from each circle. We assign a random label in $\{-1, 1\}$ to each of them. The quotient distribution $\mathcal{S}_0$ is defined by sampling a random representative in X and adding some noise. In practice, we add Gaussian noise on each of the D circles and, then, project each of them on the unit circle. This is the training augmentation $\mathbb{P}_S$. We finally add additional small Gaussian noise. Fig. 7 shows a projection of the dataset for $D = 2$. In our experiments we use $D = 6$. We also build

(a) $\mathrm{SO}(2)$ rotates both circles with frequency 1

(b) $\mathrm{SO}(2)$ rotates the largest circle (horizontal) with frequency 1 and the smallest (vertical) with frequency 2.

Figure 7: $3D$ projection of $\mathrm{SO}(2)$ synthetic datasets for $D = 2$.

a similar dataset for $\mathrm{O}(2)$. Here, each circle is replaced by a pair of circles such that the action of the reflection moves the points from one to the other. Each pair of circles is isomorphic to the group $\mathrm{O}(2)$ itself. We fix two points per pair such that we still have 2^{D-1} representative points but embedded in a $4D$-dimensional space.

Discrete Symmetries We consider a group G of N discrete rotations and, possibly, reflections. We build a dataset as described for $\mathrm{O}(2)$ using $D = N$ pairs, each associated with a different frequency from 1 to $F = \lfloor N/2 \rfloor$. We associate all the 2^{F-1} representative points in X with the label $+1$. If we do not require symmetry to reflections (i.e., $G = \mathrm{C}_N$), we generate 2^{F-1} new representative points by rotating those in X by π/N and associating them with the label -1. If symmetry to reflections is necessary ($G = \mathrm{D}_N$), we generate $3 \cdot 2^{F-1}$ new representative points by i) rotating X by π/N, ii) by mirroring it or iii) by doing both. In the first two cases, we associate the points with the label -1, in the last with $+1$. Note that a model equivariant to D_{2N} or C_{2N} will be invariant to (all or part of) the transformations we used to generate the labels and, therefore, will not be able to distinguish the two classes.

12.2 Dependency on the group size $|H|$

In this section we study the effect of the size $|H|$ of the equivariance group on the generalization error. In particular, we hypothesize that the generalization errors is proportional to the quantity $\frac{1}{\sqrt{|H|}}$ as previously observed in Sokolić et al. [2017a], so we investigate the correlation between these two terms in different datasets.

We consider the synthetic datasets with continuous and discrete symmetries (for different frequencies F or rotation orders M) in Fig. 8 and the images datasets (MNIST and CIFAR10) in Fig. 9. In

(a) $G = \mathrm{SO}(2)$, $H = C_N$ (b) $G = \mathrm{O}(2)$, $H = D_N$ or C_N

(c) $G = C_M$, $H = C_N$ (d) $G = D_M$, $H = D_N$ or C_N

Figure 8: Empirical generalization error (GE) vs $\frac{1}{\sqrt{|H|}}$ for different synthetic datasets, with continuous or discrete symmetries G. In the continuous symmetries case (first row), F indicates the maximum rotational frequency in the data and, therefore, each color (F) corresponds to a different dataset. In the discrete symmetries case (second row), M indicates the the number of rotations in the discrete symmetry. The visualizations shows a strong correlation between the two terms. The correlation is weaker on low frequency $\mathrm{O}(2)$ datasets because of the different behaviours of D_N and C_{2N}.

all cases, we observe a strong correlation. However, we note that the slope of the lines varies over different versions of the same dataset or when changing the training set size m. In particular, in Fig. 8(first line), the slope grows when increasing the frequency F in the data. In the discrete symmetries case, Fig. 8(second line), the slope increases with the symmetry group's size $|G|$ but then decreases at the largest values of $|G|$. In the $\mathrm{O}(2)$ synthetic data (Fig. 8(b)), the linear correlation is weaker for low values of F as the generalization does not depend only on the size $|H|$ of the group anymore. Indeed, the groups C_{2N} and D_N have both size $|H| = 2N$. However, when the data only contains low frequency features, considering 2 times more rotations (C_{2N}) becomes eventually unnecessary while introducing reflection equivariance (D_N) allows the model to generalize over a whole new set of transformations. Overall, the

12.3 PAC-Bayesian Bound

In this section, we focus on the study of the de-randomized PAC-Bayes bound from Sec. 4 in different contexts. As reported in Nagarajan and Kolter [2019], bounds based on the spectral norm of the weight matrices tend to grow with the dataset size m in practice. Our perturbation bound falls in this same category and, therefore, we observe a similar behaviour. Moreover, such bounds are generally vacuous, i.e. greater than 1. For these reasons, we do not expect the bound derived to accurately predict the generalization error or describe the effect of different training sizes. Instead, we are

(a) MNIST, $G = \mathrm{SO}(2)$ (b) MNIST, $G = \mathrm{O}(2)$

(c) CIFAR10, $G = \mathrm{SO}(2)$ (d) CIFAR10, $G = \mathrm{O}(2)$

Figure 9: Empirical generalization error (GE) vs $\frac{1}{\sqrt{|H|}}$ on transformed MNIST and CIFAR10 datasets. All settings show a strong correlation between the two terms.

interested in how well the bound correlates with the generalization error and explains the effect of equivariance. Therefore, we study the correlation between the estimated bound and the generalization error in different settings and datasets when different equivariance groups H are used.

In Fig. 10 we look at the correlation between our bound and the generalization error on $\mathrm{O}(2)$-MNIST and on the $\mathrm{SO}(2)$ symmetric synthetic dataset for different training set sizes. Nevertheless, we observe a good correlation between them when considering a fixed training set size m, i.e. by looking at each colored sequence independently. This is the relation we are most interested in and which we now explore further.

In Fig. 2(first row) and 3(first row) we have already observed this correlation on 6 versions of the $\mathrm{O}(2)$ and the $\mathrm{SO}(2)$ symmetric synthetic datasets. In Fig. 11, we repeat a similar study on the other three types of datasets.

We notice that the results on the synthetic datasets (both with continuous symmetries as in Fig. 2 and 3 and with discrete symmetries as in the first row of Fig. 11) show a better linear correlation with the generalization error. Conversely, the results on the image datasets are in log scale on the Y axis, suggesting a superlinear scaling of the bound with the generalization error. In light of the observations in Supplementary 12.2, this also implies that the bound does not scale as $\frac{1}{\sqrt{|H|}}$ on these two datasets.

It also follows that the alternative bounds considered in Supplementary 11 are not worse here. However, in the synthetic dataset with discrete symmetries, the same observation from Section 5 apply. Indeed, in Fig. 12, we repeat the experiments in Fig. 11(b) but use the alternative bound from Supplementary 11. As already shown in Fig. 2(second row) and 3(second row) for the continuous symmetry synthetic datasets, this bound does not capture the effect of different equivariance groups H.

(a) SO(2) Synthetic Data (b) O(2) MNIST

Figure 10: Bound (B) from Theorem 9.1 vs Generalization Error (GE) of different H-equivariant models ($H = C_N$ or $H = D_N$) on O(2)-MNIST and on the SO(2) synthetic dataset. Different colors represent different training set sizes m. As expected, the bound does not explain the effect of the training set size and as it grows with m.

12.4 PAC-Bayes Bound During Training

In this section, we study how the bound in Section 4.2 evolves during training. In particular, we consider a few different equivariance groups H, and we train them on the original labels as well as on random labels.

Fig. 13 reports the results of these experiments on the two of the continuous synthetic datasets. We observe that the bounds grow during training for all models. This happens using both the original and the random labels. Moreover, the bound computed with the randomly labeled training sets grows larger than the one computed on models trained on the original labels.

12.5 PAC-Bayes Bound on Randomly Labelled Data

Zhang et al. [2017] has shown that neural networks can fit randomly labelled training sets, obtaining arbitrarily bad generalization error. We expect a useful generalization bound to be consistent with this result and, therefore, be larger when computed on models trained on random labels. We have already observed in Supplementary 12.4 that the bound tends to grow faster while training on random datasets. Here, we perform a more extensive comparison on the two variations of the MNIST dataset. In Fig. 14, we compare the bound of different models trained on the original labels (circles) or on random ones (crosses). Training is performed on an augmented dataset, which means that each image is also transformed with a random element $g \in G$. As expected all models trained on the random labels have generalization error equals to 0.9 (like a random classifier). Fixing a model (same H, i.e. same color), we observe that the bound computed over models trained on random labels is always higher. On larger training sets, we obtain higher train errors; we think this is due to the limited size, and therefore, capacity of the model used.

We perform a similar experiment on the synthetic $G = \mathrm{SO}(2)$ dataset in Fig. 15. Again, we are able to fit a randomly labeled training set with equivariant models, given sufficiently large models. Since this is a binary classification task, the models trained on the random labels have $GE = 0.5$ generalization error.

Figure 11: Bound (B) from Theorem 9.1 vs Generalization Error (GE) of different H-equivariant models on different datasets. We always use $\mathbb{P}_S = \mathbb{P}_G$. In the synthetic ones, different colors represent different discrete symmetry groups G, where M is the number of rotations in $G = \mathrm{D}_M$ or $G = \mathrm{C}_M$; higher values of M define more complex datasets.

Figure 12: Alternative bound (B) from Supplementary 11 vs Generalization Error (GE) of different H-equivariant models on synthetic dataset with discrete D_M symmetries. As in the continuous symmetry cases, this bound does not capture the effect of H equivariance.

(a) $G = \mathrm{SO}(2)$ (b) $G = \mathrm{O}(2)$

Figure 13: Evolution of the PAC-Bayes Bound (B) during the training of H-equivariant models on the SO(2) and O(2) datasets with $F = 3$ and $m = 1024$ examples. Dashed lines represent the models trained on random labels.

(a) $H = C_N$ and $G = \mathrm{SO}(2)$

(b) $H = C_N$ and $G = \mathrm{O}(2)$ (c) $H = D_N$ and $G = \mathrm{O}(2)$

Figure 14: PAC-Bayes Bound (B) from Theorem 9.1 versus Generalization Error (GE) of different H-equivariant models on different G-MNIST datasets ($G = \mathrm{SO}(2)$ or $G = \mathrm{O}(2)$) when the models are trained with the real labels or with random labels. Crosses represent models trained on random labels, while circles represent models trained on the original labels. Note that for the same training setting, the bound computed on the models trained on random labels is always higher.

Figure 15: PAC-Bayes Bound (B) versus Generalization Error (GE) of different C$_N$-equivariant models on the synthetic continuous $G = \mathrm{SO}(2)$ datasets when trained with the real labels or with random labels. Crosses represent models trained on random labels, while circles represent models trained on the original labels. We used frequency $F = 3$ and $F = 4$ and $m = 1024$.