OFFLINE REINFORCEMENT LEARNING WITH DIFFERENTIABLE FUNCTION APPROXIMATION IS PROVABLY EFFICIENT

ABSTRACT

Offline reinforcement learning, which aims at optimizing sequential decision-making strategies with historical data, has been extensively applied in real-life applications. *State-Of-The-Art* algorithms usually leverage powerful function approximators (*e.g.* neural networks) to alleviate the sample complexity hurdle for better empirical performances. Despite all that, a more systematic understanding of the statistical complexity for function approximation remains lacking. Towards bridging the gap, we take a step by considering offline reinforcement learning with *differentiable function class approximation* (DFA). This function class naturally incorporates a wide range of models with nonlinear/nonconvex structures. Most importantly, we show offline RL with differentiable function approximation is provably efficient by analyzing the *pessimistic fitted Q-learning* (PFQL) algorithm, and our results provide the theoretical basis for understanding a variety of practical heuristics that rely on Fitted Q-Iteration style design. In addition, we further improve our guarantee with a tighter instance-dependent characterization. We hope our work could draw interest in studying reinforcement learning with differentiable function approximation beyond the scope of current research.

1 INTRODUCTION

Offline reinforcement learning (Lange et al., 2012; Levine et al., 2020) refers to the paradigm of learning a policy in the sequential decision making problems, where only the logged data are available and were collected from an unknown environment (*Markov Decision Process* / MDP). Inspired by the success of scalable supervised learning methods, modern reinforcement learning algorithms (*e.g.* Silver et al. (2017)) incorporate high-capacity function approximators to acquire generalization across large state-action spaces and have achieved excellent performances along a wide range of domains. For instance, there are a huge body of deep RL-based algorithms that tackle challenging problems such as the game of Go and chess (Silver et al., 2017; Schrittwieser et al., 2020), Robotics (Gu et al., 2017; Levine et al., 2018), energy control (Degrave et al., 2022) and Biology (Mahmud et al., 2018; Popova et al., 2018). Nevertheless, practitioners also noticed that algorithms with general function approximators can be quite data/sample inefficient, especially for deep neural networks where the models may require million of steps for tuning the large number of parameters they contain.[1]

On the other hand, statistical analysis has been actively conducted to understand the sample/statistical efficiency for reinforcement learning with function approximation, and fruitful results have been achieved under the respective model representations (Munos, 2003; Chen and Jiang, 2019; Yang and Wang, 2019; Du et al., 2019; Sun et al., 2019; Modi et al., 2020; Jin et al., 2020; Ayoub et al., 2020; Zanette et al., 2020; Jin et al., 2021a; Du et al., 2021; Jin et al., 2021b; Zhou et al., 2021; Xie et al., 2021; Min et al., 2021; Nguyen-Tang et al., 2021; Zanette et al., 2021; Yin et al., 2022; Uehara et al., 2022). However, most works consider *linear* model approximators (*e.g.* linear (mixture) MDPs) or its variants. While the explicit linear structures make the analysis trackable (linear problems are easier to analyze), they are unable to reveal the sample/statistical complexity behaviors of practical algorithms that apply powerful function approximations (which might have complex structures).

---

[1]Check Arulkumaran et al. (2017) and the references therein for an overview.

In addition, there is an excellent line of works tackling provably efficient offline RL with general function approximation (*e.g.* (Chen and Jiang, 2019; Xie et al., 2021; Zhan et al., 2022)). Due to the generic function approximation class considered, those complexity bounds are usually expressed in the standard worst-case fashion $O(V_{\max}^2 \sqrt{\frac{1}{n}})$ which lack the characterizations of individual instance behaviors. However, as mentioned in Zanette and Brunskill (2019), practical reinforcement learning algorithms often perform far better than what these problem-independent bounds would suggest.

These observations motivate us to consider function approximation schemes that can help address the existing limitations. In particular, in this work we consider offline reinforcement learning with *differentiable function class* approximations. Its definition is given in below.

Definition 1.1 (Parametric Differentiable Function Class). *Let $\mathcal{S}, \mathcal{A}$ be arbitrary state, action spaces and a feature map $\phi(\cdot, \cdot) : \mathcal{S} \times \mathcal{A} \to \Psi \subset \mathbb{R}^m$. The parameter space $\Theta \in \mathbb{R}^d$. Both Θ and Ψ are compact spaces. Then the parametric function class (for a model $f : \mathbb{R}^d \times \mathbb{R}^m \to \mathbb{R}$) is defined as*

$$\mathcal{F} := \{f(\theta, \phi(\cdot, \cdot)) : \mathcal{S} \times \mathcal{A} \to \mathbb{R}, \theta \in \Theta\}$$

that satisfies differentiability/smoothness condition: 1. for any $\phi \in \mathbb{R}^m$, $f(\theta, \phi)$ is third-time differentiable with respect to θ; 2. $f, \partial_\theta f, \partial_{\theta,\theta}^2 f, \partial_{\theta,\theta,\theta}^3 f$ are jointly continuous for (θ, ϕ).

Remark 1.2. *Differentiable Function Class was recently proposed for studying Off-Policy Evaluation (OPE) Problem (Zhang et al., 2022) and we adopt it here for the policy learning task. Note by the compactness of Θ, Ψ and continuity, there exists constants $C_\Theta, B_\mathcal{F}, \kappa_1, \kappa_2, \kappa_3 > 0$ that bounds: $\|\theta\|_2 \leq C_\Theta, |f(\theta, \phi(s,a))| \leq B_\mathcal{F}, \|\nabla_\theta f(\theta, \phi(s,a))\|_2 \leq \kappa_1, \|\nabla_{\theta\theta}^2 f(\theta, \phi(s,a))\|_2 \leq \kappa_2, \|\nabla_{\theta\theta\theta}^3 f(\theta, \phi(s,a))\|_2 \leq \kappa_3$ for all $\theta \in \Theta$, $s, a \in \mathcal{S} \times \mathcal{A}$.[2]*

Why consider differentiable function class (Definition 1.1)? There are two main reasons why differentiable function class is worth studying for reinforcement learning.

- Due to the limitation of statistical tools, existing analysis in reinforcement learning usually favor basic settings such as *tabular MDPs* (where the state space and action space are finite (Azar et al., 2017; Sidford et al., 2018; Agarwal et al., 2020)) or linear MDPs (Yang and Wang, 2020; Jin et al., 2020) / linear Mixture MDPs (Modi et al., 2020; Cai et al., 2020) (where the transition dynamic admits linear structures) so that well-established techniques (*e.g.* from linear regression) can be applied. In addition, subsequent extensions are often based on linear models (*e.g.* Linear Bellman Complete models (Zanette et al., 2020) and Eluder dimension (Russo and Van Roy, 2013)). Differentiable function class strictly generalizes over the previous popular choices, *i.e.* by choosing $f(\theta, \phi) = \langle \theta, \phi \rangle$ or specifying ϕ to be one-hot representations, and is far more expressive as it encompasses nonlinear approximators.

- Practically speaking, the flexibility of selecting model f provides the possibility for handling a variety of tasks. For instance, when f is specified to be neural networks, θ corresponds to the weights of each network layers and $\phi(\cdot, \cdot)$ corresponds to the state-action representations (which is induced by the network architecture). When facing with easier tasks, we can deploy simpler model f such as polynomials. Yet, our statistical guarantee is not affected by the specific choices as we can plug the concrete form of model f into Theorem 3.2 to obtain the respective bounds (we do not need separate analysis for different tasks).

1.1 RELATED WORKS

Reinforcement learning with function approximation. RL with function approximation has a long history that can date back to Bradtke and Barto (1996); Tsitsiklis and Van Roy (1996). Later, it draws significant interest for the finite sample analysis (Jin et al., 2020; Yang and Wang, 2019). Since then, people put tremendous efforts towards generalizing over linear function approximations and examples include Linear Bellman complete models (Zanette et al., 2020), Eluder dimension (Russo and Van Roy, 2013; Jin et al., 2021a), linear deterministic $Q^\star$ (Wen and Van Roy, 2013) or Bilinear class (Du et al., 2021). While those extensions are valuable, the structure conditions assumed usually make the classes hard to track beyond the linear case. For example, the practical instances of Eluder

---

[2]Here $\|\nabla_{\theta\theta\theta}^3 f(\theta, \phi(s,a))\|_2$ is defined as the 2-norm for 3-d tensor and in the finite horizon setting we simply instantiate $\mathcal{B}_\mathcal{F} = H$.

Dimension are based on the linear-in-feature (or its transformation) representations (Section 4.1 of Wen and Van Roy (2013)). As a comparison, differentiable function class contains a range of functions that are widely used in practical algorithms (Riedmiller, 2005).

Offline RL with general function approximation (GFA). Another interesting thread of work considered offline RL with general function approximation (Ernst et al., 2005; Chen and Jiang, 2019; Liu et al., 2020; Xie et al., 2021) which only imposes realizability and completeness/concentrability assumptions. The major benefit is that the function hypothesis can be arbitrary with no structural assumptions and it has been shown that offline RL with GFA is provably efficient. However, the generic form of functions in GFA makes it hard to go beyond worst-case analysis and obtain fine-grained instance-dependent learning bounds similar to those under linear cases. On the contrary, our results with DFA can be more problem adaptive by leveraging gradients and higher order information.

In addition to the above, there are more connected works. Zhang et al. (2022) first considers the differentiable function approximation (DFA) for the off-policy evaluation (OPE) task and builds the asymptotic theory, Fan et al. (2020) analyzes the *deep Q-learning* with the specific ReLu activations, and Kallus and Uehara (2020) considers semi-parametric / nonparametric methods for offline RL (as opposed to our parametric DFA in 1.1). These are nice complementary studies to our work.

1.2 OUR CONTRIBUTION

We provide the first Instance-dependent offline learning bound under non-linear function approximation. Informally, we show that (up to a lower order term) the natural complexity measure is proportional to $\sum_{h=1}^{H} \mathbb{E}_{\pi^\star,h}[\sqrt{g_\theta(s,a)^\top \Sigma_h^{-1} g_\theta(s,a)}]$ where $g_\theta(s,a) := \nabla f(\theta, \phi(s,a))$ is the gradient *w.r.t.* the parameter $\theta^\star$ at feature ϕ and $\Sigma_h = \sum_i g(s_{i,h}, a_{i,h}) g(s_{i,h}, a_{i,h})^\top$ is the Fisher information matrix of the observed data at $\widehat{\theta}$. This is achieved by analyzing the *pessimistic fitted Q-learning* (PFQL) algorithm (Theorem 3.2). In addition, we further analyze its variance-reweighting variant, which enjoys a more instance-dependent nature and can yield faster sample convergence rate. Last but not least, existing offline RL studies with tabular models, linear models and GLM models can be directly indicated by the choice of our model $\mathcal{F}$.

2 PRELIMINARIES

Episodic Markov decision process. Let $M = (\mathcal{S}, \mathcal{A}, P, r, H, d_1)$ to denote a finite-horizon *Markov Decision Process* (MDP), where $\mathcal{S}$ is the arbitrary state space and $\mathcal{A}$ is the arbitrary action space which can be infinite or continuous. The transition kernel $P_h : \mathcal{S} \times \mathcal{A} \mapsto \Delta^{\mathcal{S}}$ ($\Delta^{\mathcal{S}}$ represents a distribution over states) maps each state action (s_h, a_h) to a probability distribution $P_h(\cdot|s_h, a_h)$ and P_h can be different for different h (time-inhomogeneous). H is the planning horizon and d_1 is the initial state distribution. Besides, $r : \mathcal{S} \times \mathcal{A} \mapsto \mathbb{R}$ is the mean reward function satisfying $0 \le r \le 1$. A policy $\pi = (\pi_1, \ldots, \pi_H)$ assigns each state $s_h \in \mathcal{S}$ a probability distribution over actions by mapping $s_h \mapsto \pi_h(\cdot|s_h) \; \forall h \in [H]$ and induces a random trajectory $s_1, a_1, r_1, \ldots, s_H, a_H, r_H, s_{H+1}$ with $s_1 \sim d_1, a_h \sim \pi(\cdot|s_h), s_{h+1} \sim P_h(\cdot|s_h, a_h), \forall h \in [H]$.

Given a policy π, the V-value functions and state-action value function (Q-functions) $Q_h^\pi(\cdot, \cdot) \in \mathbb{R}^{S \times A}$ are defined as: $V_h^\pi(s) = \mathbb{E}_\pi[\sum_{t=h}^H r_t | s_h = s]$, $Q_h^\pi(s,a) = \mathbb{E}_\pi[\sum_{t=h}^H r_t | s_h, a_h = s, a]$, $\forall s, a, h \in \mathcal{S}, \mathcal{A}, [H]$. The Bellman (optimality) equations follow $\forall h \in [H], s, a \in \mathcal{S} \times \mathcal{A}$:

$$Q_h^\pi(s,a) = r_h(s,a) + \int_{\mathcal{S}} V_{h+1}^\pi(\cdot) dP_h(\cdot|s,a), \quad V_h^\pi(s) = \mathbb{E}_{a \sim \pi_h(s)}[Q_h^\pi(s,a)],$$

$$Q_h^\star(s,a) = r_h(s,a) + \int_{\mathcal{S}} V_{h+1}^\star(\cdot) dP_h(\cdot|s,a), \quad V_h^\star(s) = \max_a Q_h^\star(s,a).$$

We define Bellman operator $\mathcal{P}_h$ for any function $V \in \mathbb{R}^{\mathcal{S}}$ as $\mathcal{P}_h(V) = r_h + \int_{\mathcal{S}} V dP_h$, then $\mathcal{P}_h(V_{h+1}^\pi) = Q_h^\pi$ and $\mathcal{P}_h(V_{h+1}^\star) = Q_h^\star$. The performance measure is $v^\pi := \mathbb{E}_{d_1}[V_1^\pi] = \mathbb{E}_{\pi,d_1}\left[\sum_{t=1}^H r_t\right]$. Lastly, the induced state-action marginal occupancy measure for any $h \in [H]$ is defined to be: for any $E \subseteq \mathcal{S} \times \mathcal{A}$, $d_h^\pi(E) := \mathbb{E}[(s_h, a_h) \in E | s_1 \sim d_1, a_i \sim \pi(\cdot|s_i), s_i \sim P_{i-1}(\cdot|s_{i-1}, a_{i-1}), 1 \le i \le h]$ and $\mathbb{E}_{\pi,h}[f(s,a)] := \int_{\mathcal{S} \times \mathcal{A}} f(s,a) d_h^\pi(s,a) ds da$.

Offline Reinforcement Learning. The goal of Offline RL is to learn the policy $\pi^\star := \arg\max_\pi v^\pi$ using only the historical data $\mathcal{D} = \{(s_h^\tau, a_h^\tau, r_h^\tau, s_{h+1}^\tau)\}_{\tau \in [K]}^{h \in [H]}$. The data generating behavior policy is denoted as μ. In the offline regime, we have neither the knowledge about μ nor the access to further exploration for a different policy. The agent is asked to find a policy $\widehat{\pi}$ such that $v^\star - v^{\widehat{\pi}} \leq \epsilon$ for the given batch data $\mathcal{D}$ and a specified accuracy $\epsilon > 0$.

2.1 ASSUMPTIONS

Function approximation in offline RL requires sufficient expressiveness of $\mathcal{F}$. In fact, even under the *realizability* and *concentrability* conditions, sample efficient offline RL might not be achievable (Foster et al., 2021). Therefore, under the differentiable function setting (Definition 1.1), we make the following assumptions.

Assumption 2.1 (Realizability+Bellman Completeness). *The differentiable function class $\mathcal{F}$ in Definition 1.1 satisfies:*

- *Realizability: for optimal $Q_h^\star$, there exists $\theta_h^\star \in \Theta$ such that $Q_h^\star(\cdot, \cdot) = f(\theta_h^\star, \phi(\cdot)) \, \forall h$;*

- *Bellman Completeness: Let $\mathcal{G} := \{V(\cdot) \in \mathbb{R}^\mathcal{S} : \text{such that } \|V\|_\infty \leq H\}$. Then $\sup_{V \in \mathcal{G}} \inf_{f \in \mathcal{F}} \|f - \mathcal{P}_h(V)\|_\infty \leq \epsilon_\mathcal{F}$ for some $\epsilon_\mathcal{F} \geq 0$.*

Realizability and *Bellman Completeness* are widely adopted in the offline RL analysis with general function approximations (Chen and Jiang, 2019; Xie et al., 2021) and Assumption 2.1 states its differentiable function approximation version.

Data coverage assumption. Furthermore, in the offline regime, it is known that function approximation cannot be sample efficient for learning a ϵ-optimal policy without data-coverage assumptions when ϵ is small (*i.e.* high accuracy) (Wang et al., 2021a). In particular, we consider two types of coverage assumptions and provide guarantees for them separately.

Assumption 2.2 (Concentrability Coverage). *For any fixed policy π, define the marginal state-action occupancy ratio as $d_h^\pi(s, a)/d_h^\mu(s, a) \, \forall s, a$. Then the concentrability coefficient is defined as $C_{eff} := \sup_\pi \sup_{h \in [H]} \|d_h^\pi/d_h^\mu\|_{2, d_h^\mu}^2$, where $\|g(\cdot, \cdot)\|_{2, d^\mu} := \sqrt{\mathbb{E}_{d^\mu}[g(\cdot, \cdot)^2]}$ and $C_{eff} < \infty$.*

This is the standard coverage assumption that has has been widely assumed in (Ernst et al., 2005; Szepesvári and Munos, 2005; Chen and Jiang, 2019; Xie and Jiang, 2020a). In addition, we can also make the following stronger coverage assumption that requires the curvature of function class $\mathcal{F}$.

Assumption 2.3 (Uniform Coverage). *We have $\forall h \in [H]$, there exists $\kappa > 0$,*

- $\mathbb{E}_{\mu, h}\left[(f(\theta_1, \phi(\cdot, \cdot)) - f(\theta_2, \phi(\cdot, \cdot)))^2\right] \geq \kappa \|\theta_1 - \theta_2\|_2^2, \ \ \forall \theta_1, \theta_2 \in \Theta; \ (\star)$

- $\mathbb{E}_{\mu, h}\left[\nabla f(\theta, \phi(s, a)) \cdot \nabla f(\theta, \phi(s, a))^\top\right] \succ \kappa I, \forall \theta \in \Theta. \ (\star\star)$

In the linear function approximation regime, Assumption 2.3 reduces to 2.4 since $(\star)$ and $(\star\star)$ are identical assumptions. Therefore, 2.3 can be considered as a natural extension of 2.4 for differentiable class. Concretely, if $f(\theta, \phi) = \langle \theta, \phi \rangle$, then $(\star) \, \mathbb{E}_{\mu, h}[(f(\theta_1, \phi(\cdot, \cdot)) - f(\theta_2, \phi(\cdot, \cdot)))^2] = (\theta_1 - \theta_2)^\top \mathbb{E}_{\mu, h}[\phi(\cdot, \cdot)\phi(\cdot, \cdot)^\top](\theta_1 - \theta_2) \geq \kappa \|\theta_1 - \theta_2\|_2^2 \, \forall \theta_1, \theta_2 \in \Theta \Leftrightarrow 2.4 \Leftrightarrow (\star\star)\mathbb{E}_{\mu, h}\left[\nabla f(\theta, \phi(s, a)) \cdot \nabla f(\theta, \phi(s, a))^\top\right] \succ \kappa I$. We do point that 2.3 can be violated for function class $\mathcal{F}$ that is "not identifiable" by the data distribution μ (*i.e.*, there exists $f(\theta_1), f(\theta_2) \in \mathcal{F}, \theta_1 \neq \theta_2$ s.t. $\mathbb{E}_{\mu, h}[(f(\theta_1, \phi(\cdot, \cdot)) - f(\theta_2, \phi(\cdot, \cdot)))^2] = 0$). Nevertheless, there are representative non-linear differentiable classes (*e.g.* generalized linear model (GLM)) satisfying 2.3.

Example 2.4 (Linear function coverage assumption (Wang et al., 2021a; Min et al., 2021; Yin et al., 2022; Xiong et al., 2022)). $\Sigma_h^{\text{feature}} := \mathbb{E}_{\mu, h}\left[\phi(s, a)\phi(s, a)^\top\right] \succ \kappa I \, \forall h \in [H]$ with some $\kappa > 0$.

Example 2.5 (offline generalized linear model (Li et al., 2017; Wang et al., 2021b)). *For a known feature map $\phi : \mathcal{S} \times \mathcal{A} \mapsto \mathcal{B}_d$ and link function $f : [-1, 1] \mapsto [-1, 1]$ the class of GLM is $\mathcal{F}_{\text{GLM}} := \{(s, a) \mapsto f(\langle \phi(s, a), \theta \rangle) : \theta \in \Theta\}$ satisfying $\mathbb{E}_{\mu, h}\left[\phi(s, a)\phi(s, a)^\top\right] \succ \kappa I$. Furthermore, $f(\cdot)$ is either monotonically increasing or decreasing and $0 < \kappa \leq |f'(z)| \leq K < \infty, |f''(z)| \leq M < \infty$ for all $|z| \leq 1$ and some κ, K, M. Then $\mathcal{F}_{\text{GLM}}$ satisfies 2.3, see Appendix B.*

3   DIFFERENTIABLE FUNCTION APPROXIMATION IS PROVABLY EFFICIENT

In this section, we present our solution for offline reinforcement learning with differentiable function approximation. As a warm-up, we first analyze the *vanilla fitted Q-learning* (VFQL, Algorithm 2), which only requires the concentrability Assumption 2.2. The algorithm is presented in Appendix I.

Theorem 3.1. *Choose* $0 < \lambda \leq 1/2C_\Theta^2$ *in Algorithm 2. Suppose Assumption 2.1,2.2. Then if* $K \geq \max\left\{512\frac{\kappa_1^4}{\kappa^2}\left(\log(\frac{2Hd}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K^3}{\lambda^2})\right), \frac{4\lambda}{\kappa}\right\}$, *with probability* $1 - \delta$, *the output* $\widehat{\pi}$ *of VFQL guarantees*

$$v^\star - v^{\widehat{\pi}} \leq \sqrt{C_{\text{eff}}}H \cdot \widetilde{O}\left(\sqrt{\frac{H^2 d + \lambda C_\Theta^2}{K}} + \sqrt[4]{\frac{H^3 d\epsilon_{\mathcal{F}}}{K}}\right) + O(\sqrt{C_{\text{eff}}H^3\epsilon_{\mathcal{F}}} + H\epsilon_{\mathcal{F}})$$

If the model approximation capacity is insufficient, 3.1 will induce extra error due to the large $\epsilon_{\mathcal{F}}$. If $\epsilon_{\mathcal{F}} \to 0$, the standard statistical rate $\frac{1}{\sqrt{K}}$ can be recovered and similar results are derived with general function approximation (GFA) (Chen and Jiang, 2019; Xie and Jiang, 2020a). However, using concentrability coefficient conceals the problem-dependent structure and omits the specific information of differentiable functions in the complexity measure. Owing to this, we switch to the stronger "uniform" coverage 2.3 and analyze the *pessimistic fitted Q-learning* (PFQL, Algorithm 1) to arrive at the conclusion that offline RL with differentiable function approximation is provably efficient.

Motivation of PFQL. The PFQL algorithm mingles the two celebrated algorithmic choices: Fitted Q-Iteration (FQI) and Pessimism. However, before going into the technical details, we provide some interesting insights that motivate our analysis.

First of all, the square error loss used in FQI (Gordon, 1999; Ernst et al., 2005) naturally couples with the differentiable function class as the resulting optimization objective is more computationally tractable (since *stochastic gradient descent* (SGD) can be readily applied) comparing to other information-theoretical algorithms derived with general function approximation (*e.g.* the *maxmin* objective in Xie et al. (2021), eqn (3.2)).[3] In particular, FQI with differentiable function approximation resembles the theoretical prototype of neural FQI algorithm (Riedmiller, 2005) and DQN algorithm (Mnih et al., 2015; Fan et al., 2020) when instantiating the model f to be deep neural networks. Furthermore, plenty of practical algorithms leverage fitted-Q subroutines for updating the *critic* step (*e.g.* (Schulman et al., 2017; Haarnoja et al., 2018)) with different differentiable function choices.

In addition, we also incorporate pessimism for the design. Indeed, one of the fundamental challenges in offline RL comes from the *distributional shift*. When such a mismatch occurs, the estimated/optimized Q-function (using batch data $\mathcal{D}$) may witness severe overestimation error due to the extrapolation of model f (Levine et al., 2020). Pessimism is the scheme to mitigate the error / overestimation bias via penalizing the Q-functions at state-action locations that have high uncertainties (as opposed to the *optimism* used in the online case), and has been widely adopted (*e.g.* (Buckman et al., 2020; Kidambi et al., 2020; Jin et al., 2021b)).

Algorithm 1 description.   Inside the backward iteration of PFQL, Fitted Q-update is performed to optimize the parameter (Line 4). $\widehat{\theta}_h$ is the root of the first-order stationarity equation $\sum_{k=1}^{K}\left(f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)\right) \cdot \nabla_\theta^\top f(\theta, \phi_{h,k}) + \lambda\theta = 0$ and Σ_h is the Gram matrix with respect to $\nabla_\theta f|_{\theta=\widehat{\theta}_h}$. Note for any $s, a \in \mathcal{S} \times \mathcal{A}$, $m(s,a) := (\nabla_\theta f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(s,a)))^{-1}$ measures the effective sample size that explored s, a along the gradient direction $\nabla_\theta f|_{\theta=\widehat{\theta}_h}$, and $\beta/\sqrt{m(s,a)}$ is the estimated uncertainty at (s,a). However, the quantity $m(s,a)$ depends on $\widehat{\theta}_h$, and $\widehat{\theta}_h$ needs to be close to the true $\theta_h^\star$ (*i.e.* $\widehat{Q}_h \approx f(\widehat{\theta}_h, \phi)$ needs to be close to $Q_h^\star$) for the uncertainty estimation Γ_h to be valid, since putting a random θ into $m(s,a)$ can cause an arbitrary Γ_h that is useless (or might even deteriorate the algorithm). Such an "implicit" constraint over $\widehat{\theta}_h$ imposes the extra difficulty for the theoretical analysis due to that general differentiable functions encode nonlinear structures. As a direct comparison, in the simpler

---

[3]We mention Xie et al. (2021) has a nice practical version PSPI, but the convergence is slower (the rate $O(n^{-\frac{1}{3}})$).

linear MDP case, the uncertainty measure $\Gamma_h := \sqrt{\phi(\cdot,\cdot)^\top (\Sigma_h^{\text{linear}})^{-1} \phi(\cdot,\cdot)}$ is always valid since it does not depend on the least-square regression weight $\widehat{w}_h$ (Jin et al., 2021b).[4] Besides, the choice of β is set to be $\widetilde{O}(dH)$ in Theorem 3.2 and the extra higher order term $\widetilde{O}(\frac{1}{K})$ in Γ_h is for theoretical reason only.

Algorithm 1 Pessimistic Fitted Q-Learning (PFQL)

1: **Input:** Offline Dataset $\mathcal{D} = \left\{ \left(s_h^k, a_h^k, r_h^k, s_{h+1}^k \right) \right\}_{k,h=1}^{K,H}$. Require β. Denote $\phi_{h,k} := \phi(s_h^k, a_h^k)$.
2: **Initialization:** Set $\widehat{V}_{H+1}(\cdot) \leftarrow 0$ and $\lambda > 0$.
3: **for** $h = H, H-1, \ldots, 1$ **do**
4:    Set $\widehat{\theta}_h \leftarrow \arg\min_{\theta \in \Theta} \left\{ \sum_{k=1}^K \left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right]^2 + \lambda \cdot \|\theta\|_2^2 \right\}$
5:    Set $\Sigma_h \leftarrow \sum_{k=1}^K \nabla_\theta f(\widehat{\theta}_h, \phi_{h,k}) \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) + \lambda I_d$.
6:    Set $\Gamma_h(\cdot,\cdot) \leftarrow \beta \sqrt{\nabla_\theta f(\widehat{\theta}_h, \phi(\cdot,\cdot))^\top \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(\cdot,\cdot))} \left(+\widetilde{O}(\frac{1}{K}) \right)$
7:    Set $\bar{Q}_h(\cdot,\cdot) \leftarrow f(\widehat{\theta}_h, \phi(\cdot,\cdot)) - \Gamma_h(\cdot,\cdot)$
8:    Set $\widehat{Q}_h(\cdot,\cdot) \leftarrow \min \left\{ \bar{Q}_h(\cdot,\cdot), H - h + 1 \right\}^+$
9:    Set $\widehat{\pi}_h(\cdot \mid \cdot) \leftarrow \arg\max_{\pi_h} \left\langle \widehat{Q}_h(\cdot,\cdot), \pi_h(\cdot \mid \cdot) \right\rangle_{\mathcal{A}}, \widehat{V}_h(\cdot) \leftarrow \max_{\pi_h} \left\langle \widehat{Q}_h(\cdot,\cdot), \pi_h(\cdot \mid \cdot) \right\rangle_{\mathcal{A}}$
10: **end for**
11: **Output:** $\left\{ \widehat{\pi}_h \right\}_{h=1}^H$.

Model-Based vs. Model-Free. PFQL can be viewed as the strict generalization over the previous value iteration based algorithms, *e.g.* PEVI algorithm (Jin et al. (2021b), linear MDPs) and the VPVI algorithm (Yin and Wang (2021), tabular MDPs). On one hand, *approximate value iteration* (AVI) algorithms (Munos, 2005) are usually model-based algorithms (for instance the tabular algorithm VPVI uses empirical model $\widehat{P}$ for planning). On the other hand, FQI has the form of batch Q-learning update (*i.e.* Q-learning is a special case with batch size equals to one), therefore is more of model-free flavor. Since FQI is a concrete instantiation of the abstract AVI procedure (Munos, 2007), PFQL draws a unified view of model-based and model-free learning.

Now we are ready to state our main result for PFQL and the full proof can be found in Appendix D,E,F.

Theorem 3.2. *Let $\beta = 8dH\iota$ and choose $0 < \lambda \le 1/2C_\Theta^2$ in Algorithm 1. Suppose Assumption 2.1,2.3 with $\epsilon_{\mathcal{F}} = 0$.[5] Then if $K \ge \max\left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2Hd}{\delta}) + d\log(1 + \frac{4\kappa_1^3 \kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$, with probability $1 - \delta$, for all policy π simultaneously, the output of PFQL guarantees*

$$v^\pi - v^{\widehat{\pi}} \le \sum_{h=1}^H 8dH \cdot \mathbb{E}_\pi \left[\sqrt{\nabla_\theta^\top f(\widehat{\theta}_h, \phi(s_h, a_h)) \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\text{hot}}}{K}),$$

where ι is a Polylog term and the expectation of π is taken over s_h, a_h. In particular, if further $K \ge \max\{\widetilde{O}(\frac{(\kappa_1^2 + \lambda)^2 \kappa_2^2 \kappa_1^2 H^4 d^2}{\kappa^6}), \frac{128\kappa_1^4 \log(2d/\delta)}{\kappa^2}\}$ we have

$$0 \le v^{\pi^\star} - v^{\widehat{\pi}} \le \sum_{h=1}^H 16dH \cdot \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi(s_h, a_h)) \Sigma_h^{\star -1} \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C'_{\text{hot}}}{K}).$$

Here $\Sigma_h^\star = \sum_{k=1}^K \nabla_\theta f(\theta_h^\star, \phi(s_h^k, a_h^k)) \nabla_\theta^\top f(\theta_h^\star, \phi(s_h^k, a_h^k)) + \lambda I_d$ and the definition of higher order parameter $C_{\text{hot}}, C'_{\text{hot}}$ can be found in List A.

Corollary 3.3 (Offline Generalized Linear Models (GLM)). *Consider the GLM function class defined in 2.5. Suppose β, λ, K are defined the same as Theorem 3.2. $\epsilon_{\mathcal{F}} = 0$. Then with probability $1 - \delta$, for all policy π simultaneously, PFQL guarantees*

$$v^\pi - v^{\widehat{\pi}} \le \sum_{h=1}^H 8dH \cdot \mathbb{E}_\pi \left[\sqrt{f'(\langle \widehat{\theta}_h, \phi(s_h, a_h) \rangle)^2 \cdot \phi^\top(s_h, a_h) \Sigma_h^{-1} \phi(s_h, a_h)} \right] \cdot \iota + \widetilde{O}(\frac{C_{\text{hot}}}{K}).$$

---

[4]Here $\Sigma_h^{\text{linear}} := \sum_{k=1}^K \phi_{h,k} \phi_{h,k}^\top + \lambda I_d$.

[5]Here we assume model capacity is sufficient to make the presentation concise. If $\epsilon_{\mathcal{F}} > 0$, the complexity bound will include the term $\epsilon_{\mathcal{F}}$. We include more discussion in Appendix H.

PFQL is provably efficient. Theorem 3.2 verifies PFQL is statistically efficient. In particular, by Lemma L.5 we have $\|\nabla_\theta f(\theta_h^\star, \phi)\|_{\Sigma_h^{-1}} \lesssim \frac{2\kappa_1}{\sqrt{\kappa K}}$, resulting the main term to be bounded by $\frac{32 d H^2 \kappa_1}{\sqrt{\kappa K}}$ that recovers the standard statistical learning convergence rate $\frac{1}{\sqrt{K}}$.

Comparing to Jin et al. (2021b). Theorem 3.2 strictly subsumes the linear MDP learning bound in Jin et al. (2021b). In fact, in the linear case $\nabla_\theta f(\theta, \phi) = \nabla_\theta \langle \theta, \phi \rangle = \phi$ and 3.2 reduces to $O(dH \sum_{h=1}^H \mathbb{E}_{\pi^\star}[\sqrt{\phi(s_h, a_h)^\top (\Sigma_h^{\text{linear}})^{-1} \phi(s_h, a_h)}])$.

Faster convergence in learning. Previous studies for offline RL with general function approximation (GFA) (Chen and Jiang, 2019; Xie and Jiang, 2020b) are more of worst-case flavors as they usually rely on the *concentrability* coefficient C. The resulting learning bounds are expressed in the form[6] $O(V_{\max}^2 \sqrt{\frac{C}{n}})$ that is unable to depict the behavior of individual instances. In contrast, the guarantee with differentiable function approximation is more adaptive due to the instance-dependent structure $\sum_{h=1}^H \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi) \Sigma_h^{\star-1} \nabla_\theta f(\theta_h^\star, \phi)} \right]$. For example, when the instance satisfies $\theta^\star$ to be the local extreme of the class $\mathcal{F}$ (*i.e.* $\nabla f(\theta^\star, \phi) = 0$), the convergence rate boosts to $O(\frac{1}{K})$ as the main term in 3.2 goes to zero. To our knowledge, existing results with GFA does not enjoy this feature.

Feature representation vs. Parameters. One interesting observation from Theorem 3.2 is that the learning complexity does not depend on the feature representation dimension m but only on parameter dimension d as long as function class $\mathcal{F}$ satisfies differentiability definition 1.1 (not even in the higher order term). This seems to suggest, when changing the model f with more complex representations, the learning hardness will not grow as long as the number of parameters need to be learned does not increase. Note in the linear MDP analysis this phenomenon is not captured since the two dimensions are coupled ($d = m$). Therefore, this heuristic might help people rethink about what is the more essential element (feature representation vs. parameter space) in the *representation learning RL* regime (*e.g.* low rank MDPs (Uehara et al., 2022)).

Technical challenges with differentiable function approximation (DFA). Informally, one key step for the analysis is to bound $|f(\widehat{\theta}_h, \phi) - f(\theta_h^\star, \phi)|$. This can be estimated by the first order approximation $\nabla f(\widehat{\theta}_h, \phi)^\top \cdot (\widehat{\theta}_h - \theta_h^\star)$. However, different from the least-square value iteration (LSVI) objective (Jin et al., 2020; 2021b), the *fitted Q-update* (Line 4, Algorithm 1) no longer admits a closed-form solution for $\widehat{\theta}_h$. Instead, we can only leverage $\widehat{\theta}_h$ is a stationary point of $Z_h(\theta) := \sum_{k=1}^K \left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right] \nabla f(\theta, \phi_{h,k}) + \lambda \cdot \theta$ (since $Z_h(\widehat{\theta}_h) = 0$). To measure the difference $\widehat{\theta}_h - \theta_h^\star$, for any $\theta \in \Theta$, we do the *Vector Taylor expansion* $Z_h(\theta) - Z_h(\widehat{\theta}_h) = \Sigma_h^s(\theta - \widehat{\theta}_h) + R_K(\theta)$ (where $R_K(\theta)$ is the higher-order residuals) at the point $\widehat{\theta}_h$ with

$$\Sigma_h^s := \frac{\partial}{\partial \theta} Z_h(\theta) \Big|_{\theta = \widehat{\theta}_h} = \frac{\partial}{\partial \theta} \left(\sum_{k=1}^K \left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right] \nabla f(\theta, \phi_{h,k}) + \lambda \cdot \theta \right)_{\theta = \widehat{\theta}_h}$$

$$= \underbrace{\sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\widehat{\theta}_h, \phi_{h,k})}_{:= \Delta_{\Sigma_h^s}} + \underbrace{\sum_{k=1}^K \nabla_\theta f(\widehat{\theta}_h, \phi_{h,k}) \nabla_\theta^\top f(\widehat{\theta}_{h,k}, \phi_{h,k}) + \lambda I_d}_{:= \Sigma_h}.$$

$$(1)$$

The perturbation term $\Delta_{\Sigma_h^s}$ encodes one key challenge for solving $\widehat{\theta}_h - \theta_h^\star$ since it breaks the positive definiteness of Σ_h^s, and, as a result, we cannot invert the Σ_h^s in the Taylor expansion of Z_h. This is due to DFA (Definition 1.1) is a rich class that incorporates *nonlinear* curvatures. In the linear function approximation regime, this hurdle will not show up since $\nabla_{\theta\theta}^2 f \equiv 0$ and $\Delta_{\Sigma_h^s}$ is always invertible as long as $\lambda > 0$. Moreover, for the *off-policy evaluation* (OPE) task, one can overcome this issue by expanding over the population counterpart of Z_h at underlying true parameter of the given behavior target policy (Zhang et al., 2022).[7] However, for the policy learning task, we cannot use

---

[6]Here n is the number of samples used in the infinite horizon discounted setting and is similar to K in the episodic setting.

[7]*i.e.* expanding over $Z_h^p(\theta) := \mathbb{E}_{s,a,s'}[(f(\theta, \phi(s,a)) - r - V_{h+1}^\pi(s')) \nabla f(\theta, \phi(s,a))]$, and the corresponding $\Delta_{\Sigma_h^s}$ in $\frac{\partial}{\partial \theta} Z_h(\theta)|_{\theta = \theta_h^\pi}$ is zero by Bellman equation.

either population quantity or the true parameter $\theta_h^\star$ since we need a computable/data-based pessimism Γ_h to make the algorithm practical. Check the following section for more discussions of the analysis.

3.1 SKETCH OF THE PFQL ANALYSIS

Due to the space constraint, here we only overview the key components of the analysis. To begin with, by following the result of general MDP in Jin et al. (2021b), the suboptimality gap can be bounded by (Appendix D) $\sum_{h=1}^H 2\mathbb{E}_\pi\left[\Gamma_h(s_h, a_h)\right]$ if $|(\mathcal{P}_h \widehat{V}_{h+1} - f(\widehat{\theta}_h, \phi))(s, a)| \le \Gamma_h(s, a)$. To deal with $\mathcal{P}_h \widehat{V}_{h+1}$, by Assumption 2.1 we can leverage the *parameter Bellman operator* $\mathbb{T}$ (Definition D.1) so that $\mathcal{P}_h \widehat{V}_{h+1} = f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi)$. Next, we apply the second-order approximation to obtain $\mathcal{P}_h \widehat{V}_{h+1} - f(\widehat{\theta}_h, \phi) \approx \nabla f(\widehat{\theta}_h, \phi)^\top (\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) + \frac{1}{2}(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h)^\top \nabla_{\theta\theta}^2 f(\widehat{\theta}_h, \phi)(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h)$. Later, we use (1) to represent

$$Z_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - Z_h(\widehat{\theta}_h) = \Sigma_h^s(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) = \Sigma_h(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) + \widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})$$

by denoting $\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) = \Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})$. Now that Σ_h^{-1} is invertible thus provides the estimation (note $Z_h(\widehat{\theta}_h) = 0$)

$$\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h = \Sigma_h^{-1} \cdot Z_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \Sigma_h^{-1}\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}).$$

However, to handle the higher order terms, we need the explicit finite sample bound for $\|\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\|_2$ (or $\|\theta_h^\star - \widehat{\theta}_h\|_2$). In the OPE literature, Zhang et al. (2022) uses *asymptotic theory* (Prohorov's Theorem) to show the existence of $B(\delta)$ such that $\|\widehat{\theta}_h - \theta_h^\star\| \le \frac{B(\delta)}{\sqrt{K}}$. However, this is insufficient for *finite sample/non-asymptotic* guarantees since the abstraction of $B(\delta)$ might prevent the result from being sample efficient. For example, if $B(\delta)$ has the form $e^H \log(\frac{1}{\delta})$, then $\frac{e^H \log(\frac{1}{\delta})}{\sqrt{K}}$ is an inefficient bound since K needs to be e^H/ϵ^2 large to guarantee ϵ accuracy.

To address this technicality, we use a novel reduction to *general function approximation* (GFA) learning proposed in Chen and Jiang (2019). Concretely, we first bound the loss objective $\mathbb{E}_\mu[\ell_h(\widehat{\theta}_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})]$ via a "orthogonal" decomposition and by solving a quadratic equation. The resulting bound can be directly used to further bound $\|\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\|_2$ for obtaining efficient guarantee $\widetilde{O}(\frac{dH}{\sqrt{\kappa K}})$. During the course, the covering technique is applied to extend the finite function hypothesis in Chen and Jiang (2019) to all the differentiable functions in Definition 1.1. See Appendix G for the complete proofs. The full proof can be found in Appendix D,E,F.

4 IMPROVED LEARNING VIA VARIANCE AWARENESS

In addition to knowing the provable efficiency for differentiable function approximation (DFA), it is of great interest to understand what is the statistical limit with DFA, or equivalently, what is the "optimal" sample/statistical complexity can be achieved in DFA (measured by minimaxity criteria)? Towards this goal, we further incorporate *variance awareness* to improve our learning guarantee. Variance awareness is first designed for linear Mixture MDPs (Talebi and Maillard, 2018; Zhou et al., 2021) to achieve the near-minimax sample complexity and it uses estimated conditional variances $\mathrm{Var}_{P(\cdot|s,a)}(V_{h+1}^\star)$ to reweight each training sample in the LSVI objective.[8] Later, such a technique is leveraged by Min et al. (2021); Yin et al. (2022) to obtained the instance-dependent results. Intuitively, conditional variances $\sigma^2(s, a) := \mathrm{Var}_{P(\cdot|s,a)}(V_{h+1}^\star)$ serves as the uncertainty measure of the sample (s, a, r, s') that comes from the distribution $P(\cdot|s, a)$. If $\sigma^2(s, a)$ is large, then the distribution $P(\cdot|s, a)$ has high variance and we should put less weights in a single sample (s, a, r, s') rather than weighting all the samples equally. In the differentiable function approximation regime, the update is modified to

$$\widehat{\theta}_h \leftarrow \underset{\theta \in \Theta}{\arg\min} \left\{ \sum_{k=1}^K \frac{\left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)\right]^2}{\sigma_h^2(s_h^k, a_h^k)} + \lambda \cdot \|\theta\|_2^2 \right\}$$

---

[8]We mention Zhang et al. (2021) uses variance-aware confidence sets in a slightly different way.

with $\sigma_h^2(\cdot, \cdot)$ estimated by the offline data. Notably, empirical algorithms have also shown uncertainty reweighting can improve the performances for both online RL (Mai et al., 2022) and offline RL (Wu et al., 2021). These motivates our *variance-aware fitted Q-learning* (VAFQL) algorithm 3.

Theorem 4.1. *Suppose Assumption 2.1,2.3 with $\epsilon_{\mathcal{F}} = 0$. Let $\beta = 8d\iota$ and choose $0 < \lambda \leq 1/2C_\Theta^2$ in Algorithm 3. Then if $K \geq K_0$ and $\sqrt{d} \geq \widetilde{O}(\zeta)$, with probability $1 - \delta$, for all policy π simultaneously, the output of VAFQL guarantees*

$$v^\pi - v^{\widehat{\pi}} \leq \sum_{h=1}^{H} 8d \cdot \mathbb{E}_\pi \left[\sqrt{\nabla_\theta^\top f(\widehat{\theta}_h, \phi(s_h, a_h)) \Lambda_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{\bar{C}_{\text{hot}}}{K}),$$

where ι is a Polylog term and the expectation of π is taken over s_h, a_h. In particular, we have

$$0 \leq v^{\pi^\star} - v^{\widehat{\pi}} \leq 8d \cdot \sum_{h=1}^{H} \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi(s_h, a_h)) \Lambda_h^{\star-1} \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{\bar{C}_{\text{hot}}'}{K}).$$

Here $\Lambda_h^\star = \sum_{k=1}^{K} \nabla_\theta f(\theta_h^\star, \phi_{h,k}) \nabla_\theta^\top f(\theta_h^\star, \phi_{h,k})/\sigma_h^\star(s_h^k, a_h^k)^2 + \lambda I_d$ and the $\sigma_h^\star(\cdot, \cdot)^2 := \max\{1, \text{Var}_{P_h} V_{h+1}^\star(\cdot, \cdot)\}$. The definition of $K_0, \bar{C}_{\text{hot}}, \bar{C}_{\text{hot}}', \zeta$ can be found in List A.

In particular, to bound the error for $\boldsymbol{u}_h, \boldsymbol{v}_h$ and $\widehat{\sigma}_h^2$, we need to define an operator $\mathbb{J}$ that is similar to the *parameter Bellman operator* D.1. The Full proof of Theorem 4.1 can be found in Appendix J. Comparing to Theorem 3.2, VAFQL enjoys a net improvement of the horizon dependence since $\text{Var}_P(V_h^\star) \leq H^2$. Moreover, VAFQL provides better instance-dependent characterizations as the main term is fully depicted by the system quantities except the feature dimension d. For instance, when the system is fully deterministic (transition P_h's are deterministic), $\sigma_h^\star \approx \text{Var}_{P_h} V_{h+1}^\star(\cdot, \cdot) \equiv 0$ (if ignore the truncation) and $\Lambda^{\star-1} \to 0$. This yields a faster convergence with rate $O(\frac{1}{K})$. Lastly, when reduced to linear MDPs, 4.1 recovers the results of Yin et al. (2022) except an extra factor of $\sqrt{d}$.

On the statistical limits. To complement the study, we incorporate a minimax lower bound via a reduction to Zanette et al. (2021); Yin et al. (2022). The following theorem reveals we cannot improve over Theorem 4.1 by more than a factor of $\sqrt{d}$ in the most general cases. The full discussion is deterred to Appendix K.

Theorem 4.2 (Minimax lower bound). *Specifying the model to have linear representation $f = \langle \theta, \phi \rangle$. There exist a pair of universal constants $c, c' > 0$ such that given dimension d, horizon H and sample size $K > c'd^3$, one can always find a family of MDP instances such that for any algorithm $\widehat{\pi}$*

$$\inf_{\widehat{\pi}} \sup_{M \in \mathcal{M}} \mathbb{E}_M \left[v^\star - v^{\widehat{\pi}} \right] \geq c\sqrt{d} \cdot \sum_{h=1}^{H} \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi(\cdot, \cdot))(\Lambda_h^{\star,p})^{-1} \nabla_\theta f(\theta_h^\star, \phi(\cdot, \cdot))} \right], \quad (2)$$

where $\Lambda_h^{\star,p} = \mathbb{E} \left[\sum_{k=1}^{K} \frac{\nabla_\theta f(\theta_h^\star, \phi(s_h^k, a_h^k)) \cdot \nabla_\theta f(\theta_h^\star, \phi(s_h^k, a_h^k))^\top}{\text{Var}_h(V_{h+1}^\star)(s_h^k, a_h^k)} \right].$

5 CONCLUSION, LIMITATION AND FUTURE DIRECTIONS

In this work, we study offline reinforcement learning with differentiable function approximation and show the sample efficiency of differentiable function learning. We further improve the sample complexity with respect to the horizon dependence via a variance aware variant. However, the dependence of the parameter space still scales with d (whereas for linear function approximation this dependence is $\sqrt{d}$), and this is due to applying covering argument for the rich class of differentiable functions. For large deep models, the dimension of the parameter can be huge, therefore it would be interesting to know if certain algorithms can further improve the parameter dependence, or whether this d is essential. Also, how to relax uniform coverage assumption 2.3 is unknown under the current analysis. We leave these open questions as future work.

Lastly, the differentiable function approximation setting provides a general framework that is not confined to offline RL. Understanding the sample complexity behaviors of online reinforcement learning (Jin et al., 2020), reward-free learning and representation learning (Uehara et al., 2022) might provide new and unified views over the existing studies.

REFERENCES

Yasin Abbasi-Yadkori, Dávid Pál, and Csaba Szepesvári. Improved algorithms for linear stochastic bandits. In *Advances in Neural Information Processing Systems*, pages 2312–2320, 2011.

Alekh Agarwal, Sham Kakade, and Lin F Yang. Model-based reinforcement learning with a generative model is minimax optimal. In *Conference on Learning Theory*, pages 67–83, 2020.

Kai Arulkumaran, Marc Peter Deisenroth, Miles Brundage, and Anil Anthony Bharath. Deep reinforcement learning: A brief survey. *IEEE Signal Processing Magazine*, 34(6):26–38, 2017.

Alex Ayoub, Zeyu Jia, Csaba Szepesvari, Mengdi Wang, and Lin Yang. Model-based reinforcement learning with value-targeted regression. In *International Conference on Machine Learning*, pages 463–474. PMLR, 2020.

Mohammad Gheshlaghi Azar, Ian Osband, and Rémi Munos. Minimax regret bounds for reinforcement learning. In *Proceedings of the 34th International Conference on Machine Learning-Volume 70*, pages 263–272. JMLR. org, 2017.

Steven J Bradtke and Andrew G Barto. Linear least-squares algorithms for temporal difference learning. *Machine learning*, 22(1):33–57, 1996.

Jacob Buckman, Carles Gelada, and Marc G Bellemare. The importance of pessimism in fixed-dataset policy optimization. *arXiv preprint arXiv:2009.06799*, 2020.

Qi Cai, Zhuoran Yang, Chi Jin, and Zhaoran Wang. Provably efficient exploration in policy optimization. In *International Conference on Machine Learning*, pages 1283–1294. PMLR, 2020.

Jinglin Chen and Nan Jiang. Information-theoretic considerations in batch reinforcement learning. In *International Conference on Machine Learning*, pages 1042–1051, 2019.

Jonas Degrave, Federico Felici, Jonas Buchli, Michael Neunert, Brendan Tracey, Francesco Carpanese, Timo Ewalds, Roland Hafner, Abbas Abdolmaleki, Diego de Las Casas, et al. Magnetic control of tokamak plasmas through deep reinforcement learning. *Nature*, 602(7897):414–419, 2022.

Simon Du, Akshay Krishnamurthy, Nan Jiang, Alekh Agarwal, Miroslav Dudik, and John Langford. Provably efficient rl with rich observations via latent state decoding. In *International Conference on Machine Learning*, pages 1665–1674. PMLR, 2019.

Simon S Du, Sham M Kakade, Jason D Lee, Shachar Lovett, Gaurav Mahajan, Wen Sun, and Ruosong Wang. Bilinear classes: A structural framework for provable generalization in rl. *International Conference on Machine Learning*, 2021.

Damien Ernst, Pierre Geurts, and Louis Wehenkel. Tree-based batch mode reinforcement learning. *Journal of Machine Learning Research*, 6:503–556, 2005.

Jianqing Fan, Zhaoran Wang, Yuchen Xie, and Zhuoran Yang. A theoretical analysis of deep q-learning. In *Learning for Dynamics and Control*, pages 486–489. PMLR, 2020.

Dylan J Foster, Akshay Krishnamurthy, David Simchi-Levi, and Yunzong Xu. Offline reinforcement learning: Fundamental barriers for value function approximation. *arXiv preprint arXiv:2111.10919*, 2021.

Geoffrey J Gordon. *Approximate solutions to Markov decision processes*. Carnegie Mellon University, 1999.

Shixiang Gu, Ethan Holly, Timothy Lillicrap, and Sergey Levine. Deep reinforcement learning for robotic manipulation with asynchronous off-policy updates. In *2017 IEEE international conference on robotics and automation (ICRA)*, pages 3389–3396. IEEE, 2017.

Tuomas Haarnoja, Aurick Zhou, Pieter Abbeel, and Sergey Levine. Soft actor-critic: Off-policy maximum entropy deep reinforcement learning with a stochastic actor. In *International conference on machine learning*, pages 1861–1870. PMLR, 2018.

Chi Jin, Zhuoran Yang, Zhaoran Wang, and Michael I Jordan. Provably efficient reinforcement learning with linear function approximation. In *Conference on Learning Theory*, pages 2137–2143. PMLR, 2020.

Chi Jin, Qinghua Liu, and Sobhan Miryoosefi. Bellman eluder dimension: New rich classes of rl problems, and sample-efficient algorithms. *arXiv preprint arXiv:2102.00815*, 2021a.

Ying Jin, Zhuoran Yang, and Zhaoran Wang. Is pessimism provably efficient for offline rl? In *International Conference on Machine Learning*, pages 5084–5096. PMLR, 2021b.

Nathan Kallus and Masatoshi Uehara. Double reinforcement learning for efficient off-policy evaluation in markov decision processes. *J. Mach. Learn. Res.*, 21(167):1–63, 2020.

Rahul Kidambi, Aravind Rajeswaran, Praneeth Netrapalli, and Thorsten Joachims. Morel: Model-based offline reinforcement learning. *Advances in Neural Information Processing Systems*, 2020.

Sascha Lange, Thomas Gabel, and Martin Riedmiller. Batch reinforcement learning. In *Reinforcement learning*, pages 45–73. Springer, 2012.

Sergey Levine, Peter Pastor, Alex Krizhevsky, Julian Ibarz, and Deirdre Quillen. Learning hand-eye coordination for robotic grasping with deep learning and large-scale data collection. *The International journal of robotics research*, 37(4-5):421–436, 2018.

Sergey Levine, Aviral Kumar, George Tucker, and Justin Fu. Offline reinforcement learning: Tutorial, review, and perspectives on open problems. *arXiv preprint arXiv:2005.01643*, 2020.

Lihong Li, Yu Lu, and Dengyong Zhou. Provably optimal algorithms for generalized linear contextual bandits. In *International Conference on Machine Learning*, pages 2071–2080. PMLR, 2017.

Yao Liu, Adith Swaminathan, Alekh Agarwal, and Emma Brunskill. Provably good batch reinforcement learning without great exploration. *arXiv preprint arXiv:2007.08202*, 2020.

Mufti Mahmud, Mohammed Shamim Kaiser, Amir Hussain, and Stefano Vassanelli. Applications of deep learning and reinforcement learning to biological data. *IEEE transactions on neural networks and learning systems*, 29(6):2063–2079, 2018.

Vincent Mai, Kaustubh Mani, and Liam Paull. Sample efficient deep reinforcement learning via uncertainty estimation. *International Conference on Learning Representations*, 2022.

Yifei Min, Tianhao Wang, Dongruo Zhou, and Quanquan Gu. Variance-aware off-policy evaluation with linear function approximation. *Advances in neural information processing systems*, 2021.

Volodymyr Mnih, Koray Kavukcuoglu, David Silver, Andrei A Rusu, Joel Veness, Marc G Bellemare, Alex Graves, Martin Riedmiller, Andreas K Fidjeland, Georg Ostrovski, et al. Human-level control through deep reinforcement learning. *nature*, 518(7540):529–533, 2015.

Aditya Modi, Nan Jiang, Ambuj Tewari, and Satinder Singh. Sample complexity of reinforcement learning using linearly combined model ensembles. In *International Conference on Artificial Intelligence and Statistics*, pages 2010–2020. PMLR, 2020.

Rémi Munos. Error bounds for approximate policy iteration. In *ICML*, volume 3, pages 560–567, 2003.

Rémi Munos. Error bounds for approximate value iteration. In *Proceedings of the National Conference on Artificial Intelligence*, volume 20, page 1006. Menlo Park, CA; Cambridge, MA; London; AAAI Press; MIT Press; 1999, 2005.

Rémi Munos. Performance bounds in l_p-norm for approximate value iteration. *SIAM journal on control and optimization*, 46(2):541–561, 2007.

Thanh Nguyen-Tang, Sunil Gupta, Hung Tran-The, and Svetha Venkatesh. On finite-sample analysis of offline reinforcement learning with deep relu networks. *arXiv preprint arXiv:2103.06671*, 2021.

Mariya Popova, Olexandr Isayev, and Alexander Tropsha. Deep reinforcement learning for de novo drug design. *Science advances*, 4(7):eaap7885, 2018.

Martin Riedmiller. Neural fitted q iteration–first experiences with a data efficient neural reinforcement learning method. In *European conference on machine learning*, pages 317–328. Springer, 2005.

Daniel Russo and Benjamin Van Roy. Eluder dimension and the sample complexity of optimistic exploration. *Advances in Neural Information Processing Systems*, 26, 2013.

Julian Schrittwieser, Ioannis Antonoglou, Thomas Hubert, Karen Simonyan, Laurent Sifre, Simon Schmitt, Arthur Guez, Edward Lockhart, Demis Hassabis, Thore Graepel, et al. Mastering atari, go, chess and shogi by planning with a learned model. *Nature*, 588(7839):604–609, 2020.

John Schulman, Xi Chen, and Pieter Abbeel. Equivalence between policy gradients and soft q-learning. *arXiv preprint arXiv:1704.06440*, 2017.

Aaron Sidford, Mengdi Wang, Xian Wu, Lin Yang, and Yinyu Ye. Near-optimal time and sample complexities for solving markov decision processes with a generative model. In *Advances in Neural Information Processing Systems*, pages 5186–5196, 2018.

David Silver, Julian Schrittwieser, Karen Simonyan, Ioannis Antonoglou, Aja Huang, Arthur Guez, Thomas Hubert, Lucas Baker, Matthew Lai, Adrian Bolton, et al. Mastering the game of go without human knowledge. *nature*, 550(7676):354–359, 2017.

Wen Sun, Nan Jiang, Akshay Krishnamurthy, Alekh Agarwal, and John Langford. Model-based rl in contextual decision processes: Pac bounds and exponential improvements over model-free approaches. In *Conference on learning theory*, pages 2898–2933. PMLR, 2019.

Csaba Szepesvári and Rémi Munos. Finite time bounds for sampling based fitted value iteration. In *Proceedings of the 22nd international conference on Machine learning*, pages 880–887, 2005.

Mohammad Sadegh Talebi and Odalric-Ambrym Maillard. Variance-aware regret bounds for undiscounted reinforcement learning in mdps. In *Algorithmic Learning Theory*, pages 770–805. PMLR, 2018.

John Tsitsiklis and Benjamin Van Roy. Analysis of temporal-diffference learning with function approximation. *Advances in neural information processing systems*, 9, 1996.

Masatoshi Uehara, Xuezhou Zhang, and Wen Sun. Representation learning for online and offline rl in low-rank mdps. In *International Conference on Learning Representations*, 2022.

Ruosong Wang, Dean P Foster, and Sham M Kakade. What are the statistical limits of offline rl with linear function approximation? *International Conference on Learning Representations*, 2021a.

Yining Wang, Ruosong Wang, Simon S Du, and Akshay Krishnamurthy. Optimism in reinforcement learning with generalized linear function approximation. *International Conference on Learning Representations*, 2021b.

Zheng Wen and Benjamin Van Roy. Efficient exploration and value function generalization in deterministic systems. *Advances in Neural Information Processing Systems*, 26, 2013.

Yue Wu, Shuangfei Zhai, Nitish Srivastava, Joshua Susskind, Jian Zhang, Ruslan Salakhutdinov, and Hanlin Goh. Uncertainty weighted actor-critic for offline reinforcement learning. *International Conference on Machine Learning*, 2021.

Tengyang Xie and Nan Jiang. Batch value-function approximation with only realizability. *arXiv preprint arXiv:2008.04990*, 2020a.

Tengyang Xie and Nan Jiang. Q* approximation schemes for batch reinforcement learning: A theoretical comparison. In *Uncertainty in Artificial Intelligence*, pages 550–559, 2020b.

Tengyang Xie, Ching-An Cheng, Nan Jiang, Paul Mineiro, and Alekh Agarwal. Bellman-consistent pessimism for offline reinforcement learning. *Advances in neural information processing systems*, 2021.

Wei Xiong, Han Zhong, Chengshuai Shi, Cong Shen, Liwei Wang, and Tong Zhang. Nearly minimax optimal offline reinforcement learning with linear function approximation: Single-agent mdp and markov game. *arXiv preprint arXiv:2205.15512*, 2022.

Lin Yang and Mengdi Wang. Sample-optimal parametric q-learning using linearly additive features. In *International Conference on Machine Learning*, pages 6995–7004. PMLR, 2019.

Lin Yang and Mengdi Wang. Reinforcement learning in feature space: Matrix bandit, kernels, and regret bound. In *International Conference on Machine Learning*, pages 10746–10756. PMLR, 2020.

Ming Yin and Yu-Xiang Wang. Towards instance-optimal offline reinforcement learning with pessimism. *Advances in neural information processing systems*, 2021.

Ming Yin, Yaqi Duan, Mengdi Wang, and Yu-Xiang Wang. Near-optimal offline reinforcement learning with linear representation: Leveraging variance information with pessimism. *International Conference on Learning Representations*, 2022.

Andrea Zanette and Emma Brunskill. Tighter problem-dependent regret bounds in reinforcement learning without domain knowledge using value function bounds. In *International Conference on Machine Learning*, pages 7304–7312. PMLR, 2019.

Andrea Zanette, Alessandro Lazaric, Mykel Kochenderfer, and Emma Brunskill. Learning near optimal policies with low inherent bellman error. In *International Conference on Machine Learning*, pages 10978–10989. PMLR, 2020.

Andrea Zanette, Martin J. Wainwright, and Emma Brunskill. Provable benefits of actor-critic methods for offline reinforcement learning, 2021.

Wenhao Zhan, Baihe Huang, Audrey Huang, Nan Jiang, and Jason D Lee. Offline reinforcement learning with realizability and single-policy concentrability. *arXiv preprint arXiv:2202.04634*, 2022.

Ruiqi Zhang, Xuezhou Zhang, Chengzhuo Ni, and Mengdi Wang. Off-policy fitted q-evaluation with differentiable function approximators: Z-estimation and inference theory. *International Conference on Machine Learning*, 2022.

Zihan Zhang, Jiaqi Yang, Xiangyang Ji, and Simon S Du. Improved variance-aware confidence sets for linear bandits and linear mixture mdp. *Advances in Neural Information Processing Systems*, 34, 2021.

Dongruo Zhou, Quanquan Gu, and Csaba Szepesvari. Nearly minimax optimal reinforcement learning for linear mixture markov decision processes. In *Conference on Learning Theory*, pages 4532–4576. PMLR, 2021.

Appendix

A NOTATION LIST

$\Sigma_h^p(\theta)$	$\mathbb{E}_{\mu,h}\left[\nabla f(\theta, \phi(s,a)) \cdot \nabla f(\theta, \phi(s,a))^\top\right]$
κ	$\min_{h,\theta} \lambda_{\min}(\Sigma_h^p(\theta))$
ι	$\min_h \lambda_{\min}(\Lambda_h^p) \geq \kappa/H^2$ for any V_h
$\sigma_V^2(s,a)$	$\max\{1, \mathrm{Var}_{P_h}(V)(s,a)\}$ for any V
δ	Failure probability
K_0	$\max\left\{512\frac{\kappa_1^4}{\kappa^2}\left(\log(\frac{2Hd}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K^3}{\lambda^2})\right), \frac{4\lambda}{\kappa}\right\}$
ζ	$2\max_{s'\sim P(\cdot\mid s,a),h\in[H]} \frac{(\mathcal{P}_h V_{h+1}^\star)(s,a) - r - V_{h+1}^\star(s')}{\sigma_h^\star(s,a)}$
$C_{\mathrm{hot}} = \bar{C}_{\mathrm{hot}}$	$\frac{\kappa_1 H}{\sqrt{\kappa}} + \frac{\kappa_1^2 H^3 d^2}{\kappa} + \sqrt{\frac{d^3 H^4 \kappa_2^2 \kappa_1^2}{\kappa^3}} + \kappa_2 \max(\frac{\kappa_1}{\kappa}, \frac{1}{\sqrt{\kappa}})d^2 H^3 + \frac{d^2 H^4 \kappa_3 + \lambda \kappa_1 C_\Theta}{\kappa} + \frac{H^3 \kappa_2 d^2}{\kappa}$
$C'_{\mathrm{hot}} = \bar{C}'_{\mathrm{hot}}$	$C_{\mathrm{hot}} + \frac{\kappa_1 \kappa_2 H^4 d^2}{\kappa^{3/2}}$

B FURTHER ILLUSTRATION THAT GENERALIZED LINEAR MODEL EXAMPLE SATISFIES 2.3

Recall the definition in 2.5, then:

For $(\star\star)$,

$$\mathbb{E}_{\mu,h}\left[\nabla f(\theta, \phi(s,a)) \cdot \nabla f(\theta, \phi(s,a))^\top\right] = \mathbb{E}_{\mu,h}\left[f'(\langle\theta, \phi(s,a)\rangle)^2 \phi(\cdot,\cdot) \cdot \phi(\cdot,\cdot)^\top\right]$$
$$\succ \kappa^2 \mathbb{E}_{\mu,h}\left[\phi(\cdot,\cdot) \cdot \phi(\cdot,\cdot)^\top\right] \succ \kappa^3 I, \quad \forall \theta \in \Theta$$

For $(\star)$, by Taylor's Theorem,

$$\mathbb{E}_{\mu,h}\left[(f(\theta_1, \phi(\cdot,\cdot)) - f(\theta_2, \phi(\cdot,\cdot)))^2\right] = \mathbb{E}_{\mu,h}[f'(\theta_{s,a}, \phi(\cdot,\cdot))^2(\theta_1 - \theta_2)^\top \phi(\cdot,\cdot)\phi(\cdot,\cdot)^\top(\theta_1 - \theta_2)]$$
$$\geq \kappa^2 \mathbb{E}_{\mu,h}[(\theta_1 - \theta_2)^\top \phi(\cdot,\cdot)\phi(\cdot,\cdot)^\top(\theta_1 - \theta_2)] = \kappa^2(\theta_1 - \theta_2)^\top \mathbb{E}_{\mu,h}[\phi(\cdot,\cdot)\phi(\cdot,\cdot)^\top](\theta_1 - \theta_2) \geq \kappa^3 \|\theta_1 - \theta_2\|_2^2$$

and choose κ^3 as κ in 2.3.

C ON THE COMPUTATIONAL COMPLEXITY

Pessimistic Fitted Q-learning. For storage, at each time step $h \in [H]$ in Algorithm 1, we need to store $\widehat{\theta}_h, \Sigma_h$ and $\nabla f(\widehat{\theta}_h, \phi_{h,k})$. Therefore, the total space complexity is $O(dH + d^2 H + dKH)$. For computation, assuming $\widehat{\theta}_h$ is solved via SGD and let M denote the number of gradient steps, then the complexity is dominated by computing $\widehat{\theta}_h, \Sigma_h$ and Σ_h^{-1}, which results in $O(MH + KdH + d^3 H)$ complexity.

The space complexity and computational complexity for VAFQL has the same order as PFQL except that the constant factors are larger.

D SOME BASIC CONSTRUCTIONS

First of all, Recall in the first-order condition, we have

$$\nabla_\theta \left\{\sum_{k=1}^K \left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)\right]^2 + \lambda \cdot \|\theta\|_2^2\right\}\bigg|_{\theta=\widehat{\theta}_h} = 0, \quad \forall h \in [H].$$

Therefore, if we define the quantity $Z_h(\cdot, \cdot) \in \mathbb{R}^d$ as

$$Z_h(\theta|V) = \sum_{k=1}^{K} \left[f(\theta, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right] \nabla f(\theta, \phi_{h,k}) + \lambda \cdot \theta, \quad \forall \theta \in \Theta, \|V\|_2 \le H,$$

then we have (recall $\widehat{\theta}_h \in \text{Int}(\Theta)$)

$$Z_h(\widehat{\theta}_h | \widehat{V}_{h+1}) = 0.$$

In addition, according to Bellman completeness Assumption 2.1, for any bounded $V(\cdot) \in \mathbb{R}^{\mathcal{S}}$ with $\|V\|_\infty \le H$, $\inf_{f \in \mathcal{F}} \|f - \mathcal{P}_h(V)\|_\infty \le \epsilon_{\mathcal{F}}$, $\forall h$ (recall $\mathcal{P}_h(V) = r_h + \int_{\mathcal{S}} V dP_h$). Therefore, we can define the *parameter Bellman operator* $\mathbb{T}$ as follows.

Definition D.1 (parameter Bellman operator). *By the Bellman completeness Assumption 2.1, for any $\|V\|_\infty \le H$, we can define the* parameter Bellman operator $\mathbb{T} : V \to \theta_{\mathbb{T}V} \in \Theta$ *such that*

$$\theta_{\mathbb{T}V} = \arg\min_{\theta \in \Theta} \|f(\theta, \phi) - \mathcal{P}_h(V)\|_\infty$$

Denote $\delta_V := f(\theta_{\mathbb{T}V}, \phi) - \mathcal{P}_h(V)$, then we have $\|f(\theta_{\mathbb{T}V}, \phi) - \mathcal{P}_h(V)\|_\infty = \|\delta_V\|_\infty \le \epsilon_{\mathcal{F}}$. In particular, by realizability Assumption 2.1 it holds $\theta_{\mathbb{T}V_{h+1}^\star} = \theta_h^\star$ and this is due to $f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi) = \mathcal{P}_h(V_{h+1}^\star) = V_h^\star = f(\theta_h^\star, \phi).$[9]

D.1 SUBOPTIMALITY DECOMPOSITION

Denote $\iota_h(s, a) := \mathcal{P}_h \widehat{V}_{h+1}(s, a) - \widehat{Q}_h(s, a)$, by Jin et al. (2021b) we have the following decomposition.

Lemma D.2 (Lemma 3.1 of Jin et al. (2021b)). *Let $\widehat{\pi} = \{\widehat{\pi}_h\}_{h=1}^H$ a policy and $\widehat{Q}_h$ be any estimates with $\widehat{V}_h = \langle \widehat{Q}_h(s, \cdot), \widehat{\pi}_h(\cdot \mid s) \rangle_{\mathcal{A}}$. Then for any policy π, we have*

$$v^\pi - v^{\widehat{\pi}} = -\sum_{h=1}^{H} E_{\widehat{\pi}}[\iota_h(s_h, a_h)] + \sum_{h=1}^{H} E_\pi[\iota_h(s_h, a_h)] + \sum_{h=1}^{H} E_\pi[\langle \widehat{Q}_h(s_h, \cdot), \pi_h(\cdot \mid s_h) - \widehat{\pi}_h(\cdot \mid s_h) \rangle_{\mathcal{A}}].$$

In particular, if we choose $\widehat{\pi}_h(\cdot|s) := \arg\max_\pi \langle \widehat{Q}_h(s, \cdot), \pi(\cdot \mid s) \rangle_{\mathcal{A}}$, then

$$v^\pi - v^{\widehat{\pi}} = -\sum_{h=1}^{H} E_{\widehat{\pi}}[\iota_h(s_h, a_h)] + \sum_{h=1}^{H} E_\pi[\iota_h(s_h, a_h)].$$

Lemma D.3. *Let $\widehat{\mathcal{P}}_h$ be the general estimated Bellman operator. Suppose with probability $1 - \delta$, it holds for all $h, s, a \in [H] \times \mathcal{S} \times A$ that $|(\mathcal{P}_h \widehat{V}_{h+1} - \widehat{\mathcal{P}}_h \widehat{V}_{h+1})(s, a)| \le \Gamma_h(s, a)$, then it implies $\forall s, a, h \in \mathcal{S} \times \mathcal{A} \times [H]$, $0 \le \zeta_h(s, a) \le 2\Gamma_h(s, a)$. Furthermore, it holds for any policy π simultaneously, with probability $1 - \delta$,*

$$V_1^\pi(s) - V_1^{\widehat{\pi}}(s) \le \sum_{h=1}^{H} 2 \cdot \mathbb{E}_\pi [\Gamma_h(s_h, a_h) \mid s_1 = s].$$

Proof of Lemma D.3. This is a generic result that holds true for the general MDPs and was first raised by Theorem 4.2 of Jin et al. (2021b). Later, it is summarized in Lemma C.1 of Yin et al. (2022). □

With Lemma D.3, we need to bound the term $|\mathcal{P}_h \widehat{V}_{h+1}(s, a) - \widehat{\mathcal{P}}_h \widehat{V}_{h+1}(s, a)|$.

---

[9]Here without loss of generality we assume $Q_h^\star$ can be uniquely identified, *i.e.* there is a unique $\theta^\star$ such that $f(\theta_h^\star, \phi) = Q_h^\star$.

E   ANALYZING $|\mathcal{P}_h\widehat{V}_{h+1}(s,a) - \widehat{\mathcal{P}}_h\widehat{V}_{h+1}(s,a)|$ FOR PFQL.

Throughout this section, we suppose $\epsilon_{\mathcal{F}} = 0$, *i.e.* $f(\theta_{\mathbb{T}V}, \phi) = \mathcal{P}_h(V)$. According to the regression oracle (Line 4 of Algorithm 1), the estimated Bellman operator $\widehat{\mathcal{P}}_h$ maps $\widehat{V}_{h+1}$ to $\widehat{\theta}_h$, *i.e.* $\widehat{\mathcal{P}}_h\widehat{V}_{h+1} = f(\widehat{\theta}_h, \phi)$. Therefore (recall Definition D.1)

$$\mathcal{P}_h\widehat{V}_{h+1}(s,a) - \widehat{\mathcal{P}}_h\widehat{V}_{h+1}(s,a) = \mathcal{P}_h\widehat{V}_{h+1}(s,a) - f(\widehat{\theta}_h, \phi(s,a))$$
$$= f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\widehat{\theta}_h, \phi(s,a)) \tag{3}$$
$$= \nabla f(\widehat{\theta}_h, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right) + \mathrm{Hot}_{h,1},$$

where we apply the first-order Taylor expansion for the differentiable function f at point $\widehat{\theta}_h$ and $\mathrm{Hot}_{h,1}$ is a higher-order term. Indeed, the following Lemma E.1 bounds the $\mathrm{Hot}_{h,1}$ term with $\widetilde{O}(\frac{1}{K})$.

Lemma E.1. *Recall the definition (from the above decomposition)* $\mathrm{Hot}_{h,1} := f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\widehat{\theta}_h, \phi(s,a)) - \nabla f(\widehat{\theta}_h, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)$, *then with probability* $1 - \delta$,

$$|\mathrm{Hot}_{h,1}| \le \frac{18H^2\kappa_2(\log(H/\delta) + C_{d,\log K}) + \kappa_2\lambda C_\Theta^2}{\kappa K}, \quad \forall h \in [H].$$

Proof of Lemma E.1. By second-order Taylor's Theorem, there exists a point ξ (lies in the line segment of $\widehat{\theta}_h$ and $\theta_{\mathbb{T}\widehat{V}_{h+1}}$) such that

$$f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\widehat{\theta}_h, \phi(s,a)) = \nabla f(\widehat{\theta}_h, \phi(s,a))^\top\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right) + \frac{1}{2}\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)^\top\nabla^2_{\theta\theta}f(\xi, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)$$

Therefore, by directly applying Theorem G.2, with probability $1 - \delta$, for all $h \in [H]$,

$$|\mathrm{Hot}_{h,1}| = \frac{1}{2}\left|\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)^\top\nabla^2_{\theta\theta}f(\xi, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)\right|$$
$$\le \frac{1}{2}\kappa_2 \cdot \left\|\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right\|_2^2 \le \frac{18H^2\kappa_2(\log(H/\delta) + C_{d,\log K}) + \kappa_2\lambda C_\Theta^2}{\kappa K}$$

$\square$

E.1   ANALYZING $\nabla f(\widehat{\theta}_h, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)$ VIA Z_h.

From (3) and Lemma E.1, the problem further reduces to bounding $\nabla f(\widehat{\theta}_h, \phi(s,a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right)$. To begin with, we first provide a characterization of $\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h$. Indeed, by first-order Vector Taylor expansion (Lemma L.1), we have (note $Z_h(\widehat{\theta}_h|\widehat{V}_{h+1}) = 0$) for any $\theta \in \Theta$,

$$Z_h(\theta|\widehat{V}_{h+1}) - Z_h(\widehat{\theta}_h|\widehat{V}_{h+1}) = \Sigma^s_h(\theta - \widehat{\theta}_h) + R_K(\theta), \tag{4}$$

where $R_K(\theta)$ is the higher-order residuals and $\Sigma^s_h := \frac{\partial}{\partial\theta}Z_h(\theta|\widehat{\theta}_{h+1})\Big|_{\theta=\widehat{\theta}_h}$ with

$$\Sigma^s_h := \frac{\partial}{\partial\theta}Z_h(\theta|\widehat{V}_{h+1})\Big|_{\theta=\widehat{\theta}_h} = \frac{\partial}{\partial\theta}\left(\sum_{k=1}^K\left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s^k_{h+1})\right]\nabla f(\theta, \phi_{h,k}) + \lambda\cdot\theta\right)_{\theta=\widehat{\theta}_h}$$
$$= \underbrace{\sum_{k=1}^K\left\{\left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s^k_{h+1})\right)\cdot\nabla^2_{\theta\theta}f(\widehat{\theta}_h, \phi_{h,k})\right\}}_{:=\Delta_{\Sigma^s_h}}$$
$$+ \underbrace{\sum_{k=1}^K\nabla_\theta f(\widehat{\theta}_h, \phi_{h,k})\nabla^\top_\theta f(\widehat{\theta}_{h,k}, \phi_{h,k}) + \lambda I_d}_{:=\Sigma_h}, \tag{5}$$

here $\nabla^2 = \nabla \bigotimes \nabla$ denotes outer product of gradients.

Note $\Delta_{\Sigma_h^s}$ is not desirable since it could prevent Σ_h^s from being positive-definite (and it could cause Σ_h^s to be singular). Therefore, we first deal with $\Delta_{\Sigma_h^s}$ in below.

Lemma E.2. *With probability $1 - \delta$, for all $h \in [H]$,*

$$\frac{1}{K} \left\| \Delta_{\Sigma_h^s} \right\|_2 = \left\| \frac{1}{K} \sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\widehat{\theta}_h, \phi_h) \right\|_2$$

$$\leq 9\kappa_2 \max(\frac{\kappa_1}{\sqrt{\kappa}}, 1) \sqrt{\frac{dH^2(\log(2H/\delta) + d\log(1 + 2C_\Theta H\kappa_3 K) + C_{d,\log K})}{K}} + \frac{1}{K}.$$

Proof of Lemma E.2. **Step1:** We prove for fixed $\bar{\theta} \in \Theta$, with probability $1 - \delta$, for all $h \in [H]$,

$$\left\| \frac{1}{K} \sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \leq 9\kappa_2 \max(\frac{\kappa_1}{\sqrt{\kappa}}, 1) \sqrt{\frac{H^2(\log(2H/\delta) + C_{d,\log K})}{K}}.$$

Indeed, we have

$$\left\| \frac{1}{K} \sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2$$

$$\leq \left\| \frac{1}{K} \sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \qquad (6)$$

$$+ \left\| \frac{1}{K} \sum_{k=1}^K \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2.$$

On one hand, by Theorem G.2 with probability $1 - \delta/2$ for all $h \in [H]$

$$\left\| \frac{1}{K} \sum_{k=1}^K \left(f(\widehat{\theta}_h, \phi_{h,k}) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \leq \kappa_2 \cdot \max_{\theta,s,a} \|\nabla f(\theta, \phi(s,a))\|_2 \left\| \widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2$$

$$\leq \kappa_2 \kappa_1 \left\| \widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2 \leq \kappa_2 \kappa_1 \left(\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} + \sqrt{\frac{b_{d,K,\epsilon_\mathcal{F}}}{\kappa}} + \sqrt{\frac{2H\epsilon_\mathcal{F}}{\kappa}} \right).$$
$$(7)$$

On other hand, recall the definition of $\mathbb{T}$, we have

$$\mathbb{E}\left[(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_{h,k}) \Big| s_h^k, a_h^k \right]$$

$$= \mathbb{E}\left[f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \Big| s_h^k, a_h^k \right] \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_{h,k})$$

$$= \left((\mathcal{P}_h \widehat{V}_{h+1})(s_h^k, a_h^k) - \mathbb{E}\left[r_{h,k} + \widehat{V}_{h+1}(s_{h+1}^k) \Big| s_h^k, a_h^k \right] \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_{h,k})$$

$$= \left((\mathcal{P}_h \widehat{V}_{h+1})(s_h^k, a_h^k) - (\mathcal{P}_h \widehat{V}_{h+1}(s_{h+1}^k))(s_h^k, a_h^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_{h,k}) = 0.$$

Also, since $\left\| \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \leq H\kappa_2$, denote $\sigma^2 := K \cdot H^2\kappa_2^2$, then by Vector Hoeffding's inequality (Lemma L.2),

$$\mathbb{P}\left(\left\| \frac{1}{K} \sum_{k=1}^K \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \geq t/K \Big| \{s_h^k, a_h^k\}_{k=1}^K \right) \leq d \cdot e^{-t^2/8dKH^2\kappa_2^2} := \delta$$

which is equivalent to

$$\mathbb{P}\left(\left\| \frac{1}{K} \sum_{k=1}^K \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\bar{\theta}, \phi_h) \right\|_2 \leq \sqrt{\frac{8dH^2\kappa_2^2 \log(d/\delta)}{K}} \Big| \{s_h^k, a_h^k\}_{k=1}^K \right) \geq 1-\delta$$

Define $A = \{\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\nabla_{\theta\theta}^{2}f(\bar{\theta},\phi_{h})\right\|_{2} \leq$
$\sqrt{\frac{8dH^{2}\kappa_{2}^{2}\log(d/\delta)}{K}}\}$, then by law of total expectation $\mathbb{P}(A)=\mathbb{E}[\mathbf{1}_{A}]=\mathbb{E}[\mathbb{E}[\mathbf{1}_{A}|\{s_{h}^{k},a_{h}^{k}\}_{k=1}^{K}]]=$
$\mathbb{E}[\mathbb{P}[A|\{s_{h}^{k},a_{h}^{k}\}_{k=1}^{K}]]\geq\mathbb{E}[1-\delta]=1-\delta$, i.e. with probability at least $1-\delta/2$ (and a union bound),

$$\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\nabla_{\theta\theta}^{2}f(\bar{\theta},\phi_{h})\right\|_{2}\leq\sqrt{\frac{8dH^{2}\kappa_{2}^{2}\log(2Hd/\delta)}{K}},\ \forall h\in[H].$$

Using above and equation 6, equation 7 and a union bound, w.p. $1-\delta$, for all $h\in[H]$,

$$\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\widehat{\theta}_{h},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\nabla_{\theta\theta}^{2}f(\bar{\theta},\phi_{h})\right\|_{2}\leq 6\kappa_{2}\kappa_{1}\sqrt{\frac{H^{2}(\log(2H/\delta)+C_{d,\log K})}{\kappa K}}+\sqrt{\frac{8dH^{2}\kappa_{2}^{2}\log(2Hd/\delta)}{K}}$$

$$\leq 9\kappa_{2}\max(\frac{\kappa_{1}}{\sqrt{\kappa}},1)\sqrt{\frac{dH^{2}(\log(2H/\delta)+C_{d,\log K})}{K}}$$

Step2: we finish the proof of the lemma.

Consider the function class $\left\{f(\bar{\theta}):=\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\widehat{\theta}_{h},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\nabla_{\theta\theta}^{2}f(\bar{\theta},\phi_{h})\right\|_{2}\Big|\bar{\theta}\in\Theta\right\}$,
then by triangular inequality

$$|f(\bar{\theta}_{1})-f(\bar{\theta}_{2})|\leq\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\widehat{\theta}_{h},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\left[\nabla_{\theta\theta}^{2}f(\bar{\theta}_{1},\phi_{h})-\nabla_{\theta\theta}^{2}f(\bar{\theta}_{2},\phi_{h})\right]\right\|_{2}$$

$$\leq H\cdot\sup_{s,a}\left\|\nabla_{\theta\theta}^{2}f(\bar{\theta}_{1},\phi_{h})-\nabla_{\theta\theta}^{2}f(\bar{\theta}_{2},\phi_{h})\right\|_{2}\leq H\kappa_{3}\left\|\bar{\theta}_{1}-\bar{\theta}_{2}\right\|_{2}.$$

By Lemma L.8, the covering number $\mathcal{C}$ of the ϵ-net of the above function class satisfies $\log\mathcal{C}\leq d\log(1+\frac{2C_{\Theta}H\kappa_{3}}{\epsilon})$. By choosing $\epsilon=1/K$, by a union bound over $\mathcal{C}$ cases we obtain for all $h\in[H]$

$$\left\|\frac{1}{K}\sum_{k=1}^{K}\left(f(\widehat{\theta}_{h},\phi_{h,k})-r_{h,k}-\widehat{V}_{h+1}(s_{h+1}^{k})\right)\cdot\nabla_{\theta\theta}^{2}f(\widehat{\theta}_{h},\phi_{h})\right\|_{2}$$

$$\leq 9\kappa_{2}\max(\frac{\kappa_{1}}{\sqrt{\kappa}},1)\sqrt{\frac{dH^{2}(\log(2H/\delta)+d\log(1+2C_{\Theta}H\kappa_{3}K)+C_{d,\log K})}{K}}+\frac{1}{K}.$$

$\square$

Combing Lemma E.2 and Theorem G.2 (and a union bound), we directly have

Corollary E.3. *With probability* $1-\delta$,

$$\left\|\frac{1}{K}\Delta_{\Sigma_{h}^{s}}(\widehat{\theta}_{h}-\theta_{\mathbb{T}\widehat{V}_{h+1}})\right\|_{2}\leq\left\|\frac{1}{K}\Delta_{\Sigma_{h}^{s}}\right\|_{2}\left\|\widehat{\theta}_{h}-\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_{2}\leq\widetilde{O}(\frac{\kappa_{2}\max(\frac{\kappa_{1}}{\kappa},\frac{1}{\sqrt{\kappa}})d^{2}H^{2}}{K})$$

Here $\widetilde{O}$ *absorbs all the constants and Polylog terms.*

Now we select $\theta=\theta_{\mathbb{T}\widehat{V}_{h+1}}$ in equation 4, and denote $\widetilde{R}_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})=\Delta_{\Sigma_{h}^{s}}(\widehat{\theta}_{h}-\theta_{\mathbb{T}\widehat{V}_{h+1}})+R_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})$, then equation 4 is equivalent to

$$Z_{h}(\theta_{\mathbb{T}\widehat{V}_{h+1}}|\widehat{V}_{h+1})-Z_{h}(\widehat{\theta}_{h}|\widehat{V}_{h+1})=\Sigma_{h}^{s}(\theta_{\mathbb{T}\widehat{V}_{h+1}}-\widehat{\theta}_{h})+R_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})=\Sigma_{h}(\theta_{\mathbb{T}\widehat{V}_{h+1}}-\widehat{\theta}_{h})+\widetilde{R}_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})$$

Note $\lambda>0$ implies Σ_{h} is invertible, then we have (recall $Z_{h}(\widehat{\theta}_{h}|\widehat{\theta}_{h+1})=0$)

$$\theta_{\mathbb{T}\widehat{V}_{h+1}}-\widehat{\theta}_{h}=\Sigma_{h}^{-1}[Z_{h}(\theta_{\mathbb{T}\widehat{V}_{h+1}}|\widehat{V}_{h+1})-Z_{h}(\widehat{\theta}_{h}|\widehat{V}_{h+1})]-\Sigma_{h}^{-1}\widetilde{R}_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})$$

$$=\Sigma_{h}^{-1}[Z_{h}(\theta_{\mathbb{T}\widehat{V}_{h+1}}|\widehat{V}_{h+1})]-\Sigma_{h}^{-1}\widetilde{R}_{K}(\theta_{\mathbb{T}\widehat{V}_{h+1}})$$

Plug it back to equation 3 to get

$$
\nabla f(\widehat{\theta}_h, \phi(s,a)) \left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right)
$$
$$
= \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [Z_h(\theta_{\mathbb{T}\widehat{V}_{h+1}} | \widehat{V}_{h+1})] - \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} \widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})
$$
$$
= \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) + \lambda \theta_{\mathbb{T}\widehat{V}_{h+1}}]
$$
$$
- \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} \widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})
$$
$$
= \underbrace{\nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k})]}_{:=I}
$$
$$
- \underbrace{\nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} \left[\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) + \lambda \theta_{\mathbb{T}\widehat{V}_{h+1}} \right]}_{:=\text{Hot}_2}
$$

(8)

We will bound second term Hot_2 to have higher order $O(\frac{1}{K})$ in Section E.5 and focus on the first term. By direct decomposition,

$$
I := \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k})]
$$
$$
= \underbrace{\nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})]}_{:=I_1}
$$
$$
+ \underbrace{\nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - \widehat{V}_{h+1}(s_{h+1}^k) + V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})]}_{:=I_2}
$$
$$
+ \underbrace{\nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} \left[\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \left(\nabla_\theta^\top f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) \right) \right]}_{:=I_3}
$$

E.2 BOUNDING THE TERM I_3

We first bound the term I_3. We have the following Lemma.

Lemma E.4. *For any fixed $V(\cdot) \in \mathbb{R}^{\mathcal{S}}$ with $\|V\|_\infty \leq H$ and any fixed θ such that $\|\theta_{\mathbb{T}V} - \theta\|_2 \leq$*
$\sqrt{\frac{36H^2(\log(H/\delta)+C_{d,\log K})+2\lambda C_\Theta^2}{\kappa K}}$. *Let*

$$
\widetilde{I}_3 := \nabla f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \left[\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot \left(\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k}) \right) \right],
$$

and if $K \geq \max \left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d \log(1 + \frac{4\kappa_1 D^2 \kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$, then with probability $1 - \delta$,
(where $D = \max\{\kappa_1, \sqrt{\frac{(144dH^2\kappa_2^2(H^2\log(H/\delta)+C_{d,\log K})+8dH^2\kappa_2^2\lambda C_\Theta^2)\log(d/\delta)}{\kappa}}\}$)

$$
|\widetilde{I}_3| \leq 4\kappa_1 \sqrt{\frac{(144dH^2\kappa_2^2 \left(H^2 \log(H/\delta) + C_{d,\log K} \right) + 8dH^2\kappa_2^2\lambda C_\Theta^2) \log(d/\delta)}{\kappa^3}} \frac{1}{K} + O(\frac{1}{K^{3/2}}).
$$

Proof of Lemma E.4. Indeed, with probability $1 - \delta/2$,

$$|\widetilde{I_3}| = \left\| \nabla f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \left[\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right] \right\|$$

$$\leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_{\Sigma_h^{-1}}$$

$$\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_{\Sigma_h^{-1}}$$

where, under the condition $K \geq \max\left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d \log(1 + \frac{4\kappa_1^3 \kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$, we applied Lemma L.5 .

Next, on one hand, $\|\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})\|_2 \leq \kappa_2 \cdot \|\theta_{\mathbb{T}V} - \theta\|_2 \leq \kappa_2 \sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}}$. On the other hand,

$$\mathbb{E}\left[\left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot \left(\nabla_\theta^\top f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta^\top f(\theta, \phi_{h,k}) \right) \big| s_h^k, a_h^k \right]$$

$$= \mathbb{E}\left[\left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \big| s_h^k, a_h^k \right] \cdot \left(\nabla_\theta^\top f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta^\top f(\theta, \phi_{h,k}) \right)$$

$$= \left((\mathcal{P}_h V)(s_h^k, a_h^k) - (\mathcal{P}_h V)(s_h^k, a_h^k) \right) \cdot \left(\nabla_\theta^\top f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta^\top f(\theta, \phi_{h,k}) \right) = 0$$

Therefore by Vector Hoeffding's inequality (Lemma L.2) (also note the condition for boundedness $\left\| \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_2 \leq H\kappa_2 \cdot \|\theta_{\mathbb{T}V} - \theta\|_2 \leq H\kappa_2 \sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}}$) with probability $1 - \delta/2$,

$$\left\| \frac{1}{K} \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_2$$

$$\leq \sqrt{\frac{4d \left(H\kappa_2 \sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} \right)^2 \log(\frac{d}{\delta})}{K}}$$

$$= \sqrt{\frac{(144 d H^2 \kappa_2^2 (H^2 \log(H/\delta) + C_{d,\log K}) + 8 d H^2 \kappa_2^2 \lambda C_\Theta^2) \log(d/\delta)}{\kappa}} \cdot \frac{1}{K}$$

and this implies with probability $1 - \delta/2$,

$$\left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_2$$

$$\leq \sqrt{\frac{(144 d H^2 \kappa_2^2 (H^2 \log(H/\delta) + C_{d,\log K}) + 8 d H^2 \kappa_2^2 \lambda C_\Theta^2) \log(d/\delta)}{\kappa}}$$

choose $u = \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k}))$ in Lemma L.5, by a union bound we obtain with probability $1 - \delta$

$$|\widetilde{I_3}| = \left\| \nabla f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \left[\sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right] \right\|$$

$$\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k) \right) \cdot (\nabla_\theta f(\theta_{\mathbb{T}V}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})) \right\|_{\Sigma_h^{-1}}$$

$$\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \left(2\sqrt{\frac{(144 d H^2 \kappa_2^2 (H^2 \log(H/\delta) + C_{d,\log K}) + 8 d H^2 \kappa_2^2 \lambda C_\Theta^2) \log(d/\delta)}{\kappa^2 K}} + O(\frac{1}{K}) \right)$$

$$= 4\kappa_1 \sqrt{\frac{(144 d H^2 \kappa_2^2 (H^2 \log(H/\delta) + C_{d,\log K}) + 8 d H^2 \kappa_2^2 \lambda C_\Theta^2) \log(d/\delta)}{\kappa^3}} \frac{1}{K} + O(\frac{1}{K^{3/2}}).$$

$\square$

Lemma E.5. *Under the same condition as Lemma E.4. With probability $1 - \delta$,*

$$|I_3| \leq 4\kappa_1 \sqrt{\frac{(144dH^2\kappa_2^2\left(H^2\log(H/\delta) + D_{d,\log K} + C_{d,\log K}\right) + 8dH^2\kappa_2^2\lambda C_\Theta^2)(\log(d/\delta) + D_{d,\log K})}{\kappa^3}}\frac{1}{K} + O(\frac{1}{K^{3/2}}).$$

Here $D_{d,\log K}$:= $d \cdot \log(1 + 6C_\Theta(2\kappa_1^2 + H\kappa_2)K) + d\log(1 + 6C_\Theta H\kappa_2 K) + d\log\left(1 + 288C_\Theta\kappa_1^2(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2K^2\right) + d^2\log\left(1 + 288\sqrt{d}B\kappa_1^4K^2\right)$ = $\widetilde{O}(d^2)$ *with* $\widetilde{O}$ *absorbs Polylog terms.*

Proof of Lemma E.5. Define

$$h(V, \widetilde{\theta}, \theta) = \sum_{k=1}^{K}\left(f(\widetilde{\theta}, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k)\right)\cdot\left(\nabla_\theta f(\widetilde{\theta}, \phi_{h,k}) - \nabla_\theta f(\theta, \phi_{h,k})\right),$$

then

$$|h(V_1, \widetilde{\theta}_1, \theta_1) - h(V_2, \widetilde{\theta}_2, \theta_2)|$$

$$\leq \left|\sum_{k=1}^{K}\left([f(\widetilde{\theta}_1, \phi_{h,k}) - V_1(s_{h+1}^k)] - [f(\widetilde{\theta}_2, \phi_{h,k}) - V_2(s_{h+1}^k)]\right)\cdot\left(\nabla_\theta f(\widetilde{\theta}_1, \phi_{h,k}) - \nabla_\theta f(\theta_1, \phi_{h,k})\right)\right|$$

$$+ \left|\sum_{k=1}^{K}\left(f(\widetilde{\theta}_2, \phi_{h,k}) - r_{h,k} - V_2(s_{h+1}^k)\right)\cdot\left([\nabla_\theta f(\widetilde{\theta}_1, \phi_{h,k}) - \nabla_\theta f(\theta_1, \phi_{h,k})] - [\nabla_\theta f(\widetilde{\theta}_2, \phi_{h,k}) - \nabla_\theta f(\theta_2, \phi_{h,k})]\right)\right|$$

$$\leq K\sup_{s,a,s'}\left|[f(\widetilde{\theta}_1, \phi(s,a)) - f(\widetilde{\theta}_2, \phi(s,a))] - [V_1(s') - V_2(s')]\right|_2\cdot 2\kappa_1$$

$$+ KH\cdot\sup_{s,a}\left\|[\nabla_\theta f(\widetilde{\theta}_1, \phi(s,a)) - \nabla_\theta f(\theta_1, \phi(s,a))] - [\nabla_\theta f(\widetilde{\theta}_2, \phi(s,a)) - \nabla_\theta f(\theta_2, \phi(s,a))]\right\|_2$$

$$\leq K2\kappa_1^2\left\|\widetilde{\theta}_1 - \widetilde{\theta}_2\right\|_2 + 2K\kappa_1\|V_1 - V_2\|_\infty + HK\kappa_2\left\|\widetilde{\theta}_1 - \widetilde{\theta}_2\right\|_2 + HK\kappa_2\|\theta_1 - \theta_2\|_2$$

$$= (2\kappa_1^2 + H\kappa_2)K\left\|\widetilde{\theta}_1 - \widetilde{\theta}_2\right\|_2 + 2\kappa_1K\|V_1 - V_2\|_\infty + HK\kappa_2\|\theta_1 - \theta_2\|_2.$$

Let $\mathcal{C}_a$ be the $\frac{\epsilon/3}{(2\kappa_1^2 + H\kappa_2)K}$-covering net of $\{\theta : \|\theta\|_2 \leq C_\Theta\}$, $\mathcal{C}_V$ be the $\frac{\epsilon}{6\kappa_1 K}$-covering net of $\mathcal{V}$ defined in Lemma L.9 and $\mathcal{C}_b$ be the $\frac{\epsilon}{3H\kappa_2 K}$-covering net of $\{\theta : \|\theta\|_2 \leq C_\Theta\}$, then by Lemma L.8 and Lemma L.9,

$$\log|\mathcal{C}_a| \leq d\cdot\log(1 + \frac{6C_\Theta(2\kappa_1^2 + H\kappa_2)K}{\epsilon}), \quad \log|\mathcal{C}_b| \leq d\log(1 + \frac{6C_\Theta H\kappa_2 K}{\epsilon})$$

$$\log\mathcal{C}_V \leq d\log\left(1 + \frac{288C_\Theta\kappa_1^2(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2K^2}{\epsilon^2}\right) + d^2\log\left(1 + \frac{288\sqrt{d}B\kappa_1^4K^2}{\epsilon^2}\right).$$

Further notice with probability $1 - \delta/2$ (by Lemma L.5), for all fixed sets of parameters θ, V satisfies $\|\theta_{\mathbb{T}V} - \theta\|_2 \leq \sqrt{\frac{36H^2(\log(2H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}}$ simultaneously,

$$|I_3 - \widetilde{I}_3| \leq \left\|\nabla f(\widehat{\theta}_h, \phi(s,a))\right\|_{\Sigma_h^{-1}}\cdot\left\|h(\widehat{V}_{h+1}, \theta_{\mathbb{T}\widehat{V}_{h+1}}, \widehat{\theta}_h) - h(V, \theta_{\mathbb{T}V}, \theta)\right\|_{\Sigma_h^{-1}}$$

$$\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K})\right)\cdot\left\|h(\widehat{V}_{h+1}, \theta_{\mathbb{T}\widehat{V}_{h+1}}, \widehat{\theta}_h) - h(V, \theta_{\mathbb{T}V}, \theta)\right\|_{\Sigma_h^{-1}}$$

and $\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right\|_2 \leq \sqrt{\frac{36H^2(\log(2H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}}$ with probability $1 - \delta/2$ by Theorem G.2. Now, choosing $\epsilon = O(1/K^2)$ and by Lemma E.4 and union bound over covering instances, we obtain with probability $1 - \delta$

$$|I_3| \leq 4\kappa_1\sqrt{\frac{(144dH^2\kappa_2^2\left(H^2\log(H/\delta) + D_{d,\log K} + C_{d,\log K}\right) + 8dH^2\kappa_2^2\lambda C_\Theta^2)(\log(d/\delta) + D_{d,\log K})}{\kappa^3}}\frac{1}{K} + O(\frac{1}{K^{3/2}}).$$

$\square$

E.3 BOUNDING THE SECOND TERM I_2

In this section, we bound the term

$$I_2 := \nabla f(\widehat{\theta}_h, \phi(s,a)) \Sigma_h^{-1} [\sum_{k=1}^{K} \Big(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - \widehat{V}_{h+1}(s_{h+1}^k) + V_{h+1}^\star(s_{h+1}^k) \Big) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})].$$

The following Lemma shows that I_2 is a higher-order error term with rate $\widetilde{O}(\frac{1}{K})$.

Lemma E.6 (Bounding I_2). *If K satisfies $K \geq 512\frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K}{\lambda^2}) \right)$, and $K \geq 4\lambda/\kappa$, then with probability $1 - \delta$*

$$|I_2| \leq \widetilde{O}(\frac{\kappa_1^2 H^2 d^2}{\kappa K}) + \widetilde{O}(\frac{1}{K^{3/2}}).$$

Here $\widetilde{O}$ absorbs constants and Polylog terms.

Proof of Lemma E.6. **Step1.** Define $\eta_k(V) := f(\theta_{\mathbb{T}V}, \phi_{h,k}) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - V(s_{h+1}^k) + V_{h+1}^\star(s_{h+1}^k)$ and let $\|V(\cdot)\|_\infty \leq H$ be any fixed function such that $\sup_{s_h^k, a_h^k, s_{h+1}^k} |\eta_k(V)| \leq \widetilde{O}(\kappa_1 H^2 \sqrt{\frac{d^2}{\kappa K}})$, *i.e.* arbitrary fixed V function in the neighborhood (measured by η_k) of $V_{h+1}^\star$. Then by definition of $\mathbb{T}$ it holds $\mathbb{E}[\eta_k(V,\theta)|s_h^k, a_h^k] = 0$. Let the fixed $\theta \in \Theta$ be arbitrary and define $x_k(\theta) = \nabla_\theta f(\theta, \phi_{h,k})$. Next, define $G_h(\theta) = \sum_{k=1}^{K} \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top + \lambda I_d$, since $\|x_k\|_2 \leq \kappa_1$ and $|\eta_k| \leq \widetilde{O}(\kappa_1 H^2 \sqrt{\frac{d^2}{\kappa K}})$, by self-normalized Hoeffding's inequality (Lemma L.3), with probability $1 - \delta$ (recall $t := K$ in Lemma L.3),

$$\left\| \sum_{k=1}^{K} x_k(\theta)\eta_k(V) \right\|_{G_h(\theta)^{-1}} \leq \widetilde{O}(\kappa_1 H^2 \sqrt{\frac{d^2}{\kappa K}}) \sqrt{d\log\left(\frac{\lambda + K\kappa_1}{\lambda\delta}\right)}.$$

Step2. Define $h(V,\theta) := \sum_{k=1}^{K} x_k(\theta)\eta_k(V)$ and $H(V,\theta) := \left\| \sum_{k=1}^{K} x_k(\theta)\eta_k(V) \right\|_{G_h(\theta)^{-1}}$, then note by definition $|\eta_k(V)| \leq 2H$, which implies $\|h(V,\theta)\|_2 \leq 2KH\kappa_1$ and

$$|\eta_k(V_1) - \eta_k(V_2)| \leq |\mathcal{P}_h V_1 - \mathcal{P}_h V_2| + \|V_1 - V_2\|_\infty \leq 2\|V_1 - V_2\|_\infty$$

and

$$\|h(V_1,\theta_1) - h(V_2,\theta_2)\|_2 \leq K \max_k \left(2H\|x_k(\theta_1) - x_k(\theta_2)\|_2 + \kappa_1|\eta_k(V_1) - \eta_k(V_2)|\right)$$

$$\leq K(2H\kappa_2\|\theta_1 - \theta_2\|_2 + 2\kappa_1\|V_1 - V_2\|_\infty).$$

Furthermore,

$$\left\|G_h(\theta_1)^{-1} - G_h(\theta_2)^{-1}\right\|_2 \leq \left\|G_h(\theta_1)^{-1}\right\|_2 \|G_h(\theta_1) - G_h(\theta_2)\|_2 \left\|G_h(\theta_2)^{-1}\right\|_2$$

$$\leq \frac{1}{\lambda^2} K \sup_k \left\|\nabla f(\theta_1, \phi_{h,k}) \cdot \nabla f(\theta_1, \phi_{h,k})^\top - \nabla f(\theta_2, \phi_{h,k}) \cdot \nabla f(\theta_2, \phi_{h,k})^\top\right\|_2$$

$$\leq \frac{1}{\lambda^2} K \sup_k \left[\left\|(\nabla f(\theta_1, \phi_{h,k}) - \nabla f(\theta_2, \phi_{h,k})) \cdot \nabla f(\theta_1, \phi_{h,k})^\top\right\|_2 + \left\|\nabla f(\theta_2, \phi_{h,k}) \cdot (\nabla f(\theta_1, \phi_{h,k})^\top - \nabla f(\theta_2, \phi_{h,k})^\top)\right\|_2\right]$$

$$\leq \frac{2\kappa_1 K}{\lambda^2}\kappa_2 \|\theta_1 - \theta_2\|_2 = \frac{2\kappa_1\kappa_2 K}{\lambda^2}\|\theta_1 - \theta_2\|_2.$$

All the above imply

$$|H(V_1,\theta_1) - H(V_2,\theta_2)| \leq \sqrt{|h(V_1,\theta_1)^\top G_h(\theta_1)^{-1}h(V_1,\theta_1) - h(V_2,\theta_2)^\top G_h(\theta_2)^{-1}h(V_2,\theta_2)|}$$

$$\leq \sqrt{\|h(V_1,\theta_1) - h(V_2,\theta_2)\|_2 \cdot \frac{1}{\lambda} \cdot 2KH\kappa_1} + \sqrt{2KH\kappa_1 \cdot \|G_h(\theta_1)^{-1} - G_h(\theta_2)^{-1}\|_2 \cdot 2KH\kappa_1}$$

$$+ \sqrt{2KH\kappa_1 \cdot \frac{1}{\lambda} \cdot \|h(V_1,\theta_1) - h(V_2,\theta_2)\|_2}$$

$$\leq 2\sqrt{K(2H\kappa_2\|\theta_1 - \theta_2\|_2 + 2\kappa_1\|V_1 - V_2\|_\infty) \cdot \frac{1}{\lambda} \cdot 2KH\kappa_1} + \sqrt{2KH\kappa_1 \cdot \frac{2\kappa_1\kappa_2 K}{\lambda^2}\|\theta_1 - \theta_2\|_2 \cdot 2KH\kappa_1}$$

$$\leq \left(4\sqrt{K^3 H^2 \kappa_1\kappa_2\frac{1}{\lambda}} + \sqrt{8K^3 H^2 \kappa_1^3\kappa_2\frac{1}{\lambda^2}}\right)\sqrt{\|\theta_1 - \theta_2\|_2} + 4\sqrt{K^3\kappa_1^2 H\frac{1}{\lambda}\|V_1 - V_2\|_\infty}$$

Then a ϵ-covering net of $\{H(V,\theta)\}$ can be constructed by the union of $\frac{\epsilon^2}{4\left(4\sqrt{K^3H^2\kappa_1\kappa_2\frac{1}{\lambda}}+\sqrt{8K^3H^2\kappa_1^3\kappa_2\frac{1}{\lambda^2}}\right)^2}$-covering net of $\{\theta \in \Theta\}$ and $\frac{\epsilon^2}{4(4\sqrt{K^3\kappa_1^2H\frac{1}{\lambda}})^2}$-covering net of $\mathcal{V}$ in Lemma L.9. The covering number $\mathcal{N}_\epsilon$ satisfies

$$
\begin{aligned}
\log \mathcal{N}_\epsilon \leq & d\log\left(1+\frac{8C_\Theta\left(4\sqrt{K^3H^2\kappa_1\kappa_2\frac{1}{\lambda}}+\sqrt{8K^3H^2\kappa_1^3\kappa_2\frac{1}{\lambda^2}}\right)^2}{\epsilon^2}\right) \\
& +d\log\left(1+\frac{8C_\Theta(\kappa_1\sqrt{C_\Theta}+2\sqrt{B\kappa_1\kappa_2})^2}{\frac{\epsilon^4}{16(4\sqrt{K^3\kappa_1^2H\frac{1}{\lambda}})^4}}\right)+d^2\log\left(1+\frac{8\sqrt{d}B\kappa_1^2}{\frac{\epsilon^4}{16(4\sqrt{K^3\kappa_1^2H\frac{1}{\lambda}})^4}}\right).
\end{aligned}
$$

Step3. First note by definition in Step2

$$
\left\|\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-f(\theta_{\mathbb{T}V^\star_{h+1}},\phi_{h,k})-\widehat{V}_{h+1}(s_{h+1}^k)+V^\star_{h+1}(s_{h+1}^k)\right)\cdot\nabla_\theta^\top f(\widehat{\theta}_h,\phi_{h,k})\right\|_{\Sigma_h^{-1}}=H(\widehat{V}_{h+1},\widehat{\theta}_h)
$$

and with probability $1-\delta$

$$
\begin{aligned}
|\eta_k(\widehat{V}_{h+1})|=&|f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-f(\theta_{\mathbb{T}V^\star_{h+1}},\phi_{h,k})-\widehat{V}_{h+1}(s_{h+1}^k)+V^\star_{h+1}(s_{h+1}^k)| \\
\leq&\kappa_1\cdot\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}-\theta_h^\star\right\|_2+\left\|\widehat{V}_{h+1}-V^\star_{h+1}\right\|_\infty \\
\leq&\kappa_1\sqrt{\frac{36H^2(\log(H/\delta)+C_{d,\log K})+2\lambda C_\Theta^2}{\kappa K}}+C\left(\kappa_1H^2\sqrt{\frac{d^2}{\kappa K}}\right)=\widetilde{O}\left(\kappa_1H^2\sqrt{\frac{d^2}{\kappa K}}\right)
\end{aligned}
$$

(9)

where the second inequality uses $\theta_{\mathbb{T}V^\star_{h+1}}=\theta_h^\star$ and the third inequality uses Theorem G.2 and Theorem G.3. The last equal sign is due to $C_{d,\log K}\leq\widetilde{O}(d^2)$ (recall Lemma G.1).

Now choosing $\epsilon=O(1/K)$ in Step2 and union bound over both equation 9 and covering number in Step2, we obtain with probability $1-\delta$,

$$
H(\widehat{V}_{h+1},\widehat{\theta}_h)=\left\|\sum_{k=1}^{K}x_k(\widehat{\theta}_h)\eta_k(\widehat{V}_{h+1})\right\|_{G_h(\widehat{\theta}_h)^{-1}}\leq\widetilde{O}(\kappa_1H^2\sqrt{\frac{d^2}{\kappa K}})\sqrt{d+d^2}=\widetilde{O}(\frac{\kappa_1H^2d^2}{\sqrt{\kappa K}})
$$

(10)

where we absorb all the Polylog terms. Meanwhile, by Lemma L.5 with probability $1-\delta$,

$$
\left\|\nabla f(\widehat{\theta}_h,\phi_{s,a})\right\|_{\Sigma_h^{-1}}\leq\frac{2\kappa_1}{\sqrt{\kappa K}}+O(\frac{1}{K}).
$$

(11)

Finally, by equation 10 and equation 11 and a union bound, we have with probability $1-\delta$,

$$
\begin{aligned}
|I_2|:=&\left|\nabla f(\widehat{\theta}_h,\phi(s,a))\Sigma_h^{-1}[\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-f(\theta_{\mathbb{T}V^\star_{h+1}},\phi_{h,k})-\widehat{V}_{h+1}(s_{h+1}^k)+V^\star_{h+1}(s_{h+1}^k)\right)\cdot\nabla_\theta^\top f(\widehat{\theta}_h,\phi_{h,k})]\right| \\
\leq&\left\|\nabla f(\widehat{\theta}_h,\phi_{s,a})\right\|_{\Sigma_h^{-1}}\left\|\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi_{h,k})-f(\theta_{\mathbb{T}V^\star_{h+1}},\phi_{h,k})-\widehat{V}_{h+1}(s_{h+1}^k)+V^\star_{h+1}(s_{h+1}^k)\right)\cdot\nabla_\theta^\top f(\widehat{\theta}_h,\phi_{h,k})\right\|_{\Sigma_h^{-1}} \\
=&\left\|\nabla f(\widehat{\theta}_h,\phi_{s,a})\right\|_{\Sigma_h^{-1}}\cdot H(\widehat{V}_{h+1},\widehat{\theta}_h)\leq\left(\frac{2\kappa_1}{\sqrt{\kappa K}}+O(\frac{1}{K})\right)\cdot\widetilde{O}(\frac{\kappa_1H^2d^2}{\sqrt{\kappa K}})=\widetilde{O}(\frac{\kappa_1^2H^2d^2}{\kappa K})+\widetilde{O}(\frac{1}{K^{3/2}})
\end{aligned}
$$

where the first inequality is Cauchy–Schwarz inequality. $\qquad\square$

E.4 BOUNDING THE MAIN TERM I_1

In this section, we bound the dominate term

$$
I_1:=\nabla f(\widehat{\theta}_h,\phi(s,a))\Sigma_h^{-1}[\sum_{k=1}^{K}\left(f(\theta_{\mathbb{T}V^\star_{h+1}},\phi_{h,k})-r_{h,k}-V^\star_{h+1}(s_{h+1}^k)\right)\cdot\nabla_\theta^\top f(\widehat{\theta}_h,\phi_{h,k})].
$$

First of all, by Cauchy–Schwarz inequality, we have

$$
|I_1| \leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot \left\| \sum_{k=1}^K \left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) \right\|_{\Sigma_h^{-1}}.
$$
(12)

Then we have the following Lemma to bound I_1.

Lemma E.7. *With probability $1 - \delta$,*

$$
|I_1| \leq 4Hd \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot C_{\delta, \log K} + \widetilde{O}\left(\frac{\kappa_1}{\sqrt{\kappa}K} \right),
$$

where $C_{\delta, \log K}$ only contains Polylog terms.

Proof of Lemma E.7. **Step1.** Let the fixed $\theta \in \Theta$ be arbitrary and define $x_k(\theta) = \nabla_\theta f(\theta, \phi_{h,k})$. Next, define $G_h(\theta) = \sum_{k=1}^K \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top + \lambda I_d$, then $\|x_k\|_2 \leq \kappa_1$. Also denote $\eta_k := f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k)$, then $\mathbb{E}[\eta_k | s_h^k, a_h^k] = 0$ and $|\eta_k| \leq H$. Now by self-normalized Hoeffding's inequality (Lemma L.3), with probability $1 - \delta$ (recall $t := K$ in Lemma L.3),

$$
\left\| \sum_{k=1}^K x_k(\theta) \eta_k \right\|_{G_h(\theta)^{-1}} \leq 2H \sqrt{d \log\left(\frac{\lambda + K\kappa_1}{\lambda \delta} \right)}.
$$

Step2. Define $h(\theta) := \sum_{k=1}^K x_k(\theta) \eta_k$ and $H(\theta) := \left\| \sum_{k=1}^K x_k(\theta) \eta_k \right\|_{G_h(\theta)^{-1}}$, then note by definition $|\eta_k| \leq H$, which implies $\|h(\theta)\|_2 \leq KH\kappa_1$ and by $x_k(\theta_1) - x_k(\theta_2) = \nabla_{\theta\theta}^2 f(\xi, \phi) \cdot (\theta_1 - \theta_2)$,

$$
\|h(\theta_1) - h(\theta_2)\|_2 \leq K \max_k \left(H \|x_k(\theta_1) - x_k(\theta_2)\|_2 \right) \leq HK\kappa_2 \|\theta_1 - \theta_2\|_2.
$$

Furthermore,

$$
\left\| G_h(\theta_1)^{-1} - G_h(\theta_2)^{-1} \right\|_2 \leq \left\| G_h(\theta_1)^{-1} \right\|_2 \left\| G_h(\theta_1) - G_h(\theta_2) \right\|_2 \left\| G_h(\theta_2)^{-1} \right\|_2
$$
$$
\leq \frac{1}{\lambda^2} K \sup_k \left\| \nabla f(\theta_1, \phi_{h,k}) \cdot \nabla f(\theta_1, \phi_{h,k})^\top - \nabla f(\theta_2, \phi_{h,k}) \cdot \nabla f(\theta_2, \phi_{h,k})^\top \right\|_2
$$
$$
\leq \frac{2\kappa_1 K}{\lambda^2} \kappa_2 \|\theta_1 - \theta_2\|_2 = \frac{2\kappa_1\kappa_2 K}{\lambda^2} \|\theta_1 - \theta_2\|_2.
$$

All the above imply

$$
|H(\theta_1) - H(\theta_2)| \leq \sqrt{\left| h(\theta_1)^\top G_h(\theta_1)^{-1} h(\theta_1) - h(\theta_2)^\top G_h(\theta_2)^{-1} h(\theta_2) \right|}
$$
$$
\leq \sqrt{\|h(\theta_1) - h(\theta_2)\|_2 \cdot \frac{1}{\lambda} \cdot KH\kappa_1} + \sqrt{KH\kappa_1 \cdot \left\| G_h(\theta_1)^{-1} - G_h(\theta_2)^{-1} \right\|_2 \cdot KH\kappa_1}
$$
$$
+ \sqrt{KH\kappa_1 \cdot \frac{1}{\lambda} \cdot \|h(\theta_1) - h(\theta_2)\|_2}
$$
$$
\leq 2\sqrt{KH\kappa_2 \|\theta_1 - \theta_2\|_2 \cdot \frac{1}{\lambda} \cdot KH\kappa_1} + \sqrt{KH\kappa_1 \cdot \frac{2\kappa_1\kappa_2 K}{\lambda^2} \|\theta_1 - \theta_2\|_2 \cdot KH\kappa_1}
$$
$$
\leq \left(\sqrt{4K^2H^2\kappa_1\kappa_2/\lambda} + \sqrt{2K^3H^2\kappa_1^3\kappa_2/\lambda^2} \right) \sqrt{\|\theta_1 - \theta_2\|_2}
$$

Then a ϵ-covering net of $\{H(\theta)\}$ can be constructed by the union of $\frac{\epsilon^2}{\left(\sqrt{4K^2H^2\kappa_1\kappa_2/\lambda} + \sqrt{2K^3H^2\kappa_1^3\kappa_2/\lambda^2} \right)^2}$-covering net of $\{\theta \in \Theta\}$. By Lemma L.8, the covering number $\mathcal{N}_\epsilon$ satisfies

$$
\log \mathcal{N}_\epsilon \leq d \log \left(1 + \frac{2C_\Theta \left(\sqrt{4K^2H^2\kappa_1\kappa_2/\lambda} + \sqrt{2K^3H^2\kappa_1^3\kappa_2/\lambda^2} \right)^2}{\epsilon^2} \right) = \widetilde{O}(d)
$$

Step3. First note by definition in Step2

$$\left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) \right\|_{\Sigma_h^{-1}} = H(\widehat{\theta}_h)$$

Now choosing $\epsilon = O(1/K)$ in Step2 and union bound over the covering number in Step2, we obtain with probability $1 - \delta$,

$$H(\widehat{\theta}_h) = \left\| \sum_{k=1}^{K} x_k(\widehat{\theta}_h)\eta_k \right\|_{G_h(\widehat{\theta}_h)^{-1}} \leq 2H\sqrt{d \left[\log\left(\frac{\lambda + K\kappa_1}{\lambda\delta} \right) + \widetilde{O}(d) \right]} + O(\frac{1}{K}). \qquad (13)$$

where we absorb all the Polylog terms. Combing above with equation 12, we obtain with probability $1 - \delta$,

$$|I_1| \leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot \left\| \sum_{k=1}^{K} \left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) \right\|_{\Sigma_h^{-1}}$$

$$\leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot \left(2H\sqrt{d \left[\log\left(\frac{\lambda + K\kappa_1}{\lambda\delta} \right) + \widetilde{O}(d) \right]} + O(\frac{1}{K}) \right)$$

$$\leq 4Hd \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot C_{\delta,\log K} + \widetilde{O}(\frac{\kappa_1}{\sqrt{\kappa}K}),$$

where $C_{\delta,\log K}$ only contains Polylog terms. $\qquad \square$

E.5   ANALYZING Hot_2 IN EQUATION 8

Lemma E.8. *Recall* $\text{Hot}_2 := \nabla f(\widehat{\theta}_h, \phi(s,a))\Sigma_h^{-1} \left[\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) + \lambda\theta_{\mathbb{T}\widehat{V}_{h+1}} \right]$. *If* $K \geq$ $\max \left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K^3}{\kappa\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$, *then with probability* $1 - \delta$,

$$\left| \nabla f(\widehat{\theta}_h, \phi(s,a))\Sigma_h^{-1} \left[\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) + \lambda\theta_{\mathbb{T}\widehat{V}_{h+1}} \right] \right| \leq \widetilde{O}\left(\frac{\kappa_2 \max(\frac{\kappa_1}{\kappa}, \frac{1}{\sqrt{\kappa}})d^2 H^2 + \frac{d^2 H^3\kappa_3 + \lambda\kappa_1 C_\Theta}{\kappa}}{K} \right)$$

where $\widetilde{O}$ absorbs all the constants and Polylog terms.

Proof of Lemma E.8. **Step1:** we first show with probability $1 - \delta$

$$\left| \nabla f(\widehat{\theta}_h, \phi(s,a))\Sigma_h^{-1} \widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right| \leq \widetilde{O}(\frac{1}{K}).$$

Recall by plug in $\theta_{\mathbb{T}\widehat{V}_{h+1}}$ in equation 4, we have

$$Z_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}|\widehat{V}_{h+1}) - Z_h(\widehat{\theta}_h|\widehat{V}_{h+1}) = \frac{\partial}{\partial\theta} Z_h(\widehat{\theta}_h|\widehat{V}_{h+1})(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}), \qquad (14)$$

and by second-order Taylor's Theorem we have

$$\left\| R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_2 = \left\| Z_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}|\widehat{V}_{h+1}) - Z_h(\widehat{\theta}_h|\widehat{V}_{h+1}) - \frac{\partial}{\partial\theta} Z_h(\widehat{\theta}_h|\widehat{V}_{h+1})(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) \right\|_2$$

$$= \frac{1}{2} \left\| (\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h)^\top \frac{\partial^2}{\partial\theta\partial\theta} Z_h(\xi|\widehat{V}_{h+1})(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h) \right\|_2 \qquad (15)$$

$$\leq \frac{1}{2}\kappa_{z_2} \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right\|_2^2$$

Note

$$\frac{\partial^2}{\partial\theta\partial\theta} Z_h(\theta|\widehat{V}_{h+1}) \bigg|_{\theta=\xi} = \frac{\partial}{\partial\theta}\Sigma_h^s = \sum_{k=1}^{K} \frac{\partial}{\partial\theta} \left\{ \left(f(\xi, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\xi, \phi_{h,k}) \right\}$$

$$+ \sum_{k=1}^{K} \frac{\partial}{\partial\theta} \left(\nabla_\theta f(\xi, \phi_{h,k})\nabla_\theta^\top f(\xi, \phi_{h,k}) + \lambda I_d \right)$$

$$(16)$$

Therefore, we can bound κ_{z_2} with $\kappa_{z_2} \leq (H\kappa_3 + 3\kappa_1\kappa_2)K$ and this implies with probability $1 - \delta/2$,

$$
\begin{aligned}
\left\| R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_2 &\leq \frac{1}{2}\kappa_{z_2} \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right\|_2^2 \leq \frac{1}{2}(H\kappa_3 + 3\kappa_1\kappa_2)K \cdot \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right\|_2^2 \\
&\leq \frac{1}{2}(H\kappa_3 + 3\kappa_1\kappa_2)K \cdot \frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K} \\
&\leq \widetilde{O}((H\kappa_3 + 3\kappa_1\kappa_2)H^2 d^2/\kappa).
\end{aligned}
$$

And by Corollary E.3 with probability $1 - \delta/2$,

$$
\left\| \Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_2 \leq \widetilde{O}(1),
$$

Therefore, by Lemma L.5 and a union bound with probability $1 - \delta$,

$$
\begin{aligned}
|\nabla f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})| &= \left| \nabla f(\widehat{\theta}_h, \phi(s,a))^\top \Sigma_h^{-1} \left(\Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right) \right| \\
&\leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \left\| \Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_{\Sigma_h^{-1}} \\
&\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \left\| \Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_{\Sigma_h^{-1}} \\
&\leq \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \left(\frac{C}{\sqrt{K}} + O(\frac{1}{K}) \right) = \widetilde{O}\left(\frac{\kappa_2 \max(\frac{\kappa_1}{\kappa}, \frac{1}{\sqrt{\kappa}})d^2 H^2 + \frac{d^2 H^3 \kappa_3}{\kappa}}{K} \right)
\end{aligned}
$$

where $\widetilde{O}$ absorbs all the constants and Polylog terms. Here the last inequality uses bound for $\left\| R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_2$ and $\left\| \Delta_{\Sigma_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) \right\|_2$.

Step2: By Lemma L.5, with probability $1 - \delta$,

$$
\begin{aligned}
\left| \nabla f(\widehat{\theta}_h, \phi(s,a))\Sigma_h^{-1} \lambda \theta_{\mathbb{T}\widehat{V}_{h+1}} \right| &\leq \lambda \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_{\Sigma_h^{-1}} \\
&\leq \lambda \left(\frac{2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) \cdot \left(\frac{2C_\Theta}{\sqrt{\kappa K}} + O(\frac{1}{K}) \right) = \frac{4\lambda \kappa_1 C_\Theta}{\kappa K} + O(\frac{1}{K^{\frac{3}{2}}})
\end{aligned}
$$

$\square$

F   PROOF OF THEOREM 3.2

Now we are ready to prove Theorem 3.2. In particular, we prove the first part. Also, recall that we consider the exact Bellman completeness ($\epsilon_{\mathcal{F}} = 0$).

F.1   THE FIRST PART

Proof of Theorem 3.2 (first part). First of all, from the previous calculation (3), (8), we have

$$
\begin{aligned}
\left| \mathcal{P}_h \widehat{V}_{h+1}(s,a) - \widehat{\mathcal{P}}_h \widehat{V}_{h+1}(s,a) \right| &\leq \left| \nabla f(\widehat{\theta}_h, \phi(s,a)) \left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right) \right| + |\mathrm{Hot}_{h,1}| \\
&\leq |I_1| + |I_2| + |I_3| + |\mathrm{Hot}_{h,2}| + |\mathrm{Hot}_{h,1}|
\end{aligned}
$$

Now by Lemma E.5, Lemma E.6, Lemma E.7, Lemma E.8 and Lemma E.1 (and a union bound), with probability $1 - \delta$,

$$
\begin{aligned}
|I_3| \leq & \widetilde{O}(\sqrt{\frac{d^3 H^2 \kappa_2^2 \kappa_1^2}{\kappa^3}}) \frac{1}{K}, \\
|I_2| \leq & \widetilde{O}(\frac{\kappa_1^2 H^2 d^2}{\kappa K}) + \widetilde{O}(\frac{1}{K^{3/2}}), \\
|I_1| \leq & 4Hd \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot C_{\delta, \log K} + \widetilde{O}(\frac{\kappa_1}{\sqrt{\kappa} K}), \\
|\mathrm{Hot}_{2,h}| \leq & \widetilde{O}\left(\frac{\kappa_2 \max(\frac{\kappa_1}{\kappa}, \frac{1}{\sqrt{\kappa}}) d^2 H^2 + \frac{d^2 H^3 \kappa_3 + \lambda \kappa_1 C_\Theta}{\kappa}}{K} \right), \\
|\mathrm{Hot}_{1,h}| \leq & \widetilde{O}(\frac{H^2 \kappa_2 d^2}{\kappa}) \frac{1}{K}.
\end{aligned}
$$

Finally, Plug the above into Lemma D.3, by a union bound over all $h \in [H]$, we have with probability $1 - \delta$, for any policy π,

$$
\begin{aligned}
v^\pi - v^{\widehat{\pi}} \leq & \sum_{h=1}^{H} 2 \cdot \mathbb{E}_\pi \left[|I_1| + |I_2| + |I_3| + |\mathrm{Hot}_{h,2}| + |\mathrm{Hot}_{h,1}| \right] \\
\leq & \sum_{h=1}^{H} 8dH \mathbb{E}_\pi \left[\sqrt{\nabla^\top f(\widehat{\theta}_h, \phi(s_h, a_h)) \Sigma_h^{-1} \nabla f(\widehat{\theta}_h, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\mathrm{hot}}}{K}).
\end{aligned}
$$

where $\iota = C_{\delta, \log K}$ only contains Polylog terms and

$$
C_{\mathrm{hot}} = \frac{\kappa_1 H}{\sqrt{\kappa}} + \frac{\kappa_1^2 H^3 d^2}{\kappa} + \sqrt{\frac{d^3 H^4 \kappa_2^2 \kappa_1^2}{\kappa^3}} + \kappa_2 \max(\frac{\kappa_1}{\kappa}, \frac{1}{\sqrt{\kappa}}) d^2 H^3 + \frac{d^2 H^4 \kappa_3 + \lambda \kappa_1 C_\Theta}{\kappa} + \frac{H^3 \kappa_2 d^2}{\kappa}
$$

$\square$

F.2 THE SECOND PART

Next we prove the second part of Theorem 3.2.

Proof of Theorem 3.2 (second part). **Step1.** Choose $\pi = \pi^\star$ in the first part, we have

$$
0 \leq v^{\pi^\star} - v^{\widehat{\pi}} \leq \sum_{h=1}^{H} 8dH \cdot \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\widehat{\theta}_h, \phi(s_h, a_h)) \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\mathrm{hot}}}{K}),
$$

Next, by the triangular inequality of the norm to obtain

$$
\begin{aligned}
& \left| \left\| \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h)) \right\|_{\Sigma_h^{-1}} - \left\| \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h)) \right\|_{\Sigma_h^{-1}} \right| \\
\leq & \left\| \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h)) - \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h)) \right\|_{\Sigma_h^{-1}} \\
= & \left\| \nabla_{\theta\theta}^2 f(\xi, \phi(s_h, a_h)) \cdot \left(\widehat{\theta}_h - \theta_h^\star \right) \right\|_{\Sigma_h^{-1}},
\end{aligned}
$$

since with probability $1 - \delta$,

$$
\left\| \nabla_{\theta\theta}^2 f(\xi, \phi(s_h, a_h)) \cdot \left(\widehat{\theta}_h - \theta_h^\star \right) \right\|_2 \leq \kappa_2 \left\| \widehat{\theta}_h - \theta_h^\star \right\|_2 \leq \widetilde{O}\left(\frac{\kappa_1 \kappa_2 H^2 d}{\kappa} \sqrt{\frac{1}{K}} \right),
$$

where the last inequality uses part three of Theorem G.3. Then by a union bound and Lemma L.5,

$$
\left\| \nabla_{\theta\theta}^2 f(\xi, \phi(s_h, a_h)) \cdot \left(\widehat{\theta}_h - \theta_h^\star \right) \right\|_{\Sigma_h^{-1}} \leq \widetilde{O}\left(\frac{\kappa_1 \kappa_2 H^2 d}{\kappa^{3/2}} \cdot \frac{1}{K} \right).
$$

Step2. Next, we show with probability $1 - \delta$,

$$\left\|\nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))\right\|_{\Sigma_h^{-1}} \le 2 \left\|\nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))\right\|_{\Sigma_h^{\star-1}}.$$

First of all,

$$\left\|\frac{1}{K}\Sigma_h - \frac{1}{K}\Sigma_h^\star\right\|_2 = \left\|\frac{1}{K}\left(\sum_{k=1}^K \nabla f(\widehat{\theta}_h, \phi(s,a))\nabla f(\widehat{\theta}_h, \phi(s,a))^\top - \nabla f(\theta_h^\star, \phi(s,a))\nabla f(\theta_h^\star, \phi(s,a))^\top\right)\right\|_2$$

$$\le \sup_{s,a}\left(\left\|\left(\nabla f(\widehat{\theta}_h, \phi(s,a)) - \nabla f(\theta_h^\star, \phi(s,a))\right)\nabla f(\widehat{\theta}_h, \phi(s,a))\right\|_2\right.$$

$$+ \left.\left\|\left(\nabla f(\widehat{\theta}_h, \phi(s,a)) - \nabla f(\theta_h^\star, \phi(s,a))\right)\nabla f(\widehat{\theta}_h, \phi(s,a))\right\|_2\right)$$

$$\le 2\kappa_2\kappa_1\left\|\widehat{\theta}_h - \theta_h^\star\right\|_2 \le \widetilde{O}\left(\frac{\kappa_2\kappa_1^2 H^2 d}{\kappa}\sqrt{\frac{1}{K}}\right)$$

Second, by Lemma L.6 with probability $1 - \delta$

$$\left\|\frac{\Sigma_h^\star}{K} - \mathbb{E}_\mu[\nabla_\theta f(\theta_h^\star, \phi)\nabla_\theta f(\theta_h^\star, \phi)^\top] - \frac{\lambda}{K}\right\| \le \frac{4\sqrt{2}\kappa_1^2}{\sqrt{K}}\left(\log\frac{2d}{\delta}\right)^{1/2}$$

This implies

$$\left\|\frac{\Sigma_h^\star}{K}\right\| \le \left\|\mathbb{E}_\mu[\nabla_\theta f(\theta_h^\star, \phi)\nabla_\theta f(\theta_h^\star, \phi)^\top]\right\| + \frac{\lambda}{K} + \frac{4\sqrt{2}\kappa_1^2}{\sqrt{K}}\left(\log\frac{2d}{\delta}\right)^{1/2}$$

$$\le \kappa_1^2 + \lambda + 4\sqrt{2}\kappa_1^2\left(\log\frac{2d}{\delta}\right)^{1/2}$$

and also by *Weyl's spectrum theorem* and under the condition $K \ge \frac{128\kappa_1^4\log(2d/\delta)}{\kappa^2}$, with probability $1 - \delta$

$$\lambda_{\min}(\frac{\Sigma_h^\star}{K}) \ge \lambda_{\min}\left(\mathbb{E}_\mu[\nabla_\theta f(\theta_h^\star, \phi)\nabla_\theta f(\theta_h^\star, \phi)^\top]\right) + \frac{\lambda}{K} - \frac{4\sqrt{2}\kappa_1^2}{\sqrt{K}}\left(\log\frac{2d}{\delta}\right)^{1/2}$$

$$\ge \kappa + \frac{\lambda}{K} - \frac{4\sqrt{2}\kappa_1^2}{\sqrt{K}}\left(\log\frac{2d}{\delta}\right)^{1/2} \ge \frac{\kappa}{2}$$

then $\left\|(\frac{\Sigma_h^\star}{K})^{-1}\right\| \le \frac{2}{\kappa}$. Similarly, with probability $1 - \delta$, $\left\|(\frac{\Sigma_h}{K})^{-1}\right\| \le \frac{2}{\kappa}$. Then by Lemma L.7,

$$\left\|\nabla_\theta f(\theta_h^\star, \phi(s,a))\right\|_{K\Sigma_h^{-1}} \le \left[1 + \sqrt{\left\|K\Sigma_h^{\star-1}\right\|\left\|\Sigma_h^\star/K\right\| \cdot \left\|K\Sigma_h^{-1}\right\| \cdot \left\|\Sigma_h/K - \Sigma_h^\star/K\right\|}\right] \cdot \left\|\nabla_\theta f(\theta_h^\star, \phi(s,a))\right\|_{K\Sigma_h^{\star-1}}$$

$$\le \left[1 + \sqrt{\frac{4}{\kappa^2}O(\kappa_1^2 + \lambda)\widetilde{O}\left(\frac{\kappa_2\kappa_1^2 H^2 d}{\kappa}\sqrt{\frac{1}{K}}\right)}\right] \cdot \left\|\nabla_\theta f(\theta_h^\star, \phi(s,a))\right\|_{K\Sigma_h^{\star-1}}$$

$$\le 2\left\|\nabla_\theta f(\theta_h^\star, \phi(s,a))\right\|_{K\Sigma_h^{\star-1}}$$

as long as $K \ge \widetilde{O}(\frac{(\kappa_1^2 + \lambda)^2\kappa_2^2\kappa_1^2 H^4 d^2}{\kappa^6})$. The above is equivalently to

$$\left\|\nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))\right\|_{\Sigma_h^{-1}} \le 2\left\|\nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))\right\|_{\Sigma_h^{\star-1}}.$$

Combining Step1, Step2 and a union bound, we have with probability $1 - \delta$,

$$
\begin{aligned}
0 \le v^{\pi^\star} - v^{\widehat{\pi}} &\le \sum_{h=1}^{H} 8dH \cdot \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\widehat{\theta}_h, \phi(s_h, a_h)) \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\mathrm{hot}}}{K}) \\
&\le \sum_{h=1}^{H} 8dH \cdot \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi(s_h, a_h)) \Sigma_h^{-1} \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\mathrm{hot}}}{K}) + \widetilde{O}\left(\frac{\kappa_1 \kappa_2 H^4 d^2}{\kappa^{3/2}} \cdot \frac{1}{K} \right) \\
&\le \sum_{h=1}^{H} 16dH \cdot \mathbb{E}_{\pi^\star} \left[\sqrt{\nabla_\theta^\top f(\theta_h^\star, \phi(s_h, a_h)) \Sigma_h^{\star -1} \nabla_\theta f(\theta_h^\star, \phi(s_h, a_h))} \right] \cdot \iota + \widetilde{O}(\frac{C_{\mathrm{hot}}'}{K})
\end{aligned}
$$

where $C_{\mathrm{hot}}' = C_{\mathrm{hot}} + \frac{\kappa_1 \kappa_2 H^4 d^2}{\kappa^{3/2}}$.

□

G PROVABLE EFFICIENCY BY REDUCTION TO GENERAL FUNCTION APPROXIMATION

In this section, we bound the accuracy of the parameter difference $\left\| \widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2$ via a reduction to General Function Approximation scheme in Chen and Jiang (2019).

Recall the objective

$$
\ell_h(\theta) := \frac{1}{K} \sum_{k=1}^{K} \left[f\left(\theta, \phi(s_h^k, a_h^k)\right) - r(s_h^k, a_h^k) - \widehat{V}_{h+1}\left(s_{h+1}^k\right) \right]^2 + \frac{\lambda}{K} \cdot \|\theta\|_2^2 \tag{17}
$$

Then by definition, $\widehat{\theta}_h := \arg\min_{\theta \in \Theta} \ell_h(\theta)$ and $\theta_{\mathbb{T}\widehat{V}_{h+1}}$ satisfies $f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi) = \mathcal{P}_h \widehat{V}_{h+1} + \delta_{\widehat{V}_{h+1}}$. Therefore, in this case, we have the following lemma:

Lemma G.1. *Fix $h \in [H]$. With probability $1 - \delta$,*

$$
\mathbb{E}_\mu[\ell_h(\widehat{\theta}_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] \le \frac{36H^2(\log(1/\delta) + C_{d,\log K}) + \lambda C_\Theta^2}{K} + \sqrt{\frac{16H^3 \epsilon_\mathcal{F}(\log(1/\delta) + C_{d,\log K})}{K}} + 4H\epsilon_\mathcal{F}.
$$

where the expectation over μ is taken w.r.t. $(s_h^k, a_h^k, s_{h+1}^k)$ $k = 1, ..., K$ *only (i.e., first compute $\mathbb{E}_\mu[\ell_h(\theta)]$ for a fixed θ, then plug-in either $\widehat{\theta}_{h+1}$ or $\theta_{\mathbb{T}\widehat{V}_{h+1}}$).* *Here* $C_{d,\log(K)} := d\log(1 + 24C_\Theta(H+1)\kappa_1 K) + d\log\left(1 + 288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{\kappa_1\kappa_2/\lambda})^2 K^2\right) + d^2\log\left(1 + 288H^2\sqrt{d}\kappa_1^2 K^2/\lambda\right)$.

Proof of Lemma G.1. **Step1:** we first prove the case where $\lambda = 0$.

Indeed, fix $h \in [H]$ and any function $V(\cdot) \in \mathbb{R}^\mathcal{S}$. Similarly, define $f_V(s, a) := f(\theta_{\mathbb{T}V}, \phi) = \mathcal{P}_h V + \delta_V$. For any fixed $\theta \in \Theta$, denote $g(s, a) = f(\theta, \phi(s, a))$. Then define[10]

$$
X(g, V, f_V) := (g(s, a) - r - V(s'))^2 - (f_V(s, a) - r - V(s'))^2.
$$

Since all episodes are independent of each other, $X_k(g, V, f_V) := X(g(s_h^k, a_h^k), V(s_{h+1}^k), f_V(s_h^k, a_h^k))$ are independent r.v.s and it holds

$$
\frac{1}{K} \sum_{k=1}^{K} X_k(g, V, f_V) = \ell(g) - \ell(f_V). \tag{18}
$$

---

[10]We abuse the notation here to use either $X(g, V, f_V)$ or $X(\theta, V, f_V)$. They mean the same quantity.

Next, the variance of X is bounded by:

$$\text{Var}[X(g, V, f_V)] \leq \mathbb{E}_\mu[X(g, f, f_V)^2]$$

$$=\mathbb{E}_\mu\left[\left((g(s_h, a_h) - r_h - V(s_{h+1}))^2 - (f_V(s_h, a_h) - r_h - V(s_{h+1}))^2\right)^2\right]$$

$$=\mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h))^2(g(s_h, a_h) + f_V(s_h, a_h) - 2r_h - 2V(s_{h+1}))^2\right]$$

$$\leq 4H^2 \cdot \mathbb{E}_\mu[(g(s_h, a_h) - f_V(s_h, a_h))^2]$$

$$\leq 4H^2 \cdot \mathbb{E}_\mu\left[(g(s_h, a_h) - r_h - V(s_{h+1}))^2 - (f_V(s_h, a_h) - r_h - V(s_{h+1}))^2\right] + 8H^3\epsilon_\mathcal{F} \quad (*)$$

$$=4H^2 \cdot \mathbb{E}_\mu[X(g, f, f_V)] + 8H^3\epsilon_\mathcal{F}$$

where the step $(*)$ comes from

$$\mathbb{E}_\mu\left[(g(s_h, a_h) - r_h - V(s_{h+1}))^2 - (f_V(s_h, a_h) - r_h - V(s_{h+1}))^2\right]$$

$$=\mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h)) \cdot (g(s_h, a_h) + f_V(s_h, a_h) - 2r_h - 2V(s_{h+1}))\right]$$

$$=\mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h)) \cdot (g(s_h, a_h) - f_V(s_h, a_h) + 2f_V(s_h, a_h) - 2r_h - 2V(s_{h+1}))\right]$$

$$=\mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h))^2\right] + \mathbb{E}_\mu\left[2(g(s_h, a_h) - f_V(s_h, a_h))\mathbb{E}_{P_h}[f_V(s_h, a_h) - r_h - V(s_{h+1}) \mid s_h, a_h]\right]$$

$$\geq \mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h))^2\right] - 2H\|\delta_V\|_\infty \geq \mathbb{E}_\mu\left[(g(s_h, a_h) - f_V(s_h, a_h))^2\right] - 2H\epsilon_\mathcal{F}$$

$$(19)$$

where the last step uses law of total expectation and the definition of f_V.

Therefore, by Bernstein inequality, with probability $1 - \delta$,

$$\mathbb{E}_\mu[X(g, f, f_V)] - \frac{1}{K}\sum_{k=1}^{K} X_k(g, f, f_V)$$

$$\leq \sqrt{\frac{2\text{Var}[X(g, f, f_V)]\log(1/\delta)}{K}} + \frac{4H^2\log(1/\delta)}{3K}$$

$$\leq \sqrt{\frac{8H^2\mathbb{E}_\mu[X(g, f, f_V)]\log(1/\delta)}{K}} + \sqrt{\frac{16H^3\epsilon_\mathcal{F}\log(1/\delta)}{K}} + \frac{4H^2\log(1/\delta)}{3K}.$$

Now, if we choose $g(s, a) := f(\widehat{\theta}_h, \phi(s, a))$, then $\widehat{\theta}_h$ minimizes $\ell_h(\theta)$, therefore, it also minimizes $\frac{1}{K}\sum_{k=1}^{K} X_i(\theta, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})$ and this implies

$$\frac{1}{K}\sum_{k=1}^{K} X_k(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}) \leq \frac{1}{K}\sum_{k=1}^{K} X_k(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}) = 0.$$

Therefore, we obtain

$$\mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})] \leq \sqrt{\frac{8H^2 \cdot \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})]\log(1/\delta)}{K}} + \sqrt{\frac{16H^3\epsilon_\mathcal{F}\log(1/\delta)}{K}} + \frac{4H^2\log(1/\delta)}{3K}.$$

However, the above does not hold with probability $1 - \delta$ since $\widehat{\theta}_h$ and $\widehat{V}_{h+1} :=$ $\min\{\max_a f(\widehat{\theta}_{h+1}, \phi(\cdot, a)) - \sqrt{\nabla f(\widehat{\theta}_{h+1}, \phi(\cdot, a))^\top A \cdot \nabla f(\theta, \phi(\cdot, a))}, H\}$ (where A is certain symmetric matrix with bounded norm) depend on $\widehat{\theta}_h$ and $\widehat{\theta}_{h+1}$ which are data-dependent. Therefore, we need to further apply covering Lemma L.10 and choose $\epsilon = O(1/K)$ and a union bound to obtain with probability $1 - \delta$,

$$\mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})] \leq \sqrt{\frac{8H^2 \cdot \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})](\log(1/\delta) + C_{d,\log K})}{K}} + \frac{7H^2(\log(1/\delta) + C_{d,\log K})}{3K}$$

$$+ \sqrt{\frac{16H^3\epsilon_\mathcal{F}(\log(1/\delta) + C_{d,\log K})}{K}} + 4H\epsilon_\mathcal{F}$$

where $C_{d,\log(K)} := \log(1+24C_\Theta(H+1)\kappa_1 K)+d\log\left(1+288H^2C_\Theta(\kappa_1\sqrt{C_\Theta}+2\sqrt{\kappa_1\kappa_2/\lambda})^2K^2\right)+$ $d^2\log\left(1+288H^2\sqrt{d}\kappa_1^2K^2/\lambda\right)$.[11] Solving this quadratic equation to obtain with probability $1-\delta$,

$$\mathbb{E}_\mu[X(\widehat{\theta}_h,\widehat{V}_{h+1},f_{\widehat{V}_{h+1}})] \leq \frac{36H^2(\log(1/\delta)+C_{d,\log K})}{K}+\sqrt{\frac{16H^3\epsilon_\mathcal{F}(\log(1/\delta)+C_{d,\log K})}{K}}+4H\epsilon_\mathcal{F}$$

Now according to equation 18, by definition we finally have with probability $1-\delta$ (recall the expectation over μ is taken w.r.t. $(s_h^k, a_h^k, s_{h+1}^k)$ $k = 1, ..., K$ only)

$$\mathbb{E}_\mu[\ell_h(\widehat{\theta}_{h+1})] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] = \mathbb{E}_\mu[X(\widehat{\theta}_h,\widehat{V}_{h+1},f_{\widehat{V}_{h+1}})]$$

$$\leq \frac{36H^2(\log(1/\delta)+C_{d,\log K})}{K} + \sqrt{\frac{16H^3\epsilon_\mathcal{F}(\log(1/\delta)+C_{d,\log K})}{K}} + 4H\epsilon_\mathcal{F}. \tag{20}$$

Step2. If $\lambda > 0$, there is only extra term $\frac{\lambda}{K}\left(\left\|\widehat{\theta}_h\right\|_2 - \left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2\right) \leq \frac{\lambda}{K}\left\|\widehat{\theta}_h\right\|_2 \leq \frac{\lambda C_\Theta^2}{K}$ in addition to above. This finishes the proof.

$\square$

Theorem G.2 (Provable efficiency (Part I)). *Let $C_{d,\log K}$ be the same as Lemma G.1. Then denote* $b_{d,K,\epsilon_\mathcal{F}} := \sqrt{\frac{16H^3\epsilon_\mathcal{F}(\log(1/\delta)+C_{d,\log K})}{K}} + 4H\epsilon_\mathcal{F}$, *with probability* $1-\delta$

$$\left\|\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2 \leq \sqrt{\frac{36H^2(\log(H/\delta)+C_{d,\log K})+2\lambda C_\Theta^2}{\kappa K}} + \sqrt{\frac{b_{d,K,\epsilon_\mathcal{F}}}{\kappa}} + \sqrt{\frac{2H\epsilon_\mathcal{F}}{\kappa}}, \ \forall h \in [H].$$

Proof of Theorem G.2. Apply a union bound in Lemma G.1, we have with probability $1-\delta$,

$$\mathbb{E}_\mu[\ell_h(\widehat{\theta}_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] \leq \frac{36H^2(\log(H/\delta)+C_{d,\log K})+\lambda C_\Theta^2}{K} + b_{d,K,\epsilon_\mathcal{F}}, \ \forall h \in [H]$$

$$\Rightarrow \mathbb{E}_\mu[\ell_h(\widehat{\theta}_h) - \frac{\lambda}{K}\left\|\widehat{\theta}_h\right\|_2^2] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda}{K}\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2] \leq \frac{36H^2(\log(H/\delta)+C_{d,\log K})+2\lambda C_\Theta^2}{K} + b_{d,K,\epsilon_\mathcal{F}} \tag{21}$$

Now we prove for all $h \in [H]$,

$$\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(\cdot,\cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(\cdot,\cdot))\right)^2\right] \leq \mathbb{E}_\mu\left[\ell_h(\widehat{\theta}_h) - \frac{\lambda\left\|\widehat{\theta}_h\right\|_2^2}{K}\right] - \mathbb{E}_\mu\left[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2}{K}\right] + 2H\epsilon_\mathcal{F}. \tag{22}$$

Indeed, similar to equation 20, by definition we have

$$\mathbb{E}_\mu\left[\ell_h(\widehat{\theta}_h) - \frac{\lambda\left\|\widehat{\theta}_h\right\|_2^2}{K}\right] - \mathbb{E}_\mu\left[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2}{K}\right] = \mathbb{E}_\mu[X(\widehat{\theta}_h,\widehat{V}_{h+1},f_{\widehat{V}_{h+1}})]$$

$$=\mathbb{E}_\mu\left(\left[f\left(\widehat{\theta}_h,\phi(s_h,a_h)\right) - r_h - \widehat{V}_{h+1}(s_{h+1})\right]^2 - \left[f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(s_h,a_h)\right) - r_h - \widehat{V}_{h+1}(s_{h+1})\right]^2\right)$$

$$=\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(\cdot,\cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(\cdot,\cdot))\right)^2\right]$$

$$+\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(s_h,a_h)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(s_h,a_h))\right) \cdot \left(f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(s_h,a_h)\right) - r_h - \widehat{V}_{h+1}(s_{h+1})\right)\right]$$

$$=\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(\cdot,\cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(\cdot,\cdot))\right)^2\right]$$

$$+\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(s_h,a_h)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(s_h,a_h))\right) \cdot \mathbb{E}\left(f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(s_h,a_h)\right) - r_h - \widehat{V}_{h+1}(s_{h+1})\Big|s_h,a_h\right)\right]$$

$$\geq\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h,\phi(\cdot,\cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}},\phi(\cdot,\cdot))\right)^2\right] - 2H\epsilon_\mathcal{F}$$

---

[11]Here in our realization of Lemma L.9, we set $B = 1/\lambda$ (since $\left\|\Sigma_h^{-1}\right\|_2 \leq 1/\lambda$).

where the third identity uses μ is taken w.r.t. s_h, a_h, s_{h+1} (recall Lemma G.1) and law of total expectation. The first inequality uses the definition of $\theta_{\mathbb{T}\widehat{V}_{h+1}}$.

Now apply Assumption 2.3, we have

$$\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot))\right)^2\right] \geq \kappa \left\|\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2,$$

Combine the above with equation 21 and equation 22, we obtain the stated result. □

Theorem G.3 (Provable efficiency (Part II)). *Let $C_{d,\log K}$ be the same as Lemma G.1 and suppose $\epsilon_\mathcal{F} = 0$. Furthermore, suppose $\lambda \leq 1/2C_\Theta^2$ and $K \geq \max\left\{512\frac{\kappa_1^4}{\kappa^2}\left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K^3}{\lambda^2})\right), \frac{4\lambda}{\kappa}\right\}$. Then, with probability $1 - \delta$, $\forall h \in [H]$,*

$$\sup_{s,a}\left|f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a))\right| \leq \left(\kappa_1 H\sqrt{\frac{36H^2(\log(H^2/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^2 d\kappa_1}{\sqrt{\kappa}}\right)\sqrt{\frac{1}{K}} + O(\frac{1}{K}).$$

Furthermore, we have with probability $1 - \delta$,

$$\sup_h\left\|\widehat{V}_h - V_h^\star\right\|_\infty \leq \left(\kappa_1 H\sqrt{\frac{36H^2(\log(H^2/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^2 d\kappa_1}{\sqrt{\kappa}}\right)\sqrt{\frac{1}{K}} + O(\frac{1}{K})$$

$$= \widetilde{O}\left(\kappa_1 H^2\sqrt{\frac{d^2}{\kappa}}\sqrt{\frac{1}{K}}\right)$$

where $\widetilde{O}$ absorbs Polylog terms and higher order terms. Lastly, it also holds for all $h \in [H]$, w.p. $1 - \delta$

$$\left\|\widehat{\theta}_h - \theta_h^\star\right\|_2 \leq \left(\kappa_1 H\frac{\sqrt{72H^2(\log(H^2/\delta) + C_{d,\log K}) + 4\lambda C_\Theta^2}}{\kappa} + \frac{4H^2 d\kappa_1}{\kappa}\right)\sqrt{\frac{1}{K}} + O(\frac{1}{K})$$

$$= \widetilde{O}\left(\frac{\kappa_1 H^2 d}{\kappa}\sqrt{\frac{1}{K}}\right)$$

Proof of Theorem G.3. **Step1:** we show the first result.

We prove this by backward induction. When $h = H + 1$, by convention $f(\widehat{\theta}_h, \phi(s,a)) = f(\theta_h^\star, \phi(s,a)) = 0$ so the base case holds. Suppose for $h + 1$, with probability $1 - (H - h)\delta$, $\sup_{s,a}\left|f(\widehat{\theta}_{h+1}, \phi(s,a)) - f(\theta_{h+1}^\star, \phi(s,a))\right| \leq C_{h+1}\sqrt{\frac{1}{K}} + a(h+1)$, we next consider the case for $t = h$.

On one hand, by Theorem G.2, we have with probability $1 - \delta/2$,

$$\sup_{s,a}\left|f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a))\right|$$

$$\leq \sup_{s,a}\left|f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a))\right| + \sup_{s,a}\left|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\theta_h^\star, \phi(s,a))\right|$$

$$= \sup_{s,a}\left|\nabla f(\xi, \phi(s,a))^\top(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}})\right| + \sup_{s,a}\left|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi(s,a))\right|$$

$$\leq \kappa_1 \cdot \left\|\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2 + \sup_{s,a}\left|\mathcal{P}_{h,s,a}\widehat{V}_{h+1} - \mathcal{P}_{h,s,a}V_{h+1}^\star\right|$$

$$\leq \kappa_1\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} + \left\|\widehat{V}_{h+1} - V_{h+1}^\star\right\|_\infty,$$

Recall $\widehat{V}_{h+1}(\cdot) := \min\{\max_a f(\widehat{\theta}_{h+1}, \phi(\cdot, a)) - \Gamma_h(\cdot, a), H\}$ and $V_{h+1}^\star(\cdot) = \max_a f(\theta_{h+1}^\star, \phi(\cdot, a)) = \min\{\max_a f(\theta_{h+1}^\star, \phi(\cdot, a)), H\}$, we obtain

$$\left\|\widehat{V}_{h+1} - V_{h+1}^\star\right\|_\infty \leq \sup_{s,a}\left|f(\widehat{\theta}_{h+1}, \phi(s,a)) - f(\theta_{h+1}^\star, \phi(s,a))\right| + \sup_{h,s,a}\Gamma_h(s,a) \quad (23)$$

Note the above holds true for any generic $\Gamma_h(s, a)$. In particular, according to Algorithm 1, we specify

$$\Gamma_h(\cdot, \cdot) = dH\sqrt{\nabla_\theta f(\widehat{\theta}_h, \phi(\cdot, \cdot))^\top \Sigma_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(\cdot, \cdot))} \left(+\widetilde{O}(\frac{1}{K}) \right)$$

and by Lemma L.5, with probability $1 - \delta$,

$$\Gamma_h \leq \frac{2dH\kappa_1}{\sqrt{\kappa K}} + \widetilde{O}(\frac{1}{K})$$

and by a union bound this implies with probability $1 - (H - h + 1)\delta$,

$$\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)) \right|$$

$$\leq C_{h+1}\sqrt{\frac{1}{K}} + \kappa_1\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} + \frac{2dH\kappa_1}{\sqrt{\kappa K}} + \widetilde{O}(\frac{1}{K}) := C_h\sqrt{\frac{1}{K}} + \widetilde{O}(\frac{1}{K})$$

Solving for C_h, we obtain $C_h \leq \kappa_1 H\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + H\frac{2dH\kappa_1}{\sqrt{\kappa}}$ for all H. By a union bound (replacing δ by δ/H), we obtain the stated result.

Step2: Utilizing the intermediate result equation 23, we directly have with probability $1 - \delta$,

$$\sup_h \left\| \widehat{V}_h - V_h^\star \right\|_\infty \leq \sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)) \right| + \frac{2dH\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}),$$

where $\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)) \right|$ can be bounded using Step1.

Step3: Denote $M := \left(\kappa_1 H\sqrt{\frac{36H^2(\log(H^2/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^2 d\kappa_1}{\sqrt{\kappa}} \right)\sqrt{\frac{1}{K}} + O(\frac{1}{K})$, then by Step1 we have with probability $1 - \delta$ (here ξ is some point between $\widehat{\theta}_h$ and $\theta_h^\star$) for all $h \in [H]$

$$M^2 \geq \sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)) \right|^2$$

$$\geq \mathbb{E}_{\mu,h}[(f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)))^2] \geq \kappa \left\| \widehat{\theta}_h - \theta_h^\star \right\|_2^2$$

where the last inequality is by Assumption 2.3. Solve this to obtain the stated result.

$\square$

H WITH POSITIVE BELLMAN COMPLETENESS COEFFICIENT $\epsilon_{\mathcal{F}} > 0$

In Theorem 3.2, we consider the case where $\epsilon_{\mathcal{F}} = 0$. If $\epsilon_{\mathcal{F}} > 0$, similar guarantee can be achieved with the measurement of model misspecification. For instance, the additional error $\sqrt{\frac{16H^3\epsilon_{\mathcal{F}}(\log(1/\delta) + C_{d,\log K})}{K}} + 4H\epsilon_{\mathcal{F}}$ will show up in Lemma G.1 (as stated in the current version), $\sqrt{\frac{b_{d,K,\epsilon_{\mathcal{F}}}}{\kappa}} + \sqrt{\frac{2H\epsilon_{\mathcal{F}}}{\kappa}}$ will show up in Lemma G.2. Then the decomposition in equation 3 will incur the extra $\delta_{\widehat{V}_{h+1}}$ term with $\delta_{\widehat{V}_{h+1}}$ might not be 0. The analysis with positive $\epsilon_{\mathcal{F}} > 0$ will make the proofs more intricate but incurs no additional technical challenge. Since the inclusion of this quantity is not our major focus, as a result, we only provide the proof for the case where $\epsilon_{\mathcal{F}} = 0$ so the readers can focus on the more critical components that characterize the hardness of differentiable function class.

I VFQL AND ITS ANALYSIS

We present the *vanilla fitted Q-learning* (VFQL) Algorithm 2 as follows. For VFQL, no pessimism is used and we assume $\widehat{\theta}_h \in \Theta$ without loss of generality.

Algorithm 2 Vanilla Fitted Q-Learning (VFQL)

1: **Input:** Offline Dataset $\mathcal{D} = \left\{ \left(s_h^k, a_h^k, r_h^k, s_{h+1}^k \right) \right\}_{k,h=1}^{K,H}$. Denote $\phi_{h,k} := \phi(s_h^k, a_h^k)$.

2: **Initialization:** Set $\widehat{V}_{H+1}(\cdot) \leftarrow 0$ and $\lambda > 0$.

3: **for** $h = H, H-1, \ldots, 1$ **do**

4:      Set $\widehat{\theta}_h \leftarrow \arg\min_{\theta \in \Theta} \left\{ \sum_{k=1}^K \left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right]^2 + \lambda \cdot \|\theta\|_2^2 \right\}$

5:      Set $\widehat{Q}_h(\cdot, \cdot) \leftarrow \min \left\{ f(\widehat{\theta}_h, \phi(\cdot, \cdot)), H - h + 1 \right\}^+$

6:      Set $\widehat{\pi}_h(\cdot \mid \cdot) \leftarrow \arg\max_{\pi_h} \langle \widehat{Q}_h(\cdot, \cdot), \pi_h(\cdot \mid \cdot) \rangle_{\mathcal{A}}$, $\widehat{V}_h(\cdot) \leftarrow \max_{\pi_h} \langle \widehat{Q}_h(\cdot, \cdot), \pi_h(\cdot \mid \cdot) \rangle_{\mathcal{A}}$

7: **end for**

8: **Output:** $\{\widehat{\pi}_h\}_{h=1}^H$.

I.1   ANALYSIS FOR VFQL (THEOREM 3.1)

Recall $\iota_h(s, a) := \mathcal{P}_h \widehat{V}_{h+1}(s, a) - \widehat{Q}_h(s, a)$ and the definition of Bellman operator D.1. Note $\min\{\cdot, H - h + 1\}^+$ is a non-expansive operator, therefore we have

$$|\iota_h(s, a)| = |\mathcal{P}_h \widehat{V}_{h+1}(s, a) - \widehat{Q}_h(s, a)| = \left| \min\left\{ \mathcal{P}_h \widehat{V}_{h+1}(s, a), H - h + 1 \right\}^+ - \min\left\{ f(\widehat{\theta}_h, \phi(\cdot, \cdot)), H - h + 1 \right\}^+ \right|$$

$$\leq \left| \mathcal{P}_h \widehat{V}_{h+1}(s, a) - f(\widehat{\theta}_h, \phi(\cdot, \cdot)) \right| \leq \left| f(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - f(\widehat{\theta}_h, \phi(\cdot, \cdot)) \right| + \epsilon_{\mathcal{F}}.$$

By Lemma D.2, we have for any π,

$$v^\pi - v^{\widehat{\pi}} = -\sum_{h=1}^H E_{\widehat{\pi}}[\iota_h(s_h, a_h)] + \sum_{h=1}^H E_\pi[\iota_h(s_h, a_h)] \leq \sum_{h=1}^H E_{\widehat{\pi}}[|\iota_h(s_h, a_h)|] + \sum_{h=1}^H E_\pi[|\iota_h(s_h, a_h)|]$$

$$\leq \sum_{h=1}^H \mathbb{E}_{\widehat{\pi}}[|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) - f(\widehat{\theta}_h, \phi(\cdot, \cdot))|] + \sum_{h=1}^H \mathbb{E}_\pi[|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) - f(\widehat{\theta}_h, \phi(\cdot, \cdot))|] + 2H\epsilon_{\mathcal{F}}$$

$$\leq \sum_{h=1}^H \sqrt{\mathbb{E}_{\widehat{\pi}}[|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) - f(\widehat{\theta}_h, \phi(\cdot, \cdot))|^2]} + \sum_{h=1}^H \sqrt{\mathbb{E}_\pi[|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) - f(\widehat{\theta}_h, \phi(\cdot, \cdot))|^2]} + 2H\epsilon_{\mathcal{F}}$$

$$\leq 2\sqrt{C_{\text{eff}}} \sum_{h=1}^H \sqrt{\mathbb{E}_{\mu,h}[|f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) - f(\widehat{\theta}_h, \phi(\cdot, \cdot))|^2]} + 2H\epsilon_{\mathcal{F}}$$

$$\tag{24}$$

where the second inequality uses Cauchy inequality and the third one uses the definition of concentrability coefficient 2.2.

Next, for VFQL, there is no pessimism therefore the quantity B in Lemma L.10 is zero, hence the covering number applied in Lemma G.1 is bounded by $C_{d,\log(K)} \leq \widetilde{O}(d)$ and

$$\mathbb{E}_\mu[\ell_h(\widehat{\theta}_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] \leq \frac{36H^2(\log(1/\delta) + C_{d,\log K}) + \lambda C_\Theta^2}{K} + \sqrt{\frac{16H^3 \epsilon_{\mathcal{F}}(\log(1/\delta) + C_{d,\log K})}{K}} + 4H\epsilon_{\mathcal{F}}.$$

Now leveraging equation 21 and equation 22 in Theorem G.2 to obtain

$$\mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 \right] \leq \mathbb{E}_\mu \left[\ell_h(\widehat{\theta}_h) - \frac{\lambda \left\| \widehat{\theta}_h \right\|_2^2}{K} \right] - \mathbb{E}_\mu \left[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2^2}{K} \right] + 2H\epsilon_{\mathcal{F}}$$

$$\leq \frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{K} + b_{d,K,\epsilon_{\mathcal{F}}} + 2H\epsilon_{\mathcal{F}}$$

Plug the above into equation 24, we obtain with probability $1 - \delta$, for all policy π,

$$v^\pi - v^{\widehat\pi} \le 2\sqrt{C_{\text{eff}}}H\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{K} + b_{d,K,\epsilon_{\mathcal{F}}}} + 2H\epsilon_{\mathcal{F}} + 2H\epsilon_{\mathcal{F}}$$

$$= 2\sqrt{C_{\text{eff}}}H\sqrt{\frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{K}} + \sqrt{\frac{16H^3\epsilon_{\mathcal{F}}(\log(1/\delta) + C_{d,\log K})}{K}} + 6H\epsilon_{\mathcal{F}} + 2H\epsilon_{\mathcal{F}}$$

$$= \sqrt{C_{\text{eff}}}H \cdot \widetilde{O}\left(\sqrt{\frac{H^2d + \lambda C_\Theta^2}{K}} + \sqrt[4]{\frac{H^3 d\epsilon_{\mathcal{F}}}{K}}\right) + O(\sqrt{C_{\text{eff}}H^3\epsilon_{\mathcal{F}}} + H\epsilon_{\mathcal{F}})$$

This finishes the proof of Theorem 3.1.

J   PROOFS FOR VAFQL

In this section, we present the analysis for *variance-aware fitted Q learning* (VAFQL). Throughout the whole section, we assume $\epsilon_{\mathcal{F}} = 0$, *i.e.* the exact Bellman-Completeness holds. The algorithm is presented in the following. Before giving the proofs of Theorem 3, we first prove some useful lemmas.

Algorithm 3 Variance-Aware Fitted Q Learning (VAFQL)

1: **Input:** Split dataset $\mathcal{D} = \left\{\left(s_h^k, a_h^k, r_h^k\right)\right\}_{k,h=1}^{K,H}$ $\mathcal{D}' = \left\{\left(\bar{s}_h^k, \bar{a}_h^k, \bar{r}_h^k\right)\right\}_{k,h=1}^{K,H}$. Require β.
2: **Initialization:** Set $\widehat{V}_{H+1}(\cdot) \leftarrow 0$. Denote $\phi_{h,k} := \phi(s_h^k, a_h^k), \bar\phi_{h,k} := \phi(\bar{s}_h^k, \bar{a}_h^k)$
3: **for** $h = H, H-1, \dots, 1$ **do**
4:     Set $\boldsymbol{u}_h \leftarrow \arg\min_{\theta \in \Theta}\left\{\sum_{k=1}^{K}\left[f\left(\theta, \bar\phi_{h,k}\right) - \widehat{V}_{h+1}(\bar{s}_{h+1}^k)\right]^2 + \lambda \cdot \|\theta\|_2^2\right\}$
5:     Set $\boldsymbol{v}_h \leftarrow \arg\min_{\theta \in \Theta}\left\{\sum_{k=1}^{K}\left[f\left(\theta, \bar\phi_{h,k}\right) - \widehat{V}_{h+1}^2(\bar{s}_{h+1}^k)\right]^2 + \lambda \cdot \|\theta\|_2^2\right\}$
6:     Set $\left[\widehat{\text{Var}}_h\widehat{V}_{h+1}\right](\cdot,\cdot) = f(\boldsymbol{v}_h, \phi(\cdot,\cdot))_{[0,(H-h+1)^2]} - \left[f(\boldsymbol{u}_h, \phi(\cdot,\cdot))_{[0,H-h+1]}\right]^2$
7:     Set $\widehat\sigma_h(\cdot,\cdot)^2 \leftarrow \max\{1, \widehat{\text{Var}}_{P_h}\widehat{V}_{h+1}(\cdot,\cdot)\}$
8:     Set $\widehat\theta_h \leftarrow \arg\min_{\theta \in \Theta}\left\{\sum_{k=1}^{K}\left[f\left(\theta, \phi_{h,k}\right) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)\right]^2 / \widehat\sigma_h^2(s_h^k, a_h^k) + \lambda \cdot \|\theta\|_2^2\right\}$
9:     Set $\Lambda_h \leftarrow \sum_{k=1}^{K}\nabla f(\widehat\theta_h, \phi_{h,k})\nabla f(\widehat\theta_h, \phi_{h,k})^\top / \widehat\sigma^2(s_h^k, a_h^k) + \lambda \cdot I$,
10:    Set $\Gamma_h(\cdot,\cdot) \leftarrow \beta\sqrt{\nabla_\theta f(\widehat\theta_h, \phi(\cdot,\cdot))^\top \Lambda_h^{-1}\nabla_\theta f(\widehat\theta_h, \phi(\cdot,\cdot))}\left(+\widetilde{O}(\frac{1}{K})\right)$
11:    Set $\bar{Q}_h(\cdot,\cdot) \leftarrow f(\widehat\theta_h, \phi(\cdot,\cdot)) - \Gamma_h(\cdot,\cdot), \widehat{Q}_h(\cdot,\cdot) \leftarrow \min\left\{\bar{Q}_h(\cdot,\cdot), H-h+1\right\}^+$
12:    Set $\widehat\pi_h(\cdot \mid \cdot) \leftarrow \arg\max_{\pi_h}\left\langle\widehat{Q}_h(\cdot,\cdot), \pi_h(\cdot \mid \cdot)\right\rangle_{\mathcal{A}}, \widehat{V}_h(\cdot) \leftarrow \max_{\pi_h}\left\langle\widehat{Q}_h(\cdot,\cdot), \pi_h(\cdot \mid \cdot)\right\rangle_{\mathcal{A}}$
13: **end for**
14: **Output:** $\{\widehat\pi_h\}_{h=1}^{H}$.

J.1   PROVABLE EFFICIENCY FOR VARIANCE-AWARE FITTED Q LEARNING

Recall the objective

$$\ell_h(\theta) := \frac{1}{K}\sum_{k=1}^{K}\left[f\left(\theta, \phi(s_h^k, a_h^k)\right) - r(s_h^k, a_h^k) - \widehat{V}_{h+1}(s_{h+1}^k)\right]^2 / \widehat\sigma_h^2(s_h^k, a_h^k) + \frac{\lambda}{K} \cdot \|\theta\|_2^2$$

Then by definition, $\widehat\theta_h := \arg\min_{\theta \in \Theta}\ell_h(\theta)$ and $\theta_{\mathbb{T}\widehat{V}_{h+1}}$ satisfies $f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi) = \mathcal{P}_h\widehat{V}_{h+1}(s_{h+1}^k)$ (recall $\epsilon_{\mathcal{F}} = 0$). Therefore, in this case, we have the following lemma:

Lemma J.1. *Fix $h \in [H]$. With probability $1 - \delta$,*

$$\mathbb{E}_\mu[\ell_h(\widehat\theta_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] \le \frac{36H^2(\log(1/\delta) + C_{d,\log K}) + \lambda C_\Theta^2}{K}$$

where the expectation over μ is taken w.r.t. $(s_h^k, a_h^k, s_{h+1}^k)$ $k = 1, \dots, K$ only (i.e., first compute $\mathbb{E}_\mu[\ell_h(\theta)]$ for a fixed θ, then plug-in either $\widehat\theta_{h+1}$ or $\theta_{\mathbb{T}\widehat{V}_{h+1}}$).   Here

$$C_{d,\log(K)} := d\log(1+24C_\Theta(H+1)\kappa_1 K)+d\log\left(1+288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta}+2\sqrt{\kappa_1\kappa_2/\lambda})^2 K^2\right)+$$
$$d^2\log\left(1+288H^2\sqrt{d}\kappa_1^2 K^2/\lambda\right)+d\log(1+16C_\Theta H^2\kappa_1 K)+d\log(1+32C_\Theta H^3\kappa_1 K).$$

Proof of Lemma J.1. **Step1:** Consider the case where $\lambda = 0$. Indeed, fix $h \in [H]$ and any function $V(\cdot) \in \mathbb{R}^{\mathcal{S}}$. Similarly, define $f_V(s,a) := f(\theta_{\mathbb{T}V}, \phi) = \mathcal{P}_h V$. For any fixed $\theta \in \Theta$, denote $g(s,a) = f(\theta, \phi(s,a))$. Moreover, for any $u, v \in \Theta$, define

$$\sigma_{u,v}^2(\cdot,\cdot) := \max\{1, f(v, \phi(\cdot,\cdot))_{[0,(H-h+1)^2]} - \left[f(u, \phi(\cdot,\cdot))_{[0,H-h+1]}\right]^2\}$$

Then define (we omit the subscript u,v of $\sigma_{u,v}^2$ for the illustration purpose when there is no ambiguity)

$$X(g, V, f_V, \sigma^2) := \frac{(g(s,a) - r - V(s'))^2 - (f_V(s,a) - r - V(s'))^2}{\sigma_{u,v}^2(s,a)}.$$

Since all episodes are independent of each other, $X_k(g, V, f_V) := X(g(s_h^k, a_h^k), V(s_{h+1}^k), f_V(s_h^k, a_h^k), \sigma^2(s_h^k, a_h^k))$ are independent r.v.s and it holds

$$\frac{1}{K}\sum_{k=1}^K X_k(g, V, f_V, \sigma^2) = \ell(g) - \ell(f_V). \tag{25}$$

Next, the variance of X is bounded by

$$\mathrm{Var}[X(g, V, f_V, \sigma^2)] \le \mathbb{E}_\mu[X(g, f, f_V, \sigma^2)^2]$$
$$=\mathbb{E}_\mu\left[\left((g(s_h, a_h) - r_h - V(s_{h+1}))^2 - (f_V(s_h, a_h) - r_h - V(s_{h+1}))^2\right)^2 / \sigma^2(s_h, a_h)^2\right]$$
$$=\mathbb{E}_\mu\left[\frac{(g(s_h, a_h) - f_V(s_h, a_h))^2}{\sigma^2(s_h, a_h)} \cdot \frac{(g(s_h, a_h) + f_V(s_h, a_h) - 2r_h - 2V(s_{h+1}))^2}{\sigma^2(s_h, a_h)}\right]$$
$$\le 4H^2 \cdot \mathbb{E}_\mu[\frac{(g(s_h, a_h) - f_V(s_h, a_h))^2}{\sigma^2(s_h, a_h)}]$$
$$=4H^2 \cdot \mathbb{E}_\mu\left[\frac{(g(s_h, a_h) - r_h - V(s_{h+1}))^2 - (f_V(s_h, a_h) - r_h - V(s_{h+1}))^2}{\sigma^2(s_h, a_h)}\right] \quad (*)$$
$$=4H^2 \cdot \mathbb{E}_\mu[X(g, f, f_V, \sigma^2)]$$

$(*)$ follows from that

$$\mathbb{E}_\mu\left[\frac{f(\widehat{\theta}_h, \phi(s_h, a_h)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h))}{\sigma^2(s_h, a_h)} \cdot \mathbb{E}\left(f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h)\right) - r_h - \widehat{V}_{h+1}(s_{h+1})\Big| s_h, a_h\right)\right] = 0.$$

Therefore, by Bernstein inequality, with probability $1 - \delta$,

$$\mathbb{E}_\mu[X(g, f, f_V, \sigma^2)] - \frac{1}{K}\sum_{k=1}^K X_k(g, f, f_V, \sigma^2)$$
$$\le \sqrt{\frac{2\mathrm{Var}[X(g, f, f_V, \sigma^2)]\log(1/\delta)}{K}} + \frac{4H^2\log(1/\delta)}{3K}$$
$$\le \sqrt{\frac{8H^2\mathbb{E}_\mu[X(g, f, f_V, \sigma^2)]\log(1/\delta)}{K}} + \frac{4H^2\log(1/\delta)}{3K}.$$

Now, if we choose $g(s,a) := f(\widehat{\theta}_h, \phi(s,a))$ and $u = \boldsymbol{u}_h, v = \boldsymbol{v}_h$ from Algorithm 3, then $\widehat{\theta}_h$ minimizes $\ell_h(\theta)$, therefore, it also minimizes $\frac{1}{K}\sum_{k=1}^K X_i(\theta, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2)$ and this implies

$$\frac{1}{K}\sum_{k=1}^K X_k(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2) \le \frac{1}{K}\sum_{k=1}^K X_k(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2) = 0.$$

Thus, we obtain

$$\mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2)] \le \sqrt{\frac{8H^2 \cdot \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2)] \log(1/\delta)}{K}} + \frac{4H^2 \log(1/\delta)}{3K}.$$

However, the above does not hold with probability $1 - \delta$ since $\widehat{\theta}_h$, $\widehat{\sigma}_h^2$ and $\widehat{V}_{h+1} := \min\{\max_a f(\widehat{\theta}_{h+1}, \phi(\cdot, a)) - \sqrt{\nabla f(\widehat{\theta}_{h+1}, \phi(\cdot, a))^\top A \cdot \nabla f(\theta, \phi(\cdot, a))}, H\}$ (where A is certain symmetric matrix with bounded norm) depend on $\widehat{\theta}_h, \widehat{\theta}_{h+1}$ which are data-dependent. Therefore, we need to further apply covering Lemma L.11 and choose $\epsilon = O(1/K)$ and a union bound to obtain with probability $1 - \delta$,

$$\mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2)] \le \sqrt{\frac{8H^2 \cdot \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}}, \widehat{\sigma}_h^2)](\log(1/\delta) + C_{d,\log K})}{K}} + \frac{4H^2(\log(1/\delta) + C_{d,\log K})}{3K}.$$

where $C_{d,\log(K)} := d\log(1 + 24C_\Theta(H + 1)\kappa_1 K) + d\log\left(1 + 288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{\kappa_1\kappa_2/\lambda})^2 K^2\right) + d^2 \log\left(1 + 288H^2\sqrt{d}\kappa_1^2 K^2/\lambda\right) + d\log(1 + 16C_\Theta H^2\kappa_1 K) + d\log(1 + 32C_\Theta H^3\kappa_1 K)$ (where we let $B = 1/\lambda$ since $\left\|\Lambda_h^{-1}\right\|_2 \le 1/\lambda$). Solving this quadratic equation to obtain with probability $1 - \delta$,

$$\mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})] \le \frac{36H^2(\log(1/\delta) + C_{d,\log K})}{K}.$$

Now according to equation 25, by definition we finally have with probability $1 - \delta$ (recall the expectation over μ is taken w.r.t. $(s_h^k, a_h^k, s_{h+1}^k)$ $k = 1, ..., K$ only)

$$\mathbb{E}_\mu[\ell_h(\widehat{\theta}_{h+1})] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] = \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})] \le \frac{36H^2(\log(1/\delta) + C_{d,\log K})}{K} \quad (26)$$

where we used $f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi) = \mathcal{P}_h \widehat{V}_{h+1} = f_{\widehat{V}_{h+1}}$.

Step2. If $\lambda > 0$, there is only extra term $\frac{\lambda}{K}\left(\left\|\widehat{\theta}_h\right\|_2 - \left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2\right) \le \frac{\lambda}{K}\left\|\widehat{\theta}_h\right\|_2 \le \frac{\lambda C_\Theta^2}{K}$ in addition to above. This finishes the proof.

$\square$

Theorem J.2 (Provable efficiency for VAFQL). *Let $C_{d,\log K}$ be the same as Lemma J.1. Then, with probability $1 - \delta$*

$$\left\|\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2 \le \sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}}, \ \forall h \in [H].$$

Proof of Theorem J.2. Apply a union bound in Lemma J.1, we have with probability $1 - \delta$,

$$\mathbb{E}_\mu[\ell_h(\widehat{\theta}_h)] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}})] \le \frac{36H^2(\log(H/\delta) + C_{d,\log K}) + \lambda C_\Theta^2}{K}, \ \forall h \in [H]$$

$$\Rightarrow \mathbb{E}_\mu[\ell_h(\widehat{\theta}_h) - \frac{\lambda}{K}\left\|\widehat{\theta}_h\right\|_2^2] - \mathbb{E}_\mu[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda}{K}\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2] \le \frac{36H^2(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{K} \tag{27}$$

Now we prove for all $h \in [H]$,

$$\mathbb{E}_\mu\left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot))\right)^2\right] = \mathbb{E}_\mu\left[\ell_h(\widehat{\theta}_h) - \frac{\lambda\left\|\widehat{\theta}_h\right\|_2^2}{K}\right] - \mathbb{E}_\mu\left[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda\left\|\theta_{\mathbb{T}\widehat{V}_{h+1}}\right\|_2^2}{K}\right]. \tag{28}$$

Indeed, identical to equation 26,

$$
\mathbb{E}_\mu \left[\ell_h(\widehat{\theta}_h) - \frac{\lambda \left\| \widehat{\theta}_h \right\|_2^2}{K} \right] - \mathbb{E}_\mu \left[\ell_h(\theta_{\mathbb{T}\widehat{V}_{h+1}}) - \frac{\lambda \left\| \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2^2}{K} \right] = \mathbb{E}_\mu[X(\widehat{\theta}_h, \widehat{V}_{h+1}, f_{\widehat{V}_{h+1}})]
$$

$$
= \mathbb{E}_\mu \left(\left[f\left(\widehat{\theta}_h, \phi(s_h, a_h) \right) - r_h - \widehat{V}_{h+1}(s_{h+1}) \right]^2 / \widehat{\sigma}_h^2(s_h, a_h) - \left[f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h) \right) - r_h - \widehat{V}_{h+1}(s_{h+1}) \right]^2 / \widehat{\sigma}_h^2(s_h, a_h) \right)
$$

$$
= \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 / \widehat{\sigma}_h^2(\cdot, \cdot) \right]
$$

$$
+ \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(s_h, a_h)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h)) \right) \cdot \left(f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h) \right) - r_h - \widehat{V}_{h+1}(s_{h+1}) \right) / \widehat{\sigma}_h^2(s_h, a_h) \right]
$$

$$
= \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 / \widehat{\sigma}_h^2(\cdot, \cdot) \right]
$$

$$
+ \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(s_h, a_h)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h)) \right) \cdot \mathbb{E}\left(f\left(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s_h, a_h) \right) - r_h - \widehat{V}_{h+1}(s_{h+1}) \Big| s_h, a_h \right) / \widehat{\sigma}_h^2(s_h, a_h) \right]
$$

$$
= \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 / \widehat{\sigma}_h^2(\cdot, \cdot) \right]
$$

where the third identity uses law of total expectation and that μ is taken w.r.t. s_h, a_h, s_{h+1} only (recall Lemma J.1) so the $\widehat{\sigma}_h^2$ can be move outside of the conditional expectation.[12] The fourth identity uses the definition of $\theta_{\mathbb{T}\widehat{V}_{h+1}}$ since $f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s, a)) = \mathcal{P}_{h,s,a}\widehat{V}_{h+1}$.

Then we have

$$
\mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 / \widehat{\sigma}_h^2(\cdot, \cdot) \right]
$$

$$
\geq \mathbb{E}_\mu \left[\left(f(\widehat{\theta}_h, \phi(\cdot, \cdot)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(\cdot, \cdot)) \right)^2 \right] / H^2 \geq \frac{\kappa}{H^2} \left\| \widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2^2,
$$

where the third identity uses μ is over s_h, a_h only and the last one uses $\widehat{\sigma}_h^2(\cdot, \cdot) \leq H^2$. Combine the above with equation 27 and equation 28, we obtain the stated result. □

Theorem J.3 (Provable efficiency of VAFQL (Part II)). *Let $C_{d,\log K}$ be the same as Lemma J.1. Furthermore, suppose $\lambda \leq 1/2C_\Theta^2$ and $K \geq \max\left\{ 512\frac{\kappa_1^4}{\kappa^2}\left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1^3\kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$. Then, with probability $1 - \delta$, $\forall h \in [H]$*

$$
\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s, a)) - f(\theta_h^\star, \phi(s, a)) \right| \leq \left(\kappa_1 H \sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2dH^3\kappa_1}{\sqrt{\kappa}} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K}),
$$

Furthermore, we have with probability $1 - \delta$,

$$
\sup_h \left\| \widehat{V}_h - V_h^\star \right\|_\infty \leq \left(\kappa_1 H \sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2dH^3\kappa_1}{\sqrt{\kappa}} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K})
$$

$$
= \widetilde{O}\left(\kappa_1 H^3 \sqrt{\frac{d^2}{\kappa}} \sqrt{\frac{1}{K}} \right)
$$

where $\widetilde{O}$ absorbs Polylog terms and higher order terms. Lastly, it also holds for all $h \in [H]$, w.p. $1 - \delta$

$$
\left\| \widehat{\theta}_h - \theta_h^\star \right\|_2 \leq \left(\kappa_1 H \frac{\sqrt{72H^4(\log(H^2/\delta) + C_{d,\log K}) + 4\lambda C_\Theta^2}}{\kappa} + \frac{4H^3 d\kappa_1}{\kappa} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K})
$$

$$
= \widetilde{O}\left(\frac{\kappa_1 H^3 d}{\kappa} \sqrt{\frac{1}{K}} \right)
$$

---

[12]Recall $\widehat{\sigma}_h^2$ computed in Algorithm 3 uses an independent copy $\mathcal{D}'$.

Proof of Theorem J.3. **Step1:** we show the first result.

We prove this by backward induction. When $h = H + 1$, by convention $f(\widehat{\theta}_h, \phi(s,a)) = f(\theta_h^\star, \phi(s,a)) = 0$ so the base case holds. Suppose for $h + 1$, with probability $1 - (H - h)\delta$, $\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right| \leq C_{h+1}\sqrt{\frac{1}{K}}$, we next consider the case for $t = h$.

On one hand, by Theorem J.2, we have with probability $1 - \delta/2$,

$$\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right|$$

$$\leq \sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) \right| + \sup_{s,a} \left| f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right|$$

$$= \sup_{s,a} \left| \nabla f(\xi, \phi(s,a))^\top (\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) \right| + \sup_{s,a} \left| f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi(s,a)) \right|$$

$$\leq \kappa_1 \cdot \left\| \widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}} \right\|_2 + \sup_{s,a} \left| \mathcal{P}_{h,s,a}\widehat{V}_{h+1} - \mathcal{P}_{h,s,a}V_{h+1}^\star \right|$$

$$\leq \kappa_1 \sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} + \left\| \widehat{V}_{h+1} - V_{h+1}^\star \right\|_\infty,$$

Recall we have the form $\widehat{V}_{h+1}(\cdot) := \min\{\max_a f(\widehat{\theta}_{h+1}, \phi(\cdot, a)) - \Gamma_h(\cdot, a), H\}$ and $V_{h+1}^\star(\cdot) = \max_a f(\theta_{h+1}^\star, \phi(\cdot, a)) = \min\{\max_a f(\theta_{h+1}^\star, \phi(\cdot, a)), H\}$, we obtain

$$\left\| \widehat{V}_{h+1} - V_{h+1}^\star \right\|_\infty \leq \sup_{s,a} \left| f(\widehat{\theta}_{h+1}, \phi(s,a)) - f(\theta_{h+1}^\star, \phi(s,a)) \right| + \sup_{h,s,a} \Gamma_h(s,a) \qquad (29)$$

Note the above holds true for any generic $\Gamma_h(s,a)$. In particular, according to Algorithm 3, we specify

$$\Gamma_h(\cdot, \cdot) = d\sqrt{\nabla_\theta f(\widehat{\theta}_h, \phi(\cdot, \cdot))^\top \Lambda_h^{-1} \nabla_\theta f(\widehat{\theta}_h, \phi(\cdot, \cdot))} \left(+ \widetilde{O}(\frac{1}{K}) \right)$$

and by Lemma L.5, with probability $1 - \delta$ (note here Σ_h^{-1} is replaced by Λ_h^{-1} and $\left\| \Lambda_h^{-1} \right\|_2 \leq H^2/\kappa$),

$$\Gamma_h \leq \frac{2dH^2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K})$$

and by a union bound this implies with probability $1 - (H - h + 1)\delta$,

$$\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right|$$

$$\leq C_{h+1}\sqrt{\frac{1}{K}} + \kappa_1 \sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa K}} + \frac{2dH^2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}) := C_h\sqrt{\frac{1}{K}}.$$

Solving for C_h, we obtain $C_h \leq \kappa_1 H\sqrt{\frac{36H^4(\log(H/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + H\frac{2dH^2\kappa_1}{\sqrt{\kappa}}$ for all H. By a union bound (replacing δ by δ/H), we obtain the stated result.

Step2: Utilizing the intermediate result equation 29, we directly have with probability $1 - \delta$,

$$\sup_h \left\| \widehat{V}_h - V_h^\star \right\|_\infty \leq \sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right| + \frac{2dH^2\kappa_1}{\sqrt{\kappa K}} + O(\frac{1}{K}),$$

where $\sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right|$ can be bounded using Step1.

Step3: Denote $M := \left(\kappa_1 H\sqrt{\frac{36H^4(\log(H^2/\delta) + C_{d,\log K}) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^3 d\kappa_1}{\sqrt{\kappa}} \right)\sqrt{\frac{1}{K}} + O(\frac{1}{K})$, then by Step1 we have with probability $1 - \delta$ (here ξ is some point between $\widehat{\theta}_h$ and $\theta_h^\star$) for all $h \in [H]$

$$M^2 \geq \sup_{s,a} \left| f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right|^2$$

$$\geq \mathbb{E}_\mu[\left(f(\widehat{\theta}_h, \phi(s,a)) - f(\theta_h^\star, \phi(s,a)) \right)^2] \geq \kappa \left\| \widehat{\theta}_h - \theta_h^\star \right\|_2^2$$

where the last step is by Assumption 2.3. Solving this to obtain the stated result.

$\square$

J.2 Bounding $|\widehat{\sigma}_h^2 - \sigma_h^{\star 2}|$

Recall the definition $\sigma_h^{\star 2}(\cdot, \cdot) = \max\{1, [\mathrm{Var}_{P_h} V_{h+1}^\star](\cdot, \cdot)\}$. In this section, we bound the term $|\widehat{\sigma}_h^2 - \sigma_h^{\star 2}| := \left\| \widehat{\sigma}_h^2(\cdot, \cdot) - \sigma_h^{\star 2}(\cdot, \cdot) \right\|_\infty$ and

$$
\begin{aligned}
\boldsymbol{u}_h &= \underset{\theta \in \Theta}{\arg\min} \left\{ \frac{1}{K} \sum_{k=1}^K \left[f\left(\theta, \bar{\phi}_{h,k}\right) - \widehat{V}_{h+1}(\bar{s}_{h+1}^k) \right]^2 + \frac{\lambda}{K} \cdot \|\theta\|_2^2 \right\} \\
\boldsymbol{v}_h &= \underset{\theta \in \Theta}{\arg\min} \left\{ \frac{1}{K} \sum_{k=1}^K \left[f\left(\theta, \bar{\phi}_{h,k}\right) - \widehat{V}_{h+1}^2(\bar{s}_{h+1}^k) \right]^2 + \frac{\lambda}{K} \cdot \|\theta\|_2^2 \right\}
\end{aligned}
\tag{30}
$$

where
$$
\widehat{\sigma}_h^2(\cdot, \cdot) := \max\{1, f(\boldsymbol{v}_h, \phi(\cdot, \cdot))_{[0, (H-h+1)^2]} - \left[f(\boldsymbol{u}_h, \phi(\cdot, \cdot))_{[0, H-h+1]} \right]^2 \}
$$
and true parameters $\boldsymbol{u}_h^\star, \boldsymbol{v}_h^\star$ satisfy $f(\boldsymbol{u}_h^\star, \phi(\cdot, \cdot)) = \mathbb{E}_{P(s'|\cdot, \cdot)}[V_h^\star(s')], f(\boldsymbol{v}_h^\star, \phi) = \mathbb{E}_{P(s'|\cdot, \cdot)}[V_h^{\star 2}(s')]$. Furthermore, we define
$$
\sigma_{\widehat{V}_{h+1}}^2(\cdot, \cdot) := \max\{1, [\mathrm{Var}_{P_h} \widehat{V}_{h+1}](\cdot, \cdot)\}
$$
and the *parameter Expectation operator* $\mathbb{J} : V \in \mathbb{R}^{\mathcal{S}} \to \theta_{\mathbb{J}V} \in \Theta$ such that:
$$
f(\theta_{\mathbb{J}V}, \phi) = \mathbb{E}_{P_h}[V(s')], \quad \forall \|V\|_2 \leq \mathcal{B}_{\mathcal{F}}.
$$
Note $\theta_{\mathbb{J}V} \in \Theta$ by Bellman completeness, reward r is constant and differentiability (Definition 1.1) is an additive closed property. By definition,
$$
\begin{aligned}
|\widehat{\sigma}_h^2 - \sigma_{\widehat{V}_{h+1}}^2| &\leq |f(\boldsymbol{v}_h, \phi) - f(\theta_{\mathbb{J}\widehat{V}_{h+1}^2}, \phi)| + |f(\boldsymbol{u}_h, \phi)^2 - f(\theta_{\mathbb{J}\widehat{V}_{h+1}}, \phi)^2| \\
&\leq |f(\boldsymbol{v}_h, \phi) - f(\theta_{\mathbb{J}\widehat{V}_{h+1}^2}, \phi)| + 2H \cdot |f(\boldsymbol{u}_h, \phi) - f(\theta_{\mathbb{J}\widehat{V}_{h+1}}, \phi)|
\end{aligned}
$$

and

$$
\begin{aligned}
|\sigma_h^{\star 2} - \widehat{\sigma}_h^2| &\leq |f(\boldsymbol{v}_h^\star, \phi) - f(\boldsymbol{v}_h, \phi)| + |f(\boldsymbol{u}_h^\star, \phi)^2 - f(\boldsymbol{v}_h, \phi)^2| \\
&\leq |f(\boldsymbol{v}_h^\star, \phi) - f(\boldsymbol{v}_h, \phi)| + 2H \cdot |f(\boldsymbol{u}_h^\star, \phi) - f(\boldsymbol{v}_h, \phi)|
\end{aligned}
$$

We first give the following result.

Lemma J.4. *Suppose* $\lambda \leq 1/2C_\Theta^2$ *and* $K \geq \max\left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1^3 \kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$.
Then, with probability $1 - \delta, \forall h \in [H]$,

$$
\left\| \boldsymbol{u}_h - \theta_{\mathbb{J}\widehat{V}_{h+1}} \right\|_2 \leq \sqrt{\frac{36H^2(\log(H/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa K}}, \quad \forall h \in [H],
$$

$$
\left\| \boldsymbol{v}_h - \theta_{\mathbb{J}\widehat{V}_{h+1}^2} \right\|_2 \leq \sqrt{\frac{36H^4(\log(H/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa K}}, \quad \forall h \in [H].
$$

and

$$
\sup_{s,a} |f(\boldsymbol{u}_h, \phi(s, a)) - f(\boldsymbol{u}_h^\star, \phi(s, a))| \leq \left(\kappa_1 H \sqrt{\frac{36H^2(\log(H^2/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^2 d\kappa_1}{\sqrt{\kappa}} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K}),
$$

$$
\sup_{s,a} |f(\boldsymbol{v}_h, \phi(s, a)) - f(\boldsymbol{v}_h^\star, \phi(s, a))| \leq \left(\kappa_1 H \sqrt{\frac{36H^4(\log(H^2/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa}} + \frac{2H^3 d\kappa_1}{\sqrt{\kappa}} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K}).
$$

The above directly implies for all $h \in [H]$, *with probability* $1 - \delta$,

$$
|\sigma_h^{\star 2} - \widehat{\sigma}_h^2| \leq \left(3\kappa_1 H^2 \sqrt{\frac{36H^4(\log(H^2/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa}} + \frac{6H^4 d\kappa_1}{\sqrt{\kappa}} \right) \sqrt{\frac{1}{K}} + O(\frac{1}{K})
$$

$$
|\widehat{\sigma}_h^2 - \sigma_{\widehat{V}_{h+1}}^2| \leq 3H\kappa_1 \sqrt{\frac{36H^4(\log(H/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa K}}.
$$

Proof of Lemma J.4. In fact, the proof follows a reduction from the provable efficiency procedure conducted in Section G. This is due to the regression procedure in equation 30 is the same as the procedure equation 17 except the *parameter Bellman operator* $\mathbb{T}$ is replaced by the *parameter Expectation operator* $\mathbb{J}$ (recall here $\bar{\phi}_{h,k}$ uses the independent copy $\mathcal{D}'$ and $\widetilde{O}(d^2)$ comes from the covering argument.). Concretely, the $X(g, V, f_V)$ used in Lemma G.1 will be modified to $X(g, V, f_V) = (g(s, a) - V(s'))^2 - (f(\theta_{\mathbb{J}V}, \phi(s, a)) - V(s'))^2$ by removing reward information and the decomposition

$$\mathbb{E}\mu\left[(g(s_h, a_h) - V(s_{h+1}))^2 - (f(\theta_{\mathbb{J}V}, \phi(s_h, a_h)) - V(s_{h+1}))^2\right] = \mathbb{E}_\mu\left[(g(s_h, a_h) - f(\theta_{\mathbb{J}V}, \phi(s_h, a_h)))^2\right]$$

holds true. Then with probability $1 - \delta$,

$$\begin{aligned}
|\sigma_h^{\star 2} - \widehat{\sigma}_h^2| &\leq |f(\boldsymbol{v}_h^\star, \phi) - f(\boldsymbol{v}_h, \phi)| + 2H \cdot |f(\boldsymbol{u}_h^\star, \phi) - f(\boldsymbol{v}_h, \phi)| \\
&\leq \left(3\kappa_1 H^2 \sqrt{\frac{36H^4(\log(H^2/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa}} + \frac{6H^4 d\kappa_1}{\sqrt{\kappa}}\right) \sqrt{\frac{1}{K}} + O(\frac{1}{K}).
\end{aligned}$$

and

$$\begin{aligned}
|\widehat{\sigma}_h^2 - \sigma_{\widehat{V}_{h+1}}^2| &\leq |f(\boldsymbol{v}_h, \phi) - f(\theta_{\mathbb{J}\widehat{V}_{h+1}^2}, \phi)| + 2H \cdot |f(\boldsymbol{u}_h, \phi) - f(\theta_{\mathbb{J}\widehat{V}_{h+1}}, \phi)| \\
&\leq \kappa_1 \left\|\boldsymbol{v}_h - \theta_{\mathbb{J}\widehat{V}_{h+1}^2}\right\|_2 + 2H\kappa_1 \left\|\boldsymbol{u}_h - \theta_{\mathbb{J}\widehat{V}_{h+1}}\right\|_2 \\
&\leq 3H\kappa_1 \sqrt{\frac{36H^4(\log(H/\delta) + \widetilde{O}(d^2)) + 2\lambda C_\Theta^2}{\kappa K}}.
\end{aligned}$$

$\square$

J.3 PROOF OF THEOREM 4.1

In this section, we sketch the proof of Theorem 4.1 since the most components are identical to Theorem 3.2. We will focus on highlighting the difference for obtaining the tighter bound.

First of all, Recall in the first-order condition, we have

$$\nabla_\theta \left\{\sum_{k=1}^K \frac{\left[f(\theta, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k)\right]^2}{\widehat{\sigma}_h^2(s_h^k, a_h^k)} + \lambda \cdot \|\theta\|_2^2\right\}\Bigg|_{\theta=\widehat{\theta}_h} = 0, \quad \forall h \in [H].$$

Therefore, if we define the quantity $Z_h(\cdot, \cdot) \in \mathbb{R}^d$ as

$$Z_h(\theta|V, \sigma^2) = \sum_{k=1}^K \frac{\left[f(\theta, \phi_{h,k}) - r_{h,k} - V(s_{h+1}^k)\right]}{\sigma(s_h^k, a_h^k)} \frac{\nabla f(\theta, \phi_{h,k})}{\sigma(s_h^k, a_h^k)} + \lambda \cdot \theta, \quad \forall \theta \in \Theta, \|V\|_2 \leq H,$$

then we have

$$Z_h(\widehat{\theta}_h|\widehat{V}_{h+1}, \widehat{\sigma}_h^2) = 0.$$

According to the regression oracle (Line 8 of Algorithm 3), the estimated Bellman operator $\widehat{\mathcal{P}}_h$ maps $\widehat{V}_{h+1}$ to $\widehat{\theta}_h$, *i.e.* $\widehat{\mathcal{P}}_h\widehat{V}_{h+1} = f(\widehat{\theta}_h, \phi)$. Therefore (recall Definition D.1)

$$\begin{aligned}
&\mathcal{P}_h\widehat{V}_{h+1}(s, a) - \widehat{\mathcal{P}}_h\widehat{V}_{h+1}(s, a) = \mathcal{P}_h\widehat{V}_{h+1}(s, a) - f(\widehat{\theta}_h, \phi(s, a)) \\
&= f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s, a)) - f(\widehat{\theta}_h, \phi(s, a)) \\
&= \nabla f(\widehat{\theta}_h, \phi(s, a))\left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h\right) + \mathrm{Hot}_{h,1},
\end{aligned} \tag{31}$$

where we apply the first-order Taylor expansion for the differentiable function f at point $\widehat{\theta}_h$ and $\mathrm{Hot}_{h,1}$ is a higher-order term. Indeed, the following Lemma E.1 bounds the $\mathrm{Hot}_{h,1}$ term with $\widetilde{O}(\frac{1}{K})$.

Lemma J.5. *Recall the definition (from the above decomposition)* $\text{Hot}_{h,1} := f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi(s,a)) - f(\widehat{\theta}_h, \phi(s,a)) - \nabla f(\widehat{\theta}_h, \phi(s,a)) \left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right)$, *then with probability* $1 - \delta$,

$$|\text{Hot}_{h,1}| \leq \widetilde{O}(\frac{1}{K}), \quad \forall h \in [H].$$

Proof. The proof is identical to that of Lemma E.1 but with the help of Lemma J.2. $\square$

Next, according to the expansion of $Z_h(\theta|\widehat{V}_{h+1}, \widehat{\sigma}_h^2)$, we have

$$\nabla f(\widehat{\theta}_h, \phi(s,a)) \left(\theta_{\mathbb{T}\widehat{V}_{h+1}} - \widehat{\theta}_h \right) = I_1 + I_2 + I_3 + \text{Hot}_2, \tag{32}$$

where

$$\text{Hot}_2 := \nabla f(\widehat{\theta}_h, \phi(s,a)) \Lambda_h^{-1} \left[\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) + \lambda \theta_{\mathbb{T}\widehat{V}_{h+1}} \right]$$

$$\Delta_{\Lambda_h^s} = \sum_{k=1}^{K} \frac{\left(f(\widehat{\theta}_h, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \nabla_{\theta\theta}^2 f(\widehat{\theta}_h, \phi_{h,k})}{\widehat{\sigma}^2(s_h^k, a_h^k)}$$

$$\Lambda_h = \sum_{k=1}^{K} \frac{\nabla_\theta f(\widehat{\theta}_h, \phi_{h,k}) \nabla_\theta^\top f(\widehat{\theta}_{h,k}, \phi_{h,k})}{\widehat{\sigma}^2(s_h^k, a_h^k)} + \lambda I_d$$

$$\widetilde{R}_K(\theta_{\mathbb{T}\widehat{V}_{h+1}}) = \Delta_{\Lambda_h^s}(\widehat{\theta}_h - \theta_{\mathbb{T}\widehat{V}_{h+1}}) + R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})$$

where $R_K(\theta_{\mathbb{T}\widehat{V}_{h+1}})$ is the second order residual that is bounded by $\widetilde{O}(1/K)$ and

$$I_1 = \nabla f(\widehat{\theta}_h, \phi(s,a)) \Lambda_h^{-1} \sum_{k=1}^{K} \frac{\left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})}{\widehat{\sigma}_h^2(s_h^k, a_h^k)}$$

$$I_2 = \nabla f(\widehat{\theta}_h, \phi(s,a)) \Lambda_h^{-1} \sum_{k=1}^{K} \frac{\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - \widehat{V}_{h+1}(s_{h+1}^k) + V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})}{\widehat{\sigma}_h^2(s_h^k, a_h^k)}$$

$$I_3 = \nabla f(\widehat{\theta}_h, \phi(s,a)) \Lambda_h^{-1} \sum_{k=1}^{K} \frac{\left(f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - r_{h,k} - \widehat{V}_{h+1}(s_{h+1}^k) \right) \cdot \left(\nabla_\theta^\top f(\theta_{\mathbb{T}\widehat{V}_{h+1}}, \phi_{h,k}) - \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k}) \right)}{\widehat{\sigma}_h^2(s_h^k, a_h^k)}$$

Similar to the PFQL case, I_2, I_3, Hot_2 can be bounded to have order $O(1/K)$ via provably efficiency theorems in Section J.1 and in particular, the inclusion of $\sigma_{u,v}^2$ will not cause additional order in d.[13] Now we prove the result for the dominate term I_1.

Lemma J.6. *With probability* $1 - \delta$,

$$|I_1| \leq 4Hd \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Sigma_h^{-1}} \cdot C_{\delta, \log K} + \widetilde{O}(\frac{\kappa_1}{\sqrt{\kappa}K}),$$

where $C_{\delta, \log K}$ only contains Polylog terms.

Proof of Lemma J.6. First of all, by Cauchy–Schwarz inequality, we have

$$|I_1| \leq \left\| \nabla f(\widehat{\theta}_h, \phi(s,a)) \right\|_{\Lambda_h^{-1}} \cdot \left\| \sum_{k=1}^{K} \frac{\left(f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k) \right) \cdot \nabla_\theta^\top f(\widehat{\theta}_h, \phi_{h,k})}{\widehat{\sigma}_h^2(s_h^k, a_h^k)} \right\|_{\Lambda_h^{-1}}. \tag{33}$$

Recall that $\sigma_{u,v}^2(\cdot, \cdot) := \max\{1, f(v, \phi(\cdot, \cdot))_{[0, (H-h+1)^2]} - \left[f(u, \phi(\cdot, \cdot))_{[0, H-h+1]} \right]^2\}$.

---

[13]Note in Lemma L.11, we only have additive terms that has the same order has Lemma L.10.

Step1. Let the fixed $\theta \in \Theta$ be arbitrary and fixed u, v such that $\sigma_{u,v}^2(\cdot, \cdot) \geq \frac{1}{2}\sigma_{\boldsymbol{u}_h^\star, \boldsymbol{v}_h^\star}^2(\cdot, \cdot) = \frac{1}{2}\sigma_h^{\star 2}(\cdot, \cdot)$ and define $x_k(\theta, u, v) = \nabla_\theta f(\theta, \phi_{h,k})/\sigma_{u,v}(s_h^k, a_h^k)$. Next, define $G_{u,v}(\theta) = \sum_{k=1}^K \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top/\sigma_{u,v}^2(s_h^k, a_h^k) + \lambda I_d$, then $\|x_k\|_2 \leq \kappa_1$. Also denote $\eta_k := [f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k)]/\sigma_{u,v}(s_h^k, a_h^k)$, then $\mathbb{E}[\eta_k|s_h^k, a_h^k] = 0$ and

$$
\begin{aligned}
\mathrm{Var}[\eta_k|s_h^k, a_h^k] =& \frac{\mathrm{Var}[f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k)|s_h^k, a_h^k]}{\sigma_{u,v}^2(s_h^k, a_h^k)} \\
\leq& \frac{2\mathrm{Var}[f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi_{h,k}) - r_{h,k} - V_{h+1}^\star(s_{h+1}^k)|s_h^k, a_h^k]}{\sigma_h^{\star 2}(s_h^k, a_h^k)} \\
=& \frac{2[\mathrm{Var}_{P_h} V_{h+1}^\star](s_h^k, a_h^k)}{\sigma_h^{\star 2}(s_h^k, a_h^k)} \leq 2,
\end{aligned}
$$

then by Self-normalized Bernstein's inequality (Lemma L.4), with probability $1 - \delta$,

$$
\left\|\sum_{k=1}^K x_k(\theta, u, v)\eta_k\right\|_{G(\theta, u, v)^{-1}} \leq 16\sqrt{d\log\left(1 + \frac{K\kappa_1^2}{\lambda d}\right) \cdot \log\left(\frac{4K^2}{\delta}\right)} + 4\zeta\log\left(\frac{4K^2}{\delta}\right) \leq \widetilde{O}(\sqrt{d})
$$

where $|\eta_k| \leq \zeta$ with $\zeta = 2\max_{s,a,s'} \frac{|f(\theta_{\mathbb{T}V_{h+1}^\star}, \phi(s,a)) - r - V_{h+1}^\star(s')|}{\sigma_h^\star(s,a)}$ and the last inequality uses $\sqrt{d} \geq \widetilde{O}(\zeta)$.

Step2. Define $h(\theta, u, v) := \sum_{k=1}^K x_k(\theta, u, v)\eta_k(u, v)$ and $H(\theta, u, v) := \|h(\theta, u, v)\|_{G_{u,v}(\theta)^{-1}}$,

$$
\|h(\theta_1, u_1, v_1) - h(\theta_2, u_2, v_2)\|_2 \leq K\max_k \|(x_k \cdot \eta_k)(\theta_1, u_1, v_1) - (x_k \cdot \eta_k)(\theta_2, u_2, v_2)\|_2
$$

$$
\leq K\max_k \left\{ H\left|\frac{\nabla f(\theta_1, \phi_{h,k}) - \nabla f(\theta_2, \phi_{h,k})}{\sigma_{u_1, v_1}^2(s_h^k, a_h^k)}\right| + H\kappa_1\left|\frac{\sigma_{u_1, v_1}^2(s_h^k, a_h^k) - \sigma_{u_2, v_2}^2(s_h^k, a_h^k)}{\sigma_{u_1, v_1}^2(s_h^k, a_h^k)\sigma_{u_2, v_2}^2(s_h^k, a_h^k)}\right| \right\}
$$

$$
\leq KH\kappa_1 \|\theta_1 - \theta_2\|_2 + KH\kappa_1 \|\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2\|_2
$$

Furthermore,

$$
\left\|G_h(\theta_1, u_1, v_1)^{-1} - G_h(\theta_2, u_2, v_2)^{-1}\right\|_2 \leq \left\|G_h(\theta_1, u_1, v_1)^{-1}\right\|_2 \|G_h(\theta_1, u_1, v_1) - G_h(\theta_2, u_2, v_2)\|_2 \left\|G_h(\theta_2, u_2, v_2)^{-1}\right\|_2
$$

$$
\leq \frac{1}{\lambda^2} K\sup_k \left\|\frac{\nabla f(\theta_1, \phi_{h,k}) \cdot \nabla f(\theta_1, \phi_{h,k})^\top}{\sigma_{u_1, v_1}^2(s_h^k, a_h^k)} - \frac{\nabla f(\theta_2, \phi_{h,k}) \cdot \nabla f(\theta_2, \phi_{h,k})^\top}{\sigma_{u_2, v_2}^2(s_h^k, a_h^k)}\right\|_2
$$

$$
\leq \frac{1}{\lambda^2}\left(K\kappa_2\kappa_1 \|\theta_1 - \theta_2\|_2 + K\kappa_1^2 \|\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2\|_2\right)
$$

All the above imply

$$
|H(\theta_1, u_1, v_1) - H(\theta_2, u_2, v_2)| \leq \sqrt{|h(\theta_1, u_1, v_1)^\top G_{u_1, v_1}(\theta_1)^{-1}h(\theta_1, u_1, v_1) - h(\theta_2, u_2, v_2)^\top G_{u_2, v_2}(\theta_2)^{-1}h(\theta_2, u_2, v_2)|}
$$

$$
\leq \sqrt{\|h(\theta_1, u_1, v_1) - h(\theta_2, u_2, v_2)\|_2 \cdot \frac{1}{\lambda} \cdot KH\kappa_1} + \sqrt{KH\kappa_1 \cdot \|G_{u_1, v_1}(\theta_1)^{-1} - G_{u_2, v_2}(\theta_2)^{-1}\|_2 \cdot KH\kappa_1}
$$

$$
+ \sqrt{(KH\kappa_1 \cdot \frac{1}{\lambda}) \cdot \|h(\theta_1, u_1, v_1) - h(\theta_2, u_2, v_2)\|_2}
$$

$$
\leq 2\sqrt{KH\kappa_1(\|\theta_1 - \theta_2\|_2 + \|\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2\|_2) \cdot \frac{1}{\lambda} \cdot KH\kappa_1} + \sqrt{K^2H^2\kappa_1^2 \cdot \frac{K\kappa_1}{\lambda^2}\left(\kappa_2 \|\theta_1 - \theta_2\|_2 + \kappa_1 \|\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2\|_2\right)}
$$

$$
\leq \left(\sqrt{4K^2H^2\kappa_1^2/\lambda} + \sqrt{K^3H^2\kappa_1^3\kappa_2/\lambda^2}\right)\sqrt{\|\theta_1 - \theta_2\|_2} + \left(\sqrt{4K^2H^2\kappa_1^2/\lambda} + \sqrt{K^3H^2\kappa_1^4/\lambda^2}\right)\sqrt{\|\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2\|_2}
$$

note

$$
|\sigma_{u_1, v_1}^2(s, a) - \sigma_{u_2, v_2}^2(s, a)| \leq |f(v_1, \phi(s, a)) - f(v_2, \phi(s, a))| + 2H|f(u_1, \phi(s, a)) - f(u_2, \phi(s, a))|
$$

$$
\leq \kappa_1 \|v_1 - v_2\|_2 + 2H\kappa_1 \|u_1 - u_2\|_2,
$$

Then a ϵ-covering net of $\{H(\theta, u, v)\}$ can be constructed by the union of covering net for θ, u, v and by Lemma L.8, the covering number $\mathcal{N}_\epsilon$ satisfies (where $\widetilde{O}$ absorbs Polylog terms)

$$
\log \mathcal{N}_\epsilon \leq \widetilde{O}(d)
$$

Step3. First note by definition in Step2

$$\left\| \sum_{k=1}^{K} \frac{\left(f(\theta_{\mathbb{T}V_{h+1}^{\star}}, \phi_{h,k}) - r_{h,k} - V_{h+1}^{\star}(s_{h+1}^{k}) \right) \cdot \nabla_{\theta}^{\top} f(\widehat{\theta}_{h}, \phi_{h,k})}{\widehat{\sigma}_{h}^{2}(s_{h}^{k}, a_{h}^{k})} \right\|_{\Lambda_{h}^{-1}} = H(\widehat{\theta}_{h}, \boldsymbol{u}_{h}, \boldsymbol{v}_{h})$$

Now choosing $\epsilon = O(1/K)$ in Step2 and union bound over the covering number in Step2, we obtain with probability $1 - \delta$ (recall $\sqrt{d} \geq \widetilde{O}(\zeta)$),

$$H(\widehat{\theta}_{h}, \boldsymbol{u}_{h}, \boldsymbol{v}_{h}) \leq 16 \sqrt{d \log\left(1 + \frac{K\kappa_{1}^{2}}{\lambda d}\right) \cdot \left[\log\left(\frac{4K^{2}}{\delta}\right) + \widetilde{O}(d)\right] + 4\zeta \left[\log\left(\frac{4K^{2}}{\delta}\right) + \widetilde{O}(d)\right] + O(\frac{1}{K})}$$

$$\leq \widetilde{O}(d) + O(\frac{1}{K})$$

where we absorb all the Polylog terms. Combing above with equation 33, we obtain with probability $1 - \delta$,

$$|I_1| \leq \left\| \nabla f(\widehat{\theta}_{h}, \phi(s, a)) \right\|_{\Lambda_{h}^{-1}} \cdot H(\widehat{\theta}_{h}, \boldsymbol{u}_{h}, \boldsymbol{v}_{h})$$

$$\leq \left\| \nabla f(\widehat{\theta}_{h}, \phi(s, a)) \right\|_{\Lambda_{h}^{-1}} \cdot \left[\widetilde{O}(d) + O(\frac{1}{K}) \right]$$

$$\leq \widetilde{O} \left(d \left\| \nabla f(\widehat{\theta}_{h}, \phi(s, a)) \right\|_{\Lambda_{h}^{-1}} \right) + \widetilde{O}(\frac{\kappa_{1}}{\sqrt{\kappa}K}),$$

$\square$

Combing dominate term I_1 (via Lemma J.6) and all other higher order terms we can obtain the first result together with Lemma D.3.

The proof of the second result is also very similar to the proofs in Section F.2. Concretely, when picking $\pi = \pi^{\star}$, we can convert the quantity

$$\sqrt{\nabla_{\theta}^{\top} f(\widehat{\theta}_{h}, \phi(s_{h}, a_{h})) \Lambda_{h}^{-1} \nabla_{\theta} f(\widehat{\theta}_{h}, \phi(s_{h}, a_{h}))}$$

to

$$\sqrt{\nabla_{\theta}^{\top} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h})) \Lambda_{h}^{-1} \nabla_{\theta} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h}))}$$

using Theorem J.3, and convert

$$\sqrt{\nabla_{\theta}^{\top} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h})) \Lambda_{h}^{-1} \nabla_{\theta} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h}))}$$

to

$$\sqrt{\nabla_{\theta}^{\top} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h})) \Lambda_{h}^{\star-1} \nabla_{\theta} f(\theta_{h}^{\star}, \phi(s_{h}, a_{h}))}$$

using Lemma J.4.

K THE LOWER BOUND

Theorem K.1 (Restatement of Theorem 4.2). *Specifying the model to have linear representation* $f = \langle \theta, \phi \rangle$*. There exist a pair of universal constants* $c, c' > 0$ *such that given dimension* d*, horizon* H *and sample size* $K > c'd^3$*, one can always find a family of MDP instances such that for any algorithm* $\widehat{\pi}$

$$\inf_{\widehat{\pi}} \sup_{M \in \mathcal{M}} \mathbb{E}_{M} \left[v^{\star} - v^{\widehat{\pi}} \right] \geq c \sqrt{d} \cdot \sum_{h=1}^{H} \mathbb{E}_{\pi^{\star}} \left[\sqrt{\nabla_{\theta}^{\top} f(\theta_{h}^{\star}, \phi(\cdot, \cdot))(\Lambda_{h}^{\star, p})^{-1} \nabla_{\theta} f(\theta_{h}^{\star}, \phi(\cdot, \cdot))} \right], \quad (34)$$

where $\Lambda_{h}^{\star, p} = \mathbb{E}\left[\sum_{k=1}^{K} \frac{\nabla_{\theta} f(\theta_{h}^{\star}, \phi(s_{h}^{k}, a_{h}^{k})) \cdot \nabla_{\theta} f(\theta_{h}^{\star}, \phi(s_{h}^{k}, a_{h}^{k}))^{\top}}{\text{Var}_{h}(V_{h+1}^{\star})(s_{h}^{k}, a_{h}^{k})} \right]$.

Remark K.2. *Note Theorem 4.2 is a valid lower bound for comparison. This is because the upper bound result holds true for all model* f *such that the corresponding* $\mathcal{F}$ *satisfies Assumption 2.1, 2.3. Therefore, for the lower bound construction it suffices to find one model* f *such that the lower bound equation 34 holds. Here we simply choose the linear function approximation.*

K.1   Regarding the proof of lower bound

The proof of Theorem 4.2 can be done via a reduction to linear function approximation lower bound. In fact, it can be directly obtained from Theorem 3.5 of Yin et al. (2022), and the original proof comes from Theorem 2 of Zanette et al. (2021).

Concretely, all the proofs in Theorem 3.5 of Yin et al. (2022) follows and the only modification is to replace

$$\sqrt{\mathbb{E}_{\pi^\star}[\phi]^\top (\Lambda_h^\star)^{-1} \mathbb{E}_{\pi^\star}[\phi]} \leq \frac{1}{2} \left\| \phi\left(+1, u_h\right)\right\|_{\left(\Lambda_h^{\star,p}\right)^{-1}} + \frac{1}{2} \left\| \phi\left(-1, u_h\right)\right\|_{\left(\Lambda_h^{\star,p}\right)^{-1}}$$

in Section E.5 by

$$\mathbb{E}_{\pi^\star}\left[\sqrt{\phi(\cdot,\cdot)^\top (\Lambda_h^{\star,p})^{-1} \phi(\cdot,\cdot)}\right] = \frac{1}{2}\left\|\phi(+1, u_h)\right\|_{\left(\Lambda_h^{\star,p}\right)^{-1}} + \frac{1}{2}\left\|\phi(-1, u_h)\right\|_{\left(\Lambda_h^{\star,p}\right)^{-1}},$$

and the final result holds with $\phi(\cdot,\cdot) = \nabla_\theta f(\theta_h^\star, \phi(\cdot,\cdot))$ by the reduction $f = \langle \theta, \phi \rangle$.

L   Auxiliary lemmas

Lemma L.1 (k-th Order Mean Value Form of Taylor's Expansion)**.** *Let $k \geq 1$ be an integer and let function $f : \mathbb{R}^d \to \mathbb{R}$ be k times differentiable and continuous over the compact domain $\Theta \subset \mathbb{R}^d$. Then for any $x, \theta \in \Theta$, there exists ξ in the line segment of x and θ, such that*

$$f(x) - f(\theta) = \nabla f(\theta)^\top (x - \theta) + \frac{1}{2!}(x - \theta)^\top \nabla_{\theta\theta}^2 f(\theta)(x - \theta) + \ldots + \frac{1}{(k-1)!}\nabla^{k-1} f(\theta) \left(\bigotimes(x - \theta)\right)^{k-1}$$

$$+ \frac{1}{k!}\nabla^k f(\xi) \left(\bigotimes(x - \theta)\right)^k.$$

Here $\nabla^k f(\theta)$ denotes k-dimensional tensor and $\bigotimes$ denotes tensor product.

Lemma L.2 (Vector Hoeffding's Inequality)**.** *Let $X = (X_1, \ldots, X_d)$ be d-dimensional vector Random Variable with $E[X] = 0$ and $\|X\|_2 \leq R$. $X^{(1)}, \ldots, X^{(n)}$'s are n samples. Then with probability $1 - \delta$,*

$$\left\| \frac{1}{n} \sum_{i=1}^n X^{(i)} \right\|_2 \leq \sqrt{\frac{4dR^2}{n} \log(\frac{d}{\delta})}.$$

Proof of Lemma L.2. Since $\|X\|_2 \leq R$ implies $|X_j| \leq R$, by the univariate Hoeffding's inequality, for a fixed $j \in \{1, ..., d\}$, denote $Y_j := \frac{1}{n} \sum_{i=1}^n X_j^{(i)}$. Then with probability $1 - \delta$ (note $|X_j^{(i)}| \leq R$),

$$\mathbb{P}\left(|Y_j| \geq 2\sqrt{\frac{R^2}{n} \log(\frac{1}{\delta})}\right) \leq \delta.$$

By a union bound,

$$\mathbb{P}\left(\exists\, i\ s.t.\ |Y_j| \geq 2\sqrt{\frac{R^2}{n} \log(\frac{1}{\delta})}\right) \leq d\delta \Leftrightarrow \mathbb{P}\left(\forall\, i\ \ |Y_j| \leq 2\sqrt{\frac{R^2}{n} \log(\frac{1}{\delta})}\right) \geq 1 - d\delta$$

$$\Leftrightarrow \mathbb{P}\left(\forall\, i\ \ Y_j^2 \leq \frac{4R^2}{n} \log(\frac{1}{\delta})\right) \geq 1 - d\delta \Rightarrow \mathbb{P}\left(\|Y\|_2 \leq \sqrt{\frac{4dR^2}{n} \log(\frac{1}{\delta})}\right) \geq 1 - d\delta$$

$$\Leftrightarrow \mathbb{P}\left(\|Y\|_2 \leq \sqrt{\frac{4dR^2}{n} \log(\frac{d}{\delta})}\right) \geq 1 - \delta.$$

$\square$

Lemma L.3 (Hoeffding inequality for self-normalized martingales (Abbasi-Yadkori et al., 2011))**.** *Let $\{\eta_t\}_{t=1}^\infty$ be a real-valued stochastic process. Let $\{\mathcal{F}_t\}_{t=0}^\infty$ be a filtration, such that η_t is $\mathcal{F}_t$-measurable. Assume η_t also satisfies η_t given $\mathcal{F}_{t-1}$ is zero-mean and R-subgaussian, i.e.*

$$\forall \lambda \in \mathbb{R}, \quad \mathbb{E}\left[e^{\lambda \eta_t} \mid \mathcal{F}_{t-1}\right] \leq e^{\lambda^2 R^2 / 2}$$

Let $\{x_t\}_{t=1}^\infty$ be an $\mathbb{R}^d$-valued stochastic process where x_t is $\mathcal{F}_{t-1}$ measurable and $\|x_t\| \le L$. Let $\Lambda_t = \lambda I_d + \sum_{s=1}^t x_s x_s^\top$. Then for any $\delta > 0$, with probability $1 - \delta$, for all $t > 0$,

$$\left\| \sum_{s=1}^t x_s \eta_s \right\|_{\Lambda_t^{-1}}^2 \le 8R^2 \cdot \frac{d}{2} \log\left(\frac{\lambda + tL}{\lambda \delta} \right).$$

Lemma L.4 (Bernstein inequality for self-normalized martingales (Zhou et al., 2021)). *Let $\{\eta_t\}_{t=1}^\infty$ be a real-valued stochastic process. Let $\{\mathcal{F}_t\}_{t=0}^\infty$ be a filtration, such that η_t is $\mathcal{F}_t$-measurable. Assume η_t also satisfies*

$$|\eta_t| \le R, \mathbb{E}\left[\eta_t \mid \mathcal{F}_{t-1}\right] = 0, \mathbb{E}\left[\eta_t^2 \mid \mathcal{F}_{t-1}\right] \le \sigma^2.$$

Let $\{x_t\}_{t=1}^\infty$ be an $\mathbb{R}^d$-valued stochastic process where x_t is $\mathcal{F}_{t-1}$ measurable and $\|x_t\| \le L$. Let $\Lambda_t = \lambda I_d + \sum_{s=1}^t x_s x_s^\top$. Then for any $\delta > 0$, with probability $1 - \delta$, for all $t > 0$,

$$\left\| \sum_{s=1}^t \mathbf{x}_s \eta_s \right\|_{\mathbf{\Lambda}_t^{-1}} \le 8\sigma \sqrt{d \log\left(1 + \frac{tL^2}{\lambda d}\right) \cdot \log\left(\frac{4t^2}{\delta}\right)} + 4R \log\left(\frac{4t^2}{\delta}\right)$$

Lemma L.5. *Let $\nabla f(\theta, \phi(\cdot, \cdot)) : \mathcal{S} \times \mathcal{A} \to \mathbb{R}^d$ be a bounded function s.t. $\sup_{\theta \in \Theta} \|\nabla f(\theta, \phi(\cdot, \cdot))\|_2 \le \kappa_1$. If K satisfies*

$$K \ge \max\left\{ 512 \frac{\kappa_1^4}{\kappa^2} \left(\log(\frac{2d}{\delta}) + d \log(1 + \frac{4\kappa_1 B^2 \kappa_2 C_\Theta K^3}{\lambda^2}) \right), \frac{4\lambda}{\kappa} \right\}$$

Then with probability at least $1 - \delta$, for all $\|u\|_2 \le B$ simultaneously, it holds that

$$\|u\|_{\Sigma_h^{-1}} \le \frac{2B}{\sqrt{\kappa K}} + O(\frac{1}{K})$$

where $\Sigma_h = \sum_{k=1}^K \nabla f(\widehat{\theta}_h, \phi(s_h^k, a_h^k)) \cdot \nabla f(\widehat{\theta}_h, \phi(s_h^k, a_h^k))^\top + \lambda I_d$.

Proof of Lemma L.5. For a fixed θ, define $\bar{G} = \sum_{k=1}^K \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top + \lambda I_d$, and $G = \mathbb{E}_\mu[\nabla f(\theta, \phi(s_h, a_h)) \cdot \nabla f(\theta, \phi(s_h, a_h))^\top]$, then by Lemma H.5. of Min et al. (2021), as long as

$$K \ge \max\{512 \kappa_1^4 \left\|G^{-1}\right\|^2 \log(\frac{2d}{\delta}), 4\lambda \left\|G^{-1}\right\|_2\}, \tag{35}$$

then with probability $1 - \delta$, for all $u \in \mathbb{R}^d$ simultaneously, $\|u\|_{\bar{G}^{-1}} \le \frac{2}{\sqrt{K}} \|u\|_{G^{-1}}$. As a corollary, if we constraint u to the subspace $\|u\|_2 \le B$, then we have: with probability $1 - \delta$, for all $\{u \in \mathbb{R}^d : \|u\|_2 \le B\}$ simultaneously,

$$\|u\|_{\bar{G}^{-1}} \le \frac{2}{\sqrt{K}} \|u\|_{G^{-1}} = \frac{2}{\sqrt{K}} \sqrt{u^\top G^{-1} u} \le \frac{2B\sqrt{\|G^{-1}\|_2}}{\sqrt{K}}. \tag{36}$$

Next, for any θ, define

$$h_u(\theta) := \|u\|_{\bar{G}^{-1}} = \sqrt{u^\top \bar{G}^{-1} u} = \sqrt{u^\top \left(\sum_{k=1}^K \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top + \lambda I_d \right)^{-1} u}$$

and $\bar{G}(\theta) = \sum_{k=1}^K \nabla f(\theta, \phi(s_h^k, a_h^k)) \cdot \nabla f(\theta, \phi(s_h^k, a_h^k))^\top + \lambda I_d$, we have for any θ_1, θ_2

$$\left\| \bar{G}(\theta_1) - \bar{G}(\theta_2) \right\|_2 \le \left\| \sum_{k=1}^K \left(\nabla f(\theta_1, \phi(s_h^k, a_h^k)) - \nabla f(\theta_2, \phi(s_h^k, a_h^k)) \right) \cdot \nabla f(\theta_1, \phi(s_h^k, a_h^k))^\top \right\|$$

$$+ \left\| \sum_{k=1}^K \nabla f(\theta_2, \phi(s_h^k, a_h^k)) \left(\nabla f(\theta_1, \phi(s_h^k, a_h^k)) - \nabla f(\theta_2, \phi(s_h^k, a_h^k)) \right)^\top \right\|$$

$$\le K \kappa_2 \kappa_1 \|\theta_1 - \theta_2\|_2 + K \kappa_2 \kappa_1 \|\theta_1 - \theta_2\|_2 \le 2K \kappa_2 \kappa_1 \|\theta_1 - \theta_2\|_2.$$

Use the basic inequality for $a, b > 0 \Rightarrow |\sqrt{a} - \sqrt{b}| \leq \sqrt{|a-b|}$,

$$\sup_u |h_u(\theta_1) - h_u(\theta_2)| \leq \sup_u \sqrt{\left|u^\top \left(\bar{G}(\theta_1)^{-1} - \bar{G}(\theta_2)^{-1}\right) u\right|} \leq \sqrt{B^2 \cdot \left\|\bar{G}(\theta_1)^{-1} - \bar{G}(\theta_2)^{-1}\right\|_2}$$

$$\leq \sqrt{B^2 \cdot \left\|\bar{G}(\theta_1)^{-1}\right\|_2 \left\|\bar{G}(\theta_1) - \bar{G}(\theta_2)\right\|_2 \left\|\bar{G}(\theta_2)^{-1}\right\|_2}$$

$$\leq \sqrt{B^2 \frac{1}{\lambda} 2K\kappa_2\kappa_1 \left\|\theta_1 - \theta_2\right\|_2 \frac{1}{\lambda}} = \sqrt{\frac{2B^2 K \kappa_1 \kappa_2 \left\|\theta_1 - \theta_2\right\|_2}{\lambda^2}}$$

Therefore, the ϵ-covering net of $\{h(\theta) : \theta \in \Theta\}$ is implies by the $\frac{\lambda^2 \epsilon^2}{2KB^2\kappa_1\kappa_2}$-covering net of $\{\theta : \theta \in \Theta\}$, so by Lemma L.8, the covering number $\mathcal{N}_\epsilon$ satisfies

$$\log \mathcal{N}_\epsilon \leq d\log(1 + \frac{4B^2 K \kappa_1 \kappa_2 C_\Theta}{\lambda^2 \epsilon^2}).$$

Select $\theta = \widehat{\theta}_h$. Choose $\epsilon = O(1/K)$ and by a union bound over equation 36 to get with probability $1 - \delta$, for all $\|u\|_2 \leq B$ (note By Assumption 2.3 $\left\|G^{-1}\right\|_2 \leq 1/\kappa$),

$$\|u\|_{\Sigma_h^{-1}} \leq \frac{2B}{\sqrt{\kappa K}} + O(\frac{1}{K})$$

if (union bound over the condition equation 35)

$$K \geq \max\left\{512\frac{\kappa_1^4}{\kappa^2}\left(\log(\frac{2d}{\delta}) + d\log(1 + \frac{4\kappa_1 B^2 \kappa_2 C_\Theta K^3}{\lambda^2})\right), \frac{4\lambda}{\kappa}\right\}$$

where this condition is satisfied by the Lemma statement.

$\square$

Lemma L.6. *let $\phi : \mathcal{S} \times \mathcal{A} \to \mathbb{R}^d$ satisfies $\|\phi(s,a)\| \leq C$ for all $s, a \in \mathcal{S} \times \mathcal{A}$. For any $K > 0, \lambda > 0$, define $\bar{G}_K = \sum_{k=1}^K \phi(s_k, a_k)\phi(s_k, a_k)^\top + \lambda I_d$ where (s_k, a_k)'s are i.i.d samples from some distribution ν. Then with probability $1 - \delta$,*

$$\left\|\frac{\bar{G}_K}{K} - \mathbb{E}_\nu\left[\frac{\bar{G}_K}{K}\right]\right\| \leq \frac{4\sqrt{2}C^2}{\sqrt{K}}\left(\log\frac{2d}{\delta}\right)^{1/2}.$$

Proof of Lemma L.6. See Lemma H.5 of Yin et al. (2022) or Lemma H.4 of Lemma Min et al. (2021) for details. $\square$

Lemma L.7 (Lemma H.4 in Yin et al. (2022))**.** *Let Λ_1 and $\Lambda_2 \in \mathbb{R}^{d \times d}$ are two positive semi-definite matrices. Then:*

$$\left\|\Lambda_1^{-1}\right\| \leq \left\|\Lambda_2^{-1}\right\| + \left\|\Lambda_1^{-1}\right\| \cdot \left\|\Lambda_2^{-1}\right\| \cdot \|\Lambda_1 - \Lambda_2\|$$

and

$$\|\phi\|_{\Lambda_1^{-1}} \leq \left[1 + \sqrt{\left\|\Lambda_2^{-1}\right\| \|\Lambda_2\| \cdot \left\|\Lambda_1^{-1}\right\| \cdot \|\Lambda_1 - \Lambda_2\|}\right] \cdot \|\phi\|_{\Lambda_2^{-1}}.$$

for all $\phi \in \mathbb{R}^d$.

L.1 COVERING ARGUMENTS

Lemma L.8. *(Covering Number of Euclidean Ball) For any $\epsilon > 0$, the ϵ-covering number of the Euclidean ball in $\mathbb{R}^d$ with radius $R > 0$ is upper bounded by $(1 + 2R/\epsilon)^d$.*

Lemma L.9. *Define $\mathcal{V}$ to be the class mapping $\mathcal{S}$ to $\mathbb{R}$ with the parametric form*

$$V(\cdot) := \min\{\max_a f(\theta, \phi(\cdot, a)) - \sqrt{\nabla f(\theta, \phi(\cdot, a))^\top A \cdot \nabla f(\theta, \phi(\cdot, a))}, H\}$$

where the parameter spaces are $\{\theta : \|\theta\|_2 \leq C_\Theta\}$ and $\{A : \|A\|_2 \leq B\}$. Let $\mathcal{N}_\epsilon^\mathcal{V}$ be the covering number of ϵ-net with respect to l_∞ distance, then we have

$$\log \mathcal{N}_\epsilon^\mathcal{V} \leq d\log\left(1 + \frac{8C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2}\right) + d^2\log\left(1 + \frac{8\sqrt{d}B\kappa_1^2}{\epsilon^2}\right).$$

Proof of Lemma L.9.

$$\sup_s |V_1(s) - V_2(s)|$$

$$\leq \sup_{s,a} \left| f(\theta_1, \phi(s,a)) - \sqrt{\nabla f(\theta_1, \phi(s,a))^\top A_1 \cdot \nabla f(\theta_1, \phi(s,a))} - f(\theta_2, \phi(s,a)) + \sqrt{\nabla f(\theta_2, \phi(s,a))^\top A_2 \cdot \nabla f(\theta_2, \phi(s,a))} \right|$$

$$= \sup_{s,a} \left| \nabla f(\xi, \phi(s,a)) \cdot (\theta_1 - \theta_2) - \sqrt{\nabla f(\theta_1, \phi(s,a))^\top A_1 \cdot \nabla f(\theta_1, \phi(s,a))} + \sqrt{\nabla f(\theta_2, \phi(s,a))^\top A_2 \cdot \nabla f(\theta_2, \phi(s,a))} \right|$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + \sup_{s,a} \left| \sqrt{\nabla f(\theta_1, \phi(s,a))^\top A_1 \cdot \nabla f(\theta_1, \phi(s,a))} - \sqrt{\nabla f(\theta_2, \phi(s,a))^\top A_2 \cdot \nabla f(\theta_2, \phi(s,a))} \right|$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + \sup_{s,a} \sqrt{\left| [\nabla f(\theta_1, \phi(s,a)) - \nabla f(\theta_2, \phi(s,a))]^\top A_1 \cdot \nabla f(\theta_1, \phi(s,a)) \right|}$$

$$+ \sup_{s,a} \sqrt{\left| \nabla f(\theta_2, \phi(s,a))^\top (A_1 - A_2) \cdot \nabla f(\theta_1, \phi(s,a)) \right|} + \sup_{s,a} \sqrt{\left| \nabla f(\theta_2, \phi(s,a))^\top A_2 \cdot [\nabla f(\theta_1, \phi(s,a)) - \nabla f(\theta_2, \phi(s,a))] \right|}$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 2 \sup_{s,a} \sqrt{\|\nabla f(\theta_1, \phi(s,a)) - \nabla f(\theta_2, \phi(s,a))\|_2 \cdot B \cdot \kappa_1} + \sqrt{\kappa_1^2 \|A_1 - A_2\|_2}$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 2 \sup_{s,a} \sqrt{\|\nabla f(\theta_1, \phi(s,a)) - \nabla f(\theta_2, \phi(s,a))\|_2 \cdot B \cdot \kappa_1} + \sqrt{\kappa_1^2 \|A_1 - A_2\|_2}$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 2 \sup_{s,a} \sqrt{\|\nabla f(\theta_1, \phi(s,a))\|_2 \cdot \|\theta_1 - \theta_2\|_2 \cdot B \cdot \kappa_1} + \sqrt{\kappa_1^2 \|A_1 - A_2\|_2}$$

$$\leq \kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 2 \sqrt{\kappa_2 \cdot \|\theta_1 - \theta_2\|_2 \cdot B \cdot \kappa_1} + \sqrt{\kappa_1^2 \|A_1 - A_2\|_2}$$

$$\leq \left(\kappa_1 \sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2} \right) \sqrt{\|\theta_1 - \theta_2\|_2} + \kappa_1 \sqrt{\|A_1 - A_2\|_2} \leq \left(\kappa_1 \sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2} \right) \sqrt{\|\theta_1 - \theta_2\|_2} + \kappa_1 \sqrt{\|A_1 - A_2\|_F}$$

Here $\|\cdot\|_F$ is Frobenius norm. Let $\mathcal{C}_\theta$ be the $\frac{\epsilon^2}{4(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}$-net of space $\{\theta : \|\theta\|_2 \leq C_\Theta\}$ and $\mathcal{C}_w$ be the $\frac{\epsilon^2}{4\kappa_1^2}$-net of the space $\{A : \|A\|_F \leq \sqrt{d}B\}$, then by Lemma L.8,

$$|\mathcal{C}_w| \leq \left(1 + \frac{8C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2} \right)^d, \quad |\mathcal{C}_A| \leq \left(1 + \frac{8\sqrt{d}B\kappa_1^2}{\epsilon^2} \right)^{d^2}$$

Therefore, the covering number of space $\mathcal{V}$ satisfies

$$\log \mathcal{N}_\epsilon^{\mathcal{V}} \leq \log(|\mathcal{C}_w| \cdot |\mathcal{C}_A|) \leq d \log \left(1 + \frac{8C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2} \right) + d^2 \log \left(1 + \frac{8\sqrt{d}B\kappa_1^2}{\epsilon^2} \right)$$

$\square$

Lemma L.10 (Covering of $\mathbb{E}_\mu(X(g, V, f))$)**.** *Define*

$$X(\theta, \theta') := (f(\theta, \phi(s,a)) - r - V_{\theta'}(s'))^2 - (f_{V_{\theta'}}(s,a) - r - V_{\theta'}(s'))^2,$$

where $f_V := \mathcal{P}_h V + \delta_V$ and $V(s)$ has form $V_\theta(s)$ that belongs to $\mathcal{V}$ (as defined in Lemma L.9). Here $X(\theta, \theta')$ is a function of s, a, r, s' as well, and we suppress the notation for conciseness only. Then the function class $\mathcal{H} = \{h(\theta, \theta') := \mathbb{E}_\mu[X(\theta, \theta')] \mid \|\theta\|_2 \leq C_\Theta, V_\theta \in \mathcal{V}\}$ has the covering number of $(\epsilon + 4H\epsilon_{\mathcal{F}})$-net bounded by

$$d\log(1 + \frac{24C_\Theta(H+1)\kappa_1}{\epsilon}) + d\log \left(1 + \frac{288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2} \right) + d^2 \log \left(1 + \frac{288H^2\sqrt{d}B\kappa_1^2}{\epsilon^2} \right).$$

Proof of Lemma L.10. First of all,

$$X(\theta, \theta') = f(\theta, \phi(s,a))^2 - f_{V_{\theta'}}(s,a)^2 - 2f(\theta, \phi(s,a)) \cdot (r + V_{\theta'}(s')) + 2f_{V_{\theta'}}(s,a) \cdot (r + V_{\theta'}(s')),$$

For any $(\theta_1, \theta_1'), (\theta_2, \theta_2')$,

$$
\begin{aligned}
|X(\theta_1, \theta_1') - X(\theta_2, \theta_2')| &\le |f(\theta_1, \phi(s,a))^2 - f(\theta_2, \phi(s,a))^2| \\
&+ |f_{V_{\theta_1'}}(s,a)^2 - f_{V_{\theta_2'}}(s,a)^2| + 2|f_{V_{\theta_1'}}(s,a) - f_{V_{\theta_2'}}(s,a)| \cdot (r + V_{\theta_1'}(s')) \\
&+ 2f_{V_{\theta_2'}}(s,a) \cdot |V_{\theta_1'}(s') - V_{\theta_2'}(s')| + 2|f(\theta_1, \phi(s,a)) - f(\theta_2, \phi(s,a))| \cdot (r + V_{\theta_1'}(s')) \\
&+ 2|f(\theta_2, \phi(s,a))| \cdot |V_{\theta_1'}(s') - V_{\theta_2'}(s')| \\
&\le 2H \cdot |f(\theta_1, \phi(s,a)) - f(\theta_2, \phi(s,a))| + 2H \cdot |f_{V_{\theta_1'}}(s,a) - f_{V_{\theta_2'}}(s,a)| \\
&+ 4H \cdot |V_{\theta_1'}(s') - V_{\theta_2'}(s')| + 4(H+1) \cdot |f(\theta_1, \phi(s,a)) - f(\theta_2, \phi(s,a))| \\
&\le (6H+1) \cdot |f(\theta_1, \phi(s,a)) - f(\theta_2, \phi(s,a))| + 2H \max_{s'} |V_{\theta_1'}(s') - V_{\theta_2'}(s')| + 4H\epsilon_{\mathcal{F}} \\
&+ 4H \cdot |V_{\theta_1'}(s') - V_{\theta_2'}(s')| \\
&\le (6H+1)\|\nabla f(\xi, \phi(s,a))\|_2 \cdot \|\theta_1 - \theta_2\|_2 + 6H \left\| V_{\theta_1'} - V_{\theta_2'} \right\|_\infty + 4H\epsilon_{\mathcal{F}} \\
&\le (6H+1)\kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 6H \left\| V_{\theta_1'} - V_{\theta_2'} \right\|_\infty + 4H\epsilon_{\mathcal{F}}
\end{aligned}
$$

where the second inequality comes from $f_V = \mathcal{P}_h V + \delta_V$. Note the above holds true for all s, a, r, s', therefore it implies

$$
\begin{aligned}
|\mathbb{E}_\mu[X(\theta_1, \theta_1')] - \mathbb{E}_\mu[X(\theta_2, \theta_2')]| &\le \sup_{s,a,s'} |X(\theta_1, \theta_1') - X(\theta_2, \theta_2')| \\
&\le (6H+1)\kappa_1 \cdot \|\theta_1 - \theta_2\|_2 + 6H \left\| V_{\theta_1'} - V_{\theta_2'} \right\|_\infty + 4H\epsilon_{\mathcal{F}}
\end{aligned}
$$

Now let $\mathcal{C}_1$ be the $\frac{\epsilon}{12(H+1)\kappa_1}$-net of $\{\theta : \|\theta\|_2 \le C_\Theta\}$ and $\mathcal{C}_2$ be the $\epsilon/6H$-net of $\mathcal{V}$, applying Lemma L.8 and Lemma L.9 to obtain

$$
\log|\mathcal{C}_1| \le d\log(1 + \frac{24C_\Theta(H+1)\kappa_1}{\epsilon}), \quad \log|\mathcal{C}_2| \le d\log\left(1 + \frac{288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2}\right) + d^2\log\left(1 + \frac{288H^2\sqrt{d}B\kappa_1^2}{\epsilon^2}\right)
$$

which implies the covering number of $\mathcal{H}$ to be bounded by

$$
\log|\mathcal{C}_1| \cdot |\mathcal{C}_2| \le d\log(1 + \frac{24C_\Theta(H+1)\kappa_1}{\epsilon}) + d\log\left(1 + \frac{288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2}\right) + d^2\log\left(1 + \frac{288H^2\sqrt{d}B\kappa_1^2}{\epsilon^2}\right).
$$

$\square$

Lemma L.11. *Denote* $\sigma_{u,v}^2(\cdot, \cdot) := \max\{1, f(v, \phi(\cdot, \cdot))_{[0, (H-h+1)^2]} - \left[f(u, \phi(\cdot, \cdot))_{[0, H-h+1]}\right]^2\}$ *and define*

$$
\bar{X}(\theta, \theta', u, v) := \frac{(f(\theta, \phi(s,a)) - r - V_{\theta'}(s'))^2 - (f_{V_{\theta'}}(s,a) - r - V_{\theta'}(s'))^2}{\sigma_{u,v}^2(s,a)},
$$

where $f_V := \mathcal{P}_h V$ *and* $V(s)$ *has form* $V_\theta(s)$ *that belongs to* $\mathcal{V}$ *(as defined in Lemma L.9). Here* $\bar{X}(\theta, \theta', u, v)$ *is a function of* s, a, r, s' *as well, and we suppress the notation for conciseness only. Then the function class* $\mathcal{H} = \{h(\theta, \theta', u, v) := \mathbb{E}_\mu[\bar{X}(\theta, \theta', u, v)] \mid \|\theta\|_2 \le C_\Theta, V_\theta \in \mathcal{V}\}$ *has the covering number of* ϵ*-net bounded by*

$$
d\log(1 + \frac{24C_\Theta(H+1)\kappa_1}{\epsilon}) + d\log\left(1 + \frac{288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B\kappa_1\kappa_2})^2}{\epsilon^2}\right) + d^2\log\left(1 + \frac{288H^2\sqrt{d}B\kappa_1^2}{\epsilon^2}\right)
$$

$$
+ d\log(1 + \frac{16C_\Theta H^2\kappa_1}{\epsilon}) + d\log(1 + \frac{32C_\Theta H^3\kappa_1}{\epsilon})
$$

Proof of Lemma L.11. Recall $\sigma_{u,v}^2(\cdot, \cdot) := \max\{1, f(v, \phi(\cdot, \cdot))_{[0, (H-h+1)^2]} - \left[f(u, \phi(\cdot, \cdot))_{[0, H-h+1]}\right]^2\}$, and since $\max$, truncation are non-expansive operations, then we can achieve for any s, a

$$
\begin{aligned}
|\sigma_{u_1, v_1}^2(s,a) - \sigma_{u_2, v_2}^2(s,a)| &\le |f(v_1, \phi(s,a)) - f(v_2, \phi(s,a))| + 2H |f(u_1, \phi(s,a)) - f(u_2, \phi(s,a))| \\
&\le \kappa_1 \|v_1 - v_2\|_2 + 2H\kappa_1 \|u_1 - u_2\|_2,
\end{aligned}
$$

Hence

$$
\left| \bar{X}(\theta_1, \theta_1', u_1, v_1) - \bar{X}(\theta_2, \theta_2', u_2, v_2) \right| = \left| \frac{X(\theta_1, \theta_1')}{\sigma_{u_1, v_1}^2} - \frac{X(\theta_2, \theta_2')}{\sigma_{u_2, v_2}^2} \right|
$$

$$
\leq \left| \frac{X(\theta_1, \theta_1') - X(\theta_2, \theta_2')}{\sigma_{u_1, v_1}^2} \right| + \left| \frac{X(\theta_2, \theta_2')}{\sigma_{u_1, v_1}^2 \sigma_{u_2, v_2}^2} \left(\sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2 \right) \right|
$$

$$
\leq \left| X(\theta_1, \theta_1') - X(\theta_2, \theta_2') \right| + 2H^2 \left| \sigma_{u_1, v_1}^2 - \sigma_{u_2, v_2}^2 \right|
$$

$$
\leq \left| X(\theta_1, \theta_1') - X(\theta_2, \theta_2') \right| + 2H^2 \kappa_1 \left\| v_1 - v_2 \right\|_2 + 4H^3 \kappa_1 \left\| u_1 - u_2 \right\|_2
$$

$$
\leq (6H + 1)\kappa_1 \cdot \left\| \theta_1 - \theta_2 \right\|_2 + 6H \left\| V_{\theta_1'} - V_{\theta_2'} \right\|_\infty + 2H^2 \kappa_1 \left\| v_1 - v_2 \right\|_2 + 4H^3 \kappa_1 \left\| u_1 - u_2 \right\|_2
$$

Note the above holds true for all s, a, r, s', therefore it implies

$$
\left| \mathbb{E}_\mu[\bar{X}(\theta_1, \theta_1', u_1, v_1)] - \mathbb{E}_\mu[\bar{X}(\theta_2, \theta_2', u_2, v_2)] \right|
$$

$$
\leq (6H + 1)\kappa_1 \cdot \left\| \theta_1 - \theta_2 \right\|_2 + 6H \left\| V_{\theta_1'} - V_{\theta_2'} \right\|_\infty + 2H^2 \kappa_1 \left\| v_1 - v_2 \right\|_2 + 4H^3 \kappa_1 \left\| u_1 - u_2 \right\|_2
$$

and similar to Lemma L.10, the covering number of ϵ-net will be bounded by

$$
d\log(1 + \frac{24C_\Theta(H+1)\kappa_1}{\epsilon}) + d\log\left(1 + \frac{288H^2 C_\Theta(\kappa_1\sqrt{C_\Theta} + 2\sqrt{B}\kappa_1\kappa_2)^2}{\epsilon^2} \right) + d^2\log\left(1 + \frac{288H^2\sqrt{d}B\kappa_1^2}{\epsilon^2} \right)
$$

$$
+ d\log(1 + \frac{16C_\Theta H^2 \kappa_1}{\epsilon}) + d\log(1 + \frac{32C_\Theta H^3 \kappa_1}{\epsilon})
$$

Comparing to Lemma L.10, the last two terms are incurred by covering u, v arguments.

$\square$