SGDA WITH SHUFFLING: FASTER CONVERGENCE FOR NONCONVEX-PŁ MINIMAX OPTIMIZATION

ABSTRACT

Stochastic gradient descent-ascent (SGDA) is one of the main workhorses for solving finite-sum minimax optimization problems. Most practical implementations of SGDA randomly reshuffle components and sequentially use them (*i.e.*, without-replacement sampling); however, there are few theoretical results on this approach for minimax algorithms, especially outside the easier-to-analyze (strongly-)monotone setups. To narrow this gap, we study the convergence bounds of SGDA with random reshuffling (**SGDA-RR**) for smooth nonconvex-nonconcave objectives with Polyak-Łojasiewicz (PŁ) geometry. We analyze both simultaneous and alternating SGDA-RR for *nonconvex-PŁ* and *primal-PŁ-PŁ* objectives, and obtain convergence rates faster than with-replacement SGDA. Our rates also extend to mini-batch SGDA-RR, recovering known rates for full-batch gradient descent-ascent (GDA). Lastly, we present a comprehensive lower bound for two-time-scale GDA, which matches the full-batch rate for primal-PŁ-PŁ case.

1 INTRODUCTION

A finite-sum minimax optimization problem aims to solve the following:

$$\min_{\boldsymbol{x} \in \mathbb{R}^{d_x}} \max_{\boldsymbol{y} \in \mathbb{R}^{d_y}} f(\boldsymbol{x}; \boldsymbol{y}) := \frac{1}{n} \sum_{i=1}^{n} f_i(\boldsymbol{x}; \boldsymbol{y}), \tag{1}$$

where f_i denotes the i-th component function. In plain language, we would like to minimize the average of n component functions for $\boldsymbol{x}$, while maximizing it for $\boldsymbol{y}$ given $\boldsymbol{x}$. There are many important areas in modern machine learning that fall within this framework, including generative adversarial networks (GANs) (Goodfellow et al., 2014; Heusel et al., 2017), adversarial training (Sinha et al., 2018; Madry et al., 2018), multi-agent reinforcement learning (MARL) (Li et al., 2019), AUC maximization (Ying et al., 2016; Yuan et al., 2021), and many more. In most cases, the objective f is usually nonconvex-nonconcave, *i.e.*, neither convex in $\boldsymbol{x}$ nor concave in $\boldsymbol{y}$. Since general nonconvex-nonconcave problems are known to be intractable, we would like to tackle problems with some additional structures, such as smoothness and Polyak-Łojasiewicz (PŁ) condition(s). We present the detailed settings, *nonconvex-PŁ* and *primal-PŁ-PŁ* (or, *PŁ(Φ)-PŁ*), for our analysis in Section 2.

One of the simplest and most popular algorithms to solve the problem (1) would be *stochastic gradient descent-ascent* (**SGDA**). This naturally extends the idea of stochastic gradient descent (SGD) used for minimization problems. Given an initial iterate $(\boldsymbol{x}_0; \boldsymbol{y}_0)$, at time $t \in \mathbb{N}$, SGDA (randomly) chooses an index $i(t) \in \{1, \dots, n\}$ and accesses the $i(t)$-th component to perform a pair of updates

$$\begin{bmatrix} \boldsymbol{x}_t = \boldsymbol{x}_{t-1} - \alpha \nabla_1 f_{i(t)}(\boldsymbol{x}_{t-1}; \boldsymbol{y}_{t-1}), \\ \boldsymbol{y}_t = \boldsymbol{y}_{t-1} + \beta \nabla_2 f_{i(t)}(\boldsymbol{x}'; \boldsymbol{y}_{t-1}), \end{bmatrix} \quad \text{where } \boldsymbol{x}' = \begin{cases} \boldsymbol{x}_{t-1}, & (\textit{simSGDA}), \text{ or} \\ \boldsymbol{x}_t, & (\textit{altSGDA}). \end{cases}$$

Here, $\alpha > 0$ and $\beta > 0$ are the step sizes and ∇_j denotes the gradient with respect to j-th argument for $f_{i(t)}$ ($j = 1, 2$). As shown in the update equations above, there are two widely used versions of SGDA: *simultaneous SGDA* (**simSGDA**), and *alternating SGDA* (**altSGDA**).

In such stochastic gradient methods, there are two main categories of sampling schemes for the component indices $i(t)$. One way is to sample $i(t)$ independently (in time) and uniformly at random from $\{1, \dots, n\}$, which is called *with-replacement sampling*. This scheme is widely adopted in theory papers because it makes analysis of stochastic methods amenable: the noisy gradients $\nabla f_{i(t)}$

are independent over time t and are unbiased estimators of the full-batch gradient ∇f. In contrast, the vast majority of practical implementations employ *without-replacement sampling*, indicating a huge theory-practice gap. In without-replacement sampling, we sample each index precisely once at each epoch. Perhaps the most popular of such schemes is *random reshuffling* (**RR**), which uniformly randomly shuffles the order of indices at the beginning of every epoch. Unfortunately, it is well-known that without-replacement methods are much more difficult to analyze theoretically, largely because the sampled indices in each epoch are no longer independent of each other.

Interestingly, for minimization problems, several recent works overcome this obstacle and show that SGD using without-replacement sampling leads to faster convergence, given that the number of epochs is large enough (Nagaraj et al., 2019; Rajput et al., 2020; Ahn et al., 2020; Mishchenko et al., 2020; Nguyen et al., 2021; Yun et al., 2021; 2022). On the other hand, for minimax problems like (1), the majority of the studies still assume with-replacement sampling and/or rely on the independent unbiased gradient assumption (Nouiehed et al., 2019; Lin et al., 2020; Guo et al., 2020; Yan et al., 2020; Yang et al., 2020; Loizou et al., 2021; Beznosikov et al., 2022). There are very few known results on minimax optimization algorithms using without-replacement sampling, and even most of the existing ones take advantage of (strong-)convexity (in $\boldsymbol{x}$) and/or (strong-)concavity (in $\boldsymbol{y}$) of the problem (Das et al., 2022; Yu et al., 2022; Maheshwari et al., 2022). A detailed comparative analysis of these works is conducted in Section 4.

Putting all these issues into consideration, our main question is the following.

> *Does SGDA using without-replacement component sampling provably converge fast,*
> *even on smooth nonconvex-nonconcave objective f with PŁ structures?*

1.1 SUMMARY OF OUR CONTRIBUTIONS

To answer the question above, we analyze the convergence of SGDA with random reshuffling (**SGDA-RR**, Algorithm 1). We analyze both the simultaneous and alternating versions of SGDA-RR and prove convergence theorems for the following two regimes.

- If $-f(\boldsymbol{x}; \boldsymbol{y})$ satisfies μ_2-PŁ condition in $\boldsymbol{y}$ (**nonconvex-PŁ**) and component function f_i's are L-smooth, then we prove that SGDA-RR with step size ratio $r = \beta/\alpha \gtrsim L^2/\mu_2^2$ converges to ε-stationarity in expectation in $\mathcal{O}\left(rL\varepsilon^{-2} + r^{3/2}L\varepsilon^{-3}\right)$ epochs (Theorem 1).

- If we further assume μ_1-PŁ condition on the primal function $\Phi(\cdot) := \max_{\boldsymbol{y}} f(\cdot; \boldsymbol{y})$ (**primal-PŁ-PŁ**, or **PŁ(Φ)-PŁ**), we prove that SGDA-RR with step size ratio $r \gtrsim L^2/\mu_2^2$ converges within ε-tolerance in expectation after $\tilde{\mathcal{O}}\left(\frac{Lr}{\mu_1}\log(\varepsilon^{-1}) + L(\frac{r}{\mu_1})^{3/2}\varepsilon^{-1}\right)$ epochs (Theorem 2).

As will be discussed in Section 4, the rates shown above are *faster* than existing results on with-replacement SGDA. In fact, Theorems 1 & 2 are special cases ($b = 1$) of our extended theorems (Theorems 4 & 5, Appendix A) that analyze *mini-batch* SGDA-RR of batch size b; by setting $b = n$, we also recover known convergence rates for full-batch gradient descent ascent (GDA). Hence, our analysis covers the entire spectrum between vanilla SGDA-RR ($b = 1$) and GDA ($b = n$), for both simultaneous and alternating versions.

- Additionally, we provide complexity lower bounds for minimizing strongly-convex-strongly-concave (SC-SC) functions using two-time-scale simultaneous GDA with arbitrary fixed step size ratio $r = \beta/\alpha$. Perhaps surprisingly, we find that the lower bound for SC-SC functions matches the convergence upper bound for a much larger class of primal-PŁ-PŁ functions when the step size ratio satisfies $r \gtrsim L^2/\mu_2^2$ (Theorem 3).

2 PROBLEM SETUP

2.1 NOTATION

In our problem (1), the domain of every f_i is $\mathcal{Z} = \mathcal{X} \times \mathcal{Y}$, where $\mathcal{X} = \mathbb{R}^{d_x}$, $\mathcal{Y} = \mathbb{R}^{d_y}$, and $\mathcal{Z} = \mathbb{R}^d$: we concern unconstrained problems for simplicity. We denote the Euclidean norm and

---

Algorithm 1 simSGDA/altSGDA-RR

1: **Given:** The number of components n; the number of epochs K; step sizes $\alpha, \beta > 0$
2: **Initialize:** $(\boldsymbol{x}_0^1; \boldsymbol{y}_0^1) \in \mathbb{R}^{d_x} \times \mathbb{R}^{d_y}$
3: **for** $k \in [K]$ **do**
4:      Sample $\sigma_k \sim \mathrm{Unif}(\mathbb{S}_n)$ ▷ RR: uniformly randomly shuffle the indices every epoch
5:      **for** $i \in [n]$ **do**
6:          $\boldsymbol{x}_i^k = \boldsymbol{x}_{i-1}^k - \alpha \nabla_1 f_{\sigma_k(i)}(\boldsymbol{x}_{i-1}^k; \boldsymbol{y}_{i-1}^k)$
7:          **if** simSGDA-RR **then**
8:              $\boldsymbol{y}_i^k = \boldsymbol{y}_{i-1}^k + \beta \nabla_2 f_{\sigma_k(i)}(\boldsymbol{x}_{i-1}^k; \boldsymbol{y}_{i-1}^k)$ ▷ simultaneous update: $\boldsymbol{x}$ & $\boldsymbol{y}$
9:          **else if** altSGDA-RR **then**
10:              $\boldsymbol{y}_i^k = \boldsymbol{y}_{i-1}^k + \beta \nabla_2 f_{\sigma_k(i)}(\boldsymbol{x}_i^k; \boldsymbol{y}_{i-1}^k)$ ▷ alternating update: $\boldsymbol{x} \to \boldsymbol{y}$
11:      $(\boldsymbol{x}_0^{k+1}; \boldsymbol{y}_0^{k+1}) = (\boldsymbol{x}_n^k; \boldsymbol{y}_n^k)$

---

the standard inner product by $\|\cdot\|$ and $\langle \cdot, \cdot \rangle$, respectively. We often use an abbreviated notation $\boldsymbol{z} = (\boldsymbol{x}; \boldsymbol{y}) \in \mathcal{Z}$ for $\boldsymbol{x} \in \mathcal{X}$ and $\boldsymbol{y} \in \mathcal{Y}$. Even when $\boldsymbol{z}$ or $(\boldsymbol{x}; \boldsymbol{y})$ is followed by superscripts and/or subscripts, we use the symbols interchangeably; *e.g.*, $\boldsymbol{z}_i^k = (\boldsymbol{x}_i^k; \boldsymbol{y}_i^k)$. Note that we split the arguments for minimization ($\boldsymbol{x}$, called *primal variable*) and for maximization ($\boldsymbol{y}$, called *dual variable*) by a semicolon (';'). We use ∇_1 and ∇_2 to denote the gradients with respect to first and second argument, respectively.[1] Given a closed set $\mathcal{S} \in \mathbb{R}^d$, we denote the set of all projection(s) of $\boldsymbol{v} \in \mathbb{R}^d$ onto $\mathcal{S}$, *i.e.*, the nearest point(s) in $\mathcal{S}$ from $\boldsymbol{v}$, by $\Pi_{\mathcal{S}}(\boldsymbol{v}) := \arg\min_{\boldsymbol{w} \in \mathcal{S}} \|\boldsymbol{v} - \boldsymbol{w}\|$. For a positive integer N, we denote $[N] := \{1, 2, 3, \ldots, N\}$. Let the set $\mathbb{S}_N$ be a symmetric group of degree N. That is, each *permutation* $\sigma \in \mathbb{S}_N$ is a bijection from $[N]$ to itself, or equivalently, a re-arrangement of $[N]$. Lastly, we use the usual $\mathcal{O}/\Omega/\Theta$ notation for bounds, where $\tilde{\mathcal{O}}, \tilde{\Omega}$, and $\tilde{\Theta}$ are for hiding some logarithmic factors.

2.2 Algorithms: (two-time-scale) simSGDA-RR & altSGDA-RR

As we explained in Section 1, we consider simSGDA and altSGDA combined with RR, a without-replacement sampling scheme. We call them **simSGDA-RR** and **altSGDA-RR**, respectively. We present a detailed description of the methods in Algorithm 1. For completeness, we also provide an extended version that uses mini-batches of size ≥ 1 (Algorithm 2) in Appendix A. For comparison, we call the SGDA algorithms using *with-replacement* sampling by just **simSGDA** and **altSGDA**.

The quantities $\alpha, \beta > 0$ are step sizes associated with primal variable $\boldsymbol{x}$ and dual variable $\boldsymbol{y}$, respectively. We use two separate symbols α and β to allow the two step sizes to be different. Such algorithms are commonly called *two-time-scale* algorithms, and they are adopted in nonconvex minimax optimization problems (Heusel et al., 2017; Lin et al., 2020; Yang et al., 2020). In fact, a recent result (Li et al., 2022) shows that having $\alpha \neq \beta$ is sometimes *necessary* for convergence.

2.3 Assumptions and definitions

To define the function classes that we are interested in solving, we introduce a few assumptions.

Assumption 1 (Component smoothness)**.** *Every i-th component $f_i : \mathcal{Z} \to \mathbb{R}$ is L-**smooth**, i.e., f_i is differentiable and ∇f_i is L-Lipschitz continuous: $\|\nabla f_i(\boldsymbol{z}) - \nabla f_i(\boldsymbol{z}')\| \leq L \|\boldsymbol{z} - \boldsymbol{z}'\|$. As a result, for any $\boldsymbol{z}, \boldsymbol{z}' \in \mathcal{Z}$, $f_i(\boldsymbol{z}') - f_i(\boldsymbol{z}) \leq \langle \nabla f_i(\boldsymbol{z}), \boldsymbol{z}' - \boldsymbol{z} \rangle + \frac{L}{2} \|\boldsymbol{z}' - \boldsymbol{z}\|^2$.*

Since f_i's are all L-smooth, the average f of f_i's is also L-smooth.

The next one controls the discrepancy between the objective function f and its components f_i's. This is similar to Assumption 2 of Nguyen et al. (2021), adapted to minimax problems.

Assumption 2 (Component gradient variance)**.** *There exist constants $A, B \geq 0$ such that, for any $\boldsymbol{z} = (\boldsymbol{x}; \boldsymbol{y}) \in \mathcal{Z}$ and $j \in \{1, 2\}$, we have $\frac{1}{n} \sum_{i=1}^n \|\nabla_j f_i(\boldsymbol{z}) - \nabla_j f(\boldsymbol{z})\|^2 \leq A \|\nabla_j f(\boldsymbol{z})\|^2 + B$.*

In Assumption 2, letting $A = 0$ retrieves a common assumption that the variance of component gradients is uniformly bounded; thus, our assumption is a relaxation. Also notice that $A = B = 0$ if $n = 1$.

---

[1]We adopt this notation in order to avoid the possibility that an expression $\nabla_{\boldsymbol{x}} g(\boldsymbol{x}; \boldsymbol{y}(\boldsymbol{x}))$ is misunderstood as the *total* derivative with respect to $\boldsymbol{x}$ when $\boldsymbol{y}$ is dependent to $\boldsymbol{x}$; *e.g.*, see Section 3.1.

Assumption 3. *For a function $f : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$, the* **primal function** $\Phi : \mathcal{X} \to \mathbb{R}$ *is well-defined as $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}' \in \mathcal{Y}} f(\boldsymbol{x}; \boldsymbol{y}')$. For each $\boldsymbol{x} \in \mathcal{X}$, the set $\mathcal{Y}_{\boldsymbol{x}}^* := \arg\max_{\boldsymbol{y}' \in \mathcal{Y}} f(\boldsymbol{x}; \boldsymbol{y}')$ is non-empty and closed. Moreover, we assume $\Phi(\boldsymbol{x})$ is bounded below by $\Phi^* = \inf_{\boldsymbol{x}' \in \mathcal{X}} \Phi(\boldsymbol{x}') > -\infty$.*

Because the sets $\mathcal{Y}_{\boldsymbol{x}}^*$ is assumed to be non-empty and closed, the set of projection $\Pi_{\mathcal{Y}_{\boldsymbol{x}}^*}(\boldsymbol{y})$ is also well-defined for each $\boldsymbol{x} \in \mathcal{X}$ and $\boldsymbol{y} \in \mathcal{Y}$.

We now add an additional structure to our objective function, which is called Polyak-Łojasiewicz (PŁ) condition. A function $g : \mathbb{R}^d \to \mathbb{R}$ is said to be μ-**PŁ** if it has a minimum value g^* and satisfies

$$\|\nabla g(\boldsymbol{t})\|^2 \geq 2\mu(g(\boldsymbol{t}) - g^*). \quad (\forall\, \boldsymbol{t} \in \mathbb{R}^d)$$

This condition is studied in the context of machine learning in many recent papers, *e.g.*, Karimi et al. (2016); Nouiehed et al. (2019); Yang et al. (2020); Guo et al. (2020), and more. Note that every μ-strongly convex[2] function satisfies μ-PŁ condition, whereas a PŁ function does not need to be convex. Hence, μ-PŁ is a strict generalization of μ-strong convexity. In addition, every stationary point of a PŁ function is a global optimum, which is a benign property for optimization.

We are interested in the case where our objective function $f(\boldsymbol{x}; \boldsymbol{y})$ has such a structure in terms of $\boldsymbol{y}$ (Assumption 4). Sometimes, we further assume the primal function Φ is also PŁ (Assumption 5). We emphasize that we do not necessarily assume the PŁ conditions for the individual f_i's.

Assumption 4 (*$\boldsymbol{y}$-side PŁ*). *For each (fixed) $\boldsymbol{x} \in \mathcal{X}$, $-f(\boldsymbol{x}; \cdot)$ is μ_2-PŁ, i.e., for every $(\boldsymbol{x}; \boldsymbol{y}) \in \mathcal{Z}$, $\|\nabla_2 f(\boldsymbol{x}; \boldsymbol{y})\|^2 \geq 2\mu_2(\Phi(\boldsymbol{x}) - f(\boldsymbol{x}; \boldsymbol{y}))$, where Φ is the primal function associated with f.*

Assumption 5 (*Primal PŁ, or PŁ(Φ)*). *The primal function $\Phi(\cdot) = \max_{\boldsymbol{y}'} f(\boldsymbol{x}; \boldsymbol{y}')$ of f is μ_1-PŁ, i.e., for every $\boldsymbol{x} \in \mathcal{X}$, $\|\nabla \Phi(\boldsymbol{x})\|^2 \geq 2\mu_1(\Phi(\boldsymbol{x}) - \Phi^*)$, where $\Phi^* = \min_{\boldsymbol{x}} \Phi(\boldsymbol{x})$ is well-defined.*

We say the function f is **nonconvex-PŁ** when it satisfies Assumption 4. Since we do not assume any convexity/concavity, it is generally hard to reach global optima. Due to the $\boldsymbol{y}$-side PŁ condition, we can guarantee that the primal function Φ is differentiable and even L_Φ-smooth with $L_\Phi \leq L + L^2/\mu_2$ (see Proposition 9 in Appendix B). Since the problem (1) can be reformulated as the minimization problem of Φ (when we can always find $\boldsymbol{y}$ well that maximizes $f(\boldsymbol{x}; \boldsymbol{y})$ given $\boldsymbol{x}$), we could aim to find an approximate first-order stationary point of Φ, by making the norm of the gradient of Φ small.

On top of that, if f satisfies both Assumptions 4 and 5, the function is said to be **primal-PŁ-PŁ**, or **PŁ(Φ)-PŁ** for short.[3] In this case, we directly aim not only to decrease the primal function Φ associated with the objective function f but also to increase the function value $f(\boldsymbol{x}; \boldsymbol{y})$ in terms of $\boldsymbol{y}$. To evaluate how close we are to our goal, we define a *potential function* V_λ later in Section 3. When we attain $V_\lambda(\boldsymbol{x}^*, \boldsymbol{y}^*) = 0$, it implies that we arrive at a global minimax point: $f(\boldsymbol{x}^*, \boldsymbol{y}^*) = \Phi(\boldsymbol{x}^*) = \Phi^*$. The function V_λ enables us to develop a unified analysis for nonconvex-PŁ and PŁ(Φ)-PŁ objective functions; we discuss this in greater detail in Section 3.

3 Main results

Based on the assumptions stated in the previous section, we present the convergence results for both smooth nonconvex-PŁ objectives and smooth PŁ(Φ)-PŁ objectives. Before stating the main theorems, we first introduce the most important tool for our analyses: the *potential function*.

3.1 Potential function V_λ

For our convergence analyses, we utilize a function $V_\lambda : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$ defined as

$$V_\lambda(\boldsymbol{x}; \boldsymbol{y}) := \lambda[\Phi(\boldsymbol{x}) - \Phi^*] + [\Phi(\boldsymbol{x}) - f(\boldsymbol{x}; \boldsymbol{y})], \tag{2}$$

where $\lambda > 1$ is a constant. We borrow inspiration from Yang et al. (2020) and Das et al. (2022) to come up with this function, although the placement of λ of ours is different from what they

---

[2]We say a function $g : \mathbb{R}^d \to \mathbb{R}$ is μ-strongly convex for some $\mu > 0$ if it holds $g(\boldsymbol{x}') \geq g(\boldsymbol{x}) + \langle \nabla g(\boldsymbol{x}), \boldsymbol{x}' - \boldsymbol{x} \rangle + (\mu/2) \|\boldsymbol{x}' - \boldsymbol{x}\|^2$ $(\forall \boldsymbol{x}, \boldsymbol{x}')$; we say g is μ-strongly concave if $-g$ is μ-strongly convex.

[3]The PŁ(Φ)-PŁ condition is much weaker than **two-sided PŁ** condition assuming "$\boldsymbol{x}$-side" PŁ condition: see Proposition 10. As pointed out by Guo et al. (2020), there exist a PŁ(Φ)-PŁ function $g(\boldsymbol{x}; \boldsymbol{y})$ that is not $\boldsymbol{x}$-side μ-PŁ for any $\mu > 0$ but even strongly *concave* in $\boldsymbol{x}$.

use. In fact, the convergence to a neighborhood of a global minimax point (if it exists) implies the reduction of this function. For each $\boldsymbol{x}$, a non-negative term $\Phi(\boldsymbol{x}) - f(\boldsymbol{x}; \boldsymbol{y})$ gets smaller as $\boldsymbol{y}$ makes $f(\boldsymbol{x}; \boldsymbol{y})$ larger. The term becomes zero when $\boldsymbol{y} = \boldsymbol{y}^*(\boldsymbol{x})$ for some $\boldsymbol{y}^*(\boldsymbol{x}) \in \mathcal{Y}_{\boldsymbol{x}}^*$, since $\Phi(\boldsymbol{x}) = f(\boldsymbol{x}; \boldsymbol{y}^*(\boldsymbol{x}))$. Also, another non-negative term $\Phi(\boldsymbol{x}) - \Phi^*$ gets smaller as $\boldsymbol{x}$ makes $\Phi(\boldsymbol{x})$ smaller. Thus, as $(\boldsymbol{x}; \boldsymbol{y})$ approaches to a minimax optimal point, $V_\lambda(\boldsymbol{x}; \boldsymbol{y})$ decreases to near zero. In general, V_λ is not guaranteed to attain exact zero, especially when the objective function $f(\boldsymbol{x}; \boldsymbol{y})$ is nonconvex in $\boldsymbol{x}$ (*e.g.*, f is nonconvex-PŁ). Nevertheless, the potential function is still useful for deriving our convergence results.

3.2 MAIN THEOREMS: UPPER BOUNDS OF CONVERGENCE RATES

Now, we present our main results. We provide a detailed comparison of our theorems against existing results in Section 4, and we give a sketch of proofs in Section 6. The detailed proofs are presented in Appendices C and D. We remark that both Theorems 1 and 2 are special cases (for mini-batch size $b = 1$) of their *mini-batch* extensions: Theorems 4 and 5 in Appendix A.

Theorem 1 (Nonconvex-PŁ). *Suppose that f satisfies Assumptions 1, 2, 3, and 4. Let $\kappa_2 = L/\mu_2$, where μ_2 is PŁ constant of $-f(\boldsymbol{x}; \cdot)$ at all $\boldsymbol{x}$. Let $\lambda = 4$. Choose the step sizes α and β such that*

$$\beta = \min\left\{ \frac{1}{6L\sqrt{n(n+A)}}, \ \mathcal{O}\left(\left(\frac{V_\lambda(\boldsymbol{z}_0^1)}{Bn^2 K}\right)^{\frac{1}{3}}\right) \right\} \quad and \quad \alpha = \frac{\beta}{r},$$

for some $r \geq 14\kappa_2^2$. Then, both simSGDA-RR and altSGDA-RR (Algorithm 1) satisfy

$$\min_{k \in [K]} \mathbb{E}\left[\left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \right] \leq \mathcal{O}\left(\frac{rLV_\lambda(\boldsymbol{z}_0^1)}{K}\sqrt{1 + \frac{A}{n}} + r\left(\frac{L^2 B V_\lambda(\boldsymbol{z}_0^1)^2}{nK^2}\right)^{\frac{1}{3}} \right).$$

Upper bound on epoch complexity. To achieve ε-stationarity in terms of the primal function, *i.e.*, $\min_{k \in [K]} \mathbb{E}\left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \leq \varepsilon^2$, a sufficient number of epochs (denoted by K_ε) is

$$K_\varepsilon = \mathcal{O}\left(\frac{rLV_\lambda(\boldsymbol{z}_0^1)}{\varepsilon^2}\sqrt{1 + \frac{A}{n}} + \frac{r^{3/2}L\sqrt{B}V_\lambda(\boldsymbol{z}_0^1)}{\sqrt{n}\varepsilon^3} \right).$$

Theorem 2 (PŁ(Φ)-PŁ). *Suppose that f satisfies Assumptions 1, 2, 3, 4, and 5. Let $\kappa_1 = L/\mu_1$ and $\kappa_2 = L/\mu_2$, where μ_1 and μ_2 are PŁ constants of $\Phi(\cdot)$ and $-f(\boldsymbol{x}; \cdot)$ (at all $\boldsymbol{x}$), respectively. Let $\lambda = 4$. Choose appropriate step sizes α and β such that*

$$\beta = \min\left\{ \frac{1}{6L\sqrt{n(n+A)}}, \ \tilde{\mathcal{O}}\left(\frac{\kappa_2^2}{\mu_1 nK}\right) \right\} \quad and \quad \alpha = \frac{\beta}{r},$$

for some $r \geq 14\kappa_2^2$. Then, both simSGDA-RR and altSGDA-RR (Algorithm 1) satisfy

$$\mathbb{E}[V_\lambda(\boldsymbol{z}_0^{K+1})] \leq \mathcal{O}\left(V_\lambda(\boldsymbol{z}_0^1) \cdot \exp\left(-\frac{K}{12\kappa_1 r\sqrt{1 + \frac{A}{n}}}\right) \right) + \tilde{\mathcal{O}}\left(\frac{\kappa_1^2 r^3 B}{\mu_1 nK^2}\right).$$

Upper bound on epoch complexity. To achieve ε^2-accuracy on expectation of $V_\lambda(\boldsymbol{z}_n^K)$, *i.e.*, $\mathbb{E}[V_\lambda(\boldsymbol{z}_n^K)] \leq \varepsilon^2$, a sufficient number of epochs (denoted by K_ε') is

$$K_\varepsilon' = \mathcal{O}\left(\kappa_1 r\sqrt{1 + \frac{A}{n}} \log\frac{V_\lambda(\boldsymbol{z}_0^1)}{\varepsilon} \right) + \tilde{\mathcal{O}}\left(\frac{\kappa_1 r^{3/2}}{\varepsilon}\sqrt{\frac{B}{\mu_1 n}} \right).$$

Remark on step size ratio. In both theorems, we use the step sizes of ratio $r = \beta/\alpha \gtrsim \kappa_2^2$. It is common to use such a step size scheme $r = \Theta(\kappa_2^2)$ to analyze two-time-scale (S)GDA for nonconvex minimax problems (Jin et al., 2020; Lin et al., 2020; Yang et al., 2020).

Remark on the parameter λ. In our convergence analyses, we arbitrarily choose $\lambda = 4$ which makes the numerical calculations easier. The value of $\lambda > 0$ does not matter for the equivalence between the equation $V_\lambda(\boldsymbol{x}^*; \boldsymbol{y}^*) = 0$ and global minimax condition (Proposition 11). Also, the choice of λ in both theorems can be arbitrary as long as $\lambda > 1$; our logic does not fall apart if other appropriate step sizes for that λ is chosen. That is to say, we can show that the sequence $V_\lambda(\boldsymbol{z}_0^k)$ almost monotonically decreases, ignoring some small variance terms.

4 COMPARISON WITH RELATED WORKS

Now, we compare and contrast our main results with other existing works on minimax algorithms.

4.1 COMPARISON WITH STOCHASTIC WITH-REPLACEMENT SETTING

First, we confirm that SGDA with random reshuffling (RR) has *faster* convergence rates (*i.e.*, less gradient complexity) than SGDA based on with-replacement sampling. To make the comparisons fair and easy, we simply let $r = \beta/\alpha = \Theta(\kappa_2^2)$, $A = 0$, and $B = \tau^2$.

Lin et al. (2020) obtain a convergence rate for with-replacement simSGDA with $r = \Theta(\kappa_2^2)$, run on nonconvex & μ_2-strongly-concave problems with a convex *bounded* constraint set $\mathcal{Y}$ for dual variable $\boldsymbol{y}$. Their *gradient* complexity to achieve $\min_{t \in [T]} \mathbb{E}\left[\|\nabla\Phi(\boldsymbol{x}_t)\|^2\right] \leq \varepsilon^2$ (where T is the number of iterations) is written as

$$\mathcal{O}\left(\frac{\kappa_2^2 L\Delta_\Phi + \kappa_2 L^2 D^2}{\varepsilon^2} \max\left\{1, \frac{\kappa_2\tau^2}{\varepsilon^2}\right\}\right), \qquad \text{(Lin et al. (2020, Theorem 4.9))}$$

where $\kappa_2 = L/\mu_2$, $\Delta_\Phi = \Phi(\boldsymbol{x}_0) - \Phi^*$, $D = \operatorname{diam}\mathcal{Y}$, and τ^2 is the variance of the (unbiased) stochastic gradient oracles. Ignoring parameters except for ε, this reduces to $\mathcal{O}(\varepsilon^{-4})$. On the contrary, our Theorem 1 has a better gradient complexity in terms of ε, with help of random reshuffling. Our gradient complexity is just the same as the epoch complexity K_ε multiplied by the number of iterations per epoch (n for Algorithm 1):

$$\mathcal{O}\left(\frac{\kappa_2^2 L V_\lambda(\boldsymbol{z}_0^1)}{\varepsilon^2}\left(1 + \frac{\kappa_2\tau\sqrt{n}}{\varepsilon}\right)\right), \qquad \text{(Ours, from Theorem 1)}$$

or simply $\mathcal{O}(\varepsilon^{-3})$, which is better than $\mathcal{O}(\varepsilon^{-4})$ of with-replacement simSGDA. Our bound has three more strengths: (i) our analysis does not require strong concavity in $\boldsymbol{y}$, which is a strictly stronger assumption than requiring $\boldsymbol{y}$-side PL condition; (ii) our analysis does not require the constraint set $\mathcal{Y}$ to be bounded; (iii) our result can easily extend to the case of *any* mini-batch sizes, whereas Lin et al. (2020) need a particular choice of mini-batch size $M = \mathcal{O}(\kappa_2\tau^2/\varepsilon)$ to ensure convergence.

For PŁ(Φ)-PŁ objective, Yang et al. (2020) obtain a convergence rate for with-replacement altSGDA with $r = \Theta(\kappa_2^2)$.[4] They apply diminishing step sizes ($\mathcal{O}(1/t)$, $t \in \mathbb{N}$) to derive a gradient complexity bound $\mathcal{O}\left(\frac{\kappa_1\kappa_2^4\tau^2}{\mu_1\varepsilon^2}\right)$ to achieve $\mathbb{E}[V_\lambda(\boldsymbol{z}_T)] \leq \varepsilon^2$ (Yang et al., 2020, Theorem 3.3). One can apply the step sizes depending on the total number of iterations to their analysis and derive a similar complexity with only deterioration in logarithmic factor. On the other hand, our gradient complexity for altSGDA-RR *and* simSGDA-RR can be written as

$$\tilde{\mathcal{O}}\left(\frac{\kappa_1\kappa_2^3\tau\sqrt{n}}{\varepsilon\sqrt{\mu_1}}\right). \qquad \text{(Ours, from Theorem 2)}$$

This is a better complexity in ε and κ_2, even without the requirement of diminishing step size.

4.2 COMPARISON WITH OTHER WORKS ON STOCHASTIC WITHOUT-REPLACEMENT SETTING

One of the most relevant works to this paper is Das et al. (2022, Theorem 3). The authors obtain a similar convergence rate to us for two-sided PŁ objective, based on linearization of gradients, but for a dissimilar algorithm which they refer to as *AGDA-RR*. The algorithm can be also thought of as *epoch-wise*-alternating SGDA-RR, whereas our algorithm (altSGDA-RR) can be called as *step-wise*-alternating SGDA-RR. In epoch k, their algorithm (*i*) performs updates only on $\boldsymbol{x}$ ($\boldsymbol{x}_0^k, \ldots, \boldsymbol{x}_n^k$) while fixing $\boldsymbol{y}$ to $\boldsymbol{y}_0^k$, and then (*ii*) performs updates only on $\boldsymbol{y}$ ($\boldsymbol{y}_0^k, \ldots, \boldsymbol{y}_n^k$) while fixing $\boldsymbol{x}$ to $\boldsymbol{x}_0^{k+1} = \boldsymbol{x}_n^k$. We believe that our step-wise algorithm is closer to practice, especially when n is large. Because of the distinction between algorithms, the proof techniques are also different.

Xie et al. (2021, Theorem 3) present a convergence rate of *CD-MA*, an extension of simSGDA to the cross-device federated learning setup, on nonconvex-PŁ setting. Their convergence result for

---

[4]Although they consider two-sided PŁ problems, their analysis applies to PŁ(Φ)-PŁ problems as well.

CD-MA also assumes mini-batch sampling by random reshuffling. As a consequence, they yield a rate analogous to our Theorem 1 if we reduce their result to the single-machine setup. Nevertheless, our convergence bound contains a term that shrink with the number of components or mini-batches, whereas theirs does not. For a more detailed comparison, please refer to Appendix F.

There are also some works on RR-based (constrained) minimax optimization algorithms other than SGDA, but for convex-concave problems. Maheshwari et al. (2022) present *OGDA-RR*, a gradient-free RR-based optimistic GDA algorithm. Yu et al. (2022) study *stochastic proximal point with RR*, consisting of double-loop epochs. Their analyses exploit convex-concavity and Lipschitz continuity of their objective, based on the arguments by Nagaraj et al. (2019). This enables a direct usage of duality gap, the difference between primal function $\Phi(\cdot)$ and dual function $\Psi(\cdot) = \min_{\boldsymbol{x}} f(\boldsymbol{x}; \cdot)$, as a criterion for optimality. On the contrary, our work rely on a different structure of the functions, which in turn differentiate the constructions of convergence rates.

4.3 COMPARISON WITH DETERMINISTIC SETTING

Here, we compare our rates with (full-batch two-time-scale) *gradient descent ascent* (GDA):

$$
\begin{bmatrix} \boldsymbol{x}_k = \boldsymbol{x}_{k-1} - \alpha \nabla_1 f(\boldsymbol{x}_{k-1}; \boldsymbol{y}_{k-1}), \\ \boldsymbol{y}_k = \boldsymbol{y}_{k-1} + \beta \nabla_2 f(\boldsymbol{x}'; \boldsymbol{y}_{k-1}), \end{bmatrix} \quad \text{where } \boldsymbol{x}' = \begin{cases} \boldsymbol{x}_{k-1}, & (simGDA), \text{ or} \\ \boldsymbol{x}_k, & (altGDA). \end{cases}
$$

It uses the whole information of the objective f at every iteration without any noise. For comparison with GDA, we utilize our extended theorems for arbitrary mini-batch size b dividing n (Theorems 4 and 5, Appendix A). By letting $b = n$ and matching our iterate $\boldsymbol{z}_0^k = (\boldsymbol{x}_0^k; \boldsymbol{y}_0^k)$ to a GDA iterate $\boldsymbol{z}_k = (\boldsymbol{x}_k; \boldsymbol{y}_k)$, our results reduce to upper convergence bounds for simGDA and altGDA.

For nonconvex-PŁ problems (Theorems 1 & 4), the convergence rate and epoch complexity become

$$
\min_{k \in [K]} \|\nabla \Phi(\boldsymbol{x}_k)\|^2 \leq \mathcal{O}\left(\frac{\kappa_2^2 L V_\lambda(\boldsymbol{z}_1)}{K} \right); \quad i.e., \quad K_\varepsilon = \mathcal{O}\left(\frac{\kappa_2^2 L V_\lambda(\boldsymbol{z}_1)}{\varepsilon^2} \right), \tag{3}
$$

when $r = \Theta(\kappa_2^2)$. This is similar to a known rate of simGDA with $r = \Theta(\kappa_2^2)$ for nonconvex-strongly-concave problems by Lin et al. (2020, Theorem 4.4) as a special case. Their iteration complexity is written as $\mathcal{O}((\kappa_2^2 L \Delta_\Phi + \kappa_2 L^2 D^2)/\varepsilon^2)$, where the symbols are already defined in Section 4.1. To see how the two bounds compare in terms of the factors other than ε, notice that we have $\Phi(\boldsymbol{x}) - f(\boldsymbol{x}; \boldsymbol{y}) \leq \frac{L}{2} \|\boldsymbol{y} - \boldsymbol{y}^*(\boldsymbol{x})\|^2$ for any $(\boldsymbol{x}; \boldsymbol{y})$, due to L-smoothness of $-f$. Here, $\boldsymbol{y}^*(\boldsymbol{x})$ is an element of $\mathcal{Y}_{\boldsymbol{x}}^* = \arg\max_{\boldsymbol{y}} f(\boldsymbol{x}; \boldsymbol{y})$. Thus, we have $V_\lambda(\boldsymbol{z}_1) = \lambda[\Phi(\boldsymbol{x}_1) - \Phi^*] + [\Phi(\boldsymbol{x}_1) - f(\boldsymbol{z}_1)] \leq \lambda \Delta_\Phi + L D^2/2$. As a result, we could *loosely* translate our iteration complexity (3) to $\mathcal{O}((\kappa_2^2 L \Delta_\Phi + \kappa_2^2 L^2 D^2)/\varepsilon^2)$. We suspect that the discrepancy in terms of κ_2 comes from the fact that our analysis does not require the (strong) concavity in terms of $\boldsymbol{y}$ or a bounded constraint $\mathcal{Y}$: these requirements made a considerable difference in proofs.

For PŁ(Φ)-PŁ problems (Theorems 2 & 5), the convergence rate and epoch complexity reduce to

$$
V_\lambda(\boldsymbol{z}_{K+1}) \leq V_\lambda(\boldsymbol{z}_1) \cdot \exp\left(-\frac{K}{C \kappa_1 \kappa_2^2} \right); \quad i.e., \quad K_\varepsilon' = \mathcal{O}\left(\kappa_1 \kappa_2^2 \log(1/\varepsilon) \right) \tag{4}
$$

where $r = \Theta(\kappa_2^2)$ and C is a numerical constant. This recovers the linear convergence by Yang et al. (2020, Theorem 3.2) as a special case, where they prove convergence of altGDA with step size ratio $r = \Theta(\kappa_2^2)$ for two-sided PŁ problem. Following the proof of (Yang et al., 2020, Theorem 3.2), one can show that the bound (4) indeed implies the actual convergence to a global minimax point $\boldsymbol{z}^*$, in the sense that we can achieve $\|\boldsymbol{z}_k - \boldsymbol{z}^*\| \leq \varepsilon$ in $\mathcal{O}\left(\kappa_1 \kappa_2^2 \log(1/\varepsilon) \right)$ iterations.

5 LOWER BOUND FOR TWO-TIME-SCALE SIMGDA

As an extension of the discussion from Section 4.3, we characterize a lower complexity bound of two-time-scale simGDA with arbitrary step size ratio $r = \beta/\alpha$ for smooth strongly-convex-strongly-concave (SC-SC) cases. At least for $r \gtrsim \kappa_2^2$, our lower bound matches the upper complexity bound of GDA for a much wider class of problems: smooth PŁ(Φ)-PŁ problems,[5] which is quite surprising.

---

[5] strongly-convex-strongly-concave $\subset$ two-sided PŁ $\subset$ PŁ(Φ)-PŁ $\subset$ nonconvex-PŁ.

For a smooth PŁ(Φ)-PŁ problem, simGDA with step size ratio at least $r = \Omega(\kappa_2^2)$ has an upper complexity bound $K = \mathcal{O}(\kappa_1 r \log(1/\varepsilon))$ for a *global* ε-convergence $V_\lambda(\boldsymbol{z}_K) \leq \varepsilon^2$ in terms of potential function. This means that the lowest complexity is $\mathcal{O}(\kappa_1 \kappa_2^2 \log(1/\epsilon))$, achieved when $r = \Theta(\kappa_2^2)$. On the other hand, for a L-smooth μ-SC-SC problem with saddle point $\boldsymbol{z}^*$, it is well-known that the one-time-scale ($\alpha = \beta$) simGDA has a tight upper/lower complexity $K = \Theta(\kappa^2 \log(1/\varepsilon))$ to achieve $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2 \leq \varepsilon^2$, where $\kappa = L/\mu$ (*e.g.*, Das et al. (2022, Theorem C.1)). The difference of complexity bounds in condition number ($\kappa_1 \kappa_2^2$ v.s. κ^2) is somewhat questionable because, at least in smooth minimization problems, strongly convex problems and PŁ problems have identical gradient descent (GD) iteration complexity $\mathcal{O}(\kappa \log(1/\varepsilon))$ (Karimi et al., 2016, Theorem 1).

One could ask where the discrepancy in terms of κ comes from: is it due to the criteria ($V_\lambda(\boldsymbol{z}_K)$ v.s. $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2$) for ε-accuracy, (*ii*) the function classes (PŁ(Φ)-PŁ v.s. SC-SC), or (*iii*) the step size ratios ($\Omega(\kappa_2^2)$ v.s. 1)? We answer the question by showing the following theorem: the discrepancy in κ comes from the step size ratio difference. We defer the proof to Appendix E.

Theorem 3 (Lower bound, ratio-specific). *Consider a class $\mathcal{F}(L, \mu_1, \mu_2)$ of functions $f(\boldsymbol{x}; \boldsymbol{y})$ with two arguments $\boldsymbol{x}$ and $\boldsymbol{y}$, which is L-smooth, μ_1-strongly-convex in $\boldsymbol{x}$, and μ_2-strongly-concave in $\boldsymbol{y}$. Suppose $\kappa_1 = L/\mu_1 \geq c$ and $\kappa_2 = L/\mu_2 \geq c$ for a constant $c > 1$. Then, for any step size ratio $r = \beta/\alpha > 0$, there exists a function $f \in \mathcal{F}(L, \mu_1, \mu_2)$ with a unique saddle point $\boldsymbol{z}^*$, for which two-time-scale simGDA with any step sizes $(\alpha, \beta) = (\beta/r, \beta)$ requires at least*

$$
K = \begin{cases} \Omega\left(\kappa_1 r \log(1/\varepsilon)\right), & \text{if } r \geq \kappa_2/c, \\ \Omega\left(\kappa_1 \kappa_2 \log(1/\varepsilon)\right), & \text{if } c/\kappa_1 \leq r \leq \kappa_2/c, \\ \Omega((\kappa_2/r) \log(1/\varepsilon)), & \text{if } 0 < r \leq c/\kappa_1 \end{cases}
$$

iterations to achieve either $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2 \leq \varepsilon^2$ or $V_\lambda(\boldsymbol{z}_K) \leq \varepsilon^2$.

Thanks to Theorem 3, we can say from Theorem 5 that for any step size ratio $r \gtrsim \kappa_2^2$, we have a *tight* upper bound on the iteration complexity $K = \mathcal{O}(\kappa_1 r \log(1/\varepsilon))$ of two-time-scale simGDA for general PŁ(Φ)-PŁ problems. Note that Theorem 3 also subsumes the existing lower bound of the one-time-scale ($r = 1$) GDA for SC-SC problems.

Given the tightness of bounds for $r \gtrsim \kappa_2^2$, a natural next step is to discuss $1 \lesssim r \lesssim \kappa_2^2$. A recent work by Li et al. (2022) also discusses the step size ratio of simGDA. In Li et al. (2022, Theorem 4.1), the authors construct a y-side strongly-concave function[6] and show that simGDA with a step size ratio $r \leq \kappa_2$ is *impossible* to converge. The *necessity* of $r \gtrsim \kappa_2$ implied by this theorem also applies to the PŁ(Φ)-PŁ case. Thus, there is no hope for showing an upper convergence bound of simGDA with $1 \lesssim r \lesssim \kappa_2$ for general nonconvex-PŁ problems. We remark that their theorem does not contradict nor subsume Theorem 3 because we consider a much smaller function class (SC-SC) to construct the lower bounds.

On the *sufficiency* of $r \gtrsim \kappa_2$ for convergence, Li et al. (2022, Theorem 4.2) show that simGDA with $r \geq c\kappa$ (for some $c > 1$) can *locally* converge at the iteration complexity $\mathcal{O}(\kappa_1 r \log(1/\varepsilon))$ for some nonconvex-strongly-concave problems, which matches the bound in Theorem 3. Our upper bounds (Theorems 4 and 5) do require $r \gtrsim \kappa_2^2$, which may look suboptimal, but we claim that our results are not necessarily weaker. One reason is that our convergence guarantee is *global*, *i.e.*, independent of the initialization. Another reason is that their analysis is only valid when a *differential Stackelberg equilibrium*[7] exists, whereas a general PŁ(Φ)-PŁ function may not have such an equilibrium (for an example, see Proposition 13 in Appendix B).

As far as we know, it is still an interesting open problem whether a *global* convergence bound for simGDA on nonconvex-PŁ problems can be shown when the step size ratio r is between $\Omega(\kappa_2)$ and $\mathcal{O}(\kappa_2^2)$.

6 PROOF SKETCH

In this section, we briefly sketch our proofs. This section is mainly about proofs for our upper bounds (Section 3). For the proof of full-batch lower bounds (Section 5), please check Appendix E.

---

[6] $g(x; y) = -\frac{L}{2}x^2 + Lxy - \frac{\mu}{2}y^2$, where $L/\mu > 1$: its primal function is strongly convex.

[7] Loosely speaking, a differential Stackelberg equilibrium is a stationary point $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ where $f(\boldsymbol{x}^*; \cdot)$ is locally strongly concave near $\boldsymbol{y}^*$ and $\Phi(\cdot)$ is locally strongly convex near $\boldsymbol{x}^*$.

We simply consider the proofs of Theorem 1 and 2 for **sim**SGDA-RR, which is a fully stochastic case (mini-batches of size $b = 1$). The proofs for altSGDA-RR can be done with slight modifications.

We start the proof by aggregating all updates throughout an epoch to obtain an "epoch-wise" update:

$$\boldsymbol{x}_0^{k+1} = \boldsymbol{x}_0^k - n\alpha\boldsymbol{g}^k, \quad \boldsymbol{g}^k = \frac{1}{n}\sum_{i=1}^n \nabla_1 f_{\sigma_k(i)}(\boldsymbol{z}_{i-1}^k),$$
$$\boldsymbol{y}_0^{k+1} = \boldsymbol{y}_0^k + n\beta\boldsymbol{h}^k, \quad \boldsymbol{h}^k = \frac{1}{n}\sum_{i=1}^n \nabla_2 f_{\sigma_k(i)}(\boldsymbol{z}_{i-1}^k).$$

The reason is that the sampled components in each epoch are dependent to each other, so that it is much harder to deal with each iteration individually. The strategy of update-aggregation is quite general for analysis of optimization algorithms involving without-replacement sampling (Ahn et al., 2020; Mishchenko et al., 2020; Nguyen et al., 2021; Das et al., 2022). We assume that the intermediate iterates $\boldsymbol{z}_1^k, \ldots, \boldsymbol{z}_n^k$ stay close to the starting iterate $\boldsymbol{z}_0^k$ of an epoch k, which can be ensured by small step sizes. Then, we can approximate the aggregated epoch of SGDA-RR as a step of simGDA applied to $f = \frac{1}{n}\sum_{i=1}^n f_i$, with approximations of $\boldsymbol{g}^k \approx \nabla_1 f(\boldsymbol{z}_0^k)$ and $\boldsymbol{h}^k \approx \nabla_2 f(\boldsymbol{z}_0^k)$.

With Assumptions 1 and 4, note that the primal function $\Phi(\cdot)$ is $(L+L^2/\mu_2)$-smooth (Proposition 9). Applying this and L-smoothness of $-f$, we can have the following inequality (Lemma 16):

$$V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k) \le -\left((\lambda+1)/2\right) n\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + (\lambda+1)n\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$
$$+ (n\alpha/2)\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 - (n\beta/2)\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2$$
$$+ (\lambda+1/2)n\alpha \left\|\boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 + (n\beta/2)\left\|\boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k)\right\|^2.$$

Hence, to guarantee the fast decrease of $V_\lambda(\boldsymbol{z}_0^k)$, it is important to control the "noise" terms for GDA approximations, $\left\|\boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$ and $\left\|\boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k)\right\|^2$, in the last line of inequality above. By applying the tools for without-replacement sampling (Proposition 14), we can actually upper-bound the conditional expectations of both noise terms by

$$2L^2 n(n+A)\left(\alpha^2\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 + \beta^2\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2\right) + 2L^2 n(\alpha^2+\beta^2)B. \quad \text{(Lemma 17 \& 18)}$$

Then, by taking advantage of several properties of smooth nonconvex-PŁ functions (*e.g.*, Propositions 7, 8, and 9) and some small-step-size assumptions (*e.g.*, $\beta = \mathcal{O}(1/nL)$, $\beta/\alpha = r \gtrsim \kappa_2^2$), we eventually have

$$\mathbb{E}\left[V_\lambda(\boldsymbol{z}_0^{k+1})\right] - \mathbb{E}\left[V_\lambda(\boldsymbol{z}_0^k)\right] \le -n\alpha\mathbb{E}\left[\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2\right] - (L\kappa_2 n\alpha/2)\mathbb{E}\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right] + C\alpha^3,$$

where $C \ge 0$ is a constant (with respect to k) depending on L, n, B, and $r = \beta/\alpha$. (Lemma 20). We note that the step size ratio $r \gtrsim \kappa_2^2$ is crucial for showing that the coefficient in front of the term $\mathbb{E}\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right]$ is non-positive: even if it is possible with $r \lesssim \kappa_2^2$, we must assume that κ_2 upper-bounded by a positive numerical constant, which is not desirable for showing convergence bounds. Thus, we expect that a different proof strategy should be applied to avoid the requirement $r \gtrsim \kappa_2^2$ on the step size ratio.

The proofs of Theorems 1 and 2 diverge from here. The rest of the proof is mostly about choosing appropriate step sizes and solving the recurrence inequalities. Readers can find the full proofs for more general mini-batch sizes ($b \ge 1$) in Appendix C (for simSGDA-RR) and D (for altSGDA-RR).

7 CONCLUSION

We investigated two-time-scale stochastic algorithms based on without-replacement component sampling, called simSGDA-RR and altSGDA-RR, for solving smooth nonconvex finite-sum minimax optimization problems. We established convergence rates under the $\boldsymbol{y}$-side PŁ condition (nonconvex-PŁ) and, additionally, the primal PŁ condition (PŁ(Φ)-PŁ). We ascertain that the SGDA-RR can achieve a faster rate than its with-replacement counterpart, which agrees with the existing theory on without-replacement SGD for minimization. Lastly, we provided lower bounds for two-time-scale simGDA with any fixed step size ratio r, demonstrating that the full-batch upper bound with $r \gtrsim \kappa_2^2$ for PŁ(Φ)-PŁ functions is tight. Possible future directions include widening our results beyond sim/altSGDA (*e.g.*, extra-gradient or optimistic GDA) and beyond RR (*e.g.*, single/adversarial shuffling). As also discussed in Section 5, an interesting open question remains open: can we identify tight convergence rates for stochastic (with-/without-replacement) and/or deterministic GDA with step size ratio r satisfying $\kappa_2 \lesssim r \lesssim \kappa_2^2$, for general nonconvex-PŁ problems?

REFERENCES

Kwangjun Ahn, Chulhee Yun, and Suvrit Sra. SGD with shuffling: optimal rates without component convexity and large epoch requirements. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin (eds.), *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*, 2020. URL https://proceedings.neurips.cc/paper/2020/hash/cb8acb1dc9821bf74e6ca9068032d623-Abstract.html.

Aleksandr Beznosikov, Eduard Gorbunov, Hugo Berard, and Nicolas Loizou. Stochastic gradient descent-ascent: Unified theory and new efficient methods. *CoRR*, abs/2202.07262, 2022. URL https://arxiv.org/abs/2202.07262.

Aniket Das, Bernhard Schölkopf, and Michael Muehlebach. Sampling without replacement leads to faster rates in finite-sum minimax optimization. *CoRR*, abs/2206.02953, 2022. doi: 10.48550/arXiv.2206.02953. URL https://doi.org/10.48550/arXiv.2206.02953.

Ian J. Goodfellow, Jean Pouget-Abadie, Mehdi Mirza, Bing Xu, David Warde-Farley, Sherjil Ozair, Aaron C. Courville, and Yoshua Bengio. Generative adversarial nets. In Zoubin Ghahramani, Max Welling, Corinna Cortes, Neil D. Lawrence, and Kilian Q. Weinberger (eds.), *Advances in Neural Information Processing Systems 27: Annual Conference on Neural Information Processing Systems 2014, December 8-13 2014, Montreal, Quebec, Canada*, pp. 2672–2680, 2014. URL https://proceedings.neurips.cc/paper/2014/hash/5ca3e9b122f61f8f06494c97b1afccf3-Abstract.html.

Zhishuai Guo, Zhuoning Yuan, Yan Yan, and Tianbao Yang. Fast objective and duality gap convergence for non-convex strongly-concave min-max problems. *CoRR*, abs/2006.06889, 2020. URL https://arxiv.org/abs/2006.06889.

Martin Heusel, Hubert Ramsauer, Thomas Unterthiner, Bernhard Nessler, and Sepp Hochreiter. Gans trained by a two time-scale update rule converge to a local nash equilibrium. In Isabelle Guyon, Ulrike von Luxburg, Samy Bengio, Hanna M. Wallach, Rob Fergus, S. V. N. Vishwanathan, and Roman Garnett (eds.), *Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, December 4-9, 2017, Long Beach, CA, USA*, pp. 6626–6637, 2017. URL https://proceedings.neurips.cc/paper/2017/hash/8a1d694707eb0fefe65871369074926d-Abstract.html.

Roger A Horn and Charles R Johnson. *Matrix Analysis*. Cambridge University Press, Cambridge, England, 2 edition, October 2012.

Chi Jin, Praneeth Netrapalli, and Michael I. Jordan. What is local optimality in nonconvex-nonconcave minimax optimization? In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pp. 4880–4889. PMLR, 2020. URL http://proceedings.mlr.press/v119/jin20e.html.

Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In Paolo Frasconi, Niels Landwehr, Giuseppe Manco, and Jilles Vreeken (eds.), *Machine Learning and Knowledge Discovery in Databases - European Conference, ECML PKDD 2016, Riva del Garda, Italy, September 19-23, 2016, Proceedings, Part I*, volume 9851 of *Lecture Notes in Computer Science*, pp. 795–811. Springer, 2016. doi: 10.1007/978-3-319-46128-1_50. URL https://doi.org/10.1007/978-3-319-46128-1_50.

Haochuan Li, Farzan Farnia, Subhro Das, and Ali Jadbabaie. On convergence of gradient descent ascent: A tight local analysis. In Kamalika Chaudhuri, Stefanie Jegelka, Le Song, Csaba Szepesvári, Gang Niu, and Sivan Sabato (eds.), *International Conference on Machine Learning, ICML 2022, 17-23 July 2022, Baltimore, Maryland, USA*, volume 162 of *Proceedings of Machine Learning Research*, pp. 12717–12740. PMLR, 2022. URL https://proceedings.mlr.press/v162/li22e.html.

Shihui Li, Yi Wu, Xinyue Cui, Honghua Dong, Fei Fang, and Stuart Russell. Robust multi-agent reinforcement learning via minimax deep deterministic policy gradient. In *The Thirty-Third AAAI Conference on Artificial Intelligence, AAAI 2019, The Thirty-First Innovative Applications of Artificial Intelligence Conference, IAAI 2019, The Ninth AAAI Symposium on Educational Advances in Artificial Intelligence, EAAI 2019, Honolulu, Hawaii, USA, January 27 - February 1, 2019*, pp. 4213–4220. AAAI Press, 2019. doi: 10.1609/aaai.v33i01.33014213. URL https://doi.org/10.1609/aaai.v33i01.33014213.

Tianyi Lin, Chi Jin, and Michael I. Jordan. On gradient descent ascent for nonconvex-concave minimax problems. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pp. 6083–6093. PMLR, 2020. URL http://proceedings.mlr.press/v119/lin20a.html.

Nicolas Loizou, Hugo Berard, Gauthier Gidel, Ioannis Mitliagkas, and Simon Lacoste-Julien. Stochastic gradient descent-ascent and consensus optimization for smooth games: Convergence analysis under expected co-coercivity. In Marc'Aurelio Ranzato, Alina Beygelzimer, Yann N. Dauphin, Percy Liang, and Jennifer Wortman Vaughan (eds.), *Advances in Neural Information Processing Systems 34: Annual Conference on Neural Information Processing Systems 2021, NeurIPS 2021, December 6-14, 2021, virtual*, pp. 19095–19108, 2021. URL https://proceedings.neurips.cc/paper/2021/hash/9f96f36b7aae3b1ff847c26ac94c604e-Abstract.html.

Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. In *6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Conference Track Proceedings*. OpenReview.net, 2018. URL https://openreview.net/forum?id=rJzIBfZAb.

Chinmay Maheshwari, Chih-Yuan Chiu, Eric Mazumdar, Shankar Sastry, and Lillian Ratliff. Zeroth-order methods for convex-concave min-max problems: Applications to decision-dependent risk minimization. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera (eds.), *Proceedings of The 25th International Conference on Artificial Intelligence and Statistics*, volume 151 of *Proceedings of Machine Learning Research*, pp. 6702–6734. PMLR, 28–30 Mar 2022. URL https://proceedings.mlr.press/v151/maheshwari22a.html.

Konstantin Mishchenko, Ahmed Khaled, and Peter Richtárik. Random reshuffling: Simple analysis with vast improvements. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin (eds.), *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*, 2020. URL https://proceedings.neurips.cc/paper/2020/hash/c8cc6e90ccbff44c9cee23611711cdc4-Abstract.html.

Dheeraj Nagaraj, Prateek Jain, and Praneeth Netrapalli. SGD without replacement: Sharper rates for general smooth convex functions. In Kamalika Chaudhuri and Ruslan Salakhutdinov (eds.), *Proceedings of the 36th International Conference on Machine Learning, ICML 2019, 9-15 June 2019, Long Beach, California, USA*, volume 97 of *Proceedings of Machine Learning Research*, pp. 4703–4711. PMLR, 2019. URL http://proceedings.mlr.press/v97/nagaraj19a.html.

Lam M. Nguyen, Quoc Tran-Dinh, Dzung T. Phan, Phuong Ha Nguyen, and Marten van Dijk. A unified convergence analysis for shuffling-type gradient methods. *J. Mach. Learn. Res.*, 22: 207:1–207:44, 2021. URL http://jmlr.org/papers/v22/20-1238.html.

Maher Nouiehed, Maziar Sanjabi, Tianjian Huang, Jason D. Lee, and Meisam Razaviyayn. Solving a class of non-convex min-max games using iterative first order methods. In Hanna M. Wallach, Hugo Larochelle, Alina Beygelzimer, Florence d'Alché-Buc, Emily B. Fox, and Roman Garnett (eds.), *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, December 8-14, 2019, Vancouver, BC, Canada*, pp. 14905–14916, 2019. URL https://proceedings.neurips.cc/paper/2019/hash/25048eb6a33209cb5a815bff0cf6887c-Abstract.html.

Shashank Rajput, Anant Gupta, and Dimitris S. Papailiopoulos. Closing the convergence gap of SGD without replacement. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pp. 7964–7973. PMLR, 2020. URL http://proceedings.mlr.press/v119/rajput20a.html.

Aman Sinha, Hongseok Namkoong, and John Duchi. Certifiable distributional robustness with principled adversarial training. In *International Conference on Learning Representations*, 2018. URL https://openreview.net/forum?id=Hk6kPgZA-.

Jiahao Xie, Chao Zhang, Zebang Shen, Weijie Liu, and Hui Qian. Efficient cross-device federated learning algorithms for minimax problems. *arXiv e-prints*, pp. arXiv–2105, 2021. URL https://arxiv.org/abs/2105.14216.

Yan Yan, Yi Xu, Qihang Lin, Wei Liu, and Tianbao Yang. Optimal epoch stochastic gradient descent ascent methods for min-max optimization. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin (eds.), *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*, 2020. URL https://proceedings.neurips.cc/paper/2020/hash/3f8b2a81da929223ae025fcec26dde0d-Abstract.html.

Junchi Yang, Negar Kiyavash, and Niao He. Global convergence and variance reduction for a class of nonconvex-nonconcave minimax problems. In Hugo Larochelle, Marc'Aurelio Ranzato, Raia Hadsell, Maria-Florina Balcan, and Hsuan-Tien Lin (eds.), *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*, 2020. URL https://proceedings.neurips.cc/paper/2020/hash/0cc6928e741d75e7a92396317522069e-Abstract.html.

Yiming Ying, Longyin Wen, and Siwei Lyu. Stochastic online AUC maximization. In Daniel D. Lee, Masashi Sugiyama, Ulrike von Luxburg, Isabelle Guyon, and Roman Garnett (eds.), *Advances in Neural Information Processing Systems 29: Annual Conference on Neural Information Processing Systems 2016, December 5-10, 2016, Barcelona, Spain*, pp. 451–459, 2016. URL https://proceedings.neurips.cc/paper/2016/hash/c52f1bd66cc19d05628bd8bf27af3ad6-Abstract.html.

Yaodong Yu, Tianyi Lin, Eric V. Mazumdar, and Michael I. Jordan. Fast distributionally robust learning with variance-reduced min-max optimization. In Gustau Camps-Valls, Francisco J. R. Ruiz, and Isabel Valera (eds.), *International Conference on Artificial Intelligence and Statistics, AISTATS 2022, 28-30 March 2022, Virtual Event*, volume 151 of *Proceedings of Machine Learning Research*, pp. 1219–1250. PMLR, 2022. URL https://proceedings.mlr.press/v151/yu22a.html.

Zhuoning Yuan, Yan Yan, Milan Sonka, and Tianbao Yang. Large-scale robust deep AUC maximization: A new surrogate loss and empirical studies on medical image classification. In *2021 IEEE/CVF International Conference on Computer Vision, ICCV 2021, Montreal, QC, Canada, October 10-17, 2021*, pp. 3020–3029. IEEE, 2021. doi: 10.1109/ICCV48922.2021.00303. URL https://doi.org/10.1109/ICCV48922.2021.00303.

Chulhee Yun, Suvrit Sra, and Ali Jadbabaie. Open problem: Can single-shuffle SGD be better than reshuffling SGD and GD? In Mikhail Belkin and Samory Kpotufe (eds.), *Conference on Learning Theory, COLT 2021, 15-19 August 2021, Boulder, Colorado, USA*, volume 134 of *Proceedings of Machine Learning Research*, pp. 4653–4658. PMLR, 2021. URL http://proceedings.mlr.press/v134/open-problem-yun21a.html.

Chulhee Yun, Shashank Rajput, and Suvrit Sra. Minibatch vs local SGD with shuffling: Tight convergence bounds and beyond. In *The Tenth International Conference on Learning Representations, ICLR 2022, Virtual Event, April 25-29, 2022*. OpenReview.net, 2022. URL https://openreview.net/forum?id=LdlwBP2mlq.

CONTENTS

A   MINI-BATCH SGDA-RR AND CONVERGENCE RATES

Here, we present an algorithm that extends simSGDA-RR and altSGDA-RR by using mini-batches of size $b \geq 1$. For simplicity, we assume that the number of components n is an integer multiple of the mini-batch size b for simplicity; *i.e.*, for some integer $q \geq 1$, $n = bq$. One can easily extend this to the case when n does not need to be a multiple of b (namely, $n = b(q-1) + s$, where $q \geq 1$, $s \in [b]$) so that there are $q - 1$ mini-batches of size b and one more mini-batch of size $s \leq b$.

Algorithm 2 Mini-batch simSGDA/altSGDA-RR

1: **Given:** The number of components n; the size of mini-batches b (a divisor of n); the number of epochs K; step sizes $\alpha, \beta > 0$
2: **Initialize:** $(\boldsymbol{x}_0^1; \boldsymbol{y}_0^1) \in \mathbb{R}^{d_x} \times \mathbb{R}^{d_y}$
3: **for** $k \in [K]$ **do**
4:     Sample $\sigma_k \sim \mathrm{Unif}(\mathbb{S}_n)$ ▷ RR: uniformly randomly shuffle the indices every epoch
5:     **for** $t \in [n/b]$ **do**
6:         $\mathcal{B}_t^k := \{\sigma_k(b(t-1)+1), \cdots, \sigma_k(bt)\}$ ▷ Mini-batch : a set of component indices
7:         $\boldsymbol{x}_t^k = \boldsymbol{x}_{t-1}^k - \frac{\alpha}{b} \sum_{i \in \mathcal{B}_t^k} \nabla_1 f_i(\boldsymbol{x}_{t-1}^k; \boldsymbol{y}_{t-1}^k)$
8:         **if** simSGDA-RR **then**
9:             $\boldsymbol{y}_t^k = \boldsymbol{y}_{t-1}^k + \frac{\beta}{b} \sum_{i \in \mathcal{B}_t^k} \nabla_2 f_i(\boldsymbol{x}_{t-1}^k; \boldsymbol{y}_{t-1}^k)$ ▷ simultaneous update: $\boldsymbol{x}$ & $\boldsymbol{y}$
10:        **else if** altSGDA-RR **then**
11:            $\boldsymbol{y}_t^k = \boldsymbol{y}_{t-1}^k + \frac{\beta}{b} \sum_{i \in \mathcal{B}_t^k} \nabla_2 f_i(\boldsymbol{x}_t^k; \boldsymbol{y}_{t-1}^k)$ ▷ alternating update: $\boldsymbol{x} \to \boldsymbol{y}$
12:    $(\boldsymbol{x}_0^{k+1}; \boldsymbol{y}_0^{k+1}) = (\boldsymbol{x}_{n/b}^k; \boldsymbol{y}_{n/b}^k)$

Next, we illustrate the generalized versions of our main results (Theorems 1 and 2) for Algorithm 2 with mini-batches of size $b \geq 1$. Let us assume $n \geq 2$ because the case $n = 1$ trivially boils down to simGDA or altGDA. We defer the proofs for simultaneous updates to Appendix C. We present the parts that change in the proof for alternating updates in Appendix D.

Theorem 4 (Nonconvex-PŁ, mini-batch SGDA-RR). *Suppose that f satisfies Assumptions 1, 2, 3, and 4. Let $\lambda = 4$. Choose the step sizes α and β by $\alpha = \beta/r$ for some $r \geq 14\kappa_2^2$ and*

$$\beta = b \cdot \min \left\{ \frac{1}{6Ln\sqrt{1 + \frac{n-b}{n-1} \cdot \frac{A}{n}}}, \frac{1}{c} \left(\frac{\cdot V_\lambda(\boldsymbol{z}_0^1)}{Ln^2(\frac{n-b}{n-1})BK} \right)^{\frac{1}{3}} \right\},$$

for some numerical constant $c > 0$. Then, mini-batch simSGDA-RR and altSGDA-RR with mini-batch size b (a divisor of n) satisfy

$$\min_{k \in [K]} \mathbb{E} \left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \leq \frac{1}{K} \sum_{k=1}^K \mathbb{E} \left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2$$

$$\leq \frac{6rLV_\lambda(\boldsymbol{z}_0^1)}{K} \sqrt{1 + \left(\frac{n-b}{n-1} \right) \frac{A}{n}} + cr \left(\frac{L^2 B V_\lambda(\boldsymbol{z}_0^1)^2}{nK^2} \cdot \frac{n-b}{n-1} \right)^{1/3}.$$

Theorem 5 (PŁ(Φ)-PŁ, mini-batch SGDA-RR). *Suppose that f satisfies Assumptions 1, 2, 3, 4, and 5. Let $\lambda = 4$. Choose the step sizes α and β by $\alpha = \beta/r$ for some $r \geq 14\kappa_2^2$ and*

$$\beta = b \cdot \min \left\{ \frac{1}{6Ln\sqrt{1 + \frac{n-b}{n-1} \cdot \frac{A}{n}}}, \frac{2r}{\mu_1 nK} \max \left\{ 1, \ \log \left(\frac{V_\lambda(\boldsymbol{z}_0^1)\mu_1 nK^2}{8c^3\kappa_1^2 r^3 \left(\frac{n-b}{n-1} \right) B} \right) \right\} \right\},$$

for some numerical constant $c > 0$. Then, mini-batch simSGDA-RR and altSGDA-RR with mini-batch size b (a divisor of n) satisfy

$$\mathbb{E}[V_\lambda(\boldsymbol{z}_0^{K+1})] \leq \mathcal{O} \left(V_\lambda(\boldsymbol{z}_0^1) \cdot \exp \left(-\frac{K}{12\kappa_1 r \sqrt{1 + \frac{n-b}{n-1} \frac{A}{n}}} \right) \right) + \tilde{\mathcal{O}} \left(\frac{\kappa_1^2 r^3 B}{\mu_1 nK^2} \right) \cdot \frac{n-b}{n-1}.$$

As a side remark, some works consider a sampling method called *b-minibatch sampling* where all the elements in each mini-batch are distinct (*i.e.*, without-replacement component sampling per mini-batch), *e.g.*, Loizou et al. (2021, Definition 2.1). However, there is a significant gap between this method and ours: any two distinct mini-batches sampled by the *b*-minibatch sampling can intersect to each other (*i.e.*, mini-batches are sampled with replacement), whereas in each epoch of our Algorithm 2, all the mini-batches are mutually disjoint.

B TECHNICAL PROPOSITIONS

Throughout this appendix, we use $\mathcal{X} = \mathbb{R}^{d_x}$ and $\mathcal{Y} = \mathbb{R}^{d_y}$.

B.1 FUNCTION CLASSES: PŁ CONDITION, SMOOTHNESS, AND MORE

Proposition 6 ($\kappa \geq 1$). *Let g be an L-smooth function which is bounded below by g^*. Then, for any* $\boldsymbol{x}$,

$$\|\nabla g(\boldsymbol{x})\|^2 \leq 2L \left[g(\boldsymbol{x}) - g^* \right].$$

If g is μ-PŁ as well, then $\mu \leq L$. Consequently, the condition number $\kappa := L/\mu$ of g is ≥ 1.

Proof. Since g is L-smooth, for any $\boldsymbol{x}$ and $\boldsymbol{y}$,

$$g^* \leq g(\boldsymbol{y}) \leq g(\boldsymbol{x}) + \langle \nabla g(\boldsymbol{x}), \boldsymbol{y} - \boldsymbol{x} \rangle + \frac{L}{2} \|\boldsymbol{y} - \boldsymbol{x}\|^2. \tag{5}$$

Now define a convex quadratic function $h_x(\boldsymbol{y})$ of $\boldsymbol{y}$ as

$$h_x(\boldsymbol{y}) := g(\boldsymbol{x}) + \langle \nabla g(\boldsymbol{x}), \boldsymbol{y} - \boldsymbol{x} \rangle + \frac{L}{2} \|\boldsymbol{y} - \boldsymbol{x}\|^2.$$

Since its gradient is

$$\nabla h_x(\boldsymbol{y}) = \nabla g(\boldsymbol{x}) + L(\boldsymbol{y} - \boldsymbol{x}),$$

$\boldsymbol{y}^* := \boldsymbol{x} - \frac{1}{L} \nabla g(\boldsymbol{x})$ is a minimum of h_x. Plugging $\boldsymbol{y} = \boldsymbol{y}^*$ to the equation (5), we get

$$g^* \leq g(\boldsymbol{x}) + \left\langle \nabla g(\boldsymbol{x}), -\frac{1}{L} \nabla g(\boldsymbol{x}) \right\rangle + \frac{L}{2} \left\| -\frac{1}{L} \nabla g(\boldsymbol{x}) \right\|^2 = g(\boldsymbol{x}) - \frac{1}{2L} \|\nabla g(\boldsymbol{x})\|^2.$$

Rearranging the terms,

$$\|\nabla g(\boldsymbol{x})\|^2 \leq 2L \left[g(\boldsymbol{x}) - g^* \right].$$

If we additionally utilize PŁ inequality with $g^* := \min g(\boldsymbol{x})$,

$$\|\nabla g(\boldsymbol{x})\|^2 \geq 2\mu \left[g(\boldsymbol{x}) - g^* \right],$$

we directly yield $\mu \leq L$ and thus $\kappa = L/\mu \geq 1$. $\qquad\square$

Definition 1 ([Karimi et al. (2016)](#)). *Consider $g : \mathcal{X} \to \mathbb{R}$. Let $\boldsymbol{x}_p \in \Pi_{\mathcal{X}^*}(\boldsymbol{x})$ be a projection of $\boldsymbol{x}$ onto the optimal set $\mathcal{X}^* = \arg\min_{\boldsymbol{x} \in \mathcal{X}} g(\boldsymbol{x})$.*

(1) We say g satisfies μ-strong convexity (SC) if $g(\boldsymbol{x}') \geq g(\boldsymbol{x}) + \langle \nabla g(\boldsymbol{x}), \boldsymbol{x}' - \boldsymbol{x} \rangle + \frac{\mu}{2} \|\boldsymbol{x}' - \boldsymbol{x}\|^2$ for any $\boldsymbol{x}, \boldsymbol{x}' \in \mathcal{X}$.

(2) We say g satisfies μ-restricted secant inequality (RSI) if $\langle \nabla g(\boldsymbol{x}), \boldsymbol{x} - \boldsymbol{x}_p \rangle \geq \mu \|\boldsymbol{x}_p - \boldsymbol{x}\|^2$ for any $\boldsymbol{x} \in \mathcal{X}$.

(3) We say g satisfies μ-error bound (EB) condition if $\|\nabla g(\boldsymbol{x})\| \geq \mu \|\boldsymbol{x}_p - \boldsymbol{x}\|$ for any $\boldsymbol{x} \in \mathcal{X}$.

(4) We say g satisfies μ-quadratic growth (QG) condition if $g(\boldsymbol{x}) - \min_{\boldsymbol{x}'} g(\boldsymbol{x}') \geq \frac{\mu}{2} \|\boldsymbol{x}_p - \boldsymbol{x}\|^2$ for any $\boldsymbol{x} \in \mathcal{X}$.

Proposition 7. *From Definition [1](#), The following implications are true.*

- *μ-SC implies μ-PŁ and μ-RSI.*

- *μ-PŁ implies μ-QG and μ-EB.*

- *μ-RSI implies μ-EB.*

- *μ-EB and L-smoothness together imply (μ^2/L)-PŁ.*

Proof. Most of the proofs are originated from Karimi et al. (2016, Theorem 2).

(SC $\Rightarrow$ PŁ) Substitute $\boldsymbol{x}$ to $\boldsymbol{x}_p$ and $\boldsymbol{x}'$ to $\boldsymbol{x}$, respectively, from Definition 1.(1).

(PŁ $\Rightarrow$ QG & EB) See the proof in Karimi et al. (2016, Theorem 2)

(SC $\Rightarrow$ RSI) We know μ-SC $\Rightarrow$ μ-PŁ $\Rightarrow$ μ-QG. From Definition 1.(1) & 1.(4),

$$\langle \nabla g(\boldsymbol{x}), \boldsymbol{x} - \boldsymbol{x}_p \rangle \overset{\text{SC}}{\geq} g(\boldsymbol{x}) - g(\boldsymbol{x}_p) + \frac{\mu}{2} \|\boldsymbol{x}_p - \boldsymbol{x}\|^2$$

$$\overset{\text{QG}}{\geq} \frac{\mu}{2} \|\boldsymbol{x}_p - \boldsymbol{x}\|^2 + \frac{\mu}{2} \|\boldsymbol{x}_p - \boldsymbol{x}\|^2 = \mu \|\boldsymbol{x}_p - \boldsymbol{x}\|^2 .$$

This implies μ-RSI.

(RSI $\Rightarrow$ EB) See the proof in Karimi et al. (2016, Theorem 2).

(EB & smooth $\Rightarrow$ PŁ) We use $\nabla g(\boldsymbol{x}_p) = \boldsymbol{0}$. By L-smoothness and μ-EB condition,

$$g(\boldsymbol{x}) - g(\boldsymbol{x}_p) \overset{\text{smooth}}{\leq} \langle \nabla g(\boldsymbol{x}_p), \boldsymbol{x} - \boldsymbol{x}_p \rangle + \frac{L}{2} \|\boldsymbol{x} - \boldsymbol{x}_p\|^2 = \frac{L}{2} \|\boldsymbol{x} - \boldsymbol{x}_p\|^2$$

$$\overset{\text{EB}}{\leq} \frac{L}{2\mu^2} \|\nabla g(\boldsymbol{x})\|^2 .$$

This implies (μ^2/L)-PŁ condition on g. $\qquad\square$

Proposition 8 (Lipschitz continuity-like property of $\boldsymbol{y}^*(\boldsymbol{x})$). *For an L-smooth function $g : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$, suppose $-g(\boldsymbol{x}; \cdot)$ is μ_2-PŁ. Let $\kappa_2 = L/\mu_2$.*

Consider any $\boldsymbol{x}_0, \boldsymbol{x}_1 \in \mathcal{X}$. For any $\boldsymbol{y}_0^* \in \mathcal{Y}_{\boldsymbol{x}_0}^* = \arg\max_{\boldsymbol{y} \in \mathcal{Y}} g(\boldsymbol{x}_0; \boldsymbol{y})$, there exists a $\boldsymbol{y}_1^* \in \mathcal{Y}_{\boldsymbol{x}_1}^* = \arg\max_{\boldsymbol{y} \in \mathcal{Y}} g(\boldsymbol{x}_1; \boldsymbol{y})$ such that $\|\boldsymbol{y}_0^* - \boldsymbol{y}_1^*\| \leq \kappa_2 \|\boldsymbol{x}_0 - \boldsymbol{x}_1\|$.

In fact, it is enough to choose $\boldsymbol{y}_1^*$ as a projection of $\boldsymbol{y}_0^*$ onto the set $\mathcal{Y}_{\boldsymbol{x}_1}^*$, namely, $\boldsymbol{y}_1^* \in \Pi_{\mathcal{Y}_{\boldsymbol{x}_1}^*}(\boldsymbol{y}_0^*)$.

Proof. We borrow the proof from Nouiehed et al. (2019, Lemma A.3).

Recall $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}' \in \mathcal{Y}} g(\boldsymbol{x}; \boldsymbol{y}')$. By PŁ inequality and smoothness of g,

$$2\mu_2 \left(\Phi(\boldsymbol{x}_1) - g(\boldsymbol{x}_1; \boldsymbol{y}_0^*) \right) \leq \|\nabla_2 f(\boldsymbol{x}_1; \boldsymbol{y}_0^*)\|^2$$

$$= \|\nabla_2 g(\boldsymbol{x}_1; \boldsymbol{y}_0^*) - \nabla_2 g(\boldsymbol{x}_0; \boldsymbol{y}_0^*)\|^2 \leq L^2 \|\boldsymbol{x}_1 - \boldsymbol{x}_0\|^2 .$$

The second equality applies $\nabla_2 g(\boldsymbol{x}_0; \boldsymbol{y}_0^*) = \boldsymbol{0}$, since $\boldsymbol{y}_0^* \in \arg\max_{\boldsymbol{y}} g(\boldsymbol{x}_0; \boldsymbol{y})$.

Moreover, note that $-g(\boldsymbol{x}_1; \cdot)$ satisfies μ_2-QG condition ($\because$ Proposition 7). To apply this, we utilize our choice of $\boldsymbol{y}_1^*$:

$$\Phi(\boldsymbol{x}_1) - g(\boldsymbol{x}_1; \boldsymbol{y}_0^*) \geq \frac{\mu_2}{2} \|\boldsymbol{y}_1^* - \boldsymbol{y}_0^*\|^2 .$$

As a result, we have $\mu_2^2 \|\boldsymbol{y}_0^* - \boldsymbol{y}_1^*\|^2 \leq L^2 \|\boldsymbol{x}_0 - \boldsymbol{x}_1\|^2$. This completes the proof. $\qquad\square$

Proposition 9 (Smoothness of primal function). *Consider the same function g as Proposition 8. Then, the function $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}' \in \mathcal{Y}} g(\boldsymbol{x}; \boldsymbol{y}')$ is differentiable with*

$$\nabla \Phi(\boldsymbol{x}) = \nabla_1 g(\boldsymbol{x}; \boldsymbol{y}^*(\boldsymbol{x})), \quad \text{regardless of the choice of } \boldsymbol{y}^*(\boldsymbol{x}) \in \arg\max_{\boldsymbol{y}' \in \mathcal{Y}} g(\boldsymbol{x}; \boldsymbol{y}').$$

Moreover, Φ is $L(\kappa_2 + 1)$-smooth, where $\kappa_2 = L/\mu_2$.

Proof. This is already proved in Lemma A.5 of Nouiehed et al. (2019). However, we present a bit different proof without using second-order Taylor expansion. To start, recall $\mathcal{Y}_{\boldsymbol{x}}^* := \arg\max_{\boldsymbol{y} \in \mathcal{Y}} g(\boldsymbol{x}; \boldsymbol{y})$. That is, we could choose any $\boldsymbol{y}^*(\boldsymbol{x}) \in \mathcal{Y}_{\boldsymbol{x}}^*$.

We first show the differentiability of Φ. Fix a unit vector $\boldsymbol{u} \in \mathcal{X} = \mathbb{R}^{d_x} : \|\boldsymbol{u}\| = 1$. Let any $h > 0$. We first claim that there exists a path $\boldsymbol{p} : (-h, h] \to \mathcal{Y} = \mathbb{R}^{d_y}$ which is continuous at $t = 0$ and

$\boldsymbol{p}(t) \in \mathcal{Y}^*_{(\boldsymbol{x}+t\boldsymbol{u})}$. In fact, let $\boldsymbol{p}(t)$ be a projection of $\boldsymbol{y}^*(\boldsymbol{x})$ (that we chose) onto the set $\mathcal{Y}^*_{(\boldsymbol{x}+t\boldsymbol{u})}$. Then, $\boldsymbol{p}(0) = \boldsymbol{y}^*(\boldsymbol{x})$, and by Proposition 8, we have $\|\boldsymbol{p}(0) - \boldsymbol{p}(t)\| \leq \kappa_2 \|\boldsymbol{x} - (\boldsymbol{x} + t\boldsymbol{u})\| = \kappa_2 t$. This shows the continuity of $\boldsymbol{p}(t)$ at $t = 0$. Now, note that there exists a $t_1 \in (0, h)$ such that,

$$
\begin{aligned}
\Phi(\boldsymbol{x} + h\boldsymbol{u}) - \Phi(\boldsymbol{x}) &= g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(h)) - g(\boldsymbol{x}; \boldsymbol{p}(0)) \\
&= \{g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(h)) - g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(0))\} + \{g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(0)) - g(\boldsymbol{x}; \boldsymbol{p}(0))\} \\
&\geq 0 + \langle \nabla_1 g(\boldsymbol{x} + t_1\boldsymbol{u}; \boldsymbol{p}(0)), h\boldsymbol{u} \rangle,
\end{aligned}
$$

by mean value theorem (applied to the first argument). We have the inequality at the last line because $g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(h)) \geq g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(0))$, since $\boldsymbol{p}(h) \in \mathcal{Y}^*_{(\boldsymbol{x}+h\boldsymbol{u})}$. With a similar logic, there exists a $t_2 \in (0, h)$ such that,

$$
\begin{aligned}
\Phi(\boldsymbol{x} + h\boldsymbol{u}) - \Phi(\boldsymbol{x}) &= g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(h)) - g(\boldsymbol{x}; \boldsymbol{p}(0)) \\
&= \{g(\boldsymbol{x} + h\boldsymbol{u}; \boldsymbol{p}(h)) - g(\boldsymbol{x}; \boldsymbol{p}(h))\} + \{g(\boldsymbol{x}; \boldsymbol{p}(h)) - g(\boldsymbol{x}; \boldsymbol{p}(0))\} \\
&\leq \langle \nabla_1 g(\boldsymbol{x} + t_2\boldsymbol{u}; \boldsymbol{p}(h)), h\boldsymbol{u} \rangle + 0.
\end{aligned}
$$

To combine these two inequalities into a single line,

$$
\langle \nabla_1 g(\boldsymbol{x} + t_1\boldsymbol{u}; \boldsymbol{p}(0)), \boldsymbol{u} \rangle \leq \frac{\Phi(\boldsymbol{x} + h\boldsymbol{u}) - \Phi(\boldsymbol{x})}{h} \leq \langle \nabla_1 g(\boldsymbol{x} + t_2\boldsymbol{u}; \boldsymbol{p}(h)), \boldsymbol{u} \rangle.
$$

Using the continuity of $p(\cdot)$ and $\nabla_1 g(\cdot; \cdot)$ ($\because g$ has Lipschitz continuous gradient), we can deduce that the directional derivative of Φ in a direction $\boldsymbol{u}$ (denoted by $D_{\boldsymbol{u}}\Phi$) is in fact

$$
D_{\boldsymbol{u}}\Phi(\boldsymbol{x}) = \langle \nabla_1 g(\boldsymbol{x}; \boldsymbol{y}^*(\boldsymbol{x})), \boldsymbol{u} \rangle,
$$

by taking the limit $h \to 0+$. Since $\boldsymbol{u}$ is arbitrary, we can conclude that $\nabla\Phi(\boldsymbol{x}) = \nabla_1 g(\boldsymbol{x}; \boldsymbol{y}^*(\boldsymbol{x}))$.

The proof of Lipschitz smoothness of Φ exactly follows the proof by Nouiehed et al. (2019). Consider any $\boldsymbol{x}_0, \boldsymbol{x}_1 \in \mathcal{X}$. As in Proposition 8, choose any $\boldsymbol{y}_0^* \in \mathcal{Y}^*_{\boldsymbol{x}_0}$ and $\boldsymbol{y}_1^* \in \Pi_{\mathcal{Y}^*_{\boldsymbol{x}_1}}(\boldsymbol{y}_0^*)$. Then,

$$
\begin{aligned}
\|\nabla\Phi(\boldsymbol{x}_0) - \nabla\Phi(\boldsymbol{x}_1)\| &= \|\nabla_1 g(\boldsymbol{x}_0; \boldsymbol{y}_0^*) - \nabla_1 g(\boldsymbol{x}_1; \boldsymbol{y}_1^*)\| \\
&\leq \|\nabla_1 g(\boldsymbol{x}_0; \boldsymbol{y}_0^*) - \nabla_1 g(\boldsymbol{x}_1; \boldsymbol{y}_0^*)\| + \|\nabla_1 g(\boldsymbol{x}_1; \boldsymbol{y}_0^*) - \nabla_1 g(\boldsymbol{x}_1; \boldsymbol{y}_1^*)\| \\
&\leq L\{\|\boldsymbol{x}_0 - \boldsymbol{x}_1\| + \|\boldsymbol{y}_0^* - \boldsymbol{y}_1^*\|\} \\
&\leq L(1 + \kappa_2)\|\boldsymbol{x}_0 - \boldsymbol{x}_1\|.
\end{aligned}
$$

The last inequality holds because of Proposition 8. $\qquad\square$

Proposition 10 ($\boldsymbol{x}$-side PŁ $\Rightarrow$ primal PŁ). *Suppose $g : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$ is L-smooth and two-sided PŁ with constants μ_1 and μ_2. Then, g satisfies primal PŁ condition: the function $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}' \in \mathcal{Y}} g(\boldsymbol{x}; \boldsymbol{y}')$ is μ_1-PŁ. As a result, a smooth two-sided PŁ function is PŁ(Φ)-PŁ.*

Proof. See Lemma A.3 of Yang et al. (2020). $\qquad\square$

Definition 2. *Consider $g : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$. Then, the point $(\boldsymbol{x}^*; \boldsymbol{y}^*) \in \mathcal{X} \times \mathcal{Y}$ is called*

    (i) a stationary point of g if $\nabla_1 g(\boldsymbol{x}^*; \boldsymbol{y}^*) = \nabla_2 g(\boldsymbol{x}^*; \boldsymbol{y}^*) = 0$.

    (ii) a saddle point of g if $g(\boldsymbol{x}^*; \boldsymbol{y}) \leq g(\boldsymbol{x}^*; \boldsymbol{y}^*) \leq g(\boldsymbol{x}; \boldsymbol{y}^*)$ for all $\boldsymbol{x}, \boldsymbol{y}$.

    (iii) a global minimax point of g if $g(\boldsymbol{x}^*; \boldsymbol{y}) \leq g(\boldsymbol{x}^*; \boldsymbol{y}^*) \leq \max_{\boldsymbol{y}'} g(\boldsymbol{x}; \boldsymbol{y}')$ for all $\boldsymbol{x}, \boldsymbol{y}$.

    (iv) a global maximin point of g if $\min_{\boldsymbol{x}'} g(\boldsymbol{x}'; \boldsymbol{y}) \leq g(\boldsymbol{x}^*; \boldsymbol{y}^*) \leq g(\boldsymbol{x}; \boldsymbol{y}^*)$ for all $\boldsymbol{x}, \boldsymbol{y}$.

Proposition 11. *Consider a function $g : \mathcal{X} \times \mathcal{Y} \to \mathbb{R}$.*

    (1) In general, a saddle point of g is a global minimax/maximin point.

(2) Let $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}} g(\boldsymbol{x}; \boldsymbol{y})$ and $\Phi^* := \min_x \Phi(\boldsymbol{x})$ be well-defined. Let $\lambda > 0$ be a constant. In general, a point $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ is a global minimax point of g if and only if

$$V_\lambda(\boldsymbol{x}^*; \boldsymbol{y}^*) := \lambda[\Phi(\boldsymbol{x}) - \Phi^*] + [\Phi(\boldsymbol{x}) - g(\boldsymbol{x}; \boldsymbol{y})] = 0.$$

(3) If g is smooth nonconvex-PŁ, then a global minimax point is a stationary point.

(4) If g is PŁ(Φ)-PŁ, then there exists a global minimax point $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ of g. As a result, if g is also smooth, then the point $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ is a stationary point.

(5) If g is smooth two-sided PŁ, every stationary point is a saddle point. As a result, there exists a saddle point $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ of g.

In particular, smooth two-sided PŁ functions enjoy the "minimax theorem," which establishes "minimax = maximin."

Proof. (1) (saddle point $\Rightarrow$ global minimax & global maximin) This is straightforward by the definitions: for any $\boldsymbol{x}$ and $\boldsymbol{y}$,

$$\min_{\boldsymbol{x}'} g(\boldsymbol{x}'; \boldsymbol{y}) \le g(\boldsymbol{x}^*; \boldsymbol{y}) \le g(\boldsymbol{x}^*; \boldsymbol{y}^*) \le g(\boldsymbol{x}; \boldsymbol{y}^*) \le \max_{\boldsymbol{y}'} g(\boldsymbol{x}; \boldsymbol{y}').$$

(2) (global minimax $\iff V_\lambda = 0$) The terms $\Phi(\boldsymbol{x}) - \Phi^*$ and $\Phi(\boldsymbol{x}) - g(\boldsymbol{x}; \boldsymbol{y})$ are non-negative. Hence, $V_\lambda(\boldsymbol{x}; \boldsymbol{y})$ is non-negative, and $V_\lambda(\boldsymbol{x}^*; \boldsymbol{y}^*) = 0$ if and only if $\Phi^* = \Phi(\boldsymbol{x}^*) = g(\boldsymbol{x}^*; \boldsymbol{y}^*)$, which is equivalent to the global minimax point condition.

(3) (smooth nonconvex-PŁ: global minimax $\Rightarrow$ stationary) Suppose $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ is a global minimax point. Since $g(\boldsymbol{x}^*; \boldsymbol{y}) \le g(\boldsymbol{x}^*; \boldsymbol{y}^*)$ for any $\boldsymbol{y}$, $\Phi(\boldsymbol{x}^*) = \max_{\boldsymbol{y}} g(\boldsymbol{x}^*; \boldsymbol{y}) = g(\boldsymbol{x}^*; \boldsymbol{y}^*)$. Thus, Φ has a minimum $g(\boldsymbol{x}^*; \boldsymbol{y}^*)$ at $\boldsymbol{x} = \boldsymbol{x}^*$. By Proposition 9, $\Phi(\cdot)$ is a differentiable function and we have

$$\nabla_1 g(\boldsymbol{x}^*; \boldsymbol{y}^*) = \nabla\Phi(\boldsymbol{x}^*) = 0.$$

Also, since a differentiable function $g(\boldsymbol{x}^*; \boldsymbol{y})$ has a maximum at $\boldsymbol{y} = \boldsymbol{y}^*$, we also have $\nabla_2 g(\boldsymbol{x}^*; \boldsymbol{y}^*) = 0$. Therefore, $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ is a stationary point.

(4) (PŁ(Φ)-PŁ: $\exists$ global minimax) Let $\boldsymbol{x}^* \in \arg\min_{\boldsymbol{x}} \Phi(\boldsymbol{x})$ and $\boldsymbol{y}^* \in \arg\max_{\boldsymbol{y}} f(\boldsymbol{x}^*; \boldsymbol{y})$. Then, $f(\boldsymbol{x}^*, \boldsymbol{y}^*) = \Phi(\boldsymbol{x}^*) = \Phi^*$. as noted in (2), $(\boldsymbol{x}^*, \boldsymbol{y}^*)$ is a global minimax point. By (3), it is in fact a stationary point, when g is smooth as well.

(5) (smooth two-sided PŁ: stationary $\Rightarrow$ saddle) Let $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ be a stationary point. By PŁ inequalities, for any $\boldsymbol{x}$ and $\boldsymbol{y}$,

$$0 = \|\nabla_2 g(\boldsymbol{x}^*; \boldsymbol{y}^*)\|^2 \ge 2\mu_2(\max_{\boldsymbol{y}} g(\boldsymbol{x}^*; \boldsymbol{y}) - g(\boldsymbol{x}^*; \boldsymbol{y}^*)) \ge 0,$$

$$0 = \|\nabla_1 g(\boldsymbol{x}^*; \boldsymbol{y}^*)\|^2 \ge 2\mu_1(g(\boldsymbol{x}^*; \boldsymbol{y}^*) - \min_{\boldsymbol{x}} g(\boldsymbol{x}; \boldsymbol{y}^*)) \ge 0.$$

Since $\mu_1, \mu_2 > 0$, these imply $\max_{\boldsymbol{y}} g(\boldsymbol{x}^*; \boldsymbol{y}) = g(\boldsymbol{x}^*; \boldsymbol{y}^*) = \min_{\boldsymbol{x}} g(\boldsymbol{x}; \boldsymbol{y}^*)$. Thus, $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ is a saddle point. Note that (4) and Proposition 10 together proves the existence of a stationary point of g. Therefore, there must exists a saddle point, which is also pointed out by Guo et al. (2020, Lemma 8). This concludes the proof. $\qquad\square$

We remark that, in the proof above, (3) is false for general (nonconvex-nonconcave) functions. Only *local* minimax point can ensure stationarity (Jin et al., 2020). As remarked by Jin et al. (2020) (Figure 2 of their paper), the function $xy - \cos(y)$ has non-stationary global minimax points $(0, \pm\pi)$.

The following two propositions are for showing that general two-sided PŁ function may not have a *differential Stackelberg equilibrium* defined as Li et al. (2022, Definition 3.1).

Proposition 12. *Let g be a μ-PŁ function on $\mathbb{R}^n$. Consider a full-rank matrix $\boldsymbol{M} \in \mathbb{R}^{n \times m}$ ($m \ge n$). Then $g(\boldsymbol{M}\boldsymbol{y})$ is also a PŁ function in $\boldsymbol{y} \in \mathbb{R}^m$.*

Proof. By the μ-PŁ condition of g, for any $\boldsymbol{x} \in \mathbb{R}^n$,

$$\|\nabla g(\boldsymbol{x})\|^2 \ge 2\mu[g(\boldsymbol{x}) - g^*] \quad \text{where} \quad g^* := \min_{\boldsymbol{x}' \in \mathbb{R}^n} g(\boldsymbol{x}').$$

Note that we have $\min_{\boldsymbol{y}\in\mathbb{R}^m} g(\boldsymbol{My}) = \min_{\boldsymbol{x}'\in\mathbb{R}^n} g(\boldsymbol{x}') = g^*$. Let θ be the smallest singular value of $\boldsymbol{M}$. Since $\boldsymbol{MM}^\top$ is an invertible symmetric $(m \times m)$ matrix, θ must be nonzero. Now let $h(\boldsymbol{y}) = g(\boldsymbol{My})$. Thus,

$$\|\nabla h(\boldsymbol{y})\|^2 = \left\|\boldsymbol{M}^\top \nabla g(\boldsymbol{My})\right\|^2$$
$$\geq \theta^2 \|\nabla g(\boldsymbol{My})\|^2 \geq 2\mu\theta^2 \left[g(\boldsymbol{My}) - g^*\right] = 2\mu\theta^2 \left(h(\boldsymbol{y}) - \min_{\boldsymbol{y}'\in\mathbb{R}^m} h(\boldsymbol{y})\right).$$

This implies h is $(\mu\theta^2)$-PŁ. $\qquad\square$

Proposition 13. *Consider a twice continuously differentiable strongly-convex-strongly-concave function $h : \mathbb{R}^r \times \mathbb{R}^s \to \mathbb{R}$. That is, for some constants $\mu_1, \mu_2 > 0$, $h(\boldsymbol{x}; \boldsymbol{y})$ is μ_1-strongly-convex in $\boldsymbol{x}$ and $-h(\boldsymbol{x}; \boldsymbol{y})$ is μ_2-strongly-convex in $\boldsymbol{y}$. Let $(\boldsymbol{x}^*; \boldsymbol{y}^*)$ be the unique stationary point of h. Of course, it is a **differential Stackelberg equilibrium** of h. That is, if the hessian matrix $\nabla^2 h(\boldsymbol{x}^*; \boldsymbol{y}^*)$ at that point is written as*

$$\nabla^2 h(\boldsymbol{x}^*; \boldsymbol{y}^*) = \begin{bmatrix} \nabla^2_{1,1} h(\boldsymbol{x}^*; \boldsymbol{y}^*) & \nabla^2_{1,2} h(\boldsymbol{x}^*; \boldsymbol{y}^*) \\ \nabla^2_{2,1} h(\boldsymbol{x}^*; \boldsymbol{y}^*) & \nabla^2_{2,2} h(\boldsymbol{x}^*; \boldsymbol{y}^*) \end{bmatrix} = \begin{bmatrix} \boldsymbol{C} & \boldsymbol{B} \\ \boldsymbol{B}^\top & -\boldsymbol{A} \end{bmatrix},$$

then $\boldsymbol{A}$ and $\boldsymbol{C} - \boldsymbol{B}\boldsymbol{A}^{-1}\boldsymbol{B}^\top$ are both positive definite matrices. Consider a function $g : \mathbb{R}^p \times \mathbb{R}^q \to \mathbb{R}$ defined by $g(\boldsymbol{x}; \boldsymbol{y}) = h(\boldsymbol{Mx}; \boldsymbol{Ny})$ where $\boldsymbol{M} \in \mathbb{R}^{r \times p}$, $\boldsymbol{N} \in \mathbb{R}^{s \times q}$ are full-rank matrices and $p > r$ and $q > s$. Then g is two-sided PŁ. Moreover, each stationary point of g cannot be a Stackelberg equilibrium.

Proof. Because of Proposition 7 and 12, g is clearly a two-sided PŁ function.

If $(\boldsymbol{x}; \boldsymbol{y})$ is a stationary point of g, then it must be an element of an affine set $\{(\boldsymbol{x}; \boldsymbol{y}) \in \mathbb{R}^p \times \mathbb{R}^q : \boldsymbol{Mx} = \boldsymbol{x}^*; \boldsymbol{Ny} = \boldsymbol{y}^*\}$. This is because

$$\nabla g(\boldsymbol{x}; \boldsymbol{y}) = \begin{bmatrix} \nabla_1 g(\boldsymbol{x}; \boldsymbol{y}) \\ \nabla_2 g(\boldsymbol{x}; \boldsymbol{y}) \end{bmatrix} = \begin{bmatrix} \boldsymbol{M}^\top \nabla_1 h(\boldsymbol{Mx}; \boldsymbol{Ny}) \\ \boldsymbol{N}^\top \nabla_2 h(\boldsymbol{Mx}; \boldsymbol{Ny}) \end{bmatrix} = \boldsymbol{0}$$

if and only if $\nabla_1 h(\boldsymbol{Mx}; \boldsymbol{Ny}) = \boldsymbol{0}$ and $\nabla_2 h(\boldsymbol{Mx}; \boldsymbol{Ny}) = \boldsymbol{0}$, being equivalent to $\boldsymbol{Mx} = \boldsymbol{x}^*$ and $\boldsymbol{Ny} = \boldsymbol{y}^*$. Furthermore, the hessian of g at $(\boldsymbol{x}; \boldsymbol{y})$ is

$$\nabla^2 g(\boldsymbol{x}; \boldsymbol{y}) = \begin{bmatrix} \boldsymbol{M}^\top \nabla^2_{1,1} h(\boldsymbol{Mx}; \boldsymbol{Ny}) \boldsymbol{M} & \boldsymbol{M}^\top \nabla^2_{1,2} h(\boldsymbol{Mx}; \boldsymbol{Ny}) \boldsymbol{N} \\ \boldsymbol{N}^\top \nabla^2_{2,1} h(\boldsymbol{Mx}; \boldsymbol{Ny}) \boldsymbol{M} & \boldsymbol{N}^\top \nabla^2_{2,2} h(\boldsymbol{Mx}; \boldsymbol{Ny}) \boldsymbol{N} \end{bmatrix}$$
$$= \begin{bmatrix} \boldsymbol{M}^\top \boldsymbol{C} \boldsymbol{M} & \boldsymbol{M}^\top \boldsymbol{B} \boldsymbol{N} \\ (\boldsymbol{M}^\top \boldsymbol{B} \boldsymbol{N})^\top & -\boldsymbol{N}^\top \boldsymbol{A} \boldsymbol{N} \end{bmatrix}.$$

Since the rank of a $q \times q$ matrix $\boldsymbol{N}^\top \boldsymbol{A} \boldsymbol{N}$ is (at most) $s < q$, it cannot be even invertible. This implies the stationary point $(\boldsymbol{x}; \boldsymbol{y})$ cannot be a Stackelberg equilibrium. $\qquad\square$

B.2 WITHOUT-REPLACEMENT SAMPLING

In this subsection, we provide a useful proposition for analysis of mini-batching approach under without-replacement sampling. We consider the case of mutually disjoint mini-batches in a whole epoch, not only applying without-replacement sampling to each individual mini-batch.

Consider a collection of n vectors $\boldsymbol{v}_1, \ldots, \boldsymbol{v}_n \in \mathbb{R}^d$. Suppose we randomly sample a permutation $\sigma : [n] \to [n]$; *i.e.*, $\sigma \sim \text{Unif}(\mathbb{S}_n)$. Define

$$\boldsymbol{m} = \frac{1}{n} \sum_{i=1}^n \boldsymbol{v}_i \text{ (sample mean)} \quad \text{and} \quad \tau^2 = \frac{1}{n} \sum_{i=1}^n \|\boldsymbol{v}_i - \boldsymbol{m}\|^2 \text{ (sample variance)}.$$

Fix any $b \in [n]$ and let $n = b(q-1) + s$ for some integers $q \geq 1$ and $s \in [b]$. Now, divide the indices $[n]$ into q batches, with exactly b items per batch (except for the last batch, when $s < b$), as

follows:

$$\mathcal{W}_1 = \{\sigma(1), \cdots, \sigma(b)\},$$
$$\mathcal{W}_2 = \{\sigma(b+1), \cdots, \sigma(2b)\},$$
$$\vdots$$
$$\mathcal{W}_{q-1} = \{\sigma(b(q-2)+1), \cdots, \sigma(b(q-1))\},$$
$$\mathcal{W}_q = \{\sigma(b(q-1)+1), \cdots, \sigma(n)\}.$$

For each batch $\mathcal{W}_t$, define

$$\boldsymbol{w}_t = \frac{1}{|\mathcal{W}_t|} \sum_{i \in \mathcal{W}_t} \boldsymbol{v}_i \quad \text{(batch mean)}.$$

For any $k \in [q-1]$, define

$$\boldsymbol{m}_k := \frac{1}{k} \sum_{t=1}^{k} \boldsymbol{w}_t \quad \text{(accumulative average of batch means over } 1 \le t \le k).$$

Of course, we may simply take $\boldsymbol{m}_q = \boldsymbol{m}$ (deterministically) for $k = q$. Then, due to the randomness of σ, we can obtain the mean (vector) and the variance (scalar) of $\boldsymbol{m}_k$ as follows.

Proposition 14 (Without-replacement sampling). *Given the setup above, for any $k < q$ and $n > 1$,*

$$\mathbb{E}[\boldsymbol{m}_k] = \boldsymbol{m} \quad \text{and} \quad \mathbb{E}\left[\|\boldsymbol{m}_k - \boldsymbol{m}\|^2\right] = \frac{(n-bk)}{bk(n-1)}\tau^2.$$

(Of course, if $k = q$ or $n = 1 = q$, $\mathbb{E}[\|\boldsymbol{m}_q - \boldsymbol{m}\|^2] = 0$ since $\boldsymbol{m}_q = \boldsymbol{m}$.)

Remark. As a special case, if $n = bq$ (namely, b divides n and $r = b$), then for any $k \le q$,

$$\mathbb{E}\left[\|\boldsymbol{m}_k - \boldsymbol{m}\|^2\right] = \frac{(q-k)}{k(n-1)}\tau^2.$$

If we further assume $b = s = 1$ and $q = n$, this proposition exactly recovers Lemma 1 of Mishchenko et al. (2020).

Proof of Proposition 14. Since σ is a uniformly randomly sampled permutation, it is easy to obtain that

$$\mathbb{E}[\boldsymbol{v}_{\sigma(i)}] = \mathbb{E}[\boldsymbol{w}_t] = \mathbb{E}[\boldsymbol{m}_k] = \boldsymbol{m},$$

for any $i \in [n]$, $t \in [q]$, and $k \in [q]$.

The covariances between $\boldsymbol{v}_{\sigma(i)}$'s can be deduced from the proof by Mishchenko et al. (2020, Lemma 1) as follows:

$$\mathrm{Cov}(\boldsymbol{v}_{\sigma(i)}, \boldsymbol{v}_{\sigma(j)}) := \mathbb{E}\left[\langle \boldsymbol{v}_{\sigma(i)} - \boldsymbol{m}, \boldsymbol{v}_{\sigma(j)} - \boldsymbol{m}\rangle\right] = \begin{cases} -\frac{\tau^2}{n-1}, & \text{if } i \ne j, \\ \tau^2 & \text{if } i = j. \end{cases}$$

Thus, for each $t \in [q]$, the variance of $\boldsymbol{w}_t$ is obtained as

$$\mathbb{E}\left[\|\boldsymbol{w}_t - \boldsymbol{m}\|^2\right] = \mathbb{E}\left[\left\|\frac{1}{|\mathcal{W}_t|} \sum_{i \in \mathcal{W}_t} (\boldsymbol{v}_i - \boldsymbol{m})\right\|^2\right]$$

$$= \frac{1}{|\mathcal{W}_t|^2}\left\{\sum_{i \in \mathcal{W}_t} \mathbb{E}\left[\|\boldsymbol{v}_i - \boldsymbol{m}\|^2\right] + \sum_{\substack{i,j \in \mathcal{W}_t \\ i \ne j}} \mathrm{Cov}(\boldsymbol{v}_i, \boldsymbol{v}_j)\right\}$$

$$= \frac{1}{|\mathcal{W}_t|^2}\left\{|\mathcal{W}_t|\tau^2 + |\mathcal{W}_t|(|\mathcal{W}_t| - 1)\left(-\frac{\tau^2}{n-1}\right)\right\} = \frac{n - |\mathcal{W}_t|}{|\mathcal{W}_t|(n-1)}\tau^2,$$

which can also be directly deduced by Lemma 1 of Mishchenko et al. (2020). We notice that this does not depends on the size of the batch $\mathcal{W}_t$.

Next, we look at the covariances between distinct $\boldsymbol{w}_t$'s. For a pair of distinct integers $t, u \in [q]$, by the bi-linearity of covariance,

$$
\begin{aligned}
\mathrm{Cov}(\boldsymbol{w}_t, \boldsymbol{w}_u) &= \frac{1}{|\mathcal{W}_t| \cdot |\mathcal{W}_u|} \sum_{(i,j) \in \mathcal{W}_t \times \mathcal{W}_u} \mathrm{Cov}(\boldsymbol{v}_i, \boldsymbol{v}_j) \\
&= \frac{1}{|\mathcal{W}_t| \cdot |\mathcal{W}_u|} \sum_{(i,j) \in \mathcal{W}_t \times \mathcal{W}_u} \left(-\frac{\tau^2}{n-1} \right) = -\frac{\tau^2}{n-1}.
\end{aligned}
$$

The second equality holds because $\mathcal{W}_t$ and $\mathcal{W}_u$ are a disjoint set of integers whenever $t \neq u$.

Now, fix any $k \in [q-1]$. Note that, by our mini-batching strategy, $|\mathcal{W}_t| = b$ for every $t < q$. Therefore, by definition of $\boldsymbol{m}_k$,

$$
\begin{aligned}
\mathbb{E}\left[\|\boldsymbol{m}_k - \boldsymbol{m}\|^2 \right] &= \mathbb{E}\left[\left\| \frac{1}{k} \sum_{t=1}^{k} (\boldsymbol{w}_t - \boldsymbol{m}) \right\|^2 \right] \\
&= \frac{1}{k^2} \left\{ \sum_{t=1}^{k} \mathbb{E}\left[\|\boldsymbol{w}_t - \boldsymbol{m}\|^2 \right] + \sum_{\substack{t,u \in [k] \\ t \neq u}} \mathrm{Cov}(\boldsymbol{w}_t, \boldsymbol{w}_u) \right\} \\
&= \frac{1}{k^2} \left\{ k \cdot \left(\frac{n-b}{b(n-1)} \tau^2 \right) + k(k-1) \cdot \left(-\frac{\tau^2}{n-1} \right) \right\} = \frac{n-bk}{bk(n-1)} \tau^2.
\end{aligned}
$$

$\square$

B.3 BASIC RECURRENCE INEQUALITY

In this subsection, we present a basic result on a recurrence inequality. It serves as a stepping-stone of our convergence bound, particularly at the end of the proof (Appendix C.4).

Proposition 15. *Let $\{a_k\}_{k=1}^{\infty}$ is a sequence of non-negative numbers satisfying the following recurrence inequality:*

$$a_{k+1} \leq (1-b\eta)a_k + c\eta^{m+1},$$

where b, c, and η are non-negative real numbers such that $b\eta \in (0,1)$, and m is a non-negative integer. Then, for any integer $K \geq 1$, we have

$$a_{K+1} \leq (1-b\eta)^K a_1 + c\eta^m / b.$$

Proof. We proceed an induction on $K = 0, 1, 2, \cdots$. Note that

$$a_1 \leq (1-b\eta)^0 a_1 + c\eta^m / b.$$

This shows the case when $K = 0$. On the other hand, if $K \geq 1$, by an inductive assumption,

$$
\begin{aligned}
a_{K+1} &\leq (1-b\eta)a_K + c\eta^{m+1} \\
&\leq (1-b\eta) \cdot \left((1-b\eta)^{K-1} a_1 + c\eta^m / b \right) + c\eta^{m+1} \\
&= (1-b\eta)^K a_1 + c\eta^m / b.
\end{aligned}
$$

$\square$

C Proofs for (mini-batch) simultaneous SGDA with random reshuffling

In this appendix, we provide a convergence analysis for the mini-batch **sim**SGDA-RR (Algorithm 2) on both general nonconvex-PŁ problems and PŁ(Φ)-PŁ problems. The two cases mostly share the same proof strategies; they only diverge at the end of the proofs. The proof is long; we divide the proof into 4 subsections of this appendix. You can find the proof sketch in Section 6 of this paper.

We again remark that we assume that the mini-batch size b divides the number of components n (namely, $q := n/b$ is a positive integer) for simplicity: thus, readers who just want to read proofs for fully stochastic case (*i.e.*, $b = 1$) can substitute every q to n. Also, there is no problem in treating any fraction with a positive numerator and a zero denominator as $+\infty$. Moreover, we simply regard $(q-1)/(n-1) = 1$ when $n = 1$ ($\because q = 1 = n$ in this case).

The proof for the alternating counterpart (altSGDA-RR) can be done with some modifications illustrated in Appendix D.

C.1 Epoch-wise representations and bounding noise terms

We start the proof by aggregating all updates throughout an epoch to obtain an "epoch-wise" update equation. The reason is that the sampled components in each epoch are dependent to each other, so that it is much harder to deal with each iteration individually. At iteration $t \in [n/b] = [q]$ of epoch $k \in [K]$, we use a mini-batch

$$\mathcal{B}_t^k := \{\sigma_k(b(t-1)+1), \ldots, \sigma_k(bt)\}.$$

Denote the averaged (partial) gradients evaluated at a point $\boldsymbol{z} = (\boldsymbol{x}; \boldsymbol{y})$ over the mini-batch as

$$\boldsymbol{g}_t^k(\boldsymbol{z}) := \frac{1}{b}\sum_{i\in\mathcal{B}_t^k} \nabla_1 f_i(\boldsymbol{z}), \quad \boldsymbol{h}_t^k(\boldsymbol{z}) := \frac{1}{b}\sum_{i\in\mathcal{B}_t^k} \nabla_2 f_i(\boldsymbol{z}).$$

By Assumption 1, $\boldsymbol{g}_t^k$ and $\boldsymbol{h}_t^k$ are L-Lipschitz continuous. Computing the average of them over a whole epoch $(\boldsymbol{z}_0^k, \cdots, \boldsymbol{z}_{q-1}^k)$, we define

$$\boldsymbol{g}^k := \frac{1}{q}\sum_{t=1}^{q} \boldsymbol{g}_t^k(\boldsymbol{z}_{t-1}^k), \quad \boldsymbol{h}^k := \frac{1}{q}\sum_{t=1}^{q} \boldsymbol{h}_t^k(\boldsymbol{z}_{t-1}^k).$$

Then, by summing up the updates in the epoch k, we can summarize the epoch as follows.

$$\boldsymbol{x}_0^{k+1} = \boldsymbol{x}_0^k - q\alpha\boldsymbol{g}^k, \quad \boldsymbol{y}_0^{k+1} = \boldsymbol{y}_0^k + q\beta\boldsymbol{h}^k. \tag{simSGDA-RR}$$

We may assume that the intermediate iterates $\boldsymbol{z}_1^k, \ldots, \boldsymbol{z}_q^k$ stay close to the starting iterate $\boldsymbol{z}_0^k$ of an epoch k, which results from, *e.g.*, small step sizes. Then, we can approximate the aggregated epoch of SGDA-RR as a step of simGDA applied to $f = \frac{1}{n}\sum_{i=1}^n f_i$: $\boldsymbol{g}^k \approx \nabla_1 f(\boldsymbol{z}_0^k)$, $\boldsymbol{h}^k \approx \nabla_2 f(\boldsymbol{z}_0^k)$. In other words,

$$\boldsymbol{x}_0^{k+1} \approx \boldsymbol{x}_0^k - q\alpha\nabla_1 f(\boldsymbol{z}_0^k), \quad \boldsymbol{y}_0^{k+1} \approx \boldsymbol{y}_0^k + q\beta\nabla_2 f(\boldsymbol{z}_0^k), \tag{$\approx$simGDA}$$

With Assumptions 1, 3 and 4, we can yield a naive (but complicated) upper bound of the gap $V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k)$, only applying the smoothness of Φ and $-f$, without any assumptions on step sizes.

Lemma 16. *Suppose that Assumptions 1, 3 and 4 hold. Let $\kappa_2 = L/\mu_2$, where μ_2 is PŁ constant of $-f(\boldsymbol{x}; \cdot)$. Then, the mini-batch simSGDA-RR satisfies that*

$$
\begin{aligned}
V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k) \leq & -\left(\frac{\lambda+1}{2}\right)q\alpha\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + (\lambda+1)q\alpha\left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 \\
& + \frac{q\alpha}{2}\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 - \frac{q\beta}{2}\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2 \\
& + \left(\lambda+\frac{1}{2}\right)q\alpha\left\|\boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 + \frac{q\beta}{2}\left\|\boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k)\right\|^2 \\
& - \left[\lambda - \{(\lambda+1)(\kappa_2+1)+1\}Lq\alpha\right]\frac{q\alpha}{2}\left\|\boldsymbol{g}^k\right\|^2 - (1-Lq\beta)\frac{q\beta}{2}\left\|\boldsymbol{h}^k\right\|^2.
\end{aligned}
\tag{6}
$$

Proof. By definition of V_λ, the following equation holds:

$$V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k) = (\lambda + 1) \left[\Phi(\boldsymbol{x}_0^{k+1}) - \Phi(\boldsymbol{x}_0^k) \right] + \left[f(\boldsymbol{z}_0^k) - f(\boldsymbol{z}_0^{k+1}) \right]. \tag{7}$$

First, we seek for an upper bound of $\Phi(\boldsymbol{x}_0^{k+1}) - \Phi(\boldsymbol{x}_0^k)$. By Proposition 9, Φ is $L(\kappa_2 + 1)$-smooth. Hence, we have

$$
\begin{aligned}
\Phi(\boldsymbol{x}_0^{k+1}) - \Phi(\boldsymbol{x}_0^k) &\leq \left\langle \nabla\Phi(\boldsymbol{x}_0^k), \boldsymbol{x}_0^{k+1} - \boldsymbol{x}_0^k \right\rangle + \frac{L(\kappa_2+1)}{2} \left\| \boldsymbol{x}_0^{k+1} - \boldsymbol{x}_0^k \right\|^2 \\
&= -q\alpha \left\langle \nabla\Phi(\boldsymbol{x}_0^k), \boldsymbol{g}^k \right\rangle + \frac{L(\kappa_2+1)}{2} q^2\alpha^2 \left\| \boldsymbol{g}^k \right\|^2 \\
&= -\frac{q\alpha}{2} \left\{ \left\| \nabla\Phi(\boldsymbol{x}_0^k) \right\|^2 + \left\| \boldsymbol{g}^k \right\|^2 - \left\| \nabla\Phi(\boldsymbol{x}_0^k) - \boldsymbol{g}^k \right\|^2 \right\} + \frac{L(\kappa_2+1)}{2} q^2\alpha^2 \left\| \boldsymbol{g}^k \right\|^2 \\
&= -\frac{q\alpha}{2} \left\| \nabla\Phi(\boldsymbol{x}_0^k) \right\|^2 + \frac{q\alpha}{2} \left\| \nabla\Phi(\boldsymbol{x}_0^k) - \boldsymbol{g}^k \right\|^2 - \frac{q\alpha}{2} (1 - L(\kappa_2+1)q\alpha) \left\| \boldsymbol{g}^k \right\|^2 \\
&\leq -\frac{q\alpha}{2} \left\| \nabla\Phi(\boldsymbol{x}_0^k) \right\|^2 + q\alpha \left\| \nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + q\alpha \left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \\
&\quad - \frac{q\alpha}{2} (1 - L(\kappa_2+1)q\alpha) \left\| \boldsymbol{g}^k \right\|^2.
\end{aligned}
\tag{8}
$$

The third line is due to polarization equality[8] and the last inequality applies Young's inequality.[9]

Next, applying Assumption 1, L-smoothness of $-f(\cdot; \cdot)$ yields an upper bound of $f(\boldsymbol{z}_0^k) - f(\boldsymbol{z}_0^{k+1})$.

$$
\begin{aligned}
f(\boldsymbol{z}_0^k) - f(\boldsymbol{z}_0^{k+1}) &\leq -\left\langle \nabla f(\boldsymbol{z}_0^k), \boldsymbol{z}_0^{k+1} - \boldsymbol{z}_0^k \right\rangle + \frac{L}{2} \left\| \boldsymbol{z}_0^{k+1} - \boldsymbol{z}_0^k \right\|^2 \\
&= -\left\langle \nabla_1 f(\boldsymbol{z}_0^k), \boldsymbol{x}_0^{k+1} - \boldsymbol{x}_0^k \right\rangle - \left\langle \nabla_2 f(\boldsymbol{z}_0^k), \boldsymbol{y}_0^{k+1} - \boldsymbol{y}_0^k \right\rangle \\
&\quad + \frac{L}{2} \left\| \boldsymbol{x}_0^{k+1} - \boldsymbol{x}_0^k \right\|^2 + \frac{L}{2} \left\| \boldsymbol{y}_0^{k+1} - \boldsymbol{y}_0^k \right\|^2 \\
&= q\alpha \left\langle \nabla_1 f(\boldsymbol{z}_0^k), \boldsymbol{g}^k \right\rangle - q\beta \left\langle \nabla_2 f(\boldsymbol{z}_0^k), \boldsymbol{h}^k \right\rangle + \frac{L}{2} q^2\alpha^2 \left\| \boldsymbol{g}^k \right\|^2 + \frac{L}{2} q^2\beta^2 \left\| \boldsymbol{h}^k \right\|^2 \\
&= \frac{q\alpha}{2} \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 - \frac{q\alpha}{2} \left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \frac{q\alpha}{2} (1 + Lq\alpha) \left\| \boldsymbol{g}^k \right\|^2 \\
&\quad - \frac{q\beta}{2} \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \frac{q\beta}{2} \left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 - \frac{q\beta}{2} (1 - Lq\beta) \left\| \boldsymbol{h}^k \right\|^2.
\end{aligned}
\tag{9}
$$

The last equality is due to polarization equality. Lastly, substituting (8) and (9) to (7) finishes the proof. $\qquad\square$

We remark that, the last two terms of the inequality (6) can be simply ignored by applying small enough step sizes. However, the terms in the third line of (6) are non-negatives terms related to the "noise" of approximation $\boldsymbol{g}^k \approx \nabla_1 f(\boldsymbol{z}_0^k)$, $\boldsymbol{h}^k \approx \nabla_2 f(\boldsymbol{z}_0^k)$. Hence, it is important to control the noise terms $\left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2$ and $\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2$ to guarantee a fast decrease of $V_\lambda(\boldsymbol{z}_0^k)$.

Lemma 17. *For mini-batch simSGDA-RR, define*

$$G_k := \frac{1}{q} \sum_{t=1}^q \left\| \boldsymbol{z}_{t-1}^k - \boldsymbol{z}_0^k \right\|^2. \tag{10}$$

With Assumption 1, then

$$\left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \leq L^2 G_k \quad and \quad \left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \leq L^2 G_k.$$

As a side remark, $G_k = 0$ when $q = 1$ and, in particular, $n = 1$.

---

[8]For any $\boldsymbol{a}, \boldsymbol{b} \in \mathbb{R}^d$, $2\langle \boldsymbol{a}, \boldsymbol{b} \rangle = \|\boldsymbol{a}\|^2 + \|\boldsymbol{b}\|^2 - \|\boldsymbol{a} - \boldsymbol{b}\|^2$.
[9]For any $\boldsymbol{a}, \boldsymbol{b} \in \mathbb{R}^d$, $\|\boldsymbol{a} + \boldsymbol{b}\|^2 \leq 2\|\boldsymbol{a}\|^2 + 2\|\boldsymbol{b}\|^2$.

Proof. Recall that $\frac{1}{q}\sum_{t=1}^{q} \boldsymbol{g}_t^k(\boldsymbol{z}) = \nabla_1 f(\boldsymbol{z})$ and $\frac{1}{q}\sum_{t=1}^{q} \boldsymbol{h}_t^k(\boldsymbol{z}) = \nabla_2 f(\boldsymbol{z})$. By Lipschitz continuity and Jensen's inequality,[10]

$$
\left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 = \left\| \frac{1}{q}\sum_{t=1}^{q} [\boldsymbol{g}_t^k(\boldsymbol{z}_{t-1}^k) - \boldsymbol{g}_t^k(\boldsymbol{z}_0^k)] \right\|^2
$$

$$
\leq \frac{1}{q}\sum_{t=1}^{q} \left\| \boldsymbol{g}_t^k(\boldsymbol{z}_{t-1}^k) - \boldsymbol{g}_t^k(\boldsymbol{z}_0^k) \right\|^2 \leq \frac{L^2}{q}\sum_{t=1}^{q} \left\| \boldsymbol{z}_{t-1}^k - \boldsymbol{z}_0^k \right\|^2.
$$

Similarly,

$$
\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \leq \frac{L^2}{q}\sum_{t=1}^{q} \left\| \boldsymbol{z}_{t-1}^k - \boldsymbol{z}_0^k \right\|^2.
$$

This concludes the proof. $\qquad\square$

Thanks to the lemma, it suffices to bound the term G_k. One can notice that it also represents how far the intermediate iterates $\boldsymbol{z}_t^k$ are from the pivot $\boldsymbol{z}_0^k$ in average. Before moving on, we define an algorithm-specific symbol denoting a conditional expectation.

Definition 3. *We denote a conditional expectation of a random variable X given all iterates of the first $k-1$ epochs by $\mathbb{E}_k[X] = \mathbb{E}[X | \boldsymbol{z}_0^1, \boldsymbol{z}_1^1, \ldots, \boldsymbol{z}_n^{k-1}]$. In particular, if $k = 1$, it boils down to a conditional expectation given only the initial iterate $\boldsymbol{z}_0^1$.*

We get an upper bound of a (conditional) expectation $\mathbb{E}_k[G_k]$ in the following lemma, which extends a lemma of Nguyen et al. (2021, Lemma 6) to our minimax problems.

Lemma 18. *Suppose that Assumptions 1 and 2 hold. Assume that the permutation σ_k is sampled uniformly at random from $\mathbb{S}_n$. Then, for any step sizes α, β satisfying $\alpha^2 + \beta^2 \leq \frac{1}{3q(q-1)L^2}$, the iterates $\{\boldsymbol{z}_t^k\}_{t=0}^{q-1}$ of the k-th epoch of mini-batch simSGDA-RR satisfies (for $n > 1$)*

$$
\mathbb{E}_k G_k \leq 2\left(q^2 + \frac{q(q-1)}{n-1}A \right)\left(\alpha^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \beta^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right) + \frac{2q(q-1)}{n-1}(\alpha^2 + \beta^2)B.
$$

Proof. Note that $G_k = 0$ when $q = 1$ by its definition. From now, we may assume $q > 1$ and $n > 1$ in this proof. By summing the first $t \in [q-1]$ updates of the k-th epoch of mini-batch simSGDA-RR, we have

$$
\boldsymbol{x}_t^k = \boldsymbol{x}_0^k - t\alpha\left(\frac{1}{t}\sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_{j-1}^k) \right), \quad \boldsymbol{y}_t^k = \boldsymbol{y}_0^k + t\beta\left(\frac{1}{t}\sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_{j-1}^k) \right).
$$

Then we can bound the following squared distance.

$$
\left\| \boldsymbol{x}_t^k - \boldsymbol{x}_0^k \right\|^2 = \alpha^2 t^2 \left\| \frac{1}{t}\sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_{j-1}^k) \right\|^2
$$

$$
\leq 3\alpha^2 t^2 \left[\left\| \frac{1}{t}\sum_{j=1}^{t} [\boldsymbol{g}_j^k(\boldsymbol{z}_{j-1}^k) - \boldsymbol{g}_j^k(\boldsymbol{z}_0^k)] \right\|^2 + \left\| \frac{1}{t}\sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \right]
$$

$$
\leq 3\alpha^2 t \sum_{j=1}^{t} \left\| \boldsymbol{g}_j^k(\boldsymbol{z}_{j-1}^k) - \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) \right\|^2 + 3\alpha^2 t^2 \left[\left\| \frac{1}{t}\sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \right]
$$

$$
\leq 3\alpha^2 L^2 t \cdot \sum_{j=1}^{t} \left\| \boldsymbol{z}_{j-1}^k - \boldsymbol{z}_0^k \right\|^2 + 3\alpha^2 t^2 \left[\left\| \frac{1}{t}\sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \right]
$$

---

[10]For any n vectors $a_1, \cdots, a_n$, $\left\| \frac{1}{n}\sum_{j=1}^{n} a_j \right\|^2 \leq \frac{1}{n}\sum_{j=1}^{n} \|a_j\|^2$.

$$\leq 3\alpha^2 L^2 t \cdot q G_k + 3\alpha^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \right]. \tag{11}$$

The second and third lines are due to Jensen's inequality. The fourth line is due to L-Lipschitz continuity of $\boldsymbol{g}_j^k$. Likewise,

$$\left\| \boldsymbol{y}_t^k - \boldsymbol{y}_0^k \right\|^2 \leq 3\beta^2 L^2 t \cdot q G_k + 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]. \tag{12}$$

Summing up (11) and (12),

$$\left\| \boldsymbol{z}_t^k - \boldsymbol{z}_0^k \right\|^2 = \left\| \boldsymbol{x}_t^k - \boldsymbol{x}_0^k \right\|^2 + \left\| \boldsymbol{y}_t^k - \boldsymbol{y}_0^k \right\|^2$$

$$\leq 3(\alpha^2 + \beta^2)L^2 t q G_k + 3\alpha^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \right]$$

$$+ 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]. \tag{13}$$

Taking (conditional) expectation $\mathbb{E}_k$ (given $\boldsymbol{z}_0^k$) to inequality (13),

$$\mathbb{E}_k \left\| \boldsymbol{z}_t^k - \boldsymbol{z}_0^k \right\|^2$$

$$\overset{(13)}{\leq} 3(\alpha^2 + \beta^2)L^2 t q \cdot (\mathbb{E}_k[G_k]) + 3\alpha^2 t^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + 3\beta^2 t^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2$$

$$+ 3\alpha^2 t^2 \mathbb{E}_k \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + 3\beta^2 t^2 \mathbb{E}_k \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2. \tag{14}$$

Here we take advantage of the without-replacement sampling. Putting $\nabla_s f_i(\boldsymbol{z}_0^k) \mapsto \boldsymbol{v}_i$ ($s \in \{1, 2\}$), one can realize a correspondence between the quantities that arise from our algorithm and the symbols in Appendix B.2: for $s = 1$ ($\nabla_1 f_i(\boldsymbol{z}_0^k) \mapsto \boldsymbol{v}_i$),

$$\boldsymbol{m} = \nabla_1 f(\boldsymbol{z}_0^k), \quad \tau^2 \leq A \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + B, \quad \boldsymbol{w}_t = \boldsymbol{g}_t^k(\boldsymbol{z}_0^k), \quad \boldsymbol{m}_t = \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k),$$

and for $s = 2$ ($\nabla_2 f_i(\boldsymbol{z}_0^k) \mapsto \boldsymbol{v}_i$),

$$\boldsymbol{m} = \nabla_2 f(\boldsymbol{z}_0^k), \quad \tau^2 \leq A \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + B, \quad \boldsymbol{w}_t = \boldsymbol{h}_t^k(\boldsymbol{z}_0^k), \quad \boldsymbol{m}_t = \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k).$$

The upper bounds of τ^2's come from Assumption 2. Then by Proposition 14, for any $t \leq q$,

$$t^2 \mathbb{E}_k \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{g}_j^k(\boldsymbol{z}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \leq \frac{t(q-t)}{n-1} \left(A \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + B \right),$$

$$t^2 \mathbb{E}_k \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \leq \frac{t(q-t)}{n-1} \left(A \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + B \right).$$

Putting these to the inequality (14),

$$\mathbb{E}_k \left\| \boldsymbol{z}_t^k - \boldsymbol{z}_0^k \right\|^2 \leq 3(\alpha^2 + \beta^2) \left[L^2 t q \mathbb{E}_k[G_k] + \frac{t(q-t)}{n-1} B \right]$$

$$+ 3 \left(\alpha^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \beta^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right) \left[t^2 + \frac{t(q-t)}{n-1} A \right].$$

Taking an average of the inequality above over $0 \leq t \leq q - 1$,

$$
\begin{aligned}
\mathbb{E}_k G_k &= \frac{1}{q} \sum_{t=0}^{q-1} \mathbb{E}_k \left\| \boldsymbol{z}_t^k - \boldsymbol{z}_0^k \right\|^2 \\
&\leq \frac{3q(q-1)}{2}(\alpha^2 + \beta^2) L^2 \mathbb{E}_k G_k + (\alpha^2 + \beta^2) \frac{q^2 - 1}{2(n-1)} B \\
&\quad + \left(\alpha^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \beta^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right) \left(\frac{(q-1)(2q-1)}{2} + \frac{q^2 - 1}{2(n-1)} A \right), \quad (15)
\end{aligned}
$$

where we used the facts $\sum_{t=0}^{q-1} t = \frac{q(q-1)}{2}$, $\frac{1}{q} \sum_{t=0}^{q-1} t^2 = \frac{(q-1)(2q-1)}{6}$, and $\frac{1}{q} \sum_{t=0}^{q-1} \frac{t(q-t)}{n-1} = \frac{q^2-1}{6(n-1)}$. Since we assumed that $\alpha^2 + \beta^2 \leq \frac{1}{3q(q-1)L^2}$, we have $1 \leq 2\left(1 - \frac{3q(q-1)L^2}{2}(\alpha^2 + \beta^2)\right)$. Using this,

$$
\begin{aligned}
\mathbb{E}_k G_k &\leq 2\left(1 - \frac{3q(q-1)L^2}{2}(\alpha^2 + \beta^2)\right) \mathbb{E}_k G_k \\
&\overset{(15)}{\leq} \left((q-1)(2q-1) + \frac{q^2-1}{(n-1)} A\right) \left(\alpha^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \beta^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2\right) + \frac{q^2-1}{n-1}(\alpha^2 + \beta^2) B \\
&\leq 2\left(q^2 + \frac{q(q-1)}{n-1} A\right) \left(\alpha^2 \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 + \beta^2 \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2\right) + \frac{2q(q-1)}{n-1}(\alpha^2 + \beta^2) B,
\end{aligned}
$$

where the last inequality used $(q-1)(2q-1) \leq 2q^2$ and $q + 1 \leq 2q$ for $q \geq 1$. $\qquad \square$

C.2 RECURRENCE INEQUALITIES FOR GENERAL SMOOTH NONCONVEX-PŁ OBJECTIVE

Subsequently, we obtain recurrence inequalities about (expected) potential function $\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^k)]$ for nonconvex-PŁ problem. Since PŁ(Φ)-PŁ problem is a subclass of nonconvex-PŁ problem, the recurrence relations can serve as stepping-stones of our convergence rates.

We introduce some assumptions on *small* step sizes which enable us to get rid of a few troublesome terms from our bound. On top of that, combining the PŁ condition (Assumption 4) with Lemmas 16, 17, and 18, we eventually obtain a much more concise bound on the expected per-epoch change of V_λ. This simple recurrence inequality becomes the key to proving our convergence bounds.

Lemma 19. *Suppose that Assumptions 1, 2, 3, and 4 hold. Assume that the step sizes α and β satisfy*

$$
\alpha \leq \frac{\lambda}{\{(\lambda+1)(\kappa_2+1)+1\}qL}, \quad \beta \leq \frac{1}{qL}, \quad \alpha^2 + \beta^2 \leq \frac{1}{3q(q-1)L^2}, \quad (16)
$$

and the condition

$$
C_0 := q\beta - 2L^2 q \left(q^2 + \frac{q(q-1)}{n-1} A\right) \left((2\lambda+1)\alpha + \beta\right)\beta^2 \geq 0
$$

as well. Then, the iterates of mini-batch simSGDA-RR satisfy

$$
\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^{k+1})] - V_\lambda(\boldsymbol{z}_0^k) \leq -C_1 \left\| \nabla\Phi(\boldsymbol{x}_0^k) \right\|^2 - C_2 \left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right] + C_3
$$

where

$$
C_1 = \left(\frac{\lambda-1}{2}\right) q\alpha - 2L^2 q \left(q^2 + \frac{q(q-1)}{n-1} A\right) \left((2\lambda+1)\alpha + \beta\right)\alpha^2,
$$

$$
\begin{aligned}
C_2 &= \mu_2 C_0 - 2(\lambda+2)L\kappa_2 q\alpha - 4L^3\kappa_2 q \left(q^2 + \frac{q(q-1)}{n-1} A\right) \left((2\lambda+1)\alpha + \beta\right)\alpha^2 \\
&= \mu_2 q\beta - 2(\lambda+2)L\kappa_2 q\alpha - 2L^2\mu_2 q \left(q^2 + \frac{q(q-1)}{n-1} A\right) \left((2\lambda+1)\alpha + \beta\right) \left(2\kappa_2^2\alpha^2 + \beta^2\right),
\end{aligned}
$$

$$
C_3 = \left(\frac{L^2 q^2 (q-1)}{n-1}\right) \left((2\lambda+1)\alpha + \beta\right)(\alpha^2 + \beta^2) B.
$$

Proof. The first two inequalities of (16) eliminate the last two terms on the right-hand side of the inequality in Lemma 16. In addition, applying Lemma 17 to Lemma 16 as well, we have

$$V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k) \leq -\left(\frac{\lambda+1}{2}\right) q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + (\lambda+1)q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$

$$+ \frac{q\alpha}{2}\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 - \frac{q\beta}{2}\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2 + \frac{(2\lambda+1)\alpha+\beta}{2}qL^2 G_k. \quad (17)$$

If we take the conditional expectation $\mathbb{E}_k$ and apply Lemma 18 (which requires the third inequality of (16) to hold) to (17)

$$\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^{k+1})] - V_\lambda(\boldsymbol{z}_0^k)$$

$$\leq -\left(\frac{\lambda+1}{2}\right) q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + (\lambda+1)q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$

$$+ \frac{1}{2}\left[q\alpha + 2L^2 q \left(q^2 + \frac{q(q-1)}{n-1}A\right)((2\lambda+1)\alpha+\beta)\alpha^2\right]\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$

$$- \frac{1}{2}\underbrace{\left[q\beta - 2L^2 q \left(q^2 + \frac{q(q-1)}{n-1}A\right)((2\lambda+1)\alpha+\beta)\beta^2\right]}_{C_0}\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2$$

$$+ \underbrace{\left(\frac{L^2 q^2(q-1)}{n-1}\right)((2\lambda+1)\alpha+\beta)(\alpha^2+\beta^2)B}_{C_3}. \quad (18)$$

It is now left to bound terms in (18) using the tools developed so far. First, recall that $\Phi(\boldsymbol{x}) := \max_{\boldsymbol{y}'\in\mathcal{Y}} f(\boldsymbol{x};\boldsymbol{y}')$. Since $-f(\boldsymbol{x};\boldsymbol{y})$ is μ_2-PŁ in $\boldsymbol{y}$, we have

$$-\left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2 \leq -2\mu_2(\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)). \quad (19)$$

Given any $\boldsymbol{x}$, $\nabla\Phi(\boldsymbol{x}) = \nabla_1 f(\boldsymbol{x};\boldsymbol{y}^*(\boldsymbol{x}))$ for any $\boldsymbol{y}^*(\boldsymbol{x}) \in \arg\max_{\boldsymbol{y}'\in\mathcal{Y}} f(\boldsymbol{x};\boldsymbol{y}')$ by Proposition 9. Besides, $-f(\boldsymbol{x};\cdot)$ satisfies QG condition with constant μ_2 by Proposition 7. Thus, by choosing $\boldsymbol{y}^*(\boldsymbol{x}_0^k)$ to be the projection of $\boldsymbol{y}_0^k$ onto $\arg\max_{\boldsymbol{y}'\in\mathcal{Y}} f(\boldsymbol{x}_0^k;\boldsymbol{y}')$,

$$\left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 \leq L^2 \left\|\boldsymbol{y}^*(\boldsymbol{x}_0^k) - \boldsymbol{y}_0^k\right\|^2 \leq 2L\kappa_2\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right]. \quad (20)$$

Here, the first inequality applies L-Lipschitz continuity of $\nabla_1 f(\boldsymbol{x}_0^k;\cdot)$, implied by Assumption 1. On top of that, applying the Young's inequality to the term $\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$,

$$\left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 \leq 2\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + 2\left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$

$$\overset{(20)}{\leq} 2\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + 4L\kappa_2\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right] \quad (21)$$

By applying inequalities (19), (20), and (21) to the bound (18), we conclude the proof. □

In Lemma 19, we saw that if step sizes are chosen to satisfy certain conditions, then we can simplify the per-epoch progress a great deal. It is now left to choose appropriate step sizes and parameters (*e.g.*, λ) so as to make sure not only that α and β meet the *small* step size conditions (16) but also that the constants C_0, C_1, C_2, and C_3 are positive.

Lemma 20. *Suppose that Assumptions 1, 2, 3 and 4 hold. Let $\lambda = 4$ and assume that*

$$0 < \beta \leq \frac{1}{6L\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \quad \alpha = \frac{\beta}{r}, \quad \text{where } r \geq 14\kappa_2^2.$$

Then these satisfy all the inequalities (16) and the terms defined in Lemma 19 satisfy

$$C_0 > 0, \quad C_1 > q\alpha, \quad C_2 > L\kappa_2 q\alpha/2, \quad C_3 \geq 0.$$

Consequently, due to the recurrence inequality in Lemma 19, mini-batch simSGDA-RR satisfies, for some numerical constant $c > 0$,

$$\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^{k+1})] - V_\lambda(\boldsymbol{z}_0^k) \leq -q\alpha\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 - (L\kappa_2 q\alpha/2)\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right]$$

$$+ (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3. \quad (\star)$$

Please note that we mark the recurrence inequality above with a special symbol ($\star$) because this inequality is the exact point where the proofs of Theorems 4 and 5 start to deviate.

Proof. Regardless of $A \geq 0$, we have

$$\beta \leq \frac{1}{6Lq} \quad \text{and} \quad \alpha \leq \frac{1}{6Lqr} \leq \frac{1}{84L\kappa_2^2 q}. \tag{22}$$

This is enough to guarantee that the inequalities (16) hold with $\lambda = 4$. Since $C_0 > C_2/\mu_2$, it is enough to show $C_2 > 0$ to prove that $C_0 > 0$. Applying $\lambda = 4$, $\kappa_2 \geq 1$, and $\beta/\alpha = r \geq 14\kappa_2^2$,

$$\begin{aligned}
\frac{C_1}{q\alpha} &= \frac{3}{2} - 2L^2 \left(q^2 + \frac{q(q-1)}{n-1}A \right) (9+r)\alpha^2 \\
&\geq \frac{3}{2} - \frac{2}{6^2} \cdot \frac{9+r}{r^2} \geq \frac{3}{2} - \frac{2 \cdot 23}{6^2 \cdot 14^2} > 1, \\
\frac{C_2}{\mu_2 q\beta} &= 1 - \frac{12\kappa_2^2}{r} - 2L^2 \left(q^2 + \frac{q(q-1)}{n-1}A \right) \left(\frac{9}{r} + 1 \right) \left(\frac{2\kappa_2^2}{r^2} + 1 \right) \beta^2 \\
&\geq 1 - \frac{12}{14} - \frac{2}{6^2} \left(\frac{9}{14\kappa_2^2} + 1 \right) \left(\frac{2}{14^2\kappa_2^2} + 1 \right) \geq \frac{2}{14} - \frac{2 \cdot 23 \cdot 198}{6^2 \cdot 14^3} > \frac{1}{2 \cdot 14}.
\end{aligned}$$

Thus, $C_1 > q\alpha$ and

$$C_2 > \frac{\mu_2 q\beta}{2 \cdot 14} = \frac{\mu_2 qr\alpha}{2 \cdot 14} \geq L\kappa_2 q\alpha/2.$$

Then we conclude the proof by bounding the term C_3. We can already check from the definition that $C_3 \geq 0$. We can upper-bound C_3 by

$$C_3 = \left(\frac{L^2 q^2(q-1)}{n-1} \right) (9+r)(1+r^2)B\alpha^3 \leq (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1} \right) B\alpha^3,$$

for some numerical constant $c > 0$. $\qquad \square$

C.3 Convergence rates for smooth nonconvex-PŁ problem

In this subsection, we show the convergence bound of general smooth nonconvex-PŁ problems in terms of $\min_{k \in [K]} \mathbb{E}\left[\left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \right]$. From the inequality ($\star$) in Lemma 20, we can simply ignore the second term

$$-(L\kappa_2 q\alpha/2)\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k) \right] \leq 0$$

of the right-hand side because $\Phi(\boldsymbol{x}) \geq f(\boldsymbol{x}; \boldsymbol{y})$ for any $(\boldsymbol{x}; \boldsymbol{y})$. In other words, we may deal with the inequality

$$\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^{k+1})] - V_\lambda(\boldsymbol{z}_0^k) \leq -q\alpha \left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1} \right) B\alpha^3. \tag{nc-PŁ}$$

Plugging $q = n/b$, we eventually show the convergence rate (Theorem 4). (Recall that b is the size of mini-batches.)

Theorem 21 (Equivalent to Theorem 4, for simSGDA-RR). *Suppose that f satisfies Assumptions 1, 2, 3, and 4 are satisfied. Let $\lambda = 4$. Choose the step sizes α and β by $\alpha = \beta/r$ for some $r \geq 14\kappa_2^2$ and*

$$\beta = \min \left\{ \frac{1}{6L\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \frac{1}{c} \left(\frac{V_\lambda(\boldsymbol{z}_0^1)}{L^2 q^2 (\frac{q-1}{n-1})BK} \right)^{\frac{1}{3}} \right\},$$

for some numerical constant $c > 0$. Then, mini-batch simSGDA-RR satisfies

$$\begin{aligned}
\min_{k \in [K]} \mathbb{E}\left[\left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \right] &\leq \frac{1}{K} \sum_{k=1}^K \mathbb{E}\left[\left\| \nabla \Phi(\boldsymbol{x}_0^k) \right\|^2 \right] \\
&< \frac{6rLV_\lambda(\boldsymbol{z}_0^1)}{K}\sqrt{1 + \left(\frac{q-1}{n-1} \right)\frac{A}{q}} + cr \left(\frac{L^2 B V_\lambda(\boldsymbol{z}_0^1)^2}{qK^2} \cdot \frac{q-1}{n-1} \right)^{1/3}.
\end{aligned}$$

Proof. To replace the conditional expectations with unconditional expections, we take expaction to the both side of the inequality (nc-PŁ):

$$\mathbb{E}[V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k)] \leq -q\alpha\mathbb{E}\left[\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2\right] + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3.$$

Rearranging the terms and taking a sum from $k = 1$ to $k = K$, we have

$$q\alpha\sum_{k=1}^K \mathbb{E}\left[\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2\right] \leq \mathbb{E}[V_\lambda(\boldsymbol{z}_0^1) - V_\lambda(\boldsymbol{z}_0^{K+1})] + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3 K.$$

Dividing both sides by $qK\alpha$, we get the following. Note that V_λ is non-negative.

$$\frac{1}{K}\sum_{k=1}^K \mathbb{E}\left[\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2\right] \leq \frac{V_\lambda(\boldsymbol{z}_0^1)}{qK\alpha} + (cr)^3 L^2 \left(\frac{q(q-1)}{n-1}\right) B\alpha^2$$

Since our choice of step sizes implies that

$$\alpha = \min\left\{\frac{1}{6rL\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \frac{1}{cr}\left(\frac{V_\lambda(\boldsymbol{z}_0^1)}{L^2 Bq^2(\frac{q-1}{n-1})K}\right)^{\frac{1}{3}}\right\}.$$

Using the inequality $\max\{a, b\} \leq a + b$ (for $a, b \geq 0$), we eventually have

$$\frac{1}{K}\sum_{k=1}^K \mathbb{E}\left[\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2\right] < \frac{6rLV_\lambda(\boldsymbol{z}_0^1)}{K}\sqrt{1 + \left(\frac{q-1}{n-1}\right)\frac{A}{q}} + cr\left(\frac{L^2 B\,V_\lambda(\boldsymbol{z}_0^1)^2}{qK^2}\cdot\frac{q-1}{n-1}\right)^{1/3}.$$

$\square$

C.4 Convergence rates for smooth PŁ(Φ)-PŁ problem

In this subsection, we prove the convergence bound of PŁ(Φ)-PŁ problems in terms of $\mathbb{E}\left[V_\lambda(\boldsymbol{z}_0^{K+1})\right]$.

Unlike the previous subsection, we additionally utilize Assumption 5 stating that $f(\boldsymbol{x}; \boldsymbol{y})$ satisfies primal PŁ condition, namely, the primal function $\Phi(\boldsymbol{x}) = \max_{\boldsymbol{y}'} f(\boldsymbol{x}; \boldsymbol{y}')$ is a μ_1-PŁ function. With this assumption, we yield another recurrence inequality from the inequlity ($\star$). We note that it uses the μ_1-PŁ condition for Φ ($\because$ Proposition 10) but not necessarily for $f(\cdot; \boldsymbol{y})$.

Lemma 22. *Suppose that f satisfies Assumptions 1, 2, 3, 4, and 5. Then, with the same choice of $\lambda = 4$ and the same condition of the step sizes α and β as in Lemma 20, the mini-batch simSGDA-RR satisfies that, for some numerical constant $c > 0$,*

$$\mathbb{E}_k[V_\lambda(\boldsymbol{z}_0^{k+1})] \leq (1 - \mu_1 q\alpha/2)V_\lambda(\boldsymbol{z}_0^k) + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3. \tag{2PŁ}$$

Proof. Since the primal function Φ is a μ_1-PŁ function,

$$-\left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 \leq -2\mu_1\left[\Phi(\boldsymbol{x}_0^k) - \Phi^*\right].$$

Also, since $\mu_1 \leq L$ and $\kappa_2 \geq 1$, we know that $-L\kappa_2 \leq -\mu_1$. Applying these to the inequality ($\star$), we have

$$\mathbb{E}_k\left[V_\lambda(\boldsymbol{z}_0^{k+1})\right] - V_\lambda(\boldsymbol{z}_0^k)$$

$$\leq -(2\mu_1 q\alpha/\lambda)\cdot\lambda\left[\Phi(\boldsymbol{x}_0^k) - \Phi^*\right] - (\mu_1 q\alpha/2)\left[\Phi(\boldsymbol{x}_0^k) - f(\boldsymbol{z}_0^k)\right] + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3$$

$$= -(\mu_1 q\alpha/2)\cdot V_\lambda(\boldsymbol{z}_0^k) + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1}\right) B\alpha^3,$$

since $\lambda = 4$. By re-arranging the terms, we conclude the proof. $\square$

Of course, the multiplier $1 - \mu_1 q\alpha/2$ have a value between 0 and 1. To see why, note that from Equation (22),

$$0 < \mu_1 q\alpha/2 \le \mu_1 q \cdot \frac{1}{2 \cdot 84 L \kappa_2^2 q} = \frac{1}{168 \kappa_1 \kappa_2^2} < 1.$$

Theorem 23 (Equivalent to Theorem 5, for simSGDA-RR). *Assume that f satisfies Assumptions 1, 2, 3, 4, and 5. Let $\lambda = 4$. Choose the step sizes by $\alpha = \beta/r$ for some $r \ge 14\kappa_2^2$ and*

$$\beta = \min \left\{ \frac{1}{6L\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \ \frac{2r}{\mu_1 qK} \max \left\{ 1, \ \log \left(\frac{V_\lambda(z_0^1)\mu_1 q K^2}{8(cr)^3 \kappa_1^2 \left(\frac{q-1}{n-1}\right) B} \right) \right\} \right\},$$

for some numerical constant $c > 0$. Then, mini-batch simSGDA-RR satisfies

$$\mathbb{E}[V_\lambda(z_n^K)] \le \mathcal{O}\left(V_\lambda(z_0^1) \cdot \exp \left(-\frac{K}{12\kappa_1 r \sqrt{1 + \left(\frac{q-1}{n-1}\right)\frac{A}{q}}} \right) \right) + \tilde{\mathcal{O}}\left(\frac{\kappa_1^2 r^3 B}{\mu_1 q K^2} \right) \cdot \frac{q-1}{n-1}.$$

Proof. To replace the conditional expectations with unconditional expactions, we take expaction to the both side of the inequality (2PŁ):

$$\mathbb{E}\left[V_\lambda(z_0^{k+1})\right] \le (1 - \mu_1 q\alpha/2)\mathbb{E}\left[V_\lambda(z_0^k)\right] + (cr)^3 L^2 \left(\frac{q^2(q-1)}{n-1} \right) B\alpha^3.$$

Unrolling the recurrence inequality (Proposition 15) and using the facts $\beta = 14\kappa_2^2\alpha$, we have

$$\mathbb{E}[V_\lambda(z_n^K)] \le (1 - \mu_1 q\alpha/2)^K V_\lambda(z_0^1) + \frac{2 \cdot (cr)^3 L^2}{\mu_1 q\alpha} \left(\frac{q^2(q-1)}{n-1} \right) B\alpha^3$$

$$\le \exp(-\mu_1 qK\alpha/2)V_\lambda(z_0^1) + 2(cr)^3 \mu_1 \kappa_1^2 \left(\frac{q(q-1)}{n-1} \right) B\alpha^2. \tag{23}$$

Note that, in the inequality above, the second term of the right hand side becomes zero when $q = 1$. In that case, we can prove exponential decay of $\mathbb{E}[V_\lambda(z_0^k)]$. Thus, we simply assume $q > 1$ hereafter.

Case 1: If K is as large as

$$K > \frac{\kappa_1 r^{3/2}}{\sqrt{\mu_1}} \cdot \sqrt{\frac{8c^3 eB}{V_\lambda(z_0^1)q} \left(\frac{q-1}{n-1} \right)}, \quad (e = \exp(1))$$

we have a step size α as

$$\alpha = \min \left\{ \frac{1}{6Lr\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \ \frac{2}{\mu_1 qK} \log (\clubsuit) \right\}, \quad \text{where } \clubsuit = \frac{V_\lambda(z_0^1)\mu_1 q K^2}{8(cr)^3 \kappa_1^2 \kappa_2^6 \left(\frac{q-1}{n-1}\right) B}.$$

Due to the lower bound of epoch size K, the fraction $\clubsuit$ inside the log factor is indeed greater than $e > 1$, which guarantees the step size positive. Putting this to the inequality (23) and using the fact that $\max\{a, b\} \le a + b$ (for $a, b \ge 0$), we eventually have

$$\mathbb{E}\left[V_\lambda(z_n^K)\right]$$

$$\le V_\lambda(z_0^1) \cdot \exp \left(-\frac{K}{12\kappa_1 r \sqrt{1 + \left(\frac{q-1}{n-1}\right)\frac{A}{q}}} \right) + \frac{2 \cdot 8(cr)^3 \kappa_1^2 B}{\mu_1 q K^2} \left(\frac{q-1}{n-1} \right) \left[1 + \log^2 (\clubsuit)\right]$$

$$= V_\lambda(z_0^1) \cdot \exp \left(-\frac{K}{12\kappa_1 r \sqrt{1 + \left(\frac{q-1}{n-1}\right)\frac{A}{q}}} \right) + \tilde{\mathcal{O}}\left(\frac{\kappa_1^2 r^3 B}{\mu_1 q K^2} \right) \cdot \frac{q-1}{n-1}.$$

Case 2: Otherwise, the log factor might have negative value when K is too small. However, in this case we have

$$V_\lambda(z_0^1) \le \frac{8(cr)^3 e\kappa_1^2 B}{\mu_1 qK^2} \cdot \frac{q-1}{n-1}; \quad \alpha = \min\left\{ \frac{1}{84L\kappa_2^2\sqrt{q^2 + \frac{q(q-1)}{n-1}A}}, \frac{2}{\mu_1 qK} \right\}.$$

Putting these to the inequality (23), we have

$$\mathbb{E}\left[V_\lambda(z_n^K)\right] \le \frac{8(cr)^3 e\kappa_1^2 B}{\mu_1 qK^2}\left(\frac{q-1}{n-1}\right)\left[\exp(-\mu_1 qK\alpha/2) + \frac{1}{e}\cdot(\mu_1 qK\alpha/2)^2\right]$$

$$\le \frac{8(cr)^3 e\kappa_1^2 B}{\mu_1 qK^2}\left(\frac{q-1}{n-1}\right) = \mathcal{O}\left(\frac{\kappa_1^2 r^3 B}{\mu_1 qK^2}\right)\cdot\frac{q-1}{n-1}.$$

The inequlity in the last line is due to the fact that $e^{-t} + t^2/e \le 1$ for each $t \in (0,1]$, and that $\mu_1 qK\alpha/2 \in (0,1]$.

In either case, the conclusion does not change:

$$\mathbb{E}\left[V_\lambda(z_n^K)\right] \le \mathcal{O}\left(V_\lambda(z_0^1)\cdot\exp\left(-\frac{K}{12\kappa_1 r\sqrt{1 + \left(\frac{q-1}{n-1}\right)\frac{A}{q}}}\right)\right) + \tilde{\mathcal{O}}\left(\frac{\kappa_1^2 r^3 B}{\mu_1 qK^2}\right)\cdot\frac{q-1}{n-1}.$$

$\square$

D PROOFS FOR (MINI-BATCH) ALTERNATING SGDA-RR: FOCUSING ON CHANGES IN THE PROOF

In this appendix, we prove the same convergence rates for altSGDA-RR as the simultaneous update counterpart. Since most of the steps in the proof is similar to those in Appendix C, we only describe which steps change in the proof.

D.1 EPOCH-WISE REPRESENTATIONS AND BOUNDING NOISE TERMS

To analyze altSGDA-RR, we modify the notation for epoch-wise updates. The only change is that an update $\boldsymbol{y}_t^k \mapsto \boldsymbol{y}_{t+1}^k$ uses $\boldsymbol{x}_{t+1}^k$ instead of $\boldsymbol{x}_t^k$. Hence, the definition of $\boldsymbol{h}^k$ should be modified. Recall that

$$\boldsymbol{g}_t^k(\boldsymbol{z}) := \frac{1}{b} \sum_{i \in \mathcal{B}_t^k} \nabla_1 f_i(\boldsymbol{z}), \quad \boldsymbol{h}_t^k(\boldsymbol{z}) := \frac{1}{b} \sum_{i \in \mathcal{B}_t^k} \nabla_2 f_i(\boldsymbol{z}),$$

where $\mathcal{B}_t^k$ is a mini-batch of size b formed at iteration t of epoch k. Then, at epoch k, by re-definition of $\boldsymbol{h}^k$,

$$\boldsymbol{g}^k := \frac{1}{q} \sum_{t=1}^q \boldsymbol{g}_t^k(\boldsymbol{x}_{t-1}^k; \boldsymbol{y}_{t-1}^k), \quad \boldsymbol{h}^k := \frac{1}{q} \sum_{t=1}^q \boldsymbol{h}_t^k(\boldsymbol{x}_t^k; \boldsymbol{y}_{t-1}^k).$$

$$\boldsymbol{x}_0^{k+1} = \boldsymbol{x}_0^k - q\alpha \boldsymbol{g}^k, \quad \boldsymbol{y}_0^{k+1} = \boldsymbol{y}_0^k + q\beta \boldsymbol{h}^k. \qquad \text{(altSGDA-RR)}$$

We still approximate this epoch-wise update rule to a full-batch simultaneous GDA update ($\approx$simGDA) with step sizes $q\alpha$ and $q\beta$. Again, we control the "noise" terms $\left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2$ and $\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2$ not to be large. Because of the modification of $\boldsymbol{h}^k$, we have a different result for $\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2$ as follows.

Lemma 24. *For mini-batch altSGDA-RR, recall that*

$$G_k := \frac{1}{q} \sum_{t=1}^q \left\| \boldsymbol{z}_{t-1}^k - \boldsymbol{z}_0^k \right\|^2.$$

If we have Assumption 1, then we have

$$\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \le L^2 G_k + L^2 q\alpha^2 \left\| \boldsymbol{g}^k \right\|^2, \quad \text{whereas} \quad \left\| \boldsymbol{g}^k - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 \le L^2 G_k. \quad (24)$$

Proof. Because of L-Lipschitz continuity of $\boldsymbol{h}_t^k(\cdot; \cdot)$,

$$\left\| \boldsymbol{h}^k - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 = \left\| \frac{1}{q} \sum_{t=1}^q \left[\boldsymbol{h}_t^k(\boldsymbol{x}_t^k; \boldsymbol{y}_{t-1}^k) - \boldsymbol{h}_t^k(\boldsymbol{x}_0^k; \boldsymbol{y}_0^k) \right] \right\|^2$$

$$\le \frac{1}{q} \sum_{t=1}^q \left\| \boldsymbol{h}_t^k(\boldsymbol{x}_t^k; \boldsymbol{y}_{t-1}^k) - \boldsymbol{h}_t^k(\boldsymbol{x}_0^k; \boldsymbol{y}_0^k) \right\|^2$$

$$\le \frac{L^2}{q} \sum_{t=1}^q \left\| \boldsymbol{z}_{t-1}^k - \boldsymbol{z}_0^k \right\|^2 + \frac{L^2}{q} \left\| \boldsymbol{x}_q^k - \boldsymbol{x}_0^k \right\|^2 = L^2 G_k + L^2 q\alpha^2 \left\| \boldsymbol{g}^k \right\|^2.$$

The last ineqaulity holds because $\boldsymbol{x}_q^k = \boldsymbol{x}_0^{k+1}$. □

D.2 BOUNDING NOISE TERMS: A BIT DIFFERENT PROOF OF LEMMA 18

We notice that the same result as Lemma 18 holds not only for simultaneous updates but also alternating updates, even though it is not very straightforward. We need to reflect the changes from the previous subsection. That is, we have to be careful when we expand the term $\left\| \boldsymbol{y}_t^k - \boldsymbol{y}_0^k \right\|^2$

($0 \leq t \leq q-1$). Unlike the inequality (11) (in the original proof), we have

$$\left\| \boldsymbol{y}_t^k - \boldsymbol{y}_0^k \right\|^2 = \beta^2 t^2 \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{x}_j^k; \boldsymbol{y}_{j-1}^k) \right\|^2$$

$$\leq 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \left[\boldsymbol{h}_j^k(\boldsymbol{x}_j^k; \boldsymbol{y}_{j-1}^k) - \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) \right] \right\|^2 + \left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]$$

$$\leq 3\beta^2 t \sum_{j=1}^{t} \left\| \boldsymbol{h}_j^k(\boldsymbol{x}_j^k; \boldsymbol{y}_{j-1}^k) - \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) \right\|^2 + 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]$$

$$\leq 3\beta^2 L^2 t \left(\sum_{j=1}^{t} \left\| \boldsymbol{z}_{j-1}^k - \boldsymbol{z}_0^k \right\|^2 + \left\| \boldsymbol{x}_t^k - \boldsymbol{x}_0^k \right\|^2 \right) + 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]$$

$$\leq 3\beta^2 L^2 t \sum_{j=1}^{t} \left\| \boldsymbol{z}_j^k - \boldsymbol{z}_0^k \right\|^2 + 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right]$$

$$\leq 3\beta^2 L^2 t \cdot q G_k + 3\beta^2 t^2 \left[\left\| \frac{1}{t} \sum_{j=1}^{t} \boldsymbol{h}_j^k(\boldsymbol{z}_0^k) - \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 \right].$$

The second and third inequality holds by Jensen's inequality, and the last inequality holds because $t \leq q-1$. The resulting upper bound is identical to the inequality (12). Proving this inequality above suffices to show that the conclusion of Lemma 18 also holds for altSGDA-RR, because we eventually take an average along $0 \leq t \leq q-1$ and the other steps in the proof does not utilize the "order" (either simultaneous or alternating) of updates.

D.3 RECURRENCE INEQUALITIES FOR GENERAL SMOOTH NONCONVEX-PŁ OBJECTIVE

In the proof for simSGDA-RR, we applied Lemma 16, Lemma 18, and the "small-step-size" assumptions (three inequalities in (16)) to deduce Lemma 19. However, due to Lemma 24 that we obtained for altSGDA-RR, we need slightly different assumptions on step sizes rather than (16).

Fortunately, we notice that the Lemma 16 also holds for altSGDA-RR, with a modified version of $\boldsymbol{h}^k$. This is because the proof of the lemma does not utilize step-wise updates, while the discrepancy between simultaneous and alternating updates only appear in the step-wise updates. Thus, we have the same result as Lemma 19.

Lemma 25. *Suppose that Assumptions 1, 2, 3, and 4 hold. Modify the inequalities (16) (from Lemma 19) by*

$$\lambda - \left\{ (\lambda+1)(\kappa_2+1) + 1 \right\} Lq\alpha - L^2 q\alpha\beta \geq 0, \quad \beta \leq \frac{1}{qL}, \quad \alpha^2 + \beta^2 \leq \frac{1}{3q(q-1)L^2}. \quad (25)$$

(In fact, only the first one is different.) Then, the result of Lemma 19 still holds for mini-batch altSGDA-RR.

Proof. We first apply Lemma 24 to the general bound resulted from Lemma 16:

$$V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k)$$

$$\leq - \left(\frac{\lambda+1}{2} \right) q\alpha \left\| \nabla\Phi(\boldsymbol{x}_0^k) \right\|^2 + (\lambda+1)q\alpha \left\| \nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2$$

$$+ \frac{q\alpha}{2} \left\| \nabla_1 f(\boldsymbol{z}_0^k) \right\|^2 - \frac{q\beta}{2} \left\| \nabla_2 f(\boldsymbol{z}_0^k) \right\|^2 + \frac{(2\lambda+1)\alpha + \beta}{2} qL^2 G_k$$

$$- \left[\lambda - \left\{ (\lambda+1)(\kappa_2+1) + 1 \right\} Lq\alpha - L^2 q\alpha\beta \right] \frac{q\alpha}{2} \left\| \boldsymbol{g}^k \right\|^2 - (1 - Lq\beta) \frac{q\beta}{2} \left\| \boldsymbol{h}^k \right\|^2. \quad (26)$$

Hence, the first two inequalities of (25) eliminates the last two terms on the right side of the inequality (26) above:

$$V_\lambda(\boldsymbol{z}_0^{k+1}) - V_\lambda(\boldsymbol{z}_0^k) \leq - \left(\frac{\lambda+1}{2}\right) q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k)\right\|^2 + (\lambda+1)q\alpha \left\|\nabla\Phi(\boldsymbol{x}_0^k) - \nabla_1 f(\boldsymbol{z}_0^k)\right\|^2$$
$$+ \frac{q\alpha}{2} \left\|\nabla_1 f(\boldsymbol{z}_0^k)\right\|^2 - \frac{q\beta}{2} \left\|\nabla_2 f(\boldsymbol{z}_0^k)\right\|^2 + \frac{(2\lambda+1)\alpha + \beta}{2} qL^2 G_k.$$

This is identical to the inequality (17) in the proof of Lemma 19. From this point on, the rest of proof is exactly identical to Lemma 19. □

Lemma 25 establishes that altSGDA-RR also satisfies a concise bound on the expected per-epoch change of V_λ, albeit under a slightly different set of assumptions (25) on step sizes. Using this result, we can prove the convergence rates for altSGDA-RR that are exactly the same as simSGDA-RR.

D.4   SMALL STEP SIZE ASSUMPTIONS

It is left to show an altSGDA-RR counterpart for Lemma 20 which establishes the general recurrence inequality (⋆). In fact, the same choice of step sizes as simSGDA-RR, namely

$$0 < \beta \leq \frac{1}{6L\sqrt{q^2 + \frac{q(q-1)}{n-1}A}} \quad \text{and} \quad \alpha = \frac{\beta}{r} \quad \text{where} \quad r \geq 14\kappa_2^2,$$

actually meets the newly introduced conditions (25). Among three inequalities, the only one that needs to be checked is

$$\lambda - \{(\lambda+1)(\kappa_2+1) + 1\} Lq\alpha - L^2 q\alpha\beta > 0.$$

Note that, regardless of $A \geq 0$,

$$\beta \leq \frac{1}{6Lq} \quad \text{and} \quad \alpha \leq \frac{1}{6Lqr} \leq \frac{1}{84L\kappa_2^2 q}$$

In this case,

$$\lambda - \{(\lambda+1)(\kappa_2+1) + 1\} Lq\alpha - L^2 q\alpha\beta$$
$$\geq 4 - (11\kappa_2 + L\beta)Lq\alpha \geq 4 - \left(11\kappa_2 + \frac{1}{6}\right) \cdot \frac{1}{84\kappa_2^2} > 0.$$

Therefore, there is no need to modify our choices of λ and the step sizes α, β for analysis of altSGDA-RR, and the rest of the proof for simSGDA-RR goes through.

E PROOFS FOR LOWER BOUND OF TWO-TIME-SCALE GDA

In this appendix, we illustrate a comprehensive lower bound for full-batch two-time-scale GDA which is specific to the choice of step size ratio (Theorem 3). The proof of Theorem 3 is at the end of this appendix.

Before we start the proof, we define a class of smooth strongly-convex-strongly concave functions.

Definition 4. *Let $\mathcal{F}(L, \mu_1, \mu_2)$ be the class of functions $f(\boldsymbol{x}; \boldsymbol{y})$ with two arguments $\boldsymbol{x}$ and $\boldsymbol{y}$ of any dimension, which is L-smooth, μ_1-strongly-convex in $\boldsymbol{x}$, and μ_2-strongly-concave in $\boldsymbol{y}$. Let $\kappa_1 = L/\mu_1 \geq 1$ and $\kappa_2 = L/\mu_2 \geq 1$ be condition numbers of the function class. Denote the (unique) saddle (or, global minimax) point by $\boldsymbol{z}^* = (\boldsymbol{x}^*; \boldsymbol{y}^*)$.*

We restate and prove the Theorem 3 for reader's convenience.

Theorem 26 (Restatement of Theorem 3). *Suppose $\kappa_1 \geq c$ and $\kappa_2 \geq c$ for some constant $c > 1$. Then, for each step size ratio $r > 0$, there exists a function $f \in \mathcal{F}(L, \mu_1, \mu_2)$ for which two-time-scale GDA with any step sizes α and β of ratio $r = \beta/\alpha$ requires*

$$
K = \begin{cases} \Omega\left(\kappa_1 r \log(1/\varepsilon)\right), & \text{if } r \geq \kappa_2/c, \\ \Omega\left(\kappa_1 \kappa_2 \log(1/\varepsilon)\right), & \text{if } c/\kappa_1 \leq r \leq \kappa_2/c, \\ \Omega((\kappa_2/r) \log(1/\varepsilon)), & \text{if } 0 < r \leq c/\kappa_1 \end{cases}
$$

iterations to achieve either $\|\boldsymbol{z}_k - \boldsymbol{z}^*\|^2 \leq \varepsilon^2$ or $V_\lambda(\boldsymbol{z}_K) \leq \varepsilon^2$.

Proof. The proof is done in case by case, suggesting a "bad" function for each of 4 different regimes of step size ratio r: *(1)* $\mu_1/\mu_2 \leq r \leq \kappa_2/c$, *(2)* $c/\kappa_1 \leq r \leq \mu_1/\mu_2$, *(3)* $r \geq \kappa_2/c$, and *(4)* $0 < r \leq c/\kappa_1$. Readers might notice the similarities of the proofs for *(1)*↔*(2)* and *(3)*↔*(4)*.

Case 1. ($\mu_1/\mu_2 \leq r \leq \kappa_2/c$). Consider

$$
f^{(1)}(v, x; y) := \frac{\mu_1}{2} v^2 + \frac{r\mu_2}{2} x^2 - \frac{\mu_2}{2} y^2 + \ell xy,
$$

where $\ell^2 = L^2 - r\mu_2^2 - L\mu_2|r - 1| \geq 0$. Applying Proposition 28, it can be shown that $f^{(1)} \in \mathcal{F}(L, \mu_1, \mu_2)$. Also, $\boldsymbol{z}^* = (0, 0; 0)$ is its unique saddle point. Note that, the GDA on $f^{(1)}$ can be written as

$$
v_{t+1} = \left(1 - \frac{\beta\mu_1}{r}\right) v_t, \quad \begin{bmatrix} x_{t+1} \\ y_{t+1} \end{bmatrix} = \underbrace{\begin{bmatrix} 1 - \beta\mu_2 & -\beta\ell/r \\ \beta\ell & 1 - \beta\mu_2 \end{bmatrix}}_{\boldsymbol{A}} \begin{bmatrix} x_t \\ y_t \end{bmatrix} = \boldsymbol{A} \begin{bmatrix} x_t \\ y_t \end{bmatrix}.
$$

Also, the eigenvalues τ of $\boldsymbol{A}$ is

$$
\begin{aligned}
\tau &= 1 - \beta\mu_2 \pm \sqrt{(1 - \beta\mu_2)^2 - ((1 - \beta\mu_2)^2 + \beta^2\ell^2/r)} \\
&= 1 - \beta\mu_2 \pm \frac{\beta\ell}{\sqrt{r}}\sqrt{-1}.
\end{aligned}
$$

The spectral radius (*i.e.*, maximum absolute eigenvalue) is

$$
\rho(\boldsymbol{A}) = \sqrt{(1 - \beta\mu_2)^2 + \beta^2\ell^2/r}.
$$

Since the eigenvalues are complex conjugates of each other (the magnitudes are the same), both eigenvalues have magnitude $\rho(\boldsymbol{A})$. Then, by Proposition 27, $\rho(\boldsymbol{A}) < 1$ is necessary for convergence. To this end, we need $\beta > 0$ satisfying $\beta < 2\mu_2 r/(r\mu_2^2 + \ell^2)$.

To guarantee $\|(v_k, x_k; y_k) - (0, 0; 0)\|^2 \leq \varepsilon^2$, we need a large enough k to have $w_k^2 \leq \mathcal{O}(\varepsilon^2)$. Such a k is required to be at least $\Omega\left(\frac{r}{\beta\mu_1} \log(1/\varepsilon)\right)$. Now note that, since $\mu_1/\mu_2 \leq r \leq \kappa_2/c$ and $\kappa_2 \geq c$,

$$
\frac{1}{\beta} > \frac{r\mu_2^2 + \ell^2}{2\mu_2 r} = \frac{L^2 - L\mu_2|r - 1|}{2\mu_2 r} = \frac{L^2}{2\mu_2 r}\left(1 - \frac{|r - 1|}{\kappa_2}\right) \geq \frac{L^2}{2\mu_2 r}\left(1 - \frac{1}{c}\right).
$$

The last inequality is true by minimizing $\left(1 - \frac{|r-1|}{\kappa_2}\right)$ for $r \in [\mu_1/\mu_2, \kappa_2/c]$. If $r \geq 1$, it has smaller value when r is larger: by taking $r = \kappa_2/c$, we have $1 - \frac{\kappa_2/c - 1}{\kappa_2} \geq 1 - \frac{1}{c}$. Otherwise ($r < 1$), which

is possible only when $\mu_1 < \mu_2$, the term has smaller value when r is smaller: by taking $r = \mu_1/\mu_2$, we have $1 + \frac{\mu_1/\mu_2 - 1}{\kappa_2} = 1 + \frac{\mu_1 - \mu_2}{L} \geq 1 - \frac{1}{\kappa_2} \geq 1 - \frac{1}{c}$. Thus, we eventually need $\Omega\left(\frac{L^2}{\mu_1\mu_2}\log(1/\varepsilon)\right)$ iterations.

Case 2. $(c/\kappa_1 \leq r \leq \mu_1/\mu_2)$. Consider

$$f^{(2)}(x; y, w) = f_r^{(2)}(x; y, w) := \frac{\mu_1}{2}x^2 - \frac{\mu_1}{2r}y^2 + \tilde{\ell}xy - \frac{\mu_2}{2}w^2,$$

where $\tilde{\ell}^2 = L^2 - \mu_1^2/r - L\mu_1|1 - 1/r| \geq 0$. Applying Proposition 28, it can be shown that $f^{(2)} \in \mathcal{F}(L, \mu_1, \mu_2)$, and $\boldsymbol{z}^* = (0; 0, 0)$ is its unique saddle point. Note that, the GDA on $f^{(2)}$ can be written as

$$\begin{bmatrix} x_{t+1} \\ y_{t+1} \end{bmatrix} = \underbrace{\begin{bmatrix} 1 - \beta\mu_1/r & -\beta\ell/r \\ \beta\ell & 1 - \beta\mu_1/r \end{bmatrix}}_{\boldsymbol{B}} \begin{bmatrix} x_t \\ y_t \end{bmatrix} = \boldsymbol{B}\begin{bmatrix} x_t \\ y_t \end{bmatrix}, \quad w_{t+1} = (1 - \beta\mu_2)w_t.$$

Also, the eigenvalues τ of $\boldsymbol{B}$ is

$$\tau = 1 - \beta\mu_1/r \pm \sqrt{(1 - \beta\mu_1/r)^2 - ((1 - \beta\mu_1/r)^2 + \beta^2\ell^2/r)}$$
$$= 1 - \frac{\beta\mu_1}{r} \pm \frac{\beta\ell}{\sqrt{r}}\sqrt{-1}.$$

The spectral radius is
$$\rho(\boldsymbol{B}) = \sqrt{(1 - \beta\mu_1/r)^2 + \beta^2\ell^2/r}.$$

Since the eigenvalues are complex conjugates of each other (the magnitudes are the same), both eigenvalues have magnitude $\rho(\boldsymbol{A})$. Then, by Proposition 27, $\rho(\boldsymbol{A}) < 1$ is necessary for convergence. To this end, we need $\beta > 0$ satisfying $\beta < 2\mu_1/(\mu_1^2/r + \ell^2)$.

To guarantee $\|(x_k; y_k, w_k) - (0; 0, 0)\|^2 \leq \varepsilon^2$, we need a large enough k to have $w_k^2 \leq \mathcal{O}(\varepsilon^2)$. Such a k is required to be at least $\Omega\left(\frac{1}{\beta\mu_2}\log(1/\varepsilon)\right)$. Now note that, since $c/\kappa_1 \leq r \leq \mu_1/\mu_2$ and $\kappa_1 \geq c$,

$$\frac{1}{\beta} > \frac{\mu_1^2/r + \ell^2}{2\mu_1} = \frac{L^2 - L\mu_1|1 - 1/r|}{2\mu_1} = \frac{L^2}{2\mu_1}\left(1 - \frac{|1 - 1/r|}{\kappa_1}\right) \geq \frac{L^2}{2\mu_1}\left(1 - \frac{1}{c}\right).$$

The last inequality is true by minimizing $\left(1 - \frac{|1 - 1/r|}{\kappa_1}\right)$ for $r \in [c/\kappa_1, \mu_1/\mu_2]$. If $1 > 1/r$, which is possible only when $\mu_1 > \mu_2$, it has smaller value when r is larger: by taking $r = \mu_1/\mu_2$, we have $1 - \frac{1 - \mu_2/\mu_1}{\kappa_1} = 1 - \frac{\mu_1 - \mu_2}{L} \geq 1 - \frac{1}{\kappa_1} \geq 1 - \frac{1}{c}$. Otherwise $(1 < 1/r)$, the term has smaller value when r is smaller: by taking $r = c/\kappa_1$, we have $1 + \frac{1 - \kappa_1/c}{\kappa_1} \geq 1 - \frac{1}{c}$. Thus, we eventually need $\Omega\left(\frac{L^2}{\mu_1\mu_2}\log(1/\varepsilon)\right)$ iterations.

Case 3. $(r \geq \kappa_2/c)$. Consider $f^{(3)}(x; y) = \frac{\mu_1}{2}x^2 - \frac{L}{2}y^2$. Clearly, $f^{(3)} \in \mathcal{F}(L, \mu_1, L) \subset \mathcal{F}(L, \mu_1, \mu_2)$ and $\boldsymbol{z}^* = (0, 0)$ is its unique saddle point. The GDA on $f^{(3)}$ can be written as

$$x_{k+1} = \left(1 - \frac{\beta\mu_1}{r}\right)x_k, \quad y_{k+1} = (1 - \beta L)y_k.$$

To guarantee $\|(x_k, y_k) - (0, 0)\|^2 \leq \varepsilon^2$, we need a large enough k to have $x_k^2 \leq \mathcal{O}(\varepsilon^2)$. Such a k is required to be at least $\Omega\left(\frac{r}{\beta\mu_1}\log(1/\varepsilon)\right)$. Also, we need $\beta < 2/L$ to guarantee $y_k \to 0$ (i.e., otherwise, it diverges). Combining these facts, we eventually need $\Omega\left(\frac{Lr}{\mu_1}\log(1/\varepsilon)\right)$ iterations.

Case 4. $(0 < r \leq c/\kappa_1)$. Consider $f^{(4)}(x; y) = \frac{L}{2}x^2 - \frac{\mu_2}{2}y^2$. Clearly, $f^{(4)} \in \mathcal{F}(L, L, \mu_2) \subset \mathcal{F}(L, \mu_1, \mu_2)$ and $\boldsymbol{z}^* = (0, 0)$ is its unique saddle point. The GDA on $f^{(4)}$ can be written as

$$x_{k+1} = \left(1 - \frac{\beta L}{r}\right)x_k, \quad y_{k+1} = (1 - \beta\mu_2)y_k.$$

To guarantee $\|(x_k, y_k) - (0,0)\|^2 \leq \varepsilon^2$, we need a large enough k to have $y_k^2 \leq \mathcal{O}(\varepsilon^2)$. Such a k is required to be at least $\Omega\left(\frac{1}{\beta\mu_2}\log(1/\varepsilon)\right)$. Also, we need $\beta < 2r/L$ to guarantee $x_k \to 0$ (i.e., otherwise, it diverges). Combining these facts, we eventually need $\Omega\left(\frac{L}{r\mu_2}\log(1/\varepsilon)\right)$ iterations.

Lastly, we note that the lower iteration complexity bound in terms of the potential function V_λ is equivalent to the complexity in terms of squared distance norm from the (unique) saddle point z^*, up to constant factors. This is proved in Lemma 29 that we defer its proof. $\qquad\square$

Here are the postponed proofs from the proof above.

Proposition 27. *For a square matrix $A \in \mathbb{R}^{m \times m}$ and a sequence of m-dimensional vectors (v_k), the matrix iteration $v_{k+1} = Av_k$ converges to $v_k \to 0$ if and only if the spectral radius (i.e., maximum absolute eigenvalue) of $\rho(A)$ of A is less than 1. Furthermore, its convergence speed is characterized by $\mathcal{O}((\rho(A)+\varepsilon)^k)$ for any (arbitrarily small) $\varepsilon > 0$.*

Proof. See Horn & Johnson (2012, Theorem 5.6.10-12). $\qquad\square$

Proposition 28. *Let μ_1, μ_2, and L be positive numbers such that $L \geq \max\{\mu_1, \mu_2\}$. Consider a quadratic function f on $\mathbb{R} \times \mathbb{R}$ defined by*

$$f(x; y) = \frac{\mu_1}{2}x^2 - \frac{\mu_2}{2}y^2 + \ell xy, \quad \text{where } \ell^2 \leq L^2 - \mu_1\mu_2 - L|\mu_1 - \mu_2|.$$

Then, $f \in \mathcal{F}(L, \mu_1, \mu_2)$, and its unique saddle point is $z^* = (0,0)$.

For example, if $\mu_1 \geq \mu_2$, $\ell^2 = (L - \mu_1)(L + \mu_2)$ is enough to guarantee L-smoothness.

Proof. The strong-convex-strong-concavity is trivially true. Note that the gradient and hessian of f is

$$\nabla f(x; y) = H[x \; y]^\top, \quad H = \begin{bmatrix} \mu_1 & \ell \\ \ell & -\mu_2 \end{bmatrix}.$$

Since H is a non-singular matrix, f has a unique stationary point at origin. By Proposition 11, it is also a unique saddle & global minimax point.

For any two distinct points $z_1 = (x_1; y_1)$ and $z_2 = (x_2; y_2)$ in $\mathbb{R} \times \mathbb{R}$,

$$\frac{\|\nabla f(z_1) - \nabla f(z_2)\|}{\|z_1 - z_2\|} = \frac{\|H(z_1 - z_2)\|}{\|z_1 - z_2\|} \leq \|H\|_2,$$

where $\|H\|_2$ is spectral norm (i.e., maximum singular value) of H. We want to show that $\|HH^\top\|_2 \leq L^2$. To show this, it is enough to verify the following two inequalities:

$$\det(L^2 I - HH^\top) = L^4 - (\mu_1^2 + \mu_2^2 + 2\ell^2)L^2 + (\mu_1\mu_2 + \ell^2)^2 \geq 0,$$
$$\text{trace}(H)/2 = (\mu_1^2 + \mu_2^2)/2 + \ell^2 \leq L^2.$$

Let $\ell^2 = L^2 - \mu_1\mu_2 + b$ for some b. Plugging this ℓ^2 into both inequalities, we get

$$\det(L^2 I - HH^\top) = b^2 - (\mu_1 - \mu_2)^2 L^2 \geq 0, \text{ and } b \leq -(\mu_1 - \mu_2)^2/2,$$

respectively. One can check that $b = -L|\mu_1 - \mu_2|$ is the largest choice of b that satisfies both inequalities above. This proves the proposition. $\qquad\square$

Subsequently, we show that if our convergence rate is exponential, then the iteration complexity in terms of $\|z - z^*\|^2$ is equivalent to that in terms of $V_\lambda(z) = \lambda[\Phi(x) - \Phi^*] + [\Phi(x) - f(z)]$ for PŁ(Φ)-PŁ problem, up to constant factors. This also applies to the function class $\mathcal{F}(L, \mu_1, \mu_2)$ since it is a subclass of PŁ(Φ)-PŁ ($\because$ Propositions 7 and 10).

Lemma 29. *Suppose $f(x; y)$ is an L-smooth function satisfying y-side μ_2-PŁ condition and primal μ_1-PŁ condition (i.e., PŁ(Φ)-PŁ). Suppose $z^* = (x^*; y^*)$ is a global minimax point of f. Then, it satisfies*

$$\frac{\lambda\mu_1\mu_2^2}{2(\lambda\mu_1\mu_2 + 2L^2)}\|z - z^*\|^2 \leq V_\lambda(z) \leq \frac{(\lambda+1)L^3}{\mu_2^2}\|z - z^*\|^2.$$

We remark that the second inequality also holds for general smooth nonconvex-PŁ problems.

Proof. Let $\kappa_1 = L/\mu_1$ and $\kappa_2 = L/\mu_2$ be condition numbers. By the conditions of f (smoothness and PŁ conditions), for any x and y,

$$\frac{\mu_1}{2} \|x - x^*\|^2 \overset{\text{Prop. 10}}{\leq} \Phi(x) - \Phi^* \overset{\text{Prop. 9}}{\leq} \frac{L(\kappa_2 + 1)}{2} \|x - x^*\|^2,$$

$$\frac{\mu_2}{2} \|y - y^*(x)\|^2 \overset{\text{Ass. 4}}{\leq} \Phi(x) - f(x; y) \overset{\text{Ass. 1}}{\leq} \frac{L}{2} \|y - y^*(x)\|^2,$$

where $y^*(x)$ is a projection of y to $\arg\max_{y'} f(x; y')$. In particular, $y^*(x^*) = y^*$. Since $y^*(x)$ is a function of x and can differ from y^*, we need to bound the term $\|y - y^*(x)\|^2$ using $\|x - x^*\|^2$ and $\|y - y^*\|^2$. To upper-bound the term $\|y - y^*(x)\|^2$, note that,

$$\|y - y^*(x)\|^2 \leq (\|y - y^*(x^*)\| + \|y^*(x) - y^*(x^*)\|)^2$$
$$\leq (\|y - y^*\| + \kappa_2 \|x - x^*\|)^2$$
$$\leq (1 + \kappa_2^2) \left(\|y - y^*\|^2 + \|x - x^*\|^2\right).$$

The first inequality holds by triangle inequality, the second inequality holds by Proposition 8, and the last inequality holds by Cauchy-Schwarz inequality.[11] To lower-bound in a similar way, note that for any constant $a > 0$,

$$\|y - y^*\|^2 \leq (\|y - y^*(x)\| + \|y^*(x) - y^*(x^*)\|)^2$$
$$\leq \left(\|y - y^*(x)\| + \frac{\kappa_2}{\sqrt{a}} \cdot \sqrt{a} \|x - x^*\|\right)^2$$
$$\leq \left(1 + \frac{\kappa_2^2}{a}\right) \left(\|y - y^*(x)\|^2 + a \|x - x^*\|^2\right).$$

$$\therefore \|y - y^*(x)\|^2 \geq \frac{1}{1 + \kappa_2^2/a} \|y - y^*\|^2 - a \|x - x^*\|^2.$$

Now we can prove the inequalities in the lemma. We first show the second one. Applying $\kappa_2 \geq 1$ multiple times,

$$V_\lambda(x; y) = \lambda[\Phi(x) - \Phi^*] + [\Phi(x) - f(z)]$$
$$\leq \frac{\lambda L(\kappa_2 + 1)}{2} \|x - x^*\|^2 + \frac{L}{2} \|y - y^*(x)\|^2$$
$$\leq \left(\frac{\lambda L(\kappa_2 + 1)}{2} + \frac{L(1 + \kappa_2^2)}{2}\right) \|x - x^*\|^2 + \frac{L(1 + \kappa_2^2)}{2} \|y - y^*\|^2$$
$$\leq (\lambda + 1)L\kappa_2^2 \left(\|x - x^*\|^2 + \|y - y^*\|^2\right) = \frac{(\lambda + 1)L^3}{\mu_2^2} \|z - z^*\|^2.$$

To show the first inequality of the lemma, let $a = \frac{\lambda \mu_1}{2\mu_2}$.

$$V_\lambda(x; y) \geq \frac{\lambda \mu_1}{2} \|x - x^*\|^2 + \frac{\mu_2}{2} \|y - y^*(x)\|^2$$
$$\geq \left(\frac{\lambda \mu_1}{2} - \frac{\mu_2 a}{2}\right) \|x - x^*\|^2 + \frac{\mu_2}{2(1 + \kappa_2^2/a)} \|y - y^*\|^2$$
$$\geq \frac{\lambda \mu_1}{4} \|x - x^*\|^2 + \frac{\lambda \mu_1}{4(a + \kappa_2^2)} \|y - y^*\|^2$$
$$\geq \frac{\lambda \mu_1}{4(a + \kappa_2^2)} \left(\|x - x^*\|^2 + \|y - y^*\|^2\right) = \frac{\lambda \mu_1 \mu_2^2}{2(\lambda \mu_1 \mu_2 + 2L^2)} \|z - z^*\|^2.$$

This concludes the proof. □

---

[11] $(ax + by)^2 \leq (a^2 + b^2)(x^2 + y^2)$ for real numbers a, b, x, y.

The equivalence of iteration complexities for achieving $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2 \le \varepsilon^2$ or $V_\lambda(\boldsymbol{z}_K) \le \varepsilon^2$ is quite straightforward from this lemma, as long as the convergence speed is exponential. For example, suppose we have a upper convergence bound $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2 \le a \exp(-K/r)$ for some constants $a, r > 0$. This implies a upper iteration complexity bound $K = \mathcal{O}(r \log(1/\varepsilon))$ sufficient to achieve $\|\boldsymbol{z}_K - \boldsymbol{z}^*\|^2 \le \varepsilon^2$. Then by Lemma 29, we also have $V_\lambda(\boldsymbol{z}_K)^2 \le a' \exp(-K/r)$ where $a' = a(\lambda + 1)L^3/\mu_2^2$ is also a constant. This implies a lower iteration complexity bound $K = \mathcal{O}(r \log(1/\varepsilon))$ as well, sufficient to achieve $V_\lambda(\boldsymbol{z}_K)^2 \le \varepsilon^2$. The other way of complexity translation operates with a similar logic.

F   OMITTED COMPARISON WITH RELATED WORKS

F.1   COMPARISON WITH XIE ET AL. (2021)

To specialize Xie et al. (2021, Theorem 3) to the single-machine setup and discuss their results in terms of our notation, we need to replace their symbols

$$(T, S, K, \sigma_1^2, \sigma_2^2, G_1^2, G_2^2, L_{12}, L_f, \mu, L_\Phi, \mathcal{L}_0, \eta_t, \gamma_t)$$

with the following symbols from our notation:

$$(K, 1, n, 0, 0, B, B, L, L, \mu_2, L(\kappa_2 + 1), V_\lambda(\boldsymbol{z}_0^1), \alpha, \beta),$$

and also put $A = 0$ (there analysis only applies uniformly bounded component variance per machine). Then we can *naively* translate the bound of Xie et al. (2021, Theorem 3) to our language as

$$\min_{k \in [K]} \mathbb{E}\left[\|\Phi(\boldsymbol{x})\|^2\right] \overset{(?)}{\leq} \mathcal{O}\left(\frac{\kappa_2 L V_\lambda(\boldsymbol{z}_0^1)}{K} + \kappa_2^2 \left(\frac{L^2 B V_\lambda(\boldsymbol{z}_0^1)^2}{K^2}\right)^{1/3}\right).$$

To the best of our knowledge, however, we believe there may be a mistake in the proof of Xie et al. (2021, Appendix C.4). From the inequalities in the last page of their paper, we notice that the term $\frac{40 L_{12}^2 \mathcal{L}_0}{\mu^2 \gamma K T}$ might be missing in a step, where γ is chosen to be the minimum of several terms including $\frac{1}{87 L_f K}$. Thus, as far as we can tell, it seems inevitable that this omitted term would lead to an additional term $\frac{3480 L_f L_{12}^2 \mathcal{L}_0}{\mu^2 T}$ in the final bound. By combining this to their bound and re-translating it, we eventually have

$$\min_{k \in [K]} \mathbb{E}\left[\|\Phi(\boldsymbol{x})\|^2\right] \leq \mathcal{O}\left(\frac{\kappa_2^2 L V_\lambda(\boldsymbol{z}_0^1)}{K} + \kappa_2^2 \left(\frac{L^2 B V_\lambda(\boldsymbol{z}_0^1)^2}{K^2}\right)^{1/3}\right),$$

since their L_{12}^2/μ^2 translates to our κ_2^2. Therefore, their result actually shows the same dependency on condition number κ_2 as our Theorem 1. Nevertheless, comparing the terms related to the component-wise variance B, ours is better. In the second term in the bound above does not shrink even when the number of iterations (per machine & per communication) grows. In our case (Theorem 1), however, the dominant term (in K) can be briefly written as $\mathcal{O}((\frac{B}{nK^2})^{1/3})$ which can diminish with large n, *i.e.*, the number of iterations per epoch.