ANOCE: Analysis of Causal Effects with Multiple Mediators via Constrained Structural Learning

Abstract

In the era of causal revolution, identifying the causal effect of an exposure on the outcome of interest is an important problem in many areas, such as epidemics, medicine, genetics, and economics. Under a general causal graph, the exposure may have a direct effect on the outcome and also an indirect effect regulated by a set of mediators. An analysis of causal effects that interprets the causal mechanism contributed through mediators is hence challenging but on demand. To the best of our knowledge, there are no feasible algorithms that give an exact decomposition of the indirect effect on the level of individual mediators, due to common interaction among mediators in the complex graph. In this paper, we establish a new statistical framework to comprehensively characterize causal effects with multiple mediators, namely, ANalysis Of Causal Effects (ANOCE), with a newly introduced definition of the mediator effect, under the linear structure equation model. We further propose a constrained causal structure learning method by incorporating a novel identification constraint that specifies the temporal causal relationship of variables. The proposed algorithm is applied to investigate the causal effects of 2020 Hubei lockdowns on reducing the spread of the coronavirus in Chinese major cities out of Hubei.

1 Introduction

In the era of causal revolution, identifying the causal effect of an exposure on the outcome of interest is an important problem in many areas, such as epidemics (Hernán, 2004), medicine (Hernán et al., 2000), education (Card, 1999), and economics (Panizza & Presbitero, 2014). Under a general causal graph, the exposure may have a direct effect on the outcome and also an indirect effect regulated by a set of mediators (or intermediate variables). For instance, during the outbreak of Coronavirus disease 2019 (COVID-19), the Chinese government has taken extreme measures to stop the virus spreading such as locking Wuhan down on Jan 23rd, 2020, followed by 12 other cities in Hubei, known as the "2020 Hubei lockdowns". This approach (viewed as the exposure), directly blocked infected people leaving from Hubei; and also stimulated various quarantine measures taken by cities outside of Hubei (as the mediators), which further decreased the migration countrywide in China, and thus indirectly control the spread of COVID-19. Quantifying the causal effects of 2020 Hubei lockdowns on reducing the COVID-19 spread regulated by different cities outside Hubei is challenging but of great interest for the current COVID-19 crisis. An analysis of causal effects that interprets the causal mechanism contributed via individual mediators is thus very important.

Many recent efforts have been made on studying causal effects that are regulated by mediators. Chakrabortty et al. (2018) specified the individual mediation effect in a sparse high-dimensional causal graphical model. However, the sum of marginal individual mediation effect is not equal to the effect of all mediators considered jointly (i.e. the indirect effect) due to the common interaction among mediators (VanderWeele & Vansteelandt, 2014). Here, 'interaction' means that there exists at least one mediator that is regulated by other mediator(s) (see Figure 1b for illustration), in contrast to the simple 'parallel' case (shown in Figure 1a). Vansteelandt & Daniel (2017) considered an exact decomposition of the indirect effect with a two-mediator setting based on the conditional densities of mediators, while there was no feasible algorithm provided to solve their proposed expressions yet. Therefore, a new framework with a computational friendly algorithm that gives an exact decomposition of the indirect effect on the level of individual mediators is desired under the complex causal network.

(a) A causal graph with parallel mediators.   (b) A causal graph with interacted mediators.

Figure 1: Illustration of causal graphs with different types of mediators, where A is the exposure, $\{M_1, \cdots, M_p\}$ are mediators, and Y is the outcome of interest.

To estimate the underlying causal network, structure learning algorithms of the directed acyclic graph (DAG) are widely used. Popular methods such as the PC algorithm (Spirtes et al., 2000) that uses conditional independence tests to examine the existence of edges between each pair of variables, require strong assumptions and thus have no guarantee in the finite sample regime. Recently, Zheng et al. (2018) opened up another class of causal discovery methods by directly formulating a pure optimization problem over real metrics with a novel characteristic of the acyclicity. Yu et al. (2019) further extended Zheng et al. (2018)'s work with a deep generative model, and showed better performance on the structure learning with weaker assumptions on the noise. See more follow-up works in Lachapelle et al. (2019) and Zhu & Chen (2019). However, the current cutting-edge methods neglect the temporal causal relationship among variables, and thus cannot appropriately represent the causal network with pre-specified exposure and outcome.

In this paper, we consider establishing a new statistical framework to comprehensively characterize causal effects with multiple mediators, namely, ANalysis Of Causal Effects (ANOCE), under the linear structure equation model (LSEM). Specifically, we propose two causal effects on the level of individual mediators, the natural direct effect and the natural indirect effect for a mediator, denoted as DM and IM, respectively. Our proposed DM can be interpreted as the direct effect of a particular mediator on the outcome that is not regulated by other mediators, while the IM is the indirect effect of the mediator controlled by its descendant mediators. We prove that the DM is valid in the sense that it exactly decomposes the indirect effect of the exposure on the outcome, followed by an ANOCE table to explain different sources of causal effects. To bridge the cutting-edge graphical learning approaches with the temporal causal relationship of variables, we extend the variational auto-encoder (VAE) framework in Yu et al. (2019) with a novel identification constraint that specifies the topological order of the exposure and the outcome. The proposed constrained VAE algorithm is then used to estimate causal effects defined in our ANOCE table, named as 'ANOCE-CVAE'.

Our contributions can be summarized in the following three aspects:
• 1). Conceptually, we define different sources of causal effects through mediators with a newly introduced definition of direct and indirect mediator effects, and give an exact decomposition of the indirect effect on the level of individual mediators, under the linear structure equation model.
• 2). Methodologically, we incorporate the background knowledge (the temporal causal relationship among variables) when using an optimization approach to the causal discovery. Such prior knowledge can be generalized for any measured variable and on the possible set of their parents. Our proposed constrained structural learning can be easily extended to other score-based algorithms.
• 3). Practically, extensive simulations are conducted to demonstrate the empirical validity of the proposed algorithm and its competitive performance among existing causal discovery algorithms. Our method is applied to investigate the causal effects of 2020 Hubei lockdowns on reducing the COVID-19 spread in China, by quantifying the individual effect for each city.

2 BACKGROUND AND RELATED WORK

2.1 GRAPH TERMINOLOGY

Consider a graph $\mathcal{G} = (X, E)$ with a node set X and an edge set E. There is at most one edge between any pair of nodes. If there is an edge between X_i and X_j, then X_i and X_j are adjacent. A node X_i is said to be a parent of X_j if there is a directed edge from X_i to X_j. Let the set of all parents of node X_j in $\mathcal{G}$ as $PA_{X_j}(\mathcal{G})$. A path from X_i to X_j in $\mathcal{G}$ is a sequence of distinct vertices, $\pi \equiv \{a_0, a_1, \cdots, a_L\} \subset V$ such that $a_0 = X_i$, and $a_L = X_j$. A directed path from X_i to X_j is a

path between X_i and X_j where all edges are directed toward X_j. A directed cycle is formed by the directed path from X_i to X_j together with the directed edge X_j to X_i. A directed graph that does not contain directed cycles is called a directed acyclic graph (DAG). A directed graph is acyclic if and only if it has a topological ordering. Suppose a DAG $\mathcal{G} = (X, E)$ that characterizes the causal relationship among $|X| = d$ nodes, where $X = [X_1, X_2, \cdots, X_d]^\top$ represents a random vector and an edge $X_i \rightarrow X_j$ means that X_i is a direct cause of X_j. Let $B = \{b_{i,j}\}_{1 \le i \le d, 1 \le j \le d}$ be a $d \times d$ matrix, where $b_{i,j}$ is the weight of the edge $X_i \rightarrow X_j \in E$, and $b_{i,j} = 0$ otherwise. Then, we say that $\mathcal{G} = (X, B)$ is a weighted DAG with the node set X and the weighted adjacency matrix B (the edge set E is nested in B).

2.2 Related Work

Our work connects to the literature of the causal graphical model. Pearl et al. (2009) provided a comprehensive review of recent advances in the analysis of causes and counterfactuals using 'do-operator' by graphical methods. Later, Maathuis et al. (2009) started to use an unknown DAG without hidden variables to estimate the causal effects from the high-dimensional observational data. Nandy et al. (2017) extended the work of Maathuis et al. (2009) with the linear structure equation model, followed by the individual mediation effect defined in Chakrabortty et al. (2018). All of these models rely on the PC algorithm to search the Markov equivalence class of the partial DAG, and usually require strong assumptions due to the computational limit. Our ANOCE is established under the same causal structure of Chakrabortty et al. (2018) but without sparsity and normality assumptions.

Wide literature on causal discovery can be summarized in three classes. The first type focuses on local conditional independence tests to find a causal skeleton and then determine the orientation of edges, such as the well-known PC algorithm (Spirtes et al., 2000; Kalisch & Bühlmann, 2007). However, these methods have no guarantee in the finite sample regime because of the hardness of testing conditional independence when the variables are continuous (Shah & Peters, 2018). The second class specifies properly functional causal models with additional assumptions on data distribution, including the ICA-LiNGAM (Shimizu et al., 2006) and the causal additive model (CAM) (Bühlmann et al., 2014). The last class, the score-based method, includes the greedy equivalence search (GES) (Chickering, 2002) and the fast GES (fGES) that use for example Bayesian scores in searching a space of causal models. Recently, Zheng et al. (2018) opened up another track of score-based methods by constructing an optimization with an acyclicity constraint under the LSEM, i.e. the NOTEARS. A follow-up work using a VAE parameterized by a graph neural network that generalizes LSEM was proposed in Yu et al. (2019) with a more computational friendly constraint, namely DAG-GNN. Also see Zhu & Chen (2019) and Lachapelle et al. (2019) for other cutting-edge structural learning methods. Our ANOCE-CVAE algorithm extends (though not limits to) the DAG-GNN for an illustration, since it can handle non-linear cases and does not require the normality assumption of the noise.

3 Analysis of Causal Effects

3.1 Statistical Framework and Assumptions

Let A be the exposure/treatment, $M = [M_1, M_2, \cdots, M_p]^\top$ be mediators with dimension p, and Y be the outcome of interest. Suppose there exists a weighted DAG $\mathcal{G} = (X, B)$ that characterizes the causal relationship among $X = [A, M^\top, Y]^\top$, where the dimension of X is $d = p + 2$. Let $Y^*(A = a, M = m)$ be the potential outcome that would be observed after receiving treatment a and setting mediators as m, and $M^*(A = a)$ be the potential mediators that would be observed after receiving treatment a. As standard in the causal inference (Rosenbaum & Rubin, 1983), we assume that there is no unmeasured confounder: (A1) the effect of the treatment A on the outcome Y is unconfounded, i.e. $Y^*(A = a, M = m) \perp A, \forall a, m$; (A2) the effect of the treatment A on the mediators M is unconfounded, i.e. $M^*(A = a) \perp A, \forall a$; (A3) the effect of the mediators M on the outcome Y is unconfounded given the treatment A, i.e. $Y^*(A = a, M = m) \perp M|A, \forall a, m$. In addition, as standard in the graphical causal discovery, we also make the Markov condition, the faithfulness condition, causal sufficiency assumption, and the linear structural equation model (LSEM) such that $X = [A, M^\top, Y]^\top$ characterized by the pair $(\mathcal{G}, \epsilon)$ is generated by

$$X = B^\top X + \epsilon, \tag{1}$$

where ϵ is a random vector of jointly independent error variables.

Denote all directed paths in $\mathcal{G}$ that start with the exposure A and end with the outcome Y as set $\{\pi_{AY}(\mathcal{G})\}$. If there exists at least one directed path $\pi^* \in \{\pi_{AY}(\mathcal{G})\}$ such that the length of π^* is

larger than 2, we say there is an interaction among mediators, as shown in Figure 1b; otherwise, we call mediators are 'parallel' as shown in Figure 1a. In this paper, we consider all possible causal structures with multiple mediators under assumptions (A1-A3).

We next give the total effect (TE), the natural direct effect that is not mediated by mediators (DE), and the natural indirect effect that is regulated by mediators (IE) defined in Pearl et al. (2009).

Definition 3.1 *(Pearl et al., 2009)*

$$TE = \partial E\{Y|do(A = a)\}/\partial a = E\{Y|do(A = a + 1)\} - E\{Y|do(A = a)\},$$
$$DE = E\{Y|do(A = a + 1, M = m^{(a)})\} - E\{Y|do(A = a)\},$$
$$IE = E\{Y|do(A = a, M = m^{(a+1)})\} - E\{Y|do(A = a)\},$$

where $do(A = a)$ is a mathematical operator to simulate physical interventions that hold A constant as a while keeping the rest of the model unchanged, which corresponds to remove edges into A and replace A by the constant a in $\mathcal{G}$. Here, $m^{(a)}$ is the value of M if setting $do(A = a)$, and $m^{(a+1)}$ is the value of M if setting $do(A = a + 1)$. Refer to Pearl et al. (2009) for more details of 'do-operator'.

Note that in the assumed linear model, the slope of the line is the same everywhere; for convenience and simplicity, we use a and $a + 1$ to present the change of the treatment of 1 in the definition.

3.2 NATURAL DIRECT AND INDIRECT EFFECT FOR INDIVIDUAL MEDIATORS

We first give the definition of the natural direct effect for an individual mediator (DM).

Definition 3.2 *Natural direct effect for M_i:*

$$
\begin{aligned}
DM_i = & \Big[E\{M_i|do(A = a + 1)\} - E\{M_i|do(A = a)\}\Big] \\
& \times \Big[E\{Y|do(A = a, M_i = m_i^{(a)} + 1, \Omega_i = o_i^{(a)})\} - E\{Y|do(A = a)\}\Big],
\end{aligned}
\tag{2}
$$

where $m_i^{(a)}$ is the value of M_i when setting $do(A = a)$, $\Omega_i = M \setminus M_i$ is the set of mediators except M_i, and $o_i^{(a)}$ is the value of Ω_i when setting $do(A = a)$.

Remark 3.1 *From Definition 3.2, the natural direct effect for M_i is the product of the total effect of the treatment A on the mediator M_i and the direct effect of the mediator M_i on the outcome Y. The second multiplier is in line with the classical meaning of 'natural' in the causal inference literature (Pearl et al., 2009). Thus, the DM can be interpreted as the causal effect through a particular mediator from the treatment on the outcome that is not regulated by its descendent mediators.*

The natural indirect effect for an individual mediator (IM) can be defined similarly.

Definition 3.3 *Natural indirect effect for M_i:*

$$
\begin{aligned}
IM_i = & \Big[E\{M_i|do(A = a + 1)\} - E\{M_i|do(A = a)\}\Big] \\
& \times \Big[E\{Y|do(A = a, M_i = m_i^{(a)} + 1)\} - E\{Y|do(A = a, M_i = m_i^{(a)} + 1, \Omega_i = o_i^{(a)})\}\Big].
\end{aligned}
$$

Remark 3.2 *The second multiplier in the IM_i captures the indirect effect of a particular mediator on the outcome regulated by its descendent mediators. We show the individual mediation effect in Chakrabortty et al. (2018) can be decomposed into the DM and the IM in Section F in the appendix.*

Next, we give explicit expressions of defined causal effects under the LSEM. Specifically, we can write the linear structural model 1 under assumptions (A1-A3) as

$$
\begin{bmatrix} A \\ M \\ Y \end{bmatrix} = B^\top \begin{bmatrix} A \\ M \\ Y \end{bmatrix} + \epsilon = \begin{bmatrix} 0 & \mathbf{0}_{p \times 1} & 0 \\ \boldsymbol{\alpha} & B_M^\top & 0 \\ \gamma & \boldsymbol{\beta}^\top & 0 \end{bmatrix} \begin{bmatrix} A \\ M \\ Y \end{bmatrix} + \begin{bmatrix} \epsilon_A \\ \epsilon_{M_p} \\ \epsilon_Y \end{bmatrix},
\tag{3}
$$

where γ is a scalar, $\boldsymbol{\alpha}$, $\boldsymbol{\beta}$, and $\mathbf{0}_{p \times 1}$ are $p \times 1$ vectors, B_M is a $p \times p$ matrix, and $\epsilon \equiv [\epsilon_A, \epsilon_M^\top, \epsilon_Y]^\top$. Here, γ presents the weight of the edge $A \to Y$, the i-th element of $\boldsymbol{\alpha}$ corresponds to the weight of the edge $A \to M_i$, and the i-th element of $\boldsymbol{\beta}$ is the weight of the edge $M_i \to Y$. Note that by

assumptions (A1-A3), we have the exposure A has no parents and the outcome Y has no descendants, so equivalently, the first row and the last column of $B^\top$ are all zeros (i.e., the first column and the last row of B are all zeros). Next, we obtain expressions of causal effects under the LSEM in the following theorem. The proof can be found in Section G.1 of the appendix.

Theorem 3.1 *Under assumptions (A1-A3) and Model 1, we have:*
1). the natural direct effect is $DE = \gamma$;
2). the natural indirect effect is $IE = \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1} \boldsymbol{\alpha}$, where I_p is a $p \times p$ identity matrix;
3). the total effect of A on Y is $TE = \gamma + \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1} \boldsymbol{\alpha}$;
4). the natural direct effect of M_i on Y is $DM_i = \boldsymbol{\beta}_i \{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i$, where $\boldsymbol{\beta}_i$ is the i-th element of $\boldsymbol{\beta}$ corresponding to the weight of $M_i \to Y$, and $\{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i$ is the i-th element of $(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}$ as the total effect of A on M_i, i.e. $E\{M_i | do(A = a + 1)\} - E\{M_i | do(A = a)\}$.

Remark 3.3 *One may refer to section A in the appendix for the invertibility of $I_p - B_M^\top$. Also, a toy example is provided in section E to illustrate how to manually compute the causal effects defined above. Note that there is no explicit expression of the IM due to the complex interaction among mediators, while we provide its theoretical form in Section G.2 with its numerical form in Section B.*

Based on the result 2) and 4) in Theorem 3.1, the IE can be presented as an additive form of DMs, as shown in Theorem 3.2. Thus, the proposed natural direct effect of individual mediators is valid in the sense that it exactly decomposes the indirect effect of the exposure on the outcome.

Theorem 3.2 *Under assumptions (A1-A3) and Model 1, the IE can be decomposed through DMs:*

$$IE = \sum_{i=1}^p DM_i.$$

3.3 ANALYSIS OF CAUSAL EFFECTS TABLE

Based on the result $TE = DE + IE$ in Pearl et al. (2009) and Theorem 3.2, we summarize the defined causal effects and their relationship in Table 1 for the analysis of causal effects (ANOCE). Firstly, the causal effect of A on Y has two sources, the direct effect from A and the indirect effect via p mediators M ($M_1, \cdots, M_p$). Next, the direct source has the degree of freedom ($d.f.$) as 1, while the indirect source has $d.f.$ as p from p mediators. Note the true $d.f.$ of the indirect effect may be smaller than p, since A may not be regulated by all mediators. Then, the causal effect for the direct source is the DE and for the indirect source is the IE, where the IE can be further decomposed into p DMs and each component corresponds to the natural direct effect for a specific mediator. The last row in the table shows that the DE and the IE compose the total effect TE with $d.f.$ as $p + 1$.

Table 1: Table of Analysis of Causal Effects (ANOCE Table).

Source		Degree of freedom		Causal effects	
Direct effect from A		1		DE	
Indirect effect via M		p		IE	
	M_1		1		DM_1
	M_2		1		DM_2
	$\vdots$		$\vdots$		$\vdots$
	M_p		1		DM_p
Total		$1 + p$		TE	

4 CONSTRAINED STRUCTURAL LEARNING FOR ANOCE

We next estimate the weighted adjacency matrix B with our causal framework under the LSEM to calculate causal effects. To better capture the sampling distribution faithful to the DAG, we consider a deep generative model that generalizes the LSEM instead of using a regression that heavily relies on assumptions of noise (see more discussion in Section 2.2). Specifically, the LSEM 1 can be rewritten as $(I_{p+2} - B^\top) X = \epsilon$, where I_{p+2} is a $(p+2) \times (p+2)$ identity matrix. Inversely, we have $X = (I_{p+2} - B^\top)^{-1} \epsilon$. Following the VAE architecture in Yu et al. (2019), we treat the random error ϵ as the independent latent variables to generate X, by two multilayer perceptrons as the encoder and the decoder, with weights denoted as θ. We adopt their acyclicity constraint on B as,

$$h_1(B) \equiv \text{tr}\big[(I_{p+2} + tB \bullet B)^{p+2}\big] - (p+2) = 0,$$

where $tr(\cdot)$ is the trace of a matrix, t is a hyperparameter that depends on an estimation of the largest eigenvalue of B, and $\bullet$ denotes for the element-wise square.

Next, to incorporate the background knowledge of the temporal causal relationship among variables, we propose an identification constraint that indicates the topological order of the exposure and the outcome. As mentioned in Equation 3, under assumptions (A1-A3), the exposure A has no parents, i.e. $PA_A(\mathcal{G}) = \varnothing$, and the outcome Y has no descendants, i.e. $Y \notin PA_X(\mathcal{G})$. Or equivalently, we have the first column and the last row of B should equal to zero. Therefore, the matrix B must satisfy

$$h_2(B) \equiv \sum_{i=1}^{p+2} |b_{i,1}| + \sum_{j=2}^{p+2} |b_{p+2,j}| = 0, \qquad (4)$$

where $b_{i,j}$ is the element of the matrix B in i-th row and j-th column. The above constraint forces the topological order of the exposure as 1 while the outcome as $p + 2$, under which the DAG is searched within a restricted regime. The prior knowledge in 4 can be generalized for any measured variable and on the possible set of their parents, by connecting the topological order to the weighted matrix B.

Following Yu et al. (2019), the objective function is the evidence lower bound with two constraints:

$$\begin{cases} \min_{B,\theta} \quad f(B,\theta) = \frac{1}{p+2} \sum_{i=1}^{p+2} D_{KL}\{q(\epsilon|X_i)||p(\epsilon)\} - E_{q(\epsilon|X_i)}\{\log p(X_i|\epsilon)\}, \\ s.t. \quad h_1(B) = 0 \quad \text{and} \quad h_2(B) = 0, \end{cases} \qquad (5)$$

where $D_{KL}(\cdot||\cdot)$ is the Kullback-Leibler divergence, $p(\epsilon)$ is the prior distribution of ϵ, $q(\epsilon|X_i)$ is the reconstructed empirical posterior distribution of ϵ, and $p(X_i|\epsilon)$ is the likelihood function. Then, we have the loss function based on the augmented Lagrangian as

$$L_{c,d}(B,\theta,\lambda_1,\lambda_2) = f(B,\theta) + \lambda_1 h_1(B) + \lambda_2 h_2(B) + c|h_1(B)|^2 + d|h_2(B)|^2, \qquad (6)$$

where λ_1 and λ_2 are Lagrange multipliers, and c and d are penalty terms. To minimize the loss in 6 and satisfy both $h_1(B) = 0$ and $h_2(B) = 0$, we simultaneously update λ_1 and λ_2 and increase c and d to infinity, by modifying the basic technique in Yu et al. (2019). Here, the minimization can be solved using a blackbox stochastic optimization such as 'Adam' in Kingma & Ba (2014). Denote the estimated matrix as $\widehat{B}$ from the above constrained structural learning. Under Theorem 3.1, we can estimate causal effects in the ANOCE table based on the learned $\widehat{B}$. We name the above algorithm as ANOCE-CVAE, with a detail pseudocode provided in Section B.

Remark 4.1 *We incorporate the temporal causal relationship among variables when using an optimization approach to the causal discovery. Such constrained structural learning is not limited to the VAE framework and can be extended to any score-based algorithms. For instance, one can add constraint 4 into the objective function in Zheng et al. (2018) or the reward in Zhu & Chen (2019).*

5 EXPERIMENTS

We conduct extensive simulation studies to investigate the proposed method on learning causal effects with multiple mediators, followed by a comparison to the popular structural learning algorithms.

5.1 SIMULATION STUDIES

Scenarios are generated as follows. In Scenario 1 to 3, we fix the dimension of the mediators M as $p = 10$ while increasing the complexity of the underlying true graph to examine the sensitivity of our algorithm to sparsity. Specifically, Scenario 1 represents the simplest causal graph with only one edge ($A \rightarrow Y$) shown in Figure 2a; and Scenario 2 has a fully connected causal graph with independent mediators (corresponding to the parallel case, i.e. $B_M = \mathbf{0}_{p \times p}$) illustrated in Figure 2d. In Scenario 3, we consider interacted mediators such that $B_M \neq \mathbf{0}_{p \times p}$, as demonstrated in Figure 2g. For Scenario 4, we allow $p = 30$ with interacted mediators to examine the stability of our method under the high-dimensional setting. Here, $\{A, M, Y\}$ is generated from Model 1 with normal noises in Scenario 1-4. We also set $A \in \{-1, 1\}$ in Scenario 4 to show that our algorithm is capable to handle both discrete and continuous exposure, denoted as Scenario 4^*. The sample size n is chosen from $\{50, 500\}$ to be consistent with the scale of our real data analysis. More details of the simulation setting and the implementation are reported in Section C in the appendix to save space.

The averaged estimated matrix $\widehat{B}^\top$ by the ANOCE-CVAE under different settings is illustrated in Figure 2 over 100 replications. The numerical results are summarized in Table 2 (for Scenario 1 to 3)

(a) True. (b) $n = 50$ (c) 500 (d) True. (e) $n = 50$ (f) 500 (g) True. (h) $n = 50$ (i) 500

(j) True. (k) $n = 50$ (l) $n = 500$ (m) $n = 50$ (n) $n = 500$

Figure 2: The averaged estimated $\widehat{B}^\top$ under the ANOCE-CVAE: (a-c) correspond to Scenario 1; (d-f) correspond to Scenario 2; (g-i) correspond to Scenario 3; (j) is the shared true DAG for Scenario 4 and 4^*; (k-l) are results under Scenario 4; (m-n) are results under Scenario 4^*.

(a) True. (b) ANOCE (c) PC (d) ICA (e) NOTEARS (f) DAG-GNN

Figure 3: The averaged estimated $\widehat{B}^\top$ for Scenario 3 under different methods with $n = 500$.

and Table 3 (for Scenario 4 and 4^*) in the appendix, including the bias of the estimated TE, DE, IE, DM and IM for each mediator with their standard error. It can be observed that our proposed method could correctly identify most of the edges in the causal graph when $n = 500$ in almost all cases. Based on Table 2 and 3, the estimated causal effects are close to the true values as the sample size increases, indicating the good performance of our proposed method on identifying the causal effects regardless of the sparsity, the distribution of the exposure, and the dimension of mediators.

5.2 COMPARISON

We compare our approach against the PC (Spirtes et al., 2000), the ICA-LiNGAM (Shimizu et al., 2006), the NOTEARS (Zheng et al., 2018), and the DAG-GNN (Yu et al., 2019), using a moderate setting (Scenario 3). Default hyper-parameters of these implementations are used unless otherwise stated. Here, to be consistent with the NOTEARS and the DAG-GNN, we also use a graph threshold as 0.3 to prune the noise edges. The averaged estimated matrix $\widehat{B}^\top$ over 100 replications under different methods is shown in Figure 3 with $n = 500$. All the numerical results are reported in Table 4 of the appendix, including the false discovery rate, the true positive rate, and the structural Hamming distance. Figure 3 shows our algorithm performs the best among five methods, followed by the DAG-GNN. The results of the comparison study not only support the choice of the extension on the score-based method (by comparing the results of the DAG-GNN and the NOTEARS with other methods), but also validate the necessity of our method on the causal discovery with the background knowledge (by comparing the results of our method with the DAG-GNN).

6 REAL DATA ANALYSIS: COVID-19 OUTBREAK

From early Jan 2020 to late Feb 2020 (the Spring Festival period), COVID-19 spread to every province-level division of China, exacerbated by the Chinese new year migration and human to human transmission. The Chinese Government locked Hubei down on Jan 24th, which directly blocked infected people leaving from Hubei, and also indirectly control the spread of COVID-19. Here, for simplicity, we attribute the causal effects of the measures taken by cities outside of Hubei to the original and main action of interest, i.e. Hubei lockdowns. Thus, the COVID-19 example satisfies the considered causality framework for studying the causal mediator effects.

(a) The estimated weighted matrix for $B^\top$.   (b) The estimated (in)direct effects for selected cities.

Figure 4: The analysis of the causal effects of 2020 Hubei lockdowns on reducing the COVID-19 spread in China regulated by Chinese major cities outside Hubei.

We collect the data from the National Health Commission (NHC) of China and Baidu Qianxi for analysis. Specifically, let the exposure A as if Hubei is on lockdown, 0 for unlocked (before and on Jan 23rd), and 1 for locked (on and after Jan 24th). We select 30 candidate cities outside Hubei that contain most potential infected people, as mediators M. The daily migration scale index (MSI) of each city is used as the value of each mediator, which is the migration magnitude of large groups of people from one geographical area to another (Chen et al., 2020) and is comparable among cities. Lastly, we use the daily increase rate of confirmed cases out of Hubei to characterize the severity of the virus spreading with a one-week delay (due to the diagnose and incubation period of COVID-19 (Lauer et al., 2020)): $Y_t = \frac{\text{Confirmed cases out of Hubei}_{t+8} - \text{Confirmed cases out of Hubei}_{t+7}}{\text{Confirmed cases out of Hubei}_{t+7}}$. Here, the time t starts from Jan 12th to Feb 20th, 2020, since Jan 19th, 2020 is the earliest date with an available number of confirmed cases out of Hubei (to compute $Y_{t=1}$ on Jan 12th), and after Feb 20th, 2020, the pandemic was under control outside Hubei with the evidence of the work resumption in China. The final dataset yields a total of 38 records. More details of data collection can be found in Section D of the appendix.

The proposed algorithm is applied to the COVID-19 data with 100 replications by setting different random seeds in the neural network. The estimated weighted adjacency matrix is shown in Figure 4a, with the detailed ANOCE table reported in Table 5 in the appendix. The total effect of 2020 Hubei lockdowns on the daily increase rate of confirmed cases outside Hubei as -0.497, where the direct effect is -0.078 and the indirect effect is -0.419. In other words, by locking Hubei down, China successfully reduced 49.7% of the daily new cases outside Hubei; 84% of which is the indirect effect contributed via the reduced migration of cities (the mediators) out of Hubei, and the rest 16% owes to the direct effect of Hubei lockdowns since infected people were constrained in Hubei after the lockdown. Thus, the lockdown is effective in reducing the COVID-19 spread in China.

The total indirect effect of the lockdown can be further broken down by cities' direct effects (DMs, corresponds to the intensity of transmission within a particular city). We compare cities' DMs with their associated indirect effects (IMs, describes the secondary migration from a particular city to other places) in Figure 4b, where a positive effect means spreading the virus while negative means control. Note that the selected 30 cities are ordered by their cumulative MSI during the data period. ● 1). From Figure 4b, the majority of cities have a negative DM (colored in blue), which implies the infection within cities outside Hubei have been effectively controlled under the lockdown. ● 2). There are more cities with a positive IMs (red), which is in line with the intuition that the secondary migration among cities may exacerbate the pandemic. ● 3). The positive effects (red) are more likely located at the first 20 nodes, which corresponds to the cities with large MSI, while the last 10 cities with relatively small MSI are almost all blue. This accords with the migration peak among big cities during the Spring Festival period that aggravated the spread of the virus.

7   CONCLUSION

We conclude our paper with the following discussions. First, our proposed method can be extended to a non-linear model via a transformation on the outcome, and a generalized definition of the DM from a graphical perspective is given in Section F.2 of the appendix without the LSEM assumption. Second, due to possibly unmeasured confounders in our real data, such as cities' features and periodic effect, we may consider extending our model with a new topological order that contains confounders for a wider utility, such as forcing the topological order of k confounders as 1 to k followed by the exposure as $1 + k$. Third, our proposed identification constraint can be generalized to other background knowledge and can be applied to any score-based structural learning methods.

REFERENCES

Chen Avin, Ilya Shpitser, and Judea Pearl. Identifiability of path-specific effects. 2005.

Peter Bühlmann, Jonas Peters, Jan Ernest, et al. Cam: Causal additive models, high-dimensional order search and penalized regression. *The Annals of Statistics*, 42(6):2526–2556, 2014.

David Card. The causal effect of education on earnings. In *Handbook of labor economics*, volume 3, pp. 1801–1863. Elsevier, 1999.

Abhishek Chakrabortty, Preetam Nandy, and Hongzhe Li. Inference for individual mediation effects and interventional effects in sparse high-dimensional causal graphical models. *arXiv preprint arXiv:1809.10652*, 2018.

Huijie Chen, Ye Chen, Baijun Sun, Ping Wang, Lihai Wen, Zhiyong Lian, Ying Lu, Ying Qi, Shuo Zhao, Linlin Zhang, et al. Correlation between the migration scale index and the number of new confirmed novel coronavirus pneumonia cases in china. 2020.

David Maxwell Chickering. Optimal structure identification with greedy search. *Journal of machine learning research*, 3(Nov):507–554, 2002.

Miguel Angel Hernán. A definition of causal effect for epidemiological research. *Journal of Epidemiology & Community Health*, 58(4):265–271, 2004.

Miguel Ángel Hernán, Babette Brumback, and James M Robins. Marginal structural models to estimate the causal effect of zidovudine on the survival of hiv-positive men. *Epidemiology*, pp. 561–570, 2000.

Markus Kalisch and Peter Bühlmann. Estimating high-dimensional directed acyclic graphs with the pc-algorithm. *Journal of Machine Learning Research*, 8(Mar):613–636, 2007.

Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.

Sébastien Lachapelle, Philippe Brouillard, Tristan Deleu, and Simon Lacoste-Julien. Gradient-based neural dag learning. *arXiv preprint arXiv:1906.02226*, 2019.

Stephen A Lauer, Kyra H Grantz, Qifang Bi, Forrest K Jones, Qulu Zheng, Hannah R Meredith, Andrew S Azman, Nicholas G Reich, and Justin Lessler. The incubation period of coronavirus disease 2019 (covid-19) from publicly reported confirmed cases: estimation and application. *Annals of internal medicine*, 2020.

Marloes H Maathuis, Markus Kalisch, Peter Bühlmann, et al. Estimating high-dimensional intervention effects from observational data. *The Annals of Statistics*, 37(6A):3133–3164, 2009.

Preetam Nandy, Marloes H Maathuis, Thomas S Richardson, et al. Estimating the effect of joint interventions from observational data in sparse high-dimensional settings. *The Annals of Statistics*, 45(2):647–674, 2017.

Ugo Panizza and Andrea F Presbitero. Public debt and economic growth: is there a causal effect? *Journal of Macroeconomics*, 41:21–41, 2014.

Adam Paszke, Sam Gross, Soumith Chintala, Gregory Chanan, Edward Yang, Zachary DeVito, Zeming Lin, Alban Desmaison, Luca Antiga, and Adam Lerer. Automatic differentiation in pytorch. 2017.

Judea Pearl et al. Causal inference in statistics: An overview. *Statistics surveys*, 3:96–146, 2009.

Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983.

Rajen D Shah and Jonas Peters. The hardness of conditional independence testing and the generalised covariance measure. *arXiv preprint arXiv:1804.07203*, 2018.

Shohei Shimizu, Patrik O Hoyer, Aapo Hyvärinen, and Antti Kerminen. A linear non-gaussian acyclic model for causal discovery. *Journal of Machine Learning Research*, 7(Oct):2003–2030, 2006.

Pater Spirtes, Clark Glymour, Richard Scheines, Stuart Kauffman, Valerio Aimale, and Frank Wimberly. Constructing bayesian network models of gene expression networks from microarray data. 2000.

Tyler VanderWeele and Stijn Vansteelandt. Mediation analysis with multiple mediators. *Epidemiologic methods*, 2(1):95–115, 2014.

Stijn Vansteelandt and Rhian M Daniel. Interventional effects for mediation analysis with multiple mediators. *Epidemiology (Cambridge, Mass.)*, 28(2):258, 2017.

Sewall Wright. Correlation and causation. *Journal of agricultural research*, 20(7):557–585, 1921.

Yue Yu, Jie Chen, Tian Gao, and Mo Yu. Dag-gnn: Dag structure learning with graph neural networks. *arXiv preprint arXiv:1904.10098*, 2019.

Xun Zheng, Bryon Aragam, Pradeep K Ravikumar, and Eric P Xing. Dags with no tears: Continuous optimization for structure learning. In *Advances in Neural Information Processing Systems*, pp. 9472–9483, 2018.

Fei Zhou, Ting Yu, Ronghui Du, Guohui Fan, Ying Liu, Zhibo Liu, Jie Xiang, Yeming Wang, Bin Song, Xiaoying Gu, et al. Clinical course and risk factors for mortality of adult inpatients with covid-19 in wuhan, china: a retrospective cohort study. *The Lancet*, 2020.

Shengyu Zhu and Zhitang Chen. Causal discovery with reinforcement learning. *arXiv preprint arXiv:1906.04477*, 2019.

A ADDITIONAL GRAPH TERMINOLOGY

Given the node set, the weighted DAG can be uniquely determined by its weighted adjacency matrix, i.e., there is a one-to-one transformation between $\mathcal{G}$ and B. Suppose the graph nodes X in $\mathcal{G}$ are sorted in its topological order (corresponding to elementary transformation of the matrix), then the matrix B is strictly upper triangular with the diagonal elements as 0. Therefore, for an identity matrix I with the same dimension as B, $I - B^\top$ is invertible since all its diagonal elements are 1 (positive).

B ALGORITHM: ANOCE-CVAE

The first part of the ANOCE-CVAE algorithm is on learning causal DAG from the observational data in the constrained space, by minimizing the loss function in Equation 6 using blackbox stochastic optimization solvers. Here, to minimize the loss in 6 and satisfy both $h_1(B) = 0$ and $h_2(B) = 0$, we simultaneously update λ_1 and λ_2 and increase c and d to infinity, by modifying the basic technique in Yu et al. (2019), corresponding to Part One.II.A.b and Part One.II.B in Algorithm 1. The second part is to estimate causal effects in the ANOCE table from the learned causal structure, based on the results from Theorem 3.1. Here, we numerically calculate the natural indirect effect for mediator IM based on Corollary F.1 in step IV of the second part.

Algorithm 1 Analysis of Causal Effects via Constrained VAE (ANOCE-CVAE)

Global: Dataset $X = \{A, M, Y\}$, sample size n, dimension of mediators p, max iteration K,
      number of epoch H, original learning rate r_0, tolerance of constraint to zero δ,
      parameter update bound U, tuning parameters ρ and ω, and penalty terms c and d;
Local: mean and standard variance of ϵ μ_ϵ and σ_ϵ, mean and standard variance of X μ_X and σ_X,
      weights in multilayer perceptrons of encoder and decoder $\theta = \{W^{(1)}, W^{(2)}, W^{(3)}, W^{(4)}\}$,
      Lagrange multipliers λ_1 and λ_2, penalty terms c and d, $(p+2) \times (p+2)$ matrix B,
      Loss function L, old and new values of the first constraint h_1^{old} and h_1^{new},
      old and new values of the second constraint h_2^{old} and h_2^{new}, and learning rate r;
Output: estimated matrix $\widehat{B}$, total effect TE, natural direct and indirect effect
      DE and IE, and natural direct and indirect effect for mediator DM and IM.

Part One: Generate matrix $\widehat{B}$ via Constrained Variational Auto-Encoder;
  I. Initialization: $\lambda_1 \leftarrow 0$; $\lambda_2 \leftarrow 0$; $c \leftarrow 1$; $d \leftarrow 1$; $r \leftarrow r_0$; $B = \mathbf{0}_{(p+2)\times(p+2)}$; $h_1^{\text{old}} \leftarrow \infty$; $h_2^{\text{old}} \leftarrow \infty$;
  II. For step k, $k = 1, \cdots, K$:
      A. While $c \times d < U$:
          a). For epoch i, $i = 1, \cdots, H$:
             1. Build Encoder $(\mu_\epsilon, \sigma_\epsilon) \leftarrow (I_{p+2} - B^\top) MLP\{X, W^{(1)}, W^{(2)}\}$;
             2. Build Decoder $(\mu_X, \sigma_X) \leftarrow MLP\{(I_{p+2} - B^\top)^{-1}\epsilon, W^{(3)}, W^{(4)}\}$;
             3. Calculate values of constraints $h_1^{\text{new}} \leftarrow h_1(B)$ and $h_2^{\text{new}} \leftarrow h_2(B)$,
               and the loss function $L \leftarrow L_{c,d}(B, W^{(1)}, W^{(2)}, W^{(3)}, W^{(4)}, \lambda_1, \lambda_2)$;
             4. Use backward to update parameters $\{B, W^{(1)}, W^{(2)}, W^{(3)}, W^{(4)}\}$;
             5. Update learning rate r;
          b). If $h_1^{\text{new}} > \rho h_1^{\text{old}}$ and $h_2^{\text{new}} > \rho h_2^{\text{old}}$: $c \leftarrow c \times \omega$; $d \leftarrow d \times \omega$;
             Elseif $h_1^{\text{new}} > \rho h_1^{\text{old}}$ and $h_2^{\text{new}} < \rho h_2^{\text{old}}$: $c \leftarrow c \times \omega$;
             Elseif $h_1^{\text{new}} < \rho h_1^{\text{old}}$ and $h_2^{\text{new}} > \rho h_2^{\text{old}}$: $d \leftarrow d \times \omega$;
             Else: Break;
      B. $h_1^{\text{old}} \leftarrow h_1^{\text{new}}$; $h_2^{\text{old}} \leftarrow h_2^{\text{new}}$; $\lambda_1 \leftarrow \lambda_1 \times h_1^{\text{new}}$; $\lambda_2 \leftarrow \lambda_2 \times h_2^{\text{new}}$;
      C. If $h_1^{\text{new}} < \delta$ and $h_2^{\text{new}} < \delta$: Break;
  III. Output $\widehat{B} \leftarrow B$;

Algorithm 2 ANOCE-CVAE (cont.)

Part Two: Estimate causal effects in ANOCE based on matrix $\widehat{B}$;

I. According to Equation 3:
    A. Get $\widehat{\gamma}$ as the direct effect DE;
    B. Get $\widehat{\alpha}$ as the effect of A on M, $\widehat{\beta}$, and the inside matrix $\widehat{B}_M$;
II. Get $\widehat{\zeta} \equiv (I_p - B_M^\top)^{-1}\widehat{\alpha}$ that represents the causal effect of A on M;
III. Get $\widehat{\beta}^\top \widehat{\zeta}$ that represents the total natural indirect effect IE;
    For each mediator M_i, $i = 1, \cdots, p$: Define the natural direct effect for M_i as $DM[i] = \widehat{\alpha}[i]\widehat{\zeta}[i]$;
IV. Get the natural indirect effect for mediator:
    For each mediator M_i, $i = 1, \cdots, p$:
    A. Delete M_i from the matrix $\widehat{B}$ and get $\widehat{B_i'}$;
    B. Repeat step II. with reduced matrix $\widehat{B_i'}$ and get $\widehat{\beta'}$ and $\widehat{\zeta'}$;
    C. Calculate the effect difference as the total mediation effect $\widehat{\beta}^\top \widehat{\zeta} - \widehat{\beta'}^\top \widehat{\zeta'}$
    D. Define the natural indirect effect for M_i as $IM[i] = \{\widehat{\beta}^\top \widehat{\zeta} - \widehat{\beta'}^\top \widehat{\zeta'}\} - DM[i]$;
V. Define the total effect $TE = \widehat{\gamma} + \widehat{\beta}^\top \widehat{\zeta}$.

C   Additional Simulation Studies

In this section, we give more details on simulation studies to investigate the finite sample performance of the proposed method for learning causal effects with multiple mediators, in comparison to the popular causal discovery methods, including the PC, the ICA-LiNGAM, the NOTEARS, and the DAG-GNN. The computing infrastructure used is a virtual machine in the compute engine of Google Cloud Platform with 8 processor cores and 32GB memory. The average runtime for each result is around 1 to 2 hours.

C.1   Data Generation

We first generate a random DAG by the Erdős-Reńyi model with the expected node degree as 2, and then remove all in-edges (from precedent nodes) of the first node as A; and remove all out-edges (from descendent nodes) of the last node as Y; then the remained nodes are the mediators M. Edges in DAGs for all scenarios are randomly assigned with weights ($w \in \{-1, 1\}$ with equal probability) to obtain the weighted adjacency matrix B. The experiment data sets are generated from Model 3, where the error variables in $\epsilon \equiv [\epsilon_A, \epsilon_M^\top, \epsilon_Y]^\top$ independently follow a normal distribution with mean 0 and noise 0.5. Here, we add a baseline of 1.0 on the outcome Y. Note that the gaussian exposure in Scenario 4 and the binary exposure Scenario 4* have the same mean and noise and thus their results are comparable.

C.2   Implementation of ANOCE-CVAE

For each setting, we apply the proposed method with 100 replications. Here, the ANOCE-CVAE is implemented based on PyTorch (Paszke et al., 2017), using Adam (Kingma & Ba, 2014) to minimize the loss function in Equation 6. We set the batch size as 25 for $n = 50$ and 100 for $n = 500$ with hidden nodes as p^2, the initial learning rate as 0.003 with an update rule as $r \leftarrow r/\{log(c) + log(d) + 0.01\}$ where c and d are penalty terms for two constraints, and the parameter update bound as $U = 10^{20}$, for all settings. Following the recommendation of Yu et al. (2019), we find that tuning parameters $\rho = 0.25$ and $\omega = 10$ also work well in our settings, and we adopt the Huber-norm regularization of B for a better convergence. Here, the variational posterior and the likelihood are parameterized as Gaussian with unit noise to approximate the underlying true model.

Table 2: The bias and standard error of the estimated causal effects for Scenario 1 to 3 ($p = 10$).

Results	Scen. 1 (simplest)		Scen. 2 (parallel)		Scen. 3 (interacted)	
	$n = 50$	$n = 500$	$n = 50$	$n = 500$	$n = 50$	$n = 500$
TE	0.19 (0.02)	0.01 (0.00)	0.66 (0.05)	0.03 (0.02)	0.15 (0.05)	0.05 (0.03)
DE	0.23 (0.02)	0.01 (0.00)	0.78 (0.02)	0.12 (0.06)	0.78 (0.03)	0.15 (0.06)
IE	0.04 (0.01)	0.00 (0.00)	1.44 (0.04)	0.08 (0.06)	0.63 (0.04)	0.20 (0.04)
DM_1	0.01 (0.00)	0.00 (0.00)	0.69 (0.03)	0.02 (0.01)	0.48 (0.04)	0.06 (0.02)
DM_2	0.00 (0.00)	0.00 (0.00)	0.43 (0.03)	0.01 (0.01)	0.00 (0.00)	0.00 (0.00)
DM_3	0.00 (0.00)	0.00 (0.00)	0.65 (0.02)	0.01 (0.02)	0.00 (0.00)	0.00 (0.00)
DM_4	0.00 (0.00)	0.00 (0.00)	0.65 (0.03)	0.02 (0.01)	0.20 (0.02)	0.06 (0.01)
DM_5	0.01 (0.00)	0.00 (0.00)	0.68 (0.03)	0.02 (0.01)	0.44 (0.02)	0.10 (0.01)
DM_6	0.01 (0.00)	0.00 (0.00)	0.67 (0.02)	0.02 (0.02)	0.27 (0.04)	0.03 (0.01)
DM_7	0.00 (0.00)	0.00 (0.00)	0.62 (0.03)	0.03 (0.01)	0.67 (0.03)	0.16 (0.03)
DM_8	0.00 (0.00)	0.00 (0.00)	0.66 (0.03)	0.01 (0.01)	0.59 (0.03)	0.14 (0.03)
DM_9	0.00 (0.00)	0.00 (0.00)	0.65 (0.02)	0.01 (0.01)	0.00 (0.00)	0.00 (0.00)
DM_{10}	0.01 (0.00)	0.00 (0.00)	0.66 (0.03)	0.01 (0.01)	0.10 (0.02)	0.05 (0.01)
IM_1	0.00 (0.00)	0.00 (0.00)	0.08 (0.01)	0.08 (0.01)	0.76 (0.03)	0.16 (0.04)
IM_2	0.00 (0.00)	0.00 (0.00)	0.09 (0.01)	0.01 (0.01)	0.00 (0.00)	0.00 (0.00)
IM_3	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)	0.03 (0.01)	0.00 (0.00)	0.00 (0.00)
IM_4	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)	0.05 (0.01)	0.16 (0.01)	0.03 (0.01)
IM_5	0.00 (0.00)	0.00 (0.00)	0.06 (0.01)	0.12 (0.01)	0.02 (0.01)	0.03 (0.00)
IM_6	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)	0.04 (0.01)	0.66 (0.02)	0.09 (0.02)
IM_7	0.00 (0.00)	0.00 (0.00)	0.09 (0.01)	0.00 (0.01)	0.05 (0.02)	0.04 (0.01)
IM_8	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)	0.02 (0.01)	0.11(0.02)	0.06 (0.01)
IM_9	0.00 (0.00)	0.00 (0.00)	0.03 (0.01)	0.01 (0.01)	0.00 (0.00)	0.00 (0.00)
IM_{10}	0.00 (0.00)	0.00 (0.00)	0.07 (0.01)	0.08 (0.01)	0.03 (0.01)	0.00 (0.00)

C.3 ADDITIONAL RESULTS OF ANOCE-CVAE

The results are summarized in Table 2 (for Scenario 1 to 3) and Table 3 (for Scenario 4 and 4^*), including the bias of the estimated TE, DE, IE, DM and IM for each mediator with their standard error, over 100 replications. Note that due to limited space, we save the numerical results of the IM in Table 3. From the results in Table 2, it is clear that the estimated TE, DE, IE, DM and IM for each mediator are close to the true values as the sample size increases in Scenario 1 to 3, which indicates the good performance of our proposed method on identifying the causal effects regardless of the sparsity. With the expected node degree increasing, one can observe a slightly larger bias and standard error of the estimated causal effects as expected, as shown in Table 2. Based on Table 3, the results of Scenario 4 and Scenario 4^* are merely identical under different sample sizes, indicating our proposed method can handle either discrete or continuous exposure. In addition, by comparing the results of Scenario 3 and 4 where we fix the expected node degree as 2, one can observe a slightly larger bias of the estimated causal effects but of a similar small scale, as the dimension of mediators p increases, which implies the stability of our method under the high-dimensional setting.

Table 3: The bias and standard error of the estimated causal effects for Scenario 4 and 4^* ($p = 30$).

Results	Scen. 4 (continuous)		Scen. 4^* (binary)	
	$n = 50$	$n = 500$	$n = 50$	$n = 500$
TE	0.29 (0.04)	0.20 (0.02)	0.09 (0.03)	0.05 (0.01)
DE	0.90 (0.01)	0.28 (0.02)	0.85 (0.01)	0.25 (0.02)
IE	1.19 (0.04)	0.48 (0.03)	0.94 (0.03)	0.30 (0.02)
DM_1	0.12 (0.02)	0.08 (0.01)	0.18 (0.02)	0.07 (0.01)
DM_2	0.61 (0.02)	0.02 (0.01)	0.45 (0.02)	0.02 (0.01)
DM_3	0.00 (0.01)	0.03 (0.01)	0.01 (0.01)	0.02 (0.01)
DM_4	0.31 (0.03)	0.25 (0.02)	0.09 (0.02)	0.07 (0.01)
DM_5	0.01 (0.01)	0.07 (0.01)	0.05 (0.03)	0.10 (0.02)
DM_6	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_7	0.37 (0.02)	0.19 (0.02)	0.60 (0.04)	0.18 (0.02)
DM_8	0.02 (0.01)	0.00 (0.01)	0.05 (0.01)	0.02 (0.02)
DM_9	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{10}	0.02 (0.01)	0.09 (0.01)	0.14 (0.01)	0.15 (0.01)
DM_{11}	0.00 (0.01)	0.00 (0.00)	0.01 (0.01)	0.00 (0.00)
DM_{12}	0.02 (0.00)	0.01 (0.01)	0.06 (0.01)	0.01 (0.01)
DM_{13}	0.25 (0.02)	0.11 (0.02)	0.51 (0.04)	0.16 (0.03)
DM_{14}	0.02 (0.01)	0.10 (0.01)	0.09 (0.02)	0.16 (0.01)
DM_{15}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{16}	0.09 (0.01)	0.10 (0.02)	0.18 (0.03)	0.13 (0.02)
DM_{17}	0.03 (0.01)	0.07 (0.02)	0.04 (0.03)	0.06 (0.03)
DM_{18}	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{19}	0.02 (0.00)	0.01 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{20}	0.01 (0.00)	0.00 (0.00)	0.08 (0.01)	0.01 (0.01)
DM_{21}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{22}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{23}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{24}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{25}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{26}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{27}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{28}	0.01 (0.01)	0.02 (0.01)	0.01 (0.01)	0.01 (0.00)
DM_{29}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)
DM_{30}	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)	0.00 (0.00)

| (a) True. | (b) ANOCE | (c) PC | (d) ICA | (e) NOTEARS | (f) DAG-GNN |

Figure 5: The averaged estimated $\widehat{B}^{\top}$ for Scenario 3 under different methods with $n = 50$.

| (a) True. | (b) ANOCE | (c) PC | (d) ICA | (e) NOTEARS | (f) DAG-GNN |

Figure 6: The averaged estimated $\widehat{B}^{\top}$ for Scenario 3^* under different methods with $n = 50$.

| (a) True. | (b) ANOCE | (c) PC | (d) ICA | (e) NOTEARS | (f) DAG-GNN |

Figure 7: The averaged estimated $\widehat{B}^{\top}$ for Scenario 3^* under different methods with $n = 500$.

C.4 ADDITIONAL COMPARISON STUDIES

This section provides more results on comparison studies against the existing methods. Here, we also include a binary setting by replacing the gaussian exposure in Scenario 3 with the binary exposure as Scenario 3^*. Default hyper-parameters of these implementations are used unless otherwise stated. The estimated graphs (after pruning) are evaluated by three metrics: the false discovery rate (FDR), the true positive rate (TPR), and the structural Hamming distance (SHD, the smallest number of edge additions, deletions, and reversals to convert the estimated graph into the true DAG). Here, the SHD takes into account both false positives and negatives and a lower SHD indicates a better estimate of the causal graph. We illustrate the averaged estimated matrix $\widehat{B}^{\top}$ (with a graph threshold as 0.3) over 100 replications under different methods in Figure 5 for Scenario 3 with $n = 50$, 6 for Scenario 3^* with $n = 50$, and 7 for Scenario 3^* with $n = 500$. Note the results of Scenario 3 with $n = 500$ is reported in Figure 3. The FDR, TPR, and SHD of the averaged estimated matrix $\widehat{B}^{\top}$ with their standard deviation are reported in Table 4 over 100 replications under different methods.

From Table 4, it is clear that our algorithm performs best among five methods, followed by the DAG-GNN and the NOTEARS. Under the ANOCE-CVAE, the FDR and SHD for the averaged estimated matrix $\widehat{B}^{\top}$ are close to 0 and the TPR approaches to 1 in all scenarios, when $n = 500$. Another support is illustrated in Figure 3 and 7, where the averaged estimated matrix $\widehat{B}^{\top}$ under the ANOCE-CVAE is approximately the same as the ground true graph $B^{\top}$ in all scenarios when $n = 500$. However, the traditional methods (the PC and the ICA-LiNGAM) can hardly recognize the true causal pattern even under $n = 500$. In addition, it can be observed that under the ANOCE-CVAE, the first row and the last column of the estimated $\widehat{B}^{\top}$ are all zeros with different sample sizes, which corresponds to our identification constraint successfully satisfied; while other methods fail to identify the treatment and the outcome when the sample size is small from Figure 5 and 6.

Table 4: The empirical results for Scenario 3 and 3* under different causal discovery methods.

Methods		Scen. 3 (continuous)		Scen. 3* (binary)	
		$n = 50$	$n = 500$	$n = 50$	$n = 500$
ANOCE-CVAE	FDR	0.05 (0.07)	0.00 (0.09)	0.05 (0.10)	0.08 (0.12)
	TPR	0.78 (0.14)	1.00 (0.08)	0.78 (0.12)	1.00 (0.10)
	SHD	6.0 (3.3)	0.0 (3.6)	6.0 (3.6)	2.0 (5.0)
PC	FDR	0.47 (0.08)	0.48 (0.08)	0.36 (0.08)	0.44 (0.10)
	TPR	0.43 (0.07)	0.70 (0.05)	0.61 (0.06)	0.65 (0.07)
	SHD	18.0 (2.5)	17.0 (3.2)	13.0 (2.4)	13.0 (2.7)
ICA-LiNGAM	FDR	1.00 (0.15)	0.92 (0.11)	0.86 (0.09)	1.00 (0.07)
	TPR	0.00 (0.06)	0.04 (0.06)	0.04 (0.04)	0.00 (0.05)
	SHD	23.0 (2.2)	23.0 (2.0)	22.0 (1.6)	26.0 (2.0)
NOTEARS	FDR	0.00 (0.10)	0.10 (0.05)	0.09 (0.11)	0.20 (0.06)
	TPR	0.52 (0.08)	0.39 (0.06)	0.43 (0.07)	0.35 (0.05)
	SHD	11.0 (2.2)	14.0 (1.3)	13.0 (2.1)	15.0 (1.1)
DAG-GNN	FDR	0.21 (0.10)	0.15 (0.06)	0.29 (0.09)	0.15 (0.06)
	TPR	0.65 (0.13)	0.74 (0.06)	0.43 (0.13)	0.74 (0.08)
	SHD	12.0 (3.8)	9.0 (2.2)	17.0 (3.3)	9.0 (2.1)

Figure 8: A direct acyclic graph illustrating the temporal causal relationship of the spreading network of the coronavirus outbreak under 2020 Hubei lockdowns in China.

D ADDITIONAL REAL DATA RESULTS

In this section, we provide additional real data analysis on the COVID-19.

D.1 DATA COLLECTION

To better characterize the causality of the virus spreading under the Hubei lockdowns in China, we assume: 1) Hubei was the centre of the COVID-19 outbreak in China (Zhou et al., 2020); 2) the decreased migration outside Hubei was largely stimulated by the lockdown; 3) individual who departed to one destination would not return to the original departure due to travel restrictions in China. Under the above assumptions, it is reasonable to use a temporal causal relationship to describe the spread of COVID-19 under 2020 Hubei lockdowns as in Figure 8.

Next, we give more details on how to present components in Figure 8 with appropriate variables. First, we set the exposure A as if Hubei is on lockdown, 0 for unlocked (before and on Jan 23rd) and 1 for locked (on and after Jan 24th). To select the candidate cities that contain most potential infected people, we rank cities outside Hubei by their received Wuhan (identified over 60% cases in China reported by the NHC) migration between Jan 1st, 2020 and Jan 22nd, 2020 (before the lockdown), and choose the top 30 cities (account for 69.17% of total Wuhan migration) as mediators M, to control the noise. We use the daily migration scale index (MSI) of each city as the value of each mediator, defined in Baidu Qianxi to describe the migration magnitude. By noticing the following facts: 1) it took usually 2 days to diagnose the COVID-19; 2) the estimated median incubation period is 5 days (Lauer et al., 2020); the outcome of interest Y is defined as the daily increase rate of confirmed cases out of Hubei with a one-week (7=2+5) delay. We delete extreme data points (two outbreaks at jails), and the final dataset yields a total of 38 records.

(b) The partial heat map of correlation matrix between cities and the treatment A or the outcome Y.

(a) The cumulative confirmed cases of selected cities.

Figure 9: Additional figures for the real data analysis of the COVID-19 spread.

D.2 INITIAL DATA ANALYSIS

Figure 9a demonstrates the location of selected cities on the Chinese map with the color representing its cumulative confirmed cases by March 1st, 2020. It can be observed that the selected mediators are either located around Hubei province or are big central cities such as Beijing that has a large population and migration scale. In addition, we provide the partial heat map of the correlation matrix between cities and the treatment A or the outcome Y as illustrated in Figure 9b. Here, one could observe that cities' MSIs are highly positive correlated with the lockdown while are highly negative correlated with the daily increase rate of confirmed cases.

D.3 ADDITIONAL RESULTS

Table 5 lists all the numerical facts of the selected cities, including the population (million), the cumulative migration scale index (MSI) during the data period (Jan 12th to Feb 20th, 2020), the ratio of received Wuhan migration between Jan 1st to Jan 22nd, 2020 (before the lockdown), the cumulative confirmed cases by March 1st, 2020, and cities' direct (DM) and indirect effect (IM). Note the selected 30 cities in Table 5 are sorted according to their cumulative MSI, and its order is used as the order of the mediators in this paper. It can be seen from Table 5 that the population, the cumulative MSI, the ratio of received Wuhan migration, and the confirmed cases are highly correlated among selected cities as expected.

Besides the general causal pattern of cities' estimated DMs and IMs stated in the main text, we provide more interpretation on the level of the individual city. Here, we compare three groups of cities that are of similar scale (population and migration) or geographic position, to specify our results. First, we compare the results between Beijing and Shanghai, where both cities have comparable population scale and are the center of politic or economic in China (the correlation between the MSI of Beijing and the outcome yields the same value as of Shanghai as 0.50). It can be seen from Table 5 that Beijing and Shanghai have similar positive indirect effects as 0.247 and 0.235, respectively, while Shanghai has a slightly higher direct effect on controlling the virus as -0.069, which is possibly due to the smaller MSI of Shanghai. Second, we compare three cities located at Guangdong province (see Figure 10a and 10b), including Shenzhen, Guangzhou, and Dongguan. All three cities have positive effects, among which Guangzhou yields the highest direct effect as 0.847, followed by Dongguan. The effect size of these cities agrees with their correlation coefficients with the outcome, where the correlation between Guangzhou and Y is 0.61, followed by Dongguan as 0.58 and Shenzhen as 0.50. The last comparison is among cities in southeastern China (see Figure 10a and 10b), including Suzhou, Hangzhou, and Wenzhou, all of which have a negative direct effect on the virus control and a positive indirect effect for the virus spread. Here, Wenzhou achieves the largest absolute value of the negative direct effect as -0.650, which conforms to its strict local shelter in home order after the Hubei lockdowns, where the correlation between Wenzhou and the outcome achieves the highest value as 0.83.

Table 5: Analysis of causal effects of 2020 Hubei lockdowns on reducing the COVID-19 spread in China regulated by Chinese major cities outside Hubei.

	Source				$d.f.$	Causal effects	
	Direct effect from the Hubei lockdowns				1	$DE = -0.078$	
	Inirect effect via Chinese major cities outside Hubei				$p = 30$	$IE = -0.419$	
Index	City name	Population	Ratio of Wuhan	Confirmed Cases	Cumulative MSI	DM	IM
1	Shenzhen	10.36	3.02%	418	275.42	1	0.021	0.002
2	Guangzhou	12.70	3.22%	346	270.38	1	0.847	0.176
3	Beijing	23.00	5.50%	414	260.02	1	-0.003	0.247
4	Chengdu	14.05	2.57%	143	233.62	1	-0.201	-0.004
5	Shanghai	23.02	4.09%	338	232.80	1	-0.069	0.235
6	Dongguan	8.22	1.14%	99	211.99	1	0.257	0.003
7	Suzhou	10.47	1.42%	87	171.46	1	-0.481	0.553
8	Xian	10.00	1.87%	120	136.56	1	0.015	-0.089
9	Hangzhou	8.70	1.77%	169	135.73	1	-0.335	0.144
10	Zhengzhou	10.14	3.04%	157	132.58	1	-0.019	0.309
11	Chongqing	30.17	4.97%	576	128.98	1	0.313	0.050
12	Changsha	7.04	5.21%	242	126.78	1	0.399	0.222
13	Nanjing	8.00	1.78%	93	103.87	1	-0.210	-0.100
14	Kunming	6.43	1.24%	53	95.35	1	0.186	-0.075
15	Tianjin	12.94	1.02%	136	90.50	1	-0.092	-0.372
16	Hefei	5.70	2.11%	174	87.28	1	-0.182	0.249
17	Nanning	6.66	1.01%	55	76.63	1	0.114	0.152
18	Wenzhou	9.12	1.02%	504	60.38	1	-0.650	0.200
19	Nanchang	5.04	2.15%	230	46.95	1	-0.109	0.095
20	Zhoukou	8.95	1.50%	76	42.01	1	0.045	-0.043
21	Fuyang	7.60	1.25%	155	39.26	1	0.031	-0.023
22	Shangqiu	7.36	1.12%	91	35.48	1	0.030	-0.057
23	Yueyang	5.48	2.31%	156	29.56	1	0.006	-0.050
24	Zhumadian	7.23	2.34%	139	28.86	1	-0.070	-0.075
25	Changde	5.72	1.05%	82	28.49	1	0.002	-0.003
26	Nanyang	10.26	1.91%	156	27.71	1	-0.039	-0.106
27	Yichun	5.42	0.90%	106	24.91	1	-0.051	-0.085
28	Xinyang	6.11	5.00%	274	24.78	1	-0.080	-0.027
29	Anqing	5.31	1.57%	83	23.61	1	-0.022	-0.012
30	Jiujiang	4.73	2.06%	118	22.29	1	-0.072	-0.047
/	Selected cities	295.95	69.17%	5790	3204.24	30	-0.419	/
	Total					$1 + p = 31$	$TE = -0.497$	

We summarize all different sources of causal effects in Table 5 as the ANOCE table of 2020 Hubei lockdowns on reducing the COVID-19 spread in China. Note that due to cities' different levels of control measures on the coronavirus outside Hubei as well as other possible confounders, there are some inconsistency between cities' DMs and their cumulative MSIs. We leave the extension with confounders for further investigation.

One can refer to Figure 10a and 10b for cities' DMs and IMs on the Chinese map. To check the reasonability of our results, we plot the estimated weighted matrix in Figure 4a, where the first node (indexed 0) represents the Hubei lockdowns, the last node (indexed 31) is the daily increase rate out of Hubei, and the middle 30 nodes (indexed 1-30) correspond to 30 selected cities in Figure 4b. From Figure 4a, we can observe: ● 1). The color of the first column is almost all blue, indicating that locking Hubei down can reduce the migration of selected cities; ● 2). An approximate red upper

(a) The estimated direct effect of cities (DMs).

(b) The estimated indirect effect of cities (IMs).

(c) The estimated DAG for the coronavirus data.

(d) The spreading network among cities that received most of the Wuhan migration during the data period ($2.57\% \sim 5.50\%$, presented by the color of nodes).

Figure 10: Additional analysis of the causal effects of 2020 Hubei lockdowns on reducing the COVID-19 spread in China regulated by Chinese major cities outside Hubei.

triangular among first 20 nodes implies a migration trend with positive effects among central cities with large MSI, i.e. relatively smaller cities tend to have positive effects on other cities with relatively larger MSI; • 3). An approximate red lower triangular among last 10 nodes indicates a weaker migration trend with an opposite direction among non-central cities with small MSI; • 4). There is also an almost all blue rectangle in the right top of the estimated matrix, showing non-central cities tend to have negative effects on central cities. All the above findings are accords with the migration trend during the Spring Festival and intensive mutual communications among central cities in China.

In addition, we provide the estimated DAG for the coronavirus data in Figure 10c as the complete spreading network among major cities outside Hubei, where the exposure A, mediators M (cities' corresponding index can be found in Table 5), and the outcome Y are colored in red, blue and green, respectively. It can be observed that the in-degree is larger than the out-degree for nodes with small index, while an opposite rule is applied for nodes with large index. This finding is consistent with the migration trend identified in our main text. Lastly, we give the spreading network among cities that received most Wuhan migration during the data period, including Beijing, Shanghai, Guangzhou, Shenzhen, Chengdu, Chongqing, Zhengzhou, Changsha, and Xinyang, plus Wuhan, in Figure 10d, to illustrate the partial interaction trend among cities.

E   TOY EXAMPLE

Example E.1 *Here, we use a toy example of a weighted DAG $\mathcal{G}^{toy}$ under the LSEM given in Figure 11 to better demonstrate our definitions.*

In $\mathcal{G}^{toy}$, $p = 4$ mediators are included, where $M_1 \leftarrow 0.2A + \epsilon_1$, $M_2 \leftarrow 0.2M_1 + 0.5M_4 + \epsilon_2$, $M_3 \leftarrow 0.4A + \epsilon_3$, $M_4 \leftarrow 0.5M_3 + \epsilon_4$, $Y = 0.5A + 1.1M_2 + 0.7M_3$. There are 4 directed path from A to Y:

Figure 11: A weighted causal DAG $\mathcal{G}^{toy}$.

1) '$A \to Y$' with length 1;
2) '$A \to M_3 \to Y$' with length 2;
3) '$A \to M_1 \to M_2 \to Y$' with length 3;
4) '$A \to M_3 \to M_4 \to M_2 \to Y$' with length 4.

Since there exists 3 directed path $\pi^* \in \{\pi_{AY}(\mathcal{G}^{toy})\}$ such that the length of π^* is larger than 2, we have the mediators in $\mathcal{G}^{toy}$ are interacted.

From the weighted DAG $\mathcal{G}^{toy}$, we have the direct effect of A on Y is $\gamma^{toy} = 0.5$, $\boldsymbol{\alpha}^{toy} \equiv [0.5, 0, 0.4, 0]^\top$, $\boldsymbol{\beta}^{toy} \equiv [0, 1.1, 0.7, 0]^\top$, and

$$B_M^{\top \, toy} = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0.2 & 0 & 0 & 0.5 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0.5 \end{bmatrix}.$$

Note one may recover $\mathcal{G}^{toy}$ from the weight matrix B^{toy} as long as the order of the vertices in B^{toy} is given. Then, we have

$$(I_p - B_M^{\top \, toy})^{-1} = \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.2 & 1 & 0.25 & 0.5 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0.5 & 1 \end{bmatrix}.$$

Thus, the indirect effect of A on Y is

$$\begin{aligned} &\boldsymbol{\beta}^{toy\,\top}(I_p - B_M^{\top \, toy})^{-1}\boldsymbol{\alpha}^{toy} \\ =&[0, 1.1, 0.7, 0] \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0.2 & 1 & 0.25 & 0.5 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0.5 & 1 \end{bmatrix} [0.5, 0, 0.4, 0]^\top \\ =&[0, 1.1, 0.7, 0][0.5, 0.2, 0.4, 0.2]^\top \\ =&0 + 0.22 + 0.28 + 0 = 0.5. \end{aligned} \tag{E.1}$$

From the Equation E.1 and Theorem 3.1, we have the DM of M_2 and M_3 is 0.22 and 0.28, respectively, while other DM are 0. Note that there is no explicit expression of the natural indirect effect through the directed path from A to Y (IM) due to the complex interaction among mediators, while we provide its theoretical form in Equation G.6 based on the path method in Wright (1921) and Nandy et al. (2017). Specifically, for each directed path from A to Y, we have:

1) the direct effect through '$A \to Y$' (DE): 0.5;
2) the effect of path '$A \to M_3 \to Y$': $0.4 \times 0.7 = 0.28$;
3) the effect of path '$A \to M_1 \to M_2 \to Y$': $0.5 \times 0.2 \times 1.1 = 0.11$;
4) the effect of path '$A \to M_3 \to M_4 \to M_2 \to Y$': $0.4 \times 0.5 \times 0.5 \times 1.1 = 0.11$;
5) so the indirect effect of A on Y (IE) is $0.11 + 0.11 + 0.28 = 0.5$;
6) and the total effect of A on Y (TE) is $0.5 + 0.5 = 1.0$;
7) the indirect effect for M_1 (IM_1) corresponds to the effect of path '$A \to M_1 \to M_2 \to Y$', thus is

0.11;

8) the indirect effect for M_2 (IM_2) is zero since there is no path first goes through M_2 followed by other mediators;

9) the indirect effect for M_3 (IM_3) corresponds to the effect of path '$A \to M_3 \to M_4 \to M_2 \to Y$', thus is 0.11;

10) the indirect effect for M_4 (IM_4) corresponds to the effect of path '$A \to M_3 \to M_4 \to M_2 \to Y$', thus is 0.11.

We can calculate the last edge-specific effect directly from its definition. Since there is no $M_1 \to Y$ and $M_4 \to Y$ in $\mathcal{G}^{toy}$, we have $LE_1^{toy} = LE_4^{toy} = 0$. By deleting $M_2 \to Y$ in $\mathcal{G}^{toy}$, the total effect reduced by $0.11 + 0.11 = 0.22$, so $LE_2^{toy} = 0.22$; similarly, after deleting $M_3 \to Y$, the total effect decreases 0.28, thus $LE_3^{toy} = 0.28$. One may notice that the last edge-specific effects are equal to the DMs, and the additive of the last edge-specific effects is exact the last step of calculation $\boldsymbol{\beta}^{toy\top}(I_p - B_M^{\top\ toy})^{-1}\boldsymbol{\alpha}^{toy}$ in Equation E.1.

F CONNECTION TO LITERATURE

We establish the connection between our proposed method to the literature from three different angles. First, we show that the individual mediation effect defined in Chakrabortty et al. (2018) can be decomposed into our defined DM and IM. Next, we give an equivalent definition of the DM through a type of special edge (last edge) in the causal graph. Lastly, we prove that the proposed DM is consistent with the interventional effect via a particular mediator defined in Vansteelandt & Daniel (2017) under the LSEM.

F.1 FROM INDIVIDUAL MEDIATION VIEWPOINT

Chakrabortty et al. (2018) defined the individual mediation effect under the LSEM as follows.

Definition F.1 *(Chakrabortty et al., 2018) Individual mediation effect for M_i:*

$$
\begin{aligned}
\eta_i = &\Big[E\{M_i|do(A = a + 1)\} - E\{M_i|do(A = a)\} \Big] \\
&\times \Big[E\{Y|do(M_i = m_i + 1)\} - E\{Y|do(M_i = m_i)\} \Big].
\end{aligned}
\tag{F.1}
$$

In the following theorem, we show that the summation of η_i is strictly larger than the IE if the mediators are not parallel. The proof is given in Section G.2.

Theorem F.1 *If there exists at least one directed path $\pi^* \in \{\pi_{AY}(\mathcal{G})\}$ such that the length of π^* is larger than 2, and the element in B is nonnegative, then*

$$
\sum \eta_i > IE.
\tag{F.2}
$$

Remark F.1 *From the above theorem, it is clear that the mediator effect defined in Chakrabortty et al. (2018) is not appropriate for interpreting the decomposition of the indirect effect, when there exists interaction among mediators (a common situation as described in the introduction). Here, we keep the condition that the element in B is nonnegative, as the multiple count mediation effects in Chakrabortty et al. (2018) may cancel out in some cases and their summation would equal to the IE by chance.*

Inspired by the proof of Theorem F.1, the mediator effect η_i can be decomposed into two parts, the natural direct and indirect effect for i-th mediator, as shown in the following corollary.

Corollary F.1 *Under assumptions (A1-A3) and Model 1, we have*

$$
\eta_i = DM_i + IM_i.
\tag{F.3}
$$

Remark F.2 *Corollary F.1 together with the definition of η_i in Chakrabortty et al. (2018) provides a feasible way to numerically calculate the natural indirect effect IM_i. Specifically, by deleting the mediator M_i in the causal graph, the reduced treatment effect corresponds to η_i, then $IM_i = \eta_i - DM_i$, where the explicit expression of the DM_i is provided in Theorem 3.1. See more implementation details in Section B.*

F.2 FROM GRAPHICAL PERSPECTIVE

Next, we give the definition of the edge-specific effect following Avin et al. (2005). Suppose a directed edge of interest as $X_i \to X_j$ in a weighted DAG $\mathcal{G}$. Define a new weighted DAG $\mathcal{G}'_{i,j}$ by deleting the directed edge $X_i \to X_j$ in $\mathcal{G}$, i.e. $\mathcal{G}'_{i,j} \equiv \mathcal{G} \setminus (X_i \to X_j)$.

Definition F.2 *(Avin et al., 2005) Edge-specific effect:*

$$ET(X_i, X_j) = TE_{\mathcal{G}} - TE_{\mathcal{G}'_{i,j}}, \tag{F.4}$$

where $TE_{\mathcal{G}}$ means the total effect in graph $\mathcal{G}$.

We next give an equivalent definition of our proposed DM from a graphical perspective. Let the edge in $\mathcal{G}$ that starts with i-th mediator and ends with node Y, i.e. $M_i \to Y$, as the i-th last edge. Denote the graph $\mathcal{G}$ deleting the i-th last edge ($M_i \to Y$) as $\mathcal{G}'_i$. We define the ith last edge-specific effect as

Definition F.3 *Last edge-specific effect for M_i:*

$$LE_i = \begin{cases} TE_{\mathcal{G}} - TE_{\mathcal{G}'_i}, & \text{if there exists edge } M_i \to Y \text{ in } \mathcal{G}; \\ 0, & \text{otherwise}. \end{cases} \tag{F.5}$$

By Theorem 3.1, we have $(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}$ is the causal effect of A on M. Let $\zeta \equiv (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}$, with its i-th element $\zeta_i \equiv \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i$. Next, we show that the i-th last edge-specific effect can be presented as $\boldsymbol{\beta}_i \zeta_i$ under the LSEM in the following theorem, where $\boldsymbol{\beta}_i$ is the i-th element of the vector $\boldsymbol{\beta}$ and corresponds to the weight of the edge $M_i \to Y$. The proof can be found in Section G.3.

Theorem F.2 *Under assumptions (A1-A3) and Model 1, we have*

$$LE_i = \boldsymbol{\beta}_i \zeta_i. \tag{F.6}$$

Based on Theorem 3.1, the natural direct effect of M_i on Y can be expressed as $DM_i = \boldsymbol{\beta}_i \zeta_i$. Thus, with the result of Theorem F.2, it is easy to show the following corollary.

Corollary F.2 *Under assumptions (A1-A3) and Model 1, the natural direct effect of M_i is equal to the i-th last edge-specific effect:*

$$LE_i = DM_i = \boldsymbol{\beta}_i \zeta_i. \tag{F.7}$$

Remark F.3 *Here, both definitions describe the direct impact of one mediator M_i on the outcome. The natural direct effect of a particular mediator M_i can be understood as the influence when removing the direct edge between M_i and Y. Thus, we have the equivalence between two definitions.*

Then, we can decompose the total natural indirect effect into p last edge-specific effects or p DMs as the following additive form, based on Theorem 3.2 and Corollary F.2.

Corollary F.3 *Under assumptions (A1-A3) and Model 1, we have*

$$IE = \boldsymbol{\beta}^\top \zeta = \sum_{i=1}^p \boldsymbol{\beta}_i \zeta_i = \sum_{i=1}^p DM_i = \sum_{i=1}^p LE_i. \tag{F.8}$$

In fact, based on the uniqueness of each last edge, the natural indirect effect can be decomposed into p last edge-specific effect regardless of the LSEM setting through the graphical perspective. We give the following intuitive conclusion. The proof can be found in Section G.4.

Theorem F.3 *The IE can be decomposed through LEs as:*

$$IE = \sum_{i=1}^p LE_i. \tag{F.9}$$

Remark F.4 *One can view the last edge-specific effect as the generalized definition of the natural direct effect for mediator without the LSEM assumption.*

F.3 FROM INTERVENTIONAL EFFECT LEVEL

Finally, we show the consistency of our defined DM to the interventional effect via a particular mediator defined in Vansteelandt & Daniel (2017) under the LSEM.

Definition F.4 *(Vansteelandt & Daniel, 2017) Under assumptions (A1-A3), the interventional effect via M_i is*

$$\xi_i = \sum_{m_1 \in \mathbb{M}_1} \cdots \sum_{m_p \in \mathbb{M}_p} \left[E(Y|A=a, M_i=m_i, \Omega_i=o_i)P(\Omega_i=o_i|A=a) \right.$$

$$\left. \times \left\{ P(M_i=m_i|A=a+1) - P(M_i=m_i|A=a) \right\} \right], \tag{F.10}$$

where $\mathbb{M}_i$ is the support of M_i, $o_i = [m_1, \cdots, m_{i-1}, m_{i+1}, \cdots, m_p]$, $P(M=m|A=a)$ is the probability of $M=m$ when setting $A=a$.

Theorem F.4 *Under assumptions (A1-A3) and Model 1, we have*

$$DM_i = \xi_i,$$

Remark F.5 *The proof can be found in Section G.5. Based on Definition 3.2 and Equation F.10, both the proposed DM and the effect defined in Vansteelandt & Daniel (2017) contain the information of the causal effect of A on the mediator M_i, i.e. $P(M_i=m_i|A=a+1) - P(M_i=m_i|A=a)$.*

G TECHNICAL PROOFS

G.1 PROOF OF THEOREM 3.1

Proof G.1 *In this proof, we will give the explicit expressions of causal effects defined under the LSEM. First, Equation 3 is equivalent to*

$$\begin{cases} A \equiv \epsilon_A, \\ M = \boldsymbol{\alpha}A + B_M^\top M + \epsilon_M, \\ Y = \gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y. \end{cases} \tag{G.1}$$

Based on $M = \boldsymbol{\alpha}A + B_M^\top M + \epsilon_M$, by moving $B_M^\top M$ to the left-hand side, we have

$$(I_p - B_M^\top)M = \boldsymbol{\alpha}A + \epsilon_M.$$

Suppose the mediators are sorted in the topological order (a series of elementary transformation of the matrix), then the matrix $B_M^\top$ is strictly upper triangular with the diagonal element as 0. Thus, we have $I_p - B_M^\top$ is invertible, then $I_p - B_M^\top$ under its original order should be also invertible (any invertible matrix after elementary transformation is still invertible).

Therefore, we can rewrite M as a purely function of A plus the error term as follows.

$$M = (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}A + (I_p - B_M^\top)^{-1}\epsilon_M. \tag{G.2}$$

Then we replace mediators in Equation G.1 with Equation G.2 and obtain

$$\begin{cases} A & \equiv \epsilon_A, \\ M & = (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}A + (I_p - B_M^\top)^{-1}\epsilon_M, \\ Y & = \gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y \\ & = \gamma A + \{\boldsymbol{\beta}^\top(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}A + \{\boldsymbol{\beta}^\top(I_p - B_M^\top)^{-1}\epsilon_M + \epsilon_Y\}. \end{cases} \tag{G.3}$$

Next, we show how to get the explicit expressions of $E\{Y|do(A=a)\}$ under the LSEM. Following the results in Rosenbaum & Rubin (1983), under the assumption (A2), we have $P\{M|do(A=a)\} = P(M|A=a)$, and thus,

$$E\{M|do(A=a)\} = E(M|A=a).$$

Similarly, we can get $E\{Y|do(A = a)\} = E(Y|A = a)$ under the assumption (A1), and $E\{Y|do(A = a, M = m)\} = E(Y|A = a, M = m)$ under the assumption (A3).

Based on above results and Equation G.3, we have

$$
\begin{aligned}
E\{Y|do(A = a)\} &= E\{Y|A = a\} \\
&= E\{\gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y | A = a\} \\
&= \gamma a + \boldsymbol{\beta}^\top E\{M|A = a\} \\
&= \gamma a + \boldsymbol{\beta}^\top E\{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha} A + (I_p - B_M^\top)^{-1}\epsilon_M | A = a\} \\
&= \gamma a + \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1}\boldsymbol{\alpha} a,
\end{aligned}
\tag{G.4}
$$

where the first '=' is held under the assumption (A1), the second and forth '=' are given by Equation G.3 that $Y = \gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y$ and $M = (I_p - B_M^\top)^{-1}\boldsymbol{\alpha} A + (I_p - B_M^\top)^{-1}\epsilon_M$.

Following the same calculation procedure of $E\{Y|do(A = a)\}$, we next give the natural direct effect under assumptions (A1-A3) and Model 1 as

$$
\begin{aligned}
DE &= E\{Y|do(A = a + 1, M = m^{(a)})\} - E\{Y|do(A = a)\} \\
&= \{\gamma(a + 1) + \boldsymbol{\beta}^\top m^{(a)}\} - \{\gamma a + \boldsymbol{\beta}^\top m^{(a)}\} \\
&= \gamma,
\end{aligned}
$$

where the first '=' is given by the definition of the DE.

Similarly, the natural indirect effect is

$$
\begin{aligned}
IE &= E\{Y|do(A = a, M = m^{(a+1)})\} - E\{Y|do(A = a)\} \\
&= \{\gamma a + \boldsymbol{\beta}^\top m^{(a+1)}\} - \{\gamma a + \boldsymbol{\beta}^\top m^{(a)}\} \\
&= \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}(a + 1) - \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1}\boldsymbol{\alpha} a \\
&= \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}.
\end{aligned}
$$

Thus, the total effect of A on Y is

$$
TE = E\{Y|do(A = a + 1)\} - E\{Y|do(A = a)\} = DE + IE = \gamma + \boldsymbol{\beta}^\top (I_p - B_M^\top)^{-1}\boldsymbol{\alpha}.
$$

Finally, we give the expression for the natural direct effect of M_i on Y under the LSEM. Based on the assumption (A2) and Equation G.2, we have

$$
\begin{aligned}
&E\{M_i|do(A = a + 1)\} - E\{M_i|do(A = a)\} \\
&= E\{M_i|A = a + 1\} - E\{M_i|A = a\} \\
&= \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i(a + 1) - \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i a \\
&= \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i,
\end{aligned}
\tag{G.5}
$$

where $\{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i$ is the i-th element of the vector $(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}$.

Then, based on $Y = \gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y$ and the assumption (A3), we have,

$$
\begin{aligned}
&E\{Y|do(A = a, M_i = m_i^{(a)} + 1, \Omega_i = o_i^{(a)})\} - E\{Y|do(A = a)\} \\
&= E\{Y|A = a, M_i = m_i^{(a)} + 1, \Omega_i = o_i^{(a)}\} - E\{Y|A = a\} \\
&= \gamma a + \boldsymbol{\beta}^\top \begin{bmatrix} m_1^{(a)} \\ \vdots \\ m_i^{(a)} + 1 \\ \vdots \\ m_p^{(a)} \end{bmatrix} - \gamma a - \boldsymbol{\beta}^\top \begin{bmatrix} m_1^{(a)} \\ \vdots \\ m_i^{(a)} \\ \vdots \\ m_p^{(a)} \end{bmatrix} = \boldsymbol{\beta}^\top \boldsymbol{1}_i = \beta_i,
\end{aligned}
$$

where $\boldsymbol{1}_i$ is a $p \times 1$ vector with the i-th element as 1 while others qual to 0, and β_i is the i-th element of the vector $\boldsymbol{\beta}$.

Thus, we have

$$
\begin{aligned}
DM_i =& \Big[E\{M_i|do(A=a+1)\} - E\{M_i|do(A=a)\} \Big] \\
& \times \Big[E\{Y|do(A=a, M_i=m_i^{(a)}+1, \Omega_i=o_i^{(a)})\} - E\{Y|do(A=a)\} \Big], \\
=& \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i \times \boldsymbol{\beta}_i \\
=& \boldsymbol{\beta}_i \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i. \qquad \square
\end{aligned}
$$

G.2 PROOF OF THEOREM F.1

Proof G.2 *1. If there is no directed path $\pi^* \in \{\pi_{AY}(\mathcal{G})\}$ such that the length of π^* is larger than 2, i.e. the length of $\pi^* \in \{\pi_{AY}(\mathcal{G})\}$ is either 1 or 2. Here, the path with length 1 corresponds to $A \to Y$, and paths with length 2 are $A \to M_i \to Y$ with possibly $i = 1, \cdots, p$. Thus, there is no interaction among mediators.*

By the definition of the LSEM, we have $B_M = \boldsymbol{0}_{p \times p}$, where $\boldsymbol{0}_{p \times p}$ is a $p \times p$ zero matrix. Following the path method (the causal effect of X_i on X_j along a directed path from $X_i \to X_j$ in $\mathcal{G}$ can be calculated by multiplying all edge weights along the path) illustrated in Wright (1921) and Nandy et al. (2017), we could obtain $\sum \eta_i = \sum_i \boldsymbol{\beta}_i \boldsymbol{\alpha}_i = IE$. (See a toy example provided in section E to illustrate how to use the path method to manually compute the causal effects.)

2. If there exists at least one directed path $\pi^* \in \{\pi_{AY}(\mathcal{G})\}$ such that the length of π^* is larger than 2, and the element in B is nonnegative, we have $B_M \neq \boldsymbol{0}_{p \times p}$. Without loss of generality, suppose there exists $M_i \in M$ with a set of directed path that starts with A, contains M_i, then goes through other mediators, and ends with Y, denoted each path in such set as $\pi_{i,j} = \{A \to \cdots \to M_i \cdots \to \cdots \to Y\}$ for $j = 1, \cdots, n_i$, where n_i is the size of such path set for M_i, and the weights of edges in $\pi_{i,j}$ is positive. Note the set $\{\pi_{i,j}\}$ excludes the paths end with $M_i \to Y$.

Let $e_{\pi_{i,j}}$ denote the causal effect of A on Y through directed path $\pi_{i,j}$. Based on the path method in Wright (1921) and Nandy et al. (2017) with the definition of IM_i, we have its theoretical form as

$$
IM_i = \sum_{j=1}^{n_i} e_{\pi_{i,j}}. \tag{G.6}
$$

By Equation G.5 and the definition of η_i, we have its first multiplier as

$$
E\{M_i|do(A=a+1)\} - E\{M_i|do(A=a)\} = \{(I_p - B_M^\top)^{-1}\boldsymbol{\alpha}\}_i,
$$

which is also the first multiplier in both DM_i and IM_i.

And the second multiplier of η_i can be expressed as

$$
\begin{aligned}
& E\{Y|do(M_i=m_i+1)\} - E\{Y|do(M_i=m_i)\} \\
=& E\{Y|do(M_i=m_i^{(a)}+1)\} - E\{Y|do(M_i=m_i^{(a)})\} \\
=& E\{Y|do(A=a, M_i=m_i^{(a)}+1)\} - E\{Y|do(A=a, M_i=m_i^{(a)})\}, \\
=& E\{Y|do(A=a, M_i=m_i^{(a)}+1)\} - E\{Y|do(A=a)\},
\end{aligned}
$$

where $m_i^{(a)}$ is the value of M_i when setting $do(A=a)$. Here, the first '=' is valid since m_i can be arbitrary number, and the second and third '=' are based on the equivalent interventions.

Based on the technique of plus and minus the same term, we decompose the second multiplier of η_i into two parts as follows

$$
\begin{aligned}
& E\{Y|do(A=a, M_i=m_i^{(a)}+1)\} - E\{Y|do(A=a)\} \\
=& \underbrace{\Big[E\{Y|do(A=a, M_i=m_i^{(a)}+1, \Omega_i=o_i^{(a)})\} - E\{Y|do(A=a)\} \Big]}_{\text{the second multiplier of } DM_i} \\
& + \underbrace{\Big[E\{Y|do(A=a, M_i=m_i^{(a)}+1)\} - E\{Y|do(A=a, M_i=m_i^{(a)}+1, \Omega_i=o_i^{(a)})\} \Big]}_{\text{the second multiplier of } IM_i}
\end{aligned}
$$

$$
\tag{G.7}
$$

where $\Omega_i = M \setminus M_i$ is the sets of mediators except M_i, and $o_i^{(a)}$ is the value of Ω_i when setting $do(A = a)$. Here, the first term in the above equation corresponds to the second multiplier of DM_i, while the second term is the second multiplier of IM_i.

Thus, the summation of η_i is

$$\sum \eta_i = \sum_i \left\{ \left[E\{M_i | do(A = a+1)\} - E\{M_i | do(A = a)\} \right] \right.$$
$$\left. \times \left[E\{Y | do(M_i = m_i + 1)\} - E\{Y | do(M_i = m_i)\} \right] \right\}$$
$$= \sum_i \{DM_i + IM_i\} = \sum_i DM_i + \sum_i IM_i = IE + \sum_i \sum_{j=1}^{n_i} e_{\pi_{i,j}},$$

where the first '=' is from Definition F.1, the second '=' is given by Equation G.7 and Definition 3.2 and 3.3, and the last '=' comes from Theorem 3.2 and the theoretical form of IM in Equation G.6.

Here, we have $e_{\pi_{i,j}} > 0$ since the weights of edges in $\pi_{i,j}$ is positive based on the path method in Wright (1921) and Nandy et al. (2017). Then, $\sum_i \sum_{j=1}^{n_i} e_{\pi_{i,j}}$ is also strictly larger than 0. Therefore, we have

$$\sum \eta_i > IE. \qquad \square$$

G.3 PROOF OF THEOREM F.2

Proof G.3 *1. If there doesn't exist edge $M_i \to Y$ in $\mathcal{G}$, then by definition we have $\boldsymbol{\beta}_i = 0$. Thus, $LE_i = \boldsymbol{\beta}_i \zeta_i = 0$.*

2. If there exists edge $M_i \to Y$ in $\mathcal{G}$. Suppose there is a directed path set with size m_i associated to the edge $M_i \to Y$, where each directed path $\tilde{\pi}_{i,j}$ starts with node A and ends with $M_i \to Y$, denoted as $\tilde{\pi}_{i,j} = \{A \to \cdots \to \cdots \to M_i \to Y\}$ for $j = 1, \cdots, m_i$.

Let $e_{\tilde{\pi}_{i,j}}$ denote the causal effect of A on Y through directed path $\tilde{\pi}_{i,j}$, $e_{\tilde{\pi}_{i,j}}^{(A,M_i)}$ be the causal effect of A on M_i through directed path $\tilde{\pi}_{i,j}$, and $e^{(M_i,Y)}$ is the causal effect of M_i on Y through directed edge $M_i \to Y$. Following the path method in Wright (1921) and Nandy et al. (2017), we have $e_{\tilde{\pi}_{i,j}} = e_{\tilde{\pi}_{i,j}}^{(A,M_i)} e^{(M_i,Y)}$.

Then the i-th last edge-specific effect is equal to the summation of the effect through each path $\tilde{\pi}_{i,j}$, i.e.,

$$LE_i = \sum_{j=1}^{n_i} e_{\tilde{\pi}_{i,j}} = \sum_{j=1}^{n_i} e_{\tilde{\pi}_{i,j}}^{(A,M_i)} e^{(M_i,Y)} = e^{(M_i,Y)} \sum_{j=1}^{n_i} e_{\tilde{\pi}_{i,j}}^{(A,M_i)}.$$

Here, by the similar argument based on the path method, we have $e^{(M_i,Y)} = \boldsymbol{\beta}_i$ and $\sum_{j=1}^{n_i} e_{\tilde{\pi}_{i,j}}^{(A,M_i)}$ as the total causal effect of A on M_i.

Recall that $\zeta_i \equiv \{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i$ is the causal effect of A on M_i. Therefore, the i-th LE is the product of the causal effect of A on M_i and the causal effect of M_i on Y, i.e.,

$$LE_i = \boldsymbol{\beta}_i \zeta_i. \qquad \square$$

G.4 PROOF OF THEOREM F.3

Proof G.4 *Given a general DAG $\mathcal{G}$ with nodes $\{A, M, Y\}$, let the union of all directed paths that contain the i-th last edge as $\tau_i = \{\pi : A \to \cdots \to M_i \to Y\}, i = 1, \cdots p$. Here, we have $\tau_i = \{\tilde{\pi}_{i,j}\}_{1 \le j \le m_j}$ established in Section G.3. It is clear that the union set of τ_i in $\mathcal{G}$ is equal to the set of all directed paths start with A and end with node Y (except $A \to Y$) in $\mathcal{G}$ as*

$$\bigcup_i \tau_i = \{\pi_{AY}(\mathcal{G})\} \setminus \{A \to Y\}.$$

Also, based on the uniqueness of each last edge, τ_i is pairwise disjoint, i.e.

$$\tau_i \bigcap \tau_j = \varnothing, \qquad \forall i \neq j.$$

Since the IE is defined as the total causal effect of A on Y that goes through mediators, we have the IE equal to the causal effect that goes through the set $\{\pi_{AY}(\mathcal{G})\} \setminus \{A \to Y\}$, i.e. the IE equal to the causal effect that goes through set $\bigcup_i \tau_i$. Based on the mutual disjoint property of τ_i, we have the IE is exactly the summation of the causal effect through τ_i. Lastly, from the definition of LE_i, we have

$$IE = \sum_{i=1}^{p} LE_i. \qquad \square$$

G.5 PROOF OF THEOREM F.4

Proof G.5 *The proof of the consistency of our defined DM to the interventional effect ξ_i can be completed based on Equation 3 under assumptions (A1-A3) and Model 1.*

Recall the definition in Equation F.10, we have

$$\xi_i = \sum_{m_1 \in \mathbb{M}_1} \cdots \sum_{m_p \in \mathbb{M}_p} \left[E(Y|A = a, M_i = m_i, \Omega_i = o_i) P(\Omega_i = o_i | A = a) \right.$$

$$\left. \times \left\{ P(M_i = m_i | A = a + 1) - P(M_i = m_i | A = a) \right\} \right].$$

$$= \sum_{m_1 \in \mathbb{M}_1} \cdots \sum_{m_p \in \mathbb{M}_p} \left\{ E(Y|A = a, M_i = m_i, \Omega_i = o_i) P(\Omega_i = o_i | A = a) P(M_i = m_i | A = a + 1) \right.$$

$$\left. - E(Y|A = a, M_i = m_i, \Omega_i = o_i) P(\Omega_i = o_i | A = a) P(M_i = m_i | A = a) \right\}.$$

Given $A = a$, the value of M_i is $m_i^{(a)}$ and Ω_i takes $o_i^{(a)}$; while when setting $A = a + 1$, the value of M_i is $m_{a+1}^{(i)}$. Therefore, we have $P(M_i = m_i | A = a) = 1$ if $m_i = m_i^{(a)}$ otherwise is 0, and $P(\Omega_i = o_i | A = a) = 1$ if $o_i = o_i^{(a)}$ otherwise is 0.

Under assumptions (A1-A3), we have

$$\xi_i = E(Y|A = a, M_i = m_i^{(a+1)}, \Omega_i = o_i^{(a)}) - E(Y|A = a, M_i = m_i^{(a)}, \Omega_i = o_i^{(a)}).$$

Then, based on the LSEM that $Y = \gamma A + \boldsymbol{\beta}^\top M + \epsilon_Y$, we can further obtain that

$$\xi_i = \gamma a + \boldsymbol{\beta}^\top \begin{bmatrix} m_1^{(a)} \\ \vdots \\ m_i^{(a+1)} \\ \vdots \\ m_p^{(a)} \end{bmatrix} - \gamma a - \boldsymbol{\beta}^\top \begin{bmatrix} m_1^{(a)} \\ \vdots \\ m_i^{(a)} \\ \vdots \\ m_p^{(a)} \end{bmatrix} = \boldsymbol{\beta}_i \{ m_i^{(a+1)} - m_i^{(a)} \}.$$

From Equation G.2, we have

$$\xi_i = \boldsymbol{\beta}_i \left[\{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i (a + 1) - \{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i a \right] = \boldsymbol{\beta}_i \{(I_p - B_M^\top)^{-1} \boldsymbol{\alpha}\}_i.$$

Thus, under assumptions (A1-A3) and Model 1, we have

$$DM_i = \xi_i. \qquad \square$$