WHEN DO MODELS GENERALIZE? A PERSPECTIVE FROM DATA-ALGORITHM COMPATIBILITY

ABSTRACT

One of the major open problems in machine learning is to characterize generalization in the overparameterized regime, where most traditional generalization bounds become inconsistent (Nagarajan and Kolter, 2019). In many scenarios, their failure can be attributed to obscuring the crucial interplay between the training algorithm and the underlying data distribution. To address this issue, we propose a concept named compatibility, which quantitatively characterizes generalization in a both data-relevant and algorithm-relevant manner. By considering the entire training trajectory and focusing on early-stopping iterates, compatibility exploits the data and the algorithm information and is therefore a more suitable notion for generalization. We validate this by theoretically studying compatibility under the setting of solving overparameterized linear regression with gradient descent. Specifically, we perform a data-dependent trajectory analysis and derive a sufficient condition for compatibility in such a setting. Our theoretical results demonstrate that in the sense of compatibility, generalization holds with significantly weaker restrictions on the problem instance than the previous last iterate analysis.

1 INTRODUCTION

Although deep neural networks achieve great success in practice (Silver et al., 2017; Devlin et al., 2019; Brown et al., 2020), their remarkable generalization ability is still among the essential mysteries in the deep learning community. One of the most intriguing features of deep neural networks is overparameterization, which confers a level of tractability to the training problem, but leaves traditional generalization theories failing to work. In the overparameterized regime, both the training algorithm and the data distribution play essential roles in generalization analysis (Jiang et al., 2020). For instance, a line of work (Zhang et al., 2021; Nagarajan and Kolter, 2019) highlights the role of the algorithm by showing that the algorithm-irrelevant uniform convergence bounds can become inconsistent in deep learning regimes. Another line of work (Bartlett et al., 2019; Tsigler and Bartlett, 2020) on benign overfitting emphasizes the role of data distribution via profound analysis of specific overparameterized models.

Despite the significant role of data and algorithm in generalization analysis, existing theories usually focus on either the data factor (*e.g.*, uniform convergence (Nagarajan and Kolter, 2019) and last iterate analysis (Bartlett et al., 2019; Tsigler and Bartlett, 2020)) or the algorithm factor (*e.g.*, stability-based bounds (Hardt et al., 2016)), but do not explain how they work well together in practice. In this sense, a natural question arises:

How to incorporate both data factor and algorithm factor into generalization analysis?

To gain insight into the interplay between data and algorithms, we provide motivating examples of a synthetic overparameterized linear regression task and a classification task on the corrupted MNIST dataset in figure 1. In both scenarios, the final iterate with less algorithmic information, which may include the algorithm type (*e.g.*, GD or SGD), hyperparameters (*e.g.*, learning rate, number of epochs), generalizes much worse than the early stopping solutions (see the Blue Line). In the linear regression case, the generalization error of the final iterate can be more than $\times 100$ larger than that of the early stopping solution. In the MNIST case, the final interpolator has 19.9% test error, much higher than the 2.88% test error of the best solution on the GD trajectory. Therefore, the almost ubiq-

(a) Linear Regression

(b) Corrupted MNIST

Figure 1: **(a) The training plot for linear regression with spectrum $\lambda_i = 1/i^2$ using GD.** Note that the axes are in the log scale. **(b) The training plot of CNN on corrupted MNIST with 20% label noise using SGD.** Both models successfully learn the useful features in the initial phase of training, but it takes a long time for them to fit the noise in the dataset. The observations demonstrate the power of data-dependent trajectory analysis, since the early stopping solution on the trajectory generalizes well but the final iterate fails to generalize. See Appendix C for details.

uitous strategy of early stopping is a key ingredient in generalization analysis for overparameterized models, whose benefits have been demonstrated both theoretically and empirically (Yao et al., 2007; Ali et al., 2019; Li et al., 2020b; Ji et al., 2021). By focusing on the entire optimization trajectory and performing data-dependent trajectory analysis, both data information and the dynamics of the training algorithm can be exploited to yield consistent generalization bounds.

To analyze the data-dependent trajectory, we introduce a new concept named **data-algorithm-compatibility**, which jointly characterizes the role of the data and the algorithm in generalization analysis. Informally speaking, an algorithm is compatible with a data distribution if the best iterate on the training trajectory reaches zero excess risk as the sample size goes to infinity.

The significance of compatibility comes in three folds. Firstly, compatibility incorporates both data and algorithm factors into generalization analysis, and brings new messages into generalization in the overparameterization regime (see Definition 3.1). Secondly, compatibility serves as a minimal condition for generalization, without which one cannot expect to find a consistent solution via standard learning procedures. Consequently, compatibility holds with only mild assumptions and applies to a wide range of problem instances (see Theorem 4.1). Thirdly, compatibility captures the algorithmic significance of early stopping in generalization. By exploiting the algorithm information along the entire trajectory, we arrive at better generalization bounds than the final iterate analysis (see Table 1 and 2 for examples).

To theoretically validate compatibility, we study it under overparameterized linear regression setting. Analysis of the overparameterized linear regression is a reasonable starting point to study compatibility for more complex models like deep neural networks, since many phenomena of the high dimensional non-linear model are also observed in the linear regime (*e.g.*, Figure 1). Furthermore, the recent neural tangent kernel (NTK) framework demonstrates that very wide neural networks trained using gradient descent with appropriate random initialization can be approximated by kernel regression in a reproducing kernel Hilbert space, which rigorously establishes a close relationship between overparameterized linear regression and deep neural network training (Jacot et al., 2018; Arora et al., 2019).

Specifically, we investigate solving overparameterized linear regression using gradient descent with constant step size, and prove that under some mild regularity conditions, gradient descent is compatible with overparameterized linear regression if the effective dimensions of the feature covariance matrix are asymptotically bounded by the sample size. In this setting, the assumptions needed for generalization in the sense of compatibility are significantly weaker than that in the final iterate analysis (Bartlett et al., 2019), which demonstrates the benefits of data-relevant and algorithm-relevant generalization analysis.

We summarize our contributions as follows:

- We formalize the notion of compatibility, which highlights the interaction between data and algorithm and serves as a minimal condition for generalization.

- We derive a sufficient condition for compatibility in solving overparameterized linear regression with gradient descent. Our theory substantiates the meaningfulness of compatibility by showing that generalization in the sense of compatibility typically requires much weaker restrictions in the problem instance.

- Technically, we derive time-variant generalization bounds for overparameterized linear regression via data-dependent trajectory analysis. Empirically, various experiment results verify the motivation of compatibility and demonstrate the benefits of early stopping.

2 RELATED WORKS

Data-Dependent Techniques mainly focus on the data distribution condition for generalization. One of the most popular bounds among data-dependent bounds is uniform convergence (Koltchinskii and Panchenko, 2000; Bartlett et al., 2017; Zhou et al., 2020; Zhang et al., 2021). However, recent works(Nagarajan and Kolter, 2019; Negrea et al., 2020) point out that uniform convergence may not be powerful enough to explain generalization, because it may only yield inconsistent bound in even linear regression cases. Another line of works investigates benign overfitting, which mainly involves the last iterate generalization (Bartlett et al., 2019; Zou et al., 2021; Tsigler and Bartlett, 2020; Li et al., 2020c; Wang and Thrampoulidis, 2021; Frei et al., 2022).

Algorithm-Dependent Techniques measure the role of the algorithmic information in generalization. A line of works derives generalization bounds via algorithm stability (Hardt et al., 2016; Feldman and Vondrák, 2018; Mou et al., 2018; Feldman and Vondrák, 2019; Bousquet et al., 2020; Li et al., 2020a; Lei and Ying, 2020; Bassily et al., 2020; Teng et al., 2021). A parallel line of works analyzes the implicit bias of algorithmic information (Bousquet and Elisseeff, 2002; Soudry et al., 2018; Shah et al., 2020; Hu et al., 2020; Lyu and Li, 2020; Lyu et al., 2021), which are mainly based on analyzing a specific data distribution (*e.g.*, linear separable).

Other Generalization Techniques. Besides the techniques discussed above, there are many other approaches, including PAC-Bayes (Parrado-Hernández et al., 2012; Dziugaite and Roy, 2017; Neyshabur et al., 2018), information theory (Russo and Zou, 2016; Xu and Raginsky, 2017; Banerjee and Montúfar, 2021), and compression-based bounds (Arora et al., 2018).

Early Stopping has the potential to improve generalization (Zhang et al., 2021) and is effective under noisy label regimes (Li et al., 2020b; Bai et al., 2021). On the theoretical side, Ji et al. (2021) prove that early stopping helps the shallow ReLU neural networks to be consistent. Another line of research focuses on the signal for early stopping (Prechelt, 2012; Forouzesh and Thiran, 2021).

3 COMPATIBILITY

In this section, we formally define compatibility between the data distribution and the training algorithm, starting from the basic notations.

3.1 NOTATIONS

Data Distribution. Let $\mathcal{D}$ denote the population distribution and $\boldsymbol{z} \sim \mathcal{D}$ denote a data point sampled from distribution $\mathcal{D}$. Usually, $\boldsymbol{z}$ contains a feature and its corresponding response. Besides, we denote the dataset with n samples as $\boldsymbol{Z} \triangleq \{\boldsymbol{z}_i\}_{i \in [n]}$, where $\boldsymbol{z}_i \sim \mathcal{D}$ are i.i.d. sampled from distribution $\mathcal{D}$.

Loss and Excess Risk. Let $\ell(\boldsymbol{\theta}; \boldsymbol{z})$ denote the loss on sample $\boldsymbol{z}$ with parameter $\boldsymbol{\theta} \in \mathbb{R}^p$. The corresponding population loss is defined as $L(\boldsymbol{\theta}; \mathcal{D}) \triangleq \mathbb{E}_{\boldsymbol{z} \sim \mathcal{D}} \ell(\boldsymbol{\theta}; \boldsymbol{z})$. When the context is clear, we omit the dependency on $\mathcal{D}$ and denote the population loss by $L(\boldsymbol{\theta})$. Our goal is to find the optimal parameter $\boldsymbol{\theta}^*$ which minimizes the population loss, i.e., $L(\boldsymbol{\theta}^*) = \min_{\boldsymbol{\theta}} L(\boldsymbol{\theta})$. Measuring how a parameter $\boldsymbol{\theta}$ approaches $\boldsymbol{\theta}^*$ relies on a term *excess risk* $R(\boldsymbol{\theta})$, defined as $R(\boldsymbol{\theta}) \triangleq L(\boldsymbol{\theta}) - L(\boldsymbol{\theta}^*)$.

Algorithm. Let $\mathcal{A}(\cdot)$ denote an algorithm that takes training data $\boldsymbol{Z}$ as input and outputs a sequence of parameters $\{\boldsymbol{\theta}_n^{(t)}\}_{t \geq 0}$. The algorithm can be either deterministic or stochastic, *e.g.*, variants of (S)GD.

3.2 Definitions of Compatibility

Based on the above notations, we introduce the notion of compatibility between data distribution and algorithm in Definition 3.1. Informally, compatibility measures whether a consistent excess risk can be reached along the training trajectory. Note that we omit the role of the loss function in the definition, although the algorithm depends on the loss function.

Definition 3.1 (**Compatibility**). *Given a loss function $\ell(\cdot)$ with corresponding excess risk $R(\cdot)$, a data distribution $\mathcal{D}$ is **compatible** with an algorithm $\mathcal{A}$ if there exists nonempty sets T_n of $\mathbb{N}$, such that $\sup_{t \in T_n} R(\boldsymbol{\theta}_n^{(t)})$ converges to zero in probability as sample size n tends to infinity, where the randomness comes from the sampling of training data $\boldsymbol{Z}$ from distribution $\mathcal{D}$ and the execution of algorithm $\mathcal{A}$. That is to say, $(\mathcal{D}, \mathcal{A})$ is compatible if*

$$\sup_{t \in T_n} R(\boldsymbol{\theta}_n^{(t)}) \xrightarrow{P} 0 \quad as \quad n \to \infty. \tag{1}$$

We call $\{T_n\}_{n>0}$ the compatibility region of $(\mathcal{D}, \mathcal{A})$. The distribution $\mathcal{D}$ is allowed to change with n. In this case, $\mathcal{D}$ should be understood as a sequence of distributions $\{\mathcal{D}_n\}_{n \geq 1}$. We also allow the dimension of model parameter $\boldsymbol{\theta}$ to be infinity or to grow with n. We omit this dependency on n when the context is clear.

Compatibility serves as a minimal condition for generalization, since if a data distribution is incompatible with the algorithm, one cannot expect to reach a small excess risk even if we allow for *arbitrary* early stopping. However, we remark that considering only the minimal excess risk is insufficient for a practical purpose, as one cannot exactly find the t that minimizes $R(\boldsymbol{\theta}_n^{(t)})$ due to the noise in the validation set. Therefore, it is meaningful to consider a region of time t on which the excess risk is consistent as in Definition 3.1. The larger the compatibility region is, the more robust the algorithm will be to the noise in its execution.

Comparisons with Other Notions. Compared to classic definitions of learnability, *e.g.*, PAC learning, the definition of compatibility is data-specific and algorithm-specific, and is thus a more fine-grained notion. Compared to the concept of *benign* proposed in the recent paper (Bartlett et al., 2019), which studies whether the excess risk at $t = \infty$ converges to zero in probability as the sample size goes to infinity, compatibility only requires that there exists a time to derive a consistent excess risk. We will show later in Section 4.2 that in the overpamameterized linear regression setting, there exist cases such that the problem instance is compatible but not benign.

4 Compatibility Analysis of Overparameterized Linear Regression with Gradient Descent

This paper mainly analyzes compatibility in the overparameterized linear regression regime. We first introduce the data distribution, loss, and training algorithm, and then present the main theorem, which provides a sufficient condition for compatibility in this setting.

4.1 Preliminaries for Overparameterized Linear Regression

Data Distribution. Let $(\boldsymbol{x}, y) \in \mathbb{R}^p \times \mathbb{R}$ denote the feature vector and the response, following a joint distribution $\mathcal{D}$. Let $\boldsymbol{\Sigma} \triangleq \mathbb{E}[\boldsymbol{x}\boldsymbol{x}^\top]$ denote the feature covariance matrix, whose eigenvalue decomposition is $\boldsymbol{\Sigma} = \boldsymbol{V}\boldsymbol{\Lambda}\boldsymbol{V}^\top = \sum_{i>0} \lambda_i \boldsymbol{v}_i \boldsymbol{v}_i^\top$ with decreasing eigenvalues $\lambda_1 \geq \lambda_2 \geq \cdots$. We make the following assumptions on the distribution of the feature vector.

Assumption 1 (Assumptions on feature distribution). *We assume that*

1. $\mathbb{E}[\boldsymbol{x}] = 0$.

2. $\lambda_1 > 0, \sum_{i>0} \lambda_i < C$ *for some absolute constant C.*

3. *Let $\tilde{\boldsymbol{x}} = \boldsymbol{\Lambda}^{-\frac{1}{2}} \boldsymbol{V}^\top \boldsymbol{x}$. The random vector $\tilde{\boldsymbol{x}}$ has independent σ_x-subgaussian entries.*[1]

Loss and Excess Risk. We choose square loss as the loss function ℓ, i.e. $\ell(\boldsymbol{\theta}, (\boldsymbol{x}, y)) = 1/2(y - \boldsymbol{x}^\top \boldsymbol{\theta})^2$. The corresponding population loss is denoted by $L(\boldsymbol{\theta}) = \mathbb{E}\ell(\boldsymbol{\theta}, (\boldsymbol{x}, y))$ and the optimal parameter is denoted by $\boldsymbol{\theta}^* \triangleq \operatorname{argmin}_{\boldsymbol{\theta} \in \mathbb{R}^p} L(\boldsymbol{\theta})$. We assume that $\|\boldsymbol{\theta}^*\| < C$ for some absolute constant C. If there are multiple such minimizers, we choose an arbitrary one and fix it thereafter. We focus on the excess risk of parameter $\boldsymbol{\theta}$, defined as

$$R(\boldsymbol{\theta}) = L(\boldsymbol{\theta}) - L(\boldsymbol{\theta}^*) = \frac{1}{2}\mathbb{E}(y - \boldsymbol{x}^\top \boldsymbol{\theta})^2 - \frac{1}{2}\mathbb{E}(y - \boldsymbol{x}^\top \boldsymbol{\theta}^*)^2 = \frac{1}{2}(\boldsymbol{\theta} - \boldsymbol{\theta}^*)^\top \boldsymbol{\Sigma}(\boldsymbol{\theta} - \boldsymbol{\theta}^*).$$

Let $\varepsilon = y - \boldsymbol{x}^\top \boldsymbol{\theta}^*$ denote the noise in data point $(\boldsymbol{x}, y)$. The following assumptions involve the conditional distribution of the noise.

Assumption 2 (Assumptions on noise distribution). *We assume that*

1. *The conditional noise $\varepsilon | \boldsymbol{x}$ has zero mean.*

2. *The conditional noise $\varepsilon | \boldsymbol{x}$ is σ_y-subgaussian.*

Note that both Assumption 1 and Assumption 2 are mild and commonly considered in the related literatures (Bartlett et al., 2019; Tsigler and Bartlett, 2020; Zou et al., 2021).

Training Set. Given a training set $\{(\boldsymbol{x}_i, y_i)\}_{1 \leq i \leq n}$ with n pairs independently sampled from the population distribution $\mathcal{D}$, we define $\boldsymbol{X} \triangleq (\boldsymbol{x}_1, \cdots, \boldsymbol{x}_n)^\top \in \mathbb{R}^{n \times p}$ as the feature matrix, $\boldsymbol{Y} \triangleq (y_1, \cdots, y_n)^\top \in \mathbb{R}^n$ as the corresponding noise vector, and $\boldsymbol{\varepsilon} \triangleq \boldsymbol{Y} - \boldsymbol{X}\boldsymbol{\theta}^*$ as the residual vector. Let the singular value decomposition (SVD) of $\boldsymbol{X}$ be $\boldsymbol{X} = \boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}} \boldsymbol{W}^\top$, with $\tilde{\boldsymbol{\Lambda}} = \operatorname{diag}\{\mu_1 \cdots, \mu_n\} \in \mathbb{R}^{n \times n}$, $\mu_1 \geq \cdots \geq \mu_n$.

We consider the overparameterized regime where the feature dimension is larger than the sample size, namely, $p > n$. In this regime, we assume that $\operatorname{rank}(\boldsymbol{X}) = n$ almost surely as in Bartlett et al. (2019).

Assumption 3 (Linear independent training set). *For any $n < p$, we assume that the features in the training set $\{\boldsymbol{x}_1, \boldsymbol{x}_2, \cdots, \boldsymbol{x}_n\}$ is linearly independent almost surely.*

Algorithm. Given the dataset $(\boldsymbol{X}, \boldsymbol{Y})$, define the empirical loss function as $\hat{L}(\boldsymbol{\theta}) \triangleq \frac{1}{2n}\|\boldsymbol{X}\boldsymbol{\theta} - \boldsymbol{Y}\|^2$. We choose full-batch gradient descent on the empirical risk with a constant learning rate λ as the algorithm $\mathcal{A}$ in the previous template. In this case, the update rule for the optimization trajectory $\{\boldsymbol{\theta}_t\}_{t \geq 0}$ is formulated as

$$\boldsymbol{\theta}_{t+1} = \boldsymbol{\theta}_t - \frac{\lambda}{n}\boldsymbol{X}^\top(\boldsymbol{X}\boldsymbol{\theta}_t - \boldsymbol{Y}). \tag{2}$$

Without loss of generality, we consider zero initialization $\boldsymbol{\theta}_0 = \boldsymbol{0}$ in this paper. In this case, for a sufficiently small learning rate λ, $\boldsymbol{\theta}_t$ converges to the *min-norm interpolator* $\hat{\boldsymbol{\theta}} = \boldsymbol{X}^\top(\boldsymbol{X}\boldsymbol{X}^\top)^{-1}\boldsymbol{Y}$ as t goes to infinity, which was well studied previously (Bartlett et al., 2019). This paper takes one step further and discuss the excess risk along the entire training trajectory $\{R(\boldsymbol{\theta}_t)\}_{t \geq 0}$.

Effective Rank and Effective Dimensions. We define the effective rank of the feature matrix $\boldsymbol{\Sigma}$ as $r(\boldsymbol{\Sigma}) \triangleq \frac{\sum_{i > 0} \lambda_i}{\lambda_1}$. Our results on compatibility depend on two notions of effective dimension of the feature covariance $\boldsymbol{\Sigma}$, defined as

$$\begin{aligned} k_0 &\triangleq \min\left\{l \geq 0 : \lambda_{l+1} \leq \frac{c_0 \sum_{i > l} \lambda_i}{n}\right\}, \\ k_1 &\triangleq \min\left\{l \geq 0 : \lambda_{l+1} \leq \frac{c_1 \sum_{i > 0} \lambda_i}{n}\right\}, \end{aligned} \tag{3}$$

where c_0, c_1 are constants independent of the dimension p, sample size n, and time t [2]. We omit the dependency of k_0, k_1 on $c_0, c_1, n, \boldsymbol{\Sigma}$ when the context is clear.

---

[1] A random variable X is σ-subgaussian if $\mathbb{E}[e^{\lambda X}] \leq e^{\frac{\lambda^2 \sigma^2}{2}}$ for any λ.

[2] Constants may depend on σ_x, and we omit this dependency thereafter for clarity.

4.2 COMPATIBILITY FOR OVERPARAMETERIZED LINEAR REGRESSION WITH GRADIENT DESCENT

Next, we present the main result of this section, which provides a clean condition for compatibility between gradient descent and overparameterized linear regression.

Theorem 4.1 (**Compatibility for Overparameterized Linear Regression with Gradient Descent**). *Consider the overparameterized linear regression setting defined in section 4.1. Let Assumption 1,2 and 3 hold. Assume the learning rate satisfies $\lambda = O\left(\frac{1}{\text{Tr}(\Sigma)}\right)$. Then under the condition that $k_0 = O(n), k_1 = o(n), r(\Sigma) = o(n)$, gradient descent is compatible with overparameterized linear regression in the region $T_n = \left(\omega\left(\frac{1}{\lambda}\right), o\left(\frac{n}{\lambda}\right)\right)$, namely,*

$$\sup_{t \in T_n} R(\boldsymbol{\theta}_t) \xrightarrow{P} 0 \quad as \quad n \to \infty.$$

Furthermore, if the feature dimension $p = \infty$ and the data distribution does not change with n, then the condition $k_0 = O(n)$ alone suffices for compatibility.

The proof of Theorem 4.1 is given in Appendix A and sketched in Section 5. The theorem shows that gradient descent is compatible with overparameterized linear regression under some mild regularity conditions on the learning rate, effective rank and effective dimensions. The condition on the learning rate is natural for optimizing a smooth objective. We conjecture that the condition $k_0 = O(n)$ can not be removed in general cases, since the effective dimension k_0 characterizes the concentration of the singular values of the data matrix X and plays a crucial role in the excess risk of the gradient descent dynamics. We discuss extensions to the kernel regression setting in Appendix B.5.

Comparison with Benign Overfitting. The recent paper Bartlett et al. (2019) studies overparameterized linear regression and gives the condition for min-norm interpolator to generalize. They prove that the feature covariance Σ is benign iff

$$k_0 = o(n), \quad R_{k_0}(\Sigma) \triangleq \frac{(\sum_{i > k_0} \lambda_i)^2}{\sum_{i > k_0} \lambda_i^2} = \omega(n), \quad r(\Sigma) = o(n)$$

As discussed in Section 3.2, benign problem instance also satisfies compatibility, since benign overfitting requires a stronger condition on k_0 and an additional assumption on $R_{k_0}(\Sigma)$. The following example shows that this inclusion relationship is strict.

Example 4.1. *Under the same assumption as in Theorem 4.1, if the spectrum of Σ satisfies*

$$\lambda_k = \frac{1}{k^\alpha},$$

for some $\alpha > 1$, we derive that $k_0 = \Theta(n)$. Therefore, this problem instance satisfies compatibility, but does not satisfy benign overfitting.

Example 4.1 shows the existence of a case where the early stopping solution can generalize but interpolating solution fails to generalize. Therefore, compatibility can characterize generalization for a much wider range of problem instances. [3]

Comparisons with Stability-Based Bounds and Uniform Convergence Bounds. Theorem 5.1 cannot be directly implied by off-the-shelf stability-based generalization bounds (Hardt et al., 2016; Feldman and Vondrák, 2019) or uniform convergence bounds (Koltchinskii and Panchenko, 2000; Bartlett et al., 2017). The main reason is that both methods rely on a *high probability* bound of the parameter norm, which requires a nontrivial data-dependent and algorithm-dependent analysis. Even if this can be done, both methods will provably give looser bounds with smaller compatibility regions than that in Theorem 5.1. See Appendix B.3 for a detailed comparison with stability-based bounds and Appendix B.4 for discussions on uniform convergence bounds.

---

[3]It is worth mentioning that Bartlett et al. (2019) and this paper study the generalization behavior of different models. The comparison here and in the following sections aims to validate the meaningfulness of compatibility analysis, instead of beating their results.

5 PROOF SKETCH AND TECHNIQUES

In this section, we sketch the proof for Theorem 4.1, and present the major techniques.

5.1 A TIME VARIANT BOUND

We further introduce an additional type of effective dimension besides k_0, k_1, which is time variant and is utilized to track the optimization dynamics.

Definition 5.1 (Effective Dimensions). *Given a feature covariance matrix Σ, define the effective dimension k_2 as*

$$k_2 \triangleq \min \left\{ l \geq 0 : \sum_{i>l} \lambda_i + n\lambda_{l+1} \leq c_2 c(t,n) \sum_{i>0} \lambda_i \right\}, \tag{4}$$

where c_2 is a constant independent of the dimension p, sample size n, and time t. The term $c(t,n)$ is a function to be discussed later. When the context is clear, we omit its dependencies on $c_2, c(t,n), n, \Sigma$ and only denote it by k_2.

Based on the effective rank and effective dimensions defined above, we provide a time-variant bound in Theorem 5.1 for overparameterized linear regression, which further leads to the compatibility argument in Theorem 4.1. Compared to the existing bound (Bartlett et al., 2019), Theorem 5.1 focuses on investigating the role of training epoch t in the excess risk, and is of independent interest.

Theorem 5.1 (Time Variant Bound). *Suppose Assumption 1, 2 and 3 hold. Fix a function $c(t,n)$. Given $\delta \in (0,1)$, assume that $k_0 \leq \frac{n}{c}$, $\log \frac{1}{\delta} \leq \frac{n}{c}$, $0 < \lambda \leq \frac{1}{c \sum_{i>0} \lambda_i}$ for a large enough constant c. Then with probability at least $1 - \delta$, we have for any $t \in \mathbb{N}$,*

$$R(\boldsymbol{\theta}_t) \lesssim B(\boldsymbol{\theta}_t) + V(\boldsymbol{\theta}_t),$$

where

$$B(\boldsymbol{\theta}_t) = \|\boldsymbol{\theta}^*\|^2 \left(\frac{1}{\lambda t} + \|\Sigma\| \max \left\{ \sqrt{\frac{r(\Sigma)}{n}}, \frac{r(\Sigma)}{n}, \sqrt{\frac{\log(\frac{1}{\delta})}{n}} \right\} \right),$$

$$V(\boldsymbol{\theta}_t) = \sigma_y^2 \log \left(\frac{1}{\delta} \right) \left(\frac{k_1}{n} + \frac{k_2}{c(t,n)n} + c(t,n) \left(\frac{\lambda t}{n} \sum_{i>0} \lambda_i \right)^2 \right).$$

Theorem 5.1 first decomposes $R(\boldsymbol{\theta}_t)$ into the bias term and the variance term, and bounds them with $B(\boldsymbol{\theta}_t)$ and $V(\boldsymbol{\theta}_t)$, respectively. We note that there is a tradeoff between $B(\boldsymbol{\theta}_t)$ and $V(\boldsymbol{\theta}_t)$, where $B(\boldsymbol{\theta}_t)$ decreases with time t while $V(\boldsymbol{\theta}_t)$ increases with time t.

The $c(t,n)$ Principle. It is worth emphasizing that our bound holds for arbitrary positive function $c(t,n)$. Therefore, one can fine-tune the generalization bound by choosing a proper $c(t,n)$. In the subsequent sections, we show how to derive consistent risk bounds for different time t, based on different choices of $c(t,n)$.

5.2 VARYING t, CONSTANT $c(t,n)$

Theorem 5.1 provides an excess risk upper bound uniformly for $t \in \mathbb{N}$. However, it is still non-trivial to derive Theorem 4.1, where the remaining question is to decide the term $c(t,n)$. The following corollary shows the generalization bound when setting $c(t,n)$ to a constant.

Corollary 5.1 (Constant $c(t,n)$). *Let Assumption 1, 2 and 3 hold. Fix a constant $c(t,n)$. Suppose $k_0 = O(n)$, $k_1 = o(n)$, $r(\Sigma) = o(n)$, $\lambda = O\left(\frac{1}{\sum_{i>0} \lambda_i} \right)$. Then there exists a sequence of positive constants $\{\delta_n\}_{n \geq 0}$ which converge to 0, such that with probability at least $1 - \delta_n$, the excess risk is consistent for $t \in \left(\omega\left(\frac{1}{\lambda} \right), o\left(\frac{n}{\lambda} \right) \right)$, i.e.*

$$R(\boldsymbol{\theta}_t) = o(1).$$

Table 1: **Comparisons of excess risk bound with Bartlett et al. (2019) and Zou et al. (2021).** We provide four types of feature covariance with eigenvalues λ_k, including *Inverse Polynomial* ($\lambda_k = \frac{1}{k^\alpha}$, $\alpha > 1$), *Inverse Log Polynomial* ($\lambda_k = \frac{1}{k \log^\beta(k+1)}$, $\beta > 1$), *Constant* ($\lambda_k = \frac{1}{n^{1+\varepsilon}}$, $1 \le k \le n^{1+\varepsilon}$, $\varepsilon > 0$), and *Piecewise Constant* ($\lambda_k = \frac{1}{s}$ if $1 \le k \le s$ and $\lambda_k = \frac{1}{d-s}$ if $s + 1 \le k \le d$, where $s = n^r, d = n^q, 0 < r \le 1, q > 1$). In light of these bounds, ours outperforms Bartlett et al. (2019) in all the cases, and outperforms Zou et al. (2021) in Constant / Piecewise Constant cases if $\varepsilon < \frac{1}{2}$ and $q < \min\{2-r, \frac{3}{2}\}$. We refer to Appendix B for more details.

DISTRIBUTIONS	OURS	BARTLETT ET AL. (2019)	ZOU ET AL. (2021)
INVERSE POLYNOMIAL	$O\left(n^{-\min\left\{\frac{\alpha-1}{\alpha}, \frac{1}{2}\right\}}\right)$	$O(1)$	$O\left(n^{-\frac{\alpha-1}{\alpha}}\right)$
INVERSE LOG POLYNOMIAL	$O\left(\frac{1}{\log^\beta n}\right)$	$o(1)$	$O\left(\frac{1}{\log^\beta n}\right)$
CONSTANT	$O\left(n^{-\frac{1}{2}}\right)$	$O\left(n^{-\min\left\{\varepsilon, \frac{1}{2}\right\}}\right)$	$O\left(n^{-\min\{\varepsilon, 1\}}\right)$
PIECEWISE CONSTANT	$O\left(n^{-\min\left\{1-r, \frac{1}{2}\right\}}\right)$	$O\left(n^{-\min\left\{1-r, q-1, \frac{1}{2}\right\}}\right)$	$O\left(n^{-\min\{1-r, q-1\}}\right)$

Furthermore, for any positive constant δ, with probability at least $1 - \delta$, the minimal excess risk on the training trajectory can be bounded as

$$\min_t R(\boldsymbol{\theta}_t) \lesssim \frac{\max\{\sqrt{r(\boldsymbol{\Sigma})}, 1\}}{\sqrt{n}} + \frac{\max\{k_1, 1\}}{n}.$$

Lemma 5.1 below shows that $k_1 = o(n)$ always holds for fixed distribution. Therefore, combining Corollary 5.1 and Lemma 5.1 completes the proof of Theorem 4.1.

Lemma 5.1. *For any fixed (i.e. independent of sample size n) feature covariance $\boldsymbol{\Sigma}$ satisfying assumption 1, we have $k_1(n) = o(n)$.*

Example Distributions We apply the bound in Corollary 5.1 to several data distributions. These distributions are widely discussed in (Bartlett et al., 2019; Zou et al., 2021). We also derive the existing excess risk bounds, which focus on the min-norm interpolator (Bartlett et al., 2019) and one-pass SGD iterate (Zou et al., 2021), of these distributions and compare them with our theorem. The results are summarized in Table 1, which shows that the bound in Corollary 5.1 outperforms previous results for a general class of distributions. We defer the details to Appendix B.

5.3 VARYING t, VARYING $c(t, n)$

Although setting $c(t, n)$ to a constant as in Corollary 5.1 suffices to prove Theorem 4.1, in this section we show that the choice of $c(t, n)$ can be much more flexible. Specifically, we provide a concrete example and demonstrate that by setting $c(t, n)$ to a non-constant, Theorem 5.1 can indeed produce larger compatibility regions.

Example 5.1. *Under the same conditions as Theorem 5.1, let $\boldsymbol{\Sigma}$ denote the feature covariance matrix. If the spectrum of $\boldsymbol{\Sigma}$ satisfies $\lambda_k = \frac{1}{k^\alpha}$ for some $\alpha > 1$, we set $c(t, n) = \Theta\left(n^{\frac{\alpha+1-2\alpha\tau}{2\alpha+1}}\right)$ for a given $\frac{\alpha+1}{2\alpha} \le \tau \le \frac{3\alpha+1}{2\alpha+2}$. Then for $t = \Theta(n^\tau)$, we derive that $V(\boldsymbol{\theta}_t) = O\left(n^{\frac{2\alpha\tau-3\alpha+2\tau-1}{2\alpha+1}}\right)$.*

Example 5.1 shows that by choosing $c(t, n)$ as a non-constant, we exploit the full power of Theorem 5.1, and extend the compatibility region to $t = o\left(n^{\frac{3\alpha+1}{2\alpha+2}}\right) = \omega(n)$. In this example, Theorem 5.1 outperforms all $O\left(\frac{t}{n}\right)$-type bounds, which become vacuous when $t = \omega(n)$.

6 EXPERIMENTS

In this section, we provide numerical studies of overparameterized linear regression problems. The empirical results **(a.) demonstrate the benefits of trajectory analysis underlying the definition**

Table 2: **The effective dimension k_1, the optimal early stopping excess risk, and the min-norm excess risk for different feature distributions.** The table shows that early stopping solutions generalize significantly better than min-norm interpolators, and reveals a positive correlation between the effective dimension k_1 and excess risk of early stopping solution. We calculate the 95% confidence interval for each excess risk.

DISTRIBUTIONS	k_1	OPTIMAL EXCESS RISK	MIN-NORM EXCESS RISK
$\lambda_i = \frac{1}{i}$	$\Theta(n)$	2.399 ± 0.0061	24.071 ± 0.2447
$\lambda_i = \frac{1}{i^2}$	$\Theta\left(n^{\frac{1}{2}}\right)$	0.214 ± 0.0050	43.472 ± 0.6463
$\lambda_i = \frac{1}{i^3}$	$\Theta\left(n^{\frac{1}{3}}\right)$	0.077 ± 0.0005	10.401 ± 0.2973
$\lambda_i = \frac{1}{i \log(i+1)}$	$\Theta\left(\frac{n}{\log n}\right)$	0.697 ± 0.0053	89.922 ± 0.9591
$\lambda_i = \frac{1}{i \log^2(i+1)}$	$\Theta\left(\frac{n}{\log^2 n}\right)$	0.298 ± 0.0054	82.413 ± 0.9270
$\lambda_i = \frac{1}{i \log^3(i+1)}$	$\Theta\left(\frac{n}{\log^3 n}\right)$	0.187 ± 0.0047	38.145 ± 0.5862

of compatibility, since the optimal excess risk along the algorithm trajectory is significantly lower than that of the min-norm interpolator **(b.) validate the statements in Corollary 5.1**, since the optimal excess risk is lower when the eigenvalues of feature covariance decay faster. We refer to Appendix C for additional results and discussions.

Setup. We consider six overparameterized linear regression instances with input dimension $p = 1000$ and sample size $n = 100$. The feature vectors are independently sampled from zero-mean Gaussian distributions, whose covariances are diagonal with entries $\lambda_i = \frac{1}{i}, \frac{1}{i^2}, \frac{1}{i^3}, \frac{1}{i \log(i+1)}, \frac{1}{i \log^2(i+1)}, \frac{1}{i \log^3(i+1)}$, respectively. For each feature $\boldsymbol{x}$, we construct the response as $y = \boldsymbol{x}^\top \boldsymbol{\theta}^* + \varepsilon$, where $\boldsymbol{\theta}^*$ is sampled from a p-dimensional standard Gaussian distribution for each instance, and ε is sampled from a standard Gaussian distribution. We conduct gradient descent with learning rate $\lambda = 0.001$ on the above instances. We also compute min-norm excess risk via its closed form, with 1×10^{-4} weight decay on parameter $\boldsymbol{\theta}$ to avoid numerical instability. Note that the regularization will only reduce the final excess and will not jeopardize our final conclusion's correctness.

Observation One: Early stopping solution along the training trajectory generalizes significantly better than the min-norm interpolator. We calculate the excess risk of optimal early stopping solutions and min-norm interpolators from 1000 independent trials and list the results in Table 2. The results illustrate that the early stopping solution on the algorithm trajectory enjoys much better generalization properties. This observation corroborates the meaningfulness of compatibility and the importance of data-dependent training trajectory in generalization analysis.

Observation Two: The faster covariance spectrum decays, the lower optimal excess risk is. Table 2 also illustrates a positive correlation between the decaying rate of λ_i and the generalization performance of the early stopping solution. This accords with Theorem 5.1, showing that the excess risk is better for a smaller effective dimension k_1, where small k_1 indicates a faster-decaying eigenvalue λ_i. We additionally note that such a phenomenon also illustrates the difference between min-norm and early stopping solutions in linear regression, since Bartlett et al. (2019) demonstrate that the min-norm solution is not consistent when the eigenvalues decay too fast. By comparison, early stopping solutions do not suffer from this restriction.

7 CONCLUSION

In this paper, we propose the concept of data-algorithm compatibility, and study compatibility for overparameterized linear regression with gradient descent. Our theoretical and empirical results demonstrate that compatibility eases the assumptions and broadens the scope of generalization by fully exploiting the data information and the algorithm information. Despite linear cases in this paper, compatibility can be a much more general concept. Therefore, we believe this paper will motivate more work on data-dependent trajectory analysis.

REFERENCES

Ali, A., Kolter, J. Z., and Tibshirani, R. J. (2019). A continuous-time view of early stopping for least squares regression. In Chaudhuri, K. and Sugiyama, M., editors, *The 22nd International Conference on Artificial Intelligence and Statistics, AISTATS 2019, 16-18 April 2019, Naha, Okinawa, Japan*, volume 89 of *Proceedings of Machine Learning Research*, pages 1370–1378. PMLR.

Arora, S., Du, S. S., Hu, W., Li, Z., and Wang, R. (2019). Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In Chaudhuri, K. and Salakhutdinov, R., editors, *Proceedings of the 36th International Conference on Machine Learning, ICML 2019, 9-15 June 2019, Long Beach, California, USA*, volume 97 of *Proceedings of Machine Learning Research*, pages 322–332. PMLR.

Arora, S., Ge, R., Neyshabur, B., and Zhang, Y. (2018). Stronger generalization bounds for deep nets via a compression approach. In Dy, J. G. and Krause, A., editors, *Proceedings of the 35th International Conference on Machine Learning, ICML 2018, Stockholmsmässan, Stockholm, Sweden, July 10-15, 2018*, volume 80 of *Proceedings of Machine Learning Research*, pages 254–263. PMLR.

Bai, Y., Yang, E., Han, B., Yang, Y., Li, J., Mao, Y., Niu, G., and Liu, T. (2021). Understanding and improving early stopping for learning with noisy labels. *CoRR*, abs/2106.15853.

Banerjee, P. K. and Montúfar, G. (2021). Information complexity and generalization bounds. *CoRR*, abs/2105.01747.

Bartlett, P. L., Foster, D. J., and Telgarsky, M. (2017). Spectrally-normalized margin bounds for neural networks. In Guyon, I., von Luxburg, U., Bengio, S., Wallach, H. M., Fergus, R., Vishwanathan, S. V. N., and Garnett, R., editors, *Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, December 4-9, 2017, Long Beach, CA, USA*, pages 6240–6249.

Bartlett, P. L., Long, P. M., Lugosi, G., and Tsigler, A. (2019). Benign overfitting in linear regression. *CoRR*, abs/1906.11300.

Bassily, R., Feldman, V., Guzmán, C., and Talwar, K. (2020). Stability of stochastic gradient descent on nonsmooth convex losses. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M., and Lin, H., editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*.

Bousquet, O. and Elisseeff, A. (2002). Stability and generalization. *J. Mach. Learn. Res.*, 2:499–526.

Bousquet, O., Klochkov, Y., and Zhivotovskiy, N. (2020). Sharper bounds for uniformly stable algorithms. In Abernethy, J. D. and Agarwal, S., editors, *Conference on Learning Theory, COLT 2020, 9-12 July 2020, Virtual Event [Graz, Austria]*, volume 125 of *Proceedings of Machine Learning Research*, pages 610–626. PMLR.

Brown, T. B., Mann, B., Ryder, N., Subbiah, M., Kaplan, J., Dhariwal, P., Neelakantan, A., Shyam, P., Sastry, G., Askell, A., Agarwal, S., Herbert-Voss, A., Krueger, G., Henighan, T., Child, R., Ramesh, A., Ziegler, D. M., Wu, J., Winter, C., Hesse, C., Chen, M., Sigler, E., Litwin, M., Gray, S., Chess, B., Clark, J., Berner, C., McCandlish, S., Radford, A., Sutskever, I., and Amodei, D. (2020). Language models are few-shot learners. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M., and Lin, H., editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*.

Devlin, J., Chang, M., Lee, K., and Toutanova, K. (2019). BERT: pre-training of deep bidirectional transformers for language understanding. In Burstein, J., Doran, C., and Solorio, T., editors, *Proceedings of the 2019 Conference of the North American Chapter of the Association for Computational Linguistics: Human Language Technologies, NAACL-HLT 2019, Minneapolis, MN, USA, June 2-7, 2019, Volume 1 (Long and Short Papers)*, pages 4171–4186. Association for Computational Linguistics.

Dziugaite, G. K. and Roy, D. M. (2017). Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. In Elidan, G., Kersting, K., and Ihler, A. T., editors, *Proceedings of the Thirty-Third Conference on Uncertainty in Artificial Intelligence, UAI 2017, Sydney, Australia, August 11-15, 2017*. AUAI Press.

El Karoui, N. (2010). The spectrum of kernel random matrices. *The Annals of Statistics*, 38(1):1–50.

Feldman, V. and Vondrák, J. (2018). Generalization bounds for uniformly stable algorithms. In Bengio, S., Wallach, H. M., Larochelle, H., Grauman, K., Cesa-Bianchi, N., and Garnett, R., editors, *Advances in Neural Information Processing Systems 31: Annual Conference on Neural Information Processing Systems 2018, NeurIPS 2018, December 3-8, 2018, Montréal, Canada*, pages 9770–9780.

Feldman, V. and Vondrák, J. (2019). High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. In Beygelzimer, A. and Hsu, D., editors, *Conference on Learning Theory, COLT 2019, 25-28 June 2019, Phoenix, AZ, USA*, volume 99 of *Proceedings of Machine Learning Research*, pages 1270–1279. PMLR.

Forouzesh, M. and Thiran, P. (2021). Disparity between batches as a signal for early stopping. In Oliver, N., Pérez-Cruz, F., Kramer, S., Read, J., and Lozano, J. A., editors, *Machine Learning and Knowledge Discovery in Databases. Research Track - European Conference, ECML PKDD 2021, Bilbao, Spain, September 13-17, 2021, Proceedings, Part II*, volume 12976 of *Lecture Notes in Computer Science*, pages 217–232. Springer.

Frei, S., Chatterji, N. S., and Bartlett, P. L. (2022). Benign overfitting without linearity: Neural network classifiers trained by gradient descent for noisy linear data. *CoRR*, abs/2202.05928.

Hardt, M., Recht, B., and Singer, Y. (2016). Train faster, generalize better: Stability of stochastic gradient descent. In Balcan, M. and Weinberger, K. Q., editors, *Proceedings of the 33nd International Conference on Machine Learning, ICML 2016, New York City, NY, USA, June 19-24, 2016*, volume 48 of *JMLR Workshop and Conference Proceedings*, pages 1225–1234. JMLR.org.

Hu, W., Xiao, L., Adlam, B., and Pennington, J. (2020). The surprising simplicity of the early-time learning dynamics of neural networks. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M., and Lin, H., editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*.

Jacot, A., Hongler, C., and Gabriel, F. (2018). Neural tangent kernel: Convergence and generalization in neural networks. In Bengio, S., Wallach, H. M., Larochelle, H., Grauman, K., Cesa-Bianchi, N., and Garnett, R., editors, *Advances in Neural Information Processing Systems 31: Annual Conference on Neural Information Processing Systems 2018, NeurIPS 2018, December 3-8, 2018, Montréal, Canada*, pages 8580–8589.

Ji, Z., Li, J. D., and Telgarsky, M. (2021). Early-stopped neural networks are consistent. In Ranzato, M., Beygelzimer, A., Dauphin, Y. N., Liang, P., and Vaughan, J. W., editors, *Advances in Neural Information Processing Systems 34: Annual Conference on Neural Information Processing Systems 2021, NeurIPS 2021, December 6-14, 2021, virtual*, pages 1805–1817.

Jiang, Y., Neyshabur, B., Mobahi, H., Krishnan, D., and Bengio, S. (2020). Fantastic generalization measures and where to find them. In *8th International Conference on Learning Representations, ICLR 2020, Addis Ababa, Ethiopia, April 26-30, 2020*. OpenReview.net.

Koltchinskii, V. and Panchenko, D. (2000). Rademacher processes and bounding the risk of function learning. In *High dimensional probability II*, pages 443–457. Springer.

Lei, Y. and Ying, Y. (2020). Fine-grained analysis of stability and generalization for stochastic gradient descent. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pages 5809–5819. PMLR.

Li, J., Luo, X., and Qiao, M. (2020a). On generalization error bounds of noisy gradient methods for non-convex learning. In *8th International Conference on Learning Representations, ICLR 2020, Addis Ababa, Ethiopia, April 26-30, 2020*. OpenReview.net.

Li, M., Soltanolkotabi, M., and Oymak, S. (2020b). Gradient descent with early stopping is provably robust to label noise for overparameterized neural networks. In Chiappa, S. and Calandra, R., editors, *The 23rd International Conference on Artificial Intelligence and Statistics, AISTATS 2020, 26-28 August 2020, Online [Palermo, Sicily, Italy]*, volume 108 of *Proceedings of Machine Learning Research*, pages 4313–4324. PMLR.

Li, Z., Su, W., and Sejdinovic, D. (2020c). Benign overfitting and noisy features. *CoRR*, abs/2008.02901.

Liang, T. and Rakhlin, A. (2018). Just interpolate: Kernel "ridgeless" regression can generalize. *CoRR*, abs/1808.00387.

Lyu, K. and Li, J. (2020). Gradient descent maximizes the margin of homogeneous neural networks. In *8th International Conference on Learning Representations, ICLR 2020, Addis Ababa, Ethiopia, April 26-30, 2020*. OpenReview.net.

Lyu, K., Li, Z., Wang, R., and Arora, S. (2021). Gradient descent on two-layer nets: Margin maximization and simplicity bias. In Ranzato, M., Beygelzimer, A., Dauphin, Y. N., Liang, P., and Vaughan, J. W., editors, *Advances in Neural Information Processing Systems 34: Annual Conference on Neural Information Processing Systems 2021, NeurIPS 2021, December 6-14, 2021, virtual*, pages 12978–12991.

Mei, S. and Montanari, A. (2022). The generalization error of random features regression: Precise asymptotics and the double descent curve. *Communications on Pure and Applied Mathematics*, 75(4):667–766.

Mohri, M., Rostamizadeh, A., and Talwalkar, A. (2012). *Foundations of Machine Learning*. Adaptive computation and machine learning. MIT Press.

Mou, W., Wang, L., Zhai, X., and Zheng, K. (2018). Generalization bounds of SGLD for non-convex learning: Two theoretical viewpoints. In Bubeck, S., Perchet, V., and Rigollet, P., editors, *Conference On Learning Theory, COLT 2018, Stockholm, Sweden, 6-9 July 2018*, volume 75 of *Proceedings of Machine Learning Research*, pages 605–638. PMLR.

Nagarajan, V. and Kolter, J. Z. (2019). Uniform convergence may be unable to explain generalization in deep learning. In Wallach, H. M., Larochelle, H., Beygelzimer, A., d'Alché-Buc, F., Fox, E. B., and Garnett, R., editors, *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, December 8-14, 2019, Vancouver, BC, Canada*, pages 11611–11622.

Negrea, J., Dziugaite, G. K., and Roy, D. (2020). In defense of uniform convergence: Generalization via derandomization with an application to interpolating predictors. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pages 7263–7272. PMLR.

Neyshabur, B., Bhojanapalli, S., and Srebro, N. (2018). A pac-bayesian approach to spectrally-normalized margin bounds for neural networks. In *6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Conference Track Proceedings*. OpenReview.net.

Parrado-Hernández, E., Ambroladze, A., Shawe-Taylor, J., and Sun, S. (2012). Pac-bayes bounds with data dependent priors. *J. Mach. Learn. Res.*, 13:3507–3531.

Prechelt, L. (2012). Early stopping - but when? In Montavon, G., Orr, G. B., and Müller, K., editors, *Neural Networks: Tricks of the Trade - Second Edition*, volume 7700 of *Lecture Notes in Computer Science*, pages 53–67. Springer.

Russo, D. and Zou, J. (2016). Controlling bias in adaptive data analysis using information theory. In Gretton, A. and Robert, C. C., editors, *Proceedings of the 19th International Conference on Artificial Intelligence and Statistics, AISTATS 2016, Cadiz, Spain, May 9-11, 2016*, volume 51 of *JMLR Workshop and Conference Proceedings*, pages 1232–1240. JMLR.org.

Shah, H., Tamuly, K., Raghunathan, A., Jain, P., and Netrapalli, P. (2020). The pitfalls of simplicity bias in neural networks. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M., and Lin, H., editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*.

Shankar, V., Roelofs, R., Mania, H., Fang, A., Recht, B., and Schmidt, L. (2020). Evaluating machine accuracy on imagenet. In *Proceedings of the 37th International Conference on Machine Learning, ICML 2020, 13-18 July 2020, Virtual Event*, volume 119 of *Proceedings of Machine Learning Research*, pages 8634–8644. PMLR.

Silver, D., Schrittwieser, J., Simonyan, K., Antonoglou, I., Huang, A., Guez, A., Hubert, T., Baker, L., Lai, M., Bolton, A., Chen, Y., Lillicrap, T. P., Hui, F., Sifre, L., van den Driessche, G., Graepel, T., and Hassabis, D. (2017). Mastering the game of go without human knowledge. *Nat.*, 550(7676):354–359.

Soudry, D., Hoffer, E., Nacson, M. S., and Srebro, N. (2018). The implicit bias of gradient descent on separable data. In *6th International Conference on Learning Representations, ICLR 2018, Vancouver, BC, Canada, April 30 - May 3, 2018, Conference Track Proceedings*. OpenReview.net.

Stock, P. and Cissé, M. (2018). Convnets and imagenet beyond accuracy: Understanding mistakes and uncovering biases. In Ferrari, V., Hebert, M., Sminchisescu, C., and Weiss, Y., editors, *Computer Vision - ECCV 2018 - 15th European Conference, Munich, Germany, September 8-14, 2018, Proceedings, Part VI*, volume 11210 of *Lecture Notes in Computer Science*, pages 504–519. Springer.

Teng, J., Ma, J., and Yuan, Y. (2021). Towards understanding generalization via decomposing excess risk dynamics. *CoRR*, abs/2106.06153.

Tsigler, A. and Bartlett, P. L. (2020). Benign overfitting in ridge regression. *arXiv preprint arXiv:2009.14286*.

Wang, K. and Thrampoulidis, C. (2021). Benign overfitting in binary classification of gaussian mixtures. In *IEEE International Conference on Acoustics, Speech and Signal Processing, ICASSP 2021, Toronto, ON, Canada, June 6-11, 2021*, pages 4030–4034. IEEE.

Xu, A. and Raginsky, M. (2017). Information-theoretic analysis of generalization capability of learning algorithms. In Guyon, I., von Luxburg, U., Bengio, S., Wallach, H. M., Fergus, R., Vishwanathan, S. V. N., and Garnett, R., editors, *Advances in Neural Information Processing Systems 30: Annual Conference on Neural Information Processing Systems 2017, December 4-9, 2017, Long Beach, CA, USA*, pages 2524–2533.

Yao, Y., Rosasco, L., and Caponnetto, A. (2007). On early stopping in gradient descent learning. *Constructive Approximation*, 26(2):289–315.

Yun, S., Oh, S. J., Heo, B., Han, D., Choe, J., and Chun, S. (2021). Re-labeling imagenet: From single to multi-labels, from global to localized labels. In *IEEE Conference on Computer Vision and Pattern Recognition, CVPR 2021, virtual, June 19-25, 2021*, pages 2340–2350. Computer Vision Foundation / IEEE.

Zhang, C., Bengio, S., Hardt, M., Recht, B., and Vinyals, O. (2021). Understanding deep learning (still) requires rethinking generalization. *Commun. ACM*, 64(3):107–115.

Zhou, L., Sutherland, D. J., and Srebro, N. (2020). On uniform convergence and low-norm interpolation learning. In Larochelle, H., Ranzato, M., Hadsell, R., Balcan, M., and Lin, H., editors, *Advances in Neural Information Processing Systems 33: Annual Conference on Neural Information Processing Systems 2020, NeurIPS 2020, December 6-12, 2020, virtual*.

Zou, D., Wu, J., Braverman, V., Gu, Q., and Kakade, S. M. (2021). Benign overfitting of constant-stepsize SGD for linear regression. In Belkin, M. and Kpotufe, S., editors, *Conference on Learning Theory, COLT 2021, 15-19 August 2021, Boulder, Colorado, USA*, volume 134 of *Proceedings of Machine Learning Research*, pages 4633–4635. PMLR.

Supplementary Materials

A PROOFS FOR THE MAIN RESULTS

We first sketch the proof in section A.1 and give some preliminary lemmas A.2. The following sections A.3, A.4 and A.5 are devoted to the proof of Theorem 5.1. The proof of Theorem 4.1 is given in A.6.

A.1 PROOF SKETCH

We start with a standard bias-variance decomposition following Bartlett et al. (2019), which derives that the time-variant excess risk $R(\boldsymbol{\theta}_t)$ can be bounded by a bias term and a variance term. We refer to Appendix A.3 for more details.

For the bias part, we first decompose it into an optimization error and an approximation error. For the optimization error, we use the spectrum analysis to bound it with $O\left(1/t\right)$ where t denotes the time. For the approximation error, we bound it with $O\left(1/\sqrt{n}\right)$ where n denotes the sample size, inspired by Bartlett et al. (2019). We refer to Appendix A.4 for more details.

For the variance part, a key step is to bound the term $(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top)^t$, where $\boldsymbol{X}$ is the feature matrix. The difficulty arises from the different scales of the eigenvalues of $\boldsymbol{X}\boldsymbol{X}^\top$, where the largest eigenvalue has order $\Theta(n)$ while the smallest eigenvalue has order $O(1)$, according to Lemma 10 in Bartlett et al. (2019). To overcome this issue, we divide the matrix $\boldsymbol{X}\boldsymbol{X}^\top$ based on whether its eigenvalues is larger than $c(t,n)$, which is a flexible term dependent on time t and sample size n. Therefore, we split the variance term based on eigenvalues of covariance matrix $\boldsymbol{\Sigma}$ (leading to the k_1-related term) and based on the eigenvalues of $\boldsymbol{X}\boldsymbol{X}^\top$ (leading to the k_2-related term). We refer to Appendix A.5 for more details.

A.2 PRELIMINARIES

Notations. We use O, o, Ω, ω as asymptotic notations, with their usual meaning. We use $\lesssim$ with the same meaning as the asymptotic notation O. We use $\|\boldsymbol{x}\|$ to denote the ℓ_2 norm for vector $\boldsymbol{x}$ and $\|\boldsymbol{A}\|$ to denote the operator norm for matrix $\boldsymbol{A}$. We allow the vector to belong to a countably infinite-dimensional Hilbert space $\mathcal{H}$, and with a slight abuse of notation, we use $\mathbb{R}^\infty$ interchangeably with $\mathcal{H}$. In this case, $x^\top z$ denotes inner product and $xz^\top$ denotes tensor product for $x, z \in \mathcal{H}$ as in Bartlett et al. (2019).

The following result comes from Bartlett et al. (2019), which bounds the eigenvalues of $\boldsymbol{X}\boldsymbol{X}^\top$.

Lemma A.1. *(Lemma 10 in Bartlett et al. (2019)) For any σ_x, there exists a constant c, such that for any $0 \le k < n$, with probability at least $1 - e^{-\frac{n}{c}}$,*

$$\mu_{k+1} \le c\left(\sum_{i>k} \lambda_i + \lambda_{k+1}n\right). \tag{5}$$

This implies that as long as the step size λ is small than a threshold independent of sample size n, gradient descent is stable.

Corollary A.1. *There exists a constant c, such that with probability at least $1 - e^{-\frac{n}{c}}$, for any $0 \le \lambda \le \frac{1}{c\sum_{i>0} \lambda_i}$ we have*

$$\boldsymbol{O} \preceq \boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X} \preceq \boldsymbol{I}. \tag{6}$$

Proof. The right hand side of the inequality is obvious since $\lambda > 0$. For the left hand side, we have to show that the eigenvalues of $\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}$ is non-negative. since $\boldsymbol{X}^\top\boldsymbol{X}$ and $\boldsymbol{X}\boldsymbol{X}^\top$ have the same non-zero eigenvalues, we know that with probability at least $1 - e^{-\frac{n}{c}}$, the smallest eigenvalue of $\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}$ can be lower bounded by

$$\mu_p \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) = 1 - \frac{\lambda}{n} \mu_1$$

$$\geq 1 - c\lambda \left(\frac{\sum_{i>0} \lambda_i}{n} + \lambda_{k+1} \right)$$

$$\geq 1 - 2c\lambda \sum_{i>0} \lambda_i \tag{7}$$

$$\geq 0.$$

where the second inequality uses lemma A.1, and the last inequality holds if $\lambda \leq \frac{1}{2c \sum_{i>0} \lambda_i}$. □

A.3 PROOF FOR THE BIAS-VARIANCE DECOMPOSITION

Let $\boldsymbol{X}^\dagger$ denote the Moore–Penrose pseudoinverse of matrix $\boldsymbol{X}$. The following lemma gives a closed form expression for $\boldsymbol{\theta}_t$.

Lemma A.2. *The dynamics of $\{\boldsymbol{\theta}_t\}_{t\geq0}$ satisfies*

$$\boldsymbol{\theta}_t = \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t (\boldsymbol{\theta}_0 - \boldsymbol{X}^\dagger \boldsymbol{Y}) + \boldsymbol{X}^\dagger \boldsymbol{Y}. \tag{8}$$

Proof. We prove the lemma using induction. The equality holds at $t = 0$ as both sides are $\boldsymbol{\theta}_0$. Recall that $\boldsymbol{\theta}_t$ is updated as

$$\boldsymbol{\theta}_{t+1} = \boldsymbol{\theta}_t + \frac{\lambda}{n} \boldsymbol{X}^\top (\boldsymbol{Y} - \boldsymbol{X} \boldsymbol{\theta}_t). \tag{9}$$

Suppose that the dynamic holds up to the t-th step. Plug the expression for $\boldsymbol{\theta}_t$ into the above recursion and note that $\boldsymbol{X}^\top \boldsymbol{X} \boldsymbol{X}^\dagger = \boldsymbol{X}^\top$, we get

$$\begin{aligned}
\boldsymbol{\theta}_{t+1} &= \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \boldsymbol{\theta}_t + \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{Y} \\
&= \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^{t+1} (\boldsymbol{\theta}_0 - \boldsymbol{X}^\dagger \boldsymbol{Y}) + \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \boldsymbol{X}^\dagger \boldsymbol{Y} + \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{Y} \\
&= \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^{t+1} (\boldsymbol{\theta}_0 - \boldsymbol{X}^\dagger \boldsymbol{Y}) + \boldsymbol{X}^\dagger \boldsymbol{Y}.
\end{aligned} \tag{10}$$

which finishes the proof. □

Next we prove two identities which will be used in further proof.

Lemma A.3. *The following two identities hold for any matrix X and non-negative integer t:*

$$\boldsymbol{I} - \boldsymbol{X}^\dagger \boldsymbol{X} + \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \boldsymbol{X}^\dagger \boldsymbol{X} = \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t, \tag{11}$$

$$\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \right] \boldsymbol{X}^\dagger \boldsymbol{X} \boldsymbol{X}^\top = \boldsymbol{X}^\top \left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X} \boldsymbol{X}^\top \right)^t \right]. \tag{12}$$

Proof. Note that $\boldsymbol{X}^\top \boldsymbol{X} \boldsymbol{X}^\dagger = \boldsymbol{X}^\top$, we can expand the left hand side of the first identity above using binomial theorem and eliminate the pseudo-inverse $\boldsymbol{X}^\dagger$:

$$\boldsymbol{I} - \boldsymbol{X}^\dagger \boldsymbol{X} + \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^t \boldsymbol{X}^\dagger \boldsymbol{X}$$

$$= \boldsymbol{I} - \boldsymbol{X}^\dagger \boldsymbol{X} + \sum_{k=0}^t \binom{t}{k}\left(-\frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^k \boldsymbol{X}^\dagger \boldsymbol{X}$$

$$= \boldsymbol{I} - \boldsymbol{X}^\dagger \boldsymbol{X} + \boldsymbol{X}^\dagger \boldsymbol{X} + \sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\right)^k (\boldsymbol{X}^\top \boldsymbol{X})^{k-1}\boldsymbol{X}^\top \boldsymbol{X}\boldsymbol{X}^\dagger \boldsymbol{X} \tag{13}$$

$$= \boldsymbol{I} + \sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\right)^k (\boldsymbol{X}^\top \boldsymbol{X})^k$$

$$= \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^t.$$

The second identity can be proved in a similar way:

$$\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^t\right]\boldsymbol{X}^\dagger \boldsymbol{X}\boldsymbol{X}^\top$$

$$= -\sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^k \boldsymbol{X}^\dagger \boldsymbol{X}\boldsymbol{X}^\top$$

$$= -\sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\right)^k (\boldsymbol{X}^\top \boldsymbol{X})^{k-1}\boldsymbol{X}^\top \boldsymbol{X}\boldsymbol{X}^\dagger \boldsymbol{X}\boldsymbol{X}^\top$$

$$= -\sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\right)^k (\boldsymbol{X}^\top \boldsymbol{X})^{k-1}\boldsymbol{X}^\top \boldsymbol{X}\boldsymbol{X}^\top \tag{14}$$

$$= -\sum_{k=1}^t \binom{t}{k}\left(-\frac{\lambda}{n}\right)^k \boldsymbol{X}^\top (\boldsymbol{X}\boldsymbol{X}^\top)^k$$

$$= \boldsymbol{X}^\top \left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top\right)^t\right].$$

$\square$

We are now ready to prove the main result of this section.

Lemma A.4. *The excess risk at the t-th epoch can be upper bounded as*

$$R(\boldsymbol{\theta}_t) \leq \boldsymbol{\theta}^{*\top}\boldsymbol{B}\boldsymbol{\theta}^* + \boldsymbol{\varepsilon}^\top \boldsymbol{C}\boldsymbol{\varepsilon}, \tag{15}$$

where

$$\boldsymbol{B} = \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^t \boldsymbol{\Sigma}\left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top \boldsymbol{X}\right)^t, \tag{16}$$

$$\boldsymbol{C} = (\boldsymbol{X}\boldsymbol{X}^\top)^{-1}\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top\right)^t\right]\boldsymbol{X}\boldsymbol{\Sigma}\boldsymbol{X}^\top\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top\right)^t\right](\boldsymbol{X}\boldsymbol{X}^\top)^{-1}, \tag{17}$$

which characterizes bias term and variance term in the excess risk. Furthermore, there exists constant c such that with probability at least $1 - \delta$ over the randomness of $\boldsymbol{\varepsilon}$, we have

$$\boldsymbol{\varepsilon}^\top \boldsymbol{C}\boldsymbol{\varepsilon} \leq c\sigma_y^2 \log\frac{1}{\delta}\operatorname{Tr}[\boldsymbol{C}]. \tag{18}$$

Proof. First note that $\boldsymbol{X}\boldsymbol{X}^\top$ is invertible by Assumption 3. Express the excess risk as follows

$$
\begin{aligned}
R(\boldsymbol{\theta}_t) &= \frac{1}{2}\mathbb{E}[(y - \boldsymbol{x}^\top\boldsymbol{\theta}_t)^2 - (y - \boldsymbol{x}^\top\boldsymbol{\theta}^*)^2] \\
&= \frac{1}{2}\mathbb{E}[(y - \boldsymbol{x}^\top\boldsymbol{\theta}^* + \boldsymbol{x}^\top\boldsymbol{\theta}^* - \boldsymbol{x}^\top\boldsymbol{\theta}_t)^2 - (y - \boldsymbol{x}^\top\boldsymbol{\theta}^*)^2] \\
&= \frac{1}{2}\mathbb{E}[(\boldsymbol{x}^\top(\boldsymbol{\theta}_t - \boldsymbol{\theta}^*))^2 + 2(y - \boldsymbol{x}^\top\boldsymbol{\theta}^*)(\boldsymbol{x}^\top\boldsymbol{\theta}^* - \boldsymbol{x}^\top\boldsymbol{\theta}_t)] \\
&= \frac{1}{2}\mathbb{E}[\boldsymbol{x}^\top(\boldsymbol{\theta}_t - \boldsymbol{\theta}^*)]^2.
\end{aligned}
\tag{19}
$$

Recall that $\boldsymbol{\theta}_0 = 0$ and $\boldsymbol{Y} = \boldsymbol{X}\boldsymbol{\theta}^* + \boldsymbol{\varepsilon}$ and we can further simplify the formula for $\boldsymbol{\theta}_t$ in lemma A.2:

$$
\begin{aligned}
\boldsymbol{\theta}_t &= \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t(\boldsymbol{\theta}_0 - \boldsymbol{X}^\dagger\boldsymbol{Y}) + \boldsymbol{X}^\dagger\boldsymbol{Y} \\
&= \left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right]\boldsymbol{X}^\dagger(\boldsymbol{X}\boldsymbol{\theta}^* + \boldsymbol{\varepsilon}).
\end{aligned}
\tag{20}
$$

Plug it into the above expression for $R(\boldsymbol{\theta}_t)$, we have

$$
\begin{aligned}
R(\boldsymbol{\theta}_t) &= \frac{1}{2}\mathbb{E}\left[\boldsymbol{x}^\top\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right]\boldsymbol{X}^\dagger(\boldsymbol{X}\boldsymbol{\theta}^* + \boldsymbol{\varepsilon}) - \boldsymbol{x}^\top\boldsymbol{\theta}^*\right]^2 \\
&= \frac{1}{2}\mathbb{E}\left[\boldsymbol{x}^\top\left(\boldsymbol{X}^\dagger\boldsymbol{X} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\boldsymbol{X}^\dagger\boldsymbol{X} - \boldsymbol{I}\right)\boldsymbol{\theta}^*\right. \\
&\quad\left. + \boldsymbol{x}^\top\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right]\boldsymbol{X}^\dagger\boldsymbol{\varepsilon}\right]^2.
\end{aligned}
\tag{21}
$$

Applying lemma A.3, we obtain

$$
\begin{aligned}
R(\boldsymbol{\theta}_t) &= \frac{1}{2}\mathbb{E}\left[-\boldsymbol{x}^\top\left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\boldsymbol{\theta}^* + \boldsymbol{x}^\top\boldsymbol{X}^\top\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top\right)^t\right](\boldsymbol{X}\boldsymbol{X}^\top)^{-1}\boldsymbol{\varepsilon}\right]^2 \\
&\leq \mathbb{E}\left[\boldsymbol{x}^\top\left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\boldsymbol{\theta}^*\right]^2 + \mathbb{E}\left[\boldsymbol{x}^\top\boldsymbol{X}^\top\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^\top\right)^t\right](\boldsymbol{X}\boldsymbol{X}^\top)^{-1}\boldsymbol{\varepsilon}\right]^2 \\
&:= \boldsymbol{\theta}^{*\top}\boldsymbol{B}\boldsymbol{\theta}^* + \boldsymbol{\varepsilon}^\top\boldsymbol{C}\boldsymbol{\varepsilon}.
\end{aligned}
\tag{22}
$$

which proves the first claim in the lemma. The second part of the theorem directly follows from lemma 18 in Bartlett et al. (2019). □

A.4 PROOF FOR THE BIAS UPPER BOUND

The next lemma guarantees that the sample covariace matrix $\frac{1}{n}\boldsymbol{X}^\top\boldsymbol{X}$ concentrates well around $\boldsymbol{\Sigma}$.

Lemma A.5. *(Lemma 35 in Bartlett et al. (2019)) There exists constant c such that for any $0 < \delta < 1$ with probability as least $1 - \delta$,*

$$
\left\|\boldsymbol{\Sigma} - \frac{1}{n}\boldsymbol{X}^\top\boldsymbol{X}\right\| \leq c\|\boldsymbol{\Sigma}\|\max\left\{\sqrt{\frac{r(\boldsymbol{\Sigma})}{n}}, \frac{r(\boldsymbol{\Sigma})}{n}, \sqrt{\frac{\log(\frac{1}{\delta})}{n}}, \frac{\log(\frac{1}{\delta})}{n}\right\}.
\tag{23}
$$

The following inequality will be useful in our proof to characterize the decaying rate of the bias term with t.

Lemma A.6. *For any positive semidefinite matrix $\boldsymbol{P}$ which satisfies $\|\boldsymbol{P}\| \leq 1$, we have*

$$\|\boldsymbol{P}(1 - \boldsymbol{P})^t\| \leq \frac{1}{t}. \tag{24}$$

Proof. Assume without loss of generality that $\boldsymbol{P}$ is diagonal. Then it suffices to consider seperately each eigenvalue σ of $\boldsymbol{P}$, and show that $\sigma(1 - \sigma)^t \leq \frac{1}{t}$.

In fact, by AM-GM inequality we have

$$\sigma(1 - \sigma)^t \leq \frac{1}{t} \left[\frac{t\sigma + (1 - \sigma)t}{t + 1} \right]^{t+1} \leq \frac{1}{t}, \tag{25}$$

which completes the proof. $\qquad\square$

Next we prove the main result of this section.

Lemma A.7. *There exists constant c such that if $0 \leq \lambda \leq \frac{1}{c \sum_{i>0} \lambda_i}$, then for any $0 < \delta < 1$, with probability at least $1 - \delta$ the following bound on the bias term holds for any t*

$$\boldsymbol{\theta}^{*\top} \boldsymbol{B} \boldsymbol{\theta}^* \leq c \|\boldsymbol{\theta}^*\|^2 \left(\frac{1}{\lambda t} + \|\boldsymbol{\Sigma}\| \max \left\{ \sqrt{\frac{r(\boldsymbol{\Sigma})}{n}}, \frac{r(\boldsymbol{\Sigma})}{n}, \sqrt{\frac{\log(\frac{1}{\delta})}{n}}, \frac{\log(\frac{1}{\delta})}{n} \right\} \right). \tag{26}$$

Proof. The bias can be decomposed into the following two terms

$$
\begin{aligned}
\boldsymbol{\theta}^{*\top} \boldsymbol{B} \boldsymbol{\theta}^* &= \boldsymbol{\theta}^{*\top} \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \left(\boldsymbol{\Sigma} - \frac{1}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \boldsymbol{\theta}^* \\
&+ \boldsymbol{\theta}^{*\top} \left(\frac{1}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^{2t} \boldsymbol{\theta}^*.
\end{aligned}
\tag{27}
$$

For sufficiently small learning rate λ as given by corollary A.1, we know that with high probability

$$\left\| \boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right\| \leq 1, \tag{28}$$

which together with lemma A.5 gives a high probability bound on the first term:

$$
\begin{aligned}
&\boldsymbol{\theta}^{*\top} \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \left(\boldsymbol{\Sigma} - \frac{1}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^t \boldsymbol{\theta}^* \\
&\leq c \|\boldsymbol{\Sigma}\| \|\boldsymbol{\theta}^*\|^2 \max \left\{ \sqrt{\frac{r(\boldsymbol{\Sigma})}{n}}, \frac{r(\boldsymbol{\Sigma})}{n}, \sqrt{\frac{\log(\frac{1}{\delta})}{n}}, \frac{\log(\frac{1}{\delta})}{n} \right\}.
\end{aligned}
\tag{29}
$$

For the second term, invoke lemma A.6 with $\boldsymbol{P} = \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X}$ and we get

$$
\begin{aligned}
\boldsymbol{\theta}^{*\top} \left(\frac{1}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^{2t} \boldsymbol{\theta}^* &\leq \frac{1}{\lambda} \|\boldsymbol{\theta}^*\|^2 \left\| \left(\frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right) \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}^\top \boldsymbol{X} \right)^{2t} \right\| \\
&\leq \frac{1}{2\lambda t} \|\boldsymbol{\theta}^*\|^2.
\end{aligned}
\tag{30}
$$

Putting these two bounds together gives the proof for the main theorem.

$\qquad\square$

A.5 PROOF FOR THE VARIANCE UPPER BOUND

Recall that $\boldsymbol{X} = \boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^{\top}$ is the singular value decomposition of data matrix $\boldsymbol{X}$, where $\boldsymbol{U} = (\boldsymbol{u}_1, \cdots, \boldsymbol{u}_n)$, $\boldsymbol{W} = (\boldsymbol{w}_1, \cdots, \boldsymbol{w}_n)$, $\tilde{\boldsymbol{\Lambda}} = \text{diag}\{\mu_1, \cdots, \mu_n\}$ with $\mu_1 \geq \mu_2 \geq \cdots \mu_n$.

Recall that

$$
\begin{aligned}
k_0 &= \min\{l \geq 0 : \lambda_{l+1} \leq \frac{c_0 \sum_{i>l} \lambda_i}{n}\}, \\
k_1 &= \min\{l \geq 0 : \lambda_{l+1} \leq \frac{c_1 \sum_{i>0} \lambda_i}{n}\}, \\
k_2 &= \min\{l \geq 0 : \sum_{i>l} \lambda_i + n\lambda_{l+1} \leq c_2 c(t,n) \sum_{i>0} \lambda_i\}\},
\end{aligned}
\tag{31}
$$

for some constant c_0, c_1, c_2 and function $c(t,n)$.

We further define

$$
k_3 = \min\{l \geq 0 : \mu_{l+1} \leq c_3 c(t,n) \sum_{i>0} \lambda_i\},
\tag{32}
$$

for some constant c_3.

The next lemma shows that we can appropriately choose constants to ensure that $k_3 \leq k_2$ holds with high probability, and in some specific cases we have $k_2 \leq k_1$.

Lemma A.8. *For any function $c(t,n)$ and constant c_2, there exists constants c, c_3, such that $k_3 \leq k_2$ with probability at least $1 - e^{-\frac{n}{c}}$. Furthermore, if $c(t,n)$ is a positive constant function, for any c_1, there exists c_2 such that $k_2 \leq k_1$.*

Proof. According to lemma A.1, there exists a constant c, with probability at least $1 - e^{-\frac{n}{c}}$ we have

$$
\mu_{k_2+1} \leq c(\sum_{i>k_2} \lambda_i + n\lambda_{k_2+1}) \leq c c_2 c(t,n) \sum_{i>0} \lambda_i.
\tag{33}
$$

Therefore, we know that $k_3 \leq k_2$ for $c_3 = c c_2$.

By the definition of k_1, we have

$$
\sum_{i>k_1} \lambda_i + n\lambda_{k_1+1} \leq (c_1 + 1) \sum_{i>0} \lambda_i,
\tag{34}
$$

which implies that $k_2 \leq k_1$ for $c_2 = \frac{c_1+1}{c(t,n)}$, if $c(t,n)$ is a positive constant. □

Next we bound $\text{Tr}[\boldsymbol{C}]$, which implies an upper bound on the variance term.

Theorem A.1. *There exist constants c, c_0, c_1, c_2 such that if $k_0 \leq \frac{n}{c}$, then with probability at least $1 - e^{-\frac{n}{c}}$, the trace of the variance matrix C has the following upper bound*

$$
\text{Tr}[\boldsymbol{C}] \leq c\left(\frac{k_1}{n} + \frac{k_2}{c(t,n)n} + c(t,n)\left(\frac{\lambda t}{n}\sum_{i>0}\lambda_i\right)^2\right).
\tag{35}
$$

Proof. We divide the eigenvalues of $\boldsymbol{X}\boldsymbol{X}^{\top}$ into two groups based on whether they are greater than $c_3 c(t,n) \sum_{i>0} \lambda_i$. The first group consists of $\mu_1 \cdots \mu_{k_3}$, and the second group consists of $\mu_{k_3+1} \cdots \mu_n$. For $1 \leq j \leq k_3$, we have

$$
1 - \left(1 - \frac{\lambda}{n}\mu_j\right)^t \leq 1.
\tag{36}
$$

Therefore we have the following upper bound on $\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}\boldsymbol{X}^{\top}\right)^t\right]^2$:

$$\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}\boldsymbol{X}^\top \right)^t \right]^2$$

$$= \boldsymbol{U}\mathrm{diag}\left\{ \left[1 - \left(1 - \frac{\lambda}{n}\mu_1 \right)^t \right]^2 \cdots \left[1 - \left(1 - \frac{\lambda}{n}\mu_n \right)^t \right]^2 \right\} \boldsymbol{U}^\top$$

$$\preceq \boldsymbol{U}\mathrm{diag}\left\{ \overbrace{1, \cdots 1}^{k_3 \text{ times}}, \overbrace{\left[1 - \left(1 - \frac{\lambda}{n}\mu_{k_3+1} \right)^t \right]^2, \cdots \left[1 - \left(1 - \frac{\lambda}{n}\mu_n \right)^t \right]^2}^{n-k_3 \text{ times}} \right\} \boldsymbol{U}^\top \tag{37}$$

$$= \boldsymbol{U}\mathrm{diag}\left\{ \overbrace{1, \cdots 1}^{k_3 \text{ times}}, \overbrace{0, \cdots 0}^{n-k_3 \text{ times}} \right\} \boldsymbol{U}^\top$$

$$+ \boldsymbol{U}\mathrm{diag}\left\{ \overbrace{0, \cdots 0}^{k_3 \text{ times}}, \overbrace{\left[1 - \left(1 - \frac{\lambda}{n}\mu_{k_3+1} \right)^t \right]^2, \cdots \left[1 - \left(1 - \frac{\lambda}{n}\mu_n \right)^t \right]^2}^{n-k_3 \text{ times}} \right\} \boldsymbol{U}^\top.$$

For positive semidefinite matrices $\boldsymbol{P}, \boldsymbol{Q}, \boldsymbol{R}$ which satisfies $\boldsymbol{Q} \preceq \boldsymbol{R}$, it holds that $\mathrm{Tr}[\boldsymbol{P}\boldsymbol{Q}] \le \mathrm{Tr}[\boldsymbol{P}\boldsymbol{R}]$. It implies the following upperbound of $\mathrm{Tr}[\boldsymbol{C}]$:

$$\mathrm{Tr}[\boldsymbol{C}]$$

$$= \mathrm{Tr}\left[\left[\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n} \boldsymbol{X}\boldsymbol{X}^\top \right)^t \right]^2 \left(\boldsymbol{X}\boldsymbol{X}^\top \right)^{-2} \boldsymbol{X}\boldsymbol{\Sigma}\boldsymbol{X}^\top \right]$$

$$\le \underbrace{\mathrm{Tr}\left[\boldsymbol{U}\mathrm{diag}\left\{ \overbrace{1, \cdots 1}^{k_3 \text{ times}}, \overbrace{0, \cdots 0}^{n-k_3 \text{ times}} \right\} \boldsymbol{U}^\top \left(\boldsymbol{X}\boldsymbol{X}^\top \right)^{-2} \boldsymbol{X}\boldsymbol{\Sigma}\boldsymbol{X}^\top \right]}_{①}$$

$$+ \underbrace{\mathrm{Tr}\left[\boldsymbol{U}\mathrm{diag}\left\{ \overbrace{0, \cdots 0}^{k_3 \text{ times}}, \overbrace{\left[1 - \left(1 - \frac{\lambda}{n}\mu_{k_3+1} \right)^t \right]^2, \cdots \left[1 - \left(1 - \frac{\lambda}{n}\mu_n \right)^t \right]^2}^{n-k_3 \text{ times}} \right\} \boldsymbol{U}^\top \left(\boldsymbol{X}\boldsymbol{X}^\top \right)^{-2} \boldsymbol{X}\boldsymbol{\Sigma}\boldsymbol{X}^\top \right]}_{②}. \tag{38}$$

Bounding ①

Noticing $\boldsymbol{X} = \boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^\top$ and $\boldsymbol{\Sigma} = \sum_{i\geq 1}\lambda_i \boldsymbol{v}_i \boldsymbol{v}_i^\top$, we express the first term as sums of eigenvector products,

$$
\begin{aligned}
① &= \mathrm{Tr}\left[\boldsymbol{U}\mathrm{diag}\left\{\overbrace{1,\cdots 1}^{k_3 \text{ times}},\ \overbrace{0,\cdots 0}^{n-k_3 \text{ times}}\right\}\boldsymbol{U}^\top\left(\boldsymbol{X}\boldsymbol{X}^\top\right)^{-2}\boldsymbol{X}\boldsymbol{\Sigma}\boldsymbol{X}^\top\right] \\
&= \mathrm{Tr}\left[\boldsymbol{U}\mathrm{diag}\left\{\overbrace{1,\cdots 1}^{k_3 \text{ times}},\ \overbrace{0,\cdots 0}^{n-k_3 \text{ times}}\right\}\boldsymbol{U}^\top\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{-2}\boldsymbol{U}^\top\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^\top\boldsymbol{\Sigma}\boldsymbol{W}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{U}^\top\right] \\
&= \mathrm{Tr}\left[\mathrm{diag}\left\{\overbrace{1,\cdots 1}^{k_3 \text{ times}},\ \overbrace{0,\cdots 0}^{n-k_3 \text{ times}}\right\}\tilde{\boldsymbol{\Lambda}}^{-1}\boldsymbol{W}^\top\boldsymbol{\Sigma}\boldsymbol{W}\right] \\
&= \sum_{i\geq 1}\lambda_i\,\mathrm{Tr}\left[\mathrm{diag}\left\{\overbrace{1,\cdots 1}^{k_3 \text{ times}},\ \overbrace{0,\cdots 0}^{n-k_3 \text{ times}}\right\}\tilde{\boldsymbol{\Lambda}}^{-1}\boldsymbol{W}^\top\boldsymbol{v}_i\boldsymbol{v}_i^\top\boldsymbol{W}\right] \\
&= \sum_{i\geq 1}\sum_{1\leq j\leq k_3}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 .
\end{aligned}
\tag{39}
$$

Next we divide the above summation into $1 \leq i \leq k_1$ and $i > k_1$. For the first part, notice that

$$
\begin{aligned}
\sum_{1\leq j\leq k_3}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 &\leq \sum_{1\leq j\leq n}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 \\
&= \lambda_i \boldsymbol{v}_i^\top\left(\sum_{1\leq j\leq n}\frac{1}{\mu_j}\boldsymbol{w}_j\boldsymbol{w}_j^\top\right)\boldsymbol{v}_i \\
&= \lambda_i \boldsymbol{v}_i^\top\boldsymbol{W}\tilde{\boldsymbol{\Lambda}}^{-1}\boldsymbol{W}^\top\boldsymbol{v}_i \\
&= \lambda_i \boldsymbol{v}_i^\top\boldsymbol{W}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{U}^\top\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{-2}\boldsymbol{U}^\top\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^\top\boldsymbol{v}_i \\
&= \lambda_i^2\tilde{\boldsymbol{x}}_i^\top\left(\boldsymbol{X}\boldsymbol{X}^\top\right)^{-2}\tilde{\boldsymbol{x}}_i ,
\end{aligned}
\tag{40}
$$

where $\tilde{\boldsymbol{x}}_i$ is defined as $\tilde{\boldsymbol{x}}_i = \frac{\boldsymbol{X}\boldsymbol{v}_i}{\sqrt{\lambda_i}} = \frac{\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^\top\boldsymbol{v}_i}{\sqrt{\lambda_i}}$.

From the proof of lemma 11 in Bartlett et al. (2019), we know that for any σ_x, there exists a constant c_0 and c such that if $k_0 \leq \frac{n}{c}$, with probability at least $1 - e^{-\frac{n}{c}}$ the first part can be bounded as

$$
\sum_{1\leq i\leq k_1}\sum_{1\leq j\leq k_3}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 \leq \sum_{1\leq i\leq k_1}\lambda_i^2\tilde{\boldsymbol{x}}_i(\boldsymbol{X}\boldsymbol{X}^\top)^{-2}\tilde{\boldsymbol{x}}_i \leq c\frac{k_1}{n},
\tag{41}
$$

which gives a bound for the first part.

For the second part we interchange the order of summation and have

$$
\begin{aligned}
\sum_{i\geq k_1}\sum_{1\leq j\leq k_3}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 &= \sum_{1\leq j\leq k_3}\sum_{i\geq k_1}\frac{\lambda_i}{\mu_j}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 \\
&\leq \frac{1}{c_3 c(t,n)\sum_{i>0}\lambda_i}\sum_{1\leq j\leq k_3}\sum_{i\geq k_1}\lambda_i\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 \\
&= \frac{\lambda_{k_1+1}}{c_3 c(t,n)\sum_{i>0}\lambda_i}\sum_{1\leq j\leq k_3}\sum_{i\geq k_1}\left(\boldsymbol{v}_i^\top\boldsymbol{w}_j\right)^2 \\
&\leq \frac{\lambda_{k_1+1}}{c_3 c(t,n)\sum_{i>0}\lambda_i}\sum_{1\leq j\leq k_3}1 \\
&= \frac{\lambda_{k_1+1}k_3}{c_3 c(t,n)\sum_{i>0}\lambda_i} \\
&\leq c\frac{k_3}{c(t,n)n} .
\end{aligned}
\tag{42}
$$

for c large enough.

Putting 41 and 42 together, and noting that $k_3 \leq k_2$ with high probability as given in lemma A.8, we know there exists a constant c that with probability at least $1 - e^{-\frac{n}{c}}$,

$$① \leq c\frac{k_1}{n} + c\frac{k_2}{c(t,n)n}. \tag{43}$$

Bounding ②

Similar to the first step in bounding ①, we note that

$$
② = \mathrm{Tr}\left[\boldsymbol{U}\mathrm{diag}\left\{ \overbrace{0,\cdots 0,}^{k_3 \text{ times}} \overbrace{\left[1 - \left(1 - \frac{\lambda}{n}\mu_{k_3+1}\right)^t\right]^2, \cdots, \left[1 - \left(1 - \frac{\lambda}{n}\mu_n\right)^t\right]^2}^{n-k_3 \text{ times}} \right\} \right.
$$
$$
\left. \boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{-2}\boldsymbol{U}^\top\boldsymbol{U}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{W}^\top\boldsymbol{\Sigma}\boldsymbol{W}\tilde{\boldsymbol{\Lambda}}^{\frac{1}{2}}\boldsymbol{U}^\top \right]
$$
$$
= \mathrm{Tr}\left[\mathrm{diag}\left\{ \overbrace{0,\cdots 0,}^{k_3 \text{ times}} \overbrace{\frac{1}{\mu_{k_3+1}}\left[1 - \left(1 - \frac{\lambda}{n}\mu_{k_3+1}\right)^t\right]^2, \cdots, \frac{1}{\mu_n}\left[1 - \left(1 - \frac{\lambda}{n}\mu_n\right)^t\right]^2}^{n-k_3 \text{ times}} \right\} \right.
$$
$$
\left. \boldsymbol{W}^\top\boldsymbol{\Sigma}\boldsymbol{W} \right]. \tag{44}
$$

From Bernoulli's inequality and the definition of k_3, for any $k_3 + 1 \leq j \leq n$, we have

$$\frac{1}{\mu_k}\left[1 - \left(1 - \frac{\lambda}{n}\mu_k\right)^t\right]^2 \leq \frac{1}{\mu_k}\left(\frac{\lambda}{n}\mu_k t\right)^2 = \left(\frac{\lambda t}{n}\right)^2\mu_k \leq c_3\left(\frac{\lambda t}{n}\right)^2 c(t,n)\sum_{i>0}\lambda_i, \tag{45}$$

Hence,

$$
② \leq c_3 c(t,n)\left(\frac{\lambda t}{n}\right)^2\sum_{i>0}\lambda_i\,\mathrm{Tr}[\boldsymbol{W}^\top\boldsymbol{\Sigma}\boldsymbol{W}]
$$
$$
= c_3 c(t,n)\left(\frac{\lambda t}{n}\sum_{i>0}\lambda_i\right)^2. \tag{46}
$$

Putting things together

From the bounds for ① and ② given above, we know that there exists a constant c such that with probability at least $1 - e^{-\frac{n}{c}}$, the trace of the variance matrix C has the following upper bound

$$\mathrm{Tr}[C] \leq c\left(\frac{k_1}{n} + \frac{k_2}{c(t,n)n} + c(t,n)\left(\frac{\lambda t}{n}\sum_{i>0}\lambda_i\right)^2\right). \tag{47}$$

□

Proof of theorem 5.1. Lemma A.4, A.7 and Theorem A.1 gives the complete proof. Note that the high probability events in the proof are independent of the epoch number t, and this implies that the theorem holds uniformly for all $t \in \mathbb{N}$. □

A.6 PROOF OF COMPATIBILITY RESULTS

Corollary A.2 (Corollary 5.1 restated). *Let Assumption 1, 2 and 3 hold. Fix a constant $c(t, n)$. Suppose $k_0 = O(n)$, $k_1 = o(n)$, $r(\Sigma) = o(n)$, $\lambda = O\left(\frac{1}{\sum_{i>0} \lambda_i}\right)$. Then there exists a sequence of positive constants $\{\delta_n\}_{n \geq 0}$ which converge to 0, such that with probability at least $1 - \delta_n$, the excess risk is consistent for $t \in \left(\omega\left(\frac{1}{\lambda}\right), o\left(\frac{n}{\lambda}\right)\right)$, i.e.*

$$R(\boldsymbol{\theta}_t) = o(1).$$

Furthermore, for any positive constant δ, with probability at least $1 - \delta$, the minimal excess risk on the training trajectory can be bounded as

$$\min_t R(\boldsymbol{\theta}_t) \lesssim \frac{\max\{\sqrt{r(\Sigma)}, 1\}}{\sqrt{n}} + \frac{\max\{k_1, 1\}}{n}.$$

Proof. According to Lemma A.7, with probability at least $1 - \frac{\delta_n}{2}$, the following inequality holds for all t:

$$B(\boldsymbol{\theta}_t) \lesssim \left(\frac{1}{\lambda t} + \max\left\{\sqrt{\frac{r(\Sigma)}{n}}, \frac{r(\Sigma)}{n}, \sqrt{\frac{\log(\frac{1}{\delta_n})}{n}}, \frac{\log(\frac{1}{\delta_n})}{n}\right\}\right). \tag{48}$$

If δ_n is chosen such that $\log \frac{1}{\delta_n} = o(n)$, we have that with probability at least $1 - \frac{\delta_n}{2}$, we have for all $t = \omega\left(\frac{1}{\lambda}\right)$:

$$B(\boldsymbol{\theta}_t) = o(1), \tag{49}$$

in the sample size n.

When $c(t, n)$ is a constant, we have $k_2 \leq k_1$ with high probability as given in lemma A.8. Therefore, according to Lemma A.4 and Theorem A.1, we know that if $\log \frac{1}{\delta_n} = O(n)$, with probability at least $1 - \frac{\delta_n}{2}$, the following bound holds for all t:

$$V(\boldsymbol{\theta}_t) \lesssim \log\left(\frac{1}{\delta_n}\right)\left(\frac{k_1}{n} + \frac{\lambda^2 t^2}{n^2}\right). \tag{50}$$

Since $k_1 = o(n)$, $t = o\left(\frac{n}{\lambda}\right)$, we have $\frac{k_1}{n} + \frac{\lambda^2 t^2}{n^2} = o(1)$. Therefore, there exists a mildly decaying sequence of δ_n with $\log\left(\frac{1}{\delta_n}\right)\left(\frac{k_1}{n} + \frac{\lambda^2 t^2}{n^2}\right) = o(1)$, i.e.,

$$V(\boldsymbol{\theta}_t) = o(1). \tag{51}$$

To conclude, δ_n can be chosen such that

$$\log\left(\frac{1}{\delta_n}\right) = \omega(1), \log\left(\frac{1}{\delta_n}\right) = O(n), \log\left(\frac{1}{\delta_n}\right) = O\left(\frac{1}{\frac{k_1}{n} + \frac{\lambda^2 t^2}{n^2}}\right), \tag{52}$$

and then with probability at least $1 - \delta_n$, the excess risk is consistent for all $t \in \left(\omega\left(\frac{1}{\lambda}\right), o\left(\frac{n}{\lambda}\right)\right)$:

$$R(\boldsymbol{\theta}_t) = B(\boldsymbol{\theta}_t) + V(\boldsymbol{\theta}_t) = o(1). \tag{53}$$

This completes the proof for the first claim. The second claim follows from Equation 48 and 50 by setting $t = \Theta\left(\frac{\sqrt{n}}{\lambda}\right)$. □

Lemma A.9 (Lemma 5.1 restated). *For any fixed (i.e. independent of sample size n) feature covariance Σ satisfying assumption 1, we have $k_1(n) = o(n)$.*

Proof. Suppose there exists constant c, such that $k_1(n) \geq cn$. By definition of k_1, we know that $\lambda_l \geq \frac{c_1 \sum_{i>0} \lambda_i}{n}$ holds for $1 \leq l \leq k_1(n)$. Hence we have

$$\sum_{l=\lfloor cn2^i \rfloor + 1}^{\lfloor cn2^{i+1} \rfloor} \lambda_l \gtrsim \frac{c_1 \sum_{i>0} \lambda_i}{n2^{i+1}} cn2^i \gtrsim \sum_{i>0} \lambda_i. \tag{54}$$

summing up all l leads to a contradiction since $\sum_{i>0} \lambda_i < \infty$, which finishes the proof. □

Theorem A.2 (Theorem 4.1 restated). *Consider the overparameterized linear regression setting defined in section 4.1. Let Assumption 1,2 and 3 hold. Assume the learning rate satisfies $\lambda = O(\frac{1}{\text{Tr}(\boldsymbol{\Sigma})})$. Then under the condition that $k_0 = O(n), k_1 = o(n), r(\boldsymbol{\Sigma}) = o(n)$, gradient descent is compatible with overparameterized linear regression in the region $T_n = \left(\omega\left(\frac{1}{\lambda}\right), o\left(\frac{n}{\lambda}\right)\right)$, namely,*

$$\sup_{t \in T_n} R(\boldsymbol{\theta}_t) \xrightarrow{P} 0 \quad as \quad n \to \infty.$$

Furthermore, if the feature dimension $p = \infty$ and the data distribution does not change with n, then the condition $k_0 = O(n)$ alone suffices for compatibility.

Proof. According to Corollary 5.1 and Lemma 5.1, for any $\varepsilon > 0$, there exists $\{\delta_n\}_{n>0}$ and N such that for any sample size $n > N$, we have

$$\Pr\left[\left|\sup_{t \in T_n} R(\boldsymbol{\theta}_t)\right| > \varepsilon\right] \leq \delta_n. \tag{55}$$

Let $n \to \infty$ shows that $\sup_{t \in T_n} R(\boldsymbol{\theta}_t)$ converges to 0 in probability, which completes the proof. □

B EXAMPLES AND DISCUSSIONS

B.1 EXAMPLE DISTRIBUTIONS

We apply our bound in Corollary 5.1 to several examples. In each example, we show the data distribution, the time interval, and the corresponding generalization bound. These distributions are widely discussed in (Bartlett et al., 2019; Zou et al., 2021).

Example B.1. *Under the same conditions as Theorem 5.1, let $\boldsymbol{\Sigma}$ denote the feature covariance matrix. We show the following examples:*

1. **(Inverse Polynominal).** *If the spectrum of $\boldsymbol{\Sigma}$ satisfies*

$$\lambda_k = \frac{1}{k^\alpha},$$

for some $\alpha > 1$, we derive that $k_0 = \Theta(n)$, $k_1 = \Theta\left(n^{\frac{1}{\alpha}}\right)$. Therefore, $\min_t V(\boldsymbol{\theta}_t) = O\left(n^{\frac{1-\alpha}{\alpha}}\right)$ and

$$\min_t R(\boldsymbol{\theta}_t) = O\left(n^{-\min\left\{\frac{\alpha-1}{\alpha}, \frac{1}{2}\right\}}\right).$$

2. **(Inverse Log-Polynominal).** *If the spectrum of $\boldsymbol{\Sigma}$ satisfies*

$$\lambda_k = \frac{1}{k \log^\beta(k+1)},$$

for some $\beta > 1$, we derive that $k_0 = \Theta\left(\frac{n}{\log n}\right)$, $k_1 = \Theta\left(\frac{n}{\log^\beta n}\right)$. Therefore, $\min_t V(\boldsymbol{\theta}_t) = O\left(\frac{1}{\log^\beta n}\right)$ and

$$\min_t R(\boldsymbol{\theta}_t) = O\left(\frac{1}{\log^\beta n}\right).$$

3. **(Constant).** *If the spectrum of $\boldsymbol{\Sigma}$ satisfies*

$$\lambda_k = \frac{1}{n^{1+\varepsilon}}, 1 \leq k \leq n^{1+\varepsilon},$$

for some $\varepsilon > 0$, we derive that $k_0 = 0$, $k_1 = 0$. Therefore, $\min_t V(\boldsymbol{\theta}_t) = O\left(\frac{1}{n}\right)$ and

$$\min_t R(\boldsymbol{\theta}_t) = O\left(\frac{1}{\sqrt{n}}\right).$$

4. **(Piecewise Constant).** *If the spectrum of Σ satisfies*

$$\lambda_k = \begin{cases} \frac{1}{s} & 1 \le k \le s, \\ \frac{1}{d-s} & s+1 \le k \le d, \end{cases}$$

where $s = n^r, d = n^q, 0 < r \le 1, q \ge 1$. We derive that $k_0 = n^r$, $k_1 = n^r$. Therefore, $\min_t V(\boldsymbol{\theta}_t) = O(n^{r-1})$ and

$$\min_t R(\boldsymbol{\theta}_t) = O\left(n^{-\min\left\{1-r, \frac{1}{2}\right\}}\right).$$

B.2 COMPARISONS WITH BENIGN OVERFITTING RESULTS

We summarize the results in Bartlett et al. (2019); Zou et al. (2021) and our results in Table 1, and provide a detailed comparison with them below.

Comparison to Bartlett et al. (2019). In this seminal work, the authors study the excess risk of the min-norm interpolator. As discussed before, the min-norm interpolator is the convergence point of gradient descent in the overparameterized linear regression setting. One of the main results in Bartlett et al. (2019) is providing a tight bound for the variance part[4] in excess risk as

$$V(\hat{\boldsymbol{\theta}}) = O\left(\frac{k_0}{n} + \frac{n}{R_{k_0}(\boldsymbol{\Sigma})}\right), \tag{56}$$

where $\hat{\boldsymbol{\theta}} = \boldsymbol{X}^\top (\boldsymbol{X}\boldsymbol{X}^\top)^{-1}\boldsymbol{Y}$ denotes the min-norm interpolator, and $R_k(\boldsymbol{\Sigma}) = (\sum_{i>k} \lambda_i)^2 / (\sum_{i>k} \lambda_i^2)$ denote another type of effective rank.

By introducing the time factor, Theorem 5.1 improves over Equation (56) in at least two aspects. Firstly, Theorem 5.1 guarantees the consistency of the gradient descent dynamics for a broad range of step number t, while Bartlett et al. (2019) only study the limiting behavior of the dynamics of $t \to \infty$. Secondly, Theorem 5.1 implies that the excess risk of early stopping gradient descent solution can be much better than the min-norm interpolator. Compared to the bound in Equation (56), the bound in Corollary 5.1 (a.) replaces k_0 with a much smaller quantity k_1; and (b.) drops the second term involving $R_{k_0}(\boldsymbol{\Sigma})$. Therefore, we can derive a consistent bound for overparameterized linear regression for an early stopping solution, even though the excess risk of limiting point (min-norm interpolator) can be $\Omega(1)$ (See the first example in B.1). We can further derive a data-dependent time interval in which the bound is consistent, which cannot be directly obtained in Bartlett et al. (2019).

Comparison to Zou et al. (2021). Zou et al. (2021) study a different setting, which focuses on the one-pass stochastic gradient descent solution of linear regression. The authors prove a bound for the excess risk as

$$R(\bar{\boldsymbol{\theta}}_t) = O\left(\frac{k_1}{n} + \frac{n\sum_{i>k_1}\lambda_i^2}{(\sum_{i>0}\lambda_i)^2}\right), \tag{57}$$

where $\bar{\boldsymbol{\theta}}_t$ denotes the parameter obtained using stochastic gradient descent (SGD) with constant step size at epoch t. Similar to our bound, Equation 57 also uses the effective dimension k_1 to characterize the variance term. However, we emphasize that Zou et al. (2021) derive the bound in a pretty different scenario from ours, which is one-pass SGD scenario. During the one-pass SGD training, we use a fresh data point to perform stochastic gradient descent in each epoch, and therefore they set $t = \Theta(n)$ by default. As a comparison, we apply the standard full-batch gradient descent, and thus the time can be more flexible. Besides, our results in Corollary 5.1 improve the bound in Equation (57) by dropping the second term. We refer to the third and fourth example in Example B.1 for more details, where[5] our bound outperforms Zou et al. (2021) when $\varepsilon < 1/2$ or $q < \min\{2 - r, 3/2\}$.

---

[4]We do not compare the bias component in the excess risk bound here since our bound for the bias component follows Bartlett et al. (2019).

[5]Due to the bias term in Theorem 5.1, the overall excess risk bound cannot surpass the order $O(1/\sqrt{n})$, which leads to the cases that Zou et al. (2021) outperforms our bound. However, we note that such differences come from the intrinsic property of GD and SGD, which may be unable to avoid in the GD regimes.

B.3 COMPARISONS WITH STABILITY-BASED BOUNDS

In this section, we show that Theorem 5.1 gives provably better upper bounds than the stability-based method. We cite a result from Teng et al. (2021), which uses stability arguments to tackle overparameteried linear regression under similar assumptions.

Theorem B.1 (modified from Theorem 1 in Teng et al. (2021)). *Under the overparameterized linear regression settings, assume that $\|x\| \leq 1$, $|\varepsilon| \leq V$, $w = \frac{\theta^{*,\top} x}{\sqrt{\theta^{*,\top} \Sigma \theta^*}}$ is σ_w^2-subgaussian. Let $B_t = \sup_{\tau \in [t]} \|\theta_t\|$. the following inequality holds with probability at least $1 - \delta$:*

$$R(\theta_t) = \tilde{O}\left(\max\{1, \theta^{*,\top} \Sigma \theta^* \sigma_w^2, (V + B_t)^2\} \sqrt{\frac{\log(4/\delta)}{n}} + \frac{\|\theta^*\|^2}{\lambda t} + \frac{\lambda t (V + B_t)^2}{n}\right). \quad (58)$$

Theorem B.1 applies the general stability-based results (Hardt et al., 2016; Feldman and Vondrák, 2019) in the overparameterized linear regression setting, by replacing the bounded Lipschitz condition with the bounded domain condition. A fine-grained analysis (Lei and Ying, 2020) may remove the bounded Lipschitz condition, but it additionally requires zero noise or decaying learning rate, which is different from our setting. We omit the excess risk decomposition technique adopted in Teng et al. (2021) for presentation clarity.

Theorem B.1 can not directly yield the stability argument in Theorem 4.1, since obtaining a high probability bound of B_t requires a delicate trajectory analysis and is a non-trivial task. Therefore, data-irrelevant methods such as stability-based bounds can not be directly applied to our setting. Even if one can replace B_t in Equation 58 with its expectation that is easier to handle (this modification will require adding concentration-related terms, and make the bound in Equation 58 looser), we can still demonstrate that Theorem 5.1 is tighter than the corresponding stability-based analysis by providing a lower bound on $\mathbb{E}[B_t^2]$, which will imply a lower bound on the righthand side of Equation 58.

Theorem B.2. *Let Assumption 1, 2, 3 holds. Suppose $\lambda = O\left(\frac{1}{\sum_{i>0} \lambda_i}\right)$. Suppose the conditional variance of the noise $\varepsilon | x$ is lower bounded by σ_ε^2. There exists constant c, such that with probability at least $1 - ne^{-\frac{n}{c}}$, we have for $t = o(n)$,*

$$\mathbb{E}\|\theta_t\|^2 = \Omega\left(\frac{\lambda^2 t^2}{n}\left(\sum_{i > k_0} \lambda_i\right)\right) \quad (59)$$

First we prove the following lemma, bounding the number of large μ_i.

Lemma B.1. *Suppose $t = o(n)$. Let l denote the number of μ_i, such that $\mu_i = \Omega\left(\frac{n}{t}\right)$. Then with probability at least $1 - ne^{-\frac{n}{c}}$, we have $l = O(t)$.*

Proof. According to Lemma A.1, we know that with probability at least $1 - ne^{-\frac{n}{c}}$, Equation 5 holds for all $0 \leq k \leq n - 1$. Conditioned on this, we have

$$\frac{n}{t} l \lesssim \sum_{k=1}^{l} \mu_i \lesssim \sum_{k=1}^{l} (\sum_{i \geq k} \lambda_i + \lambda_k n) \lesssim (l + n) \sum_{i > 0} \lambda_i \lesssim l + n. \quad (60)$$

Since $t = o(n)$, we have $l = O(t)$ as claimed. $\square$

We also need the result from Bartlett et al. (2019), which gives a lowerbound of μ_n.

Lemma B.2. *(Lemma 10 in Bartlett et al. (2019)) For any σ_x, there exists a constant c, such that with probability at least $1 - e^{-\frac{n}{c}}$ we have,*

$$\mu_n \geq c\left(\sum_{i > k_0} \lambda_i\right). \quad (61)$$

We are now ready to prove Theorem B.2.

Proof. We begin with the calculation of $\|\boldsymbol{\theta}_t\|^2$. By Lemma A.2, the conditional unbiasedness of noise in Assumption 2 and the noise variance lower bound, we have

$$
\begin{aligned}
\mathbb{E}\|\boldsymbol{\theta}_t\|^2 &= \left\|\left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t(\boldsymbol{\theta}_0 - \boldsymbol{X}^\dagger\boldsymbol{Y}) + \boldsymbol{X}^\dagger Y\right\|^2 \\
&= \mathbb{E}\left\|\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger(\boldsymbol{X}\boldsymbol{\theta}^* + \varepsilon)\right\|^2 \\
&= \mathbb{E}\left\|\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger\boldsymbol{X}\boldsymbol{\theta}^*\right\|^2 + \mathbb{E}\left\|\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger\varepsilon\right\|^2 \\
&\geq \mathbb{E}\left\|\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger\varepsilon\right\|^2 \\
&= \mathbb{E}\operatorname{Tr}\left[\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger\varepsilon\varepsilon^\top\boldsymbol{X}^{\dagger,\top}\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\right] \\
&\geq \sigma_\varepsilon^2\mathbb{E}\operatorname{Tr}\left[\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\boldsymbol{X}^\dagger\boldsymbol{X}^{\dagger,\top}\left(\boldsymbol{I} - \left(\boldsymbol{I} - \frac{\lambda}{n}\boldsymbol{X}^\top\boldsymbol{X}\right)^t\right)\right] \\
&= \sigma_\varepsilon^2\sum_{i=1}^n\frac{[1 - (1 - \frac{\lambda}{n}\mu_i)^t]^2}{\mu_i}.
\end{aligned}
\tag{62}
$$

When $\mu_i = o\left(\frac{n}{t}\right)$, we have

$$
1 - (1 - \frac{\lambda}{n}\mu_i)^t = 1 - 1 + \frac{\lambda}{n}\mu_i t + O\left(\left(\frac{\lambda}{n}\mu_i t\right)^2\right) = \Theta\left(\frac{\lambda}{n}\mu_i t\right).
\tag{63}
$$

Plugging it into Equation 62 and then use Lemma B.1, B.2, we know that under the high probability event in Lemma B.1 and B.2,

$$
\mathbb{E}\|\boldsymbol{\theta}_t\|^2 = \Omega\left((n - l)\frac{\lambda^2}{n^2}\mu_n t^2\right) = \Omega\left(\frac{\lambda^2}{n}\mu_n t^2\right) = \Omega\left(\frac{\lambda^2 t^2}{n}\left(\sum_{i>k_0}\lambda_i\right)\right)
\tag{64}
$$

$\square$

Therefore, the stability-based bound, i.e., the right hand side of Equation 58, can be lower bounded in expectation as $\Omega\left(\frac{\lambda^3 t^3}{n^2}\sum_{i>k_0}\lambda_i\right)$. This implies that the stability-based bound is vacuous when $t = \Omega\left(\frac{n^{\frac{2}{3}}\left(\sum_{i>k_0}\lambda_i\right)^{-\frac{1}{3}}}{\lambda}\right)$. Thus, stability-based methods will provably yield smaller compatibility region than $\left(\omega\left(\frac{1}{\lambda}\right), o\left(\frac{n}{\lambda}\right)\right)$ in Theorem 4.1 when $\sum_{i>k_0}\lambda_i$ is not very small, as demonstrated in the examples below.

Example B.2. *Let Assumption 1, 2, 3 holds. Assume without loss of generality that $\lambda = \Theta(1)$. We have the following examples:*

1. **(Inverse Polynominal).** *If the spectrum of Σ satisfies*
$$
\lambda_k = \frac{1}{k^\alpha},
$$
for some $\alpha > 1$, we derive that $k_0 = \Theta(n)$, $\sum_{i>k_0}\lambda_i = \Theta(\frac{1}{n^{\alpha-1}})$. Therefore, the stability bound in Theorem B.1 is vacuous when
$$
t = \Omega\left(n^{\frac{\alpha+1}{3}}\right),
$$
which is outperformed by the compatibility region in Theorem 5.1 when $\alpha < 2$.

2. **(Inverse Log-Polynominal).** *If the spectrum of Σ satisfies*

$$\lambda_k = \frac{1}{k \log^\beta (k+1)},$$

for some $\beta > 1$, we derive that $k_0 = \Theta\left(\frac{n}{\log n}\right), \sum_{i>k_0} \lambda_i = \tilde{\Theta}(1)$. Therefore, the stability bound in Theorem B.1 is vacuous when

$$t = \tilde{\Omega}\left(n^{\frac{2}{3}}\right),$$

which is outperformed by the compatibility region in Theorem 5.1.

3. **(Constant).** *If the spectrum of Σ satisfies*

$$\lambda_k = \frac{1}{n^{1+\varepsilon}}, 1 \le k \le n^{1+\varepsilon},$$

for some $\varepsilon > 0$, we derive that $k_0 = 0$, $\sum_{i>k_0} \lambda_i = 1$. Therefore, the stability bound in Theorem B.1 is vacuous when

$$t = \Omega\left(n^{\frac{2}{3}}\right),$$

which is outperformed by the compatibility region in Theorem 5.1.

4. **(Piecewise Constant).** *If the spectrum of Σ satisfies*

$$\lambda_k = \begin{cases} \frac{1}{s} & 1 \le k \le s, \\ \frac{1}{d-s} & s+1 \le k \le d, \end{cases}$$

where $s = n^r, d = n^q, 0 < r \le 1, q \ge 1$, we derive that $k_0 = n^r$, $\sum_{i>k_0} \lambda_i = 1$. Therefore, the stability bound in Theorem B.1 is vacuous when

$$t = \Omega\left(n^{\frac{2}{3}}\right),$$

which is outperformed by the compatibility region in Theorem 5.1.

B.4 COMPARISONS WITH UNIFORM CONVERGENCE BOUNDS

We give a standard bound on the Rademacher complexity of linear models.

Theorem B.3 (Theorem in Mohri et al. (2012)). *Let $S \subseteq \{\boldsymbol{x} : \|x\|_2 \le r\}$ be a sample of size n and let $\mathcal{H} = \{x \mapsto \langle w, x \rangle : \|w\|_2 \le \Lambda\}$. Then, the empirical Rademacher complexity of $\mathcal{H}$ can be bounded as follows:*

$$\hat{\mathcal{R}}_S(\mathcal{H}) \le \sqrt{\frac{r^2 \Lambda^2}{n}}. \tag{65}$$

Furthermore, Talagrand's Lemma (See Lemma 5.7 in Mohri et al. (2012)) indicates that

$$\hat{\mathcal{R}}_S(l \circ \mathcal{H}) \le L\hat{\mathcal{R}}_S(\mathcal{H}) = \frac{\Theta(\Lambda^2)}{\sqrt{n}}, \tag{66}$$

where $L = \Theta(\Lambda)$ is the Lipschitz coefficient of the square loss function l in our setting. Therefore, the Rademacher generalization bound is vacuous when $\Lambda = \Omega(n^{\frac{1}{4}})$. By Theorem B.2, we know that $\mathbb{E}\|\boldsymbol{\theta}_t\|^2 = \Omega(n^{\frac{1}{2}})$ when $t = \Omega\left(\frac{n^{\frac{3}{4}}}{\lambda\left(\sum_{i>k_0} \lambda_i\right)^{\frac{1}{2}}}\right)$. A similar comparison as in Example B.2 can demonstrate that uniform stability arguments will provably yield smaller compatibility region than that in Theorem 5.1 for example distributions.

B.5 DISCUSSIONS ON EXTENSIONS TO KERNEL REGRESSION

In this section, we discuss extending the analysis in overparameterized linear regression to kernel regression setting.

Let $\mathcal{H}$ denote a infinite dimensional Hilbert space equipped with inner product $\langle \cdot, \cdot \rangle_{\mathcal{H}}$, and $\phi : \mathbb{R}^d \to \mathcal{H}$ denote a feature map. Consider the following class of functions:

$$\mathcal{F} = \{f : \mathbb{R}^d \to \mathbb{R} | f(\boldsymbol{x}) = \langle \boldsymbol{\theta}, \phi(\boldsymbol{x}) \rangle_{\mathcal{H}}\}. \tag{67}$$

Let $(\boldsymbol{x}, y) \in \mathbb{R}^d \times \mathbb{R}$ denote the data vector and the response, following a joint distribution $\mathcal{D}$. The goal of kernel regression is to find a function f parameterized by $\boldsymbol{\theta}$, that minimizes the following population loss

$$L(\boldsymbol{\theta}) = \frac{1}{2} \mathbb{E}\left(y - f(\boldsymbol{x})\right)^2 = \frac{1}{2} \mathbb{E}\left(y - \langle \boldsymbol{\theta}, \phi(\boldsymbol{x}) \rangle_{\mathcal{H}}\right)^2 \tag{68}$$

Therefore, kernel regression is equivalent to solving a linear regression problem on transformed data $(\phi(\boldsymbol{x}), y)$. By replacing $\boldsymbol{x}$ with $\phi(\boldsymbol{x})$, the notations and results in Section 4 can be naturally extended to the kernel regression case, which is detailed as follows.

Given a dataset $\{(\boldsymbol{x}_i, y_i)\}_{1 \le i \le n}$ sampled i.i.d from $\mathcal{D}$, consider the following dynamics of gradient descent analogous to Equation (2),

$$\boldsymbol{\theta}_{t+1} = \boldsymbol{\theta}_t - \frac{\lambda}{n} \phi(\boldsymbol{X})^\top (\phi(\boldsymbol{X}) \boldsymbol{\theta}_t - \boldsymbol{Y}), \tag{69}$$

where $\phi(\boldsymbol{X}) = (\phi(\boldsymbol{x}_1), \cdots, \phi(\boldsymbol{x}_n))^\top$ and $\boldsymbol{Y} = (y_1, \cdots, y_n)^\top$. Note that this dynamic takes place in the feature space $\mathcal{H}$. The following corollary characterizes the compatibility between kernel regression and gradient descent.

Corollary B.1. *Assume the distribution of $(\phi(\boldsymbol{x}), y)$ satisfies Assumptions 1, 2, 3, and does not change with sample size n. Let $\boldsymbol{\Sigma}_\phi = \mathbb{E}\left[\phi(\boldsymbol{x})\phi(\boldsymbol{x})^\top\right]$ denote the feature covariance. Then under the condition that the effective dimension $k_0(\boldsymbol{\Sigma}_\phi) = O(n)$ and the learning rate $\lambda = O(\frac{1}{\text{Tr}(\boldsymbol{\Sigma}_\phi)})$, gradient descent using 69 is compatible with kernel regression.*

Proof. The corollary follows from Theorem 4.1, by noting that its proof holds for feature vectors in an infinite dimensional Hilbert space. $\square$

We give several remarks regarding the difference between the generalization analysis of kernel regression and overparameterized linear regression.

Firstly, Corollary B.1 assumes $\phi(\boldsymbol{x})$ has subgaussian i.i.d entries after normalization (see Assumption 1.3), which can be hard to satisfy due to the non-linearity of feature mapping $\phi(\cdot)$.

Secondly, Corollary B.1 focuses on gradient descent in the feature space $\mathcal{H}$. Another perspective is to apply the Representer theorem (Mohri et al., 2012) to express the weight $\boldsymbol{\theta}$ into a linear combination of transformed inputs as $\boldsymbol{\theta} = \sum_{1 \le j \le n} \alpha_j \phi(\boldsymbol{x}_j)$, and analyze the gradient descent dynamics of the n-dimensional vector $\boldsymbol{\alpha} = (\alpha_1, \cdots, \alpha_n)$. Although they both converge to the min-norm solution, these two gradient descent dynamics are different, and Theorem 4.1 is not directly applicable to gradient descent on $\boldsymbol{\alpha}$.

Thirdly, previous works (El Karoui, 2010; Liang and Rakhlin, 2018; Mei and Montanari, 2022) prove that the spectrum of feature covariance $\boldsymbol{\Sigma}_\phi$ has an approximately linear relationship with the spectrum of data covariance $\boldsymbol{\Sigma} = \mathbb{E}\left[\boldsymbol{x}\boldsymbol{x}^\top\right]$ when feature mapping $\phi(\cdot)$ corresponds to an inner product type kernel. This can be used to derive a more intrinsic compatibility condition using only data distribution rather than feature distribution. We leave a more refined analysis for data-algorithm compatibility in the kernel regression regime for future works.

B.6 CALCULATIONS IN SECTION B.1

We calculate the quantities $r(\Sigma), k_0, k_1, k_2$ for the example distributions in B.1. The results validate that k_1 is typically a much smaller quantity than k_0, and k_1 serves as a proxy for k_2 in the constant $c(t, n)$ setting.

1. **Calculations for $\lambda_k = \frac{1}{k^\alpha}, \alpha > 1$.**

   Define $r_k(\boldsymbol{\Sigma}) = \frac{\sum_{i>k} \lambda_i}{\lambda_{k+1}}$ as in Bartlett et al. (2019). Since $\sum_{i>k} \frac{1}{i^\alpha} = \Theta(\frac{1}{k^{\alpha-1}})$, we have $r_k(\boldsymbol{\Sigma}) = \Theta\left(\frac{\frac{1}{k^{\alpha-1}}}{\frac{1}{k^\alpha}}\right) = \Theta(k)$. Hence, $k_0 = \Theta(n)$ [6], and the conditions of theorem 5.1 is satisfied.

   As $\sum_{i>0} \lambda_i < \infty$, By its definition we know that k_1 is the smallest l such that $\lambda_{l+1} = O(\frac{1}{n})$. Therefore, $k_1 = \Theta(n^{\frac{1}{\alpha}})$. Similarly, $k_2 = \Theta(n^{\frac{1}{\alpha}})$.

2. **Calculations for $\lambda_k = \frac{1}{k \log^\beta(k+1)}, \beta > 1$.**

   $\sum_{i>k} \frac{1}{i \log^\beta(i+1)} = \Theta(\int_k^\infty \frac{1}{x \log^\beta x} dx) = \Theta(\frac{1}{\log^{\beta-1} k})$, which implies $r_k(\boldsymbol{\Sigma}) = k \log k$. Solving $k_0 \log k_0 \geq \Theta(n)$, we have $k_0 = \Theta(\frac{n}{\log n})$.

   By the definition of k_1, we know that k_1 is the smallest l such that $l \log^\beta(l+1) \geq \Theta(n)$. Therefore, $k_1 = \Theta(\frac{n}{\log^\beta n})$. $k_2 = \Theta(\frac{n}{\log^\beta n})$ by similar calculations.

3. **Calculations for $\lambda_i = \frac{1}{n^{1+\varepsilon}}, 1 \leq i \leq n^{1+\varepsilon}, \varepsilon > 0$.**

   Since $r_0(\boldsymbol{\Sigma}) = n^{1+\varepsilon}$, we have $k_0 = 0$. By the definition of k_1 and k_2, we also have $k_1 = k_2 = 0$.

4. **Calculations for $\lambda = \begin{cases} \frac{1}{s} & 1 \leq k \leq s \\ \frac{1}{d-s} & s+1 \leq k \leq d \end{cases}, s = n^r, d = n^q, 0 < r \leq 1, q \geq 1$.**

   For $0 \leq k < s$, $r_k(\boldsymbol{\Sigma}) = \Theta(\frac{1}{\frac{1}{s}}) = \Theta(n^r) = o(n)$, while $r_s(\boldsymbol{\Sigma}) = \frac{1}{\frac{1}{d-s}} = \Theta(n^q) = \omega(n)$. Therefore, $k_0 = s = n^r$.

   Similarly, noting that $\lambda_k = \frac{1}{n^r} = \omega(n)$ for $0 \leq k < s$ and $\lambda_s = \Theta(\frac{1}{n^d}) = o(n)$, we know that $k_1 = s = n^r$. Similarly, $k_2 = s = n^r$.

B.7 Calculations for $\lambda_k = \frac{1}{k^\alpha}, \alpha > 1$ in Section 5.3

Set $c(t, n) = \frac{1}{n^\beta}$, where $\beta > 0$ will be chosen later. First we calculate k_2 under this choice of $c(t, n)$. Note that $\sum_{i>k} \frac{1}{i^\alpha} = \Theta\left(\frac{1}{k^{\alpha-1}}\right)$. Therefore, k_2 is the smallest k such that $\frac{1}{k^{\alpha-1}} + \frac{n}{k^\alpha} = O(\frac{1}{n^\beta})$. For the bound on $V(\theta_t)$ to be consistent, we need $k_2 = o(n)$. Hence, $\frac{1}{k^{\alpha-1}} = O(\frac{n}{k^\alpha})$, which implies $k_2 = n^{\frac{\beta+1}{\alpha}}$.

Plugging the value of $c(t, n)$ and k_2 into our bound, we have

$$V(\theta_t) = O\left(n^{(\frac{1}{\alpha}-1)+(\frac{1}{\alpha}+1)\beta} + n^{2\tau-\beta-2}\right)$$

which attains its minimum $\Theta(n^{\frac{2\alpha\tau-3\alpha+2\tau-1}{2\alpha+1}})$ at $\beta = \Theta\left(\frac{2\alpha\tau-\alpha-1}{2\alpha+1}\right)$.

For $V(\theta) = O(1)$, we need $\tau \leq \frac{3\alpha+1}{2\alpha+2}$. For $\beta \geq 0$, we need $\tau \geq \frac{\alpha+1}{2\alpha}$. Putting them together gives the range of t in which the above calculation applies.

C Additional Experiment Results

C.1 Details and Discussions for Linear Regression Experiments

In this section, we provide the experiment details for linear regression experiments in Figure 1 and present additional empirical results.

The linear regression experiment in 1 follows the setting described in section 6. We run gradient descent for 10^7 epochs. Although the final iterate does not interpolate the training data, the results suffice to demonstrate the gap between the early-stopping and final-iterate excess risk. The training plot for different covariances are given in Figure 2.

---

[6]The calculations for k_0, k_1 and k_2 in this section only apply when n is sufficiently large.

Figure 2: **The training plot for overparameterized linear regression with different covariances using GD.** Figure (b) is copied from Figure 1.

Next, we provide the experiment results of sample size $n = 50$ and $n = 200$ analogous to those in Section 6. The settings are the same as described in the main text, except for the sample size. The optimal excess risk and min-norm excess risk are provided in Table 3 and Table 4. The tables indicate that the three observations given in section 6 hold for different sample size n.

Table 3: **The effective dimension k_1, the optimal early stopping excess risk and min-norm excess risk for different feature distributions, with sample size $n = 50$.** The table shows that early stopping solution generalizes significantly better than min-norm interpolator, and reveals a positive correlation between the effective dimension k_1 and generalization performance of early stopping solution. We calculate the 95% confidence interval for the excess risk.

DISTRIBUTIONS	k_1	OPTIMAL EXCESS RISK	MIN-NORM EXCESS RISK
$\lambda_i = \frac{1}{i}$	$\Theta(n)$	2.515 ± 0.0104	12.632 ± 0.1602
$\lambda_i = \frac{1}{i^2}$	$\Theta(n^{\frac{1}{2}})$	0.269 ± 0.0053	50.494 ± 0.9378
$\lambda_i = \frac{1}{i^3}$	$\Theta(n^{\frac{1}{3}})$	0.083 ± 0.0011	13.208 ± 0.4556
$\lambda_i = \frac{1}{i \log(i+1)}$	$\Theta(\frac{n}{\log n})$	0.808 ± 0.0090	46.706 ± 0.6983
$\lambda_i = \frac{1}{i \log^2(i+1)}$	$\Theta(\frac{n}{\log^2 n})$	0.381 ± 0.0076	74.423 ± 1.1472
$\lambda_i = \frac{1}{i \log^3(i+1)}$	$\Theta(\frac{n}{\log^3 n})$	0.233 ± 0.0052	43.045 ± 0.8347

C.2 DETAILS FOR MNIST EXPERIMENTS

In this section, we provide the experiment details and additional results in MNIST dataset.

The MNIST experiment details are described below. We create a noisy version of MNIST with label noise rate 20%, i.e. randomly perturbing the label with probability 20% for each training data, to simulate the label noise which is common in real datasets, e.g ImageNet (Stock and Cissé, 2018; Shankar et al., 2020; Yun et al., 2021). We do not inject noise into the test data. We provide the plot for different levels of label noise in Figure 3. We also present the corresponding test error of the best early stopping iterate and the final iterate in Table 5.

We choose a standard four layer convolutional neural network as the classifier. We use a vanilla SGD optimizer without momentum or weight decay. The initial learning rate is set to 0.5. Learning rate is decayed by 0.98 every epoch. Each model is trained for 300 epochs. The training batch size

Table 4: **The effective dimension k_1, the optimal early stopping excess risk and min-norm excess risk for different feature distributions, with sample size $n = 200$.** The table shows that early stopping solution generalizes significantly better than min-norm interpolator, and reveals a positive correlation between the effective dimension k_1 and generalization performance of early stopping solution. We calculate the 95% confidence interval for the excess risk.

DISTRIBUTIONS	k_1	OPTIMAL EXCESS RISK	MIN-NORM EXCESS RISK
$\lambda_i = \frac{1}{i}$	$\Theta(n)$	2.173 ± 0.0065	52.364 ± 0.4009
$\lambda_i = \frac{1}{i^2}$	$\Theta(n^{\frac{1}{2}})$	0.161 ± 0.0039	36.855 ± 0.4833
$\lambda_i = \frac{1}{i^3}$	$\Theta(n^{\frac{1}{3}})$	0.068 ± 0.0012	8.1990 ± 0.2313
$\lambda_i = \frac{1}{i \log(i+1)}$	$\Theta(\frac{n}{\log n})$	0.628 ± 0.0034	152.70 ± 1.1073
$\lambda_i = \frac{1}{i \log^2(i+1)}$	$\Theta(\frac{n}{\log^2 n})$	0.241 ± 0.0036	83.550 ± 0.7596
$\lambda_i = \frac{1}{i \log^3(i+1)}$	$\Theta(\frac{n}{\log^3 n})$	0.146 ± 0.0108	33.469 ± 0.4540

(a) 0% label noise   (b) 10% label noise   (c) 20% label noise

(d) 30% label noise   (e) 40% label noise   (f) 50% label noise

Figure 3: **The training plot for corrupted MNIST with different levels of label noise using SGD.** Figure (c) is copied from Figure 1.

is set to 1024, and the test batch size is set to 1000. We choose the standard cross entropy loss as the loss function.

Table 5: **The test error of optimal stopping iterate and final iterate on MNIST dataset with different levels of label noise.** The results demonstrate that stopping iterate can have significantly better generalization performance than interpolating solutions for real datasets.

NOISE LEVEL	OPTIMAL TEST ERROR	FINAL TEST ERROR
0%	1.07%	1.13%
10%	1.75%	10.16%
20%	2.88%	19.90%
30%	2.18%	27.94%
40%	2.57%	35.15%
50%	2.71%	42.95%