Investigating the Effects of Plant Diversity on Soil Thermal Diffusivity using Physics-Informed Neural Networks

Gideon Stein[12] & Sai Karthikeya Vemuri[1],Yuanyuan Huang[2], Anne Ebeling[3],
Nico Eisenhauer[2], Maha Shadaydeh[1], Joachim Denzler[12]
[1] Computer Vision Group, Friedrich Schiller University Jena, Germany
[2] German Centre for Integrative Biodiversity Research (iDiv), Halle-Jena-Leipzig, Germany
[3] Institute of Ecology and Evolution, Friedrich Schiller University Jena, Germany
`{gideon.stein,sai.karthikeya.vemuri }@cuni-jena.de`

Abstract

The influence of plant diversity on the stability of ecosystems is well-reported in the literature. However, the exact mechanisms responsible for this effect are still a topic of debate. Recently, soil temperature stability was proposed as one possible candidate for such a mechanism. To further evaluate this hypothesis, we investigate the relationship between plant diversity and the thermal diffusivity of the soil during the very dry and hot summer of 2018 in Central Europe. By leveraging Physics-Informed Neural Networks and a 30-minute resolution soil temperature dataset from the Jena Experiment, we find an inverse relationship between plant diversity and the thermal diffusivity of the associated soil. With this, we provide support for the idea of plant diversity as a natural protection against climate-related ecosystem change.

1 Introduction

The intricate effects of plant diversity on the stability of ecosystems are well reported in the literature (Isbell et al., 2015), (Loreau et al., 2021), (Tilman et al., 2006). Determining the exact mechanisms that are responsible for these effects is, however, still in progress (De Boeck et al., 2018), (Gross et al., 2014), (Loreau & de Mazancourt, 2013). Recently, (Huang et al., 2024) showed that plant diversity substantially affects soil temperature stability, which in turn influences various mechanisms in the soil. Soil temperature drives a multitude of processes in the ecosystem (Wildung et al., 1975), (Onwuka, 2018), (Guy, 1999), (Nievola et al., 2017). Therefore, stabilizing the abiotic environment of these processes can lead to a generally more stable system. Besides increased biomass, which provides a natural shield against radiation, it was hypothesized that Soil Organic Carbon (SOC), which was reported to be increased through plant diversity (Chen et al., 2020), (Chen et al., 2018), (Spohn et al., 2023) also plays a key role concerning soil temperature stability. SOC is known to influence the thermal diffusivity of the soil (Zhu et al., 2019), (Chen et al., 2012) and, through this, increases soil temperature stability. While this chain is physically well supported, no data-driven studies have investigated this relationship's magnitude.

To close this gap, we perform a data-driven analysis of a multi-depth, long-term, 30-minute resolution soil temperature dataset collected at Europe's largest plant diversity experiment site, the Jena Experiment (Weisser et al., 2017). With this data source, we aim to uncover the relationship between plant diversity and the thermal diffusivity of the corresponding soil. For this, contrary to previous approaches that estimate soil thermal conductivity such as (Lukiashchenko & Arkhangelskaya, 2018) or (Arkhangelskaya & Lukyashchenko, 2018), we leverage Physics-Informed Neural Networks (Raissi et al., 2019) to estimate the soil thermal diffusivity constants for selected summer intervals of 2018, a specifically hot and dry year in Central Europe. Physics-Informed Neural Networks (PINNs) provide a versatile framework to perform data-driven Deep Learning while also incorporating physical knowledge. In this study, we use PINNs in an inverse problem setting to infer the thermal diffusivity of the soil from measurements of temperature at different depths, combined

with the partial-differential equation for heat conduction. To our knowledge, this is the first study that directly investigates the relationship between plant diversity and heat diffusivity.

Our results suggest an inverse relationship between plant diversity and the thermal diffusivity of the associated soil during the selected intervals. More specifically, we find that the thermal diffusivity of soil is reduced with increasing plant diversity, effectively suggesting a more stable soil system. With this, we provide data-driven evidence for the stabilizing effects of plant diversity on soil temperature as suggested in (Huang et al., 2024) and support the idea of plant diversity as an effective shield against climate-related ecosystem change.

2 DATA

The Jena Experiment (Weisser et al., 2017), situated near Jena, Germany, is a long-term grassland study field site that has operated since 2002. The main site consists of 88 unique plots with different plant diversity levels from 0 (bare ground) to 60 species. The plant diversity level is maintained through frequent weeding year-around. Further, since the field site holds a gradient of soil conditions depending on the distance to the Saale River (east side of the field), the field site is subdivided into four bigger blocks that group the plots according to these conditions. In all plots, soil temperature measurements are taken every minute in depths 5 and 15 cm. Further, in Block 2, additional measurements for the surface temperature and 60 cm depth are taken. We, therefore, focus on data from this block as marked in Figure 1 since every additional measurement can improve the estimation of heat diffusivity. Additionally to these measurements, a weather station is installed in the center of the field site from which we extract the general climatic conditions.

For the scope of this paper, we focus on 2018 (at this time, the hottest year in Central Europe since 1881) and aggregate the data to a 30-minute resolution for each plot individually. To compare the impact of air temperature and general soil moisture availability, we select four three-day intervals (Figure 2) from the summer of 2018 (June to August). Each interval maximizes or minimizes air temperature at 2 m height or soil moisture at 16cm depth (measured under the climate station), which we call *hot*, *cold*, *wet*, and *dry*. Further, we exclude the measurements in 60 cm depth since they show very little variance for the short intervals we selected. While the selected intervals in which we estimate heat diffusivity are sparse so far, we believe they provide a first good insight into the relationship under investigation.

Figure 1: The Jena Experiment field site (left) and the measurements taken in each plot of block 2 (right). Plots with a diversity level of -1 specify uncontrolled plots not considered in this study (together with diversity levels 60 and 0 since these provide only a single sample plot in block 2). In total, we consider 19 plots covering the full diversity gradient.

3 METHOD

Heat conduction in solids can be described by a partial differential equation called *heat conduction equation*, which is given as:

$$\frac{\partial T}{\partial t} = K \frac{\partial^2 T}{\partial z^2}, \qquad (1)$$

where T is the temperature, z is the spatial coordinate (only depth, in our case since we have a single measurement position per plot), and t is the time index. K is called the *thermal diffusivity* of the soil. This equation is valid for soils, given that the soil is homogeneous and isotropic which we assume here to be approximately true (since we select short intervals that we individually model). While these are arguably strong assumptions, they allow for a first and, more importantly, general picture of the relationship between plant diversity and heat diffusivity since *K* characterizes the overall thermal properties of soil. As it is a measure of the rate of heat transfer inside the soil, the higher the thermal diffusivity, the faster heat flows through the soil (Carlslaw & Jaeger, 1959).

Physics-Informed Neural Networks(PINNs) (Raissi et al., 2019) offers a paradigm to combine physical knowledge, in the form of differential equations, with Deep Learning. It involves constructing a multi-objective loss that fuses physics-related and data-related constraints. PINNs can be used to solve both forward and inverse problems involving differential equations. Applications and effectiveness of PINNs for inverse problems are discussed in the literature (Cuomo et al., 2022; Lu et al., 2021; Karniadakis et al., 2021). This study uses PINNs to estimate the soil thermal diffusivity K of a specific plot and a specific interval from multi-depth soil temperature measurements. In other words, we solve the inverse problem involving Equation 1 and the soil temperature data measurements as depicted in Figure 1. Importantly, next to obtaining an estimate for K, PINNs also provide an opportunity to interpolate smoothly between depths and time, which is highly beneficial for practitioners as they can match soil temperature measurements with other measurements at the exact same depth and time.

To this end, we construct a feed-forward neural network whose input is the time index t and the depth z. Its output is the amplitude and the phase shift for a diurnal sinusoidal cycle of 12 hours that is evaluated at time t to predict the corresponding soil temperature T. Consider the neural network T(θ,z,t), where θ denotes the model parameters that must be trained. The corresponding objective of the training is given as:

$$\min_{\theta, K} \underbrace{\left\| \frac{\partial T(\theta, z, t)}{\partial t} - K \frac{\partial^2 T(\theta, z, t)}{\partial z^2} \right\|}_{L_{physics}} + \underbrace{\| T(\theta, z, t) - T_{true}(z, t) \|}_{L_{data}}, \qquad (2)$$

where L_{phyics} is the loss term that incorporates physics knowledge (Equation 1) while L_{data} guarantees a proper prediction of data samples. The PINN is trained to optimize the values of K and θ to find the best representation, satisfying the heat conduction equations and minimizing the error on the data samples. The domain to which the equation and data are restricted is $z \in [0, 15cm]$ and $t \in [0, 3days]$.

We train a PINN for different plots and time intervals, as described in the previous section, to infer the corresponding thermal diffusivities. Concerning Hyperparameters, we use AdamW ((Loshchilov & Hutter, 2018)) and train for 200k Epochs, starting with a learning rate of $1e^{-4}$ that is halved after every 50k epochs for all our experiments. We perform full gradient descent (no batching) during all experiments. Further, we use sin as our activation function of choice. Finally, we deploy networks with ten hidden layers and a hidden dimension of 256. We use 432 collocation points (three times as much as the available data points). Finally, since we find that the PINNs do not always converge to a meaningful solution or sometimes cannot produce a reasonably low loss, we, in total, run two independent samples for each data interval and mean the resulting K between all valid runs. We filter runs that result in an MSE loss higher than 1.0 (loss distribution that we use to determine this threshold is given in **??**) and runs that produce $K's$ that are negative or very close to zero (mathematically possible but physically implausible). We believe that further regularization terms might help restrict the solution space for K in the future to improve consistency even further.

4 RESULTS

Figure 2: Top-Left: data samples selected to train PINNs Equation 2 independently. Top-Right: The resulting relationship between plant diversity and thermal diffusivity. The thermal diffusivity constant K is generally reduced with diversity. Bottom-Left: The temporal variance of K for individual plots. Bottom-Right: A cherry-picked example of model predictions. The model is able to interpolate smoothly between the data samples (dashed lines). However, it also produces some artifacts close to the 15cm depth.

We report the resulting thermal diffusivity constants and how they relate to plant diversity in Figure 2, alongside the general climatic conditions during the selected intervals and a selected example of a model fit. We found an inverse relation between plant diversity and thermal diffusivity ($p \leq 0.0002$), tested via a linear mixed effects model with random intercepts, (Oberg & Mahoney, 2007)). These findings are consistent with the results in (Huang et al., 2024) and provide an explanation for the increased soil temperature stability of the soil systems with high plant diversity. Further, we find that the temporal variance of thermal diffusivity of individual plots seems to differ strongly. However, contrary to what we expected based on the results of (Huang et al., 2024), this variance is not generally reduced through plant diversity.(Figure 2, bottom-left). Further evidence might provide a clearer picture here as our sample size is limited, and the expected effect might show up only over a longer period of time. Thirdly, we provide predictions of some example PINNs (Plot 2A08 for the *hot* interval) in Figure 2 to depict the learned representation of the system. We find that the PINN can properly predict the data samples while providing reasonable interpolations between depths(We display all depths from 0-15 cm). Finally, since the network architecture we deployed is quite powerful, we evaluated whether our PINNs can find a nonsensical solution to Equation 2 or abuse hidden bias in the data. For this, we performed a permutation test where we shuffled the phase of the Fourier-transformed data and again estimated the thermal diffusivity. The results can be found in the **??**. We find that no meaningful patterns can be learned in this setup and there is no relationship between plant diversity and K ($p \leq 0.607$).

5 CONCLUSION

In this work, we investigated how plant diversity affects the thermal diffusivity of the corresponding soil. Our first findings show an inverse relationship between plant diversity and thermal diffusivity, confirming this previously hypothesized (most likely indirect, (Huang et al., 2024)) mechanism that governs the stability of soil temperature. These findings support the notion that plant diversity

can serve as a natural shield against climate change-related ecosystem stress. Further, we confirm the applicability of PINNs to ecological hypotheses. Here, we have to, however, also note that PINNs have their own specific intricates that have to be kept in mind. Most notably, we found that optimizing for Equation 2 can sometimes lead to models that have low loss but fail to interpolate for all depths inside the domain properly. Such an effect is visible in Figure 2(bottom-right) where temperature interpolations are sometimes outside of the boundary values. We believe that additional regularization might help to overcome this imprecision in the future.

Finally, while this study provides a first insight into this relationship, it simplifies the soil system, ignoring additional interactions between depth, climatic conditions, soil moisture, phenological cycles, and thermal diffusivity. We plan on investigating these relationships in the future to extend our understanding of the soil system's intricacies. Possible options to incorporate more physical knowledge would be to leverage soil moisture data, use more complex PDEs like the Richardson-Richards equation (Bandai & Ghezzehei, 2022) or heat and moisture transfer models Künzel (1995). Finally, for future work, we plan to extend our work to the full 20 years of data that the Jena experiment provides. With this, we hope to have sufficient data to evaluate how the relationship between thermal diffusivity and plant diversity changes over time and, ultimately, through a broad spectrum of climatic conditions.

6 Acknowledgments

We gratefully recognize the support of iDiv (German Centre of Integrative Biodiversity Research), which is funded by the German Research Foundation (DFG – FZT 118, 202548816). Gideon Stein is funded by the iDiv flexpool (No 06203674-22). Maha Shadaydeh is funded by the Carl Zeiss Foundation within the scope of the program line "Breakthroughs: Exploring Intelligent Systems" for "Digitization—explore the basics (No P2017-01-003), use applications".

References

Tatiana Arkhangelskaya and Ksenia Lukyashchenko. Estimating soil thermal diffusivity at different water contents from easily available data on soil texture, bulk density, and organic carbon content. *Biosystems Engineering*, 168:83–95, April 2018. ISSN 1537-5110. doi: 10.1016/j.biosystemseng. 2017.06.011. URL https://www.sciencedirect.com/science/article/pii/S1537511016305621.

T. Bandai and T. A. Ghezzehei. Forward and inverse modeling of water flow in unsaturated soils with discontinuous hydraulic conductivities using physics-informed neural networks with domain decomposition. *Hydrology and Earth System Sciences*, 26(16):4469–4495, 2022. doi: 10.5194/hess-26-4469-2022. URL https://hess.copernicus.org/articles/26/4469/2022/.

HS Carlslaw and JC Jaeger. Conduction of heat in solids, oxford. 1959.

Shiping Chen, Wantong Wang, Wenting Xu, Yang Wang, Hongwei Wan, Dima Chen, Zhiyao Tang, Xuli Tang, Guoyi Zhou, Zongqiang Xie, Daowei Zhou, Zhouping Shangguan, Jianhui Huang, Jin-Sheng He, Yanfen Wang, Jiandong Sheng, Lisong Tang, Xinrong Li, Ming Dong, Yan Wu, Qiufeng Wang, Zhiheng Wang, Jianguo Wu, F. Stuart Chapin, and Yongfei Bai. Plant diversity enhances productivity and soil carbon storage. *Proceedings of the National Academy of Sciences*, 115(16):4027–4032, April 2018. doi: 10.1073/pnas.1700298114. URL https://www.pnas.org/doi/abs/10.1073/pnas.1700298114.

Xinli Chen, Han Y. H. Chen, Chen Chen, Zilong Ma, Eric B. Searle, Zaipeng Yu, and Zhiqun Huang. Effects of plant diversity on soil carbon in diverse ecosystems: a global meta-analysis. *Biological Reviews*, 95(1):167–183, 2020. ISSN 1469-185X. doi: 10.1111/brv.12554. URL https://onlinelibrary.wiley.com/doi/abs/10.1111/brv.12554.

YingYing Chen, Kun Yang, WenJun Tang, Jun Qin, and Long Zhao. Parameterizing soil organic carbon's impacts on soil porosity and thermal parameters for Eastern Tibet grasslands. *Science China Earth Sciences*, 55(6):1001–1011, June 2012. ISSN 1869-1897. doi: 10.1007/s11430-012-4433-0. URL https://doi.org/10.1007/s11430-012-4433-0.

Salvatore Cuomo, Vincenzo Schiano Di Cola, Fabio Giampaolo, Gianluigi Rozza, Maziar Raissi, and Francesco Piccialli. Scientific machine learning through physics–informed neural networks: Where we are and what's next. *Journal of Scientific Computing*, 92(3):88, Jul 2022.

Hans J. De Boeck, Juliette M. G. Bloor, Juergen Kreyling, Johannes C. G. Ransijn, Ivan Nijs, Anke Jentsch, and Michaela Zeiter. Patterns and drivers of biodiversity–stability relationships under climate extremes. *Journal of Ecology*, 106(3):890–902, 2018. ISSN 1365-2745. doi: 10. 1111/1365-2745.12897. URL https://onlinelibrary.wiley.com/doi/abs/10. 1111/1365-2745.12897. _eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1111/1365-2745.12897.

Kevin Gross, Bradley J. Cardinale, Jeremy W. Fox, Andrew Gonzalez, Michel Loreau, H. Wayne Polley, Peter B. Reich, and Jasper van Ruijven. Species Richness and the Temporal Stability of Biomass Production: A New Analysis of Recent Biodiversity Experiments. *The American Naturalist*, 183(1):1–12, January 2014. ISSN 0003-0147. doi: 10.1086/673915. URL https://www.journals.uchicago.edu/doi/10.1086/673915.

Charles Guy. The Influence of Temperature Extremes on Gene Expression, Genomic Structure, and the Evolution of Induced Tolerance in Plants. In *Plant Responses to Environmental Stresses*. Routledge, 1999. ISBN 978-0-203-74315-7. Num Pages: 52.

Yuanyuan Huang, Gideon Stein, Olaf Kolle, Karl Kübler, Ernst-Detlef Schulze, Hui Dong, David Eichenberg, Gerd Gleixner, Anke Hildebrandt, Markus Lange, Christiane Roscher, Holger Schielzeth, Bernhard Schmid, Alexandra Weigelt, Wolfgang W. Weisser, Maha Shadaydeh, Joachim Denzler, Anne Ebeling, and Nico Eisenhauer. Enhanced stability of grassland soil temperature by plant diversity. *Nature Geoscience*, 17(1):44–50, January 2024. ISSN 1752-0908. doi: 10.1038/s41561-023-01338-5. URL https://www.nature.com/articles/s41561-023-01338-5.

Forest Isbell, Dylan Craven, John Connolly, Michel Loreau, Bernhard Schmid, Carl Beierkuhnlein, T. Martijn Bezemer, Catherine Bonin, Helge Bruelheide, Enrica de Luca, Anne Ebeling, John N. Griffin, Qinfeng Guo, Yann Hautier, Andy Hector, Anke Jentsch, Jürgen Kreyling, Vojtěch Lanta, Pete Manning, Sebastian T. Meyer, Akira S. Mori, Shahid Naeem, Pascal A. Niklaus, H. Wayne Polley, Peter B. Reich, Christiane Roscher, Eric W. Seabloom, Melinda D. Smith, Madhav P. Thakur, David Tilman, Benjamin F. Tracy, Wim H. van der Putten, Jasper van Ruijven, Alexandra Weigelt, Wolfgang W. Weisser, Brian Wilsey, and Nico Eisenhauer. Biodiversity increases the resistance of ecosystem productivity to climate extremes. *Nature*, 526(7574):574–577, October 2015. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature15374. URL http://www.nature.com/articles/nature15374.

George Em Karniadakis, Ioannis G. Kevrekidis, Lu Lu, Paris Perdikaris, Sifan Wang, and Liu Yang. Physics-informed machine learning. *Nature Reviews Physics*, 3(6):422–440, Jun 2021. ISSN 2522-5820. doi: 10.1038/s42254-021-00314-5. URL https://doi.org/10.1038/s42254-021-00314-5.

Hartwig M Künzel. Simultaneous heat and moisture transport in building components. *One-and two-dimensional calculation using simple parameters. IRB-Verlag Stuttgart*, 65, 1995.

Michel Loreau and Claire de Mazancourt. Biodiversity and ecosystem stability: a synthesis of underlying mechanisms. *Ecology Letters*, 16(s1):106–115, 2013. ISSN 1461-0248. doi: 10.1111/ele.12073. URL https://onlinelibrary.wiley.com/doi/abs/10. 1111/ele.12073.

Michel Loreau, Matthieu Barbier, Elise Filotas, Dominique Gravel, Forest Isbell, Steve J. Miller, Jose M. Montoya, Shaopeng Wang, Raphaël Aussenac, Rachel Germain, Patrick L. Thompson, Andrew Gonzalez, and Laura E. Dee. Biodiversity as insurance: from concept to measurement and application. *Biological Reviews*, 96(5):2333–2354, October 2021. ISSN 1464-7931. doi: 10. 1111/brv.12756. URL https://doi.org/10.1111/brv.12756. Publisher: John Wiley & Sons, Ltd.

Ilya Loshchilov and Frank Hutter. Decoupled Weight Decay Regularization. September 2018. URL https://openreview.net/forum?id=Bkg6RiCqY7.

Lu Lu, Raphael Pestourie, Wenjie Yao, Zhicheng Wang, Francesc Verdugo, and Steven G Johnson. Physics-informed neural networks with hard constraints for inverse design. *SIAM Journal on Scientific Computing*, 43(6):B1105–B1132, 2021. doi: 10.1137/21M1397908.

K. I. Lukiashchenko and T. A. Arkhangelskaya. Modelling Thermal Diffusivity of Differently Textured Soils. *Eurasian Soil Science*, 51(2):183–189, February 2018. ISSN 1556-195X. doi: 10.1134/S1064229318020084. URL https://doi.org/10.1134/S1064229318020084.

Catarina C. Nievola, Camila P. Carvalho, Victória Carvalho, and Edson Rodrigues. Rapid responses of plants to temperature changes. *Temperature: Multidisciplinary Biomedical Journal*, 4(4):371–405, November 2017. ISSN 2332-8940. doi: 10.1080/23328940.2017.1377812. URL https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5800372/.

Ann L. Oberg and Douglas W. Mahoney. Linear Mixed Effects Models. In Walter T. Ambrosius (ed.), *Topics in Biostatistics*, Methods in Molecular Biology™, pp. 213–234. Humana Press, Totowa, NJ, 2007. ISBN 9781597455305. doi: 10.1007/978-1-59745-530-5_11. URL https://doi.org/10.1007/978-1-59745-530-5_11.

Brown Mang Onwuka. Effects of soil temperature on some soil properties and plant growth. *Advances in Plants & Agriculture Research*, 8(1):34–37, January 2018. ISSN 23736402. doi: 10.15406/apar.2018.08.00288. URL https://medcraveonline.com/APAR/effects-of-soil-temperature-on-some-soil-properties-and-plant-growth.html.

M. Raissi, P. Perdikaris, and G. E. Karniadakis. Physics-informed neural networks: A deep learning framework for solving forward and inverse problems involving nonlinear partial differential equations. *Journal of Computational Physics*, 378:686–707, February 2019. ISSN 0021-9991. doi: 10.1016/j.jcp.2018.10.045. URL https://www.sciencedirect.com/science/article/pii/S0021999118307125.

Marie Spohn, Sumanta Bagchi, Lori A. Biederman, Elizabeth T. Borer, Kari Anne Bråthen, Miguel N. Bugalho, Maria C. Caldeira, Jane A. Catford, Scott L. Collins, Nico Eisenhauer, Nicole Hagenah, Sylvia Haider, Yann Hautier, Johannes M. H. Knops, Sally E. Koerner, Lauri Laanisto, Ylva Lekberg, Jason P. Martina, Holly Martinson, Rebecca L. McCulley, Pablo L. Peri, Petr Macek, Sally A. Power, Anita C. Risch, Christiane Roscher, Eric W. Seabloom, Carly Stevens, G. F. (Ciska) Veen, Risto Virtanen, and Laura Yahdjian. The positive effect of plant diversity on soil carbon depends on climate. *Nature Communications*, 14(1):6624, October 2023. ISSN 2041-1723. doi: 10.1038/s41467-023-42340-0. URL https://www.nature.com/articles/s41467-023-42340-0.

David Tilman, Peter B. Reich, and Johannes M. H. Knops. Biodiversity and ecosystem stability in a decade-long grassland experiment. *Nature*, 441(7093):629–632, June 2006. ISSN 1476-4687. doi: 10.1038/nature04742. URL https://www.nature.com/articles/nature04742. Number: 7093 Publisher: Nature Publishing Group.

Wolfgang W Weisser, Christiane Roscher, Sebastian T Meyer, Anne Ebeling, Guangjuan Luo, Eric Allan, Holger Beßler, Romain L Barnard, Nina Buchmann, François Buscot, and others. Biodiversity effects on ecosystem functioning in a 15-year grassland experiment: Patterns, mechanisms, and open questions. *Basic and Applied Ecology*, 23:1–73, 2017. Publisher: Elsevier.

R. E. Wildung, T. R. Garland, and R. L. Buschbom. The interdependent effects of soil temperature and water content on soil respiration rate and plant root decomposition in arid grassland soils. *Soil Biology and Biochemistry*, 7(6):373–378, November 1975. ISSN 0038-0717. doi: 10.1016/0038-0717(75)90052-8. URL https://www.sciencedirect.com/science/article/pii/0038071775900528.

Dan Zhu, Philippe Ciais, Gerhard Krinner, Fabienne Maignan, Albert Jornet Puig, and Gustaf Hugelius. Controls of soil organic matter on soil thermal dynamics in the northern high latitudes. *Nature Communications*, 10(1):3172, July 2019. ISSN 2041-1723. doi: 10.1038/s41467-019-11103-1. URL https://www.nature.com/articles/s41467-019-11103-1.