Unifying Width-Reduced Methods for Quasi-Self-Concordant Optimization

Abstract

We provide several algorithms for constrained optimization of a large class of convex problems, including softmax, ℓ_p regression, and logistic regression. Central to our approach is the notion of width reduction, a technique which has proven immensely useful in the context of maximum flow [Christiano et al., STOC'11] and, more recently, ℓ_p regression [Adil et al., SODA'19], in terms of improving the iteration complexity from $O(m^{1/2})$ to $\tilde{O}(m^{1/3})$, where m is the number of rows of the design matrix, and where each iteration amounts to a linear system solve. However, a considerable drawback is that these methods require both problem-specific potentials and individually tailored analyses.

As our main contribution, we initiate a new direction of study by presenting the first *unified* approach to achieving $m^{1/3}$-type rates. Notably, our method goes beyond these previously considered problems to more broadly capture *quasi-self-concordant* losses, a class which has recently generated much interest and includes the well-studied problem of logistic regression, among others. In order to do so, we develop a unified width reduction method for carefully handling these losses based on a more general set of potentials. Additionally, we directly achieve $m^{1/3}$-type rates in the constrained setting without the need for any explicit acceleration schemes, thus naturally complementing recent work based on a ball-oracle approach [Carmon et al., NeurIPS'20].

1 Introduction

We study a class of constrained optimization problems of the following form:

$$\min_{\boldsymbol{Ax}=\boldsymbol{b}} \sum_i \boldsymbol{f}\big((\boldsymbol{Px})_i\big) \tag{1}$$

for convex $\boldsymbol{f} : \mathbb{R} \to \mathbb{R}$, where $\boldsymbol{A} \in \mathbb{R}^{d \times n}, \boldsymbol{b} \in \mathbb{R}^d, \boldsymbol{P} \in \mathbb{R}^{m \times n}$, with $d \le n \le m$. Specifically, we are interested in the case where $\boldsymbol{f}$ satisfies a certain higher-order smoothness-like condition known as M-quasi-self-concordance (q.s.c.), i.e., $|\boldsymbol{f}'''(\boldsymbol{x})| \le M\boldsymbol{f}''(\boldsymbol{x})$ for all $\boldsymbol{x} \in \mathbb{R}$. Several problems of significant interest in machine learning and numerical methods meet this condition, including logistic regression [Bac10, KSJ18], as well as softmax (often used to approximate ℓ_∞ regression) [Nes05, CKM+11, EV19, Bul20] and (regularized) ℓ_p regression [BCLL18, AKPS19].

A very useful optimization technique, first introduced by [CKM+11] for faster approximate maximum flow and later by [CMMP13] for regression, is that of width reduction, whereby they used it to improve the iteration complexity dependence on m, the number of rows of the design matrix from $O(m^{1/2})$ to $\tilde{O}(m^{1/3})$, and where each iteration requires a linear system solve. Later work by [AKPS19] for high-accuracy ℓ_p regression, building on an $O(m^{1/2})$-iteration result from [BCLL18], again showed

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

how width reduction could lead to improved $\tilde{O}(m^{1/3})$-iteration algorithms. As a drawback, however, these approaches rely on potential methods and analyses specifically tailored to each problem.

Building on these results, we present the first *unified* approach to achieving $m^{1/3}$-type rates, at the heart of which lies a more general width reduction scheme. Notably, our method goes beyond these previously considered problems to capture *quasi-self-concordant* losses, thereby further including well-studied problems such as logistic regression, among others. By doing so, we directly achieve $m^{1/3}$-type rates in the constrained setting without relying on explicit acceleration schemes [MS13], thus complementing recent work based on a ball-oracle approach [CJJ+20]. We additionally note that, given the ways in which our results achieve improvements similar to those of [CJJ+20], we believe our work hints at a deeper, though to our knowledge not yet fully understood, connection between the techniques of width reduction and Monteiro-Svaiter acceleration.

1.1 Main results and applications

We first present in Section 3 a width reduced method for obtaining a crude approximation to (1) for quasi-self-concordant $\boldsymbol{f}$. At a high level, our algorithm returns an approximate solution $\tilde{\boldsymbol{x}}$ that both satisfies the linear constraints and is bounded in ℓ_∞-norm by $O(R)$, where R is a bound on the norm of the optimal solution. Following from Theorem 3.3, the result below shows how, for the problem of minimizing softmax (parameterized by $\nu > 0$), we can bound the norm of the solution by $(1 + \nu)R$.

Theorem 1.1. *Let $\boldsymbol{x}^\star$ denote the optimum of $\min_{\boldsymbol{A}\boldsymbol{x}=\boldsymbol{b}} smax_\nu(\boldsymbol{P}\boldsymbol{x})$. Algorithm 1 when applied to the function $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}$ with $\epsilon = \nu$, returns $\tilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\tilde{\boldsymbol{x}} = \boldsymbol{b}$, and*

$$smax_\nu(\boldsymbol{P}\tilde{\boldsymbol{x}}) \le (1 + \widetilde{O}(\nu))smax_\nu(\boldsymbol{P}\boldsymbol{x}^\star),$$

in at most $\widetilde{O}(m^{1/3}\nu^{-5/3})$ calls to a linear system solver.

As a consequence of Theorem 1.1 when taking $\nu = \Omega\left(\epsilon / \log^{O(1)}(m)\right)$, we have by Theorem 5.2 a $(1 + \epsilon)$ approximate solution to the problem of ℓ_∞ regression with $\tilde{O}(m^{1/3}\epsilon^{-5/3})$ calls to a linear system solver.

Further, we show the following result that can utilize the approximate solution returned by Theorem 1.1 as an initial point for achieving a high-accuracy solution. We also present in Appendix A a natural extension of our results to minimizing generalized-self-concordant (g.s.c.) functions.

Theorem 1.2. *For M-q.s.c. $\boldsymbol{f}$, $\epsilon > 0$, and $\boldsymbol{x}^{(0)}$ such that $\boldsymbol{A}\boldsymbol{x}^{(0)} = \boldsymbol{b}$ and $\|\boldsymbol{x}^{(0)}\|_\infty \le R$, Algorithm 2 finds $\tilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\tilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\boldsymbol{f}(\tilde{\boldsymbol{x}}) \le \boldsymbol{f}(\boldsymbol{x}^\star) + \epsilon$ in $\widetilde{O}\left(MRm^{1/3}\log(MR)\log\left(\frac{\boldsymbol{f}(\boldsymbol{x}^{(0)}) - \boldsymbol{f}(\boldsymbol{x}^\star)}{\epsilon}\right)\right)$ calls to a linear system solver.*

Resulting from the theorem above, as detailed in Section 5 are guarantees given by Theorems 5.4 and 5.5 which establish convergence rates of $\widetilde{O}(p^2\mu^{-1/(p-2)}m^{1/3}R)$ and $\widetilde{O}(m^{1/3}R)$, respectively, for μ-regularized ℓ_p regression and logistic regression. We emphasize that the latter is, to our knowledge, the first such use of width reduction for directly solving constrained logistic regression problems.

1.2 Related Works

Quasi-self-concordance and higher-order smoothness.   [Bac10] showed how to analyze Newton's method for quasi-self-concordant functions, with an emphasis on its application to logistic regression. Later, notions of local, or Hessian, stability which follow from quasi-self-concordance gave rise to methods with better dependence on various conditioning parameters [KSJ18, CJJ+20]. In the work by [KSJ18], the authors show how a trust-region-based Newton method [NW06] achieves linear convergence for locally stable functions without requiring, e.g., strong convexity. Meanwhile, after noting that quasi-self-concordance implies Hessian stability, [CJJ+20] further improve the dependence on the distance to the optimum by leveraging Monteiro-Svaiter acceleration [MS13], which has proved useful in the context of near-optimal methods for higher-order acceleration [ASS19, GDG+19]. However, in general these methods, which assume higher-order smoothness, require access to an oracle which minimizes a higher-order Taylor expansion model, though in some cases this may be relaxed to requiring linear system solves [Bul20].

Width reduction and ℓ_p regression. The technique of width-reduction first came to prominence in seminal work by [CKM$^+$11] for achieving faster approximate maximum flow, being the first to achieve an improved $m^{1/3}$ dependence. At a high level, the idea behind the approach is to solve a sequence of weighted ℓ_2-minimizing flow problems, whereby at each iteration one of two cases occurs: either the proposed step is added to the current solution (a "flow" step) along with the weights, or else there exist some set of coordinates that exceed a certain threshold, and so their weights are updated accordingly (a "width reduction" step). Several works have since adapted this approach to regression problems [CMMP13, AKPS19, EV19, ABKS21] and matrix scaling [AZLOW17].

In addition to their importance in machine learning, regression methods capture several fundamental problems in scientific computing and signal processing. A recent line of work initiated by [BCLL18] showed how to attain high-accuracy solutions for ℓ_p regression using $O_p(m^{1/2-1/p})$ linear system solves, thus going beyond what is achievable via self-concordance. Building on this work, [AKPS19] showed how width reduction could be applied to this setting to achieve, as in the case of approximate maximum flow [CKM$^+$11], a similar improvement from $O_p(m^{1/2})$ to $O_p(m^{1/3})$ (for $p \to \infty$). Further developments by [KPSW19, AS20] for graph problems showed almost-linear time solutions for ℓ_p regression for $p \approx \sqrt{\log(n)}$ which have since been a critical part of recent advances in high-accuracy maximum flow on unit-capacity graphs [LS20, KLS20].

Accelerated methods. Recent developments by [CJJ$^+$20] have shown several advantages that arise in the case of unconstrained minimization for L-smooth M-quasi-self-concordant problems. By considering a certain ball oracle method (whereby each call to the oracle returns the minimizer of the function inside an ℓ_2 ball of radius r), [CJJ$^+$20] implement an accelerated scheme which returns a solution to the unconstrained smooth convex minimization problem in $(R/r)^{2/3}$ calls to the oracle, where R is the initial ℓ_2-norm distance to the optimum, and they further show a matching lower bound under this oracle model. While this approach transfers its difficulty to implementing the oracle, a key insight from their work involves showing this can be done efficiently for smooth quasi-self-concordant functions when r is sufficiently small, where the allowed size depends on the quasi-self-concordance parameter. One limitation to the results of [CJJ$^+$20] is that they apply directly to *unconstrained* optimization problems and require the function to be smooth, and so we complement these results in the quasi-self-concordant setting by establishing comparable rates for directly optimizing a large class of *constrained* convex problems without requiring smoothness of the function.

1.3 Outline of the Paper

After establishing the potential functions at the heart of our width reduction techniques, we present in Section 3 our oracle for roughly approximating a solution to problem (1). We then show in Section 4 how we may attain a high-accuracy solution by using the crude approximation as a starting point. Here, the key idea is to considering a sequence of optimization problems inside ℓ_∞ balls of manageable size, similar to [CMTV17, CJJ$^+$20]. As in the case of the crude oracle, our primary advantage comes from carefully handling a pair of coupled potentials which are amenable to the large class of quasi-self-concordant problems, and in Section 5 we further show how our results may be applied to several problems of interest, including logistic and ℓ_p regression. The missing proofs can be found in the appendix.

2 Preliminaries

Notation: We use boldface lowercase letters to denote vectors or functions and boldface uppercase letters for matrices. Scalars are non-bold alphabets. Our functions are univariate and we overload function notation to act on a vector coordinate-wise, i.e. $\boldsymbol{f}(\boldsymbol{x}) = \sum_i \boldsymbol{f}(\boldsymbol{x}_i)$. The notation $\boldsymbol{x} \geq \boldsymbol{y}$ for vectors refers to entry-wise inequality. Refer to the algorithm boxes for definitions of certain algorithm specific parameters that appear in Lemma/Theorem statements.

2.1 Quasi Self-Concordance

Definition 2.1 (g.s.c. and q.s.c.). *Let $\boldsymbol{f} : \mathbb{R} \to \mathbb{R}$ be a thrice differentiable function with continuous third derivative, and let $\nu > 0$ and $M > 0$ be two constants. We say that $\boldsymbol{f}$ is (M, ν)-generalized*

self-concordant (g.s.c.) if

$$\forall x, \quad |\boldsymbol{f}'''(x)| \le M\boldsymbol{f}''(x)^{\frac{\nu}{2}}.$$

When $\nu = 2$, we get the following condition,

$$\forall x, \quad |\boldsymbol{f}'''(x)| \le M\boldsymbol{f}''(x),$$

and we call such functions M-quasi-self-concordant (q.s.c.).

2.2 Problem

Recall that we are solving the following problem.

$$\min_{\boldsymbol{Ax=b}} \sum_i \boldsymbol{f}\big((\boldsymbol{Px})_i\big),$$

$\boldsymbol{A} \in \mathbb{R}^{d\times n}, \boldsymbol{b} \in \mathbb{R}^d, \boldsymbol{P} \in \mathbb{R}^{m\times n}, d \le n, m \ge n, \boldsymbol{f}$ is convex, M-q.s.c. and, for $\boldsymbol{w} \ge \boldsymbol{w}_0 \ge 0$, $\boldsymbol{f}''(\boldsymbol{w})$ is coordinate-wise monotonic. We can ignore the case when $\boldsymbol{f}''$ is constant since that corresponds to a quadratic problem which we know how to solve directly via linear system solves.

Assumptions on the optimum $\boldsymbol{x}^\star$

We assume that $R \in \mathbb{R}_{>0}$ is such that the optimum $\boldsymbol{x}^\star \stackrel{\text{def}}{=} \arg\min_{\boldsymbol{Ax=b}} \boldsymbol{f}(\boldsymbol{Px})$ satisfies

$$\|\boldsymbol{Px}^\star\|_\infty \le R. \tag{2}$$

We now define potentials that we will track in the algorithm.

2.3 Potentials

Definition 2.2 (Dual Potential). *For a weights vector $\boldsymbol{w} \in \mathbb{R}^m_{\ge 0}$, we define a potential*

$$\Phi(\boldsymbol{w}) = \sum_i \Phi(\boldsymbol{w}_i) = \sum_i \boldsymbol{f}''(\boldsymbol{w}_i).$$

We define the following linear regression problem that we will solve at each step of our algorithm, as our oracle.

Definition 2.3 (Resistances and Oracle). *For a weights vector $\boldsymbol{w} \in \mathbb{R}^m_{\ge 0}$ and $\epsilon > 0$, define resistances $\boldsymbol{r} \in \mathbb{R}^m_{\ge 0}$ and a corresponding potential Ψ as,*

$$\boldsymbol{r}_i = \frac{1}{R^2}\left(\boldsymbol{f}''(\boldsymbol{w}_i) + \frac{\epsilon\Phi(\boldsymbol{w})}{m}\right),$$

$$\Psi(\boldsymbol{r}) = \min_{\boldsymbol{A\Delta=b}} \sum_i \boldsymbol{r}_i(\boldsymbol{P\Delta})_i^2.$$

We have the following relation between our two potentials Φ and Ψ.

Lemma 2.4. *For $\epsilon > 0$, resistances $\boldsymbol{r}$ (Definition 2.3), with corresponding weights $\boldsymbol{w}$, we have*

$$\Psi(\boldsymbol{r}) \le (1 + \epsilon)\Phi(\boldsymbol{w}),$$

and for $\|\boldsymbol{P}\|_{\min} = \min_{\boldsymbol{Ax=b}} \|\boldsymbol{Px}\|_2, \|\boldsymbol{A}\|$ the operator norm of $\boldsymbol{A}$,

$$\Psi(\boldsymbol{r}) \ge \frac{\epsilon\Phi(\boldsymbol{w})}{mR^2} \frac{\|\boldsymbol{P}\|_{\min}^2\|\boldsymbol{b}\|_2^2}{\|\boldsymbol{A}\|^2} \stackrel{\text{def}}{=} \Phi(\boldsymbol{w})L.$$

3 Algorithm and Analysis for a Crude Solution for Q.S.C. Functions

In this section, we give an algorithm for solving Problem (1) to a crude approximation i.e., we return a solution $\widetilde{x}$ such that $A\widetilde{x} = b$ (satisfies subspace constraints) and $\|P\widetilde{x}\|_\infty$ is bounded. We will see later in our applications how this translates into a constant or polynomial approximation guarantee to the function value for some functions. In the next section we will see how we can use the guarantees of the solution returned as a starting solution and boost it to an ϵ approximate solution.

Our algorithm is based on a multiplicative weight update scheme with width reduction. Such algorithms have only been used for ℓ_p-regression, $p = 1$ or $p \in [2, \infty]$. Here, we are able to extend the analysis to q.s.c. functions and also give a unified analysis for the known cases of ℓ_p-regression (Refer to Section 5 to see how we apply this algorithm to these special functions). We note that we can extend this analysis to other general-self-concordant functions. We have deferred these cases to the appendix.

3.1 Algorithm and Analysis

Our proof is based on tracking of two potentials, Ψ (Definition 2.3) and Φ (Definition 2.2) that depend on the weights. We first show how these potentials change with weight updates corresponding to a flow step and a width reduction step in the algorithm. We next show that if our algorithm runs for at most $K = \widetilde{O}(m^{1/3})$ width reduction steps, then after $T = \widetilde{O}(m^{1/3})$ flow steps we can bound Φ. Further, using the relation between Φ and Ψ (Lemma 2.4) and appropriately chosen parameters, we show that we cannot have more than K width reduction steps. The key part of the analysis lies in the growth of Φ with respect to both, a flow and width step.

Algorithm 1 Width Reduced Algorithm for M-q.s.c. Functions with Φ q.s.c.

1: **procedure** QSC-MWU(A, b, P, M, R, ϵ)
2: $\quad x^{(0,0)} = 0, w^{(0,0)} = w_0$ ($\Phi'(w)$ monotonic for $w \geq w_0$, $\Phi(w_0) > 0$)
3: $\quad \tau \leftarrow \widetilde{\Theta}\left(m^{1/3}\epsilon^{-2/3}\right)$
4: $\quad \alpha \leftarrow \widetilde{\Theta}\left(m^{-1/3}M^{-1}\epsilon^{2/3}\right)$
5: $\quad t = 0, k = 0, \quad T = \alpha^{-1}M^{-1}\epsilon^{-1} = \widetilde{\Theta}(m^{1/3}\epsilon^{-5/3})$
6: $\quad$ **while** $t \leq T$ **do**
7: $\qquad r_i^{(t,k)} \leftarrow \frac{1}{R^2}\left(f''(w_i^{(t,k)}) + \frac{\epsilon\Phi(w^{(t,k)})}{m}\right)$ $\qquad\qquad\qquad$ ▷ Resistances
8: $\qquad \widetilde{\Delta} \leftarrow \arg\min_{A\Delta = b} \quad \sum_i r_i(P\Delta)_i^2$ $\qquad\qquad\qquad\qquad$ ▷ Oracle
9: $\qquad$ **if** $\left\|P\widetilde{\Delta}\right\|_\infty \leq R\tau$ **then** $\qquad\qquad\qquad\qquad\qquad$ ▷ Flow Step
10: $\qquad\quad x^{(t+1,k)} \leftarrow x^{(t,k)} + \widetilde{\Delta}$
11: $\qquad\quad w^{(t+1,k)} \leftarrow w^{(t,k)} + \frac{\epsilon\alpha}{R}|P\widetilde{\Delta}|$
12: $\qquad\quad t \leftarrow t + 1$
13: $\qquad$ **else**
14: $\qquad\quad$ **for** Indices i such that $|P\widetilde{\Delta}|_i \geq R\tau$ **do** $\qquad\qquad$ ▷ Width Reduction
15: $\qquad\qquad$ **if** f'' is non-decreasing in w **then**[1]
16: $\qquad\qquad\quad w^{(t,k+1)}$ is such that $r_i^{(t,k+1)} \leftarrow (1 + \epsilon)r_i^{(t,k)}$
17: $\qquad\qquad$ **else**
18: $\qquad\qquad\quad w^{(t,k+1)}$ is such that $r_i^{(t,k+1)} \leftarrow \frac{1}{1+\epsilon}r_i^{(t,k)}$
19: $\qquad\quad k \leftarrow k + 1$
20: $\quad$ **return** $x^{(T,k)}/T$

---

[1]We will see later how such weight/resistance changes can be realized for some special cases.

Changes in Ψ and Φ

Lemma 3.1. *Let Ψ be as defined in [2.3]. After t flow steps and k width reduction steps, we have,*

$$\Psi(\boldsymbol{r}^{(t,k)}) \geq \Psi(\boldsymbol{r}^{(0,0)})\left(1 + \frac{\epsilon^2\tau^2}{2(1+\epsilon)^2 m}\right)^k \qquad \text{if } \boldsymbol{f}'' \text{ non-decreasing in } \boldsymbol{w},$$

$$\Psi(\boldsymbol{r}^{(t,k)}) \leq \Psi(\boldsymbol{r}^{(0,0)})\left(1 - \frac{\epsilon^2\tau^2}{4(1+\epsilon)^2 m}\right)^k \qquad \text{if } \boldsymbol{f}'' \text{ non-increasing in } \boldsymbol{w}.$$

Lemma 3.2. *Suppose $\boldsymbol{f}$ is M-q.s.c. Let α and τ be such that, $\alpha\tau \leq M^{-1}$. After t flow steps and k width reduction steps our potential Φ satisfies,*

$$\Phi(\boldsymbol{w}^{(t,k)}) \leq \left(1 + \epsilon(1+\epsilon)^2\alpha M\right)^t \left(1 + \epsilon(1+\epsilon)\tau^{-1}\right)^k \Phi(\boldsymbol{w}_0) \quad \text{if } \boldsymbol{f}'' \text{ non-decreasing in } \boldsymbol{w},$$

$$\Phi(\boldsymbol{w}^{(t,k)}) \geq \left(1 - \epsilon(1+\epsilon)^2\alpha M\right)^t \left(1 - \epsilon(1+\epsilon)\tau^{-1}\right)^k \Phi(\boldsymbol{w}_0) \quad \text{if } \boldsymbol{f}'' \text{ non-increasing in } \boldsymbol{w}.$$

Runtime Bound

Theorem 3.3. *Let $\epsilon > 0$, $\boldsymbol{f}$ be M-q.s.c. After $T \leq \frac{\alpha^{-1}}{M\epsilon} = \widetilde{\Theta}(m^{1/3}\epsilon^{-5/3})$ flow steps and $K \leq \tau = \widetilde{\Theta}(m^{1/3}\epsilon^{-2/3})$ width reduction steps, Algorithm [1] returns $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$, $\|\boldsymbol{P}\widetilde{\boldsymbol{x}}\|_\infty \leq RM\|\boldsymbol{w}^{(T,K)}\|_\infty$, where $\boldsymbol{w}^{(T,K)}$ is the final weights vector that satisfies,*

$$\Phi(\boldsymbol{w}^{(T,K)}) \leq \Phi(\boldsymbol{w}_0)e^{1+4\epsilon} \qquad \text{if } \boldsymbol{f}'' \text{ is non-decreasing in } \boldsymbol{w},$$

$$\Phi(\boldsymbol{w}^{(T,K)}) \geq \Phi(\boldsymbol{w}_0)e^{-(1+4\epsilon)} \qquad \text{if } \boldsymbol{f}'' \text{ is non-increasing in } \boldsymbol{w}.$$

Proof. We show the case when $\boldsymbol{f}''$ is a non-decreasing function. The other case follows similarly. We set,

$$\tau \leftarrow \widetilde{\Theta}\left(m^{1/3}\epsilon^{-2/3}\right) \quad \alpha \leftarrow \widetilde{\Theta}\left(m^{-1/3}M^{-1}\epsilon^{2/3}\right).$$

After $T = \frac{\alpha^{-1}}{M\epsilon}$ flow steps and $K = \tau$ width reduction steps, from Lemma [3.2], we have,

$$\Phi(\boldsymbol{w}^{(T,K)}) \leq \left(1 + \epsilon(1+\epsilon)^2\alpha M\right)^T \left(1 + \epsilon(1+\epsilon)\tau^{-1}\right)^K \Phi(\boldsymbol{w}_0)$$

$$\leq \Phi(\boldsymbol{w}_0)e^{\epsilon(1+\epsilon)^2\alpha MT + \epsilon(1+\epsilon)\tau^{-1}K} \leq \Phi(\boldsymbol{w}_0)e^{(1+4\epsilon)}.$$

We will now show we cannot have more width steps. Throughout the algorithm, we have $\Phi(\boldsymbol{w}^{(t,k)}) \leq \Phi(\boldsymbol{w}_0)e^{1+4\epsilon}$. From Lemma [2.4] we always have $\Psi(\boldsymbol{r}^{(0,0)}) \geq \Phi(\boldsymbol{w}_0)L$ and $\Psi(\boldsymbol{r}^{(T,K)}) \leq (1 + \epsilon)\Phi(\boldsymbol{w}^{(T,K)}) \leq (1+\epsilon)e^{1+4\epsilon}\Phi(\boldsymbol{w}_0)$. Thus, from Lemma [3.1], we must have,

$$(1 + \epsilon)e^{1+4\epsilon}\Phi(\boldsymbol{w}_0) \geq L\Phi(\boldsymbol{w}_0)\left(1 + \frac{\epsilon^2\tau^2}{2(1+\epsilon)^2 m}\right)^K,$$

From the definition of τ, we note that K has to be less than τ for the above bound to be satisfied. Next, let $\widetilde{\Delta}^{(t)}$ denote the solution of our oracle at iteration t of the flow step. From the $\boldsymbol{x}$ and $\boldsymbol{w}$ update in the algorithm,

$$|\boldsymbol{P}\widetilde{\boldsymbol{x}}| = \left|\sum_t \boldsymbol{P}\widetilde{\Delta}^{(t)}\right|\epsilon\alpha M \leq \frac{\epsilon\alpha}{R}\sum_t \left|\boldsymbol{P}\widetilde{\Delta}^{(t)}\right|RM \leq \boldsymbol{w}^{(T,K)}RM.$$

This concludes our proof. $\square$

 ## 4 Boosting to a High-Accuracy Solution for Q.S.C. Functions

In this section, we will give a width reduced multiplicative weights update algorithm that, given a starting solution $\boldsymbol{x}^{(0)}$ that satisfies $\|\boldsymbol{x}^{(0)}\|_\infty \leq R$ and $\boldsymbol{A}\boldsymbol{x}^{(0)} = \boldsymbol{b}$, finds $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\boldsymbol{f}(\widetilde{\boldsymbol{x}}) \leq (1+\epsilon)\boldsymbol{f}(\boldsymbol{x}^\star)$ for any q.s.c. function $\boldsymbol{f}$. We would like to mention here that for the algorithms in this section, it is key that we have a starting solution satisfying our subspace constraints and have ℓ_∞-norm bounded by R. Thus, the algorithms here may be of independent interest if such a starting solution is available. We can use Algorithm 1 with $\epsilon = 1$ to obtain such a solution.

For any $\boldsymbol{x}$, we define a residual problem. We will show how it is sufficient to solve the residual problem approximately $\log(\epsilon^{-1})$ times to obtain our high-accuracy solution. Similar approaches have been applied to specific functions such as soft-max [AZLOW17] and ℓ_p-regression [AKPS19]. We unify these approaches and give a version that works for any q.s.c. function.

We further note that our residual problem is to optimize a simple quadratic objective inside an infinity box. The difficulty lies in solving such ℓ_∞ box constraints fast. We use a binary search followed by a width reduced multiplicative weights routine analogous to [CKM+11] to solve our residual problem.

Definition 4.1 (Residual Problem). *We define the residual objective at any $\boldsymbol{x}$ satisfying $\|\boldsymbol{P}\boldsymbol{x}\|_\infty \leq R$ as,*

$$res(\Delta) = \nabla \boldsymbol{f}(\boldsymbol{x})^\top \boldsymbol{P}\Delta - e^{-1}(\boldsymbol{P}\Delta)^\top \nabla^2 \boldsymbol{f}(\boldsymbol{x})\boldsymbol{P}\Delta,$$

and the residual problem as,

$$\max_\Delta \quad res(\Delta)$$
$$s.t. \quad \boldsymbol{A}\Delta = 0, \quad and \quad \|\boldsymbol{P}\Delta - \boldsymbol{z}\|_\infty \leq \frac{1}{2M}. \tag{3}$$

Here $\boldsymbol{z}$ is a vector that depends on $\boldsymbol{x}$, and is defined as,

$$\boldsymbol{z}_i = \begin{cases} \left(-\frac{1}{2M} + R + (\boldsymbol{P}\boldsymbol{x})_i\right) \in [-\frac{1}{2M}, 0)], & if \ (\boldsymbol{P}\boldsymbol{x})_i - \frac{1}{2M} < -R \\ \left(-R + (\boldsymbol{P}\boldsymbol{x})_i + \frac{1}{2M}\right) \in (0, \frac{1}{2M}], & if \ (\boldsymbol{P}\boldsymbol{x})_i + \frac{1}{2M} > R \\ 0, & otherwise. \end{cases}$$

We note that any solution Δ satisfying the above box constraint satisfies $\|\boldsymbol{P}\Delta\|_\infty \leq M^{-1}$ and $\|\boldsymbol{P}\boldsymbol{x} - e^{-2}\boldsymbol{P}\Delta\|_\infty \leq R$.

Lemma 4.2. *[Iterative Refinement] Let $\boldsymbol{f}$ be M-q.s.c. and $\widetilde{\Delta}^{(t)}$ a κ-approximate solution to the residual problem at $\boldsymbol{x}^{(t)}$ (Problem (3)). Starting from $\boldsymbol{x}^{(0)}$ such that $\boldsymbol{A}\boldsymbol{x}^{(0)} = \boldsymbol{b}$, $\|\boldsymbol{x}^{(0)}\|_\infty \leq R$, and iterating as $\boldsymbol{x}^{(t+1)} = \boldsymbol{x}^{(t)} - e^{-2}\widetilde{\Delta}^{(t)}$, after at most $O\left(\kappa M R \log\left(\frac{\boldsymbol{f}(\boldsymbol{x}^{(0)}) - \boldsymbol{f}(\boldsymbol{x}^\star)}{\epsilon}\right)\right)$ iterations we get $\boldsymbol{x}$ such that $\boldsymbol{A}\boldsymbol{x} = \boldsymbol{b}$ and, $\boldsymbol{f}(\boldsymbol{x}) \leq \boldsymbol{f}(\boldsymbol{x}^\star) + \epsilon$.*

4.1 Approximately Solving the Residual Problem

Binary Search

Lemma 4.3. *Let ν be such that $\boldsymbol{f}(\boldsymbol{x}^{(t)}) - \boldsymbol{f}(\boldsymbol{x}^\star) \in (\nu/2, \nu]$ and $\Delta^\star$ denote the optimum of the residual problem at $\boldsymbol{x}^{(t)}$. Then, $res(\Delta^\star) \in \left(\frac{\nu}{8MR}, e^2\nu\right]$.*

From the above lemma we can do a binary search in the range $\left(\frac{\nu}{8MR}, e^2\nu\right]$. Let us start with the assumption the residual problem has a solution between $(\zeta/2, \zeta]$.

Lemma 4.4. *Let ζ be such that $res(\Delta^\star) \in (\zeta/2, \zeta]$ and $\Delta^\star$ the optimum of the residual problem. Then, $(\boldsymbol{P}\Delta^\star)^\top \nabla^2 \boldsymbol{f}(\boldsymbol{x})\boldsymbol{P}\Delta^\star \leq e \cdot \zeta$.*

Using Width Reduction

We will show that Algorithm 3 returns Δ such that $\|\boldsymbol{P}\Delta - \boldsymbol{z}\|_\infty \leq \frac{1}{2M}$ and $res(\Delta) \geq \frac{1}{400}\zeta$.

Lemma 4.5. *Let ζ be such that $res(\Delta^\star) \in (\zeta/2, \zeta]$. Algorithm 3 returns $\boldsymbol{y}$ such that $\boldsymbol{A}\boldsymbol{y} = 0$, $\|\boldsymbol{P}\boldsymbol{y} - \boldsymbol{z}\|_\infty \leq \frac{1}{2M}$ and $res(\boldsymbol{y}) \geq \frac{1}{400}res(\Delta^\star)$ in $O(m^{1/3})$ calls to a linear system solver.*

Algorithm 2 Boosting to ϵ-approximation

1: **procedure** QSC-MIN($\boldsymbol{A}, \boldsymbol{b}, \boldsymbol{P}, \boldsymbol{x}_0, M, \epsilon$) such that $\boldsymbol{A}\boldsymbol{x}_0 = \boldsymbol{b}, \|\boldsymbol{x}_0 - \boldsymbol{x}^\star\|_\infty \le 2R$)
2: $\quad$ $\boldsymbol{x}^{(0)} = \boldsymbol{x}_0, \tau \leftarrow m^{1/3}, \alpha \leftarrow m^{-1/3}$
3: $\quad$ **for** $i \le O(MR \log \epsilon^{-1})$ **do**
4: $\quad\quad$ **for** $\nu \in \big(\epsilon, \boldsymbol{f}(\boldsymbol{x})\big]$ **do** $\qquad\qquad\qquad$ ▷ Decrease ν by 2 in each iteration
5: $\quad\quad\quad$ **for** $\zeta \in \big(\frac{\nu}{8MR}, e^2\nu\big]$ **do** $\qquad\qquad$ ▷ Decrease ζ by 2 in each iteration
6: $\quad\quad\quad\quad$ $\boldsymbol{y}_{\zeta,\nu} \leftarrow MWU(\boldsymbol{A}, \boldsymbol{P}, \boldsymbol{x}^{(i)}, M, \zeta)$
7: $\quad\quad$ $\boldsymbol{x}^{(i+1)} \leftarrow \boldsymbol{x}^{(i)} - e^{-2} \arg\min_{\boldsymbol{y}_{\zeta,\nu}} \boldsymbol{f}(\boldsymbol{x} - e^{-2}\boldsymbol{y}_{\zeta,\nu})$

Algorithm 3

1: **procedure** MWU($\boldsymbol{A}, \boldsymbol{P}, \boldsymbol{x}, M, \zeta$)
2: $\quad$ $\boldsymbol{y}^{(0)} = 0, \boldsymbol{w}^{(0)} = \frac{\zeta}{m}$
3: $\quad$ $t = 0$
4: $\quad$ $\boldsymbol{A}' = \left[\boldsymbol{A}^\top, \boldsymbol{P}^\top \nabla \boldsymbol{f}(\boldsymbol{x})\right]^\top, \boldsymbol{b} = [0, \frac{\zeta}{2}]$
5: $\quad$ **while** $\|\boldsymbol{w}\|_1 \le 10\zeta$ **do**
6: $\quad\quad$ $\widetilde{\Delta} \leftarrow \arg\min_{\boldsymbol{A}'\Delta = \boldsymbol{b}'} \sum_j \boldsymbol{f}''(\boldsymbol{x}_j)(\boldsymbol{P}\Delta)_j^2 + 4M^2 \sum_j \left(\boldsymbol{w}_j^{(t)} + \frac{\|\boldsymbol{w}^{(t)}\|_1}{m}\right)(\boldsymbol{P}\Delta - \boldsymbol{z})_j^2$
7: $\quad\quad$ **if** $2M\left\|\boldsymbol{P}\widetilde{\Delta} - \boldsymbol{z}\right\|_\infty \le \tau$ **then** $\qquad\qquad\qquad\qquad\qquad$ ▷ Flow Step
8: $\quad\quad\quad$ $\boldsymbol{y}^{(t+1)} \leftarrow \boldsymbol{y}^{(t)} + \widetilde{\Delta}$
9: $\quad\quad\quad$ $\boldsymbol{w}^{(t+1)} \leftarrow \boldsymbol{w}^{(t)}\left(1 + \frac{1}{2}\alpha M|\boldsymbol{P}\widetilde{\Delta} - \boldsymbol{z}|\right)$
10: $\quad\quad$ **else**
11: $\quad\quad\quad$ **for** Indices i such that $2M|\boldsymbol{P}\widetilde{\Delta} - \boldsymbol{z}|_i \ge \tau$ **do**
12: $\quad\quad\quad\quad$ $\boldsymbol{w}_i^{(t+1)} \leftarrow 2\boldsymbol{w}_i^{(t)}$ $\qquad\qquad\qquad\qquad\qquad\qquad$ ▷ Width Step
13: $\quad\quad$ $t \leftarrow t + 1$
14: $\quad$ **return** $\frac{\boldsymbol{y}^{(t)}}{100t}$

We now state the main result of the section which follows directly from Lemmas 4.2, 4.3 and 4.5

Theorem 4.6. *For $\epsilon > 0$, M-q.s.c. function $\boldsymbol{f}$ and, $\boldsymbol{x}^{(0)}$ such that $\boldsymbol{A}\boldsymbol{x}^{(0)} = \boldsymbol{b}$, $\|\boldsymbol{x}^{(0)}\|_\infty \le R$, Algorithm 2 finds $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\boldsymbol{f}(\widetilde{\boldsymbol{x}}) - \boldsymbol{f}(\boldsymbol{x}^\star) \le \epsilon$ in*
$$\widetilde{O}\left(MRm^{1/3}\log(MR)\log\left(\frac{\boldsymbol{f}(\boldsymbol{x}^{(0)}) - \boldsymbol{f}(\boldsymbol{x}^\star)}{\epsilon}\right)\right) \text{ calls to a linear system solver.}$$

5 Application to some Special Functions

We now provide an overview of our methods as applied to various quasi-self-concordant functions.

5.1 Sum of exponential, soft-max and ℓ_∞ regression

The function soft-max with parameter ν is defined as $smax_\nu(\boldsymbol{P}\boldsymbol{x}) = \nu \log\left(\sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}\right)$ and is $1/\nu$-q.s.c. We will start with assuming that at the optimum, for $R \ge \Omega\big((\log m)^{-1}\big)$, $smax_\nu(\boldsymbol{P}\boldsymbol{x}^\star) \le R$.

We will apply Algorithm 1 to $\sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}$ which is also $1/\nu$-q.s.c. We can use the following weight update step for the width reduction step : $\boldsymbol{w}_i^{(t,k+1)} \leftarrow \boldsymbol{w}_i^{(t,k)} + \nu \log(1 + \epsilon)$.

Theorem 5.1. *Let $\boldsymbol{x}^\star$ denote the optimum of $smax_\nu$. Algorithm 1 when applied to the function $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}$ with $\epsilon = \nu$, returns $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$, and*
$$smax_t(\boldsymbol{P}\widetilde{\boldsymbol{x}}) \le (1 + \widetilde{O}(\nu))smax_\nu(\boldsymbol{P}\boldsymbol{x}^\star),$$
in at most $\widetilde{O}(m^{1/3}\nu^{-5/3})$ calls to a linear system solver.

Proof. We know that $smax_\nu(\boldsymbol{P}\widetilde{\boldsymbol{x}}) \leq \|\boldsymbol{P}\widetilde{\boldsymbol{x}}\|_\infty + \nu \log m$. From Lemma 3.3, we get that $\widetilde{\boldsymbol{x}}$ is obtained in at most $\widetilde{O}(m^{1/3}\nu^{-5/3})$ calls to a linear system solver satisfying $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$. Further, we also have, $\|\boldsymbol{P}\widetilde{\boldsymbol{x}}\|_\infty \leq MR\|\boldsymbol{w}^{(T,K)}\|_\infty \leq R\frac{\|\boldsymbol{w}^{(T,K)}\|_\infty}{\nu}$. We will now bound $\frac{\|\boldsymbol{w}^{(T,K)}\|_\infty}{\nu}$. We note that $\Phi(\boldsymbol{w}^{(T,K)}) \leq \Phi(\boldsymbol{w}_0)e^{1+4\nu}$. For $\boldsymbol{w}_0 = 0$,

$$\Phi(\boldsymbol{w}^{(T,K)}) = \frac{1}{\nu^2}\sum_i e^{\frac{w_i^{(T,K)}}{\nu}} = \Phi(\boldsymbol{w}_0)\sum_i e^{\frac{w_i^{(T,K)}}{\nu}} \leq \Phi(\boldsymbol{w}_0)e^{1+4\nu}.$$

Therefore, we must have $\boldsymbol{w}^{(T,K)} \leq (1+4\nu)\nu$. Our bound is,

$$smax_\nu(\boldsymbol{P}\widetilde{\boldsymbol{x}}) \leq (1+4\nu)R + \nu\log m \leq \Big(1 + \widetilde{O}(\nu)\Big)R,$$

for $R \geq \Omega(1/\log m)$. We can now do a binary search on R as follows to obtain,

$$smax_\nu(\boldsymbol{P}\widetilde{\boldsymbol{x}}) \leq (1 + \widetilde{O}(\nu))smax_\nu(\boldsymbol{P}\boldsymbol{x}^\star)$$

Binary search on R: Let R_0 denote the value $\|\boldsymbol{P}\boldsymbol{x}^\star\|_\infty = R_0$. Now, for any $R \geq R_0$, we get an $\widetilde{\boldsymbol{x}}$ giving objective value at most $R(1 + 4\epsilon)$. For any $R < R_0$, as long as R is such that the plane $\boldsymbol{A}\boldsymbol{x} = \boldsymbol{b}$ has at least one point with infinity norm at most R, we will get a feasible solution to our problem. However, the objective value guarantee of $R(1 + 4\epsilon)$ may not hold. Since the optimum is R_0, the solution returned in such cases must give an objective value larger than R_0. We can thus do a binary search on R and reach $O(\epsilon)$ close to the value R_0. This will require running our algorithm $O\big(\log(R_0\epsilon^{-1})\big)$ times. In the end we can return the $\boldsymbol{x}$ which gives the smallest objective values among all these runs. $\qquad\square$

Theorem 5.2. *Let $\boldsymbol{x}^\star$ denote the optimum of the ℓ_∞-regression problem, $\min_{\boldsymbol{A}\boldsymbol{x}=\boldsymbol{b}} \|\boldsymbol{P}\boldsymbol{x}\|_\infty$. Algorithm 1 when applied to the function $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i\Big(e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}} + e^{\frac{-(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}\Big)$ for $\nu = \Omega\Big(\frac{\epsilon}{\log m}\Big)$, returns $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and*

$$\|\boldsymbol{P}\widetilde{\boldsymbol{x}}\|_\infty \leq (1+\epsilon)\|\boldsymbol{P}\boldsymbol{x}^\star\|_\infty,$$

in at most $\widetilde{O}(m^{1/3}\epsilon^{-5/3})$ calls to a linear system solve.

Theorem 5.3. *For $\delta > 0$, let $\overline{\boldsymbol{x}}$ be the solution returned by Algorithm 1 (with $\epsilon = 1$) applied to $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x})_i}{\nu}}$. Now, Algorithm 2 with starting solution $\boldsymbol{x}^{(0)} = \overline{\boldsymbol{x}}$, applied to $\boldsymbol{f}$ finds $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\sum_i e^{\frac{(\boldsymbol{P}\widetilde{\boldsymbol{x}})_i}{\nu}} \leq (1+\delta)\sum_i e^{\frac{(\boldsymbol{P}\boldsymbol{x}^\star)_i}{\nu}}$ in at most $O\Big(m^{1/3}R^2\nu^{-2}\log\big(\frac{m}{\delta}\big)\Big)$ calls to a linear system solver.*

5.2 p-Norm Regression

We will solve, $\min_{\boldsymbol{A}\boldsymbol{x}=\boldsymbol{b}}\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \|\boldsymbol{P}\boldsymbol{x}\|_p^p + \mu\|\boldsymbol{P}\boldsymbol{x}\|_2^2$, for $p \geq 3$ which is $p\mu^{-1/(p-2)}$-q.s.c. w.r.t. its argument. We first apply Algorithm 1 to this function and use the returned solution as the starting point of Algorithm 2. We can use the following weight update step for the width reduction step : $\boldsymbol{w}_i^{(t,k+1)} \leftarrow (1+\epsilon)^{1/(p-2)}\boldsymbol{w}_i^{(t,k)}$.

Theorem 5.4. *For $\delta > 0$ and $p \geq 3$, let $\overline{\boldsymbol{x}}$ be the solution returned by Algorithm 1 (with $\epsilon = 1$) applied to $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \|\boldsymbol{P}\boldsymbol{x}\|_p^p + \mu\|\boldsymbol{P}\boldsymbol{x}\|_2^2$. Now, Algorithm 2 with starting solution $\boldsymbol{x}^{(0)} = \overline{\boldsymbol{x}}$, applied to $\boldsymbol{f}$ finds $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\boldsymbol{f}(\boldsymbol{P}\widetilde{\boldsymbol{x}}) \leq \boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}^\star) + \delta$ in at most $O\Big(p^2\mu^{-1/(p-2)}m^{1/3}R\log\Big(\frac{pmR}{\mu\delta}\Big)\Big)$ calls to a linear system solver.*

5.3 Logistic Regression

We will consider the function $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i \log(1 + e^{(\boldsymbol{P}\boldsymbol{x})_i})$ which is 1-q.s.c. w.r.t its argument. We will use Algorithm 1 with the following weight update for the width reduction step which reduces the resistance by a factor of $(1+\epsilon)$: $\boldsymbol{w}_i^{(t,k+1)} \leftarrow \boldsymbol{w}_i^{(t,k)} + 0.9\epsilon$

Theorem 5.5. *For $\delta > 0$, let $\overline{\boldsymbol{x}}$ be the solution returned by Algorithm 1 (with $\epsilon = 1$) applied to $\boldsymbol{f}(\boldsymbol{P}\boldsymbol{x}) = \sum_i \log(1 + e^{(\boldsymbol{P}\boldsymbol{x})_i})$. Now, Algorithm 2 with starting solution $\boldsymbol{x}^{(0)} = \overline{\boldsymbol{x}}$, applied to $\boldsymbol{f}$ finds $\widetilde{\boldsymbol{x}}$ such that $\boldsymbol{A}\widetilde{\boldsymbol{x}} = \boldsymbol{b}$ and $\sum_i \log(1 + e^{(\boldsymbol{P}\widetilde{\boldsymbol{x}})_i}) \leq \sum_i \log(1 + e^{(\boldsymbol{P}\boldsymbol{x}^\star)_i}) + \delta$ in at most $O\Big(m^{1/3}R\log\Big(\frac{mR}{\delta}\Big)\Big)$ calls to a linear system solver.*

References

[ABKS21] Deeksha Adil, Brian Bullins, Rasmus Kyng, and Sushant Sachdeva. Almost-linear-time weighted ℓ_p-norm solvers in slightly dense graphs via sparsification, 2021.

[AKPS19] Deeksha Adil, Rasmus Kyng, Richard Peng, and Sushant Sachdeva. Iterative refinement for ℓ_p-norm regression. In *Proceedings of the Thirtieth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1405–1424. SIAM, 2019.

[AS20] Deeksha Adil and Sushant Sachdeva. Faster p-norm minimizing flows, via smoothed q-norm problems. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 892–910. SIAM, 2020.

[ASS19] Yossi Arjevani, Ohad Shamir, and Ron Shiff. Oracle complexity of second-order methods for smooth convex optimization. *Mathematical Programming*, 178(1):327–360, 2019.

[AZLOW17] Zeyuan Allen-Zhu, Yuanzhi Li, Rafael Oliveira, and Avi Wigderson. Much faster algorithms for matrix scaling. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 890–901. IEEE, 2017.

[Bac10] Francis Bach. Self-concordant analysis for logistic regression. *Electronic Journal of Statistics*, 4:384–414, 2010.

[BCLL18] Sébastien Bubeck, Michael B Cohen, Yin Tat Lee, and Yuanzhi Li. An homotopy method for lp regression provably beyond self-concordance and in input-sparsity time. In *Proceedings of the 50th Annual ACM SIGACT Symposium on Theory of Computing*, pages 1130–1137, 2018.

[Bul20] Brian Bullins. Highly smooth minimization of non-smooth problems. In *Conference on Learning Theory*, pages 988–1030. PMLR, 2020.

[CJJ+20] Yair Carmon, Arun Jambulapati, Qijia Jiang, Yujia Jin, Yin Tat Lee, Aaron Sidford, and Kevin Tian. Acceleration with a ball optimization oracle. In H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 19052–19063. Curran Associates, Inc., 2020.

[CKM+11] Paul Christiano, Jonathan A Kelner, Aleksander Madry, Daniel A Spielman, and Shang-Hua Teng. Electrical flows, laplacian systems, and faster approximation of maximum flow in undirected graphs. In *Proceedings of the forty-third annual ACM symposium on Theory of computing*, pages 273–282, 2011.

[CMMP13] Hui Han Chin, Aleksander Madry, Gary L Miller, and Richard Peng. Runtime guarantees for regression problems. In *Proceedings of the 4th conference on Innovations in Theoretical Computer Science*, pages 269–282, 2013.

[CMTV17] Michael B Cohen, Aleksander Madry, Dimitris Tsipras, and Adrian Vladu. Matrix scaling and balancing via box constrained newton's method and interior point methods. In *2017 IEEE 58th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 902–913. IEEE, 2017.

[EV19] Alina Ene and Adrian Vladu. Improved convergence for ℓ_1 and ℓ_∞ regression via iteratively reweighted least squares. In *International Conference on Machine Learning*, pages 1794–1801. PMLR, 2019.

[GDG+19] Alexander Gasnikov, Pavel E. Dvurechensky, Eduard A. Gorbunov, Evgeniya A. Vorontsova, Daniil Selikhanovych, César A. Uribe, Bo Jiang, Haoyue Wang, Shuzhong Zhang, Sébastien Bubeck, Qijia Jiang, Yin Tat Lee, Yuanzhi Li, and Aaron Sidford. Near optimal methods for minimizing convex functions with lipschitz p-th derivatives. In *COLT*, pages 1392–1393, 2019.

[KLS20] Tarun Kathuria, Yang P Liu, and Aaron Sidford. Unit capacity maxflow in almost $m^{4/3}$ time. In *2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS)*, pages 119–130. IEEE, 2020.

[KPSW19] Rasmus Kyng, Richard Peng, Sushant Sachdeva, and Di Wang. Flows in almost linear time via adaptive preconditioning. In *Proceedings of the 51st Annual ACM SIGACT Symposium on Theory of Computing*, pages 902–913, 2019.

[KSJ18] Sai Praneeth Karimireddy, Sebastian U Stich, and Martin Jaggi. Global linear convergence of newton's method without strong-convexity or lipschitz gradients. *arXiv preprint arXiv:1806.00413*, 2018.

[LS20] Yang P Liu and Aaron Sidford. Faster energy maximization for faster maximum flow. In *Proceedings of the 52nd Annual ACM SIGACT Symposium on Theory of Computing*, pages 803–814, 2020.

[MS13] Renato DC Monteiro and Benar Fux Svaiter. An accelerated hybrid proximal extragradient method for convex optimization and its implications to second-order methods. *SIAM Journal on Optimization*, 23(2):1092–1125, 2013.

[Nes05] Yu Nesterov. Smooth minimization of non-smooth functions. *Mathematical programming*, 103(1):127–152, 2005.

[NW06] Jorge Nocedal and Stephen Wright. *Numerical optimization*. Springer Science & Business Media, 2006.

[STD19] Tianxiao Sun and Quoc Tran-Dinh. Generalized self-concordant functions: a recipe for newton-type methods. *Mathematical Programming*, 178(1):145–213, 2019.