DENOISING DIFFUSION SAMPLERS

ABSTRACT

Denoising diffusion models are a popular class of generative models providing state-of-the-art results in many domains. One adds gradually noise to data using a diffusion to transform the data distribution into a Gaussian distribution. Samples from the generative model are then obtained by simulating an approximation of the time-reversal of this diffusion initialized by Gaussian samples. Practically, the intractable score terms appearing in the time-reversed process are approximated using score matching techniques. We explore here a similar idea to sample approximately from unnormalized probability density functions and estimate their normalizing constants. We consider a process where the target density diffuses towards a Gaussian. Denoising Diffusion Samplers (DDS) are obtained by approximating the corresponding time-reversal. While score matching is not applicable in this context, we can leverage many of the ideas introduced in generative modeling for Monte Carlo sampling. Existing theoretical results from denoising diffusion models also provide theoretical guarantees for DDS. We discuss the connections between DDS, optimal control and Schrödinger bridges and finally demonstrate DDS experimentally on a variety of challenging sampling tasks.

1 INTRODUCTION

Let π be a probability density on $\mathbb{R}^d$ of the form

$$\pi(x) = \frac{\gamma(x)}{Z}, \qquad Z = \int_{\mathbb{R}^d} \gamma(x) \mathrm{d}x, \tag{1}$$

where $\gamma : \mathbb{R}^d \to \mathbb{R}^+$ can be evaluated pointwise but the normalizing constant Z is intractable. We are here interested in both estimating Z and obtaining approximate samples from π.

A large variety of Monte Carlo techniques has been developed in the literature to address this problem. In particular Annealed Importance Sampling (AIS) (Neal, 2001) and its Sequential Monte Carlo (SMC) extensions (Del Moral et al., 2006) are often regarded as the "gold standard" approach to compute normalizing constants. These methods rely on the introduction of a sequence of annealed versions of π which are explored using a combination of Markov chain Monte Carlo (MCMC), importance sampling and resampling steps.

Variational techniques are a popular alternative to MCMC and SMC where one considers a flexible family of easy-to-sample distributions q^θ whose parameters are optimized by minimizing a suitable metric w.r.t. π, typically the reverse Kullback–Leibler discrepancy $\mathrm{KL}(q^\theta || \pi)$. Typical choices for q^θ include mean-field approximation (Wainwright & Jordan, 2008) or more sophisticated techniques such as normalizing flows (Papamakarios et al., 2019). To be able to model complex variational distributions, it has often proven useful to model $q^\theta(x)$ as the marginal of an auxiliary extended distribution; i.e. $q^\theta(x) = \int q^\theta(x, u) \mathrm{d}u$. As this marginal is typically intractable, θ is then learned by minimizing a discrepancy measure between $q^\theta(x, u)$ and an extended target distribution $p^\theta(x, u) = \pi(x) p^\theta(u|x)$ where $p^\theta(u|x)$ is an auxiliary conditional distribution (Agakov & Barber, 2004).

Over recent years, variational techniques have also been used fruitfully to optimize the algorithmic parameters of Monte Carlo techniques. For example, AIS can be thought of a procedure where $q^\theta(x, u)$ is the joint distribution of a Markov chain defined by a sequence of MCMC kernels whose final state is x while $p^\theta(x, u)$ is the corresponding AIS extended target distribution (Neal, 2001). The parameters θ of these kernels can then be optimized by minimizing $\mathrm{KL}(q^\theta || p^\theta)$ using stochastic

gradient descent (Wu et al., 2020; Geffner & Domke, 2021; Thin et al., 2021; Zhang et al., 2021; Doucet et al., 2022; Geffner & Domke, 2022).

In this work, we follow an approach inspired by Denoising Diffusion Probabilistic Models (DDPM), a powerful class of generative models (Sohl-Dickstein et al., 2015; Ho et al., 2020; Song et al., 2021b). In this context, one adds noise progressively to data using a diffusion to transform the complex data distribution into a Gaussian distribution. The time-reversal of this diffusion can then be used to transform a Gaussian sample into a sample from the target. While superficially similar to Langevin dynamics, this process mixes fast even in high dimensions as it inherits the mixing properties of the forward diffusion (De Bortoli et al., 2021b). However, as the time-reversal involves the derivatives of the logarithms of the intractable marginal densities of the forward diffusion, these so-called scores are practically approximated using score matching techniques. If the score estimation error is small, the approximate time-reversal still enjoys remarkable theoretical properties (De Bortoli, 2022; Chen et al., 2022; Lee et al., 2022).

These results motivate us to introduce Denoising Diffusion Samplers (DDS). Like DDPM, we consider a forward diffusion which progressively transforms the target π into a Gaussian distribution. This defines an extended target distribution $p(x, u) = \pi(x)p(u|x)$. DDS are obtained by approximating the time-reversal of this diffusion using a process of distribution $q^\theta(x, u)$. What distinguishes DDS from DDPM is that we cannot simulate sample paths from the diffusion we want to time-reverse, as we cannot sample its initial state x from π. Hence score matching ideas cannot be used to approximate the score terms.

We focus on minimizing $\mathrm{KL}(q^\theta||p)$, equivalently maximizing an Evidence Lower Bound (ELBO), as in variational inference. We leverage a representation of this KL discrepancy based on the introduction of a suitable auxiliary reference process that provides low variance estimate of this objective and its gradient. We can exploit the many similarities between DDS and DDPM to leverage some of the ideas developed in generative modeling for Monte Carlo sampling. This includes using the probability flow ordinary differential equation (ODE) (Song et al., 2021b) to derive novel normalizing flows and the use of underdamped Langevin diffusions as a forward noising diffusion (Dockhorn et al., 2022). The implementation of these samplers requires designing numerical integrators for the resulting stochastic differential equations (SDE) and ODE. In particular, it is practically highly desirable to preserve a valid ELBO in discrete time as, contrary to DDPM, we are here interested in evaluating the normalizing constant Z. To achieve this, DDS requires designing a numerical integrator for the auxiliary reference process preserving exactly its marginals distributions as well as an integrator for the approximate time-reversal inducing a distribution absolutely continuous w.r.t. the distribution of the discretized reference process. This prevents the use of integrators such as the standard Euler–Maryuama scheme for overdamped and underdamped Langevin noising diffusions for the reference process as well as the use of the integrator used in (Dockhorn et al., 2022) for the time reversal. Finally we compare experimentally DDS to state-of-the-art Monte Carlo methods.

2   DENOISING DIFFUSION SAMPLERS: CONTINUOUS TIME

We start here by formulating DDS in continuous-time to gain insight on the structure of the time-reversal we want to approximate. We introduce $\mathcal{C} = C([0, T], \mathbb{R}^d)$ the space of continuous functions from $[0, T]$ to $\mathbb{R}^d$ and $\mathcal{B}(\mathcal{C})$ the Borel sets on $\mathcal{C}$. We will consider in this section path measures which are probability measures on $(\mathcal{C}, \mathcal{B}(\mathcal{C}))$, see Léonard (2014a) for a formal definition. Numerical integrators are discussed in the following section.

2.1   FORWARD DIFFUSION AND ITS TIME-REVERSAL

Consider the forward noising diffusion given by an Ornstein–Uhlenbeck process[1]

$$\mathrm{d}x_t = -\beta_t x_t \mathrm{d}t + \sigma\sqrt{2\beta_t}\mathrm{d}B_t, \qquad x_0 \sim \pi, \tag{2}$$

where $(B_t)_{t \in [0,T]}$ is a d-dimension Brownian motion and $t \to \beta_t$ is a non-decreasing positive function. This diffusion induces the path measure $\mathcal{P}$ on the time interval $[0, T]$ and the marginal density of x_t is denoted p_t. The transition density of this diffusion is given by $p_{t|0}(x_t|x_0) = $

---

[1]This is referred to as a Variance Preserving diffusion by Song et al. (2021b).

$\mathcal{N}(x_t; \sqrt{1 - \lambda_t}x_0, \sigma^2\lambda_t I)$ where $\lambda_t = 1 - \exp\left(-2\int_0^t \beta_s \mathrm{d}s\right)$. From now on, we will always consider a scenario where $\int_0^T \beta_s \mathrm{d}s \gg 1$ so that $\lambda_T \approx 1$ and $p_T(x_T) \approx \mathcal{N}(x_T; 0, \sigma^2 I)$. We can thus think of (2) as transporting approximately the target density π to this Gaussian density.

From (Haussmann & Pardoux, 1986), its time-reversal $(y_t)_{t \in [0,T]} = (x_{T-t})_{t \in [0,T]}$, where equality is here in distribution, satisfies

$$\mathrm{d}y_t = \beta_{T-t}\left\{y_t + 2\sigma^2\nabla\ln p_{T-t}(y_t)\right\}\mathrm{d}t + \sigma\sqrt{2\beta_{T-t}}\mathrm{d}W_t, \qquad y_0 \sim p_T, \qquad (3)$$

where $(W_t)_{t \in [0,T]}$ is another d-dimensional Brownian motion. By definition this time-reversal starts from $y_0 \sim p_T(y_0) \approx \mathcal{N}(y_0; 0, \sigma^2 I)$ and is such that $y_T \sim \pi$. This suggests that if we could approximately simulate the diffusion (3), then we would obtain approximate samples from π.

However, putting this idea in practice requires being able to approximate the intractable scores $(\nabla\ln p_t(x))_{t \in [0,T]}$. To achieve this, DDPM rely on score matching techniques (Hyvärinen, 2005; Vincent, 2011) as it is easy to sample from (2). This is impossible in our scenario as sampling from (2) requires sampling $x_0 \sim \pi$ which is impossible by assumption.

2.2 REFERENCE DIFFUSION AND VALUE FUNCTION

In our context, it will useful to introduce a *reference* process defined by the diffusion following (2) but initialized at $p_0^{\mathrm{ref}}(x_0) = \mathcal{N}(x_0; 0, \sigma^2 I)$ rather than $\pi(x_0)$ thus ensuring that the marginals of the resulting path measure $\mathcal{P}^{\mathrm{ref}}$ all satisfy $p_t^{\mathrm{ref}}(x_t) = \mathcal{N}(x_t; 0, \sigma^2 I)$. From the chain rule for KL for path measures (see e.g. (Léonard, 2014b, Theorem 2.4) and (De Bortoli et al., 2021a, Proposition 24)), one has $\mathrm{KL}(\mathcal{Q}||\mathcal{P}^{\mathrm{ref}}) = \mathrm{KL}(q_0||p_0^{\mathrm{ref}}) + \mathbb{E}_{x_0 \sim q_0}(\mathrm{KL}(\mathcal{Q}(\cdot|x_0)||\mathcal{P}^{\mathrm{ref}}(\cdot|x_0)))$. Thus $\mathcal{P}$ can be identified as the path measure minimizing the KL discrepancy w.r.t. $\mathcal{P}^{\mathrm{ref}}$ over the set of path measures $\mathcal{Q}$ with marginal $q_0(x_0) = \pi(x_0)$ at time $t = 0$, i.e. $\mathcal{P} = \arg\min_{\mathcal{Q}}\left\{\mathrm{KL}(\mathcal{Q}||\mathcal{P}^{\mathrm{ref}}) : q_0 = \pi\right\}$.

A time-reversal representation of $\mathcal{P}^{\mathrm{ref}}$ is given by $(y_t)_{t \in [0,T]} = (x_{T-t})_{t \in [0,T]}$ satisfying

$$\mathrm{d}y_t = -\beta_{T-t}y_t\mathrm{d}t + \sigma\sqrt{2\beta_{T-t}}\mathrm{d}W_t, \qquad y_0 \sim p_0^{\mathrm{ref}}, \qquad (4)$$

as $\beta_{T-t}y_t + 2\sigma^2\nabla\ln p_{T-t}^{\mathrm{ref}}(y_t) = -\beta_{T-t}y_t$ and we can rewrite the time-reversal (3) of $\mathcal{P}$ as

$$\mathrm{d}y_t = -\beta_{T-t}\left\{y_t - 2\sigma^2\nabla\ln\phi_{T-t}(y_t)\right\}\mathrm{d}t + \sigma\sqrt{2\beta_{T-t}}\mathrm{d}W_t, \qquad y_0 \sim p_T, \qquad (5)$$

where $\phi_t(x) = p_t(x)/p_t^{\mathrm{ref}}(x)$ is a so-called value function which satisfies a Kolmogorov equation such that $\phi_t(x_t) = \mathbb{E}_{\mathcal{P}^{\mathrm{ref}}}[\phi_0(x_0)|x_t]$, the expectation being w.r.t. the law of the reference process.

2.3 LEARNING THE TIME-REVERSAL

To approximate the time-reversal (3) of $\mathcal{P}$, consider a path measure $\mathcal{Q}^\theta$ whose time-reversal is induced by

$$\mathrm{d}y_t = \beta_{T-t}\left\{y_t + 2\sigma^2 s_\theta(T - t, y_t)\right\}\mathrm{d}t + \sigma\sqrt{2\beta_{T-t}}\mathrm{d}W_t, \qquad y_0 \sim \mathcal{N}(0, \sigma^2 I), \qquad (6)$$

so that $y_t \sim q_{T-t}^\theta$. To obtain $s_\theta(t, x) \approx \nabla\ln p_t(x)$, we parameterize $s_\theta(t, x)$ by a neural network whose parameters are obtained by minimizing

$$\mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}) = \mathrm{KL}(p_T||\mathcal{N}(0, \sigma^2 I)) + \mathbb{E}_{y_0 \sim \mathcal{N}(0, \sigma^2 I)}[\mathrm{KL}(\mathcal{Q}^\theta(\cdot|y_0)||\mathcal{P}(\cdot|y_0))] \qquad (7)$$

$$= \mathrm{KL}(p_T||\mathcal{N}(0, \sigma^2 I)) + \sigma^2\mathbb{E}_{\mathcal{Q}^\theta}\left[\int_0^T \beta_{T-t}||s_\theta(T - t, y_t) - \nabla\ln p_{T-t}(y_t)||^2\mathrm{d}t\right],$$

where we have used the chain rule for KL then Girsanov's formula (see e.g. (Klebaner, 2012)). This expression of the KL is reminiscent of the expression obtained in (Song et al., 2021a, Theorem 1) in the context of DDPM. However, the expectation appearing in (7) is here w.r.t. $\mathcal{Q}^\theta$ and not w.r.t. $\mathcal{P}$ and we cannot get rid of the intractable scores $(\nabla\ln p_t(x))_{t \in [0,T]}$ using score matching ideas.

Instead, taking inspiration from (5), we reparameterize the time-reversal of $\mathcal{Q}^\theta$ using

$$\mathrm{d}y_t = -\beta_{T-t}\left\{y_t - 2\sigma^2 f_\theta(T - t, y_t)\right\}\mathrm{d}t + \sigma\sqrt{2\beta_{T-t}}\mathrm{d}W_t, \qquad y_0 \sim \mathcal{N}(0, \sigma^2 I), \qquad (8)$$

i.e. $f_\theta(t, x) = s_\theta(t, x) - \nabla \ln p_t^{\text{ref}}(x) = s_\theta(t, x) + x/\sigma^2$. This reparameterization allows us to express $\text{KL}(\mathcal{Q}^\theta || \mathcal{P})$ in a compact form.

Proposition 1. *The Radon-Nikodym derivative $\frac{\mathrm{d}\mathcal{Q}^\theta}{\mathrm{d}\mathcal{P}^{\text{ref}}}(y_{[0,T]})$ satisfies under $\mathcal{Q}^\theta$*

$$\ln\left(\frac{\mathrm{d}\mathcal{Q}^\theta}{\mathrm{d}\mathcal{P}^{\text{ref}}}\right) = \sigma^2 \int_0^T \beta_{T-t} ||f_\theta(T-t, y_t)||^2 \mathrm{d}t + \sigma \int_0^T \sqrt{2\beta_{T-t}} f_\theta(T-t, y_t)^\top \mathrm{d}W_t. \tag{9}$$

From the identity $\text{KL}(\mathcal{Q}^\theta || \mathcal{P}) = \text{KL}(\mathcal{Q}^\theta || \mathcal{P}^{\text{ref}}) + \mathbb{E}_{y_T \sim q_0^\theta}\left[\ln\left(\frac{p_0^{\text{ref}}(y_T)}{p_0(y_T)}\right)\right]$, *it follows that*

$$\text{KL}(\mathcal{Q}^\theta || \mathcal{P}) = \mathbb{E}_{\mathcal{Q}^\theta}\left[\sigma^2 \int_0^T \beta_{T-t} ||f_\theta(T-t, y_t)||^2 \mathrm{d}t + \ln\left(\frac{\mathcal{N}(y_T; 0, \sigma^2 I)}{\pi(y_T)}\right)\right]. \tag{10}$$

For θ minimizing (10), approximate samples from π can be obtained by simulating (8) and returning $y_T \sim q_0^\theta$. We can obtain an unbiased estimate of Z via the following importance sampling identity

$$\hat{Z} = \frac{\gamma(y_T)}{\mathcal{N}(y_T; 0, \sigma^2 I)} \frac{\mathrm{d}\mathcal{P}^{\text{ref}}}{\mathrm{d}\mathcal{Q}^\theta}(y_{[0,T]}), \qquad y_{[0,T]} \sim \mathcal{Q}^\theta. \tag{11}$$

where the second term can be computed directly from (9).

2.4 CONTINUOUS-TIME NORMALIZING FLOW

To approximate the log likelihood of DDPM, it was proposed by Song et al. (2021b) to rely on the probability flow ODE. This is an ODE admitting the same marginal distributions $(p_t)_{t \in [0,T]}$ as the noising diffusion given by $\mathrm{d}x_t = -\beta_t\{x_t + \sigma^2 \nabla \ln p_t(x_t)\}\mathrm{d}t$. We use here this ODE to design a continuous-time normalizing flow to sample from π. For θ such that $f_\theta(t, x) \approx \nabla \ln \phi_t(x)$ (obtained by minimization of the KL in (10)), we have $x_t + \sigma^2 \nabla \ln p_t(x) \approx \sigma^2 f_\theta(t, x)$. So it is possible to sample approximately from π by integrating the following ODE over $[0, T]$

$$\mathrm{d}y_t = \sigma^2 \beta_{T-t} f_\theta(T-t, y_t)\mathrm{d}t, \qquad y_0 \sim \mathcal{N}(0, \sigma^2 I). \tag{12}$$

If denote by $\bar{q}_{T-t}^\theta$ the distribution of y_t then $y_T \sim \bar{q}_0^\theta$ is an approximate sample from π. We can use this normalizing flow to obtain an unbiased estimate of Z using importance sampling $\hat{Z} = \gamma(y_T)/\bar{q}_0^\theta(y_T)$ for $y_T \sim \bar{q}_0^\theta$. Indeed, contrary to the proposal q_0^θ induced by (8), we can compute pointwise $\bar{q}_0^\theta$ using the instantaneous change of variables formula (Chen et al., 2018) such that $\ln \bar{q}_0^\theta(y_T) = \ln \mathcal{N}(y_0; 0, \sigma^2 I) - \sigma^2 \int_0^T \beta_{T-t} \nabla \cdot f_\theta(T-t, y_t)\mathrm{d}t$ and where $(y_t)_{t \in [0,T]}$ arises from the ODE (12). The expensive divergence term can be estimated using the Hutchinson estimator (Grathwohl et al., 2018; Song et al., 2021b).

2.5 EXTENSION TO UNDERDAMPED LANGEVIN DYNAMICS

For DDPM, it has been proposed to extend the original state $x \in \mathbb{R}^d$ by a momentum variable $p \in \mathbb{R}^d$. One then diffuses the data distribution augmented by a Gaussian distribution on the momentum using an underdamped Langevin dynamics targeting $\mathcal{N}(x; 0, \sigma^2 I)\mathcal{N}(p; 0, M)$ where M is a positive definite mass matrix (Dockhorn et al., 2022). It was demonstrated empirically that the resulting scores are smoother, hence easier to estimate and this leads to improved performance. We adapt here this approach to Monte Carlo sampling; see Section B for more details.

We diffuse $\bar{\pi}(x, p) = \pi(x)\mathcal{N}(p; 0, M)$ using an underdamped Langevin dynamics, i.e.

$$\mathrm{d}x_t = M^{-1} p_t \mathrm{d}t, \quad \mathrm{d}p_t = -\frac{x_t}{\sigma^2}\mathrm{d}t - \beta_t p_t \mathrm{d}t + \sqrt{2\beta_t} M^{1/2} \mathrm{d}B_t, \quad (x_0, p_0) \sim \bar{\pi}. \tag{13}$$

The resulting path measure on $[0, T]$ is denoted $\mathcal{P}$ and the marginal of (x_t, p_t) is denoted η_t.

Here, the reference process $\mathcal{P}^{\text{ref}}$ is defined by the diffusion (13) initialized using $x_0 \sim \mathcal{N}(0, \sigma^2 I), p_0 \sim \mathcal{N}(0, M)$. This ensures that $\eta_t^{\text{ref}}(x_t, p_t) = \mathcal{N}(x_t; 0, \sigma^2 I)\mathcal{N}(p_t; 0, M)$ for all t and the time-reversal process $(y_t, q_t)_{t \in [0,T]} = (x_{T-t}, p_{T-t})_{t \in [0,T]}$ (where equality is in distribution) of this stationary diffusion satisfies

$$\mathrm{d}y_t = -M^{-1} q_t \mathrm{d}t, \quad \mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t} q_t \mathrm{d}t + \sqrt{2\beta_{T-t}} M^{1/2} \mathrm{d}W_t. \tag{14}$$

Algorithm 1 DDS Training

1: **Input:** $\sigma > 0, \gamma : \mathbb{R}^d \to \mathbb{R}^+, (\beta_k)_{k=1}^K \in (\mathbb{R}^+)^K, \theta \in \mathbb{R}^p, \lambda > 0$
2: **for** $i = 1, \dots,$ training iterations **do**
3:     **for** $n = 1, \dots, N$ **do**   # Iterate over training batch size.
4:         Sample $y_{0,n} \sim \mathcal{N}(0, \sigma^2 I)$ and set $r_0 = 0$
5:         **for** $k = 0, \dots, K - 1$ **do**   # Iterate over integration steps.
6:             Let $\lambda_{K-k} := 1 - \sqrt{1 - \alpha_{K-k}}$
7:             $y_{k+1,n} = \sqrt{1 - \alpha_{K-k}} y_{k,n} + 2\sigma^2 \lambda_{K-k} f_\theta(K-k, y_{k,n}) + \sigma\sqrt{\alpha_{K-k}} \varepsilon_{k,n}, \ \varepsilon_{k,n} \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0, I)$
8:             $r_{k+1,n} = r_{k,n} + \frac{2\sigma^2 \lambda_{K-k}^2}{\alpha_{K-k}} \|f_\theta(K - k, y_{k,n})\|^2$
9:         **end for**
10:         $\theta \leftarrow \theta - \lambda \nabla_\theta \frac{1}{N} \sum_{n=1}^N \left(r_{K,n} + \ln\left(\frac{\mathcal{N}(y_{K,n}; 0, \sigma^2 I)}{\pi(y_{K,n})} \right) \right)$
11:     **end for**
12: **end for**
13: **Return:** f_θ

Using manipulations identical to Section 2.2, the time-reversal of $\mathcal{P}$ can be also be written for $\phi_t(x, p) := \eta_t(x, p)/\eta_t^{\text{ref}}(x, p)$ as $\mathrm{d}y_t = -M^{-1} q_t \mathrm{d}t$ and

$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t} q_t \mathrm{d}t + 2\beta_{T-t} \nabla_{q_t} \ln \phi_{T-t}(y_t, q_t)\mathrm{d}t + \sqrt{2\beta_{T-t}} M^{1/2} \mathrm{d}W_t. \qquad (15)$$

To approximate $\mathcal{P}$, we will consider a parameterized path measure $\mathcal{Q}^\theta$ whose time reversal is defined for $(y_0, q_0) \sim \mathcal{N}(y_0; 0, I)\mathcal{N}(q_0; 0, M)$ by $\mathrm{d}y_t = -M^{-1} q_t \mathrm{d}t$ and

$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t} q_t \mathrm{d}t + 2\beta_{T-t} M f_\theta(T - t, y_t, q_t)\mathrm{d}t + \sqrt{2\beta_{T-t}} M^{1/2} \mathrm{d}W_t. \qquad (16)$$

We can then express the KL of interest in a compact way similar to Proposition 1, as can be seen in Appendix B.1.1, where we also provide normalising flow based formulations.

3   Denoising Diffusion Samplers: Discrete-Time

We introduce here discrete-time integrators for the SDEs and ODEs introduced in Section 2. Contrary to DDPM, we not only need an integrator for $\mathcal{Q}_\theta$ but also for $\mathcal{P}^{\text{ref}}$ to be able to compute an approximation of the Radon-Nikodym derivative $\mathrm{d}\mathcal{Q}_\theta/\mathrm{d}\mathcal{P}^{\text{ref}}$. Additionally this integrator needs to be carefully designed as explained below. For sake of simplicity, we consider a constant discretization step $\delta > 0$ such that $K = T/\delta$ is an integer. In the indices, we write k for $t_k = k\delta$ to simplify notation; e.g. x_k for $x_{k\delta}$ and p_k for $p_{k\delta}$.

3.1   Integrator for $\mathcal{P}^{\text{ref}}$

Proposition 1 shows that learning the time reversal in continuous-time can be achieved by minimising the objective $\mathrm{KL}(\mathcal{Q}^\theta || \mathcal{P})$ given in (10). This expression is obtained by using the identity $\mathrm{KL}(\mathcal{Q}^\theta || \mathcal{P}) = \mathrm{KL}(\mathcal{Q}^\theta || \mathcal{P}^{\text{ref}}) + \mathbb{E}_{y_T \sim q_0^\theta}\left[\ln\left(p_0^{\text{ref}}(y_T)/p_0(y_T) \right) \right]$. This is also equivalent to maximizing the corresponding ELBO $\mathbb{E}_{\mathcal{Q}^\theta}[\ln \hat{Z}]$ for the unbiased estimate $\hat{Z}$ of Z given in (11)

An Euler–Maruyama (EM) discretisation of the corresponding SDEs is obviously applicable. However, it is problematic as established below.

Proposition 2. *Consider integrators of $\mathcal{P}^{\text{ref}}$ and $\mathcal{Q}^\theta$ leading to approximations $p^{\text{ref}}(x_{0:K})$ and $q^\theta(x_{0:K})$. Then $\mathrm{KL}(q^\theta || p^{\text{ref}}) + \mathbb{E}_{y_K \sim q_0^\theta}\left[\ln\left(p_0^{\text{ref}}(y_K)/\pi(y_K) \right) \right]$ is guaranteed to be non-negative and $\mathbb{E}_{\mathcal{Q}^\theta}[\ln \hat{Z}]$ is an ELBO for $\hat{Z} = (\gamma(y_K) p^{\text{ref}}(y_{0:K}))/(\mathcal{N}(y_K; 0, \sigma^2) q^\theta(y_{0:K}))$ if one has $p_K^{\text{ref}}(y_K) = p_0^{\text{ref}}(y_K)$. The EM discretisation of $\mathcal{P}^{\text{ref}}$ does* not *satisfy this condition.*

A direct consequence of this result is that the estimator for $\ln Z$ that uses the EM discretisation can be such that $\mathbb{E}_{\mathcal{Q}^\theta}[\ln \hat{Z}] \geq \ln Z$ as observed empirically in Table 5. To resolve this issue, we derive in the next section an integrator for $\mathcal{P}^{\text{ref}}$ which ensures that $p_k^{\text{ref}}(y) = p_0^{\text{ref}}(y)$ for all k.

3.2 ORNSTEIN–UHLENBECK

We can integrate exactly (4). This is given by $y_0 \sim \mathcal{N}(0, \sigma^2 I)$ and

$$y_{k+1} = \sqrt{1 - \alpha_{K-k}} y_k + \sigma \sqrt{\alpha_{K-k}} \varepsilon_k, \quad \varepsilon_k \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0, I), \tag{17}$$

for $\alpha_k = 1 - \exp\left(-2 \int_{(k-1)\delta}^{k\delta} \beta_s \mathrm{d}s\right)$ [2]. This defines the discrete-time reference process. We propose the following exponential type integrator (De Bortoli, 2022) for (8) initialized using $y_0 \sim \mathcal{N}(0, \sigma^2 I)$

$$y_{k+1} = \sqrt{1 - \alpha_{K-k}} y_k + 2\sigma^2 (1 - \sqrt{1 - \alpha_{K-k}}) f_\theta(K-k, y_k) + \sigma \sqrt{\alpha_{K-k}} \varepsilon_k, \tag{18}$$

where $\varepsilon_k \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0, I)$. Equations (17) and (18) define the time reversals of the reference process $p^{\text{ref}}(x_{0:K})$ and proposal $q^\theta(x_{0:K})$. For its time reversal, we write $q^\theta(y_{0:K}) = \mathcal{N}(y_0; 0, \sigma^2 I) \prod_{k=1}^{K} q^\theta_{k-1|k}(y_{K-k+1}|y_{K-k})$ abusing slightly notation, and similarly for $p^{\text{ref}}(y_{0:K})$. While we could adapt the KL decomposition developed in (Sohl-Dickstein et al., 2015; Ho et al., 2020) for DDPM to the reverse KL (see Appendix A.5), we found empirically that it is very unstable in our context. Instead, we will be relying on the discrete-time counterpart of (9).

Proposition 3. *The log density ratio between $q^\theta(y_{0:K})$ and $p^{\text{ref}}(y_{0:K})$ satisfies*

$$\ln\left(\frac{q^\theta(y_{0:K})}{p^{\text{ref}}(y_{0:K})}\right) = 2\sigma^2 \sum_{k=1}^{K} \frac{\lambda_k^2}{\alpha_k} ||f_\theta(k, y_{K-k})||^2 + 2\sigma \sum_{k=1}^{K} \frac{\lambda_k}{\sqrt{\alpha_k}} f_\theta(k, y_{K-k})^\top \varepsilon_k \tag{19}$$

where $\lambda_k := 1 - \sqrt{1 - \alpha_k}$ and ε_k is defined through (18). In particular for $y_{0:K} \sim q^\theta(y_{0:K})$, hence $\varepsilon_k \overset{\text{i.i.d.}}{\sim} \mathcal{N}(0, I)$, one obtains from $\mathrm{KL}(q^\theta||p) = \mathrm{KL}(q^\theta||p^{\text{ref}}) + \mathbb{E}_{q_0^\theta}\left[\ln\left(\frac{\pi(y_K)}{\mathcal{N}(y_K; 0, \sigma^2 I)}\right)\right]$ that

$$\mathrm{KL}(q^\theta||p) = \mathbb{E}_{q^\theta}\left[2\sigma^2 \sum_{k=1}^{K} \frac{\lambda_k^2}{\alpha_k} ||f_\theta(k, y_{K-k})||^2 + \ln\left(\frac{\mathcal{N}(y_K; 0, \sigma^2 I)}{\pi(y_K)}\right)\right]. \tag{20}$$

We compute an unbiased gradient of this objective using the reparameterization trick and the JAX software package (Bradbury et al., 2018). The training procedure is summarized in Algorithm 1 in Appendix D. Unfortunately, contrary to DDPM, $q^\theta_{k|K}$ is not available in closed form for $k < K - 1$ so we can neither mini-batches over the time index k without having to simulate the process until the minimum sampled time nor reparameterize x_k as in Ho et al. (2020). Once we obtain the parameter θ minimizing (20), DDS samples from q^θ using (18). The final sample y_K has a distribution q_0^θ approximating π by design. By using importance sampling, we obtain an unbiased estimate of the normalizing constant $\hat{Z} = (\gamma(y_K) p^{\text{ref}}(y_{0:K}))/(\mathcal{N}(y_K; 0, \sigma^2) q^\theta(y_{0:K}))$ for $y_{0:K} \sim q^\theta(y_{0:K})$. Finally Appendices B.3 and B.5 extend this approach to provide a similar sketch for the underdamped dynamics proposed in (2.5). In this context, the proposed integrators relies on a leapfrog scheme (Leimkuhler & Matthews, 2016).

3.3 THEORETICAL GUARANTEES

Motivated by DDPM, bounds on the total variation between the target distribution and the distribution of the samples generated by a time-discretization of an approximate time reversal of a forward noising diffusion have been first obtained in (De Bortoli et al., 2021b) then refined in (De Bortoli, 2022; Chen et al., 2022; Lee et al., 2022) and extended to the Wasserstein metric. These results are directly applicable to DDS because their proofs rely on assumptions on the score approximation error but not on the way these approximations are learned, i.e. via score matching for DDPM and reverse KL for DDS. For example, the main result in Chen et al. (2022) shows that if the true scores are L-Lipschitz, the $L_2(p_t)$ error on the scores is bounded and other mild integrability assumptions then DDS outputs samples ϵ-close in total variation to π in $O(L^2 d/\epsilon^2)$ time steps. As pointed out by the authors, this is truly remarkable as it matches state-of-the-art complexity bounds for Langevin Monte Carlo algorithm for sampling targets satisfying a log-Sobolev inequality. However, we emphasize that the assumption on the approximation error for score estimates is less realistic in the DDS setting than in the DDPM setting as we do not observe realizations of the forward diffusion.

---

[2]Henceforth, it is assumed that α_k can be computed exactly.

Figure 1: Training loss per hyperparameter: PIS (left) vs DDS (right).

Figure 2: $\ln Z$ estimate (median plus upper/lower quartiles) as a function of number of steps K - a) Funnel , b) LGCP, c) Logistic Ionosphere dataset. Yellow Dotted line is MF-VI and dashed magenta is the gold standard.

3.4 INTEPRETATION, RELATED WORK AND EXTENSIONS

DDS as KL and path integral control. The reverse KL we minimize can be expressed as

$$\text{KL}(q^\theta||p) = \mathbb{E}_{q^\theta}\left[\left\{\ln\left(\frac{\mathcal{N}(x_0;0,\sigma^2 I)}{\pi(x_0)}\right) + \sum_{k=1}^{K}\ln\left(\frac{q^\theta_{k-1|k}(x_{k-1}|x_k)}{p^{\text{ref}}_{k-1|k}(x_{k-1}|x_k)}\right)\right\}\right]. \quad (21)$$

This objective is a specific example of KL control problem (Kappen et al., 2012). In continuous-time, (10) corresponds to a path integral control problem; see e.g. (Kappen & Ruiz, 2016).

Heng et al. (2020) use KL control so as to sample from a target π and estimate Z. However, their algorithm relies on $p^{\text{ref}}(x_{0:K})$ being defined by a discretized non-homogeneous Langevin dynamics such that $p_K(x_K)$ is typically not approximating a known distribution. Additionally it approximates the value functions ϕ_k using regression using simple linear/quadratic functions. Finally it relies on a good initial estimate of $p(x_{0:K})$ obtained through SMC. This limits the applicability of this methodology to a restricted class of models.

Connections to Schrödinger Bridges. The Schrödinger Bridge problem (Léonard, 2014a) takes the following form in discrete time. Given a reference density $p^{\text{ref}}(x_{0:K})$, we want to find the density $p(x_{0:K})$ such that $p = \arg\min_q\left\{\text{KL}(q||p^{\text{ref}}) : q_0 = \mu_0, \ q_K = \mu_K\right\}$ where μ_0, μ_K are prescribed distributions. This problem can be solved using iterative proportional fitting (IPF) which is defined by the following recursion with initialization $p^0 = p^{\text{ref}}$

$$p^{2n+1} = \arg\min_q\left\{\text{KL}(q||p^{2n}) : q_K = \mu_K\right\}, \ p^{2n+2} = \arg\min_q\left\{\text{KL}(q||p^{2n+1}) : q_0 = \mu_0\right\} \quad (22)$$

For $\mu_0(x_0) = \pi(x_0)$ and $\mu_K(x_K) = \mathcal{N}(x_K;0,\sigma^2 I)$, IPF would converge in one iteration if we had $p^{\text{ref}}_K(x_K) = \mathcal{N}(x_K;0,\sigma^2 I)$. As $p^{\text{ref}}_K(x_K) \approx \mathcal{N}(x_K;0,\sigma^2 I)$, we can thus think of DDS as approximating the solution to this SB as pointed out by De Bortoli et al. (2021b).

For $q_0 = \pi$ and $q_K = \delta_0$, IPF also converges in one iteration for $p^{\text{ref}}_K(x_K) = \delta_0(x_K)$. Designing such a reference process is simple, set e.g. a pinned Brownian motion $p^{\text{ref}}_K(x_{0:K}) = \delta_0(x_K)\prod_{k=0}^{K-1}\mathcal{N}(x_k;x_{k+1},\delta\sigma^2 I)$. This was discussed in discrete-time in (Beghi, 1996) and in

Figure 3: $\ln Z$ estimate as a function of number of steps K - a) Logistic Sonar dataset, b) Brownian motion, c) NICE. Yellow dotted line is MF-VI and dashed magenta is the gold standard.

continuous-time in (Follmer, 1984; Dai Pra, 1991; Tzen & Raginsky, 2019). Algorithmically, Zhang et al. (2021) proposed approximating the solution by some projection on some eigenfunctions. In parallel, Barr et al. (2020), Vargas et al. (2021) and Zhang & Chen (2022) approximated this SB by using a neural network parameterization of the gradient of the logarithm of the corresponding value function trained by minimizing a reverse KL. In this context, an Euler–Maruyama discretization of $\mathcal{P}^{\text{ref}}$ preserves the marginal distributions of this process as required by Proposition 2 so no alternative integrator has to be developed. We will adopt the Path Integral Sampler (PIS) terminology proposed by Zhang & Chen (2022) for this approach. DDS can thus be seen as an alternative to PIS which relies on a reference dynamics corresponding to an overdamped or underdamped Ornstein–Uhlenbeck process instead of relying on a reference dynamics corresponding to a pinned Brownian motion. Empirically this significantly improves numerical stability of the training procedure; see Figure 1. The use of a pinned Brownian motion is also not amenable to the construction of normalizing flows.

4 EXPERIMENTS

In our implementation, f_θ follows the PIS-GRAD network proposed in (Zhang & Chen, 2022): $f_\theta(k, x) = \text{NN}_1(k, x; \theta) + \text{NN}_2(k; \theta) \circ \nabla \ln \pi(x)$. Across all experiments we use a two layer architecture with 64 hidden units each (for both networks), as in Zhang & Chen (2022), with the exception of the NICE (Dinh et al., 2014) target where we use 3 layers with 512, 256, and 64 units respectively. The final layers are initialised to 0 in order to make the path regularisation term null. We use $\alpha_k^{1/2} \propto \alpha_{\max}^{1/2} \cos^2\left(\frac{\pi}{2} \frac{1-k/K+s}{1+s}\right)$ with $s=0.008$ as in (Nichol & Dhariwal, 2021). We found that detaching the target score stabilised optimization in both approaches without affecting the final result. We adopt this across experiments, an ablation of this feature can be seen in Appendix C.8.1.

Across all tasks we compare DDS to SMC (Del Moral et al., 2006; Zhou et al., 2016), PIS (Barr et al., 2020; Vargas et al., 2021; Zhang & Chen, 2022), and Mean Field-VI (MF-VI) with a Gaussian variational distribution. We also compare DDS to AIS (Neal, 2001) and optimized variants of AIS using score matching (MCD) (Doucet et al., 2022; Geffner & Domke, 2022) for two standard Bayesian models. Finally we explore a task introduced in (Doucet et al., 2022) that uses a pre-trained normalising flow as a target. Within this setting we propose a benchmarking criterion that allows us to assess mode collapse in high dimensions and explore the benefits of incorporating inductive biases into f_θ. We carefully tuned the hyper-parameters of all algorithms (e.g. step size, diffusion coefficient, and such), details can be found in Appendix C.1. Experiments for the underdamped approach are presented in Appendix C.10. Finally training time can be found in Appendix C.4

4.1 BENCHMARKING TARGETS

We first discuss two standard target distributions which are often used to benchmark methods; see e.g. (Neal, 2003; Arbel et al., 2021; Heng et al., 2020; Zhang & Chen, 2022). Results are presented in Figure 2.

Funnel Distribution: This 10-dimensional challenging distribution is given by $\gamma(x_{0:9}) = \mathcal{N}(x_0; 0, \sigma_f^2)\mathcal{N}(0, \exp(x_0)I)$, where $\sigma_f^2 = 9$ (Neal, 2003). [3].

---

[3]Zhang & Chen (2022) erroneously considered the more favourable $\sigma_f = 1$ scenario for PIS but used $\sigma_f = 3$ for other methods. This explains the significant differences between their results and ours.

Log Gaussian Cox process: This model arises in spatial statistics (Møller et al., 1998). We use a $d = M \times M = 1600$ grid, resulting in the unnormalized target density $\gamma(x) = \mathcal{N}(x; \mu, K) \prod_{i \in [1:M]^2} \exp(x_i y_i - a \exp(x_i))$.

4.2 BAYESIAN MODELS

We explore two standard Bayesian models and compare them with standard AIS and MCD benchmarks (See Appendix C.7) in addition to the SMC and VI benchmarks presented so far. Results for Ionosphere can be found in the 3rd pane of Figure 2 whilst SONAR and Brownian are in Figure 3.

Logistic Regression: We set $x \sim \mathcal{N}(0, \sigma_w^2 I), y_i \sim \text{Bernoulli}(\text{sigmoid}(x^\top u_i))$. This Bayesian logistic model is evaluated on two datasets, Ionosphere ($d = 32$) and Sonar ($d = 61$).

Brownian Motion: We consider a discretised Brownian motion with a Gaussian observation model and a latent volatility as a time series model, $d = 32$. This model was obtained from the Sountsov et al. (2020) software package and is specified in more detail in Appendix C.2.

4.3 MODE COLLAPSE IN HIGH DIMENSIONS

Normalizing Flow Evaluation: Following Doucet et al. (2022) we train NICE (Dinh et al., 2014) on a down-sampled $d = 14 \times 14$ variant of MNIST (LeCun & Cortes, 2010) and use the trained model as our target. As we can generate samples from our target we chose to evaluate the methods samples by measuring the Sinkhorn distance between true and estimated samples. This constitutes of a new evaluation criteria for this benchmark which allows to asses mode collapse for samplers in high dimensional settings. Results can be seen in Table 1 and the third pane of Figure 3.

Method	DDS	PIS	SMC	MCD
$\ln Z$ Estimate	-3.204 ± 0.645	-3.933 ± 0.754	-4.255 ± 2.043	-6.25
$\mathcal{W}^2_{\gamma=0.01}(\pi_{\text{true}}, \hat{\pi})$	658.079	658.778	750.245	NA

Table 1: Results on NICE target, we performed 30 runs with different seeds for each approach. For MCD we used the results from (Doucet et al., 2022).

5 DISCUSSION

We have explored the use of DDPM ideas to develop algorithms to sample unnormalized probability distributions and estimate their normalizing constants. We have compared experimentally the resulting Denoising Diffusion Samplers to state-of-the-art Monte Carlo methods. We found overall that DDS are competitive with SMC and numerically more stable than PIS. However, this comes at the cost of a non-negligible training time compared to SMC. When accounting for this training time, SMC often provide better performance on simple target distributions. However, in the challenging multimodal NICE example, even a carefully tuned SMC sampler using HMC transitions was not competitive to DDS. This is despite the fact that DDS (and PIS) is prone to mode dropping as any method relying on the reverse KL.

Overall, we believe that DDS are a class of algorithms worth considering and further developing. Contrary to SMC methods (Maddison et al., 2017; Le et al., 2018; Naesseth et al., 2018), it provides an objective that can be easily optimized using the reparameterization trick. Additionally, for high-dimensional applications, SMC suffer from high memory requirements as it relies on a large collection of interacting particles. DDS does not suffer from such limitations and, once it has been trained, it allows easy parallel simulation of multiple samples. Moreoever, there has much work devoted to improving successfully DDPM over the past two years including, among many others, modified forward noising mechanisms (Hoogeboom & Salimans, 2022), denoising diffusion implicit models (Song et al., 2020) and sophisticated numerical integrators (Karras et al., 2022). It is highly likely that some of these techniques will lead to more powerful DDS. Advances in KL and path integral control might also be adapted to DDS to provide better training procedures; see e.g. Thalmeier et al. (2020).

REFERENCES

Felix V Agakov and David Barber. An auxiliary variational method. In *Advances in Neural Information Processing Systems*, 2004.

Michael Arbel, Alex Matthews, and Arnaud Doucet. Annealed flow transport Monte Carlo. In *International Conference on Machine Learning*, 2021.

Ariel Barr, Willem Gispen, and Austen Lamacraft. Quantum ground states from reinforcement learning. In *Mathematical and Scientific Machine Learning*, pp. 635–653. PMLR, 2020.

Alessandro Beghi. On the relative entropy of discrete-time Markov processes with given end-point densities. *IEEE Transactions on Information Theory*, 42(5):1529–1535, 1996.

James Bradbury, Roy Frostig, Peter Hawkins, Matthew James Johnson, Chris Leary, Dougal Maclaurin, George Necula, Adam Paszke, Jake VanderPlas, Skye Wanderman-Milne, and Qiao Zhang. JAX: composable transformations of Python+NumPy programs, 2018. URL http://github.com/google/jax.

Ricky TQ Chen, Yulia Rubanova, Jesse Bettencourt, and David K Duvenaud. Neural ordinary differential equations. *Advances in Neural Information Processing Systems*, 31, 2018.

Sitan Chen, Sinho Chewi, Jerry Li, Yuanzhi Li, Adil Salim, and Anru R. Zhang. Sampling is as easy as learning the score: theory for diffusion models with minimal data assumptions. *arXiv preprint arXiv:22209.11215*, 2022.

Paolo Dai Pra. A stochastic control approach to reciprocal diffusion processes. *Applied Mathematics and Optimization*, 23(1):313–329, 1991.

Valentin De Bortoli. Convergence of denoising diffusion models under the manifold hypothesis. *arXiv preprint arXiv:2208.05314*, 2022.

Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. Diffusion schrödinger bridge with applications to score-based generative modeling. In *Advances in Neural Information Processing Systems*, 2021a.

Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. Diffusion Schrödinger bridge with applications to score-based generative modeling. In *Advances in Neural Information Processing Systems*, 2021b.

Pierre Del Moral, Arnaud Doucet, and Ajay Jasra. Sequential Monte Carlo samplers. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*, 68(3):411–436, 2006.

Laurent Dinh, David Krueger, and Yoshua Bengio. Nice: Non-linear independent components estimation. *arXiv preprint arXiv:1410.8516*, 2014.

Tim Dockhorn, Arash Vahdat, and Karsten Kreis. Score-based generative modeling with critically-damped Langevin diffusion. In *International Conference on Learning Representations*, 2022.

Arnaud Doucet, Will Sussman Grathwohl, Alexander G de G Matthews, and Heiko Strathmann. Score-based diffusion meets annealed importance sampling. In *Advances in Neural Information Processing Systems*, 2022.

H. Follmer. An entropy approach to the time reversal of diffusion processes. *Lecture Notes in Control and Information Sciences*, 69:156–163, 1984.

Tomas Geffner and Justin Domke. MCMC variational inference via uncorrected Hamiltonian annealing. In *Advances in Neural Information Processing Systems*, 2021.

Tomas Geffner and Justin Domke. Langevin diffusion variational inference. *arXiv preprint arXiv:2208.07743*, 2022.

Will Grathwohl, Ricky TQ Chen, Jesse Bettencourt, Ilya Sutskever, and David Duvenaud. Ffjord: Free-form continuous dynamics for scalable reversible generative models. In *International Conference on Learning Representations*, 2018.

Klaus Greff, Raphaël Lopez Kaufman, Rishabh Kabra, Nick Watters, Christopher Burgess, Daniel Zoran, Loic Matthey, Matthew Botvinick, and Alexander Lerchner. Multi-object representation learning with iterative variational inference. In *International Conference on Machine Learning*, pp. 2424–2433. PMLR, 2019.

Ulrich G Haussmann and Etienne Pardoux. Time reversal of diffusions. *The Annals of Probability*, 14(3):1188–1205, 1986.

Jeremy Heng, Adrian N Bishop, George Deligiannidis, and Arnaud Doucet. Controlled sequential Monte Carlo. *The Annals of Statistics*, 48(5):2904–2929, 2020.

Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. In *Advances in Neural Information Processing Systems*, 2020.

Emiel Hoogeboom and Tim Salimans. Blurring diffusion models. *arXiv preprint arXiv:2209.05557*, 2022.

Aapo Hyvärinen. Estimation of non-normalized statistical models by score matching. *The Journal of Machine Learning Research*, 6:695–709, 2005.

Hilbert J Kappen, Vicenç Gómez, and Manfred Opper. Optimal control as a graphical model inference problem. *Machine learning*, 87(2):159–182, 2012.

Hilbert Johan Kappen and Hans Christian Ruiz. Adaptive importance sampling for control and inference. *Journal of Statistical Physics*, 162(5):1244–1266, 2016.

Tero Karras, Miika Aittala, Timo Aila, and Samuli Laine. Elucidating the design space of diffusion-based generative models. *arXiv preprint arXiv:2206.00364*, 2022.

Diederik P. Kingma and Jimmy Ba. Adam: A method for stochastic optimization. In *International Conference on Learning Representations*, 2015.

Diederik P Kingma and Max Welling. Auto-encoding variational Bayes. *arXiv preprint arXiv:1312.6114*, 2013.

Fima C Klebaner. *Introduction to Stochastic Calculus with Applications*. Imperial College Press, 2012.

Tuan Anh Le, Maximilian Igl, Tom Rainforth, Tom Jin, and Frank Wood. Auto-encoding sequential Monte Carlo. In *International Conference on Learning Representations*, 2018.

Yann LeCun and Corinna Cortes. MNIST handwritten digit database. http://yann.lecun.com/exdb/mnist/, 2010. URL http://yann.lecun.com/exdb/mnist/.

Yann LeCun, Léon Bottou, Yoshua Bengio, and Patrick Haffner. Gradient-based learning applied to document recognition. *Proceedings of the IEEE*, 86(11):2278–2324, 1998.

Holden Lee, Jianfeng Lu, and Yixin Tan. Convergence of score-based generative modeling for general data distributions. *arXiv preprint arXiv:2209.12381*, 2022.

Ben Leimkuhler and Charles Matthews. *Molecular Dynamics with Deterministic and Stochastic Numerical Methods*. Springer, 2016.

Christian Léonard. A survey of the Schrödinger problem and some of its connections with optimal transport. *Discrete and Continuous Dynamical Systems-Series A*, 34(4):1533–1574, 2014a.

Christian Léonard. Some properties of path measures. In *Séminaire de Probabilités XLVI*, pp. 207–230. Springer, 2014b.

Chris J Maddison, John Lawson, George Tucker, Nicolas Heess, Mohammad Norouzi, Andriy Mnih, Arnaud Doucet, and Yee Teh. Filtering variational objectives. In *Advances in Neural Information Processing Systems*, 2017.

Luke Metz, Niru Maheswaranathan, Ruoxi Sun, C Daniel Freeman, Ben Poole, and Jascha Sohl-Dickstein. Using a thousand optimization tasks to learn hyperparameter search strategies. *arXiv preprint arXiv:2002.11887*, 2020.

Jesper Møller, Anne Randi Syversveen, and Rasmus Plenge Waagepetersen. Log Gaussian Cox processes. *Scandinavian Journal of Statistics*, 25(3):451–482, 1998. ISSN 03036898, 14679469.

Christian A Naesseth, Scott W Linderman, Rajesh Ranganath, and David M Blei. Variational sequential Monte Carlo. In *Artificial Intelligence and Statistics*, 2018.

Radford M Neal. Annealed importance sampling. *Statistics and Computing*, 11(2):125–139, 2001.

Radford M. Neal. Slice sampling. *The Annals of Statistics*, 31:705–767, 06 2003.

Alexander Quinn Nichol and Prafulla Dhariwal. Improved denoising diffusion probabilistic models. In *International Conference on Machine Learning*, pp. 8162–8171. PMLR, 2021.

George Papamakarios, Eric Nalisnick, Danilo Jimenez Rezende, Shakir Mohamed, and Balaji Lakshminarayanan. Normalizing flows for probabilistic modeling and inference. *arXiv preprint arXiv:1912.02762*, 2019.

Paavo Parmas, Carl Edward Rasmussen, Jan Peters, and Kenji Doya. Pipps: Flexible model-based policy search robust to the curse of chaos. In *International Conference on Machine Learning*, pp. 4065–4074. PMLR, 2018.

Jascha Sohl-Dickstein, Eric Weiss, Niru Maheswaranathan, and Surya Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In *International Conference on Machine Learning*, 2015.

Jiaming Song, Chenlin Meng, and Stefano Ermon. Denoising diffusion implicit models. *arXiv preprint arXiv:2010.02502*, 2020.

Yang Song, Conor Durkan, Iain Murray, and Stefano Ermon. Maximum likelihood training of score-based diffusion models. In *Advances in Neural Information Processing Systems*, 2021a.

Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2021b.

Pavel Sountsov, Alexey Radul, and contributors. Inference gym, 2020. URL `https://pypi.org/project/inference_gym`.

Dominik Thalmeier, Hilbert J Kappen, Simone Totaro, and Vicenç Gómez. Adaptive smoothing for path integral control. 2020.

Achille Thin, Nikita Kotelevskii, Alain Durmus, Eric Moulines, Maxim Panov, and Arnaud Doucet. Monte Carlo variational auto-encoders. In *International Conference on Machine Learning*, 2021.

Belinda Tzen and Maxim Raginsky. Theoretical guarantees for sampling and inference in generative models with latent diffusions. In *Conference on Learning Theory*, pp. 3084–3114. PMLR, 2019.

Francisco Vargas, Andrius Ovsianas, David Fernandes, Mark Girolami, Neil Lawrence, and Nikolas Nüsken. Bayesian learning via neural Schrödinger-Föllmer flows. *arXiv preprint arXiv:2111.10510*, 2021.

Paul Vicol, Luke Metz, and Jascha Sohl-Dickstein. Unbiased gradient estimation in unrolled computation graphs with persistent evolution strategies. In *International Conference on Machine Learning*, pp. 10553–10563. PMLR, 2021.

Pascal Vincent. A connection between score matching and denoising autoencoders. *Neural Computation*, 23(7):1661–1674, 2011.

Martin J. Wainwright and Michael I. Jordan. Graphical Models, Exponential Families, and Variational Inference. *Foundations and Trends® in Machine Learning*, 1(1–2):1–305, November 2008. URL `https://www.nowpublishers.com/article/Details/MAL-001`.

Hao Wu, Jonas Köhler, and Frank Noé. Stochastic normalizing flows. In *Advances in Neural Information Processing Systems*, 2020.

Benjamin Zhang, Tuhin Sahai, and Youssef Marzouk. Sampling via controlled stochastic dynamical systems. In *I (Still) Can't Believe It's Not Better! NeurIPS 2021 Workshop*, 2021.

Guodong Zhang, Kyle Hsu, Jianing Li, Chelsea Finn, and Roger Grosse. Differentiable annealed importance sampling and the perils of gradient noise. In *Advances in Neural Information Processing Systems*, 2021.

Qinsheng Zhang and Yongxin Chen. Path integral sampler: a stochastic control approach for sampling. In *International Conference on Learning Representations*, 2022.

Yan Zhou, Adam M Johansen, and John AD Aston. Toward automatic model comparison: an adaptive sequential monte carlo approach. *Journal of Computational and Graphical Statistics*, 25 (3):701–726, 2016.

A APPENDIX

A.1 PROOF OF PROPOSITION 1

The Radon-Nikodym derivative expression (9) directly follows from an application of Girsanov theorem (see e.g. Klebaner (2012)). The path measures $\mathcal{P}$ and $\mathcal{P}^{\text{ref}}$ are induced by two diffusions following the same dynamics but with different initial conditions so that

$$
\begin{aligned}
\mathcal{P}(\omega) &= \mathcal{P}(\omega|y_T)\pi(y_T) \\
&= \mathcal{P}^{\text{ref}}(\omega|y_T)\pi(y_T) \\
&= \mathcal{P}^{\text{ref}}(\omega|y_T)\mathcal{N}(y_T; 0, \sigma^2 I)\frac{\pi(y_T)}{\mathcal{N}(y_T; 0, \sigma^2 I)} \\
&= \mathcal{P}^{\text{ref}}(\omega)\frac{\pi(y_T)}{\mathcal{N}(y_T; 0, \sigma^2 I)}.
\end{aligned}
$$

So it follows directly that $\mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}) = \mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}^{\text{ref}}) + \mathbb{E}_{y_T \sim q_0^\theta}\left[\ln\left(\frac{p_0^{\text{ref}}(y_T)}{p_0(y_T)}\right)\right]$. As W_t is a Brownian motion under $\mathcal{Q}^\theta$, the final expression (10) for $\mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P})$ follows.

A.2 COMPARING DDS AND PIS DRIFTS

The optimal drift for both PIS and DDS can be expressed as a logarithmic derivative of the value function:

$$
b_{\text{DDS}}(x,t) = -\beta_{T-t}(x - 2\sigma^2\nabla_x\ln\phi(x,t)), \quad b_{\text{PIS}} = \sigma^2\nabla_x\ln\phi(x,t) \tag{23}
$$

where ϕ is the corresponding value for DDS or PIS respectively. In particular for PIS $\ln\phi$ is given by:

$$
\ln\phi_{\text{PIS}}(x,t) = \ln\mathbb{E}_{Z\sim\mathcal{N}(0,I)}\left[\frac{d\pi}{\mathcal{N}(0,\sigma^2 TI)}(x + \sqrt{T-t}Z)\right], \tag{24}
$$

whilst for DDS we have a very similar expression

$$
\ln\phi_{\text{DDS}}(x,t) = \ln\mathbb{E}_{Z\sim\mathcal{N}(0,I)}\left[\frac{d\pi}{\mathcal{N}(0,\sigma^2 I)}\left(e^{-\int_0^{T-t}\beta_s ds}x + \left(1 - e^{-2\int_0^{T-t}\beta_s ds}\right)^{1/2}Z\right)\right], \tag{25}
$$

we can see now that both take the general form,

$$
\ln\mathbb{E}_{Z\sim\mathcal{N}(0,I)}\left[\frac{d\pi}{\mathcal{N}(0,cI)}(ax + \sqrt{b}Z)\right]. \tag{26}
$$

Thus in this section we will obtain analytic formulations of the above expectation which immediately provides us with $\nabla_x\ln\phi(x,t)$ for both DDS and PIS and subsequently its optimal drifts.

Lemma 1. *Let* $f(x) = \frac{\mathcal{N}(x;\alpha,\Sigma)}{\mathcal{N}(x;0,cI)}$ *then it follows that:*

$$\nabla_{\tilde{x}} \ln \mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[f(a\tilde{x} + \sqrt{b}Z) \right]$$

$$= a \left(\Sigma^{-1}\alpha + (c^{-1}I - \Sigma^{-1}) \left(\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1} (b\Sigma^{-1}\alpha - a\tilde{x}) \right) \right)$$

Proof. Firstly notice:

$$\nabla_{\tilde{x}} \ln \mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[f(ax + \sqrt{b}Z) \right] = \frac{\mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[\nabla_{\tilde{x}} f(a\tilde{x} + \sqrt{b}Z) \right]}{\mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[f(a\tilde{x} + \sqrt{b}Z) \right]}$$

$$= a \frac{\mathbb{E}_{Y \sim \mathcal{N}(a\tilde{x},b)} \left[\nabla f(Y) \right]}{\mathbb{E}_{Y \sim \mathcal{N}(a\tilde{x},b)} \left[f(Y) \right]}.$$

For brevity $x = a\tilde{x}$

$$\mathbb{E}_{Y \sim \mathcal{N}(x,b)} \left[f(Y) \right] = \frac{c^{d/2}}{|\Sigma|^{1/2}} \mathbb{E}_{Y \sim \mathcal{N}(x,b)} \left[e^{\frac{1}{2}\left(c^{-1}||Y||^2 - ||Y - \alpha||^2_{\Sigma^{-1}}\right)} \right]$$

$$= \frac{c^{d/2}}{(2\pi b)^{d/2}|\Sigma|^{1/2}} \int \left[e^{\frac{1}{2}\left(c^{-1}||y||^2 - ||y - \alpha||^2_{\Sigma^{-1}} - b^{-1}||y - x||^2\right)} \right] dy$$

Let $C_1 = \left(\frac{c^{d/2}}{(2\pi b)^{d/2}|\Sigma|^{1/2}}\right)^{-1}$ and $C_2 = e^{-\frac{1}{2cb}(||x||^2 + cb||\alpha||^2_{\Sigma^{-1}})}$.

$$= \frac{1}{C_1} \int \left[e^{\frac{1}{2cb}\left(b||y||^2 - cb||y||^2_{\Sigma^{-1}} + 2cby^\top\Sigma^{-1}\alpha - cb||\alpha||^2_{\Sigma^{-1}} - c(||y||^2 - 2y^\top x + ||x||^2)\right)} \right] dy$$

$$= \frac{e^{-\frac{1}{2cb}(c||x||^2 + cb||\alpha||^2_{\Sigma^{-1}})}}{C_1} \int \left[e^{\frac{1}{2cb}\left(b||y||^2 - cb||y||^2_{\Sigma^{-1}} + 2cby^\top\Sigma^{-1}\alpha - c(||y||^2 - 2y^\top x)\right)} \right] dy$$

$$= \frac{C_2}{C_1} \int \left[e^{\frac{1}{2b}\left(y^\top\left(\left(\frac{b}{c} - 1\right)I - b\Sigma^{-1}\right)y + 2y^\top\left(b\Sigma^{-1}\alpha - x\right)\right)} \right] dy$$

$$= \frac{C_2 e^{\frac{1}{2}||\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1/2}(b\Sigma^{-1}\alpha - x)||^2}}{C_1} \int \left[e^{-\frac{1}{2b}\left(||y - \left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1/2}(b\Sigma^{-1}\alpha - x)||^2_{\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)}\right)} \right] dy$$

$$= \frac{c^{d/2}C_2 e^{\frac{1}{2b}||\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1/2}(b\Sigma^{-1}\alpha - x)||^2}}{|\Sigma|^{1/2}|\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}|^{1/2}} = \frac{c^{d/2}e^{\frac{1}{2b}\left(||\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1/2}(b\Sigma^{-1}\alpha - x)||^2 - ||x||^2 - b||\alpha||^2_{\Sigma^{-1}}\right)}}{|\left(1 - \frac{b}{c}\right)\Sigma + b|^{1/2}}$$

Now let $g(x,t) = e^{\frac{1}{2b}\left(||\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1/2}(b\Sigma^{-1}\alpha - x)||^2 - ||x||^2 - b||\alpha||^2_{\Sigma^{-1}}\right)}$ (where b will depend on t).
Then we wish to compute:

$$\mathbb{E}_{Y \sim \mathcal{N}(x,b)} \left[\nabla f(Y) \right] = \Sigma^{-1}\alpha \mathbb{E}_{Y \sim \mathcal{N}(x,b)} \left[f(Y) \right] + (c^{-1}I - \Sigma^{-1})\mathbb{E}_{Y \sim \mathcal{N}(x,b)} \left[Y f(Y) \right]$$

$$= \frac{\Sigma^{-1}\alpha c^{d/2}g(x,t)}{|\left(1 - \frac{b}{c}\right)\Sigma + b|^{1/2}} + \frac{(c^{-1}I - \Sigma^{-1})\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1}(b\Sigma^{-1}\alpha - x)c^{d/2}g(x,t)}{|\left(1 - \frac{b}{c}\right)\Sigma + b|^{1/2}}$$

combining both results yields:

$$\nabla_{\tilde{x}} \ln \mathbb{E}_{Y \sim \mathcal{N}(a\tilde{x},b)} \left[f(Y) \right]$$

$$= a \left(\Sigma^{-1}\alpha + (c^{-1}I - \Sigma^{-1}) \left(\left(\left(1 - \frac{b}{c}\right)I + b\Sigma^{-1}\right)^{-1} (b\Sigma^{-1}\alpha - a\tilde{x}) \right) \right)$$

$\square$

Corollary 1. *For DDS consider an OU reference process of the form:*

$$dx_t = -x_t dt + \sqrt{2}dw_t, \quad x_0 \sim \mathcal{N}(0,1) \tag{27}$$

Then the optimal half bridge drift to the target $\mathcal{N}(\alpha, 1)$ *is given by:*

$$b(x,t) = -x + 2e^{t-T}\alpha \tag{28}$$

Proof. Here we set $a = e^{-(T-t)}$, $b = 1 - e^{-2(T-t)}$, $c = 1$ and $\Sigma = 1$. and thus we have :

$$\nabla_x \ln \phi(x,t) = \nabla_{\tilde{x}} \ln \mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[f(ax + \sqrt{b}Z) \right] = e^{t-T}\alpha \tag{29}$$

$\square$

Corollary 2. *For PIS consider a pinned Brownian motion reference process of the form:*

$$\mathrm{d}x_t = dw_t, \;\; x_0 \sim \delta_0 \tag{30}$$

Then the optimal half bridge drift to the target $\mathcal{N}(\alpha, 1)$ is given by:

$$b(x,t) = \frac{T(\alpha + x) - x}{T(T-t) + t} \tag{31}$$

Proof. Here we set $a = 1$, $b = T - t$, $c = T$ and $\Sigma = 1$. and thus after simplifying we have :

$$\nabla_x \ln \phi(x,t) = \nabla_{\tilde{x}} \ln \mathbb{E}_{Z \sim \mathcal{N}(0,1)} \left[f(\tilde{x} + \sqrt{1-t}Z) \right]$$
$$= \frac{T\alpha}{T(T-t)+t} - \frac{(1-T)x}{T(T-t)+t} = \frac{T(\alpha+x)-x}{T(T-t)+t}$$

$\square$

From Corollaries 1 and 2 we can observe that the DDS drifts magnitude scales down the target mean via e^{-T} which is much smaller than PIS whose scaling is $\mathcal{O}(\frac{1}{T})$, this suggests that the DDS drifts diffuse more smoothly and stably towards the target distribution as we have seen empirically whilst the PIS drift can take on larger values resulting on difficulties at train time.

A.3 PROOF OF PROPOSITION 2

Proof. Let us express the discrete time version of reference process as

$$p^{\mathrm{ref}}(y_{0:K}) = p_0^{\mathrm{ref}}(y_0) \prod_{k=0}^{K-1} p_{k+1|k}^{\mathrm{ref}}(y_{k+1}|y_k) \tag{32}$$

and denote by p_k^{ref} the corresponding marginal density of y_k which satisfies

$$p_{k+1}(y_{k+1}) = \int p_{k+1|k}^{\mathrm{ref}}(y_{k+1}|y_k) \; p_k(y_k) \mathrm{d}y_k. \tag{33}$$

The backward decomposition of this joint distribution is given by

$$p^{\mathrm{ref}}(y_{0:K}) = p_K^{\mathrm{ref}}(y_K) \prod_{k=0}^{K-1} p_{k|k+1}^{\mathrm{ref}}(y_k|y_{k+1}), \tag{34}$$

where

$$p_{k|k+1}^{\mathrm{ref}}(y_k|y_{k+1}) = \frac{p_{k+1|k}^{\mathrm{ref}}(y_{k+1}|y_k)p_k(y_k)}{p_{k+1}(y_{k+1})}. \tag{35}$$

If our chosen integrator induces a transition kernel $p_{k+1|k}^{\mathrm{ref}}(y_{k+1}|y_k)$ which is such that $p_K^{\mathrm{ref}}(y_K) = p_0^{\mathrm{ref}}(y_K)$, then

$$p^{\mathrm{ref}}(y_{0:K}) \frac{\pi(y_K)}{p_0^{\mathrm{ref}}(y_K)} = \pi(y_K) \prod_{k=0}^{K-1} p_{k|k+1}^{\mathrm{ref}}(y_k|y_{k+1}) \tag{36}$$

is a valid (normalised) probability density. Hence it follows that

$$\text{KL}(q^\theta(y_{0:K})||p^{\text{ref}}(y_{0:K})) + \mathbb{E}_{y_K \sim q_0^\theta}\left[\ln\left(\frac{p_0^{\text{ref}}(y_K)}{\pi(y_K)}\right)\right] \tag{37}$$

$$=\text{KL}\left(q^\theta(y_{0:K})\middle|\middle|p^{\text{ref}}(y_{0:K})\frac{\pi(y_K)}{p_0^{\text{ref}}(y_K)}\right) \geq 0.$$

If the integrator does not preserve the marginals we have that

$$p^{\text{ref}}(y_{0:K})\frac{\pi(y_K)}{p_0^{\text{ref}}(y_K)} = \frac{p_K^{\text{ref}}(y_K)}{p_0^{\text{ref}}(y_K)}\pi(y_K)\prod_{k=0}^{K-1}p_{k|k+1}^{\text{ref}}(y_k|y_{k+1}). \tag{38}$$

This is not a probability density and thus the objective is no longer guaranteed to be positive and as a result the expectation of our estimator of $\ln Z$ is not a guaranteed lower bound for $\ln Z$. Finally a simple calculation shows that the Euler discretisation does not preserve the invariant distribution of $\mathcal{P}^{\text{ref}}$ in DDS thus making our objective no longer an ELBO in discrete time. $\qquad\square$

A.4 PROOF OF PROPOSITION 3

We have

$$\ln\left(\frac{q^\theta(y_{0:K})}{p(y_{0:K})}\right) = \ln\left(\frac{q^\theta(y_{0:K})}{p^{\text{ref}}(y_{0:K})}\right) + \ln\left(\frac{\mathcal{N}(y_0; 0, \sigma^2 I)}{\pi(y_0)}\right). \tag{39}$$

Now by construction, we have from (17) that $p_{k-1|k}^{\text{ref}}(y_{K-k+1}|y_{K-k}) = \mathcal{N}(y_{K-k+1}; \sqrt{1-\alpha_k}y_{K-k}, \sigma^2\alpha_k I)$ and from (18) we obtain $q_{k-1|k}^\theta(y_{K-k+1}|y_{K-k}) = \mathcal{N}(y_{K-k+1}; \sqrt{1-\alpha_k}y_{K-k} + 2\sigma^2(1-\sqrt{1-\alpha_k})f_\theta(k, y_{K-k}), \sigma^2\alpha_k I)$.

It follows that

$$\ln\left(\frac{q^\theta(y_{0:K})}{p^{\text{ref}}(y_{0:K})}\right) \tag{40}$$

$$=\ln\left(\frac{q_K(y_0)}{p_K^{\text{ref}}(y_0)}\right) + \sum_{k=1}^{K}\ln\left(\frac{q_{k-1|k}^\theta(y_{K-k+1}|y_{K-k})}{p_{k-1|k}^{\text{ref}}(y_{K-k+1}|y_{K-k})}\right)$$

$$=\sum_{k=1}^{K}\frac{1}{2\alpha_k\sigma^2}\Big[||y_{K-k+1} - \sqrt{1-\alpha_k}y_{K-k}||^2$$

$$- ||y_{K-k+1} - \sqrt{1-\alpha_k}y_{K-k} - 2\sigma^2(1-\sqrt{1-\alpha_k})f(k, y_{K-k})||^2\Big],$$

where we have exploited the fact that $q_K(y_0) = p_K^{\text{ref}}(y_0) = \mathcal{N}(y_0; 0, \sigma^2 I)$.

Now using $\epsilon_k := \frac{1}{\sigma\sqrt{\alpha_k}}(y_{K-k+1} - \sqrt{1-\alpha_k}y_{K-k} - 2\sigma^2(1-\sqrt{1-\alpha_k})f(k, y_{K-k}))$, we can rewrite (40) as

$$\ln\left(\frac{q^\theta(y_{0:K})}{p^{\text{ref}}(y_{0:K})}\right) \tag{41}$$

$$=\frac{1}{2}\sum_{k=1}^{K}\Big[||\epsilon_k + 2\sigma\frac{(1-\sqrt{1-\alpha_k})}{\sqrt{\alpha_k}}f_\theta(k, y_{K-k})||^2 - ||\epsilon_k||^2\Big]$$

$$=2\sigma^2\sum_{k=1}^{K}\frac{(1-\sqrt{1-\alpha_k})^2}{\alpha_k}||f_\theta(k, y_{K-k})||^2 + 2\sigma\sum_{k=1}^{K}\frac{(1-\sqrt{1-\alpha_k})}{\sqrt{\alpha_k}}f_\theta(k, y_{K-k})^\top\epsilon_k.$$

Now (19) follows directly from (39) and (72). Note we have for $\delta \ll 1$ that $2\sigma^2\frac{(1-\sqrt{1-\alpha_k})^2}{\alpha_k} \approx \sigma^2\beta_k\delta$ and $2\sigma\frac{(1-\sqrt{1-\alpha_k})}{\sqrt{\alpha_k}} \approx \sqrt{2\beta_k\delta\delta}$ as expected from (9).

Finally the final expression (20) of the KL follows now from the fact that $\mathbb{E}_{q^\theta}[f_\theta(k, y_{K-k})^\top\epsilon_k] = \mathbb{E}_{q_k^\theta}[\mathbb{E}_{q_{k-1|k}^\theta}[f_\theta(k, y_{K-k})^\top\epsilon_k|y_{K-k}]] = \mathbb{E}_{q_k^\theta}[f_\theta(k, y_{K-k})^\top\mathbb{E}_{q_{k-1|k}^\theta}[\epsilon_k|y_{K-k}]] = 0.$

A.5   ALTERNATIVE KULLBACK–LEIBLER DECOMPOSITION

A KL decomposition similar in spirit to the one developed for DDPM (Ho et al., 2020) can be be derived. It leverages the fact that

$$p_{k-1|k,0}(x_{k-1}|x_k, x_0) = \mathcal{N}(x_k; \tilde{\mu}_k(x_k, x_0), \sigma^2 \tilde{\beta}_k I) \tag{42}$$

for $\tilde{\mu}_k(x_k, x_0) = \frac{\sqrt{\bar{\alpha}_{k-1}} \beta_k}{1 - \bar{\alpha}_k} x_0 + \frac{\sqrt{\bar{\alpha}_k}(1 - \bar{\alpha}_{k-1})}{1 - \bar{\alpha}_k} x_k, \tilde{\beta}_k = \frac{1 - \bar{\alpha}_{k-1}}{1 - \bar{\alpha}_k} \beta_k$.

Proposition 4. *The reverse Kullback–Leibler discrepancy* $\mathrm{KL}(q^\theta || p)$ *satisfies*

$$\mathrm{KL}(q^\theta || p) = \mathbb{E}_{q^\theta} \Big[\mathrm{KL}(q_K(x_K) || p_{K|0}(x_K | x_0)) + \sum_{k=2}^{K} \mathrm{KL}(q_{k-1|k}^\theta(x_{k-1}|x_k) || p_{k-1|0,k}(x_{k-1}|x_0, x_k))$$

$$+ \mathrm{KL}(q_{0|1}^\theta(x_0|x_1) || \pi(x_0)) \Big].$$

So for $q_{k-1|k}^\theta(x_{k-1}|x_k) = \mathcal{N}(x_{k-1}; \sqrt{1 - \beta_k} x_k + \sigma^2 \beta_k f_\theta(k, x_k), \sigma^2 \beta_k I)$, *the terms* $\mathrm{KL}(q_{k-1|k}^\theta(x_{k-1}|x_k) || p_{k-1|0,k}(x_{k-1}|x_0, x_k))$ *are KL between two Gaussian distributions and can be calculated analytically.*

We found this elegant decomposition to be however numerically unstable and prone to diverging in our reverse KL setting. Empirically we observed the term $\mathrm{KL}(q_{0|1}^\theta(x_0|x_1) || \pi(x_0))$ becoming unstable once approaching the true value of $\ln Z$ this happened since the term $x_1 = x_1^\theta$ has attached gradients and is able to become large making the term $\mathrm{KL}(q_{0|1}^\theta(x_0|x_1) || \pi(x_0))$ unstable. Furthermore we notice that this formulation of the KL has no term that prevents x_1 from becoming large thus leading to numerical instabilities, note that this is not an issue with DDPM since in their setting x_1 is fixed and has no gradients.

Proof. The reverse KL can be decomposed as follows

$$\mathrm{KL}(q^\theta || p) = \mathbb{E}_{q^\theta} \Big[\ln \Big(\frac{q^\theta(x_{0:K})}{p(x_{0:K})} \Big) \Big] = \mathbb{E}_{q^\theta} \Big[\ln \Big(\frac{q^\theta(x_{0:K})}{\pi(x_0) p(x_{1:K}|x_0)} \Big) \Big]$$

$$= \mathbb{E}_{q^\theta} \Big[\ln \Big(\frac{q^\theta(x_{0:K})}{p(x_{1:K}|x_0)} \Big) \Big] - \mathbb{E}_{q^\theta} [\ln \pi(x_0)]$$

$$= L(\theta) - \mathbb{E}_{q^\theta} [\ln \pi(x_0)]$$

where

$$L(\theta) = \mathbb{E}_{q^\theta} \Big[\ln \Big(\frac{q^\theta(x_{0:K})}{p(x_{1:K}|x_0)} \Big) \Big] = \mathbb{E}_{q^\theta} \Big[\ln q_K(x_K) + \sum_{k=1}^{K} \ln \Big(\frac{q_{k-1|k}^\theta(x_{k-1}|x_k)}{p_{k|k-1}(x_k|x_{k-1})} \Big) \Big]$$

Now using the identity for $k \geq 2$

$$p_{k-1,k|0}(x_{k-1}, x_k | x_0) = p_{k-1|0}(x_{k-1}|x_0) p_{k|k-1}(x_k|x_{k-1})$$

$$= p_{k-1|0,k}(x_{k-1}|x_0, x_k) p_{k|0}(x_k|x_0),$$

we can rewrite $L(\theta)$ as

$$L(\theta) = \mathbb{E}_{q^\theta} \Big[\ln q_K(x_K) + \sum_{k=2}^{K} \ln \Big(\frac{q_{k-1|k}^\theta(x_{k-1}|x_k)}{p_{k-1|0,k}(x_{k-1}|x_0, x_k)} \cdot \frac{p_{k-1|0}(x_{k-1}|x_0)}{p_{k|0}(x_k|x_0)} \Big) + \ln \Big(\frac{q_{0|1}^\theta(x_0|x_1)}{p_{1|0}(x_1|x_0)} \Big) \Big]$$

$$= \mathbb{E}_{q^\theta} \Big[\ln \Big(\frac{q_K(x_K)}{p_{K|0}(x_K|x_0)} \Big) + \sum_{k=2}^{K} \ln \Big(\frac{q_{k-1|k}^\theta(x_{k-1}|x_k)}{p_{k-1|0,k}(x_{k-1}|x_0, x_k)} \Big) + \ln q_{0|1}^\theta(x_0|x_1) \Big]$$

$$= \mathbb{E}_{q^\theta} [\mathrm{KL}(q_K(x_K) || p_{K|0}(x_K|x_0))] + \sum_{k=2}^{K} \mathrm{KL}(q_{k-1|k}^\theta(x_{k-1}|x_k) || p_{k-1|0,k}(x_{k-1}|x_0, x_k))$$

$$+ \ln q_{0|1}^\theta(x_0|x_1) \Big]$$

The result now follows directly.   □

B UNDERDAMPED LANGEVIN DYNAMICS

In the generative modeling context, it has been proposed to extend the original state $x \in \mathbb{R}^d$ by a momentum variance $p \in \mathbb{R}^d$. One then diffuses in this extended space using an underdamped Langevin dynamics (Dockhorn et al., 2022) targeting $\mathcal{N}(x; 0, \sigma^2 I)\mathcal{N}(p; 0, M)$. It was demonstrated empirically that the resulting scores one needs to estimate are smoother and this led to improved performance. We adapt this approach to Monte Carlo sampling. This adaptation is non trivial and in particular requires to design carefully numerical integrators.

B.1 CONTINUOUS TIME

We now consider an augmented target distribution $\pi(x)\mathcal{N}(0, M)$ where M is a positive definite mass matrix. We then diffuse this extended target using the following underdamped Langevin dynamics, i.e.

$$\mathrm{d}x_t = M^{-1}p_t\mathrm{d}t, \tag{43}$$
$$\mathrm{d}p_t = -\frac{x_t}{\sigma^2}\mathrm{d}t - \beta_t p_t\mathrm{d}t + \sqrt{2\beta_t}M^{1/2}\mathrm{d}B_t,$$

where $x_0 \sim \pi, p_0 \sim \mathcal{N}(0, M)$. The resulting path measure on $[0, T]$ is denoted again $\mathcal{P}$. From (Haussmann & Pardoux, 1986), the time-reversal process is also a diffusion satisfying

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \tag{44}$$
$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t + \beta_{T-t}q_t\mathrm{d}t + 2\beta_{T-t}M\nabla_{q_t}\ln\eta_{T-t}(y_t, q_t)\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t,$$

for $(y_0, q_0) \sim \eta_T$ where η_t denotes the density of (x_t, p_t) under (43).

Now consider a reference process $\mathcal{P}^{\mathrm{ref}}$ on $[0, T]$ defined by the forward process (43) initialized using $x_0 \sim \mathcal{N}(0, \sigma^2), p_0 \sim \mathcal{N}(0, M)$. In this case one can check that $\eta_t^{\mathrm{ref}}(x_t, p_t) = \mathcal{N}(x_t; 0, \sigma^2 I)\mathcal{N}(p_t; 0, M)$ and the time-reversal process of $\mathcal{P}^{\mathrm{ref}}$ satisfies

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \tag{45}$$
$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t + \beta_{T-t}q_t\mathrm{d}t - 2\beta_{T-t}q_t\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t$$
$$= \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t}q_t\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t,$$

as $\nabla_q \ln\eta_t^{\mathrm{ref}}(y, q) = \nabla_q \ln(\mathcal{N}(y; 0, \sigma^2)\mathcal{N}(q; 0, M)) = -M^{-1}q$.

Hence it follows that the time-reversal (44) of $\mathcal{P}$ can be also be written as

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \tag{46}$$
$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t}q_t\mathrm{d}t + 2\beta_{T-t}M\nabla_{q_t}\ln\phi_{T-t}(y_t, q_t)\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t,$$

where $\phi_t(x, p) := \eta_t(x, p)/\eta_t^{\mathrm{ref}}(x, p)$.

To approximate $\mathcal{P}$, we consider a parameterized path measure $\mathcal{Q}^\theta$ whose time reversal is defined for $(y_0, q_0) \sim \mathcal{N}(y_0; 0, I)\mathcal{N}(q_0; 0, M)$ by $\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t$ and

$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t - \beta_{T-t}q_t\mathrm{d}t + 2\beta_{T-t}Mf_\theta(T-t, y_t, q_t)\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t. \tag{47}$$

B.1.1 KL DIVERGENCE FOR UNDERDAMPED

In this section we present the continuous time KL for the underdamped sampler as well as its normalising flow based estimators for $\ln Z$.

Proposition 5. *The Radon-Nikodym derivative $\frac{\mathrm{d}\mathcal{Q}^\theta}{\mathrm{d}\mathcal{P}^{\mathrm{ref}}}(y_{[0,T]}, q_{[0,T]})$ satisfies under $\mathcal{Q}^\theta$*

$$\ln\left(\frac{\mathrm{d}\mathcal{Q}^\theta}{\mathrm{d}\mathcal{P}^{\mathrm{ref}}}\right) = \int_0^T \beta_{T-t}||f_\theta(T-t, y_t, q_t)||_M^2\mathrm{d}t + \int_0^T \sqrt{2\beta_{T-t}}M^{1/2}f_\theta(T-t, y_t, q_t)^\top\mathrm{d}W_t, \tag{48}$$

where $||x||_A^2 := x^\top A x$. From $\mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}) = \mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}^{\mathrm{ref}}) + \mathbb{E}_{Q^\theta}\left[\ln\left(\frac{p_0^{\mathrm{ref}}(y_T, q_T)}{\eta_0(y_T, q_T)}\right)\right]$, it follows that

$$\mathrm{KL}(\mathcal{Q}^\theta||\mathcal{P}) = \mathbb{E}_{\mathcal{Q}^\theta}\left[\int_0^T \beta_{T-t}||f_\theta(T-t, y_t, q_t)||_M^2\mathrm{d}t + \ln\left(\frac{\mathcal{N}(y_T; 0, \sigma^2 I)}{\pi(y_T)}\right)\right]. \tag{49}$$

Similarly to Section 2.4, after having learned θ by minimizing (49), we can design a continuous-time normalizing flow to approximate π. In practice, this suggests that once we have learned an approximation $f_\theta(t, x, p) \approx \nabla_p \ln \phi_t(x, p)$ by minimization of (49) then we can construct a proposal using the following ODE initialized at $(y_0, q_0) \sim \mathcal{N}(y_0; 0, \sigma^2 I)\mathcal{N}(q_0; 0, M)$

$$\mathrm{d}y_t = -M^{-1}q_t \mathrm{d}t, \quad \mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t + \beta_{T-t}Mf_\theta(T - t, y_t, q_t)\mathrm{d}t. \tag{50}$$

The resulting sample (y_T, q_T) has a distribution $\sim \bar{q}_0$ close to $\pi(y_T)\mathcal{N}(q_T; 0, M)$ and $\bar{q}_0$ can be computed pointwise via the instantaneous change of variables formula.

B.2 Normalizing flow through ordinary differential equation

The following ODE that gives exactly the same marginals η_T as the SDE (43) defining $\mathcal{P}$

$$\mathrm{d}x_t = M^{-1}p_t\mathrm{d}t, \tag{51}$$

$$\mathrm{d}p_t = -\frac{x_t}{\sigma^2}\mathrm{d}t - \beta_t p_t \mathrm{d}t - \beta_t M \nabla_{p_t} \ln \eta_t(x_t, p_t)\mathrm{d}t,$$

$$= -\frac{x_t}{\sigma^2}\mathrm{d}t - \beta_t M \nabla_{p_t} \ln \phi_t(x_t, p_t)\mathrm{d}t,$$

Thus if we integrate its time reversal from 0 to T starting from $(y_0, q_0) \sim \eta_T$

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \tag{52}$$

$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t + \beta_{T-t}M\nabla_{q_t} \ln \phi_{T-t}(y_t, q_t)\mathrm{d}t,$$

then we would obtain at time T a sample $(y_T, q_T) \sim \pi(y_T)\mathcal{N}(q_T; 0, M)$.

In practice, this suggests that once we have learned an approximation $f_{\theta^*}(t, x, p) \approx \nabla_p \ln \phi_t(x, p)$ by minimization of the ELBO then we can construct a proposal using

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \tag{53}$$

$$\mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t + \beta_{T-t}Mf_{\theta^*}(T - t, y_t, q_t)\mathrm{d}t,$$

for $(y_0, q_0) \sim \mathcal{N}(y_0; 0, \sigma^2 I)\mathcal{N}(q_0; 0, M)$. The resulting sample $(y_T, q_T) \sim \bar{q}_0$ will have distribution close to $\pi \times \mathcal{N}(0, M)$. Again it is possible to compute pointwise the distribution of this sample.

B.3 From continuous time to discrete time

We now need to come up with discrete-time integrator for the time-reversal of $\mathcal{P}^{\mathrm{ref}}$ given by (45) and the time-reversal of $\mathcal{Q}^\theta$ given by (47). Let us start with (45). We split it into the two components

$$\mathrm{d}y_t = -M^{-1}q_t\mathrm{d}t, \qquad \mathrm{d}q_t = \frac{y_t}{\sigma^2}\mathrm{d}t, \tag{54}$$

and

$$\mathrm{d}y_t = 0, \qquad \mathrm{d}q_t = -\beta_{T-t}q_t\mathrm{d}t + \sqrt{2\beta_{T-t}}M^{1/2}\mathrm{d}W_t. \tag{55}$$

We will compose these transitions. To obtain (y_{k+1}, q_{k+1}) from (y_k, q_k), we first integrate (54). To do this, consider the Hamiltonian equation $\mathrm{d}x_t = M^{-1}p_t\mathrm{d}t, \mathrm{d}p_t = -\frac{x_t}{\sigma^2}\mathrm{d}t$ which preserves $\mathcal{N}(y; 0, \sigma^2 I)\mathcal{N}(q; 0, M)$ as invariant distribution. We can integrate this ODE exactly over an interval of length δ and denote its solution $\Phi(x, p)$; see Section B.4 for details. We use its inverse $\Phi^{-1}(x, p) = \Phi_{\mathrm{flip}}(x, p) \circ \Phi(x, p) \circ \Phi_{\mathrm{flip}}(x, p)$ where $\Phi_{\mathrm{flip}}(x, p) = (x, -p)$ so that $(y_{k+1}, q'_k) = \Phi^{-1}(y_k, q_k)$. Then we integrate exactly (55) using

$$q_{k+1} = \sqrt{1 - \alpha_{K-k}}q'_k + \sqrt{\alpha_{k-1}}M^{1/2}\epsilon_k. \tag{56}$$

We have thus design a transition of the form

$$p^{\mathrm{ref}}(y_{k+1}, q_{k+1}, q'_k | y_k, q_k) = \delta_{\Phi^{-1}(y_k, q_k)}(y_{k+1}, q'_k)\mathcal{N}(q_{k+1}; \sqrt{1 - \alpha_{K-k}}q'_k; \alpha_{K-k}M) \tag{57}$$

Now to integrate (47), we split it in three parts. We first integrate (54) using exactly the same integrator $(y_{k+1}, q'_k) = \Phi^{-1}(y_k, q_k)$. Then we integrate

$$\mathrm{d}q_t = 2\beta_{T-t}Mf_\theta(T - t, y_t, q_t)\mathrm{d}t \tag{58}$$

using

$$q_k'' = q_k' + 2(1 - \sqrt{1 - \alpha_{K-k}}) M f_\theta(K - k, y_{k+1}, q_k'), \tag{59}$$

where we abuse notation and write $f_\theta(K - k, y_{k+1}, q_k')$ instead of $f_\theta((K - k)\delta, y_{k+1}, q_k')$. Finally, we integrate the OU part using (56) but replacing q_k' by with q_k''. So the final transition we get is

$$q^\theta(y_{k+1}, q_{k+1}, q_k', q_k'' | y_k, q_k) = \delta_{\Phi^{-1}(y_k, q_k)}(y_{k+1}, q_k') \delta_{q_k' + 2(1 - \sqrt{1 - \alpha_{K-k}}) M f_\theta(K-k, y_{k+1}, q_k')}(q_k'')$$

$$\times \mathcal{N}(q_{k+1}; \sqrt{1 - \alpha_{K-k}} q_k''; \alpha_{K-k} M).$$

However, we can integrate q_k'' analytically to obtain

$$q^\theta(y_{k+1}, q_{k+1}, q_k' | y_k, q_k) = \delta_{\Phi^{-1}(y_k, q_k)}(y_{k+1}, q_k') \tag{60}$$

$$\times \mathcal{N}(q_{k+1}; \sqrt{1 - \alpha_{K-k}}(q_k' + 2(1 - \sqrt{1 - \alpha_{K-k}}) M f_\theta(K - k, y_{k+1}, q_k')); \alpha_{K-k} M). \tag{61}$$

B.4 EXACT SOLUTION OF HAMILTONIAN EQUATION

Consider the Hamiltonian equation and $M = I$, we do have for

$$\mathrm{d}x_t = p_t \mathrm{d}t, \qquad \mathrm{d}p_t = -\frac{x_t}{\sigma^2} \mathrm{d}t \tag{62}$$

that the solution can be exactly written as $(x_t, p_t) = \Phi_t(x_0, p_0)$ defined through

$$x_t = x_0 \cos(t/\sigma) + p_0 \sigma \sin(t/\sigma) \tag{63}$$

$$p_t = -\frac{x_0}{\sigma} \sin(t/\sigma) + p_0 \cos(t/\sigma) \tag{64}$$

This is the so-called harmonic oscillator.

Now the inverse of the Hamiltonian flow satisfies $\Phi^{-1}(x, p) = \Phi_{\mathrm{flip}}(x, p) \circ \Phi(x, p) \circ \Phi_{\mathrm{flip}}(x, p)$ (see e.g. Leimkuhler & Matthews (2016)) so we can simply integrate

$$\mathrm{d}y_t = -q_t \mathrm{d}t, \qquad \mathrm{d}q_t = \frac{y_t}{\sigma^2} \mathrm{d}t \tag{65}$$

using $(y_{k+1}, q_k') = \Phi_\tau^{-1}(y_k, q_k)$.

This gives

$$y_{k+1} = y_k \cos(\tau/\sigma) - q_k \sigma \sin(\tau/\sigma) \tag{66}$$

$$q_k' = -(-\frac{y_k}{\sigma} \sin(\tau/\sigma) - q_k \cos(t/\sigma))$$

$$= \frac{y_k}{\sigma} \sin(\tau/\sigma) + q_k \cos(t/\sigma) \tag{67}$$

Obviously, if we have $M = mI$ then

$$\mathrm{d}x_t = \frac{p_t}{m} \mathrm{d}t, \qquad \mathrm{d}p_t = -\frac{x_t}{\sigma^2} \mathrm{d}t \tag{68}$$

then we use a reparameterization $\tilde{p}_t = p_t/m$ and

$$\mathrm{d}x_t = \tilde{p}_t \mathrm{d}t, \qquad \mathrm{d}\tilde{p}_t = -\frac{1}{m} \frac{x_t}{\sigma^2} \mathrm{d}t, \tag{69}$$

and the solution is as above with σ being replaced by $\sigma\sqrt{m}$.

So for $\tilde{q}_k = q_k/m$, we have

$$y_{k+1} = y_k \cos(\tau/(\sqrt{m}\sigma)) - \tilde{q}_k \sigma \sqrt{m} \sin(\tau/(\sqrt{m}\sigma))$$

$$\tilde{q}_k' = \frac{y_k}{\sqrt{m}\sigma} \sin(\tau/(\sqrt{m}\sigma)) + \tilde{q}_k \cos(t/(\sqrt{m}\sigma))$$

so as $q_k = m\tilde{q}_k$ and similarly $q_k' = m\tilde{q}_k'$, we finally obtain

$$y_{k+1} = y_k \cos(\tau/(\sqrt{m}\sigma)) - q_k \frac{\sigma}{\sqrt{m}} \sin(\tau/(\sqrt{m}\sigma)) \tag{70}$$

$$q_k' = \frac{\sqrt{m}}{\sigma} y_k \sin(\tau/(\sqrt{m}\sigma)) + q_k \cos(\tau/(\sqrt{m}\sigma)) \tag{71}$$

B.5 DERIVATION OF UNDERDAMPENED KL IN DISCRETE TIME

In this section we derive the discrete

Proposition 6. *The log density ratio* $\mathrm{lr} = \ln(q^\theta(y_{0:K}, q_{0:K}, q'_{0:K})/p^{\mathrm{ref}}(y_{0:K}, q_{0:K}, q'_{0:K}))$ *equals*

$$\mathrm{lr} = \sum_{k=1}^{K} 2\left[\frac{\kappa_k^2}{\alpha_k}||f_\theta(k, y_{K-k+1}, q'_{K-k})||_M^2 + M^{1/2}\frac{\kappa_k}{\sqrt{\alpha_k}}f_\theta(k, y_{K-k+1}, q'_{K-k})^\top\varepsilon_k\right] \tag{72}$$

where $\kappa_k := \sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})$ *and* ε_k *is obtained as a function of* q_{k+1} *and* q''_k *when describing the integrator for* $\mathcal{Q}^\theta$. *In particular for* $y_{0:K}, q_{0:K}, q'_{0:K} \sim q^\theta(\cdot)$, *hence* $\varepsilon_k \overset{\mathrm{i.i.d.}}{\sim} \mathcal{N}(0, I)$, *one obtains*

$$\mathrm{KL}(q^\theta||p) = 2\mathbb{E}_{q^\theta}\left[\sum_{k=1}^{K}\frac{\kappa_k^2}{\alpha_k}\left[||f_\theta(k, y_{K-k+1}, q'_{K-k})||_M^2 + \ln\left(\frac{\mathcal{N}(y_K; 0, \sigma^2 I)}{\pi(y_K)}\right)\right]\right]. \tag{73}$$

The intergrator has been designed so that the ratio between the transitions of the proposal (60) and the reference process (57) is well-defined as the deterministic parts are identical in the two transitions so cancel and

$$\frac{q^\theta(y_{k+1}, q_{k+1}, q'_k|y_k, q_k)}{p^{\mathrm{ref}}(y_{k+1}, q_{k+1}, q'_k|y_k, q_k)} \tag{74}$$
$$= \frac{\mathcal{N}(q_{k+1}; \sqrt{1-\alpha_{K-k}}(q'_k + 2(1-\sqrt{1-\alpha_{K-k}})Mf_\theta(K-k, y_{k+1}, q'_k)); \alpha_{K-k}M)}{\mathcal{N}(q_{k+1}; \sqrt{1-\alpha_{K-k}}q'_k; \alpha_{K-k}M)}.$$

The calculations to compute the Radon-Nikodym derivative are now very similar to what we did in the proof of Proposition 3.

We have

$$\ln\left(\frac{q^\theta(y_{0:K}, q_{0:K}, q'_{0:K})}{p^{\mathrm{ref}}(y_{0:K}, q_{0:K}, q'_{0:K})}\right) \tag{75}$$
$$= \ln\left(\frac{q_K(y_0, q_0)}{p_K^{\mathrm{ref}}(y_0, q_0)}\right) + \sum_{k=1}^{K}\ln\left(\frac{q^\theta_{k-1|k}(y_{K-k+1}, q_{K-k+1}, q'_{K-k}|y_{K-k}, q_{K-k})}{p^{\mathrm{ref}}_{k-1|k}(y_{K-k+1}, q_{K-k+1}, q'_{K-k}|y_{K-k}, q_{K-k})}\right)$$
$$= \sum_{k=1}^{K}\Big[||q_{K-k+1} - \sqrt{1-\alpha_k}q'_{K-k}||^2_{(2\alpha_k M)^{-1}}$$
$$- ||q_{K-k+1} - \sqrt{1-\alpha_k}(q'_{K-k} + 2(1-\sqrt{1-\alpha_k})Mf_\theta(k, y_{K-k+1}, q'_{K-k}))||^2_{(2\alpha_k M)^{-1}}\Big]$$

where we have exploited the fact that $q_K(y_0, q_0) = p_K^{\mathrm{ref}}(y_0, q_0) = \mathcal{N}(y_0; 0, \sigma^2 I)\mathcal{N}(q_0; 0, M)$.

Now let us introduce

$$\epsilon_k := \frac{M^{-1/2}}{\sqrt{\alpha_k}}(q_{K-k+1} - \sqrt{1-\alpha_k}q'_{K-k} - 2\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})Mf_\theta(k, y_{K-k}, q'_{K-k})) \tag{76}$$

hence

$$\frac{M^{-1/2}}{\sqrt{\alpha_k}}(q_{K-k+1} - \sqrt{1-\alpha_k}q'_{K-k})$$
$$= \epsilon_k + \frac{2M^{-1/2}}{\sqrt{\alpha_k}}\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})Mf_\theta(k, y_{K-k+1}, q'_{K-k})$$
$$= \epsilon_k + \frac{2M^{1/2}}{\sqrt{\alpha_k}}\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})f_\theta(k, y_{K-k+1}, q'_{K-k}).$$

We can rewrite (75) as

$$
\ln\left(\frac{q^\theta(y_{0:K}, q_{0:K}, q'_{0:K})}{p^{\mathrm{ref}}(y_{0:K}, q_{0:K}, q'_{0:K})}\right) \tag{77}
$$

$$
= \frac{1}{2}\sum_{k=1}^{K}\left[||\epsilon_k + \frac{2M^{1/2}}{\sqrt{\alpha_k}}\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})f_\theta(k, y_{K-k+1}, q'_{K-k}))||^2 - ||\epsilon_k||^2\right]
$$

$$
= \frac{1}{2}\sum_{k=1}^{K}\left[||f_\theta(k, y_{K-k+1}, q'_{K-k}))||^2_{4M\frac{(1-\alpha_k)(1-\sqrt{1-\alpha_k})^2}{\alpha_k}}\right.
$$

$$
\left. + \frac{4M^{1/2}}{\sqrt{\alpha_k}}\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})f_\theta(k, y_{K-k+1}, q'_{K-k}))^\top \varepsilon_k\right]
$$

$$
= \sum_{k=1}^{K}\left[||f_\theta(k, y_{K-k+1}, q'_{K-k}))||^2_{2M\frac{(1-\alpha_k)(1-\sqrt{1-\alpha_k})^2}{\alpha_k}}\right.
$$

$$
\left. + 2M^{1/2}\frac{\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})}{\sqrt{\alpha_k}}f_\theta(k, y_{K-k+1}, q'_{K-k}))^\top \varepsilon_k\right]
$$

where we note that $\frac{\sqrt{1-\alpha_k}(1-\sqrt{1-\alpha_k})}{\sqrt{\alpha_k}} \approx \alpha_k/2 \approx \beta_{k\delta}\delta$.

C EXPERIMENTS

In this section we incorporate additional experiments and ablations as well as further experimental detail. For both PIS and DDS we created a grid with $\delta = \frac{K}{T} = 0.05$ and values of $T \in \{3.4, 6.4, 12.8, 25.6\}$ and corresponding number of steps $K \in \{64, 128, 128, 256\}$. When tuning PIS we explore different values of σ which in practice, when discretised, amounts to the same effect as changing δ. For DDS we apply the cosine schedule directly on the number of steps however we ensure that $\sum_k \alpha_k = \alpha_{\max}T$ such that we have a similar scaling as in PIS. Finally both PIS and DDS where trained with Adam (Kingma & Ba, 2015) to at most 11000 iterations, although in most cases most both converged in less than 6000 iterations. Across all experiments used a sample size of 300 to estimate the ELBO at train time and of 2000 to for the reported IS estimator of Z for all methods.

C.1 TUNING HYPER-PARAMETERS

For both DDS and PIS we explore a grid of 25 hyper-parameter values. For PIS we search for the best performing value of the volatility coefficient over 25 values, depending on the task we vary the end points of the drift however we noticed that PIS leads to numerical instabilities for $\gamma > 4$ thus we never search for values larger than this. For DDS we searched across 5 values for σ and 5 values for $\alpha_{\max}$, this led to a total of 25 combinations we explored for each experiment. Finally for SMC we searched over 3 of its different step sizes leading to a total of $5^3 = 125$.

C.2 BROWNIAN MOTION MODEL WITH GAUSSIAN OBSERVATION NOISE

The model is given by

$$
\begin{aligned}
\alpha_{\mathrm{inn}} &\sim \mathrm{LogNormal}(0, 2), \\
\alpha_{\mathrm{obs}} &\sim \mathrm{LogNormal}(0, 2), \\
x_1 &\sim \mathcal{N}(0, \alpha_{\mathrm{inn}}), \\
x_i &\sim \mathcal{N}(x_{i-1}, \alpha_{\mathrm{inn}}), \quad i = 2, \ldots 20 \\
y_i &\sim \mathcal{N}(x_i, \alpha_{\mathrm{obs}}), \quad i = 1, \ldots 30
\end{aligned}
$$

The goal is to perform inference over the variables $\alpha_{\mathrm{inn}}, \alpha_{\mathrm{obs}}$ and $\{x_i\}_{i=1}^{30}$ given the observations $\{y_i\}_{i=1}^{10} \cup \{y_i\}_{i=20}^{30}$.

Figure 4: Magnitude comparison of trained networks. Both trained to convergence.

	Training Times per Iteration - seconds (LGCP)			
	$k = 64$	$k = 128$	$k = 256$	$k = 512$
DDS	0.104 ± 0.0004	0.205 ± 0.0004	0.406 ± 0.0004	0.809 ± 0.0004
PIS	0.104 ± 0.0004	0.205 ± 0.0003	0.406 ± 0.0005	0.808 ± 0.0006

Table 2: Training time per ELBO gradient update on 300 samples.

C.3   DRIFT MAGNITUDE

In Figure 4 we compare the magnitudes of the NN drifts learned by PIS and DDS on a $\mathcal{N}(6, \sigma)$ target. The drifts where sampled on randomly evaluated trajectories from each of the learned samplers. We observe the PIS drift to have large magnitudes with higher variance more prone to numerical instabilities, whilst the OU drift has a notably lower magnitude with significantly less variance. We conjecture this is the reason that PIS becomes unstable at training time for a large number of steps across several of our PIS experiments, many of which did not converge due numerical instabilities. For simplicity in this experiment we set α_k to be uniform.

C.4   TRAINING TIME

We evaluate the training times of DDS and PIS on an 8-chip TPU circuit and average over 99 runs. Table 2 shows running times per training iteration. In total we trained for 11000 iterations thus the total training times are 19 minutes; 37 minutes; 1 hour and 14 minutes; 2 hours and 26 minutes for $k = 64, 128, 256, 512$ respectively.

C.5   VARIATIONAL ENCODER IN LATENT SPACE

Following Arbel et al. (2021) we explore estimating the posterior of a pre-trained Variational Auto Encoder (VAE) (Kingma & Welling, 2013). We found all approaches to reach a very small error for $\ln Z$ very quickly however as seen with PIS across experiments results became unstable for large T and thus many hyper-parameters when tuning the volatility for PIS led to the loss exploding.

C.6   GENERATED IMAGES

In this section we provide some of the generated samples for the normalising flow evaluation task. In Figure 6 we can observe how both PIS and DDS are able to generate more image classes than SMC due to mode collapse. Out of the 3 approaches we can see that DDS mode collapses the least whilst SMC generates the higher quality images. Using a neural network with inductive biases such as a convolutional neural network (LeCun et al., 1998) should improve the image quality over the

Figure 5: Results on pretrained VAE from Arbel et al. (2021).

Figure 6: a) DDS , b) PIS , c) SMC. We can see how SMC mode collapses significantly as it only seems to have 5 image classes. For PIS we identified 7 classes whilst for DDS we found 9 out of 10.

simple feed-forward networks we have used, we believe this is the reason behind the low quality in the images as both minimising the path KL and sampling high quality images is a difficult task for a small feed-forward network.

C.7 COMPARISON TO AIS AND MCD

We compare our results to the AIS and MCD baselines presented in Geffner & Domke (2022). Results can be seen in Tables 4 and 3, we can see that both DDS and PIS outperform LDVI across all number of steps and tasks with DDS outperforming PIS on higher values of k due to inconssitent and unstable behaviour presented by PIS.

C.8 EULER MAYURAMA VS EXPONENTIAL INTEGRATOR

In this section we demonstrate empirically how using the Euler Mayurama integrator directly on the DDS objective leads to overestimating $\ln Z$. We compare on the Funnel dataset for which we know $\ln Z$ to be 0 and the Ionosphere dataset which is small and for which we have a reliable estimate of $\ln Z$ via a long run SMC. Results in Table 5 show case how for lower values of $k = 64, 128$ the EM based approach significantly overestimates $\ln Z$ and overall does not perform well.

	Ionosphere						Sonar					
	ULA	MCD	UHA	LDVI	DDS	PIS	ULA	MCD	UHA	LDVI	DDS	PIS
$k = 64$	-113.8	-112.5	-112.8	-112.1	-111.7	-111.6	-115.3	-111.1	-111.9	-109.7	-109.4	-108.9
$k = 128$	-113.1	-112.2	-112.3	-111.9	-111.6	-112.0	-113.5	-110.2	-110.6	-109.1	-108.9	-109.3
$k = 256$	-112.7	-112.1	-112.1	-111.7	-111.6	-113.0	-112.1	-109.7	-109.7	-108.9	-108.9	-108.9

Table 3: Comparison of DDS and PIS to AIS based approaches on logistic regression models.

	Brownian Motion					
	ULA	MCD	UHA	LDVI	DDS	PIS
$k = 64$	-0.7	0.1	0.1	- 0.5	1.0	1.0
$k = 128$	-0.3	0.2	0.4	0.7	1.1	1.0
$k = 256$	-0.1	0.5	0.6	0.9	1.0	0.7

Table 4: Comparison of DDS and PIS to AIS based approaches on Brownian motion timeseries target.

	Funnel			Ionosphere		
	Exponential Integrator	Euler Mayurama	Ground Truth	Exponential Integrator	Euler Mayurama	Ground Truth
$k = 64$	-0.206 ± 0.059	2.425 ± 0.266		-111.693 ± 0.169	-106.364 ± 0.371	
$k = 128$	-0.176 ± 0.099	2.011 ± 0.532	0	-111.587 ± 0.136	-111.158 ± 0.399	-111.560
$k = 256$	-0.221 ± 0.076	-0.086 ± 0.124		-111.575 ± 0.163	-112.271 ± 0.366	
$k = 512$	-0.176 ± 0.068	-0.154 ± 0.066		-111.582 ± 0.136	-112.912 ± 0.353	

Table 5: Comparing EM vs Exponential integrators on Funnel and Ionosphere datasets. We can see how EM significantly overestimates $\ln Z$.

C.8.1   DETACHED GRADIENT ABLATIONS

In this section we perform an ablation over our proposed modification of the PIS-Grad network architecture. We compare the same architecture with and without the gradient attached. We found that detaching gradient led to a favourable performance as well as more stable training. Results are presented in table 6 and were carried out for $K = 128$ and using the best found diffusion coefficient from each task.

We chose to explore this feature as it is known that optimization through unrolled computation graphs can be chaotic and introduce exploding/vanishing gradients (Parmas et al., 2018; Metz et al., 2020) which can numerically introduce bias. This has been successfully applied in related prior work (Greff et al., 2019) where detached scores are provided as features to the recognition networks. Alternative approaches exist based on smoothing the chaotic loss-landscape (Vicol et al., 2021) but we leave this for future work.

C.9   PROBABILITY FLOW ODE

Figures 7 and 8 show samples obtained from the probability flow ODE of a trained DDS model. We can see that the probability flow ODE is able to perfectly sample from a uni-modal Gaussian, matters became more challenging with multi-modal distributions. Initially we discretised the ODE (12) using the same type of integrators as the SDEs to obtain $y_{k+1} = y_k + \sigma^2(1 - \sqrt{1 - \alpha_{K-k}})f_\theta(K - k, y_k)$ for $y_0 \sim \mathcal{N}(0; \sigma^2 I)$. Unfortunately under this discretisation we found the probability flow ODE to become stiff with more complex distributions such as the mixture of Gaussian's, resulting in strange effects in the samples (matching the modes but completely wrong shapes). By using a Heun integrator as proposed in Karras et al. (2022) we were able to obtain better results, however we also found we needed to increase the network size to improve the results as well as train with a learning rate decay of 0.99, with these extra features we were able to simulate a prob flow ODE that roughly matched the marginal densities of the SDE. However even with these fixes we still found the prob flow based estimator of $\ln Z$ to drastically overestimate, this is not entirely surprising as in discrete time this estimator is not guaranteed to be a lower bound.

C.10   UNDERDAMPED OU RESULTS

We perform some additional experiments with the underdamped OU reference process. Similar to the damped setting we parametrise the network as:

$$f_\theta(k, x, p) = \mathrm{NN}_1(k, x, p; \theta) + \mathrm{NN}_2(k; \theta)\nabla \ln \pi(x). \tag{78}$$

Unfortunately unlike in DDS and PIS we cannot directly aid the update/proposal for y_t with $\nabla \ln \pi(y)$ thus this naive parametrisation is not the ideal inductive bias.

Method	Funnel	LGCP	VAE	Sonar	Ionosphere	Brownian	NICE
PIS Grad Detach	-0.268 ± 0.07	502.002 ± 0.957	-110.025 ± 0.0704	-109.349 ± 0.356	-111.602 ± 0.145	0.960 ± 0.145	-3.985 ± 0.349
PIS Grad	-0.268 ± 0.09	501.273 ± 0.798	-110.033 ± 0.0647	-109.377 ± 0.361	-111.705 ± 0.190	-5.793 ± 0.499	-3.927 ± 0.666

Table 6: Results for PIS grad with and without detaching the gradient.

Figure 7: a) Target distribution (MoG), b) DDS SDE samples, c) trained probability flow ODE.

Results in Figures 9 and 10 show some initial simulation for this approach. The results perform worse than DDS and PIS, however they are still better than VI, we believe that future work exploring better inductive biases for the network $f_\theta(k, x, p)$ could narrow the performance gap in this approach. Additionally we highlight that in Figure 9 (Funnel Distribution) we can see the under-damped approach overestimates $\ln Z$ for $k = 64$. We believe this is due to the more stiff numerics of the second order system at a small number of time-steps.

C.11 BOX PLOTS FOR MAIN RESULTS

For completeness in this section we also present our results via box plots as can be seen in Figures 11 and 12.

D DENOISING DIFFUSION SAMPLERS IN DISCRETE-TIME: FURTHER DETAILS

D.1 FORWARD PROCESS AND ITS TIME REVERSAL

In discrete-time, we consider the following discrete-time "forward" Markov process

$$p(x_{0:K}) = \pi(x_0) \prod_{k=1}^{K} p_{k|k-1}(x_k|x_{k-1}), \tag{79}$$

Following (17), we use $p_{k|k-1}(x_k|x_{k-1}) = \mathcal{N}(x_k; \sqrt{1 - \alpha_k} x_{k-1}, \sigma^2 \alpha_k I)$. The parameters (α_k) are such that $p_K(x_K) \approx \mathcal{N}(x_K; 0, \sigma^2 I)$.

We propose here to sample approximately from π by approximating the ancestral sampling scheme for (79) corresponding to the backward decomposition

$$p(x_{0:K}) = p_K(x_K) \prod_{k=1}^{K} p_{k-1|k}(x_{k-1}|x_k), \text{ with } p_{k-1|k}(x_{k-1}|x_k) = \frac{p_{k-1}(x_{k-1})p_{k|k-1}(x_k|x_{k-1})}{p_k(x_k)}, \tag{80}$$

where $p_0(x_0) = \pi(x_0)$. If we could thus sample $x_K \sim p_K(\cdot)$ then $x_{k-1} \sim p_{k-1|k}(\cdot|x_k)$ for $k = K, ..., 1$ then x_0 would indeed be a sample from π. However as the marginal densities $(p_k)_{k=1}^{K}$ are not available in closed-form, we cannot implement exactly this ancestral sampling procedure.

Figure 8: a) Target distribution $\mathcal{N}(6, 1)$, b) DDS SDE c) Trained DDS probability flow ODE.

Figure 9: $\ln Z$ estimate as a function of number of steps K - a) Funnel , b) LGCP, c) Logistic Ionosphere dataset. Yellow Dotted line is MF-VI and dashed magenta is the gold standard.

Figure 10: $\ln Z$ estimate as a function of number of steps K - a) Logistic Sonar dataset, b) Brownian motion, c) NICE. Yellow dotted line is MF-VI and dashed magenta is the gold standard.

As we have by design $p_K(x_K) \approx \mathcal{N}(x_K; 0, \sigma^2 I)$, we can simply initialize the ancestral sampling procedure by a Gaussian sample. The approximation of the backward Markov kernels $p_{k-1|k}(x_{k-1}|x_k)$ is however more involved.

D.2 REFERENCE PROCESS AND BELLMAN RECURSION

We introduce the "simple" reference process $p^{\text{ref}}(x_{0:K})$ defined by

$$p^{\text{ref}}(x_{0:K}) = \mathcal{N}(x_0; 0, \sigma^2 I)p(x_{1:K}|x_0) = \mathcal{N}(x_0; 0, \sigma^2 I)\prod_{k=1}^{K} p_{k|k-1}(x_k|x_{k-1}) \qquad (81)$$

which is designed by construction to admits the marginal distributions $p_k^{\text{ref}}(x_k) = \mathcal{N}(x_k; 0, \sigma^2 I)$ for all k. It can be easily verified using the chain rule for KL that the extended target process p is the distribution minimizing the reverse (or forward) KL discrepancy w.r.t. p^{ref} over the set of path measures $q(x_{0:K})$ with marginal $q_0(x_0) = \pi(x_0)$ at the initial timem i.e.

$$p = \arg\min_q \left\{ \text{KL}(q||p^{\text{ref}}) : q_0 = \pi \right\}.$$

Figure 11: $\ln Z$ estimate as a function of number of steps K - a) Funnel , b) LGCP, c) Logistic Ionosphere dataset. Yellow Dotted line is MF-VI and dashed magenta is the gold standard.

The intractable backward Markov densities $p_{k-1|k}$ one wants to approximate can also be rewritten as twisted versions of the tractable backward Markov densities $p^{\mathrm{ref}}_{k-1|k}$ by some value functions $(\phi_k)^K_{k=0}$.

Proposition 7. *We have for $k = 1, ..., K$*

$$p_{k-1|k}(x_{k-1}|x_k) = \frac{\phi_{k-1}(x_{k-1})p^{\mathrm{ref}}_{k-1|k}(x_{k-1}|x_k)}{\phi_k(x_k)}, \quad for \quad \phi_k(x_k) = \frac{p_k(x_k)}{p^{\mathrm{ref}}_k(x_k)}. \tag{82}$$

The value functions $(\phi_k)^K_{k=1}$ satisfy a forward Bellman type equation

$$\phi_k(x_k) = \int \phi_{k-1}(x_{k-1})p^{\mathrm{ref}}_{k-1|k}(x_{k-1}|x_k)\mathrm{d}x_{k-1}, \qquad \phi_0(x_0) = \frac{\pi(x_0)}{p^{\mathrm{ref}}_0(x_0)}. \tag{83}$$

It follows that

$$\phi_k(x_k) = \int \phi_0(x_0) \; p^{\mathrm{ref}}_{0|k}(x_0|x_k)\mathrm{d}x_0. \tag{84}$$

Proof. To establish (82), we use Bayes' rule

$$\begin{aligned}p_{k-1|k}(x_{k-1}|x_k) &= \frac{p_{k-1}(x_{k-1})p_{k|k-1}(x_k|x_{k-1})}{p_k(x_k)} \\ &= \frac{\phi_{k-1}(x_{k-1})p^{\mathrm{ref}}_{k-1}(x_{k-1})p_{k|k-1}(x_k|x_{k-1})}{\phi_k(x_k)p^{\mathrm{ref}}_k(x_{k-1})} \\ &= \frac{\phi_{k-1}(x_{k-1})p^{\mathrm{ref}}_{k-1|k}(x_{k-1}|x_k)}{\phi_k(x_k)},\end{aligned}$$

where we have used the fact that $p_k(x_k) = \phi_k(x_k)p^{\mathrm{ref}}_k(x_k)$, $p^{\mathrm{ref}}_{k|k-1}(x_k|x_{k-1}) = p_{k|k-1}(x_k|x_{k-1})$ and Bayes' rule again. Now we have $\int p_{k-1|k}(x_{k-1}|x_k)\mathrm{d}x_{k-1} = 1$ for any x_k, so it follows directly from the expression of this transition kernel that the value function satisfies

$$\phi_k(x_k) = \int \phi_{k-1}(x_{k-1})p^{\mathrm{ref}}_{k-1|k}(x_{k-1}|x_k)\mathrm{d}x_{k-1}, \qquad \phi_0(x_0) = \frac{\pi(x_0)}{p^{\mathrm{ref}}_0(x_0)}. \tag{85}$$

By iterating this recursion, it follows that

$$\phi_k(x_k) = \int \phi_0(x_0) \; p^{\mathrm{ref}}_{0|k}(x_0|x_k)\mathrm{d}x_0. \tag{86}$$

$\square$

D.3 APPROXIMATING THE BACKWARD KERNELS

In practice, we want to approximate the backward Markov transition densities $p_{k-1|k}$. From (82), this would require not only to approximate the value functions - Monte Carlo estimates based on (84) are high variance - but also to sample from the resulting twisted kernel which is difficult. However,

Figure 12: $\ln Z$ estimate as a function of number of steps K - a) Logistic Sonar dataset, b) Brownian motion, c) NICE. Yellow dotted line is MF-VI and dashed magenta is the gold standard.

if we select $\beta_k \approx 0$, then we have $\phi_{k-1}(x) \approx \phi_k(x)$ and by a Taylor expansion of $\nabla \ln \phi_k$ around x_k we obtain

$$p_{k-1|k}(x_{k-1}|x_k) = p_{k-1|k}^{\text{ref}}(x_{k-1}|x_k) \exp(\ln \phi_{k-1}(x_{k-1}) - \ln \phi_k(x_k))$$
$$\approx \mathcal{N}(x_{k-1}; \sqrt{1-\alpha_k}x_k + \sigma^2(1-\sqrt{1-\alpha_k})\nabla \ln \phi_k(x_k)). \qquad (87)$$

By differentiating the identity (84) w.r.t. x_k, we obtain the following expression for ϕ_k

$$\nabla \ln \phi_k(x_k) = \int \nabla \ln p_{0|k}^{\text{ref}}(x_0|x_k) \; p_{0|k}(x_0|x_k)\mathrm{d}x_0. \qquad (88)$$

Alternatively, we can also use $\nabla \ln \phi_k(x_k) = \nabla \ln p_k(x_k) - \frac{x}{\sigma^2}$ where $\nabla \ln p_k(x_k)$ can be easily shown to satisfy

$$\nabla \ln p_k(x_k) = \int \nabla \ln p_{k|0}(x_k|x_0) \; p_{0|k}(x_0|x_k)\mathrm{d}x_0. \qquad (89)$$

In score-based generative modeling, the conditional expectation appearing in (89) is reformulated as the solution to a regression problem, leveraging the fact that we can easily obtain samples from $\pi(x_0)p_{k|0}(x_k|x_0)$ as π is the data distribution in this context. In the Monte Carlo sampling context considered here, we could sample approximately from $p_{0|k}(x_0|x_k) \propto \pi(x_0)p_{k|0}(x_k|x_0)$ using MCMC so as to approximate the conditional expectations in (88) and (89) but this would defeat the purpose of the proposed methodology. We propose instead to approximate $\nabla \ln \phi_k$ by minimizing a suitable criterion.

In practice, we will consider a distribution $q^\theta(x_{0:K})$ approximating $p(x_{0:K})$ of the form

$$q^\theta(x_{0:K}) = \mathcal{N}(x_K; 0, \sigma^2 I) \prod_{k=1}^{K} q_{k-1|k}^\theta(x_{k-1}|x_k), \qquad (90)$$

i.e. we select $q_K^\theta(x_K) = q_K(x_K) = \mathcal{N}(x_K; 0, \sigma^2 I)$ as $p_K(x_K) \approx \mathcal{N}(x_K; 0, \sigma^2 I)$. We want $q_{k-1|k}^\theta(x_{k-1}|x_k)$ to approximate $p_{k-1|k}(x_{k-1}|x_k)$, i.e. $f_\theta(k, x_k) \approx \nabla \ln \phi_k(x_k)$. Inspired by (87), we will consider an approximation of the form

$$q_{k-1|k}^\theta(x_{k-1}|x_k) = \mathcal{N}(x_{k-1}; \sqrt{1-\alpha_k}x_k + \sigma^2(1-\sqrt{1-\alpha_k})f_\theta(k, x_k), \sigma^2\alpha_k I). \qquad (91)$$

where $f_\theta(k, x_k)$ is differentiable w.r.t. θ.

D.4 FITTED HYPERPARAMETERS

To aid reproducibility we report all fitted hyperparameters for each of our methods and PIS across all experiments in Tables 7 and 8.

	funnel			lgcp			ion		
	DDS	PIS	UDMP	DDS	PIS	UDMP	DDS	PIS	UDMP
$K = 64$	$\sigma = 1.075, \alpha = 1.075$	$\sigma = 1.068$	$\sigma = 1.85, \alpha = 1.67, m = 0.9$	$\sigma = 2.1, \alpha = 1.50$	$\sigma = 1.068$	$\sigma = 1.1, \alpha = 2.5, m = 0.4$	$\sigma = 0.688, \alpha = 1.463$	$\sigma = 0.253$	$\sigma = 0.6, \alpha = 3.85, m = 0.600$
$K = 128$	$\sigma = 1.075, \alpha = 0.6875$	$\sigma = 0.416$	$\sigma = 1.85, \alpha = 3.7, m = 0.9$	$\sigma = 2.1, \alpha = 0.75$	$\sigma = 0.742$	$\sigma = 1.4, \alpha = 2.5, m = 0.4$	$\sigma = 0.3, \alpha = 1.075$	$\sigma = 0.09$	$\sigma = 0.6, \alpha = 3.85, m = 0.600$
$K = 256$	$\sigma = 1.85, \alpha = 0.3$	$\sigma = 0.742$	$\sigma = 1.075, \alpha = 2.5, m = 0.9$	$\sigma = 2.1, \alpha = 0.900$	$\sigma = 0.579$	$\sigma = 1.4, \alpha = 4.5, m = 0.4$	$\sigma = 0.3, \alpha = 0.688$	$\sigma = 0.416$	$\sigma = 0.6, \alpha = 3.85, m = 1.0$
$K = 512$	$\sigma = 1.463, \alpha = 0.3$	$\sigma = 0.253$	$\sigma = 0.688, \alpha = 3.7, m = 0.9$	$\sigma = 2.1, \alpha = 1.500$	$\sigma = 0.416$	$\sigma = 1.7, \alpha = 4.5, m = 0.4$	$\sigma = 0.688, \alpha = 0.688$	$\sigma = 0.253$	$\sigma = 0.6, \alpha = 3.85, m = 1.0$

Table 7: Fitted hyperparameters for funnel, lgcp and ion experiments.

	lr_sonar			vae			brownian			nice		
	DDS	PIS	UDMP	DDS	PIS	UDMP	DDS	PIS	UDMP	DDS	PIS	UDMP
$K = 64$	$\sigma = 0.3, \alpha = 1.650$	$\sigma = 0.253$	$\sigma = 1.15, \alpha = 1.7, m = 2.2$	$\sigma = 0.61, \alpha = 2.2$	$\sigma = 0.253$	NA	$\sigma = 0.1, \alpha = 2.35$	$\sigma = 0.084$	$\sigma = 0.115, \alpha = 4.8, m = 2.2$	$\sigma = 1.5, \alpha = 2.125$	$\sigma = 0.75$	$\sigma = 1.2, \alpha = 1.0, m = 0.9$
$K = 128$	$\sigma = 0.3, \alpha = 1.2$	$\sigma = 0.253$	$\sigma = 0.55, \alpha = 1.7, m = 3.1$	$\sigma 0.61, \alpha = 1.670$	$\sigma = 0.2523$	NA	$\sigma = 0.1, \alpha = 1.8$	$\sigma = 0.093$	$\sigma = 0.115, \alpha = 4.8, m = 2.2$	$\sigma = 1.5, \alpha = 1.75$	$\sigma = 0.588$	$\sigma = 1.2, \alpha = 1.0m = 1.65$
$K = 256$	$\sigma = 0.3, \alpha = 0.75$	$\sigma = 0.253$	$\sigma = 0.55, \alpha = 2.9, m = 2.2$	$\sigma = 0.61, \alpha = 1.140$	$\sigma = 0.506$	NA	$\sigma = 0.1, \alpha = 2.35$	$\sigma = 0.043$	$\sigma = 0.115, \alpha = 3.75, m = 2.2$	$\sigma = 1.5, \alpha = 1.75$	$\sigma = 0.425$	$\sigma = 1.2, \alpha = 2.5, m = 0.9$
$K = 512$	$\sigma = 0.3, \alpha = 0.75$	$\sigma = 0.253$	$\sigma = 0.55, \alpha = 2.9, m = 3.1$	$\sigma = 0.61, \alpha = 1.670$	$\sigma = 0.416$	NA	$\sigma = 0.1, \alpha = 1.8$	$\sigma = 0.0408$	$\sigma = 0.115, \alpha = 4.8, m = 2.2$	$\sigma = 1.5, \alpha = 2.5$	$\sigma = 0.263$	$\sigma = 1.2, \alpha = 2.5, m = 1.65$

Table 8: Fitted hyperparameters for sonar, brownian, vae and nice experiments.