Low-Rank Extragradient Method for Nonsmooth and Low-Rank Matrix Optimization Problems

Abstract

Low-rank and nonsmooth matrix optimization problems capture many fundamental tasks in statistics, machine learning. While significant progress has been made in recent years in developing efficient methods for *smooth* low-rank optimization problems that avoid maintaining high-rank matrices and computing expensive high-rank SVDs, advances for nonsmooth problems have been slow paced.

In this paper we consider standard convex relaxations for such problems. Mainly, we prove that under a natural *generalized strict complementarity* condition and under the relatively mild assumption that the nonsmooth objective can be written as a maximum of smooth functions, the *extragradient method*, when initialized with a "warm-start" point, converges to an optimal solution with rate $O(1/t)$ while requiring only two *low-rank* SVDs per iteration. We give a precise trade-off between the rank of the SVDs required and the radius of the ball in which we need to initialize the method. We support our theoretical results with empirical experiments on several nonsmooth low-rank matrix recovery tasks, demonstrating the correct convergence of the method from simple initializations and using SVDs with rank that matches that of the optimal (low-rank) solution.

1 Introduction

Low-rank and nonsmooth matrix optimization problems have many important applications in statistics, machine learning, and related fields, such as *sparse PCA* [20, 33], *robust PCA* [27, 32, 2, 9, 37], *phase synchronization* [40, 6, 28], *community detection and stochastic block models* [1] [1], *low-rank and sparse covariance matrix recovery* [34], *robust matrix completion* [21, 10], and more. For many of these problems, convex relaxations, in which one replaces the nonconvex low-rank constraint with a trace-norm constraint, have been demonstrated in numerous papers to be highly effective both in theory (under suitable assumptions) and empirically (see references above). These convex relaxations can be formulated as the following general nonsmooth optimization problem:

$$\min_{\mathbf{X} \in \mathcal{S}_n} g(\mathbf{X}), \tag{1}$$

where $g : \mathbb{S}^n \to \mathbb{R}$ is convex but nonsmooth, and $\mathcal{S}_n = \{\mathbf{X} \in \mathbb{S}^n \mid \mathrm{Tr}(\mathbf{X}) = 1, \ \mathbf{X} \succeq 0\}$ is the spectrahedron in $\mathbb{S}^n$, $\mathbb{S}^n$ being the space of $n \times n$ real symmetric matrices.

Problem (1), despite being convex, is notoriously difficult to solve in large scale. The simplest and most general approach applicable to it is the *projected subgradient method* [3, 7], which requires on each iteration to compute a Euclidean projection onto the spectrahedron $\mathcal{S}_n$, which in worst case

---

[1] in [40, 6, 28] and [1] the authors consider SDPs with linear objective function and affine constraints of the form $\mathcal{A}(\mathbf{X}) = \mathbf{b}$. By incorporating the linear constraints into the objective function via a ℓ_2 penalty term of the form $\lambda \|\mathcal{A}(\mathbf{X}) - \mathbf{b}\|_2, \lambda > 0$, we obtain a nonsmooth objective function.

amounts to $O(n^3)$ runtime per iteration. In many applications $g(\mathbf{X})$ follows a composite model, i.e., $g(\mathbf{X}) = h(\mathbf{X}) + w(\mathbf{X})$, where $h(\cdot)$ is convex and smooth and $w(\cdot)$ is convex and nonsmooth but admits a simple structure (e.g., nonsmooth regularizer). For such composite objectives, without the spectrahedron constraint, proximal methods such as FISTA [4] or splitting methods such as ADMM [31] are often very effective. However, with the spectrahderon constraint, all such methods require on each iteration to apply a subprocedure (e.g., computing the proximal mapping) which in worst case amounts to at least $O(n^3)$ runtime. A third type of off-the-shelf methods include those which are based on the *conditional gradient method* and adapted to nonsmooth problems, see for instance [30, 18, 35, 25]. The advantage of such methods is that no expensive high-rank SVD computations are needed. Instead, only a single leading eigenvector computation (i.e., a rank-one SVD) per iteration is required. However, whenever the number of iterations is not small, these methods still require to store high-rank matrices in memory, even when the optimal solution is low-rank. Thus, to conclude, standard first-order methods for Problem (1) require in worst case $\Omega(n^3)$ runtime per iteration and/or to store high-rank matrices in memory.

In the recent works [16, 17], it was established that for smooth objective functions, the high-rank SVD computations required for Euclidean projections onto the spectrahedron in standard gradient methods can be replaced with low-rank SVDs in the close proximity of a low-rank optimal solution. These results depend on the existence of eigen-gaps in the gradient of the optimal solution, which we refer to as a *generalized strict complementarity condition*. These results also hinge on a unique property of the Euclidean projection onto the spectrahedron. The projection onto the spectrahedron of a matrix $\mathbf{X} \in \mathbb{S}^n$, which admits an eigen-decomposition $\mathbf{X} = \sum_{i=1}^{n} \lambda_i \mathbf{v}_i \mathbf{v}_i^\top$, is given by

$$\Pi_{\mathcal{S}_n}[\mathbf{X}] = \sum_{i=1}^{n} \max\{0, \lambda_i - \lambda\} \mathbf{v}_i \mathbf{v}_i^\top, \tag{2}$$

where $\lambda \in \mathbb{R}$ is the unique scalar satisfying $\sum_{i=1}^{n} \max\{0, \lambda_i - \lambda\} = 1$. This operation thus truncates all the eigenvalues that are smaller than λ, while leaving the eigenvectors unchanged, thereby returning a matrix with rank equal to the number of eigenvalues greater than λ. Importantly, when the projection of $\mathbf{X}$ onto $\mathcal{S}_n$ is of rank r, only the first r components in the eigen-decomposition of $\mathbf{X}$ are required to compute it in the first place, and thus, only a rank-r SVD of $\mathbf{X}$ is required. In other words and simplifying, [16, 17] show that under strict complementary, at the proximity of an optimal solution of rank r, the exact Euclidean projection equals the rank-r truncated projection given by:

$$\widehat{\Pi}_{\mathcal{S}_n}^r[\mathbf{X}] := \Pi_{\mathcal{S}_n}\left[\sum_{i=1}^{r} \lambda_i \mathbf{v}_i \mathbf{v}_i^\top\right]. \tag{3}$$

Extending the results of [16, 17] to the nonsmooth setting is difficult since the smoothness assumption is critical to the analysis. Moreover, while [16, 17] rely on certain eigen-gaps in the gradients at optimal points, for nonsmooth problems, since the subdifferential set is often not a singleton, it is not likely that a similar eigen-gap property holds for all subgradients of an optimal solution.

In this paper we show that under the mild assumption that Problem (1) can be formulated as a smooth convex-concave saddle-point problem, i.e., the nonsmooth term can be written as a maximum over (possibly infinite number of) smooth convex functions, we can obtain results in the the spirit of [16, 17]. Concretely, we show that if a generalized strict complementarity (GSC) assumption holds for a low-rank optimal solution, the *extragradient method* for smooth convex-concave saddle-point problems [22, 29] (see Algorithm 1 below), when initialized in the proximity of the optimal solution, converges with its original convergence rate of $O(1/t)$, while requiring only two low-rank SVDs per iteration[2]. It is important to recall that while the extragradient method requires two SVDs per iteration, it has the benefit of a fast $O(1/t)$ convergence rate, while simpler saddle-point methods such as mirror-descent-based only achieve a $O(1/\sqrt{t})$ rate [7].

Our contributions can be summarized as follows:

- We prove that even under (standard) strict complementarity, the projected subgradient method, when initialized with a "warm-start", may produce iterates with rank higher than

---

[2]note that the extradgradient method computes two projected-gradient steps on each iteration, and thus two SVDs are needed per iteration.

that of the optimal solution. This phenomena further motivates our saddle-point approach. See Lemma 5.

- We suggest a generalized strict complementarity (GSC) condition for saddle-point problems and prove that when $g(\cdot)$ admits a highly popular saddle-point structure (one which captures all applications we mentioned in this paper), GSC for an optimal solution to Problem (1) implies GSC (with the same parameters) for a corresponding optimal solution of the equivalent saddle-point problem (the other direction always holds). See Section 3.

- **Main result:** we prove that for a smooth convex-concave saddle-point problem and an optimal solution which satisfies GSC, the extragradient method, when initialized with a "warm-start", converges with rate $O(1/t)$ while requiring only two low-rank SVDs per iteration. Moreover, we prove GSC facilitates a precise and powerful tradeoff: increasing the rank of SVD computations (beyond the rank of the optimal solution) can significantly increase the radius of the ball in which the method needs to be initialized. See Theorem 1.

- We present extensive numerical evidence that demonstrate both the plausibility of the GSC assumption in various tasks, and more importantly, demonstrate that indeed the extragradient method with simple initialization converges correctly (i.e., produces exactly the same sequences of iterates) when only rank($\mathbf{X}^*$)-SVDs are used to compute the projections (as in (3)), instead of full-rank SVDs, where $\mathbf{X}^*$ is an optimal solution. See Section 5.

1.1 Additional related work

Since, as in the works [16, 17] mentioned before which deal with smooth objectives, strict complementarity plays a key role in our analysis, we refer the interested reader also to the recent works [15, 38, 13, 19] which also exploit this property for efficient smooth and convex optimization over the spectrahedron.

Besides convex relaxations such as Problem (1), considerable advances have been made in the past several yeas in developing efficient *nonconvex* methods with global convergence guarantees for low-rank matrix problems. In [36] the authors consider semidefinite programs and prove that under a smooth manifold assumption on the constraints, such methods converge to the optimal global solution. In [23] the authors prove global convergence of factorized nonconvex gradient descent from a "warm-start" initialization point for non-linear smooth minimization on the positive semidefinite cone. Very recently, [8] has established, under statistical conditions, fast convergence results from "warm-start" initialization of nonconvex first-order methods, when applied to nonsmooth nonconvex matrix recovery problems which are based on the explicit factorization of the low-rank matrix. A result of similar flavour concerning nonsmooth and nonconvex formulation of robust recovery of low-rank matrices from random linear measurements was presented in [24]. Finally, several recent works have considered nonconvex low-rank regularizers which result in nonconvex nonsmooth optimization problems, but guarantee convergence only to a stationary point [26, 39].

2 Strict complementarity for nonsmooth optimization and difficulty of applying low-rank projected subgradient steps

The analysis of the nonsmooth Problem (1) naturally depends on certain subgradients of an optimal solution which, in many aspects, behave like the gradients of smooth functions. The existence of such a subgradient is guaranteed from the first-order optimality condition for constrained convex minimization problems:

Lemma 1 (first-order optimality condition, see [3]). *Let $g : \mathbb{S}^n \to \mathbb{R}$ be a convex function. Then $\mathbf{X}^* \in \mathcal{S}_n$ minimizes g over $\mathcal{S}_n$ if and only if there exists a subgradient $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ such that $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \geq 0$ for all $\mathbf{X} \in \mathcal{S}_n$.*

For some some $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ which satisfies the first-order optimality condition for an optimal solution $\mathbf{X}^*$, if the multiplicity of the smallest eigenvalue equals $r^* = \text{rank}(\mathbf{X}^*)$, then it can be shown that the optimal solution satisfies a strict complementarity assumption. The equivalence between a standard strict complementarity assumption of some low-rank optimal solution of a smooth optimization problem over the spectrahedron and an eigen-gap in the gradient of the optimal solution was established in [38]. We generalize this equivalence to also include nonsmooth problems. The proof follows similar arguments and is given in Appendix A.1.

Definition 1 (strict complementarity). *An optimal solution $\mathbf{X}^* \in \mathcal{S}_n$ of rank r^* for Problem (1) satisfies the strict complementarity assumption with parameter $\delta > 0$, if there exists an optimal solution of the dual problem[3] $(\mathbf{Z}^*, s^*) \in \mathbb{S}^n \times \mathbb{R}$ such that $\text{rank}(\mathbf{Z}^*) = n - r^*$, and $\lambda_{n-r^*}(\mathbf{Z}^*) \geq \delta$.*

Lemma 2. *Let $\mathbf{X}^* \in \mathcal{S}_n$ be a rank-r^* optimal solution to Problem (1). $\mathbf{X}^*$ satisfies the (standard) strict complementarity assumption with parameter $\delta > 0$ if and only if there exists a subgradient $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ such that $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \geq 0$ for all $\mathbf{X} \in \mathcal{S}_n$ and $\lambda_{n-r^*}(\mathbf{G}^*) - \lambda_n(\mathbf{G}^*) \geq \delta$.*

Throughout this paper we assume a weaker and more general assumption than strict complementarity. Namely, we assume generalized strict complementarity (GSC) holds with respect to at least one subgradient of the optimal solution for which the first-order optimality condition holds.

Assumption 1 (generalized strict complementarity). *We say an optimal solution $\mathbf{X}^*$ to Problem (1) satisfies the generalized strict complementarity assumption with parameters r, δ, if there exists a subgradient $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ such that $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \geq 0$ for all $\mathbf{X} \in \mathcal{S}_n$ and $\lambda_{n-r}(\mathbf{G}^*) - \lambda_n(\mathbf{G}^*) \geq \delta$.*

In [16] the author presents several characteristic properties of the gradient of the optimal solution in optimization problems over the spectrahedron. Using the existence of subgradients which satisfy the condition in Lemma 1, we can extend these properties also to the nonsmooth setting. The following lemma shows that GSC with parameters (r, δ) for some $\delta > 0$ (Assumption 1) is a sufficient condition for the optimal solution to be of rank at most r. The proof follows immediately from the proof of the analogous Lemma 7 in [16], by replacing the gradient of the optimal solution with a subgradient for which the first-order optimality condition holds.

Lemma 3. *Let $\mathbf{X}^*$ be an optimal solution to Problem (1) and write its eigen-decomposition as $\mathbf{X}^* = \sum_{i=1}^{r^*} \lambda_i \mathbf{v}_i \mathbf{v}_i^T$. Then, any subgradient $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ which satisfies $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \geq 0$ for all $\mathbf{X} \in \mathcal{S}_n$, admits an eigen-decomposition such that the set of vectors $\{\mathbf{v}_i\}_{i=1}^{r^*}$ is a set of leading eigenvectors of $(-\mathbf{G}^*)$ which corresponds to the eigenvalue $\lambda_1(-\mathbf{G}^*) = -\lambda_n(\mathbf{G}^*)$. Furthermore, there exists at least one such subgradient.*

One motivation for assuming (standard) strict complementarity (Assumption 1 with parameters $r = \text{rank}(\mathbf{X}^*)$ and $\delta > 0$) is that it guarantees a certain notion of robustness of the problem to small perturbations in the parameters. It is well known (see for instance [3]) that a projected subgradient step from $\mathbf{X}^*$ with respect to a subgradient $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ for which the the first-order optimality condition holds, returns the optimal solution $\mathbf{X}^*$ itself. This implies that $\text{rank}\left(\Pi_{\mathcal{S}_n}[\mathbf{X}^* - \eta \mathbf{G}^*]\right) = \text{rank}(\mathbf{X}^*)$. Without (standard) strict complementarity however, a small change in the parameters could result in a higher rank matrix. This is captured in the following lemma which is analogous to Lemma 3 in [17], where again the proof is straightforward from the proof in [17] by replacing the gradient of the optimal solution with a subgradient for which the first-order optimality condition holds.

Lemma 4. *Let $\mathbf{X}^*$ be an optimal solution of rank r^* to Problem (1). Let $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ be a subgradient at $\mathbf{X}^*$ such that $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \geq 0$ for all $\mathbf{X} \in \mathcal{S}_n$. Then, $\lambda_{n-r^*}(\mathbf{G}^*) = \lambda_n(\mathbf{G}^*)$ if and only if for any arbitrarily small $\zeta > 0$ it holds that $\text{rank}\left(\Pi_{(1+\zeta)\mathcal{S}_n}[\mathbf{X}^* - \eta \mathbf{G}^*]\right) > r^*$, where $\eta > 0$, $(1+\zeta)\mathcal{S}_n = \{(1+\zeta)\mathbf{X} \mid \mathbf{X} \in \mathcal{S}_n\}$, and $\Pi_{(1+\zeta)\mathcal{S}_n}[\cdot]$ denotes the Euclidean projection onto the set $(1+\zeta)\mathcal{S}_n$.*

2.1 The challenge of applying low-rank projected subgradient steps

We now demonstrate the difficulty of replacing the full-rank SVD computations required in projected subgradient steps over the spectrahedron, with their low-rank SVD counterparts when attempting to solve Problem (1). We prove that a projected subgradient step from a point arbitrarily close to a low-rank optimal solution — even one that satisfies strict complementarity, may result in a higher rank matrix. The problem on which we demonstrate this phenomena is a well known convex formulation of the *sparse PCA* problem [12]. The proof is given in Appendix A.2.

Lemma 5 (failure of low-rank subgradient descent on sparse PCA). *Consider the problem $\min_{\mathbf{X} \in \mathcal{S}_n} \{g(\mathbf{X}) := -\langle \mathbf{z}\mathbf{z}^\top + \mathbf{z}_\perp \mathbf{z}_\perp^\top, \mathbf{X} \rangle + \frac{1}{2k}\|\mathbf{X}\|_1\}$, where $\mathbf{z} = (1/\sqrt{k}, \ldots, 1/\sqrt{k}, 0, \ldots, 0)^\top$ is supported on the first k entries, $\mathbf{z}_\perp = (0, \ldots, 0, 1/\sqrt{n-k}, \ldots, 1/\sqrt{n-k})^\top$ is supported on the*

---

[3]Denote $q(\mathbf{Z}, s) = \min_{\mathbf{X} \in \mathbb{S}^n} \{g(\mathbf{X}) + s(1 - \text{Tr}(\mathbf{X})) - \langle \mathbf{Z}, \mathbf{X} \rangle\}$. The dual problem of Problem (1) can be written as: $\max_{\{\mathbf{Z} \succeq 0, \ s \in \mathbb{R}\}} \{q(\mathbf{Z}, s) \mid (\mathbf{Z}, s) \in \text{dom}(q)\}$.

178 *last $n-k$ entries, and $k \le n/4$. Then, $\mathbf{z}\mathbf{z}^\top$ is a rank-one optimal solution for which strict complemen-*
179 *tarity holds. However, for any $\eta < \frac{2}{3}$ and any $\mathbf{v} \in \mathbb{R}^n$ such that $\|\mathbf{v}\| = 1$, support($\mathbf{v}$) $\subseteq$ support($\mathbf{z}$),*
180 *and $\langle \mathbf{z}, \mathbf{v} \rangle^2 = 1 - \frac{1}{2}\|\mathbf{v}\mathbf{v}^\top - \mathbf{z}\mathbf{z}^\top\|_F^2 \ge 1 - \frac{1}{2k^2}$, it holds that rank $\left(\Pi_{\mathcal{S}_n}[\mathbf{v}\mathbf{v}^\top - \eta\mathbf{G}_{\mathbf{v}\mathbf{v}^\top}]\right) > 1$,*
181 *where $\mathbf{G}_{\mathbf{v}\mathbf{v}^\top} = -\mathbf{z}\mathbf{z}^\top - \mathbf{z}_\perp\mathbf{z}_\perp^\top + \frac{1}{2k}\mathrm{sign}(\mathbf{v}\mathbf{v}^\top) \in \partial g(\mathbf{v}\mathbf{v}^\top)$.*

182 Note that the subgradient of the ℓ_1-norm which we choose for the projected subgradient step simply
183 corresponds to the sign function, which is arguably the most natural choice.

3 From nonsmooth to saddle-point problems

185 To circumvent the difficulty demonstrated in Lemma 5 in incorporating low-rank SVDs into standard
186 subgradient methods for solving Problem (1), we propose tackling the nonsmooth problem with
187 saddle-point methods.

188 We assume the nonsmooth Problem (1) can be written as a maximum of smooth functions, i.e.,
189 $g(\mathbf{X}) = \max_{\mathbf{y} \in \mathcal{K}} f(\mathbf{X}, \mathbf{y})$, where $\mathcal{K}$ is some compact and convex set onto which it is efficient to
190 compute Euclidean projections. We assume $f(\cdot, \mathbf{y})$ is convex for all $\mathbf{y} \in \mathcal{K}$ and $f(\mathbf{X}, \cdot)$ is concave
191 for all $\mathbf{X} \in \mathcal{S}_n$. That is, we rewrite Problem (1) as the following equivalent saddle-point problem:

$$\min_{\mathbf{X} \in \mathcal{S}_n} \max_{\mathbf{y} \in \mathcal{K}} f(\mathbf{X}, \mathbf{y}). \tag{4}$$

192 Finding an optimal solution to problem (4) is equivalent to finding a saddle-point $(\mathbf{X}^*, \mathbf{y}^*) \in \mathcal{S}_n \times \mathcal{K}$
193 such that for all $\mathbf{X} \in \mathcal{S}_n$ and $\mathbf{y} \in \mathcal{K}$, $f(\mathbf{X}^*, \mathbf{y}) \le f(\mathbf{X}^*, \mathbf{y}^*) \le f(\mathbf{X}, \mathbf{y}^*)$.

194 We make a standard assumption that $f(\cdot, \cdot)$ is smooth with respect to all the components. That
195 is, we assume there exist $\beta_X, \beta_y, \beta_{Xy}, \beta_{yX} \ge 0$ such that for any $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{S}_n$ and $\mathbf{y}, \tilde{\mathbf{y}} \in \mathcal{K}$ the
196 following four inequalities hold: $\|\nabla_\mathbf{X} f(\mathbf{X}, \mathbf{y}) - \nabla_\mathbf{X} f(\tilde{\mathbf{X}}, \mathbf{y})\|_F \le \beta_X \|\mathbf{X} - \tilde{\mathbf{X}}\|_F$, $\|\nabla_\mathbf{y} f(\mathbf{X}, \mathbf{y}) - $
197 $\nabla_\mathbf{y} f(\mathbf{X}, \tilde{\mathbf{y}})\|_2 \le \beta_y \|\mathbf{y} - \tilde{\mathbf{y}}\|_2$, $\|\nabla_\mathbf{X} f(\mathbf{X}, \mathbf{y}) - \nabla_\mathbf{X} f(\mathbf{X}, \tilde{\mathbf{y}})\|_F \le \beta_{Xy}\|\mathbf{y} - \tilde{\mathbf{y}}\|_2$, $\|\nabla_\mathbf{y} f(\mathbf{X}, \mathbf{y}) - $
198 $\nabla_\mathbf{y} f(\tilde{\mathbf{X}}, \mathbf{y})\|_2 \le \beta_{yX}\|\mathbf{X} - \tilde{\mathbf{X}}\|_F$, where $\nabla_\mathbf{X} f = \frac{\partial f}{\partial \mathbf{X}}$ and $\nabla_\mathbf{y} f = \frac{\partial f}{\partial \mathbf{y}}$.

199 We denote by β the full Lipschitz parameter of the gradient, that is for any $\mathbf{X}, \tilde{\mathbf{X}} \in \mathcal{S}_n$ and $\mathbf{y}, \tilde{\mathbf{y}} \in \mathcal{K}$,

$$\|(\nabla_\mathbf{X} f(\mathbf{X}, \mathbf{y}), -\nabla_\mathbf{y} f(\mathbf{X}, \mathbf{y})) - (\nabla_\mathbf{X} f(\tilde{\mathbf{X}}, \tilde{\mathbf{y}}), -\nabla_\mathbf{y} f(\tilde{\mathbf{X}}, \tilde{\mathbf{y}}))\|_F \le \beta \|(\mathbf{X}, \mathbf{Y}) - (\tilde{\mathbf{X}}, \tilde{\mathbf{y}})\|_F.$$

200 The relationship between the full Lipschitz parameter β and its components $\beta_X, \beta_y, \beta_{Xy}, \beta_{yX}$ can
201 be written as $\beta = \sqrt{2} \max\left\{\sqrt{\beta_X^2 + \beta_{yX}^2}, \sqrt{\beta_y^2 + \beta_{Xy}^2}\right\}$. See proof in Appendix B.

202 The following lemma highlights a connection between the gradient of a saddle-point of (4) and
203 subgradients of an optimal solution to (1) for which the first order optimality condition holds. One of
204 the connections we will be interested in, is that GSC for Problem (1) implies GSC (with the same
205 parameters) for Problem (4). However, to prove this specific connection we require an additional
206 structural assumption on the objective function $g(\cdot)$. We note that this assumption holds for all
207 applications mentioned in this paper.

Assumption 2. *$g(\mathbf{X})$ is of the form $g(\mathbf{X}) = h(\mathbf{X}) + \max_{\mathbf{y} \in \mathcal{K}} \mathbf{y}^\top(\mathcal{A}(\mathbf{X}) - \mathbf{b})$, where $h(\cdot)$ is smooth*
209 *and convex, and $\mathcal{A}$ is a linear map.*

Lemma 6. *If $(\mathbf{X}^*, \mathbf{y}^*)$ is a saddle-point of Problem (4) then $\mathbf{X}^*$ is an optimal solution to Problem*
211 *(1), $\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*) \in \partial g(\mathbf{X}^*)$, and for all $\mathbf{X} \in \mathcal{S}_n$ it holds that $\langle \mathbf{X} - \mathbf{X}^*, \nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*) \rangle \ge 0$.*
212 *Conversely, under Assumption 2, if $\mathbf{X}^*$ is an optimal solution to Problem (1), then there exists $\mathbf{y}^* \in$*
213 *$\arg\max_{\mathbf{y} \in \mathcal{K}} f(\mathbf{X}^*, \mathbf{y})$ such that $(\mathbf{X}^*, \mathbf{y}^*)$ is a saddle-point of Problem (4), and $\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*) = \mathbf{G}^*$*
214 *for some $\mathbf{G}^* \in \partial g(\mathbf{X}^*)$ which satisfies $\langle \mathbf{X} - \mathbf{X}^*, \mathbf{G}^* \rangle \ge 0$ for all $\mathbf{X} \in \mathcal{S}_n$.*

215 The proof is given in Appendix C.1. The connection between the gradient of an optimal solution to
216 the saddle-point problem and a subgradient of a corresponding optimal solution in the equivalent
217 nonsmooth problem established in Lemma 6, naturally leads to the formulation of the following
218 generalized strict complementarity assumption for saddle-point problems.

Assumption 3 (generalized strict complementarity for saddle-points). *We say a saddle-point*
220 *$(\mathbf{X}^*, \mathbf{y}^*) \in \mathcal{S}_n \times \mathcal{K}$ of Problem (4) with $\mathrm{rank}(\mathbf{X}^*) = r^*$ satisfies the generalized strict complementar-*
221 *ity assumption with parameters $r \ge r^*, \delta > 0$, if $\lambda_{n-r}(\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*)) - \lambda_n(\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*)) \ge \delta$.*

Remark 1. *Note that under Assumption 2, due to Lemma 6, GSC with parameters r, δ for some optimal solution $\mathbf{X}^*$ to Problem (1) implies GSC with parameters r, δ to a corresponding saddle-point $(\mathbf{X}^*, \mathbf{y}^*)$ of Problem (4). Nevertheless, Assumption 2 is not necessary for proving our convergence results for Problem (4), which are directly stated in terms of Assumption 3.*

4 Projected extragradient method with low-rank projections

In this section we formally state and prove our main result: the projected extragradient method for the saddle-point Problem (4), when initialized in the proximity of a saddle-point which satisfies GSC (Assumption 3), converges with its original $O(1/t)$ rate while requiring only two low-rank SVD computations per iteration.

Algorithm 1 Projected extragradient method for saddle-point problems (see also [22, 29])

Input: sequence of step-sizes $\{\eta_t\}_{t \geq 1}$
Initialization: $(\mathbf{X}_1, \mathbf{y}_1) \in \mathcal{S}_n \times \mathcal{K}$
for $t = 1, 2, ...$ **do**
$\quad \mathbf{Z}_{t+1} = \Pi_{\mathcal{S}_n}[\mathbf{X}_t - \eta_t \nabla_{\mathbf{X}} f(\mathbf{X}_t, \mathbf{y}_t)]$
$\quad \mathbf{w}_{t+1} = \Pi_{\mathcal{K}}[\mathbf{y}_t + \eta_t \nabla_{\mathbf{y}} f(\mathbf{X}_t, \mathbf{y}_t)]$
$\quad \mathbf{X}_{t+1} = \Pi_{\mathcal{S}_n}[\mathbf{X}_t - \eta_t \nabla_{\mathbf{X}} f(\mathbf{Z}_{t+1}, \mathbf{w}_{t+1})]$
$\quad \mathbf{y}_{t+1} = \Pi_{\mathcal{K}}[\mathbf{y}_t + \eta_t \nabla_{\mathbf{y}} f(\mathbf{Z}_{t+1}, \mathbf{w}_{t+1})]$
end for

Theorem 1 (main theorem). *Fix an optimal solution $(\mathbf{X}^*, \mathbf{y}^*) \in \mathcal{S}_n \times \mathcal{K}$ to Problem (4). Let $\tilde{r}$ denote the multiplicity of $\lambda_n(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*))$ and for any $r \geq \tilde{r}$ define $\delta(r) = \lambda_{n-r}(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*) - \lambda_n(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*))$. Let $\{(\mathbf{X}_t, \mathbf{y}_t)\}_{t \geq 1}$ and $\{(\mathbf{Z}_t, \mathbf{w}_t)\}_{t \geq 2}$ be the sequences of iterates generated by Algorithm 1 with a fixed step-size $\eta = \min\left\{\frac{1}{2\sqrt{\beta_X^2 + \beta_{yX}^2}}, \frac{1}{2\sqrt{\beta_y^2 + \beta_{Xy}^2}}, \frac{1}{\beta_X + \beta_{Xy}}, \frac{1}{\beta_y + \beta_{yX}}\right\}$. Assume the initialization $(\mathbf{X}_1, \mathbf{y}_1)$ satisfies $\|(\mathbf{X}_1, \mathbf{y}_1) - (\mathbf{X}^*, \mathbf{y}^*)\|_F \leq R_0(r)$, where*

$$R_0(r) := \frac{\eta}{(2 + \sqrt{2})\left(1 + (2 + \sqrt{2})\eta \max\{\beta_X, \beta_{Xy}\}\right)} \max\left\{\frac{\sqrt{\tilde{r}}\delta(r - \tilde{r} + 1)}{2}, \frac{\delta(r)}{(1 + 1/\sqrt{\tilde{r}})}\right\}.$$

Then, for all $t \geq 1$, the projections $\Pi_{\mathcal{S}_n}[\mathbf{X}_t - \eta\nabla_{\mathbf{X}} f(\mathbf{X}_t, \mathbf{y}_t)]$ and $\Pi_{\mathcal{S}_n}[\mathbf{X}_t - \eta\nabla_{\mathbf{X}} f(\mathbf{Z}_{t+1}, \mathbf{w}_{t+1})]$ can be replaced with their rank-r truncated counterparts (see (3)) without changing the sequences $\{(\mathbf{X}_t, \mathbf{y}_t)\}_{t \geq 1}$ and $\{(\mathbf{Z}_t, \mathbf{w}_t)\}_{t \geq 2}$, and for any $T \geq 0$ it holds that

$$\min_{t \in [T]} \max_{\mathbf{y} \in \mathcal{K}} f(\mathbf{Z}_{t+1}, \mathbf{y}) - \max_{t \in [T]} \min_{\mathbf{X} \in \mathcal{S}_n} f(\mathbf{X}, \mathbf{w}_{t+1})$$

$$\leq \frac{D^2 \max\left\{\sqrt{\beta_X^2 + \beta_{yX}^2}, \sqrt{\beta_y^2 + \beta_{Xy}^2}, \frac{1}{2}(\beta_X + \beta_{Xy}), \frac{1}{2}(\beta_y + \beta_{yX})\right\}}{T},$$

where $D := \sup_{(\mathbf{X}, \mathbf{y}), (\mathbf{Z}, \mathbf{w}) \in \mathcal{S}_n \times \mathcal{K}} \|(\mathbf{X}, \mathbf{y}) - (\mathbf{Z}, \mathbf{w})\|_F$.

Remark 2. *Note that Theorem 1 implies that if standard strict complementarity holds, that is Assumption 3 holds with $r = r^* = \text{rank}(\mathbf{X}^*)$ and some $\delta > 0$, then only rank-r^* SVDs are required so that Algorithm 1 converges with the guaranteed convergence rate of $O(1/t)$, when initialized with a "warm-start". Furthermore, by using SVDs of rank $r > r^*$, with moderately higher values of r, we can increase the radius of the ball in which Algorithm 1 needs to be initialized quite significantly.*

The complete proof of Theorem 1 is given in Appendix D. Below we give a short sketch of the proof.

Proof sketch for Theorem 1. The convergence rate of Algorithm 1 is well known in the literature (see Appendix D.1 for completeness). Thus, we focus on the novel part of the low-rank projections. For simplicity consider the case $r = \tilde{r}$. From (2) it can be deduced that for any $\mathbf{P} \in \mathbb{S}^n$ it holds that $\text{rank}(\Pi_{\mathcal{S}_n}[\mathbf{P}]) \leq \tilde{r}$ if and only if the condition $\sum_{i=1}^{\tilde{r}} \lambda_i(\mathbf{P}) \geq 1 + \tilde{r}\lambda_{\tilde{r}+1}(\mathbf{P})$ holds. Denote $\mathbf{P}^* = \mathbf{X}^* - \eta\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*)$. From Lemma 6 and Lemma 3 it follows that

$$\forall i \leq \text{rank}(\mathbf{X}^*): \; \lambda_i(\mathbf{P}^*) = \lambda_i(\mathbf{X}^*) - \eta\lambda_n(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*));$$
$$\forall i > \text{rank}(\mathbf{X}^*): \; \lambda_i(\mathbf{P}^*) = -\eta\lambda_{n-i+1}(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*)).$$

Using this and the fact that $\lambda_{n-i+1}(\nabla f_{\mathbf{X}}(\mathbf{X}^*, \mathbf{y}^*)) = \lambda_n(\nabla f_{\mathbf{X}}(\mathbf{X}^*, \mathbf{y}^*))$ for all $i \leq \tilde{r}$ we have,

$$\sum_{i=1}^{\tilde{r}} \lambda_i(\mathbf{P}^*) = \sum_{i=1}^{r^*} \lambda_i(\mathbf{X}^*) - \eta \sum_{i=1}^{\tilde{r}} \lambda_n(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*)) = \text{Tr}(\mathbf{X}^*) - \eta \tilde{r} \lambda_n(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*))$$

$$= 1 - \eta \tilde{r} \lambda_{n-\tilde{r}}(\nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*)) + \eta \tilde{r} \delta(\tilde{r}) = 1 + \tilde{r} \lambda_{\tilde{r}+1}(\mathbf{P}^*) + \eta \tilde{r} \delta(\tilde{r}). \quad (5)$$

Thus, $\mathbf{P}^* = \mathbf{X}^* - \eta \nabla_{\mathbf{X}} f(\mathbf{X}^*, \mathbf{y}^*)$ not only satisfies the condition that ensures its projection onto $\mathcal{S}_n$ is of rank at most $\tilde{r}$, but it satisfies it with a positive slack of $\eta \tilde{r} \delta(\tilde{r})$. Also, for any $(\mathbf{X}, \mathbf{y})$ sufficiently close to $(\mathbf{X}^*, \mathbf{y}^*)$, using the smoothness of f, we have that $\mathbf{P} = \mathbf{X} - \eta \nabla_{\mathbf{X}} f(\mathbf{X}, \mathbf{y})$ is close to $\mathbf{P}^*$. Thus, by applying perturbation bounds for the eigenvalues of symmetric matrices to Eq. (5) and using the fact that the positive slack $\eta \tilde{r} \delta(\tilde{r})$ in the RHS of (5) allows to absorb sufficiently small errors, we can establish that for such points $(\mathbf{X}, \mathbf{y})$ it holds that $\text{rank}(\Pi_{\mathcal{S}_n}[\mathbf{P}]) \leq \tilde{r}$. A similar argument shows that the second primal step in Algorithm 1: $\mathbf{X}_{t+1} = \Pi_{\mathcal{S}_n}[\mathbf{X}_t - \eta \nabla \mathbf{X} f(\mathbf{Z}_{t+1}, \mathbf{w}_{t+1})]$, also results in matrices of rank at most $\tilde{r}$ when $\mathbf{Z}_t, \mathbf{w}_t$ are also in the proximity of $\mathbf{X}^*, \mathbf{y}^*$. Finally, we prove a complementary argument, that when initialized in the proximity of $(\mathbf{X}^*, \mathbf{y}^*)$, the iterates of Algorithm 1 stay in the proximity of $(\mathbf{X}^*, \mathbf{y}^*)$ throughout all iterations.

□

Remark 3. *A downside of considering the saddle-point formulation* (4) *when attempting to solve Problem* (1) *that arises from Theorem 1, is that not only do we need a "warm-start" initialization for the original primal matrix variable* $\mathbf{X}$, *in the saddle-point formulation we need a "warm-start" for the primal-dual pair* $(\mathbf{X}, \mathbf{y})$. *Nevertheless, as we demonstrate extensively in Section 5, it seems that very simple initialization schemes work very well in practice.*

4.1 Efficiently-computable certificates for correctness of low-rank projections

Since Theorem 1 only applies in some neighborhood of an optimal solution, it is of interest to have a procedure for verifying if the rank-r truncated projection of a given point indeed equals the exact Euclidean projection. In particular, from a practical point of view, it does not matter whether the conditions of Theorem 1 hold. In practice, as long as the truncated projection equals the exact projection, we are guaranteed that Algorithm 1 converges correctly with rate $O(1/t)$, without needing to verify any other condition. Luckily, the expression in (2) which characterizes the structure of the Euclidean projection onto the spectrahedron, yields exactly such a verification procedure. As already noted in [16], for any $\mathbf{X} \in \mathbb{S}^n$, we have $\widehat{\Pi}_{\mathcal{S}_n}^r[\mathbf{X}] = \Pi_{\mathbf{S}_n}[\mathbf{X}]$ if an only if the condition $\sum_{i=1}^r \lambda_i(\mathbf{X}) \geq 1 + r \cdot \lambda_{r+1}(\mathbf{X})$. holds. Note that verifying this condition simply requires increasing the rank of SVD computation by one, i.e., computing a rank-$(r+1)$ SVD of the matrix to project rather than a rank-r SVD.

5 Empirical evidence

The goal of this section is to bring empirical evidence in support of our theoretical approach. We consider various tasks that take the form of minimizing a composite objective, i.e., the sum of a smooth convex function and a nonsmooth convex function, where the nonsmoothness comes from either an ℓ_1-norm or ℓ_2-norm regularizer / penalty term, over a τ-scaled spectrahedron. In all cases the nonsmooth objective can be written as a saddle-point with function $f(\mathbf{X}, \mathbf{y})$ which is linear in $\mathbf{y}$ and in particular satisfies Assumption 2.

The tasks considered include 1. sparse PCA, 2. robust PCA, 3. low-rank and sparse recovery, 4. phase synchronization, and 5. linearly-constrained low-rank estimation, under variety of parameters. Due to lack of space many of the results are deferred to Appendix F.

For all tasks considered we generate random instances, and examine the sequences of iterates generated by Algorithm 1 $\{(\mathbf{X}_t, \mathbf{y}_t)\}_{t \geq 1}, \{(\mathbf{Z}_t, \mathbf{w}_t)\}_{t \geq 2}$, when initialized with simple initialization procedures. Out of both sequences generated, we choose our candidate for the optimal solution to be the iterate for which the dual-gap, which is a certificate for optimality, is smallest. See Appendix E.

We consider several different types of nonsmooth low-rank matrix recovery problems where the goal is to recover a ground-truth low-rank matrix $\mathbf{M}_0 \in \mathbb{S}^n$ from some noisy observation of it $\mathbf{M} = \mathbf{M}_0 + \mathbf{N}$, where $\mathbf{N} \in \mathbb{S}^n$ is a noise matrix. We measure the signal-to-noise ratio (SNR) as $\|\mathbf{M}_0\|_F^2 / \|\mathbf{N}\|_F^2$.

In all experiments we measure the relative initialization error by $\left\|\frac{\mathrm{Tr}(\mathbf{M}_0)}{\tau}\mathbf{X}_1 - \mathbf{M}_0\right\|_F^2 \Big/ \|\mathbf{M}_0\|_F^2$,
and similarly we measure the relative recovery error by $\left\|\frac{\mathrm{Tr}(\mathbf{M}_0)}{\tau}\mathbf{X}^* - \mathbf{M}_0\right\|_F^2 \Big/ \|\mathbf{M}_0\|_F^2$. Note that
in some of the experiments we take $\tau < \mathrm{Tr}(\mathbf{M}_0)$ to prevent the method from overfitting the noise.
In addition, we measure the (standard) strict complementarity parameter which corresponds to the
eigen-gap $\mathrm{gap}(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*)) := \lambda_{n-r}(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*)) - \lambda_n(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*))$, $r = \mathrm{rank}(\mathbf{M}_0)$.

In all experiments we use SVDs of rank $r = \mathrm{rank}(\mathbf{M}_0)$ to compute the projections in Algorithm
1 according to the truncated projection given in (3). To certify the correctness of these low-rank
projections (that is, that they equal the exact Euclidean projection) we confirm that the inequality
$\sum_{i=1}^r \lambda_i(\mathbf{P}_j) \geq \tau + r \cdot \lambda_{r+1}(\mathbf{P}_j)$ always holds for $\mathbf{P}_1 = \mathbf{X}_t - \eta\nabla_{\mathbf{X}}f(\mathbf{X}_t, \mathbf{Y}_t)$ and $\mathbf{P}_2 = \mathbf{X}_t - $
$\eta\nabla_{\mathbf{X}}f(\mathbf{Z}_{t+1}, \mathbf{W}_{t+1})$ (see also Section 4.1). Indeed, we can now already state our main observation
from the experiments:

> *In all tasks considered and for all random instances generated, throughout all iterations of Algorithm
> 1, when initialized with a simple "warm-start" strategy and when computing only rank-r truncated
> projections, $r = \mathrm{rank}(\mathbf{M}_0)$, the truncated projections of $\mathbf{P}_1 = \mathbf{X}_t - \eta\nabla_{\mathbf{X}}f(\mathbf{X}_t, \mathbf{Y}_t)$ and $\mathbf{P}_2 = $
> $\mathbf{X}_t - \eta\nabla_{\mathbf{X}}f(\mathbf{Z}_{t+1}, \mathbf{W}_{t+1})$ equal their exact full-rank counterparts. That is, Algorithm 1, using
> only rank-r SVDs, computed exactly the same sequences of iterates it would have computed if using
> full-rank SVDs.*

Aside from the above observation, in the sequel we demonstrate that all models considered indeed
satisfy that: 1. the returned solution is of the same rank as the ground truth matrix and satisfies
the strict complementarity condition with non-negligible parameter (measured by the eigengap
$\lambda_{n-r}(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*)) - \lambda_n(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*)))$, 2. the recovery error of the returned solution indeed
improves significantly over the error of the initialization point.

Sparse PCA: We consider the sparse PCA problem in a well known convex formulation taken
form [12] and its equivalent saddle-point formulation:

$$\min_{\substack{\mathrm{Tr}(\mathbf{X})=1, \\ \mathbf{X}\succeq 0}} \langle\mathbf{X}, -\mathbf{M}\rangle + \lambda\|\mathbf{X}\|_1 = \min_{\substack{\mathrm{Tr}(\mathbf{X})=1, \\ \mathbf{X}\succeq 0}} \max_{\|\mathbf{Y}\|_\infty \leq 1} \{\langle\mathbf{X}, -\mathbf{M}\rangle + \lambda\langle\mathbf{X}, \mathbf{Y}\rangle\},$$

where $\mathbf{M} = \mathbf{z}\mathbf{z}^\top + \frac{c}{2}(\mathbf{N} + \mathbf{N}^\top)$ is a noisy observation of a rank-one matrix $\mathbf{z}\mathbf{z}^\top$, with $\mathbf{z}$ being
a sparse unit vector. Each entry $\mathbf{z}_i$ is chosen to be 0 with probability 0.9 and $U\{1, \ldots, 10\}$ with
probability 0.1, and then we normalize $\mathbf{z}$ to be of unit norm. We test the results obtained when adding
different magnitudes of Gaussian or uniform noise. We set the signal-to-noise ratio (SNR) to be a
constant. Thus, we set the noise level to $c = \frac{2}{\mathrm{SNR}\cdot\|\mathbf{N}+\mathbf{N}^\top\|_F}$ for our choice of SNR. We initialize the
$\mathbf{X}$ variable with the rank-one approximation of $\mathbf{M}$. That is, we take $\mathbf{X}_1 = \mathbf{u}_1\mathbf{u}_1^\top$, where $\mathbf{u}_1$ is the top
eigenvector of $\mathbf{M}$. For the $\mathbf{Y}$ variable we initialize it with $\mathbf{Y}_1 = \mathrm{sign}(\mathbf{X}_1)$ which is a subgradient of
$\|\mathbf{X}_1\|_1$. We set the step-size to $\eta = 1/(2\lambda)$ and we set the number of iterations to $T = 1000$ and for
any set of parameters we average the measurements over 10 i.i.d. runs.

Table 1: Numerical results for the sparse PCA problem. $\mathbf{N}_{ij} \sim U[0, 1]$, SNR $= 0.05$.

dimension (n)	100	200	400	600
λ	0.04	0.02	0.01	0.0067
initialization error	1.7456	1.7494	1.7566	1.7625
recovery error	0.0425	0.0244	0.0149	0.0100
dual gap	2.0×10^{-9}	5.8×10^{-6}	4.5×10^{-4}	0.0018
$\mathrm{gap}(\nabla_{\mathbf{X}}f(\mathbf{X}^*, \mathbf{y}^*))$	0.7092	0.7854	0.8340	0.8622

Robust PCA: We consider the robust PCA problem [27] in the following formulation:

$$\min_{\substack{\mathrm{Tr}(\mathbf{X})=\tau, \\ \mathbf{X}\succeq 0}} \|\mathbf{X} - \mathbf{M}\|_1 = \min_{\substack{\mathrm{Tr}(\mathbf{X})=\tau, \\ \mathbf{X}\succeq 0}} \max_{\|\mathbf{Y}\|_\infty \leq 1} \langle\mathbf{X} - \mathbf{M}, \mathbf{Y}\rangle,$$

where $\mathbf{M} = r\mathbf{Z}_0\mathbf{Z}_0^\top + \frac{1}{2}(\mathbf{N} + \mathbf{N}^\top)$ is a sparsely-corrupted observation of some rank-r matrix $\mathbf{Z}_0\mathbf{Z}_0^\top$. We choose $\mathbf{Z}_0 \in \mathbb{R}^{n \times r}$ to be a random unit Frobenius norm matrix. For $\mathbf{N} \in \mathbb{R}^{n \times n}$, we choose each entry to be 0 with probability $1 - 1/\sqrt{n}$ and otherwise 1 or -1 with equal probability. We initialize the $\mathbf{X}$ variable with the projection $\mathbf{X}_1 = \Pi_{\{\mathrm{Tr}(\mathbf{X})=\tau, \, \mathbf{x} \succeq 0\}}[\mathbf{M}]$, and the $\mathbf{Y}$ variable with $\mathbf{Y}_1 = \mathrm{sign}(\mathbf{X}_1 - \mathbf{M})$. We vary the rank of $\mathbf{Z}_0$, which in turn determines the step-size η. We set the trace bound to $\tau = 0.95 \cdot \mathrm{Tr}(r\mathbf{Z}_0\mathbf{Z}_0^\top)$, and the number of iterations to $T = 30,000$. For every set of parameters we average the measurements over 10 i.i.d. runs.

Table 2: Numerical results for the robust PCA problem. $\mathrm{rank}(\mathbf{Z}_0\mathbf{Z}_0^\top) = 10$, step-size $\eta = 1$.

dimension (n)	100	200	400	600
SNR	0.0229	0.0077	0.0026	0.0014
initialization error	1.5729	1.6485	1.6317	1.5949
recovery error	0.0079	0.0081	0.0073	0.0065
dual gap	0.0139	0.0338	0.1533	0.3561
gap($\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*)$)	1.7945	16.9890	48.9799	82.2727

Linearly-constrained low-rank estimation: Consider the following penalized formulation:

$$\min_{\substack{\mathrm{Tr}(\mathbf{X})=1, \\ \mathbf{X} \succeq 0}} \langle \mathbf{X}, -\mathbf{M} \rangle + \lambda \|\mathcal{A}(\mathbf{X}) - \mathbf{b}\|_2 = \min_{\substack{\mathrm{Tr}(\mathbf{X})=1, \\ \mathbf{X} \succeq 0}} \max_{\|\mathbf{y}\|_2 \leq 1} \langle \mathbf{X}, -\mathbf{M} \rangle + \lambda \langle \mathcal{A}(\mathbf{X}) - \mathbf{b}, \mathbf{y} \rangle,$$

where $\mathbf{M} = \mathbf{z}_0\mathbf{z}_0^\top + \frac{c}{2}(\mathbf{N} + \mathbf{N}^\top)$ is they noisy observation of some rank-one matrix $\mathbf{z}_0\mathbf{z}_0^\top$ such that $\|\mathbf{z}_0\|_2 = 1$ and the noise matrix is chosen $\mathbf{N} \sim \mathcal{N}(0, \mathbf{I}_n)$. We take $\mathcal{A}(\mathbf{X}) = (\langle \mathbf{A}_i, \mathbf{X} \rangle, \ldots, \langle \mathbf{A}_i, \mathbf{X} \rangle)^\top$ with matrices $\mathbf{A}_1, \ldots, \mathbf{A}_m \in \mathbb{S}^n$ of the form $\mathbf{A}_i = \mathbf{v}_i\mathbf{v}_i^\top$ such that $\mathbf{v}_i \sim \mathcal{N}(0, 1)$. We take $\mathbf{b} \in \mathbb{R}^m$ such that $b_i = \langle \mathbf{A}_i, \mathbf{z}_0\mathbf{z}_0^\top \rangle$. We initialize the $\mathbf{X}$ variable with the rank-one approximation of $\mathbf{M}$. That is, we take $\mathbf{X}_1 = \mathbf{u}_1\mathbf{u}_1^\top$, where $\mathbf{u}_1$ is the top eigenvector of $\mathbf{M}$. The $\mathbf{y}$ variable is initialized with $\mathbf{y}_1 = (\mathcal{A}(\mathbf{X}_1) - \mathbf{b})/\|\mathcal{A}(\mathbf{X}_1) - \mathbf{b}\|_2$. We set the number of constraints to $m = n$, the penalty parameter to $\lambda = 2$, and the step-size to $\eta = 1/(2\lambda)$. We set the number of iterations in each experiment to $T = 2000$ and for each value of n we average the measurements over 10 i.i.d. runs.

6 Discussion

This work expands upon a line of research that aims to harness the ability of convex relaxations to produce low-rank and high-quality solutions to important low-rank matrix optimization problems, while insisting on methods that, at least locally, store and manipulate only low-rank matrices. Focusing on the challenging case of nonsmooth objective functions and following our evidence for the difficulties of obtaining such a result for subgradient methods (Lemma 5), we consider tackling nonsmooth objectives via saddle-point formulations. We prove that indeed under a generalized strict complementarity condition, a state-of-the-art method for convex-concave saddle-point problems converges locally while storing and manipulating only low-rank matrices. Extensive experiments over several tasks demonstrate that our conceptual approach of utilizing low-rank projections for more efficient optimization is not only of theoretical merit, but indeed seems to work well in practice.

Table 3: Numerical results for the linearly-constrained low-rank matrix estimation problem.

dimension (n)	100	200	400	600
SNR	0.15	0.075	0.04	0.027
initialization error	0.1219	0.1324	0.1242	0.1228
recovery error	0.0437	0.0617	0.0685	0.0735
dual gap	5.3×10^{-11}	5.0×10^{-12}	8.5×10^{-12}	2.3×10^{-11}
gap($\nabla_\mathbf{X} f(\mathbf{X}^*, \mathbf{y}^*)$)	0.2941	0.3409	0.4690	0.5069
$\|\mathcal{A}(\mathbf{X}^*) - \mathbf{b}\|_2$	0.0080	0.0082	0.0079	0.0073

References

[1] Emmanuel Abbe. Community detection and stochastic block models: recent developments. *The Journal of Machine Learning Research*, 18(1):6446–6531, 2017.

[2] Praneeth Netrapalli, UN Niranjan, Sujay Sanghavi, Animashree Anandkumar and Prateek Jain. Non-convex robust pca. *Advances in Neural Information Processing Systems*, page 1107–1115, 2014.

[3] Amir Beck. First-order methods in optimization. *SIAM*, 2017.

[4] Amir Beck and Marc Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM journal on imaging sciences*, 2(1):183–202, 2009.

[5] Dimitri P Bertsekas. Nonlinear programming. *Athena Scientific*, 1999.

[6] Afonso S. Bandeira, Nicolas Boumal, and Amit Singer. Tightness of the maximum likelihood semidefinite relaxation for angular synchronization. *Mathematical Programming*, 163(1-2):145–167, 2017.

[7] Sébastien Bubeck. Convex optimization: Algorithms and complexity. *Foundations and Trends in Machine Learning*, 8(3-4):231–357, 2015.

[8] Vasileios Charisopoulos, Yudong Chen, Damek Davis, Mateo Díaz, Lijun Ding, and Dmitriy Drusvyatskiy. Low-rank matrix recovery with composite optimization: good conditioning and rapid convergence. *Foundations of Computational Mathematics*, pages 1–89, 2021.

[9] PXinyang Yi, Dohyung Park, Yudong Chen and Constantine Caramanis. Fast algorithms for robust pca via gradient descent. *Advances in Neural Information Processing Systems*, page 4152–4160, 2016.

[10] Yudong Chen, Huan Xu, Constantine Caramanis, and Sujay Sanghavi. Robust matrix completion and corrupted columns. In *Proceedings of the 28th International Conference on Machine Learning (ICML-11)*, pages 873–880. Citeseer, 2011.

[11] Frank H. Clarke. *Optimization and Nonsmooth Analysis*. SIAM, 1990.

[12] Alexandre d'Aspremont, Laurent El Ghaoui, Michael I Jordan, and Gert RG Lanckriet. A direct formulation for sparse pca using semidefinite programming. *SIAM review*, 49(3):434–448, 2007.

[13] Lijun Ding, Jicong Fan, and Madeleine Udell. $k$fw: A frank-wolfe style algorithm with stronger subproblem oracles, 2020.

[14] Francisco Facchinei and Jong-Shi Pang. *Finite-Dimensional Variational Inequalities and Complementarity Problems*, volume II. Springer-Verlag New York, 2003.

[15] Dan Garber. Linear convergence of frank-wolfe for rank-one matrix recovery without strong convexity. *arXiv preprint arXiv:1912.01467*, 2019.

[16] Dan Garber. On the convergence of projected-gradient methods with low-rank projections for smooth convex minimization over trace-norm balls and related problems. *SIAM Journal on Optimization*, 2019.

[17] Dan Garber. On the convergence of stochastic gradient descent with low-rank projections for convex low-rank matrix problems. *Conference on Learning Theory, COLT*, 125:1666–1681, 2020.

[18] Dan Garber and Atara Kaplan. Fast stochastic algorithms for low-rank and nonsmooth matrix problems. *The 22nd International Conference on Artificial Intelligence and Statistics, AISTATS*, 89:286–294, 2019.

[19] Dan Garber and Atara Kaplan. On the efficient implementation of the matrix exponentiated gradient algorithm for low-rank matrix optimization. *arXiv preprint arXiv:2012.10469*, 2020.

[20] Hui Zou, Trevor Hastie, and Robert Tibshirani. Sparse principal component analysis. *Journal of Computational and Graphical Statistics*, 15:2006, 2004.

[21] Olga Klopp, Karim Lounici, and Alexandre B Tsybakov. Robust matrix completion. *Probability Theory and Related Fields*, 169(1):523–564, 2017.

[22] G.M. Korpelevich. The extragradient method for finding saddle points and other problems. *Matecon*, 12:747–756, 1976.

[23] Srinadh Bhojanapalli, Anastasios Kyrillidis, and Sujay Sanghavi. Dropping convexity for faster semi-definite optimization. In *29th Annual Conference on Learning Theory*, volume 49 of *Proceedings of Machine Learning Research*, pages 530–582, 2016.

[24] Xiao Li, Zhihui Zhu, Anthony Man-Cho So, and Rene Vidal. Nonconvex robust low-rank matrix recovery. *SIAM Journal on Optimization*, 30(1):660–686, 2020.

[25] Francesco Locatello, Alp Yurtsevert, Olivier Fercoq, and Volkan Cevhert. Stochastic frank-wolfe for composite convex minimization. *Advances in Neural Information Processing Systems*, 32, 2019.

[26] Canyi Lu, Jinhui Tang, Shuicheng Yan, and Zhouchen Lin. Generalized nonconvex nonsmooth low-rank minimization. In *Proceedings of the IEEE conference on computer vision and pattern recognition*, pages 4130–4137, 2014.

[27] Emmanuel J. Candès, Xiaodong Li, Yi Ma and John Wright. Robust principal component analysis? *Journal of the ACM*, 58, 2009.

[28] Adel Javanmard, Andrea Montanari, and Federico Ricci-Tersenghi. Phase transitions in semidefinite relaxations. *Proceedings of the National Academy of Sciences*, 113(16):E2218–E2223, 2016.

[29] Arkadi Nemirovski. Prox-method with rate of convergence o(1/t) for variational inequalities with lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM J. on Optimization*, 15(1):229–251, 2005.

[30] Gergely Odor, Yen-Huan Li, Alp Yurtsever, Ya-Ping Hsieh, Quoc Tran-Dinh, Marwa El Halabi, and Volkan Cevher. Frank-wolfe works for non-lipschitz continuous gradient objectives: scalable poisson phase retrieval. In *2016 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 6230–6234. Ieee, 2016.

[31] Stephen Boyd, Neal Parikh, Eric Chu, Borja Peleato, and Jonathan Eckstein. Distributed optimization and statistical learning via the alternating direction method of multipliers. *Foundations and Trends in Machine Learning*, 3(1):1–122, 2011.

[32] John Wright, Arvind Ganesh, Shankar Rao, Yigang Peng and Yi Ma. Robust principal component analysis: Exact recovery of corrupted low-rank matrices via convex optimization. *Advances in Neural Information Processing Systems 22*, pages 2080–2088, 2009.

[33] Julien Mairal, Francis Bach, Jean Ponce, and Guillermo Sapiro. Online learning for matrix factorization and sparse coding. *Journal of Machine Learning Research*, 11:19–60, 2010.

[34] Emile Richard, Pierre-Andr'e Savalle and Nicolas Vayatis. Estimation of simultaneously sparse and low rank matrices. *Proceedings of the 29th International Conference on Machine Learning*, 2012.

[35] Kiran K Thekumparampil, Prateek Jain, Praneeth Netrapalli, and Sewoong Oh. Projection efficient subgradient method and optimal nonsmooth frank-wolfe method. In *Advances in Neural Information Processing Systems*, volume 33, pages 12211–12224. Curran Associates, Inc., 2020.

[36] Nicolas Boumal, Vladislav Voroninski, and Afonso S. Bandeira. Deterministic guarantees for burer-monteiro factorizations of smooth semidefinite programs. *Communications on Pure and Applied Mathematics*, 73(3):581 – 608, 2020.

[37] Cun Mu, Yuqian Zhang, John Wright and Donald Goldfarb. Scalable robust matrix recovery: Frank-wolfe meets proximal methods. *SIAM Journal on Scientific Computing*, 38(5):A3291–A3317, 2016.

[38] Lijun Ding, Yingjie Fei, Qiantong Xu and Chengrun Yang. Spectral frank-wolfe algorithm: Strict complementarity and linear convergence. *ICML*, 2020.

[39] Quanming Yao, James T Kwok, Taifeng Wang, and Tie-Yan Liu. Large-scale low-rank matrix learning with nonconvex regularizers. *IEEE transactions on pattern analysis and machine intelligence*, 41(11):2628–2643, 2018.

[40] Yiqiao Zhong and Nicolas Boumal. Near-optimal bounds for phase synchronization. *SIAM Journal on Optimization*, 28(2):989–1016, 2018.