A VARIATIONAL CONDITION FOR
MINIMAL-RESIDUAL LATENT REPRESENTATIONS

ABSTRACT

Autoencoders are a useful unsupervised-learning architecture that can be used to
build surrogate models of systems governed by partial differential equations. In
this article, we address two key questions underpinning this procedure: whether
the reconstructed output satisfies the partial differential equation, and whether
other latent vectors not corresponding to the encoding of any training data sat-
isfy the same equation. Our results spell out some relevant conditions, and clarify
the different impact of three main design decisions (architecture, training criterion,
and choice of training solutions) on the final result.

1 INTRODUCTION

The use of neural networks for modelling partial differential equation (PDE) solutions is a rapidly ex-
panding branch in scientific machine learning (see, among others, Sirignano & Spiliopoulos (2018);
Raissi et al. (2017); Li et al. (2020)). One of the architectures used for the task is autoencoders (AEs,
Takeishi & Kalousis (2021)), as they allow for unsupervised training and only require a sample of
solutions to produce a surrogate model for the PDE.

Basic AEs, however, are not physics informed, and there is no mechanism to ensure that the recon-
structed solutions satisfy the original equation to an acceptable degree. Whilst a good reconstruction
error guarantees that the AE output is close to the input in some norm, the constraint it places on
the residual of the reconstructed solution is unclear. Furthermore, using the network's decoder as
a general emulator presupposes that sampling other latent points not in the image of the training
set also leads to acceptable solutions to the equation; the conditions under which this holds true are
also presently unclear. In this paper, we formulate answers to both questions using a variational
approach.

2 SETUP

We consider an AE mapping a family of inputs $\{\bar{\boldsymbol{u}}\}_k$ to the outputs:

$$\bar{\boldsymbol{v}} \equiv \boldsymbol{v}(\bar{\boldsymbol{u}}; \boldsymbol{\theta}) = \boldsymbol{d}(\boldsymbol{e}(\bar{\boldsymbol{u}}; \boldsymbol{\theta_e}); \boldsymbol{\theta_d}) = \boldsymbol{d}(\bar{z}; \boldsymbol{\theta_d}) \tag{1}$$

where $\boldsymbol{e}$ and $\boldsymbol{d}$ represent the AE's encoder and decoder section, respectively. We study the scenario
when the $\{\bar{\boldsymbol{u}}\}_k$ approximate solutions of a partial differential equation (PDE):

$$O(u, \nabla u) = 0 \tag{2}$$

i.e. the components $\bar{u}^i$ of $\bar{\boldsymbol{u}}$ can be written as:

$$\bar{u}^i = \bar{u}(x^i) + \epsilon_d h^i \tag{3}$$

where $\bar{u}(x)$ is a solution of (2) and the second term represents the *discretization error* arising from
the representation of $\bar{u}$ as a vector $\bar{\boldsymbol{u}}$ (such as any round-off error or truncation error from the nu-
merical integration of (2)), and $\epsilon_d \ll 1$ typically.

The AE is trained via the requirement that the *reconstruction error*, or the difference between $\bar{\boldsymbol{u}}$ and
the corresponding $\bar{\boldsymbol{v}}$, is as low as possible, so that we can write:

$$\bar{\boldsymbol{v}} = \bar{\boldsymbol{u}} + \epsilon_r \hat{\boldsymbol{u}} \tag{4}$$

where again ϵ_r should be much smaller than one.

If we define the equation residual as:

$$R[u] = ||O(u, \nabla u)||_2 \tag{5}$$

as in equation (12), and introduce $R_d[\boldsymbol{u}]$ as its approximation on discrete data (obtained, e.g., using numerical derivatives), the two questions we aim to answer can then be cast in the form:

1. How much is $R_d[\bar{\boldsymbol{v}}]$, i.e. to what extent do the $\{\bar{\boldsymbol{v}}\}_k$ approximate valid solutions?

2. How much is $R_d[\boldsymbol{d}(\boldsymbol{z} \notin \bar{\boldsymbol{z}}; \boldsymbol{\theta_d})]$, i.e. can the network generate approximations to novel solutions? In other words, how much of the latent space encodes solutions of equation (2)?

The answers to the above questions are a crucial requirement in the process of using the AE as a surrogate solver for equation (2).

3 RESULTS

Using the expression for the second variation of $R^2[u]$ discussed in the Appendix, we can prove the following theorem.

Theorem 1: If $\epsilon_d \ll \epsilon_r \ll 1$, the residual of the reconstructed solutions, $R_d[\bar{\boldsymbol{v}}]$, can be written as:

$$R_d^2[\bar{\boldsymbol{v}}] = \epsilon_r^2 \sum_i a_i \left[\frac{\partial^2 s}{\partial u^2} + 2 \sum_n D^n \frac{\partial s}{\partial u} + \sum_{n,m} D^m D^n s \right]_{u=u_i} (\hat{u}^i)^2 + O(\epsilon_r^3) \tag{6}$$

where $\sum_i a_i f^i$ represents any suitable quadrature formula for the function discretized by f^i. This result follows from equation (18), combined with the fact that:

$$R_d^2[\bar{\boldsymbol{v}}] = R_d^2[\bar{\boldsymbol{u}} + \epsilon_r \hat{\boldsymbol{u}}] = R_d^2[u(x^i) + \epsilon_r \hat{u}^i] \tag{7}$$

where the term containing the discretization error has been dropped because $\epsilon_d \ll \epsilon_r$.

Equation (6) results in two conditions for lowering the residual $R[\bar{\boldsymbol{v}}]$: either decrease the reconstruction error (e.g. by adding more degrees of freedom to the network to obtain more faithful reconstructions), or minimize the expression:

$$\left[\frac{\partial^2 s}{\partial u^2} + 2 \sum_n D^n \frac{\partial s}{\partial u} + \sum_{n,m} D^m D^n s \right]_{u=u_i} \tag{8}$$

The former condition relates to the AE architecture and training details. The latter, on the other hand, depends solely on how the input set $\{\bar{\boldsymbol{u}}\}_k$ has been chosen. In other words, choosing the training solutions in regions where the equation residual $R[u]$ is comparatively less sensitive to the functional details of u results in reconstructed data with a small residual. Vice versa, if the $\{\bar{\boldsymbol{u}}\}_k$ represent solutions in the functional neighborhood of which the residual rises sharply, the $\{\bar{\boldsymbol{v}}\}_k$ will exhibit higher PDE violations. This result is broadly intuitive, but equation (8) quantifies the condition for selecting input data leading to minimal PDE violations on the reconstructed data.

We can use a similar approach to quantify $R[\boldsymbol{d}(\boldsymbol{z}; \boldsymbol{\theta_d})]$ in the neighborhood of the latent points $\bar{\boldsymbol{z}}$ encoding the input data. If:

$$\boldsymbol{z} = \bar{\boldsymbol{z}} + \epsilon_z \hat{\boldsymbol{z}} \tag{9}$$

we can write:

$$R_d^2[\boldsymbol{d}(\boldsymbol{z}; \boldsymbol{\theta_d})] = R_d^2[\boldsymbol{d}(\bar{\boldsymbol{z}} + \epsilon_z \hat{\boldsymbol{z}}; \boldsymbol{\theta_d})] = R_d^2[\bar{\boldsymbol{v}} + \epsilon_z \tilde{\boldsymbol{z}} + O(\epsilon_z^2)] \tag{10}$$

where $\tilde{\boldsymbol{z}}$ depends on the first derivatives of the decoder function with respect to the latent vector $\boldsymbol{z}$. To leading order, therefore, this expression can be written in a form similar to (6), where the functional variation includes a term proportional to ϵ_r and a term proportional to ϵ_z, both multiplied by the expression in (8). Choosing training data where this expression is low guarantees that not only the reconstructed data $\bar{\boldsymbol{v}} \equiv \boldsymbol{d}(\bar{\boldsymbol{z}}; \boldsymbol{\theta_d})$ has low residual, but so does also any decoded output from the neighborhood of $\bar{\boldsymbol{z}}$.

URM STATEMENT

The authors acknowledge that at least one key author of this work meets the URM criteria of ICLR 2023 Tiny Papers Track.

REFERENCES

M. Kot. *A First Course in the Calculus of Variations*. Student Mathematical Library. American Mathematical Society, 2014. ISBN 9781470414955. URL https://books.google.co.uk/books?id=UBi8BAAAQBAJ.

Zongyi Li, Nikola Kovachki, Kamyar Azizzadenesheli, Burigede Liu, Kaushik Bhattacharya, Andrew Stuart, and Anima Anandkumar. Fourier neural operator for parametric partial differential equations. *arXiv preprint arXiv:2010.08895*, 2020.

Maziar Raissi, Paris Perdikaris, and George Em Karniadakis. Physics Informed Deep Learning (Part I): Data-driven Solutions of Nonlinear Partial Differential Equations. *arXiv e-prints*, art. arXiv:1711.10561, November 2017. doi: 10.48550/arXiv.1711.10561.

Justin Sirignano and Konstantinos Spiliopoulos. DGM: A deep learning algorithm for solving partial differential equations. *Journal of Computational Physics*, 375:1339–1364, December 2018. doi: 10.1016/j.jcp.2018.08.029.

Naoya Takeishi and Alexandros Kalousis. Physics-integrated variational autoencoders for robust and interpretable generative modeling. *CoRR*, abs/2102.13156, 2021. URL https://arxiv.org/abs/2102.13156.

A VARIATIONS OF THE EQUATION RESIDUAL

Given a partial differential equation of the form:

$$O(u, \nabla u) = 0 \tag{11}$$

where u is a field over space and time, $u : \Omega \subseteq \mathbb{R}^d \to \mathbb{R}$, and ∇u indicates generically its derivatives, the *residual* of (11) is defined as:

$$r(u, \nabla u) \equiv O(u, \nabla u) \tag{12}$$

and represents the extent to which a function violates (11). If $\mathcal{F}$ is a suitably regular function space, $r : \mathcal{F} \to \mathcal{F}$. A functional $R : \mathcal{F} \to \mathbb{R}$ can then be defined as:

$$R[u] = ||r(u, \nabla u)||_2 \tag{13}$$

or

$$R^2[u] = \int r^2(u, \nabla u) \mathrm{d}x \tag{14}$$

For ease of notation, we introduce $S[u] \equiv R^2[u]$ and $s(u, \nabla u) \equiv r^2(u, \nabla u)$, as well as $\partial/\partial x^i \equiv \partial_i$. Let us first note that $S[u] = 0$ iff u is a solution of (11). Furthermore, if u is a solution and $\bar{u} = u + \epsilon \hat{u}$, with $\epsilon \ll 1$ and $\hat{u}$ an arbitrary function, then

$$\delta S = \epsilon \int \left[\frac{\partial s}{\partial u} + \sum_n \sum_{i_1 \cdots i_n} (-1)^n \partial_{i_1} \cdots \partial_{i_n} \frac{\partial s}{\partial(\partial_{i_1} \ldots \partial_{i_n} u)} \right] \hat{u} \mathrm{d}x = 0 \tag{15}$$

or

$$\frac{\partial s}{\partial u} + \sum_n D^n s = 0 \tag{16}$$

where we have introduced the further notation

$$D^n = (-1)^n \sum_{i_1 \cdots i_n} \partial_{i_1} \cdots \partial_{i_n} \frac{\partial}{\partial(\partial_{i_1} \ldots \partial_{i_n} u)} \tag{17}$$

Equation (15) follows from the fact that, as S is non-negative, its zeros are also minima, so S is stationary there and its first variation δS vanishes. Notice that the expression in (15) assumes that the field $\hat{u}$ vanishes at the boundary of Ω.

If $S[u] = 0 = \delta S[u]$, it follows that, around solutions, the second variation is the leading-order contribution to $S[u]$. This quantity can be written as:

$$\delta^2 S \quad = \quad \epsilon^2 \int \left[\frac{\partial^2 s}{\partial u^2} + 2 \sum_n D^n \frac{\partial s}{\partial u} + \sum_{n,m} D^m D^n s \right] \hat{u}^2 \mathrm{d}x \tag{18}$$

For more information on the variation procedure, see e.g. Kot (2014).