A Finite-Particle Convergence Rate
for Stein Variational Gradient Descent

Jiaxin Shi[*]   JIAXINS@STANFORD.EDU
Stanford University

Lester Mackey   LMACKEY@MICROSOFT.COM
Microsoft Research New England

Abstract

We provide a first finite-particle convergence rate for Stein variational gradient descent (SVGD). Specifically, whenever the target distribution satisfies Talagrand's T_1 inequality, SVGD with n particles and an appropriate step size sequence drives the kernel Stein discrepancy to zero at an order $1/\sqrt{\log \log n}$ rate. We suspect that the dependence on n can be improved, and we hope that our explicit, non-asymptotic proof strategy will serve as a template for future refinements.

1. Introduction

Stein variational gradient descent [SVGD, 14] is an algorithm for approximating a target probability distribution P on $\mathbb{R}^d$ with a collection of n particles. Given an initial particle approximation $\mu_0^n = \frac{1}{n} \sum_{i=1}^{n} \delta_{x_i}$ with locations $x_i \in \mathbb{R}^d$, SVGD (Algorithm 1) iteratively evolves the particle locations to provide a more faithful approximation of the target P by performing optimization in the space of probability measures. SVGD has demonstrated encouraging results for a wide variety of inferential tasks, including approximate inference [14, 22, 23], generative modeling [21], and reinforcement learning [10, 16].

Algorithm 1 Stein Variational Gradient Descent [14]: SVGD(μ_0, r)

Input: Target P with density p, kernel k, step sizes $(\epsilon_s)_{s \geq 0}$, approximating measure μ_0, rounds r
for $s = 0, \cdots, r-1$ **do**
    Let μ_{s+1} be the distribution of $X^s + \epsilon_s \int k(x, X^s) \nabla \log p(x) + \nabla_x k(x, X^s) d\mu_s(x)$ for $X^s \sim \mu_s$.
Output: Updated approximation μ_r of the target P

Despite the popularity of SVGD, relatively little is known about its approximation quality. A first analysis by Liu [13, Thm. 3.3] showed that *continuous SVGD*—that is, Algorithm 1 initialized with a continuous distribution μ_0^∞ in place of the discrete particle approximation μ_0^n—converges to P in kernel Stein discrepancy [KSD, 4, 8, 15]. KSD convergence is also known to imply weak convergence under various conditions on the target P and the SVGD kernel k [1, 3, 8, 11]. Follow-up work by Korba et al. [12], Salim et al. [17], Sun et al. [19] sharpened the result of Liu with path-independent constants, weaker smoothness conditions, and explicit rates of convergence. In addition, Duncan et al. [6] analyzed the continuous-time limit of continuous SVGD to provide

---

[*] Part of this work was done at Microsoft Research New England.

conditions for exponential convergence. However, each of these analyses applies only to continuous SVGD and not to the finite-particle algorithm used in practice.

To bridge this gap, Liu [13, Thm. 3.2] showed that n-particle SVGD converges to continuous SVGD in bounded-Lipschitz distance but only under boundedness assumptions violated by most applications of SVGD. To provide a more broadly applicable proof of convergence, Gorham et al. [9, Thm. 7] showed that n-particle SVGD converges to continuous SVGD in 1-Wasserstein distance under assumptions commonly satisfied in SVGD applications. However, both convergence results are asymptotic, providing neither explicit error bounds nor rates of convergence. Korba et al. [12, Prop. 7] explicitly bounded the expected squared Wasserstein distance between n-particle and continuous SVGD but only under the assumption of bounded $\nabla \log p$, an assumption that rules out all strongly log concave or dissipative distributions and all distributions for which the KSD is currently known to control weak convergence [1, 3, 8, 11]. In addition, Korba et al. [12] do not provide a unified bound for the convergence of n-particle SVGD to P.

In this work, we derive a first unified convergence bound for finite-particle SVGD to its target. To achieve this, we first bound the 1-Wasserstein discretization error between finite-particle and continuous SVGD under assumptions commonly satisfied in SVGD applications and compatible with KSD weak convergence control (see Theorem 1). We next bound KSD in terms of 1-Wasserstein distance and SVGD moment growth to explicitly control KSD discretization error in Theorem 2. Finally, Theorem 3 combines our results with the established KSD analysis of continuous SVGD to arrive at an explicit KSD error bound for n-particle SVGD.

2. Notation and Assumptions

Throughout, we fix a target distribution P in the set $\mathcal{P}$ of probability measures on $\mathbb{R}^d$, a nonnegative step size sequence $(\epsilon_s)_{s \geq 0}$, and a reproducing kernel $k : \mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ with reproducing kernel Hilbert space (RKHS) $\mathcal{H}$ and product RKHS $\mathcal{H}^d \triangleq \bigotimes_{i=1}^d \mathcal{H}$ [2]. For all $\mu, \nu \in \mathcal{P}$, we let $W_1(\mu, \nu) \triangleq \inf_{X \sim \mu, Z \sim \nu} \mathbb{E}[\|Z - X\|_2]$ denote the 1-Wasserstein distance between them and introduce the shorthand $m_\mu \triangleq \mathbb{E}_\mu[\|\cdot\|]_2$, $m_{\mu,P} \triangleq \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2]$, and $M_{\mu,P} \triangleq \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2^2]$. We allow each of these quantities to take on the value ∞ when the random variables are not suitably integrable. We further define the Kullback-Leibler divergence as $\mathrm{KL}(\mu\|\nu) \triangleq \mathbb{E}_\mu[\log(\frac{d\mu}{d\nu})]$ when μ is absolutely continuous with respect to ν, denoted by $\mu \ll \nu$, and as ∞ otherwise.

Our analysis will make use of the following standard assumptions on the SVGD kernel and target distribution.

Assumption 1 (Lipschitz, mean-zero score function) *The target distribution $P \in \mathcal{P}$ has a differentiable density p with an L-Lipschitz score function $s_p \triangleq \nabla \log p$, i.e., $\|s_p(x) - s_p(y)\|_2 \leq L\|x - y\|_2$ for all $x, y \in \mathbb{R}^d$. Moreover, $\mathbb{E}_P[s_p] = 0$ and $s_p(x^\star) = 0$ for some $x^\star \in \mathbb{R}^d$.*

Assumption 2 (Bounded kernel derivatives) *For any multi-index $I = (I_1, I_2, \ldots, I_d)$ with $|I| \triangleq \sum_{i=1}^d I_i \leq 2$, we have $\sup_{x \in \mathbb{R}^d} \left(D_x^I D_y^I k(x, y)|_{y=x} \right) \leq \kappa^2$. Here, D^I is the differential operator defined as $D_x^I = \frac{d^{|I|}}{dx_1^{I_1} dx_2^{I_2} \ldots dx_d^{I_d}}$.*

Assumption 3 (Decaying kernel derivatives) *There exists $\gamma > 0$ such that*

$$\sup_{x,y \in \mathbb{R}^d, \|x-y\|_2 \geq r} \|\nabla_x k(x, y)\|_2 \leq \gamma/r.$$

To leverage the continuous SVGD convergence rates of Salim et al. [17], we additionally assume that the target P satisfies Talagrand's T_1 inequality [20, Def. 22.1]. Remarkably, Villani [20, Thm. 22.10] showed that Assumption 4 is *equivalent* to P being a sub-Gaussian distribution. Hence, this mild assumption holds for all strongly log concave P [18, Def. 2.9], all P satisfying the log Sobolev inequality [20, Thm. 22.17], and all *distantly dissipative P* for which KSD is known to control weak convergence [8, Def. 4].

Assumption 4 (Talagrand's T_1 inequality [20, Def. 22.1]) $\mathbb{E}_P[\|\cdot\|_2] < \infty$, *and there exists* $\lambda > 0$ *such that for all* μ *with* $\mathbb{E}_\mu[\|\cdot\|_2] < \infty$

$$W_1(\mu, P) \leq \sqrt{2\mathrm{KL}(\mu\|P)/\lambda}.$$

Finally we make use of the following notation specific to the SVGD algorithm.

Definition 1 (Stein operator) *For any vector-valued function* $g : \mathbb{R}^d \to \mathbb{R}^d$, *the* Langevin Stein operator [7] *for P satisfying Assumption 1 is defined by*

$$(\mathcal{T}_P g)(x) \triangleq \langle s_p(x), g(x) \rangle + \nabla \cdot g(x).$$

Definition 2 (Vector-valued Stein operator) *For any function* $h : \mathbb{R}^d \to \mathbb{R}$, *the* vector-valued Langevin Stein operator [15] *for P satisfying Assumption 1 is defined by*

$$(\mathcal{A}_P h)(x) \triangleq s_p(x)h(x) + \nabla h(x)$$

with components

$$(\mathcal{A}_P^j h)(x) \triangleq (s_p(x))_j \, h(x) + \nabla_j h(x).$$

Definition 3 (SVGD transport map and pushforward) *The* SVGD transport map [14] *for a kernel k, target P satisfying Assumption 1, approximating $\mu \in \mathcal{P}$, and step size ϵ is given by*

$$T_{\mu,\epsilon}(x) \triangleq x + \epsilon \mathbb{E}_{X \sim \mu}[(\mathcal{A}_P k(\cdot, x))(X)].$$

Moreover, the SVGD pushforward $\Phi_\epsilon(\mu)$ *represents the distribution of* $T_{\mu,\epsilon}(X)$ *when* $X \sim \mu$.

Definition 4 (Kernel Stein discrepancy) *For a target P satisfying Assumption 1 and measures* $\mu, \nu \in \mathcal{P}$, *we define the* Langevin kernel Stein discrepancy [KSD, 4, 8, 15] *with base kernel k by*[1]

$$\mathrm{KSD}_P(\mu, \nu) \triangleq \sup_{\|g\|_{\mathcal{H}^d} \leq 1} \mathbb{E}_\mu[\mathcal{T}_P g] - \mathbb{E}_\nu[\mathcal{T}_P g].$$

3. Wasserstein Discretization Error of SVGD

Our first main result concerns the discretization error of SVGD and shows that n-particle SVGD does not stray too far from its continuous SVGD limit when its step sizes are chosen appropriately.

---

1. Prior work [4, 8, 15] only defined $\mathrm{KSD}_P(\mu, \nu)$ for the case $\nu = P$. Definition 4 extends this definition to general ν.

Theorem 1 (Wasserstein discretization error of SVGD) *Suppose Assumptions 1, 2, and 3 hold. For any $\mu_0^n, \mu_0^\infty \in \mathcal{P}$, the Algorithm 1 outputs $\mu_r^n = \mathrm{SVGD}(\mu_0^n, r)$ and $\mu_r^\infty = \mathrm{SVGD}(\mu_0^\infty, r)$ satisfy*

$$W_1(\mu_r^n, \mu_r^\infty) \le W_1(\mu_0^n, \mu_0^\infty) \exp(b_{r-1}(A + B \exp(C b_{r-1})))$$

for $b_{r-1} \triangleq \sum_{s=0}^{r-1} \epsilon_s$, $A = (c_1 + c_2)(1 + m_P)$, $B = c_1 m_{\mu_0^n, P} + c_2 m_{\mu_0^\infty, P}$, and $C = \kappa^2(3L + d)$. Here $c_1 = \max(\sqrt{d}\kappa^2 L, \sqrt{d}\kappa^2 L \|x^\star\|_2 + d\kappa^2)$ and $c_2 = \kappa^2 L + d\kappa^2 + (\gamma + \sqrt{d}\kappa^2)(1 + \|x^\star\|_2)$.

The proof of Theorem 1 in Appendix A relies on two lemmas. The first, due to Gorham et al. [9], shows that the one-step SVGD pushforward Φ_ϵ (Definition 3) is pseudo-Lipschitz with respect to the 1-Wasserstein distance whenever the score function $\nabla \log p$ and kernel k fulfill a commonly-satisfied pseudo-Lipschitz condition.

Lemma 1 (Wasserstein pseudo-Lipschitzness of SVGD [9, Lem. 12]) *For $P \in \mathcal{P}$ with differentiable density p, suppose that the following pseudo-Lipschitz bounds hold*

$$\sup_{z \in \mathbb{R}^d} \|\nabla_z(s_p(x)k(x,z) + \nabla_x k(x,z))\|_{\mathrm{op}} \le c_1(1 + \|x\|_2),$$

$$\sup_{x \in \mathbb{R}^d} \|\nabla_x(s_p(x)k(x,z) + \nabla_x k(x,z))\|_{\mathrm{op}} \le c_2(1 + \|z\|_2).$$

for some constants $c_1, c_2 > 0$. Then, for any $\mu, \nu \in \mathcal{P}$,

$$W_1(\Phi_\epsilon(\mu), \Phi_\epsilon(\nu)) \le W(\mu, \nu)(1 + \epsilon c_{\mu,\nu}),$$

where Φ_ϵ is the one-step SVGD pushforward (Definition 3) and $c_{\mu,\nu} = c_1(1 + m_\mu) + c_2(1 + m_\nu)$.

The second lemma, proved in Appendix B, controls the growth of the first and second absolute moments under SVGD.

Lemma 2 (SVGD moment growth) *Suppose Assumptions 1 and 2 hold, and let $C = \kappa^2(3L+d)$. Then the SVGD output μ_r of Algorithm 1 with $b_{r-1} \triangleq \sum_{s=0}^{r-1} \epsilon_s$ satisfies*

$$m_{\mu_r} \le m_{\mu_0, P} \prod_{s=0}^{r-1}(1 + \epsilon_s C) + m_P \le m_{\mu_0, P} \exp(C b_{r-1}) + m_P,$$
$$M_{\mu_r, P} \le M_{\mu_0, P} \prod_{s=0}^{r-1}(1 + \epsilon_s C)^2 \le M_{\mu_0, P} \exp(2C b_{r-1}).$$

4. KSD Discretization Error of SVGD

Our next result translates the Wasserstein error bounds of Theorem 1 into KSD error bounds.

Theorem 2 (KSD discretization error of SVGD) *Suppose Assumptions 1 and 2 hold. For any $\mu_0^n, \mu_0^\infty \in \mathcal{P}$, the Algorithm 1 outputs $\mu_r^n = \mathrm{SVGD}(\mu_0^n, r)$ and $\mu_r^\infty = \mathrm{SVGD}(\mu_0^\infty, r)$ satisfy*

$$\begin{aligned}
\mathrm{KSD}_P(\mu_r^n, \mu_r^\infty) &\le \kappa(d + L) w_{0,n} \exp(b_{r-1}(A + B \exp(C b_{r-1}))) \\
&\quad + d^{1/4} \kappa L \sqrt{2 M_{\mu_0^\infty, P} w_{0,n}} \exp(b_{r-1}(2C + A + B \exp(C b_{r-1}))/2)
\end{aligned}$$

for $w_{0,n} \triangleq W_1(\mu_0^n, \mu_0^\infty)$ and A, B, C defined as in Theorem 1.

Our proof of Theorem 2 relies on the following lemma, proved in Appendix C, that shows that the KSD is controlled by the 1-Wasserstein distance.

Lemma 3 (KSD-Wasserstein bound) *Suppose Assumptions 1 and 2 hold. For any $\mu, \nu \in \mathcal{P}$,*

$$\mathrm{KSD}_P(\mu, \nu) \leq \kappa(d+L)W_1(\mu, \nu) + L\mathbb{E}_{(X\sim\mu, Y\sim\nu)\perp\!\!\!\perp Z\sim P}[\|Y-Z\|_2 \min(2\kappa, \sqrt{d}\kappa\|X-Y\|_2)]$$
$$\leq \kappa(d+L)W_1(\mu, \nu) + d^{1/4}\kappa L\sqrt{2M_{\nu,P}W_1(\mu, \nu)}.$$

Proof of Theorem 2 The result follows directly from Lemma 3, Lemma 2, and Theorem 1. ∎

5. A Finite-particle Convergence Rate for SVGD

To establish our main SVGD convergence result, we combine Theorems 1 and 2 with the following descent lemma for continuous SVGD error due to Salim et al. [17] which shows that continuous SVGD decreases the KL divergence to P and drives the KSD to P to zero.

Lemma 4 (Continuous SVGD descent lemma [17, Thm. 3.2]) *Suppose Assumptions 1, 2, and 4 hold, and consider the outputs $\mu_r^\infty = \mathrm{SVGD}(\mu_0^\infty, r)$ and $\mu_{r+1}^\infty = \mathrm{SVGD}(\mu_0^\infty, r+1)$ of Algorithm 1 with $\mu_0^\infty \ll P$. If $\max_{0\leq s\leq r}\epsilon_s \leq R_{\alpha,2}$ for some $\alpha > 1$ and*

$$R_{\alpha,p} \triangleq \min\left(\frac{p}{\kappa^2(L+\alpha^2)}, (\alpha-1)(1+L\mathbb{E}_{\mu_0^\infty}[\|\cdot - x^\star\|_2] + 2L\sqrt{2\mathrm{KL}(\mu_0^\infty\|P)/\lambda})\right) \text{ for } p \in \{1, 2\},$$

then

$$\mathrm{KL}(\mu_{r+1}^\infty\|P) - \mathrm{KL}(\mu_r^\infty\|P) \leq -\epsilon_r\left(1 - \frac{\kappa^2(L+\alpha^2)}{2}\epsilon_r\right)\mathrm{KSD}_P(\mu_r^\infty, P)^2. \tag{1}$$

By summing the result (1) over $r = 0, \ldots, t$, we obtain the following corollary.

Corollary 1 *Under the assumptions and notation of Lemma 4, suppose $\max_{0\leq r\leq t}\epsilon_r \leq R_{\alpha,1}$ for some $\alpha > 1$, and let $\pi_r \triangleq \frac{c(\epsilon_r)}{\sum_{r=0}^t c(\epsilon_r)}$ for $c(\epsilon) \triangleq \epsilon\left(1 - \frac{\kappa^2(L+\alpha^2)}{2}\epsilon\right)$. Since $\frac{\epsilon}{2} \leq c(\epsilon) < \epsilon$, we have*

$$\sum_{r=0}^t \pi_r\mathrm{KSD}_P(\mu_r^\infty, P)^2 \leq \frac{1}{\sum_{r=0}^t c(\epsilon_r)}\mathrm{KL}(\mu_0^\infty\|P) \leq \frac{2}{\sum_{r=0}^t \epsilon_r}\mathrm{KL}(\mu_0^\infty\|P).$$

Finally, we arrive at our main result that bounds the approximation error of n-particle SVGD in terms of the chosen step size sequence and the initial discretization error $W_1(\mu_0^n, \mu_0^\infty)$.

Theorem 3 (KSD error of finite-particle SVGD) *Suppose Assumptions 1, 2, 3, and 4 hold, fix any $\mu_0^\infty \ll P$ and $\mu_0^n \in \mathcal{P}$, and let $w_{0,n} \triangleq W_1(\mu_0^n, \mu_0^\infty)$. If $\max_{0\leq r\leq t}\epsilon_r \leq R_{\alpha,1}$ for $\alpha > 1$ and $R_{\alpha,1}$ defined in Lemma 4, then the Algorithm 1 outputs $\mu_r^n = \mathrm{SVGD}(\mu_0^n, r)$ satisfy*

$$\min_{0\leq r\leq t}\mathrm{KSD}_P(\mu_r^n, P) \leq \sum_{r=0}^t \pi_r\mathrm{KSD}_P(\mu_r^n, P) \leq a_t + \left(\frac{2}{b_t}\mathrm{KL}(\mu_0^\infty\|P)\right)^{1/2}, \tag{2}$$

for π_r as defined in Lemma 4, (A, B, C) as defined in Theorem 1, $b_t \triangleq \sum_{r=0}^t \epsilon_r$, and

$$a_t \triangleq \kappa(d+L)w_{0,n}\exp(b_t(A + B\exp(Cb_t)))$$
$$+ d^{1/4}\kappa L\sqrt{2M_{\mu_0^\infty, P}w_{0,n}}\exp(b_t(2C + A + B\exp(Cb_t))/2).$$

Proof By the triangle inequality and Theorem 2 we have

$$|\text{KSD}_P(\mu_r^n, P) - \text{KSD}_P(\mu_r^\infty, P)| \leq \text{KSD}_P(\mu_r^n, \mu_r^\infty) \leq a_{r-1}$$

for each r. Therefore

$$\sum_{r=0}^t \pi_r (\text{KSD}_P(\mu_r^n, P) - a_{r-1})^2 \leq \sum_{r=0}^t \pi_r \text{KSD}_P(\mu_r^\infty, P)^2 \leq \frac{2}{b_t} \text{KL}(Q_0^\infty \| P), \qquad (3)$$

where the last inequality follows from Corollary 1. Moreover, by Jensen's inequality,

$$\sum_{r=0}^t \pi_r (\text{KSD}_P(\mu_r^n, P) - a_{r-1})^2 \geq \left(\sum_{r=0}^t \pi_r \text{KSD}_P(\mu_r^n, P) - \sum_{r=0}^t \pi_r a_{r-1} \right)^2. \qquad (4)$$

Combining (3) and (4), we have

$$\sum_{r=0}^t \pi_r \text{KSD}_P(\mu_r^n, P) \leq \sum_{r=0}^t \pi_r a_{r-1} + \left(\frac{2}{b_t} \text{KL}(\mu_0^\infty \| P) \right)^{1/2}.$$

We finish the proof by noticing that $\sum_{r=0}^t \pi_r a_{r-1} \leq \max_{0 \leq r \leq t} a_{r-1} = a_{t-1} \leq a_t$. ∎

The following corollary, proved in Appendix D, provides an explicit SVGD convergence rate by choosing the step size sequence to balance the three terms on the right-hand side of (2).

Corollary 2 (A finite-particle convergence rate for SVGD) *Instantiate the notation and assumptions of Theorem 3, and define the shorthand* $\psi_{B,C}(x, y, \beta) \triangleq \frac{1}{C} \log(\frac{1}{B}(\frac{1}{\beta} \log \frac{1}{x} - y))$. *If*

$$b_t = \sum_{r=0}^t \epsilon_r = \min\left(\psi_{B,C}\left(w_{0,n} \sqrt{\log\log \tfrac{1}{w_{0,n}}}, A, \beta_1 \right), \psi_{B,C}\left(w_{0,n} \log\log \tfrac{1}{w_{0,n}}, A + 2C, \beta_2 \right) \right),$$

$$\text{for } \beta_1 \triangleq \max\left(1, \psi_{B,C}\left(w_{0,n} \sqrt{\log\log \tfrac{1}{w_{0,n}}}, A, 1 \right) \right) \text{ and}$$

$$\beta_2 \triangleq \max\left(1, \psi_{B,C}\left(w_{0,n} \log\log \tfrac{1}{w_{0,n}}, A + 2C, 1 \right) \right),$$

then,

$$\min_{0 \leq r \leq t} \text{KSD}_P(\mu_r^n, P) = O\left(1 \Big/ \sqrt{\log\log \tfrac{1}{w_{0,n}}} \right).$$

If, in addition, $\mu_0^n = \frac{1}{n} \sum_{i=1}^n \delta_{x_i}$ *for* $x_i \overset{i.i.d.}{\sim} \mu_0^\infty$, *then*

$$\min_{0 \leq r \leq t} \text{KSD}_P(\mu_r^n, P) = O\left(\frac{1}{\sqrt{\log\log n}} \right).$$

References

[1] Alessandro Barp, Carl-Johann Simon-Gabriel, Mark Girolami, and Lester Mackey. Targeted separation and convergence with kernel discrepancies. *arXiv preprint arXiv:2209.12835*, 2022.

[2] Alain Berlinet and Christine Thomas-Agnan. *Reproducing kernel Hilbert spaces in probability and statistics*. Springer Science & Business Media, 2011.

[3] Wilson Ye Chen, Lester Mackey, Jackson Gorham, François-Xavier Briol, and Chris Oates. Stein points. In *International Conference on Machine Learning*, pages 844–853, 2018.

[4] Kacper Chwialkowski, Heiko Strathmann, and Arthur Gretton. A kernel test of goodness of fit. In *International Conference on Machine Learning*, pages 2606–2615, 2016.

[5] Richard Mansfield Dudley. The speed of mean Glivenko-Cantelli convergence. *The Annals of Mathematical Statistics*, 40(1):40–50, 1969.

[6] Andrew Duncan, Nikolas Nüsken, and Lukasz Szpruch. On the geometry of Stein variational gradient descent. *arXiv preprint arXiv:1912.00894*, 2019.

[7] Jackson Gorham and Lester Mackey. Measuring sample quality with Stein's method. In *Advances in Neural Information Processing Systems*, pages 226–234, 2015.

[8] Jackson Gorham and Lester Mackey. Measuring sample quality with kernels. In *International Conference on Machine Learning*, pages 1292–1301, 2017.

[9] Jackson Gorham, Anant Raj, and Lester Mackey. Stochastic Stein discrepancies. *Advances in Neural Information Processing Systems*, 33:17931–17942, 2020.

[10] Tuomas Haarnoja, Haoran Tang, Pieter Abbeel, and Sergey Levine. Reinforcement learning with deep energy-based policies. In *International Conference on Machine Learning*, pages 1352–1361, 2017.

[11] Jonathan Huggins and Lester Mackey. Random feature Stein discrepancies. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, pages 1903–1913. 2018.

[12] Anna Korba, Adil Salim, Michael Arbel, Giulia Luise, and Arthur Gretton. A non-asymptotic analysis for Stein variational gradient descent. *Advances in Neural Information Processing Systems*, 33, 2020.

[13] Qiang Liu. Stein variational gradient descent as gradient flow. In *Advances in Neural Information Processing Systems*, pages 3115–3123, 2017.

[14] Qiang Liu and Dilin Wang. Stein variational gradient descent: A general purpose Bayesian inference algorithm. *Advances in Neural Information Processing Systems*, 29:2378–2386, 2016.

[15] Qiang Liu, Jason Lee, and Michael Jordan. A kernelized Stein discrepancy for goodness-of-fit tests. In *International Conference on Machine Learning*, pages 276–284, 2016.

[16] Yang Liu, Prajit Ramachandran, Qiang Liu, and Jian Peng. Stein variational policy gradient. In *33rd Conference on Uncertainty in Artificial Intelligence*, 2017.

[17] Adil Salim, Lukang Sun, and Peter Richtarik. A convergence theory for SVGD in the population limit under Talagrand's inequality T1. In *International Conference on Machine Learning*, pages 19139–19152. PMLR, 2022.

[18] Adrien Saumard and Jon A Wellner. Log-concavity and strong log-concavity: a review. *Statistics surveys*, 8:45, 2014.

[19] Lukang Sun, Avetik Karagulyan, and Peter Richtarik. Convergence of Stein variational gradient descent under a weaker smoothness condition. *arXiv preprint arXiv:2206.00508*, 2022.

[20] Cédric Villani. *Optimal transport: old and new*, volume 338. Springer, 2009.

[21] Dilin Wang and Qiang Liu. Learning to draw samples: With application to amortized MLE for generative adversarial learning. *arXiv preprint arXiv:1611.01722*, 2016.

[22] Dilin Wang, Zhe Zeng, and Qiang Liu. Stein variational message passing for continuous graphical models. In *International Conference on Machine Learning*, pages 5219–5227, 2018.

[23] Jingwei Zhuo, Chang Liu, Jiaxin Shi, Jun Zhu, Ning Chen, and Bo Zhang. Message passing Stein variational gradient descent. In *International Conference on Machine Learning*, pages 6018–6027, 2018.

Appendix A. Proof of Theorem 1: Wasserstein discretization error of SVGD

We first show that the pseudo-Lipschitzness conditions of Lemma 1 hold given our assumptions. According to Assumption 2, we have

$$\nabla_{z_i} k(x, z) = \langle \nabla_{z_i} k(z, \cdot), k(x, \cdot) \rangle_{\mathcal{H}} \leq (\nabla_{z_i} \nabla_{z'_i} k(z, z')|_{z'=z} k(x, x))^{1/2} \leq \kappa^2,$$

$$\|\nabla_z k(x, z)\|_2 = \sqrt{\sum_{i=1}^d (\nabla_{z_i} k(x, z))^2} \leq \sqrt{d}\kappa^2.$$

$$\nabla_{z_j} \nabla_{x_i} k(x, z) = \langle \nabla_{z_j} k(z, \cdot), \nabla_{x_i} k(x, \cdot) \rangle_{\mathcal{H}} \leq (\nabla_{z_j} \nabla_{z'_j} k(z, z')|_{z'=z} \nabla_{x_i} \nabla_{x'_i} k(x, x')|_{x'=x})^{1/2} \leq \kappa^2,$$

$$\|\nabla_z \nabla_x k(x, z)\|_{\mathrm{op}} \leq \|\nabla_z \nabla_x k(x, z)\|_F = \sqrt{\sum_{i=1}^d \sum_{j=1}^d (\nabla_{z_j} \nabla_{x_i} k(x, z))^2} \leq d\kappa^2,$$

Recall that $x^\star$ satisfies $s_p(x^\star) = 0$ by Assumption 1. Then, by the triangle inequality, the definition of $\|\cdot\|_{\mathrm{op}}$, and Cauchy-Schwartz,

$$\begin{aligned}
&\|\nabla_z (s_p(x) k(x, z) + \nabla_x k(x, z))\|_{\mathrm{op}} \\
&= \|(s_p(x) - s_p(x^\star)) \nabla_z k(x, z)^\top\|_{\mathrm{op}} + \|\nabla_z \nabla_x k(x, z)\|_{\mathrm{op}} \\
&\leq \sup_{\|u\|_2 \leq 1} (\|s_p(x) - s_p(x^\star)\|_2 |\nabla_z k(x, z)^\top u|) + d\kappa^2 \\
&\leq L \|x - x^\star\|_2 \|\nabla_z k(x, z)\|_2 + d\kappa^2 \\
&\leq \sqrt{d} \kappa^2 L (\|x\|_2 + \|x^\star\|_2) + d\kappa^2 \\
&\leq \max(\sqrt{d} \kappa^2 L, \sqrt{d} \kappa^2 L \|x^\star\|_2 + d\kappa^2)(1 + \|x\|_2).
\end{aligned}$$

Letting $c_1 = \max(\sqrt{d} \kappa^2 L, \sqrt{d} \kappa^2 L \|x^\star\|_2 + d\kappa^2)$ and taking supremum over z proves the first pseudo-Lipschitzness condition. Similarly, we have

$$\begin{aligned}
&\|\nabla_x (s_p(x) k(x, z) + \nabla_x k(x, z))\|_{\mathrm{op}} \\
&= \|\nabla s_p(x)\|_{\mathrm{op}} k(x, z) + \|(s_p(x) - s_p(x^\star)) \nabla_x k(x, z)^\top\|_{\mathrm{op}} + \|\nabla_x^2 k(x, z)\|_{\mathrm{op}} \\
&\leq \kappa^2 L + L \|x - x^\star\|_2 \|\nabla_x k(x, z)\|_2 + d\kappa^2,
\end{aligned} \tag{5}$$

where we used

$$k(x, z) = \langle k(x, \cdot), k(z, \cdot) \rangle_{\mathcal{H}} \leq \sqrt{k(x, x) k(z, z)} \leq \kappa^2,$$

$$\|\nabla_x^2 k(x, z)\|_{\mathrm{op}} \leq \|\nabla_x^2 k(x, z)\|_F = \sqrt{\sum_{|I|=2} (D_x^I k(x, z))^2} \leq d\kappa^2.$$

The last inequality is due to

$$|D_x^I k(x, z)| = |\langle D_x^I k(x, \cdot), k(z, \cdot) \rangle_{\mathcal{H}}| \leq \sqrt{k(z, z) D_x^I D_{x'}^I k(x, x')|_{x'=x}} \leq \kappa^2.$$

According to Assumption 3, there exists $\gamma > 0$ such that $\|\nabla_x k(x, z)\|_2 \leq \gamma \|x - z\|_2^{-1}$. Therefore, for $\|x - z\|_2 \geq 1$,

$$\begin{aligned}
\|x - x^\star\|_2 \|\nabla_x k(x, z)\|_2 &\leq \gamma \|x - x^\star\|_2 / \|x - z\|_2 \\
&\leq \gamma (1 + \|z - x^\star\|_2 / \|x - z\|_2) \\
&\leq \gamma (1 + \|z - x^\star\|_2).
\end{aligned} \tag{6}$$

And for $\|x - z\|_2 < 1$,

$$\|x - x^\star\|_2 \|\nabla_x k(x, z)\|_2 \leq \sqrt{d}\kappa^2(\|x - z\|_2 + \|z - x^\star\|_2)$$
$$\leq \sqrt{d}\kappa^2(1 + \|z - x^\star\|_2). \tag{7}$$

Combining (6) and (7),

$$\|x - x^\star\|_2 \|\nabla_x k(x, z)\|_2 \leq (\gamma + \sqrt{d}\kappa^2)(1 + \|z - x^\star\|_2)$$
$$\leq (\gamma + \sqrt{d}\kappa^2)(1 + \|x^\star\|_2 + \|z\|_2). \tag{8}$$

Plugging (8) back into (5), we can show the second pseudo-Lipschitzness condition holds for $c_2 = \kappa^2 L + d\kappa^2 + (\gamma + \sqrt{d}\kappa^2)(1 + \|x^\star\|_2)$.

Now that the pseudo-Lipschitzness conditions hold, by repeated application of Lemma 1 and the inequality $(1 + x) \leq e^x$, we have

$$W_1(\mu_{r+1}^n, \mu_{r+1}^\infty) = W_1(\Phi_{\epsilon_r}(\mu_r^n), \Phi_{\epsilon_r}(\mu_r^\infty)) \leq (1 + \epsilon_r C_r)W(\mu_r^n, \mu_r^\infty)$$
$$\leq W_1(\mu_0^n, \mu_0^\infty) \prod_{s=0}^{r}(1 + \epsilon_s C_s) \leq W_1(\mu_0^n, \mu_0^\infty)\exp\left(\sum_{s=0}^{r}\epsilon_s C_s\right) \tag{9}$$

for $C_s = c_1(1 + m_{\mu_s^n}) + c_2(1 + m_{\mu_s^\infty})$.

Using the result from Lemma 2, we have

$$C_{s+1} \leq A + B\exp(\kappa^2(3L + d)b_s)$$

for $A = (c_1 + c_2)(1 + m_P)$ and $B = c_1 m_{\mu_0^n, P} + c_2 m_{\mu_0^\infty, P}$. Therefore

$$\sum_{s=0}^{r}\epsilon_s C_s \leq \max_{0 \leq s \leq r} C_s \sum_{s=0}^{r}\epsilon_s$$
$$\leq b_r(A + B\exp(\kappa^2(3L + d)b_{r-1}))$$
$$\leq b_r(A + B\exp(\kappa^2(3L + d)b_r)).$$

Plugging this back into (9) proves the result.

Appendix B. Proof of Lemma 2: SVGD moment growth

From Assumption 2 we know

$$|k(y, x)| \leq |\langle k(y, \cdot), k(x, \cdot)\rangle_{\mathcal{H}}| \leq \|k(y, \cdot)\|_{\mathcal{H}}\|k(x, \cdot)\|_{\mathcal{H}} = \sqrt{k(y, y)k(x, x)} \leq \kappa^2,$$
$$|D_y^I k(y, x)| = |\langle D_y^I k(y, \cdot), k(x, \cdot)\rangle_{\mathcal{H}}| \leq \sqrt{k(x, x)D_y^I D_{y'}^I k(y, y')|_{y'=y}} \leq \kappa^2,$$
$$\left\|\nabla_y^2 k(y, x)\right\|_{op} \leq \left\|\nabla_y^2 k(y, x)\right\|_F = \sqrt{\sum_{|I|=2}(D_y^I k(y, x))^2} \leq d\kappa^2.$$

The last inequality implies

$$\|\nabla_y k(y, x) - \nabla_z k(z, x)\|_2 \leq d\kappa^2\|y - z\|_2.$$

Let μ be any probability measure. Using the above results, Jensen's inequality and the facts that $\mathbb{E}_P[s_p(\cdot)] = 0$ and $\mathbb{E}_{Z \sim P}[(\mathcal{A}_P k(\cdot, x))(Z)] = 0$, we have

$$
\begin{aligned}
&\|T_{\mu,\epsilon}(x) - x\|_2 \\
&\leq \epsilon \|\mathbb{E}_{X \sim \mu}[(\mathcal{A}_P k(\cdot, x))(X)]\|_2 \\
&= \epsilon \|\mathbb{E}_{X \sim \mu}[(\mathcal{A}_P k(\cdot, x))(X)] - \mathbb{E}_{Z \sim P}[(\mathcal{A}_P k(\cdot, x))(Z)]\|_2 \\
&= \epsilon \|\mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[k(Z, x)(s_p(X) - s_p(Z)) + (k(X, x) - k(Z, x))(s_p(X) - \mathbb{E}_P[s_p(\cdot)]) \\
&\quad + (\nabla_X k(X, x) - \nabla_Z k(Z, x))]\|_2 \\
&\leq \epsilon \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[|k(Z, x)|\|s_p(X) - s_p(Z)\|_2 + (|k(X, x)| + |k(Z, x)|)\|s_p(X) - \mathbb{E}_P[s_p(\cdot)]\|_2 \\
&\quad + \|\nabla_X k(X, x) - \nabla_Z k(Z, x)\|_2] \\
&\leq \epsilon \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\kappa^2 L \|X - Z\|_2 + 2\kappa^2 \|s_p(X) - \mathbb{E}_{Y \sim P}[s_p(Y)]\|_2 + d\kappa^2 \|X - Z\|_2] \\
&\leq \epsilon \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\kappa^2(L + d)\|X - Z\|_2] + \epsilon \cdot 2\kappa^2 L \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Y \sim P}[\|X - Y\|_2] \\
&= \epsilon \kappa^2 (3L + d) \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2] \\
&= \epsilon C m_{\mu,P}. \tag{10}
\end{aligned}
$$

The last step used the definitions $m_{\mu,P} \triangleq \mathbb{E}_{X \sim \mu \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2]$ and $C = \kappa^2(3L + d)$. Then, applying triangle inequality and (10), we have

$$
\begin{aligned}
m_{\mu_{r+1},P} &= \mathbb{E}_{X \sim \mu_{r+1} \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2] \\
&= \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|T_{\mu_r,\epsilon_r}(X) - Z\|_2] \\
&\leq \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|T_{\mu_r,\epsilon_r}(X) - X\|_2 + \|X - Z\|_2] \\
&\leq \epsilon_r C \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2] + \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2] \\
&= (1 + \epsilon_r C) m_{\mu_r,P}. \tag{11} \\
M_{\mu_{r+1},P} &= \mathbb{E}_{X \sim \mu_{r+1} \perp\!\!\!\perp Z \sim P}[\|X - Z\|_2^2] \\
&= \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|T_{\mu_r,\epsilon_r}(X) - Z\|_2^2] \\
&\leq \mathbb{E}_{X \sim \mu_r \perp\!\!\!\perp Z \sim P}[\|T_{\mu_r,\epsilon_r}(X) - X\|_2^2 + 2\|T_{\mu_r,\epsilon_r}(X) - X\|_2 \|X - Z\|_2 + \|X - Z\|_2^2] \\
&\leq (\epsilon_r^2 C^2 + 2\epsilon_r C) m_{\mu_r,P}^2 + M_{\mu_r,P} \\
&\leq (1 + 2\epsilon_r C + \epsilon_r^2 C^2) M_{\mu_r,P} \\
&\leq (1 + \epsilon_r C)^2 M_{\mu_r,P}. \tag{12}
\end{aligned}
$$

where the second last step used Jensen's inequality $m_{\mu_r,P}^2 \leq M_{\mu_r,P}$. Then, we repeatedly apply (11) and (12) together with the inequality $1 + x \leq e^x$ to get

$$
\begin{aligned}
M_{\mu_r,P} &\leq M_{\mu_0,P} \prod_{s=0}^{r-1}(1 + \epsilon_s C)^2 \leq M_{\mu_0,P} \exp(2C \sum_{s=0}^{r-1} \epsilon_s) \leq M_{\mu_0,P} \exp(2C b_{r-1}), \\
m_{\mu_r} &= \mathbb{E}_{\mu_r}[\|X\|_2] \leq m_{\mu_r,P} + \mathbb{E}_P[\|Z\|_2] \leq m_{\mu_0,P} \prod_{s=0}^{r-1}(1 + \epsilon_s C) + m_P \\
&\leq m_{\mu_0,P} \exp(C b_{r-1}) + m_P.
\end{aligned}
$$

Appendix C. Proof of Lemma 3: KSD-Wasserstein bound

Our proof generalizes that of Gorham and Mackey [8, Lem. 18]. Consider any $g \in \mathcal{H}^d$ satisfying $\|g\|_{\mathcal{H}^d}^2 \triangleq \sum_{i=1}^d \|g_i\|_{\mathcal{H}}^2 \leq 1$. From Assumption 2 we know

$$\|g(x)\|_2^2 \leq \sum_{i=1}^d \kappa^2 \|g_i\|_{\mathcal{H}}^2 \leq \kappa^2, \tag{13}$$

$$\|\nabla g(x)\|_{op}^2 \leq \|\nabla g(x)\|_F^2 = \sum_{i=1}^d \sum_{j=1}^d |\nabla_{x_i} g_j(x)|^2 \leq \sum_{i=1}^d \sum_{j=1}^d \kappa^2 \|g_j\|_{\mathcal{H}}^2 \leq d\kappa^2, \text{ and} \tag{14}$$

$$\|\nabla(\nabla \cdot g(x))\|_2^2 = \sum_{i=1}^d \left(\sum_{j=1}^d \nabla_{x_i} \nabla_{x_j} g_j(x) \right)^2 \leq d \sum_{i=1}^d \sum_{j=1}^d |\nabla_{x_i} \nabla_{x_j} g_j(x)|^2 \leq d^2 \kappa^2.$$

Suppose X, Y, Z are distributed so that (X, Y) is a 1-Wasserstein optimal coupling of (μ, ν) and Z is independent of (X, Y). Since s_p is L-Lipschitz with $\mathbb{E}_P[s_p] = 0$ (Assumption 1), g is bounded (13), and g and $\nabla \cdot g$ are Lipschitz (14), repeated use of Cauchy-Schwarz gives

$$\mathbb{E}_\mu[\mathcal{T}_P g] - \mathbb{E}_\nu[\mathcal{T}_P g]$$
$$= \mathbb{E}[\nabla \cdot g(X) - \nabla \cdot g(Y)] + \mathbb{E}[\langle s_p(X) - s_p(Y), g(X) \rangle] + \mathbb{E}[\langle s_p(Y) - s_p(Z), g(X) - g(Y) \rangle]$$
$$\leq \kappa(d + L) W_1(\mu, \nu) + L \mathbb{E}[\|Y - Z\|_2 \min(2\kappa, \sqrt{d}\kappa\|X - Y\|_2)].$$

Since our choice of g was arbitrary, the first advertised result now follows from the definition of KSD (Definition 4). The second claim then follows from Cauchy-Schwarz and the inequality $\min(a, b)^2 \leq ab$ for $a, b \geq 0$, since

$$\mathbb{E}[\|Y - Z\|_2 \min(2\kappa, \sqrt{d}\kappa\|X - Y\|_2)] \leq M_{\nu,P}^{1/2} \mathbb{E}[\min(2\kappa, \sqrt{d}\kappa\|X - Y\|_2)^2]^{1/2}$$
$$\leq \sqrt{2M_{\nu,P}} d^{1/4} \kappa \mathbb{E}[\|X - Y\|_2]^{1/2} = \sqrt{2M_{\nu,P} W_1(\mu, \nu)} d^{1/4} \kappa.$$

Appendix D. Proof of Corollary 2: A finite-particle convergence rate for SVGD

We first state and prove a useful lemma.

Lemma 5 *Suppose f is a monotonic decreasing function. Let $x = f(\beta)$ for $\beta = \max(1, f(1))$. Then $x \leq f(x)$.*

Proof If $\beta = 1 \geq f(1)$, then $x = f(\beta) = f(1) \leq 1$. Because f is monotonic decreasing, $f(x) \geq f(1) = x$. Otherwise, $\beta = f(1) > 1$ and hence $x = f(\beta) < f(1) = \beta$. Again, due to monotonicity, $f(x) > f(\beta) = x$. Combining the two cases finishes the proof. ∎

Now we are ready to prove Corollary 2. Let $b_t^{(1)} = \psi_{B,C}(w_{0,n} \sqrt{\log \log \frac{1}{w_{0,n}}}, A, \beta_1)$ and $b_t^{(2)} = \psi_{B,C}(w_{0,n} \log \log \frac{1}{w_{0,n}}, A + 2C, \beta_2)$. According to Lemma 5,

$$b_t^{(1)} \leq \psi_{B,C}(w_{0,n} \sqrt{\log \log \frac{1}{w_{0,n}}}, A, b_t^{(1)}).$$

Rearranging the terms, we have

$$w_{0,n} \exp(b_t^{(1)}(A + B \exp(C b_t^{(1)}))) \leq \frac{1}{\sqrt{\log \log \frac{1}{w_{0,n}}}}. \tag{15}$$

Similarly, we have $b_t^{(2)} \leq \psi_{B,C}(w_{0,n} \log \log \frac{1}{w_{0,n}}, A + 2C, b_t^{(2)})$ and

$$\sqrt{w_{0,n}} \exp(b_t^{(2)}(2C + A + B \exp(Cb_t^{(2)}))/2) \leq \frac{1}{\sqrt{\log \log \frac{1}{w_{0,n}}}}. \tag{16}$$

Since $b_t = \min(b_t^{(1)}, b_t^{(2)})$, we can substitute b_t for $b_t^{(1)}$ and $b_t^{(2)}$ in (15) and (16), respectively. Also, from the definition of b_t we know $\frac{1}{b_t} = O\left(\frac{1}{\log \log \frac{1}{w_{0,n}}}\right)$. Plugging them into (2) proves the first claim. The second claim follows from the $O(n^{-1/d})$ convergence rate of an empirical measure to its population measure in 1-Wasserstein distance [5].