Fast Extra Gradient Methods for Smooth Structured Nonconvex-Nonconcave Minimax Problems

Abstract

Modern minimax problems, such as generative adversarial network and adversarial training, are often under a nonconvex-nonconcave setting, and developing an efficient method for such setting is of interest. Recently, two variants of the extragradient (EG) method are studied in that direction. First, a two-time-scale variant of the EG, named EG+, was proposed under a smooth structured nonconvex-nonconcave setting, with a slow $\mathcal{O}(1/k)$ rate on the squared gradient norm, where k denotes the number of iterations. Second, another variant of EG with an anchoring technique, named extra anchored gradient (EAG), was studied under a smooth convex-concave setting, yielding a fast $\mathcal{O}(1/k^2)$ rate on the squared gradient norm. Built upon EG+ and EAG, this paper proposes a two-time-scale EG with anchoring, named fast extragradient (FEG), that has a fast $\mathcal{O}(1/k^2)$ rate on the squared gradient norm for smooth structured nonconvex-nonconcave problems. This paper further develops its backtracking line-search version, named FEG-A, for the case where the problem parameters are not available. The stochastic analysis of FEG is also provided.

1 Introduction

Recently, nonconvex-nonconcave minimax problems have received an increased attention in the optimization community and the machine learning community due to their applications to generative adversarial network [9] and adversarial training [22]. In this paper, we consider a smooth structured nonconvex-nonconcave minimax problem:

$$\min_{\boldsymbol{x} \in \mathbb{R}^{d_x}} \max_{\boldsymbol{y} \in \mathbb{R}^{d_y}} f(\boldsymbol{x}, \boldsymbol{y}), \tag{1}$$

where $f : \mathbb{R}^{d_x} \times \mathbb{R}^{d_y} \to \mathbb{R}$ is smooth and is possibly nonconvex in $\boldsymbol{x}$ for fixed $\boldsymbol{y}$, and possibly nonconcave in $\boldsymbol{y}$ for fixed $\boldsymbol{x}$. We construct an efficient (first-order) method, using a saddle gradient operator $\boldsymbol{F} := (\nabla_x f, -\nabla_y f)$, for finding a first-order stationary point of the problem (1).

So far little is known under the nonconvex-nonconcave setting, compared to the convex-concave setting. Recent works [4, 7, 18, 19, 21, 37, 39] studied extragradient-type methods [15, 34] for minimax problems under various structured nonconvex-nonconcave settings. In other words, they consider various non-monotone conditions on $\boldsymbol{F}$, such as the Minty variational inequality (MVI) condition [4], the weak MVI condition [7], and the negative comonotonicity [1].[1] Among them, this paper focuses on the negative comonotonicity condition for a Lipschitz continuous $\boldsymbol{F}$. To the best of

Submitted to 35th Conference on Neural Information Processing Systems (NeurIPS 2021). Do not distribute.

---

[1]Relations between the conditions on $\boldsymbol{F}$ considered in this paper is summarized in Figure 1.

our knowledge, the following two-time-scale variant of the extragradient method, named EG+:

$$\boldsymbol{z}_{k+1/2} = \boldsymbol{z}_k - \frac{\alpha_k}{\beta}\boldsymbol{F}\boldsymbol{z}_k,$$
$$\boldsymbol{z}_{k+1} = \boldsymbol{z}_k - \alpha_k\boldsymbol{F}\boldsymbol{z}_{k+1/2}, \tag{EG+}$$

is the only known (explicit)[2] method, using $\boldsymbol{F}$, that converges under the considered setting[3] [7], where $\boldsymbol{z}_k := (\boldsymbol{x}_k, \boldsymbol{y}_k)$. The EG+, however, has a slow $\mathcal{O}(1/k)$ rate on the squared gradient norm.

Meanwhile, under the smooth convex-concave setting, recent works [6, 14, 16, 35, 38] suggest that Halpern-type [11] (or anchoring) methods, performing a convex combination of an initial point $\boldsymbol{z}_0$ and the last updated point $\boldsymbol{z}_k$ at each iteration, has a fast $\mathcal{O}(1/k^2)$ rate in terms of the squared gradient norm. In particular, [38] developed the following anchoring variant of the extragradient method, named extra anchored gradient (EAG):

$$\boldsymbol{z}_{k+1/2} = \boldsymbol{z}_k + \beta_k(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \alpha_k\boldsymbol{F}\boldsymbol{z}_k,$$
$$\boldsymbol{z}_{k+1} = \boldsymbol{z}_k + \beta_k(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \alpha_k\boldsymbol{F}\boldsymbol{z}_{k+1/2}. \tag{EAG}$$

This is the first (explicit) method with a fast $\mathcal{O}(1/k^2)$ rate on the squared gradient norm, when $\boldsymbol{F}$ satisfies both the Lipschitz continuity and the monotonicity. [38] also showed that such $\mathcal{O}(1/k^2)$ rate is optimal for first-order methods using a Lipschitz continuous and monotone $\boldsymbol{F}$.

Built upon both EG+ and EAG, this paper studies the following class of two-time-scale anchored extragradient methods, named fast extragradient (FEG):

$$\boldsymbol{z}_{k+1/2} = \boldsymbol{z}_k + \beta_k(\boldsymbol{z}_0 - \boldsymbol{z}_k) - (1 - \beta_k)(\alpha_k + 2\rho_k)\boldsymbol{F}\boldsymbol{z}_k,$$
$$\boldsymbol{z}_{k+1} = \boldsymbol{z}_k + \beta_k(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \alpha_k\boldsymbol{F}\boldsymbol{z}_{k+1/2} - (1 - \beta_k)2\rho_k\boldsymbol{F}\boldsymbol{z}_k. \tag{Class FEG}$$

Note that (Class FEG) reuses the $\boldsymbol{F}\boldsymbol{z}_k$ term in the $\boldsymbol{z}_{k+1}$ update, unlike standard extragradient-type methods. The FEG (with appropriately chosen step coefficients α_k, β_k and ρ_k discussed later) has an $\mathcal{O}(1/k^2)$ rate on the squared gradient norm, under the Lipschitz continuity and the negative comonotonicity conditions on $\boldsymbol{F}$. To the best of our knowledge, this is the first accelerated method under the nonconvex-nonconcave setting. The FEG also has value under the smooth convex-concave setting. First, when $\boldsymbol{F}$ is Lipschitz continuous and monotone, the rate bound of FEG is about 27/4 times smaller than that of EAG. Also note that the rate bound of FEG is only about four times larger than the $\mathcal{O}(1/k^2)$ lower complexity bound of first-order methods under such setting [38], further closing the gap between the lower and upper complexity bounds. Second, when $\boldsymbol{F}$ is cocoercive, FEG has a rate faster than that of a version of Halpern iteration [11] in [6].

We also develop an adaptive variant of FEG, named FEG-A, which updates its parameters, α_k and ρ_k in (Class FEG), adaptively using a backtracking line-search [2, 20, 26]. FEG requires the knowledge of the two problem parameters for the Lipschitz continuity and the comonotonicity of $\boldsymbol{F}$. However, those global parameters can be conservative, and in practice, they are even usually unknown. For such cases, the FEG-A adaptively and locally estimates the problem parameters, while preserving the fast rate $\mathcal{O}(1/k^2)$ on the squared gradient norm for smooth structured nonconvex-nonconcave minimax problems.

Lastly, we study a stochastic version of FEG, named S-FEG, which uses an unbiased stochastic estimate of $\boldsymbol{F}\boldsymbol{z}$, i.e., $\tilde{\boldsymbol{F}}\boldsymbol{z} = \boldsymbol{F}\boldsymbol{z} + \xi$, instead of $\boldsymbol{F}\boldsymbol{z}$ in FEG, where ξ denotes a stochastic noise. For a Lipschitz continuous and monotone $\boldsymbol{F}$, we provide a convergence analysis in terms of the expected squared gradient norm. In specific, we show that the S-FEG is stable with a rate $\mathcal{O}(1/k^2) + \mathcal{O}(\epsilon)$, when the noise variance decreases in the order of $\mathcal{O}(\epsilon/k)$, while being unstable otherwise due to error accumulation. This is similar to the convergence behavior of a stochastic version of Nesterov's fast gradient method [30, 31], observed in [5], for smooth convex minimization.

Our main contributions are summarized as follows.

- We propose the FEG method that has an accelerated convergence rate $\mathcal{O}(1/k^2)$ on the squared gradient norm for smooth structured nonconvex-nonconcave minimax problems.

---

[2]A proximal point method converges under the negative comonotonicity [1], but such implicit method is not preferable over explicit methods in practice due to its implicit nature.

[3]The EG+ was originally shown to work under the weak MVI condition of $\boldsymbol{F}$, which is weaker than the negative comonotonicity.

- We present that the FEG method has a rate faster than those of the EAG and the Halpern iteration for smooth convex-concave problems.

- We construct a backtracking line-search version of FEG, named FEG-A, for the case where the Lipschitz constant and comonotonicity parameters of $\boldsymbol{F}$ are unavailable.

- We analyze a stochastic version of FEG, named S-FEG, for smooth convex-concave problems.

2 Related work

2.1 Methods for convex-concave minimax problems

The extragradient method [15] is one of the widely used methods for solving smooth convex-concave minimax problems (see, *e.g.*, [4, 7, 18, 19, 21, 37, 39] for its extensions and applications). In terms of the *duality gap*, $\max_{\boldsymbol{y}' \in \mathcal{Y}} f(\boldsymbol{x}, \boldsymbol{y}') - \min_{\boldsymbol{x}' \in \mathcal{X}} f(\boldsymbol{x}', \boldsymbol{y})$, where $\mathcal{X}$ and $\mathcal{Y}$ are compact[4] domains, the ergodic iterate of the extragradient-type methods [27, 32] have an $\mathcal{O}(1/k)$ rate. Such $\mathcal{O}(1/k)$ rate on the duality gap is order-optimal for the first-order methods [29, 33], leaving no room for improvement. On the other hand, the last iterate of the extragradient method has a slower $\mathcal{O}(1/\sqrt{k})$ rate on the duality gap, under an additional assumption that $\boldsymbol{F}$ has a Lipschitz derivative [8]. In terms of the *squared gradient norm*, $\|\boldsymbol{F}\boldsymbol{z}\|^2$, the best iterate of the extragradient-type methods [15, 34] have an $\mathcal{O}(1/k)$ rate [35, 36, 38]. The last iterate of the extragradient method also has a rate $\mathcal{O}(1/k)$, when $\boldsymbol{F}$ is further assumed to have a Lipschitz derivative [8]. Unlike the duality gap, the $\mathcal{O}(1/k)$ rate on the squared gradient norm is not optimal [38]. From now on throughout this paper, we mainly study and compare the convergence rates on the squared gradient norm, which still has room for improvement in convex-concave problems, and has meaning for nonconvex-nonconcave minimax problems, unlike the duality gap.

Recently, [6, 14, 16, 35, 38] found that Halpern-type [11] (or anchoring) methods yield a fast $\mathcal{O}(1/k^2)$ rate in terms of the squared gradient norm for minimax problems. [14, 16] showed that the (implicit) Halpern iteration [11] with appropriately chosen step coefficients has an $\mathcal{O}(1/k^2)$ rate on the squared norm of a monotone $\boldsymbol{F}$. Then, for a cocoercive $\boldsymbol{F}$, an (explicit) version of the Halpern iteration was studied in [6, 14] that has the same fast rate. In addition, [6] constructed a double-loop version of the Halpern iteration for a Lipschitz continuous and monotone $\boldsymbol{F}$, which has a rate $\tilde{\mathcal{O}}(1/k^2)$ on the squared gradient norm, slower than the rate $\mathcal{O}(1/k^2)$. While this is promising compared to the $\mathcal{O}(1/k)$ rate of the extragradient methods on the squared gradient norm [35, 36, 38], the computational complexity due to its double-loop nature and a relatively slow rate remained a problem. Very recently, [38] proposed the extra anchored gradient (EAG) method, which is the first (explicit) method with a fast $\mathcal{O}(1/k^2)$ rate for smooth convex-concave minimax problems, *i.e.*, for Lipschitz continuous and monotone operators. In addition, [38] proved that the EAG is order-optimal by showing that the lower complexity bound of first-order methods is $\mathcal{O}(1/k^2)$.

2.2 Methods for nonconvex-nonconcave minimax problems

Some recent literature considered relaxing the monotonicity condition of the saddle gradient operator to tackle modern nonconvex-nonconcave minimax problems. For example, the Minty variational inequality (MVI) condition, *i.e.*, there exists $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$ satisfying $\langle \boldsymbol{F}\boldsymbol{z}, \boldsymbol{z} - \boldsymbol{z}_* \rangle \geq 0$ for all $\boldsymbol{z} \in \mathbb{R}^d$ where $\boldsymbol{Z}_*(\boldsymbol{F}) := \{\boldsymbol{z}_* \in \mathbb{R}^d : \boldsymbol{F}\boldsymbol{z}_* = \boldsymbol{0}\}$, is studied in [4, 17, 18, 19]. This condition is also studied under the name, the coherence, in [21, 37, 39]. Moreover, [7] considered a weaker condition, named the weak MVI condition, *i.e.*, for some $\rho < 0$, there exists $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$ satisfying

Figure 1: Relations between the conditions on $\boldsymbol{F}$.

---

[4] The convergence analysis on the duality gap of the extragradient type methods are generalized under the unbounded domain assumption in [23, 24, 25].

$\langle \boldsymbol{F}\boldsymbol{z}, \, \boldsymbol{z} - \boldsymbol{z}_* \rangle \geq \rho \|\boldsymbol{F}\boldsymbol{z}\|^2$ for all $\boldsymbol{z} \in \mathbb{R}^d$. The weak MVI condition is implied by the negative comonotonicity [1] or, equivalently, the (positive) cohypomonotonicity [3]. The comonotonicity will be further discussed in the upcoming section.

For L-Lipschitz continuous $\boldsymbol{F}$, [4, 37] showed that the extragradient-type methods have an $\mathcal{O}(1/k)$ rate on the squared gradient norm under the MVI condition, and [7] developed the (EG+) method under the weak MVI condition (and thus under the negative comonotonicty), which also has an $\mathcal{O}(1/k)$ rate on the squared gradient norm. To the best of our knowledge, there is no known accelerated method for the nonconvex-nonconcave setting; our proposed FEG method is the first method to have a fast $\mathcal{O}(1/k^2)$ rate under the nonconvex-nonconcave setting. The convergence rates of the existing methods and the FEG on the squared gradient norm are summarized in Table 1.

Table 1: Comparison of the convergence rates of the existing extragradient-type methods and the FEG, with respect to the squared gradient norm, for smooth structured minimax problems, under various assumptions on the Lipschitz continuous saddle gradient operator $\boldsymbol{F}$.

Method		Convex-concave		Nonconvex-nonconcave		
		Cocoercive	Monotone	Negative comonotone	MVI	Weak MVI
Normal	EG [4, 37]	$\mathcal{O}(1/k)$	$\mathcal{O}(1/k)$		$\mathcal{O}(1/k)$	
	EG+ [7]	$\mathcal{O}(1/k)$	$\mathcal{O}(1/k)$	$\mathcal{O}(1/k)$	$\mathcal{O}(1/k)$	$\mathcal{O}(1/k)$
Accelerated	Halpern [11, 6]	$\mathcal{O}(1/k^2)$	$\tilde{\mathcal{O}}(1/k^2)$			
	EAG [38]	$\mathcal{O}(1/k^2)$	$\mathcal{O}(1/k^2)$			
	FEG (this paper)	$\mathcal{O}(1/k^2)$	$\mathcal{O}(1/k^2)$	$\mathcal{O}(1/k^2)$		

3 Preliminaries

The followings are the two main assumptions for the saddle gradient operator $\boldsymbol{F}$ of the smooth structured nonconvex-nonconcave problem (1). Under such assumptions, we develop efficient methods that find a first-order stationary point $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$ where $\boldsymbol{Z}_*(\boldsymbol{F}) := \{\boldsymbol{z}_* \in \mathbb{R}^d : \boldsymbol{F}\boldsymbol{z}_* = \boldsymbol{0}\}$.

Assumption 1 (L-Lipschitz continuity)**.** *For some $L \in (0, \infty)$, $\boldsymbol{F}$ satisfies*

$$\|\boldsymbol{F}\boldsymbol{z} - \boldsymbol{F}\boldsymbol{z}'\| \leq L\|\boldsymbol{z} - \boldsymbol{z}'\|, \quad \forall \boldsymbol{z}, \boldsymbol{z}' \in \mathbb{R}^d.$$

Assumption 2 (ρ-Comonotonicity)**.** *For some $\rho \in \mathbb{R}$, $\boldsymbol{F}$ satisfies*

$$\langle \boldsymbol{F}\boldsymbol{z} - \boldsymbol{F}\boldsymbol{z}', \, \boldsymbol{z} - \boldsymbol{z}' \rangle \geq \rho \|\boldsymbol{F}\boldsymbol{z} - \boldsymbol{F}\boldsymbol{z}'\|^2, \quad \forall \boldsymbol{z}, \boldsymbol{z}' \in \mathbb{R}^d.$$

The ρ-comonotonicity consists of three cases depending on the choice of ρ; the negative comonotonicity when $\rho < 0$, the monotonicity when $\rho = 0$, and the cocoercivity when $\rho > 0$. The negative comonotonicity is weaker than the other two, and is the main focus of this paper. The following is an examplary nonconvex-nonconcave condition that is stronger than the negative comonotonicity.

Example 1. *Let f be twice continuously differentiable and γ-smooth. Further assume that f satisfies*

$$\nabla_{\boldsymbol{x}\boldsymbol{x}}^2 f + \nabla_{\boldsymbol{x}\boldsymbol{y}}^2 f(\eta\boldsymbol{I} - \nabla_{\boldsymbol{y}\boldsymbol{y}}^2 f)^{-1}\nabla_{\boldsymbol{y}\boldsymbol{x}}^2 f \succeq \alpha\boldsymbol{I},$$
$$-\nabla_{\boldsymbol{y}\boldsymbol{y}}^2 f + \nabla_{\boldsymbol{y}\boldsymbol{x}}^2 f(\eta\boldsymbol{I} + \nabla_{\boldsymbol{x}\boldsymbol{x}}^2 f)^{-1}\nabla_{\boldsymbol{x}\boldsymbol{y}}^2 f \succeq \alpha\boldsymbol{I},$$

for some $\alpha \geq 0$ and $\eta > \gamma$, named α-interaction dominate condition in [10]. Then, the saddle gradient of f satisfies the negative comonotonicity. (See Appendix A.1.)

We next present our proposed FEG, and illustrate that the FEG outperforms existing methods such as EG+, EAG, and the Halpern iteration, for each three comonoticity case, respectively.

4 Fast extragradient (FEG) method for Lipschitz continuous and comonotone operators

This section considers an instance of (Class FEG) with $\alpha_k = \frac{1}{L}$, $\beta_k = \frac{1}{k+1}$, and $\rho_k = \rho$ for all $k \geq 0$. The resulting method, named FEG, is illustrated in Algorithm 1, which has an $\mathcal{O}(1/k^2)$ fast rate with respect to the squared gradient norm.

Algorithm 1 Fast extragradient (FEG) method

Input: $\boldsymbol{z}_0 \in \mathbb{R}^d$, $L \in (0, \infty)$, $\rho \in \left(-\frac{1}{2L}, \infty\right)$
for $k = 0, 1, \ldots$ **do**

$$\boldsymbol{z}_{k+1/2} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \left(1 - \frac{1}{k+1}\right)\left(\frac{1}{L} + 2\rho\right)\boldsymbol{F}\boldsymbol{z}_k$$

$$\boldsymbol{z}_{k+1} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \frac{1}{L}\boldsymbol{F}\boldsymbol{z}_{k+1/2} - \left(1 - \frac{1}{k+1}\right)2\rho\boldsymbol{F}\boldsymbol{z}_k.$$

end for

Theorem 4.1. *For the L-Lipschitz continuous and ρ-comonotone operator $\boldsymbol{F}$ with $\rho > -\frac{1}{2L}$ and for any $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$, the sequence $\{\boldsymbol{z}_k\}_{k \geq 0}$ generated by FEG satisfies, for all $k \geq 1$,*

$$\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{4\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{\left(\frac{1}{L} + 2\rho\right)^2 k^2}. \tag{2}$$

The following example shows that the bound (2) of the FEG is exact for $\rho = 0$ and $k = 4l + 2$. The bound (2) is not known to be exact in general, and we leave finding the exact bound as future work.

Example 2. *Let $f : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ be $f(x, y) = Lxy$. Its saddle gradient operator and solution are $\boldsymbol{F}(x, y) = (Ly, -Lx)$ and $\boldsymbol{z}_* = (0, 0)$, respectively. For the initial point $\boldsymbol{z}_0 = (x_0, y_0) = (1, 0)$, the sequence $\{\boldsymbol{z}_k\}_{k \geq 0}$ generated by FEG satisfies $\boldsymbol{z}_{4l+2} = \left(0, \frac{1}{2l+1}\right)$ for all $l \geq 0$. Hence, $\|\boldsymbol{F}\boldsymbol{z}_{4l+2}\|^2 = \frac{L^2}{(2l+1)^2} = \frac{4L^2\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{(4l+2)^2}$ for all $l \geq 0$.*

We next compare the rate bound (2) with existing analyses for the three cases $-\frac{1}{2L} < \rho < 0$, $\rho = 0$, and $\rho > 0$.

4.1 Comparison to EG+ under the negative comonotonicity ($\rho < 0$)

Under the negative comonotonicity with $-\frac{1}{8L} < \rho < 0$, the (EG+) method with $\alpha_k = \frac{1}{2L}$ and $\beta = \frac{1}{2}$ has an $\mathcal{O}(1/k)$ rate on the squared gradient norm. To the best of our knowledge, this is the best known rate, and the FEG has a faster $\mathcal{O}(1/k^2)$ rate with a wider region of convergence $-\frac{1}{2L} < \rho < 0$.

4.2 Comparison to EAG under the monotonicity ($\rho = 0$)

For an L-Lipschitz continuous and monotone operator $\boldsymbol{F}$, [38] proposed two EAG methods, named EAG-C and EAG-V, with same $\beta_k = \frac{1}{k+2}$ but with different choices of α_k. EAG-C sets α_k to be a constant $\frac{1}{8L}$ for all $k \geq 0$ in (EAG), and has a large constant 260 in its convergence rate, $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{260L^2\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{(k+1)^2}$ for all $k \geq 0$. On the other hand, while EAG-V requires a complicated recursive update for $\{\alpha_k\}$, $\alpha_{k+1} = \frac{\alpha_k}{1 - \alpha_k^2 L^2}\left(1 - \frac{(k+2)^2}{(k+1)(k+3)}\alpha_k^2 L^2\right)$ for all $k \geq 0$, with $\alpha_0 = \frac{0.618}{L}$, its rate has a smaller constant 27.

The FEG takes a constant $\alpha_k = \frac{1}{L}$, unlike EAG-V, but has an even smaller constant 4 in its convergence rate $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{4L^2\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{k^2}$ for $\rho = 0$. Therefore, the FEG with $\rho = 0$ has about 260/4-times and 27/4-times faster convergence rate compared to those of EAG-C and EAG-V, respectively. Furthermore, the rate bound of FEG with $\rho = 0$ is only about 4-times larger than the lower complexity bound of first-order methods under the considered setting [38], reducing the gap between the lower and upper complexity bounds from 27 to 4.

4.3 Comparison to the Halpern iteration under the cocoercivity ($\rho > 0$)

For a ρ-cocoercive operator $\boldsymbol{F}$, an (explicit) version of Halpern iteration [11], studied in [6], has a fast rate, $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{\rho^2 k^2}$. Note that while the ρ-cocoercivity implies the $\frac{1}{\rho}$-Lipschitz continuity, there is case where the ρ-cocoercive (and thus Lipschitz continuous) operator has a Lipschitz constant L smaller than $\frac{1}{\rho}$. Since $L \leq \frac{1}{\rho}$, the FEG has a rate $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{4\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{(1/L + 2\rho)^2 k^2} = \frac{4\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{9\rho^2 k^2}$ that

is faster than that of Halpern iteration. However, if we take into account that the FEG requires computing the saddle gradient twice per iteration, unlike Halpern iteration studied in [6], the FEG method has a slower rate in terms of the number of gradient computations. If we narrow down to the case $L < \frac{1}{2\rho}$, the FEG has a faster rate, $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{4\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{(1/L + 2\rho)^2 k^2} < \frac{\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{4\rho^2 k^2}$. For such case, the FEG has a rate faster than that of the Halpern iteration, even in terms of the number of gradient computations.

5   FEG with backtracking line-search

The FEG requires the knowledge of the two global parameters L and ρ for Lipschitz continuity and comonotonicity, respectively. Those global parameters are often difficult to compute in practice and can be locally conservative. To handle these two disadvantages, we employ the backtracking line-search technique [2, 20, 26] in FEG, which adaptively decreases the two parameters for step size, τ and η, to satisfy the both conditions, the local $\frac{1}{\tau}$-Lipschitz continuity and the $\frac{\eta - \tau}{2}$-comonotonicity. A pseudocode of the resulting method, named FEG-A, is illustrated in Algorithm 2. For a detailed description of the FEG-A, see Algorithm 4 in Appendix B.1.

Algorithm 2 Fast extragradient method with adaptive step size (FEG-A)

Input: $\boldsymbol{z}_0 \in \mathbb{R}^d$, $\tau_{-1} \in \left(\frac{1-\delta}{L}, \infty\right)$, $\eta_0 \in \left(\frac{(1-\delta)^2}{L} + (1-\delta)2\rho, \infty\right)$, $\delta \in (0, 1)$
Find the smallest nonnegative integer i_0 such that $\hat{\boldsymbol{z}} = \boldsymbol{z}_0 - \hat{\tau}\boldsymbol{F}\boldsymbol{z}_0$ satisfies $\hat{\tau}\|\boldsymbol{F}\hat{\boldsymbol{z}} - \boldsymbol{F}\boldsymbol{z}_0\| \leq \|\hat{\boldsymbol{z}} - \boldsymbol{z}_0\|$ where $\hat{\tau} = \tau_{-1}(1-\delta)^{i_0}$.
$\tau_0 = \tau_{-1}(1-\delta)^{i_0}$, $\boldsymbol{z}_1 = \boldsymbol{z}_0 - \tau_0\boldsymbol{F}\boldsymbol{z}_0$.
for $k = 1, 2, \ldots$ **do**
  $i_k = j_k = 0$.
  Increase each i_k and j_k one by one until

$$\hat{\boldsymbol{z}}_{k+1/2} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \left(1 - \frac{1}{k+1}\right)\eta_{k-1}(1-\delta)^{j_k}\boldsymbol{F}\boldsymbol{z}_k \qquad \text{and}$$

$$\hat{\boldsymbol{z}}_{k+1} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \tau_{k-1}(1-\delta)^{i_k}\boldsymbol{F}\boldsymbol{z}_{k+1/2}$$
$$- \left(1 - \frac{1}{k+1}\right)\left(\eta_{k-1}(1-\delta)^{j_k} - \tau_{k-1}(1-\delta)^{i_k}\right)\boldsymbol{F}\boldsymbol{z}_k$$

  satisfy both conditions,

$$\|\boldsymbol{F}\hat{\boldsymbol{z}}_{k+1} - \boldsymbol{F}\hat{\boldsymbol{z}}_{k+1/2}\| \leq \frac{1}{\tau_{k-1}(1-\delta)^{i_k}}\|\hat{\boldsymbol{z}}_{k+1} - \hat{\boldsymbol{z}}_{k+1/2}\| \qquad \text{and}$$

$$\langle \boldsymbol{F}\hat{\boldsymbol{z}}_{k+1} - \boldsymbol{F}\boldsymbol{z}_k, \hat{\boldsymbol{z}}_{k+1} - \boldsymbol{z}_k \rangle \leq \frac{\eta_{k-1}(1-\delta)^{j_k} - \tau_{k-1}(1-\delta)^{i_k}}{2}\|\boldsymbol{F}\hat{\boldsymbol{z}}_{k+1} - \boldsymbol{F}\boldsymbol{z}_k\|^2.$$

  $\tau_k = \tau_{k-1}(1-\delta)^{i_k}$, $\eta_k = \eta_{k-1}(1-\delta)^{j_k}$, $\boldsymbol{z}_{k+1} = \hat{\boldsymbol{z}}_{k+1}$.
end for

The following lemma shows that each of the decreasing sequences $\{\tau_k\}_{k \geq 0}$ and $\{\eta_k\}_{k \geq 0}$ of FEG-A is lower bounded and thus FEG-A is well-defined.

Lemma 5.1. *For the L-Lipschitz and ρ-comonotone operator F and a given constant $\delta \in (0, 1)$, the sequences $\{\tau_k\}_{k \geq 0}$ and $\{\eta_k\}_{k \geq 0}$ generated by FEG-A are lower bounded by $\frac{1-\delta}{L}$ and $\frac{(1-\delta)^2}{L} + (1 - \delta)2\rho$, respectively.*

The FEG-A method also has the following $\mathcal{O}(1/k^2)$ rate with respect to the squared gradient norm.

Theorem 5.1. *For the L-Lipschitz and ρ-comonotone operator F with $\rho > -\frac{1-\delta}{2L}$ and for any $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$, the sequence $\{\boldsymbol{z}_k\}_{k \geq 0}$ generated by FEG-A satisfies, for all $k \geq 1$,*

$$\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq \frac{4\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{((k-1)(1-\delta) + 1)^2 \left(\frac{1-\delta}{L} + 2\rho\right)^2}.$$

6 FEG under stochastic setting

When exactly computing $\boldsymbol{F z}$ is expensive in practice, one usually instead consider its stochastic estimate for computational efficiency (see, *e.g.*, [12, 13, 21, 28, 35, 37, 39]). This section also considers using a stochastic oracle in FEG for smooth convex-concave problems. In specific, this section assumes that we only have access to a noisy saddle gradient oracle, $\tilde{\mathbf{F}} \boldsymbol{z}_{k/2} = \boldsymbol{F z}_{k/2} + \xi_{k/2}$, where $\{\xi_{k/2}\}_{k \geq 0}$ are independent random variables satisfying $\mathbb{E}[\xi_{k/2}] = 0$ and $\mathbb{E}[\|\xi_{k/2}\|^2] = \sigma_{k/2}^2$ for all $k \geq 0$. Under this setting, we study a stochastic first-order method, named stochastic fast extragradient (S-FEG) method, illustrated in Algorithm 3.

Algorithm 3 Stochastic fast extragradient (S-FEG) method

Input: $\boldsymbol{z}_0 \in \mathbb{R}^d$, $L \in (0, \infty)$, $\rho \in \left(-\frac{1}{2L}, \infty\right)$
for $k = 0, 1, \dots$ **do**

$$\boldsymbol{z}_{k+1/2} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \left(1 - \frac{1}{k+1}\right)\frac{1}{L}\tilde{\mathbf{F}}\boldsymbol{z}_k$$

$$\boldsymbol{z}_{k+1} = \boldsymbol{z}_k + \frac{1}{k+1}(\boldsymbol{z}_0 - \boldsymbol{z}_k) - \frac{1}{L}\tilde{\mathbf{F}}\boldsymbol{z}_{k+1/2}$$

end for

The following theorem provides an upper bound of the expected squared gradient norm for the S-FEG.

Theorem 6.1. *Let* $\tilde{\mathbf{F}}\boldsymbol{z}_{k/2} = \boldsymbol{F z}_{k/2} + \xi_{k/2}$*, where* $\{\xi_{k/2}\}_{k \geq 0}$ *are independent random variables satisfying* $\mathbb{E}[\xi_{k/2}] = 0$ *and* $\mathbb{E}[\|\xi_{k/2}\|^2] = \sigma_{k/2}^2$ *for all* $k \geq 0$*. Then, for the L-Lipschitz continuous and monotone operator F and for any* $\boldsymbol{z}_* \in \boldsymbol{Z}_*(\boldsymbol{F})$*, the sequence* $\{\boldsymbol{z}_k\}_{k \geq 0}$ *generated by S-FEG satisfies*

$$\mathbb{E}[\|\boldsymbol{F z}_k\|^2] \leq \frac{4L^2\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{k^2} + \frac{6}{k^2}\left[\sigma_0^2 + \sum_{l=1}^{k-1}(l^2\sigma_l^2 + (l+1)^2\sigma_{l+1/2}^2)\right] \tag{3}$$

for all $k \geq 1$*. Furthermore, if* $\sigma_k^2 \leq \frac{\epsilon}{6k}$ *and* $\sigma_{k+1/2}^2 \leq \frac{\epsilon}{6(k+1)}$ *for all* $k \geq 0$*, then the bound* (3) *reduces to*

$$\mathbb{E}[\|\boldsymbol{F z}_k\|^2] \leq \frac{4L^2\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|^2}{k^2} + \epsilon$$

for all $k \geq 1$*.*

Here, we needed the noise variance $\sigma_{k/2}^2$ to decrease in the order of $\mathcal{O}(1/k)$ so that the stochastic error of the S-FEG does not accumulate. Otherwise, if $\sigma_{k/2}^2$ is a constant for all k, the error accumulates with rate $\mathcal{O}(k)$. In short, the S-FEG will suffer from error accumulation, unless the stochastic error decreases with rate $\mathcal{O}(1/k)$. Such error accumulation behavior also appears in a stochastic version of Nesterov's fast gradient method [30, 31] for smooth convex minimization [5]. Similar to [5], we believe that adjusting the step coefficients of the S-FEG can make the S-FEG become relatively stable even with a constant noise, which we leave as future work.

7 Convergence analysis with nonincreasing potential lemma

We analyze FEG and FEG-A by finding a nonincreasing potential function in a form $V_k = a_k\|\boldsymbol{F z}_k\|^2 - b_k\langle\boldsymbol{F z}_k, \boldsymbol{z}_0 - \boldsymbol{z}_k\rangle$ in the lemma below. The convergence analyses of EAG and Halpern iteration are also based on such potential function [6, 38].

Lemma 7.1. *Let* $\{\boldsymbol{z}_k\}_{k \geq 0}$ *be the sequence generated by* (Class FEG) *with* $\{\alpha_k\}_{k \geq 0}$*,* $\{\beta_k\}_{k \geq 0}$*,* $\{L_k\}_{k \geq 0} \subset (0, \infty)$ *and* $\{\rho_k\}_{k \geq 0} \subset \mathbb{R}$*, satisfying* $\alpha_0 \in (0, \infty)$*,* $\alpha_k \in \left(0, \frac{1}{L_k}\right]$*,* $\beta_0 = 1$*,* $\{\beta_k\}_{k \geq 1} \subseteq (0, 1)$ *for all* $k \geq 1$*, and*

$$\frac{(1 - \beta_{k+1})}{2\beta_{k+1}}(\alpha_{k+1} + 2\rho_{k+1}) - \rho_{k+1} \leq \frac{1}{2\beta_k}(\alpha_k + 2\rho_k) - \rho_k$$

227 *for all $k \geq 0$. Assume that the following conditions are satisfied.*

$$\|\boldsymbol{F}\boldsymbol{z}_1 - \boldsymbol{F}\boldsymbol{z}_0\| \leq L_0\|\boldsymbol{z}_1 - \boldsymbol{z}_0\|$$
$$\|\boldsymbol{F}\boldsymbol{z}_{k+1} - \boldsymbol{F}\boldsymbol{z}_{k+1/2}\| \leq L_k\|\boldsymbol{z}_{k+1} - \boldsymbol{z}_{k+1/2}\| \quad \text{for all } k \geq 1,$$
$$\langle \boldsymbol{F}\boldsymbol{z}_{k+1} - \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_{k+1} - \boldsymbol{z}_k \rangle \geq \rho_k\|\boldsymbol{F}\boldsymbol{z}_{k+1} - \boldsymbol{F}\boldsymbol{z}_k\|^2 \quad \text{for all } k \geq 1.$$

228 *Then a potential function*

$$V_k = a_k\|\boldsymbol{F}\boldsymbol{z}_k\|^2 - b_k \langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_0 - \boldsymbol{z}_k \rangle$$

229 *with $a_0 = \frac{\alpha_0(L_0^2\alpha_0^2 - 1)}{2}$, $b_0 = 0$, $b_1 = 1$,*

$$a_k = \frac{b_k(1 - \beta_k)}{2\beta_k}(\alpha_k + 2\rho_k) - b_k\rho_k \quad \text{and} \quad b_{k+1} = \frac{b_k}{1 - \beta_k}$$

230 *for all $k \geq 1$ satisfies $V_k \leq V_{k-1}$ for all $k \geq 1$.*

231 Based on the above potential lemma, we next provide a convergence analysis of FEG. The analyses
232 for the convergence rate of FEG-A and S-FEG, *i.e.*, the proofs of Theorem 5.1 and Theorem 6.1, are
233 similar to that of FEG and are provided in Appendix C.2 and Appendix D.3.

7.1 Convergence analysis for FEG

235 *Proof of Theorem 4.1.* Recall that FEG is equivalent to (Class FEG) with $\alpha_k = \frac{1}{L}$, $\beta_k = \frac{1}{k+1}$, and
236 $\rho_k = \rho$. It is straightforward to verify that the given $\{\alpha_k\}_{k \geq 0}$ and $\{\beta_k\}_{k \geq 0}$ satisfy the conditions in
237 Lemma 7.1 with $L_k = L$ for all $k \geq 0$. Since

$$a_k = \frac{b_k(1 - \beta_k)}{2\beta_k}(\alpha_k + 2\rho_k) - b_k\rho_k = \frac{k^2}{2}\left(\frac{1}{L} + 2\rho\right) - k\rho \qquad \text{and}$$

$$b_k = \frac{1}{1 - \beta_{k-1}}b_{k-1} = \left(\prod_{i=1}^{k-1}\frac{1}{1 - \beta_i}\right)b_1 = k,$$

238 Lemma 7.1 implies that

$$0 = V_1 \geq V_k = \left(\frac{k^2}{2}\left(\frac{1}{L} + 2\rho\right) - k\rho\right)\|\boldsymbol{F}\boldsymbol{z}_k\|^2 - k\langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_0 - \boldsymbol{z}_k \rangle.$$

239 Therefore,

$$\frac{k^2}{2}\left(\frac{1}{L} + 2\rho\right)\|\boldsymbol{F}\boldsymbol{z}_k\|^2 \leq k\langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_0 - \boldsymbol{z}_k \rangle + k\rho\|\boldsymbol{F}\boldsymbol{z}_k\|^2$$
$$= k\langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_0 - \boldsymbol{z}_* \rangle + k\langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_* - \boldsymbol{z}_k \rangle + k\rho\|\boldsymbol{F}\boldsymbol{z}_k\|^2$$
$$\leq k\langle \boldsymbol{F}\boldsymbol{z}_k, \, \boldsymbol{z}_0 - \boldsymbol{z}_* \rangle \qquad (\because \rho\text{-comonotonicity of } \boldsymbol{F})$$
$$\leq k\|\boldsymbol{F}\boldsymbol{z}_k\|\|\boldsymbol{z}_0 - \boldsymbol{z}_*\|.$$

240 The desired result follows directly by dividing both sides by $\frac{k^2}{2}\left(\frac{1}{L} + 2\rho\right)\|\boldsymbol{F}\boldsymbol{z}_k\|$. $\qquad\square$

8 Discussion: first-order methods for Lipschitz continuous operators

242 Throughout this paper, we studied and constructed efficient methods in a class of first-order methods:

$$\boldsymbol{z}_k \in \boldsymbol{z}_0 + \text{span}\{\boldsymbol{F}\boldsymbol{z}_0, \cdots, \boldsymbol{F}\boldsymbol{z}_k\}$$

243 denoted by $\mathcal{A}$, for smooth structured nonconvex-nonconcave problems. We observed that all existing
244 first-order methods, including the FEG, required an additional condition, such as the negative
245 comonoticity, on a Lipschitz continuous $\boldsymbol{F}$ to guarantee convergence. One would then be curious
246 whether or not there exists an (efficient) method in class $\mathcal{A}$ that guarantees convergence without any
247 additional condition on a Lipschitz continuous $\boldsymbol{F}$. Unfortunately, the following lemma states that
248 there exists a *worst-case* smooth example that none of the methods in $\mathcal{A}$ can find its stationary point.
249 The corresponding smooth function is illustrated in Figure 2.

Figure 2: A smooth worst-case example $f(x,y)$ (4) with $L = R = 1$ for first-order methods. any sequence $\{\boldsymbol{z}_k\}_{k\geq 0}$ generated by a first-order method in class $\mathcal{A}$ starting from $(0,0)$ is contained in the line $x = y$.

Lemma 8.1. *Let us consider the following L-smooth function $f : \mathbb{R}^2 \to \mathbb{R}$ for some $L, R > 0$:*

$$f(x,y) = \begin{cases} \frac{R}{2} & \text{for } x < y - \sqrt{\frac{R}{L}} \\ -\frac{L}{2}(x-y)^2 - \sqrt{LR}(x-y) & \text{for } y - \sqrt{\frac{R}{L}} \leq x < y \\ \frac{L}{2}(x-y)^2 - \sqrt{LR}(x-y) & \text{for } y \leq x < y + \sqrt{\frac{R}{L}} \\ -\frac{R}{2} & \text{for } y + \sqrt{\frac{R}{L}} < x. \end{cases} \tag{4}$$

The sequence $\{\boldsymbol{z}_k\}_{k\geq 0}$ generated by any first-order method in class $\mathcal{A}$ with $\boldsymbol{z}_0 = (0,0)$ satisfies $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 = 2LR$ for all $k \geq 0$.

Proof. *$\boldsymbol{F}$ satisfies $\boldsymbol{F}(x,y) = (-\sqrt{LR}, -\sqrt{LR})$ whenever $x = y$. Hence, for all sequences $\{\boldsymbol{z}_k\}_{k\geq 0}$ satisfying $\boldsymbol{z}_0 = (0,0)$ and $\boldsymbol{z}_k \in \boldsymbol{z}_0 + \text{span}\{\boldsymbol{F}\boldsymbol{z}_0, \cdots, \boldsymbol{F}\boldsymbol{z}_k\}$ for all $k \geq 0$, we have that $\{\boldsymbol{z}_k\}_{k\geq 0} \subseteq \{\boldsymbol{z} = (x,y) \in \mathbb{R}^2 | x = y\}$; thus, $\|\boldsymbol{F}\boldsymbol{z}_k\|^2 = 2LR$ for all $k \geq 0$.* $\square$

The lemma implies that one should consider a class of methods, other than the class $\mathcal{A}$, to guarantee finding a stationary point of any smooth problem, which we leave as future work. We also leave finding additional conditions for a Lipschitz continuous $\boldsymbol{F}$, weaker than the weak MVI condtion and the negative comonotonicity, which guarantee convergence or its accelerated rate, respectively.

9 Conclusion

This paper proposed a two-time-scale and anchored extragradient method, named FEG, for smooth structured nonconvex-nonconcave problems. The proposed FEG has an accelerated $\mathcal{O}(1/k^2)$ rate, with respect to the squared gradient norm, for the Lipschitz continuous and negative comonotone operators for the first time. The FEG also has value for smooth convex-concave problems, compared to existing works. We further studied its backtracking line-search version, named FEG-A, for the smooth structured nonconvex-nonconcave problems and studied its stochastic version, named S-FEG, for smooth convex-concave problems. We leave extending this work to stochastic, composite, or more general nonconvex-nonconcave settings as future work.

References

[1] H. H. Bauschke, W. M. Moursi, and X. Wang. Generalized monotone operators and their averaged resolvents. *Mathematical Programming*, 2020.

[2] A. Beck and M. Teboulle. A fast iterative shrinkage-thresholding algorithm for linear inverse problems. *SIAM J. Imaging Sci.*, 2(1):183–202, 2009.

[3] P. L. Combettes and T. Pennanen. Proximal methods for cohypomonotone operators. *SIAM J. Control Optim.*, 43(2):731–42, 2004.

[4] C. D. Dang and G. Lan. On the convergence properties of non-Euclidean extragradient methods for variational inequalities with generalized monotone operators. *Computational Optimization and Applications*, 60(2):277–310, 2015.

[5] O. Devolder. Stochastic first order methods in smooth convex optimization, 2011.

[6] J. Diakonikolas. Halpern iteration for near-optimal and parameter-free monotone inclusion and strong solutions to variational inequalities. In *Conference on Learning Theory*, pages 1428–1451. PMLR, 2020.

[7] J. Diakonikolas, C. Daskalakis, and M. Jordan. Efficient methods for structured nonconvex-nonconcave min-max optimization. In *International Conference on Artificial Intelligence and Statistics*, pages 2746–2754. PMLR, 2021.

[8] N. Golowich, S. Pattathil, C. Daskalakis, and A. Ozdaglar. Last iterate is slower than averaged iterate in smooth convex-concave saddle point problems. In *Conference on Learning Theory (COLT)*, 2020.

[9] I. J. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio. Generative adversarial networks. In *Neural Info. Proc. Sys.*, 2014.

[10] B. Grimmer, H. Lu, P. Worah, and V. Mirrokni. The landscape of the proximal point method for nonconvex-nonconcave minimax optimization, 2020.

[11] B. Halpern. Fixed points of nonexpanding maps. *Bull. Amer. Math. Soc*, 73:957–961, 1967.

[12] Y.-G. Hsieh, F. Iutzeler, J. Malick, and P. Mertikopoulos. On the convergence of single-call stochastic extra-gradient methods. In *Neural Info. Proc. Sys.*, 2019.

[13] A. Juditsky, A. Nemirovski, and C. Tauvel. Solving variational inequalities with stochastic mirror-prox algorithm. *Stochastic Systems*, 1(1):17–58, 2011.

[14] D. Kim. Accelerated proximal point method for maximally monotone operators. *Mathematical Programming*, pages 1–31, 2021.

[15] G. M. Korpelevich. An extragradient method for finding saddle points and other problems. *Ekonomika i Mateaticheskie Metody*, 12(4):747–56, 1976.

[16] F. Lieder. On the convergence rate of the Halpern-iteration. *Optimization Letters*, 2020.

[17] Q. Lin, M. Liu, H. Rafique, and T. Yang. Solving weakly-convex-weakly-concave saddle-point problems as weakly-monotone variational inequality. *arXiv preprint arXiv:1810.10207*, 5, 2018.

[18] M. Liu, Y. Mroueh, J. Ross, W. Zhang, X. Cui, P. Das, and T. Yang. Towards better understanding of adaptive gradient algorithms in generative adversarial nets. In *International Conference on Learning Representations*, 2020.

[19] Y. Malitsky. Golden ratio algorithms for variational inequalities. *Mathematical Programming*, 2020.

[20] Y. Malitsky and T. Pock. A first-order primal-dual algorithm with linesearch. *SIAM Journal on Optimization*, 28(1):411–432, 2018.

[21] P. Mertikopoulos, B. Lecouat, H. Zenati, C.-S. Foo, V. Chandrasekhar, and G. Piliouras. Optimistic mirror descent in saddle-point problems: going the extra (gradient) mile. In *Proc. Intl. Conf. on Learning Representations*, 2019.

[22] A. Mądry, A. Makelov, L. Schmdit, D. Tsipras, and A. Vladu. Towards deep learning models resistant to adversarial attacks. In *Proc. Intl. Conf. on Learning Representations*, 2018.

[23] A. Mokhtari, A. Ozdaglar, and S. Pattathil. A unified analysis of extra-gradient and optimistic gradient methods for saddle point problems: Proximal point approach. In *Proc. Intl. Conf. Artificial Intelligence and Stat. (AISTATS)*, 2020.

[24] R. D. C. Monteiro and B. F. Svaiter. On the complexity of the hybrid proximal extragradient method for the iterates and the ergodic mean. *SIAM J. Optim.*, 20(6):2755–87, 2010.

[25] R. D. C. Monteiro and B. F. Svaiter. Complexity of variants of Tseng's modified FB splitting and Korpelevich's methods for hemivariational inequalities with applications to saddle-point and convex optimization problems. *SIAM Journal on Optimization*, 21(4):1688–1720, 2011.

[26] M. C. Mukkamala, P. Ochs, T. Pock, and S. Sabach. Convex-concave backtracking for inertial bregman proximal gradient algorithms in nonconvex optimization. *SIAM Journal on Mathematics of Data Science*, 2(3):658–682, 2020.

[27] A. Nemirovski. Prox-method with rate of convergence $O(1/t)$ for variational inequalities with Lipschitz continuous monotone operators and smooth convex-concave saddle point problems. *SIAM J. Optim.*, 15(1):229–51, 2004.

[28] A. Nemirovski, A. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM J. Optim.*, 19(4):1574–609, 2009.

[29] A. Nemirovski and D. Yudin. *Problem complexity and method efficiency in optimization*. wiley, 1983.

[30] Y. Nesterov. A method for unconstrained convex minimization problem with the rate of convergence $O(1/k^2)$. *Dokl. Akad. Nauk. USSR*, 269(3):543–7, 1983.

[31] Y. Nesterov. Smooth minimization of non-smooth functions. *Mathematical Programming*, 103(1):127–52, May 2005.

[32] Y. Nesterov. Dual extrapolation and its applications to solving variational inequalities and related problems. *Mathematical Programming*, 109(2-3):319–44, March 2007.

[33] Y. Ouyang and Y. Xu. Lower complexity bounds of first-order methods for convex-concave bilinear saddle-point problems. *Mathematical Programming*, 2019.

[34] L. D. Popov. A modification of the Arrow-Hurwicz method for search of saddle points. *Mathematical notes of the Academy of Sciences of the USSR*, 28(5):845–8, November 1980.

[35] E. K. Ryu, K. Yuan, and W. Yin. ODE analysis of stochastic gradient methods with optimism and anchoring for minimax problems and GANs, 2019. arxiv 1905.10899.

[36] M. V. Solodov and B. F. Svaiter. A hybrid approximate extragradient–proximal point algorithm using the enlargement of a maximal monotone operator. *Set-Valued Analysis*, 7(4):323–345, 1999.

[37] C. Song, Z. Zhou, Y. Zhou, Y. Jiang, and Y. Ma. Optimistic dual extrapolation for coherent non-monotone variational inequalities. In *Neural Info. Proc. Sys.*, 2020.

[38] T. Yoon and E. K. Ryu. Accelerated algorithms for smooth convex-concave minimax problems with $\mathcal{O}(1/k^2)$ rate on squared gradient norm. In *Proc. Intl. Conf. Mach. Learn*, 2021.

[39] Z. Zhou, P. Mertikopoulos, N. Bambos, S. Boyd, and P. Glynn. Stochastic mirror descent in variationally coherent optimization problems. In *Neural Info. Proc. Sys.*, 2017.