Sequential Correlations Change In-Context Learning: Effective Context Length and Architectural Mismatch

Mary Letey   MARYLETEY@FAS.HARVARD.EDU
John A. Paulson School of Engineering and Applied Sciences, Harvard University;
Kempner Institute for the Study of Natural and Artificial Intelligence, Harvard University

Yue M. Lu   YUELU@SEAS.HARVARD.EDU
Cengiz Pehlevan[*]   CPEHLEVAN@SEAS.HARVARD.EDU
John A. Paulson School of Engineering and Applied Sciences, Harvard University

Jacob Zavatone-Veth[*]   JZAVATONEVETH@FAS.HARVARD.EDU
Society of Fellows and Center for Brain Science, Harvard University

Abstract

Modern sequence models have a striking capacity for in-context learning (ICL); they can perform new tasks based only on examples given in the prompt. Understanding how this ability emerges requires theory that captures important properties of natural data. Linear regression has served as a useful sandbox for ICL theory, but existing work has largely focused on prompts with independent examples. In this work, we extend this setting to sequentially correlated data, a basic feature of real sequences. We present a solvable model based on linear attention and test our predictions on realistic transformer architectures. We identify two distinct effects: First, when the query token is independent of the context, within-context correlations induce an effective context length: correlated prompts behave like shorter i.i.d. prompts. Second, when the query is also correlated with its context, test error is reduced, particularly for softmax attention when compared to linear attention. These results suggest that correlated prompts alter not only the effective sample size of in-context learning, but also which attention architectures are best matched to the task.

1. Introduction

In-context learning (ICL) is an important and useful capability of modern sequence models (Brown et al., 2020; Von Oswald et al., 2023; Wei et al., 2022a). In this setting, a model performs a task implicitly from few examples without parameter updates. Regression has been a useful controlled setting for studying how sequence models can achieve this ability. In-context regression requires a model to predict the label of a final query based on a prompt of input-output examples. The appeal of this setup is that it is simple enough to admit explicit analyses, yet rich enough to differentiate architectures and learning mechanisms (Akyürek et al., 2023; Von Oswald et al., 2023; Zhang et al., 2023; Lu et al., 2025; Zhang et al., 2024b; Oko et al., 2024; Vasudeva et al., 2025; Letey et al., 2026).

A standard simplifying assumption in the in-context regression literature is that the examples within a prompt are independent. This assumption is analytically convenient, but might not reflect real datasets. This has prompted the study of sequential correlations in other, non-regression in-context data settings, including Markov chains and dynamical

---

[*] CP and JZ-V jointly supervised this work.

Figure 1: ICL test loss against token correlation strength for three attention architectures. We highlight two findings. (1) ICL performance on correlated data with an independent query (purple) is well approximated by uncorrelated data at an effective context length (blue) given by eq (1). (2) When the query token is correlated with the preceding context (green), ICL error is reduced in all models as they utilise this correlation for inference. The dashed theory curves are derived for reduced-parameter linear attention; experimental details given in Appendix A.

systems (Edelman et al., 2024; Nichani et al., 2024; Cole et al., 2025; Park et al., 2025). For regression, the effects of sequential correlations have yet to be analysed. We know that sequential correlations matter even for classical ridge regression, where they change the asymptotic risk and invalidate estimators designed for i.i.d. data (Atanasov et al., 2025). Thus, it is natural to ask how they affect in-context regression, particularly as attention architectures were designed to operate on structured sequences (Vaswani et al., 2017).

In this work, we study a solvable model of in-context linear regression with sequentially correlated tokens. We begin from a reduced linear-attention model analysed in prior works (Zhang et al., 2024a; Wu et al., 2024; Lu et al., 2025; Letey et al., 2026; Bordelon et al., 2025), and perturb the usual i.i.d. setup by introducing a sequence correlation kernel for the context tokens, together with an optional correlation between the test query and the preceding context. This gives a controlled setting in which we can separate the effects of correlations *within* the context, and correlations *between* the query and the context.

Figure 1 summarises our findings. First, when correlations are confined to the context and the query remains independent, their effect is equivalent to an effective context-length reduction: correlated prompts behave like shorter i.i.d. prompts. Second, when the query token is correlated with its preceding context, ICL error decreases substantially, because the model can exploit this extra statistical structure to improve inference. Third, these settings reveal an architectural mismatch: the introduction of additional query-dependent statistics broadens the performance gap between linear and softmax attention architectures. An additional technical contribution is we identify a regime in which strong sequential correlations lead to the breakdown of the high-dimensional treatment originally formulated in Lu et al. (2025) When correlations persist on a length scale that grows with the context length, the effective number of independent examples may no longer scale proportionally with dimension, and the moments of the linear attention estimator need not concentrate.

Related Works

Here we summarise key empirical and theoretical results from previous studies of ICL.

ICL in practice. Early works showed that language models demonstrate an ability to learn new tasks, without weight updates, given a few examples provided "in context" (Radford et al., 2019; Brown et al., 2020; Chowdhery et al., 2022; Touvron et al., 2023; Achiam et al., 2023; Bubeck et al., 2023). This phenomenon seemed to emerge with scale (Wei et al., 2022a, 2023) and to be robust to task complexity (Wei et al., 2022b; DeepSeek-AI et al., 2025). These observations led to much interest in understanding ICL (Dong et al., 2024), particularly regarding the question of how unknown tasks are solved at test time. Many empirical works attempted to answer this question, focusing on mechanisms (Von Oswald et al., 2023; Li et al., 2024a; Akyürek et al., 2023; Akyürek et al., 2024; Kirsch et al., 2024; Olsson et al., 2022; Tong and Pehlevan, 2025; Li et al., 2024b) and data structures (Bietti et al., 2023; Chan et al., 2022; Raventós et al., 2023; Garg et al., 2022; Liu et al., 2025; Singh et al., 2023; Goddard et al., 2025; Liu et al., 2024; Vasudeva et al., 2025; Todd et al., 2026) that enable ICL.

ICL in theory. Recent theoretical works have sought to answer similar questions: what algorithms could ICL be implementing, and what is the effect of data? Simple models, such as shallow networks or linear attention, have proven particularly useful testbeds for understanding what happens at inference time (Chandra et al., 2024; Fu et al., 2023; Shen et al., 2025; Gozeten et al., 2025; Mahankali et al., 2023; Barnfield et al., 2026; Ahn et al., 2023; Zhang et al., 2025; Demir and Dogan, 2025). Many works have been able to show through constructive approaches that transformers or simple architectures are able to implement general algorithms, such as gradient descent, to solve tasks in-context (Bai et al., 2023; Reddy, 2024; Elmoznino et al., 2025; Lee et al., 2025; Zhang et al., 2023). Many theoretical works have explicitly considered sequential structures, such as dynamical systems (Li et al., 2023; Cole et al., 2025) and Markov chains (Edelman et al., 2024; Park et al., 2025; Nichani et al., 2024), in addition to more complex synthetic tasks (Tao et al., 2025; He et al., 2024; Ye et al., 2026; Cole et al., 2026).

ICL in regression. Moving away from reasoning and language-like tasks, regression has been a very useful sandbox for ICL research. By focusing on simpler function classes, one can derive sharper results, including sample sufficiency (Oko et al., 2024), gradient-descent-like behaviour (Wu et al., 2024; Zhang et al., 2024a,b; Bordelon et al., 2025; Fu et al., 2024), and finite-sample error formulas. (Lu et al., 2025; Letey et al., 2026). We build on this to isolate the effect of sequential correlations on ICL for linear regression, and to use this setting as a probe of how attention architectures respond to such correlations.

2. Theory and Data Setup

In-context linear regression. We study a correlated-token version of the solvable linear-regression ICL model induced by a reduced linear-attention block, studied by Lu et al. (2025); Zhang et al. (2023). A context consists of a sequence $\{\boldsymbol{x}_1, y_1, \ldots, \boldsymbol{x}_\ell, y_\ell, \boldsymbol{x}_{\text{test}}\}$ with

$$y_a = \boldsymbol{x}_a^\top \boldsymbol{w} + \varepsilon_a, \qquad y_{\text{test}} = \boldsymbol{x}_{\text{test}}^\top \boldsymbol{w} + \varepsilon_{\text{test}},$$

for task vector $\boldsymbol{w} \in \mathbb{R}^d$, and independent Gaussian noise $\boldsymbol{\varepsilon} \sim \mathcal{N}(0, \rho I_\ell)$, $\varepsilon_{\text{test}} \sim \mathcal{N}(0, \rho)$. The goal is for the model's output on this sequence to be close to y_{test}, thus performing the

correct regression inference given examples in the context. Throughout, we write $X \in \mathbb{R}^{\ell \times d}$ for the matrix whose ath row is $\boldsymbol{x}_a^\top$.

Sequential correlations. We model sequential structure within the context as linear dependence between the tokens, following previous work in the ridge regression setting (Atanasov et al., 2025; Moniri and Hassani, 2025). Under this model, the mean-zero Gaussian tokens $\boldsymbol{x}_1, \ldots, \boldsymbol{x}_\ell$ have second moments given by

$$\mathbb{E}[\boldsymbol{x}_a \boldsymbol{x}_b^\top] = K_{ab} I_d / d, \qquad \mathbb{E}[\boldsymbol{x}_{\text{test}} \boldsymbol{x}_a^\top] = \boldsymbol{k}_{\text{test}\,a} I_d / d$$

i.e., $K \in \mathbb{R}^{\ell \times \ell}$ controls correlations across context positions, and the feature-feature covariance is given by I_d / d. The case of independently sampled tokens is described by $K = I_\ell$. The query correlations are controlled by $\boldsymbol{k}_{\text{test}} \in \mathbb{R}^\ell$ which specifies how strongly the query is coupled to each preceding context token; for an independent query, $\boldsymbol{k}_{\text{test}} = \boldsymbol{0}$.

Correlation summaries and correlation length. We normalise such that $\text{tr}(K) = \ell$. The theory depends on K and $\boldsymbol{k}_{\text{test}}$ through the summary statistics

$$k_0 := \boldsymbol{k}_{\text{test}}^\top \boldsymbol{k}_{\text{test}}, \qquad k_1 := \boldsymbol{k}_{\text{test}}^\top K \boldsymbol{k}_{\text{test}}, \qquad k_2 := \text{tr}(K^2) / \ell.$$

For general K and $\boldsymbol{k}_{\text{test}}$, we can think of k_0, k_1, k_2 as measures of token correlation strength. The quantity k_2 is a bulk measure of context correlations: under our chosen normalisation $\text{tr}(K) = \ell$, it is minimised by the uncorrelated case $K = I_\ell$, where $k_2 = 1$, and increases as the spectrum of K becomes more anisotropic or concentrated. The query-dependent quantity k_0 measures the overall strength of the direct query-context coupling, while the mixed quantity k_1 measures how strongly that query-correlation profile is amplified by the correlated structure of the context itself.

These quantities become more intuitive if we take a specific kernel as an example: the exponential kernel $K_{ab} = \exp(-|a - b| / \xi)$ and $\boldsymbol{k}_{\text{test}\,a} = \exp(-(\ell + 1 - a) / \xi)$. Because K and $\boldsymbol{k}_{\text{test}}$ only depend on *distances* between tokens, the natural parameter of correlation "strength" is correlation *length* ξ. This case provides an intuitive sandbox, as the ICL summary statistics above are directly related to the correlation length by $k_2, k_0 \approx \xi$, $k_1 \approx \xi^2$. We thus choose to use the exponential kernel for the experiments in all figures, and save a more detailed analysis with non-exponential kernels to future work.

Reduced linear-attention predictor. Following prior work on linear attention, the next-token prediction for y_{test} made by linear attention can be well-approximated by

$$\hat{y}_{\text{test}} := \text{tr}\left(\Gamma H^\top\right), \quad \Gamma \in \mathbb{R}^{d \times (d+1)}, \quad H := \boldsymbol{x}_{\text{test}} \left[\frac{d}{\ell} (X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top X \quad \frac{1}{\ell} (X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top (X\boldsymbol{w} + \boldsymbol{\varepsilon}) \right].$$

The parameter matrix Γ here is defined in terms of components of the value, key, and query matrices from full-parameter linear attention. This is the predictor that we will study in the theory, and is the "reduced model" referred to on the leftmost panel of Figure 1.

Given a training batch of n sequences, corresponding to n such data matrices H for the reduced model, the optimal parameters and corresponding test loss can be written as

$$\Gamma^* = \left(\frac{n}{d} \lambda I_{d(d+1)} + \sum_{\mu=1}^n H^\mu \otimes H^\mu \right)^{-1} \sum_{\mu=1}^n y_{\text{test}}^\mu H^\mu, \quad \mathcal{E}_{\text{ICL}}(\Gamma^*) = \mathbb{E}[(y_{\text{test}}^{\text{new}} - \langle H^{\text{new}}, \Gamma^* \rangle)^2].$$

High-dimensional scaling. We study the matrix Γ^*, and functions of it, in a high-dimensional limit using random-matrix techniques as in Lu et al. (2025). We work in the standard proportional regime, with $d \to \infty$ with $\ell \propto d$. We assume the task signal remains identifiable, e.g. $\boldsymbol{w}^\top \boldsymbol{w} = \Theta(d)$.

We analytically study the case of *weak correlations* with respect to context length ℓ: for the computation of theory curves we take $k_2, k_0, k_1 = \Theta_\ell(1)$. We do not study stronger correlations in this work, corresponding to cases such as $k_2 = o(\ell)$ or $k_2 \propto \ell$, and do not attempt to characterise all of the ways in which the theory's predictions break down. In stronger-correlation regimes, concentration of the parameter matrix Γ^* and its corresponding ICL error can fail (see Figure 2), and a full characterisation is left for future work, with some further discussion in Section 3 and Appendix D.

Figure 2: Simulations of $\mathcal{E}_{\text{ICL}}(\Gamma^*)$ for several scaling regimes, compared with theory (Prop. 2).

3. Effective in-context sample size

Working in the proportional limit described above, we can analyse the random parameter matrix Γ^*. The full result is given in Appendix E. We present its first implication.

Effective context length formula

Consider tokens $\boldsymbol{x}_1, \ldots, \boldsymbol{x}_\ell$ that are weakly correlated, *i.e.* $k_2 = \Theta(1)$, and an independent query $\boldsymbol{x}_{\text{test}}$, *i.e.* $\boldsymbol{k}_{\text{test}} = \boldsymbol{0}$. Then the correlated-ICL performance at context length ℓ matches the uncorrelated-ICL performance at the lower effective context length

$$\ell_{\text{eff}} := \frac{1 + \rho}{k_2 + \rho}\ell, \tag{1}$$

where by Jensen's inequality $k_2 = \text{tr}(K^2)/\ell \geq (\text{tr}(K)/\ell)^2 = 1$, so $\ell_{\text{eff}} \leq \ell$.

Eq (1) makes precise how sequential correlations reduce the usable information content of the prompt. Since k_2 increases as the context covariance becomes more anisotropic, stronger or more persistent correlations decrease the effective context length ℓ_{eff}. For stationary kernels, where k_2 grows with the correlation scale, this recovers the intuitive statement that longer-ranged correlations make fewer context examples effectively independent. The simplicity of the resulting formula is itself notable: the reduction depends linearly on the single scalar k_2, rather than on the full spectrum of K. This contrasts with classical correlated linear regression, where test error typically depends more delicately on the spectrum, and sometimes the eigenvectors, of the design covariance.

Eq (1) also hints why the strongly-correlated regime is analytically difficult. Our theory here assumes that the correlation summary k_2 remains $\Theta(1)$ as $\ell, d \to \infty$ with $\ell \propto d$. But once correlations persist across more and more of the context, we instead have either $k_2 = o(\ell)$ or $k_2 \propto \ell$. In this regime, ℓ_{eff} no longer scales proportionally with ℓ, but remains only $o(\ell)$ or $\Theta(1)$ respectively. Thus, even as the nominal context length increases, we are no longer in the standard linear-regression regime where samples are proportional to

dimension. We discuss this further in Appendix D, as these regimes are where the standard high-dimensional concentration underlying our asymptotic analysis may break down.

4. Gain from query correlations

We now turn to the second effect in Figure 1: correlations between the test query and its preceding context can provide additional predictive signal and thus reduce the ICL error. To isolate this contribution, define the *query-correlation gain* as

$$\Delta_{\text{query}} := \mathcal{L}_{\text{ICL}}(\ell, k_2 \,; \boldsymbol{k}_{\text{test}} = 0) - \mathcal{L}_{\text{ICL}}(\ell, k_2, k_0, k_1),$$

namely, the reduction in ICL error obtained by introducing query-context correlations while holding the bulk context kernel K fixed. Here $\mathcal{L}_{\text{ICL}}$ denotes the ICL error of a model (not necessarily linear attention) on this correlated linear regression task.

Figure 3: Query effects depend on architecture.

In Figure 3, we find that Δ_{query} is nonnegative (up to finite-sample fluctuations) and vanishes in the uncorrelated case $K = I_\ell$. Thus, unlike bulk context correlations, which reduce effective sample size and hurt ICL performance, query correlations provide additional predictive signal that can in principle be exploited for inference.

In our theory, this effect is controlled not only by the bulk statistic k_2, but also by the query-dependent summaries k_0 and k_1, which increase with stronger correlations. The resulting closed-form expression for Δ_{query} is substantially more complicated than the effective-sample law in eq. (1); we therefore omit it from the main text. Empirically, Δ_{query} is relatively modest for linear-attention architectures but much larger for architectures that use softmax attention. This suggests that softmax attention is better matched to sequential inference, as it can form query-dependent weights over the most informative tokens rather than compressing correlations into the second-order summaries available to linear attention.

5. Conclusion

This work is a first step toward a theory of ICL for structured, sequential data. By separating bulk context correlations from query-context correlations, we show that prompt structure can affect ICL in two qualitatively different ways: bulk context correlations reduce effective sample size, while query-context correlations provide useful signal for inference, highlighting an architectural mismatch between linear and softmax attention. The latter effect is especially important: softmax attention appears to have a distinctive ability to exploit sequential information, which is not captured by the linear-attention theory. Developing theory that explains this advantage, and extending it beyond linear regression to richer sequential and dynamical tasks, will be necessary for understanding ICL in more realistic data regimes.

Acknowledgements

M.L. is supported by a Graduate Fellowship from the Kempner Institute for the Study of Natural and Artificial Intelligence. Y.M.L. gratefully acknowledges support from a Harvard College Professorship, the Harvard FAS Dean's Fund for Promising Scholarship, and DARPA grant DIAL-FP-038. C.P. was supported by an NSF CAREER Award (IIS-2239780), DARPA grants DIAL-FP-038 and AIQ-HR00112520041, the Simons Collaboration on the Physics of Learning and Neural Computation, and the William F. Milton Fund from Harvard University. J.Z.-V. was supported by a Junior Fellowship from the Harvard Society of Fellows. This work has been made possible in part by a gift from the Chan Zuckerberg Initiative Foundation to establish the Kempner Institute for the Study of Natural and Artificial Intelligence.

References

Josh Achiam, Steven Adler, Sandhini Agarwal, Lama Ahmad, Ilge Akkaya, Florencia Leoni Aleman, Diogo Almeida, Janko Altenschmidt, Sam Altman, Shyamal Anadkat, et al. Gpt-4 technical report. *arXiv preprint arXiv:2303.08774*, 2023.

Kwangjun Ahn, Xiang Cheng, Hadi Daneshmand, and Suvrit Sra. Transformers learn to implement preconditioned gradient descent for in-context learning, 2023.

Ekin Akyürek, Dale Schuurmans, Jacob Andreas, Tengyu Ma, and Denny Zhou. What learning algorithm is in-context learning? investigations with linear models. In *The Eleventh International Conference on Learning Representations*, 2023. URL https://openreview.net/forum?id=0g0X4H8yN4I.

Ekin Akyürek, Bailin Wang, Yoon Kim, and Jacob Andreas. In-context language learning: Architectures and algorithms, 2024. URL https://arxiv.org/abs/2401.12973.

Alexander Atanasov, Jacob A Zavatone-Veth, and Cengiz Pehlevan. Risk and cross validation in ridge regression with correlated samples. In *Forty-second International Conference on Machine Learning*, 2025. URL https://openreview.net/forum?id=GMwKpJ9TiR.

Yu Bai, Fan Chen, Huan Wang, Caiming Xiong, and Song Mei. Transformers as statisticians: Provable in-context learning with in-context algorithm selection. In A. Oh, T. Naumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 57125–57211. Curran Associates, Inc., 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/file/b2e63e36c57e153b9015fece2352a9f9-Paper-Conference.pdf.

Nicholas Barnfield, Subhabrata Sen, and Pragya Sur. Multi-layer cross-attention is provably optimal for multi-modal in-context learning, 2026. URL https://arxiv.org/abs/2602.04872.

Alberto Bietti, Vivien Cabannes, Diane Bouchacourt, Herve Jegou, and Leon Bottou. Birth of a transformer: A memory viewpoint. In A. Oh, T. Naumann,

A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 1560–1588. Curran Associates, Inc., 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/file/0561738a239a995c8cd2ef0e50cfa4fd-Paper-Conference.pdf.

Blake Bordelon, Mary I. Letey, and Cengiz Pehlevan. Theory of scaling laws for in-context regression: Depth, width, context and time, 2025. URL https://arxiv.org/abs/2510.01098.

Tom Brown, Benjamin Mann, Nick Ryder, Melanie Subbiah, Jared D Kaplan, Prafulla Dhariwal, Arvind Neelakantan, Pranav Shyam, Girish Sastry, Amanda Askell, et al. Language models are few-shot learners. *Advances in neural information processing systems*, 33:1877–1901, 2020.

Sébastien Bubeck, Varun Chandrasekaran, Ronen Eldan, Johannes Gehrke, Eric Horvitz, Ece Kamar, Peter Lee, Yin Tat Lee, Yuanzhi Li, Scott Lundberg, Harsha Nori, Hamid Palangi, Marco Tulio Ribeiro, and Yi Zhang. Sparks of artificial general intelligence: Early experiments with gpt-4, 2023. URL https://arxiv.org/abs/2303.12712.

Stephanie C. Y. Chan, Adam Santoro, Andrew K. Lampinen, Jane X. Wang, Aaditya Singh, Pierre H. Richemond, Jay McClelland, and Felix Hill. Data distributional properties drive emergent in-context learning in transformers, 2022.

Pritam Chandra, Tanmay Kumar Sinha, Kabir Ahuja, Ankit Garg, and Navin Goyal. Towards analyzing self-attention via linear neural network, 2024. URL https://openreview.net/forum?id=4fVuBf5HE9.

Aakanksha Chowdhery, Sharan Narang, Jacob Devlin, Maarten Bosma, Gaurav Mishra, Adam Roberts, Paul Barham, Hyung Won Chung, Charles Sutton, Sebastian Gehrmann, Parker Schuh, Kensen Shi, Sasha Tsvyashchenko, Joshua Maynez, Abhishek Rao, Parker Barnes, Yi Tay, Noam Shazeer, Vinodkumar Prabhakaran, Emily Reif, Nan Du, Ben Hutchinson, Reiner Pope, James Bradbury, Jacob Austin, Michael Isard, Guy Gur-Ari, Pengcheng Yin, Toju Duke, Anselm Levskaya, Sanjay Ghemawat, Sunipa Dev, Henryk Michalewski, Xavier Garcia, Vedant Misra, Kevin Robinson, Liam Fedus, Denny Zhou, Daphne Ippolito, David Luan, Hyeontaek Lim, Barret Zoph, Alexander Spiridonov, Ryan Sepassi, David Dohan, Shivani Agrawal, Mark Omernick, Andrew M. Dai, Thanumalayan Sankaranarayana Pillai, Marie Pellat, Aitor Lewkowycz, Erica Moreira, Rewon Child, Oleksandr Polozov, Katherine Lee, Zongwei Zhou, Xuezhi Wang, Brennan Saeta, Mark Diaz, Orhan Firat, Michele Catasta, Jason Wei, Kathy Meier-Hellstern, Douglas Eck, Jeff Dean, Slav Petrov, and Noah Fiedel. Palm: Scaling language modeling with pathways, 2022. URL https://arxiv.org/abs/2204.02311.

Frank Cole, Yulong Lu, Tianhao Zhang, and Yuxuan Zhao. In-context learning of linear dynamical systems with transformers: Error bounds and depth-separation, 2025. URL https://arxiv.org/abs/2502.08136.

Frank Cole, Dixi Wang, Yineng Chen, Yulong Lu, and Rongjie Lai. In-context operator learning on the space of probability measures, 2026. URL https://arxiv.org/abs/2601.09979.

DeepSeek-AI, Daya Guo, Dejian Yang, Haowei Zhang, Junxiao Song, Ruoyu Zhang, Runxin Xu, Qihao Zhu, Shirong Ma, Peiyi Wang, Xiao Bi, Xiaokang Zhang, Xingkai Yu, Yu Wu, Z. F. Wu, Zhibin Gou, Zhihong Shao, Zhuoshu Li, Ziyi Gao, Aixin Liu, Bing Xue, Bingxuan Wang, Bochao Wu, Bei Feng, Chengda Lu, Chenggang Zhao, Chengqi Deng, Chenyu Zhang, Chong Ruan, Damai Dai, Deli Chen, Dongjie Ji, Erhang Li, Fangyun Lin, Fucong Dai, Fuli Luo, Guangbo Hao, Guanting Chen, Guowei Li, H. Zhang, Han Bao, Hanwei Xu, Haocheng Wang, Honghui Ding, Huajian Xin, Huazuo Gao, Hui Qu, Hui Li, Jianzhong Guo, Jiashi Li, Jiawei Wang, Jingchang Chen, Jingyang Yuan, Junjie Qiu, Junlong Li, J. L. Cai, Jiaqi Ni, Jian Liang, Jin Chen, Kai Dong, Kai Hu, Kaige Gao, Kang Guan, Kexin Huang, Kuai Yu, Lean Wang, Lecong Zhang, Liang Zhao, Litong Wang, Liyue Zhang, Lei Xu, Leyi Xia, Mingchuan Zhang, Minghua Zhang, Minghui Tang, Meng Li, Miaojun Wang, Mingming Li, Ning Tian, Panpan Huang, Peng Zhang, Qiancheng Wang, Qinyu Chen, Qiushi Du, Ruiqi Ge, Ruisong Zhang, Ruizhe Pan, Runji Wang, R. J. Chen, R. L. Jin, Ruyi Chen, Shanghao Lu, Shangyan Zhou, Shanhuang Chen, Shengfeng Ye, Shiyu Wang, Shuiping Yu, Shunfeng Zhou, Shuting Pan, S. S. Li, Shuang Zhou, Shaoqing Wu, Shengfeng Ye, Tao Yun, Tian Pei, Tianyu Sun, T. Wang, Wangding Zeng, Wanjia Zhao, Wen Liu, Wenfeng Liang, Wenjun Gao, Wenqin Yu, Wentao Zhang, W. L. Xiao, Wei An, Xiaodong Liu, Xiaohan Wang, Xiaokang Chen, Xiaotao Nie, Xin Cheng, Xin Liu, Xin Xie, Xingchao Liu, Xinyu Yang, Xinyuan Li, Xuecheng Su, Xuheng Lin, X. Q. Li, Xiangyue Jin, Xiaojin Shen, Xiaosha Chen, Xiaowen Sun, Xiaoxiang Wang, Xinnan Song, Xinyi Zhou, Xianzu Wang, Xinxia Shan, Y. K. Li, Y. Q. Wang, Y. X. Wei, Yang Zhang, Yanhong Xu, Yao Li, Yao Zhao, Yaofeng Sun, Yaohui Wang, Yi Yu, Yichao Zhang, Yifan Shi, Yiliang Xiong, Ying He, Yishi Piao, Yisong Wang, Yixuan Tan, Yiyang Ma, Yiyuan Liu, Yongqiang Guo, Yuan Ou, Yuduan Wang, Yue Gong, Yuheng Zou, Yujia He, Yunfan Xiong, Yuxiang Luo, Yuxiang You, Yuxuan Liu, Yuyang Zhou, Y. X. Zhu, Yanhong Xu, Yanping Huang, Yaohui Li, Yi Zheng, Yuchen Zhu, Yunxian Ma, Ying Tang, Yukun Zha, Yuting Yan, Z. Z. Ren, Zehui Ren, Zhangli Sha, Zhe Fu, Zhean Xu, Zhenda Xie, Zhengyan Zhang, Zhewen Hao, Zhicheng Ma, Zhigang Yan, Zhiyu Wu, Zihui Gu, Zijia Zhu, Zijun Liu, Zilin Li, Ziwei Xie, Ziyang Song, Zizheng Pan, Zhen Huang, Zhipeng Xu, Zhongyu Zhang, and Zhen Zhang. Deepseek-r1: Incentivizing reasoning capability in llms via reinforcement learning. *arXiv*, 2025. URL https://arxiv.org/abs/2501.12948.

Samet Demir and Zafer Dogan. How data mixing shapes in-context learning: Asymptotic equivalence for transformers with mlps. In D. Belgrave, C. Zhang, H. Lin, R. Pascanu, P. Koniusz, M. Ghassemi, and N. Chen, editors, *Advances in Neural Information Processing Systems*, volume 38, pages 119346–119376. Curran Associates, Inc., 2025. URL https://proceedings.neurips.cc/paper_files/paper/2025/file/ad184b4e096284a55cf66e1d4eeb29df-Paper-Conference.pdf.

Qingxiu Dong, Lei Li, Damai Dai, Ce Zheng, Jingyuan Ma, Rui Li, Heming Xia, Jingjing Xu, Zhiyong Wu, Tianyu Liu, Baobao Chang, Xu Sun, Lei Li, and Zhifang Sui. A survey on in-context learning, 2024. URL https://arxiv.org/abs/2301.00234.

Benjamin L. Edelman, Ezra Edelman, Surbhi Goel, Eran Malach, and Nikolaos Tsilivis. The evolution of statistical induction heads: In-context learning markov chains, 2024. URL https://arxiv.org/abs/2402.11004.

Eric Elmoznino, Tom Marty, Tejas Kasetty, Leo Gagnon, Sarthak Mittal, Mahan Fathi, Dhanya Sridhar, and Guillaume Lajoie. In-context learning and occam's razor, 2025. URL https://arxiv.org/abs/2410.14086.

Deqing Fu, Tian-Qi Chen, Robin Jia, and Vatsal Sharan. Transformers learn higher-order optimization methods for in-context learning: A study with linear models, 2023.

Deqing Fu, Tian-Qi Chen, Robin Jia, and Vatsal Sharan. Transformers learn to achieve second-order convergence rates for in-context linear regression, 2024. URL https://arxiv.org/abs/2310.17086.

Shivam Garg, Dimitris Tsipras, Percy Liang, and Gregory Valiant. What can transformers learn in-context? a case study of simple function classes. In Alice H. Oh, Alekh Agarwal, Danielle Belgrave, and Kyunghyun Cho, editors, Advances in Neural Information Processing Systems, 2022. URL https://openreview.net/forum?id=flNZJ2eOet.

Chase Goddard, Lindsay M. Smith, Vudtiwat Ngampruetikorn, and David J. Schwab. When can in-context learning generalize out of task distribution?, 2025. URL https://arxiv.org/abs/2506.05574.

Halil Alperen Gozeten, M. Emrullah Ildiz, Xuechen Zhang, Mahdi Soltanolkotabi, Marco Mondelli, and Samet Oymak. Test-time training provably improves transformers as in-context learners, 2025. URL https://arxiv.org/abs/2503.11842.

Tianyu He, Darshil Doshi, Aritra Das, and Andrey Gromov. Learning to grok: Emergence of in-context learning and skill composition in modular arithmetic tasks. In A. Globerson, L. Mackey, D. Belgrave, A. Fan, U. Paquet, J. Tomczak, and C. Zhang, editors, Advances in Neural Information Processing Systems, volume 37, pages 13244–13273. Curran Associates, Inc., 2024. doi: 10.52202/079017-0422. URL https://proceedings.neurips.cc/paper_files/paper/2024/file/17d60fef592086d1a5cb136f1946df59-Paper-Conference.pdf.

Louis Kirsch, James Harrison, Jascha Sohl-Dickstein, and Luke Metz. General-purpose in-context learning by meta-learning transformers, 2024.

Jin Hwa Lee, Andrew Kyle Lampinen, Aaditya K Singh, and Andrew M Saxe. Distinct computations emerge from compositional curricula in in-context learning. In Workshop on Spurious Correlation and Shortcut Learning: Foundations and Solutions, 2025. URL https://openreview.net/forum?id=oo5TNikeJl.

Mary Letey, Jacob A Zavatone-Veth, Yue M. Lu, and Cengiz Pehlevan. Pretrain-test task alignment governs generalization in in-context learning. In The Fourteenth International Conference on Learning Representations, 2026. URL https://openreview.net/forum?id=KZLeg0MQ2r.

Jiaoda Li, Yifan Hou, Mrinmaya Sachan, and Ryan Cotterell. What do language models learn in context? the structured task hypothesis, 2024a. URL https://arxiv.org/abs/2406.04216.

Yingcong Li, M. Emrullah Ildiz, Dimitris Papailiopoulos, and Samet Oymak. Transformers as algorithms: Generalization and stability in in-context learning, 2023.

Yingcong Li, Ankit Singh Rawat, and Samet Oymak. Fine-grained analysis of in-context linear estimation: Data, architecture, and beyond. In A. Globerson, L. Mackey, D. Belgrave, A. Fan, U. Paquet, J. Tomczak, and C. Zhang, editors, *Advances in Neural Information Processing Systems*, volume 37, pages 138324–138364. Curran Associates, Inc., 2024b. URL https://proceedings.neurips.cc/paper_files/paper/2024/file/f9dc462382fef56d58279e75de2438f3-Paper-Conference.pdf.

Binwen Liu, Peiyu Xu, Quan Yuan, and Yihong Chen. Probing in-context learning: Impact of task complexity and model architecture on generalization and efficiency, 2025. URL https://arxiv.org/abs/2505.06475.

Toni J.b. Liu, Nicolas Boulle, Raphaël Sarfati, and Christopher Earls. Llms learn governing principles of dynamical systems, revealing an in-context neural scaling law. In *Proceedings of the 2024 Conference on Empirical Methods in Natural Language Processing*, page 15097–15117. Association for Computational Linguistics, 2024. doi: 10.18653/v1/2024.emnlp-main.842. URL http://dx.doi.org/10.18653/v1/2024.emnlp-main.842.

Yue M. Lu, Mary Letey, Jacob A. Zavatone-Veth, Anindita Maiti, and Cengiz Pehlevan. Asymptotic theory of in-context learning by linear attention. *Proceedings of the National Academy of Sciences*, 122(28):e2502599122, 2025. doi: 10.1073/pnas.2502599122. URL https://www.pnas.org/doi/abs/10.1073/pnas.2502599122.

Arvind Mahankali, Tatsunori B. Hashimoto, and Tengyu Ma. One step of gradient descent is provably the optimal in-context learner with one layer of linear self-attention, 2023. URL https://arxiv.org/abs/2307.03576.

Behrad Moniri and Hamed Hassani. Asymptotics of linear regression with linearly dependent data. In Necmiye Ozay, Laura Balzano, Dimitra Panagou, and Alessandro Abate, editors, *Proceedings of the 7th Annual Learning for Dynamics & Control Conference*, volume 283 of *Proceedings of Machine Learning Research*, pages 72–85. PMLR, 04–06 Jun 2025. URL https://proceedings.mlr.press/v283/moniri25a.html.

Eshaan Nichani, Alex Damian, and Jason D. Lee. How transformers learn causal structure with gradient descent. In Ruslan Salakhutdinov, Zico Kolter, Katherine Heller, Adrian Weller, Nuria Oliver, Jonathan Scarlett, and Felix Berkenkamp, editors, *Proceedings of the 41st International Conference on Machine Learning*, volume 235 of *Proceedings of Machine Learning Research*, pages 38018–38070. PMLR, 21–27 Jul 2024. URL https://proceedings.mlr.press/v235/nichani24a.html.

Kazusato Oko, Yujin Song, Taiji Suzuki, and Denny Wu. Pretrained transformer efficiently learns low-dimensional target functions in-context, 2024. URL https://arxiv.org/abs/2411.02544.

Catherine Olsson, Nelson Elhage, Neel Nanda, Nicholas Joseph, Nova DasSarma, Tom Henighan, Ben Mann, Amanda Askell, Yuntao Bai, Anna Chen, Tom Conerly, Dawn

Drain, Deep Ganguli, Zac Hatfield-Dodds, Danny Hernandez, Scott Johnston, Andy Jones, Jackson Kernion, Liane Lovitt, Kamal Ndousse, Dario Amodei, Tom Brown, Jack Clark, Jared Kaplan, Sam McCandlish, and Chris Olah. In-context learning and induction heads. *Transformer Circuits Thread*, 2022. URL https://transformer-circuits.pub/2022/in-context-learning-and-induction-heads/index.html.

Core Francisco Park, Ekdeep Singh Lubana, Itamar Pres, and Hidenori Tanaka. Competition dynamics shape algorithmic phases of in-context learning, 2025. URL https://arxiv.org/abs/2412.01003.

Alec Radford, Jeffrey Wu, Rewon Child, David Luan, Dario Amodei, Ilya Sutskever, et al. Language models are unsupervised multitask learners. *OpenAI blog*, 1(8):9, 2019.

Allan Raventós, Mansheej Paul, Feng Chen, and Surya Ganguli. Pretraining task diversity and the emergence of non-bayesian in-context learning for regression. In A. Oh, T. Naumann, A. Globerson, K. Saenko, M. Hardt, and S. Levine, editors, *Advances in Neural Information Processing Systems*, volume 36, pages 14228–14246. Curran Associates, Inc., 2023. URL https://proceedings.neurips.cc/paper_files/paper/2023/file/2e10b2c2e1aa4f8083c37dfe269873f8-Paper-Conference.pdf.

Gautam Reddy. The mechanistic basis of data dependence and abrupt learning in an in-context classification task. In *The Twelfth International Conference on Learning Representations*, 2024. URL https://openreview.net/forum?id=aN4Jf6Cx69.

Zhaiming Shen, Alexander Hsu, Rongjie Lai, and Wenjing Liao. Understanding in-context learning on structured manifolds: Bridging attention to kernel methods, 2025. URL https://arxiv.org/abs/2506.10959.

Aaditya K. Singh, Stephanie C. Y. Chan, Ted Moskovitz, Erin Grant, Andrew M. Saxe, and Felix Hill. The transient nature of emergent in-context learning in transformers, 2023.

Tao Tao, Darshil Doshi, Dayal Singh Kalra, Tianyu He, and Maissam Barkeshli. (how) can transformers predict pseudo-random numbers?, 2025. URL https://arxiv.org/abs/2502.10390.

Eric Todd, Jannik Brinkmann, Rohit Gandikota, and David Bau. In-context algebra, 2026. URL https://arxiv.org/abs/2512.16902.

William L. Tong and Cengiz Pehlevan. MLPs learn in-context on regression and classification tasks. In *The Thirteenth International Conference on Learning Representations*, 2025. URL https://openreview.net/forum?id=MbX0t1rUlp.

Hugo Touvron, Thibaut Lavril, Gautier Izacard, Xavier Martinet, Marie-Anne Lachaux, Timothée Lacroix, Baptiste Rozière, Naman Goyal, Eric Hambro, Faisal Azhar, et al. Llama: Open and efficient foundation language models. *arXiv preprint arXiv:2302.13971*, 2023.

Bhavya Vasudeva, Deqing Fu, Tianyi Zhou, Elliott Kau, Youqi Huang, and Vatsal Sharan. Transformers learn low sensitivity functions: Investigations and implications. In

International Conference on Learning Representations, volume 2025, pages 3565–3594, 2025.

Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.

Johannes Von Oswald, Eyvind Niklasson, Ettore Randazzo, João Sacramento, Alexander Mordvintsev, Andrey Zhmoginov, and Max Vladymyrov. Transformers learn incontext by gradient descent. In Andreas Krause, Emma Brunskill, Kyunghyun Cho, Barbara Engelhardt, Sivan Sabato, and Jonathan Scarlett, editors, *Proceedings of the 40th International Conference on Machine Learning*, volume 202 of *Proceedings of Machine Learning Research*, pages 35151–35174. PMLR, 23–29 Jul 2023. URL https://proceedings.mlr.press/v202/von-oswald23a.html.

Jason Wei, Yi Tay, Rishi Bommasani, Colin Raffel, Barret Zoph, Sebastian Borgeaud, Dani Yogatama, Maarten Bosma, Denny Zhou, Donald Metzler, Ed H. Chi, Tatsunori Hashimoto, Oriol Vinyals, Percy Liang, Jeff Dean, and William Fedus. Emergent abilities of large language models. *Transactions on Machine Learning Research*, 2022a. ISSN 2835-8856. URL https://openreview.net/forum?id=yzkSU5zdwD.

Jason Wei, Xuezhi Wang, Dale Schuurmans, Maarten Bosma, Ed H. Chi, Quoc Le, and Denny Zhou. Chain of thought prompting elicits reasoning in large language models. *CoRR*, abs/2201.11903, 2022b. URL https://arxiv.org/abs/2201.11903.

Jerry Wei, Jason Wei, Yi Tay, Dustin Tran, Albert Webson, Yifeng Lu, Xinyun Chen, Hanxiao Liu, Da Huang, Denny Zhou, and Tengyu Ma. Larger language models do in-context learning differently, 2023. URL https://arxiv.org/abs/2303.03846.

Jingfeng Wu, Difan Zou, Zixiang Chen, Vladimir Braverman, Quanquan Gu, and Peter Bartlett. How many pretraining tasks are needed for in-context learning of linear regression? In *The Twelfth International Conference on Learning Representations*, 2024. URL https://openreview.net/forum?id=vSh5ePa0ph.

Qilin Ye, Deqing Fu, Robin Jia, and Vatsal Sharan. Transformers provably learn algorithmic solutions for graph connectivity, but only with the right data, 2026. URL https://arxiv.org/abs/2510.19753.

Ruiqi Zhang, Spencer Frei, and Peter L. Bartlett. Trained transformers learn linear models in-context. *Journal of Machine Learning Research*, 25(49):1–55, 2024a. URL http://jmlr.org/papers/v25/23-1042.html.

Ruiqi Zhang, Jingfeng Wu, and Peter L. Bartlett. In-context learning of a linear transformer block: Benefits of the mlp component and one-step gd initialization, 2024b.

Yedi Zhang, Aaditya K. Singh, Peter E. Latham, and Andrew Saxe. Training dynamics of in-context learning in linear attention, 2025. URL https://arxiv.org/abs/2501.16265.

Yufeng Zhang, Fengzhuo Zhang, Zhuoran Yang, and Zhaoran Wang. What and how does in-context learning learn? bayesian model averaging, parameterization, and generalization, 2023. URL https://arxiv.org/abs/2305.19420.

Appendix A. Experimental details

The data structure follows the Section 2 setup. For all experiments we choose a stationary kernel, *i.e.*, distance-dependent correlations, given by

$$K_{ab} := \exp(-|a-b|/\xi). \tag{2}$$

Here the correlation length ξ is manifestly part of the definition of K. This definition is convenient to choose due to the clarity of its correlation length parameter, as well as the fact that our Gaussian model with this kernel is equivalent to an AR(1) process which allows for efficient sampling. When query correlations are present, *i.e.*, $\boldsymbol{k}_{\text{test}} \neq \boldsymbol{0}$, we choose matching structure

$$(\boldsymbol{k}_{\text{test}})_a := \exp(-|\ell + 1 - a|/\xi). \tag{3}$$

This parameter ξ is what is referred to on the x-axes of all figures.

Data parameters. A batch of data contains sequences of length ℓ with one query each. We train in an offline manner with n total sequences. Each sequence is defined by a task vector $\boldsymbol{w}$. As in Lu et al. (2025); Letey et al. (2026); Raventós et al. (2023), the total number of task vectors in the pretraining batch may differ from n. The total number of unique task vectors in the batch is called k, and each unique vector is sampled i.i.d. from $\mathcal{N}(\boldsymbol{0}, I_d)$. For all experiments, we take

$$d = 32\,, \quad n = 4096 = 4d^2\,, \quad \ell = 128 = 4d\,, \quad k = 320 = 10d$$

matching the asymptotic scalings of these variables derived in Lu et al. (2025). We take label noise $\rho = 0.01$.

Architecture details. The experiments train sequence models on the above linear regression sequence data with sequentially correlated inputs. The architectures considered include softmax-only layers, softmax+MLP transformer-like architectures, and pure linear attention. Note that we distinguish between the reduced model (where the optimal solution is given directly by Γ^*) from *trainable* fully-parameterised linear attention. Softmax models embed inputs into width $d+1$ with a dedicated label channel, initialise input embeddings with scale proportional to $\sqrt{(d+1)}$, initialise query/key maps on the input channels, and initialise value/output maps to primarily use the label channel with small Gaussian noise. MLP blocks use GELU and are initialised close to zero. Linear-attention models operate directly on concatenated $(\boldsymbol{x}_i, y_i)$ tokens of dimension d+1, with query/key/value matrices initialised near identity. Models are trained by minimising mean-squared error on the held-out query label only. Optimisation uses Optax AdamW-style updates with weight decay `lamb`; the total number of gradient steps is set to T (something large enough to reach training loss convergence), but we employ early stopping and report the test loss of the best checkpointed-model. We use a linear learning-rate warmup from 0 to `max_lr` over the first 10% of training, with learning rate remaining constant after warmup. We use exponential moving average weights for evaluation. Minibatching is used to divide the full dataset of n sequences; we find that smaller batch sizes perform better for softmax attention architectures. Figures 1 and 3 show mean best-test loss over different seeds, where seeds control train data batch, testing samples, and batch divisions. A summary of our training hyperparameters is

$$\texttt{max\_lr} = 0.001\,, \quad \texttt{lamb} = 0.0001\,, \quad T = 10000\,, \quad \text{batch size} = 64 = 2d\,, \quad \#\text{ seeds} = 5.$$

Appendix B. Detailed setup

Here we re-iterate the data and model setup given in the main document with more details fleshed out. We leave the token design matrix general, *i.e.*,

$$\boldsymbol{x}_a \sim_K \mathcal{N}(0, \Sigma/d)$$

where the sample-sample correlations are given by positive semi-definite K as

$$\mathbb{E}_X[X_{ai}X_{bj}] = \frac{1}{d}K_{ab}\Sigma_{ij} \qquad \text{for } a,b \in [\ell] \text{ and } i,j \in [d]. \tag{4}$$

The label noise we take to be sequentially uncorrelated

$$\boldsymbol{\varepsilon} \sim \mathcal{N}(0, \rho I_{\ell+1}) \qquad \text{independent of all } X \text{ and } \boldsymbol{x}_{\text{test}}.$$

The query $\boldsymbol{x}_{\text{test}}$ may be correlated with the contexts as

$$\mathbb{E}[\boldsymbol{x}_{\text{test}}|X] = X^\top \boldsymbol{m}, \qquad \text{Var}(\boldsymbol{x}_{\text{test}}|X) = \frac{1}{d}(1 - k_{-1})\Sigma$$

for

$$\boldsymbol{m} \equiv K^{-1}\boldsymbol{k}_{\text{test}} \in \mathbb{R}^\ell, \qquad k_{-1} \equiv \boldsymbol{k}_{\text{test}}^\top K^{-1}\boldsymbol{k}_{\text{test}} \leq 1.$$

This is equivalent to sampling, as in eq (4), using $\ell+1 \times \ell+1$ positive-semidefinite correlation kernel

$$K_{\text{query}} = \begin{bmatrix} K & \boldsymbol{k}_{\text{test}} \\ \boldsymbol{k}_{\text{test}}^\top & 1 \end{bmatrix}.$$

As a final note, this entire data setup is equivalent to

$$X = \frac{1}{\sqrt{d}}\sqrt{K}Z\sqrt{\Sigma}, \qquad Z_{si} \sim_{\text{i.i.d.}} \mathcal{N}(0,1).$$

B.1. Notation and Assumptions

We will consider a standard high-dimensional proportional limit, as in (Letey et al., 2026; Lu et al., 2025). The context length scales as

$$\alpha \equiv \frac{\ell}{d} = \Theta_{d,\ell}(1).$$

The usual high dimensional assumptions on Σ are such that

$$\text{tr}(\Sigma), \ \text{tr}\left(\Sigma \boldsymbol{w}\boldsymbol{w}^\top\right) = \Theta(d).$$

These are a succinct summary of conditions on both Σ and $\boldsymbol{w}$. The first condition $\text{tr}(\Sigma) = \Theta(d)$ ensures that there is reasonable signal to estimate in the tokens; note that after this section we will always assume $\Sigma = I_d$. The second condition $\text{tr}\left(\Sigma \boldsymbol{w}\boldsymbol{w}^\top\right) = \Theta(d)$ is equivalent to $\|\boldsymbol{w}\|^2 = \Theta(d)$ ensuring that there is enough recoverable task signal. Given these assumptions, everything will be written in normalised $\Theta_d(1)$ quantities

$$\text{tr}[\Sigma] := \frac{1}{d}\text{tr}(\Sigma), \qquad \text{tr}\left[\Sigma \boldsymbol{w}\boldsymbol{w}^\top\right] := \frac{1}{d}\text{tr}\left(\Sigma \boldsymbol{w}\boldsymbol{w}^\top\right).$$

These need to be extended to assumptions about K. We will take

$$\text{tr}[K] = \frac{1}{\ell}\text{tr}(K) = 1$$

The final thing to reason about is the contributions from K^2 and from $\boldsymbol{k}_{\text{test}}$. This is where the notion of "correlation length" captured in K and $\boldsymbol{k}_{\text{test}}$ becomes important, as the terms

$$\text{tr}\left(K^2\right), \quad k_{-1} \equiv \boldsymbol{k}_{\text{test}}^\top K^{-1}\boldsymbol{k}_{\text{test}}, \qquad k_0 \equiv \boldsymbol{k}_{\text{test}}^\top \boldsymbol{k}_{\text{test}}, \qquad k_1 \equiv \boldsymbol{k}_{\text{test}}^\top K\boldsymbol{k}_{\text{test}}$$

will appear in the computation.

Example: the exponential kernel It is often convenient and natural to choose K to be PSD and Toeplitz, *i.e.*,

$$K_{st} = c(|s - t|).$$

This means the correlations between $\boldsymbol{x}_s$ and $\boldsymbol{x}_t$ are stationary: they only depend on their relative distance in the sequence. A particularly nice example of such a matrix is one with

$$c(\tau) = \exp(-\tau/\xi).$$

As we've seen from Figure 2, there are various cases that may occur.

- Finite correlations: $\xi = \Theta_\ell(1)$. Then $k_0, k_1, k_2 = \Theta(1)$.

- Subleading correlations: Have $\xi = o(\ell)$. If $\xi^2 = o(\ell)$, then $k_0, k_1, k_2 = o(\ell)$; else, have $k_0, k_2 = o(\ell)$, $k_1 = O(\ell)$.

- Proportional correlations: $\xi \propto \ell$. Then $k_0, k_2 = \Theta(\ell)$, $k_1 = \Theta(\ell^2)$.

Appendix C. Population loss

We will be considering linear features

$$\boldsymbol{y} = X\boldsymbol{w} + \boldsymbol{\varepsilon}, \quad y_{\text{test}} = \boldsymbol{x}_{\text{test}}^\top \boldsymbol{w} + \mathcal{N}(0, \rho).$$

For each such sequence of $X, \boldsymbol{y}, \boldsymbol{x}_{\text{test}}, y_{\text{test}}$, define a data matrix

$$H = \boldsymbol{x}_{\text{test}} \left[\tfrac{d}{\ell}(X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top X \quad \tfrac{1}{\ell}(X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top (X\boldsymbol{w} + \boldsymbol{\varepsilon}) \right] \in \mathbb{R}^{d \times (d+1)}.$$

The predictor we will use for y_{test} is given by

$$\hat{y}_{\text{test}} = \langle \Gamma, H \rangle = \operatorname{tr}\left(\Gamma H^\top \right)$$

for parameters $\Gamma \in \mathbb{R}^{d \times (d+1)}$. Before we consider ICL error at optimal parameters Γ^*, we can compute a population formula for

$$\mathcal{E}_{\text{ICL}}(\Gamma) = \mathbb{E}_{\text{new data}} \left[(y_{\text{test}}^{\text{new}} - \langle \Gamma, H^{\text{new}} \rangle)^2 \right].$$

This formula will be different depending on which asymptotic treatment of k_2, k_0, k_1 we are considering.

Notation. We will use the convention of row-wise vectorisation, and so have

$$\operatorname{Vec}(vu^\top) = v \otimes u \in \mathbb{R}^{\dim(v) \times \dim(u)}, \qquad \operatorname{Vec}(vu^\top)\operatorname{Vec}(vu^\top)^\top = (vv^\top) \otimes (\boldsymbol{u}\boldsymbol{u}^\top)$$

Lemma 1 *Suppose we have weak correlations, i.e.,* $k_2, k_0, k_1 = \Theta(1)$. *For* $\boldsymbol{w} \sim \mathcal{N}(\boldsymbol{0}, C)$, *writing* $\operatorname{tr}[C] = \operatorname{tr}(C)/d$, *with*

$$\rho_1 \equiv \operatorname{tr}[C] + \rho, \qquad \rho_2 \equiv k_2 \operatorname{tr}[C] + \rho$$

*have **population ICL risk***

$$\mathcal{E}_{\text{ICL}}(\Gamma) = \rho + \operatorname{tr}[C] - 2\frac{1}{d}\operatorname{tr}\left(\Gamma A^\top\right) + \frac{1}{d}\operatorname{tr}\left(\Gamma B\Gamma^\top\right) + \frac{1}{d^2}\operatorname{tr}\left(\mathcal{T}\operatorname{vec}(\Gamma)\operatorname{vec}(\Gamma)^\top\right) \quad (5)$$

for

$$A \equiv \begin{bmatrix} C + \frac{1}{\alpha}k_0 \operatorname{tr}[C]I_d & \boldsymbol{0} \end{bmatrix}, \qquad B \equiv \begin{bmatrix} C + \frac{\rho_2}{\alpha}I_d & \boldsymbol{0} \\ \boldsymbol{0} & \rho_1^2 \end{bmatrix}$$

$$\mathcal{T} \equiv \frac{k_1 \operatorname{tr}[C] + \rho k_0}{\alpha^2} \operatorname{vec}\left(\begin{bmatrix} I & \boldsymbol{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} I & \boldsymbol{0} \end{bmatrix}\right)^\top$$

$$+ \frac{k_0}{\alpha}\left(\operatorname{vec}\left(\begin{bmatrix} I & \boldsymbol{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} C & \boldsymbol{0} \end{bmatrix}\right)^\top + \operatorname{vec}\left(\begin{bmatrix} C & \boldsymbol{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} I & \boldsymbol{0} \end{bmatrix}\right)^\top\right)$$

Appendix D. Breakdown of the high-dimensional analysis for strong correlations

In the subsequent appendices (starting with Appendix E), we will use random-matrix-theoretic techniques to derive an asymptotically-precise formula for $\mathcal{E}_{\mathrm{ICL}}(\Gamma^*)$ in the high-dimensional limit. This analysis hinges on our ability to derive what are known as *deterministic equivalents* for the high-dimensional behaviour of random matrices related to the optimal parameter matrix Γ^* of our reduced linear attention model. Before we embark on the analysis, in this Appendix we derive a condition on the correlations K that is necessary for the random matrix analysis to go through as in Lu et al. (2025). This condition is violated when the correlations are strong, which shows that this approach does not obviously extend to the strong correlation regime.

We recall from Lu et al. (2025) that we can write the optimal parameter matrix in vectorized form as

$$\mathrm{vec}(\Gamma^*) = \left(\frac{n}{d}\lambda I_{d(d+1)} + \sum_{\mu=1}^{n} \mathrm{vec}(H^\mu) \otimes \mathrm{vec}(H^\mu)\right)^{-1} \sum_{\mu=1}^{n} y_{\mathrm{test}}^\mu \,\mathrm{vec}(H^\mu). \tag{6}$$

Starting from this formula, a key ingredient of the high-dimensional analysis is (as we will see in the sequel), the assumption that quadratic forms

$$\mathrm{vec}(H^\mu)^\top G\,\mathrm{vec}(H^\mu), \tag{7}$$

for G a possibly-random matrix that is independent of H^μ, concentrate to their expectations in the high-dimensional limit. To gain some insight into whether this assumption is justified, we consider the test case $G = I$, which corresponds to considering the norms $\|\mathrm{vec}(H^\mu)\|^2$. If the norms do not concentrate, then for the class of matrices G that appear in the computation, neither should the more general quadratic forms.

To determine when the norms concentrate, suppress the index $\mu \in [n]$, and recall that we have

$$H = \boldsymbol{x}_{\mathrm{test}}\left[\tfrac{d}{\ell}(X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top X \quad \tfrac{1}{\ell}(X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top(X\boldsymbol{w} + \boldsymbol{\varepsilon})\right].$$

We will simplify our analysis here, as our goal is to focus on intuition, by ignoring the noise terms given by $\boldsymbol{\varepsilon}$; these do not affect the argument. Thus we write

$$\mathrm{vec}(H) = \boldsymbol{v} \otimes \boldsymbol{x}_{\mathrm{test}} + \text{ignored noise contribution}$$

where

$$\boldsymbol{v} = \begin{bmatrix} \tfrac{d}{\ell}X^\top X\boldsymbol{w} \\ \tfrac{1}{\ell}\boldsymbol{w}^\top X^\top X\boldsymbol{w} \end{bmatrix}.$$

We thus have that

$$\frac{1}{d^2}\|\mathrm{vec}(H)\|^2 = \frac{1}{d^2}\|\boldsymbol{x}_{\mathrm{test}}\|^2\|\boldsymbol{v}\|^2$$

$$= \frac{1}{d^2}\|\boldsymbol{x}_{\mathrm{test}}\|^2\left(\frac{d^2}{\ell^2}\boldsymbol{w}^\top X^\top X X^\top X\boldsymbol{w} + \frac{1}{\ell^2}(\boldsymbol{w}^\top X^\top X\boldsymbol{w})^2\right)$$

$$\overset{d}{=} \left(\frac{1}{d}\boldsymbol{m}^\top K^{1/2}ZZ^\top K^{1/2}\boldsymbol{m} + 2\frac{\sqrt{1-k_{-1}}}{d}\boldsymbol{z}^\top Z^\top K^{1/2}\boldsymbol{m} + \frac{1-k_{-1}}{d}\|\boldsymbol{m}z\|^2\right)$$

$$\times \left(\frac{1}{d^2\ell^2}\boldsymbol{w}^\top Z^\top KZZ^\top KZ\boldsymbol{w} + \frac{1}{d^4\ell^2}(\boldsymbol{w}^\top Z^\top KZ\boldsymbol{w})^2\right),$$

where $\stackrel{d}{=}$ means equivalent in distribution. Here Z is an $\ell \times d$ matrix with i.i.d. $\mathcal{N}(0,1)$ entries and $\boldsymbol{z} \sim \mathcal{N}(0, I_d)$ for the query (independent of Z).

Upon analysis of each of these terms, which we omit for brevity, we find that the problematic term is

$$\frac{1}{d^2 \ell^2} \boldsymbol{w}^\top Z^\top K Z Z^\top K Z \boldsymbol{w} \stackrel{d}{=} \|\boldsymbol{w}\|^2 (Z^\top K Z Z^\top K Z)_{11}$$

$$\stackrel{d}{=} \frac{\|\boldsymbol{w}\|^2}{d} \frac{1}{\ell^2} \sum_{a,b=1}^{\ell} \kappa_a \kappa_b Z_{a1} Z_{b1} \frac{1}{d} \sum_{i=1}^{d} Z_{ai} Z_{bi}$$

$$= \frac{\|\boldsymbol{w}\|^2}{d} \frac{1}{d} \left(\frac{1}{\ell} \sum_{a=1}^{\ell} \kappa_a Z_{a1}^2 \right)^2 + \frac{\|\boldsymbol{w}\|^2}{d} \frac{1}{\ell^2} \sum_{a,b=1}^{\ell} \kappa_a \kappa_b Z_{a1} Z_{b1} \frac{1}{d} \sum_{i=2}^{d} Z_{ai} Z_{bi}.$$

We have arrived at this expression by exploiting the fact that the distribution of the matrix Z is invariant under both left and right rotation to perform two changes of basis: (1) we let K have orthogonal eigendecomposition $K = O \operatorname{diag}(\kappa_1, \ldots, \kappa_\ell) O^\top$, and (2) we choose a basis such that $\boldsymbol{w} = (\|\boldsymbol{w}\|, 0, \ldots, 0)^\top$.

Now we must analyse this expression to provide a condition on K, specifically its eigenvalues $\kappa_1, \ldots, \kappa_\ell$, for when this term does or does not concentrate. We focus on the second term in the sum, supposing for simplicity that $\|\boldsymbol{w}\|^2 / d = 1$:

$$A := \frac{1}{\ell^2} \sum_{a,b=1}^{\ell} \kappa_a \kappa_b Z_{a1} Z_{b1} \frac{1}{d} \sum_{i=2}^{d} Z_{ai} Z_{bi}.$$

It has moments given by

$$\mathbb{E}[A] = \frac{d-1}{d} \frac{1}{\ell^2} \sum_{a=1}^{\ell} \kappa_a^2,$$

$$\operatorname{var}[A] = \frac{2}{\ell^4 d^2} (d-1) \left(\sum_{a=1}^{\ell} \kappa_a^2 \right)^2 + \frac{2}{\ell^4 d^2} (d^2 - 1) \sum_{a=1}^{\ell} \kappa_a^4,$$

and so the square of the coefficient of variation of A is therefore

$$\frac{\operatorname{var}[A]}{\mathbb{E}[A]^2} = 2 \left[\frac{1}{d-1} + \frac{d^2 - 1}{(d-1)^2} \frac{\sum_{a=1}^{\ell} \kappa_a^4}{\left(\sum_{a=1}^{\ell} \kappa_a^2 \right)^2} \right]$$

$$= 2[1 + \mathcal{O}(d^{-1})] \frac{\sum_{a=1}^{\ell} \kappa_a^4}{\left(\sum_{a=1}^{\ell} \kappa_a^2 \right)^2} + \mathcal{O}(d^{-1}).$$

We now notice that the K-dependence here

$$\frac{\sum_{a=1}^{\ell} \kappa_a^4}{\left(\sum_{a=1}^{\ell} \kappa_a^2 \right)^2} = \frac{\operatorname{tr}(K^4)}{\operatorname{tr}(K^2)^2}$$

looks like some sort of participation ratio. Indeed if $K = I_\ell$ (uncorrelated case, we know this concentrates) then

$$\frac{\operatorname{tr}(K^4)}{\operatorname{tr}(K^2)^2} = \frac{1}{\ell} \to 0,$$

while if $K = \mathbf{1}\mathbf{1}^\top$ (maximally correlated case, here $\boldsymbol{x}_1 = \ldots = \boldsymbol{x}_\ell$) then

$$\frac{\mathrm{tr}(K^4)}{\mathrm{tr}(K^2)^2} = 1.$$

In the maximally-correlated case $K = \mathbf{1}\mathbf{1}^\top$, we see that $A = Z_{11}^2 \frac{1}{d} \sum_{i=2}^d Z_{1i}^2$ tends in distribution to a χ^2 random variable with 1 degree of freedom (as $\frac{1}{d} \sum_{i=2}^d Z_{1i}^2 \to 1$ almost surely), which clearly does not concentrate. This gives us an intuitive heuristic for concentration of $\| \mathrm{vec}(H) \|$:

Necessary condition for concentration

Quadratic forms in $\mathrm{vec}(H)$ converge to a deterministic high-dimensional limit only if

$$\frac{\mathrm{tr}(K^4)}{\mathrm{tr}(K^2)^2} \to 0. \tag{8}$$

For the exponential case $K_{ab} = \exp(-|a - b|/\xi)$, we can estimate

$$\frac{\mathrm{tr}(K^4)}{\mathrm{tr}(K^2)^2} \sim \frac{\xi}{\ell}$$

at large ℓ, which gives the condition we expected. We see that for correlations that persist on the same order as the context length, we do not have concentration of $\mathrm{vec}(H)$, and thus we do not have concentration of $\mathcal{E}_{\mathrm{ICL}}(\Gamma^*)$. An extension of great interest to us would be to analyse this nonconcentration for a wider range of K choices, *e.g.* how does this participation ratio behave for other non-exponential (or even non-Toeplitz) choices of K?

Remark. It is worth noting that the case of $k_2 = o(\ell)$ does not violate this condition (8). Yet, the formula we shall derive will assume that $k_2 = \Theta(1)$. This might potentially be too restrictive of an assumption; indeed Figure 2 shows great match between simulation and our theory curve, so long as $k_1 = o(\ell)$ as well. We choose to study the $k_0, k_1, k_2 = \Theta(1)$ branch only as this is the branch where the test matrices $A, B, \mathcal{T}$ defined in Lemma 1 have order 1 entries. Indeed, under numerical investigation, this population loss formula appears to be accurate for the $k_0, k_1, k_2 = o(\ell)$ case. However, we choose not to study this branch in this work due to the elements of $A, B, \mathcal{T}$ becoming extensive in ℓ.

Appendix E. Asymptotic formula for ICL error in the weak-correlations case

Here we present a deterministic formula for $\mathcal{E}_{\mathrm{ICL}}(\Gamma^*)$ in the weak-correlations case. Again, the optimal parameters are given by eq. (6). The explicit ridge λ is a parameter that adds regularisation to this solution; it is common to take the ridgeless limit $\lambda \to 0$ in similar works.

Here we have n sample sequences, giving n different $H^\mu \in \mathbb{R}^{d \times (d+1)}$ data embeddings, where the sequence length is of course $\ell + 1$ (ℓ main tokens and one query). We will also introduce a new parameter k that controls the number of regression task vectors $\boldsymbol{w}$ that define the labels in these sequences. The phenomenology surrounding k is not discussed in this work for space reasons, but we include it to best match previous work on this model

for consistency (Lu et al., 2025; Letey et al., 2026; Raventós et al., 2023). We sample k task vectors $\boldsymbol{w}_j \sim_{\text{i.i.d.}} \mathcal{N}(\boldsymbol{0}, I_d)$ and these k vectors are distributed evenly across the n sequences. This serves to limit the "task diversity" of the training batch, allowing us to judge how much of the true task distribution $\mathcal{N}(\boldsymbol{0}, I_d)$ the model is truly learning in-context.

We consider the same asymptotic treatment as derived in Lu et al. (2025), namely $d \to \infty$ with

$$\alpha := \frac{\ell}{d}, \quad \tau := \frac{n}{d^2}, \quad \kappa := \frac{k}{d}$$

all taken to be $\Theta(1)$ as $d \to \infty$.

An important quantity to define is the Stieltjes transform of the task sample covariance. This will be given by

$$\mathcal{M}_\kappa(z) := \lim_{d \to \infty, k = \kappa d} \frac{1}{d} \operatorname{tr}\left(\left(\frac{1}{k} \sum_{j=1}^{k} \boldsymbol{w}_j \boldsymbol{w}_j^\top + z I_d \right)^{-1} \right) \tag{9}$$

We will also use

$$\mathcal{M}'_\kappa(z) := \frac{\mathrm{d}}{\mathrm{d}z} \mathcal{M}_\kappa(z).$$

These quantities will appear throughout our formula.

The explicit ridge parameter λ will be modulated by the fact that we have finite samples, and will appear in our formula through an *effective* ridge defined implicitly by

$$\tilde{\lambda} \mathcal{M}_\kappa\left(\tilde{\lambda} + \frac{\rho_2}{\alpha} \right) + \frac{\lambda \tau}{\tilde{\lambda}} = 1 - \tau$$

We will sometimes write $\mathcal{M} := \mathcal{M}_\kappa(\sigma), \quad \mathcal{M}' := \mathcal{M}'_\kappa(\sigma)$ for shorthand, where

$$\sigma = \tilde{\lambda} + \frac{\rho_2}{\alpha}, \quad \tilde{\sigma} = \sigma - \frac{k_0}{\alpha}.$$

Finally, we are only considering the case where the sequential correlations defined by $K, \boldsymbol{k}_{\text{test}}$ are sufficiently weak, *i.e.*,

$$k_2 := \frac{1}{\ell} \operatorname{tr}\left(K^2 \right), \quad k_0 := \boldsymbol{k}_{\text{test}}^\top \boldsymbol{k}_{\text{test}}, \quad k_1 := \boldsymbol{k}_{\text{test}}^\top K \boldsymbol{k}_{\text{test}}$$

are all $\Theta(1)$ as $d, \ell \to \infty$. These terms will appear in the constants

$$\rho_1 \equiv 1 + \rho, \quad \rho_2 \equiv k_2 + \rho, \quad \phi_1 \equiv \frac{1}{\alpha^2}(k_1 + \rho k_0), \quad \phi_2 \equiv \frac{1}{\alpha} k_0.$$

We present the main formula below. All proofs are given in subsequent sections.

Proposition 2 *Consider isotropic tokens and tasks, i.e., $\Sigma = I_d = C$, with sequential correlations in the tokens given by PSD $K \in \mathbb{R}^{\ell \times \ell}$ and $\boldsymbol{k}_{\text{test}} \in \mathbb{R}^\ell$ for $\boldsymbol{k}_{\text{test}}^\top K^{-1} \boldsymbol{k}_{\text{test}} \leq 1$. We then have that ICL error for a model defined by Γ^* concentrates as $d \to \infty$ in the above limit, i.e.,*

$$\mathcal{E}_{\text{ICL}}(\Gamma^*) \simeq e_{\text{ICL}}^{\text{corr}}(\alpha, \kappa, \tau, \rho, k_0, k_1, k_2)$$

for

$$e_{\text{ICL}}^{\text{corr}}(\alpha, \kappa, \tau, \rho, k_0, k_1, k_2) = e_{\text{ICL}}^{\text{uncorr}}\left(\frac{1 + \rho}{k_2 + \rho} \alpha, \kappa, \tau, \rho \right) + e_{\text{query}}(\alpha, \kappa, \tau, \rho, k_0, k_1, k_2)$$

where

$$e_{\text{query}} = -2\phi_2 + q_{\text{query}}(\mathcal{M} + \tilde{\lambda}\mathcal{M}') + \phi_2(2\sigma - \phi_2)\mathcal{M}' + 2\phi_2\tilde{\sigma}\mathcal{M}$$

$$+ \ \boldsymbol{m}_2^\top S^\top M \, S\boldsymbol{m}_2 - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + 2(1 + \phi_2)\boldsymbol{m}_1^\top S\boldsymbol{m}_2 + (\phi_1 + 2\phi_2)\left(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1^\top S\boldsymbol{m}_2\right)^2$$

$$+ \ \frac{\rho_2}{\alpha}\Big[q_{\text{query}}(\mathcal{M} + \tilde{\lambda}\mathcal{M}') + \phi_2(2\sigma - \phi_2)\mathcal{M}' + \boldsymbol{m}_2^\top S^\top M \, S\boldsymbol{m}_2 - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + 2\phi_2\mathcal{M}\Big].$$

for

$$\boldsymbol{m}_1 = \begin{bmatrix} \mathcal{M} \\ \phi_2(1 - \sigma\mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_2 = \begin{bmatrix} 1 - \tilde{\sigma}\mathcal{M} \\ \phi_2(1 - \tilde{\sigma} + \sigma\tilde{\sigma}\mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_3 = \begin{bmatrix} \mathcal{M} + \tilde{\sigma}\mathcal{M}' \\ \phi_2(1 - \sigma\mathcal{M} - \tilde{\sigma}\mathcal{M} - \sigma\tilde{\sigma}\mathcal{M}') \end{bmatrix}$$

$$M = \begin{bmatrix} -\mathcal{M}' & \phi_2(\mathcal{M} + \sigma\mathcal{M}') \\ \phi_2(\mathcal{M} + \sigma\mathcal{M}') & \phi_2^2(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}') \end{bmatrix}, \quad S = \begin{bmatrix} \mathcal{M} & 1 + \phi_2(1 - \sigma\mathcal{M}) \\ 1 + \phi_2(1 - \sigma\mathcal{M}) & -\phi_1 + \phi_2^2(1 - \sigma + \sigma^2\mathcal{M}) \end{bmatrix}^{-1}$$

$$q_{\text{query}} = \frac{1}{\tau - (1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2\mathcal{M}')}\left((2\frac{k_0}{\alpha}\Big(\tilde{\lambda}(\mathcal{M} + \sigma\mathcal{M}') + (1 - 2\sigma\mathcal{M})\Big) + \frac{k_0^2}{\alpha^2}\Big(\mathcal{M} - \tilde{\lambda}\mathcal{M}'\Big)\right.$$

$$\left. + \ \boldsymbol{m}_2^\top S\boldsymbol{m}_2 - \tilde{\lambda}\Big(-2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + \boldsymbol{m}_2^\top S^\top M S\boldsymbol{m}_2\Big)\right).$$

Appendix F. Proofs of Population Averages

We begin for necessary y_{test}, H averages needed for terms in the population loss formula in Lemma 1. *A language model was used at various stages of these proofs to help with the Wick expansions of the 6th order moments in terms of traces. All expressions were subsequently carefully verified by the authors both analytically and numerically.*

Lemma 3 *Suppose we have $k_2, k_0, k_1 = \Theta(1)$ i.e., weak correlations. Write $\boldsymbol{v} = \Sigma\boldsymbol{w}$. Then we have*

$$\mathbb{E}[y_{\text{test}}^2] = \text{tr}\Big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\Big] + \rho$$

$$\mathbb{E}[y_{\text{test}}H] \approx \frac{1}{d}\Big[\boldsymbol{v}\boldsymbol{v}^\top + \tfrac{1}{\alpha}k_0\,\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big]\Sigma \quad \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\boldsymbol{v}\Big]$$

$$\mathbb{E}[\text{vec}(H)\text{vec}(H)^\top] \approx \frac{1}{d}\Sigma \otimes \begin{bmatrix} \boldsymbol{v}\boldsymbol{v}^\top + \big(k_2\,\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\Sigma/\alpha & \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\boldsymbol{v} \\ \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\boldsymbol{v}^\top & \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big))^2 \end{bmatrix} + \frac{1}{d^2}\mathcal{X}$$

for $d(d+1) \times d(d+1)$ tensor $\mathcal{X}$ given by

$$\mathcal{X} = \frac{1}{\alpha^2}\Big(k_1\,\text{tr}[\Sigma W] + \rho k_0\Big)\,\text{vec}\left(\begin{bmatrix}\Sigma & \boldsymbol{0}\end{bmatrix}\right)\text{vec}\left(\begin{bmatrix}\Sigma & \boldsymbol{0}\end{bmatrix}\right)^\top$$

$$+ \ \frac{k_0}{\alpha}\Big(\text{vec}\left(\begin{bmatrix}\Sigma & \boldsymbol{0}\end{bmatrix}\right)\text{vec}\left(\Big[S \quad \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\boldsymbol{v}\Big]\right)^\top$$

$$+ \ \text{vec}\left(\Big[S \quad \big(\text{tr}\big[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\big] + \rho\big)\boldsymbol{v}\Big]\right)\text{vec}\left(\begin{bmatrix}\Sigma & \boldsymbol{0}\end{bmatrix}\right)^\top\Big).$$

The "$\approx$" means subleading terms in ℓ, d are neglected.

Proof Let's go term by term.

Label-label term. Conditioning on X, we have

$$\mathbb{E}[\boldsymbol{x}_{\text{test}}\boldsymbol{x}_{\text{test}}^\top \mid X] = \frac{1}{d}(1 - k_{-1})\Sigma + (X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top$$

and so

$$\mathbb{E}[y_{\text{test}}^2] = \mathbb{E}[\text{tr}\left(\boldsymbol{x}_{\text{test}}\boldsymbol{x}_{\text{test}}^\top \boldsymbol{w}\boldsymbol{w}^\top\right) + \rho]$$

$$= \frac{1}{d}(1 - k_{-1})\mathbb{E}_X\left[\text{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right)\right] + \mathbb{E}_X\left[\text{tr}\left((X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \boldsymbol{w}\boldsymbol{w}^\top\right)\right] + \rho$$

$$= (1 - k_{-1} + k_{-1})\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right] + \rho.$$

Label-feature term. For

$$\boldsymbol{b} = X^\top(X\boldsymbol{w} + \boldsymbol{\varepsilon}), \quad c = (X\boldsymbol{w} + \boldsymbol{\varepsilon})^\top(X\boldsymbol{w} + \boldsymbol{\varepsilon})$$

we have

$$y_{\text{test}}H = \boldsymbol{x}_{\text{test}}(\boldsymbol{x}_{\text{test}}^\top\boldsymbol{w} + \varepsilon_{\text{test}})\left[\frac{d}{\ell}\boldsymbol{b}^\top \quad \frac{1}{\ell}c\right]$$

and so

$$\mathbb{E}[y_{\text{test}}H] = \frac{1}{d}(1 - k_{-1})\Sigma\boldsymbol{w}\begin{bmatrix}\frac{d}{\ell}\mathbb{E}_X[\boldsymbol{b}]\\ \frac{1}{\ell}\mathbb{E}_X[c]\end{bmatrix}^\top + \left[\frac{d}{\ell}\mathbb{E}_X[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top\boldsymbol{w}\boldsymbol{b}^\top] \quad \frac{1}{\ell}\mathbb{E}_X[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top\boldsymbol{w}\,c]\right]$$

Have

$$\frac{d}{\ell}\mathbb{E}_X[\boldsymbol{b}] = \text{tr}[K]\boldsymbol{v}$$

$$\frac{1}{\ell}\mathbb{E}_X[c] = \text{tr}[K]\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right] + \rho$$

$$\frac{d}{\ell}\mathbb{E}_X[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top\boldsymbol{w}\boldsymbol{b}^\top] = \frac{1}{d}\cdot\frac{1}{\ell}(k_{-1}\,\text{tr}(K) + k_0)\boldsymbol{v}\boldsymbol{v}^\top + \frac{1}{d}\cdot\frac{1}{\ell}k_0\,\text{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right)\Sigma$$

$$\approx \frac{1}{d}\cdot\left(k_{-1}\,\text{tr}[K]\boldsymbol{v}\boldsymbol{v}^\top + \frac{1}{\alpha}k_0\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right]\Sigma\right)$$

$$\frac{1}{\ell}\mathbb{E}_X[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top\boldsymbol{w}\,c] = \frac{1}{d}\cdot\rho k_{-1}\boldsymbol{v} + \frac{1}{d}\cdot k_{-1}\,\text{tr}[K]\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right]\boldsymbol{v} + \frac{2}{d^2}\frac{k_0}{\ell}\boldsymbol{v}\boldsymbol{v}^\top\boldsymbol{w}$$

$$\approx \frac{1}{d}\cdot k_{-1}\left(\text{tr}[K]\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right] + \rho\right)\boldsymbol{v}$$

Combining gives

$$\mathbb{E}[y_{\text{test}}H] = \frac{1}{d}\left[\text{tr}[K]\boldsymbol{v}\boldsymbol{v}^\top + \frac{1}{\alpha}k_0\,\text{tr}\left[\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right]\Sigma \quad \rho_1\boldsymbol{v}\right]$$

Feature-feature term Using $\boldsymbol{b}, c$ as above we can write more easily

$$\text{Vec}(H)\text{Vec}(H)^\top = (\boldsymbol{x}_{\text{test}}\boldsymbol{x}_{\text{test}}^\top) \otimes \left(\begin{bmatrix}\frac{d}{\ell}\boldsymbol{b}\\ \frac{1}{\ell}c\end{bmatrix}\begin{bmatrix}\frac{d}{\ell}\boldsymbol{b}\\ \frac{1}{\ell}c\end{bmatrix}^\top\right)$$

Taking conditional expectation over $\boldsymbol{x}_{\text{test}}$ for fixed X, our expression simplifies as

$$\mathbb{E}\left[\text{Vec}(H)\text{Vec}(H)^\top\right] = \frac{1}{d}(1 - k_{-1})\Sigma \otimes \mathbb{E}_{X,\epsilon}\left[\boldsymbol{u}\boldsymbol{u}^\top\right] + \mathbb{E}_{X,\epsilon}\left[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes (\boldsymbol{u}\boldsymbol{u}^\top)\right].$$

where I'm writing $\boldsymbol{u}^\top = \left[\frac{d}{\ell}\boldsymbol{b}^\top \quad \frac{1}{\ell}c\right]$ for convenience. Now let's start with the $\mathbb{E}[\boldsymbol{u}\boldsymbol{u}^\top]$ term.

We have

$$\mathbb{E}[\boldsymbol{b}\boldsymbol{b}^\top] = \mathbb{E}[X^\top X \boldsymbol{w}\boldsymbol{w}^\top X^\top X] + \mathbb{E}[X^\top \boldsymbol{\varepsilon}\boldsymbol{\varepsilon}^\top X]$$

$$= \frac{1}{d^2}\left(\mathrm{tr}(K^2) + \mathrm{tr}(K)^2\right)\Sigma\boldsymbol{w}\boldsymbol{w}^\top\Sigma + \frac{1}{d^2}\mathrm{tr}(K^2)\,\mathrm{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right)\Sigma + \frac{1}{d}\rho\,\mathrm{tr}(K)\Sigma$$

$$\mathbb{E}[c\boldsymbol{b}] = \mathbb{E}[\boldsymbol{w}^\top X^\top X \boldsymbol{w} X^\top X \boldsymbol{w}] + 2\mathbb{E}[\boldsymbol{\varepsilon}^\top X \boldsymbol{w} X^\top \boldsymbol{\varepsilon}] + \mathbb{E}[\boldsymbol{\varepsilon}^\top \boldsymbol{\varepsilon} X^\top X \boldsymbol{w}]$$

$$= \left(\frac{1}{d^2}\left(\mathrm{tr}(K)^2 + 2\,\mathrm{tr}(K^2)\right)\mathrm{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right) + \frac{1}{d}\rho(\ell+2)\,\mathrm{tr}(K)\right)\Sigma\boldsymbol{w}$$

$$\mathbb{E}[c^2] = \mathbb{E}[\boldsymbol{w}^\top X^\top X \boldsymbol{w}\boldsymbol{w}^\top X^\top X \boldsymbol{w}] + 2\mathbb{E}[\boldsymbol{w}^\top X^\top X \boldsymbol{w}\boldsymbol{\varepsilon}^\top \boldsymbol{\varepsilon}] + 4\mathbb{E}[\boldsymbol{w}^\top X^\top \boldsymbol{\varepsilon}\boldsymbol{\varepsilon}^\top X \boldsymbol{w}] + \mathbb{E}[\boldsymbol{\varepsilon}^\top \boldsymbol{\varepsilon}\boldsymbol{\varepsilon}^\top \boldsymbol{\varepsilon}]$$

$$= \frac{1}{d^2}\left(\mathrm{tr}(K)^2 + 2\,\mathrm{tr}(K^2)\right)\mathrm{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right)^2 + \frac{1}{d}\rho(2\ell+4)\,\mathrm{tr}(K)\,\mathrm{tr}\left(\Sigma\boldsymbol{w}\boldsymbol{w}^\top\right) + \rho^2(\ell^2 + 2\ell).$$

This simplifies as

$$\frac{d^2}{\ell^2}\mathbb{E}[\boldsymbol{b}\boldsymbol{b}^\top] \approx \mathrm{tr}[K]^2(\Sigma\boldsymbol{w})(\Sigma\boldsymbol{w})^\top + \frac{1}{\alpha}\left(\rho\,\mathrm{tr}[K] + \mathrm{tr}\left[K^2\right]\mathrm{tr}[\Sigma W]\right)\Sigma$$

$$\frac{d}{\ell^2}\mathbb{E}[c\boldsymbol{b}] \approx \mathrm{tr}[K](\mathrm{tr}[K]\,\mathrm{tr}[\Sigma W] + \rho)\Sigma\boldsymbol{w}$$

$$\frac{1}{\ell^2}\mathbb{E}[c^2] \approx (\rho + \mathrm{tr}[K]\,\mathrm{tr}[\Sigma W])^2$$

upon making the above high-dimensional assumptions.

For the higher order term,

$$\mathbb{E}\left[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes (\boldsymbol{u}\boldsymbol{u}^\top)\right] = \begin{bmatrix} \frac{d^2}{\ell^2}\mathbb{E}[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes \boldsymbol{b}\boldsymbol{b}^T] & \frac{d}{\ell^2}\mathbb{E}[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes c\boldsymbol{b}] \\ \frac{d}{\ell^2}\mathbb{E}[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes c\boldsymbol{b}^T] & \frac{1}{\ell^2}\mathbb{E}[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes c^2] \end{bmatrix}$$

Write

$$\boldsymbol{v} := \Sigma\boldsymbol{w}, \qquad S := \boldsymbol{v}\boldsymbol{v}^\top, \qquad k_{-1} \equiv \mathrm{tr}(KM) = \boldsymbol{k}_{\mathrm{test}}^\top K^{-1}\boldsymbol{k}_{\mathrm{test}}, \quad k_0 \equiv \boldsymbol{k}_{\mathrm{test}}^\top\boldsymbol{k}_{\mathrm{test}}, \quad k_1 \equiv \boldsymbol{k}_{\mathrm{test}}^\top K\boldsymbol{k}_{\mathrm{test}}$$

For large ℓ, d and the chosen high-dimensional assumptions, the Isserlis expansion gives

$$\frac{d^2}{\ell^2}\mathbb{E}\left[((X^\top m)(X^\top m)^\top) \otimes (\boldsymbol{b}\boldsymbol{b}^\top)\right]_{(I,i),(J,j)} \approx \frac{1}{d}k_{-1}\left(\Sigma_{IJ}S_{ij}\,\mathrm{tr}[K]^2 + \frac{1}{\alpha}\left(\mathrm{tr}[\Sigma W]\,\mathrm{tr}\left[K^2\right] + \rho\,\mathrm{tr}[K]\right)\Sigma_{IJ}\Sigma_{ij}\right)$$
$$+ \frac{1}{d^2}\mathcal{X}_{(I,i)(J,j)}$$

with

$$\mathcal{X}_{(I,i)(J,j)} \equiv \frac{1}{\alpha^2}k_1\,\mathrm{tr}[\Sigma W]\Sigma_{Ii}\Sigma_{Jj} + \frac{1}{\alpha}\mathrm{tr}[K]k_0(\Sigma_{Ii}S_{Jj} + \Sigma_{Jj}S_{Ii}) + \rho\frac{1}{\alpha^2}k_0\Sigma_{Ii}\Sigma_{Jj}$$

$$= \frac{1}{\alpha^2}(k_1\,\mathrm{tr}[\Sigma W] + \rho k_0)\Sigma_{Ii}\Sigma_{Jj} + \frac{1}{\alpha}k_0\,\mathrm{tr}[K](\Sigma_{Ii}S_{Jj} + \Sigma_{Jj}S_{Ii}).$$

Now for the second term,

$$\mathbb{E}[(X^\top\boldsymbol{m})(X^\top\boldsymbol{m})^\top \otimes c\boldsymbol{b}]_{(I,i),J} = \mathbb{E}[c(X^\top\boldsymbol{m})_I(X^\top\boldsymbol{m})_J b_i]$$
$$= \mathbb{E}[(X_{sk}w_k + \varepsilon_s)(X_{sn}w_n + \varepsilon_s)X_{tI}X_{pJ}M_{tp}X_{qi}(X_{qj}w_j + \varepsilon_q)]$$
$$= M_{tp}W_{kn}w_j\mathbb{E}[X_{sk}X_{sn}X_{tI}X_{pJ}X_{qi}X_{qj}] + (\ell+2)\rho M_{tp}w_j\mathbb{E}[X_{tI}X_{pJ}X_{qi}X_{qj}]$$

Expanding all the Wick terms gives

$$\frac{d}{\ell^2}\mathbb{E}\left[((X^\top m)(X^\top m)^\top) \otimes (c b)\right]_{(I,i),J} \approx \frac{1}{d}k_{-1}(\mathrm{tr}[\Sigma W] + \rho)\Sigma_{IJ}v_i + \frac{1}{d^2}\mathcal{X}_{(I,i)(J,d+1)}$$

where

$$\mathcal{X}_{(I,i)(J,d+1)} \equiv \frac{1}{\alpha}k_0(\mathrm{tr}[\Sigma W] + \rho)\Sigma_{Ii}v_J.$$

Finally, the last term gives
$$\frac{1}{\ell^2}\mathbb{E}\Big[((X^\top m)(X^\top m)) \otimes c^2\Big]_{IJ} \approx \frac{1}{d}k_{-1}(\mathrm{tr}[\Sigma W] + \rho)^2 \Sigma_{IJ}$$
Gathering everything together gives the required formula. $\blacksquare$

Appendix G. Proof of Proposition 2

This entire section is an extended calculation that serves as a complete proof of the deterministic formula for $\mathcal{E}_{\mathrm{ICL}}(\Gamma^*)$ given in Proposition 2. The methodology in this section is to analyse the random object Γ^*, which contains randomness through the particular training batch $(H^\mu, y_{\mathrm{test}}^\mu)_{\mu=1}^n$, as a resolvent of a larger random matrix, and the necessary $\mathrm{tr}(\Gamma A^\top)$ and $\mathrm{tr}(\Gamma^\top B\Gamma)$ terms as traces of a resolvent against deterministic test matrices. Throughout this entire section we will take $\Sigma = I_d$.

This section will be relatively light on detail as the full calculation setup can be found in both Lu et al. (2025) and Letey et al. (2026). For a more rigorous analysis, with error terms bounded properly, see Lu et al. (2025); here we will simply write $\approx$ when terms can be dropped due to the negligibility in high dimensions.

We care about the MSE loss (computed by Theorem 1) for the optimal parameters

$$\mathrm{vec}(\Gamma^*) = \left(\frac{n}{d}\lambda I + \sum_{\mu=1}^n \mathrm{vec}(H^\mu)\mathrm{vec}(H^\mu)^\top \right)^{-1} \sum_{\mu=1}^n y_{\mathrm{test}}^\mu \mathrm{vec}(H^\mu) \tag{10}$$

for $\mu \in [n]$ denoting the index for the n training sample sequences. Define

$$\boldsymbol{z}_\mu = \begin{bmatrix} y_{\mathrm{test}}^\mu/d \\ \mathrm{vec}(H^\mu)/\sqrt{d} \end{bmatrix} \in \mathbb{R}^{d(d+1)+1}$$

and construct extended resolvent

$$G_{\mathrm{ext}}(\pi) = \frac{1}{\sum_{\mu\in[n]} \boldsymbol{z}_\mu \boldsymbol{z}_\mu^\top + \pi B_{\mathrm{ext}} + \tau\lambda I} . \tag{11}$$

for some test matrix $B_{\mathrm{ext}} \in \mathbb{R}^{(d(d+1)+1)\times(d(d+1)+1)}$.

It's important to note that even though we're studying sequentially-correlated data, the vectors $\boldsymbol{z}_\mu$s are still independent of each other. We can thus apply the "leave-one-out" or cavity method here over the μ index. We end up with

$$\sum_{\mu\in[n]} \frac{1}{1 + \boldsymbol{z}_\mu^\top G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu} G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu \boldsymbol{z}_\mu^\top + G_{\mathrm{ext}}(\pi B_{\mathrm{ext}} + \tau\lambda I) = I. \tag{12}$$

For fixed task $\boldsymbol{w}_\mu$ we can apply Lemma 3 to compute

$$\mathbb{E}_{X,\boldsymbol{x}_{\mathrm{test}},\boldsymbol{\varepsilon}}[\boldsymbol{z}\boldsymbol{z}^\top] = \frac{1}{d^2}\Upsilon(\boldsymbol{w})$$

for

$$\Upsilon(\boldsymbol{w}) \equiv \begin{bmatrix} \mathrm{tr}[\boldsymbol{w}\boldsymbol{w}^\top] + \rho & \frac{1}{\sqrt{d}}\mathrm{vec}\left(\begin{bmatrix} \boldsymbol{w}\boldsymbol{w}^\top + \frac{1}{\alpha}k_0\,\mathrm{tr}[\boldsymbol{w}\boldsymbol{w}^\top]I_d & \rho_1(\boldsymbol{w})\boldsymbol{w} \end{bmatrix} \right) \\ \frac{1}{\sqrt{d}}\mathrm{vec}\left(\begin{bmatrix} \boldsymbol{w}\boldsymbol{w}^\top + \frac{1}{\alpha}k_0\,\mathrm{tr}[\boldsymbol{w}\boldsymbol{w}^\top]I_d & \rho_1(\boldsymbol{w})\boldsymbol{w} \end{bmatrix} \right)^\top & I_d \otimes E(\boldsymbol{w}) + \frac{1}{d}\mathcal{X}(\boldsymbol{w}) \end{bmatrix}$$

where

$$E(\boldsymbol{w}) \equiv \begin{bmatrix} \boldsymbol{w}\boldsymbol{w}^\top + \rho_2(\boldsymbol{w})I_d/\alpha & \rho_1(\boldsymbol{w})\boldsymbol{w} \\ \rho_1(\boldsymbol{w})\boldsymbol{w}^\top & \rho_1(\boldsymbol{w})^2 \end{bmatrix}, \quad \rho_1(\boldsymbol{w}) = \mathrm{tr}[\boldsymbol{w}\boldsymbol{w}^\top] + \rho, \quad \rho_2(\boldsymbol{w}) = \mathrm{tr}[\boldsymbol{w}\boldsymbol{w}^\top]k_2 + \rho$$

and $d(d+1) \times d(d+1)$ tensor

$$
\begin{aligned}
\mathcal{X}(\boldsymbol{w}) \equiv \; & \frac{1}{\alpha^2}\Big(\operatorname{tr}\big[\boldsymbol{w}\boldsymbol{w}^\top\big]k_1 + \rho k_0\Big)\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)^\top \\
& + \frac{1}{\alpha}k_0\Big(\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)\operatorname{vec}\left(\begin{bmatrix}\boldsymbol{w}\boldsymbol{w}^\top & \mathbf{0}\end{bmatrix}\right)^\top + \operatorname{vec}\left(\begin{bmatrix}\boldsymbol{w}\boldsymbol{w}^\top & \mathbf{0}\end{bmatrix}\right)\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)^\top\Big) \\
& + \frac{1}{\alpha}\rho_1(\boldsymbol{w})k_0\Big(\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)\operatorname{vec}\left(\boldsymbol{w}\boldsymbol{e}_{d+1}^\top\right)^\top + \operatorname{vec}\left(\boldsymbol{w}\boldsymbol{e}_{d+1}^\top\right)\operatorname{vec}\left(\begin{bmatrix}I_d & \mathbf{0}\end{bmatrix}\right)^\top\Big).
\end{aligned}
$$

The quadratic form $\boldsymbol{z}_\mu^\top G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu$ concentrates for fixed tasks $\boldsymbol{w}_\mu$ (assuming weak correlations, so that the condition noted in Appendix D is not violated), so we have

$$
\boldsymbol{z}_\mu^\top G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu \simeq \chi^\mu(\boldsymbol{w}_\mu) \tag{13}
$$

where

$$
\chi^\mu(\boldsymbol{w}_\mu) \equiv \frac{1}{d^2}\operatorname{tr}\left([G_{\mathrm{ext}}^\mu]_{\backslash 0} \cdot \left[I \otimes E(\boldsymbol{w}_\mu) + \frac{1}{d}\mathcal{X}(\boldsymbol{w}_\mu)\right]\right). \tag{14}
$$

Replacing $\boldsymbol{z}_\mu^\top G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu$ in (12) with $\chi^\mu(\boldsymbol{w}_\mu)$ gives

$$
\sum_{\mu \in [n]} \frac{1}{1 + \chi^\mu(\boldsymbol{w}_\mu)} G_{\mathrm{ext}}^{[\mu]} \boldsymbol{z}_\mu \boldsymbol{z}_\mu^\top + G_{\mathrm{ext}}(\pi B_{\mathrm{ext}} + \tau \lambda I) \simeq I. \tag{15}
$$

In this equation we will also replace $\boldsymbol{z}_\mu \boldsymbol{z}_\mu^\top$ with its conditional expectation over $X, \boldsymbol{x}_{\mathrm{test}}, \boldsymbol{\varepsilon}$, giving

$$
\frac{\tau}{n} \sum_{\mu \in [n]} \frac{1}{1 + \chi^\mu(\boldsymbol{w}_\mu)} G_{\mathrm{ext}}^{[\mu]} \Upsilon(\boldsymbol{w}_\mu) + G_{\mathrm{ext}}(\pi B_{\mathrm{ext}} + \tau \lambda I) \simeq I \tag{16}
$$

where recall $\tau = n/d^2$.

In high dimensions and for large n, there is negligible difference between $\sum_{\nu \neq \mu}$ and $\sum_\mu$. Thus, we replace G_{ext}^μ by G_{ext}, and $\chi^\mu(\boldsymbol{w}_\mu)$ by

$$
\chi(\boldsymbol{w}_\mu) \equiv \frac{1}{d^2}\operatorname{tr}\left([G_{\mathrm{ext}}]_{\backslash 0} \cdot \left[I \otimes E(\boldsymbol{w}_\mu) + \frac{1}{d}\mathcal{X}(\boldsymbol{w}_\mu)\right]\right). \tag{17}
$$

So finally we have the expression for G_{ext}

$$
G_{\mathrm{ext}}\left(\frac{\tau}{n} \sum_{\mu \in [n]} \frac{1}{1 + \chi(\boldsymbol{w}_\mu)} \Upsilon(\boldsymbol{w}_\mu) + \pi B_{\mathrm{ext}} + \tau \lambda I\right) \simeq I. \tag{18}
$$

Exploit finiteness of training task set. So far we are summing over n task vectors, but really only n/k of these are unique. Thus, we can simplify (18) as

$$
G_{\mathrm{ext}}\left(\frac{\tau}{k} \sum_{j \in [k]} \frac{1}{1 + \chi(\boldsymbol{w}_j)} \Upsilon(\boldsymbol{w}_j) + \pi B_{\mathrm{ext}} + \tau \lambda I\right) \simeq I. \tag{19}
$$

We replace $\chi(\boldsymbol{w}_j)$, which is self-averaging in $\boldsymbol{w}_j$, with its mean

$$
\widehat{\chi}_{\mathrm{ave}} \equiv \frac{1}{k} \sum_{j \in [k]} \chi(\boldsymbol{w}_j). \tag{20}
$$

To clean up the sums over the tasks $\boldsymbol{w}_1, \ldots \boldsymbol{w}_k$ we use the that

$$\frac{1}{k} \sum_{j \in [k]} \operatorname{tr}\left[\boldsymbol{w}_j \boldsymbol{w}_j^\top\right] = \operatorname{tr}[R_{\mathrm{tr}}]$$

$$\frac{1}{k} \sum_{j \in [k]} \left(\rho + \operatorname{tr}\left[\boldsymbol{w}_j \boldsymbol{w}_j^\top\right]\right) \boldsymbol{w}_j \simeq (\rho + \operatorname{tr}[R_{\mathrm{tr}}]) \, \boldsymbol{b}_{\mathrm{tr}}$$

$$\frac{1}{k} \sum_{j \in [k]} \left(\rho + \operatorname{tr}\left[\boldsymbol{w}_j \boldsymbol{w}_j^\top\right]\right)^2 \simeq (\rho + \operatorname{tr}[R_{\mathrm{tr}}])^2$$

for

$$\boldsymbol{b}_{\mathrm{tr}} \equiv \frac{1}{k} \sum_{j \in [k]} \boldsymbol{w}_j, \qquad R_{\mathrm{tr}} \equiv \frac{1}{k} \sum_{j \in [k]} \boldsymbol{w}_j \boldsymbol{w}_j^\top.$$

We thus have that

$$\frac{1}{k} \sum_{j \in [k]} E(\boldsymbol{w}_j) \approx B_{\mathrm{tr}} \equiv \begin{bmatrix} R_{\mathrm{tr}} + \rho_2 I_d / \alpha & \rho_1 \boldsymbol{b}_{\mathrm{tr}} \\ \rho_1 \boldsymbol{b}_{\mathrm{tr}}^\top & \rho_1^2 \end{bmatrix}$$

$$\frac{1}{k} \sum_{j \in [k]} \mathcal{X}(\boldsymbol{w}_j) \approx \Phi$$

for

$$\Phi \equiv \frac{1}{\alpha^2} (\operatorname{tr}[R_{\mathrm{tr}}] k_1 + \rho k_0) \operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right)^\top$$

$$+ \frac{1}{\alpha} k_0 \left(\operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} & \mathbf{0} \end{bmatrix}\right)^\top + \operatorname{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} & \mathbf{0} \end{bmatrix}\right) \operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right)^\top \right)$$

$$+ \frac{1}{\alpha} \rho_1 k_0 \left(\operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right) \operatorname{vec}\left(\boldsymbol{b}_{\mathrm{tr}} \boldsymbol{e}_{d+1}^\top\right)^\top + \operatorname{vec}\left(\boldsymbol{b}_{\mathrm{tr}} \boldsymbol{e}_{d+1}^\top\right) \operatorname{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right)^\top \right)$$

where

$$\rho_1 = \operatorname{tr}[R_{\mathrm{tr}}] + \rho, \qquad \rho_2 = \operatorname{tr}\left[K^2\right] \operatorname{tr}[R_{\mathrm{tr}}] + \rho.$$

Finally, we have

$$\widehat{\chi}_{\mathrm{ave}} = \frac{1}{d^2} \operatorname{tr}\left([G_{\mathrm{ext}}]_{\backslash 0} \cdot \left[I \otimes B_{\mathrm{tr}} + \frac{1}{d} \Phi \right] \right). \tag{21}$$

Thus, after averaging over $X, \boldsymbol{x}_{\mathrm{test}}, \boldsymbol{\varepsilon}$ in the extended resolvent G_{ext}, we have a deterministic equivalent $G_{\mathrm{ext}} \simeq \mathcal{G}_{\mathrm{ext}}$ (still depending on random task quantities $R_{\mathrm{tr}}, \boldsymbol{b}_{\mathrm{tr}}$) defined by self-consistent equations

$$[\mathcal{G}_{\mathrm{ext}}]_{\backslash 0} = \left(\frac{\tau}{1 + \chi_\pi} I_d \otimes B_{\mathrm{tr}} + \frac{1}{d} \frac{\tau}{1 + \chi_\pi} \Phi + \pi \Pi + \tau \lambda I_d \otimes I_{d+1} \right)^{-1} \tag{22}$$

$$\chi_\pi = \frac{1}{d^2} \operatorname{tr}\left([\mathcal{G}_{\mathrm{ext}}]_{\backslash 0} \left(I_d \otimes B_{\mathrm{tr}} + \frac{1}{d} \Phi \right) \right) \tag{23}$$

where the full matrix is given by

$$\mathcal{G}_{\mathrm{ext}}(\pi)^{-1} \equiv \frac{\tau}{1 + \chi_\pi} \begin{bmatrix} \rho_1 & \frac{1}{\sqrt{d}} \operatorname{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} + \frac{1}{\alpha} k_0 \operatorname{tr}[R_{\mathrm{tr}}] I_d & \rho_1 \boldsymbol{b}_{\mathrm{tr}} \end{bmatrix}\right)^\top \\ \frac{1}{\sqrt{d}} \operatorname{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} + \frac{1}{\alpha} k_0 \operatorname{tr}[R_{\mathrm{tr}}] I_d & \rho_1 \boldsymbol{b}_{\mathrm{tr}} \end{bmatrix}\right) & I_d \otimes B_{\mathrm{tr}} + \frac{1}{d} \Phi \end{bmatrix}$$

$$+ \pi B_{\mathrm{ext}} + \tau \lambda I \tag{24}$$

Note that up until this point, the results have matched the previous analysis from Lu et al. (2025) and Letey et al. (2026) with the exception of ρ_2 (depending on k_2) and the low-rank term Φ (depending on k_0 and k_1).

The low-rank term is given by

$$\Phi = \phi_1 1_+ 1_+^\top + \phi_2(1_+ r^\top + r 1_+^\top) + \phi_3(1_+ \mu^\top + \mu 1_+^\top)$$

$$= \begin{bmatrix} 1_+ & v \end{bmatrix} \begin{bmatrix} \phi_1 & 1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} 1_+^\top \\ v^\top \end{bmatrix}$$

$$= T\phi T^\top$$

for

$$1_+ \equiv \text{vec}\left(\begin{bmatrix} I_d & \mathbf{0} \end{bmatrix}\right), \qquad r \equiv \text{vec}\left(\begin{bmatrix} R_{\text{tr}} & \mathbf{0} \end{bmatrix}\right), \qquad \mu \equiv \text{vec}\left(\boldsymbol{b}_{\text{tr}} e_{d+1}^\top\right), \qquad v \equiv \text{vec}\left(\begin{bmatrix} \phi_2 R_{\text{tr}} & \phi_3 \boldsymbol{b}_{\text{tr}} \end{bmatrix}\right)$$

and

$$\phi_1 \equiv \frac{1}{\alpha^2}(\text{tr}[R_{\text{tr}}]k_1 + \rho k_0), \qquad \phi_2 \equiv \frac{1}{\alpha}k_0, \qquad \phi_3 \equiv \frac{1}{\alpha}(\text{tr}[R_{\text{tr}}] + \rho)k_0.$$

G.1. Effective ridge

The effective ridge will be $\tilde{\lambda} = \lambda(1 + \chi_0)$ where χ_0 is defined as the solution to the implicit equations

$$[\mathcal{G}_{\text{ext}}]_{\backslash 0} = \left(\frac{\tau}{1 + \chi_0} I_d \otimes B_{\text{tr}} + \frac{1}{d}\frac{\tau}{1 + \chi_0}\Phi + \tau\lambda I_d \otimes I_{d+1}\right)^{-1} \tag{25}$$

$$\chi_0 = \frac{1}{d^2} \text{tr}\left([\mathcal{G}_{\text{ext}}]_{\backslash 0}\left(I_d \otimes B_{\text{tr}} + \frac{1}{d}\Phi\right)\right). \tag{26}$$

We can recast the above implicit equations using Woodbury, as

$$\frac{\tau\chi_0}{1 + \chi_0} = \frac{1}{d^2} \text{tr}\left(\left(I_d \otimes F_0 - \frac{1}{d}\Phi_C\right)\left(I_d \otimes B_{\text{tr}} + \frac{1}{d}\Phi\right)\right) \tag{27}$$

where

$$F_0 = (B_{\text{tr}} + \lambda(1 + \chi_0)I_{d+1})^{-1}$$

$$\Phi_C = (I_d \otimes F_0)\begin{bmatrix} 1_+ & v \end{bmatrix} S \begin{bmatrix} 1_+^\top \\ v^\top \end{bmatrix}(I_d \otimes F_0) \qquad \text{for some } 2 \times 2 \text{ mtx } S \text{ with } \Theta(1) \text{ elements.}$$

However, all the $1/d$ terms in (27) are subleading, e.g.

$$\frac{1}{d^3} \text{tr}\left((I_d \otimes F_0)rr^\top(I_d \otimes F_0 B_{\text{tr}})\right) = \frac{1}{d^3} \text{tr}(R_{\text{tr}} F_0 B_{\text{tr}} F_0 R_{\text{tr}}) = \mathcal{O}\left(\frac{1}{d^2}\right)$$

and so we write

$$\frac{\tau\chi_0}{1 + \chi_0} \approx \frac{1}{d^2} \text{tr}(I_d \otimes F_0 B_{\text{tr}}) = \text{tr}\left[(B_{\text{tr}} + \tilde{\lambda}I_{d+1})^{-1}B_{\text{tr}}\right]. \tag{28}$$

This can be simplified as

$$\frac{\tau\chi_0}{1 + \chi_0} = 1 - \lambda(1 + \chi_0)\text{tr}\left[(B_{\text{tr}} + \lambda(1 + \chi_0)I_{d+1})^{-1}\right]$$

$$\approx 1 - \lambda(1 + \chi_0)\text{tr}\left[\left(R_{\text{tr}} + \left(\lambda(1 + \chi_0) + \frac{\rho_2}{\alpha}\right)I_d\right)^{-1}\right]$$

after ignoring the $\boldsymbol{b}_{\text{tr}}$ terms in B_{tr}. This is where the Stieltjes transform of R_{tr} is first introduced:

$$\mathcal{M}_\kappa(w) = \lim_{d\to\infty, k\to\infty, k/d=\kappa} \frac{1}{d} \text{tr}\left((R_{\text{tr}} + wI_d)^{-1}\right).$$

Using this, we find the effective ridge self-consistency equation

$$\tilde{\lambda}\mathcal{M}_\kappa\left(\tilde{\lambda} + \frac{\rho_2}{\alpha}\right) + \frac{\lambda\tau}{\tilde{\lambda}} = 1 - \tau \tag{29}$$

G.2. Relating Γ^* to G

Recall that

$$\mathrm{vec}(\Gamma^*) = \left(\frac{n}{d}\lambda I + \sum_{\mu=1}^n \mathrm{vec}(H^\mu)\mathrm{vec}(H^\mu)^\top\right)^{-1}\sum_{\mu=1}^n y_{\mathrm{test}}^\mu \mathrm{vec}(H^\mu)$$

and so we have

$$G_{\mathrm{ext}}(0) = d\left[\begin{matrix} \frac{1}{d}\sum_\mu (y_{\mathrm{test}}^\mu)^2 + \frac{n}{d}\lambda & \frac{1}{\sqrt{d}}\sum_\mu y_{\mathrm{test}}^\mu \mathrm{vec}(H^\mu)^\top \\ \frac{1}{\sqrt{d}}\sum_\mu y_{\mathrm{test}}^\mu \mathrm{vec}(H^\mu) & \sum_\mu \mathrm{vec}(H^\mu)\mathrm{vec}(H^\mu)^\top + \frac{n}{d}\lambda I \end{matrix}\right]^{-1}$$

Using that

$$\begin{bmatrix} a & \boldsymbol{b}^\top \\ \boldsymbol{b} & D \end{bmatrix}^{-1} = \begin{bmatrix} c & -c\boldsymbol{q}^\top \\ -c\boldsymbol{q} & D^{-1} + c\boldsymbol{q}\boldsymbol{q}^\top \end{bmatrix}, \qquad c = \frac{1}{a - \boldsymbol{b}^\top D^{-1}\boldsymbol{b}}, \qquad \boldsymbol{q} = D^{-1}\boldsymbol{b}$$

we have a formula for Γ^* in terms of $G_{\mathrm{ext}}(0)$ as

$$\frac{\sqrt{d}}{\boldsymbol{e}_1^\top G_{\mathrm{ext}}(0)\boldsymbol{e}_1}G_{\mathrm{ext}}(0)\boldsymbol{e}_1 = \begin{bmatrix} \sqrt{d} \\ \mathrm{vec}(\Gamma)^* \end{bmatrix}$$

Using our above formulas, we also have

$$\frac{\sqrt{d}}{\boldsymbol{e}_1^\top \mathcal{G}_{\mathrm{ext}}(0)\boldsymbol{e}_1}\mathcal{G}_{\mathrm{ext}}(0)\boldsymbol{e}_1 = \left[\left(I_d \otimes B_{\mathrm{tr}} + \frac{1}{d}\Phi + \tilde{\lambda}I_{d(d+1)}\right)^{-1}\mathrm{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} + \frac{k_0}{\alpha}\mathrm{tr}[R_{\mathrm{tr}}]I_d & \rho_1\boldsymbol{b}_{\mathrm{tr}} \end{bmatrix}\right) \right]$$

and so

$$\mathrm{vec}(\Gamma^*) \simeq \mathrm{vec}(\Gamma_{\mathrm{de}}) \equiv \left(I_d \otimes B_{\mathrm{tr}} + \frac{1}{d}\Phi + \tilde{\lambda}I_{d(d+1)}\right)^{-1}\mathrm{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} + \frac{k_0}{\alpha}\mathrm{tr}[R_{\mathrm{tr}}]I_d & \rho_1\boldsymbol{b}_{\mathrm{tr}} \end{bmatrix}\right) = F_\Phi\boldsymbol{g}$$

for

$$F_\Phi = \left(I_d \otimes B_{\mathrm{tr}} + \frac{1}{d}\Phi + \tilde{\lambda}I_{d(d+1)}\right)^{-1}$$

$$\boldsymbol{g} = \mathrm{vec}\left(\begin{bmatrix} R_{\mathrm{tr}} + \frac{k_0}{\alpha}\mathrm{tr}[R_{\mathrm{tr}}]I_d & \rho_1\boldsymbol{b}_{\mathrm{tr}} \end{bmatrix}\right).$$

This immediately gives the linear term in the MSE error expression as

$$\mathrm{tr}\left[\Gamma^*A^\top\right] = \frac{1}{d}\mathrm{tr}\left(\Gamma^*A^\top\right) = \frac{1}{d}\mathrm{tr}\left(\mathrm{vec}(\Gamma^*)\,\mathrm{vec}(A)^\top\right) \simeq \frac{1}{d}\mathrm{tr}\left(\mathrm{vec}(\Gamma_{\mathrm{de}})\,\mathrm{vec}(A)^\top\right)$$

where we will specifically use $\mathrm{vec}(A) = (1 + \phi_2)1_+$ from Lemma 1.

For the quadratic terms we need to be a bit more careful, as the correct object to work with for high-dimensional equivalence is technically the components of $\mathcal{G}$ and not this linear representation of $\mathrm{vec}(\Gamma_{\mathrm{de}})$. This is what the parameterisation πB_{ext} is for. Take

$$B_{\mathrm{ext}} = \begin{bmatrix} 0 & 0 \\ 0 & \Pi \end{bmatrix}$$

for some $\Pi \in \mathbb{R}^{d(d+1)\times d(d+1)}$. Then

$$\frac{\mathrm{d}}{\mathrm{d}\pi}\frac{1}{c(\pi)}(\pi = 0) = \frac{1}{d}\mathrm{vec}(\Gamma^*)^\top \Pi\,\mathrm{vec}(\Gamma^*)$$

where $c(\pi) = \boldsymbol{e}_1^\top G_{\text{ext}}(\pi)\boldsymbol{e}_1$. We can safely replace
$$c(\pi) = \boldsymbol{e}_1^\top G_{\text{ext}}(\pi)\boldsymbol{e}_1 \simeq \boldsymbol{e}_1 \mathcal{G}_{\text{ext}}(\pi)\boldsymbol{e}_1.$$

By Schur complement on $\mathcal{G}_{\text{ext}}(\pi)$ we have

$$\frac{1}{\boldsymbol{e}_1\mathcal{G}_{\text{ext}}(\pi)\boldsymbol{e}_1} = \frac{\tau}{1+\chi_\pi}\rho_1 + \tau\lambda - \frac{1}{d}\frac{\tau^2}{(1+\chi_\pi)^2}\boldsymbol{g}^\top\left(\frac{\tau}{1+\chi_\pi}\left(I_d \otimes B_{\text{tr}} + \frac{1}{d}\Phi\right) + \pi\Pi + \tau\lambda I\right)^{-1}\boldsymbol{g}$$

$$= \frac{\tau}{1+\chi_\pi}\rho_1 + \tau\lambda - \frac{1}{d}\frac{\tau}{1+\chi_\pi}\boldsymbol{g}^\top\left(I_d \otimes B_{\text{tr}} + \frac{1}{d}\Phi + \pi\frac{1+\chi_\pi}{\tau}\Pi + \lambda(1+\chi_\pi)I\right)^{-1}\boldsymbol{g}.$$

Given the eventual MSE term we want from Lemma 1, we choose
$$\Pi = I_d \otimes B + \frac{1}{d}\Psi$$

for

$$\Psi = \begin{bmatrix} 1_+ & \text{vec}\left(\begin{bmatrix} I_d & \boldsymbol{0} \end{bmatrix}\right) \end{bmatrix} \begin{bmatrix} \phi_1 & \phi_2 \\ \phi_2 & 0 \end{bmatrix} \begin{bmatrix} 1_+^\top \\ \text{vec}\left(\begin{bmatrix} I_d & \boldsymbol{0} \end{bmatrix}\right)^\top \end{bmatrix} = (\phi_1 + 2\phi_2)1_+ 1_+^\top.$$

Using the same approximation as before, we will take
$$[\mathcal{G}_{\text{ext}}]_{\backslash 0} = I_d \otimes \left(\frac{\tau}{1+\chi_\pi}B_{\text{tr}} + \pi B + \tau\lambda I\right)^{-1} + \frac{1}{d} \text{ low rank terms from } \Phi \text{ and } \Psi$$

and approximate χ_π as

$$\chi_\pi = \text{tr}\left[\left(\frac{\tau}{1+\chi_\pi}B_{\text{tr}} + \pi B + \tau\lambda I\right)^{-1}B_{\text{tr}}\right].$$

As before, we find that
$$\frac{\tau\chi_0'}{(1+\chi_0)^2} = \frac{\text{tr}[F_0 B F_0 B_{\text{tr}}]}{\text{tr}[F_0 B_{\text{tr}} F_0 B_{\text{tr}}] - \tau}, \qquad F_0 = (B_{\text{tr}} + \tilde{\lambda}I)^{-1}.$$

Thus,
$$\frac{1}{d}\text{vec}(\Gamma^*)^\top \Pi \text{vec}(\Gamma^*) \simeq \frac{1}{d}\text{vec}(\Gamma_{\text{de}})^\top \Pi \text{vec}(\Gamma_{\text{de}}) - \frac{\tau\chi_0'}{(1+\chi_0)^2}\left(\rho_1 - \frac{1}{d}\boldsymbol{g}^\top \text{vec}(\Gamma_{\text{de}}) - \tilde{\lambda}\frac{1}{d}\text{vec}(\Gamma_{\text{de}})^\top \text{vec}(\Gamma_{\text{de}})\right).$$

Note that
$$\text{vec}(\Gamma_{\text{de}}) = (I_d \otimes F_0)\boldsymbol{g} - \frac{1}{d}\Phi_C \boldsymbol{g}$$

$$\Pi = I_d \otimes B_{\text{test}} + \frac{1}{d}\Psi$$

where
$$\Phi_C = ((I_d \otimes F_0)T)\begin{bmatrix} \mathcal{M} & 1 + \phi_2(1 - \sigma\mathcal{M}) \\ 1 + \phi_2(1 - \sigma\mathcal{M}) & -\phi_1 + \phi_2^2(1 - \sigma + \sigma^2\mathcal{M}) \end{bmatrix}^{-1}((I_d \otimes F_0)T)^\top$$

$$= (I_d \otimes F_0)\begin{bmatrix} 1_+ & \phi_2 r + \phi_3\mu \end{bmatrix}S\begin{bmatrix} 1_+^\top \\ \phi_2 r^\top + \phi_3\mu^\top \end{bmatrix}(I_d \otimes F_0)$$

This comes from expanding

$$F_0 = \begin{bmatrix} F & \boldsymbol{f} \\ \boldsymbol{f}^\top & f \end{bmatrix}$$

$$\frac{1}{d} \begin{bmatrix} 1_+^\top \\ r^\top \\ \mu^\top \end{bmatrix} \begin{bmatrix} 1_+ & r & \mu \end{bmatrix} = \begin{bmatrix} \mathrm{tr}[F] & \mathrm{tr}[FR_{\mathrm{tr}}] & \mathrm{tr}[\boldsymbol{fb}_{\mathrm{tr}}^\top] \\ \mathrm{tr}[FR_{\mathrm{tr}}] & \mathrm{tr}[FR_{\mathrm{tr}}^2] & \mathrm{tr}[F\boldsymbol{fb}_{\mathrm{tr}}^\top] \\ \mathrm{tr}[\boldsymbol{fb}_{\mathrm{tr}}^\top] & \mathrm{tr}[F\boldsymbol{fb}_{\mathrm{tr}}^\top] & f\,\mathrm{tr}[\boldsymbol{b}_{\mathrm{tr}}\boldsymbol{b}_{\mathrm{tr}}^\top] \end{bmatrix}$$

$$\simeq \begin{bmatrix} \mathcal{M}_\kappa(\sigma) & 1 - \sigma\mathcal{M}_\kappa(\sigma) & 0 \\ 1 - \sigma\mathcal{M}_\kappa(\sigma) & 1 - \sigma + \sigma^2\mathcal{M}_\kappa(\sigma) & 0 \\ 0 & 0 & 0 \end{bmatrix}, \qquad \sigma \equiv \frac{\rho_2}{\alpha} + \tilde{\lambda}.$$

As we can see, all the $\boldsymbol{b}_{\mathrm{tr}}, \boldsymbol{f}$ contributions are negligible compared to the resolvent $(R_{\mathrm{tr}} + \sigma I_d)^{-1}$ contributions. Using similar intuition, in the expansions of the linear and quadratic error terms, we will neglect the $\rho_1\boldsymbol{b}_{\mathrm{tr}}$ component of $\boldsymbol{g}$, as well as $\mu = \mathrm{vec}(\boldsymbol{b}_{\mathrm{tr}}\boldsymbol{e}_{d+1}^\top)$ in Φ_C.

Linear error term Following this, and writing the matrix component of $\boldsymbol{g}$ as

$$\tilde{R} = R_{\mathrm{tr}} + \frac{k_0}{\alpha}\,\mathrm{tr}[R_{\mathrm{tr}}]I_d$$

we have

$$\frac{1}{d}\,\mathrm{vec}(A)^\top\,\mathrm{vec}(\Gamma_{\mathrm{de}}) = (1 + \phi_2)\frac{1}{d}\left(1_+^\top(I \otimes F)\boldsymbol{g} - \frac{1}{d}1_+^\top\Phi_C\boldsymbol{g}\right)$$

$$\simeq (1 + \phi_2)\left(\mathrm{tr}[F\tilde{R}] - \begin{bmatrix} \mathrm{tr}[F] & \phi_2\,\mathrm{tr}[FR_{\mathrm{tr}}] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F\tilde{R}] \end{bmatrix}\right)$$

Quadratic error term Recall

$$B = \begin{bmatrix} (1 + \frac{\rho_2}{\alpha})I_d & \boldsymbol{0} \\ \boldsymbol{0}^\top & \rho_1^2 \end{bmatrix}, \qquad \Psi = (\phi_1 + 2\phi_2)1_+1_+^\top.$$

Have

$$\frac{1}{d}\boldsymbol{g}^\top\,\mathrm{vec}(\Gamma_{\mathrm{de}}) \approx \mathrm{tr}[\tilde{R}F\tilde{R}] - \begin{bmatrix} \mathrm{tr}[\tilde{R}F] & \phi_2\,\mathrm{tr}[\tilde{R}FR_{\mathrm{tr}}] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F\tilde{R}] \end{bmatrix}$$

$$\frac{1}{d}\,\mathrm{vec}(\Gamma_{\mathrm{de}})^\top\,\mathrm{vec}(\Gamma_{\mathrm{de}}) \approx \mathrm{tr}[\tilde{R}F^2\tilde{R}] - 2\begin{bmatrix} \mathrm{tr}[\tilde{R}F] & \phi_2\,\mathrm{tr}[\tilde{R}FR_{\mathrm{tr}}] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F^2\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F^2\tilde{R}] \end{bmatrix}$$

$$+ \begin{bmatrix} \mathrm{tr}[\tilde{R}F] & \phi_2\,\mathrm{tr}[\tilde{R}FR_{\mathrm{tr}}] \end{bmatrix} S^\top \begin{bmatrix} \mathrm{tr}[F^2] & \phi_2\,\mathrm{tr}[F^2R_{\mathrm{tr}}] \\ \phi_2\,\mathrm{tr}[F^2R_{\mathrm{tr}}] & \phi_2^2\,\mathrm{tr}[R_{\mathrm{tr}}F^2R_{\mathrm{tr}}] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F\tilde{R}] \end{bmatrix}$$

$$\frac{1}{d}\,\mathrm{vec}(\Gamma_{\mathrm{de}})^\top(I_d \otimes B)\,\mathrm{vec}(\Gamma_{\mathrm{de}}) \approx \left(1 + \frac{\rho_2}{\alpha}\right) \cdot \frac{1}{d}\,\mathrm{vec}(\Gamma_{\mathrm{de}})^\top\,\mathrm{vec}(\Gamma_{\mathrm{de}})$$

$$\frac{1}{\phi_1 + 2\phi_2}\frac{1}{d^2}\,\mathrm{vec}(\Gamma_{\mathrm{de}})^\top\Psi\,\mathrm{vec}(\Gamma_{\mathrm{de}}) \approx \mathrm{tr}[\tilde{R}F]^2 - 2\,\mathrm{tr}[\tilde{R}F]\begin{bmatrix} \mathrm{tr}[F] & \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F\tilde{R}] \end{bmatrix}$$

$$+ \left(\begin{bmatrix} \mathrm{tr}[F] & \phi_2\,\mathrm{tr}[FR_{\mathrm{tr}}] \end{bmatrix} S \begin{bmatrix} \mathrm{tr}[F\tilde{R}] \\ \phi_2\,\mathrm{tr}[R_{\mathrm{tr}}F\tilde{R}] \end{bmatrix}\right)^2.$$

The final step is to remember that $F = (R_{\mathrm{tr}} + \sigma)^{-1}$ and $\tilde{R} = R_{\mathrm{tr}} + k_0 I_d/\alpha$ and so we have the following dictionary of terms

$$\mathrm{tr}[F] \simeq \mathcal{M}_\kappa(\sigma)$$

$$\mathrm{tr}[F R_{\mathrm{tr}}] \simeq 1 - \sigma \mathcal{M}_\kappa(\sigma)$$

$$\mathrm{tr}\left[F \tilde{R}\right] \simeq 1 - \sigma \mathcal{M}_\kappa(\sigma) + \frac{k_0}{\alpha} \mathcal{M}_\kappa(\sigma)$$

$$\mathrm{tr}\left[F R_{\mathrm{tr}} \tilde{R}\right] \simeq \frac{k_0}{\alpha}(1 - \sigma \mathcal{M}_\kappa(\sigma)) + 1 - \sigma + \sigma^2 \mathcal{M}_\kappa(\sigma)$$

$$\mathrm{tr}\left[F \tilde{R} \tilde{R}\right] \simeq \left(\frac{k_0}{\alpha}\right)^2 \mathcal{M}_\kappa(\sigma) + 2\frac{k_0}{\alpha}(1 - \sigma \mathcal{M}_\kappa(\sigma))) + 1 - \sigma + \sigma^2 \mathcal{M}_\kappa(\sigma)$$

$$\mathrm{tr}\left[F^2\right] \simeq -\mathcal{M}'_\kappa(\sigma)$$

$$\mathrm{tr}\left[F^2 R_{\mathrm{tr}}\right] \simeq \mathcal{M}_\kappa(\sigma) + \sigma \mathcal{M}'_\kappa(\sigma)$$

$$\mathrm{tr}\left[F^2 \tilde{R}\right] \simeq \mathcal{M}_\kappa(\sigma) + \sigma \mathcal{M}'_\kappa(\sigma) - \frac{k_0}{\alpha} \mathcal{M}'_\kappa(\sigma)$$

$$\mathrm{tr}\left[F^2 R_{\mathrm{tr}} R_{\mathrm{tr}}\right] \simeq 1 - 2\sigma \mathcal{M}_\kappa(\sigma) - \sigma^2 \mathcal{M}'_\kappa(\sigma)$$

$$\mathrm{tr}\left[F^2 R_{\mathrm{tr}} \tilde{R}\right] \simeq 1 - 2\sigma \mathcal{M}_\kappa(\sigma) - \sigma^2 \mathcal{M}'_\kappa(\sigma) + \frac{k_0}{\alpha}(\mathcal{M}_\kappa(\sigma) + \sigma \mathcal{M}'_\kappa(\sigma))$$

$$\mathrm{tr}\left[F^2 \tilde{R} \tilde{R}\right] \simeq 1 - 2\sigma \mathcal{M}_\kappa(\sigma) - \sigma^2 \mathcal{M}'_\kappa(\sigma) + 2\frac{k_0}{\alpha}(\mathcal{M}_\kappa(\sigma) + \sigma \mathcal{M}'_\kappa(\sigma)) - \left(\frac{k_0}{\alpha}\right)^2 \mathcal{M}'_\kappa(\sigma)$$

So finally, if we use some shorthand

$$\tilde{\sigma} = \sigma - \frac{k_0}{\alpha}, \quad \boldsymbol{m}_1 = \begin{bmatrix} \mathcal{M} \\ \phi_2(1 - \sigma \mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_2 = \begin{bmatrix} 1 - \tilde{\sigma} \mathcal{M} \\ \phi_2(1 - \tilde{\sigma} + \sigma\tilde{\sigma}\mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_3 = \begin{bmatrix} \mathcal{M} + \tilde{\sigma}\mathcal{M}' \\ \phi_2(1 - \sigma \mathcal{M} - \tilde{\sigma}\mathcal{M} - \sigma\tilde{\sigma}\mathcal{M}') \end{bmatrix}$$

$$M = \begin{bmatrix} -\mathcal{M}' & \phi_2(\mathcal{M} + \sigma \mathcal{M}') \\ \phi_2(\mathcal{M} + \sigma \mathcal{M}') & \phi_2^2(1 - 2\sigma \mathcal{M} - \sigma^2 \mathcal{M}') \end{bmatrix}$$

and so

$$\frac{1}{d} \mathrm{vec}(A)^\top \mathrm{vec}(\Gamma_{\mathrm{de}}) \simeq (1 + \phi_2)\left(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1^\top S \boldsymbol{m}_2\right)$$

$$\frac{1}{d}\boldsymbol{g}^\top \mathrm{vec}(\Gamma_{\mathrm{de}}) \simeq 1 + \sigma - 2\tilde{\sigma} + \tilde{\sigma}^2 \mathcal{M} - \boldsymbol{m}_2^\top S \boldsymbol{m}_2$$

$$\frac{1}{d} \mathrm{vec}(\Gamma_{\mathrm{de}})^\top \mathrm{vec}(\Gamma_{\mathrm{de}}) \simeq 1 - 2\tilde{\sigma}\mathcal{M} - \tilde{\sigma}^2 \mathcal{M}' - 2\boldsymbol{m}_2^\top S \boldsymbol{m}_3 + \boldsymbol{m}_2^\top S^\top M S \boldsymbol{m}_2$$

$$\frac{1}{d^2} \mathrm{vec}(\Gamma_{\mathrm{de}})^\top \Psi \mathrm{vec}(\Gamma_{\mathrm{de}}) \simeq (\phi_1 + 2\phi_2)\left((1 - \tilde{\sigma}\mathcal{M}) - \boldsymbol{m}_1^\top S \boldsymbol{m}_2\right)^2$$

with

$$c = \frac{\tau \chi_0'}{(1 + \chi_0)^2} \simeq (1 + \frac{\rho_2}{\alpha})\frac{\mathcal{M} + \tilde{\lambda}\mathcal{M}'}{1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2 \mathcal{M}' - \tau}.$$

So finally, we have

$$e_{\mathrm{ICL}} = (1 - c)\rho_1 - 2(1 + \phi_2)(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1^\top S \boldsymbol{m}_2) + (\phi_1 + 2\phi_2)(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1 S \boldsymbol{m}_2)^2$$

$$+ \left(1 + \frac{\rho_2}{\alpha} + c\tilde{\lambda}\right)(1 - 2\tilde{\sigma}\mathcal{M} - \tilde{\sigma}^2 \mathcal{M}' - 2\boldsymbol{m}_2^\top S \boldsymbol{m}_3 + \boldsymbol{m}_2^\top S^\top M S \boldsymbol{m}_2)$$

$$+ c(1 + \sigma - 2\tilde{\sigma} + \tilde{\sigma}^2 \mathcal{M} - \boldsymbol{m}_2^\top S \boldsymbol{m}_2)$$

for

$$\rho_1 \equiv 1 + \rho, \qquad \rho_2 \equiv \mathrm{tr}[K^2] + \rho$$

$$\sigma = \tilde{\lambda} + \frac{\rho_2}{\alpha}, \quad \tilde{\sigma} = \sigma - \frac{k_0}{\alpha}, \quad \tilde{\lambda}\mathcal{M}_\kappa\left(\tilde{\lambda} + \frac{\rho_2}{\alpha}\right) + \frac{\lambda\tau}{\tilde{\lambda}} = 1 - \tau, \quad c = (1 + \frac{\rho_2}{\alpha})\frac{\mathcal{M} + \tilde{\lambda}\mathcal{M}'}{1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2\mathcal{M}' - \tau}$$

$$\boldsymbol{m}_1 = \begin{bmatrix} \mathcal{M} \\ \phi_2(1 - \sigma\mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_2 = \begin{bmatrix} 1 - \tilde{\sigma}\mathcal{M} \\ \phi_2(1 - \tilde{\sigma} + \sigma\tilde{\sigma}\mathcal{M}) \end{bmatrix}, \quad \boldsymbol{m}_3 = \begin{bmatrix} \mathcal{M} + \tilde{\sigma}\mathcal{M}' \\ \phi_2(1 - \sigma\mathcal{M} - \tilde{\sigma}\mathcal{M} - \sigma\tilde{\sigma}\mathcal{M}') \end{bmatrix}$$

$$M = \begin{bmatrix} -\mathcal{M}' & \phi_2(\mathcal{M} + \sigma\mathcal{M}') \\ \phi_2(\mathcal{M} + \sigma\mathcal{M}') & \phi_2^2(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}') \end{bmatrix}, \quad S = \begin{bmatrix} \mathcal{M} & 1 + \phi_2(1 - \sigma\mathcal{M}) \\ 1 + \phi_2(1 - \sigma\mathcal{M}) & -\phi_1 + \phi_2^2(1 - \sigma + \sigma^2\mathcal{M}) \end{bmatrix}^{-1}$$

$$\phi_1 \equiv \frac{1}{\alpha^2}(k_1 + \rho k_0) = \phi_1, \quad \phi_2 \equiv \frac{1}{\alpha}k_0 = \phi_2$$

Sanity check For $K = I_d$ have $\rho_1 = \rho_2$, $\sigma = \tilde{\sigma}$, $\phi_1 = \phi_2 = \phi_1 = \phi_2 = 0$, and all $\boldsymbol{m}, S, M$ terms don't contribute. Get

$$e_{\mathrm{ICL}} = 1 + \rho - 2(1 - \sigma\mathcal{M}) + \left(1 + \frac{1 + \rho}{\alpha}\right)(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}')$$

$$- c(\rho + \sigma - \sigma^2\mathcal{M} - \tilde{\lambda}(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}'))$$

$$= 1 + \rho - 2(1 - \sigma\mathcal{M}) + \left(1 + \frac{1 + \rho}{\alpha}\right)(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}') + \left(1 + \frac{1 + \rho}{\alpha}\right)(\mathcal{M} + \tilde{\lambda}\mathcal{M}')q$$

$$= \rho + q\mathcal{M} + (\tilde{\lambda}q - \sigma^2)\mathcal{M}' + \frac{1 + \rho}{\alpha}(1 - (q - 2\sigma)\mathcal{M} + (\tilde{\lambda}q - \sigma^2)\mathcal{M}')$$

as previously in Letey et al. (2026).

G.3. Dependence on query terms

This formula can be cleaned up a fair bit by gathering terms that depend on k_0, k_1 *i.e.*, the query correlation terms. First we write all the terms depending on c as

$$(-c)\Big(\rho_1 - \tilde{\lambda}\big(1 - 2\tilde{\sigma}\mathcal{M} - \tilde{\sigma}^2\mathcal{M}' - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + \boldsymbol{m}_2^\top S^\top M S\boldsymbol{m}_2\big) - \big(1 + \sigma - 2\tilde{\sigma} + \tilde{\sigma}^2\mathcal{M} - \boldsymbol{m}_2^\top S\boldsymbol{m}_2\big)\Big)$$

$$= \left(1 + \frac{\rho_2}{\alpha}\right)\frac{\mathcal{M} + \tilde{\lambda}\mathcal{M}'}{\tau - (1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2\mathcal{M}')}\big(\rho + \sigma - \sigma^2\mathcal{M} - \tilde{\lambda}(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}')\big)$$

$$- \left(1 + \frac{\rho_2}{\alpha}\right)\frac{\mathcal{M} + \tilde{\lambda}\mathcal{M}'}{\tau - (1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2\mathcal{M}')}\left(2\frac{k_0}{\alpha}\big(\tilde{\lambda}(\mathcal{M} + \sigma\mathcal{M}') + (1 - 2\sigma\mathcal{M})\big) + \frac{k_0^2}{\alpha^2}\big(\mathcal{M} - \tilde{\lambda}\mathcal{M}'\big)\right)$$

$$+ \left(1 + \frac{\rho_2}{\alpha}\right)\frac{\mathcal{M} + \tilde{\lambda}\mathcal{M}'}{\tau - (1 - 2\tilde{\lambda}\mathcal{M} - \tilde{\lambda}^2\mathcal{M}')}\big(\boldsymbol{m}_2^\top S\boldsymbol{m}_2 - \tilde{\lambda}\big(-2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + \boldsymbol{m}_2^\top S^\top M S\boldsymbol{m}_2\big)\big)$$

$$= \left(1 + \frac{\rho_2}{\alpha}\right)(\mathcal{M} + \tilde{\lambda}\mathcal{M}')(q_{\mathrm{old}} + q_{\mathrm{query}})$$

and the remaining ICL error terms can be expressed as

$$\rho_1 - 2(1 - \sigma\mathcal{M}) + \left(1 + \frac{\rho_2}{\alpha}\right)(1 - 2\sigma\mathcal{M} - \sigma^2\mathcal{M}')$$

$$- \frac{k_0}{\alpha}\left(2\mathcal{M} - 2\left(1 + \frac{\rho_2}{\alpha}\right)(\mathcal{M} + \sigma\mathcal{M}')\right) - \frac{k_0^2}{\alpha^2}\left(1 + \frac{\rho_2}{\alpha}\right)\mathcal{M}'$$

$$- 2\phi_2(1 - \tilde{\sigma}\mathcal{M}) + 2(1 + \phi_2)\boldsymbol{m}_1^\top S\boldsymbol{m}_2$$

$$+ \left(1 + \frac{\rho_2}{\alpha}\right)(\boldsymbol{m}_2^\top S^\top M S\boldsymbol{m}_2 - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3)$$

$$+ (\phi_1 + 2\phi_2)(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1^\top S\boldsymbol{m_2})^2.$$

Thus we can write

$$
\begin{aligned}
e_{\mathrm{ICL}}(k_0, k_1) = {}& e_{\mathrm{ICL}}(\text{independent query}) - 2\phi_2 + q_{\mathrm{query}}(\mathcal{M} + \tilde{\lambda}\mathcal{M}') + \phi_2(2\sigma - \phi_2)\mathcal{M}' \\
& + \; \boldsymbol{m}_2^\top S^\top M\, S\boldsymbol{m}_2 - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + 2\phi_2\tilde{\sigma}\mathcal{M} + 2(1+\phi_2)\boldsymbol{m}_1^\top S\boldsymbol{m}_2 \\
& + \; (\phi_1 + 2\phi_2)\left(1 - \tilde{\sigma}\mathcal{M} - \boldsymbol{m}_1^\top S\boldsymbol{m}_2\right)^2 \\
& + \frac{\rho_2}{\alpha}\left[q_{\mathrm{query}}(\mathcal{M} + \tilde{\lambda}\mathcal{M}') + \phi_2(2\sigma - \phi_2)\mathcal{M}' + \boldsymbol{m}_2^\top S^\top M\, S\boldsymbol{m}_2 - 2\boldsymbol{m}_2^\top S\boldsymbol{m}_3 + 2\phi_2\mathcal{M}\right].
\end{aligned}
$$