Score-Based Generative Models Detect Manifolds

Abstract

Score-based generative models (SGMs) need to approximate the scores $\nabla \log p_t$ of the intermediate distributions as well as the final distribution p_T of the forward process. The theoretical underpinnings of the effects of these approximations are still lacking. We find precise conditions under which SGMs are able to produce samples from an underlying (low-dimensional) *data manifold* $\mathcal{M}$. This assures us that SGMs are able to generate the "right kind of samples". For example, taking $\mathcal{M}$ to be the subset of images of faces, we find conditions under which the SGM robustly produces an image of a face, even though the relative frequencies of these images might not accurately represent the true data generating distribution. Moreover, this analysis is a first step towards understanding the generalization properties of SGMs: Taking $\mathcal{M}$ to be the set of all training samples, our results provide a precise description of when the SGM memorizes its training data.

1 Introduction

Score-based generative models, also called diffusion models ([Sohl-Dickstein et al., 2015, Song and Ermon, 2019, Song et al., 2021b, Vahdat et al., 2021]) and the related models ([Bordes et al., 2017, Ho et al., 2020, Kingma et al., 2021]) have shown great empirical success in many areas, such as image generation ([Jolicoeur-Martineau et al., 2021, Nichol and Dhariwal, 2021, Dhariwal and Nichol, 2021, Ho et al., 2022]), audio generation ([Chen et al., 2021, Kong et al., 2021, Jeong et al., 2021, Popov et al., 2021]) as well as in other applications ([Batzolis et al., 2021, De Bortoli et al., 2021, Zhou et al., 2021, Cai et al., 2020, Luo and Hu, 2021, Meng et al., 2021, Saharia et al., 2021, Li et al., 2022, Sasaki et al., 2021]). Recently some progress has been made to bridge the gap between the different approaches ([Song et al., 2021b, Huang et al., 2021]) through the framework of SDEs and reverse SDEs.

In generative modelling one is given samples $\{x^i\}_{i=1}^N$ from a measure μ_{data}. The task is to learn a measure μ_{sample} which approximates μ_{data}. The performance of a generative model can then be measured by the distance from μ_{sample} to μ_{data}. In practice however, the true data generating distribution μ_{data} is unknown. All that is known are the samples $\{x_i\}_{i=1}^n$, which can be used to define the empirical measure $\hat{\mu}_{\text{data}}$,

$$\hat{\mu}_{\text{data}} := \text{Unif}\{x^1, x^2, \ldots, x^n\}.$$

Any sample from the empirical measure $\hat{\mu}_{\text{data}}$ will be equal to a training example. Hence, while μ_{sample} being close to μ_{data} is the final goal, μ_{sample} being close to $\hat{\mu}_{\text{data}}$ implies that the generative model has memorized the training data. To summarize, a good generative model will output a measure μ_{sample} which is as close to μ_{data} as possible, while keeping some distance from $\hat{\mu}_{\text{data}}$, even though it only knows μ_{data} through $\hat{\mu}_{\text{data}}$.

Given a target measure π_0, a score-based generative model (SGM) employs two stochastic differential equations (SDEs). The first one is called the *forward SDE*

$$
\begin{array}{rcl}
dX_t & = & \beta(X_t)dt + \sigma dW_t, \\
X_0 & \sim & \pi_0.
\end{array}
\tag{1}
$$

Figure 1: *Top left*: The leftmost plot shows the true data distribution μ_{data} which is a Gaussian mixture. The heat maps show the intermediate densities p_t of X_t, followed by line plots of p_t for $t = 1$. *Bottom left*: The rightmost plot shows μ_{prior}, which is a standard Gaussian and differs from p_1. We start the reverse SDE (2) in μ_{prior}. But instead of using the real score, we introduce an approximation error and use $s(x, t) = \nabla \log p_{1-t}(x) + 3$ with a constant error of 3. Again, the heat maps show how the densities q_t of Y_t evolve backwards in time. The leftmost plot shows the resulting distribution q_1 which is used as sample distribution, $\mu_{\text{sample}} = q_1$. *Right:* The densities μ_{data} and μ_{sample} are shown for direct comparison. We see that the approximation errors in μ_{prior} and the drift lead to an incorrect sample distribution $\mu_{\text{sample}} \neq \mu_{\text{data}}$. Nevertheless, μ_{sample} is supported in the same area as μ_{data}. For details on the numerical implementation see Appendix B.

36 The marginals of X_t are denoted by π_t. The forward SDE is run until some terminal time T.
37 Furthermore, the *reverse SDE* is defined by

$$
\begin{aligned}
\mathrm{d}Y_t &= -\beta(Y_t)\mathrm{d}t + \sigma\sigma^T \nabla \log p_{T-t}(Y_t)\mathrm{d}t + \sigma\mathrm{d}B_t, \\
Y_0 &\sim q_0.
\end{aligned}
\tag{2}
$$

38 We refer to the marginals of Y_t as q_t. The samples are generated from the final distribution q_T,
39 i.e. $\mu_{\text{sample}} := q_T$. The reverse SDE has the property that if q_0 is chosen to be equal to π_T, then
40 $q_t = \pi_{T-t}$. In particular, this implies that $q_T = \pi_0$. Therefore, if we have samples from π_T, we can
41 run the reverse SDE on them to create new samples from π_0.

42 In the following we will denote by p_t the marginals of the forward SDE when started in the true data
43 generating distribution $\pi_0 = \mu_{\text{data}}$. We will denote by $\hat{p}_t$ the marginals of the forward SDE when
44 started in $\pi_0 = \hat{\mu}_{\text{data}}$. Optimally, we would like to run the algorithm using $\pi_0 = \mu_{\text{data}}$, i.e. with
45 marginals $\pi_t = p_t$. This is however not possible, since μ_{data} itself is unknown.

46 To circumvent the problem of not knowing p_T, the forward SDE is chosen such that it forgets its
47 initial condition p_0. At time T, the marginal p_T is then well approximated by a proxy distribution
48 $\mu_{\text{prior}} \approx p_T$, independently of p_0. Additionally, the marginals p_t and therefore the scores $\nabla \log p_t$
49 cannot be evaluated for $p_0 = \mu_{\text{data}}$. Therefore, the scores are replaced by a neural network $s_\theta(x, t)$,
50 which is trained via score-matching techniques [Vincent, 2011, Hyvärinen and Dayan, 2005].

51 As a result, SGMs make two approximations. The first one is in approximating p_T by μ_{prior}. The
52 second one is the approximation of $\nabla \log p_t$ by the neural net $s_\theta(x, t)$. We illustrate this in Figure 1.
53 It is important to understand how these approximations translate to the distance of μ_{sample} to μ_{data} or
54 $\hat{\mu}_{\text{data}}$.

55 Some early works already deal with these questions. In [De Bortoli et al., 2021] the total variation
56 distance between μ_{sample} and μ_{data} in bounded, whereas [Song et al., 2021a] derives bounds with
57 respect to the KL-Divergence. However, both of these works assume that the initial distribution μ_{data}
58 is rather well behaved. In particular, it is assumed that $\mu_{\text{data}}(x) > 0$ for all x. We shortly discuss this
59 assumption now.

(a) Here μ_{data} is chosen as the uniform distribution on the unit sphere $\mathcal{S}^1$. We use the Brownian motion for the forward SDE. We can compute the exact score $\nabla \log p_t(x)$. We perturb it with the vector $v = (x = 0, y = -1)$ and define the approximation $s(x, t) = \nabla \log p_t(x) + v$. Furthermore, we purposely adopt a poor approximation $\mu_{\text{prior}} \not\approx p_1$ by setting $\mu_{\text{prior}} = \mathcal{N}(m, I)$, where $m = (x = 1.5, y = 0)$. We then run the reverse SDE (6). The figure shows heat maps of the intermediate distributions q_t of the reverse SDE. At time $t = 1$ we reach $q_1 = \mu_{\text{sample}}$. We see that μ_{sample} is a distribution on $\mathcal{M} = \mathcal{S}^1$, albeit not the uniform one.

(b) The above experiment is repeated but with μ_{data} chosen to be the uniform distribution on $\mathcal{M} = \{x_i\}_{i=1}^9$, where the x_i are 9 evenly spaced points on the unit sphere $\mathcal{S}^1$. Notice that while the approximation errors cause a non-uniform distribution the training examples, μ_{sample} is still supported solely on $\mathcal{M}$ and will not generate novel samples.

Figure 2

Assuming that $\mu_{\text{data}}(x) > 0$ for any x means that one postulates that every x is a possible sample from μ_{data}. For example, this implies that even if all samples $\{x_1, \ldots, x_n\}$ of μ_{data} consist of images of human faces, we say that μ_{data} can possibly also generate images of for example furniture, animals or pure white noise. Even though it might have low probability, any combinations of pixels is a possible sample from μ_{data}. The assumption that μ_{data} is actually supported on some lower dimensional substructure $\mathcal{M}$ is well known under the name *manifold hypothesis*, see for example [Bengio et al., 2013, Pope et al., 2021]. In practice this means that $\mu_{\text{data}}(x) = 0$ for many x. This also leads to exploding scores $\nabla \log p_t$ as $t \to 0$ (see Section 5), a behaviour which also observed empirically and further underpins the relevance of the manifold assumption. We will from now on denote the support of μ_{data} as *data manifold* $\mathcal{M}$.

A fundamental question is then: What is the support of μ_{sample} and how does it compare to $\mathcal{M}$. This is interesting for multiple reasons. On the one hand, if we for example assume that $\mathcal{M}$ is the set of all images of faces, then the knowledge that μ_{sample} also has support $\mathcal{M}$ implies that, regardless of how close μ_{sample} actually is to μ_{data}, it will at least always produce an image of a face. On the other hand, we can also compare the support of μ_{sample} to that of $\hat{\mu}_{\text{data}}$. The measures μ_{sample} and μ_{data} sharing their support translates to μ_{sample} memorizing the training data and not being able to generalize. Both of these are very valuable insights into the qualities of μ_{sample} on its own. Furthermore, statistical distances like the KL-Divergence or the total variation distance are only meaningful if the support of the measures overlap to some degree, as otherwise they will equal the maximum distance value.

If two measures have the same support, they are said to be *equivalent*. Our main result is the following:

Main result. *We identify conditions under which μ_{sample} is able to learn the data manifold, that is $supp(\mu_{\text{sample}}) = supp(\mu_{\text{data}})$. Applying these results to different settings, we find precise conditions under which an SGM memorizes its training data or under which it is able to learn the right data manifold $\mathcal{M}$.*

In particular we find that for SGMs to be able to generalize, the approximation error made when approximating the training drift $\nabla \log \hat{p}_t$ has to be unbounded.

A first illustration of these results is given in Figure 2. We use a simple example, where we can perfectly evaluate the true drift $\nabla \log \pi_t$. We then choose an incorrect initial condition μ_{prior} which is far from p_T, and also add a constant error to $\nabla \log \pi_t(x)$. The initial measures π_0 are given as

the uniform distribution on the unit sphere and the uniform distribution on 9 samples from the unit sphere in Figure 2a and Figure 2 respectively. We see that the final distribution of the reverse SDE, μ_{sample}, is not the uniform distribution anymore. This is due to errors in the initial conditions and the drift. Nevertheless, μ_{sample} is still supported on the exact same subset as π_0.

We now proceed as follows. In Section 2 we will summarize some of the most popular forward SDEs that are applied in SGMs. Then, in Section 3 we discuss how the drift approximation $s_\theta(x, t)$ for SGMs is trained in most implementations. We are ready to state our main results in Section 4. In Section 5 we show why and with which speed the true drift $\nabla \log p_t$ of an SGM explodes under the manifold hypothesis. Most of the paper discusses the error in the drift approximation. But as we have discussed, we also have an error in the initial conditions. In Section 6 we will discuss how large the error in the initial conditions will be in practice.

2   Popular SDEs used in SGMs

In this section we introduce some of the most popular SDEs that are used when implementing SGMs. The first works on SGMs studied discrete forward and backward processes. Nevertheless, the transition kernels and algorithms proposed in those works can be seen as discretisations of some well-known SDEs. More recent works have studied this connection and state the algorithms in terms of SDEs ([Song et al., 2021b, Huang et al., 2021]).

Brownian Motion: The works [Song and Ermon, 2019, 2020] can be seen as a discretization of the SDE
$$\mathrm{d}X_t = \sigma(t)\mathrm{d}W_t.$$
Denoting $h(t) = \int_0^t \sigma(s)\mathrm{d}s$, the solution to the above process can be explicitly stated as a time-changed Brownian motion, $X_t = W_{h(t)}$. The time-change can help in the implementation but does not alter the qualitative behaviour of the reverse SDE. In our following analysis we therefore set $\sigma(t) = 1$. Nevertheless, our results still hold for any positive $\sigma(t)$.

Ornstein-Uhlenbeck Process: The works [Sohl-Dickstein et al., 2015, Ho et al., 2020] can be seen as a discretization of
$$\mathrm{d}X_t = -\frac{1}{2}\alpha(t)X_t\mathrm{d}t + \sqrt{\alpha(t)}\mathrm{d}W_t,$$
which is an Ornstein-Uhlenbeck process. Again, the parameters α_t are a time-change and do not influence the properties that we are investigating in this paper. Therefore, to simplify notation, we again set $\alpha(t) = 1$.

Critically Damped Langevin Dynamics (CLD):   The work [Dockhorn et al., 2021] studies a second order SDE. Here artificial velocity coordinates V_t are introduced and the system under consideration is
$$\begin{aligned}
\mathrm{d}X_t &= V_t, \\
\mathrm{d}V_t &= -X_t - 2V_t + 2\mathrm{d}W_t,
\end{aligned}$$
where $X_0 \sim p_{\text{data}}$, $V_0 \sim \mathcal{N}(0, I)$. For generation one runs the reverse SDE in X_t and V_t but discards the V coordinate at the end. The work [Dockhorn et al., 2021] also includes the parameters M and γ. We set both to 1 as they do in their numerical experiments.

3   Score approximation with a finite number of samples

We now quickly discuss how the neural network is trained to approximate $\nabla \log p_t$ and what implications this has. The score $\nabla \log p_t$ is approximated by minimizing a variant of
$$L(\theta) = \int_0^T w(t)\, \mathbb{E}_{p_t(x)}[\|\nabla \log p_t(x) - s_\theta(x, t)\|^2]\mathrm{d}t. \tag{3}$$

for some weighting function w. This is done via score matching techniques (see [Hyvärinen and Dayan, 2005, Vincent, 2011, Song et al., 2020]). However, the above expectation depends on

(a)

(b)

Figure 3: (a): Both lines correspond to the same experiment for different drifts in the reverse SDE. For both lines we started $N = 1000$ paths in the zero vector in $\mathbb{R}^{32 \times 32 \times 3}$. For the blue line we used the pretrained CIFAR-10 DDPMPP model from Song et al. [2021b], whereas for the orange line we used the true drift $\nabla \log \hat{p}_t$, which is a mixture of 50000 Gaussians, one for each training example in CIFAR-10. We then saved the distance from Y_t to the CIFAR-10 training examples, by calculating the distance to the closest example. Above we plot the average distance. We see, that while the reverse SDE run with $\hat{p}_t$ will have a distance of 0 to the training examples in the end, the SDE with the pretrained drift keeps some distance to the training examples and therefore produces novel images. (b): We evaluate $\nabla \log \hat{p}_t$ as in (a). We do the analogous experiment to Figure 2b on CIFAR-10 and add the 1-vector in $\mathbb{R}^{32 \times 32 \times 3}$ to the score, i.e. we use $s(x, t) = \nabla \log \hat{p}_t(x) + (1, 1, \ldots, 1)$. The error $e(x, t) = (1, 1, \ldots, 1)$ is bounded and therefore fulfils Assumption 1. All of the plotted images are nearly equal to an CIFAR-10 image. If we run the SDE with $\nabla \log \hat{p}_t$, we know that the final distribution should be the uniform distribution on all training examples. We see, that the error did not change the support, it just changed the distribution on the training examples. In this case it made very white images much more probable, since the one-vector corresponds to a fully white image.

p_t, which we cannot evaluate since it depends on $p_0 = \mu_{\text{data}}$. Nevertheless, we can evaluate the approximation $\hat{p}_t$,

$$\hat{p}_t(x) = \mathbb{E}_{\hat{\mu}_{\text{data}}(x_0)}[p_{t|0}(x|x_0)] = \frac{1}{N} \sum_{i=1}^{N} p_{t|0}(x|x^i) \approx p_t(x) = \mathbb{E}_{\mu_{\text{data}}}[p_{t|0}(x|x_0)]. \tag{4}$$

The surrogate loss

$$\hat{L}(\theta) = \int_0^T w(t) \, \mathbb{E}_{\hat{p}_t(x)}[\|\nabla \log \hat{p}_t(x) - s_\theta(x, t)\|^2] \mathrm{d}t, \tag{5}$$

can be evaluated and used for training. This raises the question in which way or to which degree the loss should actually be minimized. If we would minimize this loss perfectly, then $s_\theta(x, t)$ would be equal to $\nabla \log \hat{p}_t$. The reverse SDE started in an appropriate initial condition with drift $\nabla \log \hat{p}_t$ however, will produce samples from $\hat{\mu}_{\text{data}}$, which are training examples.

4   Effects of the approximations

This section contains our main results. We first state the assumptions we have to make and then the Theorems.

4.1   Error in the initial condition

The following assumption is needed for the reverse SDE to be defined even in the case when the initial distribution π_0 has a degenerate support. In Section **??** we will show that all the SDEs from Section 2 satisfy the Assumption.

143 **Assumption 1.** *Let C and $C_{S,N}$ be constants that can change from appearance to appearance.*

144   *(i) β is globally Lipschitz, i.e.$\|\beta(x) - \beta(y)\| \leq C\|x - y\|$.*

145   *(ii) β grows at most linearly, i.e. $\|\beta(x)\| \leq C(1 + \|x\|)$*

146   *(iii) X_t has a density $\pi_t \in C^1$ for every $t > 0$ and $\int_{t_0}^1 \int_{\|x\|<R} |\pi_t(x)|^2 + \|\nabla_x \pi_t(x)\|^2 \mathrm{d}x \mathrm{d}t < \infty$*
147   *for any $R > 0$ and $0 < t_0 \leq T$.*

148 *Furthermore, for each $S \in (0, T)$,*

149   *(iv) $\nabla \log \pi_t$ is locally Lipschitz, $\|\nabla \log \pi_t(x) - \nabla \log \pi_t(y)\| \leq C_{S,N} \|x - y\|$ for all $t \in [S, T]$*
150   *and $\|x\|, \|y\| \leq N$.*

151 Conditions (i)-(iii) are technical conditions on the forward SDE. They ensure that if we run a
152 solution to the forward SDE, X_t, backwards in time, then $X_t^R := X_{T-t}$ will be a solution to the
153 reverse SDE (2) on $[0, T)$. The last condition then ensures that the solutions to the reverse SDE are
154 unique, therefore we will be able to transmit the properties of X^R to any other solution Y_t of (2).
155 The following result shows that Assumption 1 can be expected to hold in practice. To simplify the
156 calculations we assume that the data manifold $\mathcal{M} = \mathrm{supp}(\mu_{\mathrm{data}})$ is contained in a ball of diameter M.
157 This is a natural assumption for many data sets. Nevertheless, we note that this assumption could be
158 weakened by additional technical effort.

159 **Lemma 1.** *Assume that the data manifold $\mathcal{M}$ is contained in a ball of radius M. Then all the*
160 *methods introduced in Section 2 fulfil Assumption 1.*

161 We are now ready to state our first main result,

162 **Theorem 1.** *Denote by π_t the marginals of the forward SDE started in π_0. Assume that Assumption*
163 *1 holds and that μ_{prior} is absolutely continuous with respect to π_T. Then the following hold.*

164   *(i) Let Y_t be a solution to (2) on $[0, T)$. The limit $Y_T := \lim_{t \to T} Y_t$ exists almost surely. We*
165   *refer to its distribution as μ_{sample}. The distribution μ_{sample} is absolutely continuous with*
166   *respect to μ_{data}. If π_T and μ_{prior} are equivalent, then so are μ_{sample} and π_0.*

167   *(ii) Furthermore, for any f-divergence D_f,*
$$D_f(\mu_{sample}|\pi_0) \leq D_f(\mu_{prior}|\pi_T) \quad \text{and} \quad D_f(\pi_0|\mu_{sample}) \leq D_f(\pi_T|\mu_{prior}).$$

168 Applying the above theorem with $\pi_T = p_T$ tells us something about the equivalence and distance
169 between μ_{sample} and μ_{data}, since $\pi_0 = p_0 = \mu_{\mathrm{data}}$. Applying the theorem with $\hat{p}_t$ tells us something
170 about the generalization capabilities of SGMs.

171 The measures π_T and μ_{prior} are normally both supported on all of $\mathbb{R}^d$ and therefore equivalent. Since
172 Assumption 1 also holds in most situations (see Lemma 1), the requirements for Theorem 1 are
173 satisfied in practice. From item (i) with $\pi_t = \hat{p}_t$ we can conclude that the error we make in the initial
174 conditions is not responsible for the generalization capacities of SGMs.

175 The second item then shows that the f-divergences between μ_{sample} and μ_{data} are bounded by the
176 f-divergences of μ_{prior} to p_T. The total variation distance and the KL-Divergence are both special
177 cases of f-divergences.

178 ## 4.2 Error in the drift

179 Given a forward SDE with marginals π_t and an approximation $s(x, t)$ to $\nabla \log \pi_t$, we define the
180 reverse SDE for the approximation s as

$$\begin{aligned} \mathrm{d}\tilde{Y}_t &= -\beta(\tilde{Y}_t)\mathrm{d}t + \sigma\sigma^T s(\tilde{Y}_t, t)\mathrm{d}t + \sigma\mathrm{d}B_t, \\ \tilde{Y}_0 &\sim q_0. \end{aligned} \tag{6}$$

181 **Assumption 2.** *We assume that the reverse SDE $\tilde{Y}_t$ has a solution on $[0, T)$. For $t < T$, we define*
182 *the Girsanov weights*

$$Z_t = \exp\left(\int_0^t \sigma^T(s(\tilde{Y}_t, t) - \nabla \log \pi_t(\tilde{Y}_t)) \cdot \mathrm{d}B_s - \frac{1}{2} \int_0^t \|\sigma^T(s(\tilde{Y}_t, t) - \nabla \log \pi_t(\tilde{Y}_t))\|^2 \mathrm{d}s \right)$$
$$\tag{7}$$

183 *and assume that the Z_t are a uniformly integrable martingale.*

We shortly discuss this assumption. The assumption that Z_t is a martingale is equivalent to the expectation of Z_t being equal to 1 for all t. The assumption that it is uniformly is more technical (see Appendix A.2), but is fulfilled for example if for some $p > 1$, $\mathbb{E}[|Z_t|^p] < \infty$ for all t. For example, if the Z_t have bounded variance, then they are uniformly integrable.

A condition that ensures that Z_t is both, a martingale and uniformly integrable, is given by *Novikov's condition* Novikov [1980]. It states, that if

$$N = \mathbb{E}\left[\exp\left(\frac{1}{2}\int_0^T \|\sigma^T(s(x,t) - \nabla \log \pi_t)\|^2 \mathrm{d}s\right)\right] < \infty, \tag{8}$$

then Z_t is a uniformly integrable martingale.

Using Assumption 2 we can now state

Theorem 2. *Assume that Assumption 2 holds. Assume furthermore that Assumption 1 holds with $\nabla \log \pi_t$ replaced by $s(x,t)$.*

Then $\tilde{Y}_T = \lim_{t \to T} \tilde{Y}_t$ is well defined. Moreover, its distribution is equivalent to the distribution of Y_T. In particular, if $\|s(x,t) - \nabla \log \hat{p}_t\|$ is bounded, then Assumption 2 holds, and the SGM has memorized its training data.

Putting Theorem 1 and Theorem 2 together, we can conclude that, if both Assumptions hold, μ_{sample} is equivalent to π_0. Therefore, if Assumption 2 holds for $\pi_t = p_t$, we have a positive statement and know that μ_{sample} will have the exact same support as μ_{data}, i.e. it has learned the data manifold $\mathcal{M}$.

If the Assumption would hold for $\pi_t = \hat{p}_t$, we would however just memorize the training data, see Figure 2b or Figure 3b for a visualization. However, empirically it has been shown that SGMs are able to create novel samples (see, Figure 3a or Dhariwal and Nichol [2021]). Therefore, we can deduce that Assumption 2 is violated in practice.

SGMs are trained by minimizing $\hat{L}(\theta)$ (see (5)). On the other hand, if $L(\theta)$ is minimized well enough for Z_t to be fulfil Assumption 2, then the SGM is collapsed onto its training data. This is especially interesting since

$$L(\theta) = \mathbb{E}_{Y_0 \sim \hat{p}_0}[\log(Z_T)].$$

We evaluated N from (8) on CIFAR-10, once for the difference between the $s_\theta(x,t)$ from Song et al. [2021b] so and $\nabla \log \hat{p}_t$, and once by just using a perturbed drift with a constant error, $s(x,t) = \nabla \log \hat{p}_t + \frac{1}{2}(1,1,\ldots,1)$, see Figure 4. The reverse SDE using the drift $\nabla \log \hat{p}_t + \frac{1}{2}(1,1,\ldots,1)$, is equivalent to $\hat{\mu}_{\text{data}}$, as we know from Theorem 2 and have also already observed in Figure 3b. Figure 4 confirms that Z_t is indeed a uniformly integrable martingale and therefore fulfils Assumption 2.

In future work we believe that further understanding the characteristics of Z_t, how they relate to the minimization of $L(\theta)$ and generalization is crucial to the understanding of SGMs and their empirical success. Recent works also study the question on how the neural network architecture and parametrization is related to the boundedness of the output Kim et al. [2022]. The relationship between these choices and the properties of Z_t is also an interesting research avenue. Lastly, the distribution of Z_t is very heavy-tailed. Most of the samples are very small with a few extremely large samples in between. Therefore one needs many samples from Z_t to understand its characteristics. Finding robust estimators for Z_t or its expectation could help their usage in the training or evaluation procedure.

5 Drift Explosion under manifold hypothesis

First, we note that all SDEs in Section 2 are linear SDEs. Therefore, their transition kernels are Gaussian ([Pavliotis, 2014, Section 3.7]):

$$\pi_t(Z_t = z|Z_0 = z_0) = \mathcal{N}(z; m_t(z_0), \Sigma_t).$$

The explicit form of m_t and Σ_t differ for each of the SDEs and can be found in Appendix C.1. We remark that Σ_t does not depend on the initial condition z_0. The transition kernel above gives us the distribution of the SDE started in a single point z_0. Since we start the SDE in μ_{data} we need to average over μ_{data} to get the marginal at time t:

$$p_t(z) = \int_{\mathbb{R}^d} \mathcal{N}(z; m_t(z_0), \Sigma_t)\mu_{\text{data}}(z_0)\mathrm{d}z_0. \tag{9}$$

Figure 4: Left: We simulated the reverse SDE on CIFAR-10, once with the pretrained CIFAR-10 DDPMPP model s_θ from Song et al. [2021b] and once with a perturbed drift $s(x,t) = \nabla \log \hat{p}_t + \frac{1}{2}(1,1,\ldots,1)$. We then evaluated the integral (8) numerically. For the perturbed drift, the integral does not seem to explode as $t \to 1$, implying that Z_t is a martingale. We see that for the DDPMPP drift, the integral explodes, therefore we can not infer that Z_t is a martingale. We used $N = 12000$ simulations from both of the SDEs to generate this plot. Right: We again ran the two SDEs with the drifts as in the left Figure. This time, we measured the distance to the empirical drift $\|s(x,t) - \nabla \log \hat{p}_t(x)\|$. We repeated the experiment $N = 2560$ times and plotted the mean distance. For the constant perturbation we also of course get a constant distance. The distance of the true drift to $\nabla \log \hat{p}_t$ is initially very small but explodes as $t \to 1$. From our results we know that this explosion is necessary for the SGM to generalize.

We can also compute the additional drift in the reverse SDE (see Appendix C.2),

$$\nabla \log p_t = \frac{\nabla p_t(z)}{p_z(z)} = \Sigma_t^{-1}(z - \mathbb{E}[m_t(Z_0)|Z_t = z]). \tag{10}$$

The drift is intuitive, especially for the Brownian Motion, where $m_t(z) = z$ and $\Sigma_t = t$. In that case, $\mathbb{E}[Z_0|Z_t = z]$ is the expected position from which the SDE was started on the manifold, if it is at position z at time t. For a point z which has distance $\delta > 0$ to $\mathcal{M}$, $\|z - \mathbb{E}[Z_0|Z_t = z]\| > \delta$, since Z_0 is in $\mathcal{M}$ almost surely. Therefore, we can infer that $\nabla \log p_t(z)$ explodes like $\frac{\delta}{t}$ as $t \to 0$. A similar analysis can be conducted for the other two SDEs using the formulas for the transition kernels in Appendix C.1. In the Brownian motion case with $\sigma = 1$, the forward SDE can be represented as $X_t = X_0 + B_t$. Since B_t has distribution $\mathcal{N}(0, tI_d)$, we can expect the distance of $X_t - X_0$ to be $\mathbb{E}[\|X_t - X_0\|] = \mathbb{E}[\|B_t\|] \approx \sqrt{t}$. Therefore, under the manifold hypothesis $\nabla \log p_t(Y_t)$ will behave like $\frac{\sqrt{t}}{t} = \frac{1}{\sqrt{t}}$. In particular, it will explode along a typical path of Y_t as $t \to 0$. This behaviour is also observed empirically. This argument can be generalized to arbitrary SDEs. If σ_t^2 is the variance of p_t, then we can expect $\nabla \log p_t(Y_t)$ to be of order $\frac{1}{\sigma_t}$. Furthermore, $\nabla \log p_t(x)$ will point towards the data manifold for small t. The drift $\nabla \log p_t$ then acts like a *support matching force*, where the force grows to infinity as $t \to 0$, absorbing all the SDE paths onto the manifold.

6 Distance from p_T to μ_{prior}

We have seen in Theorem 1 that the distance between μ_{sample} and μ_{data} is directly related to the distance between p_T and μ_{prior}. The OU-process and the CLD are well-known SDEs in the Markov Chain Monte Carlo community. They form building blocks of algorithms like Langevin MCMC ([Roberts and Tweedie, 1996, Welling and Teh, 2011]) or Hamiltonian Monte Carlo ([Neal et al., 2011]) and the speed at which they converge to their stationary distribution is well studied ([Roberts and Tweedie, 1996, Markowich and Villani, 2000, Bou-Rabee et al., 2010, Eberle et al., 2019, Cao et al., 2019]). It is known that their intermediate distributions p_t will converge to $\mu_{\text{prior}} = \mathcal{N}(0, I_d)$ at an exponential rate.

The Brownian motion however does not converge to a stationary distribution and therefore one has to choose a different μ_{prior}^T for each T to approximate p_T. We now study the distance of μ_{prior}^T to p_T. If we transfer the suggestions of [Song et al., 2021b, Appendix C] to our setting, they propose to set μ_{prior}^T to $\mathcal{N}(0, T)$. However, p_t has the same mean as μ_{data} and has variance $\text{Var}[\mu_{\text{data}}] + T$. Loosely

speaking, in contrast to the other two SDEs, the Brownian Motion does not forget its initial mean and covariance. As we see in the following lemma, we can decrease the distance between p_T and μ_{prior}^T by choosing the mean and covariance of μ_{prior}^T equal to the mean and covariance of p_T.

Lemma 2. *Let p_T be the T-time marginal of the Brownian motion process X_t of Section 2. The following minimization problem*

$$\min_{m,C} KL(p_T \mid \mathcal{N}(m, C))$$

is minimized by $m_T = \mathbb{E}[\mu_{data}]$ and $C_T = \text{Cov}[\mu_{data}] + TI_d$. If we restrict the covariance to be a multiple of the identity matrix, i.e.

$$\min_{m,c} KL(p_T \mid \mathcal{N}(m, cI_d)),$$

the problem is solved by choosing m as above and c as $c = \mathbb{E}[\|X_t - m\|^2] = trace(C_T)$.

This result is a slight variation on the well known fact that the KL-projection in the second argument matches its moments. We prove it in Appendix E.2. Equivalently to choosing the mean of μ_{prior}^T equal to the mean of p_T, $\mathbb{E}[p_T] = \mathbb{E}[p_0]$, one can normalize the training data to have mean 0 and then also centre μ_{prior}^T. However, the covariance matrix will not be equal to I_d after common normalization methods.

The next lemma bounds the distance of p_T and μ_{prior}^T in the Brownian motion case.

Lemma 3. *Let p_t be the time t-marginal of a Brownian motion with initial condition μ_{data}. Denote by $c_i, i = 1, \ldots, d$, the eigenvalues of $\text{Cov}(\mu_{data})$. Let μ_{prior}^T be any normal distribution. Then*

$$KL(p_T | \mu_{prior}^T) \leq \frac{1}{2} \log \left(\frac{\prod_{i=1}^{d}(c_i + T)}{T^d} \right).$$

This lemma is proven by explicitly bounding the distance of p_T to the optimal normal distribution from Lemma 2, see Appendix E.2.

7 Broader impact

The results deepen the understanding of score-based generative models. As such, they can be seen as a step towards improving the quality of generative models. Therefore the possible negative societal impacts are the same ones that apply to generative modelling in general. First, generative models can be used to create synthetic data that is hard to distinguish from real data (for example images or videos), see [Mirsky and Lee, 2021]. Second, generative models can learn and reproduce biases that are prevalent in the training data ([Esser et al., 2020]). Last, depending on the application, generative models might be used to do creative work that was previously done by humans.

8 Conclusion

We conducted a theoretical study of some properties of SGMs. We found explicit conditions under which the sample measure μ_{sample} is equivalent to the true data generating distribution μ_{data}. Under these conditions we can guarantee, that the SGM generates samples that could also be samples from μ_{data}. Furthermore, each sample that can be generated by μ_{data} also has positive probability under μ_{sample}, meaning that the full support is covered.

Since one can not actually access the full support of μ_{data}, but only a finite number of training examples $\{x_i\}_{i=1}^N$, our results can be applied to find conditions under which the SGM memorizes its training data. We believe that this observation provides a first step towards understanding the generalization capabilities of SGMs.

References

G. Batzolis, J. Stanczuk, C. Schönlieb, and C. Etmann. Conditional image generation with score-based diffusion models. *CoRR*, abs/2111.13606, 2021. URL https://arxiv.org/abs/2111.13606.

Y. Bengio, A. Courville, and P. Vincent. Representation learning: A review and new perspectives. *IEEE transactions on pattern analysis and machine intelligence*, 35(8):1798–1828, 2013.

F. Bordes, S. Honari, and P. Vincent. Learning to generate samples from noise through infusion training. *arXiv preprint arXiv:1703.06975*, 2017.

N. Bou-Rabee, M. Hairer, and E. Vanden-Eijnden. Non-asymptotic mixing of the MALA algorithm. *IMA Journal of Numerical Analysis*, 33, 08 2010. doi: 10.1093/imanum/drs003.

R. Cai, G. Yang, H. Averbuch-Elor, Z. Hao, S. J. Belongie, N. Snavely, and B. Hariharan. Learning gradient fields for shape generation. In A. Vedaldi, H. Bischof, T. Brox, and J. Frahm, editors, *Computer Vision - ECCV 2020 - 16th European Conference, Glasgow, UK, August 23-28, 2020, Proceedings, Part III*, volume 12348 of *Lecture Notes in Computer Science*, pages 364–381. Springer, 2020. doi: 10.1007/978-3-030-58580-8_22. URL https://doi.org/10.1007/978-3-030-58580-8_22.

Y. Cao, J. Lu, and L. Wang. On explicit L^2-convergence rate estimate for underdamped Langevin dynamics. *arXiv preprint arXiv:1908.04746*, 2019.

N. Chen, Y. Zhang, H. Zen, R. J. Weiss, M. Norouzi, and W. Chan. Wavegrad: Estimating gradients for waveform generation. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021. URL https://openreview.net/forum?id=NsMLjcFaO8O.

M. Costa and T. Cover. On the similarity of the entropy power inequality and the Brunn-Minkowski inequality (corresp.). *IEEE Transactions on Information Theory*, 30(6):837–839, 1984.

V. De Bortoli, J. Thornton, J. Heng, and A. Doucet. Diffusion schrödinger bridge with applications to score-based generative modeling. *Advances in Neural Information Processing Systems*, 34, 2021.

P. Dhariwal and A. Nichol. Diffusion models beat gans on image synthesis. *Advances in Neural Information Processing Systems*, 34, 2021.

T. Dockhorn, A. Vahdat, and K. Kreis. Score-based generative modeling with critically-damped Langevin diffusion. *CoRR*, abs/2112.07068, 2021. URL https://arxiv.org/abs/2112.07068.

A. Eberle, A. Guillin, and R. Zimmer. Couplings and quantitative contraction rates for langevin dynamics. *The Annals of Probability*, 47(4):1982–2010, 2019.

P. Esser, R. Rombach, and B. Ommer. A note on data biases in generative models. In *NeurIPS 2020 Workshop on Machine Learning for Creativity and Design*, 2020. URL https://arxiv.org/abs/2012.02516.

U. G. Haussmann and E. Pardoux. Time reversal of diffusions. *The Annals of Probability*, pages 1188–1205, 1986.

J. Ho, A. Jain, and P. Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 33:6840–6851, 2020.

J. Ho, C. Saharia, W. Chan, D. J. Fleet, M. Norouzi, and T. Salimans. Cascaded diffusion models for high fidelity image generation. *Journal of Machine Learning Research*, 23(47):1–33, 2022.

C.-W. Huang, J. H. Lim, and A. C. Courville. A variational perspective on diffusion-based generative models and score matching. *Advances in Neural Information Processing Systems*, 34, 2021.

A. Hyvärinen and P. Dayan. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4), 2005.

M. Jeong, H. Kim, S. J. Cheon, B. J. Choi, and N. S. Kim. Diff-tts: A denoising diffusion model for text-to-speech. In H. Hermansky, H. Cernocký, L. Burget, L. Lamel, O. Scharenborg, and P. Motlícek, editors, *Interspeech 2021, 22nd Annual Conference of the International Speech Communication Association, Brno, Czechia, 30 August - 3 September 2021*, pages 3605–3609. ISCA, 2021. doi: 10.21437/Interspeech.2021-469. URL https://doi.org/10.21437/Interspeech.2021-469.

A. Jolicoeur-Martineau, R. Piché-Taillefer, I. Mitliagkas, and R. T. des Combes. Adversarial score matching and improved sampling for image generation. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021. URL `https://openreview.net/forum?id=eLfqMl3z3lq`.

I. Karatzas and S. Shreve. *Brownian motion and stochastic calculus*, volume 113. Springer Science & Business Media, 2012.

D. Kim, S. Shin, K. Song, W. Kang, and I. Moon. Soft truncation: A universal training technique of score-based diffusion model for high precision score estimation. In K. Chaudhuri, S. Jegelka, L. Song, C. Szepesvári, G. Niu, and S. Sabato, editors, *International Conference on Machine Learning, ICML 2022, 17-23 July 2022, Baltimore, Maryland, USA*, volume 162 of *Proceedings of Machine Learning Research*, pages 11201–11228. PMLR, 2022. URL `https://proceedings.mlr.press/v162/kim22i.html`.

D. P. Kingma, T. Salimans, B. Poole, and J. Ho. Variational diffusion models. *arXiv preprint arXiv:2107.00630*, 2021.

A. Klenke. *Probability theory: a comprehensive course*. Springer Science & Business Media, 2013.

Z. Kong, W. Ping, J. Huang, K. Zhao, and B. Catanzaro. Diffwave: A versatile diffusion model for audio synthesis. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021. URL `https://openreview.net/forum?id=a-xFK8Ymz5J`.

A. Krizhevsky, G. Hinton, et al. Learning multiple layers of features from tiny images. 2009.

C. Léonard. Stochastic derivatives and generalized h-transforms of Markov processes. *arXiv preprint arXiv:1102.3172*, 2011.

H. Li, Y. Yang, M. Chang, S. Chen, H. Feng, Z. Xu, Q. Li, and Y. Chen. Srdiff: Single image super-resolution with diffusion probabilistic models. *Neurocomputing*, 479:47–59, 2022. doi: 10.1016/j.neucom.2022.01.029. URL `https://doi.org/10.1016/j.neucom.2022.01.029`.

F. Liese and I. Vajda. On divergences and informations in statistics and information theory. *IEEE Transactions on Information Theory*, 52(10):4394–4412, 2006.

S. Luo and W. Hu. Diffusion probabilistic models for 3d point cloud generation. In *IEEE Conference on Computer Vision and Pattern Recognition, CVPR 2021, virtual, June 19-25, 2021*, pages 2837–2845. Computer Vision Foundation / IEEE, 2021. URL `https://openaccess.thecvf.com/content/CVPR2021/html/Luo_Diffusion_Probabilistic_Models_for_3D_Point_Cloud_Generation_CVPR_2021_paper.html`.

P. A. Markowich and C. Villani. On the trend to equilibrium for the Fokker-Planck equation: an interplay between physics and functional analysis. *Mat. Contemp*, 19:1–29, 2000.

C. Meng, Y. Song, J. Song, J. Wu, J. Zhu, and S. Ermon. Sdedit: Image synthesis and editing with stochastic differential equations. *CoRR*, abs/2108.01073, 2021. URL `https://arxiv.org/abs/2108.01073`.

Y. Mirsky and W. Lee. The creation and detection of deepfakes: A survey. *ACM Computing Surveys (CSUR)*, 54(1):1–41, 2021.

R. M. Neal et al. Mcmc using hamiltonian dynamics. *Handbook of markov chain monte carlo*, 2(11): 2, 2011.

A. Q. Nichol and P. Dhariwal. Improved denoising diffusion probabilistic models. In *International Conference on Machine Learning*, pages 8162–8171. PMLR, 2021.

A. A. Novikov. On conditions for uniform integrability of continuous non-negative martingales. *Theory of Probability & Its Applications*, 24(4):820–824, 1980.

G. A. Pavliotis. *Stochastic processes and applications: diffusion processes, the Fokker-Planck and Langevin equations*, volume 60. Springer, 2014.

S. Peluchetti. Non-denoising forward-time diffusions. 2021.

P. Pope, C. Zhu, A. Abdelkader, M. Goldblum, and T. Goldstein. The intrinsic dimension of images and its impact on learning. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021. URL `https://openreview.net/forum?id=XJk19XzGq2J`.

V. Popov, I. Vovk, V. Gogoryan, T. Sadekova, and M. A. Kudinov. Grad-tts: A diffusion probabilistic model for text-to-speech. In M. Meila and T. Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning, ICML 2021, 18-24 July 2021, Virtual Event*, volume 139 of *Proceedings of Machine Learning Research*, pages 8599–8608. PMLR, 2021. URL `http://proceedings.mlr.press/v139/popov21a.html`.

O. Rioul. Information theoretic proofs of entropy power inequalities. *IEEE Transactions on Information Theory*, 57(1):33–55, 2010.

G. O. Roberts and R. L. Tweedie. Exponential convergence of langevin distributions and their discrete approximations. *Bernoulli*, pages 341–363, 1996.

C. Saharia, J. Ho, W. Chan, T. Salimans, D. J. Fleet, and M. Norouzi. Image super-resolution via iterative refinement. *CoRR*, abs/2104.07636, 2021. URL `https://arxiv.org/abs/2104.07636`.

H. Sasaki, C. G. Willcocks, and T. P. Breckon. UNIT-DDPM: UNpaired image translation with denoising diffusion probabilistic models. *CoRR*, abs/2104.05358, 2021. URL `https://arxiv.org/abs/2104.05358`.

J. Sohl-Dickstein, E. Weiss, N. Maheswaranathan, and S. Ganguli. Deep unsupervised learning using nonequilibrium thermodynamics. In *International Conference on Machine Learning*, pages 2256–2265. PMLR, 2015.

Y. Song and S. Ermon. Generative modeling by estimating gradients of the data distribution. In H. M. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. B. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019, December 8-14, 2019, Vancouver, BC, Canada*, pages 11895–11907, 2019. URL `https://proceedings.neurips.cc/paper/2019/hash/3001ef257407d5a371a96dcd947c7d93-Abstract.html`.

Y. Song and S. Ermon. Improved techniques for training score-based generative models. *Advances in neural information processing systems*, 33:12438–12448, 2020.

Y. Song, S. Garg, J. Shi, and S. Ermon. Sliced score matching: A scalable approach to density and score estimation. In *Uncertainty in Artificial Intelligence*, pages 574–584. PMLR, 2020.

Y. Song, C. Durkan, I. Murray, and S. Ermon. Maximum likelihood training of score-based diffusion models. *Advances in Neural Information Processing Systems*, 34:1415–1428, 2021a.

Y. Song, J. Sohl-Dickstein, D. P. Kingma, A. Kumar, S. Ermon, and B. Poole. Score-based generative modeling through stochastic differential equations. In *9th International Conference on Learning Representations, ICLR 2021, Virtual Event, Austria, May 3-7, 2021*. OpenReview.net, 2021b. URL `https://openreview.net/forum?id=PxTIG12RRHS`.

A. Vahdat, K. Kreis, and J. Kautz. Score-based generative modeling in latent space. *Advances in Neural Information Processing Systems*, 34, 2021.

P. Vincent. A connection between score matching and denoising autoencoders. *Neural computation*, 23(7):1661–1674, 2011.

M. Welling and Y. W. Teh. Bayesian learning via stochastic gradient langevin dynamics. In *Proceedings of the 28th international conference on machine learning (ICML-11)*, pages 681–688. Citeseer, 2011.

L. Zhou, Y. Du, and J. Wu. 3d shape generation and completion through point-voxel diffusion. In *2021 IEEE/CVF International Conference on Computer Vision, ICCV 2021, Montreal, QC, Canada, October 10-17, 2021*, pages 5806–5815. IEEE, 2021. doi: 10.1109/ICCV48922.2021.00577. URL `https://doi.org/10.1109/ICCV48922.2021.00577`.

A Stochastic prerequisites

In this section we give a formal introduction to some of the concepts used in this work. For a more rigorous treatment, see for example [Klenke, 2013] or [Karatzas and Shreve, 2012].

A.1 Equivalence of measures / Girsanov Theorem

First we define absolute continuity of measures. Let μ and ν be two measures on $(\Omega, \mathcal{F})$, where $\mathcal{F}$ is a σ-algebra.

Definition 1. *We say that μ is absolutely continuous with respect to ν if $\mu(A) = 0$ for any $A \in \mathcal{F}$ such that $\nu(A) = 0$. We also denote this by $\mu \ll \nu$.*

Two measures μ and ν are equivalent if $\mu \ll \nu$ and $\nu \ll \mu$. Loosely speaking, we can say that $\mu \ll \nu$ if the support of μ is contained in the support of ν and they are equivalent if they share the same support.

The Radon-Nikodym theorem tells us that if $\mu \ll \nu$, then under mild conditions there exists a density $\frac{\mathrm{d}\mu}{\mathrm{d}\nu} : \Omega \to \mathbb{R}$ such that $\mu(A) = \int_A \frac{\mathrm{d}\mu}{\mathrm{d}\nu}(\omega)\mathrm{d}\nu(\omega)$. Therefore, we can obtain μ through a reweighting of ν. One specific instance of this is the Girsanov Theorem. Assume we are given the solutions to two SDEs in $\mathbb{R}^d$,

$$\mathrm{d}Y_t = b(t, Y_t)\mathrm{d}t + \sigma(t, Y_t)\mathrm{d}W_t \tag{11}$$

and

$$\mathrm{d}\tilde{Y}_t = b(t, \tilde{Y}_t)\mathrm{d}t + \sigma(t, \tilde{Y}_t)e(t, \tilde{Y}_t)\mathrm{d}t + \sigma(t, \tilde{Y}_t)\mathrm{d}B_t. \tag{12}$$

Both of these induce a measure on the space of continuous functions $\Omega = C([0, T], \mathbb{R}^d)$. We denote them by $\mathbb{P}$ and $\tilde{\mathbb{P}}$ respectively. Then the Girsanov Theorem equips us with conditions under which the measures $\mathbb{P}$ and $\tilde{\mathbb{P}}$ are equivalent. Furthermore, in case of equivalence we get a formula for the density of $\tilde{\mathbb{P}}$ with respect to $\mathbb{P}$. The relative density is given as

$$Z_T = \exp\left(\int_0^T e(s, Y_s)\mathrm{d}W_s - \frac{1}{2}\int_0^T \|e(s, Y_s)\|^2\mathrm{d}s\right).$$

For a full statement of the Girsanov Theorem and under which conditions it holds, see [Karatzas and Shreve, 2012, Section 3.5].

A.2 Uniform integrability

Since we are treating the case where the drift explodes as $t \to T$ we end up with densities

$$Z_t = \exp\left(\int_0^t e(s, Y_s)\mathrm{d}W_s - \frac{1}{2}\int_0^t \|e(s, Y_s)\|^2\mathrm{d}s\right). \tag{13}$$

on $C([0, t], \mathbb{R}^d)$, but not with a density on $C([0, T], \mathbb{R}^d)$. Uniform integrability is exactly the condition one needs to extend these local densities.

Definition 2. *A family $\{X_\alpha\}$ of random variables is called uniformly integrable if*

$$\sup_\alpha \mathbb{E}\left[|X_\alpha|\, 1_{\{|X_\alpha|>s\}}\right] \to 0$$

as $s \to \infty$.

In the proof of Theorem 2 we implicitly use the following two results which we here state as a lemma. The filtration $\mathcal{F}_t$ is defined as in the proof of Theorem 2.

Lemma 4. *Assume the Z_t in (13) form a uniformly integrable martingale on $[0, T)$. Then,*

- *the limit $\lim_{t \to T} Z_t$ exists in L^1. We denote this limit by Z.*

- *Furthermore, $\tilde{\mathbb{P}}$ is absolutely continuous with respect to $\mathbb{P}$ on $\mathcal{F} = \sigma(\cup_{t<T}\mathcal{F}_t)$ with density Z.*

514 *Proof.* Both of these results are standard. The first one can for example be found in [Karatzas and
515 Shreve, 2012, Section 1.3.B]. For the second one we compute that for any $A \in \mathcal{F}_s$,

$$\mathbb{E}_{\mathbb{P}}[1_A Z] = \mathbb{E}_{\mathbb{P}}[1_A \lim_{t \to T} Z_t] = \lim_{t \to T} \mathbb{E}_{\mathbb{P}}[1_A Z_t] = \mathbb{E}_{\mathbb{P}}[1_A Z_s] = \mathbb{E}_{\tilde{\mathbb{P}}}[1_A],$$

516 where we used L^1 convergence in the second equality and the martingale property of Z_s in the third
517 equality. Therefore Z is a density of $\tilde{\mathbb{P}}$ with respect to $\mathbb{P}$ on each $\mathcal{F}_s$ for $s < T$. Therefore Z is also a
518 density of $\tilde{\mathbb{P}}$ with respect to $\mathbb{P}$ on $\mathcal{F}$ which concludes the proof. $\qquad\square$

B Numerics

520 All numerical experiments can be run on a consumer grade computer within a few minutes.

B.1 Figure 1

522 We first discuss the top left figure of Figure 1. We set $p_0 = \mu_{\text{data}}$ to a mixture of two Gaussian
523 $\mathcal{N}(-2, \frac{1}{100})$ and $\mathcal{N}(2, \frac{1}{100})$ with weights $w_1 = \frac{1}{3}$ and $w_2 = \frac{2}{3}$ respectively. Then, we draw $N =$
524 $5\,000\,000$ samples from μ_{data}, denoted by Y_0^n, $n = 1, \ldots, N$. An Euler-Maruyama discretization of
525 the Brownian motion propagates these samples from time $t = 0$ to $t = 1$ by

$$X_{i+1}^n = X_i^n + \sqrt{dt} Z_i^n,$$

526 where $Z_n^i \sim \mathcal{N}(0, 1)$ are i.i.d. random variables, independent of X_m^j for $m \leq n$ and $j = 1, \ldots, N$.
527 The time index i runs from 0 to $I = 2000$ and dt is set to $dt = \frac{1}{I}$. The initial samples $\{X_0^n\}_{n=1}^N$
528 are used to create the left line plot of p_0 and the final samples $\{X_I^n\}_{n=1}^N$ are used to create the right
529 line plot of p_1 using kernel density estimation. The $\{X_i^n\}_{n=1}^N$ are approximate samples from $p_{i/I}$.
530 Therefore, we create histograms using $\{X_i^n\}_{n=1}^N$ to approximate $p_{i/I}$. The height of the histogram
531 bars corresponds to the square root of the colour intensity in the heat map. The horizontal axis in
532 the heat map stands for the time t, whereas the vertical axis stands for the position x. At location
533 (t, x) we plot an estimate of $\sqrt{p_t(x)}$. We apply the square root since it improves the contrast in areas
534 where $p_t(x)$ is close to 0 and makes it more visible where $p_t(x) > 0$ to the observer.

535 For the bottom left figure we show the same plots, just for the reverse SDE (6) instead of the forward
536 SDE. Since the initial distribution is a Gaussian mixture we can exactly calculate p_t using

$$p_t(x) = w_1 \mathcal{N}(x; m_1, s_1^2 + t) + w_2 \mathcal{N}(x; m_2, s_2^2 + t), \tag{14}$$

537 where we use $\mathcal{N}(x; m, v)$ for the probability density function of a normal distribution with mean
538 m and variance v, evaluated at x. With the above expression of p_t one could compute an analytical
539 representation of $\nabla \log p_t$. We use automatic differentiation instead. The reverse SDE (6) is simulated
540 with a disturbance $e(x, t) = 1$ and initial condition $q_0 = \mu_{\text{prior}} = \mathcal{N}(0, 1)$. The Euler-Maruyama
541 method is run with the same step size $dt = \frac{1}{I}$. More precisely, the one step transition kernel of the
542 discretized reverse SDE is

$$Y_{i+1}^n = Y_i^n + dt \left(\nabla \log p_{1 - \frac{i}{I}}(Y_i^n) + 1 \right) + \sqrt{dt} \, \tilde{Z}_i^n, \tag{15}$$

543 where $\tilde{Z}_n^i \sim \mathcal{N}(0, 1)$ are i.i.d. random variables, independent of Y_m^j for $m \leq n$ and $j = 1, \ldots, N$.
544 The plots are created in the same way as for the upper left plot, except that we reverse the time axis to
545 plot p_t and q_{1-t} directly underneath each other.

546 On the right side we plot the same kernel density estimates already plotted on the left side as μ_{sample}
547 and μ_{data} into the same plot for comparison.

B.2 Figure 2b and 2a

549 Figure 2b is created by setting μ_{data} to be the uniform distribution on $M = 9$ equally spaced samples
550 $\{x_i\}_{i=1}^M$ on the unit sphere $\mathcal{S}^1$. This can also be viewed as a Gaussian mixture with 9 components
551 where each component having mean x_i and variance 0. Therefore, we can again explicitly calculate
552 p_t for $t > 0$ as in (14),

$$p_t(y) = \frac{1}{M} \sum_{i=1}^M \mathcal{N}(y; x_i, t).$$

	$m_t(z_0)$	Σ_t
Brownian Motion	z_0	tI_d
OU-Process	$\exp(-t)z_0$	$(1 - \exp(-2t))I_d$

Table 1: The mean and covariance of the Gaussian transition kernels of the Brownian Motion and the Ornstein-Uhlenbeck process SDEs.

The score $\nabla \log p_t(y)$ is evaluated using automatic differentiation. The reverse SDE 6 is simulated with $q_0 = \mathcal{N}((x = -1.5, y = 0), I_2)$ and $e((x, y), t) = (x = 0, y = -1)$. For the numerical simulation, we again use the Euler-Maruyama scheme. We use a step width of $dt = \frac{0.9}{1000} \approx \frac{1}{1000}$ for $t = [0, 0.9]$, $dt = \frac{0.09}{1000} \approx \frac{1}{10\,000}$ for $t \in [0.9, 0.99]$ and of $dt = \frac{1}{100\,000}$ for $t \in [0.99, 1]$. For a simulation of $N = 50\,000$ paths of the reverse SDE, we start by drawing $Y_0^n = A_n + Z_n$, where A_n are i.i.d. uniformly distributed on $\{x_i\}_{i=1}^M$ and $Z_n \sim \mathcal{N}(0, 1)$ i.i.d.. The $\{A_n\}$ and $\{Z_n\}$ are also independent from each other. We then propagate the Y_0^n similarly to (15), except that we use a different values for dt depending on t. This leads to approximate samples Y_t^i from q_t. At the displayed times t we plot the function

$$h_t(x) = \sum_{i=1}^N k(x, Y_t^i),$$

where k is an unnormalized Gaussian kernel with a very small bandwidth parameter,

$$k(x, y) = \exp(-1000\|x - y\|^2).$$

Normalizing h_t gives us a density estimate of q_t. We plot these estimates as heat maps for different values of t.

For Figure 2a we follow the same steps as for Figure 2b, except that μ_{data} is set to the uniform distribution over $M = 256$ evenly spaced samples from the unit sphere $\mathcal{S}^1$.

B.3 Figure 3

For Figure 3 we used the DDPMPP model from the Github repository for the paper from Song et al. [2021b]. Then we evaluated the true score using (4), where the sum runs through the $N = 50000$ training examples of CIFAR-10, Krizhevsky et al. [2009]. A similar experiment using the true marginals $\hat{\mu}_{\text{data}}$ has also been conducted in Peluchetti [2021].

C Studying the forward Densities

C.1 Transition kernels

The transition kernels $p(z_0, \cdot)$ for the SDEs from Section 2 are of the from

$$q_0(z_0, \cdot) = \mathcal{N}(m_t(z_0), \Sigma_t),$$

where m_t and Σ_t are given in Table 1 for the Brownian Motion and the Ornstein-Uhlenbeck process. The form of the transition kernels for the CLD are more involved. They can be found in Dockhorn et al. [2021, Appendix B.1].

Lemma 5. *The marginal densities $p_t(x)$ of the SDEs treated in Section 2 depend smoothly on x and t.*

Proof. One can combine the form of m_t and Σ_t and the explicit representation of p_t in (9) to see this. More generally, for the Brownian motion and the OU-Process this is a result of the Hörmander theorem. For the CLD it is a result of hypocoercivity. $\square$

C.2 Form of the drift

We now prove that we can represent the drift as in (10).

Lemma 6. *Assume that p_t has the form (9), i.e.*

$$p_t(z) = \int_{\mathbb{R}^d} \mathcal{N}(z; m_t(z_0), \Sigma_t) \mu_{data}(z_0) \mathrm{d}z_0.$$

Then (10) holds true, i.e.

$$\nabla \log p_t = \frac{\nabla p_t(z)}{p_z(z)} = \Sigma_t^{-1}(z - \mathbb{E}[m_t(Z_0)|Z_t = z]).$$

Proof.

$$\nabla p_t(z)$$

$$= \frac{1}{\sqrt{\det(2\pi\Sigma_t)}} \int_{\mathbb{R}^d} \nabla \exp\left(-\frac{1}{2}(z - m_t(z_0))\Sigma_t^{-1}(z - m_t(z_0))\right) \mu_{\text{data}}(z_0)\mathrm{d}z_0$$

$$= \frac{1}{\sqrt{\det(2\pi\Sigma_t)}} \int_{\mathbb{R}^d} \Sigma_t^{-1}(z - m_t(z_0)) \exp\left(-\frac{1}{2}(z - m_t(z_0))\Sigma_t^{-1}(z - m_t(z_0))\right) \mu_{\text{data}}(z_0)\mathrm{d}z_0$$

$$= \Sigma_t^{-1} z p_t(z)$$

$$- \frac{1}{\sqrt{\det(2\pi\Sigma_t)}} \int_{\mathbb{R}^d} \Sigma_t^{-1} m_t(z_0) \exp\left(-\frac{1}{2}(z - m_t(z_0))\Sigma_t^{-1}(z - m_t(z_0))\right) \mu_{\text{data}}(z_0)\mathrm{d}z_0.$$

If we now divide everything by p_t it cancels in the first summand. In the second summand we get the formula for the conditional expectation (see, for example [Klenke, 2013, Section 8.2]). □

D Reverse SDEs: The general case

One can also treat more general forward SDEs than we did in Section 1. This leads to a more complicated form of the reverse SDE. Our Theorems do not use the specific form of the forward SDEs and therefore also hold in the general case. We denote the forward SDE by

$$\begin{aligned} \mathrm{d}X_t &= \beta(t, X_t)\mathrm{d}t + \sigma(t, X_t)\mathrm{d}W_t, \\ X_0 &\sim \mu_{\text{data}}, \end{aligned} \quad (16)$$

where μ_{data} is supported on $\mathcal{M} \subset \mathbb{R}^d$ and W_t is a $\mathbb{R}^r$ valued Brownian motion. The drift b maps from $\mathbb{R} \times \mathbb{R}^d$ to $\mathbb{R}^d$. The dispersion coefficient σ maps from $\mathbb{R} \times \mathbb{R}^d$ to the $d \times r$-matrices. The time-reversed process $Y_t := X_{T-t}$ is then a solution to

$$\begin{aligned} \mathrm{d}Y_t &= b(t, Y_t)\mathrm{d}t + \sigma(T - t, Y_t)\mathrm{d}B_t, \\ Y_0 &\sim q_0, \end{aligned} \quad (17)$$

with

$$b_i(t, y) = -\beta(T - t, y) + \frac{\sum_j \nabla_j(a_{ij}(T - t, y)p_{T-t}(y))}{p_{T-t}(y)}, \quad a(t, y) = \sigma(t, y)\sigma(t, y)^T,$$

and $q_0 = p_T$, see Haussmann and Pardoux [1986]. This simplifies to the case discussed in Section 1 if $r = d$, σ is is a multiple of the identity matrix and β is independent of the time t. In Assumption 1 we treat the case where the SDEs are of the form written-out in Section 1. The items $(i) - (iii)$ need to be replaced by their more general counterpart as found in Haussmann and Pardoux [1986, Section 2]. In the last item (iv), $\nabla \log p_t$ needs to be replaced by $\frac{\sum_j \nabla_j(a_{ij}(T-t, y)p_{T-t}(y))}{p_{T-t}(y)}$.

E Proofs

E.1 Proofs of the theorems

We now give proofs of our main results and briefly summarize the key steps in an intuitive way. In our study we would like to include the case when μ_{data} is degenerate and supported on a low-dimensional

substructure $\mathcal{M}$. As we have seen in Section **??**, this can lead to an exploding drift in the reverse SDE as $t \to T$. Nevertheless, in order to understand the properties of μ_{data}, it is crucial to study the properties of solutions to the reverse SDE at time $t = T$. This is where the main mathematical difficulties come from. The proofs are mostly independent of the specific form of the forward SDE and hold for more general forward/backward SDEs than those stated in Section 1, see Appendix D.

E.1.1 Theorem 1

We now proceed with proving Theorem 1.

Proof. Let P be the measure on $\Omega = C([0,T], \mathbb{R}^n)$ induced by the forward SDE (1) started in $p_0 = \mu_{\text{data}}$. P has marginals p_t. Denote by X_t the canonical projections $X_t(\omega) = \omega(t)$ for $\omega \in \Omega$. We define Q through

$$\frac{\mathrm{d}Q}{\mathrm{d}P}(\omega) = \frac{\mathrm{d}\mu_{\text{prior}}}{\mathrm{d}\pi_T}(\omega(T)).$$

By the data processing inequality we obtain (see [Liese and Vajda, 2006, Theorem 14]),

$$KL(q_T | \mu_{\text{data}}) \le KL(Q|P) = KL(\mu_{\text{prior}} | \pi_T). \tag{18}$$

It remains to prove that by running Q backwards we obtain a solution to (2) started in μ_{prior}. We denote the generator of the reverse SDE (2) by $\mathcal{L}$. Denote by Q^R and P^R the time reversals of Q and P. Our assumption are such that P^R is a Markov process solving the martingale problem for $\mathcal{L}$ (see [Haussmann and Pardoux, 1986, Theorem 2.1]). A short calculation shows that Q^R is still Markov (see, for example [Léonard, 2011, Proposition 4.2]). Furthermore for $f \in C_c^\infty(\mathbb{R}^n)$,

$$\mathbb{E}_{Q^R}\left[f(X_t) - f(X_s) - \int_s^t \mathcal{L}f(X_r)\mathrm{d}r | X_s \right]$$

$$= \frac{\mathbb{E}_{P^R}\left[\left(f(X_t) - f(X_s) - \int_s^t \mathcal{L}f(X_r)\mathrm{d}r \right) \frac{\mathrm{d}\mu_{\text{prior}}}{\mathrm{d}p_T}(X_0) | X_s \right]}{\mathbb{E}_{P^R}\left[\frac{\mathrm{d}\mu_{\text{prior}}}{\mathrm{d}p_T}(X_0) | X_s \right]}$$

$$= \frac{\mathbb{E}_{P^R}\left[\left(f(X_t) - f(X_s) - \int_s^t \mathcal{L}f(X_r)\mathrm{d}r \right) | X_s \right] \mathbb{E}_{P^R}\left[\frac{\mathrm{d}\mu_{\text{prior}}}{\mathrm{d}p_T}(X_0) | X_s \right]}{\mathbb{E}_{P^R}\left[\frac{\mathrm{d}\mu_{\text{prior}}}{\mathrm{d}p_T}(X_0) | X_s \right]}$$

$$= \mathbb{E}_{P^R}\left[\left(f(X_t) - f(X_s) - \int_s^t \mathcal{L}f(X_r)\mathrm{d}r \right) | X_s \right] = 0.$$

In the second equality we used the Markov property of P^R. In the last one we used that P^R solves the martingale problem for $\mathcal{L}$. Therefore also Q^R solves the martingale problem for $\mathcal{L}$. Denote by Y_t a solution to (2) on $[0, T)$. Since solutions to (2) are unique in law on $[0, S]$ for $S < T$ (see [Karatzas and Shreve, 2012, Section 5.2]) and the solutions are continuous, the law of Y is equal to Q^R on $[0, T)$. But the paths of Q^R are continuous on $[0, T]$. Therefore, Y can be extended to $[0, T]$, i.e. the limit $Y_T := \lim_{t \to T} Y_t$ exists almost surely and its distribution is equal to the T-time marginal distribution of Q^R, which is the 0-time marginal of Q. We denote the marginals of Q and P by Q_t and P_t. Since Q is absolutely continuous with respect to P, $Q_0 = \mu_{\text{sample}}$ is absolutely continuous with respect to $P_0 = \mu_{\text{data}}$. Analogously, if μ_{prior} and p_T are equivalent, then so are P and Q and therefore P_0 and Q_0. This proves (i).

(ii) is a consequence of the data processing inequality for f-divergences ([Liese and Vajda, 2006, Theorem 14]), analogous to (18). $\qquad \square$

The main idea of this proof is that we look at the forward SDE for X_t first. It induces a distribution $\mathbb{P}$ over all continuous paths in $\Omega = C([0,T], \mathbb{R}^d)$. If we reverse the time direction of this distribution on Ω, we get a solution to the reverse SDE, started in p_T. This reverse solution is well behaved as $t \to T$, since $Y_T = X_0$. The solution for a different initial condition $\mu_{\text{prior}} \ne p_T$ is obtained by reweighting $\mathbb{P}$. This does not change the qualitative behaviour of Y_T, which still exists and is well defined. We then use a uniqueness result to see that any solution of (17) inherits these properties.

 E.1.2   Theorem 2

641   *Proof.* Denote the space $C([0, T], \mathbb{R}^d)$ by Ω. We also define the natural filtration $\mathcal{F}_t = \sigma(x(s)|s \leq t)$.
642   We denote the distribution of Y on Ω by $\mathbb{P}$. We define $\tilde{\mathbb{P}}$ by reweighting $\mathbb{P}$ with Z_t on $\mathcal{F}_t$. By Girsanov
643   theorem (see [Karatzas and Shreve, 2012, Section 3.5]) we know that the canonical process under $\tilde{\mathbb{P}}$
644   is a solution to (6) on $[0, T)$. Since Z_t is uniformly integrable, its limit $Z := \lim_{t \to T} Z_t$ exists in L^1.
645   Furthermore $\tilde{\mathbb{P}}$ is absolutely continuous with respect to $\mathbb{P}$ on $\mathcal{F} = \sigma(\cup_{t<T} \mathcal{F}_t) = \sigma(x(t)|t < T)$ with
646   density Z. We define $x(T) := \lim_{t \to T} x(t)$. Then the event

$$\mathcal{A} = \{x(T) := \lim_{t \to T} x(t) \text{ exists and } x(T) \in \mathcal{M}\}$$

647   has probability 1 under $\mathbb{P}$ (see Theorem 1) and therefore also under $\tilde{\mathbb{P}}$. Furthermore, $x(T)$ is
648   measurable with respect to $\mathcal{F}$. Therefore the distributions of $x(T)$ under $\mathbb{P}$ and $\tilde{\mathbb{P}}$ are equivalent. The
649   canonical process under $\tilde{\mathbb{P}}$ is therefore a solution of (6), with the property that its time T-marginal is
650   well defined and equivalent to the time T-marginal of (2). We can use uniqueness in law on $[0, S)$ for
651   any $S < T$ and extend it to $[0, T]$ as in the proof of Theorem 1. This shows that every solution to (6)
652   has the desired properties.

653   Finally we show that if e is bounded, it fulfils Assumptions 2. We define by $H_t = \int_0^t \|e(s, \hat{Y}_s)\| \mathrm{d}s$.
654   Then there is a Brownian motion W_t such that we can write Z_t as

$$Z_t = \exp\left(W_{H_t} - \frac{1}{2}H_t\right).$$

655   Since e is bounded by M, H_t is bounded by TM. In particular, one can view $Z_t = \mathbb{E}[Z_{TM}|\mathcal{F}_{H_t}]$.
656   Therefore Z_t is uniformly integrable since it can be viewed as a family of conditional expectations.
657 $\square$

658   Here we essentially applied the Girsanov Theorem on $[0, T)$. Using the uniform integrability of the
659   Girsanov weights Z_t, we are able to extend it to $[0, T]$. Therefore, we can infer that the distribution
660   of Y and $\hat{Y}$ are actually equivalent on the whole path space $C([0, T], \mathbb{R}^d)$. In particular, their time
661   T-marginals will be equivalent too, which is the claim of the theorem.

662   **E.2   Proof of the Lemmas**

663   We start by proving Lemma 1.

664   *Proof.* The forward drifts are $\beta(x) = 0$, $\beta(x) = -\frac{\alpha}{2}x$ and $\beta(x, v) = (v, -x - 2v)$ for the Brownian
665   Motion, the OU-Process and the Critically Damped Langevin Dynamics respectively. In particular,
666   these are all linear maps and therefore fulfil conditions (i) and (ii) of Assumption 1.

667   We show in Appendix C.1 that $\log p_t$ is C^∞ in t and x for $t > 0$. Therefore we can integrate p_t
668   and its derivative over compact sets, implying that condition (iii) holds. Furthermore, the Hessian
669   w.r.t. (x, t) is continuous and obtains its maximum and minimum on the compact set $[S, T] \times B_N$,
670   where B_N is the ball of diameter N around the origin. Therefore the gradient $\nabla \log p_t$ is Lipschitz
671   on $[S, T] \times B_N$, which proves (iv). $\square$

672   We now prove Lemma 2.

673   *Proof.* We have that $X_0 \sim \mu_{\text{data}}$. Denote the mean and covariance of μ_{data} by a and C respectively.
674   We define

$$n_t = \mathcal{N}(m_t, V_t)$$

675   for some functions m_t and V_t. If V_t would not have full rank, n_t would be a degenerate distribution.
676   Since $p_t > 0$ almost everywhere for $t > 0$, the KL divergence from p_t to n_t would be infinite.
677   We can therefore restrict V_t to be an invertible matrix. We denote the entropy of p by H, $H(p) =$
678   $-\int \log(p)p\mathrm{d}x$.

We can write X_t as $X_t = X_0 + \sqrt{t}Z$ where $Z \sim \mathcal{N}(0, I_d)$. Now for the KL-divergence at time t it holds that

$$KL(p_t|n_t) = -H(p_t) - \int \log(n_t(x))p_t(x)\mathrm{d}x$$

$$= -H(p_t) + \frac{d}{2}\log(2\pi) + \frac{1}{2}\log\left(\det(V_t)\right) + \frac{1}{2}\int (x - m_t)^T V_t^{-1}(x - m_t)\, p_t(x)\mathrm{d}x.$$

We compute

$$\int (x - m_t)^T V_t^{-1}(x - m_t)\, p_t(x)\mathrm{d}x$$

$$= \mathbb{E}[(X_t - m_t)^T V_t^{-1}(X_t - m_t)] = \mathbb{E}[(X_0 + \sqrt{t}Z - m_t)^T V_t^{-1}(X_0 + \sqrt{t}Z - m_t)]$$

$$= \mathbb{E}[(X_0 - a)^T V_t^{-1}(X_0 - a)] + (a - m_t)^T V_t^{-1}(a - m_t) + t\mathbb{E}[Z^T V_t^{-1} Z]$$

$$= \mathrm{tr}(CV_t^{-1}) + t\,\mathrm{tr}(I_d V_t^{-1}) + (a - m_t)^T V_t^{-1}(a - m_t).$$

We added and subtracted a in the second equality. In the last equality we used that $\mathbb{E}[X^T AX] = \mathrm{tr}(\mathrm{Cov}(X)A)$ if X is a centred random variable. Substituting this into our prior calculation, we arrive at

$$KL(p_t|n_t)$$
$$= -H(p_t) + \frac{1}{2}\log\left(\det(2\pi V_t)\right) + \frac{1}{2}\left(\mathrm{tr}((C + tI_d)V_t^{-1}) + (a - m_t)^T V_t^{-1}(a - m_t)\right).$$

The mean function m_t only appears in the last term. Therefore, we decrease the KL-divergence by setting $m_t = a$. We now optimize for the covariance matrix V_t. The $H(p_t)$ term is independent of V_t. Consequently, to optimize the $KL(p_t|n_t)$ with respect to V_t, it is enough to optimize

$$L(V_t^{-1}) = \log\left(\det(V_t)\right) + \mathrm{tr}((C + tI_d)V_t^{-1})$$
$$= -\log\left(\det(V_t^{-1})\right) + \mathrm{tr}((C + tI_d)V_t^{-1}),$$

where we used that $\det(A^{-1}) = \det(A)^{-1}$ to rewrite the loss in terms of V_t^{-1}. Since $-\log(\det(\cdot))$ is convex on the positive semidefinite matrices and the other summands are linear, L is convex in V_t^{-1}. We take the gradient of L with respect to the Frobenius inner product and obtain

$$\nabla_{V_t^{-1}} L(V_t^{-1}) = -V_t + C + tI_d.$$

Setting the gradient to 0 has the unique solution

$$V_t = C + tI_d,$$

which is a local and global minimum due to the convexity of L.

If we restrict V_t to be of the form $V_t = v_t I_d$, the loss becomes

$$L(v_t^{-1}) = -d\log(v_t^{-1}) + v_t^{-1}\mathrm{tr}(C) + v_t^{-1}td.$$

By setting the derivative of this to zero, we arrive at

$$v_t = \mathrm{tr}(C)/d + t.$$

$\square$

We now proceed to prove Lemma 3. For that we first need the following proposition.

Proposition 1. *Let p_t be the time t-marginal of a Brownian motion started in μ_{data}. Assume that the support of μ_{data} is contained in a ball of radius M. Let $m_t = \mathbb{E}[\mu_{data}]$ and $C_t = Cov[\mu_{data}] + tI_d$ be the optimal mean and covariance operator from Lemma 2. Denote $n_t = \mathcal{N}(m_t, V_t)$. Then $KL(p_t|n_t) \to 0$ as $t \to \infty$.*

Proof. For $t > 0$, the KL-divergence is given by

$$KL(p_t|n_t) = -H(p_t) + \frac{1}{2}\log\left(\det(2\pi V_t)\right) + \frac{1}{2}\mathrm{tr}((C + tI_d)V_t^{-1})$$

$$= -H(p_t) + \frac{1}{2}\log\left(\det(2\pi V_t)\right) + \frac{d}{2} \qquad (19)$$

$$= -H(p_t) + \frac{d}{2}\log(2\pi) + \frac{1}{2}\log\left(\prod_{i=1}^{d}(c_i + d)\right) + \frac{d}{2}.$$

702 We used that

$$\det(C + tI_d) = \prod_{i=1}^{d}(c_i + t), \tag{20}$$

703 where c_i are the eigenvalues of $\mathrm{Cov}[\mu_{\mathrm{data}}]$.

704 Now,

$$-H(p_t) = \mathbb{E}_{X_t}[\log p_t(X_t)] = \mathbb{E}_{X_t}\left[\log\left(\mathbb{E}_{X_0}\left[(2\pi t)^{-d/2}\exp\left(-\frac{1}{2t}\|X_t - X_0\|^2\right)\right]\right)\right].$$

705 We bound

$$\exp\left(-\frac{1}{2t}\|X_t - X_0\|^2\right) \le \exp\left(-\frac{1}{2t}(\|X_t\|^2 - 2\|X_t\|M)\right).$$

706 We write X_t as $X_t = X_0 + \sqrt{t}Z$ where $Z \sim \mathcal{N}(0, I_d)$. Then,

$$\begin{aligned}
\log&\left(\mathbb{E}_{X_0}\left[(2\pi t)^{-d/2}\exp\left(-\frac{1}{2t}\|X_t - X_0\|^2\right)\right]\right)\\
&\le \log((2\pi t)^{-d/2}) - \frac{1}{2t}(\|X_t\|^2 - 2\|X_t\|M)\\
&\le -\frac{d}{2}\log(2\pi) - \frac{1}{2}\log(t^d) - \frac{1}{2t}\|X_t\|^2 + \frac{1}{t}(\|M\| + \sqrt{t}\|Z\|)M.
\end{aligned} \tag{21}$$

707 We use that

$$\mathbb{E}[\|X_t\|^2] = \mathbb{E}[\|X_0\|^2] + t\mathbb{E}[\|Z\|^2] = \mathbb{E}[\|X_0\|^2] + t\mathbb{E}[\|Z\|^2] = m_0^2 + v_0 + td,$$

708 where $v_0 = \mathbb{E}[\|X_0 - \mathbb{E}[\mu_{\mathrm{data}}]\|^2] = \mathrm{tr}(C_0)$. Since Z is centred and independent of X_t, the cross-term
709 $\mathbb{E}[\langle X_t, Z\rangle] = 0$ vanishes. We now take the expectation over the right hand side of (21) to obtain

$$\mathbb{E}[-\frac{1}{2t}(\|X_t\|^2 - \|X_t\|M)] \le \frac{1}{2t}\left(-td - m_0 - v_0 + 2\sqrt{t}\mathbb{E}[\|Z\|]M + 2M^2\right).$$

710 Putting it all together we have that

$$KL(p_t|n_t) \le \frac{1}{2}\log\left(\frac{\prod_{i=1}^{d}(c_i + t)}{t^d}\right) + \frac{1}{2t}\left(-m_0 - v_0 + 2\sqrt{t}\mathbb{E}[\|Z\|]M + 2M^2\right) \to 0,$$

711 as $t \to \infty$. $\qquad\square$

712 We are now ready to prove Lemma 3.

713 *Proof.* We again use (19). We now take the derivative of the KL-divergence using De Bruijn's identity
714 for multivariate random variables (see [Rioul, 2010], [Costa and Cover, 1984]):

$$\frac{\mathrm{d}}{\mathrm{d}t}H(p_t) = \frac{1}{2}\mathbb{E}_{p_t}[\|\nabla\log p_t(X)\|^2] = \frac{1}{2t^2}\mathbb{E}[\|X_t - \mathbb{E}[X_0|X_t]\|^2], \tag{22}$$

715 where we applied Lemma 6. Since $\mathbb{E}[X_0|X_t]$ is the L^2-orthogonal projection of X_0 to the $\sigma(X_t)$-
716 measurable random variables and X_t is $\sigma(X_t)$-measurable, we have that

$$\frac{1}{2t^2}\mathbb{E}[\|X_t - \mathbb{E}[X_0|X_t]\|^2] = \frac{1}{2t^2}\left(\mathbb{E}[\|X_t - X_0\|^2] - \mathbb{E}[\|X_0 - \mathbb{E}[X_0|X_t]\|^2]\right) = \frac{d}{2t} - \frac{A_t}{2t^2}.$$

717 For $A_t = \mathbb{E}[\|X_0 - \mathbb{E}[X_0|X_t]\|^2]$ it holds that

$$0 \le A_t \le \mathbb{E}[\|X_0\|^2] = \mathrm{tr}(\mathrm{Cov}(\mu_{\mathrm{data}})),$$

718 since the conditional expectation is a contraction in L^2. We use (20) again to arrive at

$$\frac{\mathrm{d}}{\mathrm{d}t}KL(p_t|n_t) = -\frac{1}{2}\left(\frac{d}{t} - \frac{A_t}{t^2}\right) + \frac{1}{2}\sum_{i=1}^{d}\frac{1}{t + c_i} \ge -\frac{1}{2}\frac{d}{t} + \frac{1}{2}\sum_{i=1}^{d}\frac{1}{t + c_i} =: h'(t), \tag{23}$$

where c_i are the eigenvalues of $\text{Cov}[\mu_{\text{data}}] + tI_d$. Integrating the right hand side we obtain

$$h(t) = \frac{1}{2} \log \left(\frac{\prod_{i=1}^{d}(c_i + t)}{t^d} \right)$$

as a possible antiderivative. Define $g(t) = h(t) - KL(p_t|n_t)$. Let us now assume that the support of μ_{data} is contained in a ball of radius M. Then, by Proposition 1, $\lim_{t\to\infty} KL(p_t|n_t) = \lim_{t\to\infty} h(t) = 0$ and therefore $\lim g(t) \to 0$. Equation (23) implies that $g'(t) \leq 0$ for all $t > 0$. Assume that there is a s and an $\epsilon > 0$ such that $g(s) \leq -\epsilon < 0$. Since $g'(t) \leq 0$ for all $t > 0$ we can then conclude that $g(t) \leq -\epsilon < 0$ for all $t \geq s$. In particular, $g(t) \not\to 0$ as $t \to \infty$, which is a contradiction. Therefore the statement of the Lemma holds for μ_{data}.

Now let μ_{data} be any measure. Let X be a random variable that is distributed according to μ_{data} and Z be a standard normal random variable, $Z \sim \mathcal{N}(0, I_d)$. Then $X + \sqrt{t}Z$ has distribution p_t. We define $X^M := X1_{\{\|X\| \leq M\}}$ and $X_t^M = X^M + \sqrt{t}Z$. Denote the distribution of X_t^M by p_t^M. The distribution of X^M is supported on a ball of radius M. Furthermore, $X^M + \sqrt{t}Z \to X + \sqrt{t}Z$ almost surely and therefore $p_t^M \to p_t$ weakly for any t. By lower semi-continuity of the KL-divergence,

$$KL(p_t|n_t) \leq \liminf_{M\to\infty} KL(p_t^M|n_t) \leq \liminf_{M\to\infty} \frac{1}{2} \log \left(\frac{\prod_{i=1}^{d}(c_i^M + t)}{t^d} \right), \tag{24}$$

where c_i^M are the eigenvalues of $\text{Cov}[\mu_{\text{data}}^M] + tI_d$. Since $|X^M| \leq |X|$ and X has first and second moments one can apply dominated convergence to see that the covariance matrices converge, $\text{Cov}(\mu_{\text{data}}^M) \to \text{Cov}(\mu_{\text{data}})$. Especially

$$\lim_{M\to\infty} \prod_{i=1}^{d}(c_i^M + t) = \lim_{M\to\infty} \det(\text{Cov}(\mu_{\text{data}}^M)) \to \det(\text{Cov}(\mu_{\text{data}})) = \prod_{i=1}^{d}(c_i + t).$$

Combining this with (24) concludes the proof. $\qquad\qquad\square$