Archimedes Meets Privacy:
On Privately Estimating Quantiles in High Dimensions Under Minimal Assumptions

Abstract

The last few years have seen a surge of work on high dimensional statistics under privacy constraints, mostly following two main lines of work: the "worst case" line, which does not make any distributional assumptions on the input data; and the "strong assumptions" line, which assumes that the data is generated from specific families, e.g., subgaussian distributions. In this work we take a middle ground, obtaining new differentially private algorithms with polynomial sample complexity for estimating quantiles in high-dimensions, as well as estimating and sampling points of high Tukey depth, all working under very mild distributional assumptions.

From the technical perspective, our work relies upon deep robustness results in the convex geometry literature, demonstrating how such results can be used in a private context. Our main object of interest is the (convex) floating body (FB), a notion going back to Archimedes, which is a robust and well studied high-dimensional analogue of the interquantile range of a distribution. We show how one can privately, and with polynomially many samples, (a) output an approximate interior point of the FB – e.g., "a typical user" in a high-dimensional database – by leveraging the robustness of the Steiner point of the FB; and at the expense of polynomially many more samples, (b) produce an approximate uniform sample from the FB, by constructing a private noisy projection oracle.

1 Introduction

Computing statistics of large, complex high-dimensional datasets under privacy constraints is a fundamental challenge in modern data science. In this work, we study *the sample complexity* of several different tasks related to estimating the quantiles of a d-dimensional distribution $\mathcal{D}$, from having sample access to it, under privacy constraints. Our first focus is on estimating quantiles of fixed marginals. For a random vector $X \in \mathbb{R}^d$, the quantiles of X along direction θ are the quantiles of the (one-dimensional) marginal $\langle X, \theta \rangle$, which we denote by $Q_q(\langle X, \theta \rangle)$ for $q \in [0, 1]$. Our second focus is on estimating a convex region called the floating body of a d-dimensional distribution, an analogue of the 1-dimensional interquantile range exhibiting rich mathematical properties, investigated over decades of research in convex geometry and sharing connections with the work of Archimedes (see, e.g., the survey [NSW19]). Formally, the q-floating body of a random vector $X \in \mathbb{R}^d$ is given by

$$F_q(X) = \bigcap_{\theta \in \mathbb{S}^{d-1}} \left\{ x \in \mathbb{R}^d : \langle x, \theta \rangle \leq Q_q(\langle X, \theta \rangle) \right\}, \tag{1}$$

and in statistical language corresponds to the points of so-called *Tukey-depth* at least $1 - q$ [NSW19]. When clear from context, we also denote by $F_q(\mathcal{D})$ the floating body of the random vector $X \sim \mathcal{D}$.

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

Similar to their 1-dimensional analogues, the quantiles in high-dimensions are preferred to other descriptive statistics and location estimators due to their robustness properties [Hub81, MR97] and high breakdown points [DG07, Alo03, TP16]. As opposed to other statistics, marginal quantiles can be estimated under the mere minimal distributional assumption of having non-trivial mass around the location of the quantiles (see e.g. [AMB19, Lemma 3]). Moreover, the floating body can also be estimated in the Hausdorff distance for log-concave measures and stable laws [AR20].

Besides statistical guarantees, an essential property of an estimator in modern data science is to be private. For instance, many large healthcare datasets contain sensitive patient information [DE13, ZLZ+17, FLJ+14, YKÖ17] and it is crucial for an estimator to balance the trade-off between protecting the privacy of the input while producing accurate statistical results.

Differential privacy (DP) is the leading notion in quantifying privacy guarantees of a randomized algorithm over an input dataset. In this work, we focus on algorithms satisfying *pure* DP guarantees. Given a parameter $\varepsilon > 0$, ε-differential privacy guarantees that the probability of a given output cannot change more than a multiplicative factor e^ε after the arbitrary change of the input data of one data item (e.g., a single user record in a large database). More generally, ε-DP can be defined as follows with respect to the Hamming distance $d_H(X, X') := |\{i \in [n] | X_i \neq X'_i\}|$, defined for two n-tuples of samples $X = (X_1, \ldots, X_n), X' = (X'_1, \ldots, X'_n) \in (\mathbb{R}^d)^n$.

Definition 1 (ε-DP). A randomized algorithm $\mathcal{A}$ is ε-differential private if for all subsets $S \in \mathcal{F}$ of the output space $(\Omega, \mathcal{F})$ and n-tuples of samples $X_1, X_2 \in (\mathbb{R}^d)^n$, it holds that

$$\mathbb{P}(\mathcal{A}(X_1) \in S) \leq e^{\varepsilon d_H(X_1, X_2)} \mathbb{P}(\mathcal{A}(X_2) \in S). \tag{2}$$

While there has been a lot of work on differential privacy, it usually analyzes algorithms under worst-case assumptions. The most closely related to our work is the line of research on privately outputting an interior point in high dimensions for a worst-case input [BMNS19, KMST20, SS21, KSS20]. Only recently there has been a lot of statistical work analyzing differentially private estimators under distributional assumptions for various statistical tasks [BCS15, DJW18, BCSZ18b, KSSU19, KLSU19, SU19, ZKKW20, CKS20, TVGZ20, KSU20]. On the topic of the present work, a few years ago [TVGZ20] established the exact trade-off between privacy and accuracy of median estimation in one-dimension. One of the main motivations of the present work is to make a first step towards understanding the high-dimensional counterparts of these privacy-accuracy tradeoffs, by proposing the floating body as a natural estimate under privacy constraints in high dimensions. In particular, the convexity of the floating body allows us to invoke powerful robustness theorems, originating from decades of research in convex geometry. It is this combination, of classical theorems with the modern framework of DP, which yields several new and non-trivial results.

1.1 Our Contributions

The minimal assumptions. It is known that even in one dimension, private quantile estimation is impossible for arbitrary measures [TVGZ20]. Thus, some assumptions are required. We make an arguably natural assumption, which is a high-dimensional analogue of the one-dimensional minimal distributional assumption considered in the aforementioned work. In short, we make the following assumptions: (a) the quantiles are bounded, (b) a mild non-degeneracy assumption to ensure the q-floating body has non-empty interior, and (c), (d) are mild concentration assumptions.

Definition 2. We say that a distribution $\mathcal{D}$ supported on $\mathbb{R}^d$ has a q-admissible law with parameters $R_{\max}, R_{\min}, r, L > 0$, if, for every $\theta \in \mathbb{S}^{d-1}$, the following four conditions hold:

    1. $Q_q(\langle X, \theta \rangle) \in [-R_{\max}, R_{\max}]$.

    2. For some $c \in \mathbb{R}^d$, it holds that $|Q_q(\langle X, \theta \rangle) - \langle c, \theta \rangle| \geq R_{\min}$.

    3. The density f_θ of $\langle X, \theta \rangle$ satisfies $f_\theta(t) \geq L$ whenever $t \in [Q_q(\langle X, \theta \rangle) - r, Q_q(\langle X, \theta \rangle) + r]$.

    4. If $\{X_i\}_{i=1}^n$ are *i.i.d.* as $\mathcal{D}$, then with high probability $\|X_i\|_2 \leq \text{poly}(n, d)$, for all $i \in [n]$.[1]

The set of all such distributions is denoted by $A_q(R_{\max}, R_{\min}, r, L)$.

---

[1]We remark that condition 4 is purely technical and, since the degree of the polynomial is arbitrary, almost does not restrict generality.

Private many-quantiles estimation. Let us introduce a useful convention: we identify a fixed vector $X = (X_1, \ldots, X_n) \in (\mathbb{R}^d)^n$ with a random vector in $\mathbb{R}^d$, chosen uniformly from $(X_1, \ldots, X_n)$. In other words we identify a sample X with its empirical distribution. Now, in [AR20] it was established that for any symmetric log-concave measure $\mathcal{D}$, if one has n i.i.d. samples $X = (X_1, \ldots, X_n) \in (\mathbb{R}^d)^n$ from $\mathcal{D}$, then the empirical measure on the samples satisfies for $Y \sim \mathcal{D}$ with probability at least 0.9,

$$\delta_q(X, Y) := \sup_{\theta \in \mathbb{S}^{d-1}} |Q_q(\langle X, \theta \rangle) - Q_q(\langle Y, \theta \rangle)| \leq \alpha, \tag{3}$$

for some $n = \tilde{O}(d/\alpha^2)$. Here, and throughout the paper, $\tilde{O}$ means we omit logarithmic factors, in all parameters. One can easily generalize the above result for any q-admissible law (as described in Definition 2) (see Supplementary material).

Note that for any subset of M directions, simply taking the empirical q-quantiles of the projection, for each of the M directions, works under the standard concentration results with $n = \tilde{O}(M/\alpha^2)$ samples (using $\tilde{O}(1/\alpha^2)$ samples per direction). We note that for each specific direction $\tilde{O}(1/\alpha^2)$ samples are necessary (see e.g. [TVGZ20, Proposition 3.7]). The above results lead us to an interesting high-dimensional phenomenon: without privacy constraints, *simultaneously* estimating $M > d$ marginal quantiles in high-dimensions requires less samples than naively projecting and calculating the empirical quantile on each direction.

Our first contribution is to reveal the private analogue of the above observation. Consider the task of private estimation of M quantiles, i.e. of $Q_q(\langle Y, \theta_i \rangle), Y \sim \mathcal{D}$ for M different directions $(\theta_i)_{i=1}^M$. The "naive" approach would be, as before, to first project along the different directions, but then, instead of calculating the (non-private) empirical quantiles of the projected points, one would apply the optimal private algorithm of [TVGZ20] to each projection, separately.[2] This would lead to the guarantee in (3) with sample complexity (assuming, for simplicity, *here and throughout the contribution section* all parameters besides M, ε, α are constant)

$$n = O\left(\frac{M}{\alpha^2} + \frac{M}{\varepsilon \alpha}\right). \tag{4}$$

Our first main result is an improved private rate for the estimation of M quantiles which appropriately privatizes the high-dimensional result of [AR20].

Theorem 3 (informal)**.** *There exists an ε-differentially private algorithm that draws*

$$n = \tilde{O}\left(\frac{d}{\alpha^2} + \frac{M}{\varepsilon \alpha}\right)$$

samples from any admissible distribution $Y \sim \mathcal{D}$ and produces an estimate $\hat{m}$ that satisfies $\max_{i=1,\ldots,M} |Q_q(\langle Y, \theta_i \rangle) - \hat{m}_i| \leq \alpha$ *with probability at least* 0.9.

Notice that the rate described in Theorem 3 is always at least as good as the rate in (4), and that it is strictly better when $M > d$ and $\varepsilon > \alpha$. This constitutes indeed a private analogue of the high-dimensional phenomenon mentioned above; privately estimating $M > d$ can require much less samples than simply projecting and applying the one-dimensional machinery.

Private interior point. Next, we attempt to privately produce point estimates of the floating body $F_q(Y)$ itself. Our first task is to output an interior point of $F_q(Y)$, a relatively easy task without privacy concerns [AR20], with a clear statistical motivation of outputting a "typical datapoint" in high-dimensions. We show that with polynomial in d samples one can achieve such a guarantee with small error. There has been closely related (but not fully comparable) work on this task in the worst case regime, which we discuss in depth in Section 1.2.

For our result we leverage the Steiner point of the floating body, denoted $S(F_q(Y))$ (see (7) for the definition). The Steiner point has been widely studied in the context of *Lipschitz selection* [BL00]. In the problem of Lipschitz selection, one looks to consistently select a point from the interior of each

---

[2]Note that the algorithm from [TVGZ20] is technically only designed for median estimation, but with the natural modifications the same algorithm and analysis can be straightforwardly extended for q-quantile estimation.

convex body, in a way which is robust to perturbations of the body. The Steiner point is celebrated for being the optimal Lipschitz selector in this setting. Just recently, this construction proved to be instrumental in the resolution of the *chasing nested convex sets* problem [BKL$^+$20, Sel20, AGTG21], a problem in online learning, related to Lipschitz selection. It is precisely this optimal robustness that makes the Steiner point insensitive to individual sample points, a desirable property in the design of DP algorithms. To the best of our knowledge, this is the first time the Steiner point and its properties have been used it to build differentially private estimators, and we believe it might be useful in other similar problems.

Theorem 4 (informal). *There exists an ε-differentially private algorithm that draws*

$$n = \tilde{O}\left(\frac{d}{\alpha^2} + \frac{d^{2.5}}{\varepsilon\alpha}\right)$$

samples from any admissible distribution $Y \sim \mathcal{D}$ and produces an estimate $\hat{m} \in \mathbb{R}^d$ that satisfies $\|\hat{m} - S(F_q(Y))\|_2 \leq \alpha$ with probability at least 0.9.

In comparison, the best known dependence of n on the dimension in the worst case literature is of order $\tilde{O}(d^4)$ for pure ε-DP [KSS20] and of order $\tilde{O}(d^{2.5})$ for the weaker notion of approximate (ε, δ)-DP [BMNS19] [3] . Thus, Theorem 4 achieves essentially the same dependence as the best known approximate DP result while enjoying the stronger guarantees of pure DP. We note again however that the results are not fully comparable; see Section 1.2 for a thorough discussion.

Private sampling from the floating body. Outputting an, a-priori, arbitrary point from the floating body comes with possible drawbacks, for example biasing the output point towards or away from its boundary. This can lead to inaccurate statistical conclusions if a more detailed description of the "typical datapoints" is needed. For this reason, we undertake the harder task of outputting a *uniform sample*.

The *sampling from convex bodies* problem has a long and rich history with many deep results, see [KLS95, KLS97, LV06] for some prominent examples. In many cases, the computation preformed by the sampling algorithm reduces to iteratively projecting points into the convex body, along the trajectory of an appropriate Markov chain, [BEL18, Leh21].

It is precisely the idea described above which serves as the working engine behind our private sampling algorithm. To cope with privacy constraints, we capitalize on known robustness properties of the projection operator (see (8)) on convex bodies, as proved in [AW93]. Remarkably, when the projected point is fixed and one considers the convex body as the variable, the projection operator is known to be 1/2-Hölder with respect to the Hausdorff distance. This allows us to build a noisy projection oracle for the floating body, in a private fashion. Combining the noisy oracle with known results leads to a new sampling algorithm, which operates under pure differential privacy, and with guarantees in the 1-Wasserstein distance, W_1 (see the appendix for the definition).

Theorem 5 (informal). *Let $\mu_{F_q(Y)}$ be the uniform measure on the floating body. There exists an ε-differentially private algorithm that draws*

$$n = \tilde{O}\left(\frac{d^2}{\alpha^4} + \frac{d^4}{(\varepsilon\alpha)^2}\right)$$

samples from any admissible distribution $Y \sim \mathcal{D}$, and whose output, $\hat{\mu}_{F_q(Y)}$, satisfies $\frac{1}{\sqrt{d}}W_1(\hat{\mu}_{F_q(Y)}, \mu_{F_q(Y)}) \leq \alpha$.

Remark that the $\frac{1}{\sqrt{d}}$ normalizing factor in Theorem 5 serves as the correct scale for the W_1 metrics which typically scales as square-root of the dimension, see [Leh21] for further discussion on this.

1.2 Further Comparison with [BMNS19, KSS20]

The most closely related work to our point estimators of the floating body (Theorems 4 and 5) are by Beimel et al. [BMNS19] and Kaplan et al. [KSS20]. Both of these works address under worst-case input the private *interior point* problem: namely, given a (worst case) set of n points $x_1, \ldots, x_n$ in

---

[3]see Supplementary material for a definition and comparison between the different privacy notions.

d dimensions, the task is to privately output a point that lies within the convex hull of these points. While the non-private analogue is trivial (since one can always output, say, x_1), the problem becomes interestingly non-trivial when privacy considerations are introduced. To achieve such guarantees, the set of points is assumed to be a subset of a finite grid of possible points U^d (where U is some finite universe). This is a necessary condition for the existence of successful private estimators under worst-case assumptions [BNS13, BNSV15]), whereas this assumption is not required in our case. On the other hand, due to the use of the extension lemma, our algorithms do not have any explicit bound on the running time (see Section 1.3 below), unlike the situation in the worst case works, whose running time is generally exponential in d.

The first work [BMNS19] solves the private center point problem under approximate (ε, δ)-differential privacy, a weaker privacy guarantee than ε-differential privacy, assuming that the dataset is of size at least some $n = \tilde{O}(d^{2.5} \log^*(|U|))$ (in the $\tilde{O}$ term here we suppress dependencies in ε, δ, and lower order terms). The second work [KSS20] solves the problem under pure differential privacy (as in our paper) for $n = \tilde{O}(d^4 \log |U|)$, and obtains an improved $O(n^d)$ running time using $O(d^4 \log |U|)$ samples under approximate DP. The $n = \tilde{O}(d^4 \log |U|)$ bound is currently the best known sample complexity under pure DP for this worst-case task. In both [BMNS19] and [KSS20], the authors' approach is to output a point of high Tukey-depth with respect to the set of the n points, which can be interpreted as outputting an element of the q-floating body of the n points for some appropriate, and perhaps data-dependent, value of q.

While our considered settings are similar but incomparable, we recall that our Theorem 4 obtains pure ε-differentially private guarantees that for any q outputs a point of the q-floating body with $\tilde{O}(d^{2.5})$ samples. Finally, we note that matching the sample complexity of the best (ε, δ)-approximate private algorithm with an ε-private algorithm has been challenging in the privacy literature, e.g., in the context of high-dimensional Gaussian mean estimation [KLSU19, KSU20], and in fact provably impossible in the worst-case context for many problems of interest; see [BMNS19] for more details.

1.3 General Approach

Our approach towards building all ε-private algorithms in this work is based on an appropriate use of a Lipschitz extension tool called the Extension Lemma [BCSZ18a, BCSZ18b]. This lemma was also the main tool behind the optimal private median estimator in [TVGZ20]. One of the main hurdles in constructing sample-efficient ε-differentially private algorithms under distributional assumptions is that the privacy constraint needs to hold for all input data-sets, while the accuracy is based on the behavior of the algorithm on "with-high-probability" or typical data-sets. The Extension Lemma completely resolves this issue and allows the algorithm designer to focus on ensuring privacy solely on the typical data-sets[4] Yet, the present work . Then a (privacy-preserving) Lipschitz Extension argument takes care of extending the private algorithm on typical-inputs to a private algorithm on all possible inputs, while maintaining the original algorithm's output on the typical inputs (see Proposition 1 for more details). Leveraging this tool, our results are based on a two-step procedure.

1. First we consider the (non-private) estimator of interest. For example, this is the Steiner point of the floating body for Theorem 4 and the projection of a point to the floating body for Theorem 5. Then we define a *typical set* of possible input datasets $X = (X_1, \ldots, X_n) \in (\mathbb{R}^d)^n$ which is realized with high probability over all inputs drawn from admissible distributions. Following this we establish Lipschitzness properties of the non-private estimator seen as a function of the input X *with respect to the Hamming distance between two inputs* of the (non-private) estimator, when defined only on inputs from the typical set. Such robustness properties can be non-trivial and to do so we use interesting tools from convex geometry to establish them for the Steiner point [BL00], and for the projection of a point [AW93]. Notably, none of these estimators is (non-trivially) Lipschitz with respect to the Hamming distance of the input, unless the input is on the typical set. Using now the Lipschitzness properties combined with classic differential privacy ideas we can "privatize" the estimator by applying the (flattened) Laplace mechanism [TVGZ20], while ensuring high accuracy.

---

[4]While this work is entirely focused on sample complexity guarantees, it is worth mentioning that the Extension Lemma comes in principle with no explicit termination time guarantee (see [BCSZ18a] and the Supplementary material).

2. With the Extension Lemma, we extend our estimator to be defined and private on all possible inputs. The Extension Lemma also guarantees that the estimator remains identical when the input is from the typical set. These allow to obtain a globally private algorithm with the same accuracy guarantee.

2 Getting Started

2.1 Extension Lemma

Let us consider an arbitrary ε-differentially private algorithm defined on n samples in $\mathbb{R}^d$ as input and belonging in some set $\mathcal{H} \subseteq (\mathbb{R}^n)^d$. Then the Extension Lemma guarantees that the algorithm *can be always extended* to a 2ε-differentially private algorithm defined for arbitrary input data in $(\mathbb{R}^n)^d$ with the property that if the input data belongs in $\mathcal{H}$, the distribution of output values is exactly the same with the original algorithm. We note that the result in [BCSZ18a] is "generic", in the sense of applying to any input metric space and output probability space, but here we present it for simplicity only on inputs from $\mathbb{R}^n$ and equipped with the Hamming distance d_H. Formally the result is as follows.

Proposition 1 (The Extension Lemma, Proposition 2.1, [BCSZ18a])**.** *Let $\hat{\mathcal{A}}$ be an ε-differentially private algorithm designed for input from $\mathcal{H} \subseteq (\mathbb{R}^d)^n$ with arbitrary output measure space $(\Omega, \mathcal{F})$. Then there exists a randomized algorithm $\mathcal{A}$ defined on the whole input space $(\mathbb{R}^d)^n$ with the same output space which is 2ε-differentially private and satisfies that for every $X \in \mathcal{H}$, $\mathcal{A}(X) \overset{d}{=} \hat{\mathcal{A}}(X)$.*

2.2 Quantiles and Floating Bodies

For X a random variable over $\mathbb{R}$, and $q \in (0, 1)$, the q-*quantile* of X, which we denote $Q_q(X)$, is the smallest number in $\mathbb{R}$ satisfying $\mathbb{P}(X \leq Q_q(X)) \geq q$. Our main object of interest is a high-dimensional analogue of the quantile, called the convex floating body, defined in (1). Notice, that for $q < \frac{1}{2}$, $F_q(X)$ is necessarily empty and unfortunately it may sometimes by empty for $q \geq \frac{1}{2}$. Assumption 2. in Definition 2 ensures that $F_q(\mathcal{D}) \neq \emptyset$ and moreover that $F_q(\mathcal{D})$ contains a ball of radius $R_{\min}$, centered at c (see Supplementary material for more details).

2.3 Useful Distances

The following pseudo-metric simply measures the maximum difference between the q-quantiles of two (random) vectors.

Definition 6 (q-distance)**.** Let $q \in (\frac{1}{2}, 1)$ and let $A \subset \mathbb{S}^{d-1}$. If X and Y are two random vectors in $\mathbb{R}^d$, their q-distance, with respect to A is defined by $\delta_q(X, Y; A) := \sup_{\theta \in A} |Q_q(\langle X, \theta \rangle) - Q_q(\langle Y, \theta \rangle)|$. If $A = \mathbb{S}^{d-1}$, we abbreviate $\delta_q(X, Y; A) = \delta_q(X, Y)$.

In the sequel, we will consider two types of set A, in Definition 6. First, finite sets of directions, which correspond to privately estimating a finite (but potentially large) number of quantiles. Second the whole set of possible directions $A = \mathbb{S}^{d-1}$. In this case where $\delta_q(X, Y; A) = \delta_q(X, Y)$ bounds on $\delta_q(X, Y)$ are strong enough to yield allow estimation of high-dimensional statistics of $F_q(X)$, such as the Steiner point of the body or the projection operator to the body.

To further elaborate on this remark, let us recall that there is another natural convex geometric way to measure distances between convex bodies, *the Hausdorff distance*. For two convex bodies $K, K' \subset \mathbb{R}^d$, their Hausdorff distance is $\delta_{\text{Haus}}(K_1, K_2) := \inf\{t > 0 | K \subset K' + tB_d \text{ and } K' \subset K + tB_d\}$, where B_d is the unit Euclidean ball in $\mathbb{R}^d$. The following connection between δ_{Haus} and δ_q holds.

Lemma 1. [Bru18, Lemma 5] *Let $q \in (\frac{1}{2}, 1)$ and let X and Y be two random vectors in $\mathbb{R}^d$. Suppose that $F_q(X)$ contains a ball of radius $R_{\min}$ and is contained in a ball of radius $R_{\max}$, for some $R_{\max}, R_{\min} > 0$. If $\delta_q(X, Y) \leq \frac{R_{\min}}{2}$ then, $\delta_{\text{Haus}}(F_q(X), F_q(Y)) \leq \frac{3R_{\max}}{R_{\min}} \delta_q(X, Y)$.*

3 General Hölder Queries

We are now ready to present our main result. In the introduction, we highlighted three results of our work, all of which follow from a simple meta-theorem resulting in a differentially private algorithm for querying generic Hölder statistics of the floating body with respect to the δ_q norm. Since our result is general, it requires certain notation.

Approximate hölder queries Let $p \in [1, \infty]$, $h \in (0, 1]$, and $M \in \mathbb{N}$. Denote now by $\mathcal{C}_d$ the space of all convex bodies in $\mathbb{R}^d$, and assign M to be the dimension of the output space, equipped with the ℓ_p-norm. We say that a map $f : \mathcal{C}_d \to \mathbb{R}^M$ is h-Hölder (with constant $K > 0$, and with respect to the set $A \subset \mathbb{S}^{d-1}$), or simply Lipschitz when $h = 1$, if,

$$\|f(F_q(X)) - f(F_q(Y))\|_p \leq K \delta_q(X, Y; A)^h, \tag{5}$$

where X and Y are random vectors. It turns out that many of the queries we study, like the Steiner point, are Hölder with respect to the Hausdorff distance, not δ_q metric. In light of Lemma 1, it will sometimes be convenient to restrict the domain in which (5) holds. Thus, we say that f is *approximate h-Hölder*, with respect to $r_1, R_1, R_2 > 0$ if for all X, Y, such that for some $a \in \mathbb{R}^d$ it holds $B(a, R_1) \subseteq F_q(X) \subseteq B(a, R_2)$ and $\delta_q(F_q(X), F_q(Y)) \leq r_1$, (5) holds. Since our private algorithm will be extended from a restricted algorithm on a typical set (recall the plan from Section 1.3) using Proposition 1, there will be no loss of privacy when considering approximate Hölder functions, as long as the desired conditions hold with high probability over the sample.

The main result. We are now prepared to state our main theorem. All results mentioned in the preceding sections will follow by working with suitable approximate Hölder functions.

Theorem 7. *Fix $q \in (1/2, 1)$, $A \subset \mathbb{S}^{d-1}$, and assume that $\mathcal{D} \in A_q(R_{\max}, R_{\min}, r, L, B)$, with $f : \mathcal{C}_d \to \mathbb{R}^M$ an approximate h-Hölder function with constant $K > 0$, with respect to A. Then, there is an ε-differentially private algorithm $\mathcal{A}$ which for input $X = (X_1, \ldots, X_n)$, sampled i.i.d from $\mathcal{D}$, satisfies for all $\alpha < \frac{\min\{r, R_{\min}\}}{2}$ that $\mathbb{P}(\|\mathcal{A}(X) - f(F_q(\mathcal{D}))\|_p \leq \alpha) \geq 1 - \delta$, for some n satisfying*

$$n = \tilde{O}\left(\frac{K^{2/h}\left(d + \log\left(\frac{4}{\delta}\right)\right)}{\alpha^{2/h} L^2} + \frac{WK^{1/h}(\log\left(\frac{1}{\delta}\right)^{1/h} M^{1/h})}{(\varepsilon\alpha)^{1/h} L} + \frac{WM^{1/h}}{(\varepsilon \min\{r, R_{\min}\})^{1/h} L} \right),$$

where $W = \min\{d, \log|A|\} + \log\left(\frac{1}{\delta}\right)$.

The obtained rate may seem complicated; this is to be expected, given the generality of our results and the number of parameters. In the next section we demonstrate several concrete uses of Theorem 7, which show how the rate simplifies in various interesting settings.

4 Applications

Here we describe the main applications of Theorem 7. As mentioned we apply the theorem using suitable approximate Hölder functions.

Simultaneous estimation of quantiles Fix $q \in (0, 1)$ and let $A \subset \mathbb{S}^{d-1}$, with $|A| = M$. Define the multiple M-query function which on a random vector X, equals $f_A(X) = \{Q_q(\langle X, \theta \rangle)\}_{\theta \in A}$.

It is immediately seen that f_A is a Lipschitz function[5], with constant 1, in the L_∞ ($p = \infty$) norm:

$$\|f_A(X) - f_A(Y)\|_\infty = \max_{\theta \in A} |Q_q(\langle X, \theta \rangle) - Q_q(\langle Y, \theta \rangle)| = \delta_q(X, Y; A). \tag{6}$$

We thus have the following result.

Corollary 8. *Let $\mathcal{D} \in A_q(R_{\max}, R_{\min}, r, L)$ be an admissible measure on $\mathbb{R}^d$. Then, there is an ε-differentially private algorithm $\mathcal{A}$ which for input $X = (X_1, \ldots, X_n)$, sampled i.i.d from $\mathcal{D}$, satisfies for all $\alpha < \frac{\min\{r, R_{\min}\}}{2}$, $\mathbb{P}(\|\mathcal{A}(X) - f_A(\mathcal{D})\|_\infty \leq \alpha) \geq 0.9$, for*

$$n = \tilde{O}\left(\frac{d}{\alpha^2 L^2} + \frac{M}{\varepsilon L \alpha} + \frac{M}{\varepsilon L \min\{r, R_{\min}\}} \right).$$

---

[5]Strictly speaking, f_A is not a function of the floating body, but of the sample itself. However, with a trivial adaption, it still fits nicely within our framework.

Proof. The observation in (6) shows that f_A is a Lipschitz function with constant 1. The result now follows by invoking Theorem 7 for the L_∞ norm, with $K, h = 1$, $\delta = 0.1$, and $W = \log(M)$. $\square$

Privately returning an interior point: the Steiner point. Given Theorem 7 it will be enough to show that one can select a point from a convex body in a Lipschitz way. This naturally leads to the Steiner point, a widely studied object in Lipschitz selection.

If $K \subset \mathbb{R}^d$ is a convex body, we define its Steiner point by

$$S(K) := d \int_{\mathbb{S}^{d-1}} \theta h_K(\theta) d\sigma, \tag{7}$$

where σ is the normalized Haar measure on $\mathbb{S}^{d-1}$, and the support function of K, $h_K : \mathbb{S}^{d-1} \to \mathbb{R}$, is defined by $h_K(\theta) = \max_{x \in K} \langle x, \theta \rangle$.

The following result follows from well-known results in convex geometry and from Lemma 1.

Lemma 2. *The Steiner point $S : \mathcal{C}_d \to \mathbb{R}^d$ is an approximate Lipschitz function, with constant $\sqrt{d} \frac{3 R_{\min}}{R_{\min}}$, which satisfies $S(K) \in K$ for every convex body K.*

We immediately get the following corollary to Theorem 7.

Corollary 9. *Let $\mathcal{D} \in A_q(R_{\max}, R_{\min}, r, L)$ be an admissible measure on $\mathbb{R}^d$. Then, there is an ε-differentially private algorithm $\mathcal{A}$ which for input $X = (X_1, \ldots, X_n)$, sampled i.i.d from $\mathcal{D}$, satisfies for all $\alpha < \frac{\min\{r, R_{\min}\}}{2}$, $\mathbb{P}(\|\mathcal{A}(X) - S(F_q(\mathcal{D}))\|_2 \le \alpha) \ge 0.9$, for some*

$$n = \tilde{O}\left(d^2 \frac{R_{\max}^2}{R_{\min}^2 \alpha^2 L^2} + d^{2.5} \frac{R_{\max}}{R_{\min}} \left(\frac{1}{\varepsilon L \alpha} + \frac{1}{\varepsilon L \min\{r, R_{\min}\}}\right)\right).$$

Proof. Consider the Steiner point $S : \mathcal{C}_d \to \mathbb{R}^d$. Lemma 2 states that S is an approximate Lipschitz function with constant $\sqrt{d} \frac{3 R_{\min}}{R_{\min}}$, for every $X, Y \in \mathcal{H}_C$. The result now follows by invoking Theorem 7 for the Euclidean norm, with $K = \sqrt{d} \frac{3 R_{\min}}{R_{\min}}$, $M = d$, $h = 1$, $\delta = 0.1$, and $W = \tilde{O}(d)$. $\square$

Private projection and sampling. Let $K \subset \mathbb{R}^d$ be a convex body. Define the projection operator, $P_K : \mathbb{R}^d \to \mathbb{R}^d$ by,

$$P_K(x) = \arg \min_y \{y \in K | \|y - x\|\}. \tag{8}$$

We shall establish that, for admissible distributions $\mathcal{D}$ and for any point x, one can privately estimate $P_{F_q(\mathcal{D})}(x)$ with polynomially many samples. This follows by coupling a classic result in convex geometry, [AW93, Proposition 5.3], concerning the stability of projections with Lemma 1.

Lemma 3. *Fix $x \in \mathbb{R}^d$ with $\|x\| \le R_{\max}$ and consider $P_K(x) : \mathcal{C}_d \to \mathbb{R}^d$. Then $P_K(x)$ is an approximate $\frac{1}{2}$-Hölder function with constant $\frac{\text{poly}(R_{\max})}{R_{\min}}$.*

Now, to sample from the body K, for $\eta > 0$, we define the following discretized Langevin process:

$$X_{t+1} = P_K(X_t + \eta g_t), \quad X_0 = 0,$$

where $\{g_t\}_{t \ge 0}$ are *i.i.d.* standard Gaussians. It is well known (see for example [Leh21, Theorem 2]) that this process mixes rapidly, in the Wasserstein distance. By applying the known results about the mixing time of the Langevin process, and by taking account of the inherent noise introduced by the privacy constraints, we prove the following result.

Corollary 10. *Let $\mathcal{D} \in A_q(R_{\max}, R_{\min}, r, L)$ be an admissible measure on $\mathbb{R}^d$ and let U_q be a random vector which is uniform on $F_q(\mathcal{D})$. Then, there is an ε-differentially private algorithm $\mathcal{A}$ which for input $X = (X_1, \ldots, X_n)$, sampled i.i.d from $\mathcal{D}$, satisfies for all $\alpha < \frac{\min\{r, R_{\min}\}}{2}$, $\frac{1}{\sqrt{d}} W_1(\mathcal{A}(X), U_q) \le \alpha$, for some*

$$n = \tilde{O}\left(\text{poly}\left(\frac{R_{\max}}{R_{\min}}\right)\left(\frac{d^2}{\alpha^4 L^2} + \frac{d^4}{\varepsilon^2 \alpha^2 L} + \frac{d^4}{\varepsilon^2 \min\{r, R_{\min}\}^2 L}\right)\right).$$

5 Proof Sketch of Theorem 7

To provide intuition to the reader, we now briefly describe how one can use the two-stage procedure from Section 1.3 to obtain Theorem 7, from which all our applications follow. We fix an approximate h-Hölder query $f(F_q(Y)), Y \sim \mathcal{D}$ and assume for simplicity that it is Hölder w.r.t. $\delta_q(X, Y)$, that is $A = \mathbb{S}^{d-1}$.

Obtaining a "good" (non-private) estimator Our first step is to obtain a (non-private) estimate of the query. For that, we sample n independent points $\{X_i\}_{i=1}^n \sim \mathcal{D}$, and define X the uniform empirical measure over $(X_1, \ldots, X_n)$, for which we compute $f(F_q(X))$. In terms of accuracy, by an appropriate generalization of the arguments in [AR20] to apply for any admissible distribution we have $\delta_q(X, Y) \leq (\alpha/K)^{1/h}$ for some $n = \tilde{O}(dK^{1/h}/\alpha^{2/h})$. Notice now that, by admissibility, and the discussion in Section 2.2 it can be easily checked that $F_q(Y)$ satisfies the necessary geometric condition; it contains a ball and is contained in a ball, both of "controlled" radius. Hence, the approximate Hölderness implies $\|f(F_q(X)) - f(F_q(Y))\|_p \leq K\delta_q(X, Y)^h = \alpha$.

Designing the private estimator on a typical set. Our goal now turns to design a private, yet accurate estimate, of $f(F_q(X))$. To do this, in principle we would desire $f(F_q(X))$ to be Lipschitz (with a "good" constant, say Λ_f) with respect to the Hamming distance on X. Indeed, with such a property the Laplace mechanism [DR14] produces an ε-DP estimate of $f(F_q(X))$ with error Λ_f/ε. Unfortunately, such a property cannot exist in general; as mentioned, in many cases of interest, f is only known to be Lipschitz with respect to δ_q under certain conditions; for example, when $F_q(X)$ contains a ball and is contained in a ball. So, we must impose some restrictions on the input X.

For this reason, we design an appropriate "typical" high-probability set $\mathcal{H} \subseteq (\mathbb{R}^d)^n$, such that $f(F_q(X))$, restricted to $\mathcal{H}$ is Lipschitz, with respect to the Hamming distance on the input X. The properties of the typical set will allow that for all $X, Y \in \mathcal{H}$ unless the Hamming distance $d_H(X, Y)$ is "large" it holds (a) $\|f(F_q(X)) - f(F_q(Y))\|_p \leq K\delta_q(X, Y)^h$ and (b) $\delta_q(X, Y) \leq \frac{C}{Ln}d_H(X, Y)$. These results together imply $\|f(F_q(X)) - f(F_q(Y))\|_p \leq KC^h/(L^h n^h)d_H(X, Y)$. An application of a variant of the Laplace mechanism, called the flattened Laplacian mechanism, [BCSZ18b, TVGZ20], now applies, and gives an ε-DP, yet accurate, estimate of $f(F_q(X))$ when $X \in \mathcal{H}$. The accuracy guarantee of the flattened Laplacian mechanism results from a careful multivariate integral calculation.

Let us now describe the typical set, $\mathcal{H}$. It is built as the intersection of two conditions, each one happening with high-probability. The first condition is that, in every direction, the quantiles are appropriately bounded; a condition which is satisfied by merit of the empirical distribution of X being close to the population distribution $\mathcal{D}$ in the δ_q distance. As discussed, this condition enforces property (a) above. The second condition is more intricate, as it requires that, in every direction, the quantile is close, in several scales, to a non-negligible fraction of the points. This reduces the sensitivity of the quantile to the individual sample points and we use it to establish property (b). The proof that the second condition holds with high-probability *for any admissible distribution* is a combination of a net-argument and an appropriate use of the one-dimensional result in [TVGZ20, Lemma B.3].

Extension Lemma Finally, a direct application of the Extension Lemma 1 extends the flattened Laplacian mechanism from the previous paragraph to a 2ε-DP estimator on the whole $(\mathbb{R}^d)^n$, while remaining the same on inputs from $\mathcal{H}$. Since $\mathcal{H}$ happens with high probability, the result follows.

6 Conclusion

Our work suggests several future directions. First, our estimators come with no polynomial termination time guarantee since they are based on applying the generic Extension Lemma. Yet, in the one dimensional case, [TVGZ20] showed that the Extension Lemma can be computed in polynomial time when applied to the empirical median on an appropriate typical set. Given the analogies between our works, it is an important open problem to see if the high-dimensional estimators we propose can also be implemented in polynomial-time. Second, our work leverages classical robust convex geometric tools to construct private algorithms, which allow to demonstrate non-trivial high-dimensional phenomena. It will be interesting to see if these newly introduced tools in the privacy literature can be further used to reveal other non-trivial high-dimensional results.

References

[AGTG21]   CJ Argue, Anupam Gupta, Ziye Tang, and Guru Guruganesh. Chasing convex bodies with linear competitive ratio. *Journal of the ACM (JACM)*, 68(5):1–10, 2021.

[Alo03]   Greg Aloupis. Geometric measures of data depth. In *Data Depth: Robust Multivariate Analysis, Computational Geometry and Applications*, volume 72 of *DIMACS Series in Discrete Mathematics and Theoretical Computer Science*, pages 147–158, 2003.

[AMB19]   Marco Avella-Medina and Victor-Emmanuel Brunel. Differentially private sub-gaussian location estimators. *arXiv preprint arXiv:1906.11923*, 2019.

[AR20]   Joseph Anderson and Luis Rademacher. Efficiency of the floating body as a robust measure of dispersion. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 364–377. SIAM, 2020.

[AW93]   Hédy Attouch and Roger J.-B. Wets. Quantitative stability of variational systems. II. A framework for nonlinear conditioning. *SIAM J. Optim.*, 3(2):359–381, 1993.

[BCS15]   Christian Borgs, Jennifer T. Chayes, and Adam Smith. Private graphon estimation for sparse graphs. In *Proceedings of the 28th International Conference on Neural Information Processing Systems - Volume 1*, NIPS'15, page 1369–1377, Cambridge, MA, USA, 2015. MIT Press.

[BCSZ18a]   Christian Borgs, Jennifer Chayes, Adam Smith, and Ilias Zadik. Private algorithms can always be extended. *arXiv preprint arXiv:1810.12518*, 2018.

[BCSZ18b]   Christian Borgs, Jennifer Chayes, Adam Smith, and Ilias Zadik. Revealing network structure, confidentially: Improved rates for node-private graphon estimation. In *2018 IEEE 59th Annual Symposium on Foundations of Computer Science (FOCS)*, pages 533–543. IEEE, 2018.

[BEL18]   Sébastien Bubeck, Ronen Eldan, and Joseph Lehec. Sampling from a log-concave distribution with projected Langevin Monte Carlo. *Discrete Comput. Geom.*, 59(4):757–783, 2018.

[BKL$^+$20]   Sébastien Bubeck, Bo'az Klartag, Yin Tat Lee, Yuanzhi Li, and Mark Sellke. Chasing nested convex bodies nearly optimally. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1496–1508. SIAM, 2020.

[BL00]   Yoav Benyamini and Joram Lindenstrauss. *Geometric nonlinear functional analysis. Vol. 1*, volume 48 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2000.

[BMNS19]   Amos Beimel, Shay Moran, Kobbi Nissim, and Uri Stemmer. Private center points and learning of halfspaces. In *Proceedings of the Thirty-Second Conference on Learning Theory*, volume 99 of *Proceedings of Machine Learning Research*, pages 269–282. PMLR, 2019.

[BNS13]   Amos Beimel, Kobbi Nissim, and Uri Stemmer. Private learning and sanitization: Pure vs. approximate differential privacy. In *Approximation, Randomization, and Combinatorial Optimization. Algorithms and Techniques*, pages 363–378. Springer, 2013.

[BNSV15]   Mark Bun, Kobbi Nissim, Uri Stemmer, and Salil P. Vadhan. Differentially private release and learning of threshold functions. In *IEEE 56th Annual Symposium on Foundations of Computer Science, FOCS 2015*, pages 634–649, 2015.

[Bru18]   Victor-Emmanuel Brunel. Methods for estimation of convex sets. *Statist. Sci.*, 33(4):615–632, 2018.

[CKS20]   Clément L Canonne, Gautam Kamath, and Thomas Steinke. The discrete Gaussian for differential privacy. *Advances in Neural Information Processing Systems*, 33:15676–15688, 2020.

[DE13] Fida Kamal Dankar and Khaled El Emam. Practicing differential privacy in health care: A review. *Trans. Data Priv.*, 6(1):35–67, 2013.

[DG07] PL Davies and Ursula Gather. The breakdown point—examples and counterexamples. *REVSTAT Statistical Journal*, 5(1):1–17, 2007.

[DJW18] John C. Duchi, Michael I. Jordan, and Martin J. Wainwright. Minimax optimal procedures for locally private estimation. *J. Amer. Statist. Assoc.*, 113(521):182–201, 2018.

[DR14] Cynthia Dwork and Aaron Roth. The algorithmic foundations of differential privacy. *Foundations and Trends in Theoretical Computer Science*, 9(3–4):211–407, 2014.

[FLJ$^+$14] Matthew Fredrikson, Eric Lantz, Somesh Jha, Simon Lin, David Page, and Thomas Ristenpart. Privacy in pharmacogenetics: An end-to-end case study of personalized warfarin dosing. In *Proceedings of the 23rd USENIX Security Symposium*, pages 17–32, 2014.

[Hub81] Peter J. Huber. *Robust statistics*. Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1981.

[KLS95] Ravi Kannan, László Lovász, and Miklós Simonovits. Isoperimetric problems for convex bodies and a localization lemma. *Discrete & Computational Geometry*, 13(3):541–559, 1995.

[KLS97] Ravi Kannan, László Lovász, and Miklós Simonovits. Random walks and an $o^*(n^5)$ volume algorithm for convex bodies. *Random Structures & Algorithms*, 11(1):1–50, 1997.

[KLSU19] Gautam Kamath, Jerry Li, Vikrant Singhal, and Jonathan Ullman. Privately learning high-dimensional distributions. In *Conference on Learning Theory, COLT 2019*, volume 99 of *Proceedings of Machine Learning Research*, pages 1853–1902, 2019.

[KMST20] Haim Kaplan, Yishay Mansour, Uri Stemmer, and Eliad Tsfadia. Private learning of halfspaces: Simplifying the construction and reducing the sample complexity. In *Advances in Neural Information Processing Systems*, volume 33, pages 13976–13985, 2020.

[KSS20] Haim Kaplan, Micha Sharir, and Uri Stemmer. How to Find a Point in the Convex Hull Privately. In *36th International Symposium on Computational Geometry (SoCG 2020)*, volume 164 of *Leibniz International Proceedings in Informatics (LIPIcs)*, pages 52:1–52:15, 2020.

[KSSU19] Gautam Kamath, Or Sheffet, Vikrant Singhal, and Jonathan Ullman. Differentially private algorithms for learning mixtures of separated gaussians. In *Advances in Neural Information Processing Systems 32: Annual Conference on Neural Information Processing Systems 2019, NeurIPS 2019*, pages 168–180, 2019.

[KSU20] Gautam Kamath, Vikrant Singhal, and Jonathan Ullman. Private mean estimation of heavy-tailed distributions. *CoRR*, abs/2002.09464, 2020.

[Leh21] Joseph Lehec. The Langevin monte carlo algorithm in the non-smooth log-concave case. *arXiv preprint arXiv:2101.10695*, 2021.

[LV06] László Lovász and Santosh Vempala. Simulated annealing in convex bodies and an $o^*(n^4)$ volume algorithm. *Journal of Computer and System Sciences*, 72(2):392–417, 2006.

[MR97] M. Markatou and E. Ronchetti. *Robust inference: the approach based on influence functions*, volume 15 of *Handbook of Statist.* North-Holland, Amsterdam, 1997.

[NSW19] Stanislav Nagy, Carsten Schütt, and Elisabeth M Werner. Halfspace depth and floating body. *Statistics Surveys*, 13:52–118, 2019.

[Sel20] Mark Sellke. Chasing convex bodies optimally. In *Proceedings of the Fourteenth Annual ACM-SIAM Symposium on Discrete Algorithms*, pages 1509–1518. SIAM, 2020.

[SS21] Menachem Sadigurschi and Uri Stemmer. On the sample complexity of privately learning axis-aligned rectangles. In *Advances in Neural Information Processing Systems*, volume 34, pages 28286–28297, 2021.

[SU19] Adam Sealfon and Jonathan Ullman. Efficiently estimating erdos-renyi graphs with node differential privacy. *CoRR*, abs/1905.10477, 2019.

[TP16] Pingfan Tang and Jeff M. Phillips. The robustness of estimator composition. In *Advances in Neural Information Processing Systems 29: Annual Conference on Neural Information Processing Systems 2016*, pages 929–937, 2016.

[TVGZ20] Christos Tzamos, Emmanouil-Vasileios Vlatakis-Gkaragkounis, and Ilias Zadik. Optimal private median estimation under minimal distributional assumptions. *Advances in Neural Information Processing Systems*, 33:3301–3311, 2020.

[YKÖ17] Buket Yüksel, Alptekin Küpçü, and Öznur Özkasap. Research issues for privacy and security of electronic health services. *Future Gener. Comput. Syst.*, 68:1–13, 2017.

[ZKKW20] Huanyu Zhang, Gautam Kamath, Janardhan Kulkarni, and Zhiwei Steven Wu. Privately learning Markov random fields. In *Proceedings of the 37th International Conference on Machine Learning*, ICML'20, 2020.

[ZLZ$^+$17] Jiajun Zhang, Xiaohui Liang, Zhikun Zhang, Shibo He, and Zhiguo Shi. Re-dpoctor: Real-time health data releasing with w-day differential privacy. In *2017 IEEE Global Communications Conference, GLOBECOM 2017, Singapore, December 4-8, 2017*, pages 1–6. IEEE, 2017.