Learning Weighted Representations for Generalization Across Designs

Abstract

Predictive models that generalize well under distributional shift are often desirable and sometimes crucial to machine learning applications. One example is the estimation of treatment effects from observational data, where a subtask is to predict the effect of a treatment on subjects that are systematically different from those who received the treatment in the data. A related kind of distributional shift appears in unsupervised domain adaptation, where we are tasked with generalizing to a distribution of inputs that is different from the one in which we observe labels. We pose both of these problems as prediction under a shift in *design*. Popular methods for overcoming distributional shift are often heuristic or rely on assumptions that are rarely true in practice, such as having a well-specified model or knowing the policy that gave rise to the observed data. Other methods are hindered by their need for a pre-specified metric for comparing observations, or by poor asymptotic properties. In this work, we devise a family of algorithms to address these issues, by jointly learning a representation and a re-weighting of observed data. We show that our algorithms minimize an upper bound on the generalization error under design shift, and verify the effectiveness of this approach in causal effect estimation.

1 Introduction

A long-term goal in artificial intelligence is for agents to learn how to *act*. This endeavor relies on accurately predicting and optimizing for the outcomes of actions, and fundamentally involves estimating counterfactuals—what would have happened if the agent acted differently? In many applications, such as the treatment of patients in hospitals, experimentation is infeasible or impractical, and we are forced to learn from *biased*, observational data. Doing so requires adjusting for the *distributional shift* between groups of patients that received different treatments. A related kind of distributional shift arises in domain adaptation, the goal of which is to learn predictive models for a target domain, observing ground truth only in a source domain. In this work, we pose both domain adaptation and treatment effect estimation as special cases of prediction across shifting *designs*, referring to changes in both action policy and feature domain. We separate policy from domain as we wish to make *causal* statements about the policy, but not about the domain.

Prediction of outcomes under interventions has a long history in clinical applications, most often to verify the efficacy of a medical treatment A by asking "how would the patient have responded under an alternative treatment, B"? To answer this question, the gold-standard solution continues to be the randomized controlled trial, in which treatment is administered randomly to patients selected according to some critera. However, most clinical trials are underpowered for estimating fine-grained individual-level effects. Moreover, many interventions cannot be performed in a randomized fashion due to ethical or practical considerations. Fortunately, vast amounts of observational (non-randomized) clinical data is available in the form of electronic health records, lab tests, and insurance data. Learning from observational data to predict the counterfactual outcome under treatment B for a patient who received A, one must adjust for the fact that treatment A was systematically given to patients of different characteristics from those who received treatment B. We call this *predicting under a shift in policy*. Furthermore, if all of our observational data comes from hospital P, but we wish to predict counterfactuals for patients in hospital Q, with a population that differs from P, an additional source of distributional shift is at play. We call this *a shift in domain.*

Together, we refer to the combination of domain and policy as the *design*. The design for which we observe ground truth is called the *source*, and the design of interest the *target*.

Broadly speaking, existing methods for generalization under distributional shift rely on one or more of the following concepts: a) learning shift-invariant representations of data (Ajakan et al., 2014), b) sample re-weighting (Shimodaira, 2000), or c) unit matching and assignment (Kallus, 2016).

Representation learning approaches attempt to extract only information from the input that is invariant to a change in design *and* predictive of the variable of interest. Such representations are typically learned by fitting deep neural networks in which activations of deeper layers are regularized to be distributionally similar across designs (Ajakan et al., 2014; Long et al., 2015). With some exceptions, see e.g. Long et al. (2015), the theoretical properties of these models are not well understood.

Re-weighting methods correct for distributional shift by assigning higher weight, in the objective of interest, to samples from the source design that are representative of the target design, often using importance sampling. This idea has been well studied in, for example, the causal inference (Rosenbaum & Rubin, 1983), domain adaptation (Shimodaira, 2000) and reinforcement learning (Precup et al., 2001) literature. For example, in causal effect estimation, importance sampling is equivalent to re-weighting units by the probability of observed treatments. Re-weighting, with knowledge of importance sampling weights, often leads to asymptotically unbiased estimators of the target outcome, but may suffer from high variance in finite samples (Swaminathan & Joachims, 2015).

A considerably larger hurdle, however, is that the optimal weights are rarely known in practice. There are variety of methods to learn these weights. Weights can be estimated as the inverse of estimated propensities (Rosenbaum & Rubin, 1983; Freedman & Berk, 2008) but this plug-in approach can lead to highly unstable estimates. More stable methods attempt to learn weights by minimizing distributional distance metrics (Gretton et al., 2009; Kallus, 2016; Kallus; Zubizarreta, 2015). Closely related, matching (Stuart, 2010) produces weights by finding units in the source design that are close in some metric to units in the target design. Specifying either distributional or unit-wise metric is a challenging task, especially if the input space is high-dimensional where no metric incorporating all features can ever be made small. This necessitates heuristics such as first performing variable selection and then finding a matching in the selected covariates. Unfortunately, the theoretical properties of such procedures are largely unknown.

We address the problem of *generalization under design shift*—prediction of outcomes for previously unseen units, in a design observed without supervision, based on observational, labeled samples from a different design. We show that this problem encompasses both standard counterfactual prediction and unsupervised domain adaptation, as well as the more general case of predicting outcomes under domain shift. To overcome limitations of existing methods, we propose to combine the intuition behind shift-invariant representation learning and re-weighting methods by jointly learning a representation Φ of the input space, and a weighting function $w(\Phi)$ to minimize a) the re-weighted empirical risk and b) a re-weighted measure of distributional shift between designs. By letting w depend on Φ we alleviate the problem of choosing a metric by which to optimize unit weights, as Φ extracts information predictive of the outcome. At the same time, our theory guarantees that this will remain a uniform bound on true risk in the target design. This leads to a general algorithmic framework, and a natural bound on the generalization error under design shift. We make the following contributions.

Main contributions

- We bring together two techniques used to overcome distributional shift between designs—re-weighting and representation learning, with complementary robustness properties, generalizing existing methods based on either technique.

- We give finite-sample generalization bounds for prediction under design shift, assuming that we have access to neither importance sampling weights *nor* a well-specified model, and develop an algorithmic framework to minimize these bounds.

- We propose a neural network architecture that jointly learns a representation of the input and a weighting function to improve balance across changing settings.

- We apply our proposed algorithm to the task of predicting causal effects from observational data, achieving state-of-the art results on a widely used benchmark.

2   PREDICTING OUTCOMES UNDER DESIGN SHIFT

The goal of this work is to accurately predict outcomes of interventions $T \in \mathcal{T}$ in contexts $X \in \mathcal{X}$ drawn from a *target design* $p_\pi(X, T)$. The outcome of intervening with $t \in \mathcal{T}$ is the *potential outcome* $Y(t) \in \mathcal{Y}$ (Imbens & Rubin, 2015, Ch. 1–2), which has a stationary distribution $p_t(Y \mid X)$ given context X. For example, in the classical binary setting $Y(1)$ represents the outcome under treatment and $Y(0)$ the outcome under control. The target design consists of two components: the *target policy* $p_\pi(T \mid X)$, which describes how one intends to map observations of contexts (such as patient prognostics) to interventions (such as pharmacological treatments) and the *target domain* $p_\pi(X)$, which describes the population of contexts to which the policy will be applied. The target design is known to us only through m unlabeled samples $(x_1, t_1), \ldots, (x_m, t_m)$ from $p_\pi(X, T)$. Outcomes are only available to us in labeled samples from a source domain: $(x'_1, t'_1, y'_1), \ldots, (x'_n, t'_n, y'_n)$, where (x'_i, t'_i) are draws from a source design $p_\mu(X, T)$ and $y'_i = y'_i(t'_i)$ is a draw from $p_T(Y \mid X)$, corresponding only to the *factual* outcome $Y(T)$ of the treatment administered. Like the target design, the source design consists of a domain of contexts for which we have data and a policy, which describes the (unknown) historical administration of treatment in the data. Only the factual outcomes of the treatments administered are observed, while the counterfactual outcomes $y'_i(t)$ for $t \neq t'_i$ are, naturally, unobserved. Our focus is the *observational* or *off-policy* setting, in which interventions in the source design are performed non-randomly as a function of X, $p_\mu(T \mid X) \neq p_\mu(T)$. This encapsulates both the covariate shift often observed between treated and control populations in observational studies and the covariate shift between the domain of the study and the domain of an eventual wider intervention. Examples of this problem are plentiful: in addition to the example given in the introduction, consider predicting the return of an advertising policy based on the historical results of a different policy, applied to a different population of customers. We stress that we are interested in the *causal effect* of an intervention T on Y, conditioned on X. As such, we cannot think of X and T as a single variable.

Without additional assumptions, it is impossible to deduce the effect of an intervention based on observational data alone (Pearl, 2009), as it amounts disentangling correlation and causation. Crucially, for any unit i, we can observe the potential outcome $y'_i(t)$ of at most one intervention t. To make prediction of outcomes and causal effects possible, we make the following standard assumptions.

Assumption 1 (Consistency, ignorability and overlap). *For any unit i, assigned to intervention t_i, we observe $Y_i = Y(t_i)$. Further, $\{Y(t)\}_{t \in \mathcal{T}}$ and the data-generating process $p_\mu(X, T, Y)$ satisfy*

$$\text{strong ignorability: } \{Y(t)\}_{t \in \mathcal{T}} \perp\!\!\!\perp T \mid X, \quad \text{and overlap: } \Pr_{p_\pi}(p_\mu(T \mid X) > 0) = 1.$$

Assumption 1 is a sufficient condition for *causal identifiability* (Rosenbaum & Rubin, 1983). Ignorability is also known as the *no hidden confounders* assumption, indicating that all variables that cause both T and Y are assumed to be measured. Under ignorability therefore, any domain shift in $p(X)$ cannot be due to variables that causally influence T and Y, other than through X. Under Assumption 1, potential outcomes equals conditional expectations: $\mathbb{E}[Y(t) \mid X = x] = \mathbb{E}[Y \mid X = x, T = t]$, and we may predict $Y(t)$ by regression. We further assume common domain support, $\forall x \in \mathcal{X} : p_\pi(X = x) > 0 \Rightarrow p_\mu(X = x) > 0$. Finally, we adopt the notation $p(x) := p(X = x)$.

2.1   RE-WEIGHTED RISK MINIMIZATION

A common approach to predicting potential outcomes is to learn hypotheses $f : \mathcal{X} \times \mathcal{T} \to \mathcal{Y}$ such that $f(x, t)$ approximate $\mathbb{E}[Y \mid X = x, T = t]$. Recall that under Assumption 1, this conditional expectation is equal to the (possibly counterfactual) potential outcome $Y(t)$, conditioned on X. In this work, the goal is to ensure that these hypotheses are accurate under a design p_π that deviates from the data-generating process, p_μ. This is unlike the supervised learning setting in that the distribution p_π to which we wish to generalize is different from that which generated the training data, p_μ. Our setting also differs from that normally considered in causal effect estimation, in that we focus on prediction for unseen units, rather than counterfactual estimation for observed units.

We measure the (in)ability of f to predict outcomes under π, using the expected risk,

$$R_\pi(f) := \mathbb{E}_{x,t,y \sim p_\pi}[\ell_f(x, t, y)] \tag{1}$$

based on a sample from μ, $D_\mu^n = \{(x_i, t_i, y_i) \sim p_\mu; i = 1, ..., n\}$. Here, $\ell_f(x, t, y) := L(f(x, t), y)$ is an appropriate loss function, such as the squared loss, $L(y, y') := (y - y')^2$ or the log-loss,

depending on application. As outcomes under the target design p_π are not observed, even through a finite sample, we cannot directly estimate (1) using the empirical risk under p_π. A common way to resolve this is to use *importance sampling* (Shimodaira, 2000)—the observation that if p_μ and p_π have common support, with $w^*(x,t) = p_\pi(x,t)/p_\mu(x,t)$,

$$R_\mu^{w^*}(f) := \mathbb{E}_{x,t,y \sim p_\mu}[w^*(x,t)\ell_f(x,t,y)] = R_\pi(f) \,. \tag{2}$$

Hence, with access to w^*, an unbiased estimator of $R_\pi(f)$ may be obtained by reweighting the (factual) empirical risk under μ,

$$\hat{R}_\mu^{w^*}(f) := \frac{1}{n}\sum_{i=1}^n w^*(x_i,t_i)\ell_f(x_i,t_i,y_i) \,. \tag{3}$$

Unfortunately, importance sampling weights can be very large when p_π is large and p_μ small, resulting in large variance in $\hat{R}_\mu^w(f)$ (Swaminathan & Joachims, 2015). More importantly, $p_\pi(x,t)$ is rarely known in practice, and neither is w^*. In principle, however, any re-weighting function w with the following property yields a valid risk under the re-weighted distribution p_μ^w.

Definition 1. *A function $w : \mathcal{X} \times \mathcal{T} \to \mathbb{R}_+$ is a valid re-weighting of p_μ if*

$$\mathbb{E}_{x,t \sim p_\mu}[w(x,t)] = 1 \ \text{ and } \ p_\mu(x,t) > 0 \Rightarrow w(x,t) > 0.$$

We denote the re-weighted density $p_\mu^w(x,t) := w(x,t)p_\mu(x,t)$.

A natural candidate in place of w^* is an estimate $\hat{w}^*$ of the importance sampling weights. In this work, we adopt a different strategy, learning re-weighting functions w from observational data, that minimize an upper bound on the risk under π.

Prediction under shifting design is of interest in several tasks, perhaps the most common of which are domain adaptation and counterfactual estimation. In this work, we focus on the latter, observing that the estimation of causal effects is an important special case.

2.2 CONDITIONAL TREATMENT EFFECT ESTIMATION

An important special case of our setting is when treatments are binary, $T \in \{0,1\}$, often interpreted as treating ($T = 1$) or not treating ($T = 0$) a unit, and the domain is fixed across designs, $p_\mu(x) = p_\pi(x)$. This is the classical setting for estimating treatment effects—the effect of choosing one intervention over another (Morgan & Winship, 2014).[1] The effect of an intervention T in context X, is measured by the *conditional average treatment effect* (CATE),

$$\tau(x) = \mathbb{E}\left[Y(1) - Y(0) \mid X = x\right].$$

Predicting τ for unobserved units involves prediction of both potential outcomes. In a clinical setting, this is necessary to assess which medication should be administered to a certain individual. Historically, the (population) *average treatment effect*, ATE $= \mathbb{E}_{x \sim p}[\tau(x)]$, has received comparatively much more attention (Rosenbaum & Rubin, 1983), but is inadequate for personalized decision making. Using predictors $f(x,t)$ of potential outcomes $Y(t)$ in contexts $X = x$, we can estimate the CATE by $\hat{\tau}(x) = f(x,1) - f(x,0)$ and measure the quality using the mean squared error (MSE),

$$\text{MSE}_\pi(\hat{\tau}) = \mathbb{E}_p\left[(\hat{\tau}(x) - \tau(x))^2\right] \tag{4}$$

In Section 4, we argue that estimating CATE from observational data requires overcoming distributional shift with respect to the treat-all and treat-none policies, in predicting each respective potential outcomes, and show how this can be used to give generalization bounds for CATE.

3 RELATED WORK

The central problem in learning predictors that generalize across designs, is to adjust for the distributional shift arising from changes in policy and domain. These types of shift have typically been

---

[1] Notions of causal effects exist also for the non-binary case, but these are not considered here.

treated separately, the former in counterfactual estimation (Morgan & Winship, 2014) and reinforcement learning (Precup et al., 2001), and the latter in domain adaptation.

Our proposed framework fits into the family of regression or covariate adjustment methods (Pearl, 2009; Rubin, 2005) for counterfactual estimation. These methods focus on estimating expected potential outcomes, under the assumption that relevant variables[2] have been measured. A large body of work has shown that under the additional assumption of having a well-specified model, various regression methods are asymptotically consistent (Chernozhukov et al., 2017; Athey & Imbens, 2016; Belloni et al., 2014; Van Der Laan & Rubin, 2006). However, consistency results like these provide no insight in the case of model misspecification.

Under model misspecification, ordinary regression may suffer from (unnecessarily) large bias when generalizing across designs. A common way to alleviate this is the *importance sampling* principle, see Section 2. This idea is used in so-called propensity-score methods (Austin, 2011), that uses treatment assignment probabilities (propensities) to re-weight samples for causal effect estimation, and more generally in re-weighted regression, performed to reduce bias under a misspecified model (Swaminathan & Joachims, 2015). A major drawback of these methods is the assumption that the design density is known. To address this, others (Gretton et al., 2009; Kallus, 2016), have proposed *learning* weights to minimize the distributional distance between samples under p_π and p_μ, but rely on specifying the data representation a priori.

In high-dimensional settings, kernel methods such as the re-weighting method of Gretton et al. (2009) are often outperformed by using representations learned from data (Bengio et al., 2013). Johansson et al. (2016); Shalit et al. (2017) proposed learning representations for counterfactual inference, inspired by work in unsupervised domain adaptation (Mansour et al., 2009). However, the generalization bounds of Johansson et al. (2016) do not apply to general hypotheses and the bounds of Shalit et al. (2017) and Long et al. (2015) are loose—even if infinite samples are available.

In this work, *we combine ideas from representation learning and sample re-weighting*, generalizing previous work, to give finite-sample learning bounds that are tight in the asymptotic limit under a well-specified model, and best-in-class otherwise. These bounds give rise to algorithms with good empirical performance and added robustness over previous work.

4   GENERALIZATION UNDER DESIGN SHIFT

We proceed to derive a bound on the risk in predicting outcomes under a target design $p_\pi(T, X)$, based on unlabeled samples from p_π and labeled samples from a source design $p_\mu(T, X)$. Our result combines distribution matching and re-weighting, resulting in a tighter bound than the closest related work (Shalit et al., 2017). The predictors we consider take the form of a composition $f(x, t) = h(\Phi(x), t)$ where Φ is a representation of x and h an hypothesis. We first state a result for the general design shift setting. Then, we show how this result can be used to bound the error in prediction of treatment effects. Finally, in Section 5 we give a result about the asymptotic properties of the minimizers of the upper bound.

Our bounds build on the intuition that if either a) π is close to a re-weighting of μ, or b) the true outcome is a simple function of x and t, the gap between the target error and the re-weighted source error is small. This can be formalized using integral probability metrics (IPM) (Sriperumbudur et al., 2009) that measure distance between distributions w.r.t. a normed vector space of functions $\mathcal{H}$.

Definition 2. *The integral probability metric (IPM) distance, associated with a normed vector space of functions $\mathcal{H}$, between distributions p and q is,*

$$\text{IPM}_\mathcal{H}(p, q) := \sup_{h \in \mathcal{H}} |\mathbb{E}_p[h] - \mathbb{E}_q[h]|$$

Important examples of IPMs include the Wasserstein distance, for which $\mathcal{H}$ is the family of functions with Lipschitz constant at most 1, and the Maximum Mean Discrepancy for which $\mathcal{H}$ are functions in the norm-1 ball in a reproducing kernel Hilbert space. We can now state the following result.

---

[2]Specifically all confounding variables, causal of both treatment and outcome.

Lemma 1. *For hypotheses f with loss ℓ_f such that $\ell_f/\|\ell_f\|_{\mathcal{H}} \in \mathcal{H}$, and p_μ, p_π with common support, there exists a valid re-weighting w of p_μ, see Definition 1, such that,*

$$R_\pi(f) \le R_\mu^w(f) + \|\ell_f\|_{\mathcal{H}}\mathrm{IPM}_{\mathcal{H}}(p_\pi, p_\mu^w) \le R_\mu(f) + \|\ell_f\|_{\mathcal{H}}\mathrm{IPM}_{\mathcal{H}}(p_\pi, p_\mu) . \quad (5)$$

The first inequality is tight for importance sampling weights, $w(x,t) = p_\pi(x,t)/p_\mu(x,t)$. The second inequality is not tight for general f, even if $\ell_f/\|\ell_f\|_{\mathcal{H}} \in \mathcal{H}$, unless $p_\pi = p_\mu$.

Lemma 1 implies that there exist weighting functions $w(x,t)$ that achieve a tighter bound than uniform weights $w(x,t) = 1$. While importance sampling weights result in a tight bound in expectation, they are not known in general, and can result in large variance in finite samples (Cortes et al., 2010). Instead, we will search for a weighting function w, that minimizes a finite-sample version of Lemma 1, but first we introduce the notion of representation learning to combat distributional shift.

The idea of learning representations to minimize distributional shift in representation space, and thus the source-target error gap, has been applied to domain adaptation (Ajakan et al., 2014), algorithmic fairness (Zemel et al., 2013) and counterfactual prediction (Shalit et al., 2017), to name a few. We follow the setup of Shalit et al. (2017), and consider learning twice-differentiable, invertible representations $\Phi : \mathcal{X} \to \mathcal{Z}$, where $\mathcal{Z}$ is the representation space, and $\Psi : \mathcal{Z} \to \mathcal{X}$ is the inverse representation, such that $\Psi(\Phi(x)) = x$ for all x. Let $\mathcal{C}$ denote space of such representation functions. For a design π, we let $p_{\pi,\Phi}(z,t)$ be the distribution induced by Φ over $\mathcal{Z} \times \mathcal{T}$, with $p_{\pi,\Phi}^w(z,t) := p_{\pi,\Phi}(z,t)w(\Psi(z),t)$ its reweighted form and $\hat{p}_{\pi,\Phi}^w$ its reweighted empirical form, following our previous notation. Let $\mathcal{G} \subseteq \{h : \mathcal{Z} \times \mathcal{T} \to \mathcal{Y}\}$ denote a set of hypotheses $h(\Phi,t)$ operating on the representation Φ and let $\mathcal{F}$ the space of all compositions, $\mathcal{F} = \{f = h(\Phi(x),t) : h \in \mathcal{G}, \Phi \in \mathcal{C}\}$. We can now relate the expected target risk $R_\pi(f)$ to the re-weighted empirical source risk $\hat{R}_\mu^w(f)$.

Theorem 1. *Given is a labeled sample $(x_1, t_1, y_1), ..., (x_n, t_n, y_n)$ from p_μ, and an unlabeled sample $(x'_1, t'_1), ..., (x'_m, t'_m)$ from p_π, with corresponding empirical measures $\hat{p}_\mu$ and $\hat{p}_\pi$. Suppose that Φ is a twice-differentiable, invertible representation, that $h(\Phi,t)$ is an hypothesis, and $f = h(\Phi(x),t) \in \mathcal{F}$. Define $m_t(x) = \mathbb{E}_Y[Y \mid X = x, T = t]$, let $\ell_{h,\Phi}(\Psi(z),t) := L(h(z,t), m_t(\Psi(z)))$ where L is the squared loss, $L(y,y') = (y - y')^2$, and assume that there exists a constant $B_\Phi > 0$ such that $\ell_{h,\Phi}/B_\Phi \in \mathcal{H} \subseteq \{h : \mathcal{Z} \times \mathcal{T} \to \mathcal{Y}\}$, where $\mathcal{H}$ is a reproducing kernel Hilbert space of a kernel, k such that $k((z,t),(z,t)) < \infty$. Finally, let w be a valid re-weighting of $p_{\mu,\Phi}$. Then with probability at least $1 - 2\delta$,*

$$R_\pi(f) \le \hat{R}_\mu^w(f) + B_\Phi\mathrm{IPM}_{\mathcal{H}}(\hat{p}_{\pi,\Phi}, \hat{p}_{\mu,\Phi}^w) + V_\mu(w, \ell_f)\frac{\mathcal{C}_{n,\delta}^{\mathcal{F}}}{n^{3/8}} + \mathcal{D}_\delta^{\Phi,\mathcal{H}}\left(\frac{1}{\sqrt{m}} + \frac{1}{\sqrt{n}}\right) + \sigma_Y^2 \quad (6)$$

where $\mathcal{C}_{n,\delta}^{\mathcal{F}}$ measures the capacity of $\mathcal{F}$ and has only logarithmic dependence on n, $\mathcal{D}_{m,n,\delta}^{\mathcal{H}}$ measures the capacity of $\mathcal{H}$, σ_Y^2 is the expected variance in potential outcomes, and

$$V_\mu(w, \ell_f) = \max(\sqrt{\mathbb{E}_{p_\mu}[w^2(x,t)\ell_f^2(x,t)]}, \sqrt{\mathbb{E}_{\hat{p}_\mu}[w^2(x,t)\ell_f^2(x,t)]}) .$$

A similar bound exists where $\mathcal{H}$ is the family of functions Lipschitz constant at most 1, but with worse sample complexity.

See Appendix A.2 for a proof that involves applying finite-sample generalization bounds to the first inequality in Lemma 1, as well as moving to the space induced by the representation Φ.

Using uniform weights $w(x,t) = 1$ in (6), results in a bound similar to that of Shalit et al. (2017) and Long et al. (2015). For $\pi \ne \mu$, minimizing the resulting bound results biased hypotheses, even in the asymptotical limit, as the IPM term does not vanish when the sample size increases. This is a rather undesirable property, as even a k-nearest-neighbor classifiers are consistent in the limit of infinite samples. Instead, we consider minimizing (6) with respect to w, improving the tightness of the bound. Recall from Lemma 1 that there exist weights w for which $R_\pi(f) = R_\mu^w(f)$.

Theorem 1 indicates, as noted by for example Cortes et al. (2010), that even though importance sampling weights w^* yield estimators with small bias, they can suffer from high variance due to the factor $V_\mu(w, \ell_f)$. The factor B_Φ is not known in general as it depends on the true outcome, and is determined by $\|\ell_f\|_{\mathcal{H}}$ as well as the determinant of the Jacobian of Ψ, see the appendix. Qualitatively, B_Φ mesures the joint complexity of Φ and $\ell(r,t)$. In practice, Shalit et al. (2017) substituted a hyperparameter α for B_Φ, but discussed the difficulties of selecting its value without

access to counterfactual labels. In our experiments, we explore a heuristic for adaptively choosing α, based on measures of complexity of the observed held-out loss as a function of the input.

We note that Theorem 1 is immediately applicable to the case of unsupervised domain adaptation in which there is only a single potential outcome of interest, $\mathcal{T} = \{0\}$. In this case, $p_\mu(T \mid X) = p_\pi(T \mid X)$, and our only source of shift is in the domain $p(X)$.

Conditional Average Treatment Effects A simple argument shows that the error in predicting the conditional average treatment effect, $\mathrm{MSE}(\hat{\tau})$ can be bounded by the sum of risks under the constant treat-all and treat-none policies. As in Section 2.2, we consider the case of a fixed domain $p_\pi(X) = p_\mu(X)$ and binary treatment $\mathcal{T} = \{0, 1\}$. Let $R_{\pi_t}(f)$ denote the risk under the constant policy π_t such that $\forall x \in \mathcal{X}: p_{\pi_t}(T = t \mid X = x) = 1$.

Proposition 1. *We have with* $\mathrm{MSE}(\hat{\tau})$ *as in* (4) *and* $R_t(f)$ *the risk under the constant policy* π_t,

$$\mathrm{MSE}(\hat{\tau}) \leq 2(R_{\pi_1}(f) + R_{\pi_0}(f)) - 4\sigma^2 \tag{7}$$

where σ *is such that* $\forall t \in \mathcal{T}, x \in \mathcal{X}, \sigma_Y(x, t) \geq \sigma$ *and* $\sigma_Y(x, t)$ *is standard deviation of* $Y(t)$ *conditioned on* $X = x$. *Further, for each* $t \in \{0, 1\}$ *we have,*

$$R_{\pi_t}(f) = \underbrace{\mathbb{E}_\mu[\ell_f(x, t) \mid T = t]p_\mu(t)}_{Factual} + \underbrace{\mathbb{E}_\mu[\ell_f(x, t) \mid T = 1 - t]p_\mu(1 - t)}_{Counterfactual} . \tag{8}$$

The proof involves the relaxed triangle inequality and the law of total probability. By Proposition 1, we can apply Theorem 1 to R_{π_1} and R_{π_0} separately, to obtain a bound on $\mathrm{MSE}(\tau)$. For brevity, we refrain from stating the full result, but emphasize that if follows from Theorem 1. In Section 6, we evaluate our framework in treatment effect estimation, minimizing this bound.

5 JOINT LEARNING OF REPRESENTATIONS AND SAMPLE WEIGHTS

Motivated by the theoretical insights of Section 4, we propose to jointly learn a representation $\Phi(x)$, a re-weighting $w(x, t)$ and an hypothesis $h(\Phi, t)$ by minimizing a bound on the risk under the target design, see (6). This approach improves on previous work in two ways: it alleviates the bias of Shalit et al. (2017) when sample sizes are large, see Section 4, and it increases the flexibility of the balancing method of Gretton et al. (2009) by learning the representation to balance.

For notational brevity, we let $w_i = w(x_i, t_i)$. Recall that $\hat{p}_{\pi, \Phi}^w$ is the re-weighted empirical distribution of representations Φ under p_π. The training objective of our algorithm is the RHS of (6), with hyperparameters $\beta = (\alpha, \lambda_h, \lambda_w)$ substituted for model (and representation) complexity terms,

$$\mathcal{L}_\pi(h, \Phi, w; \beta) = \underbrace{\frac{1}{n} \sum_{i=1}^{n} w_i \ell_h(\Phi(x_i), t_i) + \frac{\lambda_h}{\sqrt{n}} \mathcal{R}(h)}_{\mathcal{L}_\pi^h(h, \Phi, w; D, \alpha, \lambda_h)} + \underbrace{\alpha \, \mathrm{IPM}_G(\hat{p}_{\pi, \Phi}, \hat{p}_{\mu, \Phi}^w) + \lambda_w \frac{\|w\|_2}{n}}_{\mathcal{L}_\pi^w(\Phi, w; D, \alpha, \lambda_w)} \tag{9}$$

where $\mathcal{R}(h)$ is a regularizer of h, such as ℓ_2-regularization. We can show the following result.

Theorem 2. *Suppose* $\mathcal{H}$ *is a reproducing kernel Hilbert space given by a bounded kernel. Suppose weak overlap holds in that* $\mathbb{E}[(p_\pi(x, t)/p_\mu(x, t))^2] < \infty$. *Then,*

$$\min_{h, \Phi, w} \mathcal{L}_\pi(h, \Phi, w; \beta)] \leq \min_{f \in \mathcal{F}} R_\pi(f) + O_p(1/\sqrt{n} + 1/\sqrt{m}) .$$

Consequently, under the assumptions of Thm. 1, for sufficiently large α *and* λ_w,

$$R_\pi(\hat{f}_n) \leq \min_{f \in \mathcal{F}} R_\pi(f) + O_p(1/n^{3/8} + 1/\sqrt{m}).$$

In words, the minimizers of (9) *converge to the representation and hypothesis that minimize the counterfactual risk, in the limit of infinite samples.*

While minimizing $\mathcal{L}_\pi(h, \Phi, w; \beta)$ directly over h, Φ and w is justified by Theorem 2, we note that adjusting w to minimize the empirical risk term serves little purpose, as it may result in overemphasizing "easy" training examples, especially if α is small. Instead, as a heuristic, we split the

Context Repres. Hypothesis Weighted risk Treatment

Figure 1: Architecture for predicting outcomes under design shift. A re-weighting function w is fit jointly with a representation Φ and hypothis h of the potential outcomes, to minimize a bound on the target risk. Dashed lines are not back-propagated through. Regularization penalties not shown.

Table 1: Causal effect estimation on IHDP. CATE error, target prediction error and std errors. Lower is better.

	RMSE($\hat{\tau}$)	$\hat{R}_\pi(f)$
OLS	$2.3 \pm .11$	$1.1 \pm .05$
OLS-IPW	$2.4 \pm .11$	$1.2 \pm .05$
Random For.	$6.6 \pm .30$	$4.1 \pm .18$
Causal For.	$3.8 \pm .18$	$1.8 \pm .08$
BART	$2.3 \pm .10$	$1.7 \pm .07$
IPM-WNN	$1.2 \pm .12$	$.65 \pm .02$
CFRW	$.76 \pm .02$	$.46 \pm .01$
RCFR Adapt. $\alpha, w = 1$	$.80 \pm .06$	$.45 \pm .03$
RCFR Oracle α	$.65 \pm .04$	$.38 \pm .01$
RCFR Adapt. α	$.67 \pm .05$	$.37 \pm .01$

objective in two, see (9), and use only the IPM term and regularizer to learn w. In short, we solve the following alternating minimization problem.

$$h^k, \Phi^k = \arg\min_{h,\Phi} \ \mathcal{L}_\pi^h(h, \Phi, w; D, \alpha, \lambda_h), \quad w^{k+1} = \arg\min_w \ \mathcal{L}_\pi^w(\Phi^k, w; D, \alpha, \lambda_w) \quad (10)$$

In principle, the weighting function $w(x, t)$ could be represented as one free parameter per training point, as it is only used to learn the model, not for prediction. However, we propose to let w be a parametric function of $\Phi(x)$. Doing so enables us to both control the smoothness of the weights more flexibly, and lets us evaluate weights on a hold-out set, for example to perform early stopping. An example architecture for this framework, is presented in Figure 1. As noted previously, estimating treatment effects involves predicting under the two constant policies—treat everyone and treat noone. In Section 6, we evaluate our framework in this task.

As noted by Shalit et al. (2017), choosing hyperparameters for prediction outcomes is fundamentally difficult, as we cannot observe ground truth for counterfactuals. In this work, we explore setting the balance parameter α adaptively. α is used in (9) in place of a factor measuring the complexity of the loss and representation function as functions of the input, a quantity that evolves during training. As a heuristic, we use an approximation of the Lipschitz constant of ℓ_f based on observed examples, with $f = h(\Phi(x), t)$, $\alpha_{h,\Phi} = \max_{i,j \in [n]} |\ell_f(x_i, t_i) - \ell_f(x_j, t_j)|/\|x_i - x_j\|_2$. We use a moving average over batches to encourage stability.

6 EXPERIMENTS

We evaluate our framework in the CATE estimation setting, see Section 2.2—our task is to predict the expected difference between potential outcomes conditioned on pre-treatment variables, *for a held-out sample* of the population. We compare our results to ordinary least squares (OLS) (with one regressor per outcome), OLS-IPW (reweighted OLS according to a logistic regression estimate of propensities), Random Forests, Causal Forests (Wager & Athey, 2017), BART (Chipman et al., 2010), and CFRW (Shalit et al., 2017) (with Wasserstein penalty). Finally, we use as baseline (IPM-WNN): first weights are found by IPM minimization in the input space (Gretton et al., 2009; Kallus, 2016), then used in a re-weighted neural net regression, with the same architecture as our method. We use IHDP as benchmark, a semi-synthetic binary-treatment dataset (Hill, 2011), split into training and test sets by Shalit et al. (2017). IHDP has synthesized continuous outcomes that can be used to compute the ground-truth CATE error.

Our implementtation, dubbed RCFR for *Reweighted Counterfactual Regression*, parameterizes representations $\Phi(x)$, weighting functions $w(\Phi, t)$ and hypotheses $h(\Phi, t)$ using neural networks, trained by minimizing (9). We use the RBF-kernel maximum mean discrepancy as IPM (Gretton et al., 2012). For a description of the architecture, training procedure and hyperparameters, see Appendix B. We compare results using uniform $w = 1$ and learned weights, picking the balance parameter α using either an oracle or the adaptive heuristic described in Section B. To pick other hyperparameters, we split training sets into one part used for function fitting and one used for early

Figure 2: Error in CATE estimation on IHDP as a function of re-weighting regularization strength λ_w (left) and source prediction error (right). We see in the left-hand plot that a) for small imbalance penalties α, re-weighting (low λ_w) has no effect, b) for moderate α, less uniform re-weighting (smaller λ_w) improves the error, c) for large α, weighting helps, but overall error increases. In the right-hand plot, we compare the ratio of CATE error to source error. Color represents α (see left) and size λ_w. We see that for large α, the source error is more representative of CATE error, but does not improve in absolute value without weighting. Here, α was fixed during training.

stopping and hyperparameter selection. Hyperparameters for regularizaion and early stopping is done based on the empirical risk on the held-out sample.

We present the results of our evalation on IHDP in Table 1. We see that our proposed method achieves state-of-the-art results, and that adaptively choosing α does not hurt performance much. Furthermore, we see a substantial improvement from using non-uniform sample weights.

In Figure 2 we take a closer look at the behavior of our model as we vary its hyperparameters on the IHDP dataset. Between the two plots we can draw the following conclusions: a) For moderate to large $\alpha \in [10, 100]$, we observe a marginal gain from using the IPM penalty. This confirms the observations of Shalit et al. (2017). b) For large $\alpha \in [10, 1000]$, we see a large gain from using a non-uniform re-weighting (small λ_w). c) While large α makes the factual error more representative of the counterfactual error, using it without re-weighting results in higher absolute error.

7 CONCLUSION

We have proposed a theory and an algorithmic framework for learning to predict outcomes of interventions under shifts in design—changes in both intervention policy and feature domain. The framework, combining representation learning and sample re-weighting, is used to derive generalization bounds on the risk under a target design, for hypotheses learned with supervision only in the source design. This leads to models that achieve state-of-the-art results on a benchmark for a special case of our setting—causal effect estimation.

REFERENCES

Hana Ajakan, Pascal Germain, Hugo Larochelle, François Laviolette, and Mario Marchand. Domain-adversarial neural networks. *arXiv preprint arXiv:1412.4446*, 2014.

Susan Athey and Guido Imbens. Recursive partitioning for heterogeneous causal effects. *Proceedings of the National Academy of Sciences*, 113(27):7353–7360, 2016.

Peter C Austin. An introduction to propensity score methods for reducing the effects of confounding in observational studies. *Multivariate behavioral research*, 46(3):399–424, 2011.

Alexandre Belloni, Victor Chernozhukov, and Christian Hansen. Inference on treatment effects after selection among high-dimensional controls. *The Review of Economic Studies*, 81(2):608–650, 2014.

Yoshua Bengio, Aaron Courville, and Pascal Vincent. Representation learning: A review and new perspectives. *IEEE transactions on pattern analysis and machine intelligence*, 35(8):1798–1828, 2013.

Victor Chernozhukov, Denis Chetverikov, Mert Demirer, Esther Duflo, Christian Hansen, Whitney Newey, James Robins, et al. Double/debiased machine learning for treatment and causal parameters. Technical report, 2017.

Hugh A Chipman, Edward I George, Robert E McCulloch, et al. Bart: Bayesian additive regression trees. *The Annals of Applied Statistics*, 4(1):266–298, 2010.

Corinna Cortes, Yishay Mansour, and Mehryar Mohri. Learning bounds for importance weighting. In *Advances in neural information processing systems*, pp. 442–450, 2010.

David A Freedman and Richard A Berk. Weighting regressions by propensity scores. *Evaluation Review*, 32(4):392–409, 2008.

Arthur Gretton, Alexander J Smola, Jiayuan Huang, Marcel Schmittfull, Karsten M Borgwardt, and Bernhard Schölkopf. Covariate shift by kernel mean matching. 2009.

Arthur Gretton, Karsten M Borgwardt, Malte J Rasch, Bernhard Schölkopf, and Alexander Smola. A kernel two-sample test. *Journal of Machine Learning Research*, 13(Mar):723–773, 2012.

Jennifer L Hill. Bayesian nonparametric modeling for causal inference. *Journal of Computational and Graphical Statistics*, 20(1), 2011.

Guido W Imbens and Donald B Rubin. *Causal inference in statistics, social, and biomedical sciences*. Cambridge University Press, 2015.

Fredrik Johansson, Uri Shalit, and David Sontag. Learning representations for counterfactual inference. In *International Conference on Machine Learning*, pp. 3020–3029, 2016.

Nathan Kallus. Optimal a priori balance in the design of controlled experiments. *Journal of the Royal Statistical Society: Series B (Statistical Methodology)*. doi: 10.1111/rssb.12240.

Nathan Kallus. Generalized optimal matching methods for causal inference. *arXiv preprint arXiv:1612.08321*, 2016.

Mingsheng Long, Yue Cao, Jianmin Wang, and Michael Jordan. Learning transferable features with deep adaptation networks. In *International Conference on Machine Learning*, pp. 97–105, 2015.

Yishay Mansour, Mehryar Mohri, and Afshin Rostamizadeh. Domain adaptation: Learning bounds and algorithms. *arXiv preprint arXiv:0902.3430*, 2009.

Stephen L Morgan and Christopher Winship. *Counterfactuals and causal inference*. Cambridge University Press, 2014.

Judea Pearl. *Causality*. Cambridge university press, 2009.

Doina Precup, Richard S Sutton, and Sanjoy Dasgupta. Off-policy temporal-difference learning with function approximation. In *ICML*, pp. 417–424, 2001.

Paul R Rosenbaum and Donald B Rubin. The central role of the propensity score in observational studies for causal effects. *Biometrika*, 70(1):41–55, 1983.

Donald B Rubin. Causal inference using potential outcomes: Design, modeling, decisions. *Journal of the American Statistical Association*, 100(469):322–331, 2005.

Uri Shalit, Fredrik Johansson, and David Sontag. Estimating individual treatment effect: generalization bounds and algorithms. In *International Conference on Machine Learning*, 2017.

Hidetoshi Shimodaira. Improving predictive inference under covariate shift by weighting the log-likelihood function. *Journal of statistical planning and inference*, 90(2):227–244, 2000.

Bharath K Sriperumbudur, Kenji Fukumizu, Arthur Gretton, Bernhard Schölkopf, and Gert RG Lanckriet. On integral probability metrics,\phi-divergences and binary classification. *arXiv preprint arXiv:0901.2698*, 2009.

Elizabeth A Stuart. Matching methods for causal inference: A review and a look forward. *Statistical science: a review journal of the Institute of Mathematical Statistics*, 25(1):1, 2010.

Adith Swaminathan and Thorsten Joachims. Counterfactual risk minimization: Learning from logged bandit feedback. In *International Conference on Machine Learning*, pp. 814–823, 2015.

Mark J Van Der Laan and Daniel Rubin. Targeted maximum likelihood learning. *The International Journal of Biostatistics*, 2(1), 2006.

Stefan Wager and Susan Athey. Estimation and inference of heterogeneous treatment effects using random forests. *Journal of the American Statistical Association*, (just-accepted), 2017.

Rich Zemel, Yu Wu, Kevin Swersky, Toni Pitassi, and Cynthia Dwork. Learning fair representations. In *Proceedings of the 30th International Conference on Machine Learning (ICML-13)*, pp. 325–333, 2013.

José R Zubizarreta. Stable weights that balance covariates for estimation with incomplete outcome data. *Journal of the American Statistical Association*, 110(511):910–922, 2015.

APPENDIX

A PROOFS

A.1 DEFINITIONS

Distribution re-weighting

Definition 1 (Restated). *A function $w : \mathcal{X} \times \mathcal{T} \to \mathbb{R}_+$ is a valid re-weighting of p_μ if*

$$\mathbb{E}_{x,t \sim p_\mu}[w(x,t)] = 1 \ \ and \ \ p_\mu(x,t) > 0 \Rightarrow w(x,t) > 0.$$

We denote the re-weighted density $p_\mu^w(x,t) := w(x,t)p_\mu(x,t)$.

Expected & empirical risk We let the (expected) risk of f measured by ℓ_h under p_μ be denoted

$$R_\mu(h) = \mathbb{E}_{p_\mu}[l_h(x,t)]$$

where l_h is an appropriate loss function, and the empirical risk over a sample $D_\mu = \{(x_1,t_1,y_1)...,(x_n,t_n,y_n)\}$ from p_μ

$$\hat{R}_\mu(f) = \frac{1}{n}\sum_{i=1}^n l_f(x_i,t_i,y_i)\,.$$

We use the superscript w to denote the re-weighted risks

$$R_\mu^w(f) = \mathbb{E}[w(x,t)l_f(x,t)] \quad \hat{R}_\mu^w(f) = \frac{1}{n}\sum_{i=1}^n w(x_i,t_i)l_h(x_i,t_i,y_i)$$

Definition A1 (Importance sampling). *For two distributions p,q on $\mathcal{Z}$, of common support, $\forall z \in \mathcal{Z} : p(z) > 0 \iff q(z) > 0$, we call*

$$w_{IS}(z) := \frac{q(z)}{p(z)}$$

the importance sampling *weights of p and q.*

Definition 2 (Restated). *The integral probability metric (IPM) distance, associated with the function family $\mathcal{H}$, between distributions p and q is defined by*

$$\text{IPM}_\mathcal{H}(p,q) := \sup_{h:\|h\|_\mathcal{H}=1} |\mathbb{E}_p[h] - \mathbb{E}_q[h]|$$

A.2 LEARNING BOUNDS

We begin by bounding the expected risk under a distribution p_π in terms of the expected risk under p_μ and a measure of the discrepancy between p_π and p_μ. Using definition 2 we can show the following result.

Lemma 1 (Restated). *For hypotheses f with loss ℓ_f such that $\ell_f/\|\ell_f\|_\mathcal{H} \in \mathcal{H}$, and p_μ, p_π with common support, there exists a valid re-weighting w of p_μ, see Definition 1, such that,*

$$R_\pi(f) \leq R_\mu^w(f) + \|\ell_f\|_\mathcal{H}\text{IPM}_\mathcal{H}(p_\pi, p_\mu^w) \leq R_\mu(f) + \|\ell_f\|_\mathcal{H}\text{IPM}_\mathcal{H}(p_\pi, p_\mu)\,. \quad (11)$$

The first inequality is tight for importance sampling weights, $w(x,t) = p_\pi(x,t)/p_\mu(x,t)$. The second inequality is not tight for general f, even if $\ell_f \in \mathcal{H}$, unless $p_\pi = p_\mu$.

Proof. The results follows immediately from the definition of IPM.

$$R_\pi(f) - R_\mu^w(f) = \mathbb{E}_\pi[\ell_f(x,t)] - \mathbb{E}_\mu[w(x,t)\ell_f(x,t)]$$
$$\leq \sup_{h \in \mathcal{H}_\ell} |\mathbb{E}_\pi[h(x,t)] - \mathbb{E}_\mu[w(x,t)h(x,t)]|$$
$$= \text{IPM}_{\mathcal{H}_\ell}(p_\pi, p_\mu^w)$$

Further, for importance sampling weights $w_{IS}(x,t) = \frac{\pi(t;x)}{\mu(t;x)}$, for any $h \in \mathcal{H}$,

$$\mathbb{E}_\pi[h(x,t)] - \mathbb{E}_\mu[w_{IS}(x,t)h(x,t)] = \mathbb{E}_\pi[h(x,t)] - \mathbb{E}_\mu[\frac{\pi(t;x)}{\mu(t;x)}h(x,t)]$$

$$= 0$$

and the LHS is tight. $\qquad\square$

We could apply Lemma 1 to bound the loss under a distribution q based on the weighted loss under p. Unfortunately, bounding the expected risk in terms of another expectation is not enough to reason about generalization from an empirical sample. To do that we use Corollary 2 of Cortes et al. (2010), restated as a Theorem below.

Theorem A1 (Generalization error of re-weighted loss (Cortes et al., 2010)). *For a loss function ℓ_h of any hypothesis $h \in \mathcal{H} \subseteq \{h' : \mathcal{X} \to \mathbb{R}\}$, such that $d = \mathrm{Pdim}(\{\ell_h : h \in \mathcal{H}\})$ where Pdim is the pseudo-dimension, and a weighting function $w(x)$ such that $\mathbb{E}_p[w] = 1$, with probability $1 - \delta$ over a sample $(x_1, ..., x_n)$, with empirical distribution $\hat{p}$,*

$$R_p^w(h) \le \hat{R}_p^w(h) + 2^{5/4}V_{p,\hat{p}}[w(x)l(x)]\left(\frac{d\log\frac{2ne}{d} + \log\frac{4}{\delta}}{n}\right)^{3/8}$$

with $V_{p,\hat{p}}[w(x)l(x)] = \max(\sqrt{\mathbb{E}_p[w^2(x)\ell_h^2(x)]}, \sqrt{\mathbb{E}_{\hat{p}}[w^2(x)\ell_h^2(x)]})$. With

$$\mathcal{C}_n^{\mathcal{H}} = 2^{5/4}\left(d\log\frac{2ne}{d} + \log\frac{4}{\delta}\right)^{3/8}$$

we get the simpler form

$$R_p^w(h) \le \hat{R}_p^w(h) + V_{p,\hat{p}}[w(x)l(x)]\frac{\mathcal{C}_n^{\mathcal{H}}}{n^{3/8}} .$$

We will also need the following result about estimating IPMs from finite samples from Sriperumbudur et al. (2009).

Theorem A2 (Estimation of IPMs from empirical samples (Sriperumbudur et al., 2009)). *Let M be a measurable space. Suppose k is measurable kernel such that $\sup_{x \in M} k(x,x) \le C \le \infty$ and $\mathcal{H}$ the reproducing kernel Hilbert space induced by k, with $\nu := \sup_{x \in M, f \in \mathcal{H}} f(x) < \infty$. Then, with $\hat{p}, \hat{q}$ the empirical distributions of p, q from m and n samples respectively, and with probability at least $1 - \delta$,*

$$|\mathrm{IPM}_{\mathcal{H}}(p,q) - \mathrm{IPM}_{\mathcal{H}}(\hat{p}, \hat{q})| \le \sqrt{18\nu^2 \log\frac{4}{\delta}C}\left(\frac{1}{\sqrt{m}} + \frac{1}{\sqrt{n}}\right)$$

We consider learning twice-differentiable, invertible representations $\Phi : \mathcal{X} \to \mathcal{Z}$, where $\mathcal{Z}$ is the representation space, and $\Psi : \mathcal{Z} \to \mathcal{X}$ is the inverse representation, such that $\Psi(\Phi(x)) = x$ for all x. Let $\mathcal{C}$ denote space of such representation functions. For a design π, we let $p_{\pi,\Phi}(z,t)$ be the distribution induced by Φ over $\mathcal{Z} \times \mathcal{T}$, with $p_{\pi,\Phi}^w(z,t) := p_{\pi,\Phi}(z,t)w(\Psi(z),t)$ its reweighted form and $\hat{p}_{\pi,\Phi}^w$ its reweighted empirical form, following our previous notation. Note that we do not include t in the representation itself, although this could be done in principle. Let $\mathcal{G} \subseteq \{h : \mathcal{Z} \times \mathcal{T} \to \mathcal{Y}\}$ denote a set of hypotheses $h(\Phi, t)$ operating on the representation Φ and let $\mathcal{F}$ denote the space of all compositions, $\mathcal{F} = \{f = h(\Phi(x), t) : h \in \mathcal{G}, \Phi \in \mathcal{C}\}$. We now restate and prove Theorem 1.

Theorem 1 (Restated). *Given is a labeled sample $D_\mu = \{(x_1, t_1, y_1), ..., (x_n, t_n, y_n)\}$ from p_μ, and an unlabeled sample $D_\pi = \{(x'_1, t'_1), ..., (x'_m, t'_m)\}$ from p_π, with corresponding empirical measures $\hat{p}_\mu$ and $\hat{p}_\pi$. Suppose that Φ is a twice-differentiable, invertible representation, that $h(\Phi, t)$ is an hypothesis, and $f = h(\Phi(x), t) \in \mathcal{F}$. Define $m_t(x) = \mathbb{E}_Y[Y \mid X = x, T = t]$, let $\ell_{h,\Phi}(\Psi(z), t) := L(h(z,t), m_t(\Psi(z)))$ where L is the squared loss, $L(y, y') = (y - y')^2$, and assume that there exists a constant $B_\Phi > 0$ such that $\ell_{h,\Phi}/B_\Phi \in \mathcal{H} \subseteq \{h : \mathcal{Z} \times \mathcal{T} \to \mathcal{Y}\}$, where $\mathcal{H}$ is a reproducing kernel Hilbert space of a kernel, k such that $k((z,t),(z,t)) < \infty$. Finally, let w be a valid re-weighting of $p_{\mu,\Phi}$. Then with probability at least $1 - 2\delta$,*

$$R_\pi(f) \le \hat{R}_\mu^w(f) + B_\Phi\mathrm{IPM}_{\mathcal{H}}(\hat{p}_{\pi,\Phi}, \hat{p}_{\mu,\Phi}^w) + V_\mu(w, \ell_f)\frac{\mathcal{C}_{n,\delta}^{\mathcal{F}}}{n^{3/8}} + \mathcal{D}_\delta^{\Phi,\mathcal{H}}\left(\frac{1}{\sqrt{m}} + \frac{1}{\sqrt{n}}\right) + \sigma_Y^2 \quad (12)$$

where $\mathcal{C}^{\mathcal{F}}_{n,\delta}$ measures the capacity of $\mathcal{F}$ and has only logarithmic dependence on n, $\mathcal{D}^{\mathcal{H}}_{m,n,\delta}$ measures the capacity of $\mathcal{H}$, σ^2_Y is the expected variance in potential outcomes, and

$$V_\mu(w, \ell_f) = \max(\sqrt{\mathbb{E}_{p_\mu}[w^2(x,t)\ell_f^2(x,t)]}, \sqrt{\mathbb{E}_{\hat{p}_\mu}[w^2(x,t)\ell_f^2(x,t)]}) .$$

A similar bound exists where $\mathcal{H}$ is the family of functions Lipschitz constant at most 1, but with worse sample complexity.

Proof. We have by definition

$$R_\pi(f) - R^w_\mu(f) = \mathbb{E}_\pi[\ell_f(x,t,y)] - \mathbb{E}_\mu[w(x,t)\ell_f(x,t,y)]$$

$$= \int_{x,t,y} \ell_f(x,t,y)p(y \mid t,x)(p_\pi(x,t) - p^w_\mu(x,t))dxdtdy$$

Define $\ell_{h,\Phi}(x,t) = L(h(\Phi(x),t), m_t(x))$ where $m_t(x) := E[Y \mid T = t, X = x]$. Then, with L, the squared loss, $L(y,y') = (y-y')^2$, we have,

$$\mathbb{E}_\pi[\ell_{h,\Phi}(x,t,y)] = \mathbb{E}_\pi[\ell_{h,\Phi}(x,t)] + \sigma^2_\pi$$

where $\sigma^2_\pi = \mathbb{E}_{p_\pi}[(Y - m_t(x))^2]$, and analogously for μ. We get that

$$R_\pi(f) - R^w_\mu(f) = \int_{x\in\mathcal{X},t\in\mathcal{T}} \ell_{h,\Phi}(x,t)(p_\pi(x,t) - p^w_\mu(x,t))dxdt + \sigma^2_\pi + \sigma^2_\mu$$

$$= \int_{z\in\mathcal{Z},t\in\mathcal{T}} \ell_{h,\Phi}(\Psi(z),t)(p_{\pi,\Phi}(z,t) - p^w_{\mu,\Phi}(z,t))|J_\Psi(z)|dzdt + \sigma_\pi$$

$$\leq A_\Phi \int_{z\in\mathcal{Z},t\in\mathcal{T}} \ell_{h,\Phi}(\Psi(z),t)(p_\pi(z,t) - p^w_\mu(z,t))dzdt + \sigma^2_\pi + \sigma^2_\mu$$

$$\leq \sigma^2_\pi + \sigma^2_\mu + A_\Phi\|\ell_{h,\Phi}\|_{\mathcal{H}} \sup_{h\in\mathcal{H}} \int_{z\in\mathcal{Z},t\in\mathcal{T}} h(\Psi(z),t)\left|p_{\pi,\Phi}(rz,t) - p^w_{\mu,\Phi}(z,t)\right| dzdt$$

$$= B_\Phi \cdot \mathrm{IPM}_{\mathcal{H}}(p_{\pi,\Phi}, p^w_{\mu,\Phi}) + \sigma^2_\pi + \sigma^2_\mu$$

where $J_\Psi(z)$ is the Jacobian matrix of Ψ evaluated at z and $A_\Phi \geq |J_\Psi(z)|$ for all $z \in \mathcal{Z}$, where $|J|$ is the absolute determinant of J. By application of Theorem A1 we have with probability at least $1 - \delta$,

$$R^w_\mu(f) \leq \hat{R}^w_\mu(f) + V_\mu(w,\ell)\frac{\mathcal{C}^{\mathcal{H}}_{n,\delta}}{n^{3/8}} .$$

and by applying Theorem A2, we have with probability at least $1 - \delta$,

$$\left|\mathrm{IPM}_{\mathcal{H}}(p_{\pi,\Phi}, p^w_{\mu,\Phi}) - \mathrm{IPM}_{\mathcal{H}}(\hat{p}_{\pi,\Phi}, \hat{p}^w_{\mu,\Phi})\right| \leq \sqrt{18\nu^2 \log\frac{4}{\delta}C}\left(\frac{1}{\sqrt{m}} + \frac{1}{\sqrt{n}}\right)$$

We let $\sigma^2_Y = \sigma^2_\pi + \sigma^2_\mu$ and

$$\mathcal{D}^{\Phi,\mathcal{H}}_\delta := B_\Phi\sqrt{18\nu^2 \log\frac{4}{\delta}C}$$

Combining these results, observing that $(1-\delta)^2 \geq 1 - 2\delta$, we obtain the desired result. $\qquad\square$

A.3 ASYMPTOTICS

Theorem 2 (Restated). *Suppose $\mathcal{H}$ is a reproducing kernel Hilbert space given by a bounded kernel. Suppose weak overlap holds in that $\mathbb{E}[(p_\pi(x,t)/p_\mu(x,t))^2] < \infty$. Then,*

$$\min_{h,\Phi,w} \mathcal{L}_\pi(h,\Phi,w;\beta)] \leq \min_{f\in\mathcal{F}} R_\pi(f) + O(1/\sqrt{n} + 1/\sqrt{m}) .$$

Proof. Let $f^* = \Phi^* \circ h^* \in \arg\min_{f\in\mathcal{F}} R_\pi(f)$ and let $w^*(x,t) = p_{\pi,\Phi}(\Phi^*(x),t)/p_{\mu,\Phi}(\Phi^*(x),t)$. Since $\min_{h,\Phi,w} \mathcal{L}_\pi(h,\Phi,w;\beta) \leq \mathcal{L}_\pi(h^*,\Phi^*,w^*;\beta)$, it suffices to show that $\mathcal{L}_\pi(h^*,\Phi^*,w^*;\beta) =$

$R_\pi(f^*) + O(1/\sqrt{n} + 1/\sqrt{m})$. We will work term by term:

$$\mathcal{L}_\pi(h^*, \Phi^*, w^*; \beta) = \underbrace{\frac{1}{n}\sum_{i=1}^n w_i \ell_h(\Phi(x_i), t_i)}_{\textcircled{A}} + \lambda_h \underbrace{\frac{\mathcal{R}(h)}{\sqrt{n}}}_{\textcircled{B}} + \alpha \underbrace{\text{IPM}_G(\hat{q}_\Phi, \hat{p}_\Phi^{w^k})}_{\textcircled{C}} + \lambda_w \underbrace{\frac{\|w\|_2}{n}}_{\textcircled{D}}.$$

For term $\textcircled{D}$, letting $w_i^* = w^*(x_i, t_i)$, we have that by weak overlap

$$\textcircled{D}^2 = \frac{1}{n} \times \frac{1}{n}\sum_{i=1}^n (w_i^*)^2 = O_p(1/n),$$

so that $\textcircled{D} = O_p(1/\sqrt{n})$. For term $\textcircled{A}$, under ignorability, each term in the sum in the first term has expectation equal to $R_\pi(f^*)$ and so, so by weak overlap and bounded second moments of loss, we have $\textcircled{A} = R_\pi(f^*) + O_p(1/\sqrt{n})$. For term $\textcircled{B}$, since h^* is fixed we have deterministically that $\textcircled{B} = O(1/\sqrt{n})$.

Finally, we address term $\textcircled{C}$, which when expanded can be written as

$$\sup_{\|h\|\leq 1} \left(\frac{1}{m}\sum_{i=1}^m h(\Phi^*(X_i^q)) - \frac{1}{n}\sum_{i=1}^n w_i^* h(\Phi^*(X_i^p))\right).$$

Let $X_i'^q i = 1, \ldots, n$ and $X_i''^q i = 1, \ldots, m$ be new iid replicates (ghost samples). By Jensen's inequality,

$$\mathbb{E}[\textcircled{C}^2] = \mathbb{E}[\sup_{\|h\|\leq 1}\left(\frac{1}{m}\sum_{i=1}^m h(\Phi^*(X_i^q)) - \frac{1}{n}\sum_{i=1}^n w_i^* h(\Phi^*(X_i^p)))\right)^2]$$

$$= \mathbb{E}[\sup_{\|h\|\leq 1}\left(\frac{1}{m}\sum_{i=1}^m (h(\Phi^*(X_i^q)) - \mathbb{E}[h(\Phi^*(X_i'^q))])\right.$$

$$\left. - \frac{1}{n}\sum_{i=1}^n (w_i^* h(\Phi^*(X_i^p)) - \mathbb{E}[h(\Phi^*(X_i''^q))]))\right)^2]$$

$$\leq \mathbb{E}[\sup_{\|h\|\leq 1}\left(\frac{1}{m}\sum_{i=1}^m (h(\Phi^*(X_i^q)) - h(\Phi^*(X_i'^q)))\right.$$

$$\left. - \frac{1}{n}\sum_{i=1}^n (w_i^* h(\Phi^*(X_i^p)) - h(\Phi^*(X_i''^q))))\right)^2]$$

$$\leq 2\mathbb{E}[\sup_{\|h\|\leq 1}\left(\frac{1}{m}\sum_{i=1}^m (h(\Phi^*(X_i^q)) - h(\Phi^*(X_i'^q)))\right)^2]$$

$$+ 2\mathbb{E}[\sup_{\|h\|\leq 1}\left(\frac{1}{n}\sum_{i=1}^n (w_i^* h(\Phi^*(X_i^p)) - h(\Phi^*(X_i''^q)))\right)^2]$$

Let $\xi_i(h) = h(\Phi^*(X_i^q)) - h(\Phi^*(X_i'^q))$ and let $\zeta_i(h) = w_i^* h(\Phi^*(X_i^p)) - h(\Phi^*(X_i''^q))$. Note that for every h, $\mathbb{E}[\zeta_i(h)] = \mathbb{E}[\xi_i(h)] = 0$. Moreover, $\mathbb{E}[\|\zeta_i\|^2] \leq 4E[K(\Phi^*(X_i^q), \Phi^*(X_i^q))] \leq M$. Similarly, $\mathbb{E}[\|\xi_i\|^2] \leq 2E[(w_i^*)^2]M + 2M \leq M' < \infty$ because of weak overlap. Let ζ_i' for $i = 1, \ldots, n$ be iid replicates of ζ_i (ghost sample) and let ϵ_i be iid Rademacher random variables. Because $\mathcal{H}$ is a Hilbert space, we have that $\sup_{\|h\|\leq 1}(A(h))^2 = \|A\|^2 = \langle A, A\rangle$. Therefore, by

Jensen's inequality,

$$\mathbb{E}[\sup_{\|h\|\leq 1} (\frac{1}{n}\sum_{i=1}^{n}(w_i^* h(\Phi^*(X_i^p)) - h(\Phi^*(X''^q_i))))^2]$$

$$= \mathbb{E}[\sup_{\|h\|\leq 1} (\frac{1}{n}\sum_{i=1}^{n}\zeta_i(h))^2]$$

$$= \mathbb{E}[\sup_{\|h\|\leq 1} (\frac{1}{n}\sum_{i=1}^{n}(\zeta_i(h) - \mathbb{E}[\zeta_i'(h)]))^2]$$

$$\leq \mathbb{E}[\sup_{\|h\|\leq 1} (\frac{1}{n}\sum_{i=1}^{n}(\zeta_i(h) - \zeta_i'(h)))^2]$$

$$= \mathbb{E}[\sup_{\|h\|\leq 1} (\frac{1}{n}\sum_{i=1}^{n}\epsilon_i(\zeta_i(h) - \zeta_i'(h)))^2]$$

$$\leq \frac{4}{n^2}\mathbb{E}[\sup_{\|h\|\leq 1} (\sum_{i=1}^{n}\epsilon_i\zeta_i(h))^2]$$

$$= \frac{4}{n^2}\mathbb{E}[\|\sum_{i=1}^{n}\epsilon_i\zeta_i\|^2]$$

$$= \frac{4}{n^2}\mathbb{E}[\sum_{i,j=1}^{n}\epsilon_i\epsilon_j \langle\zeta_i,\zeta_j\rangle]$$

$$= \frac{4}{n^2}\mathbb{E}[\sum_{i=1}^{n}\|\zeta_i\|^2]$$

$$= \frac{4}{n^2}\sum_{i=1}^{n}\mathbb{E}[\|\zeta_i\|^2]$$

$$\leq \frac{4M'}{n}$$

An analogous argument can be made of ξ_i's, showing that $\mathbb{E}[\text{ⓒ}^2] = O(1/n)$ and hence $\text{ⓒ} = O(1/\sqrt{n})$ by Markov's inequality. $\qquad\square$

B IMPLEMENTATION

We implemented all neural network models (IPM-WNN, RCFR) in TensorFlow as feed-forward fully-connected networks with ELU activations. All architectures have a representation with two hidden layers of 32 and 16 hidden units, and hypotheses (one for each outcome) of 1 layer of 16 hidden units. The networks were trained using stochastic gradient descent with the ADAM optimizer with a learning rate of 10^{-3}. The batch size was 128. Representations were normalized by dividing by the norm. Weight functions were implemented as 2 hidden layers of 32 units each, as functions of the representation Φ. σ in the RBF kernel was set to 1.0. λ_w was set to 0.1 and λ_h to 10^{-4}.