Adversarial Examples in Multi-Layer Random ReLU Networks

Abstract

We consider the phenomenon of adversarial examples in ReLU networks with independent gaussian parameters. For networks of constant depth and with a large range of widths (for instance, it suffices if the width of each layer is polynomial in that of any other layer), small perturbations of input vectors lead to large changes of outputs. This generalizes results of Daniely and Schacham (2020) for networks of rapidly decreasing width and of Bubeck et al (2021) for two-layer networks. The proof shows that adversarial examples arise in these networks because the functions that they compute are very close to linear. Bottleneck layers in the network play a key role: the minimal width up to some point in the network determines scales and sensitivities of mappings computed up to that point. The main result is for networks with constant depth, but we also show that some constraint on depth is necessary for a result of this kind, because there are suitably deep networks that, with constant probability, compute a function that is close to constant.

1 Introduction and Main Result

Since the phenomenon of adversarial examples was first observed in deep networks [SZS+14], there has been considerable interest in why this extreme sensitivity to small input perturbations arises in deep networks [GSS15, SSRD19, BLPR19, DS20, BCGdC21] and how it can be detected and avoided [CW17a, CW17b, FCSG17, MMS+18, QMG+19]. Building on the work of Shamir et al [SSRD19], Daniely and Schacham [DS20] prove that small perturbations (measured in the euclidean norm) can be found for any fixed input and most gaussian parameters in certain ReLU networks—those in which each layer has vanishing width relative to the previous layer—and conjectured the same result without this strong constraint on the architecture. Bubeck, Cherapanamjeri, Gidel and Tachet des Combes [BCGdC21] prove that the same phenomenon occurs in general two-layer ReLU networks, and give experimental evidence of its presence in deeper ReLU networks.

In this paper, we prove that adversarial examples also arise in deep ReLU networks with random weights for a wide variety of network architectures—those with constant depth and polynomially-related widths. The key fact underlying this phenomenon was already observed in [DS20]: a high-dimensional linear function $f(x) = w^\top x$ with input $x \neq 0$ and random parameter vector w with a uniformly chosen direction will satisfy $\|\nabla f(x)\| \|x\| \gg |f(x)|$ with high probability. This implies the existence of a nearby adversarial example for this linear function: a perturbation of x of size $|f(x)|/\|\nabla f(x)\| \ll \|x\|$ in the direction $-f(x)\nabla f(x)$ will flip the sign of $f(x)$. This observation can be extended to nonlinear functions that are locally almost linear. Indeed, it is easy to show that

for all $x, u \in \mathbb{R}^d$,

$$|f(x+u) - (f(x) + \langle u, \nabla f(x) \rangle)| \leq \|u\| \sup \left\{ \|\nabla f(x) - \nabla f(x+v)\| : v \in \mathbb{R}^d, \|v\| \leq \|u\| \right\},$$

and thus to demonstrate the existence of an adversarial example near x for a function f, it suffices to show the smoothness property:

$$\text{for all } v \in \mathbb{R}^d \text{ with } \|v\| \lesssim |f(x)|/\|\nabla f(x)\|, \qquad \|\nabla f(x) - \nabla f(x+v)\| \ll \|\nabla f(x)\|. \quad (1)$$

We show that for a deep ReLU network with random parameters and a high-dimensional input vector x, there is a relatively large ball around x where the function computed by the network is very likely to satisfy this smoothness property. Thus, adversarial examples arise in deep ReLU networks with random weights because the functions that they compute are very close to linear in this sense.

It is important to notice that ReLU networks are not smooth in a classical sense, because the nondifferentiability of the ReLU nonlinearity implies that the gradient can change abruptly. But for the smoothness condition (1), it suffices to have $\|\nabla f(x) - \nabla f(x+v)\| \leq \epsilon + \phi(\|v\|)$, for some increasing function $\phi : \mathbb{R}_+ \to \mathbb{R}_+$, provided that $\epsilon + \phi(\|v\|) \ll \|\nabla f(x)\|$. We prove an inequality like this for ReLU networks, where the ϵ term decreases with width.

Consider a network with input dimension d, $\ell+1$ layers, a single real output, and complete connections between layers. Let $d_1, \ldots, d_\ell$ denote the dimensions of the layers. The network has independent random weight matrices $W_i \in \mathbb{R}^{d_i \times d_{i-1}}$ for $i \in [\ell + 1]$, where we set $d_0 = d$ and $d_{\ell+1} = 1$. For input $x \in \mathbb{R}^d$, the network output is defined as follows:

$$f(x) = W_{\ell+1} \cdot \sigma(W_\ell \cdot \sigma(W_{\ell-1} \cdot \sigma(\cdots \sigma(W_1 \cdot x) \cdots))) \text{ where } \sigma(x)_i = \max\{x_i, 0\}$$

$$W_{\ell+1} \sim \mathcal{N}(0, I/d_\ell) \text{ and } (W_i)_{k,l} \overset{i.i.d}{\sim} \mathcal{N}(0, 1/d_{i-1}). \quad \text{(NN-DEF)}$$

Note that the scale of the parameters is chosen so that all the real-valued signals that appear throughout the network have roughly the same scale. This is only for convenience: because the ReLU is positively homogeneous (that is, for $\alpha > 0$, $\sigma(\alpha x) = \alpha \sigma(x)$), the scaling is not important for our results; the $1/d_{i-1}$ in (NN-DEF) could be replaced by any constant without affecting the ratio between the norm of an input vector and that of a perturbation required to change the sign of the corresponding output.

The following theorem is the main result of the paper.

Theorem 1.1. *Fix $\ell \in \mathbb{N}$. There are constants c_1, c_2, c_3 that depend on ℓ for which the following holds. Fix $\delta \in (0, 1)$ and let $f(\cdot)$ be an $(\ell + 1)$-layer ReLU neural network defined by (NN-DEF) with input dimension d and intermediate layers of width $\{d_i\}_{i=1}^\ell$. Suppose that the widths satisfy*

$$d_{\min} \geq c_1 (\log d_{\max})^{c_2} \log 1/\delta \text{ where } d_{\min} = \min \left\{ \{d_i\}_{i=1}^\ell, d \right\}, d_{\max} = \max \left\{ \{d_i\}_{i=1}^\ell, d \right\}.$$

Then for any fixed input $x \neq 0$, with probability at least $1 - \delta$,

$$|f(x + \eta \nabla f(x))| \geq |f(x)| \text{ and } \mathrm{sign}(f(x + \eta \nabla f(x))) \neq f(x),$$

for an η satisfying

$$\frac{\|\eta \nabla f(x)\|}{\|x\|} \leq c_3 \sqrt{\frac{\log 1/\delta}{d}}.$$

It suffices to choose $c_1 = (C_1 \ell)^{c_2}$, $c_2 = C_2 \ell$, $c_3 = C_3^\ell$, for some absolute constants C_1, C_2, C_3.

This theorem concerns networks of fixed depth, and the constants in the size of the perturbation and in the requirement on the network width are larger for deeper networks. We also prove a converse result that illustrates the need for some constraint on the depth. Theorem 3.1 shows that when the depth is allowed to grow polynomially in the input dimension d, the function computed by a random ReLU network is essentially constant, which rules out the possibility of adversarial examples.

The heart of the proof of Theorem 1.1 is to show a smoothness property like (1). It exploits a decomposition of the change of gradient between two input vectors. Define $H_i : \mathbb{R}^d \to \mathbb{R}^{d_i \times d_i}$ as $H_i(x)_{jk} = \mathbf{1}\{j = k, v_i(x)_j \geq 0\}$ with $v_i(x) = W_i \sigma(\cdots \sigma(W_1 x))$. For two input vectors $x, y \in \mathbb{R}^d$, we will see in Section 2.3 that we can decompose the change of gradient as

$$\nabla f(x) - \nabla f(y) = \sum_{j=1}^\ell W_{\ell+1} \left(\prod_{i=\ell}^{j+1} H_i(x) W_i \right) \cdot (H_j(y) - H_j(x)) W_j \cdot \left(\prod_{k=j-1}^{1} H_i(y) W_i \right).$$

70 (Here and elsewhere, indices of products of matrices run backwards, so $\prod_{i=j}^{k} M_i = I$ when $j < k$.)
71 For the jth term in the decomposition, we need to control the scale of: the gradient of the mapping
72 from the input to the output of layer j, the change in the layer j nonlinearity $H_j(x) - H_j(y)$, and the
73 gradient from layer j to the output. It turns out that controlling these quantities depends crucially
74 on the width of the narrowest layer before layer j—we call this the *bottleneck layer* for layer j.
75 This width determines the dimension of the image at layer j of a ball in the input space. In proving
76 bounds on gradients and function values that hold uniformly over pairs of nearby vectors x and y,
77 this dimension—the width of the bottleneck layer—dictates the size of a discretization (an ϵ-net)
78 that is a crucial ingredient in the proof of these uniform properties. Our analysis involves working
79 separately with the segments between these bottleneck layers. We show that for an input $x \in \mathbb{R}^d$
80 satisfying $\|x\| = \sqrt{d}$ and any y in a ball around x, with high probability $\|\nabla f(x) - \nabla f(y)\| = o(1)$,
81 but $|f(x)|$ is no more than a constant and $\|\nabla f(x)\|$ is at least a constant. This implies the existence
82 of a small $(o(\|x\|))$ perturbation of x in the direction $-f(x)\nabla f(x)$ that flips the sign of $f(x)$.

83 These results suggest several interesting directions for future work. First, our results show that
84 for high-dimensional inputs, adversarial examples are inevitable in random ReLU networks with
85 constant depth, and unlikely in networks with polynomial depth. Beyond this, we do not know
86 how the sensitivity to input perturbations decreases with depth. Similarly, both results are restricted
87 to networks with subexponential width, and it is not clear what happens for very wide networks.
88 Finally, we show that networks with random weights suffer from adversarial examples because their
89 behavior is very similar to that of random linear functions. It would be worthwhile to determine
90 whether randomly initialized trained networks retain this nearly linear behavior, and hence suffer
91 from adversarial examples for the same reason.

2 Proof of Main Theorem

93 In this section, we provide an outline of the proof of Theorem 1.1. As alluded to in Section 1, we will
94 prove our result first by showing that the gradient at x has large norm and changes negligibly in a
95 large ball around x. The first condition is established in Subsection 2.1. However, the second step is
96 much more intricate and requires carefully tracking the ball around x as it propagates through the
97 network. This analysis is carried out in Subsection 2.2 and finally, we use this result in tandem with
98 the decomposition of the gradient differences in Subsection 2.3 to show that our network is locally
99 linear. For the rest of the proof, unless otherwise stated, we consider a fixed $x \in \mathbb{R}^d$, and we assume:

$$d_{\min} \geq (C\ell \log d_{\max})^{240\ell} \log 1/\delta, \quad \|x\| = \sqrt{d} \text{ and } R := \frac{\sqrt{d_{\min}}}{(\ell \log d_{\max})^{80\ell}} = \Omega\left((\ell \log d_{\max})^{40\ell}\right).$$

100 Additionally, we randomize the activations of neurons whenever they receive an input of 0. This
101 does not change the behavior of the neural network in terms of its output or the images of the input
102 through the layers of the network but greatly simplifies our proof. For $x \in \mathbb{R}^d$, we let:

$$x_0 := x, \quad \widetilde{f}_i(x) := W_i f_{i-1}(x), \quad (D_i(y))_{j,k} = \begin{cases} 1 & \text{w.p } \frac{1}{2} \text{ if } j = k, \ y_j = 0, \\ 1 & \text{if } j = k, \ y_j > 0, \\ 0 & \text{otherwise,} \end{cases} \quad x_i = D_i(\widetilde{f}_i(x))\widetilde{f}_i(x).$$

103 Our first key observation is that the randomization in the activation units allows us the following
104 distributional equivalences, proved in Appendix A.1.

105 **Lemma 2.1.** *Let $m \in \mathbb{N}$, $\{d_i\}_{i=0}^{m} \subset \mathbb{R}^d$ and $W_i \in \mathbb{R}^{d_i \times d_{i-1}}$ be distributed such that each entry of*
106 *W_i is drawn iid from any symmetric distribution. Then, defining for $x \in \mathbb{R}^{d_0}$:*

$$\begin{aligned} h_0(x) &= x, \\ \widetilde{h}_i(x) &= W_i h_{i-1}(x) \qquad \text{where } (D_i(y))_{i,j} = \begin{cases} 1, & \text{if } j = k \text{ and } y_j > 0 \\ 1, & \text{with probability } \frac{1}{2} \text{ if } j = k, \ y_j = 0 \\ 0, & \text{otherwise} \end{cases} \\ h_i(x) &= D_i(\widetilde{h}_i(x))\widetilde{h}_i(x) \end{aligned}$$

107 *we have the distributional equivalences for any $x \neq 0$ and fixed diagonal matrices $B_1, \ldots, B_m$:*

$$W_m \prod_{j=m-1}^{1} (D_j(\widetilde{h}_j(x)) + B_j)W_j \stackrel{d}{=} W_m \prod_{j=m-1}^{1} (D_j + B_j)W_j$$

$$\left\| \prod_{j=m}^{1} (D_j(\widetilde{h}_j(x)) + B_j)W_j \right\| \overset{d}{=} \left\| \prod_{j=m}^{1} (D_j + B_j)W_j \right\|$$

$$\textit{where } (D_j)_{k,l} = \begin{cases} 1, & \textit{with probability } 1/2 \textit{ if } k = l \\ 0, & \textit{otherwise} \end{cases}$$

2.1 Concentration of Function Value and Gradient at a Fixed Point

We first present a simple lemma that shows that the gradient at x is at least a constant and that its output value is bounded. The proof crucially uses Lemma 2.1 and illustrates its use through the more involved proofs in the paper.

Lemma 2.2. *For some universal constant c, with probability at least $1 - \delta$ we have:*

$$|f(x)| \leq c2^{\ell}\sqrt{\log 1/\delta} \textit{ and } \|\nabla f(x)\| \geq \frac{1}{2^{\ell+1}}.$$

Proof. Note that

$$\nabla f(x) = W_{\ell+1} \prod_{i=\ell}^{1} D_i(\widetilde{f}_i(x))W_i \textit{ and } f(x) = \nabla f(x)x.$$

And we have from Lemma 2.1, $\nabla f(x) \overset{d}{=} \tilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \tilde{W}_i$, where

$$(D_i)_{j,k} = \begin{cases} 1 & \text{with probability } 1/2 \text{ if } j = k, \\ 0 & \text{otherwise,} \end{cases} \qquad \{W_i\}_{i=1}^{\ell+1} \overset{d}{=} \{\tilde{W}_i\}_{i=1}^{\ell+1}.$$

Therefore, it suffices to analyze the random vector $\tilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \tilde{W}_i$. We first condition on a favorable event for the D_i. Note that we have by the union bound and an application of Hoeffding's inequality (e.g., [BLM13, Theorem 2.8]) that:

$$\forall i \in [\ell] : \operatorname{Tr} D_i \geq \frac{d_i}{3} \text{ with probability at least } 1 - \delta/4,$$

since $d_i \geq c \log(4\ell/\delta)$. We now condition on the D_i and note that:

$$\left\| \widetilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \widetilde{W}_i \right\| \overset{d}{=} \left\| W_{\ell+1}^{\dagger} \prod_{i=\ell}^{1} W_i^{\dagger} \right\| \text{ where}$$

$$\forall i \in \{2, \ldots, \ell\} : W_i^{\dagger} \in \mathbb{R}^{\operatorname{Tr} D_i \times \operatorname{Tr} D_{i-1}}, \ W_{\ell+1}^{\dagger} \in \mathbb{R}^{1 \times \operatorname{Tr} D_{\ell}}, \ W_1^{\dagger} \in \mathbb{R}^{\operatorname{Tr} D_1 \times d}$$

$$(W_1^{\dagger})_{j,:} \sim \mathcal{N}(0, I/d), \ W_{\ell+1}^{\dagger} \sim \mathcal{N}(0, I/d_{\ell}), \ \forall i \in \{2, \ldots, \ell+1\} : (W_i^{\dagger})_{j,:} \sim \mathcal{N}(0, I/d_{i-1}).$$

From the above display, we obtain from Theorem C.3 applied to $f(x) = -\|x\|$:

$$\left\| W_{\ell+1}^{\dagger} \prod_{i=\ell}^{1} W_i^{\dagger} \right\| \geq \frac{1}{2} \cdot \left\| W_{\ell+1}^{\dagger} \prod_{i=\ell}^{2} W_i^{\dagger} \right\| \qquad \text{with probability at least } 1 - \delta/(4\ell)$$

$$\geq \frac{1}{2^{\ell+1}} \qquad \text{with probability at least } 1 - \delta/4 \text{ by induction,}$$

since $\min_{0 \leq i \leq \ell} d_i \geq c \log(4\ell/\delta)$. Through a similar argument, we obtain:

$$\left\| \tilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \tilde{W}_i \right\| \leq 2^{\ell+1} \tag{2}$$

with probability at least $1 - \delta/4$. We also have:

$$\left(\tilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \tilde{W}_i \ \middle| \ \left\| \tilde{W}_{\ell+1} \prod_{i=\ell}^{1} D_i \tilde{W}_i \right\| = m \right) \overset{d}{=} \operatorname{Unif}\left(m\mathbb{S}^{d-1}\right).$$

Combining this, Eq. (2) and Lemma C.5, we get that:

$$|f(x)| \leq c2^{\ell}\sqrt{\log 1/\delta}$$

with probability at least $1 - \delta/2$, for some absolute constant c. A union bound over all the preceeding events concludes the lemma. $\square$

2.2 Scale Preservation of Local Neighborhoods

In this section, we show that the image of a ball around x remains in a ball of suitable radius around the image of x projected through the various layers of the network. This will effectively allow us to decompose the neural network into various segments and analyze each of these segments independently. Here, we introduce additional notation used in the rest of the proof:

$$\forall j \in \{0\} \cup [\ell + 1] : f_{j,j}(x) = x, \quad \forall i > j : \widetilde{f}_{i,j}(x) = W_i f_{i-1,j}(x), \quad f_{i,j}(x) = D_i(\widetilde{f}_{i,j}(x))\widetilde{f}_{i,j}(x)$$

We now describe the decomposition of the neural network into consecutive segments. This decomposition is crucial for reducing the sizes of the ϵ-nets that arising in our proofs. Intuitively, we would like to exploit the smaller dimensionality of the bottleneck layers to reduce the size of the ε-net. The segments will be defined by indices $\{i_j\}_{j=1}^{m}$, denoting bottleneck layers, defined recursively from the output layer backwards with the convention that $d_0 = d$:

$$i_1 := \arg\min_{i \leq \ell} d_i, \quad \forall j > 1 \text{ s.t } i_j \geq 1, i_{j+1} := \arg\min_{j < i_j} d_j, \quad d_{\min}^i = \min_{j < i} d_j. \quad \text{(NN-DECOMP)}$$

Note, that $i_m = 0$, $d_{i_j} < d_{i_{j+1}}$ for all $j \in [m-1]$ and for all $k \in \{d_j, \ldots, d_{j-1}\}, d_k \geq d_j$. The main technical result of subsection is the following lemma whose full proof is deferred to Appendix A.2. In the proof sketch, we carry out the section of the proof where we transition between bottleneck layers in full as it is a simple illustration of how such ideas are used through the rest of the proof.

Lemma 2.3. *We have with probability at least* $1 - \delta$,

$$\forall i \in [\ell] : \|f_i(x)\| \geq \frac{1}{2^i} \cdot \sqrt{d_i},$$

$$\forall \|x' - x\| \leq R : \|\widetilde{f}_i(x) - \widetilde{f}_i(x')\| \leq (C \log d_{\max})^{i/2} \cdot \left(1 + \sqrt{\frac{d_i}{d_{\min}^i}}\right) \cdot R,$$

$$\forall \|x' - x\| \leq R : \|f_i(x) - f_i(x')\| \leq (C \log d_{\max})^{i/2} \cdot \left(1 + \sqrt{\frac{d_i}{d_{\min}^i}}\right) \cdot R.$$

Proof Sketch. The proof of the first claim is nearly identical to that of Lemma 2.2.

For the second claim, we use a gridding argument with some subtleties. Concretely, we construct a new grid over the image of $\mathbb{B}(x, R)$ whenever the number of units in a hidden layer drops below all the previous layers in the network starting from the input layer. These layers are precisely defined by the indices i_j in NN-DECOMP. We now establish the following claim inductively where we adopt the convention $i_0 = \ell + 1$.

Claim 2.4. Suppose for $j \geq 1$ and $\widetilde{R} \leq d_{\max} \cdot R$:

$$\forall y \in \mathbb{B}(x, R) : \|f_{i_j}(x) - f_{i_j}(y)\| \leq \widetilde{R}.$$

Then:

$$\forall i \in \{i_j + 1, \ldots, i_{j-1} - 1\}, y \in \mathbb{B}(x, R) : \|\widetilde{f}_i(x) - \widetilde{f}_i(y)\| \leq (C\ell \log d_{\max})^{(i-i_j)/2} \cdot \sqrt{\frac{d_i}{d_{i_j}}} \cdot \widetilde{R},$$

$$\forall y \in \mathbb{B}(x, R) : \|\widetilde{f}_{i_{j-1}}(x) - \widetilde{f}_{i_{j-1}}(y)\| \cdot \leq (C\ell \log d_{\max})^{(i_{j-1} - i_j)/2} \cdot \widetilde{R}$$

with probability at least $1 - \delta/8l$.

Proof Sketch. We start by constructing an ε-net [Ver18, Definition 4.2.1], $\mathcal{G}$, of $f_{i_j}(\mathbb{B}(x, R))$ with $\varepsilon = 1/(d_{\max}^\ell)^{32}$. Note that we may assume $|\mathcal{G}| \leq (10\widetilde{R}/\varepsilon)^{d_{i_j}}$. We will prove the statement on the grid and extend to the rest of the space. For layer $i + 1$, defining $\widetilde{x} = f_{i_j}(x)$, we have $\forall \widetilde{y} \in \mathcal{G}$:

$$\|\widetilde{f}_{i+1,i_j}(\widetilde{x}) - \widetilde{f}_{i+1,i_j}(\widetilde{y})\| = \|W_{i+1}(f_{i,i_j}(\widetilde{x}) - f_{i,i_j}(\widetilde{y}))\|$$

$$\leq \|f_{i,i_j}(\widetilde{x}) - f_{i,i_j}(\widetilde{y})\| \cdot \sqrt{\frac{d_{i+1}}{d_i}} \cdot \left(1 + \sqrt{\frac{\log 1/\delta'}{d_{i+1}}}\right)$$

with probability at least $1 - \delta'$ as before by Theorem C.3. By setting $\delta' = \delta/(16|\mathcal{G}|\ell^2)$ and noting that $d_i \geq d_{i_j}$, the conclusion holds for layer $i + 1 \leq i_j$ on $\mathcal{G}$ with probability at least $1 - \delta/(16\ell^2)$. By induction and the union bound, we get:

$$\forall i \in \{i_j + 1, \dots, i_{j-1} - 1\}, \widetilde{y} \in \mathcal{G} : \|\widetilde{f}_{i,i_j}(\widetilde{y}) - \widetilde{f}_{i,i_j}(\widetilde{x})\| \leq (C\ell \log d_{\max})^{(i-i_j)/2} \cdot \sqrt{\frac{d_i}{d_{i_j}}} \cdot \widetilde{R}$$

$$\forall \widetilde{y} \in \mathcal{G} : \|\widetilde{f}_{i_{j-1},i_j}(\widetilde{y}) - \widetilde{f}_{i_{j-1},i_j}(\widetilde{x})\| \leq (C\ell \log d_{\max})^{(i_{j-1}-i_j)/2} \cdot \widetilde{R}$$

with probability at least $1 - \delta/(16\ell^2)$. To extend to all $y \in f_{i_j}(\mathbb{B}(x,R))$, we condition on the bound on $\|W_i\|$ given by Lemma C.1 for all $i \leq i_{j-1}$ and note that $\forall y \in f_{i_j}(\mathbb{B}(x,R))$, for $\widetilde{y} = \arg\min_{z \in \mathcal{G}}\|z - y\|$, and for $i_j + 1 \leq i < i_{j-1}$,

$$\|\widetilde{f}_{i,i_j}(\widetilde{x}) - \widetilde{f}_{i,i_j}(y)\| \leq \|\widetilde{f}_{i,i_j}(\widetilde{x}) - \widetilde{f}_{i,i_j}(\widetilde{y})\| + \|\widetilde{f}_{i,i_j}(y) - \widetilde{f}_{i,i_j}(\widetilde{y})\|$$

$$\leq \|\widetilde{f}_{i,i_j}(\widetilde{x}) - \widetilde{f}_{i,i_j}(\widetilde{y})\| + \|y - \widetilde{y}\| \prod_{k=i_j+1}^{i} \|W_k\| \leq \|\widetilde{f}_{i,i_j}(\widetilde{x}) - \widetilde{f}_{i,i_j}(\widetilde{y})\| + \varepsilon \prod_{k=i_j+1}^{i} \left(C\sqrt{\frac{d_k}{d_{k-1}}}\right)$$

$$= \|\widetilde{f}_{i,i_j}(\widetilde{x}) - \widetilde{f}_{i,i_j}(\widetilde{y})\| + \varepsilon C^{i-i_j}\sqrt{\frac{d_i}{d_{i_j}}},$$

using that $d_i \geq d_{i_j}$. Similarly, for $i = i_{j-1}$, we have

$$\|\widetilde{f}_{i_{j-1},i_j}(\widetilde{x}) - \widetilde{f}_{i_{j-1},i_j}(y)\| \leq \|\widetilde{f}_{i_{j-1},i_j}(\widetilde{x}) - \widetilde{f}_{i_{j-1},i_j}(\widetilde{y})\| + \varepsilon C^{i_{j-1}-i_j}\sqrt{\frac{d_{i_{j-1}-1}}{d_{i_j}}}.$$

Our setting of ε concludes the proof of the claim. $\square$

An inductive application of Claim 2.4, a union bound and the observation that:

$$\|f_i(x) - f_i(y)\| \leq \|\widetilde{f}_i(x) - \widetilde{f}_i(y)\|$$

concludes the proof of the lemma. $\square$

The following lemma shows that few neurons have inputs of small magnitude for network input x.

Lemma 2.5. *With probability at least $1 - \delta$, for all $i \in [\ell]$, we have:*

$$\#\left\{j : |\langle (W_{i+1})_j, f_i(x)\rangle| \geq \alpha_i \frac{\|f_i(x)\|}{\sqrt{d_i}}\right\} \geq \left(1 - 2\sqrt{\frac{2}{\pi}}\alpha_i\right)d_{i+1}.$$

Proof. Follows from the fact that $\Pr(|Z| \leq c) \leq c\sqrt{2/\pi}$ for $Z \sim N(0,1)$, plus a simple application of Heoffding's Inequality. $\square$

2.3 Proving Local Gradient Smoothness

Here, we show that the gradient is locally constant completing our proof. We start by observing the following decomposition effectively allowing us to reason about deviations layer by layer:

$$\nabla f(x) - \nabla f(y) = W_{\ell+1}\left(\prod_{i=\ell}^{1} D_i(\widetilde{f}_i(x))W_i - \prod_{i=\ell}^{1} D_i(\widetilde{f}_i(y))W_i\right)$$

$$= W_{\ell+1}\left(\prod_{i=\ell}^{1} D_i(\widetilde{f}_i(x))W_i - \left(D_\ell(\widetilde{f}_\ell(x)) + (D_\ell(\widetilde{f}_\ell(y)) - D_\ell(\widetilde{f}_\ell(x)))\right)W_\ell \prod_{i=\ell-1}^{1} D_i(\widetilde{f}_i(y))W_i\right)$$

$$= W_{\ell+1}\left(D_\ell(\widetilde{f}_\ell(x))W_\ell\left(\prod_{i=\ell-1}^{1}D_i(\widetilde{f}_i(x))W_i - \prod_{i=\ell-1}^{1}D_i(\widetilde{f}_i(y))W_i\right)\right.$$

$$\left.+\ (D_\ell(\widetilde{f}_\ell(x)) - D_\ell(\widetilde{f}_\ell(y)))W_\ell\prod_{i=\ell-1}^{1}D_i(\widetilde{f}_i(y))W_i\right)$$

$$=\sum_{j=1}^{\ell}W_{\ell+1}\left(\prod_{i=\ell}^{j+1}D_i(\widetilde{f}_i(x))W_i\right)\cdot(D_j(\widetilde{f}_j(x)) - D_j(\widetilde{f}_j(y)))W_j\cdot\left(\prod_{i=j-1}^{1}D_i(\widetilde{f}_i(y))W_i\right).$$

(GD-DECOMP)

In our proof, we will bound each of the terms in expansion individually. Before we proceed, we prove a structural lemma on the spectral norm of the matrices appearing in the right-hand-side of the final expression in (GD-DECOMP), allowing us to ignore the portion of the network till the last-encountered bottleneck layer. Define, for all $i > j$, $M_{i,j}(y) = \prod_{k=i}^{j+1}D_k(\widetilde{f}_k(y))W_k$.

Lemma 2.6. *With probability at least $1 - \delta$ over the W_k, we have:*

$$\forall\|y - x\| \le R, j \in [m-1]: \mathbb{P}_{\left\{D_i(\widetilde{f}_i(y))\right\}}\left\{\left\|M_{i_j,i_{j+1}}(y)\right\| \le (C\cdot\ell\cdot\log d_{\max})^{(i_j - i_{j+1})/2}\right\} = 1,$$

where the probability is taken with respect to the $D_i(\widetilde{f}_i(y))$.

We provide the first part of the proof in full as the simplest application of ideas that find further application in the subsequent result establishing bounds on terms in NN-DECOMP; see Appendix A.3.

Proof Sketch. To start, consider a fixed $j \in [m]$ and condition on the conclusion of Lemma 2.3 up to level i_{j+1}. Now, consider an ε-net of $f_{i_{j+1}}(\mathbb{B}(x, R))$, $\mathcal{G}$, with resolution $\varepsilon = 1/(d_{\max}^{\ell})^{32}$. As before, $|\mathcal{G}| \le (d_{\max})^{48\ell d_{i_{j+1}}}$ for some constant C. We additionally will consider subsets

$$\mathcal{S} = \left\{(S_k)_{k=i_j-1}^{i_{j+1}+1} : S_k \subseteq [d_k], |S_k| \le 4d_{i_{j+1}}\right\}.$$

Note that $|\mathcal{G}|\cdot|\mathcal{S}|^2 \le (d_{\max})^{64 l d_{i_{j+1}}}$. For $y \in \mathcal{G}, S^1, S^2 \in \mathcal{S}$, consider the following matrix:

$$M_{y,S^1,S^2}^{i_j,i_{j+1}} = \prod_{k=i_j}^{i_{j+1}+1}(D_k(\widetilde{f}_{k,i_{j+1}}(y))) + (D_{S_k^1} - D_{S_k^2}))W_k \text{ where } (D_S)_{i,j} = \begin{cases}1, \text{if } i = j \text{ and } i \in S \\ 0, \text{otherwise}\end{cases}.$$

We will bound the spectral norm of $M_{y,S^1,S^2}^{i_j,i_{j+1}}$. First, note that:

$$\left\|M_{y,S^1,S^2}^{i_j,i_{j+1}}\right\| \le \left\|\widetilde{M}_{y,S^1,S^2}^{i_j,i_{j+1}}\right\| \text{ where } \widetilde{M}_{y,S^1,S^2}^{i_j,i_{j+1}} := W_{i_j}\prod_{k=i_j-1}^{i_{j+1}+1}(D_k(\widetilde{f}_{k,i_{j+1}}(y)) + (D_{S_k^1} - D_{S_k^2}))W_k$$

and observe that from Lemma 2.1:

$$\widetilde{M}_{y,S^1,S^2}^{i_j,i_{j+1}} \stackrel{d}{=} W_{i_j}\prod_{k=i_j-1}^{i_{j+1}+1}(D_k + (D_{S_k^1} - D_{S_k^2}))W_k) \text{ where } (D_k)_{i,j} = \begin{cases}1, \text{w.p } \frac{1}{2} \text{ if } i = j \\ 0, \text{otherwise}\end{cases}.$$

To bound the spectral norm, let $\mathcal{B}$ be a $1/3$-net of $\mathbb{S}^{d_{i_j}-1}$ and $v \in \mathcal{B}$:

$$\left\|v^\top\widetilde{M}_{y,S^1,S^2}^{i_j,i_{j+1}}\right\| \le \left\|v^\top W_{i_j}\prod_{k=i_j-1}^{i_{j+1}+2}(D_k + (D_{S_k^1} - D_{S_k^2}))W_k)\right\|\cdot\left(1 + \sqrt{\frac{\log 1/\delta'}{d_{i_{j+1}}}}\right)$$

$$\le \prod_{k=i_j}^{i_{j+1}}\left(1 + \sqrt{\frac{\log 1/\delta'}{d_k}}\right)$$

with probability at least $\ell\delta'$. But setting $\delta' = \delta/(16\ell^4 \cdot |\mathcal{G}| \cdot |\mathcal{S}| \cdot |\mathcal{B}|)$ yields with probability at least $1 - \delta/16$:

$$\forall y \in \mathcal{G}, S_1, S_2 \in \mathcal{S} : \left\| \widetilde{M}_{y,S^1,S^2}^{i_j,i_{j+1}} \right\| \leq (C \cdot \ell \cdot \log d_{\max})^{(i_j - i_{j+1})/2}.$$

On the event in the conclusion of Lemma 2.3, we have that $f_i(y) \neq 0$ for all $i \in [\ell], y \in \mathcal{G}$ and therefore, we have by a union bound over the discrete set $\mathcal{G}$:

$$\forall y \in \mathcal{G}, k \in \{i_{j+1}, \dots, i_j\}, m \in [d_k] : (\widetilde{f}_{k,i_j}(y))_k \neq 0$$

proving the lemma for $y \in \mathcal{G}$ as the activations are deterministic. For $y \notin \mathcal{G}$, the following claim concludes the proof of the lemma. The claim is essentially a generalization of [BCGdC21, Eq. (18)] and its proof is deferred to the appendix.

Claim 2.7. We have for all $m \in \{i_{j+1}, \dots, i_j - 1\}, y \in f_{i_j}(\mathbb{B}(x, R))$:

$$\mathbb{P}_{\left\{ D_n(\widetilde{f}_n(y)), D_n(\widetilde{f}_n(\widetilde{y})) \right\}} \left\{ \mathrm{Tr}|D_m(\widetilde{f}_{m,i_{j+1}}(y)) - D_m(\widetilde{f}_{m,i_{j+1}}(\widetilde{y}))| \leq 4d_{i_{j+1}} \right\} = 1$$

where $\widetilde{y} = \arg\min_{z \in \mathcal{G}} \|z - y\|$ with probability at least $1 - \delta'/\ell^2$.

The previous claim along with our previously established bounds establish the lemma. $\qquad\square$

Our final technical result establishes the near-linearity of f in a ball around x. The full proof is deferred to Appendix A.4 but in our proof sketch we identify sections which involve considerations unique to this lemma.

Lemma 2.8. *We have for some absolute constant C:*

$$\forall \|x - y\| \leq R, j \in [\ell] : \mathbb{P}_{\left\{ D_m(\widetilde{f}_m(x)), D_m(\widetilde{f}_m(y)) \right\}} \left\{ \|\nabla f(x) - \nabla f(y)\| \leq \left(\frac{C^\ell}{\log^\ell d_{\max}} \right) \right\} = 1$$

with probability at least $1 - \delta$.

Proof Sketch. Consider a fixed term from (GD-DECOMP); that is, consider:

$$\mathrm{Diff}_j(y) := W_{\ell+1} \left(\prod_{i=\ell}^{j+1} D_i(\widetilde{f}_i(x)) W_i \right) \cdot (D_j(\widetilde{f}_j(y)) - D_j(\widetilde{f}_j(x))) W_j \cdot \left(\prod_{k=j-1}^{1} D_i(\widetilde{f}_i(y)) W_i \right).$$

We will show with high probability that $\mathrm{Diff}_j(y)$ is small for all $y \in \mathbb{B}(x, R)$. This will then imply the lemma by a union bound and (GD-DECOMP). Let k be such that $i_k = \arg\min_{m<j} d_m$. We will condition on the weights of the network up to layer i_k. Specifically, we will assume the conclusions of Lemmas 2.3 and 2.6 up to layer i_k. We may now focus our attention solely on the segment of the network beyond layer i_k as a consequence of the following observation and Lemma 2.6:

$$\|\mathrm{Diff}_j(y)\| \leq \|\mathrm{Diff}_{j,k}(x,y)\| \cdot \|M_{i_k,0}(y)\| \quad \text{where} \tag{3}$$

$$\mathrm{Diff}_{j,k}(x,y) := W_{\ell+1} \left(\prod_{i=\ell}^{j+1} D_i(\widetilde{f}_i(x)) W_i \right) (D_j(\widetilde{f}_j(y)) - D_j(\widetilde{f}_j(x))) W_j \left(\prod_{k=j-1}^{i_k} D_i(\widetilde{f}_i(y)) W_i \right)$$

We will show for all y such that $\|y - x\| \leq R$:

$$\mathbb{P}_{\left\{ D_m(\widetilde{f}_m(x)), D_m(\widetilde{f}_m(y)) \right\}} \left\{ \|\mathrm{Diff}_{j,k}(x,y)\| \geq \frac{C^\ell}{(\ell \log d_{\max})^{3\ell}} \right\} = 0 \tag{4}$$

with probability at least $1 - \delta/(16\ell^2)$.

We have from Lemma 2.1 that the random vector H defined below is spherically symmetric and satisfies $\|H\| \leq 2^\ell$ with probability at least $1 - \delta/(16\ell^4)$:

$$W_{\ell+1} \prod_{j=\ell}^{j+1} D_i(\widetilde{f}_i(x)) W_i \overset{d}{=} \widetilde{W}_{\ell+1} \prod_{j=\ell}^{j+1} D_i \widetilde{W}_i =: H$$

As in the proof of Lemma 2.6, let $\mathcal{G}$ be an ε-net of $f_{i_k}(\mathbb{B}(x, R))$ with ε as in the proof of Lemma 2.6. We now break into two cases depending on how d_j compares to d_{i_k} and handle them separately. At this point, we have effectively reduced the multi-layer proof to the problem of analyzing deviations of activations at a fixed layer. For the remaining proof, we generalize approaches in [DS20] for the small width case and [BCGdC21] for the large width case.

Case 1: $d_j \leq d_{i_k}(\ell \log d_{\max})^{20\ell}$. In this case, the key observation already made in [DS20] is that under Lemma 2.3, the number of neurons that may actually differ at layer j is at most $d_j/\mathrm{poly}(\ell \log d_{\max})^{\ell}$ between $y \in \mathbb{B}(x, R)$ and x. Since, H is spherically distributed, $\|H(D_j(\widetilde{f}_j(x)) - D_j(\widetilde{f}_j(y)))\|$ is very small and consequently the whole term is small.

Case 2: $d_j \geq d_{i_k}(\ell \log d_{\max})^{20\ell}$. This case is analogous to [BCGdC21]. The proof is technically involved and requires careful analysis of the random vector $H(D_j(\widetilde{f}_j(x)) - D_j(\widetilde{f}_j(y)))W_j$ which is complicated in our setting due to the matrices preceeding it in (GD-DECOMP).

A union bound and an application of the triangle inequality now imply the lemma. $\qquad\square$

Proof of Theorem 1.1. On the intersection of the events in the conclusions of Lemmas 2.2, 2.3 and 2.8, we have that $\nabla f(x)$ is deterministic and furthermore, we have:

$$\|\nabla f(x)\| \geq \frac{1}{2^{\ell+1}}, \quad |f(x)| \leq c2^{\ell}\sqrt{\log 1/\delta}$$

$$\forall y \text{ s.t } \|y - x\| \leq R : \|\nabla f(x) - \nabla f(y)\| \leq \frac{1}{(\ell \log d_{\max})^{\ell}} = o(1).$$

Assume $f(x) > 0$ (the alternative is similar) and let $\eta = -\frac{2^{\ell}\log d \cdot \sqrt{\log 1/\delta}}{\|\nabla f(x)\|^2}$, we have for the point $x + \eta\nabla f(x)$, defining the function $g(t) = f(x + t\eta\nabla f(x))$:

$$f(x + \eta\nabla f(x)) = f(x) + \int_0^1 g'(t)dt = f(x) + \int_0^1 (\eta\nabla f(x))^{\top}\nabla f(x + t\eta\nabla f(x))dt$$

$$= f(x) - (1 - o(1))2^{\ell}\log d\sqrt{\log 1/\delta} \leq -f(x).$$

Our lower bounds on $\nabla f(x)$ ensure $\|\eta\nabla f(x)\| \leq R$ concluding the proof of the theorem. $\qquad\square$

3   The Impact of Depth

Theorem 1.1 relies on the depth being constant. In this section, we show that some constraint on the depth is necessary in order to ensure the existence of adversarial examples. In particular, the following result gives an example of a sufficiently deep network for which, with high probability, the output of the network will have the same sign for all input patterns.

Theorem 3.1. *Fix a sufficiently large* $d \in \mathbb{N}$, *an* $\ell \geq d^3$ *and* $(\ell d)^{20} \leq k \leq \exp(\sqrt{\ell})$, *and consider the randomly initialized neural network* (NN-DEF) *with* $d_1 = \cdots = d_{\ell} = k$. *There is a universal constant* C *such that with probability at least* 0.9,

$$\forall x, y \in \mathbb{S}^{d-1} : \frac{|f(x) - f(y)|}{|f(x)|} \leq C\sqrt{\frac{\log d}{d}}.$$

The proof involves showing that the image of $\mathbb{S}^{d-1}$ is bounded away from 0, and that the inner products between images of two input vectors throughout the network converge. The following lemma shows how the expected inner products evolve through the network. The lemma follows from computing a double integral; see [CS09, Eq. (6)]. The proof of Theorem 3.1 is in Appendix B.

Lemma 3.2. *Let* $d \in \mathbb{N}$. *Fix* $x, y \in \mathbb{R}^d$ *with* $x, y \neq 0$. *Then for* $g \sim \mathcal{N}(0, I)$:

$$\frac{\mathbb{E}\left[\max\{g^{\top}x, 0\}\max\{g^{\top}y, 0\}\right]}{\sqrt{\mathbb{E}(\max\{g^{\top}x, 0\})^2 \cdot \mathbb{E}(\max\{g^{\top}y, 0\})^2}} = \frac{\sin\theta}{\pi} + \left(1 - \frac{\theta}{\pi}\right)\cos\theta$$

where $\theta = \arccos(x^{\top}y/(\|x\|\|y\|))$.

References

[BCGdC21] Sébastien Bubeck, Yeshwanth Cherapanamjeri, Gauthier Gidel, and Rémi Tachet des Combes. A single gradient step finds adversarial examples on random two-layers neural networks. *arXiv*, 2104.03863, 2021. 1, 8, 9

[BLM13] Stéphane Boucheron, Gábor Lugosi, and Pascal Massart. *Concentration inequalities*. Oxford University Press, Oxford, 2013. 4, 26

[BLPR19] Sebastien Bubeck, Yin Tat Lee, Eric Price, and Ilya Razenshteyn. Adversarial examples from computational constraints. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 831–840. PMLR, 2019. 1

[CS09] Youngmin Cho and Lawrence Saul. Kernel methods for deep learning. In Y. Bengio, D. Schuurmans, J. Lafferty, C. Williams, and A. Culotta, editors, *Advances in Neural Information Processing Systems*, volume 22. Curran Associates, Inc., 2009. 9

[CW17a] Nicholas Carlini and David Wagner. Towards evaluating the robustness of neural networks. In *2017 IEEE Symposium on Security and Privacy (SP)*, pages 39–57, 2017. 1

[CW17b] Nicholas Carlini and David A. Wagner. Adversarial examples are not easily detected: Bypassing ten detection methods. In Bhavani M. Thuraisingham, Battista Biggio, David Mandell Freeman, Brad Miller, and Arunesh Sinha, editors, *Proceedings of the 10th ACM Workshop on Artificial Intelligence and Security, AISec@CCS 2017, Dallas, TX, USA, November 3, 2017*, pages 3–14. ACM, 2017. 1

[DS20] Amit Daniely and Hadas Shacham. Most ReLU networks suffer from ℓ^2 adversarial perturbations. In H. Larochelle, M. Ranzato, R. Hadsell, M. F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 6629–6636. Curran Associates, Inc., 2020. 1, 9

[FCSG17] Reuben Feinman, Ryan R. Curtin, Saurabh Shintre, and Andrew B. Gardner. Detecting adversarial samples from artifacts. *arXiv*, 1703.00410, 2017. 1

[GSS15] Ian J. Goodfellow, Jonathon Shlens, and Christian Szegedy. Explaining and harnessing adversarial examples. In Yoshua Bengio and Yann LeCun, editors, *3rd International Conference on Learning Representations, ICLR 2015, San Diego, CA, USA, May 7-9, 2015, Conference Track Proceedings*, 2015. 1

[MMS+18] Aleksander Madry, Aleksandar Makelov, Ludwig Schmidt, Dimitris Tsipras, and Adrian Vladu. Towards deep learning models resistant to adversarial attacks. In *International Conference on Learning Representations*, 2018. 1

[QMG+19] Chongli Qin, James Martens, Sven Gowal, Dilip Krishnan, Krishnamurthy Dvijotham, Alhussein Fawzi, Soham De, Robert Stanforth, and Pushmeet Kohli. Adversarial robustness through local linearization. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019. 1

[SSRD19] Adi Shamir, Itay Safran, Eyal Ronen, and Orr Dunkelman. A simple explanation for the existence of adversarial examples with small hamming distance. *arXiv*, 1901.10861, 2019. 1

[SZS+14] Christian Szegedy, Wojciech Zaremba, Ilya Sutskever, Joan Bruna, Dumitru Erhan, Ian Goodfellow, and Rob Fergus. Intriguing properties of neural networks. In *International Conference on Learning Representations*, 2014. 1

[Ver18] Roman Vershynin. *High-dimensional probability*, volume 47 of *Cambridge Series in Statistical and Probabilistic Mathematics*. Cambridge University Press, Cambridge, 2018. 5, 13, 21