Stability and Generalization Analysis of Gradient Methods for Shallow Neural Networks

Abstract

While significant theoretical progress has been achieved, unveiling the generalization mystery of overparameterized neural networks still remains largely elusive. In this paper, we study the generalization behavior of shallow neural networks (SNNs) by leveraging the concept of algorithmic stability. We consider gradient descent (GD) and stochastic gradient descent (SGD) to train SNNs, for both of which we develop consistent excess risk bounds by balancing the optimization and generalization via early-stopping. As compared to existing analysis on GD, our new analysis requires a relaxed overparameterization assumption and also applies to SGD. The key for the improvement is a better estimation of the smallest eigenvalues of the Hessian matrices of the empirical risks and the loss function along the trajectories of GD and SGD by providing a refined estimation of their iterates.

1 Introduction

Neural networks have achieved remarkable success in solving large-scale machine learning problems in various application domains such as computer vision and natural language processing [32]. First-order methods such as gradient descent (GD) and stochastic gradient descent (SGD) are mainstream optimization algorithms for training neural networks due to their simplicity and efficiency [11, 32, 46]. Although the associated optimization problems are nonconvex and nonsmooth, GD/SGD can still find a model with a very small or even zero training error [16, 20, 33, 38, 58, 61]. At the same time, the models found by such first-order methods has demonstrated good generalization performance on test data despite neural networks are often highly overparameterized in the sense that the number of parameters is much larger than the size of training examples [1, 2, 5].

These surprising phenomena have triggered a surge of research activities in understanding the generalization ability of neural networks. Generalization analysis typically uses complexity measures such as VC dimension, covering numbers or Rademacher complexities to develop capacity-dependent bounds [8, 9], which, however, may not explain well the generalization of overparameterized neural networks. Impressive alternatives have been proposed which include the compression approach [4], the norm-based analysis [8, 25], the PAC-Bayes analysis [21] and the neural tangent kernel (NTK) approach [5, 27]. In particular, the NTK approach shows that the overparameterization pulls the dynamic of GD on neural networks close to its counterpart on a kernelized machine with the least-square loss [5, 20], which shows how overparameterization can help both optimization and generalization. However, this approach often requires a very high overparameterization to gain useful results [6, 50, 54].

The recent appealing work [47] presents a kernel-free approach to study how overparameterization would improve the generalization for shallow neural networks (SNNs). Their basic tool is the algorithmic stability [12], which measures how the replacement of an observation would change the

algorithm output. The authors showed the excess risk of GD is controlled by an interpolating network with the shortest GD path from the initialization, which is able to recover the existing NTK-based risk bounds as an application. This result is achieved under an overparameterization assumption $m \gtrsim (\eta T)^5$, where m is the number of hidden nodes, η is the learning rate (step size) and T is the number of iterations. While this result is very interesting and impressive, the overparameterization requirement $m \gtrsim (\eta T)^5$ may still be more restrictive than that used in practice. Furthermore, the analysis in [47] is restricted to the case of the full-batch GD. One natural question thus arises:

> *Can we relax the overparameterization requirement for GD in [47] and further establish the stability and generalization of SGD for neural networks?*

In this paper, we provide an affirmative answer to the above question by establishing a refined stability analysis for the gradient methods (GD and SGD) for training SNNs. Our contributions are summarized as follows.

1. We develop excess risk bounds for GD on SNNs under a relaxed overparameterization. In more details, we show that GD can achieve the excess risk bounds of the order $O(1/\sqrt{n})$ if $m \gtrsim (\eta T)^3$, where n is the sample size. This improves the existing overparameterization condition $m \gtrsim (\eta T)^5$ [47]. Under a low noise condition, our excess risk bounds improve to $O(1/n)$.

2. One key technical novelty in relaxing the overparameterization condition for GD in [47] is to improve the existing bounds on the norm of iterate sequence $\{\mathbf{W}_t\}$. As we soon show in Section 4.1 below, this improvement is achieved by a better estimation of the smallest eigenvalue of the Hessian matrix of the empirical risk. Specifically, the analysis [47] uses $\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 = O(\sqrt{\eta t})$ to lower-bound the smallest eigenvalue at $\alpha \mathbf{W}_t + (1 - \alpha)\mathbf{W}_t^{(i)}$ by $\frac{-1}{\sqrt{m}}(\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 + 1)$, where $\alpha \in (0, 1)$ and $\{\mathbf{W}_t^{(i)}\}$ is an iterate sequence on a neighboring dataset. As a comparison, we show $\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 = O(n^{-1}(\eta t)^{\frac{3}{2}})$ which can be much better than $O(\sqrt{\eta t})$ if n is large. Furthermore, our bound depends on the training errors and would improve in a low noise condition. Under some specific cases, we can further show that $\mathbb{E}[\|\mathbf{W}_t\|_2^2] = O(1)$, which is independent of the iteration number.

3. We extend our analysis to SGD under the relaxed overparameterization condition $m \gtrsim (\eta T)^3$. As compared to GD, SGD has a computational advantage in the sense that it can achieve the same risk bounds with a less computational cost. The key analysis of SGD relies on the estimation of the Hessian spectrum of the loss over the individual training datum. This is more challenging than estimating the counterpart of the empirical risk of GD since several properties of GD do not hold for SGD such as the monotonicity of the objective functions along the optimization process. To overcome this technical hurdle, we provide a refined analysis to control the bounds of the iterates of SGD which further leads to the estimation of the Hessian spectrum of the loss.

The remaining parts of the paper are organized as follows. We present the related work in Section 2 and illustrate the formulation of the problem in Section 3. We present the main results in Section 4 and conclude the paper in Section 5.

2   Related Work

In this section, we group the related work into two categories: the related work on stability analysis and the related work on generalization analysis of neural networks.

Stability and generalization. As a fundamental concept in statistical learning theory, algorithmic stability considers how the perturbation of training examples would affect the output of an algorithm [49], which has a close connection to the learnability [43, 51]. The framework of using the concept of algorithmic stability to derive generalization bounds was established in an influential paper [12], where the uniform stability was introduced and was studied for regularization schemes. Since then, various concepts of stability have been introduced to study the generalization gaps, including the hypothesis stability [12, 22], on-average stability [31, 51], Bayes stability [37], locally elastic stability [19] and argument/model stability [34, 40]. A very successful application of stability analysis is to use it to study SGD for smooth, Lipschitz and convex problems [26], which motivates a

lot of follow-up studies on stochastic optimization [3, 10, 15, 30, 34, 35]. The smoothness assumption in [26] was recently removed by taking very small step sizes [10, 34], while the convexity assumption was relaxed to a weak convexity assumption [48]. Under a Polyak-Lojasiewicz (PL) condition, it was shown that any algorithm converge to global minima would generalize without convexity conditions [15, 35]. The trade-off between stability and optimization was studied in [17]. Other than stochastic optimization, stability has found wide applications in structured prediction [41], meta learning [42], transfer learning [31], hyperparameter optimization [7], minimax problems [23, 36, 59] and adversarial training [55]. While most of the stability analysis imply generalization bounds in expectation, recent studies show that uniform stability can yield almost optimal high-probability bounds [13, 24, 29].

Generalization analysis of Neural Networks (NNs). Generalization analysis of NNs has attracted increasing attention to understand their great success in practice. A popular approach to study the generalization of SNNs is via the uniform convergence approach, which studies the uniform generalization gaps in a hypothesis space [8, 25, 45]. However, this approach leads to capacity-based bounds which do not well explain why overparameterized models can still generalize well to testing examples [44]. To address this problem, researchers turn to other approaches such as the compression approach [4], the PAC-Bayes approach [21], the NTK approach [27] and the neural tangent random feature approach [14]. The key idea of the NTK approach is that, under sufficient overparameterization and random initialization, the dynamics of GD on SNNs is close to the dynamics of GD on a least-squares problem associated to the NTK [5, 20]. This leads to generalization bounds based on a data-dependent complexity measure, which can distinguish the difference between learning with random labels and learning with true labels [5]. Meanwhile, recent studies suggest the connection to kernels might be only good at interpreting the performance of very wide networks [6, 50, 54], much more overparameterized than those used in reality [47]. The most related work is the recent analysis of GD for SNNs without either the NTK condition or the PL condition [47]. They developed nontrivial generalization bounds under an overparameterization assumption $m \gtrsim (\eta T)^5$. Furthermore, their analysis allows for improved bounds if there is no label noise, and shows an interesting connection to NTK-based risk bounds. It should be mentioned that the analysis in [5] considers the ReLU activation function, while the discussions in [47] focus on smooth activation functions.

3 Problem Setup

Let P be a probability distribution defined on a sample space $\mathcal{Z} := \mathcal{X} \times \mathcal{Y}$, where $\mathcal{X} \subseteq \mathbb{R}^d$ and $\mathcal{Y} \subseteq \mathbb{R}$. Let $S = \{\mathbf{z}_i = (\mathbf{x}_i, y_i) : i = 1, \ldots, n\}$ be a sample drawn from P. Based on S we wish to build a model $f : \mathcal{X} \mapsto \mathbb{R}$. The performance of f can be measured by the population risk defined as

$$L(f) = \frac{1}{2} \iint_{\mathcal{X} \times \mathcal{Y}} (f(\mathbf{x}) - y)^2 dP(\mathbf{x}, y),$$

which is unknown and can be approximated by the empirical risk $L_S(f) = \frac{1}{2n} \sum_{i=1}^n (f(\mathbf{x}_i) - y_i)^2$. A minimizer of the population risk is the regression function $f_\rho(\mathbf{x}) = \mathbb{E}[y|\mathbf{x}]$, where $\mathbb{E}[\cdot|\mathbf{x}]$ denotes the conditional expectation given $\mathbf{x}$. In this paper, we consider a shallow neural network of the form

$$f_{\mathbf{W}}(\mathbf{x}) := \sum_{k=1}^m \mu_k \sigma(\langle \mathbf{w}_k, \mathbf{x} \rangle),$$

where we fix $\mu_k \in \{\frac{1}{\sqrt{m}}, -\frac{1}{\sqrt{m}}\}$, $\sigma : \mathbb{R} \mapsto \mathbb{R}$ is an activation function and $\mathbf{W} = (\mathbf{w}_1, \ldots, \mathbf{w}_m) \in \mathbb{R}^{d \times m}$ is the weight matrix. In the above formulation, $\mathbf{w}_k$ denotes the weight of the edge connecting the input to the k-th hidden node, and μ_k is the weight of the edge connecting the k-th hidden node to the output node. Here m is the number of nodes in the hidden layer and $\langle \cdot, \cdot \rangle$ denotes the inner product operator. For simplicity, we denote

$$L(\mathbf{W}) = L(f_{\mathbf{W}}) \quad \text{and} \quad L_S(\mathbf{W}) = L_S(f_{\mathbf{W}}).$$

Let $\mathbf{W}^* = \arg\min_{\mathbf{W}} L(\mathbf{W})$. We choose a minimizer of $L(\mathbf{W})$ with the smallest norm. The relative behavior of a model $\mathbf{W}$ w.r.t. $\mathbf{W}^*$ is quantified by the excess population risk $L(\mathbf{W}) - L(\mathbf{W}^*)$. We denote by $\ell(\mathbf{W}; \mathbf{z}) = \frac{1}{2}(f_{\mathbf{W}}(\mathbf{x}) - y)^2$ the loss function of $\mathbf{W}$ on a single example $\mathbf{z} = (\mathbf{x}, y)$. Two representative algorithms to minimize the empirical risk are GD and SGD.

Definition 1 (Gradient Descent). Let $\mathbf{W}_0 \in \mathbb{R}^{d \times m}$ be an initialization point. GD updates $\{\mathbf{W}_t\}$ as follows

$$\mathbf{W}_{t+1} = \mathbf{W}_t - \eta \nabla L_S(\mathbf{W}_t), \tag{3.1}$$

where $\eta > 0$ is the step size and ∇ denotes the gradient operator.

Definition 2 (Stochastic Gradient Descent). Let $\mathbf{W}_0 \in \mathbb{R}^{d \times m}$ be an initialization point. SGD updates $\{\mathbf{W}_t\}$ as follows

$$\mathbf{W}_{t+1} = \mathbf{W}_t - \eta \nabla \ell(\mathbf{W}_t; \mathbf{z}_{i_t}), \tag{3.2}$$

where i_t is drawn from the uniform distribution over $[n] := \{1, \ldots, n\}$.

We are interested in the excess population risk of models trained by GD/SGD with T iterations. We begin with the introduction of some assumptions on activations and loss functions. Assumptions 1, 2 were also imposed in [47]. We denote by $\|\cdot\|_2$ the Frobenius norm.

Assumption 1 (Activation). The activation $\phi(u)$ is continuous and twice differentiable with constant $B_\phi, B_{\phi'}, B_{\phi''} > 0$ bounding $|\phi(u)| \leq B_\phi, |\phi'(u)| \leq B_{\phi'}$ and $|\phi''(u)| \leq B_{\phi''}$ for any $u \in \mathbb{R}$.

Activation functions satisfying Assumption 1 include sigmoid and hyperbolic tangent activations [47].

Assumption 2 (Inputs, labels, and the loss function). There exists constants $C_x, C_y, C_0 > 0$ such that $\|\mathbf{x}\|_2 \leq C_x, |y| \leq C_y$ and $\ell(\mathbf{W}_0; \mathbf{z}) \leq C_0$ for any $\mathbf{x}, y$ and $\mathbf{z}$.

Our third assumption is on the regularity of the learning problems. For any $\lambda > 0$, we define

$$\mathbf{W}_\lambda^* = \arg \min_{\mathbf{W} \in \mathbb{R}^{d \times m}} \{L(\mathbf{W}) + \lambda \|\mathbf{W} - \mathbf{W}_0\|_2^2\}.$$

Assumption 3 (Regularity). Assume there exist $\alpha \in (0, 1]$ and $c_\alpha > 0$ such that

$$\Lambda_\lambda := L(\mathbf{W}_\lambda^*) - L(\mathbf{W}^*) + \lambda \|\mathbf{W}_\lambda^* - \mathbf{W}_0\|_2^2 \leq c_\alpha \lambda^\alpha.$$

Assumption 3 is related to the approximation error which characterize how well the SNNs approximate the least population risk [18, 53]. If $\|\mathbf{W}^*\|_2 = O(1)$, then it is clear that

$$L(\mathbf{W}_\lambda^*) - L(\mathbf{W}^*) + \lambda \|\mathbf{W}_\lambda^* - \mathbf{W}_0\|_2^2 \leq L(\mathbf{W}^*) - L(\mathbf{W}^*) + \lambda \|\mathbf{W}^* - \mathbf{W}_0\|_2^2 = O(\lambda) \tag{3.3}$$

and therefore Assumption 3 holds with $\alpha = 1$. Our analysis is based on the following error decomposition of the excess risk:

$$
\begin{aligned}
\mathbb{E}[L(\mathbf{W}_T)] - L(\mathbf{W}^*) = {} & \Big[\mathbb{E}[L(\mathbf{W}_T)] - \mathbb{E}[L_S(\mathbf{W}_T)]\Big] \\
& + \mathbb{E}\Big[L_S(\mathbf{W}_T) - L_S(\mathbf{W}_{\frac{1}{\eta T}}^*) - \frac{1}{\eta T}\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2\Big] \\
& + \Big[L(\mathbf{W}_{\frac{1}{\eta T}}^*) + \frac{1}{\eta T}\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2 - L(\mathbf{W}^*)\Big],
\end{aligned}
$$

where we have used $\mathbb{E}[L_S(\mathbf{W}_{\frac{1}{\eta T}}^*)] = L(\mathbf{W}_{\frac{1}{\eta T}}^*)$ due to the independence between $\mathbf{W}_{\frac{1}{\eta T}}^*$ and S. We refer to the first term $\mathbb{E}[L(\mathbf{W}_T)] - \mathbb{E}[L_S(\mathbf{W}_T)]$ as the generalization error (generalization gap) and the second term $\mathbb{E}[L_S(\mathbf{W}_T) - L_S(\mathbf{W}_{\frac{1}{\eta T}}^*)] - \frac{1}{\eta T}\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2$ as the optimization error. As in [47], we will use the on-average model stability to control the generalization error and tools in optimization theory to control the optimization error. We will use Assumption 3 to control the last term $L(\mathbf{W}_{\frac{1}{\eta T}}^*) + \frac{1}{\eta T}\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2 - L(\mathbf{W}^*)$. The on-average model stability considers the sensitivity of the output models up to the perturbation of a single example, and the sensitivity is averaged by traversing the single example throughout the sample set. Let $A(S)$ be the output model by applying an algorithm A to S.

Definition 3 (On-average Model Stability [34]). Let $S = \{\mathbf{z}_1, \ldots, \mathbf{z}_n\}$ and $S' = \{\mathbf{z}_1', \ldots, \mathbf{z}_n'\}$ be drawn independently from P. For any $i \in [n]$, define $S^{(i)} = \{\mathbf{z}_1, \ldots, \mathbf{z}_{i-1}, \mathbf{z}_i', \mathbf{z}_{i+1}, \ldots, \mathbf{z}_n\}$ as the set formed from S by replacing the i-th element with $\mathbf{z}_i'$. We say a randomized algorithm A is on-average model ϵ-stable if $\mathbb{E}_{S,S',A}\big[\frac{1}{n}\sum_{i=1}^n \|A(S) - A(S^{(i)})\|_2^2\big] \leq \epsilon^2$.

The connection between the generalization error and the on-average model stability was established in the following lemma.

Definition 4. We say a function $\mathbf{W} \mapsto g(\mathbf{W})$ is ρ-smooth if, for any $\mathbf{W}$ and $\mathbf{W}'$, we have

$$\|\nabla g(\mathbf{W}) - \nabla g(\mathbf{W}')\|_2 \leq \rho \|\mathbf{W} - \mathbf{W}'\|_2.$$

Lemma 1 (Stability and Generalization [34]). *Let A be an algorithm. If for any $\mathbf{z}$, the map $\mathbf{W} \mapsto \ell(\mathbf{W}; \mathbf{z})$ is ρ-smooth and nonnegative, then*

$$\mathbb{E}[L(A(S)) - L_S(A(S))] \leq \frac{\rho}{2n} \sum_{i=1}^{n} \mathbb{E}[\|A(S) - A(S^{(i)})\|_2^2] + \Big(\frac{2\rho \mathbb{E}[L_S(A(S))]}{n} \sum_{i=1}^{n} \mathbb{E}[\|A(S) - A(S^{(i)})\|_2^2] \Big)^{\frac{1}{2}}.$$

4 Main Results

In this section, we present our main results on the risk bounds of GD and SGD which are summarized in Table 1.

Algorithm	Excess risk bound	Low noise	overparameterization	Computation
GD [47]	$O(n^{-\frac{\alpha}{1+\alpha}})$	No	$m \gtrsim (\eta T)^5 \asymp n^{\frac{5}{\alpha+1}}$	$O(n^{\frac{\alpha+2}{\alpha+1}})$
	$O(n^{-\alpha})$	Yes	$m \gtrsim (\eta T)^5 \asymp n^5$	$O(n^2)$
GD	$O(n^{-\frac{\alpha}{1+\alpha}})$	No	$m \gtrsim (\eta T)^3 \asymp n^{\frac{3}{\alpha+1}}$	$O(n^{\frac{\alpha+2}{\alpha+1}})$
This work	$O(n^{-\alpha})$	Yes	$m \gtrsim (\eta T)^3 \asymp n^3$	$O(n^2)$
SGD	$O(n^{-\frac{\alpha}{1+\alpha}})$	No	$m \gtrsim (\eta T)^3 \asymp n^{\frac{3}{\alpha+1}}$	$O(n)$
This work	$O(n^{-\alpha})$	Yes	$m \gtrsim (\eta T)^3 \asymp n^3$	$O(n)$

Table 1: Summary of results. Low noise means $L(\mathbf{W}^*) = \inf_{\mathbf{W}} L(\mathbf{W}) = 0$. Computation means the complexity of the gradient computation, which is nT for GD and T for SGD. The results in second and third rows for GD are derived by combining Assumption 3 with the risk bounds in [47]. In particular, if $\alpha = 1$, our results indicate both GD and SGD for 2-layer SNNs with subquadratic overparametrization $m \gtrsim n^{3/2}$ can lead to optimal risk rate $O(n^{-1/2})$ while the results in [46] need superquadratic overparametrization $m \gtrsim n^{5/2}$.

4.1 Gradient Descent

We first study the excess risk of the GD algorithm for SNNs. To this end, we first develop its generalization bounds. We denote $B \asymp B'$ if there exist some universal constants c_1 and $c_2 > 0$ such that $c_1 B \leq B' \leq c_2 B$. We denote $B \gtrsim B'$ if there exists a universal constant $c > 0$ such that $B \geq cB'$. Let e be the base of the natural logarithm. Let $\rho = C_x^2 \big(B_{\phi'}^2 + B_{\phi''} B_\phi + \frac{B_{\phi''} C_y}{\sqrt{m}} \big)$ and $b = C_x^2 B_{\phi''}(B_{\phi'} C_x + C_0)$.

Theorem 2 (Generalization Error). *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}$ be produced by Eq. (3.1). If $\eta \leq 1/(2\rho)$ and*

$$m \geq 32 C_0 \eta^2 T^2 C_x^4 B_{\phi''}^2 \Big(2n^{-1} \sqrt{\rho(\rho \eta T + 2)} B_{\phi'} C_x (1 + \eta\rho) \eta e T + 1 \Big)^2, \tag{4.1}$$

then the following inequality holds for any $t \in [T]$

$$\mathbb{E}[L(\mathbf{W}_t) - L_S(\mathbf{W}_t)] \leq \Big(\frac{4e^2 \eta^2 \rho^2 t}{n^2} + \frac{4e\eta\rho}{n} \Big) \sum_{j=0}^{t-1} \mathbb{E}[L_S(\mathbf{W}_j)].$$

Remark 1. Under an assumption $m \gtrsim (\eta T)^3$, a bound similar to Theorem 2 was recently established in [47]. We relax this assumption to $m \gtrsim (\eta T)^5/n^2 + \eta^2 T^2$ in Eq. (4.1). As we will show, a typical choice is $\eta T \asymp n^{\frac{1}{1+\alpha}}$. In this case, the assumption in Eq. (4.1) becomes $m \gtrsim (\eta T)^3 n^{-\frac{2\alpha}{1+\alpha}} + \eta^2 T^2$, which is milder than the assumption $m \gtrsim (\eta T)^3$ in [47]. This improvement is achieved by a better

estimation of the smallest eigenvalue of a Hessian matrix. Indeed, the smallest eigenvalue at $\alpha \mathbf{W}_t + (1-\alpha)\mathbf{W}_t^{(i)}$ is lower bounded by $-\frac{1}{\sqrt{m}}(\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 + 1)$ (up to a constant factor), where $\alpha \in (0,1)$ and $\{\mathbf{W}_t^{(i)}\}$ is the SGD sequence on $S^{(i)}$. The analysis [47] uses $\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 = O(\sqrt{\eta t})$ to control the smallest eigenvalue. As a comparison, we show $\|\mathbf{W}_t - \mathbf{W}_t^{(i)}\|_2 = O(n^{-1}(\eta t)^{\frac{3}{2}})$ (Lemma B.1).

A key step to relax the overparameterization is to build a bound on $\mathbb{E}[\|\mathbf{W}_t - \mathbf{W}_{\frac{1}{\eta T}}^*\|_2^2]$. The existing analysis shows that $\|\mathbf{W}_t - \mathbf{W}_0\|_2^2 = O(\eta t)$ [47], which grows to infinity as we run more and more iterations. In the following lemma to be proved in Section B.2, we improve it to $\mathbb{E}[\|\mathbf{W}_t - \mathbf{W}_{\frac{1}{\eta T}}^*\|_2^2] = O(\frac{\eta^2 T}{n} \sum_{j=0}^{T-1} \mathbb{E}[L_S(\mathbf{w}_j)] + \|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2)$. In particular, if $\eta T = O(\sqrt{n})$ and $\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2 = O(1)$, this bound becomes $\mathbb{E}[\|\mathbf{W}_t - \mathbf{W}_{\frac{1}{\eta T}}^*\|_2^2] = O(1)$. This explains why we relax the overparameterization assumption from $m \gtrsim (\eta T)^5$ in [47] to $m \gtrsim (\eta T)^3$. Furthermore, the bound involves $\sum_{j=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_j)]$ which would improve if the training errors are small, which is critical to get fast rates in a low noise case. Our basic idea to prove Lemma 3 is to first control $\mathbb{E}[\|\mathbf{W}_t - \mathbf{W}_{\frac{1}{\eta T}}^*\|_2^2]$ in terms of training errors. Our novelty is to replace these training errors with testing errors by using Theorem 2, which allows us to use Eq. (4.2) to remove some terms. The proof is given in Section B.3. For simplicity we assume $\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2 \geq 1$.

Lemma 3. *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}$ be produced by Eq. (3.1). If $\eta \leq 1/(2\rho)$, Eq. (4.1) holds,*

$$\mathbb{E}[L(\mathbf{W}_s)] \geq L(\mathbf{W}_{\frac{1}{\eta T}}^*), \quad \forall s \in \{0, 1, \ldots, T-1\} \tag{4.2}$$

and

$$m \geq 8C_0 b^2 (\eta T)^3, \tag{4.3}$$

then for any $t \in [T]$ we have

$$\mathbb{E}[\|\mathbf{W}_t - \mathbf{W}_{\frac{1}{\eta T}}^*\|_2^2] \leq R_T := \left(\frac{8e^2 \rho^2 \eta^3 T^2}{n^2} + \frac{8e\eta^2 T \rho}{n}\right) \sum_{j=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_j)] + 2\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2.$$

Remark 2. We impose the assumption $\mathbb{E}[L(\mathbf{W}_s)] \geq L(\mathbf{W}_{\frac{1}{\eta T}}^*), \forall s \in \{0, 1, \ldots, T-1\}$. If this assumption does not hold, then Assumption 3 implies further

$$\min_{s \in \{0,1,\ldots,T-1\}} \mathbb{E}[L(\mathbf{W}_s)] - \mathbb{E}[L(\mathbf{W}^*)] \leq L(\mathbf{W}_{\frac{1}{\eta T}}^*) - \mathbb{E}[L(\mathbf{W}^*)] = O((\eta T)^{-\alpha}).$$

This shows the violation of Eq. (4.2) already implies a model $\mathbf{W}_t, t \in [T]$ with a very small excess risk, and therefore the assumption Eq. (4.2) does not essentially affect our results.

Remark 3. It should be mentioned that if $\|\mathbf{W}^*\|_2 = O(1)$ we can derive similar results by replacing $\mathbf{W}_{\frac{1}{\eta T}}^*$ in the analysis with $\mathbf{W}^*$ (note $\mathbf{W}^*$ already satisfies the inequality $L(\mathbf{W}^*) - L(\mathbf{W}^*) + \frac{1}{\eta T}\|\mathbf{W}^* - \mathbf{W}_0\|_2^2 = O(1/(\eta T))$ and therefore can play the role of $\mathbf{W}_{\frac{1}{\eta T}}^*$). In this case, we no longer require the assumption (4.2). Indeed, Eq. (4.2) always holds with $\mathbf{W}_{\frac{1}{\eta T}}^*$ replaced by $\mathbf{W}^*$ due to the inequality $L(\mathbf{W}_s) \geq L(\mathbf{W}^*)$.

Both bounds in Theorem 2 and Lemma 3 depend on the term $\sum_{s=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_s)]$, for which we provide a bound in the following lemma. Combined with with Assumption 3, it implies (if $m \gtrsim \eta^3 T^3$)

$$\sum_{s=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_s)] = O(TL(\mathbf{W}_{\frac{1}{\eta T}}^*) + \frac{1}{\eta}\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2^2) = O(TL(\mathbf{W}^*) + T(T\eta)^{-\alpha}).$$

If $L(\mathbf{W}^*) = 0$, we have $\sum_{s=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_s)] = O(T(T\eta)^{-\alpha})$, which explains why we can get improved bounds in a low noise case. The proof is given in Section B.3.

Lemma 4. *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}$ be produced by Eq. (3.1) with $\eta \leq 1/(2\rho)$. If Eq. (4.1), (4.2), (4.3) hold and*

$$m \geq 4\left(\frac{8e^2 \rho^2 \eta^3 T^2}{n^2} + \frac{8e\eta^2 T \rho}{n}\right)^2 \left(bT\left(\|\mathbf{W}_{\frac{1}{\eta T}}^* - \mathbf{W}_0\|_2 + \sqrt{2\eta T C_0}\right)\right)^2, \tag{4.4}$$

then

$$\sum_{s=0}^{T-1}\mathbb{E}\big[L_S(\mathbf{W}_s)\big] \le 2TL(\mathbf{W}^*_{\frac{1}{\eta T}})+\frac{2\|\mathbf{W}^*_{\frac{1}{\eta T}}-\mathbf{W}_0\|_2^2}{\eta}+\frac{4bT\big(\|\mathbf{W}^*_{\frac{1}{\eta T}}-\mathbf{W}_0\|_2+\sqrt{2\eta TC_0}\big)\|\mathbf{W}^*_{\frac{1}{\eta T}}-\mathbf{W}_0\|_2^2}{\sqrt{m}}.$$

Now we present the optimization error bounds for GD. Recall R_T is defined in Lemma 3. The proof is given in Section B.3.

Theorem 5 (Optimization Error)**.** *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}$ be produced by Eq. (3.1) with $\eta \le 1/(2\rho)$. If Eq. (4.1), (4.2) and (4.3) hold, then*

$$\mathbb{E}[L_S(\mathbf{W}_T)] \le L(\mathbf{W}^*_{\frac{1}{\eta T}}) + \frac{1}{\eta T}\|\mathbf{W}^*_{\frac{1}{\eta T}} - \mathbf{W}_0\|_2^2 + \frac{bR_T}{\sqrt{m}}\big(\|\mathbf{W}^*_{\frac{1}{\eta T}} - \mathbf{W}_0\|_2 + \sqrt{2\eta TC_0}\big).$$

Remark 4. The following optimization error bounds were established in [47]

$$L_S(\mathbf{W}_T) \le \min_{\mathbf{W}} \Big\{ L_S(\mathbf{W}) + \frac{\|\mathbf{W} - \mathbf{W}_0\|_2^2}{\eta T} + \frac{b\|\mathbf{W} - \mathbf{W}_0\|_2^3}{\sqrt{m}} \Big\} + \frac{bC_0(\eta T)^{\frac{3}{2}}}{\sqrt{m}}. \qquad (4.5)$$

A key difference between the above bound and Theorem 5 is that Eq. (4.5) involves a term $\frac{(\eta T)^{\frac{3}{2}}}{\sqrt{m}}$, while Theorem 5 involves a term $O\Big(\frac{\sqrt{\eta T}R_T}{\sqrt{m}}\Big)$. If $R_T = o(\eta T)$, then the optimization error bounds in Theorem 5 would be tighter than Eq. (4.5). Indeed, the analysis in [47] requires $m \gtrsim (\eta T)^5$ to get the following optimization error bounds

$$L_S(\mathbf{W}_T) \le \min_{\mathbf{W}} \Big\{ L_S(\mathbf{W}) + \frac{\|\mathbf{W} - \mathbf{W}_0\|_2^2}{\eta T} + \frac{b\|\mathbf{W} - \mathbf{W}_0\|_2^3}{\sqrt{m}} \Big\} + O\Big(\frac{1}{\eta T}\Big).$$

As a comparison, if $R_T = O(1)$, Theorem 5 requires the assumption $m \gtrsim (\eta T)^3$ to derive

$$\mathbb{E}[L_S(\mathbf{W}_T)] \le L(\mathbf{W}^*_{\frac{1}{\eta T}})+\frac{\|\mathbf{W}^*_{\frac{1}{\eta T}} - \mathbf{W}_0\|_2^2}{\eta T}+O\Big(\frac{\sqrt{\eta T}}{\sqrt{m}}\Big) = L(\mathbf{W}^*_{\frac{1}{\eta T}})+\frac{\|\mathbf{W}^*_{\frac{1}{\eta T}} - \mathbf{W}_0\|_2^2}{\eta T}+O\Big(\frac{1}{\eta T}\Big).$$

We can combine the above discussions on generalization and optimization error bounds together to derive the following excess risk bounds. Note the right-hand side of Eq. (4.1), (4.3) and Eq. (4.4) are of the order of $(\eta T)^3$ if $\eta T = O(n)$. The proofs of Theorem 6 and Corollary 7 are given in Section B.4.

Theorem 6 (Excess Population Risk)**.** *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}$ be produced by Eq. (3.1) with $\eta \le 1/(2\rho)$. If $\eta T = O(n)$, Eq. (4.1), (4.2), (4.3) and Eq. (4.4) hold, then*

$$\mathbb{E}[L(\mathbf{W}_T)] - L(\mathbf{W}^*) = O\Big(\frac{\eta TL(\mathbf{W}^*)}{n} + \Lambda_{\frac{1}{\eta T}}\Big),$$

where Λ_λ is defined in Assumption 3.

The bound in Theorem 6 was also obtained in [47] under the assumption $m \gtrsim (\eta T)^5$. As a direct corollary, we show that GD can achieve excess risk bounds of the order $O(n^{-\frac{\alpha}{1+\alpha}})$ in the general case, and bounds of the order $O(n^{-\alpha})$ in the case $L(\mathbf{W}^*) = 0$ which is due to the incorporation of empirical risks in the generalization bounds. The basic idea is to balance the optimization and generalization via early-stopping [28, 38, 39, 56, 57]. Similar bounds can be derived by the analysis in [47] under Assumption 3.

Corollary 7. *Let Assumption 3 hold and assumptions in Theorem 6 hold.*

(a) *If we choose $\eta T \asymp n^{\frac{1}{\alpha+1}}$ and $m \asymp (\eta T)^3 \asymp n^{\frac{3}{\alpha+1}}$, then $\mathbb{E}[L(\mathbf{W}_T)] - L(\mathbf{W}^*) = O(n^{-\frac{\alpha}{1+\alpha}})$.*

(b) *If $L(\mathbf{W}^*) = 0$, choosing $\eta T \asymp n$ and $m \asymp (\eta T)^3 \asymp n^3$ implies that $\mathbb{E}[L(\mathbf{W}_T)] = O(n^{-\alpha})$.*

Remark 5. Other than the stability analysis [47], there are some discussions on the stability analysis for nonconvex functions that can be applied to SNNs [15, 26, 35, 60]. The discussions in [26] use step sizes $\eta_t = O(1/t)$ to get meaningful stability bounds, which, however, is not sufficient for a good convergence of optimization errors. The discussions in [15, 35, 60] impose a PL condition, and their error bounds depend on a condition number which can be large in practice.

4.2 Stochastic Gradient Descent

In this section, we study the excess risk bounds of SGD. As compared to GD, the analysis of SGD is more challenging since several properties of GD do not hold for SGD. For example, the analysis in [47] relies critically on the monotonicity of the sequence $\{L_S(\mathbf{W}_t)\}$, which does not hold for SGD. Furthermore, the introduced randomness of $\{i_t\}$ increases the variance of the iterates, which increases the difficulty of controlling the norm of iterates. We first provide a crude bound on the SGD iterates to be proved in Section C.2, which would be useful for our analysis. Without loss of generality we assume $4T\eta C_0 \geq 1$. Let $R'_T = \max\{2\sqrt{T\eta C_0}, \|\mathbf{W}^*_{\frac{1}{\eta T}} - \mathbf{W}_0\|_2\}$ and $b' = C_x^2 B_{\phi''}\left(C_x B_{\phi'} + \sqrt{2C_0}\right)$.

Lemma 8 (Iterate Bound). *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}_t$ be produced by SGD. If $\eta \leq 1/(2\rho)$ and*

$$m \geq 64 C_0 (b')^2 (T\eta)^3, \tag{4.6}$$

then

$$\|\mathbf{W}_t - \mathbf{W}_0\|_2 \leq 2\sqrt{T\eta C_0}, \quad \forall t \in [T].$$

We now develop stability and generalization bounds of SGD. In particular, we are interested in generalization bounds incorporating the training errors in the analysis [31, 34, 47]. This shows how good optimization would improve generalization, which is consistent with the analysis of SGD in a convex setting [34]. Theorem 9 gives on-average model stability bounds, which imply generalization bounds in Theorem 10. The proofs of Theorem 9 and Theorem 10 are given in Section C.3. Let $S^{(i)}$ be defined as in Definition 3.

Theorem 9 (Stability Bound). *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}_t$ and $\{\mathbf{W}_t^{(i)}\}_t$ be produced by SGD with $\eta \leq 1/(2\rho)$ on S and $S^{(i)}$, respectively. If Eq. (4.6) holds and*

$$m \geq 16\eta^2 T^2 (b' R'_T)^2 (1 + 2\eta\rho)^2, \tag{4.7}$$

then for any $t \leq T - 1$ we have

$$\frac{1}{n}\sum_{i=1}^{n} \mathbb{E}\left[\|\mathbf{W}_{t+1} - \mathbf{W}_{t+1}^{(i)}\|_2^2\right] \leq \frac{8e^2\rho(1+t/n)\eta^2}{n}\sum_{j=0}^{t} \mathbb{E}[L_S(\mathbf{W}_j)].$$

Theorem 10 (Generalization Error). *Let assumptions in Theorem 9 hold. Let $\{\mathbf{W}_t\}_t$ be produced by SGD with $\eta \leq 1/(2\rho)$. Then*

$$\mathbb{E}[L(\mathbf{W}_t) - L_S(\mathbf{W}_t)] \leq \frac{4e^2\rho^2(1+t/n)\eta^2}{n}\sum_{j=0}^{t}\mathbb{E}[L_S(\mathbf{W}_j)]$$

$$+ 4e\rho\eta\left(\frac{(1+t/n)\mathbb{E}[L_S(\mathbf{W}_t)]}{n}\sum_{j=0}^{t}\mathbb{E}[L_S(\mathbf{W}_j)]\right)^{\frac{1}{2}}.$$

The iterate bound in Lemma 8 is a bit crude. In the following lemma to be proved in Section C.4, we show this bound can be improved if we consider bounds in expectation. Let

$$\Delta_t := \max_{j=0,\dots,t} \mathbb{E}[\|\mathbf{W}_j - \mathbf{W}^*_{\frac{1}{\eta T}}\|_2^2], \quad \forall t \in \mathbb{N}.$$

If $t\eta^2 = O(1)$ and $t = O(n)$, Lemma 11 shows $\Delta_t = O(\|\mathbf{W}_0 - \mathbf{W}^*_{\frac{1}{\eta T}}\|_2^2)$ which is significantly better than the bound $O(\eta t)$ in Lemma 8. This allows us to get excess risk bounds under a relaxed overparameterization. Similar to the case with GD, this upper bound depends on the training errors of SGD iterates.

Lemma 11. *Let Assumptions 1, 2 hold. Let $\{\mathbf{W}_t\}_t$ be produced by SGD with $\eta \leq 1/(2\rho)$. If Eq. (4.6), Eq. (4.7) and Eq. (4.2) hold, then*

$$\Delta_{t+1} \leq 2\|\mathbf{W}_0 - \mathbf{W}^*_{\frac{1}{\eta T}}\|_2^2 + 4\rho\eta^2\left(1 + \frac{4e^2\eta\rho\sum_{j=0}^{t}(1+j/n)}{n} + \frac{4e(t+1)^{\frac{1}{2}}(1+t/n)^{\frac{1}{2}}}{\sqrt{n}}\right)\sum_{j=0}^{t}\mathbb{E}[L_S(\mathbf{W}_j)].$$

We now consider the optimization error bounds of SGD for SNNs. In the following theorem, we give a bound on the average of the optimization errors for the sequence of SGD iterates. Recall that R'_T is defined above Lemma 8.

Theorem 12 (Optimization Error). *Let assumptions in Lemma 11 hold. Then*

$$2\eta \sum_{t=0}^{T-1} \mathbb{E}\big[L_S(\mathbf{W}_t) - L_S(\mathbf{W}^*_{\frac{1}{\eta T}})\big] \leq \mathbb{E}[\|\mathbf{W}_0 - \mathbf{W}^*_{\frac{1}{\eta T}}\|_2^2] + 2\rho\eta^2 \sum_{t=0}^{T-1} \mathbb{E}[L_S(\mathbf{W}_t)] + \frac{2T\eta b'R'_T\Delta_T}{\sqrt{m}}.$$

Finally, we develop the excess risk bounds for SGD on SNNs. Note Eq. (4.8) can be satisfied by choosing $m \asymp (\eta T)^3$ since $R'_T = O(\sqrt{\eta T})$, which matches the overparameterization requirement of GD and improves the requirement $m \gtrsim (\eta T)^5$ in [47]. The proofs of Theorem 13 and Corollary 14 are given in Section C.5.

Theorem 13 (Excess Population Risk). *Let Assumptions 1 and 2 hold. Let $\{\mathbf{W}_t\}$ be produced by (3.2) and Eq (4.2) hold. If $\eta \leq 1/(2\rho)$,*

$$m \geq \max\Big\{16\eta^2 T^2 (b'R'_T)^2 (1+2\eta\rho)^2, 64C_0(b')^2(T\eta)^3,$$

$$4\big(8b'T\rho\eta^2 R'_T\big)^2\Big(1 + \frac{4e^2\eta\rho T(1+T/n)}{n} + \frac{4eT^{\frac{1}{2}}(1+T/n)^{\frac{1}{2}}}{\sqrt{n}}\Big)^2\Big\} \quad (4.8)$$

and $T = O(n)$ then we have

$$\frac{1}{T}\sum_{t=0}^{T-1} \mathbb{E}[L(\mathbf{W}_t) - L(\mathbf{W}^*)] = O\big(\Lambda_{\frac{1}{\eta T}} + \eta L(\mathbf{W}^*)\big).$$

Corollary 14. *Let Assumption 3 hold and assumptions in Theorem 13 hold. We choose an appropriate $m \asymp (\eta T)^3$.*

(a) We can choose $\eta \asymp T^{-\frac{\alpha}{1+\alpha}}$ and $T \asymp n$ to get $\frac{1}{T}\sum_{t=0}^{T-1}\mathbb{E}[L(\mathbf{W}_t)] - L(\mathbf{W}^*) = O(n^{-\frac{\alpha}{1+\alpha}})$.

(b) If $L(\mathbf{W}^*) = 0$, we can choose $T \asymp n$ and $\eta \asymp 1$ to get $\frac{1}{T}\sum_{t=0}^{T-1}\mathbb{E}[L(\mathbf{W}_t)] = O(n^{-\alpha})$.

Remark 6. By Corollary 14, SGD achieves excess risk bounds of the same order to that of GD in Corollary 7. An advantage of SGD over GD is that it requires less computation. To illustrate this, let us consider the general case for example. In this case, GD requires $T \asymp n^{\frac{1}{1+\alpha}}$ to achieve the error bound $O(n^{-\frac{\alpha}{1+\alpha}})$. Since GD requires $O(n)$ gradient computations per iteration and therefore the total gradient computation complexity is $O(n^{\frac{2+\alpha}{1+\alpha}})$. As a comparison, SGD requires $O(n)$ gradient computations and therefore saves the computation by a factor of $O(n^{\frac{1}{1+\alpha}})$. Note Corollary 7 considers the risk for the last iterate, while Corollary 14 considers the average of risks for all iterates. The underlying reason is that GD consistently decreases the training errors along the optimization process, while SGD does not enjoy this property. Note that the overparameterization requirement becomes $m \asymp n^{\frac{3}{\alpha+1}}$ and $m \asymp n^3$ in Part (a) and Part (b), respectively.

5 Conclusion

In this paper, we present stability and generalization analysis of both GD and SGD to train neural networks. Under a regularity assumption, we show both GD and SGD can achieve excess risk bounds of the order $O(n^{-\frac{\alpha}{\alpha+1}})$, which further improve to the order $O(n^{-\alpha})$ under a low noise condition. As compared to the existing stability analysis [47], we achieve our bounds under a relaxed overparameterization assumption and extend the existing analysis on GD to SGD. Our improvement is achieved by developing sharper bounds on norm of the GD/SGD iterate sequences.

There remain several interesting questions for further discussion. The first question is whether the overparamterization requirement $m \gtrsim (\eta T)^3$ can be further improved, and whether the overparameterization requirement can be independent of T. Second, our analysis applies to SNNs with a smooth activation function. It would be very interesting to extend our analysis to SNNs with the ReLU activation function. A key challenge in this direction is to control the smallest eigenvalue of the associated Hessian matrix [47]. Third, our bounds are stated in expectation. It would be useful to develop high-probability bounds to understand the robustness of the algorithm.

References

[1] Z. Allen-Zhu, Y. Li, and Y. Liang. Learning and generalization in overparameterized neural networks, going beyond two layers. *Advances in neural information processing systems*, 32, 2019.

[2] Z. Allen-Zhu, Y. Li, and Z. Song. A convergence theory for deep learning via over-parameterization. In *International Conference on Machine Learning*, pages 242–252. PMLR, 2019.

[3] I. Amir, T. Koren, and R. Livni. Sgd generalizes better than gd (and regularization doesn't help). In *Conference on Learning Theory*, pages 63–92. PMLR, 2021.

[4] S. Arora, R. Ge, B. Neyshabur, and Y. Zhang. Stronger generalization bounds for deep nets via a compression approach. In *International Conference on Machine Learning*, pages 254–263. PMLR, 2018.

[5] S. Arora, S. Du, W. Hu, Z. Li, and R. Wang. Fine-grained analysis of optimization and generalization for overparameterized two-layer neural networks. In *International Conference on Machine Learning*, pages 322–332. PMLR, 2019.

[6] Y. Bai and J. D. Lee. Beyond linearization: On quadratic and higher-order approximation of wide neural networks. *arXiv preprint arXiv:1910.01619*, 2019.

[7] F. Bao, G. Wu, C. Li, J. Zhu, and B. Zhang. Stability and generalization of bilevel programming in hyperparameter optimization. *Advances in Neural Information Processing Systems*, 34, 2021.

[8] P. L. Bartlett, D. J. Foster, and M. J. Telgarsky. Spectrally-normalized margin bounds for neural networks. In *Advances in Neural Information Processing Systems*, pages 6240–6249, 2017.

[9] P. L. Bartlett, A. Montanari, and A. Rakhlin. Deep learning: a statistical viewpoint. *Acta numerica*, 30:87–201, 2021.

[10] R. Bassily, V. Feldman, C. Guzmán, and K. Talwar. Stability of stochastic gradient descent on nonsmooth convex losses. *Advances in Neural Information Processing Systems*, 33, 2020.

[11] L. Bottou, F. E. Curtis, and J. Nocedal. Optimization methods for large-scale machine learning. *SIAM Review*, 60(2):223–311, 2018.

[12] O. Bousquet and A. Elisseeff. Stability and generalization. *Journal of Machine Learning Research*, 2(Mar):499–526, 2002.

[13] O. Bousquet, Y. Klochkov, and N. Zhivotovskiy. Sharper bounds for uniformly stable algorithms. In *Conference on Learning Theory*, pages 610–626, 2020.

[14] Y. Cao and Q. Gu. Generalization bounds of stochastic gradient descent for wide and deep neural networks. *Advances in neural information processing systems*, 32, 2019.

[15] Z. Charles and D. Papailiopoulos. Stability and generalization of learning algorithms that converge to global optima. In *International Conference on Machine Learning*, pages 744–753, 2018.

[16] J. Chen, D. Zhou, Y. Tang, Z. Yang, Y. Cao, and Q. Gu. Closing the generalization gap of adaptive gradient methods in training deep neural networks. In *29th International Joint Conference on Artificial Intelligence, IJCAI 2020*, pages 3267–3275. International Joint Conferences on Artificial Intelligence, 2020.

[17] Y. Chen, C. Jin, and B. Yu. Stability and convergence trade-off of iterative optimization algorithms. *arXiv preprint arXiv:1804.01619*, 2018.

[18] F. Cucker and D.-X. Zhou. *Learning Theory: an Approximation Theory Viewpoint*. Cambridge University Press, 2007.

[19] Z. Deng, H. He, and W. Su. Toward better generalization bounds with locally elastic stability. In *International Conference on Machine Learning*, pages 2590–2600. PMLR, 2021.

[20] S. S. Du, X. Zhai, B. Poczos, and A. Singh. Gradient descent provably optimizes over-parameterized neural networks. In *International Conference on Learning Representations*, 2018.

[21] G. K. Dziugaite and D. M. Roy. Computing nonvacuous generalization bounds for deep (stochastic) neural networks with many more parameters than training data. *arXiv preprint arXiv:1703.11008*, 2017.

[22] A. Elisseeff, T. Evgeniou, and M. Pontil. Stability of randomized learning algorithms. *Journal of Machine Learning Research*, 6(Jan):55–79, 2005.

[23] F. Farnia and A. Ozdaglar. Train simultaneously, generalize better: Stability of gradient-based minimax learners. In *International Conference on Machine Learning*, pages 3174–3185. PMLR, 2021.

[24] V. Feldman and J. Vondrak. High probability generalization bounds for uniformly stable algorithms with nearly optimal rate. In *Conference on Learning Theory*, pages 1270–1279, 2019.

[25] N. Golowich, A. Rakhlin, and O. Shamir. Size-independent sample complexity of neural networks. In *Conference On Learning Theory*, pages 297–299. PMLR, 2018.

[26] M. Hardt, B. Recht, and Y. Singer. Train faster, generalize better: Stability of stochastic gradient descent. In *International Conference on Machine Learning*, pages 1225–1234, 2016.

[27] A. Jacot, F. Gabriel, and C. Hongler. Neural tangent kernel: Convergence and generalization in neural networks. *Advances in neural information processing systems*, 31, 2018.

[28] Z. Ji, J. Li, and M. Telgarsky. Early-stopped neural networks are consistent. *Advances in Neural Information Processing Systems*, 34, 2021.

[29] Y. Klochkov and N. Zhivotovskiy. Stability and deviation optimal risk bounds with convergence rate $o(1/n)$. *Advances in Neural Information Processing Systems*, 34, 2021.

[30] T. Koren, R. Livni, Y. Mansour, and U. Sherman. Benign underfitting of stochastic gradient descent. *arXiv preprint arXiv:2202.13361*, 2022.

[31] I. Kuzborskij and C. Lampert. Data-dependent stability of stochastic gradient descent. In *International Conference on Machine Learning*, pages 2820–2829, 2018.

[32] Y. LeCun, Y. Bengio, and G. Hinton. Deep learning. *Nature*, 521(7553):436–444, 2015.

[33] J. Lee, L. Xiao, S. Schoenholz, Y. Bahri, R. Novak, J. Sohl-Dickstein, and J. Pennington. Wide neural networks of any depth evolve as linear models under gradient descent. *Advances in neural information processing systems*, 32, 2019.

[34] Y. Lei and Y. Ying. Fine-grained analysis of stability and generalization for stochastic gradient descent. In *International Conference on Machine Learning*, pages 5809–5819, 2020.

[35] Y. Lei and Y. Ying. Sharper generalization bounds for learning with gradient-dominated objective functions. In *International Conference on Learning Representations*, 2021.

[36] Y. Lei, Z. Yang, T. Yang, and Y. Ying. Stability and generalization of stochastic gradient methods for minimax problems. In *International Conference on Machine Learning*, pages 6175–6186, 2021.

[37] J. Li, X. Luo, and M. Qiao. On generalization error bounds of noisy gradient methods for non-convex learning. In *International Conference on Learning Representations*, 2020.

[38] M. Li, M. Soltanolkotabi, and S. Oymak. Gradient descent with early stopping is provably robust to label noise for overparameterized neural networks. In *International conference on artificial intelligence and statistics*, pages 4313–4324. PMLR, 2020.

[39] J. Lin, R. Camoriano, and L. Rosasco. Generalization properties and implicit regularization for multiple passes SGM. In *International Conference on Machine Learning*, pages 2340–2348, 2016.

[40] T. Liu, G. Lugosi, G. Neu, and D. Tao. Algorithmic stability and hypothesis complexity. In *International Conference on Machine Learning*, pages 2159–2167, 2017.

[41] B. London, B. Huang, and L. Getoor. Stability and generalization in structured prediction. *The Journal of Machine Learning Research*, 17(1):7808–7859, 2016.

[42] A. Maurer. Algorithmic stability and meta-learning. *Journal of Machine Learning Research*, 6 (Jun):967–994, 2005.

[43] S. Mukherjee, P. Niyogi, T. Poggio, and R. Rifkin. Learning theory: stability is sufficient for generalization and necessary and sufficient for consistency of empirical risk minimization. *Advances in Computational Mathematics*, 25(1):161–193, 2006.

[44] V. Nagarajan and J. Z. Kolter. Uniform convergence may be unable to explain generalization in deep learning. *Advances in Neural Information Processing Systems*, 32, 2019.

[45] B. Neyshabur, Z. Li, S. Bhojanapalli, Y. LeCun, and N. Srebro. Towards understanding the role of over-parametrization in generalization of neural networks. In *International Conference on Learning Representations (ICLR)*, 2019.

[46] F. Orabona. A modern introduction to online learning. *arXiv preprint arXiv:1912.13213*, 2019.

[47] D. Richards and I. Kuzborskij. Stability & generalisation of gradient descent for shallow neural networks without the neural tangent kernel. *Advances in Neural Information Processing Systems*, 34, 2021.

[48] D. Richards and M. Rabbat. Learning with gradient descent and weakly convex losses. In *International Conference on Artificial Intelligence and Statistics*, pages 1990–1998. PMLR, 2021.

[49] W. H. Rogers and T. J. Wagner. A finite sample distribution-free performance bound for local discrimination rules. *The Annals of Statistics*, pages 506–514, 1978.

[50] M. Seleznova and G. Kutyniok. Analyzing finite neural networks: Can we trust neural tangent kernel theory? *arXiv preprint arXiv:2012.04477*, 2020.

[51] S. Shalev-Shwartz, O. Shamir, N. Srebro, and K. Sridharan. Learnability, stability and uniform convergence. *Journal of Machine Learning Research*, 11(Oct):2635–2670, 2010.

[52] N. Srebro, K. Sridharan, and A. Tewari. Smoothness, low noise and fast rates. In *Advances in Neural Information Processing Systems*, pages 2199–2207, 2010.

[53] I. Steinwart and A. Christmann. *Support Vector Machines*. Springer Science & Business Media, 2008.

[54] T. Suzuki and S. Akiyama. Benefit of deep learning with non-convex noisy gradient descent: Provable excess risk bound and superiority to kernel methods. *arXiv preprint arXiv:2012.03224*, 2020.

[55] Y. Xing, Q. Song, and G. Cheng. On the algorithmic stability of adversarial training. *Advances in Neural Information Processing Systems*, 34, 2021.

[56] Y. Yao, L. Rosasco, and A. Caponnetto. On early stopping in gradient descent learning. *Constructive Approximation*, 26(2):289–315, 2007.

[57] Y. Ying and M. Pontil. Online gradient descent learning algorithms. *Foundations of Computational Mathematics*, 8(5):561–596, 2008.

[58] C. Zhang, S. Bengio, M. Hardt, B. Recht, and O. Vinyals. Understanding deep learning requires rethinking generalization. In *International Conference on Learning Representations*, 2017.

[59] J. Zhang, M. Hong, M. Wang, and S. Zhang. Generalization bounds for stochastic saddle point problems. In *International Conference on Artificial Intelligence and Statistics*, pages 568–576. PMLR, 2021.

[60] Y. Zhou, Y. Liang, and H. Zhang. Understanding generalization error of sgd in nonconvex optimization. *Machine Learning*, 111(1):345–375, 2022.

[61] D. Zou, Y. Cao, D. Zhou, and Q. Gu. Stochastic gradient descent optimizes over-parameterized deep relu networks. *arXiv preprint arXiv:1811.08888*, 2018.