Fixing the Bethe Approximation:
How Structural Modifications in a Graph Improve Belief Propagation

Abstract

Belief propagation is an iterative method for reasoning in probabilistic graphical models. Its well-known relationship to a classical concept from statistical physics, the Bethe free energy, puts it on a solid theoretical foundation. If belief propagation fails to approximate the marginals, then this is often due to a failure of the Bethe approximation. In this work, we show how modifications in a graphical model can be a great remedy for fixing the Bethe approximation. Specifically, we analyze how the removal of edges influences and improves belief propagation, and demonstrate that this positive effect is particularly distinct for dense graphs.

1 INTRODUCTION

Message passing algorithms are an effective method for approximate inference in probabilistic graphical models (Koller and Friedman, 2009). Although they often perform well in practice, there are only few guarantees about their theoretical behavior. A remarkable milestone was the discovery of a direct connection between message passing algorithms and concepts of statistical mechanics (Yedidia et al., 2001), perhaps the most famous being the relationship between belief propagation (BP) (Pearl, 1988) and the so-called Bethe free energy (Bethe, 1935; Peierls, 1936).

One favorable property of BP is its exactness on trees. On loopy graphs, however, it frequently suffers from two major issues: first, it may fail to converge to a fixed point and thus to find reasonable estimates of the marginals. Second, the fixed points themselves may induce bad estimates of the marginals, in which case even convergence would not help.

To solve the first issue, various techniques have been developed that can improve the convergence behavior of BP; e.g., one can damp the message updates (Murphy et al., 1999)

or utilize elaborate scheduling schemes (Elidan et al., 2006; Sutton and McCallum, 2007; Knoll et al., 2015; Aksenov et al., 2020). Moreover, it depends on the properties of the graphical model (Tatikonda and Jordan, 2002; Ihler et al., 2005; Mooij and Kappen, 2007) and on the initialization of the messages (Koehler, 2019; Knoll et al., 2021; Leisenberger et al., 2021) if and to which fixed point BP converges.

The second issue might be even harder to overcome, as it is inherently linked to a failure of the Bethe approximation (Weller et al., 2014). The detrimental influence of loops can make the Bethe free energy non-convex and cause its local minima – and thus BP fixed points – to be far away from the exact marginals. Enhanced variants of free energy approximations (Yedidia et al., 2005) or loop corrections (Mooij et al., 2007) are prudent alternatives, that improve the accuracy but also increase the complexity.

In this work, we follow a different path: we aim to improve the approximation quality of the Bethe free energy itself. To address this problem, we modify the structure of the graphical model and show how this transforms the Bethe free energy in a way that moves its local minima closer to the exact marginals. In particular, we analyze the effect of removing individual edges from the graph. This loop-breaking approach enforces a 'reconvexification' of the Bethe free energy and therefore not only improves the accuracy of fixed points, but also the convergence behavior of BP.

We make a series of interesting theoretical contributions that arise from analyzing the behavior of the Bethe free energy on a 'small scale'. More precisely, we introduce a measure for the discrepancy between two different representations of the Bethe free energy, each induced by a different graphical model, and then utilize this tool to relate variations in the Bethe free energy to the characteristics of the model and the behavior of BP. Theoretically and experimentally, we address the following questions: (i) How does edge removal influence the estimated marginals? (ii) How does edge removal influence the estimated partition function? (iii) Which and how many edges – if at all – should we remove?

Submitted to the 38th Conference on Uncertainty in Artificial Intelligence (UAI 2022). **To be used for reviewing only**.

The structure of this paper is as follows: Sec. 2 summarizes all relevant background on graphical models, BP, and the Bethe approximation. Sec. 3 contains a detailed theoretical analysis and the main results. We experimentally validate our findings in Sec. 4 and conclude the paper in Sec. 5.

2 BACKGROUND

This introductory section provides a compact overview of the topics that we deal with: probabilistic graphical models, belief propagation, and the Bethe approximation. We further introduce the Ising model and discuss related work.

2.1 PROBABILISTIC GRAPHICAL MODELS

We consider an undirected graph $\mathcal{G} = (\mathbf{X}, \mathbf{E})$, where $\mathbf{X} = \{1, \ldots, N\}$ is a set of nodes and $\mathbf{E} \subseteq \{(i, j) : i, j \in \mathbf{X}\}$ is a set of edges. An edge connects two nodes if $(i, j) \in \mathbf{E}$. Note that we assume all edges to be undirected and hence $(i, j) = (j, i)$ for all pairs of connected nodes; specifically we do not count edges twice. Furthermore, $\mathcal{N}(i)$ denotes the neighborhood of node i (i.e., the set of nodes that are connected to i) and $d_i := |\mathcal{N}(i)|$ denotes the degree of i.

Let $X_1, \ldots, X_N$ be random variables (RVs) with state spaces $\mathcal{X}_1, \ldots, \mathcal{X}_N$. A probabilistic graphical model (PGM) represents a joint probability distribution $P_{\mathbf{X}}(\boldsymbol{x})$ over the RVs, where each node represents a RV[1] and edges indicate statistical dependencies between RVs. Formally, a PGM is a pair $(\mathcal{G}, \boldsymbol{\Phi})$ that associates a set of potential functions (or potentials) $\boldsymbol{\Phi} = \{\Phi_1(\boldsymbol{x_1}), \ldots, \Phi_K(\boldsymbol{x_K})\}$ with the graph $\mathcal{G}$ that are defined over joint realizations of subsets $\boldsymbol{X_1}, \ldots, \boldsymbol{X_K} \subseteq \mathbf{X}$. We shall focus on the class of binary pairwise models[2] that satisfy the following two assumptions: first, each RV has two states, i.e., $\mathcal{X}_i = \mathcal{X} = \{+1, -1\}$. Second, the potentials are defined over either one (singleton potentials $\Phi_i(x_i)$) or two (pairwise potentials $\Phi_{ij}(x_i, x_j)$) RVs. Then the joint distribution factorizes as

$$P_{\mathbf{X}}(\boldsymbol{x}) = \frac{1}{\mathcal{Z}} \prod_{(i,j) \in \mathbf{E}} \Phi_{ij}(x_i, x_j) \prod_{i=1}^{N} \Phi_i(x_i), \qquad (1)$$

where $\mathcal{Z}$ is the normalization constant or partition function.

We consider the Ising model, whose potentials have the form $\Phi_{ij}(x_i, x_j) = \exp(J_{ij} x_i x_j)$ and $\Phi_i(x_i) = \exp(\theta_i x_i)$ with $J_{ij} \in \mathbb{R}$ being the coupling strength of edge (i, j) and $\theta_i \in \mathbb{R}$ being the local field of node i. We further call an edge (i, j) *attractive* if $J_{ij} > 0$, and *repulsive* if $J_{ij} < 0$.

Then (1) takes the exponential form

$$P_{\mathbf{X}}(\boldsymbol{x}) = \frac{1}{\mathcal{Z}} \exp(-E(\boldsymbol{x})) \qquad (2)$$

with $E(\boldsymbol{x}) := \sum_{(i,j) \in \mathbf{E}} J_{ij} x_i x_j + \sum_{i=1}^{N} \theta_i x_i$ being the state energy.[3]

Finally, following the terminology in Knoll et al. (2021), we specify three types of models: *unidirectional* models do only contain attractive edges and all variables are biased towards the same state (i.e., either $\theta_i \leq 0$ or $\theta_i \geq 0$); *attractive* models do only contain attractive edges, but there may be local fields that differ in sign; *general* models may contain both attractive and repulsive edges. Note that, by definition, attractive models include unidirectional models, while general models include both unidirectional and attractive models.

2.2 BELIEF PROPAGATION

In this work, we consider two problems: first, the computation of marginal distributions where our specific interest lies in singleton marginals $P_{X_i}(x_i)$; and second, the computation of the partition function.[4] It is well known that an exact computation of these quantities is intractable (Valiant, 1979; Cooper, 1990) and even the approximation of the marginals to a certain precision is NP-hard (Dagum and Luby, 1993).

Belief propagation (BP) approximates the marginals by iteratively exchanging local statistical information between nodes in form of 'messages'. This process is governed by the recursive message update equations

$$\mu_{ij}^{(n+1)}(x_j) \propto \sum_{x_i \in \mathcal{X}} \Phi_{ij}(x_i, x_j) \Phi_i(x_i) \prod_{k \in N(i) \setminus j} \mu_{ki}^{(n)}(x_k), \qquad (3)$$

where the superscript (n) refers to the current iteration and the subscript ij refers to the direction in which a message is sent (e.g., node i sends μ_{ij} to node j). In principle, one can approximate the singleton marginals at any iteration, by multiplying all incoming messages with the local potential:

$$\tilde{P}_i^{(n)}(x_i) \propto \frac{1}{Z_i} \Phi_i(x_i) \prod_{k \in \mathcal{N}(i)} \mu_{ki}^{(n)}(x_k) \qquad (4)$$

Generally, however, marginal estimates are considered to be more accurate when they are obtained from a BP fixed point (Murphy et al., 1999; Knoll et al., 2017); more precisely, BP has converged to a fixed point μ_{ij}°, whenever an update of the form (3) does not alter the message values anymore (that is $\mu_{ij}^{(n+1)} = \mu_{ij}^{(n)}$ for all (i, j)).

---

[1]Due to the one-to-one correspondence between variables X_i and nodes i, we shall not rigorously distinguish between them; e.g., we often write $P_i(x_i)$ instead of $P_{X_i}(x_i)$.

[2]A wide range of models can be equivalently transformed into binary pairwise models, although this may increase the state space considerably (Weiss, 2000; Eaton and Ghahramani, 2013).

[3]This parameterization does often facilitate the theoretical analysis as it associates a unique parameter vector – consisting of all J_{ij} and θ_i – with each distribution (a so-called minimal representation, Wainwright et al. (2008)).

[4]Actually, these two problems are closely related as marginals can be expressed as a ratio of sub-partition functions (Weller and Jebara, 2014b).

2.3 BETHE APPROXIMATION

It is often useful to formulate marginal inference in terms of a variational problem. For this purpose, we consider some trial distribution $Q_{\mathbf{X}}(\boldsymbol{x})$ and define the Gibbs free energy as

$$\mathcal{F}(Q_{\mathbf{X}}) = \mathcal{U}(Q_{\mathbf{X}}) - \mathcal{S}(Q_{\mathbf{X}}) \qquad (5)$$

with $\mathcal{U} = \mathbb{E}_Q\left[E(\mathbf{X})\right]$ being the average energy of the model and $\mathcal{S} = -\sum_{\boldsymbol{x} \in \mathcal{X}^N} Q_{\mathbf{X}}(\boldsymbol{x}) \log Q_{\mathbf{X}}(\boldsymbol{x})$ being the entropy of $Q_{\mathbf{X}}(\boldsymbol{x})$. Let us further define the marginal polytope $\mathbb{M}$ as the set of all valid probability distributions over $\mathbf{X}$ (i.e., that satisfy all global and local marginalization and normalization constraints). If one minimizes $\mathcal{F}$ over $\mathbb{M}$, then one recovers the true distribution $P_{\mathbf{X}}(\boldsymbol{x})$ with the negative log-partition function $-\log(\mathcal{Z})$ as the functional value at the global minimum, i.e., $-\log(\mathcal{Z}) = \min_{\mathbb{M}} \mathcal{F}(Q_{\mathbf{X}}) = \mathcal{F}(P_{\mathbf{X}})$.[5]

Two aspects, however, render the minimiziation of the Gibbs free energy intractable: first, the definition of the marginal polytope by exponentially many constraints; second, the evaluation of the entropy that requires summing over exponentially many terms. The Bethe free energy approximation addresses these two issues as follows: first, it relaxes the marginal polytope $\mathbb{M}$ to the local polytope $\mathbb{L}$ that involves only local marginalization and normalization constraints of the pairwise and singleton 'pseudo-marginals' $\tilde{P}_{ij}$ and $\tilde{P}_i$:

$$\mathbb{L} = \{\tilde{P}_{ij}, \tilde{P}_i : \sum_{x_j} \tilde{P}_{ij}(x_i, x_j) = \tilde{P}_i(x_i),$$
$$\sum_{x_i, x_j} \tilde{P}_{ij}(x_i, x_j) = 1, \sum_{x_i} \tilde{P}_i(x_i) = 1; \quad (6)$$
$$(i, j) \in \mathbf{E}, i \in \mathbf{X}\}.$$

Second, it replaces the entropy $\mathcal{S}$ by an accordingly weighted sum of entropy contributions from edges and nodes. More concretely, the Bethe free energy is defined as $\mathcal{F}_B = \mathcal{U}_B - \mathcal{S}_B$ where the Bethe average energy is

$$\mathcal{U}_B = -\sum_{(i,j) \in \mathbf{E}} \sum_{x_i, x_j \in \mathcal{X}} \tilde{P}_{ij}(x_i, x_j) \log \Phi_{ij}(x_i, x_j)$$
$$-\sum_{i=1}^{n} \sum_{x_i \in \mathcal{X}} \tilde{P}_i(x_i) \log \Phi_i(x_i), \qquad (7)$$

and the Bethe entropy is

$$\mathcal{S}_B = -\sum_{(i,j) \in \mathbf{E}} \sum_{x_i, x_j \in \mathcal{X}} \tilde{P}_{ij}(x_i, x_j) \log \tilde{P}_{ij}(x_i, x_j)$$
$$+\sum_{i=1}^{n} (d_i - 1) \sum_{x_i \in \mathcal{X}} \tilde{P}_i(x_i) \log \tilde{P}_i(x_i). \qquad (8)$$

---

[5] For more details on variational inference in graphical models and the marginal polytope, we refer the reader to Wainwright et al. (2008); Mezard and Montanari (2009).

While $\mathcal{U}_B$ equals the true average energy $\mathcal{U}$ in the exact marginals, $\mathcal{S}_B$ is only an approximation to the true entropy $\mathcal{S}$ – unless the graph is a tree (Yedidia et al., 2005). To obtain locally consistent approximations to the exact marginals, one usually aims to minimize $\mathcal{F}_B$ over $\mathbb{L}$. Also, the so-called Bethe partition function $\mathcal{Z}_{\mathcal{B}}$, that is implicitly defined by $-\log(\mathcal{Z}_{\mathcal{B}}) = \min_{\mathbb{L}} \mathcal{F}_B$, provides an estimation of $\mathcal{Z}$.

Binary variables allow for a particularly simple description of the local polytope. Following the notation of Welling and Teh (2001); Weller and Jebara (2013), we define the pseudo-marginal distribution of X_i by $\tilde{P}_i(X_i = +1) = q_i$ (implying $\tilde{P}_i(X_i = -1) = 1 - q_i$) and, for any pair of connected nodes, we denote the joint pseudo-marginal probability $\tilde{P}_{ij}(X_i = +1, X_j = +1)$ by ξ_{ij}. Then the local marginalization and normalization constraints induce a full specification of the joint probability table between X_i and X_j in terms of the three parameters q_i, q_j, ξ_{ij} (Tab. 1).

Table 1: Variational joint probability table for two binary variables X_i and X_j.

$\tilde{P}_{ij}(X_i, X_j)$	$X_j = +1$	$X_j = -1$	
$X_i = +1$	ξ_{ij}	$q_i - \xi_{ij}$	q_i
$X_i = -1$	$q_j - \xi_{ij}$	$1 + \xi_{ij} - q_i - q_j$	$1 - q_i$
	q_j	$1 - q_j$	

If we assume that all probabilities are strictly positive, then ξ_{ij} is bounded by

$$\max(0, q_i + q_j - 1) < \xi_{ij} < \min(q_i, q_j). \qquad (9)$$

Inserting singleton and pairwise pseudo-marginals from Table 1 together with the Ising potentials from Sec. 2.1 into (7) and (8), the Bethe free energy becomes

$$\mathcal{F}_B = -\sum_{(i,j) \in \mathbf{E}} \left(1 + 2\left(2\xi_{ij} - q_i - q_j\right)\right) J_{ij}$$
$$+\sum_{i=1}^{n} (1 - 2q_i)\theta_i - \sum_{(i,j) \in \mathbf{E}} \mathcal{S}_{ij} + \sum_{i=1}^{n} (d_i - 1)\mathcal{S}_i, \qquad (10)$$

where the pairwise entropies are

$$\mathcal{S}_{ij} = -\xi_{ij} \log \xi_{ij}$$
$$- (1 + \xi_{ij} - q_i - q_j) \log(1 + \xi_{ij} - q_i - q_j) \qquad (11)$$
$$- (q_i - \xi_{ij}) \log(q_i - \xi_{ij}) - (q_j - \xi_{ij}) \log(q_j - \xi_{ij})$$

are the local entropies are

$$\mathcal{S}_i = -q_i \log q_i - (1 - q_i) \log(1 - q_i). \qquad (12)$$

Then, with $(\boldsymbol{q}; \boldsymbol{\xi})$ being the vector that contains all q_i and ξ_{ij}, the local polytope now takes the much simpler form

$$\mathbb{L} = \{(\boldsymbol{q}; \boldsymbol{\xi}) \in \mathbb{R}^{|\mathbf{X}| + |\mathbf{E}|} : 0 < q_i < 1, i \in \mathbf{X};$$
$$\max(0, q_i + q_j - 1) < \xi_{ij} < \min(q_i, q_j), (i,j) \in \mathbf{E}\}. \qquad (13)$$

To facilitate the task of minimizing $\mathcal{F}_B$ over $\mathbb{L}$, Welling and Teh (2001) have derived necessary conditions for local minima of $\mathcal{F}_B$. By setting the partial derivative $\frac{\partial}{\partial \xi_{ij}} \mathcal{F}_B$ to zero, they proved that the resulting quadratic equation

$$\alpha_{ij} \xi_{ij}^2 - (1 + \alpha_{ij}(q_i + q_j))\xi_{ij} + (1 + \alpha_{ij})q_i q_j = 0, \quad (14)$$

$$\text{where} \quad \alpha_{ij} = e^{4J_{ij}} - 1, \quad (15)$$

has a unique valid (i.e., inside the bounds (9)) solution

$$\xi_{ij}^*(q_i, q_j) = \frac{1}{2\alpha_{ij}} \Big((1 + \alpha_{ij}(q_i + q_j)) \\ - \sqrt{(1 + \alpha_{ij}(q_i + q_j))^2 - 4\alpha_{ij}(1 + \alpha_{ij})q_i q_j} \Big). \quad (16)$$

This means that for each edge (i, j), the only $\xi_{ij}^*(q_i, q_j)$, that can be located at a stationary point of $\mathcal{F}_B$, depends directly on q_i and q_j and may therefore be inserted in the definition of $\mathcal{F}_B$ (10). This is advantageous for two reasons: first, it considerably reduces the number of independent variables that are involved in optimizing $\mathcal{F}_B$ (i.e., from $|\mathbf{X}| + |\mathbf{E}|$ to $|\mathbf{X}|$); second, it simplifies the shape of the domain, as $\mathcal{F}_B$ is now defined on a box-constrained domain, the *Bethe box*

$$\mathbb{B} = \{ \boldsymbol{q} \in \mathbb{R}^{|\mathbf{X}|} : 0 < q_i < 1, i \in \mathbf{X} \}. \quad (17)$$

In this work, we do always refer to the Bethe free energy by $\mathcal{F}_B$, if it is defined over the local polytope or the Bethe box.

2.4 RELATED WORK

Since the seminal work of Yedidia et al. (2001), it is well known that fixed points of BP correspond one-to-one to stationary points of the Bethe free energy; moreover, stable fixed points of BP must always be associated to local minima of the Bethe free energy (Heskes, 2003). Consequently, one can try to overcome the convergence issue of BP by minimizing the Bethe free directly. To solve the problem, Welling and Teh (2001); Shin (2012) have derived gradient-based algorithms; Yuille (2002); Heskes (2006) have proposed provably convergent double-loop algorithms.

Yedidia et al. (2005) have also shown that BP is only a special case of a general class of message passing algorithms, the generalized belief propagation (GBP). Likewise, fixed points of these algorithms correspond to stationary points of the so-called Kikuchi free energies that try to approximate the true entropy by a sum over entropy contributions from larger node clusters (Kikuchi, 1951; Pelizzola, 2005). In practice, many of these methods can be prohibitively slow and may suffer in the same way as BP from non-convexity of the particular free energy approximation; i.e., they may – if at all – converge to suboptimal minima. This inspired various researchers to design free energy approximations that are convex (Wainwright et al., 2005; Globerson and Jaakkola, 2007b), some of which are related to convergent message passing algorithms (Kolmogorov and Wainwright,

2006; Globerson and Jaakkola, 2007a; Hazan and Shashua, 2008; Meltzer et al., 2009; Jancsary and Matz, 2011).

The Bethe approximation proves often to be superior to other methods in terms of a tradeoff between efficiency and accuaracy (Meshi et al., 2009). Its theoretical properties have therefore been intensely studied: Heskes (2004); Pakzad and Anantharam (2005) derived conditions for the convexity of the Bethe free energy. Chertkov and Chernyak (2006) formulated the so-called loop series expansion that directly relates the Bethe partition function to the true partition function. Others have found interesting connections between the Bethe approximation and classical graph theory (Watanabe and Fukumizu, 2009; Vontobel, 2013). Moreover, Weller and Jebara (2014a) derived an FPTAS [6] to approximate the Bethe partition function in attractive models.

Another line of research, that is in some sense complementary to variational inference, tries to approximate the graphical model itself. The classical Chow-Liu algorithm (Chow and Liu, 1968) finds a spanning tree such that the Kullback-Leibler (KL) divergence between the original distribution and the tree distribution is minimal. Furthermore, two different techniques deserve special attention: first, the 'annihilation' of small probabilities that are below a certain treshold (Jensen and Andersen, 1990); second, the deletion of one or more edges from the model (not necessarily until a spanning tree is reached). Due to its empirical success, the second method deserves special attention: Kjaerulff (1994) carefully selected edges whose removal decreases the treewidth of a graph. van Engelen (1997) studied how the removal of edges in a directed graph influences the KL divergence. Choi and Darwiche (2006) showed that a particular class of GBP, the so-called join graph propagation (Dechter et al., 2002), can be equivalently cast in terms of a procedure that consecutively deletes and recovers edges. In the past, these methods were primarily applied to perform exact inference in the approximated model. For large graphs, this does often remain a hard computational challenge.

3 THEORETICAL ANALYSIS

We shall now devote our attention to the central topic of this work: how removing edges from a graphical model influences the properties of BP. While the accuracy of the exact marginals degrades if one approximates a model by a sparser one (van Engelen, 1997), one might expect a similar behavior for the marginals estimated by BP. We show, that the opposite is the case: sparsifying the graph does often significantly improve the marginal accuracy of BP. The quality of the estimated partition function, however, tends to degrade by deviating from the original model.

In this section, we explain these phenomena theoretically. We further analyze the role of the 'optimal' edge to be

---

[6] Fully polynomial-time approximation scheme.

removed and relate this problem to the Bethe free energy. In particular, we prove an inherent relationship between global error measures on the Bethe free energy and the coupling strength of the edges. Our detailed analysis of the Bethe free energy on a 'small-scale' extends the work of Welling and Teh (2001); Weller and Jebara (2013); Weller et al. (2014) and leads to better understanding of BP in general.

3.1 PROBLEM SPECIFICATION

We briefly clarify the problem to be considered. Let $(\mathcal{G}, \mathbf{\Phi})$ be a PGM and let $(\mathcal{G}', \mathbf{\Phi}')$ be a second PGM that is obtained by removing a set of edges $\tilde{\mathbf{E}}$ (and the associated pairwise potentials) from the original model.[7] Let $\mathcal{P} := \{P_i : i \in \mathbf{X}\}$ be the set of exact (singleton) marginals on $(\mathcal{G}, \mathbf{\Phi})$ and $\mathcal{Z}$ be the partition function. Assume that we run BP on both models and obtain pseudo-marginals $\tilde{\mathcal{P}} := \{\tilde{P}_i : i \in \mathbf{X}\}$ on $(\mathcal{G}, \mathbf{\Phi})$ resp. $\tilde{\mathcal{P}}' := \{\tilde{P}_i' : i \in \mathbf{X}\}$ on $(\mathcal{G}', \mathbf{\Phi}')$, together with partition function estimates $\tilde{\mathcal{Z}}$ resp. $\tilde{\mathcal{Z}}'$.[8] Then we are interested in comparing the following quantities: first, the l^1-errors $||\mathcal{P}_{\mathbf{X}} - \tilde{\mathcal{P}}_{\mathbf{X}}||_{l^1}$ and $||\mathcal{P}_{\mathbf{X}} - \tilde{\mathcal{P}}_{\mathbf{X}}'||_{l^1}$; and second, the absolute errors $|\log \mathcal{Z} - \log \tilde{\mathcal{Z}}|$ and $|\log \mathcal{Z} - \log \tilde{\mathcal{Z}}'|$.

Ideally, we would like to remove a set of edges, such that the induced errors $||P_{\mathbf{X}} - \tilde{P}_{\mathbf{X}}'||_{l^1}$ and $|\log \mathcal{Z} - \log \tilde{\mathcal{Z}}'|$ become minimal over all subsets $\tilde{\mathbf{E}} \subseteq \mathbf{E}$. That is, we want to find the model for which BP best approximates the marginals and the partition function of the original model. This is of course an intractable problem (in particular, as we do not have access to the exact quantities). Still, it remains a crucial question whether and to what extent the removal of edges has a positive impact on the estimates. To identify edges to be deleted, we need to define an objective that contains information about the discrepancy between different graphical models. Note that a global comparison via the KL divergence and its generalizations (Minka, 2005) is prohibitive as this would involve a summation over exponentially many terms. Likewise, it is intractable to compare between different representations of the Gibbs free energy (5). To relax the problem, we focus on the analysis of local discrepancies between two models. The Bethe free energy (10) provides an ideal tool to explicitly measure these local differences.

3.2 THE BETHE ENERGY DIFFERENCE

Our main idea to make model comparison tractable lies in comparing between two different representations of the Bethe free energy. We formalize this concept as follows: assume for now that we remove a single edge (i, j) from a model $(\mathcal{G}, \mathbf{\Phi})$ and let $(\mathcal{G}^{\setminus (i,j)}, \mathbf{\Phi}^{\setminus (i,j)})$ denote the resulting

---

[7]Without loss of generality, we assume that the removal of $\tilde{\mathbf{E}}$ does not make the graph disconnected (otherwise, individual connected components can be treated separately).

[8]Note that $\tilde{P}_{\mathbf{X}}'$ and $\tilde{Z}'$ are approximations to the exact marginals and partition function in the new model $(\mathcal{G}', \mathbf{\Phi}')$.

model. Let further $\mathcal{F}_B$ resp. $\mathcal{F}_B^{\setminus (i,j)}$ be the representations of the Bethe free energy that are associated with $(\mathcal{G}, \mathbf{\Phi})$ resp. $(\mathcal{G}^{\setminus (i,j)}, \mathbf{\Phi}^{\setminus (i,j)})$. Specifically, $\mathcal{F}_B^{\setminus (i,j)}$ does not contain the pairwise energy and entropy contributions from edge (i, j), while the local entropy contributions from nodes i and j are counted once less than in the definition of $\mathcal{F}_B$ (10). Then we define the *Bethe free energy difference* $\Delta \mathcal{F}_B^{(i,j)}$ as the difference between $\mathcal{F}_B$ and $\mathcal{F}_B^{\setminus (i,j)}$, i.e.,

$$\Delta \mathcal{F}_B^{(i,j)} := \mathcal{F}_B - \mathcal{F}_B^{\setminus (i,j)}$$
$$= \overbrace{-(1 + 2(2\xi_{ij} - q_i - q_j))J_{ij}}^{:= \Delta \mathcal{U}_B^{(i,j)}} + \overbrace{\mathcal{S}_i + \mathcal{S}_j - \mathcal{S}_{ij}}^{:= I_B^{(i,j)}}, \quad (18)$$

where $\Delta \mathcal{U}_B^{(i,j)}$ is the difference in the Bethe average energy and $I_B^{(i,j)}$ is the mutual information between X_i and X_j.

Depending on whether we consider $\mathcal{F}_B$ on the local polytope $\mathbb{L}$ (13) or the Bethe box $\mathbb{B}$ (17), $\Delta \mathcal{F}_B^{(i,j)}$ is defined on slices of these objects, that is either on the sliced local polytope

$$\mathbb{L}^{(i,j)} := \{(q_i, q_j; \xi_{ij}) \in \mathbb{R}^3 : 0 < q_i, q_j < 1; \\ \max(0, q_i + q_j - 1) < \xi_{ij} < \min(q_i, q_j)\} \quad (19)$$

or the sliced Bethe box

$$\mathbb{B}^{(i,j)} := \{(q_i, q_j) \in \mathbb{R}^2 : 0 < q_i, q_j < 1\}. \quad (20)$$

It only depends on three resp. two variables and may therefore be considered as a function that contains variational information about local changes in a model when removing an edge. Moreover, it entails an effective way of measuring the local discrepancy between two graphical models, e.g., by computing an arbitrary norm of $\Delta \mathcal{F}_B^{(i,j)}$ on $\mathbb{L}^{(i,j)}$ or $\mathbb{B}^{(i,j)}$. In this work, we consider L^p-norms as the most natural choice and analyze the special cases of $p = \infty$ and $p = 2$ in Sec. 3.3 (Theorem 1, Corollary 1, and Theorem 2).

In principle, one can generalize the above idea to compare between models that result from removing multiple edges $\tilde{\mathbf{E}}$ in one step, as the associated Bethe free energy difference $\Delta \mathcal{F}_B^{\tilde{\mathbf{E}}}$ is then simply the sum over energy differences $\Delta \mathcal{F}_B^{(i,j)}$ for all (i, j) in $\tilde{\mathbf{E}}$. However, this increases both the number of variables to be integrated over and the number of edge sets to be taken into account for removal. To facilitate the theoretical and experimental analysis of edge removal, we shall therefore focus on removing edges one by one.

We now derive a series of auxiliary theorems on the mathematical properties of ξ_{ij}^* from (16) and $\Delta \mathcal{F}_B^{(i,j)}$. These results – besides being interesting in themselves – will help us in proving our main results in Sec. 3.3. All proves for Sec. 3.2 and 3.3 are contained in the Appendix B.

Lemma 1. *Let (i, j) be an edge. In the center point $(0.5, 0.5)$ of the sliced Bethe box $\mathbb{B}^{(i,j)}$, the unique ξ_{ij}^* that can be located at a stationary point of $\mathcal{F}_B$ has the form*

$$\xi_{ij}^*(0.5, 0.5) = \frac{\sigma(2 J_{ij})}{2}. \quad (21)$$

We will also have to analyze the behavior of ξ_{ij}^* if q_i and q_j approach the boundary $\partial\mathbb{B}^{(i,j)9}$ of the sliced Bethe box:

Lemma 2. *Let (i,j) be edge and let $k \in [0,1]$. The limits of ξ_{ij}^* at the boundary $\partial\mathbb{B}^{(i,j)}$ of the sliced Bethe box are*

$$\lim_{\substack{q_i \to 0 \\ q_j \to k}} \xi_{ij}^*(q_i, q_j) = 0 = \lim_{\substack{q_i \to k \\ q_j \to 0}} \xi_{ij}^*(q_i, q_j), \qquad (22)$$

$$\lim_{\substack{q_i \to 1 \\ q_j \to k}} \xi_{ij}^*(q_i, q_j) = k = \lim_{\substack{q_i \to k \\ q_j \to 1}} \xi_{ij}^*(q_i, q_j). \qquad (23)$$

Moreover, we shall prepare bounds and compute the boundary limits at $\partial\mathbb{B}^{(i,j)}$ of the mutual information $I_B^{(i,j)}$ (18):

Lemma 3. *Let (i,j) be an edge.*

(a) *In the interior of the sliced Bethe box $\mathbb{B}^{(i,j)}$, the mutual information $I_B^{(i,j)}$ is bounded by*

$$0 < 8(\xi_{ij}^* - q_i q_j)^2 \leq I_B^{(i,j)}(q_i, q_j) \leq$$
$$\leq \frac{(\xi_{ij}^* - q_i q_j)^2}{q_i(1-q_i)q_j(1-q_j)}. \qquad (24)$$

(b) *The limit of $I_B^{(i,j)}$ at the boundary $\partial\mathbb{B}^{(i,j)}$ is*

$$\lim_{(q_i,q_j) \to \partial\mathbb{B}^{(i,j)}} I_B^{(i,j)}(q_i, q_j) = 0. \qquad (25)$$

Next, we compute first-order and second-order derivatives of $\Delta\mathcal{F}_B^{(i,j)}$ on $\mathbb{B}^{(i,j)}$. The proof utilizes results from Welling and Teh (2001); Weller and Jebara (2013) (Appendix C).

Lemma 4. *Let (i,j) be an edge.*

(a) *The first-order derivatives of $\Delta\mathcal{F}_B^{(i,j)}$ on $\mathbb{B}^{(i,j)}$ are*

$$\frac{\partial}{\partial q_i}\Delta\mathcal{F}_B^{(i,j)} = 2J_{ij} + \log\left(\frac{(1-q_i)(q_i - \xi_{ij}^*)}{q_i(1 + \xi_{ij}^* - q_i - q_j)}\right). \quad (26)$$

(b) *The second-order derivatives of $\Delta\mathcal{F}_B^{(i,j)}$ on $\mathbb{B}^{(i,j)}$ are*

$$\frac{\partial^2}{\partial q_i^2}\Delta\mathcal{F}_B^{(i,j)} = \frac{q_j(1-q_j)}{T_{ij}} - \frac{1}{q_i(1-q_i)}, \qquad (27)$$

$$\frac{\partial^2}{\partial q_i q_j}\Delta\mathcal{F}_B^{(i,j)} = \frac{\partial^2}{\partial q_j q_i}\Delta\mathcal{F}_B^{(i,j)} = \frac{q_i q_j - \xi_{ij}^*}{T_{ij}}, \qquad (28)$$

$$\frac{\partial^2}{\partial q_j^2}\Delta\mathcal{F}_B^{(i,j)} = \frac{q_i(1-q_i)}{T_{ij}} - \frac{1}{q_j(1-q_j)}, \qquad (29)$$

where $T_{ij} := q_i q_j (1-q_i)(1-q_j) - (\xi_{ij}^* - q_i q_j)^2$.

The following result formulates a useful property of the Bethe free energy difference on the sliced Bethe box:

---

⁹That is, the four line segments connecting the points $(0,0) - (0,1), (0,0) - (1,0), (0,1) - (1,1)$, and $(1,0) - (1,1)$.

Lemma 5. *Let (i,j) be an edge. $\Delta\mathcal{F}_B^{(i,j)}$ has precisely one stationary point on $\mathbb{B}^{(i,j)}$, which is $(\bar{q}_i, \bar{q}_j) = (0.5, 0.5)$ and is neither a maximum nor a minimum (i.e., a saddle point).*

Lemma 5 implies that $\Delta\mathcal{F}_B^{(i,j)}$ cannot possess a maximum nor a minimum in the interior of $\mathbb{B}^{(i,j)}$. This implies that the supremum and infimum of $\Delta\mathcal{F}_B^{(i,j)}$ must lie at the boundary. Finally, we characterize regions of $\mathbb{B}^{(i,j)}$ on which $\Delta\mathcal{F}_B^{(i,j)}$ contributes always negatively to the Bethe free energy $\mathcal{F}_B$:

Lemma 6. *Let (i,j) be an edge.*

(a) *For an attractive edge, $\Delta\mathcal{F}_B^{(i,j)}(q_i, q_j)$ is negative if either both $q_i, q_i < 0.5$ or both $q_i, q_i > 0.5$.*

(b) *For a repulsive edge, $\Delta\mathcal{F}_B^{(i,j)}(q_i, q_j)$ is negative if either $q_i < 0.5$ and $q_j > 0.5$ or $q_i < 0.5$ and $q_j > 0.5$.*

3.3 MAIN RESULTS

After having prepared the technical framework in Sec. 3.2, we now proceed by presenting our main results. First, we directly relate the Bethe free energy difference to the local properties of the graphical model (Theorem 1). Then, we address the problem of an 'Bethe-optimal' edge to be deleted (Corollary 1 and Theorem 2). Finally, we conclude about the approximation quality of BP regarding the estimated partition function if edges are deleted (Theorems 3 and 4).

Theorem 1. *Let (i,j) be an arbitrary edge. Then the L^∞-norm of the Bethe free energy difference is*

$$||\Delta\mathcal{F}_B^{(i,j)}||_{L^\infty} = |J_{ij}|, \qquad (30)$$

with

$$-|J_{ij}| = \inf_{(q_i,q_j) \in \mathbb{B}^{(i,j)}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j), \qquad (31)$$

$$|J_{ij}| = \sup_{(q_i,q_j) \in \mathbb{B}^{(i,j)}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j). \qquad (32)$$

Theorem 1 reveals a monotonic dependence between the strength of the couplings and absolute changes in the Bethe free energy that are caused by local modifications in the graphical structure. In terms of edge deletion, this implies that the 'Bethe-optimal' choice of an edge to be removed from the graph is the one with the weakest coupling strength:

Corollary 1. *Suppose we aim to remove an edge from the graphical model such that the induced maximum error in the Bethe free energy is minimal. Then this 'L^∞-Bethe-optimal' edge is the one with the lowest absolute coupling strength:*

$$\underset{(i,j) \in \mathbf{E}}{\operatorname{argmin}} ||\Delta\mathcal{F}_B^{(i,j)}||_{L^\infty} = \underset{(i,j) \in \mathbf{E}}{\operatorname{argmin}} |J_{ij}| \qquad (33)$$

An analogous property holds for the L^2-error of $\mathcal{F}_B$ on $\mathbb{L}$:

Theorem 2. *Suppose we aim to remove an edge from the graphical model such that the induced mean squared error in the Bethe free energy on the local polytope $\mathbb{L}$ is minimal. Then this 'L²-Bethe-optimal' edge is the one with the lowest absolute coupling strength:*

$$\operatorname*{argmin}_{(i,j)\in\mathbf{E}} ||\Delta\mathcal{F}_B^{(i,j)}||_{L^2} = \operatorname*{argmin}_{(i,j)\in\mathbf{E}} |J_{ij}| \qquad (34)$$

Next, we conclude about the quantitative change in the Bethe partition function $\mathcal{Z}_\mathcal{B}$ if an edge is removed:

Theorem 3. *Let $\mathcal{Z}_\mathcal{B}$ be the Bethe partition function associated with some graphical model, i.e., the quanitity that satisfies $-\log(\mathcal{Z}_\mathcal{B}) = \min_\mathbb{B} \mathcal{F}_B$. Suppose we remove an (attractive or repulsive) edge from the graph. Let $\mathcal{F}_B^{\backslash(i,j)}$ be the representation of the Bethe free energy associated with the new model, together with the new Bethe partition function $\mathcal{Z}_\mathcal{B}^{\backslash(i,j)}$ that is implicitly defined by $-\log(\mathcal{Z}_\mathcal{B}^{\backslash(i,j)}) = \min_\mathbb{B} \mathcal{F}_B^{\backslash(i,j)}$. Then the following error estimate holds:*

$$\left| \log\left(\frac{\mathcal{Z}_\mathcal{B}}{\mathcal{Z}_\mathcal{B}^{\backslash(i,j)}} \right) \right| < |J_{ij}| \qquad (35)$$

Finally, we conclude about the quality of the estimated partition function if edges are removed. We consider unidirectional models (Sec. 2.1) that allow for a precise statement:

Theorem 4. *Consider a unidirectional model, i.e., where all edges are attractive and all variables are biased towards the same state. Let $\mathcal{Z}$ resp. $\mathcal{Z}_\mathcal{B}$ be the associated partition resp. Bethe partition function. Suppose we remove an arbitrary edge from the graph and let $\mathcal{Z}_\mathcal{B}^{\backslash(i,j)}$ be the Bethe partition function associated with the new model. Then the quality of the estimated partition function degrades, i.e.,*

$$|\mathcal{Z} - \mathcal{Z}_\mathcal{B}| < |\mathcal{Z} - \mathcal{Z}_\mathcal{B}^{\backslash(i,j)}|. \qquad (36)$$

Theorem 4 does not formally extend to models that contain both positive and negative local fields. Generally, however, the error between the true and the BP-estimated partition function tends to increase, the more edges we remove.

This negative result is contrasted by the positive effect of edge removal on the estimated marginals. While it is difficult even for restricted model classes to provide any theoretical guarantees, we validate and explain our statement in Sec. 4.

4 EXPERIMENTS

We now demonstrate empirically how removing edges can have an astonishingly positive impact on the approximation accuracy of BP. We perform a range of experiments on a fully connected graph on 10 vertices.[10] Further experiments including a 5×5- grid graph are contained in Appendix C.

---

[10]This allows for a computation of the exact marginals via the junction tree algorithm (Lauritzen and Spiegelhalter, 1988) and enables us to compare the approximated marginals to them.

We consider both attractive and general models (Sec. 2.1). In Sec. 4.1, we focus on attractive models and sample J_{ij} uniformly from $[0, \hat{J}]$ for $\hat{J} \in \{0.1, 0.2, \ldots, 2\}$. In Sec. 4.2, we focus on general models and sample J_{ij} uniformly from $[-\hat{J}, \hat{J}]$ for $\hat{J} \in \{0.1, 0.2, \ldots, 2\}$. For both settings, we create two scenarios: first, models with weak local fields (each θ_i is sampled uniformly from $[-0.2, 0.2]$); second, models with strong local fields (each θ_i is sampled uniformly from $[-0.5, 0.5]$). For each configuration, we create 200 models.

For each individual model, we remove edges one by one until we reach a spanning tree. We do not remove edges, whose deletion makes the graph disconnected.[11] We compare two criteria for selecting the next edge to be removed: first, the Bethe-optimal criterion (Corollary 1, Theorem 2); second, we remove edges that induce the lowest mutual information between two connected variables in the original model. More precisely: assume that we have already removed edge set $\tilde{\mathbf{E}}$ from a model; then the next edge (i,j) to be removed is the one that minimizes either of the following criteria:

$$\begin{aligned} \texttt{BETHE-OPT}: \quad & \operatorname*{argmin}_{(i,j)\in\mathbf{E}\backslash\tilde{\mathbf{E}}} |J_{ij}| \\ \texttt{CHOW-LIU}: \quad & \operatorname*{argmin}_{(i,j)\in\mathbf{E}\backslash\tilde{\mathbf{E}}} I(X_i; X_j) \end{aligned}$$

Note that by applying the second criterion, we end up in a Chow-Liu tree (Chow and Liu, 1968), i.e., the spanning tree with the lowest KL divergence from the original model.[12]

For each intermediate model during the edge deletion process, we run BP 100 times with random message initialization to approximate the marginals. For each run, we perform at most 1000 iterations. If BP has not converged, we estimate the marginals from the final iteration. We utilize a randomized message scheduling to achieve better convergence (Elidan et al., 2006). For the error evaluation, we compute the l^1-distance between the exact and estimated marginals. The results for each model are averaged over the 100 runs. Finally, the results are averaged over all 200 models, each based on a different configuration of the potentials.

4.1 ATTRACTIVE MODELS

For weak couplings, BP finds accurate marginal estimates in the original model. If the strength of the couplings increases, this favorable property suddenly disappears at some critical treshold and BP fails to approximate the marginals for larger values of $\hat{J}$ (Fig. 1). This behavior is not due to worse convergence properties of BP, but results from inaccurate BP

---

[11]This procedure corresponds to the so-called reverse-delete algorithm (Kruskal, 1956) that constructs a maximum spanning tree with respect to a given criterion.

[12]We cannot generally apply the second criterion, as the computation of the mutual information between two variables requires knowledge of the related exact singleton and pairwise marginals.

fixed points and thus inaccurate minima of the Bethe free energy (Weller et al., 2014). While the Bethe free energy is convex for weaker couplings and possesses a unique global minimum, this minimum becomes an (unstable) maximum if $\hat{J}$ increases and cannot be reached by BP any longer (Heskes, 2003; Mooij and Kappen, 2005; Knoll and Pernkopf, 2017). For even larger couplings, the landscape of the Bethe free energy becomes increasingly complex and the (possibly many[13]) Bethe minima approach the boundary of the domain, thus moving away from the exact marginals.

If we remove edges from the graph, the marginal accuracy of BP in the new model is much better than in the original model. This can be explained by a 'reconvexification' of the Bethe free energy that makes unstable maxima stable minima again and allows BP to converge to accurate fixed points. The question on how many edges we should actually remove, is a difficult one. In Fig. 1, we observe that there appears to be a 'channel' that defines an optimal number of edges to be removed. The stronger the couplings become, the more preferable is it to rely on tree approximations. For stronger local potentials, the channel tends to become narrower and edge removal loses some of its benefit (although the results are still superior in comparison to the original model). Moreover, we observe that the BETHE-OPT criterion performs slightly better than the CHOW-LIU criterion, with an increasing advantage for stronger local potentials.

4.2 GENERAL MODELS

The situation for general models is similar as in the attractice case. We can observe certain differences though (Fig. 2): first, the critical treshold of the couplings, beyond which the Bethe free energy becomes non-convex, is higher than for attractive models. Second, for models with strong local potentials, edge removal based on the BETHE-OPT criterion improves the marginal accuracy only slightly. Interestingly, the Chow-Liu tree induces strikingly accurate Bethe minima for all models. As in the attractive case, we observe that the problem becomes more difficult if both the pairwise and the local potentials become stronger at the same time.

5 CONCLUSION

We have proposed to approximate a graphical model as a 'preprocessing step' for approximate inference. We focused on the removal of single edges and showed that this can have a beneficial impact on the behavior of belief propagation.

We have exploited the relationship between belief propagation and the Bethe free energy to explain the success of such an approach. Subsequently, we have validated our

findings in an experimental study. Most importantly, our analysis contributes to an improved understanding of belief propagation and the Bethe approximation in general.

We are convinced that our observations inspire the development of further sophisticated methods that try to approximate a graphical model and improve the behavior of message passing algorithms. We believe that one logical extension lies in the modification of the local potentials to compensate for the lost information caused by edge removal.

Figure 1: Attractive models. First row: $\theta_i \in [-0.2, 0.2]$; second row: $\theta_i \in [-0.5, 0.5]$. (a) + (c): BETHE-OPT criterion; (b) + (d): CHOW LIU criterion.

Figure 2: General models. First row: $\theta_i \in [-0.2, 0.2]$; second row: $\theta_i \in [-0.5, 0.5]$. (a) + (c): BETHE-OPT criterion; (b) + (d): CHOW LIU criterion.

---

[13]The Bethe free energy may theoretically possess exponentially many local minima (Watanabe and Fukumizu, 2009; Knoll and Pernkopf, 2019).

References

Vitaly Aksenov, Dan Alistarh, and Janne H. Korhonen. Relaxed scheduling for scalable belief propagation. In *Proceedings of NeurIPS*, 2020.

Hans A. Bethe. Statistical theory of superlattices. *Proceedings of the Royal Society A*, 150(871):552–575, 1935.

Michael Chertkov and Vladimir Y. Chernyak. Loop series for discrete statistical models on graphs. *Journal of Statistical Mechanics: Theory and Experiment*, 2006(6):P06009, 2006.

Arthur Choi and Adnan Darwiche. A variational approach for approximating bayesian networks by edge deletion. In *Proceedings of UAI*, 2006.

Chao-Kong Chow and Chao-Ning Liu. Approximating discrete probability distributions with dependence trees. *IEEE Transactions on Information Theory*, 14(3):462–467, 1968.

Gregory F. Cooper. The computational complexity of probabilistic inference using Bayesian belief networks. *Artificial intelligence*, 42(2-3):393–405, 1990.

Martin J. Dagum and Michael G. Luby. Approximate probabilistic reasoning in Bayesian belief networks is NP-hard. *Artificial Intelligence*, 60(1):141–153, 1993.

Rina Dechter, Kalev Kask, and Robert Mateescu. Iterative join-graph propagation. In *Proceedings of UAI*, 2002.

Sever S. Dragomir, Marcel L. Scholz, and Jadranka Sunde. Some upper bounds for the relative entropy and applications. *Computers & Mathematics with Applications*, 39(9–10):91–100, 2000.

Fredetik Eaton and Ghahramani. Model reductions for inference: generality of pairwise, binary, and planar factor graphs. *Neural Computation*, 25(5):1213–1260, 2013.

Gal Elidan, Ian McGraw, and Daphne Koller. Residual belief propagation: Informed scheduling for asynchronous message passing. In *Proceedings of UAI*, 2006.

Amir Globerson and Tommi Jaakkola. Convergent propagation algorithms via oriented trees. In *Proceedings of UAI*, 2007a.

Amir Globerson and Tommi Jaakkola. Approximate inference using conditional entropy decompositions. In *Proceedings of AISTATS*, 2007b.

Tamir Hazan and Amnon Shashua. Convergent message-passing algorithms for inference over general graphs with convex free energies. In *Proceedings of UAI*, 2008.

Tom Heskes. Stable fixed points of loopy belief propagation are minima of the Bethe free energy. In *Proceedings of NIPS*, 2003.

Tom Heskes. On the uniqueness of loopy belief propagation fixed points. *Neural Computation*, 16(11):2379–2413, 2004.

Tom Heskes. Convexity arguments for efficient minimization of the Bethe and Kikuchi free energies. *Journal of Artificial Intelligence Research*, 26(1):153–190, 2006.

Alexander T. Ihler, John W. Fisher, and Alan S. Willsky. Loopy belief propagation: convergence and effects of message errors. *Journal of Machine Learning Research*, 6(1):905–936, 2005.

Jeremy Jancsary and Gerald Matz. Convergent decomposition solvers for tree-reweighted free energies. In *Proceedings of AISTATS*, 2011.

Frank Jensen and Stig K. Andersen. Approximations in Bayesian belief universes for knowledge-based systems. In *Proceedings of UAI*, 1990.

Ryoichi Kikuchi. A theory of cooperative phenomena. *Physical Review*, 81(6):988–1003, 1951.

Uffe Kjaerulff. Reduction of computational complexity in bayesian belief networks through removal of weak dependencies. In *Proceedings of UAI*, 1994.

Christian Knoll and Franz Pernkopf. On loopy belief propagation – Local stability analysis for non-vanishing fields. In *Proceedings of UAI*, 2017.

Christian Knoll and Franz Pernkopf. Belief propagation: accurate marginals or accurate partition function – where is the difference? In *Proceedings of UAI*, 2019.

Christian Knoll, Michael Rath, Sebastian Tschiatschek, and Franz Pernkopf. Message Scheduling Methods for Belief Propagation. In *Machine Learning and Knowledge Discovery in Databases*, pages 295–310. Springer, 2015.

Christian Knoll, Dhagash Mehta, Tianran Chen, and Franz Pernkopf. Fixed points of belief propagation – an analysis via polynomial homotopy continuation. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 40(9):2124–2136, 2017.

Christian Knoll, Adrian Weller, and Franz Pernkopf. Self-guided belief propagation – a homotopy continuation method. In *arXiv:1812.01339*, 2021.

Frederic Koehler. Fast convergence of belief propagation to global optima: beyond correlation decay. In *Proceedings of NeurIPS*, 2019.

Daphne Koller and Nir Friedman. *Probabilistic Graphical Models: Principles and Techniques*. MIT press, 2009.

Vladimir Kolmogorov and Martin J. Wainwright. Convergent tree-reweighted message-passing for energy minimization. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 28 (10):1568–1583, 2006.

Joseph B. Kruskal. On the shortest spanning subtree of a graph and the traveling salesman problem. *Proceedings of the American Mathematical Society*, 7(1):48–50, 1956.

Steffen L. Lauritzen and David J. Spiegelhalter. Local computations with probabilities on graphical structures and their application to expert systems. *Royal Statistical Society*, 50(2):157–224, 1988.

Harald Leisenberger, Christian Knoll, Richard Seeber, and Franz Pernkopf. Convergence behavior of belief propagation: Estimating regions of attraction via Lyapunov functions. In *Proceedings of UAI*, 2021.

Talya Meltzer, Amir Globerson, and Yair Weiss. Convergent message passing algorithms – a unifying view. In *Proceedings of UAI*, 2009.

Ofer Meshi, Ariel Jaimovich, Amir Globerson, and Nir Friedman. Convexifying the Bethe free energy. In *Proceedings of UAI*, 2009.

Marc Mezard and Andrea Montanari. *Information, Physics, and Computation*. Oxford University Press, 2009.

Thomas P. Minka. Divergence measures and message passing. Technical report, Microsoft Research, 2005.

Joris Mooij, Bastian Wemmenhove, Bert Kappen, and Rizzo Tommaso. Loop corrected belief propagation. In *Proceedings of AISTATS*, 2007.

Joris M. Mooij and Hilbert J. Kappen. On the properties of the Bethe approximation and loopy belief propagation on binary networks. *Journal of Statistical Mechanics: Theory and Experiment*, 2005(11):P11012, 2005.

Joris M. Mooij and Hilbert J. Kappen. Sufficient conditions for convergence of the sum–product algorithm. *IEEE Transactions on Information Theory*, 53(12):4422–4437, 2007.

Kevin P. Murphy, Yair Weiss, and Michael I. Jordan. Loopy belief propagation for approximate inference: An empirical study. In *Proceedings of UAI*, 1999.

Payam Pakzad and Venkat Anantharam. Estimation and marginalization using the Kikuchi approximation methods. *Neural Computation*, 17(8):1836–1873, 2005.

Judea Pearl. *Probabilistic Reasoning in Intelligent Systems: Networks of Plausible Inference*. Morgan Kaufmann Publishers, 1988.

Rudolf E. Peierls. Statistical theory of superlattices with unequal concentrations of the components. *Proceedings of the Royal Society A*, 154(881):207–222, 1936.

Alessandro Pelizzola. Cluster variation method in statistical physics and probabilistic graphical models. *Journal of Physics A: Mathematical and General*, 38(33):R309–R339, 2005.

Nicholas Ruozzi. The Bethe partition function of log-supermodular graphical models. In *Proceedings of NIPS*, 2012.

Jinwoo Shin. Complexity of Bethe approximation. In *Proceedings of AISTATS*, 2012.

Charles Sutton and Andrew McCallum. Improved dynamic schedules for belief propagation. In *Proceedings of UAI*, 2007.

Sekhar C. Tatikonda and Michael I. Jordan. Loopy belief propagation and Gibbs measures. In *Proceedings of UAI*, 2002.

Leslie G. Valiant. Constructing free energy approximations and generalized belief propagation algorithms. *Theoretical Computer Science*, 8(2):189–201, 1979.

Robert van Engelen. Approximating Bayesian belief networks by arc removal. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 19(8):916–920, 1997.

Pascal O. Vontobel. Counting in graph covers: A combinatorial characterization of the Bethe entropy function. *IEEE Trans. on Information Theory*, 59(9):6018–6048, 2013.

Martin J. Wainwright, Tommi S. Jaakkola, and Alan S. Willsky. A new class of upper bounds on the log partition function. *IEEE Transactions on Information Theory*, 51(7):2313–2335, 2005.

Martin J. Wainwright, Michael I. Jordan, et al. Graphical models, exponential families, and variational inference. *Foundations and Trends in Machine Learning*, 1(1–2):1–305, 2008.

Yusuke Watanabe and Kenji Fukumizu. Graph zeta function in the Bethe free energy and loopy belief propagation. In *Proceedings of NIPS*, 2009.

Yair Weiss. Correctness of local probability propagation in graphical models with loops. *Neural Computation*, 12(1):1–41, 2000.

Adrian Weller and Tony Jebara. Bethe bounds and approximating the global optimum. In *Proceedings of AISTATS*, 2013.

Adrian Weller and Tony Jebara. Approximating the Bethe partition function. In *Proceedings of UAI*, 2014a.

Adrian Weller and Tony Jebara. Clamping variables and approximate inference. In *Proceedings of NIPS*, 2014b.

Adrian Weller, Kui Tang, Tony Jebara, and David Sontag. Understanding the Bethe approximation: When and how can it go wrong? In *Proceedings of UAI*, 2014.

Max Welling and Yee W. Teh. Belief optimization for binary networks: A stable alternative to loopy belief propagation. In *Proceedings of UAI*, 2001.

Jonathan S. Yedidia, William T. Freeman, and Yair Weiss. Bethe free energy, Kikuchi approximations, and belief propagation algorithms. In *Proceedings of NIPS*, 2001.

Jonathan S. Yedidia, William T. Freeman, and Yair Weiss. Constructing free energy approximations and generalized belief propagation algorithms. *IEEE Transactions on Information Theory*, 51(7):2282–2312, 2005.

Alan L. Yuille. CCCP algorithms to minimize the Bethe and Kikuchi free energies: Convergent alternatives to belief propagation. *Neural Computation*, 14(7):1691–1722, 2002.

APPENDIX

APPENDIX A – PROOFS

PROOF OF LEMMA 1:.

Inserting $q_i = 0.5$ and $q_j = 0.5$ in (16) yields

$$
\begin{aligned}
\xi_{ij}^*(0.5, 0.5) &= \frac{1}{2\alpha_{ij}}\Big((1+\alpha_{ij}) \\
&\quad - \sqrt{(1+\alpha_{ij})^2 - \alpha_{ij}(1+\alpha_{ij})}\,\Big) = \\
&= \frac{1}{2\alpha_{ij}}\Big((1+\alpha_{ij}) - \sqrt{(1+\alpha_{ij})}\,\Big) = \\
&\stackrel{(15)}{=} \frac{1}{2\big(e^{4J_{ij}}-1\big)}\big(e^{4J_{ij}} - \sqrt{e^{4J_{ij}}}\big) = \\
&= \frac{1}{2}\Big(\frac{e^{2J_{ij}}-1}{e^{2J_{ij}}-e^{-2J_{ij}}}\Big) = \\
&= \frac{1}{4}\big(\tanh(J_{ij})+1\big) = \\
&= \frac{\sigma(2\,J_{ij})}{2}.
\end{aligned}
$$

$\square$

PROOF OF LEMMA 2:.

Note first that one can interchange the role of q_i and q_j in the definition (16) of ξ_{ij}^* due to symmetry. It is therefore sufficient to consider only the left hand side equalities in (22) and (23). Let first $q_i \to 0$ and $q_j \to k$. From (9) we know that ξ_{ij}^* is bounded from above by $\min(q_i, q_j)$. As also $\xi_{ij}^* > 0$, the first equality in (22) follows by continuity. Let now $q_i \to 1$ and $q_j \to k$. Then the limit of both the lower bound and the upper bound in (9) equals k. Consequently, ξ_{ij}^* must tend to k as well, which yields the first equality in (23). $\square$

PROOF OF LEMMA 3:.

(a) We apply Lemma 7 of Dragomir et al. (2000) to the special case of binary random variables. For the upper bound, substitute the matching combinations of singleton and pairwise probabilities from table (1) into the right-hand formula from (59):

$$
\begin{aligned}
I_B^{(i,j)}(q_i, q_j) \leq{} & \frac{(\xi_{ij}^*)^2}{q_iq_j} + \frac{(q_i - \xi_{ij}^*)^2}{q_i(1-q_j)} + \\
& + \frac{(q_j - \xi_{ij}^*)^2}{(1-q_i)q_j} + \frac{(1+\xi_{ij}^* - q_i - q_j)^2}{(1-q_i)(1-q_j)} \\
& - 1,
\end{aligned}
$$

and after a few simple algebraic manipulations we directly arrive at the desired result. Analogously, we derive the lower bound by inserting the corresponding

probabilities into the expression to the left of $I_B^{(i,j)}$ in (59):

$$
\begin{aligned}
I_B^{(i,j)}(q_i, q_j) \geq{} & \frac{1}{2}\,\big(\,|\,\xi_{ij}^* - q_iq_j\,| + \\
& + |\,(1+\xi_{ij}^* - q_i - q_j) \\
& - (1-q_i)(1-q_j)\,| + \\
& + |\,(q_i - \xi_{ij}^*) - q_i(1-q_j)\,| + \\
& + |\,(q_j - \xi_{ij}^*) - (1-q_i)q_j\,|\,)^2
\end{aligned}
$$

Depending on whether we have an attractive or an repulsive edge, we either make us of result (a) or (b) in Lemma 8 and get rid of the absolute value symbols, after which – in both cases – the last expression simplifies to

$$
\frac{1}{2}\,(4(\xi_{ij}^* - q_iq_j))^2 = 8(\xi_{ij}^* - q_iq_j)^2.
$$

(b) Recall from (reference to sliced Bethe box) that the boundary of the sliced Bethe Box is the union of four line segments

$$
\begin{aligned}
\partial\mathbb{B}^{(i,j)} ={} & \{(q_i, q_j) \in \mathbb{R}^2 \,|\, q_i = 0, 0 \leq q_j \leq 1\}\ \cup \\
& \{(q_i, q_j) \in \mathbb{R}^2 \,|\, q_i = 1, 0 \leq q_j \leq 1\}\ \cup \\
& \{(q_i, q_j) \in \mathbb{R}^2 \,|\, 0 \leq q_i \leq 1, q_j = 0\}\ \cup \\
& \{(q_i, q_j) \in \mathbb{R}^2 \,|\, 0 \leq q_i \leq 1, q_j = 1\}.
\end{aligned}
$$

Without loss of generality, we focus on the case where $q_i \to 1$ and $q_j \to k$ for some $k \in [0,1]$. The remaining cases can be treated similarly. We further note that it is sufficient to prove the equality

$$
\lim_{\substack{q_i \to 1 \\ q_j \to k}} \tilde{P}_{ij}(x_i, x_j) = \lim_{\substack{q_i \to 1 \\ q_j \to k}} \tilde{P}_i(x_i)\tilde{P}_j(x_j) \qquad (37)
$$

for all possible realizations $x_i, x_j \in \{+1, -1\}$ of X_i, X_j, as this implies statistical independence of X_i and X_j at the boundary and consequently their mutual information must vanish in the limit. For checking equality (37), we utilize Lemma 2. E.g., for $x_i = +1, x_j = -1$ we get

$$
\begin{aligned}
\lim_{\substack{q_i \to 1 \\ q_j \to k}} \tilde{P}_{ij}(+1, -1) &\stackrel{(1)}{=} \lim_{\substack{q_i \to 1 \\ q_j \to k}} (q_i - \xi_{ij}^*(q_i, q_j)) \\
&\stackrel{(23)}{=} 1 - k = \\
&= \lim_{\substack{q_i \to 1 \\ q_j \to k}} q_i(1 - q_j) \\
&\stackrel{(1)}{=} \lim_{\substack{q_i \to 1 \\ q_j \to k}} \tilde{P}_i(+1)\tilde{P}_j(-1).
\end{aligned}
$$

Analogous calculations for the three other possible combinations of x_i, x_j validate the statement.

$\square$

PROOF OF LEMMA 4:.

(a) We utilize Lemma 9 of Welling and Teh (2001). Then the first-order derivative of $\mathcal{F}_B^{\backslash(i,j)}$ on $\mathbb{B}^{\backslash(i,j)}$ is

$$\frac{\partial}{\partial q_i}\mathcal{F}_B^{\backslash(i,j)} = -2\,\theta_i + 2\sum_{k\in\mathcal{N}(i)\backslash j} J_{ij}$$
$$+ \log\left(\frac{(1-q_i)^{d_i-2}}{q_i^{d_i-2}}\right. \tag{38}$$
$$\left.\prod_{k\in\mathcal{N}(i)\backslash j}\frac{q_i - \xi_{ij}^*}{1 + \xi_{ij}^* - q_i - q_j}\right).$$

Consequently, we obtain

$$\frac{\partial}{\partial q_i}\Delta\mathcal{F}_B^{(i,j)} = \frac{\partial}{\partial q_i}(\mathcal{F}_B - \mathcal{F}_B^{\backslash(i,j)})$$
$$= \frac{\partial}{\partial q_i}\mathcal{F}_B - \frac{\partial}{\partial q_i}\mathcal{F}_B^{\backslash(i,j)}$$
$$= 2J_{ij} + \log\left(\frac{(1-q_i)(q_i - \xi_{ij}^*)}{q_i(1 + \xi_{ij}^* - q_i - q_j)}\right). \tag{39}$$

(b) We can apply Theorem 5 from Weller and Jebara (2013). Note, however, that we must also take the second derivatives of the node entropies $\mathcal{S}_i$ and $\mathcal{S}_j$ into account that are computed as

$$\frac{\partial^2}{\partial q_i^2}\mathcal{S}_i = -\frac{1}{q_i(1-q_i)}, \tag{40}$$
$$\frac{\partial^2}{\partial q_j^2}\mathcal{S}_j = -\frac{1}{q_j(1-q_j)}, \tag{41}$$

and vanish for the cross derivatives. By definition, we have $\Delta\mathcal{F}_B^{(i,j)} = \Delta\mathcal{U}_B^{(i,j)} + I_B^{(i,j)} = \Delta\mathcal{U}_B^{(i,j)} + \mathcal{S}_i + \mathcal{S}_j - \mathcal{S}_{ij}$. Note that f_{ij} from Theorem 5 corresponds precisely to $\Delta\mathcal{U}_B^{(i,j)} - \mathcal{S}_{ij}$. If we put these observations together, the statement follows immediately.

$\square$

PROOF OF LEMMA 5:.

The proof consists of two parts: in the first part, we show that $(0.5, 0.5)$ is the only stationary point of $\Delta\mathcal{F}_B^{(i,j)}$ on the sliced Bethe box. In the second part, we show that the Hessian matrix $\nabla^2\mathcal{F}_B$ evaluated in $(0.5, 0.5)$ is indefinite.

Part 1: Setting the gradient $\nabla\,\Delta\mathcal{F}_B^{(i,j)}$ with its components given by (39) to zero leads to the following nonlinear equa-

tion system:

$$2J_{ij} + \log\left(\frac{q_i - \xi_{ij}^* - q_i^2 + q_i\xi_{ij}^*}{q_i + q_i\xi_{ij}^* - q_i^2 - q_iq_j}\right) = 0$$
$$2J_{ij} + \log\left(\frac{q_j - \xi_{ij}^* - q_j^2 + q_j\xi_{ij}^*}{q_j + q_j\xi_{ij}^* - q_j^2 - q_iq_j}\right) = 0,$$

which is equivalent to

$$e^{2J_{ij}}(q_i - \xi_{ij}^* - q_i^2 + q_i\xi_{ij}^*) = q_i + q_i\xi_{ij}^* - q_i^2 - q_iq_j \tag{42}$$
$$e^{2J_{ij}}(q_j - \xi_{ij}^* - q_j^2 + q_j\xi_{ij}^*) = q_j + q_j\xi_{ij}^* - q_j^2 - q_iq_j. \tag{43}$$

By subtracting (43) from (42), we can reduce the above system to one equation

$$(e^{2J_{ij}} - 1)(q_j + q_i)(q_j - q_i)$$
$$+ (-e^{2J_{ij}}\xi_{ij}^* - e^{2J_{ij}} + \xi_{ij}^* + 1)(q_j - q_i) = 0. \tag{44}$$

Note that (44) might possess additional solutions, that do not solve the original system (42) + (43); each solution of (44), however, does also solve (42) + (43). Obviously all feasible pairs (q_i, q_j) such that $q_i = q_j$ solve equation (44) and thus also (42) + (43). Now assume that there exists a feasible solution to (44) with unequal components. If we divide (44) by $(q_j - q_i)$ and simplify, we end up with the contradictory result

$$\underbrace{(1 + \xi_{ij}^* - q_i - q_j)}_{=\tilde{P}_{ij}(X_i=-1, X_j=-1)>0}\underbrace{(1 - e^{2J_{ij}})}_{\neq 0} = 0. \tag{45}$$

Consequently, the only candidates for stationary points of $\Delta\mathcal{F}_B^{(i,j)}$ on $\mathbb{B}^{(i,j)}$ are those with equal components. This allows us to substitute q_i for q_j in either of the two equations in "Referenzen" and directly solve for q_i. In other words, we must solve

$$e^{2J_{ij}}(q_i - \xi_{ij}^* - q_i^2 + q_i\xi_{ij}^*) - (q_i + q_i\xi_{ij}^* - 2q_i^2) = 0,$$

or equivalently,

$$\xi_{ij}^* = \frac{(1 - e^{2J_{ij}})q_i + (e^{2J_{ij}} - 2)q_i^2}{-e^{2J_{ij}} - (1 - e^{2J_{ij}})q_i},$$

where we can replace ξ_{ij}^* by formula (16) (with $q_j = q_i$):

$$(1 + 2\alpha_{ij}q_i) - \sqrt{1 + 4\alpha_{ij}q_i(1-q_i)}$$
$$= 2\alpha_{ij}\underbrace{\frac{(1 - e^{2J_{ij}})q_i + (e^{2J_{ij}} - 2)q_i^2}{-e^{2J_{ij}} - (1 - e^{2J_{ij}})q_i}}_{:=A}$$

After isolating the radical and squaring both sides of the equation we get

$$(1 + \alpha_{ij})q_i^2 - (1 + 2\alpha_{ij}q_i)A + \alpha_{ij}A^2 = 0,$$

which can be further simplified to

$$\frac{e^{2J_{ij}}(e^{2J_{ij}} - 1)(2q_i - 1)(q_i - 1)^2}{(q_i + e^{2J_{ij}} - q_i e^{2J_{ij}})^2} = 0.$$

Finally, we multiply both sides by $\underbrace{\frac{(q_i + e^{2J_{ij}} - q_i e^{2J_{ij}})^2}{e^{2J_{ij}} \underbrace{(e^{2J_{ij}} - 1)}_{\neq 0} \underbrace{(q_i - 1)^2}_{\neq 0}}}_{\neq 0}$ and end up with

$$2q_i - 1 = 0 \quad \Rightarrow \quad q_i = 0.5.$$

We conclude that $(\bar{q}_i, \bar{q}_j) = (0.5, 0.5)$ is the only stationary point of $\Delta\mathcal{F}_B^{(i,j)}$.

Part 2: To compute the Hessian matrix $\nabla^2 \Delta\mathcal{F}_B^{(i,j)}$, we use the second partial derivatives in (27) - (29). Lemma 1 provides us with a simple expression of ξ_{ij}^* evaluated in $(0.5, 0.5)$. Furthermore, we make use of the relations

$$\sigma(2x)(1 - \sigma(2x)) = \frac{1}{2}\sigma'(2x) = \frac{1}{4\cosh^2(x)} \quad (46)$$

and

$$(1 - 2\sigma(2x)) = -\tanh(x). \quad (47)$$

We calculate

$$q_i q_j (1 - q_i)(1 - q_j) - (\xi_{ij}^* - q_i q_j)^2$$
$$= \frac{1}{16} - \left(\frac{1}{2}\sigma(2J_{ij}) - \frac{1}{4}\right)^2$$
$$= \frac{1}{4}\sigma(2J_{ij})(1 - \sigma(2J_{ij})) \overset{(46)}{=} \frac{1}{16\cosh^2(J_{ij})}. \quad (48)$$

Then the Hessian evaluated in $(0.5, 0.5)$ is

$$\nabla^2 \Delta\mathcal{F}_B^{(i,j)}(0.5, 0.5)$$
$$= \begin{pmatrix} \frac{1}{\sigma(2J_{ij})\left(1-\sigma(2J_{ij})\right)} - 4 & \frac{1-2\sigma(2J_{ij})}{\sigma(2J_{ij})\left(1-\sigma(2J_{ij})\right)} \\ \frac{1-2\sigma(2J_{ij})}{\sigma(2J_{ij})\left(1-\sigma(2J_{ij})\right)} & \frac{1}{\sigma(2J_{ij})\left(1-\sigma(2J_{ij})\right)} - 4 \end{pmatrix}$$
$$\overset{(47),(48)}{=} 4\begin{pmatrix} \cosh^2(J_{ij}) - 1 & -\cosh^2(J_{ij})\tanh(J_{ij}) \\ -\cosh^2(J_{ij})\tanh(J_{ij}) & \cosh^2(J_{ij}) - 1 \end{pmatrix}$$
$$= 4\sinh(J_{ij})\begin{pmatrix} \sinh(J_{ij}) & -\cosh(J_{ij}) \\ -\cosh(J_{ij}) & \sinh(J_{ij}) \end{pmatrix}.$$

Finally, we compute the leading principal minors of the Hessian, i.e., the determinants of all upper left square submatrices. The first leading principal minor is

$$|4\sinh^2(J_{ij})| > 0. \quad (49)$$

The second leading principal minor – and thus the determinant of the entire Hessian matrix – is

$$16\sinh^2(J_{ij})\underbrace{\left(\sinh^2(J_{ij}) - \cosh^2(J_{ij})\right)}_{= -1} \quad (50)$$
$$= -16\sinh^2(J_{ij}) < 0.$$

Since the leading principal minors alternate in sign, starting with a positive number, it follows that the Hessian evaluated in $(0.5, 0.5)$ is indefinite. $\qquad\square$

PROOF OF LEMMA 6:.

We only prove the statement for attractive edges as the statement for repulsive edges can be proven analogously. Moreover, we only consider the scenario that both q_i, q_j are > 0.5 (due to symmetry, the reverse scenario can be treated analogously). Let $\mathbb{B}_{>0.5}^{(i,j)} \subseteq \mathbb{B}^{(i,j)}$ be the orthant of $\mathbb{B}^{(i,j)}$ that consists of all points (q_i, q_j) with $0.5 < q_i, q_j < 1$. We know from Lemma 5 that $\Delta\mathcal{F}_B^{(i,j)}$ has no stationary point in $\mathbb{B}_{>0.5}^{(i,j)}$ and is thus bounded from above and below by the limit values of $\Delta\mathcal{F}_B^{(i,j)}$ at the boundary of $\mathbb{B}_{>0.5}^{(i,j)}$. If we can prove that all limit values of $\Delta\mathcal{F}_B^{(i,j)}$ at the boundary of $\mathbb{B}_{>0.5}^{(i,j)}$ are at most zero, then the statement follows immediately (as $\Delta\mathcal{F}_B^{(i,j)}$ is continuous in the interior of $\mathbb{B}^{(i,j)}$).

The boundary of $\mathbb{B}_{>0.5}^{(i,j)}$ consists of four line segments. Let us first consider the line segments that connect $(0.5, 1)$ to $(1, 1)$ and $(1, 0.5)$ to $(1, 1)$. In the proof of Theorem 1 it is shown[14] that

$$\lim_{\substack{q_i \to 0.5 \\ q_j \to 1}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) = 0,$$

$$\lim_{\substack{q_i \to 1 \\ q_j \to 0.5}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) = 0,$$

$$\lim_{\substack{q_i \to 1 \\ q_j \to 1}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) = -J_{ij},$$

and that $\Delta\mathcal{F}_B^{(i,j)}(q_i, q_j)$ is monotonically decreasing from $(0.5, 1)$ to $(1, 1)$ and from $(1, 0.5)$ to $(1, 1)$. Consequently the statement for these two line segments is correct.

Now consider the two remaining line segments, i.e., that connect $(0.5, 0.5)$ to $(0.5, 1)$ and $(0.5, 0.5)$ to $(1, 0.5)$. Due to symmetry, we can again focus on the first case. Let us explicitly compute the value of $\Delta\mathcal{F}_B^{(i,j)}(q_i, q_j)$ in $(0.5, 0.5)$. For that purpose, we must first evaluate the pairwise entropy $\mathcal{S}_{ij}$ in $(0.5, 0.5)$, where we utilize Lemma 1:

$$\mathcal{S}_{ij}(0.5, 0.5) = -\frac{\sigma(2J_{ij})}{2}\log\left(\frac{\sigma(2J_{ij})}{2}\right)$$
$$+ \left(\frac{\sigma(2J_{ij})}{2}\right)\log\left(\frac{\sigma(2J_{ij})}{2}\right)$$
$$- 2\left(0.5 - \frac{\sigma(2J_{ij})}{2}\right)\log\left(0.5 - \frac{\sigma(2J_{ij})}{2}\right)$$
$$= -2\left(0.5 - \frac{\sigma(2J_{ij})}{2}\right)\log\left(0.5 - \frac{\sigma(2J_{ij})}{2}\right)$$

---

[14]Note that these results follow independently from the current statement and can therefore be utilized, although Theorem 1 appears chronologically later in this work than Lemma 6.

Then we obtain

$$
\begin{aligned}
\Delta\mathcal{F}_B^{(i,j)}(0.5,0.5) =& (-1 - 2(2\xi_{ij}^*(0.5,0.5) - 0.5 - 0.5))J_{ij} \\
& + \overbrace{\mathcal{S}_i(0.5)}^{=\log(2)} + \overbrace{\mathcal{S}_j(0.5)}^{=\log(2)} - \mathcal{S}_{ij}(0.5,0.5) \\
=& -J_{ij} + 2J_{ij}(1 - \sigma(2J_{ij})) \\
& + \log(2) + \sigma(2J_{ij})\log(\frac{\sigma(2J_{ij})}{1 - \sigma(2J_{ij})}) \\
& + \log(1 - \sigma(2J_{ij})) \\
=& J_{ij} + \log(2) + \log(1 - \sigma(2J_{ij})) \\
=& \log(2e^{J_{ij}}(1 - \sigma(2J_{ij}))) \\
=& \log(\operatorname{sech}(J_{ij})) < 0.
\end{aligned}
$$

If we can finally show that $\Delta\mathcal{F}_B^{(i,j)}$ is monotonically increasing from $(0.5, 0.5)$ to $(0.5, 1)$, then the statement of the Lemma follows. By an analogous calculation as in the proof of Lemma 5 we conclude that the (one-dimensional) function $\Delta\mathcal{F}_B^{(i,j)}(0.5, q_j)$ has no stationary point for $q_j \in (0.5, 1)$ and must therefore be monotonically increasing (as $\Delta\mathcal{F}_B^{(i,j)}(0.5, 0.5) < 0$ and $\lim_{\substack{q_i \to 0.5 \\ q_j \to 1}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) = 0$). $\square$

PROOF OF THEOREM 1:.

By Lemma 5, the Bethe energy difference function does not have any local optima and hence both its greatest lower bound and least upper bound must be located at the boundary of the sliced Bethe box. As in the proof of Lemma 3 (b), we denote this boundary by $\partial\mathbb{B}^{(i,j)}$. Let further $\overline{\mathbb{B}^{(i,j)}} = \mathbb{B}^{(i,j)} \,\dot{\cup}\, \partial\mathbb{B}^{(i,j)}$ be the closure of $\mathbb{B}^{(i,j)}$ and

$$
\overline{\Delta\mathcal{F}_B^{(i,j)}}(q_i, q_j) := \lim_{(q_i,q_j) \to \overline{\mathbb{B}^{(i,j)}}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \quad (51)
$$

be the analytic continuation of $\Delta\mathcal{F}_B^{(i,j)}$ to $\overline{\mathbb{B}^{(i,j)}}$. Without loss of generality, we assume (i, j) to be an attractive edge (the calculations for a repulsive edge can be done analogously). To show (**??**), we utilize Lemma 2 and Lemma 3 (b):

$$
\begin{aligned}
& \lim_{\substack{q_i \to 0 \\ q_j \to 0}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \\
=& \lim_{\substack{q_i \to 0 \\ q_j \to 0}} \Delta\mathcal{U}_B^{(i,j)}(q_i, q_j) + \overbrace{\lim_{\substack{q_i \to 0 \\ q_j \to 0}} I_B^{(i,j)}(q_i, q_j)}^{=0} \\
=& \lim_{\substack{q_i \to 0 \\ q_j \to 0}} -(1 + 2(2\,\xi_{ij} - q_i - q_j))\,J_{ij} \\
=& -J_{ij} - \lim_{\substack{q_i \to 0 \\ q_j \to 0}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 0} - \overbrace{q_i}^{\to 0} - \overbrace{q_j}^{\to 0})) \\
=& -J_{ij},
\end{aligned}
$$

$$
\begin{aligned}
& \lim_{\substack{q_i \to 0 \\ q_j \to 1}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \\
=& -J_{ij} - \lim_{\substack{q_i \to 0 \\ q_j \to 1}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 0} - \overbrace{q_i}^{\to 0} - \overbrace{q_j}^{\to 1})) \\
=& -J_{ij} + 2J_{ij} = J_{ij},
\end{aligned}
$$

$$
\begin{aligned}
& \lim_{\substack{q_i \to 1 \\ q_j \to 0}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \\
=& -J_{ij} - \lim_{\substack{q_i \to 1 \\ q_j \to 0}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 0} - \overbrace{q_i}^{\to 1} - \overbrace{q_j}^{\to 0})) \\
=& -J_{ij} + 2J_{ij} = J_{ij},
\end{aligned}
$$

$$
\begin{aligned}
& \lim_{\substack{q_i \to 1 \\ q_j \to 1}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \\
=& -J_{ij} - \lim_{\substack{q_i \to 1 \\ q_j \to 1}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 2} - \overbrace{q_i}^{\to 1} - \overbrace{q_j}^{\to 1})) \\
=& -J_{ij}.
\end{aligned}
$$

For the remainder of the proof, we observe that $\overline{\Delta\mathcal{F}_B^{(i,j)}}$ exhibits monotonic behavior between the corner points of $\overline{\mathbb{B}^{(i,j)}}$. On the one hand, it is monotonically increasing over the two line segments that connect $(0, 0)$ to $(0, 1)$ and $(0, 0)$ to $(1, 0)$:

$$
\begin{aligned}
\overline{\Delta\mathcal{F}_B^{(i,j)}}(0, k) =& \lim_{\substack{q_i \to 0 \\ q_j \to k}} \Delta\mathcal{F}_B^{(i,j)}(q_i, q_j) \\
=& -J_{ij} - \lim_{\substack{q_i \to 0 \\ q_j \to k}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 0} - \overbrace{q_i}^{\to 0} - \overbrace{q_j}^{\to k})) \\
=& -J_{ij} + 2kJ_{ij} = J_{ij}(2k - 1) \nearrow
\end{aligned}
$$

and

$$
\begin{aligned}
\overline{\Delta\mathcal{F}_B^{(i,j)}}(k, 0) =& -J_{ij} - \lim_{\substack{q_i \to k \\ q_j \to 0}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 0} - \overbrace{q_i}^{\to k} - \overbrace{q_j}^{\to 0})) \\
=& -J_{ij} + 2kJ_{ij} = J_{ij}(2k - 1) \nearrow
\end{aligned}
$$

if k in $[0, 1]$ increases. On the other hand, it is monotonically decreasing over the two line segments that connect $(0, 1)$ to $(1, 1)$ and $(1, 0)$ to $(1, 1)$:

$$
\begin{aligned}
\overline{\Delta\mathcal{F}_B^{(i,j)}}(k, 1) =& -J_{ij} - \lim_{\substack{q_i \to k \\ q_j \to 1}} 2J_{ij}(\overbrace{2\xi_{ij}}^{\to 2k} - \overbrace{q_i}^{\to k} - \overbrace{q_j}^{\to 1})) \\
=& J_{ij} - 2kJ_{ij} = J_{ij}(1 - 2k) \searrow
\end{aligned}
$$

and

$$\overline{\Delta\mathcal{F}_B^{(i,j)}}(1,k) = -J_{ij} - \lim_{\substack{q_i \to 1 \\ q_j \to k}} 2J_{ij}\left(\overbrace{2\xi_{ij}}^{\to 2k} - \overbrace{q_i}^{\to 1} - \overbrace{q_j}^{\to k}\right))$$

$$= J_{ij} - 2kJ_{ij} = J_{ij}(1-2k) \searrow$$

if k in $[0,1]$ increases. By this we conclude that

$$\min_{(q_i,q_j)\in\mathbb{B}^{(i,j)}} \overline{\Delta\mathcal{F}_B^{(i,j)}}(q_i,q_j) = -J_{ij}$$

and

$$\max_{(q_i,q_j)\in\mathbb{B}^{(i,j)}} \overline{\Delta\mathcal{F}_B^{(i,j)}}(q_i,q_j) = J_{ij}.$$

By the definition (51) of $\overline{\Delta\mathcal{F}_B^{(i,j)}}$ and in consequence of our previous observations, the statements (31), (32), and hence (30) follow immediately. $\qquad\square$

PROOF OF COROLLARY 1:.

This is an immediate consequence of Theorem 1. $\qquad\square$

PROOF OF THEOREM 2:.

We prove that the derivative of $||\Delta\mathcal{F}_B^{(i,j)}||_{L^2}^2$ with respect to J_{ij} is larger than 0 if $J_{ij} > 0$, and smaller than 0 if $J_{ij} < 0$. Recall that we assume $\mathcal{F}_B$ to be defined on the local polytope instead of the Bethe box (this implies also that $\Delta\mathcal{F}_B^{(i,j)}$ is defined on the sliced local polytope $\mathbb{L}^{(i,j)}$ 19). We compute

$$\frac{\partial}{\partial J_{ij}}||\Delta\mathcal{F}_B^{(i,j)}||_{L^2}^2 \tag{52}$$

$$=\frac{\partial}{\partial J_{ij}} \iiint_{\mathbb{L}^{(i,j)}} (\Delta\mathcal{F}_B^{(i,j)})^2 d\xi_{ij}dq_idq_j \tag{53}$$

$$=\iiint_{\mathbb{L}^{(i,j)}} \frac{\partial}{\partial J_{ij}}(\Delta\mathcal{F}_B^{(i,j)})^2 d\xi_{ij}dq_idq_j \tag{54}$$

$$=\iiint_{\mathbb{L}^{(i,j)}} 2\cdot\Delta\mathcal{F}_B^{(i,j)}\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{F}_B^{(i,j)} d\xi_{ij}dq_idq_j \tag{55}$$

Now observe that $\frac{\partial}{\partial J_{ij}}\Delta\mathcal{F}_B^{(i,j)} = \frac{\partial}{\partial J_{ij}}(\Delta\mathcal{U}_B^{(i,j)} + I_B^{(i,j)})$ (this is just the definition of $\Delta\mathcal{F}_B^{(i,j)}$ (18)), and that the mutual information on the sliced local polytope is independent of J_{ij}. Consequently $\frac{\partial}{\partial J_{ij}}I_B^{(i,j)} = 0$. We continue with (55)

and split the integral:

$$2\iiint_{\mathbb{L}^{(i,j)}} (\Delta\mathcal{U}_B^{(i,j)} + I_B^{(i,j)})\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)} d\xi_{ij}dq_idq_j$$

$$=2\underbrace{\iiint_{\mathbb{L}^{(i,j)}} \Delta\mathcal{U}_B^{(i,j)}\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)} d\xi_{ij}dq_idq_j}_{(I.)}$$

$$+2\underbrace{\iiint_{\mathbb{L}^{(i,j)}} I_B^{(i,j)}\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)} d\xi_{ij}dq_idq_j}_{(II.)}$$

The integral $(I.)$ can be simplified, since

$$\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)} = -1 - 2(2\xi_{ij} - q_i - q_j) = \frac{1}{J_{ij}}\Delta\mathcal{U}_B^{(i,j)}$$

and therefore

$$(I.) = J_{ij}\iiint_{\mathbb{L}^{(i,j)}} (-1 - 2(2\xi_{ij} - q_i - q_j))^2 d\xi_{ij}dq_idq_j,$$

which is larger than 0 for $J_{ij} > 0$ and smaller than 0 for $J_{ij} < 0$.

It remains to prove that the integral $(II.)$ is equal to zero. It is sufficient to consider the integral over the halfspace of the sliced local polytope where $q_i < q_j$ (due to symmetry, the integral over the other halfspace where $q_i > q_j$ is equal to the first). We can split this halfspace in four 'sub-polytopes':

$$\mathbb{L}_1^{(i,j)}: \{(q_i,q_j,\xi_{ij})\in\mathbb{L}^{(i,j)}:$$
$$0 < q_j \le \frac{1}{2}, 0 < q_i \le q_j, 0 < \xi_{ij} < q_i\}$$
$$\mathbb{L}_2^{(i,j)}: \{(q_i,q_j,\xi_{ij})\in\mathbb{L}^{(i,j)}:$$
$$\frac{1}{2} \le q_j < 1, 0 < q_i \le 1 - q_j, 0 < \xi_{ij} < q_i\}$$
$$\mathbb{L}_3^{(i,j)}: \{(q_i,q_j,\xi_{ij})\in\mathbb{L}^{(i,j)}:$$
$$\frac{1}{2} \le q_j < 1, 1 - q_j \le q_i \le \frac{1}{2},$$
$$q_i + q_j - 1 < \xi_{ij} < q_i\}$$
$$\mathbb{L}_4^{(i,j)}: \{(q_i,q_j,\xi_{ij})\in\mathbb{L}^{(i,j)}:$$
$$\frac{1}{2} \le q_j < 1, \frac{1}{2} \le q_i \le q_j,$$
$$q_i + q_j - 1 < \xi_{ij} < q_i\}$$
$$\tag{56}$$

We show that, for each point $(q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)})$ in $\mathbb{L}_1^{(i,j)}$ we can find a point $(q_i^{(2)}, q_j^{(2)}, \xi_{ij}^{(2)})$ in $\mathbb{L}_2^{(i,j)}$ such that the value of the integrand $I_B^{(i,j)}\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)}$ of $(II.)$ in $(q_i^{(2)}, q_j^{(2)}, \xi_{ij}^{(2)})$ is precisely the negative of the value that $I_B^{(i,j)}\cdot\frac{\partial}{\partial J_{ij}}\Delta\mathcal{U}_B^{(i,j)}$ takes in $(q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)})$. To this end, we

map $(q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)})$ from $\mathbb{L}_1^{(i,j)}$ to $(q_i^{(1)}, 1 - q_j^{(1)}, q_i - \xi_{ij}^{(1)})$ in $\mathbb{L}_2^{(i,j)}$. $\qquad\square$

Then one can check that

$$\Delta \mathcal{U}_B^{(i,j)}(q_i^{(1)}, 1 - q_j^{(1)}, q_i - \xi_{ij}^{(1)})$$
$$= -\Delta \mathcal{U}_B^{(i,j)}(q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)})$$

and

$$I_B^{(i,j)}(q_i^{(1)}, 1 - q_j^{(1)}, q_i - \xi_{ij}^{(1)})$$
$$= I_B^{(i,j)}(q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)}),$$

and consequently

$$I_B^{(i,j)} \cdot \frac{\partial}{\partial J_{ij}} \Delta \mathcal{U}_B^{(i,j)}(q_i^{(1)}, 1 - q_j^{(1)}, q_i - \xi_{ij}^{(1)})$$
$$= -I_B^{(i,j)} \cdot \frac{\partial}{\partial J_{ij}} \Delta \mathcal{U}_B^{(i,j)}((q_i^{(1)}, q_j^{(1)}, \xi_{ij}^{(1)})).$$

As our so defined mapping is bijective, we conclude that the integral $(II.)$ over $\mathbb{L}_2^{(i,j)}$ is exactly the negative of the integral $(I.)$ over $\mathbb{L}_1^{(i,j)}$. Similarly, we can define a bijective mapping between $\mathbb{L}_3^{(i,j)}$ and $\mathbb{L}_4^{(i,j)}$ with the same result. That is, the integral $(II.)$ over the entire sliced local polytope equals to zero, which completes the proof.

PROOF OF THEOREM 3:.

If $\mathcal{Z}_\mathcal{B} = \mathcal{Z}_\mathcal{B}^{\setminus(i,j)}$, the statement is trivial and we will therefore exclude this scenario. According to Theorem 1, each edge contributes less than $\pm J_{ij}$ to the 'overall' Bethe function and removing an edge can only transpose the individual function values by less than $|J_{ij}|$. This means that for all $\boldsymbol{q} \in \mathbb{B}$,

$$|\mathcal{F}_B(\boldsymbol{q}) - \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q})| = |\Delta \mathcal{F}_B^{(i,j)}(q_i, q_j)| < |J_{ij}|. \quad (57)$$

In particular, this is also valid for the global minimizer $\boldsymbol{q_1^*}$ of $\mathcal{F}_B$ as well as for the global minimizer $\boldsymbol{q_2^*}$ of $\mathcal{F}_B^{\setminus(i,j)}$. We now distinguish between two cases:

Case 1: $\mathcal{Z}_\mathcal{B} > \mathcal{Z}_\mathcal{B}^{\setminus(i,j)}$, or equivalently, $\mathcal{F}_B(\boldsymbol{q_1^*}) < \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*})$. Then

$$0 < \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*}) - \mathcal{F}_B(\boldsymbol{q_1^*}) < \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_1^*}) - \mathcal{F}_B(\boldsymbol{q_1^*}) \overset{(57)}{<} |J_{ij}|.$$

Case 2: $\mathcal{Z}_\mathcal{B} < \mathcal{Z}_\mathcal{B}^{\setminus(i,j)}$, or equivalently, $\mathcal{F}_B(\boldsymbol{q_1^*}) > \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*})$. Then

$$0 < \mathcal{F}_B(\boldsymbol{q_1^*}) - \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*}) < \mathcal{F}_B(\boldsymbol{q_2^*}) - \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*}) \overset{(57)}{<} |J_{ij}|.$$

Together, we finally obtain

$$|J_{ij}| > |\mathcal{F}_B(\boldsymbol{q_1^*}) - \mathcal{F}_B^{\setminus(i,j)}(\boldsymbol{q_2^*})| = |\min_{\boldsymbol{q} \in \mathbb{B}} \mathcal{F}_B - \min_{\boldsymbol{q} \in \mathbb{B}} \mathcal{F}_B^{\setminus(i,j)}| =$$
$$= |\log(\mathcal{Z}_\mathcal{B}) - \log(\mathcal{Z}_\mathcal{B}^{\setminus(i,j)})| = \left| \log \left(\frac{\mathcal{Z}_\mathcal{B}}{\mathcal{Z}_\mathcal{B}^{\setminus(i,j)}} \right) \right|.$$

$\qquad\square$

PROOF OF THEOREM 4:.

Without loss of generality, we assume that all θ_i are larger than 0.5. From Lemma 10 (Knoll et al., 2021) we know, that the global minimum $-\log(\mathcal{Z}_\mathcal{B})$ of $\mathcal{F}_B$ must be located in a point of $\mathbb{B}$ where all components q_i are larger than 0.5. If we remove an edge, we obtain again a unidirectional model that satisfies the same property. By Lemma 6, we then know that the global minimum of $\mathcal{F}_B$ in the new model must be larger than the global minimum in the old model, i.e.,

$$-\log(\mathcal{Z}_\mathcal{B}) < -\log(\mathcal{Z}_\mathcal{B}^{\setminus(i,j)}) \quad (58)$$

(as the removed energy difference function $\Delta \mathcal{F}_B^{(i,j)}$ makes in the original model a negative contribution to the energy of all points $\boldsymbol{q} \in \mathbb{B}$ with $q_i, q_j > 0.5$ – in particular, to the global minimum of the new model). Finally, we apply Theorem 6 from Ruozzi (2012) which says that in attractive models, the Bethe partition function is always a lower bound to the true partition function, i.e., $-\log(\mathcal{Z}) < -\log(\mathcal{Z}_\mathcal{B})$. Together with (58), we conclude

$$|-\log(\mathcal{Z}) + \log(\mathcal{Z}_\mathcal{B})| < |-\log(\mathcal{Z}) + \log(\mathcal{Z}_\mathcal{B}^{\setminus(i,j)})|,$$

which is equivalent to the statement. $\qquad\square$

APPENDIX B – RESULTS FROM RELATED WORK

Lemma 7. *Let X, Y discrete random variables with alphabets $\mathcal{X}, \mathcal{Y}$. Their mutual information $I(X; Y)$ is bounded by*

$$\frac{1}{2} \Big(\sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} |p_{XY}(x, y) - p_X(x) p_Y(y)| \Big)^2 \leq I(X; Y) \leq$$
$$\leq \sum_{(x,y) \in \mathcal{X} \times \mathcal{Y}} \frac{p_{XY}^2(x, y)}{p_X(x) p_Y(y)} - 1. \quad (59)$$

Proof. Corollary 1 and 2 in Dragomir et al. (2000). $\qquad\square$

Lemma 8.

(a) *If (i, j) is an attractive edge (i.e., $J_{ij} > 0$), then $\xi_{ij}^* > q_i q_j$.*

(b) *If (i, j) is an repulsive edge (i.e., $J_{ij} < 0$), then $\xi_{ij}^* < q_i q_j$.*

Proof. Lemma 2 in Weller and Jebara (2013). $\qquad\square$

Lemma 9. *The first-order derivatives of the Bethe free energy $\mathcal{F}_B$ on the Bethe box $\mathbb{B}$ are*

$$\frac{\partial}{\partial q_i} \mathcal{F}_B = -2 \theta_i + 2 \sum_{j \in \mathcal{N}(i)} J_{ij} +$$
$$+ \log \Big(\frac{(1 - q_i)^{d_i - 1}}{q_i^{d_i - 1}} \prod_{j \in \mathcal{N}(i)} \frac{q_i - \xi_{ij}^*}{1 + \xi_{ij}^* - q_i - q_j} \Big). \quad (60)$$

Proof. This is an intermediate result in Welling and Teh (2001). $\quad\square$

Theorem 5. *The second partial derivatives of edge specific Bethe terms of the form*

$$f_{ij}(q_i, q_j) = -\left(1 + 2\left(2\,\xi_{ij} - q_i - q_j\right)\right) J_{ij} - \mathcal{S}_{ij} \quad (61)$$

are calculated as

$$\frac{\partial^2}{\partial q_i^2} f_{ij} = \frac{q_j(1 - q_j)}{T_{ij}}, \quad (62)$$

$$\frac{\partial^2}{\partial q_i q_j} f_{ij} = \frac{\partial^2}{\partial q_j q_i} f_{ij} = \frac{q_i q_j - \xi_{ij}^*}{T_{ij}}, \quad and \quad (63)$$

$$\frac{\partial^2}{\partial q_j^2} f_{ij} = \frac{q_i(1 - q_i)}{T_{ij}}, \quad (64)$$

where $T_{ij} \coloneqq q_i q_j (1 - q_i)(1 - q_j) - (\xi_{ij}^* - q_i q_j)^2.$

Proof. Theorem 10 in Weller and Jebara (2013). $\quad\square$

Lemma 10. *Consider a unidirectional model and assume without loss of generality that $\theta_i > 0.5$. Then, in the global minimum $\boldsymbol{q}^*$ of $\mathcal{F}_B$ all components q_i are > 0.5.*

Proof. Lemma 2 in Knoll et al. (2021). $\quad\square$

Theorem 6. *For an attractive model with binary variables, the Bethe partition function is always a lower bound on the true partition function. That is,*

$$\mathcal{Z}_\mathcal{B} \leq \mathcal{Z},$$

or equivalently,

$$\min_{\mathbb{M}} \mathcal{F} = -\log(\mathcal{Z}) \leq -\log(\mathcal{Z}_\mathcal{B}) = \min_{\mathbb{L}} \mathcal{F}_B.$$

Proof. Theorem 4.1 in Ruozzi (2012). $\quad\square$

APPENDIX C – FURTHER EXPERIMENTS

Here we present supplementary experiments that we have performed on a 5×5-grid graph. The experimental setting is the same as in Sec. 4. Again, we observe the beneficial effects of edge removal on the approximated marginals. In this sparser model, the treshold beyond which the Bethe free energy becomes non-convex and the accuracy of BP degrades is larger than for the fully connected graph considered in Sec. 4. Beyond that treshold, edge removal is particularly effective if the local potentials are weak; but also for models with strong local potentials the results are mostly superior to the marginal accuracy of BP in the original model. For attractive models with strong local potentials, we observe again the presence of a 'channel' that specifies an optimal intermediate model state.

Figure 3: Attractive models (5x5 - grid graph). First row: $\theta_i \in [-0.2, 0.2]$; second row: $\theta_i \in [-0.5, 0.5]$. (a) + (c): BETHE-OPT criterion; (b) + (d): CHOW LIU criterion.

Figure 4: General models (5x5 - grid graph). First row: $\theta_i \in [-0.2, 0.2]$; second row: $\theta_i \in [-0.5, 0.5]$. (a) + (c): BETHE-OPT criterion; (b) + (d): CHOW LIU criterion.