The Burer-Monteiro SDP method can fail even above the Barvinok-Pataki bound

Abstract

The most widely used technique for solving large-scale semidefinite programs (SDPs) in practice is the non-convex Burer-Monteiro method, which explicitly maintains a low-rank SDP solution for memory efficiency. There has been much recent interest in obtaining a better theoretical understanding of the Burer-Monteiro method. When the rank p of the SDP solution is above the Barvinok-Pataki bound (where a globally optimal solution of rank at most p is guaranteed to exist), a recent line of work established convergence to a global optimum for generic or smoothed instances of the problem. However, it was open whether there even exists an instance in this regime where the Burer-Monteiro method fails. We prove that the Burer-Monteiro method can fail for the Max-Cut SDP on n vertices when the rank is above the Barvinok-Pataki bound ($p \gtrsim \sqrt{2n}$). We provide a family of instances that have spurious local minima even when the rank $p = n/2$. Combined with existing guarantees, this settles the question of the existence of spurious local minima for the Max-Cut formulation in all ranges of the rank, and justifies the use of beyond-worst-case paradigms like smoothed analysis to obtain guarantees for the Burer-Monteiro method.

1 Introduction

Semidefinite programs (SDPs) are a powerful algorithmic tool with wide-ranging applications in combinatorial optimization, control theory, machine learning and operations research. Notably, they yield optimal approximation algorithms for NP-hard problems like Max-Cut [GW95] and other constraint satisfaction problems [Rag08]. While interior-point algorithms and the ellipsoid method give polynomial time guarantees for solving semidefinite programs, memory becomes a bottleneck for relatively modest instance sizes. This has prompted research into scalable semidefinite programming algorithms—see [MHA20] for a recent survey.

One of the most popularly used methods for solving SDPs in practice is the one pioneered by Burer and Monteiro [BM03, BM05], which explicitly constrains the rank of the SDP solution for efficiency. Consider the celebrated Goemans-Williamson SDP relaxation for Max-Cut [GW95]:

$$
\min_{X \in \mathbb{S}^{n \times n}} \langle A, X \rangle
$$
$$
\text{s.t.} \quad X_{ii} = 1 \text{ for } i \in [n], \tag{MC-SDP}
$$
$$
X \succeq 0.
$$

Here $\mathbb{S}^{n \times n}$ denotes the set of all symmetric $n \times n$ matrices and the cost matrix $A \in \mathbb{S}^n$ is the adjacency matrix of a weighted graph.[1] Instead of maintaining a solution X in $\mathbb{S}^{n \times n}$ that is also

---

[1]Note that (MC-SDP) can also be reformulated as a maximization SDP where the cost matrix is the Laplacian matrix $L = D - A$. (D is the degree matrix.)

Submitted to 36th Conference on Neural Information Processing Systems (NeurIPS 2022). Do not distribute.

Figure 1: Combined with existing guarantees, our work settles the question of the existence of spurious local minima for (MC-BM) in all ranges of the rank p.

positive semidefinite (p.s.d), the Burer-Monteiro method explicitly maintains a low-rank solution of the form $X = YY^\top$, where $Y \in \mathbb{R}^{n \times p}$, and aims to solve the following optimization problem:

$$\min_{Y \in \mathbb{R}^{n \times p}} \left\langle A, YY^\top \right\rangle$$
$$\text{s.t.} \quad ||Y_i||^2 = 1 \text{ for } i \in [n], \tag{MC-BM}$$

where $Y_i \in \mathbb{R}^p$ denotes the ith row of Y, taken as a column vector, and $|| \cdot ||$ denotes the Euclidean norm. We may also denote the objective of (MC-BM) as $\text{OBJ}(Y) := \left\langle A, YY^\top \right\rangle$. We denote the feasible region as

$$\mathcal{M}_{n,p} := \left\{ Y \in \mathbb{R}^{n \times p} : ||Y_i||^2 = 1 \text{ for } i \in [n] \right\}.$$

(When n is clear from the context or unimportant, we may just write $\mathcal{M}_p$.) Note that $\mathcal{M}_{n,p}$ is a manifold that can be viewed as a Cartesian product of n unit spheres in $\mathbb{R}^p$.

This formulation yields significant memory savings when $p \ll n$ by storing Y instead of X, and has the additional advantage of dropping the positive semidefiniteness constraint. On the other hand, the formulation in (MC-BM) is a non-convex constrained optimization problem. Local optimization methods like Riemannian gradient descent, and other heuristics for manifold optimization or constrained optimization are used to solve this non-convex problem with surprisingly good empirical results [BM03, BM05]. This motivates the following question:

When does the Burer-Monteiro method converge to a globally optimal solution?

This question has attracted much recent interest on the theoretical front. It is known that when the rank p of the SDP solution is above the so-called *Barvinok-Pataki bound* $p \gtrsim \sqrt{2n}$ (more formally, p satisfies $\frac{p(p+1)}{2} \geq n$), a globally optimal solution of rank at most p is guaranteed to exist [Bar95, Pat98]. Moreover, below this bound, there exist instances for which (MC-BM) has a spurious local minimum [WW20]. When the rank is above the Barvinok-Pataki bound, Boumal, Voroninski, and Bandeira [BVB16, BVB18] showed that for *generic* instances[2] (or smoothed instances) $A \in \mathbb{S}^n$, any second-order critical point Y of (MC-BM) is globally optimal for (MC-BM), and thus $YY^\top$ is optimal for (MC-SDP) (their result also applies for a broad class of SDPs with equality constraints). Under such conditions, known algorithms converge to second-order critical points of (MC-BM), and polynomial time convergence guarantees can be shown for smoothed instances [BBJN18, PJB18, CM19].

On the other hand, for general cost matrices A, the best known bound only guarantees convergence to a global optimum of the Burer-Monteiro method when $p > \frac{n}{2}$ [BVB18]. For general cost matrices A of (MC-BM), there is a large range for the rank $p \in (\sqrt{2n}, n/2)$ where we do not know whether the Burer-Monteiro method works. To the best of our knowledge, it was open whether there exists *any* instance (not specific to Max-Cut) where the Burer-Monteiro method fails above the Barvinok-Pataki bound. In fact, the authors of [BVB18] pose this question:

---

[2]By guarantees for generic instances, we mean that the guarantees hold for all instances (cost matrices) except a set of zero measure.

Our main theorem resolves this question by constructing an instance of (MC-BM) with a spurious local minimum (i.e., a local minimum which is not globally optimal), when p is as large as $\Theta(n)$ (note that a spurious local minimum is also a spurious second order critical point). This result is contextualized in Figure 1.

Theorem 1. *For any $n \geq 4$ and $2 \leq p \leq n/2$, there exist cost matrices A for which the associated instance of* (MC-BM) *has a spurious local minimum.*

Combining Theorem 1 with other existing results yields a clearer understanding of the optimization landscape and a characterization of the range of p for which (MC-BM) can have spurious local minima. Furthermore, our result justifies the use of beyond worst-case paradigms like smoothed analysis to obtain global convergence guarantees for the Burer-Monteiro method.

Outline of paper. We begin with Section 2 (Preliminaries) which gives necessary and sufficient conditions for local/global optimality for (MC-BM), introduces Riemannian gradient descent (necessary for our later arguments), and provides other definitions which will be used in our construction of spurious local minima for (MC-BM). In Section 3, we list the main lemmas for Theorem 1 and show how Theorem 1 follows. In Sections 4 and 5, we discuss some of these lemmas in more detail. In particular, Section 4 and 5 sketch the proofs that our construction yields a spurious second-order critical point and spurious local minimum respectively. We present potential follow-up directions in Section 6 (Conclusion).

2 Preliminaries

2.1 Riemannian derivatives

$\mathcal{M}_p$ is a smooth embedded submanifold of $\mathbb{R}^{n \times p}$ [BVB18, Prop. 1.2], and is furthermore an embedded Riemannian submanifold of $\mathbb{R}^{n \times p}$ [Bou22, Def. 3.55] if we equip the linearizations of $\mathcal{M}_p$ at each point (known as the *tangent spaces*—defined below) with (a restriction of) the inner product $\langle \cdot, \cdot \rangle$.

Proposition 1 (Tangent space [BVB18, Lem. 2.1]). *The tangent space to $\mathcal{M}_p$ at $Y \in \mathcal{M}_p$, denoted $\mathrm{T}_Y \mathcal{M}_p$, is the following subspace of $\mathbb{R}^{n \times p}$:*

$$\mathrm{T}_Y \mathcal{M}_p = \left\{ U \in \mathbb{R}^{n \times p} : \langle Y_i, U_i \rangle = 0 \, for \, i \in [n] \right\}.$$

Here, $Y_i \in \mathbb{R}^p$, $U_i \in \mathbb{R}^p$ denote the ith rows of Y and U respectively, taken as column vectors. We call an element of $\mathrm{T}_Y \mathcal{M}_p$ a tangent vector (at Y).

The Riemannian gradient at $Y \in \mathcal{M}_p$, $\mathrm{grad}\,\mathrm{OBJ}(Y)$, is the orthogonal projection of the classical Euclidean gradient at Y onto $\mathrm{T}_Y \mathcal{M}_p$, which yields the following expression (see Appendix A):

$$\mathrm{grad}\,\mathrm{OBJ}(Y) = 2(A - \mathrm{diag}(\nu))Y, \quad \text{where } \nu_i := \sum_{j=1}^n A_{ij} \langle Y_i, Y_j \rangle, \text{for all } i \in [n]. \tag{1}$$

(Note that $\nu \in \mathbb{R}^n$ is a function of Y, although we write ν instead of $\nu(Y)$ when Y is clear from context.) We take (1) to be the definition of ν from now on.

The Riemannian Hessian of OBJ at $Y \in \mathcal{M}_p$, $\mathrm{Hess}\,\mathrm{OBJ}(Y)$, is a linear, symmetric map from $\mathrm{T}_Y \mathcal{M}_p$ to $\mathrm{T}_Y \mathcal{M}_p$ given by the classical differential of (a smooth extension of) $\mathrm{grad}\,\mathrm{OBJ}(Y)$, projected to the tangent space [Bou22, Cor. 5.16]. This yields (see Appendix A):

$$\mathrm{Hess}\,\mathrm{OBJ}(Y)[U] = 2\mathrm{Proj}_Y \left((A - \mathrm{diag}(\nu))U \right)$$

for $U \in \mathrm{T}_Y \mathcal{M}_p$, where the linear map $\mathrm{Proj}_Y : \mathbb{R}^{n \times p} \to \mathrm{T}_Y \mathcal{M}_p$ denotes the orthogonal projector onto $\mathrm{T}_Y \mathcal{M}_p \subseteq \mathbb{R}^{n \times p}$, i.e., $\mathrm{Proj}_Y(Z) = \mathrm{argmin}_{U \in \mathrm{T}_Y \mathcal{M}_p} \|U - Z\|$. In what follows, $\mathrm{Hess\,OBJ}(Y)$ will only appear as part of a quadratic form, yielding the following cleaner expression:

$$\langle \mathrm{Hess\,OBJ}(Y)[U], U \rangle = \langle 2\mathrm{Proj}_Y \left((A - \mathrm{diag}(\nu))U \right), U \rangle = 2 \left\langle A - \mathrm{diag}(\nu), UU^\top \right\rangle,$$

for $U \in \mathrm{T}_Y \mathcal{M}_p$, where we used the fact that $\langle \mathrm{Proj}_Y(Z), U \rangle = \langle Z, U \rangle$ for any $Z \in \mathbb{R}^{n \times p}, U \in \mathrm{T}_Y \mathcal{M}_p$.

2.2 Necessary and sufficient conditions

Recall the definition of a local (and global) minimum:

Definition 1 (Local/global minimum). *Consider the program*

$$\min_{x \in D} f(x)$$

where $D \subseteq \mathbb{R}^d$. $x \in D$ is a local minimum if there exists $\epsilon > 0$ such that if $y \in D$ and $\|y - x\| < \epsilon$, we have $f(x) \leq f(y)$. $x \in D$ is a global minimum if $f(x) \leq f(y)$ for all $y \in D$.

The following are standard necessary conditions for local optimality, and correspond to the first and second-order critical point criteria that need to be satisfied by any local minimum of (MC-BM) (see [BVB18, Prop 2.4]).

Proposition 2 (First-order critical point [BVB18, Def. 2.3] and [WW20, Prop. 3]). *$Y \in \mathcal{M}_p$ is a first-order critical point for (MC-BM) if and only if $\mathrm{grad\,OBJ}(Y) = 2(A - \mathrm{diag}(\nu))Y = 0$. Equivalently, $Y \in \mathcal{M}_p$ is a first-order critical point if and only if there exists $\lambda \in \mathbb{R}^n$ such that $(A - \mathrm{diag}(\lambda))Y = 0$. If such a λ exists, it is unique and equal to ν given by (1).*

Proposition 3 (Second-order critical point [WW20, Prop. 4]). *A first-order critical point $Y \in \mathcal{M}_p$ is additionally a second-order critical point if and only if*

$$\langle \mathrm{Hess\,OBJ}(Y)[U], U \rangle = 2 \left\langle A - \mathrm{diag}(\nu), UU^\top \right\rangle \geq 0$$

for all $U \in \mathrm{T}_Y \mathcal{M}_p$, and where ν is given in (1). (This is equivalent to $\mathrm{Hess\,OBJ}(Y) \succeq 0$.)

We say a critical point or local minimum is *spurious* if it is not globally optimal. Finally, we characterize which first-order critical points of (MC-BM) are globally optimal. Since second-order critical points and local minima are also first-order critical points, this also provides a characterization of optimality for them.

Proposition 4 (Characterization of optimality for first-order critical points). *A first-order critical point Y of (MC-BM) is globally optimal if and only if $A - \mathrm{diag}(\nu) \succeq 0$, where ν is given in (1).*

While not framed precisely in this way, Proposition 4 follows directly from prior work (see Appendix A for details). The "if" direction involves showing that the given assumptions imply that the primal-dual pair $(YY^\top, \nu)$ satisfies the Karush–Kuhn–Tucker (KKT) conditions of (MC-SDP), implying (after a few simple additional steps) that Y is globally optimal for (MC-BM).

2.3 Riemannian gradient descent

The analogue to gradient descent for optimizing over a smooth manifold is Riemannian gradient descent (see [Bou22, Ch. 4] for an introduction), which yields analogous analyses and guarantees. It includes as part of its specification a *retraction* [Bou22, Def. 3.47]. A rectraction on $\mathcal{M}_p$ associates to each point $Y \in \mathcal{M}_p$ a map $\mathrm{R}_Y : \mathrm{T}_Y \mathcal{M}_p \to \mathcal{M}_p$ which converts movement in the tangent space to movement on the manifold $\mathcal{M}_p$. We use the natural *metric projection retraction* [Bou22, Sec. 5.12], defined for $Y \in \mathcal{M}_p, U \in \mathrm{T}_Y \mathcal{M}_p$ as

$$\mathrm{R}_Y(U) := \mathrm{argmin}_{Z \in \mathcal{M}_p} \|(Y + U) - Z\| = \mathrm{Diag}((Y + U)(Y + U)^\top)^{-1/2}(Y + U),$$

where $\mathrm{Diag}(X)$ denotes the diagonal matrix whose diagonal elements are those of X. In other words, $\mathrm{R}_Y(U)$ is $Y + U$ followed by a normalization of each row. This yields the following Riemannian gradient descent algorithm for (MC-BM):

Input: Initializer $Y^{(0)} \in \mathcal{M}_p$, step size $\eta > 0$.

For $t = 0, 1, 2, \ldots$

$$Y^{(t+1)} = \mathrm{R}_{Y^{(t)}} \left(-\eta \cdot \mathrm{grad\,OBJ}(Y^{(t)}) \right).$$

 2.4 Construction-specific definitions

 Finally, we give a few miscellaneous technical definitions which will be used in our construction of a
 spurious local minimum for (MC-BM).

 Definition 2 (Axial position). *We call the matrix*

$$\widetilde{Y} := \begin{bmatrix} I_{(n/2)} \\ -I_{(n/2)} \end{bmatrix} \in \mathcal{M}_{n,(n/2)}$$

 the axial position.

 We use the term "axial position" because when we view $\mathcal{M}_{n,(n/2)}$ as a Cartesian product of n unit
 spheres in $\mathbb{R}^{n/2}$, $\widetilde{Y}$ corresponds to placing a single unit vector on both the negative and positive sides
 of each axis in $\mathbb{R}^{n/2}$.

 The following set of matrices will be important in our construction:

 Definition 3 (Pseudo-PD, pseudo-PSD). *We say a matrix $M \in \mathbb{S}^{n \times n}$ is pseudo-PD ("pseudo positive*
 definite") if $M[i] \succ 0$ for all $i \in [n]$, where $M[i] \in \mathbb{S}^{(n-1) \times (n-1)}$ denotes the submatrix of M
 formed by removing the ith row and column. Similarly, we say that $M \in \mathbb{S}^{n \times n}$ is pseudo-PSD
 ("pseudo positive semidefinite") if $M[i] \succeq 0$ for all $i \in [n]$.

 We let $\mathbb{S}^{n \times n}_{\text{pseudo-PD}}, \mathbb{S}^{n \times n}_{\text{pseudo-PSD}}$ denote the sets of pseudo-PD and pseudo-PSD matrices respectively
 in $\mathbb{S}^{n \times n}$.

 Note that $\mathbb{S}^{n \times n}_{\text{pseudo-PSD}} \subseteq \mathbb{S}^{n \times n}_{+}$ since the submatrices of a positive semidefinite matrix are also
 positive semidefinite.

 Definition 4 (Strictly pseudo-PD, strictly pseudo-PSD). *We say a matrix is $M \in \mathbb{S}^{n \times n}$ is strictly*
 pseudo-PD if it is pseudo-PD but not positive semidefinite. We say M is strictly pseudo-PSD if it is
 pseudo-PSD but not positive semidefinite. (Note that in both cases we require $M \not\succeq 0$, not $M \not\succ 0$.)
 We let

$$\mathbb{S}^{n \times n}_{\text{strict-PD}} := \mathbb{S}^{n \times n}_{\text{pseudo-PD}} \setminus \mathbb{S}^{n \times n}_{+}, \qquad \mathbb{S}^{n \times n}_{\text{strict-PSD}} := \mathbb{S}^{n \times n}_{\text{pseudo-PSD}} \setminus \mathbb{S}^{n \times n}_{+},$$

 denote the set of strictly pseudo-PD matrices and strictly pseudo-PSD matrices respectively.

 Clearly every strictly pseudo-PD matrix is also strictly pseudo-PSD, but the converse turns out to be
 false.

3 Proof of Theorem 1

 In this section we prove Theorem 1. We focus on the case where n is even and $p = n/2$ and construct
 cost matrices for which $\widetilde{Y}$ is a spurious local minimum, since constructions for $p < n/2$ can be easily
 extracted from the former. Before this, as a warm up, we characterize those cost matrices for which
 $\widetilde{Y}$ is a spurious first and second-order critical point in the following two lemmas. An overview of the
 proofs of these lemmas is given in Section 4, and the full proofs can be found in Appendix B.

 Lemma 1 (First-order critical point characterization for $\widetilde{Y}$). *For (MC-BM) when $p = n/2$, the axial*
 position $\widetilde{Y}$ is a first-order critical point if and only if the cost matrix A takes the form

$$A = \begin{bmatrix} B & B \\ B & B \end{bmatrix} + \text{diag}(\alpha) \tag{2}$$

 for some $\alpha \in \mathbb{R}^n$ and $B \in \mathbb{S}^{\frac{n}{2} \times \frac{n}{2}}$. Furthermore, $\widetilde{Y}$ is additionally spurious if and only if $B \not\succeq 0$.

 Lemma 2 (Second-order critical point characterization for $\widetilde{Y}$). *For (MC-BM) when $p = n/2$, the*
 axial position $\widetilde{Y}$ is a second-order critical point if and only if the cost matrix A takes the form

$$A = \begin{bmatrix} P & P \\ P & P \end{bmatrix} + \text{diag}(\alpha) \tag{3}$$

 for some $\alpha \in \mathbb{R}^n$ and pseudo-PSD $P \in \mathbb{S}^{\frac{n}{2} \times \frac{n}{2}}_{\text{pseudo-PSD}}$. Furthermore, $\widetilde{Y}$ is additionally spurious if
 and only if M is strictly pseudo-PSD.

Figure 2: A spurious minimum and corresponding global minimum for $n = 4, p = 2$.

The following lemma provides sufficient conditions for $\widetilde{Y}$ to be a (spurious) local minimum. This is the most challenging of our results to prove, and we give an overview of the proof and discuss the challenges involved in Section 5. The full proof is given in Appendix C.

Lemma 3 (Local minimum condition for $\widetilde{Y}$). *For* (MC-BM) *when* $p = n/2$, *the axial position* $\widetilde{Y}$ *is a local minimum if the cost matrix A takes the form*

$$A = \begin{bmatrix} M & M \\ M & M \end{bmatrix} + \operatorname{diag}(\alpha) \tag{4}$$

for some $\alpha \in \mathbb{R}^n$ *and pseudo-PD* $M \in \mathbb{S}^{\frac{n}{2} \times \frac{n}{2}}_{\text{pseudo-PD}}$. *Furthermore,* $\widetilde{Y}$ *is additionally spurious if M is strictly pseudo-PD.*

Actualizing Lemma 3 to construct a spurious local minimum requires the existence of strictly pseudo-PD matrices, which we posit in the following proposition:

Proposition 5 (Existence of strictly pseudo-PD matrices). $\mathbb{S}^{k \times k}_{\text{strict-PD}}$ *is nonempty for any* $k > 2$.

We provide constructions of pseudo-PD matrices in Appendix D. Our main nonnegative construction takes the form $UU^\top - \epsilon I_k$, where $U \in \mathbb{R}^{k \times (k-1)}$ is a random matrix. It is shown $(UU^\top)[i] \succ 0$ for all i with high probability. One can show that when $\epsilon > 0$ is sufficiently small, the $(k-1) \times (k-1)$ principal submatrices of $UU^\top - \epsilon I_k$ remain positive definite, while $UU^\top - \epsilon I_k \not\succeq 0$ since $UU^\top$ is rank-deficient.

We use Lemma 3 and Proposition 5 to prove Theorem 1:

Proof of Theorem 1. Lemmas 3 and Proposition 5 imply that for even n such that $n \geq 4$, there exists an instance with a spurious local minimum when $p = n/2$. To construct a spurious local minimum for $p < n/2$, we can simply use the construction for $n' = 2p$ vertices and rank p. The entries of the cost matrix that correspond to additional rows of elements of $\mathcal{M}_{n,p}$ past n' can be set to 0, ensuring that these additional rows can not affect the objective value. $\square$

Our construction instantiated for $n = 4, p = 2$ is illustrated in Figure 2. In this visualization, we think of the nonnegative cost matrix as the weighted adjacency matrix of a graph, which is natural for (MC-BM). Each row of Y specifies the position of one of the vertices on the unit sphere in $\mathbb{R}^p$ (for $p = 2$, the circle shown in gray). We illustrate "heavy" (higher weight) edges with solid lines, and "light" (lower weight) edges with dashed lines. Intuitively, each edge "wants" to push its endpoints away from each other, with heavier edges pushing harder. We see by symmetry that this state is in equilibrium (each vertex is pushed equally clockwise and counter-clockwise), so the gradient is 0. Showing that this instance is indeed a local minimum is much more involved, and requires arguing that if the vertices were perturbed slightly, the heavy edges would still approximately cancel out, and the main force on the vertices would be the light edges pushing the pairs a, c and b, d back to being diametrically opposite.

4 Overview of proof of spurious second-order critical point (Lemmas 1 and 2)

In this section we provide overviews of the proofs of Lemmas 1 and 2, which characterize the cost matrices for which $\widetilde{Y}$ is a spurious/optimal first and second-order critical point respectively (see Appendix B for the full proofs).

4.1 Overview of the proof of spurious first-order criticality (Lemma 1)

Supposing the cost matrix A takes the form (2), one can set $\lambda \leftarrow \alpha$ where $\lambda \in \mathbb{R}^n$ is the multiplier from Proposition 2, and then observe $(A - \lambda)\widetilde{Y} = 0$, implying $\widetilde{Y}$ is a first-order critical point. For the other direction, one can show that if $\widetilde{Y}$ a first-order critical point with associated multiplier $\lambda \in \mathbb{R}^n$ (from Proposition 2), then $(A - \operatorname{diag}(\lambda))\widetilde{Y} = 0$ implies $A - \operatorname{diag}(\lambda)$ must take the block form $\begin{bmatrix} B & B \\ B & B \end{bmatrix}$ for some $B \in \mathbb{S}^{\frac{n}{2} \times \frac{n}{2}}$. Thus, A takes the form (2) with $\alpha \leftarrow \lambda$.

To show $\widetilde{Y}$ is additionally spurious if and only if $B \not\succeq 0$, note from above that if Y is a first-order critical point, the unique associated multiplier λ is equal to α. Recall from Proposition 2 that $\nu = \lambda$ at a first-order critical point, implying $A - \operatorname{diag}(\nu) = A - \operatorname{diag}(\alpha) = \begin{bmatrix} B & B \\ B & B \end{bmatrix}$. Such a matrix is positive semidefinite if and only if B is positive semidefinite, and Proposition 4 concludes the proof.

4.2 Overview of the proof of spurious second-order criticality (Lemma 2)

Since every second-order critical point is a first-order critical point, we assume the cost matrix A takes the form (2) and show that $\widetilde{Y}$ is additionally a second-order critical point if and only if B is additionally pseudo-PSD. Recall from Section 4.1 that when $\widetilde{Y}$ is a first-order critical point, $A - \operatorname{diag}(\nu) = \begin{bmatrix} B & B \\ B & B \end{bmatrix}$. Now, one can observe that an arbitrary element $U \in \mathrm{T}_{\widetilde{Y}}\mathcal{M}_{n/2}$ must take the following form: $U_{ii} = U_{(n/2)+i,i} = 0$ for all $i \in [n/2]$, and all other entries of U are arbitrary. Due to this, one can show that

$$\left\langle \operatorname{Hess} \operatorname{OBJ}(\widetilde{Y})[U], U \right\rangle = 2 \left\langle A - \operatorname{diag}(\nu), UU^\top \right\rangle \geq 0 \quad \text{for all } U \in \mathrm{T}_{\widetilde{Y}}\mathcal{M}_{n/2}$$

$$\Longleftrightarrow 2 \left\langle \begin{bmatrix} B & B \\ B & B \end{bmatrix}, uu^\top \right\rangle \geq 0 \quad \text{for all } u \in R_1 \cup \cdots \cup R_{n/2}, \tag{5}$$

where R_i is the subspace of $\mathbb{R}^n$ consisting of those vectors v such that $v_i = v_{(n/2)+i} = 0$. Restricting u to only lie in R_i in (5) is equivalent to the matrix $\begin{bmatrix} B[i] & B[i] \\ B[i] & B[i] \end{bmatrix} \in \mathbb{S}^{(n-2) \times (n-2)}$ being positive semidefinite, which is equivalent to $B[i]$ being positive semidefinite. (Recall that $B[i]$ denotes the submatrix of B formed by removing row i and column i.) Then (5), which takes the preceding statement over all $i \in [n/2]$, is equivalent to B being pseudo-PSD. (We note that the preceding portion of the proof is formalized in a slightly different way in Appendix B.) The second half of Lemma 2 (characterizing when the second-order critical point $\widetilde{Y}$ is spurious) follows immediately from the corresponding characterization in Lemma 1, since every second-order critical point is a first-order critical point.

5 Overview of the proof of local minimum (Lemma 3)

In this section we give an overview of the proof of Lemma 3 (see Appendix C for the full proof).

The second half of Lemma 3, which states that $\widetilde{Y}$ is spurious if M is strictly pseudo-PD, follows very quickly from the first half of Lemma 3 combined with propositions we have already given, so we do not devote space to it here.

Here, we provide an overview of the challenging first half of Lemma 3: proving that if the cost matrix A takes the form[3] $\begin{bmatrix} M & M \\ M & M \end{bmatrix}$ for $M \in \mathbb{S}^{\frac{n}{2} \times \frac{n}{2}}_{\text{pseudo}-\text{PD}}$, then $\widetilde{Y}$ is a local minimum. Recall that to prove that $\widetilde{Y}$ is a local minimum for such a cost matrix, we must show that there is a neighborhood (which has non-zero measure, by definition) surrounding $\widetilde{Y}$, such that every Y in the neighborhood satisfies $\operatorname{OBJ}(Y) \geq \operatorname{OBJ}(\widetilde{Y})$.

---

[3]We assume without loss of generality that $\alpha = 0$, since scaling the diagonal entries of A just corresponds to adding a constant to the objective value of each feasible point.

Challenges. Unfortunately, arguing about the actual value of the objective function at some point Y near $\widetilde{Y}$ is challenging, and classical techniques for proving that $\widetilde{Y}$ is a local minimum fail. For example, [WW20], which constructs spurious local minima for (MC-BM) when $p \ll \sqrt{2n}$, similarly first constructs spurious second-order critical points and then proves that they are additionally local minima. However, their proof follows because their spurious second-order critical points are non-degenerate [WW20, Def. 3], which corresponds to the rank of the Riemannian Hessian being sufficiently high. It can be shown that every spurious second-order critical point for (MC-BM) when $p = n/2$ is degenerate, meaning this approach won't work. Furthermore, there is no hope of using the positivity of a higher-order Riemannian derivative (e.g., the fourth derivative) to prove that $\widetilde{Y}$ is a local minimum, since it can be shown that all higher-order derivatives are degenerate. (See Appendix C for further discussion of these challenges.)

Thus, we provide a novel approach in which we show that there exists a neighborhood of $\widetilde{Y}$ such that every point Y in it either has $\mathrm{OBJ}(Y) = \mathrm{OBJ}(\widetilde{Y})$, or Riemannian gradient descent initialized at Y converges to some Y' with $\mathrm{OBJ}(Y') = \mathrm{OBJ}(\widetilde{Y})$. We show that Riemannian gradient descent with sufficiently small step size will only decrease the objective value for this optimization problem, so Y must have had $\mathrm{OBJ}(Y) \geq \mathrm{OBJ}(\widetilde{Y})$ to begin with.

It would be convenient to say that every Y in the neighborhood converges to $\widetilde{Y}$, but this is not actually true due to the degeneracy mentioned above. A key observation is that all *antipodal* configurations have the same objective function value, where an antipodal configuration is one where $Y_i = -Y_{(n/2)+i}$ for all $i \in [n/2]$. (Recall that $Y_i \in \mathbb{R}^{n/2}$ denotes the ith row of Y, taken as a column vector.) These antipodal configurations correspond to "flat" directions from $\widetilde{Y}$. For a given Y, it is not a priori clear which of these antipodal configurations it will converge to. So we argue that Y will converge to *some* antipodal configuration.

Proof sketch. The argument requires us to measure progress in converging to some unknown antipodal configuration. We do this with the scalar potential function

$$\Phi(Y) = \|E\|_F^2\,,$$

where $E \in \mathbb{R}^{\frac{n}{2} \times \frac{n}{2}}$ is defined such that the ith row of E is $Y_i + Y_{(n/2)+i}$. When Y is antipodal, each row of E becomes 0, and $\Phi(Y) = 0$. Otherwise, $\Phi(Y) > 0$.

The crux of our argument is two main lemmas.

Lemma 4 (Φ decreases with Riemannian gradient descent). *For $\eta > 0$ sufficiently small, there exists a sufficiently small $\beta > 0$, such that for any Y that is β-close to $\widetilde{Y}$,*

$$\Phi(Y') \leq \left(1 - K\eta\right)\Phi(Y),$$

where Y' is the configuration reached by an η-step of Riemannian gradient descent from Y, and K is a constant that depends only on M.

The precise definition of "β-close" is left to the full proof in Appendix C. Informally, it defines a natural notion of distance between feasible points.

Lemma 5 (Riemannian gradient descent stays close to $\widetilde{Y}$). *For $\eta > 0, \beta > 0$ chosen as in Lemma 4, there exists a $\beta' > 0$ such that for any Y that is β'-close to $\widetilde{Y}$, Riemannian gradient descent from Y with step size η only visits points Y' that are at least β-close to $\widetilde{Y}$.*

Together, these lemmas imply that if we start at a Y close enough to $\widetilde{Y}$, then Riemannian gradient descent from Y will stay at points close to $\widetilde{Y}$ (Lemma 5), and therefore every successive step will make progress in converging to an antipodal configuration (Lemma 4). This allows us to say that the set of points that are β'-close to $\widetilde{Y}$ certify that it is indeed a local minimum, and conclude Lemma 3.

To prove Lemma 4, we note that since Y is already close to axial, the ith index of $Y_i + Y_{(n/2)+i}$ is very close to 0 (that is, it makes up a very small fraction of the row norm). This means the diagonal entries of E are approximately 0. We show that the update to the matrix E resulting from Riemannian gradient descent looks like $M \cdot E$ with each row i projected onto the tangent plane at Y_i (to the unit

sphere in $\mathbb{R}^{n/2}$), plus another copy with each row i projected onto the tangent plane at $Y_{(n/2)+i}$. Since Y is close to the axial position $\widetilde{Y}$, these projections approximately zero out the ith term of each row i, and leave every other term intact.

Let c_i be the ith column of E. The negative gradient update looks like the projection of $M \cdot c_i$. Since the ith term of c_i is approximately 0, this effectively ignores the ith column of M in the matrix multiplication. Then the projection zeroes out the ith term of the product, which effectively ignores the ith row of M. So the update to column i of E looks like

$$M[i] \cdot c_i,$$

where we abuse notation slightly and define $M[i]$ to be the matrix M with the ith row and column set to 0. (This differs from the standard usage, in which $M[i]$ has smaller dimension than M.) Since $M[i]$ is positive definite for every i, we know that $M[i] \cdot c_i$ is partially aligned with c_i. So subtracting this update from c_i will decrease the norm of c_i. Since this occurs for every column of E, Φ decreases. The full proof (in Appendix C) focuses on bounding these approximations, and showing that they do not change the outcome too much.

To prove Lemma 5, we show that each Y_i moves at most a distance proportional to Φ on each step. Since Φ decreases geometrically, the total distance moved by Y_i over the course of Riemannian gradient descent is bounded by a geometric series. Choosing a neighborhood such that the initial Φ is sufficiently small ensures that no Y_i can ever get too far away from its axial position.

The full proofs of these lemmas can be found in Appendix C.

6   Conclusion

We show that the Burer-Monteiro method can fail for instances of the Max-Cut SDP, for rank up to $\frac{n}{2}$. To the best of our knowledge, prior to our work it was unknown whether the Burer-Monteiro method could fail for any instance of any SDP with rank above the Barvinok-Pataki bound. We settle this question and thus justify the use of smoothed analysis to obtain guarantees for the Burer-Monteiro method.

There are many interesting potential follow-up directions to this work. We provide one construction of (MC-BM) instances with spurious local minima. Our construction has, and relies on, many interesting properties and symmetries. It is possible that some of these properties are necessary, and further analyzing this construction could give insight that allows us to fully characterize spurious minima for (MC-BM) instances. Analyzing this construction could also help us understand this interesting threshold phenomenon for this for (MC-BM) when $p = n/2$ – one dimension higher and there are not only no spurious local minima, there are no spurious second-order critical points at all. Another potential direction is seeing if similar techniques can be used to construct instances with spurious local minima for SDPs with other structures (not just Max-Cut).

Lastly, we note that a limitation of our work is that it only points to the existence of local minima, and does not give a full characterization of when we can expect local minima to exist. We also note that since this is a theoretical result about optimization landscapes, we do not foresee any adverse societal impacts of our work.

References

[Bar95]   Alexander I. Barvinok. Problems of distance geometry and convex properties of quadratic maps. *Discrete & Computational Geometry*, 13:189–202, 1995.

[BBJN18] Srinadh Bhojanapalli, Nicolas Boumal, Prateek Jain, and Praneeth Netrapalli. Smoothed analysis for low-rank solutions to semidefinite programs in quadratic penalty form. In *COLT*, 2018.

[BM03]   Samuel Burer and Renato D. C. Monteiro. A nonlinear programming algorithm for solving semidefinite programs via low-rank factorization. volume 95, pages 329–357. 2003. Computational semidefinite and second order cone programming: the state of the art.

[BM05]   Samuel Burer and Renato D. C. Monteiro. Local minima and convergence in low-rank semidefinite programming. *Math. Program.*, 103(3, Ser. A):427–444, 2005.

[Bou22] Nicolas Boumal. An introduction to optimization on smooth manifolds. To appear with Cambridge University Press, May 2022.

[BVB16] Nicolas Boumal, Vlad Voroninski, and Afonso Bandeira. The non-convex burer-monteiro approach works on smooth semidefinite programs. In D. Lee, M. Sugiyama, U. Luxburg, I. Guyon, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 29. Curran Associates, Inc., 2016.

[BVB18] Nicolas Boumal, Vladislav Voroninski, and Afonso Bandeira. Deterministic guarantees for burer-monteiro factorizations of smooth semidefinite programs. *Communications on Pure and Applied Mathematics*, 73, 04 2018.

[CM19] Diego Cifuentes and Ankur Moitra. Polynomial time guarantees for the burer-monteiro method. *arXiv: Optimization and Control*, 2019.

[GW95] Michel X. Goemans and David P. Williamson. Improved approximation algorithms for maximum cut and satisfiability problems using semidefinite programming. *J. ACM*, 42(6):1115–1145, nov 1995.

[MHA20] Anirudha Majumdar, Georgina Hall, and Amir Ali Ahmadi. Recent scalability improvements for semidefinite programming with applications in machine learning, control, and robotics. *Annual Review of Control, Robotics, and Autonomous Systems*, 3(1):331–360, 2020.

[Pat98] Gábor Pataki. On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues. *Mathematics of Operations Research*, 23(2):339–358, 1998.

[PJB18] Thomas Pumir, Samy Jelassi, and Nicolas Boumal. Smoothed analysis of the low-rank approach for smooth semidefinite programs. In S. Bengio, H. Wallach, H. Larochelle, K. Grauman, N. Cesa-Bianchi, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 31. Curran Associates, Inc., 2018.

[Rag08] Prasad Raghavendra. Optimal algorithms and inapproximability results for every csp? In *Proceedings of the Fortieth Annual ACM Symposium on Theory of Computing*, STOC '08, page 245–254, New York, NY, USA, 2008. Association for Computing Machinery.

[WW20] Irène Waldspurger and Alden Waters. Rank optimality for the Burer-Monteiro factorization. *SIAM J. Optim.*, 30(3):2577–2602, 2020.