Equality Constraints in Linear Hawkes Processes

Abstract

Conditional independence is often used as a testable implication of causal models of random variables. In addition, equality constraints have been proposed to distinguish between data-generating mechanisms. We show that one can also find equality constraints in linear Hawkes processes, extending this theory to a class of continuous-time stochastic processes. This is done by proving that Hawkes process models in a certain sense satisfy the equality constraints of linear structural equation models. These results allow more refined constraint-based structure learning in this class of processes. Arguing for the existence of equality constraints leads us to new identification results for Hawkes processes. We also describe a causal interpretation of the linear Hawkes process which is closely related to its so-called cluster representation.

Keywords: equality constraints, causal identification, linear Hawkes processes, structure learning

1. Introduction

In causal inference, the question of what can be learned about the underlying structure from observational data is a classical one (Spirtes et al., 2000; Pearl, 2009). While most work studies models of random variables and without an explicit notion of time, this question has also been studied for stochastic process models. For a multivariate stochastic process, $X_t = (X_t^1, X_t^2, \ldots, X_t^n)^T$, this is often formalized by assuming that the dynamical evolution of each coordinate process, X_t^i, depends directly only on a subset of the other coordinate processes which can be represented by a directed graph which is the *structure* that methods aim to learn. As examples of this, Eichler et al. (2017); Xu et al. (2016) learn graphs describing the structure of multivariate linear Hawkes processes from dynamical observation of the process whereas Achab et al. (2017) use integrated cumulants of linear Hawkes processes for structure learning. These approaches assume that we have full observation in the sense that every coordinate process of the system is observed. Meek (2014); Mogensen et al. (2018); Mogensen (2020) consider structure learning based on tests of so-called local independence and provide methods that can recover structural information based on a marginal distribution only.

In this paper, we show that one can use equality constraints in linear Hawkes processes to distinguish between graphical structures that are indistinguishable using tests of local independence only which enables more refined structure learning algorithms. This development parallels that of structural causal models without an explicit notion of time in which equality constraints have been used for structure learning in addition to conditional independence constraints (Robins, 1986; Verma and Pearl, 1991; Robins, 1999; Tian and Pearl, 2002; Shpitser et al., 2014; Richardson et al., 2017).

Some results are not proven in the main text. Their proofs can be found in Appendix B.

Figure 1: Left: Example of observation from 5-dimensional Hawkes process, $V = \{1, 2, 3, 4, 5\}$. Colors (and vertical placement) indicate coordinate process. For each event (dot) time of occurrence is known as well as which coordinate process the event occurs in. Right: intensity processes, $\lambda_t^\alpha, \alpha \in V$, corresponding to event histories from the left plot (for simplicity only displayed events count in the intensities). The intensities are stochastic processes and from the stochastic integral in Definition 1 one can see that an α-event increases the β-intensity if $\phi_{\beta\alpha} \neq 0$.

2. Hawkes Processes

Hawkes processes have received much attention in machine learning in recent years as a tractable model of mutually exciting streams of events (Zhou et al., 2013; Luo et al., 2015; Etesami et al., 2016; Tan et al., 2018; Xu et al., 2018; Trouleau et al., 2019). The following is a brief introduction to this model class, see Laub et al. (2015), Liniger (2009), and Daley and Vere-Jones (2003) for additional background on Hawkes processes and point processes in general.

We consider a filtered probability space $(\Omega, \mathcal{F}, (\mathcal{F}_t), P)$ and an n-dimensional point process, $X_t = (X_t^1, X_t^2, \ldots, X_t^n)^T$, and let $V = \{1, 2 \ldots, n\}$. $\mathcal{F}_t$ is a filtration, that is, a nondecreasing family of σ-algebras which for each time point, $t \in \mathbb{R}$, represent the available information at that time. For each $\alpha \in V$, we say that X^α is a *coordinate process*. Each coordinate process is described by a series of *events*, or *points*, that occur at time points indexed by $\mathbb{R}$. For $\alpha \in V$, this is formalized by random variables, $\{T_i^\alpha\}_{i \in \mathbb{Z}}$, such that $T_i^\alpha < T_{i+1}^\alpha$ almost surely. For $s < t$, we let the $N_t^\alpha - N_s^\alpha$ denote the number of points (events) in the α-process in the interval $(s, t]$, that is, $N_t^\alpha - N_s^\alpha = \sum_i \mathbb{1}_{s < T_i^\alpha \leq t}$. We assume that no two events occur simultaneously. For $\alpha \in V$ the *intensity process*, λ_t^α, satisfies

$$\lambda_t^\alpha = \lim_{h \downarrow 0} \frac{1}{h} P(N_{t+h}^\alpha - N_t^\alpha = 1 \mid \mathcal{F}_t).$$

The intensity, λ_t^α, therefore describes how likely an α-event is in the immediate future $(t, t + h]$, for small h, given the past until time t.

Definition 1 (Linear Hawkes process) *We say that a point process, X, is a* linear Hawkes process *if for all $\beta \in V$ and $t \in \mathbb{R}$ the intensity is of the form*

$$\lambda_t^\beta = \mu_\beta + \sum_{\alpha \in V} \int_{-\infty}^t \phi_{\beta\alpha}(t - s) dN_s^\alpha$$

for a constant $\mu_\beta \geq 0$ and nonnegative functions $\phi_{\beta\alpha}$ for $\alpha \in V$.

We will often write *Hawkes process* instead of *linear Hawkes process*. We assume throughout for all α and β that $\phi_{\beta\alpha}$ is a continuous function on $(0, \infty)$ and zero outside this interval. We define the $n \times n$ matrix Φ such that $\Phi_{\beta\alpha} = \int_{-\infty}^{\infty} \phi_{\beta\alpha}(s)\mathrm{d}s$. For a square matrix, A, its spectral radius, $\rho(A)$, is the largest absolute value of its eigenvalues, $\lambda_1, \ldots, \lambda_m$, i.e., $\rho(A) = \max_{i=1,\ldots,m}\{|\lambda_i|\}$. If $\rho(A) < 1$ then we say that A is *Schur stable*. We assume that Φ is Schur stable and therefore we can assume the Hawkes process to have stationary increments and λ_t to be a stationary process (Jovanović et al., 2015; Bacry and Muzy, 2016). We define the $n \times n$ matrix $R = (I_n - \Phi)^{-1}$ where I_n is the $n \times n$ identity matrix. R is well-defined due to the assumption on the spectral radius of Φ and furthermore $R = \sum_{i=0}^{\infty} \Phi^i$. For a matrix, M, with rows indexed by I and columns indexed by J, we let $M_{\bar{I}\bar{J}}$ denote the submatrix indexed by $\bar{I} \subseteq I$ and $\bar{J} \subseteq J$. We let dN_t^{α} denote $N_{t+dt}^{\alpha} - N_t^{\alpha}$ where dt is the differential of t. Following Hawkes (1971), for each $\alpha \in V$ we define

$$\Lambda^{\alpha} = E(dN_t^{\alpha})/dt = E(\lambda_t^{\alpha})$$

which is independent of t due to stationarity. We let Λ denote the $n \times n$ diagonal matrix such that $\Lambda_{\alpha\alpha} = \Lambda^{\alpha}$ and we assume Λ to be positive definite.

Integrated Covariance   We define a probabilistic concept which is seen to be a cumulative measure of covariance between two coordinate processes. Let C be an $n \times n$ matrix such that for $\alpha, \beta \in V$,

$$C_{\beta\alpha}dt = \int_{-\infty}^{\infty} E(dN_t^{\beta}dN_{t+\tau}^{\alpha}) - E(dN_t^{\beta})E(dN_{t+\tau}^{\alpha})d\tau. \tag{1}$$

The entries of C contain the *integrated covariances* between pairs of coordinate processes. The following matrix equation holds (Jovanović et al., 2015; Achab et al., 2017),

$$C = R\Lambda R^T = (I_n - \Phi)^{-1}\Lambda(I_n - \Phi)^{-T}, \tag{2}$$

where $-T$ denotes transposition and inversion of a matrix.

Cluster Representation   In Definition 1, Hawkes processes are introduced by specifying their intensity processes. One can also define them using a so-called *cluster representation* (Jovanović et al., 2015). This representation lends itself to a straightforward causal interpretation as we will see.

The following describes the cluster representation which can be thought of as a simulation algorithm and it defines a Hawkes process given a set of constants $\{\mu_{\alpha}\}_{\alpha \in V}$ and a set of nonnegative functions $\{\phi_{\beta\alpha}\}_{\alpha,\beta \in V}$. For each $\alpha \in V$, *generation 0-events* are generated from a homogeneous Poisson process with rate μ_{α}. Each generation 0-event creates a *Hawkes tree*, that is, a cluster of events (the clusters are independent). The events in each cluster are generated recursively. Each event of type α at time s in a cluster have *child* events of type β according to an inhomogeneous Poisson process with rate $\phi_{\beta\alpha}(t - s)$ for $t > s$, and these processes are all independent. This is illustrated in Figure 2 in which four generation 0-events are shown, each of them starting a cluster. Parent-child relations are shown using line segments. Note that $\phi_{\beta\alpha} = 0$ means that there will be no β-children from an α-event, though there could be β-descendants further down from the α-event, e.g., if there exists γ such that $\phi_{\gamma\alpha} \neq 0$ and $\phi_{\beta\gamma} \neq 0$. The observed Hawkes process is then the

Figure 2: Same stream of events as in Figure 1 (left) showing the (unobserved) clusters of the events. Numbers indicate generation of the event and line segments indicate ancestry. For instance, the 1-2-3-4 cluster was produced from a generation 0-event of the 1-type which produced an event of the 2-type and so forth.

superposition of the events in all clusters. One should note that when observing a Hawkes process, we only have access to a set of event times and types (α_i, t_i) and not to the cluster structure, that is, the line segments in Figure 2 are unobserved.

The parameters in Φ and R have straightforward interpretations. From the cluster process definition of the Hawkes process, it follows that $\Phi_{\beta\alpha}$ is the expected number of child events of type β from an α-event. More generally, for $i = 0, 1, 2, \ldots$, $(\Phi^i)_{\beta\alpha}$ is the expected number of β-descendants in the i'th generation of a Hawkes tree rooted at an α-event. This implies that $R_{\beta\alpha}$ is the expected total number of β-descendants on such a tree (Jovanović et al., 2015).

2.1. Causal Interpretation

In this section, we will define what we mean by a *causal* Hawkes process. Mogensen (2020) defined a causal Hawkes process by requiring that the set of functions $\{\phi_{\beta\alpha}\}$ also describes how events enter the intensities in intervened systems where the trajectories of some coordinate processes are determined exogenously. We will instead use the cluster representation to give a causal interpretation based on a type of *injection intervention*. That is, the observational state of the system is given by the Hawkes process as defined above. An interventional state is defined by a set of deterministic *injections*, $\{(\alpha_i, t_i)\}_{i=1,\ldots,M}$, such that each injection is an element in $V \times \mathbb{R}$. An injection is a root in a Hawkes tree and the interventional processes are given as the superposition of all intrinsic and interventional trees. We then say that a linear Hawkes process is *causal* if a Hawkes tree rooted at an injection event has the same distribution as an intrinsic, or non-injection, Hawkes tree for all coordinate processes $\alpha \in V$, and all trees (interventional and intrinsic) are independent. In other words, the interventional events propagate in the same way through the system as the intrinsic events. This causal assumption is seen to correspond closely to the assumption of *autonomy* or *modularity* which is often used to define causal models (Pearl, 2009; Peters et al., 2017).

We note that this type of interventions is a simplistic version of interventions that are undertaken in some fields of science, e.g., in neuroscience where experimenters may intervene on neurons by means of extracellular electrical stimulation. This, however, is not expected to break the intrinsic dynamics of the nervous system nor sever causal mechanisms (Meffin et al., 2012; Komarov et al., 2019). It is also similar to how interventions are often modelled in linear control theory in which they are external forces (possibly dependent on the past of the process) that change the dynamics of the system without necessarily breaking any connections (Åström and Murray, 2008; Zabczyk, 2020).

Definition 2 (Causal effects) *We will say that $\Phi_{\beta\alpha}$ is the* direct (causal) effect *from α on β and that $R_{\beta\alpha}$ is the* total (causal) effect *from α on β. We will say that $\phi_{\beta\alpha}$ is the (causal) link function from α to β.*

The interpretation of the entries in Φ and R and the causal assumption justify the above definitions. If we inject an interventional event of type α at time $s \in \mathbb{R}$, this event on average has $\Phi_{\beta\alpha}$ events of type β as first-generation (or *direct*) descendants (when considering all $t > s$). The Hawkes tree rooted at the interventional α-event on average has a total of $R_{\beta\alpha}$ events of type β, counting the injectional event if $\alpha = \beta$.

Graphical representation A *graph* is a pair (V, E) where V is a finite set of nodes and E is a set of edges, each between a pair of (not necessarily distinct) nodes. We will use *directed mixed graphs* (DMGs), i.e., graphs such that every edge is either *directed*, $\rightarrow$, or *bidirected*, $\leftrightarrow$, and such that between any pair of nodes, α and β, there is a subset of the edges $\{\alpha \rightarrow \beta, \alpha \leftarrow \beta, \alpha \leftrightarrow \beta\}$. We use $\sim$ to denote a generic edge of either type. If an edge $\alpha \sim \beta$ is in a graph $\mathcal{G}$, we write $\alpha \sim_{\mathcal{G}} \beta$. If the edge $\alpha \rightarrow \beta$ is in the graph $\mathcal{G}$, we say that α is a *parent* of β, and we let $\mathrm{pa}_{\mathcal{G}}(\beta)$ denote the set of parents of β in the graph $\mathcal{G}$.

Given a causal Hawkes process with coordinate processes indexed by $V = \{1, 2, \ldots, n\}$, we construct its *causal graph*, $\mathcal{D} = (V, E)$, such that $\alpha \rightarrow_{\mathcal{D}} \beta$ if and only if $\phi_{\beta\alpha} \neq 0$. Using the continuity assumption of $\phi_{\beta\alpha}$, we see that if $\alpha \rightarrow \beta$ is in the causal graph, then there exists a nontrivial interval such that $\phi_{\beta\alpha} > 0$ on this interval.

A *walk* between α and β is an ordered, alternating sequence of nodes, γ_i, and edges, $\sim_j$,

$$\alpha = \gamma_0 \sim_1 \gamma_1 \sim_2 \ldots \gamma_{m-1} \sim_m \gamma_m = \beta$$

such that for each $i = 1, \ldots, m$ the edge $\sim_i$ is between γ_{i-1} and γ_i. We say that α and β are *endpoint nodes*. A *path* is a walk such that no node occurs more than once. We say that a path from α to β is *directed* if every edge is directed and points towards β, and a *directed cycle* is a directed path from α to β composed with the edge $\beta \rightarrow \alpha$. We say that a DMG is *acyclic* if it contains no directed cycles.

3. Representation as a Linear SEM

The relation between the observed quantity, C, and the parameters, (Φ, Λ), in Equation (2) is seen to be similar to the relation between the parameters and the covariance matrix of a *linear structural equation model* (linear SEM) and we will leverage this fact. There is a number of papers with identification results in this model class (Brito and Pearl, 2002; Tian, 2007, 2009; Foygel et al., 2012; Chen, 2016; Drton and Weihs, 2016; Weihs et al., 2018), though some results only apply to *acyclic* linear SEMs. Bollen (1989) provides an introduction to the model class. We give a short introduction here to describe the connection to Hawkes processes. Let $X = (X_1, \ldots, X_n)^T$ be a vector of random variables and $\varepsilon = (\varepsilon_1, \ldots, \varepsilon_n)^T$ be a vector of zero-mean noise terms. Let

$$X = BX + \varepsilon \tag{3}$$

such that the matrix B has zeros on the diagonal. We let Σ_ε denote the covariance of ε. The matrix $I_n - B$ is invertible if and only if B does not have 1 as an eigenvalue and in this case the covariance of X is

$$C_X = (I_n - B)^{-1}\Sigma_\varepsilon (I_n - B)^{-T}. \tag{4}$$

Often one will assume that some entries of B and Σ_ε are zero which can be encoded by a DMG (also known as a *mixed graph*) such that if $\alpha \to \beta$ is not in the DMG, then $B_{\beta\alpha} = 0$, and if $\alpha \leftrightarrow \beta$ ($\alpha \neq \beta$) is not in the DMG, then $(\Sigma_\varepsilon)_{\beta\alpha} = (\Sigma_\varepsilon)_{\alpha\beta} = 0$. The error terms are often assumed to be Gaussian. We note that acyclicity is sometimes assumed, i.e., that the DMG corresponding to the model is acyclic, though we are not making this assumption.

We should note that in the linear SEM literature, authors often parametrize the model in Equation 3 using a tranposition of B, $X = B^T X + \varepsilon$. We use this alternative notation as it aligns better with common notation in the Hawkes process literature.

One immediately sees the similarity between Equations (2) and (4) in that the observable left-hand sides of the equations impose similar restrictions on the model parameters, and we use this similarity to translate results to the Hawkes models. There are two questions that should be addressed. First, the diagonal of B in the linear SEM is often assumed to be zero, and we will handle this by re-writing Equation (2) slightly (Section 3.2). Second, Λ is a diagonal matrix in Equation (2) whereas there is no such restriction a priori on Σ_ε and we will see that this discrepancy corresponds to marginalization of the systems (Section 3.1).

3.1. Marginalization

If the Hawkes process is partially observed, only a proper submatrix of C in Equation (2) is available, and the coordinate processes, V, are partitioned into a set of observed processes, O, and a set of unobserved processes, U, such that $V = O \,\dot\cup\, U$. We use a *latent projection* to represent the graphical structure of the partially observed system.

Definition 3 (Latent projection, Verma and Pearl (1991); Richardson et al. (2017)) *Let $\mathcal{G} = (V, E)$ be a DMG. Its latent projection on nodes O is the DMG $\bar{\mathcal{G}} = (O, F)$ such that for $\alpha, \beta \in O$ we have $\alpha \to \beta$ in $\bar{\mathcal{G}}$ if there is a directed walk from α to β such that every non-endpoint node is in $V \setminus O$. We have $\alpha \leftrightarrow \beta$ in $\bar{\mathcal{G}}$ if there is a walk between α and β with heads at both α and β such that every non-endpoint node is a noncollider and is in $V \setminus O$.*

Note that for a walk in the above definition the set of non-endpoint nodes can be empty and therefore if $\alpha \to \beta$ is in $\mathcal{G}$, then it is also in its latent projection whenever $\alpha, \beta \in O$. We also note that the definition uses walks to allow for cycles (Mogensen and Hansen, 2020). In DAG-based models, this is not needed and the definition is often stated using paths (Richardson et al., 2017).

Using a DMG, $\mathcal{G} = (V, E)$, we will describe a parametrization of the observed integrated covariances. We let $\mathbb{M}_+^s(\mathcal{G})$ denote the set of $(n \times n)$ Schur stable matrices with nonnegative entries such that for $\alpha, \beta \in V = \{1, 2, \ldots, n\}$ and $M \in \mathbb{M}_+^s(\mathcal{G})$

$$\alpha \nrightarrow_{\mathcal{G}} \beta \Rightarrow M_{\beta\alpha} = 0.$$

We let $\mathbb{M}_{+,0}^s(\mathcal{G}) \subseteq \mathbb{M}_+^s(\mathcal{G})$ denote the subset of such matrices in which all diagonal elements are zero. We let $\mathbb{M}_+^{pd}(\mathcal{G})$ denote the set of symmetric, positive definite matrices with nonnegative entries such that for $\alpha, \beta \in V$ and $M \in \mathbb{M}_+^{pd}(\mathcal{G})$

$$\alpha \nleftrightarrow_{\mathcal{G}} \beta \Rightarrow M_{\beta\alpha} = M_{\alpha\beta} = 0.$$

Theorem 4 *Let X be a multivariate Hawkes process with stationary increments such that O are the observed coordinate processes and U are the unobserved coordinate processes and let $\mathcal{D} = (O \,\dot{\cup}\, U, E)$ be its causal graph. Then the observed integrated covariance can be written as*

$$\Sigma = (I - \Phi)^{-1}\Theta(I - \Phi)^{-T} \tag{5}$$

such that $(\Phi, \Theta) \in \mathbb{M}^s_+(\mathcal{G}) \times \mathbb{M}^{pd}_+(\mathcal{G})$ when $\mathcal{G}$ is the latent projection of $\mathcal{D}$ on O.

Proof In this proof, let (Φ, Λ) denote the parameters in the full system, and let Σ denote the full integrated covariance matrix. Let $\bar{\Sigma} = \Sigma_{OO}$ be the observed submatrix of Σ. We have that $I - \bar{\Phi}$ is invertible and

$$\bar{\Sigma} = (I - \bar{\Phi})^{-1}\Theta(I - \bar{\Phi})^{-T} \tag{6}$$

where

$$\bar{\Phi} = \Phi_{OO} + \Phi_{OU}(I_l - \Phi_{UU})^{-1}\Phi_{UO}$$
$$\Theta = (I_k - \bar{\Phi})[(I_n - \Phi)^{-1}\Lambda(I - \Phi)^{-T}]_{OO}(I_k - \bar{\Phi})^T,$$

see (Hyttinen et al., 2012). Using Schur complements, we can write (see Appendix A)

$$\Theta = \Lambda_{OO} + \Phi_{OU}((I_n - \Phi)^{-1})_{UU}\Lambda_{UU}((I_n - \Phi)^{-T})_{UU}\Phi^T_{OU}.$$

Θ is clearly positive semidefinite, and positive definite if Λ is. Then entries of $\bar{\Phi}$ and Θ are nonnegative, and Lemma 5 shows that $\bar{\Phi}$ is Schur stable. If α is not a parent of β in the latent projection then there is no directed path from α to β in $\mathcal{D}$ and it follows that $\bar{\bar{\Phi}}_{\beta\alpha} = 0$ using that $(I - A)^{-1} = \sum_{i=0}^{\infty} A^i$ when $\rho(A) < 1$. Similarly, we see that $\alpha \not\leftrightarrow_{\mathcal{G}} \beta$ implies that $\Omega_{\alpha\beta} = \Omega_{\beta\alpha} = 0$. $\blacksquare$

3.1.1. STABILITY UNDER MARGINALIZATION

We use the following lemma to show that if the full system is Schur stable, then a partially observed system is also Schur stable. This result does not hold without the assumption of nonnegative entries in the directed matrix (see Hyttinen et al. (2012) for an example which illustrates this).

Lemma 5 *Let A be a square matrix with nonnegative entries such that $\rho(A) < 1$, and let $O \,\dot{\cup}\, U$ be a partition of its rows and columns. The matrix $B = A_{OO} + A_{OU}(I - A_{UU})^{-1}A_{UO}$ has nonnegative entries and $\rho(B) < 1$.*

Proof The matrix A_{UU} is a principal submatrix of A and therefore $\rho(A_{UU}) < \rho(A) < 1$ (Horn and Johnson, 1985, Corollary 8.1.20). Therefore $(I - A_{UU})^{-1} = \sum_{i=0}^{\infty}(A_{UU})^i$ and it follows that every entry of B is nonnegative.

It holds that $I - B$ is the Schur complement of $I - A_{UU}$ in $I - A$, that is, $I - B = (I - A)/(I - A_{UU})$. The matrix $I - A$ is a nonsingular M-matrix and therefore so is $I - B$ (Berman and Plemmons, 1979, Exercise 5.8). Then every real eigenvalue of $I - B$ is positive (Berman and Plemmons, 1979, Theorem 2.3, D_{16}). Since B is nonnegative, $\rho(B)$ is an eigenvalue of B (Horn and Johnson, 1985, Theorem 8.3.1). It follows that $1 - \rho(B)$ is an eigenvalue of $I - B$ and therefore $1 - \rho(B) > 0$.

∎

The above shows that the observed integrated covariance of a linear Hawkes process can be parametrized such that the sparsity of matrices Φ and Ω is described by the latent projection, $\mathcal{G}$, of its causal graph, more precisely, $(\Phi, \Omega) \in \mathbb{M}_+^s(\mathcal{G}) \times \mathbb{M}_+^{pd}(\mathcal{G})$. On the other hand, we can ask if there for any pair, (Φ_0, Ω_0), in this set exists a linear Hawkes process such that its observed integrated covariance, Σ, is

$$\Sigma = (I - \Phi_0)^{-1}\Omega_0(I - \Phi_0)^{-T}$$

and such that the latent projection of the causal graph of the linear Hawkes process is equal to $\mathcal{G}$. This is in fact the case. For each bidirected edge one can add an additional node. Now consider the larger matrix that embeds Φ in a zero matrix. This matrix has eigenvalues zeros and the eigenvalues of Φ. Using continuity of eigenvalues we add ϵ along the directed edges from auxiliary nodes to nodes connected by bidirected edges in the original graph. Choosing a small ϵ this new, larger matrix will be Schur stable.

3.2. Normalized Effects

When Φ is a matrix of direct effects of a Hawkes process (or marginalized effects, $\bar{\Phi}$), the diagonal elements are not necessarily zero. On the other hand, in a linear SEM the diagonal of B (Equation (3)) is commonly assumed to be zero. We introduce the matrix Γ by scaling each column of $\bar{\Phi}$ by one minus its diagonal element and placing zeros on the diagonal,

$$\Gamma_{\beta\alpha} = \bar{\Phi}_{\beta\alpha}/(1 - \bar{\Phi}_{\beta\beta}) \text{ if } \beta \neq \alpha, \text{ and } \Gamma_{\beta\alpha} = 0 \text{ if } \beta = \alpha,$$

and $I - \Gamma$ is invertible (Hyttinen et al., 2012). Hyttinen et al. (2012) study linear SEMs and use the same approach to derive so-called canonical linear cyclic models from models with self-loops. We think of the above operation as a type of normalization, and if this is done in the full model (no marginalization), we will say that the normalized parameters, that is, $\Phi_{\beta\alpha}/(1 - \Phi_{\beta\beta})$, are *normalized direct (causal) effects*. It is seen that a normalized direct effect, $\Phi_{\beta\alpha}/(1 - \Phi_{\beta\beta})$, $\alpha \neq \beta$, is the average number of direct descendants of type β from an α-event, counting every direct self-triggered event in the β-process. That is, on the Hawkes tree rooted at α, $\Phi_{\beta\alpha}/(1 - \Phi_{\beta\beta})$ is the expected number of β-events on subtrees of the form $\alpha - \beta - \beta - \ldots - \beta$ for any number of β-events.

If we consider a Hawkes process and its causal graph, $\mathcal{D} = (V, E)$, we see that the latent projection of $\mathcal{D}$, $\mathcal{G} = (O, F)$, $O \subseteq V$, represents the observed covariance, $\bar{C} = C_{OO}$ in the following way. If $\alpha \rightarrow \beta$ is not in $\mathcal{G}$, then $\bar{\Phi}_{\beta\alpha} = 0$. If $\alpha \leftrightarrow \beta$ is not in $\mathcal{G}$, then $\Theta_{\beta\alpha} = \Theta_{\alpha\beta} = 0$. Omitting self-loops, this corresponds to the graphical representation of a linear SEM. This means

that the observable integrated covariances after rewriting are parametrized exactly like a linear SEM and we will use this to obtain identification results (Section 4) and equality constraints (Section 5).

While this casts the Hawkes integrated covariance as a set of equations describing the covariance matrix of a linear SEM, the Hawkes parameters are required to be nonnegative.

Corollary 6 *Let N be a multivariate Hawkes process with stationary increments such that O are the observed coordinate processes and U are the unobserved coordinate processes and let $\mathcal{D}$ be its causal graph. Then the observed integrated covariance can be written as*

$$\Sigma = (I - \Gamma)^{-1} \Omega (I - \Gamma)^{-T}$$

such that $(\Gamma, \Omega) \in \mathbb{M}_{0,+}^s(\mathcal{G}) \times \mathbb{M}_+^{pd}(\mathcal{G})$ when $\mathcal{G}$ is the latent projection of $\mathcal{D}$ on O.

Proof Consider the representation from Theorem 4. Then we have

$$\Sigma = (I - \Phi)^{-1} \Theta (I - \Phi)^{-T}.$$

Let D denote a diagonal matrix such that the α'th diagonal element equals $1 - \bar{\Phi}_{\alpha\alpha}$ and Γ is as defined above. Then

$$
\begin{aligned}
\Sigma &= (D(I - \Gamma))^{-1} \Theta (D(I - \Gamma))^{-T} \\
&= (I - \Gamma)^{-1} D^{-1} \Theta D^{-1} (I - \Gamma)^{-T}.
\end{aligned}
\tag{7}
$$

The claims all follow from Theorem 4. We should just show that Γ is a Schur stable matrix (Proposition 7). ∎

Proposition 7 *Let Φ be a nonnegative matrix such that $\rho(\Phi) < 1$. Then the normalized matrix, Γ, also satisfies $\rho(\Gamma) < 1$.*

4. Identification

Using constraints on the causal graph, we will discuss the identification of the entries of the matrices Φ and Γ. To give a formal definition of identification, consider two stationary Hawkes processes, $M_1 \in \mathbb{M}$ and $M_2 \in \mathbb{M}$, with coordinate processes V_1 and V_2, respectively, for a model $\mathbb{M}$. Let $\tilde{P}_1$ and $\tilde{P}_2$ be their marginal distributions over a common observed set of coordinate processes, O, let f be a function of such a marginal distribution, and let $\alpha, \beta \in O$. Finally, let $\Pi_{\beta\alpha}^i$ be a parameter ($\Pi_{\beta\alpha} = \Phi_{\beta\alpha}$ or $\Pi_{\beta\alpha} = \Gamma_{\beta\alpha,}$). We will say that the parameter $\Pi_{\beta\alpha}$ *is identified from* $f(\tilde{P})$ in the model $\mathbb{M}$ if $f(\tilde{P}_1) = f(\tilde{P}_2)$ implies $\Pi_{\beta\alpha}^1 = \Pi_{\beta\alpha}^2$. Note that while identification is often only discussed from the observed distribution in its entirety, we use f to distinguish simpler statistics that can be sufficient for identification. As we will see, under graphical constraints some parameters are identified from the integrated covariance matrix in which case the inferential problem is reduced to a finite-dimensional one.

Figure 3: **A**: Directed graph such that each node represents a coordinate process and an edge $\alpha \to \beta$ implies that $\Phi_{\beta\alpha} \neq 0$. **B**: Latent projection of the graph **A** on nodes $O = \{1, 2, 3, 4\}$. **C**: Graphs **B** and **C** are Markov equivalent (entail the same μ-separations), yet are distinguishable using an equality constraint on the observable matrix $\bar{C} = C_{OO}$. Loops (self-edges) are omitted from graphs **A**, **B**, and **C**.

4.1. Identification as a Linear SEM

There are several papers describing graphical conditions to ensure identifiability of parameters of a linear structural equation model. For these results, one assumes a certain graphical structure represented by a DMG which encodes sparsity in the matrices B and Σ_ε in Equation (4). Identification of a parameter means that for a fixed graphical structure and an observed covariance, C_X, this parameter is uniquely determined. The idea is now that one can use any available identification result from the (cyclic) linear structural equation toolbox to show identification of the entries in Γ such as those provided by, e.g., Foygel et al. (2012); Chen (2016); Weihs et al. (2018). This follows just from noting that if we use the latent projection of the Hawkes causal graph to encode sparsity, the problem in Corollary 6 is the same as the linear SEM problem, though with a smaller parameter space as Γ and Ω are required to be nonnegative and Γ is required to be Schur stable. While any identification result from the linear SEM problem therefore applies to the problem in Corollary 6, one can actually obtain stronger results using that the parameter space is smaller. Section 4.2 shows this by an example. We also note that one typically has to distinguish between *global* and *generic* identification of parameters.

Example 8

Consider the graph in Figure 3A and assume that this is the causal graph of a Hawkes process on nodes $V = \{1, 2, 3, 4, 5\}$ such that coordinate process 5 is unobserved. Let Γ be the 4×4 matrix of direct effects. If we apply the identification criterion of Weihs et al. (2018), we find that every $\Gamma_{\beta\alpha}$ such that $\alpha, \beta \in O$ is identified from Σ (note that $\Gamma_{\beta\alpha} = \Phi_{\beta\alpha}/(1 - \Phi_{\beta\beta})$ for $\alpha, \beta \in O$ in this example). It follows that Φ_{11}, Φ_{23}, and Φ_{33} are identified from the integrated mean and covariance.

On the other hand, one can show by direct computation that $\Phi_{21}, \Phi_{22}, \Phi_{24}, \Phi_{43}$, and $\Phi_{4,4}$ are not identified. If we consider Φ_{21}, we can understand the lack of identication from the fact that Φ_{22} cannot be identified. Intuitively, this is explained by its unobserved parent process which implies that we cannot, using only the integrated covariance, separate the contributions from coordinate processes 2 and 5 to the observed covariance.

4.2. Identification of cycles

The n-cycle, $\mathcal{D}_n$, is the directed graph on nodes $\alpha_1, \ldots, \alpha_n$ such that $\alpha_i \to \alpha_{i+1}$ for all $i = 1, \ldots, n-1$ and $\alpha_n \to \alpha_1$. When $n \geq 3$, Drton et al. (2011) show that Equation 4 has at most two

solutions when the graph is an n-cycle and that generically it has two solutions. We show in the following that at most one of them is in $\mathbb{M}_+^s(\mathcal{G}) \times \mathbb{M}_{+,0}^{pd}(\mathcal{G})$, thus, is a solution to the normalized Hawkes problem.

Proposition 9 *Let $\mathcal{G}$ be an n-cycle, $n \geq 3$. There is a unique solution to*

$$\Sigma = (I - \Gamma)^{-1} \Theta (I - \Gamma)^{-T}$$

such that $(\Gamma, \Omega) \in \mathbb{M}_{0,+}^s(\mathcal{G}) \times \mathbb{M}_+^{pd}(\mathcal{G})$.

5. Equality Constraints

It is well-known that partially observed DAG-models satisfy constraints that are not described by conditional independence some of which are known as *equality constraints*. Richardson et al. (2017) treat the general (acyclic) case while Chen et al. (2014); Chen (2016) consider equality constraints in linear SEMs. In DAG-models equality constraints are testable implications that may bring strictly more information about the underlying structure than using conditional independence alone. We will see that the analogous statement holds for the Hawkes process models, though the relevant notion of independence to compare with is that of *local independence*. Local independence is an asymmetric notion of independence of the coordinate processes of a multivariate stochastic process (Schweder, 1970; Aalen, 1987; Didelez, 2000, 2008). The following definition is specific to point processes, however, it can be extended to other classes of stochastic processes (Aalen, 1987; Didelez, 2006; Mogensen et al., 2018; Mogensen and Hansen, 2020).

Definition 10 (Local independence) *Let $X = (X^1, \ldots, X^n)^T$ be a multivariate point process with intensity processes λ and let $V = \{1, \ldots, n\}$. Let $A, B, C, D \subseteq V$. Let $\mathcal{F}_t^D$ denote the natural filtration of X^D, i.e., the filtration generated by the coordinate processes in D. We say that X^B is* locally independent *of X^A given X^C if for all $\beta \in B$ and for all $t \in \mathbb{R}$,*

$$E(\lambda_t^\beta \mid \mathcal{F}_t^{A \cup C})$$

is adapted to the filtration $\mathcal{F}_t^C$.

Local independence has been used for structure learning in stochastic process models (Meek, 2014; Mogensen et al., 2018; Thams, 2019; Mogensen, 2020), analogously to how conditional independence is used for constraint-based structure learning in classical models (Spirtes et al., 2000; Spirtes and Zhang, 2019). Structure learning based on local independence typically assumes *faithfulness*, that is, that X^B is locally independent of X^A given X^C if and only if B is μ-separated from A given C in $\mathcal{G}$, the DMG representing the model (μ-separation, or δ-separation, is a concept analogous to m-, or d-separation (Didelez, 2000, 2008; Mogensen and Hansen, 2020)). The *Markov equivalence class* of $\mathcal{G}$ is the set of DMGs that encode exactly the same set of μ-separations, hence local independencies. An obvious question is therefore whether there are constraints imposed by the structure that are not described by local independence, or μ-separation. This is indeed the case and such constraints can be used to distinguish between graphical structures that imply the same set of local independencies. In the next subsection, we show that one can use the integrated mean and covariance to find such constraints.

5.1. Using Integrated Covariance

From Equation 7 and the section on marginalization we see that every equality constraint in a linear structural equation model implied by the graphical structure imposes the same constraint on the parameters (Γ, Θ) and this allows us to find constraints in the Hawkes model. We will show by an example that these constraints may provide information about the underlying structure that is not contained in the Markov equivalence class.

Example 11 *Consider again the example illustrated by the graph in Figure 3B. If we think of the graph as representing a linear SEM, we can, e.g., use Theorem 1 and Lemma 1 of* Chen et al. (2014) *to see that the graphical structure imposes the constraint*

$$\Sigma_{14}/\Sigma_{13} = (\Sigma_{34} - \Gamma_{32}\Sigma_{24})/(\Sigma_{33} - \Gamma_{32}\Sigma_{23}),$$

*and therefore this algebraic constraint is also satisfied by the Hawkes model corresponding to the graph (note that $\Gamma_{32} = \Phi_{32}/(1 - \Phi_{33})$, and that the constraint corresponds to two ways of identifying Γ_{43}). The graphs **B** and **C** in Figure 3 (right) imply the same set of local independencies when using μ-separation. However, the above constraint is satisfied in **B** only and therefore this constraint allows us to discriminate between the two models.*

6. Conclusion

We obtained identification results and equality constraints for Hawkes process models through the similarity between the integrated Hawkes covariance and the observed covariance of a linear structural equation model. These equality constraints are useful for structure learning as they can supplement constraint-based learning based on tests of local independence. The approach in this paper finds equality constraints through a time-independent statistic which may be applicable in other classes of stochastic processes as well.

A Hawkes process is defined using the set of $\phi_{\beta\alpha}$-functions along with the μ_α-constants for α, β in the finite set V and a Hawkes process model is in that sense nonparametric. Constraints that arise from the integrated covariance offer also a dimension reduction as these constraints exist in a finite parameter space instead of a function space. This makes them more suitable to employ in data analysis as estimation of the link functions, $\{\phi_{\beta\alpha}\}$, is a challenging problem even in the case of full observation and partial observation only adds to the complexity of this task.

References

Odd O. Aalen. Dynamic modelling and causality. *Scandinavian Actuarial Journal*, 1987(3-4): 177–190, 1987.

Massil Achab, Emmanuel Bacry, Stéphane Gaïffas, Iacopo Mastromatteo, and Jean-François Muzy. Uncovering causality from multivariate Hawkes integrated cumulants. In *Proceedings of the 34th International Conference on Machine Learning (ICML)*, 2017.

Karl Johan Åström and Richard M. Murray. *Feedback systems: An introduction for scientists and engineers*. Princeton University Press, 2008.

Emmanuel Bacry and Jean-François Muzy. First- and second-order statistics characterization of Hawkes processes and non-parametric estimation. *IEEE Transactions on Information Theory*, 62 (4):2184–2202, 2016.

Abraham Berman and Robert J. Plemmons. *Nonnegative Matrices in the Mathematical Sciences*. Academic Press, 1979.

Kenneth A. Bollen. *Structural Equations with Latent Variables*. John Wiley & Sons, 1989.

Carlos Brito and Judea Pearl. Generalized instrumental variables. In *Proceedings of the 18th Conference on Uncertainty in Artificial Intelligence (UAI)*, 2002.

Bryant Chen. Identification and overidentification of linear structural equation models. In *Proceedings of the 30th Conference on Neural Information Processing Systems (NIPS)*, 2016.

Bryant Chen, Jin Tian, and Judea Pearl. Testable implications of linear structural equation models. In *Proceedings of the 28th AAAI Conference on Artificial Intelligence*, 2014.

Daryl J. Daley and David D. Vere-Jones. *An introduction to the theory of point processes*. New York: Springer, 2nd edition, 2003.

Vanessa Didelez. *Graphical Models for Event History Analysis based on Local Independence*. PhD thesis, Universität Dortmund, 2000.

Vanessa Didelez. Graphical models for composable finite Markov processes. *Scandinavian Journal of Statistics*, 34(1):169–185, 2006.

Vanessa Didelez. Graphical models for marked point processes based on local independence. *Journal of the Royal Statistical Society, Series B*, 70(1):245–264, 2008.

Mathias Drton and Luca Weihs. Generic identifiability of linear structural equation models by ancestor decomposition. *Scandinavian Journal of Statistics*, 43(4):1035–1045, 2016.

Mathias Drton, Rina Foygel, and Seth Sullivant. Global identifiability of linear structural equation models. *The Annals of Statistics*, 39(2):865–886, 2011.

Michael Eichler, Rainer Dahlhaus, and Johannes Dueck. Graphical modeling for multivariate Hawkes processes with nonparametric link functions. *Journal of Time Series Analysis*, 38:225–242, 2017.

Jalal Etesami, Negar Kiyavash, Kun Zhang, and Kushagra Singhal. Learning network of multivariate Hawkes processes: A time series approach. In *Proceedings of the 32nd Conference on Uncertainty in Artificial Intelligence (UAI)*, 2016.

Rina Foygel, Jan Draisma, and Mathias Drton. Half-trek criterion for generic identifiability of linear structural equation models. *The Annals of Statistics*, 40(3):1682–1713, 2012.

Alan G. Hawkes. Point spectra of some mutually exciting point processes. *Journal of the Royal Statistical Society, Series B*, 33(3):438–443, 1971.

Roger A. Horn and Charles R. Johnson. *Matrix Analysis*. Cambridge University Press, 1985.

Antti Hyttinen, Frederick Eberhardt, and Patrik O. Hoyer. Learning linear cyclic causal models with latent variables. *Journal of Machine Learning Research*, 13:3387–3439, 2012.

Stojan Jovanović, John Hertz, and Stefan Rotter. Cumulants of Hawkes point processes. *Physical Review E*, 91(4), 2015.

Maxim Komarov, Paola Malerba, Ryan Golden, Paul Nunez, Eric Halgren, and Maxim Bazhenov. Selective recruitment of cortical neurons by electrical stimulation. *PLoS Computational Biology*, 15(8), 2019.

Patrick J. Laub, Thomas Taimre, and Philip K. Pollett. Hawkes processes. 2015. URL https://arxiv.org/pdf/1507.02822.pdf.

Thomas Josef Liniger. *Multivariate Hawkes processes*. PhD thesis, ETH Zürich, 2009.

Dixin Luo, Hongteng Xu, Yi Zhen, Xia Ning, Hongyuan Zha, Xiaokang Yang, and Wenjun Zhang. Multi-task multi-dimensional Hawkes processes for modeling event sequences. In *Proceedings of the 24th International Joint Conference on Artificial Intelligence (IJCAI)*, 2015.

Christopher Meek. Toward learning graphical and causal process models. In *CI'14 Proceedings of the UAI 2014 Conference on Causal Inference: Learning and Prediction*, 2014.

Hamish Meffin, Bahman Tahayori, David B. Grayden, and Anthony N. Burkitt. Modeling extracellular electrical stimulation: I. Derivation and interpretation of neurite equations. *Journal of Neural Engineering*, 9(6), 2012.

Søren Wengel Mogensen. Causal screening for dynamical systems. In *Proceedings of the 36th Conference on Uncertainty in Artificial Intelligence (UAI), PMLR volume 124*, 2020.

Søren Wengel Mogensen and Niels Richard Hansen. Markov equivalence of marginalized local independence graphs. *The Annals of Statistics*, 48(1):539–559, 2020.

Søren Wengel Mogensen, Daniel Malinsky, and Niels Richard Hansen. Causal learning for partially observed stochastic dynamical systems. In *Proceedings of the 34th conference on Uncertainty in Artificial Intelligence (UAI)*, 2018.

Judea Pearl. *Causality*. Cambridge University Press, 2009.

Jonas Christopher Peters, Dominik Janzing, and Bernhard Schölkopf. *Elements of causal inference, foundations and learning algorithms*. MIT Press, 2017.

Thomas S. Richardson, Robin J. Evans, James M. Robins, and Ilya Shpitser. Nested markov properties for acyclic directed mixed graphs. 2017. URL https://arxiv.org/pdf/1701.06686.pdf.

James M. Robins. A new approach to causal inference in mortality studies with a sustained exposure period — application to control of the healthy worker survivor effect. *Mathematical Modeling*, 7 (9):1393–1512, 1986.

James M. Robins. Testing and estimation of direct effects by reparameterizing directed acyclic graphs with structural nested models. In C. Glymour and G. Cooper, editors, *Computation, Causation, and Discovery*, pages 349–405. Menlo Park, CA, Cambridge, MA: AAAI Press/The MIT Press, 1999.

Tore Schweder. Composable markov processes. *Journal of Applied Probability*, 7(2):400–410, 1970.

Ilya Shpitser, Robin J. Evans, Thomas S. Richardson, and James M. Robins. Introduction to nested Markov models. *Behaviormetrika*, 41(1):3–39, 2014.

Peter Spirtes and Kun Zhang. Search for causal models. In Marloes Maathuis, Mathias Drton, Steffen Lauritzen, and Martin Wainwright, editors, *Handbook of Graphical Models*, pages 439–462. CRC Press, 2019.

Peter Spirtes, Clark Glymour, and Richard Scheines. *Causation, Prediction, and Search*. MIT Press, 2000.

Xi Tan, Vinayak Rao, and Jennifer Neville. Nested CRP with Hawkes-Gaussian processes. In *Proceedings of the 21st International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2018.

Nikolaj Thams. Causal structure learning in multivariate point processes. Master's thesis, University of Copenhagen, 2019.

Jin Tian. A criterion for parameter identification in structural equation models. In *Proceedings of the 23rd Conference on Uncertainty in Artificial Intelligence (UAI)*, 2007.

Jin Tian. Parameter identification in a class of linear structural equation models. In *Proceedings of the 21st International Joint Conference on Artificial Intelligence (IJCAI)*, 2009.

Jin Tian and Judea Pearl. On the testable implications of causal models with hidden variables. In *Proceedings of the 18th Conference on Uncertainty in Artificial Intelligence (UAI)*, 2002.

William Trouleau, Jalal Etesami, Matthias Grossglauser, Negar Kiyavash, and Patrick Thiran. Learning Hawkes processes under synchronization noise. In *Proceedings of the 36th International Conference on Machine Learning (ICML)*, 2019.

Thomas Verma and Judea Pearl. Equivalence and synthesis of causal models. Technical Report R-150, University of California, Los Angeles, 1991.

Luca Weihs, Bill Robinson, Emilie Dufresne, Jennifer Kenkel, Kaie Kubjas, Reginald McGee II, Nhan Nguyen, Elina Robeva, and Mathias Drton. Determinantal generalizations of instrumental variables. *Journal of Causal Inference*, 6(1), 2018.

Hongteng Xu, Mehrdad Farajtabar, and Hongyuan Zha. Learning Granger causality for Hawkes processes. In *Proceedings of the 33rd International Conference on Machine Learning (ICML)*, 2016.

Hongteng Xu, Dixin Luo, Xu Chen, and Lawrence Carin. Benefits from superposed Hawkes processes. In *Proceedings of the 21st International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2018.

Jerzy Zabczyk. *Mathematical control theory: An introduction*. Birkhäuser, 2nd edition, 2020.

Ke Zhou, Hongyuan Zha, and Le Song. Learning social infectivity in sparse low-rank networks using multi-dimensional Hawkes processes. In *Proceedings of the 16th International Conference on Artificial Intelligence and Statistics (AISTATS)*, 2013.

Appendix A. Marginalization

In this section and for a matrix A, we define $A_{\bar{I}\bar{J}}^{-1} = (A^{-1})_{\bar{I}\bar{J}}$ and $A_{\bar{I}\bar{J}}^{-T} = (A^{-T})_{\bar{I}\bar{J}}$. Consider an invertible block matrix

$$M = \begin{pmatrix} A & B \\ C & D \end{pmatrix}.$$

We let $M/D = A - BD^{-1}C$ denote the *Schur complement* of block D in the matrix M. It holds that

$$M^{-1} = \begin{pmatrix} (M/D)^{-1} & -(M/D)^{-1}BD^{-1} \\ -D^{-1}C(M/D)^{-1} & D^{-1}(I + C(M/D)^{-1}BD^{-1}) \end{pmatrix}.$$

Therefore,

$$(I - \Phi)_{OO}^{-1} = ((I - \Phi)/(I - \Phi)_{UU})^{-1} = (I - \bar{\Phi})^{-1}$$

by definition of $\bar{\Phi}$. Furthermore,

$$(I - \Phi)_{OU}^{-1} = -(I - \bar{\Phi})^{-1}(I - \Phi)_{OU}(I - \Phi)((I - \Phi)_{UU})^{-1}$$

Let $I = I_n$ and $A_{\bar{I}} = A_{\bar{I}\bar{I}}$. From the above, it follows that

$$\begin{aligned}
\Theta &= (I_k - \bar{\Phi})[(I_n - \Phi)_O^{-1}\Lambda_O(I - \Phi)_O^{-T} + \\
&\quad (I - \Phi_{OU}^{-1})\Lambda_U(I - \Phi)_{UO}^{-T}](I_k - \bar{\Phi})^T \\
&= \Lambda_O + (I_k - \bar{\Phi})(I - \Phi)_{OU}^{-1}\Lambda_U((I - \Phi)_{UO}^{-T}(I_k - \bar{\Phi})^T \\
&= \Lambda_O + \\
&\quad (I - \Phi)_{OU}(I - \Phi)_U^{-1}\Lambda_U((I - \Phi)_U^{-T}(I - \Phi)_U^{-T}(I - \Phi)_{OU}^T \\
&= \Lambda_O + \Phi_{OU}(I - \Phi)_U^{-1}\Lambda_U((I - \Phi)^{-T})_U(I - \Phi)_U^{-T}\Phi_{OU}^T \\
&= \Lambda_O + \Phi_{OU}(I - \Phi)_U^{-1}\Lambda_U(I - \Phi)_U^{-T}\Phi_{OU}^T.
\end{aligned}$$

Appendix B. Proofs

Proof [Proposition 7] Assume to obtain a contradiction that $\rho(\Gamma) \geq 1$ and let $\lambda = \rho(\Gamma)$. The matrix Γ is nonnegative and therefore λ is an eigenvalue of Γ and a nonnegative eigenvector, $x \neq 0$, can be chosen (Horn and Johnson, 1985, Theorem 8.3.1), $\Gamma x = \lambda x$.

Let D_Φ denote the diagonal matrix such that $(D_\Phi)_{ii} = \Phi_{ii}$ for all i. From the definition of Γ, we have

$$\Phi = \Gamma - D_\Phi \Gamma + D_\Phi.$$

Multiplying with x and using $\Gamma x = \lambda x$, we obtain

$$\Phi x = \lambda x + (1 - \lambda) D_\Phi x$$

For all i, $(D_\Phi)_{ii} = \Phi_{ii} \leq \rho(\Phi) = 1$ (Horn and Johnson, 1985, Corollary 8.1.20). We have that $1 - \lambda \leq 0$ and that the entries of x are nonnegative and therefore

$$\Phi x \geq \lambda x + (1 - \lambda) x = x$$

where the inequality should be read entrywise. It holds that $x \neq 0$ and therefore $\Phi x \geq x$ implies that $\rho(\Phi) \geq 1$ which is a contradiction (Horn and Johnson, 1985, Theorem 8.3.2). ∎

Proof [Proposition 9] Let (G_0, Ω_0), (G_1, Ω_1) be the two solutions of a linear SEM corresponding to n-cycle (generically, they are distinct). We assume that the observed covariance is from a Schur stable system, without loss of generality, say $\rho(G_1) < 1$. Let δ_i^j denote the inverse of the i'th diagonal element of Ω_j and let λ_i^j denote the $(i, i+1)$-entry of G_j (modulo n). Drton et al. (2011) show that,

$$\delta_i^1 = \delta_i^0 + \prod_j \delta_j^0 (\prod_j (\lambda_j^0)^2 - 1)/\det(K_{-i}) \qquad \text{and}$$
$$\delta_i^0 = \delta_i^1 + \prod_j \delta_j^1 (\prod_j (\lambda_j^1)^2 - 1)/\det(K_{-i}),$$

where K is the inverse of the observed covariance, Σ, and K_{-i} denotes the submatrix of K with the i'th row and column deleted. Replacing δ_i^1 in the lower equation with the expression in the upper equation we obtain,

$$-(\prod_i \delta_i^1)(\prod_i (\lambda_i^1)^2 - 1) = (\prod_i \delta_i^0)(\prod_i (\lambda_i^0)^2 - 1). \tag{8}$$

Assume without loss of generality that $\rho(G_0) < 1$, then every eigenvalue of G_0 has an absolute value less than 1, and $\det(G_0) < 1$. Due to the structure of G_0 it holds that $\det(G_0) = \prod_i (\lambda_i^0)$ such that $\prod_i (\lambda_i^0) < 1$ and $\prod_i (\lambda_i^0)^2 < 1$. Every δ_i^0 is positive, and this means that the right-hand side of 8 is negative. Every δ_i^1 is positive and therefore we must have $(\prod_i (\lambda_i^1)^2 - 1) > 0$ and therefore $(\prod_i (\lambda_i^1)) > 1$. This means that $\det(G_1) > 1$ and therefore $\rho(G_1) \geq 1$. ∎