Undersampling is a Minimax Optimal Robustness Intervention in Nonparametric Classification

Abstract

While a broad range of techniques have been proposed to tackle distribution shift, the simple baseline of training on an *undersampled* dataset often achieves close to state-of-the-art-accuracy across several popular benchmarks. This is rather surprising, since undersampling algorithms discard excess majority group data. To understand this phenomenon, we ask if learning is fundamentally constrained by a lack of minority group samples. We prove that this is indeed the case in the setting of nonparametric binary classification. Our results show that in the worst case, an algorithm cannot outperform undersampling unless there is a high degree of overlap between the train and test distributions (which is unlikely to be the case in real-world datasets), or if the algorithm leverages additional structure about the distribution shift. In particular, in the case of label shift we show that there is always an undersampling algorithm that is minimax optimal. While in the case of group-covariate shift we show that there is an undersampling algorithm that is minimax optimal when the overlap between the group distributions is small. We also perform an experimental case study on a label shift dataset and find that in line with our theory the test accuracy of robust neural network classifiers is constrained by the number of minority samples.

1 Introduction

A key challenge facing the machine learning community is to design models that are robust to distribution shift. When there is a mismatch between the train and test distributions, current models are often brittle and perform poorly on rare examples [11, 2, 20, 10, 1]. In this paper, our focus is on group-structured distribution shifts. In the training set, we have many samples from a *majority* group and relatively few samples from the *minority* group, while during test time we are equally likely to get a sample from either group.

To tackle such distribution shifts, a naïve algorithm is one that first *undersamples* the training data by discarding excess majority group samples [14, 23] and then trains a model on this resulting dataset (see Figure 1 for an illustration of this algorithm). The samples that remain in this undersampled dataset constitute i.i.d. draws from the test distribution. Therefore, while a classifier trained on this pruned dataset cannot suffer biases due to distribution shift, this algorithm is clearly wasteful, as it discards training samples.

Figure 1: Example with linear models and linearly separable data. On the left we have the maximum margin classifier over the entire dataset, while on the right we have the maximum margin classifier over the undersampled dataset. The undersampled classifier is less biased and aligns more closely with the true boundary.

This perceived inefficiency of undersampling has led to the design of several algorithms to combat such distribution shift [6, 15, 18, 5, 17, 29, 13, 24]. In spite of this algorithmic progress, the simple baseline of training models on an undersampled dataset remains competitive. In the case of label shift, where one class label is overrepresented in the training data, this has been observed by Cui et al. [7], Cao et al. [5], and Yang and Xu [28]. While in the case of group-covariate shift, a study by Idrissi et al. [12] showed that the empirical effectiveness of these more complicated algorithms is limited.

For example, Idrissi et al. [12] showed that on the group-covariate shift CelebA dataset the worst-group accuracy of a ResNet-50 model on the undersampled CelebA dataset which *discards 97%* of the available training data is as good as methods that use all of available data such as importance-weighted ERM [19], Group-DRO [18] and Just-Train-Twice [16]. In Table 1, we report the performance of the undersampled classifier compared to the state-of-the-art-methods in the literature across several label shift and group-covariate shift datasets. We find that, although undersampling isn't always the optimal robustness algorithm, it is typically a very competitive baseline and within 1–4% the performance of the best method.

Table 1: Performance of undersampled classifier compared to the best classifier across several popular label shift and group-covariate shift datasets. When reporting worst-group accuracy we denote it by a $\star$. When available, we report the 95% confidence interval. We find that the undersampled classifier is always within 1–4% of the best performing robustness algorithm, except on the MultiNLI dataset.

Shift Type	Dataset/Paper	Test Accuracy/Worst-Group Accuracy*	
		Best	Undersampled
Label	Imb. CIFAR10 (step 10) [5]	87.81	84.59
	Imb. CIFAR100 (step 10) [5]	58.71	55.06
Group-Covariate	CelebA [12]	$86.9 \pm 1.1^\star$	$85.6 \pm 2.3^\star$
	Waterbirds [12]	$87.6 \pm 1.6^\star$	$89.1 \pm 1.1^\star$
	MultiNLI [12]	$78.0 \pm 0.7^\star$	$68.9 \pm 0.8^\star$
	CivilComments [12]	$72.0 \pm 1.9^\star$	$71.8 \pm 1.4^\star$

Inspired by the strong performance of undersampling in these experiments, we ask:

> *Is the performance of a model under distribution shift fundamentally*
> *constrained by the lack of minority group samples?*

To answer this question we analyze the *minimax excess risk*. We lower bound the minimax excess risk to prove that the performance of *any* algorithm is lower bounded only as a function of the minority

samples ($n_{\min}$). This shows that even if a robust algorithm optimally trades off between the bias and the variance, it is fundamentally constrained by the variance on the minority group which decreases only with $n_{\min}$.

For our study, we consider the well-studied setting of nonparametric binary classification [21]. By operating in this nonparametric regime we are able to study the properties of undersampling in rich data distributions, but are able to circumvent the complications that arise due to the optimization and implicit bias of parametric models. We explore two distribution shift scenarios: label shift and group-covariate shift. Under label shift, one of the labels is overrepresented in the training data, $\mathsf{P}_{\text{train}}(y = 1) \geq \mathsf{P}_{\text{train}}(y = -1)$, whereas the test samples are equally likely to come from either class. Here the class-conditional distribution $\mathsf{P}(x \mid y)$ is Lipschitz in x. Under group-covariate shift, we have two groups $\{a, b\}$ and in the training data we have more samples from the distribution $\mathsf{P}_a(x)$ than from $\mathsf{P}_b(x)$. Whereas during test time, it is equiprobable to receive samples from either group. In this case, the distribution $\mathsf{P}(y \mid x)$ is Lipschitz in x.

Our Contributions. We show that in the label shift setting there is a fundamental constraint, and that the minimax excess risk of *any robust learning method* is lower bounded by $1/n_{\min}^{1/3}$. That is, minority group samples fundamentally constrain performance under distribution shift. Furthermore, by leveraging previous results about nonparametric density estimation [9] we show a matching upper bound on the excess risk of a standard binning estimator trained on an undersampled dataset to demonstrate that undersampling is optimal.

In the case of group-covariate shift, we show that when the overlap (defined in terms of total variation distance) between the group distribution P_a and P_b is small, a similar result holds and the minimax excess risk of any robust learning algorithm is lower bounded by $1/n_{\min}^{1/3}$. We show that this lower bound is tight, by proving an upper bound on the excess risk of the binning estimator acting on the undersampled dataset.

Finally, we experimentally show in a label shift dataset (Imbalanced Binary CIFAR10) that the accuracy of popular classifiers generally follow the trends predicted by our theory. When the minority samples are increased, the accuracy of these classifiers increases drastically, whereas when the number of majority samples are increased the gains in the accuracy are marginal at best.

Taken together, our results underline the need to move beyond designing "general-purpose" robustness algorithms (like importance-weighting [5, 17, 13, 24], g-DRO [18], JTT [16], SMOTE [6], etc.) that are agnostic to the structure in the distribution shift. Our worst case analysis highlights that to successfully beat undersampling, an algorithm must leverage additional structure in the distribution shift.

2   Related Work

On several group-covariate shift benchmarks (CelebA, CivilComments, Waterbirds), Idrissi et al. [12] showed that training ResNet classifiers on an undersampled dataset either outperforms or performs as well as other popular reweighting methods like Group-DRO [18], reweighted ERM, and Just-Train-Twice [16]. They find Group-DRO performs comparably to undersampling, while both tend to outperform methods that don't utilize group information.

One classic method to tackle distribution shift is importance weighting [19], which reweights the loss of the minority group samples to yield an unbiased estimate of the loss. However, recent work [3, 27] has demonstrated the ineffectiveness of such methods when applied to overparameterized neural networks. Many followup papers [5, 29, 17, 13, 24] have introduced methods that modify the loss function in various ways to address this. However, despite this progress undersampling remains a competitive alternative to these importance weighted classifiers.

Our theory draws from the rich literature on non-parametric classification [21]. Apart from borrowing this setting of nonparametric classification, we also utilize upper bounds on the estimation error of the simple histogram estimator [9, 8] to prove our upper bounds in the label shift case. Finally, we note that to prove our minimax lower bounds we proceed by using the general recipe of reducing from estimation to testing [22, Chapter 15]. One difference from this standard framework is that our training samples shall be drawn from a different distribution than the test samples used to define the risk.

3 Setting

In this section, we shall introduce our problem setup and define the types of distribution shift that we consider.

3.1 Problem Setup

The setting for our study is nonparametric binary classification with Lipschitz data distributions. We are given n training datapoints $\mathcal{S} := \{(x_1, y_1), \ldots, (x_n, y_n)\} \in ([0, 1] \times \{-1, 1\})^n$ that are all drawn from a *train* distribution $\mathsf{P}_{\text{train}}$. During test time, the data shall be drawn from a *different* distribution P_{test}. To present a clean analysis, we study the case where the features x are bounded scalars, however, it is easy to extend our results to the high-dimensional setting.

Given a classifier $f : \mathbb{R} \to \{-1, 1\}$, we shall be interested in the test error (risk) of this classifier under the test distribution P_{test}:

$$R(f; \mathsf{P}_{\text{test}}) := \mathbb{E}_{(x,y) \sim \mathsf{P}_{\text{test}}} \left[\mathbf{1}(f(x) \neq y) \right].$$

3.2 Types of Distribution Shift

We assume that $\mathsf{P}_{\text{train}}$ consists of a mixture of two groups of unequal size, and P_{test} contains equal numbers of samples from both groups. Given a majority group distribution P_{maj} and a minority group distribution P_{min}, the learner has access to n_{maj} majority group samples and n_{min} minority group samples:

$$\mathcal{S}_{\text{maj}} \sim \mathsf{P}_{\text{maj}}^{n_{\text{maj}}} \quad \text{and} \quad \mathcal{S}_{\text{min}} \sim \mathsf{P}_{\text{min}}^{n_{\text{min}}}.$$

Here $n_{\text{maj}} > n/2$ and $n_{\text{min}} < n/2$ with $n_{\text{maj}} + n_{\text{min}} = n$. The full training dataset is $\mathcal{S} = \mathcal{S}_{\text{maj}} \cup \mathcal{S}_{\text{min}} = \{(x_1, y_1), \ldots, (x_n, y_n)\}$. We assume that the learner has access to the knowledge whether a particular sample (x_i, y_i) comes from the majority or minority group.

The test samples will be drawn from $\mathsf{P}_{\text{test}} = \frac{1}{2}\mathsf{P}_{\text{maj}} + \frac{1}{2}\mathsf{P}_{\text{min}}$, a uniform mixture over P_{maj} and P_{min}. Thus, the training dataset is an imbalanced draw from the distributions P_{maj} and P_{min}, whereas the test samples are balanced draws. We let $\rho := n_{\text{maj}}/n_{\text{min}} > 1$ denote the imbalance ratio in the training data.

We focus on two-types of distribution shifts: label shift and group-covariate shift that we describe below.

3.2.1 Label Shift

In this setting, the imbalance in the training data comes from there being more samples from one class over another. Without loss of generality, we shall assume that the class $y = 1$ is the majority class. Then, we define the majority and the minority class distributions as

$$\mathsf{P}_{\text{maj}}(x, y) = \mathsf{P}_1(x)\mathbf{1}(y = 1) \quad \text{and} \quad \mathsf{P}_{\text{min}} = \mathsf{P}_{-1}(x)\mathbf{1}(y = -1),$$

where $\mathsf{P}_1, \mathsf{P}_{-1}$ are class-conditional distributions over the interval $[0, 1]$. We assume that class-conditional distributions P_i have densities on $[0, 1]$ and that they are 1-Lipschitz: for any $x, x' \in [0, 1]$,

$$|\mathsf{P}_i(x) - \mathsf{P}_i(x')| \leq |x - x'|.$$

We denote the class of pairs of distributions $(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}})$ that satisfy these conditions by $\mathcal{P}_{\text{LS}}$.

3.2.2 Group-Covariate Shift

In this setting, we have two groups $\{a, b\}$, and corresponding to each of these groups is a distribution (with densities) over the features $\mathsf{P}_a(x)$ and $\mathsf{P}_b(x)$. We let a correspond to the majority group and b correspond to the minority group. Then, we define

$$\mathsf{P}_{\text{maj}}(x, y) = \mathsf{P}_a(x)\mathsf{P}(y \mid x) \quad \text{and} \quad \mathsf{P}_{\text{min}}(x, y) = \mathsf{P}_b(x)\mathsf{P}(y \mid x).$$

We assume that for $y \in \{-1, 1\}$, for all $x, x' \in [0, 1]$:

$$\left| \mathsf{P}(y \mid x) - \mathsf{P}(y \mid x') \right| \leq |x - x'|,$$

that is, the distribution of the label given the feature is 1-Lipschitz, and it varies slowly over the domain.

To quantify the shift between the train and test distribution, we define a notion of overlap between the group distributions P_a and P_b as follows:

$$\mathsf{Overlap}(P_a, P_b) := 1 - \mathrm{TV}(P_a, P_b).$$

Notice that when P_a and P_b have disjoint supports, $\mathrm{TV}(P_a, P_b) = 1$ and therefore $\mathsf{Overlap}(P_a, P_b) = 0$. On the other hand when $P_a = P_b$, $\mathrm{TV}(P_a, P_b) = 0$ and $\mathsf{Overlap}(P_a, P_b) = 1$. When the overlap is 1, the majority and minority distributions are identical and hence we have no shift between train and test. Observe that $\mathsf{Overlap}(P_a, P_b) = \mathsf{Overlap}(P_{\mathsf{maj}}, P_{\mathsf{min}})$ since $P(y \mid x)$ is shared across P_{maj} and P_{min}.

Given a level of overlap $\tau \in [0, 1]$ we denote the class of pairs of distributions $(P_{\mathsf{maj}}, P_{\mathsf{min}})$ with overlap at least τ by $\mathcal{P}_{\mathsf{GS}}(\tau)$. It is easy to check that, $\mathcal{P}_{\mathsf{GS}}(\tau) \subseteq \mathcal{P}_{\mathsf{GS}}(0)$ at any overlap level $\tau \in [0, 1]$.

4 Lower Bounds on the Minimax Excess Risk

In this section, we shall prove our lower bounds that show that the performance of any algorithm is constrained by the number of minority samples n_{min}. Before we state our lower bounds, we need to introduce the notion of excess risk and minimax excess risk.

Excess Risk and Minimax Excess Risk. We measure the performance of an algorithm $\mathcal{A}$ through its excess risk defined in the following way. Given an algorithm $\mathcal{A}$ that takes as input a dataset $\mathcal{S}$ and returns a classifier $\mathcal{A}^{\mathcal{S}}$, and a pair of distributions $(P_{\mathsf{maj}}, P_{\mathsf{min}})$ with $P_{\mathsf{test}} = \frac{1}{2}P_{\mathsf{maj}} + \frac{1}{2}P_{\mathsf{min}}$, the *expected excess risk* is given by

$$\mathsf{Excess\ Risk}[\mathcal{A}; (P_{\mathsf{maj}}, P_{\mathsf{min}})] := \mathbb{E}_{\mathcal{S} \sim P_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times P_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R(\mathcal{A}^{\mathcal{S}}; P_{\mathsf{test}})) - R(f^{\star}(P_{\mathsf{test}}); P_{\mathsf{test}}) \right], \quad (1)$$

where $f^{\star}(P_{\mathsf{test}})$ is the Bayes classifier that minimizes the risk $R(\cdot; P_{\mathsf{test}})$. The first term corresponds to the expected risk for the algorithm when given n_{maj} samples from P_{maj} and n_{min} samples from P_{min}, whereas the second term corresponds to the Bayes error for the problem.

Excess risk does not let us characterize the inherent difficulty of a problem, since for any particular data distribution $(P_{\mathsf{maj}}, P_{\mathsf{min}})$ the best possible algorithm $\mathcal{A}$ to minimize the excess risk would be the trivial mapping $\mathcal{A}^{\mathcal{S}} = f^{\star}(P_{\mathsf{test}})$. Therefore, to prove meaningful lower bounds on the performance of algorithms we need to define the notion of minimax excess risk [see 22, Chapter 15]. Given a class of pairs of distributions $\mathcal{P}$ define

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}) := \inf_{\mathcal{A}} \sup_{(P_{\mathsf{maj}}, P_{\mathsf{min}}) \in \mathcal{P}} \mathsf{Excess\ Risk}[\mathcal{A}; (P_{\mathsf{maj}}, P_{\mathsf{min}})], \quad (2)$$

where the infimum is over all measurable estimators $\mathcal{A}$. The minimax excess risk is the excess risk of the "best" algorithm in the worst case over the class of problems defined by $\mathcal{P}$.

4.1 Label Shift Lower Bounds

We demonstrate the hardness of the label shift problem in general by establishing a lower bound on the minimax excess risk. Below we let $c > 0$ be an absolute constant independent of problem parameters like n_{maj} and n_{min}.

Theorem 4.1. *Consider the label shift setting described in Section 3.2.1. Recall that $\mathcal{P}_{\mathsf{LS}}$ is the class of pairs of distributions $(P_{\mathsf{maj}}, P_{\mathsf{min}})$ that satisfy the assumptions in that section. The minimax excess risk over this class is lower bounded as follows:*

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\mathsf{LS}}) = \inf_{\mathcal{A}} \sup_{(P_{\mathsf{maj}}, P_{\mathsf{min}}) \in \mathcal{P}_{\mathsf{LS}}} \mathsf{Excess\ Risk}[\mathcal{A}; (P_{\mathsf{maj}}, P_{\mathsf{min}})] \geq \frac{c}{n_{\mathsf{min}}^{1/3}}. \quad (3)$$

We establish this result in Appendix B.

We show that rather surprisingly, the lower bound on the minimax excess risk scales only with the number of minority class samples $n_{\mathsf{min}}^{1/3}$, and does not depend on n_{maj}. Intuitively, this is because

any learner must predict which class-conditional distribution ($P(x \mid 1)$ or $P(x \mid -1)$) assigns higher likelihood at that x. To interpret this result, consider the extreme scenario where $n_{\text{maj}} \to \infty$ but n_{min} is finite. In this case, the learner has full information about the majority class distribution. However, the learning task continues to be challenging since any learner would be uncertain about whether the minority class distribution assigns higher or lower likelihood at any given x. This uncertainty underlies the reason why the minimax rate of classification is constrained by the number of minority samples n_{min}.

We also note that the theorem can be trivially extended to higher dimensions. In this case the exponents degrade to $1/3d$ rather than $1/3$ as is to be expected in nonparametric classification.

4.2 Group-Covariate Shift Lower Bounds

Next, we shall state our lower bound on the minimax excess risk that demonstrates the hardness of the group-covariate shift problem. In the theorem below $c > 0$ shall be an absolute constant independent of n_{maj}, n_{min} and τ.

Theorem 4.2. *Consider the group shift setting described in Section 3.2.2. Given any overlap $\tau \in [0,1]$ recall that $\mathcal{P}_{\text{GS}}(\tau)$ is the class of distributions such that $\mathsf{Overlap}(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}}) \geq \tau$. The minimax excess risk in this setting is lower bounded as follows:*

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\text{GS}}(\tau)) = \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}}) \in \mathcal{P}_{\text{GS}}(\tau)} \mathsf{Excess\ Risk}[\mathcal{A}; (\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}})]$$

$$\geq \frac{c}{(n_{\text{min}} \cdot (2 - \tau) + n_{\text{maj}} \cdot \tau)^{1/3}} \geq \frac{c}{n_{\text{min}}^{1/3}(\rho \cdot \tau + 2)^{1/3}}, \quad (4)$$

where $\rho = n_{\text{maj}}/n_{\text{min}} > 1$.

We prove this theorem in Appendix C.

We see that in the *low overlap* setting ($\tau \ll 1/\rho$), the minimax excess risk is lower bounded by $1/n_{\text{min}}^{1/3}$, and we are fundamentally constrained by the number of samples in minority group. To see why this is the case, consider the extreme example with $\tau = 0$ where P_a has support $[0, 0.5]$ and P_b has support $[0.5, 1]$. The n_{maj} majority group samples from P_a provide information about the correct label predict in the interval $[0, 0.5]$ (the support of P_a). However, since the distribution $P(y \mid x)$ is 1-Lipschitz in the worst case these samples provide very limited information about the correct predictions in $[0.5, 1]$ (the support of P_b). Thus, predicting on the support of P_b requires samples from the minority group and this results in the n_{min} dependent rate. In fact, in this extreme case ($\tau = 0$) even if $n_{\text{maj}} \to \infty$, the minimax excess risk is still bounded away from zero. This intuition also carries over to the case when the overlap is small but non-zero and our lower bound shows that minority samples are much more valuable than majority samples at reducing the risk.

On the other hand, when the overlap is high ($\tau \gg 1/\rho$) the minimax excess risk is lower bounded by $1/(n_{\text{min}}(2 - \tau) + n_{\text{maj}}\tau)^{1/3}$ and the extra majority samples are quite beneficial. This is roughly because the supports of P_a and P_b have large overlap and hence samples from the majority group are useful in helping make predictions even in regions where P_b is large. In the extreme case when $\tau = 1$, we have that $\mathsf{P}_a = \mathsf{P}_b$ and therefore recover the classic i.i.d. setting with no distribution shift. Here, the lower bound scales with $1/n^{1/3}$, as one might expect.

Identical to the label shift case, the theorem can be extended to hold in higher dimensions with the exponents being $1/3d$ rather than $1/3$.

5 Upper Bounds on the Excess Risk for the Undersampled Binning Estimator

We will show that an undersampled estimator matches the rates in the previous section showing that undersampling is an optimal robustness intervention. We start by defining the undersampling procedure and the undersampling binning estimator.

Undersampling Procedure. Given training data $\mathcal{S} := \{(x_1, y_1), \ldots, (x_n, y_n)\}$, generate a new undersampled dataset $\mathcal{S}_{\text{US}}$ by

216   • including all n_{min} samples from $\mathcal{S}_{\text{min}}$ and,

217   • including n_{min} samples from $\mathcal{S}_{\text{maj}}$ by sampling uniformly at random without replacement.

218 This procedure ensures that in the undersampled dataset $\mathcal{S}_{\text{US}}$, the groups are balanced, and that
219 $|\mathcal{S}_{\text{US}}| = 2n_{\text{min}}$.

220 The undersampling binning estimator defined next will first run this undersampling procedure to
221 obtain $\mathcal{S}_{\text{US}}$ and just uses these samples to output a classifier.

Undersampled Binning Estimator   The undersampled binning estimator $\mathcal{A}_{\text{USB}}$ takes as input a
223 dataset $\mathcal{S}$ and a positive integer K corresponding to the number of bins, and returns a classifier
224 $\mathcal{A}_{\text{USB}}^{\mathcal{S},K} : [0,1] \to \{-1,1\}$. This estimator is defined as follows:

1. First, we compute the undersampled dataset $\mathcal{S}_{\text{US}}$.

226   2. Given this dataset $\mathcal{S}_{\text{US}}$, let $n_{1,j}$ be the number of points with label $+1$ that lie in the interval
227        $I_j = [\frac{j-1}{K}, \frac{j}{K}]$. Also, define $n_{-1,j}$ analogously. Then set

$$\mathcal{A}_j = \begin{cases} 1 & \text{if } n_{1,j} > n_{-1,j}, \\ -1 & \text{otherwise.} \end{cases}$$

228   3. Define the classifier $\mathcal{A}_{\text{USB}}^{\mathcal{S},K}$ such that if $x \in I_j$ then

$$\mathcal{A}_{\text{USB}}^{\mathcal{S},K}(x) = \mathcal{A}_j. \tag{5}$$

229   Essentially in each bin I_j, we set the prediction to be the majority label among the samples
230   that fall in this bin.

231 Whenever the number of bins K is clear from the context we shall denote $\mathcal{A}_{\text{USB}}^{\mathcal{S},K}$ by $\mathcal{A}_{\text{USB}}^{\mathcal{S}}$. Below we
232 establish upper bounds on the excess risk of this simple estimator.

5.1   Label Shift Upper Bounds

234 We now establish an upper bound on the excess risk of $\mathcal{A}_{\text{USB}}$ in the label shift setting (see Sec-
235 tion 3.2.1). Below we let $c, C > 0$ be absolute constants independent of problem parameters like
236 n_{maj} and n_{min}.

Theorem 5.1. *Consider the label shift setting described in Section 3.2.1. For any* $(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}}) \in \mathcal{P}_{\text{LS}}$
238 *the expected excess risk of the Undersampling Binning Estimator (Eq. (5)) with number of bins with*
239 $K = c\lceil n_{\text{min}}^{1/3} \rceil$ *is upper bounded by*

$$\text{Excess Risk}[\mathcal{A}_{\text{USB}}; (\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}})] = \mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\text{maj}}^{n_{\text{maj}}} \times \mathsf{P}_{\text{min}}^{n_{\text{min}}}} \left[R(\mathcal{A}_{\text{USB}}^{\mathcal{S}}; \mathsf{P}_{\text{test}}) - R(f^{\star}; \mathsf{P}_{\text{test}}) \right] \leq \frac{C}{n_{\text{min}}^{1/3}}.$$

240 We prove this result in Appendix B. This upper bound combined with the lower bound in Theorem 4.1
241 shows that an undersampling approach is minimax optimal up to constants in the presence of label
242 shift.

243 We note that our analysis leaves open the possibility of better algorithms when the learner has
244 additional information about the structure of the label shift beyond Lipschitz continuity.

5.2   Group-Covariate Shift Upper Bounds

246 Next, we present our upper bounds on the excess risk of the undersampled binning estimator in the
247 group-covariate shift setting (see Section 3.2.2). In the theorem below, $C > 0$ is an absolute constant
248 independent of the problem parameters n_{maj}, n_{min} and τ.

Theorem 5.2. *Consider the group shift setting described in Section 3.2.2. For any overlap* $\tau \in [0,1]$
250 *and for any* $(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}}) \in \mathcal{P}_{\text{GS}}(\tau)$ *the expected excess risk of the Undersampling Binning Estimator*
251 *(Eq. (5)) with number of bins with* $K = \lceil n_{\text{min}}^{1/3} \rceil$ *is*

$$\text{Excess Risk}[\mathcal{A}_{\text{USB}}; (\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}})] = \mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\text{maj}}^{n_{\text{maj}}} \times \mathsf{P}_{\text{min}}^{n_{\text{min}}}} \left[R(\mathcal{A}_{\text{USB}}^{\mathcal{S}}; \mathsf{P}_{\text{test}})) - R(f^{\star}; \mathsf{P}_{\text{test}}) \right] \leq \frac{C}{n_{\text{min}}^{1/3}}.$$

We provide a proof for this theorem in Appendix C. Compared to the lower bound established in Theorem 4.2 which scales as $1/\left((2-\tau)n_{\mathsf{min}} + n_{\mathsf{maj}}\tau\right)^{1/3}$, the upper bound for the undersampled binning estimator always scales with $1/n_{\mathsf{min}}^{1/3}$ since it operates on the undersampled dataset ($\mathcal{S}_{\mathsf{US}}$).

Thus, we have shown that in the absence of overlap ($\tau \ll 1/\rho = n_{\mathsf{min}}/n_{\mathsf{maj}}$) there is an undersampling algorithm that is minimax optimal up to constants. However when there is high overlap ($\tau \gg 1/\rho$) there is a non-trivial gap between the upper and lower bounds:

$$\frac{\text{Upper Bound}}{\text{Lower Bound}} = c(\rho \cdot \tau + 2)^{1/3}.$$

6 Minority Sample Dependence in Practice

Figure 2: Convolutional neural network classifiers trained on the Imbalanced Binary CIFAR10 dataset with a 5:1 label imbalance. (Top) Models trained using the importance weighted cross entropy loss with early stopping. (Bottom) Models trained using the importance weighted VS loss [13] with early stopping. We report the average test accuracy calculated on a balanced test set over 5 random seeds. We start off with 2500 cat examples and 500 dog examples in the training dataset. We find that in accordance with our theory, for both of the classifiers adding only minority class samples (red) leads to large gain in accuracy ($\sim 6\%$), while adding majority class samples (blue) leads to little or no gain. In fact, adding majority samples sometimes hurts test accuracy due to the added bias. When we add majority and minority samples in a 5:1 ratio (green), the gain is largely due to the addition of minority samples and is only marginally higher ($< 2\%$) than adding only minority samples. The green curves correspond to the same classifiers in both the left and right panels.

Inspired by our worst-case theoretical predictions in nonparametric classification, we ask: how does the accuracy of neural network classifiers trained using robust algorithms evolve as a function of the majority and minority samples?

To explore this question, we conduct a small case study using the imbalanced binary CIFAR10 dataset [3, 24] that is constructed using the "cat" and "dog" classes. The test set consists of all of the 1000 cat and 1000 dog test examples. To form our initial train and validation sets, we take 2500 cat examples but only 500 dog examples from the official train set, corresponding to a 5:1

label imbalance. We then use 80% of those examples for training and the rest for validation. In our experiment, we either (a) add only minority samples; (b) add only majority samples; (c) add both majority and minority samples in a 5:1 ratio. We consider competitive robust classifiers proposed in the literature that are convolutional neural networks trained either by using (i) the importance weighted cross entropy loss, or (ii) the importance weighted VS loss [13]. We early stop using the importance weighted validation loss in both cases. The additional experimental details are presented in Appendix D.

Our results in Figure 2 are generally consistent with our theoretical predictions. By adding only minority class samples the test accuracy of both classifiers increases by a great extent (6%), while by adding only majority class samples the test accuracy remains constant or in some cases even decreases owing to the added bias of the classifiers. When we add samples to both groups proportionately, the increase in the test accuracy appears to largely to be due to the increase in the number of minority class samples and on the left panels, we see that the difference between adding only extra minority group samples (red) and both minority and majority group samples (green) is small. Thus, we find that the accuracy for these neural network classifiers is also constrained by the number of minority class samples.

7 Discussion

We showed that undersampling is an optimal robustness intervention in nonparametric classification in the absence of significant overlap between group distributions or without additional structure beyond Lipschitz continuity.

At a high level our results highlight the need to reason about the specific structure in the distribution shift and design algorithms that are tailored to take advantage of this structure. This would require us to step away from the common practice in robust machine learning where the focus is to design "universal" robustness interventions that are agnostic to the structure in the shift. Alongside this, our results also dictate the need for datasets and benchmarks with the propensity for transfer from training time to test time.

References

[1] M. A. Alcorn, Q. Li, Z. Gong, C. Wang, L. Mai, W.-S. Ku, and A. Nguyen. Strike (with) a pose: Neural networks are easily fooled by strange poses of familiar objects. In *Computer Vision and Pattern Recognition (CVPR)*, 2019.

[2] S. L. Blodgett, L. Green, and B. T. O'Connor. Demographic dialectal variation in social media: A case study of african-american english. In *Empirical Methods in Natural Language Processing (EMNLP)*, 2016.

[3] J. Byrd and Z. Lipton. What is the effect of importance weighting in deep learning? In *International Conference on Machine Learning (ICML)*, 2019.

[4] C. L. Canonne. A short note on an inequality between KL and TV. *arXiv preprint arXiv:2202.07198*, 2022.

[5] K. Cao, C. Wei, A. Gaidon, N. Arechiga, and T. Ma. Learning imbalanced datasets with label-distribution-aware margin loss. *Advances in Neural Information Processing Systems (NeurIPS)*, 2019.

[6] N. V. Chawla, K. W. Bowyer, L. O. Hall, and W. P. Kegelmeyer. Smote: Synthetic minority over-sampling technique. *Journal of Artificial Intelligence Research*, 2002.

[7] Y. Cui, M. Jia, T.-Y. Lin, Y. Song, and S. Belongie. Class-balanced loss based on effective number of samples. In *Computer Vision and Pattern Recognition (CVPR)*, 2019.

[8] L. Devroye and L. Györfi. *Nonparametric density estimation: the L_1 view*. Wiley Series in Probability and Mathematical Statistics, 1985.

[9] D. Freedman and P. Diaconis. On the histogram as a density estimator: L_2 theory. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 1981.

[10] T. Hashimoto, M. Srivastava, H. Namkoong, and P. Liang. Fairness without demographics in repeated loss minimization. In *International Conference on Machine Learning (ICML)*, 2018.

[11] D. Hovy and A. Søgaard. Tagging performance correlates with author age. In *Association for Computational Linguistics (ACL)*, 2015.

[12] B. Y. Idrissi, M. Arjovsky, M. Pezeshki, and D. Lopez-Paz. Simple data balancing achieves competitive worst-group-accuracy. In *Causal Learning and Reasoning*, 2022.

[13] G. R. Kini, O. Paraskevas, S. Oymak, and C. Thrampoulidis. Label-imbalanced and group-sensitive classification under overparameterization. *Advances in Neural Information Processing Systems (NeurIPS)*, 2021.

[14] M. Kubat, S. Matwin, et al. Addressing the curse of imbalanced training sets: one-sided selection. In *International Conference on Machine Learning (ICML)*, 1997.

[15] Z. Lipton, Y.-X. Wang, and A. Smola. Detecting and correcting for label shift with black box predictors. In *International Conference on Machine Learning (ICML)*, 2018.

[16] E. Z. Liu, B. Haghgoo, A. S. Chen, A. Raghunathan, P. W. Koh, S. Sagawa, P. Liang, and C. Finn. Just train twice: Improving group robustness without training group information. In *International Conference on Machine Learning (ICML)*, 2021.

[17] A. K. Menon, S. Jayasumana, A. S. Rawat, H. Jain, A. Veit, and S. Kumar. Long-tail learning via logit adjustment. In *International Conference on Learning Representations (ICLR)*, 2020.

[18] S. Sagawa, P. W. Koh, T. B. Hashimoto, and P. Liang. Distributionally robust neural networks. In *International Conference on Learning Representations (ICLR)*, 2020.

[19] H. Shimodaira. Improving predictive inference under covariate shift by weighting the log-likelihood function. *Journal of Statistical Planning and Inference*, 2000.

[20] R. Tatman. Gender and dialect bias in youtube's automatic captions. In *ACL Workshop on Ethics in Natural Language Processing*, 2017.

[21] A. B. Tsybakov. *Introduction to Nonparametric Estimation*. Springer, 2010.

[22] M. J. Wainwright. *High-dimensional statistics: A non-asymptotic viewpoint*. Cambridge University Press, 2019.

[23] B. C. Wallace, K. Small, C. E. Brodley, and T. A. Trikalinos. Class imbalance, redux. In *International Conference on Data Mining (ICDM*, 2011.

[24] K. A. Wang, N. S. Chatterji, S. Haque, and T. Hashimoto. Is importance weighting incompatible with interpolating classifiers? In *International Conference on Learning Representations (ICLR)*, 2022.

[25] L. Wasserman. Lecture notes in nonparametric classification. URL `https://www.stat.cmu.edu/~larry/=sml/nonparclass.pdf`. [Online; accessed 12-May-2022].

[26] Wikipedia contributors. Poisson binomial distribution — Wikipedia, the free encyclopedia, 2022. URL `https://en.wikipedia.org/w/index.php?title=Poisson_binomial_distribution&oldid=1071847908`. [Online; accessed 5-May-2022].

[27] D. Xu, Y. Ye, and C. Ruan. Understanding the role of importance weighting for deep learning. In *International Conference on Learning Representations (ICLR)*, 2020.

[28] Y. Yang and Z. Xu. Rethinking the value of labels for improving class-imbalanced learning. *Advances in Neural Information Processing Systems (NeurIPS)*, 2020.

[29] H.-J. Ye, H.-Y. Chen, D.-C. Zhan, and W.-L. Chao. Identifying and compensating for feature deviation in imbalanced deep learning. *arXiv preprint arXiv:2001.01385*, 2020.

A Technical Tools

In this section we avail ourselves of some technical tools that shall be used in all of the proofs below.

A.1 Reduction to lower bounds over a finite class

The lower bound on the minimax excess risk will be established via the usual route of first identifying a "hard" finite set of problem instances and then establishing the lower bound over this finite class. One difference from the usual setup in proving such lower bounds [see 22, Chapter 15] is that the training samples are drawn from an imbalanced distribution, whereas the test samples are drawn from a balanced one.

Let $\mathcal{P}$ be a class of pairs of distributions, where each element $(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \in \mathcal{P}$ is a pair of distributions over $[0,1] \times \{-1, 1\}$. As before, we let $\mathsf{P}_{\mathsf{test}}$ denote the uniform mixture over $\mathsf{P}_{\mathsf{maj}}$ and $\mathsf{P}_{\mathsf{min}}$. We let $\mathcal{V}$ denote a finite index set. Corresponding to each element $v \in \mathcal{V}$ there is a $\mathsf{P}_v = (\mathsf{P}_{v,\mathsf{maj}}, \mathsf{P}_{v,\mathsf{min}}) \in \mathcal{P}$ with $\mathsf{P}_{v,\mathsf{test}} = (\mathsf{P}_{v,\mathsf{maj}} + \mathsf{P}_{v,\mathsf{min}})/2$. Finally, also define a pair of random variables (V, S) as follows:

1. V is a uniform random variable over the set $\mathcal{V}$.
2. $(S \mid V = v) \sim \mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}$, is an independent draw of n_{maj} samples from $\mathsf{P}_{v,\mathsf{maj}}$ and n_{min} samples from $\mathsf{P}_{v,\mathsf{min}}$.

We shall let Q denote the joint distribution of the random variables (V, S), and let Q_S denote the marginal distribution of S.

With this notation in place, we now present a lemma that lower bounds the minimax excess risk in terms of quantities defined over the finite class of "hard" instances P_v.

Lemma A.1. *Let the random variables (V, S) be as defined above. The minimax excess risk is lower bounded as follows:*

$$\text{Minimax Excess Risk}(\mathcal{P}) = \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \in \mathcal{P}} \mathbb{E}_{S \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R(\mathcal{A}^S; \mathsf{P}_{\mathsf{test}}) - R(f^\star(\mathsf{P}_{\mathsf{test}}); \mathsf{P}_{\mathsf{test}}) \right]$$

$$\geq \mathfrak{R}_{\mathcal{V}} - \mathfrak{B}_{\mathcal{V}},$$

where $\mathfrak{R}_{\mathcal{V}}$ and Bayes-error $\mathfrak{B}_{\mathcal{V}}$ are defined as

$$\mathfrak{R}_{\mathcal{V}} := \mathbb{E}_{S \sim \mathsf{Q}_S} [\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} \mathsf{Q}(v|S)\mathsf{P}_{v,\mathsf{test}}}(h(x) \neq y)],$$

$$\mathfrak{B}_{\mathcal{V}} := \mathbb{E}_V [R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}})].$$

Proof. By the definition of Minimax Excess Risk,

$$\text{Minimax Excess Risk} = \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \in \mathcal{P}} \mathbb{E}_{S \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} [R(\mathcal{A}^S; \mathsf{P}_{\mathsf{test}})] - R(f^\star(\mathsf{P}_{\mathsf{test}}); \mathsf{P}_{\mathsf{test}})$$

$$\geq \inf_{\mathcal{A}} \sup_{v \in \mathcal{V}} \mathbb{E}_{S|v \sim \mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}} [R(\mathcal{A}^S; \mathsf{P}_{v,\mathsf{test}})] - R(f^\star(\mathsf{P}_{v,\mathsf{test}}); \mathsf{P}_{v,\mathsf{test}})$$

$$\geq \inf_{\mathcal{A}} \mathbb{E}_V \left[\mathbb{E}_{S|V \sim \mathsf{P}_{V,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{V,\mathsf{min}}^{n_{\mathsf{min}}}} [R(\mathcal{A}^S; \mathsf{P}_{V,\mathsf{test}})] - R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}})) \right]$$

$$= \inf_{\mathcal{A}} \mathbb{E}_V [\mathbb{E}_{S|V \sim \mathsf{P}_{V,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{V,\mathsf{min}}^{n_{\mathsf{min}}}} [R(\mathcal{A}^S; \mathsf{P}_{V,\mathsf{test}})]] - \underbrace{\mathbb{E}_V [R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}}))]}_{=\mathfrak{B}_{\mathcal{V}}}.$$

We continue lower bounding the first term as follows

$$\inf_{\mathcal{A}} \mathbb{E}_V [\mathbb{E}_{S|V \sim \mathsf{P}_{V,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{V,\mathsf{min}}^{n_{\mathsf{min}}}} [R(\mathcal{A}^S; \mathsf{P}_{V,\mathsf{test}})]] = \inf_{\mathcal{A}} \mathbb{E}_{(V,S) \sim \mathsf{Q}} [\mathbb{P}_{(x,y) \sim \mathsf{P}_{V,\mathsf{test}}}(\mathcal{A}^S(x) \neq y)]$$

$$= \inf_{\mathcal{A}} \mathbb{E}_{S \sim \mathsf{Q}_S} \mathbb{E}_{V \sim \mathsf{Q}(\cdot|S)} [\mathbb{P}_{(x,y) \sim \mathsf{P}_{V,\mathsf{test}}}(\mathcal{A}^S(x) \neq y)]$$

$$\overset{(i)}{\geq} \mathbb{E}_{S \sim \mathsf{Q}_S} [\inf_h \mathbb{E}_{V \sim \mathsf{Q}(\cdot|S)} [\mathbb{P}_{(x,y) \sim \mathsf{P}_{V,\mathsf{test}}}(h(x) \neq y)]]$$

$$= \mathbb{E}_{S \sim \mathsf{Q}_S} [\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} \mathsf{Q}(v|S)\mathsf{P}_{v,\mathsf{test}}}(h(x) \neq y)]$$

$$= \mathfrak{R}_{\mathcal{V}},$$

where (i) follows since $\mathcal{A}^S$ is a fixed classifier given the sample set S. This, combined with the previous equation block completes the proof. $\square$

A.2 The Hat Function and its Properties

In this section, we define the *hat function* and establish some of its properties. This function will be useful in defining "hard" problem instances to prove our lower bounds. Given a positive integer K the hat function is defined as

$$\phi_K(x) = \begin{cases} \left|x + \frac{1}{4K}\right| - \frac{1}{4K} & \text{for } x \in \left[-\frac{1}{2K}, 0\right], \\ \frac{1}{4K} - \left|x - \frac{1}{4K}\right| & \text{for } x \in \left[0, \frac{1}{2K}\right], \\ 0 & \text{otherwise.} \end{cases} \tag{6}$$

When K is clear from context, we omit the subscript.

Figure 3: The hat function with $K = 4$.

We first notice that this function is 1-Lipschitz and odd, so

$$\int_{-\frac{1}{2K}}^{\frac{1}{2K}} \phi_K(x) \, \mathrm{d}x = 0.$$

We also compute some other key quantities for ϕ.

Lemma A.2. *For any positive integer K,*

$$\int_{-\frac{1}{2K}}^{\frac{1}{2K}} |\phi_K(x)| \, \mathrm{d}x = \frac{1}{8K^2}.$$

Proof. We suppress K in the notation. We have that,

$$\int_{-\frac{1}{2K}}^{\frac{1}{2K}} |\phi(x)| \, \mathrm{d}x = \int_{-\frac{1}{2K}}^{0} \left|\frac{1}{4K} - \left|x + \frac{1}{4K}\right|\right| \, \mathrm{d}x + \int_{0}^{\frac{1}{2K}} \left|\left|x - \frac{1}{4K}\right| - \frac{1}{4K}\right| \, \mathrm{d}x.$$

The integrand $\left|\frac{1}{4K} - \left|x + \frac{1}{4K}\right|\right|$ over $x \in \left[-\frac{1}{2K}, 0\right]$ defines a triangle with base $\frac{1}{2K}$ and height $\frac{1}{4K}$, thus it has area $\frac{1}{16K^2}$. Therefore,

$$\int_{-\frac{1}{2K}}^{0} \left|\frac{1}{4K} - \left|x + \frac{1}{4K}\right|\right| \, \mathrm{d}x = \frac{1}{16K^2}.$$

The same holds for the second term. Thus, by adding them up we get that $\int_{-\frac{1}{2K}}^{\frac{1}{2K}} |\phi(x)| \, \mathrm{d}x = \frac{1}{8K^2}$. $\qquad\square$

Lemma A.3. *For any positive integer K,*

$$\int_{0}^{\frac{1}{K}} \log\left(\frac{1 + \phi_K(x - \frac{1}{2K})}{1 - \phi_K(x - \frac{1}{2K})}\right)\left(1 + \phi_K\left(x - \frac{1}{2K}\right)\right) \, \mathrm{d}x \le \frac{1}{3K^3}$$

and

$$\int_0^{\frac{1}{K}} \log\left(\frac{1 - \phi_K(x - \frac{1}{2K})}{1 + \phi_K(x - \frac{1}{2K})}\right)\left(1 - \phi_K\left(x - \frac{1}{2K}\right)\right)\, \mathrm{d}x \le \frac{1}{3K^3}.$$

Proof. Let us suppress K in the notation. We prove the first bound below and the second bound follows by an identical argument. We have that

$$\int_0^{\frac{1}{K}} \log\left(\frac{1 + \phi(x - \frac{1}{2K})}{1 - \phi(x - \frac{1}{2K})}\right)\left(1 + \phi\left(x - \frac{1}{2K}\right)\right)\, \mathrm{d}x$$

$$= \int_{-\frac{1}{2K}}^{\frac{1}{2K}} \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)(1 + \phi(x))\, \mathrm{d}x$$

$$= \int_0^{\frac{1}{2K}} \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)(1 + \phi(x))\, \mathrm{d}x + \int_{-\frac{1}{2K}}^0 \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)(1 + \phi(x))\, \mathrm{d}x$$

$$= \int_0^{\frac{1}{2K}} \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)(1 + \phi(x))\, \mathrm{d}x - \int_{\frac{1}{2K}}^0 \log\left(\frac{1 + \phi(-x)}{1 - \phi(-x)}\right)(1 + \phi(-x))\, \mathrm{d}x$$

$$= \int_0^{\frac{1}{2K}} \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)(1 + \phi(x))\, \mathrm{d}x + \int_0^{\frac{1}{2K}} \log\left(\frac{1 - \phi(x)}{1 + \phi(x)}\right)(1 - \phi(x))\, \mathrm{d}x,$$

where the last equality follows since ϕ is an odd function. Now, we may collect the integrands to get that,

$$\int_0^{\frac{1}{K}} \log\left(\frac{1 + \phi(x - \frac{1}{2K})}{1 - \phi(x - \frac{1}{2K})}\right)\left(1 + \phi\left(x - \frac{1}{2K}\right)\right)\, \mathrm{d}x$$

$$= 2\int_0^{\frac{1}{2K}} \log\left(\frac{1 + \phi(x)}{1 - \phi(x)}\right)\phi(x)\, \mathrm{d}x$$

$$= 2\int_0^{\frac{1}{2K}} \log\left(1 + \frac{2\phi(x)}{1 - \phi(x)}\right)\phi(x)\, \mathrm{d}x$$

$$\le 2\int_0^{\frac{1}{2K}} \frac{2\phi(x)^2}{1 - \phi(x)}\, \mathrm{d}x,$$

where the last inequality follows since $\log(1 + x) \le x$ for all x. Now we observe that $\phi(x) \le x \le \frac{1}{2}$ for $x \in [0, \frac{1}{2K}]$, and in particular, $\frac{1}{1 - \phi(x)} \le 2$. Thus,

$$\int_0^{\frac{1}{K}} \log\left(\frac{1 + \phi(x - \frac{1}{2K})}{1 - \phi(x - \frac{1}{2K})}\right)\left(1 + \phi\left(x - \frac{1}{2K}\right)\right)\, \mathrm{d}x$$

$$\le 8\int_0^{\frac{1}{2K}} \phi(x)^2\, \mathrm{d}x$$

$$\le 8\int_0^{\frac{1}{2K}} x^2\, \mathrm{d}x$$

$$= \frac{1}{3K^3}.$$

This proves the first bound. The second bound follows analogously. $\square$

B Proofs in the Label Shift Setting

Throughout this section we operate in the label shift setting (Section 3.2.1).

First, in Appendix B.1 through a sequence of lemmas we prove the minimax lower bound Theorem 4.1. Next, in Appendix B.2 we prove Theorem 5.1 which is an upper bound on the excess risk of the undersampled binning estimator (see Eq. (5)) with $\lceil n_{\min} \rceil^{1/3}$ bins by invoking previous results on nonparametric density estimation [9, 8].

B.1 Proof of Theorem 4.1

In this section, we provide a proof of the minimax lower bound in the label shift setting.

We construct the "hard" set of distributions as follows. Fix K to be an integer that will be specified in the sequel. Let the index set be $\mathcal{V} = \{-1, 0, 1\}^K \times \{-1, 0, 1\}^K$. For $v \in \mathcal{V}$, we will let $v_1 \in \{-1, 0, 1\}^K$ be the first K coordinates and $v_{-1} \in \{-1, 0, 1\}^K$ be the last K coordinates. That is, $v = (v_1, v_{-1})$.

For every $v \in \mathcal{P}$ we shall define pair of class-conditional distributions $\mathsf{P}_{v,1}$ and $\mathsf{P}_{v,-1}$ as follows: for $x \in I_j = [\frac{j-1}{K}, \frac{j}{K}]$,

$$\mathsf{P}_{v,1}(x) = 1 + v_{1,j}\phi\left(x - \frac{j + 1/2}{K}\right)$$

$$\mathsf{P}_{v,-1}(x) = 1 + v_{-1,j}\phi\left(x - \frac{j + 1/2}{K}\right),$$

where ϕ is defined in Eq. 6. Notice that $\mathsf{P}_{v,1}$ only depends on v_1 while $\mathsf{P}_{v,-1}$ only depends on v_{-1}. We continue to define We continue to define

$$\mathsf{P}_{v,\mathsf{maj}}(x, y) = \mathsf{P}_{v,1}(x)\mathbf{1}(y = 1)$$
$$\mathsf{P}_{v,\mathsf{min}}(x, y) = \mathsf{P}_{v,-1}(x)\mathbf{1}(y = -1),$$

and

$$\mathsf{P}_{v,\mathsf{test}}(x, y) = \frac{\mathsf{P}_{v,\mathsf{maj}}(x, y) + \mathsf{P}_{v,\mathsf{min}}(x, y)}{2} = \frac{\mathsf{P}_{v,1}(x)\mathbf{1}(y = 1) + \mathsf{P}_{v,-1}(x)\mathbf{1}(y = -1)}{2}.$$

Observe that in the test distribution it is equally likely for the label to be $+1$ or -1.

Recall that as described in Section A.1, V shall be a uniform random variable over $\mathcal{V}$ and $S \mid V \sim \mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}$. We shall let Q denote the joint distribution of (V, S) and let Q_S denote the marginal over S.

With this construction in place, we first show that the minimax excess risk is lower bounded by

Lemma B.1. *For any positive integers $K, n_{\mathsf{maj}}, n_{\mathsf{min}}$, the minimax excess risk is lower bounded as follows:*

$$\text{Minimax Excess Risk}(\mathcal{P}_{\mathsf{LS}})$$

$$= \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \in \mathcal{P}_{\mathsf{LS}}} \mathbb{E}_{S \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R(\mathcal{A}^S; \mathsf{P}_{\mathsf{test}}) - R(f^\star; \mathsf{P}_{\mathsf{test}})\right]$$

$$\geq \frac{1}{36K} - \frac{1}{2}\mathbb{E}_{S \sim \mathsf{Q}_S}\left[\text{TV}\left(\sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,1}, \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,-1}\right)\right]. \tag{7}$$

Proof. By invoking Lemma A.1 we get that

$$\text{Minimax Excess Risk}(\mathcal{P}_{\mathsf{LS}})$$

$$\geq \underbrace{\mathbb{E}_{S \sim \mathsf{Q}_S}\left[\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} \mathsf{Q}(v|S)\mathsf{P}_{v,\mathsf{test}}}(h(x) \neq y)\right]}_{=\mathfrak{R}_{\mathcal{V}}} - \underbrace{\mathbb{E}_V\left[R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}}))\right]}_{=\mathfrak{B}_{\mathcal{V}}}.$$

We proceed by calculating alternate expressions for $\mathfrak{R}_{\mathcal{V}}$ and $\mathfrak{B}_{\mathcal{V}}$ to get our desired lower bound on the minimax excess risk.

Calculation of $\mathfrak{R}_{\mathcal{V}}$: Immediately by Le Cam's lemma [22, Eq. 15.13], we get that

$$\mathfrak{R}_{\mathcal{V}} = \mathbb{E}_{S \sim \mathsf{Q}_S}\left[\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} \mathsf{Q}(v|S)\mathsf{P}_{v,\mathsf{test}}}(h(x) \neq y)\right]$$

$$= \frac{1}{2}\mathbb{E}_{S \sim \mathsf{Q}_S}\left[1 - \text{TV}\left(\sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,1}, \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,-1}\right)\right]. \tag{8}$$

Calculation of $\mathfrak{B}_\mathcal{V}$: Again by invoking Le Cam's lemma [22, Eq. 15.13], we get that for any class conditional distributions $\mathsf{P}_1, \mathsf{P}_{-1}$,

$$R(f^\star; \mathsf{P}_{\mathsf{test}}) = \frac{1}{2} - \frac{1}{2}\mathrm{TV}(\mathsf{P}_1, \mathsf{P}_{-1}).$$

So by taking expectations, we get that

$$\mathfrak{B}_\mathcal{V} = \mathbb{E}_V[R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}})] = \mathbb{E}_V\left[\frac{1}{2} - \frac{1}{2}\mathrm{TV}(\mathsf{P}_{V,1}, \mathsf{P}_{V,-1})\right]. \tag{9}$$

We now compute $\mathbb{E}_V[\mathrm{TV}(\mathsf{P}_{V,1}, \mathsf{P}_{V,-1})]$ as follows:

$$\begin{aligned}
\mathbb{E}_V[\mathrm{TV}(\mathsf{P}_{V,1}, \mathsf{P}_{V,-1})] &= \frac{1}{2}\mathbb{E}_V\left[\int_{x=0}^1 |\mathsf{P}_{V,1}(x) - \mathsf{P}_{V,-1}(x)| \; \mathrm{d}x\right] \\
&= \frac{1}{2}\mathbb{E}_V\left[\sum_{j=1}^K \int_{\frac{j-1}{K}}^{\frac{j}{K}} |V_{1,j} - V_{-1,j}|\left|\phi\left(x - \frac{j+1/2}{K}\right)\right| \; \mathrm{d}x\right] \\
&= \frac{1}{2}\sum_{j=1}^K \mathbb{E}_V\left[\int_{\frac{j-1}{K}}^{\frac{j}{K}} |V_{1,j} - V_{-1,j}|\left|\phi\left(x - \frac{j+1/2}{K}\right)\right| \; \mathrm{d}x\right] \\
&\stackrel{(i)}{=} \frac{1}{16K^2}\sum_{j=1}^K \mathbb{E}_V[|V_{1,j} - V_{-1,j}|],
\end{aligned}$$

where (i) follows by Lemma A.2. Observe that $V_{1,j}, V_{-1,j}$ are independent uniform random variables on $\{-1, 0, 1\}$, it is therefore straightforward to compute that

$$\mathbb{E}_V[|V_{1,j} - V_{-1,j}|] = \frac{8}{9}.$$

This yields that

$$\mathbb{E}_V\left[\mathrm{TV}(\mathsf{P}_{V,1}, \mathsf{P}_{V,-1})\right] = \frac{1}{18K}.$$

Plugging this into Eq. (9) allows us to conclude that

$$\mathfrak{B}_\mathcal{V} = \mathbb{E}_V[R(f^\star(\mathsf{P}_{V,\mathsf{test}}); \mathsf{P}_{V,\mathsf{test}})] = \frac{1}{2}\left(1 - \frac{1}{18K}\right). \tag{10}$$

Combining Eqs. (8) and (10) establishes the claimed result.

$\square$

In light of this previous lemma we now aim to upper bound the expected total variation distance in Eq. (7).

Lemma B.2. *Suppose that v is drawn uniformly from the set $\{-1, 1\}^K$, and that $S \mid v$ is drawn from $\mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}$ then,*

$$\mathbb{E}_S\left[\mathrm{TV}\left(\sum_{v\in\mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,1}, \sum_{v\in\mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,-1}\right)\right] \leq \frac{1}{18K} - \frac{1}{144K}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right).$$

484 *Proof.* Let $\psi := \mathbb{E}_S \left[\text{TV} \left(\sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \mathsf{P}_{v,1}, \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \mathsf{P}_{v,-1} \right) \right]$. Then,

$$
\begin{aligned}
\psi &= \mathbb{E}_S \left[\text{TV} \left(\sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \mathsf{P}_{v,1}, \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \mathsf{P}_{v,-1} \right) \right] \\
&= \frac{1}{2} \mathbb{E}_S \left[\int_{x=0}^1 \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \left(\mathsf{P}_{v,1}(x) - \mathsf{P}_{v,-1}(x) \right) \right| \, dx \right] \\
&= \frac{1}{2} \mathbb{E}_S \left[\sum_{j=1}^K \int_{x=\frac{j-1}{K}}^{\frac{j}{K}} \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \left(\mathsf{P}_{v,1}(x) - \mathsf{P}_{v,-1}(x) \right) \right| \, dx \right] \\
&= \frac{1}{2} \mathbb{E}_S \left[\sum_{j=1}^K \int_{x=\frac{j-1}{K}}^{\frac{j}{K}} \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) (v_{1,j} - v_{-1,j}) \phi \left(x - \frac{j+1/2}{K} \right) \right| \, dx \right],
\end{aligned}
$$

485 where the last equality is by the definition of $\mathsf{P}_{v,1}$ and $\mathsf{P}_{v,-1}$. Continuing we get that,

$$
\begin{aligned}
\psi &= \frac{1}{2} \left[\int_{x=\frac{j-1}{K}}^{\frac{j}{K}} \left| \phi \left(x - \frac{j+1/2}{K} \right) \right| \, dx \right] \mathbb{E}_S \left[\sum_{j=1}^K \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) (v_{1,j} - v_{-1,j}) \right| \right] \\
&\overset{(i)}{=} \frac{1}{16K^2} \mathbb{E}_S \left[\sum_{j=1}^K \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) (v_{1,j} - v_{-1,j}) \right| \right] \\
&= \frac{1}{16K^2} \sum_{j=1}^K \int \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) (v_{1,j} - v_{-1,j}) \right| \, d\mathsf{Q}_S(S) \\
&= \frac{1}{16K^2} \sum_{j=1}^K \int \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(v, S) (v_{1,j} - v_{-1,j}) \right| \, dS \\
&\overset{(i)}{=} \frac{1}{16K^2 |\mathcal{V}|} \sum_{j=1}^K \int \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(S \mid v) (v_{1,j} - v_{-1,j}) \right| \, dS,
\end{aligned}
$$

486 where (i) follows by the calculation in Lemma A.2 and (ii) follows since v is a uniform random
487 variable over the set $\mathcal{V}$.

488 The distributions $\mathsf{P}_{v,1}$ and $\mathsf{P}_{v,-1}$ are symmetrically defined over all intervals $I_j = [\frac{j-1}{K}, \frac{j}{K}]$, and
489 hence all of the summands in the RHS above are equal. Thus,

$$
\psi = \frac{1}{16K |\mathcal{V}|} \int \left| \sum_{v \in \mathcal{V}} \mathsf{Q}(S \mid v) (v_{1,1} - v_{-1,1}) \right| \, dS. \tag{11}
$$

490 Before we continue further, let us define

$$
\mathcal{V}^+ = \{ v \in \mathcal{V} \mid v_{1,1} > v_{-1,1} \}.
$$

For every $v \in \mathcal{V}^+$, let $\tilde{v} \in \mathcal{V}$ be such that is the same as v on all coordinates, except $\tilde{v}_{1,1} = -v_{1,1}$ and $\tilde{v}_{-1,1} = -v_{-1,1}$. Then continuing from Eq. (11) we find that,

$$\psi \overset{(i)}{=} \frac{1}{16K|\mathcal{V}|} \int \left| \sum_{v \in \mathcal{V}^+} (v_{1,1} - v_{-1,1})(\mathsf{Q}(S \mid v) - \mathsf{Q}(S \mid \tilde{v})) \right| \mathrm{d}S$$

$$\overset{(ii)}{\leq} \frac{1}{16K|\mathcal{V}|} \int \sum_{v \in \mathcal{V}^+} (v_{1,1} - v_{-1,1}) \left| \mathsf{Q}(S \mid v) - \mathsf{Q}(S \mid \tilde{v}) \right| \mathrm{d}S$$

$$= \frac{1}{16K|\mathcal{V}|} \sum_{v \in \mathcal{V}^+} (v_{1,1} - v_{-1,1}) \int \left| \mathsf{Q}(S \mid v) - \mathsf{Q}(S \mid \tilde{v}) \right| \mathrm{d}S$$

$$= \frac{1}{8K|\mathcal{V}|} \underbrace{\sum_{v \in \mathcal{V}^+} (v_{1,1} - v_{-1,1}) \mathrm{TV}(\mathsf{Q}(S \mid v), \mathsf{Q}(S \mid \tilde{v}))}_{=: \Xi}, \tag{12}$$

where (i) we use the definition of $\mathcal{V}^+$ and $\tilde{v}$, (ii) follows since $v_{1,1} > v_{-1,1}$ for $v \in \mathcal{V}^+$.

Now we further partition $\mathcal{V}^+$ into 3 sets $\mathcal{V}^{(1,0)}, \mathcal{V}^{(0,-1)}, \mathcal{V}^{(1,-1)}$ as follows

$$\mathcal{V}^{(1,0)} = \{v \in \mathcal{V} \mid v_{1,1} = 1, v_{-1,1} = 0\},$$
$$\mathcal{V}^{(0,-1)} = \{v \in \mathcal{V} \mid v_{1,1} = 0, v_{-1,1} = -1\},$$
$$\mathcal{V}^{(1,-1)} = \{v \in \mathcal{V} \mid v_{1,1} = 1, v_{-1,1} = -1\}.$$

Note that $\mathsf{Q}(S \mid v) = \mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}$, and therefore

$$\Xi = \sum_{v \in \mathcal{V}^+} (v_{1,1} - v_{-1,1}) \mathrm{TV}\left(\mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \right)$$

$$\overset{(i)}{=} \sum_{v \in \mathcal{V}^{(1,0)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \right)$$

$$+ \sum_{v \in \mathcal{V}^{(0,-1)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \right)$$

$$+ 2 \sum_{v \in \mathcal{V}^{(1,-1)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \right), \tag{13}$$

where (i) follows since $v_1, v_{-1} \in \{-1, 0, 1\}^K$ and by the definition of the sets $\mathcal{V}^{(1,0)}, \mathcal{V}^{(0,1)}$ and $\mathcal{V}^{(1,-1)}$.

Now by the Bretagnolle–Huber inequality [see 4, Corollary 4],

$$\mathrm{TV}\left(\mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \right) = \mathrm{TV}\left(\mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}}, \mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}} \right)$$

$$\leq 1 - \frac{1}{2} \exp\left(-\mathrm{KL}\left(\mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \| \mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}} \right) \right),$$

where we flip the arguments in the first step for simplicity later.

Next, by the chain rule for KL-divergence, we have that

$$\mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{\tilde{v},\mathrm{min}}^{n_{\mathrm{min}}} \| \mathsf{P}_{v,\mathrm{maj}}^{n_{\mathrm{maj}}} \times \mathsf{P}_{v,\mathrm{min}}^{n_{\mathrm{min}}}) = n_{\mathrm{maj}} \mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{maj}} \| \mathsf{P}_{v,\mathrm{maj}}) + n_{\mathrm{min}} \mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{min}} \| \mathsf{P}_{v,\mathrm{min}}).$$

Using these, let us upper bound the first term in Eq. (13) corresponding to $v \in \mathcal{V}^{(0,-1)}$. For $v \in \mathcal{V}^{(0,-1)}$, notice that $\mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{maj}} \| \mathsf{P}_{v,\mathrm{maj}}) = 0$ since $v_{1,j} = \tilde{v}_{1,j}$ for all $j \in \{1, \ldots, K\}$. For the second term, $\mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{min}} \| \mathsf{P}_{v,\mathrm{min}})$, only $v_{1,1}$ and $\tilde{v}_{1,1}$ differ, so

$$\mathrm{KL}(\mathsf{P}_{\tilde{v},\mathrm{min}} \| \mathsf{P}_{v,\mathrm{min}}) = \int_0^1 \mathsf{P}_{v,-1}(x) \log\left(\frac{\mathsf{P}_{v,-1}(x)}{\mathsf{P}_{\tilde{v},-1}(x)} \right) \mathrm{d}x$$

$$= \int_0^{\frac{1}{K}} \log\left(\frac{1 + \phi_K(x - \frac{1}{2K})}{1 - \phi_K(x - \frac{1}{2K})} \right) \left(1 + \phi_K\left(x - \frac{1}{2K} \right) \right) \mathrm{d}x$$

$$\leq \frac{1}{3K^3},$$

504 where the last inequality is a result of the calculation in Lemma A.3.

505 Therefore, we get

$$\sum_{v \in \mathcal{V}^{(0,-1)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}, \mathsf{P}_{\tilde{v},\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\tilde{v},\mathsf{min}}^{n_{\mathsf{min}}}\right) \le 9^{K-1}\left(1 - \frac{1}{2}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right)\right).$$

506 For the terms in Eq. (13) corresponding to $\mathcal{V}^{(0,-1)}, \mathcal{V}^{(1,-1)}$, we simply take the trivial bound to get

$$\sum_{v \in \mathcal{V}^{(0,-1)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}, \mathsf{P}_{\tilde{v},\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\tilde{v},\mathsf{min}}^{n_{\mathsf{min}}}\right) \le 9^{K-1},$$

$$\sum_{v \in \mathcal{V}^{(1,-1)}} \mathrm{TV}\left(\mathsf{P}_{v,\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{v,\mathsf{min}}^{n_{\mathsf{min}}}, \mathsf{P}_{\tilde{v},\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\tilde{v},\mathsf{min}}^{n_{\mathsf{min}}}\right) \le 9^{K-1}.$$

507 Plugging these bounds into Eq. (13) we get that,

$$\Xi \le 4 \cdot 9^{K-1} - \frac{9^{K-1}}{2}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right).$$

508 Now using this bound on Ξ in Eq. (12) and observing that $|\mathcal{V}| = 9^K$, we get that,

$$
\begin{aligned}
\psi &= \mathbb{E}_S\left[\mathrm{TV}\left(\sum_{v \in \mathcal{V}} Q(v \mid S)P_{v,1}, \sum_{v \in \mathcal{V}} Q(v \mid S)P_{v,-1}\right)\right]\\
&\le \frac{1}{8 \cdot 9^K K}\left(4 \cdot 9^{K-1} - \frac{9^{K-1}}{2}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right)\right)\\
&= \frac{1}{18K} - \frac{1}{144K}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right),
\end{aligned}
$$

509 completing the proof. $\qquad\square$

510 Finally, we combine Lemma B.1 and Lemma B.2 to establish the minimax lower bound in this label
511 shift setting. We recall the statement of the theorem here.

512 **Theorem 4.1.** *Consider the label shift setting described in Section 3.2.1. Recall that $\mathcal{P}_{\mathsf{LS}}$ is the class*
513 *of pairs of distributions $(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}})$ that satisfy the assumptions in that section. The minimax excess*
514 *risk over this class is lower bounded as follows:*

$$\text{Minimax Excess Risk}(\mathcal{P}_{\mathsf{LS}}) = \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\mathsf{maj}},\mathsf{P}_{\mathsf{min}}) \in \mathcal{P}_{\mathsf{LS}}} \text{Excess Risk}[\mathcal{A};(\mathsf{P}_{\mathsf{maj}},\mathsf{P}_{\mathsf{min}})] \ge \frac{c}{n_{\mathsf{min}}^{1/3}}. \quad (3)$$

515 *Proof.* By Lemma B.1 we know that,

$$\text{Minimax Excess Risk}(\mathcal{P}_{\mathsf{LS}}) \ge \frac{1}{36K} - \frac{1}{2}\mathbb{E}_{S \sim \mathsf{Q}_S}\left[\mathrm{TV}\left(\sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,1}, \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S)\mathsf{P}_{v,-1}\right)\right].$$

516 Next by the calculation in Lemma B.2 we have that

$$
\begin{aligned}
\text{Minimax Excess Risk}(\mathcal{P}_{\mathsf{LS}}) &\ge \frac{1}{36K} - \frac{1}{2}\left(\frac{1}{18K} - \frac{1}{144K}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right)\right)\\
&= \frac{1}{288K}\exp\left(-\frac{n_{\mathsf{min}}}{3K^3}\right).
\end{aligned}
$$

517 Setting $K = \lceil n_{\mathsf{min}}^{1/3}\rceil$ yields the result. $\qquad\square$

B.2 Proof of Theorem 5.1

519 In this section, we derive an upper bound on the excess risk of the undersampled binning estimator
520 $\mathcal{A}_{\mathsf{USB}}$ (Eq. (5)) in the label shift setting. Recall that given a dataset $\mathcal{S}$ this estimator first calculates
521 the undersampled dataset $\mathcal{S}_{\mathsf{US}}$, where the number of points from the minority group (n_{min}) is equal to

522 the number of points from the majority group ($n_{\min}$), and the size of the dataset is $2n_{\min}$. Throughout
523 this section, $(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}})$ shall be an arbitrary element of $\mathcal{P}_{\mathsf{LS}}$.

524 To bound the excess risk of the undersampling algorithm, we will relate it to density estimation.

525 Recall that $n_{1,j}$ denotes the number of points in $\mathcal{S}_{\mathsf{US}}$ with label $+1$ that lie in I_j, and $n_{-1,j}$ is defined
526 analogously.

527 Given a positive integer K, for $x \in I_j = [\frac{j-1}{K}, \frac{j}{K}]$, by the definition of the undersampled binning
528 estimator (Eq. (5))

$$
\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}(x) = \begin{cases} 1 & \text{if } n_{1,j} > n_{-1,j}, \\ -1 & \text{otherwise.} \end{cases}
$$

529 Recall that since we have undersampled, $\sum_j n_{1,j} = \sum_j n_{-1,j} = n_{\min}$. Therefore, define the simple
530 histogram estimators for $\mathsf{P}_1(x) = \mathsf{P}(x \mid y = 1)$ and $\mathsf{P}_{-1}(x) = \mathsf{P}(x \mid y = -1)$ as follows: for
531 $x \in I_j$,

$$
\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) := \frac{n_{1,j}}{K n_{\min}} \quad \text{and} \quad \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x) := \frac{n_{-1,j}}{K n_{\min}}.
$$

532 With this histogram estimator in place, we may define an estimator for $\eta(x) := \mathsf{P}_{\mathsf{test}}(y = 1|x)$ as
533 follows,

$$
\widehat{\eta}^{\mathcal{S}}(x) := \frac{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x)}{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x)}.
$$

534 Observe that, for $x \in I_j$

$$
\widehat{\eta}^{\mathcal{S}}(x) > 1/2 \iff n_{1,j} > n_{-1,j} \iff \mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}(x) = 1.
$$

535 Defining an estimator $\widehat{\eta}^{\mathcal{S}}$ for the $\mathsf{P}_{\mathsf{test}}(y = 1 \mid x)$ in this way will allow us to relate the excess risk of
536 $\mathcal{A}_{\mathsf{USB}}$ to the estimation error in $\widehat{\mathsf{P}}_1^{\mathcal{S}}$ and $\widehat{\mathsf{P}}_{-1}^{\mathcal{S}}$.

537 Before proving the theorem we restate it here.

538 **Theorem 5.1.** *Consider the label shift setting described in Section 3.2.1. For any $(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \in \mathcal{P}_{\mathsf{LS}}$*
539 *the expected excess risk of the Undersampling Binning Estimator (Eq. (5)) with number of bins with*
540 $K = c\lceil n_{\min}^{1/3} \rceil$ *is upper bounded by*

$$
\mathsf{Excess\ Risk}[\mathcal{A}_{\mathsf{USB}}; (\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}})] = \mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}; \mathsf{P}_{\mathsf{test}}) - R(f^\star; \mathsf{P}_{\mathsf{test}}) \right] \leq \frac{C}{n_{\min}^{1/3}}.
$$

541 *Proof.* By the definition of the excess risk

$$
\mathsf{Excess\ Risk}[\mathcal{A}_{\mathsf{USB}}; (\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}})] := \mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}; \mathsf{P}_{\mathsf{test}})) - R(f^\star; \mathsf{P}_{\mathsf{test}}) \right].
$$

542 By invoking [25, Theorem 1] we may upper bound the excess risk given a draw of $\mathcal{S}$ by

$$
R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}; \mathsf{P}_{\mathsf{test}})) - R(f^\star; \mathsf{P}_{\mathsf{test}}) \leq 2 \int \left| \widehat{\eta}^{\mathcal{S}}(x) - \eta(x) \right| \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x.
$$

543 Continuing using the definition of $\widehat{\eta}^{\mathcal{S}}$ above and because $\eta = \mathsf{P}_1/(\mathsf{P}_1 + \mathsf{P}_{-1})$ we have that,

$$R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}; \mathsf{P}_{\mathsf{test}})) - R(f^\star; \mathsf{P}_{\mathsf{test}})$$

$$= 2 \int_0^1 \left| \frac{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x)}{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x)} - \frac{\mathsf{P}_1(x)}{\mathsf{P}_1(x) + \mathsf{P}_{-1}(x)} \right| \left(\frac{\mathsf{P}_1(x) + \mathsf{P}_{-1}(x)}{2} \right) \, \mathrm{d}x$$

$$= \int_0^1 \left| \left(\frac{\mathsf{P}_1(x) + \mathsf{P}_{-1}(x)}{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x)} \right) \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right| \, \mathrm{d}x$$

$$\overset{(i)}{\leq} \int_0^1 \left| \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right| \, \mathrm{d}x + \int_0^1 \left| \frac{\mathsf{P}_1(x) + \mathsf{P}_{-1}(x)}{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x)} - 1 \right| \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) \, \mathrm{d}x$$

$$= \int_0^1 \left| \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right| \, \mathrm{d}x + \int_0^1 \left| \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x) - \mathsf{P}_1(x) - \mathsf{P}_{-1}(x) \right| \frac{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x)}{\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) + \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x)} \, \mathrm{d}x$$

$$\leq 2 \int_0^1 \left| \widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right| \, \mathrm{d}x + \int_0^1 \left| \widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x) - \mathsf{P}_{-1}(x) \right| \, \mathrm{d}x$$

$$\overset{(ii)}{\leq} 2 \sqrt{\int_0^1 \left(\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right)^2 \, \mathrm{d}x} + \sqrt{\int_0^1 \left(\widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x) - \mathsf{P}_{-1}(x) \right)^2 \, \mathrm{d}x},$$

544 where (i) follows by the triangle inequality, (ii) is by the Cauchy–Schwarz inequality.

545 Taking expectation over the samples $\mathcal{S}$ and by invoking Jensen's inequality we find that,

$$\mathsf{Excess\ Risk}(\mathcal{A}^{\mathcal{S}}; (\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}))$$

$$= \mathbb{E}_{\mathcal{S}} \left[R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}; \mathsf{P}_{\mathsf{test}})) - R(f^\star; \mathsf{P}_{\mathsf{test}}) \right]$$

$$\leq 2 \sqrt{\mathbb{E}_{\mathcal{S}} \left[\int \left(\widehat{\mathsf{P}}_1^{\mathcal{S}}(x) - \mathsf{P}_1(x) \right)^2 \, \mathrm{d}x \right]} + \sqrt{\mathbb{E}_{\mathcal{S}} \left[\int \left(\widehat{\mathsf{P}}_{-1}^{\mathcal{S}}(x) - \mathsf{P}_{-1}(x) \right)^2 \, \mathrm{d}x \right]}.$$

546 We note that $\widehat{\mathsf{P}}_j^{\mathcal{S}}$ only depends on n_{min} i.i.d. draws from class j. Thus by [9, Theorem 1.7], if
547 $K = c\lceil n_{\mathsf{min}} \rceil^{1/3}$ then

$$\mathbb{E}_{\mathcal{S}} \left[\int \left(\widehat{\mathsf{P}}_j^{\mathcal{S}}(x) - \mathsf{P}_j(x) \right)^2 \, \mathrm{d}x \right] \leq \frac{C}{n_{\mathsf{min}}^{2/3}}.$$

548 Plugging this into the previous inequality yields the desired result. $\qquad\square$

C Proof in the Group-Covariate Shift Setting

550 Throughout this section we operate in the group-covariate shift setting (Section 3.2.2).

551 First in Appendix C.1, we prove Theorem 4.2, the minimax lower bound through a sequence of
552 lemmas. Second in Appendix C.2, we prove Theorem 5.2 that upper bound on the excess risk of the
553 undersampled binning estimator with $\lceil n_{\mathsf{min}} \rceil^{1/3}$ bins.

C.1 Proof of Theorem 4.2

555 In this section, we provide a proof of the minimax lower bound in the group shift setting.

556 We construct the "hard" set of distributions as follows. Let the index set be $\mathcal{V} = \{-1, 1\}^K$. For every
557 $v \in \mathcal{V}$ define a distribution as follows: for $x \in I_j = [\frac{j-1}{K}, \frac{j}{K}]$,

$$\mathsf{P}_v(y = 1 \mid x) := \frac{1}{2} \left[1 + v_j \phi \left(x - \frac{j + 1/2}{K} \right) \right],$$

558 where ϕ is defined in Eq. 6. Given a $\tau \in [0, 1]$ we also construct the group distributions as follows:

$$\mathsf{P}_a(x) = \begin{cases} 2 - \tau & \text{if } x \in [0, 0.5) \\ \tau & \text{if } x \in [0.5, 1], \end{cases}$$

559 and let

$$P_b(x) = 2 - P_a(x).$$

560 We can verify that

$$\text{Overlap}(P_a, P_b) = 1 - \text{TV}(P_a, P_b) = 1 - \frac{1}{2}\int_{x=0}^{1} |P_a(x) - P_b(x)| \, dx = \tau.$$

561 We continue to define

$$P_{v,\text{maj}}(x, y) = P_v(y \mid x)P_a(x)$$
$$P_{v,\text{min}}(x, y) = P_v(y \mid x)P_b(x),$$

562 and

$$P_{v,\text{test}}(x, y) = P_v(y \mid x)\left(\frac{P_a(x) + P_b(x)}{2}\right).$$

563 Observe that $(P_a(x) + P_b(x))/2 = 1$, the uniform distribution over $[0, 1]$.

564 Recall that as described in Section A.1, V shall be a uniform random variable over $\mathcal{V}$ and $S \mid V \sim$
565 $P_{v,\text{maj}}^{n_{\text{maj}}} \times P_{v,\text{min}}^{n_{\text{min}}}$. We shall let Q denote the joint distribution of (V, S) and let Q_S denote the marginal
566 over S.

567 With this construction in place, we present the following lemma that lower bounds the minimax
568 excess risk by a sum of $\exp(-\text{KL}(Q(S \mid v_j = 1)\|Q(S \mid v_j = -1))$ over the intervals. Intuitively,
569 $\text{KL}(Q(S \mid v_j = 1)\|Q(S \mid v_j = -1)$ is a measure of how difficult it is to identify whether $v_j = 1$ or
570 $v_j = -1$ from the samples.

571 **Lemma C.1.** *For any positive integers $K, n_{\text{maj}}, n_{\text{min}}$ and $\tau \in [0, 1]$, the minimax excess risk is lower*
572 *bounded as follows:*

$$\text{Minimax Excess Risk}(\mathcal{P}_{\text{GS}}(\tau)) = \inf_{\mathcal{A}} \sup_{(P_{\text{maj}}, P_{\text{min}}) \in \mathcal{P}_{\text{GS}}(\tau)} \mathbb{E}_{S \sim P_{\text{maj}}^{n_{\text{maj}}} \times P_{\text{min}}^{n_{\text{min}}}} \left[R(\mathcal{A}^S; P_{\text{test}}) - R(f^\star; P_{\text{test}})\right]$$

$$\geq \frac{1}{32K^2} \sum_{j=1}^{K} \exp(-\text{KL}(Q(S \mid v_j = 1)\|Q(S \mid v_j = -1))).$$

573 *Proof.* By invoking Lemma A.1, we know that the minimax excess risk is lower bounded by

$$\text{Minimax Excess Risk}(\mathcal{P}_{\text{GS}}(\tau))$$

$$\geq \underbrace{\mathbb{E}_{S \sim Q_S}[\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} Q(v|S)P_{v,\text{test}}}(h(x) \neq y)]}_{=\mathfrak{R}_{\mathcal{V}}} - \underbrace{\mathbb{E}_V[R(f^\star(P_{V,\text{test}}); P_{V,\text{test}})]}_{=\mathfrak{B}_{\mathcal{V}}},$$

574 where V is a uniform random variable over the set $\mathcal{V}$, $S \mid V = v$ is a draw from $P_{v,\text{maj}}^{n_{\text{maj}}} \times P_{v,\text{min}}^{n_{\text{min}}}$, and
575 Q denotes the joint distribution over (V, S).

576 We shall lower bound this minimax risk in parts. First, we shall establish a lower bound on $\mathfrak{R}_{\mathcal{V}}$, and
577 then an upper bound on the Bayes risk $\mathfrak{B}_{\mathcal{V}}$.

578 **Lower bound on $\mathfrak{R}_{\mathcal{V}}$.** Unpacking $\mathfrak{R}_{\mathcal{V}}$ using its definition we get that,

$$\mathfrak{R}_{\mathcal{V}} = \mathbb{E}_{S \sim Q_S}[\inf_h \mathbb{P}_{(x,y) \sim \sum_{v \in \mathcal{V}} Q(v|S)P_{v,\text{test}}}(h(x) \neq y)]$$

$$= \mathbb{E}_{S \sim Q_S}\left[\inf_h \int_0^1 P_{\text{test}}(x)\mathbb{P}_{y \sim \sum_{v \in \mathcal{V}} Q(v|S)P_v(\cdot|x)}[h(x) \neq y] \, dx\right]$$

$$\overset{(i)}{=} \mathbb{E}_{S \sim Q_S}\left[\int_0^1 P_{\text{test}}(x)\min\left\{\sum_{v \in \mathcal{V}} Q(v \mid S)P_v(1 \mid x), \sum_{v \in \mathcal{V}} Q(v \mid S)P_v(-1 \mid x)\right\} \, dx\right]$$

$$\overset{(ii)}{=} \frac{1}{2} - \mathbb{E}_{S \sim Q_S}\left[\int_0^1 P_{\text{test}}(x)\left|\frac{1}{2} - \sum_{v \in \mathcal{V}} Q(v \mid S)P_v(1 \mid x)\right| \, dx\right]$$

$$\overset{(iii)}{=} \frac{1}{2} - \int_0^1 P_{\text{test}}(x)\mathbb{E}_{S \sim Q_S}\left[\left|\frac{1}{2} - \sum_{v \in \mathcal{V}} Q(v \mid S)P_v(1 \mid x)\right|\right] \, dx, \tag{14}$$

where (i) follows by taking h to be the pointwise minimizer over x, (ii) follows since $\mathsf{P}_v(-1 \mid x) = 1 - \mathsf{P}_v(1 \mid x)$ and $\min\{s, 1-s\} = (1 - |1-2s|)/2$ for all $s \in [0,1]$, and (iii) follows by Fubini's theorem which allows us to switch the order of the integrals.

If $x \in I_j = [\frac{j-1}{K}, \frac{j}{K}]$ for some $j \in \{1, \dots, K\}$ we let j_x denote the value of this index j. With this notation in place let us continue to upper bound integrand in the second term in the RHS above as follows:

$$\mathbb{E}_{S \sim \mathsf{Q}_S} \left[\left| \frac{1}{2} - \sum_{v \in \mathcal{V}} \mathsf{Q}(v \mid S) \mathsf{P}_v(1 \mid x) \right| \right]$$

$$\stackrel{(i)}{=} \mathbb{E}_{S \sim \mathsf{Q}_S} \left[\left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| |\mathsf{Q}(v_{j_x} = 1 \mid S) - \mathsf{Q}(v_{j_x} = -1 \mid S)| \right]$$

$$= \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \mathbb{E}_{S \sim \mathsf{Q}_S} [|\mathsf{Q}(v_{j_x} = 1 \mid S) - \mathsf{Q}(v_{j_x} = -1 \mid S)|]$$

$$\stackrel{(ii)}{=} \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \mathbb{E}_{S \sim \mathsf{Q}_S} \left[\left| \frac{\mathsf{Q}(S \mid v_{j_x} = 1) \mathsf{Q}_V(v_{j_x} = 1)}{\mathsf{Q}_S(S)} - \frac{\mathsf{Q}(S \mid v_{j_x} = -1) \mathsf{Q}_V(v_{j_x} = -1)}{\mathsf{Q}_S(S)} \right| \right]$$

$$\stackrel{(iii)}{=} \frac{1}{2} \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \mathrm{TV}(\mathsf{Q}(S \mid v_{j_x} = 1), \mathsf{Q}(S \mid v_{j_x} = -1)), \tag{15}$$

where (i) follows since $\mathsf{P}_v(1 \mid x) = (1 + v_{j_x} \phi(x - (j_x + 1/2)/K))/2$ and by marginalizing $\mathsf{Q}(v \mid S)$ over the indices $j \neq j_x$, (ii) follows by using Bayes' rule and (iii) follows since the total-variation distance is half the ℓ_1 distance. Now by the Bretagnolle–Huber inequality [see 4, Corollary 4] we get that,

$$\mathrm{TV}(\mathsf{Q}(S \mid v_{j_x} = 1), \mathsf{Q}(S \mid v_{j_x} = -1))$$
$$\leq 1 - \frac{\exp(-\mathrm{KL}(\mathsf{Q}(S \mid v_{j_x} = 1) \| \mathsf{Q}(S \mid v_{j_x} = -1)))}{2}. \tag{16}$$

Combining Eqs. (14)-(16) we get that

$$\mathfrak{R}_{\mathcal{V}}$$
$$\geq \frac{1}{2} - \frac{1}{2} \int_0^1 \mathsf{P}_{\text{test}}(x) \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \, \mathrm{d}x$$
$$+ \frac{1}{4} \int_0^1 \mathsf{P}_{\text{test}}(x) \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \exp(-\mathrm{KL}(\mathsf{Q}(S \mid v_{j_x} = 1) \| \mathsf{Q}(S \mid v_{j_x} = -1))) \, \mathrm{d}x. \tag{17}$$

Upper bound on $\mathfrak{B}_{\mathcal{V}}$: The Bayes error is
$$\mathfrak{B}_{\mathcal{V}} = \mathbb{E}_V[R(f^\star(\mathsf{P}_V); \mathsf{P}_V)]$$
$$= \mathbb{E}_V \left[\inf_f \mathbb{E}_{(x,y) \sim \mathsf{P}_{v,\text{test}}} \mathbf{1}(f(x) \neq y) \right]$$
$$= \mathbb{E}_V \left[\inf_f \int_{x=0}^1 \sum_{y \in \{-1,1\}} \mathsf{P}_{\text{test}}(x) \mathsf{P}_{V,\text{test}}(y \mid x) \mathbf{1}(f(x) = -y) \right]$$
$$= \mathbb{E}_V \left[\int_{x=0}^1 \mathsf{P}_{\text{test}}(x) \min_{y \in \{-1,1\}} \mathsf{P}_{V,\text{test}}(y \mid x) \right]$$
$$\stackrel{(i)}{=} \mathbb{E}_V \left[\frac{1}{2} \left(1 - \int_{x=0}^1 \mathsf{P}_{\text{test}}(x) |\mathsf{P}_{V,\text{test}}(1 \mid x) - \mathsf{P}_{V,\text{test}}(-1 \mid x)| \, \mathrm{d}x \right) \right]$$
$$\stackrel{(ii)}{=} \mathbb{E}_V \left[\frac{1}{2} \left(1 - \int_{x=0}^1 \mathsf{P}_{\text{test}}(x) \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \, \mathrm{d}x \right) \right]$$
$$= \frac{1}{2} - \frac{1}{2} \int_{x=0}^1 \mathsf{P}_{\text{test}}(x) \left| \phi \left(x - \frac{j_x + 1/2}{K} \right) \right| \, \mathrm{d}x, \tag{18}$$

where (i) follows since $\mathsf{P}_v(1 \mid x) = 1 - \mathsf{P}_v(-1 \mid x)$ and $\min\{s, 1-s\} = (1 - |1-2s|)/2$ for all $s \in [0,1]$, and (ii) follows by our construction of P_v above along with the fact that $\mathsf{P}_v(1 \mid x) = 1 - \mathsf{P}_v(-1 \mid x)$.

 Putting things together: Combining Eqs. (17) and (18) allows us to conclude that

Minimax Excess $\text{Risk}(\mathcal{P}_{\text{GS}}(\tau))$

$$\geq \frac{1}{4} \int_0^1 \mathsf{P}_{\text{test}}(x) \left| \phi\left(x - \frac{j_x + 1/2}{K}\right) \right| \exp(-\text{KL}(\mathsf{Q}(S \mid v_{j_x} = 1) \| \mathsf{Q}(S \mid v_{j_x} = -1))) \, \mathrm{d}x$$

$$= \frac{1}{4} \sum_{j=1}^K \int_{\frac{j-1}{K}}^{\frac{j}{K}} \mathsf{P}_{\text{test}}(x) \left| \phi\left(x - \frac{j + 1/2}{K}\right) \right| \exp(-\text{KL}(\mathsf{Q}(S \mid v_j = 1) \| \mathsf{Q}(S \mid v_j = -1))) \, \mathrm{d}x$$

$$= \frac{1}{4} \sum_{j=1}^K \exp(-\text{KL}(\mathsf{Q}(S \mid v_j = 1) \| \mathsf{Q}(S \mid v_j = -1))) \left[\int_{\frac{j-1}{K}}^{\frac{j}{K}} \mathsf{P}_{\text{test}}(x) \left| \phi\left(x - \frac{j + 1/2}{K}\right) \right| \, \mathrm{d}x \right]$$

$$\stackrel{(i)}{=} \frac{1}{32K^2} \sum_{j=1}^K \exp(-\text{KL}(\mathsf{Q}(S \mid v_j = 1) \| \mathsf{Q}(S \mid v_j = -1))),$$

where (i) follows by using Lemma A.2 along with the fact that $\mathsf{P}_{\text{test}}(x) = 1$ in our construction to show that the integral in the square brackets is equal to $1/8K^2$. This proves the result. $\square$

The next lemma upper bounds the KL divergence between $\mathsf{Q}(S \mid v_j = 1)$ and $\mathsf{Q}(S \mid v_j = -1)$ for each $j \in \{1, \ldots, K\}$. It shows that the KL divergence between these two posteriors is larger when the expected number of samples in that bin is larger.

Lemma C.2. *Suppose that v is drawn uniformly from the set $\{-1, 1\}^K$, and that $S \mid v$ is drawn from $\mathsf{P}_{v,\text{maj}}^{n_{\text{maj}}} \times \mathsf{P}_{v,\text{min}}^{n_{\text{min}}}$. Then for any $j \in \{1, \ldots, K/2\}$ and any $\tau \in [0, 1]$,*

$$\text{KL}(\mathsf{Q}(S \mid v_j = 1) \| \mathsf{Q}(S \mid v_j = -1)) \leq \frac{n_{\text{maj}}(2 - \tau) + n_{\text{min}}\tau}{3K^3},$$

and for any $j \in \{K/2 + 1, \ldots, K\}$

$$\text{KL}(\mathsf{Q}(S \mid v_j = 1) \| \mathsf{Q}(S \mid v_j = -1)) \leq \frac{n_{\text{maj}}\tau + n_{\text{min}}(2 - \tau)}{3K^3}.$$

Proof. Let us consider the case when $j = 1$. The bound for all other $j \in \{2, \ldots, K\}$ shall follow analogously.

Given samples S, let $S = (S_1, \bar{S}_1)$ be a partition where S_1 are the samples that fall in the interval I_1, and $\bar{S}_1$ be the other samples. Similarly, given a vector $v \in \{-1, 1\}$, let $v = (v_1, \bar{v}_1)$, where v_1 is the first component and $\bar{v}_1$ denotes the other components $(2, \ldots, K)$ of v.

First, we will show that

$$\mathsf{Q}(S \mid v_1) = \mathsf{Q}(S_1 \mid v_1)\mathsf{Q}(\bar{S}_1).$$

To see this, observe that

$$\mathsf{Q}(S \mid v_1) = \mathsf{Q}((S_1, \bar{S}_1) \mid v_1) = \mathsf{Q}(S_1 \mid v_1)\mathsf{Q}(\bar{S}_1 \mid v_1, S_1).$$

Further, if v is chosen uniformly over the hypercube $\{-1, 1\}^K$, then

$$\mathsf{Q}(\bar{S}_1 \mid v_1, S_1) = \sum_{\bar{v}_1} \mathsf{Q}(\bar{S}_1, \bar{v}_1 \mid v_1, S_1)$$

$$= \sum_{\bar{v}_1} \mathsf{Q}(\bar{S}_1 \mid v_1, \bar{v}_1, S_1)\mathsf{Q}(\bar{v}_1 \mid v_1, S_1)$$

$$\stackrel{(i)}{=} \sum_{\bar{v}_1} \mathsf{Q}(\bar{S}_1 \mid v_1, \bar{v}_1, S_1)\mathsf{Q}(\bar{v}_1)$$

$$\stackrel{(ii)}{=} \sum_{\bar{v}_1} \mathsf{Q}(\bar{S}_1 \mid v_1, \bar{v}_1)\mathsf{Q}(\bar{v}_1)$$

$$\stackrel{(iii)}{=} \sum_{\bar{v}_1} \mathsf{Q}(\bar{S}_1 \mid \bar{v}_1)\mathsf{Q}(\bar{v}_1)$$

$$= \mathsf{Q}(\bar{S}_1),$$

where (i) follows since by Bayes' rule

$$
\begin{aligned}
\mathsf{Q}(\bar{v}_1 \mid v_1, S_1) &= \frac{\mathsf{Q}(\bar{v}_1 \mid v_1)\mathsf{Q}(S_1 \mid v_1, \bar{v}_1)}{\mathsf{Q}(S_1 \mid v_1)} \\
&= \frac{\mathsf{Q}(\bar{v}_1)\mathsf{Q}(S_1 \mid v_1, \bar{v}_1)}{\mathsf{Q}(S_1 \mid v_1)} \qquad \text{(since } \bar{v}_1 \text{ is independent of } v_1\text{)} \\
&= \frac{\mathsf{Q}(\bar{v}_1)\mathsf{Q}(S_1 \mid v_1)}{\mathsf{Q}(S_1 \mid v_1)} = \mathsf{Q}(\bar{v}_1) \qquad \text{(the samples in } S_1 \text{ depend only on } v_1\text{)}.
\end{aligned}
$$

Inequality (ii) follows since the samples are drawn independently given $v = (v_1, \bar{v}_1)$. Finally, (iii) follows since $\bar{S}_1$ (the samples that lie outside the interval I_1) only depend on $\bar{v}_1$ since the marginal distribution of x is independent of v and the distribution of $y \mid x$ depends only on the value of v corresponding to the interval in which x lies.

Thus since, $\mathsf{Q}(S \mid v_1) = \mathsf{Q}(S_1 \mid v_1)\mathsf{Q}(\bar{S}_1)$ we have that

$$
\mathrm{KL}(\mathsf{Q}(S \mid v_1 = 1)\|\mathsf{Q}(S \mid v_1 = -1)) = \mathrm{KL}(\mathsf{Q}(S_1 \mid v_1 = 1)\|\mathsf{Q}(S_1 \mid v_1 = -1)). \tag{19}
$$

To bound this KL divergence, let us condition of the number of samples in S_1 from group a, (the majority group) $n_{1,a}$ and the number of samples from group b (the minority group), $n_{1,b}$. Now since $n_{1,a}$ and $n_{1,b}$ are independent of v_1 (which only affects the labels) we have that,

$$
\begin{aligned}
\mathsf{Q}(S_1 \mid v_1) &= \sum_{n_{1,a}, n_{1,b}} \mathsf{Q}(n_{1,a}, n_{1,b} \mid v_1)\mathsf{Q}(S_1 \mid v_1, n_{1,a}, n_{1,b}) \\
&= \sum_{n_{1,a}, n_{1,b}} \mathsf{Q}(n_{1,a}, n_{1,b})\mathsf{Q}(S_1 \mid v_1, n_{1,a}, n_{1,b}) \\
&= \mathbb{E}_{n_{1,a}, n_{1,b}} \left[\mathsf{Q}(S_1 \mid v_1, n_{1,a}, n_{1,b})\right].
\end{aligned}
$$

Therefore, by the joint convexity of the KL-divergence and by Jensen's inequality we have that,

$$
\begin{aligned}
&\mathrm{KL}(\mathsf{Q}(S_1 \mid v_1 = 1)\|\mathsf{Q}(S_1 \mid v_1 = -1)) \\
&\qquad \leq \mathbb{E}_{n_{1,a}, n_{1,b}} \left[\mathrm{KL}(\mathsf{Q}(S_1 \mid v_1 = 1, n_{1,a}, n_{1,b})\|\mathsf{Q}(S_1 \mid v_1 = -1, n_{1,a}, n_{1,b}))\right]. \tag{20}
\end{aligned}
$$

Now conditioned on $v_1, n_{1,a}$ and $n_{1,b}$, samples in S_1 are composed of 2 groups of samples $(S_{1,a}, S_{1,b})$. The samples in each group $(S_{1,a}, S_{1,b})$ are drawn independently from the distributions $\mathsf{P}_a(x \mid x \in I_1)\mathsf{P}_v(y \mid x)$ and $\mathsf{P}_b(x \mid x \in I_1)\mathsf{P}_v(y \mid x)$ respectively. Therefore,

$$
\begin{aligned}
&\mathrm{KL}(\mathsf{Q}(S_1 \mid v_1 = 1, n_{1,a}, n_{1,b})\|\mathsf{Q}(S_1 \mid v_1 = -1, n_{1,a}, n_{1,b})) \\
&\overset{(i)}{=} n_{1,a}\mathrm{KL}(\mathsf{P}_a(x \mid x \in I_1)\mathsf{P}_{v_1=1}(y \mid x)\|\mathsf{P}_a(x \mid x \in I_1)\mathsf{P}_{v_1=-1}(y \mid x)) \\
&\qquad\qquad + n_{1,b}\mathrm{KL}(\mathsf{P}_b(x \mid x \in I_1)\mathsf{P}_{v_1=1}(y \mid x)\|\mathsf{P}_b(x \mid x \in I_1)\mathsf{P}_{v_1=-1}(y \mid x)) \\
&\overset{(ii)}{=} (n_{1,a} + n_{1,b})\mathbb{E}_{x \sim \mathsf{Unif}(I_1)} \left[\mathrm{KL}(\mathsf{P}_{v_1=1}(y \mid x)\|\mathsf{P}_{v_1=-1}(y \mid x))\right] \\
&\overset{(iii)}{=} \frac{n_{1,a} + n_{1,b}}{2}\mathbb{E}_{x \sim \mathsf{Unif}(I_1)} \left[\sum_{y \in \{-1,1\}}\left(1 + y\phi\left(x - \frac{1}{2K}\right)\right)\log\left(\frac{(1 + y\phi(x - \frac{1}{2K}))}{(1 + y\phi(x - \frac{1}{2K}))}\right)\right] \\
&= \frac{n_{1,a} + n_{1,b}}{2}\sum_{y \in \{-1,1\}}\mathbb{E}_{x \sim \mathsf{Unif}(I_1)} \left[\left(1 + y\phi\left(x - \frac{1}{2K}\right)\right)\log\left(\frac{(1 + y\phi(x - \frac{1}{2K}))}{(1 + y\phi(x - \frac{1}{2K}))}\right)\right] \\
&= \frac{n_{1,a} + n_{1,b}}{2K}\sum_{y \in \{-1,1\}}\int_{x=0}^{\frac{1}{K}} \left[\left(1 + y\phi\left(x - \frac{1}{2K}\right)\right)\log\left(\frac{(1 + y\phi(x - \frac{1}{2K}))}{(1 + y\phi(x - \frac{1}{2K}))}\right)\right] \mathrm{d}x \\
&\overset{(iv)}{\leq} \frac{n_{1,a} + n_{1,b}}{3K^2}, \tag{21}
\end{aligned}
$$

where in (i) we let P_{v_1} denote the conditional distribution of y for $x \in I_1$ given v_1, (ii) follows since both P_a and P_b are constant in the interval, (iii) follows by our construction of P_v above, and finally (iv) follows by invoking Lemma A.3 that ensures that the integral is bounded by $1/3K^2$.

627   Using this bound in Eq. (20), along with Eq. (19) we get that

$$\mathrm{KL}(\mathsf{Q}(S \mid v_1 = 1)\|\mathsf{Q}(S \mid v_1 = -1)) \leq \frac{\mathbb{E}\left[n_{1,a} + n_{2,b}\right]}{3K^2}.$$

628   Now there are n_{maj} samples from group a in S and n_{min} samples from group b. Therefore,

$$\mathbb{E}\left[n_{1,a}\right] = n_{\mathsf{maj}}\mathbb{P}[\mathsf{P}_a(x \in I_1)] = \frac{n_{\mathsf{maj}}(2 - \tau)}{K},$$
$$\mathbb{E}\left[n_{1,b}\right] = n_{\mathsf{min}}\mathbb{P}[\mathsf{P}_b(x \in I_1)] = \frac{n_{\mathsf{min}}\tau}{K}.$$

629   Plugging this bound into Eq. (21) completes the proof by the first interval. An identical argument
630   holds for $j \in \{2, \ldots, K/2\}$. For $j \in \{K/2 + 1, \ldots, K\}$ the only change is that

$$\mathbb{E}\left[n_{j,a}\right] = n_{\mathsf{maj}}\mathbb{P}[\mathsf{P}_a(x \in I_j)] = \frac{n_{\mathsf{maj}}\tau}{K},$$
$$\mathbb{E}\left[n_{j,b}\right] = n_{\mathsf{min}}\mathbb{P}[\mathsf{P}_b(x \in I_j)] = \frac{n_{\mathsf{min}}(2 - \tau)}{K}.$$

631    $\square$

632   Next, we combine the previous two lemmas to establish our stated lower bound. We first restate it
633   here.

634   **Theorem 4.2.** *Consider the group shift setting described in Section 3.2.2. Given any overlap*
635   *$\tau \in [0, 1]$ recall that $\mathcal{P}_{\mathsf{GS}}(\tau)$ is the class of distributions such that* $\mathsf{Overlap}(\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}}) \geq \tau$. *The*
636   *minimax excess risk in this setting is lower bounded as follows:*

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\mathsf{GS}}(\tau)) = \inf_{\mathcal{A}} \sup_{(\mathsf{P}_{\mathsf{maj}},\mathsf{P}_{\mathsf{min}}) \in \mathcal{P}_{\mathsf{GS}}(\tau)} \mathsf{Excess\ Risk}[\mathcal{A}; (\mathsf{P}_{\mathsf{maj}}, \mathsf{P}_{\mathsf{min}})]$$
$$\geq \frac{c}{(n_{\mathsf{min}} \cdot (2 - \tau) + n_{\mathsf{maj}} \cdot \tau)^{1/3}} \geq \frac{c}{n_{\mathsf{min}}^{1/3}(\rho \cdot \tau + 2)^{1/3}}, \quad (4)$$

637   *where $\rho = n_{\mathsf{maj}}/n_{\mathsf{min}} > 1$.*

638   *Proof.* First, by Lemma C.1 we know that

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\mathsf{GS}}(\tau)) \geq \frac{1}{32K^2}\sum_{j=1}^{K}\exp(-\mathrm{KL}(\mathsf{Q}(S \mid v_j = 1)\|\mathsf{Q}(S \mid v_j = -1))).$$

639   Next, by invoking the bound on the KL divergences in the equation above by Lemma C.2 we get that

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\mathsf{GS}}(\tau))$$
$$\geq \frac{1}{64K}\left[\exp\left(-\frac{n_{\mathsf{maj}}(2 - \tau) + n_{\mathsf{min}}\tau}{3K^3}\right) + \exp\left(-\frac{n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau}{3K^3}\right)\right]$$
$$\geq \frac{1}{64K}\left[\exp\left(-\frac{n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau}{3K^3}\right)\right]$$

640   Setting $K = \lceil (n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau)^{1/3}\rceil$ and recalling that $\tau \leq 1$ we get that

$$\mathsf{Minimax\ Excess\ Risk}(\mathcal{P}_{\mathsf{GS}}(\tau))$$
$$\geq \frac{1}{64\lceil (n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau)^{1/3}\rceil}\left[\exp\left(-\frac{n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau}{3\lceil (n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau)^{1/3}\rceil^3}\right)\right]$$
$$\geq \frac{c'}{64\lceil (n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau)^{1/3}\rceil}$$
$$\geq \frac{c}{(n_{\mathsf{min}}(2 - \tau) + n_{\mathsf{maj}}\tau)^{1/3}},$$

641   which completes the proof. $\square$

C.2 Proof of Theorem 5.2

In this section, we derive an upper bound on the excess risk of the undersampled binning estimator $\mathcal{A}_{\text{USB}}$ (Eq. (5)). Recall that given a dataset $\mathcal{S}$ this estimator first calculates the undersampled dataset $\mathcal{S}_{\text{US}}$, where the number of points from the minority group (n_{min}) is equal to the number of points from the majority group (n_{min}), and the size of the dataset is $2n_{\text{min}}$. Throughout this section, $(\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}})$ shall be an arbitrary element of $\mathcal{P}_{\text{GS}}(\tau)$ for any $\tau \in [0,1]$. In this section, whenever we shall often denote Excess Risk$(\mathcal{A}; (\mathsf{P}_{\text{maj}}, \mathsf{P}_{\text{min}}))$ by simply Excess Risk$(\mathcal{A})$.

Before we proceed, we introduce some additional notation. For any $j \in \{1, \dots, K\}$ and $I_j = [\frac{j-1}{K}, \frac{j}{K}]$ let

$$q_{j,1} := \mathsf{P}_{\text{test}}(y = 1 \mid x \in I_j) = \int_{x \in I_j} \mathsf{P}(y = 1 \mid x)\mathsf{P}_{\text{test}}(x \mid x \in I_j)\,\mathrm{d}x, \tag{22a}$$

$$q_{j,1} := \mathsf{P}_{\text{test}}(y = 1 \mid x \in I_j) = \int_{x \in I_j} \mathsf{P}(y = 1 \mid x)\mathsf{P}_{\text{test}}(x \mid x \in I_j)\,\mathrm{d}x. \tag{22b}$$

For the undersampled binning estimator $\mathcal{A}_{\text{USB}}$ (defined above in Eq. (5)), define the *excess risk in an interval I_j* as follows:

$$R_j(\mathcal{A}_{\text{USB}}^{\mathcal{S}}) := p\left(y = -\mathcal{A}_j^{\mathcal{S}} \mid x \in I_j\right) - \min\left\{\mathsf{P}_{\text{test}}(y = 1 \mid x \in I_j), \mathsf{P}_{\text{test}}(y = -1 \mid x \in I_j)\right\}$$
$$= q_{j,-\mathcal{A}_j^{\mathcal{S}}} - \min\{q_{j,1}, q_{j,-1}\}.$$

The proof of the upper bound shall proceed in steps. First, in Lemma C.3 we will show that the excess risk is equal to sum the excess risk over the intervals up to a factor of $2/K$ on account of the distribution being 1-Lipschitz. Next, in Lemma C.4 we upper bound the risk over each interval. We put these two together and to upper bound the risk.

Lemma C.3. *The expected excess risk of undersampled binning estimator $\mathcal{A}_{\text{USB}}$ can be decomposed as follows*

$$\text{Excess Risk}(\mathcal{A}_{\text{USB}}) \leq \sum_{j=0}^{K-1} \mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\text{maj}}^{n_{\text{maj}}} \times \mathsf{P}_{\text{min}}^{n_{\text{min}}}} \left[R_j(\mathcal{A}_{\text{USB}}^{\mathcal{S}})\right] \cdot \mathsf{P}_{\text{test}}(I_j) + \frac{2}{K},$$

where $\mathsf{P}_{\text{test}}(I_j) := \int_{x \in I_j} \mathsf{P}_{\text{test}}(x)\,\mathrm{d}x$.

Proof. Recall that by definition, the expected excess risk is

$$\mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\text{maj}}^{n_{\text{maj}}} \times \mathsf{P}_{\text{min}}^{n_{\text{min}}}} \left[R(\mathcal{A}^{\mathcal{S}}; \mathsf{P}_{\text{test}}) - R(f^\star; \mathsf{P}_{\text{test}})\right].$$

Let us first decompose the Bayes risk $R(f^\star)$,

$$R(f^\star) = \inf_f \mathbb{E}_{(x,y) \sim \mathsf{P}_{\text{test}}} \left[\mathbf{1}(f(x) \neq y)\right]$$

$$= \inf_f \int_{x=0}^1 \sum_{y \in \{-1,1\}} \mathbf{1}(f(x) \neq y)\mathsf{P}_{\text{test}}(y \mid x)\mathsf{P}_{\text{test}}(x)\,\mathrm{d}x$$

$$= \int_{x=0}^1 \inf_{f(x) \in \{-1,1\}} \sum_{y \in \{-1,1\}} \mathbf{1}(f(x) \neq y)\mathsf{P}_{\text{test}}(y \mid x)\mathsf{P}_{\text{test}}(x)\,\mathrm{d}x$$

$$= \int_{x=0}^1 \inf_{f(x) \in \{-1,1\}} \mathsf{P}_{\text{test}}(y = -f(x) \mid x)\mathsf{P}_{\text{test}}(x)\,\mathrm{d}x$$

$$= \int_{x=0}^1 \min\left\{\mathsf{P}_{\text{test}}(y = 1 \mid x), \mathsf{P}_{\text{test}}(y = -1 \mid x)\right\} \mathsf{P}_{\text{test}}(x)\,\mathrm{d}x. \tag{23}$$

The risk of the undersampled binning algorithm $\mathcal{A}_{\text{USB}}$ is given by

$$R(\mathcal{A}_{\text{USB}}^{\mathcal{S}}) = \int_{x=0}^1 \sum_{y \in \{-1,1\}} \mathbf{1}(\mathcal{A}_{\text{USB}}^{\mathcal{S}}(x) \neq y)\mathsf{P}_{\text{test}}(y \mid x)\mathsf{P}_{\text{test}}(x)\,\mathrm{d}x$$

$$= \int_{x=0}^1 \mathsf{P}_{\text{test}}(y = -\mathcal{A}_{\text{USB}}^{\mathcal{S}}(x) \mid x)\mathsf{P}_{\text{test}}(x)\,\mathrm{d}x.$$

Next, recall that the undersampled binning estimator is constant over the intervals I_j for $j \in \{1, \ldots, K\}$ where it takes the value $\mathcal{A}_j^{\mathcal{S}}$ (to ease notation let us simply denote it by $\mathcal{A}_j$ below), and therefore

$$R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) = \sum_{j=0}^{K-1} \int_{x \in I_j} \mathsf{P}_{\mathsf{test}}(y = -\mathcal{A}_j | x) \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x.$$

This combined with Eq. (23) tells us that

$$R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) - R(f^\star)$$
$$= \sum_{j=0}^{K-1} \int_{x \in I_j} \left(\mathsf{P}_{\mathsf{test}}(y = -\mathcal{A}_j | x) - \min \left\{ \mathsf{P}_{\mathsf{test}}(y = 1 \mid x), \mathsf{P}_{\mathsf{test}}(y = -1 \mid x) \right\} \right) \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x. \quad (24)$$

Recall the definition of $q_{j,1}$ and $q_{j,-1}$ from Eqs. (22a)-(22b) above. For any $x \in I_j = [\frac{j-1}{K}, \frac{j}{K}]$, $|\mathsf{P}_{\mathsf{test}}(y \mid x) - q_{j,y}| \le 1/K$, since the distribution $\mathsf{P}_{\mathsf{test}}(y \mid x)$ is 1-Lipschitz and $q_{j,y}$ is its conditional mean. Therefore,

$$R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) - R(f^\star)$$
$$\le \sum_{j=0}^{K-1} \int_{x \in I_j} \left(q_{j,-\mathcal{A}_j} - \min \left\{ q_{j,1}, q_{j,-1} \right\} \right) \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x + \frac{2}{K} \sum_{j=0}^{K-1} \int_{x \in I_j} \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x$$
$$= \sum_{j=0}^{K-1} \int_{x \in I_j} R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x + \frac{2}{K}.$$

Taking expectation over the training samples $\mathcal{S}$ (where n_{min} samples are drawn independently from $\mathsf{P}_{\mathsf{min}}$ and n_{maj} samples are drawn independently from $\mathsf{P}_{\mathsf{maj}}$) concludes the proof. $\qquad \square$

Next we provide an upper bound on the expected excess risk is an interval $R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})$.

Lemma C.4. *For any $j \in \{1, \ldots, K\}$ with $I_j = [\frac{j-1}{K}, \frac{j}{K}]$,*

$$\mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) \right] \le \frac{c}{\sqrt{n_{\mathsf{min}} \mathsf{P}_{\mathsf{test}}(I_j)}} + \frac{c}{K},$$

where c is an absolute constant, and $\mathsf{P}_{\mathsf{test}}(I_j) := \int_{x \in I_j} \mathsf{P}_{\mathsf{test}}(x) \, \mathrm{d}x$.

Proof. Consider an arbitrary bucket $j \in \{1, \ldots, K\}$.

Let us introduce some notation that shall be useful in the remainder of the proof. Analogous to $q_{j,1}$ and $q_{j,-1}$ defined above (see Eqs. (22a)-(22b)), define $q_{j,1}^a$ and $q_{j,1}^b$ as follows:

$$q_{j,1}^a := \mathsf{P}_a(y = 1 \mid x \in I_j) = \int_{x \in I_j} \mathsf{P}(y = 1 \mid x) \mathsf{P}_a(x \mid x \in I_j) \, \mathrm{d}x, \quad (25a)$$

$$q_{j,1}^b := \mathsf{P}_b(y = 1 \mid x \in I_j) = \int_{x \in I_j} \mathsf{P}(y = 1 \mid x) \mathsf{P}_b(x \mid x \in I_j) \, \mathrm{d}x. \quad (25b)$$

Essentially, $q_{j,1}^a$ is the probability that a sample is from group a and has label 1, conditioned on the event that the sample falls in the interval I_j. Since

$$\mathsf{P}_{\mathsf{test}}(x \mid x \in I_j) = \frac{1}{2} \left[\mathsf{P}_a(x \mid x \in I_j) + \mathsf{P}_b(x \mid x \in I_j) \right],$$

therefore

$$|q_{j,1} - q_{j,1}^a| = \left| \int_{x \in I_j} \mathsf{P}(y = 1 \mid x) \mathsf{P}_{\mathsf{test}}(x \mid x \in I_j) \, \mathrm{d}x - \int_{x \in I_j} \mathsf{P}(y = 1 \mid x) \mathsf{P}_a(x \mid x \in I_j) \, \mathrm{d}x \right|$$
$$\le \frac{1}{K}. \quad (26)$$

681 This follows since $\mathsf{P}(y \mid x)$ is 1-Lipschitz and therefore can fluctuate by at most $1/K$ in the interval
682 I_j. Of course the same bound also holds for $|q_{j,1} - q_{j,1}^b|$.

683 With this notation in place let us present a bound on the expected value of $R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})$. By definition

$$R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}}) = q_{j,-\mathcal{A}_j^{\mathcal{S}}} - \min\{q_{j,1}, q_{j,-1}\}.$$

684 First, note that $q_{j,1} := \mathsf{P}_{\mathsf{test}}(y = 1 \mid x \in I_j) = 1 - q_{j,-1}$. Suppose that $q_{j,1} < 1/2$ and therefore
685 $q_{j,-1} > 1/2$ (the same bound shall hold in the other case). In this case, risk is incurred only when
686 $\mathcal{A}_j^{\mathcal{S}} = 1$. That is,

$$\mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}} \left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right] = |q_{j,-1} - q_{j,1}|\mathbb{P}_{\mathcal{S}}[\mathcal{A}_j^{\mathcal{S}} = 1]$$

$$= |1 - 2q_{j,1}|\mathbb{P}_{\mathcal{S}}[\mathcal{A}_j^{\mathcal{S}} = 1]. \tag{27}$$

687 Now by the definition of the undersampled binning estimator (see Eq. (5)), $\mathcal{A}_j^{\mathcal{S}} = 1$ only when there
688 are more samples in the interval I_j with label 1 than -1. However, we can bound the probability of
689 this happening since $q_{j,1}$ is smaller than $q_{j,-1}$.

690 Let n_j be the number of samples in the undersampled sample set $\mathcal{S}_{\mathsf{US}}$ in the interval I_j. Let $n_{1,j}$ be
691 the number of these samples with label 1, and $n_{-1,j} = n_j - n_{1,j}$ be the number of samples with
692 label -1. Further, let $n_{a,j}$ be the number of samples in from group a such that they fall in the interval
693 I_j, and define $m_{b,j}$ analogously.

694 The probability of incurring risk is given by

$$\mathbb{P}[\mathcal{A}_j = 1] = \sum_{s=1}^{2n_{\mathsf{min}}} \mathbb{P}[\mathcal{A}_j = 1 \mid n_j = s]\mathbb{P}[n_j = s], \tag{28}$$

695 where the sum is up to $2n_{\mathsf{min}}$ since the size of the undersample dataset $|\mathcal{S}_{\mathsf{US}}|$ is equal to $2n_{\mathsf{min}}$.

696 Conditioned on the event that $n_j = s$ the probability of incurring risk is

$$\mathbb{P}\left[\mathcal{A}_j = 1 \mid n_j = s\right] = \mathbb{P}\left[m_{1,j} > n_{-1,j} \mid n_j = s\right] = \mathbb{P}\left[n_{1,j} > n_j/2 \mid n_j = s\right]$$
$$= \mathbb{P}\left[n_{1,j} > s/2 \mid n_j = s\right]. \tag{29}$$

697 Now, note that $n_j = n_{a,j} + n_{b,j}$. Thus continuing, we have that

$$\mathbb{P}\left[n_{1,j} > s/2 \mid n_j = s\right] = \sum_{s' \le s} \mathbb{P}\left[n_{1,j} > s/2 \mid n_j = s, n_{b,j} = s'\right]\mathbb{P}[n_{b,j} = s']$$
$$= \sum_{s' \le s} \mathbb{P}\left[n_{1,j} > s/2 \mid n_{a,j} = s - s', n_{b,j} = s'\right]\mathbb{P}[n_{b,j} = s'].$$

698 In light of this previous equation, we want to control the probability that the number of samples with
699 label 1 in the interval I_j conditioned on the event that the number of samples from group a in this
700 interval is $s - s'$ and the number of samples from group b in this interval is s'. Recall that $q_{j,1}^a$ and
701 $q_{j,1}^b$ the probabilities of the label of the sample being 1 conditioned the event that sample is in the
702 interval I_j when it is group a and b respectively. So we define the random variables:

$$z_a[s - s'] \sim \mathsf{Bin}(s - s', q_{j,1}^a), \quad z_b[s'] \sim \mathsf{Bin}(s', q_{j,1}^b), \quad z[s] \sim \mathsf{Bin}(s, \max\{q_{j,1}^a, q_{j,1}^b\}).$$

703 Then,

$$\mathbb{P}\left[n_{1,j} > s/2 \mid n_j = s\right]$$

$$= \sum_{s' \leq s} \mathbb{P}\left[n_{1,j} > s/2 \mid n_{j,a} = s - s', n_{j,b} = s'\right] \mathbb{P}[n_{j,b} = s']$$

$$= \sum_{s' \leq s} \mathbb{P}\left[z_a[s - s'] + z_b[s']) > s/2 \mid n_{a,j} = s - s', n_{b,j} = s'\right] \mathbb{P}[n_{b,j} = s']$$

$$\leq \sum_{s' \leq s} \mathbb{P}\left[z[s] > s/2 \mid n_{a,j} = s - s', n_{b,j} = s'\right] \mathbb{P}[n_{b,j} = s']$$

$$= \sum_{s' \leq s} \mathbb{P}\left[z[s] > s/2\right] \mathbb{P}[n_{b,j} = s']$$

$$= \mathbb{P}\left[z[s] > s/2\right]$$

$$\overset{(i)}{\leq} \exp\left(-\frac{s}{2}(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2\right), \tag{30}$$

704 where (i) follows by invoking Hoeffding's inequality[22, Proposition 2.5]. Combining this with
705 Eqs. (28) and (29) we get that

$$\mathbb{P}[\mathcal{A}_j = 1] \leq \sum_{s=1}^{2n_{\min}} \exp\left(-\frac{s}{2}(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2\right) \mathbb{P}[n_j = s].$$

706 Now n_j, which is the number of samples that lands in the interval I_j is equal to $n_{a,j} + n_{b,j}$. Now each
707 of $n_{a,j}$ and $n_{b,j}$ (the number of samples in this interval from each of the groups) are random variables
708 with distributions $\mathrm{Bin}(n_{\min}, \mathsf{P}_a(I_j))$ and $\mathrm{Bin}(n_{\min}, \mathsf{P}_b(I_j))$, where $\mathsf{P}_a(I_j) = \int_{x \in I_j} \mathsf{P}_a(x) \, \mathrm{d}x$ and
709 $\mathsf{P}_b(I_j) = \int_{x \in I_j} \mathsf{P}_a(x) \, \mathrm{d}x$. Therefore, n_j is distributed as a sum of two binomial distribution and is
710 therefore Poisson binomially distributed [26]. Using the formula for the moment generating function
711 (MGF) of a Poisson binomially distributed random variable we infer that,

$$\mathbb{P}[\mathcal{A}_j = 1] \leq \left(1 - \mathsf{P}_a(I_j) + \mathsf{P}_a(I_j)\exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right)^{n_{\min}} \times$$

$$\left(1 - \mathsf{P}_b(I_j) + \mathsf{P}_b(I_j)\exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right)^{n_{\min}}.$$

712 Plugging this into Eq. (28) we get that,

$$\mathbb{E}_{\mathcal{S} \sim \mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}} \times \mathsf{P}_{\min}^{n_{\min}}}\left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right]$$

$$\leq |1 - 2q_{j,1}|\left[1 - \mathsf{P}_a(I_j) + \mathsf{P}_a(I_j)\exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right]^{n_{\min}} \times$$

$$\left[1 - \mathsf{P}_b(I_j) + \mathsf{P}_b(I_j)\exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right]^{n_{\min}}$$

$$= |1 - 2q_{j,1}|\left[1 - \mathsf{P}_a(I_j)\left(1 - \exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right)\right]^{n_{\min}} \times$$

$$\left[1 - \mathsf{P}_b(I_j)\left(1 - \exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right)\right)\right]^{n_{\min}}.$$

713 Since $|1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\}| \leq 1$,

$$1 - \exp\left(-\frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{2}\right) \geq \frac{(1 - 2\max\left\{q_{j,1}^a, q_{j,1}^b\right\})^2}{4},$$

714 and therefore

$$\mathbb{E}_{\mathcal{S}\sim\mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}}\times\mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}}\left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right]\leq |1-2q_{j,1}|\left[1-\mathsf{P}_a(I_j)\frac{(1-2\max\left\{q_{j,1}^a,q_{j,1}^b\right\})^2}{2}\right]^{n_{\mathsf{min}}}\times$$

$$\left[1-\mathsf{P}_b(I_j)\frac{(1-2\max\left\{q_{j,1}^a,q_{j,1}^b\right\})^2}{2}\right]^{n_{\mathsf{min}}}$$

$$\overset{(i)}{\leq} |1-2q_{j,1}|\left[1-\mathsf{P}_a(I_j)\frac{(1-2q_{j,1}-2\gamma)^2}{2}\right]^{n_{\mathsf{min}}}\times$$

$$\left[1-\mathsf{P}_b(I_j)\frac{(1-2q_{j,1}-2\gamma)^2}{2}\right]^{n_{\mathsf{min}}}$$

$$\overset{(ii)}{\leq} |1-2q_{j,1}|\exp\left(-n_{\mathsf{min}}(\mathsf{P}_a(I_j)+\mathsf{P}_b(I_j))\frac{(1-2q_{j,1}-2\gamma)^2}{2}\right),$$

715 where (i) follows since $|\max\{q_{j,1}^a,q_{j,1}^b\}-q_{j,1}|\leq 1/K$ by Eq. (26) and γ is such that $|\gamma|\leq 1/K$, and
716 (ii) follows since $(1+z)^b\leq\exp(bz)$. Now the RHS above is maximized when $(1-2q_{j,1}-2\gamma)^2=$
717 $\frac{c}{n_{\mathsf{min}}(\mathsf{P}_a(I_j)+\mathsf{P}_b(I_j))}$, for some constant c. Plugging this into the equation above we get that

$$\mathbb{E}_{\mathcal{S}\sim\mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}}\times\mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}}\left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right]\leq\frac{c'}{\sqrt{n_{\mathsf{min}}(\mathsf{P}_a(I_j)+\mathsf{P}_b(I_j))}}+c'|\gamma|$$

$$\leq\frac{c'}{\sqrt{n_{\mathsf{min}}(\mathsf{P}_a(I_j)+\mathsf{P}_b(I_j))}}+\frac{c'}{K}.$$

718 Finally, noting that $\mathsf{P}_{\mathsf{test}}(I_j)=(\mathsf{P}_a(I_j)+\mathsf{P}_b(I_j))/2$ completes the proof. $\qquad\square$

719 By combining the previous two lemmas we can now prove our upper bound on the risk of the
720 undersampled binning estimator. We begin by restating it.

721 **Theorem 5.2.** *Consider the group shift setting described in Section 3.2.2. For any overlap $\tau\in[0,1]$*
722 *and for any $(\mathsf{P}_{\mathsf{maj}},\mathsf{P}_{\mathsf{min}})\in\mathcal{P}_{\mathsf{GS}}(\tau)$ the expected excess risk of the Undersampling Binning Estimator*
723 *(Eq. (5)) with number of bins with $K=\lceil n_{\mathsf{min}}^{1/3}\rceil$ is*

$$\mathsf{Excess\ Risk}[\mathcal{A}_{\mathsf{USB}};(\mathsf{P}_{\mathsf{maj}},\mathsf{P}_{\mathsf{min}})]=\mathbb{E}_{\mathcal{S}\sim\mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}}\times\mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}}\left[R(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}};\mathsf{P}_{\mathsf{test}}))-R(f^\star;\mathsf{P}_{\mathsf{test}})\right]\leq\frac{C}{n_{\mathsf{min}}^{1/3}}.$$

724 *Proof.* First by Lemma C.3 we know that

$$\mathsf{Excess\ Risk}[\mathcal{A}_{\mathsf{USB}}]\leq\sum_{j=0}^{K-1}\mathbb{E}_{\mathcal{S}\sim\mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}}\times\mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}}\left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right]\cdot\mathsf{P}_{\mathsf{test}}(I_j)+\frac{2}{K}.$$

725 Next by using the bound on $\mathbb{E}_{\mathcal{S}\sim\mathsf{P}_{\mathsf{maj}}^{n_{\mathsf{maj}}}\times\mathsf{P}_{\mathsf{min}}^{n_{\mathsf{min}}}}\left[R_j(\mathcal{A}_{\mathsf{USB}}^{\mathcal{S}})\right]$ established in Lemma C.4 we get that,

$$\mathsf{Excess\ Risk}(\mathcal{A}_{\mathsf{USB}})\leq c\sum_{j=0}^{K-1}\frac{1}{\sqrt{n_{\mathsf{min}}\mathsf{P}_{\mathsf{test}}(I_j)}}\mathsf{P}_{\mathsf{test}}(I_j)+\frac{c}{K}$$

$$=\frac{c}{\sqrt{n_{\mathsf{min}}}}\sum_{j=0}^{K-1}\sqrt{\mathsf{P}_{\mathsf{test}}(I_j)}+\frac{c}{K}$$

$$\overset{(i)}{\leq}\frac{c}{\sqrt{n_{\mathsf{min}}}}\sqrt{K}\sum_{j=0}^{K-1}\mathsf{P}_{\mathsf{test}}(I_j)+\frac{c}{K}$$

$$=c\sqrt{\frac{K}{n_{\mathsf{min}}}}+\frac{c}{K}.$$

726 where (i) follows since for any vector $z\in\mathbb{R}^K$, $\|z\|_1\leq\sqrt{K}\|z\|_2$. Maximizing over K yields the
727 choice $K=\lceil n_{\mathsf{min}}^{1/3}\rceil$, completing the proof.

728 $\qquad\square$

D   Experimental Details for Figure 2

We construct our label shift dataset from the original CIFAR10 dataset. We create a binary classification task using the "cat" and "dog" classes. We use the official test examples as the balanced test set with 1000 cats and 1000 dogs. To form the initial train and validation sets, we use 2500 cat examples (half of the training set) and 500 dog examples, corresponding to a 5:1 label imbalance. We use 80% of those examples for training and the rest for validation. We are left with 2500 additional cat examples and 4500 dog examples from the original train set which we add into our training set to generate Figure 2.

We use the same convolutional neural network architecture as [3, 24] with random initializations for this dataset. We train this model using SGD for 400 epochs with batchsize 64, a constant learning rate 0.001 and momentum 0.9.

For the VS loss [13] we set $\tau = 3$ and $\gamma = 0.3$, the best hyperparameters identified by Wang et al. [24] on this dataset for this neural network architecture. The importance weights used upweight the minority class samples in the training loss and validation loss is calculated to be $\frac{\#\text{Cat Train Examples}}{\#\text{Dog Train Examples}}$.

We note that all of the experiments were performed on an internal cluster on 8 GPUs.