# BOME! Bilevel Optimization Made Easy: A Simple First-Order Approach

## Abstract

Bilevel optimization (BO) is useful for solving a variety of important machine learning problems including but not limited to hyperparameter optimization, meta-learning, continual learning, and reinforcement learning. Conventional BO methods need to differentiate through the low-level optimization process with implicit differentiation, which requires expensive calculations related to the Hessian matrix. There has been a recent quest for first-order methods for BO, but the methods proposed to date tend to be complicated and impractical for large-scale deep learning applications. In this work, we propose a simple first-order BO algorithm that depends only on first-order gradient information, requires no implicit differentiation, and is practical and efficient for large-scale non-convex functions in deep learning. We provide non-asymptotic convergence analysis of the proposed method to stationary points for non-convex objectives and present empirical results that show its superior practical performance.

## 1 Introduction

We consider the bilevel optimization (BO) problem:

$$\min_{v,\theta} f(v,\theta) \quad s.t. \quad \theta \in \arg\min_{\theta'} g(v,\theta'), \tag{1}$$

where the goal is to minimize an *outer objective* $f$ whose variables include the solution of another minimization problem w.r.t an *inner objective* $g$. The $\theta$ and $v$ are the *inner* and *outer* variables, respectively. We assume that $v \in \mathbb{R}^m, \theta \in \mathbb{R}^n$ and that $g(v, \cdot)$ attains a minimum for each $v$.

BO is useful in a variety of machine learning tasks. A canonical example is hyperparameter optimization, in which case $f$ (resp. $g$) is the validation (resp. training) loss associated with a model parameter $\theta$ and a hyperparameter $v$, and we want to find the optimal hyperparameter $v$ to minimize the validation loss $f$ when $\theta$ is determined by minimizing the training loss; see e.g., Pedregosa [42], Franceschi et al. [12]. Other applications include meta learning [12], continual learning [43], reinforcement learning [52], and adversarial learning [23]. See Liu et al. [32] for a recent survey.

BO is notoriously challenging due to its nested nature. Despite the large literature, most existing methods for BO are slow and unsatisfactory in various ways. For example, a major class of BO methods is based on direct gradient descent on the outer variable $v$ while viewing the optimal inner variable $\theta^*(v) = \arg\min_\theta g(v, \theta)$ as a (uniquely defined) function of $v$. The key difficulty is to calculate the derivative $\nabla_v \theta^*(v)$ which may require expensive manipulation of the Hessian matrix of $g$ via the implicit differentiation theorem. Another approach is to replace the low level optimization with the stationary condition $\nabla_\theta g(v, \theta) = 0$. This still requires Hessian information, and more importantly, is unsuitable for nonconvex $g$ since it allows $\theta$ to be any stationary point of $g(v, \cdot)$, not necessarily a minimizer. To the best of our knowledge, the only existing fully first-order BO

algorithms[1] are BSG-1 [15] and BVFSM with its variants [33–35]; but BSG-1 relies on a non-vanishing approximation that does not yield convergence to the correct solution in general, and BVFSM is sensitive to hyper-parameters on large-scale practical problems and lacks a complete non-asymptotic analysis for the practically implemented algorithm.

In this work, we seek a *simple and fast fully first-order* BO method that can be used with non-convex functions including those appear in deep learning applications. The idea is to reformulate (1) as a single-level constrained optimization problem using the so-called value-function-based approach [10, 7]. The constrained problem is then solved by stopping gradient on the single variable that contains the higher-order information and applying a simple first-order dynamic barrier gradient descent method based on a method of Gong et al. [16]. Our contributions are: **1)** we introduce a novel, fast yet powerful BO method by applying a modified dynamic barrier gradient descent on the value-function reformulation of BO; **2)** Theoretically, we establish the non-asymptotic convergence of our method to local stationary points (as measured by a special KKT loss) for non-convex $f$ and $g$. Importantly, to the best of our knowledge, this work is the first to establish non-asymptotic convergence rate for a fully first-order BO method. This result is also *much* beyond that of Gong et al. [16]. **3)** Empirically, the proposed method achieves better or comparable performance while being more efficient than state-of-the-art BO methods on a variety of benchmarks.

## 2  Background

This section provides a brief background on traditional BO methods. Please see Bard [2], Dempe and Zemkoho [7], Dempe [6] for overviews, and Liu et al. [32] for a survey on recent ML applications.

**Hypergradient Descent** Assume that the minimum of $g(v, \cdot)$ is unique for all $v$ so that we can write $\theta^*(v) = \arg\min_\theta g(v, \theta)$ as a function of $v$; this is known as the low-level singleton (LLS) assumption. The most straightforward approach to solving (1) is to conduct gradient descent on $f(v, \theta^*(v))$ as a function of $v$. Note that

$$\nabla_v f(v, \theta^*(v)) = \nabla_1 f(v, \theta^*(v)) + \nabla_v \theta^*(v) \nabla_2 f(v, \theta^*(v)).$$

The difficulty is to compute $\nabla_v \theta^*(v)$. From implicit function theorem, it satisfies a linear equation:

$$\nabla_{1,2} g(v, \theta^*(v)) + \nabla_{2,2} g(v, \theta^*(v)) \nabla_v \theta^*(v) = 0. \tag{2}$$

If $\nabla_{2,2} g$ is invertible, we can solve for $\nabla_v \theta^*(v)$ and obtain a gradient update rule on $v$:

$$v_{k+1} \leftarrow v_k - \xi \left( \nabla_1 f_k - \left( \nabla_{1,2} g_k \right)^\top \left( \nabla_{2,2} g_k \right)^{-1} \nabla_2 f_k \right),$$

where $k$ denotes iteration, $\nabla_1 f_k = \nabla_1 f(v_k, \theta^*(v_k))$ and similarly for the other terms. This approach is sometimes known as the *hypergradient descent*. However, hypergradient descent is computationally expensive: Besides requiring evaluation of the inner optimum $\theta^*(v_k)$, the main computational bottleneck is to solve the linear equation in (2). Methods have been developed that approximate (2) using conjugate gradient [42, 44, 17], Neumann series [29, 37], and related variants [14]. Another popular approximation approach is to replace $\nabla_v \theta^*(v)$ with $\nabla_v \theta^{(T)}(v)$, where $\theta^{(T)}(v)$ denotes the $T$-th iteration of gradient descent or other optimization steps on $g(v, \theta)$ w.r.t. $\theta$ starting from certain initialization. The gradient $\nabla_v \theta^{(T)}(v)$ can be calculated with auto-differentiation (AD) with either forward mode [11], backward mode [12, 11, 46, 28, 1] or their variants [31]. While these approaches claim to be first-order, they require many Hessian-vector or Jacobian-vector products at each iteration and are slow for large problems.

Other examples of approximation methods include a neural surrogate method which approximates $\theta^*(v)$ and its gradient $\nabla_v \theta^*(v)$ with neural networks [38] and Newton-Gaussian approximation of the Hessian matrix with covariance of gradient [15]. Both approaches introduce non-vanishing approximation error that is difficult to control. The neural surrogate method also suffers from high training cost for the neural network.

**Stationary-Seeking Methods.** An alternative method is to replace the argmin constraint in (1) with the stationarity condition $\nabla_\theta g(v, \theta) = 0$, yielding a constrained optimization:

$$\min_{v, \theta} f(v, \theta) \quad s.t. \quad \nabla_\theta g(v, \theta) = 0. \tag{3}$$

---

[1] By fully first-order, we mean methods that only require information of $f, g, \nabla f, \nabla g$, so this excludes methods that apply auto-differentiation or conjugate gradient that need multiple steps of matrix-vector computation.

Algorithms for nonlinear equality constrained optimization can then be applied [39]. The constraint in (3) guarantees only that $\theta$ is a stationary point of $g(v, \cdot)$, so it is equivalent to (1) only when $g$ is convex w.r.t. $\theta$. Otherwise, the solution of (3) can be a maximum or saddle point of $g$. This makes it problematic for deep learning, where non-convex functions are pervasive.

# 3 Method

We consider a *value function approach* [see e.g., 41, 53, 33], which yields natural first-order algorithms for non-convex $g$ and requires no computation of Hessian matrices. It is based on the observation that (1) is equivalent to the following constrained optimization (even for non-convex $g$):

$$\min_{v,\theta} f(v, \theta) \quad s.t. \quad q(v, \theta) := g(v, \theta) - g^*(v) \leq 0, \tag{4}$$

where $g^*(v) := \min_\theta g(v, \theta) = g(v, \theta^*(v))$ is known as the value function. Compared with the hypergradient approach, this formulation **does not require calculation of the implicit derivative** $\nabla_v \theta^*(v)$: Although $g^*(v)$ depends on $\theta^*(v)$, its derivative $\nabla_v g^*(v)$ does not depend on $\nabla_v \theta^*(v)$, by Danskin's theorem:

$$\nabla_v g^*(v) = \nabla_1 g(v, \theta^*(v)) + \nabla_v \theta^*(v) \nabla_2 g(v, \theta^*(v)) = \nabla_1 g(v, \theta^*(v)), \tag{5}$$

where the second term in (5) vanishes because we have $\nabla_2 g(v, \theta^*(v)) = 0$ by definition of the optimum $\theta^*(v)$. Therefore, provided that we can evaluate $\theta^*(v)$ at each iteration, solving (4) yields an algorithm for BO that requires no Hessian computation. In this work, we make use of the dynamic barrier gradient descent algorithm of Gong et al. [16] to solve (4). This is an elementary first-order algorithm for solving constrained optimization, but it applies only to a special case of the bilevel problem and must be extended to handle the general case we consider here.

**Dynamic Barrier Gradient Descent.** The idea is to iterative update the parameter $(v, \theta)$ to reduce $f$ while controlling the decrease of the constraint $q$, ensuring that $q$ decreases whenever $q > 0$. Specifically, denote $\xi$ as the step size, the update at each step is

$$(v_{k+1}, \theta_{k+1}) \leftarrow (v_k, \theta_k) - \xi \delta_k, \tag{6}$$

$$\text{where} \quad \delta_k = \arg\min_\delta \|\nabla f(v_k, \theta_k) - \delta\|^2 \quad \text{s.t.} \quad \langle \nabla q(v_k, \theta_k), \delta \rangle \geq \phi_k. \tag{7}$$

Here $\nabla f_k := \nabla_{(v,\theta)} f(v_k, \theta_k)$, $\nabla q_k := \nabla_{(v,\theta)} q(v_k, \theta_k)$, and $\phi_k \geq 0$ is a non-negative control barrier and should be strictly positive $\phi_k > 0$ in the non-stationary points of $q$: the lower bound on the inner product of $\nabla q(v_k, \theta_k)$ and $\delta_k$ ensures that the update in (6) can only decrease $q$ (when step size $\xi$ is sufficiently small) until it reaches stationary. In addition, by enforcing $\delta_k$ to be close to $\nabla f(v_k, \theta_k)$ in (7), we decrease the objective $f$ as much as possible so long as it does not conflict with descent of $q$.

Two straightforward choices of $\phi_k$ that satisfies the condition above are $\phi_k = \eta q(v_k, \theta_k)$ and $\phi_k = \eta \|\nabla q(v_k, \theta_k)\|^2$ with $\eta > 0$. We find that both choices of $\phi_k$ work well empirically and use $\phi_k = \eta \|\nabla q(v_k, \theta_k)\|^2$ as the default (see Section 6.2).

The optimization in (7) yields a simple closed form solution:

$$\delta_k = \nabla f(v_k, \theta_k) + \lambda_k \nabla q(v_k, \theta_k), \quad \text{with} \quad \lambda_k = \max\left( \frac{\phi_k - \langle \nabla f(v_k, \theta_k), \nabla q(v_k, \theta_k) \rangle}{\|\nabla q(v_k, \theta_k)\|^2}, \ 0 \right),$$

and $\lambda_k = 0$ in the case of $\|\nabla q(v_k, \theta_k)\| = 0$.

**Practical Approximation.** The main bottleneck of the method above is to calculate the $q(v_k, \theta_k)$ and $\nabla q(v_k, \theta_k)$ which requires evaluation of $\theta^*(v_k)$. In practice, we approximate $\theta^*(v_k)$ by $\theta_k^{(T)}$, where $\theta_k^{(T)}$ is obtained by running $T$ steps of gradient descent of $g(v_k, \cdot)$ w.r.t. $\theta$ starting from $\theta_k$. That is, we set $\theta_k^{(0)} = \theta_k$ and let

$$\theta_k^{(t+1)} = \theta_k^{(t)} - \alpha \nabla_\theta g(v_k, \theta_k^{(t)}), \quad t = 0, \ldots, T-1, \tag{8}$$

for some step size parameter $\alpha > 0$. We obtain an estimate of $q(v, \theta)$ at iteration $k$ by replacing $\theta^*(v_k)$ with $\theta_k^{(T)}$: $\hat{q}(v, \theta) = g(v, \theta) - g(v, \theta_k^{(T)})$.

We substitute $\hat{q}(v_k, \theta_k)$ into (7) to obtain the update direction $\delta_k$. The full procedure is summarized in Algorithm 1. Note that the $\theta_k^{(T)}$ is viewed as a constant when defining $\hat{q}(v, \theta)$ and hence no

---

**Algorithm 1** Bilevel Optimization Made Easy (BOME!)

---

**Goal**: Solve $\min_{v,\theta} f(v,\theta) \quad s.t. \quad \theta \in \arg\min g(v,\cdot)$.

**Input**: Initialization $(v_0, \theta_0)$; inner step $T$; outer and inner stepsize $\xi$, $\alpha$ (set $\alpha = \xi$ by default).

**for** iteration $k$ **do**

1. Get $\theta_k^{(T)}$ by $T$ steps of gradient descent on $g(v_k, \cdot)$ starting from $\theta_k$ (See Eq. (8)).

2. Set $\hat{q}(v, \theta) = g(v, \theta) - g(v, \theta_k^{(T)})$.

3. Update $(v, \theta) : (v_{k+1}, \theta_{k+1}) \leftarrow (v_k, \theta_k) - \xi(\nabla f(v_k, \theta_k) + \lambda_k \nabla \hat{q}(v_k, \theta_k))$

$$\text{where} \quad \lambda_k = \max \left( \frac{\phi_k - \langle \nabla f(v_k, \theta_k), \ \nabla \hat{q}(v_k, \theta_k) \rangle}{\|\nabla \hat{q}(v_k, \theta_k)\|^2}, \ 0 \right),$$

and $\phi_k = \eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$ (default), or $\phi_k = \eta \hat{q}(v_k, \theta_k)$ with $\eta > 0$.

**Remark**: 1) We treat $\theta_k^{(T)}$ as constant when taking derivative of $\hat{q}$; 2) In practice, step 3 can have separate stepsize $(\xi_v, \xi_\theta)$ and use standard optimizers like Adam [26]; 3) We use $\eta = 0.5$ and $T = 10$ by default.

**end for**

---

differentiation of $\theta_k^{(T)}$ is performed when calculating the gradient $\nabla \hat{q}$. This differs from truncated back-propagation methods [e.g., 46] which differentiate through $\theta_k^{(T)}$ as a function of $v$.

Each step of Algorithm 1 can be viewed as taking one step (starting from $v_k, \theta_k$) toward solving an approximate constrained optimization problem:

$$\min_{v,\theta} f(v,\theta) \quad s.t. \quad g(v,\theta) \leq g(v, \theta_k^{(T)}), \tag{9}$$

which can be viewed as a relaxation of the exact constrained optimization formulation (4), because $\{(v,\theta) : g(v,\theta) \leq g^*(v)\}$ is a subset of $\{(v,\theta) : g(v,\theta) \leq g(v, \theta_k^{(T)})\}$.

## 4 Analysis

We first elaborate the KKT condition of (4) (Section 4.1), then quantify the convergence of the method by how fast it meets the KKT condition. We consider both the case when $g$ satisfies the Polyak-Łojasiewicz (PL) inequality w.r.t. $\theta$, hence having a unique global optimum (Section 4.2), and when $g$ have multiple local minimum (Section 4.3).

### 4.1 KKT Conditions

Consider a general constrained optimization of form $\min f(v,\theta)$ s.t. $q(v,\theta) \leq 0$. Under proper regularity conditions known as constraint quantifications [40], the first-order KKT condition gives a necessary condition for a feasible point $(v^*, \theta^*)$ with $q(v^*, \theta^*) \leq 0$ to be a local optimum of (4): There exists a Lagrangian multiplier $\lambda^* \in [0, +\infty)$, such that

$$\nabla f(v^*, \theta^*) + \lambda^* \nabla q(v^*, \theta^*) = 0, \tag{10}$$

and $\lambda^*$ satisfies the complementary slackness condition $\lambda^* q(v^*, \theta^*) = 0$. A common regularity condition to ensure (10) is the *constant rank constraint quantification (CRCQ)* condition [20].

**Definition 1.** *A point $(v^*, \theta^*)$ is said to satisfy CRCQ with a function $h$ if the rank of the Jacobian matrix $\nabla h(v, \theta)$ is constant in a neighborhood of $(v^*, \theta^*)$.*

Unfortunately, **the KKT condition in** (10) **does not hold for the bilevel optimization in** (4). The CRCQ condition does not typically hold for this problem. This is because the minimum of $q$ is zero, and hence if $(v^*, \theta^*)$ is feasible for (4), then $(v^*, \theta^*)$ must attain the minimum of $q$, yielding $q(v^*, \theta^*) = 0$ and $\nabla q(v^*, \theta^*) = 0$ if $q$ is smooth; but we could not have $\nabla q(v, \theta) = 0$ uniformly in a neighborhood of $(v^*, \theta^*)$ (hence CRCQ fails) unless $q$ is a constant around $(v^*, \theta^*)$. In addition, if KKT (10) holds, we would have $\nabla f(v^*, \theta^*) = -\lambda^* \nabla q(v^*, \theta^*) = 0$ which happens only in the rare case when $(v^*, \theta^*)$ is a stationary point of both $f, g$.

Instead, one can establish a KKT condition of BO through the form in (3), because there is nothing special that prevents $(v^*, \theta^*)$ from satisfying CRCQ with $\nabla_\theta q = \nabla_\theta g$ (even though we just showed

143 that it is difficult to have CRCQ with $q$). Assume $f$ and $\nabla_\theta q$ are continuously differentiable, and
144 $(v^*, \theta^*)$ is a point satisfying $\nabla_\theta q(v^*, \theta^*) = 0$ and CRCQ with $\nabla_\theta q$. Then by the typical first order
145 KKT condition of (3), there exists a Lagrange multiplier $\omega^* \in \mathbb{R}^n$ such that

$$\nabla f(v^*, \theta^*) + \nabla(\nabla_\theta q(v^*, \theta^*))\omega^* = 0. \tag{11}$$

146 This condition can be viewed as the limit of a sequence of (10) in the following way: assume we
147 relax the constraint in (4) to $q(v, \theta) \leq c_k$ where $c_k$ is a sequence of positive numbers that converge
148 to zero, then we can establish (10) for each $c_k > 0$ and pass the limit to zero to yield (11).

149 **Proposition 1.** *Assume that $f$, $q$, $\nabla q$ are continuously differentiable and $\|\nabla f\|$, $f$ is bounded. For*
150 *a feasible point $(v^*, \theta^*)$ of (4) that satisfies CRCQ with $\nabla_\theta q$, if $(v^*, \theta^*)$ is the limit of a sequence*
151 *$\{(v_k, \theta_k)\}_{k=1}^\infty$ satisfying $q(v_k, \theta_k) \neq 0\ \forall k$, and there exists a sequence $\{\lambda_k\} \subset [0, \infty)$ such that*

$$\nabla f(v_k, \theta_k) + \lambda_k \nabla q(v_k, \theta_k) \to 0, \qquad\qquad q(v_k, \theta_k) \to 0,$$

152 *as $k \to +\infty$, then $(v^*, \theta^*)$ satisfies (11).*

153 This motivates us to use the following function as a measure of stationarity of the solution returned
154 by the algorithm:

$$\mathcal{K}(v, \theta) = \underbrace{\min_{\lambda \geq 0} \|\nabla f(v, \theta) + \lambda \nabla q(v, \theta)\|^2}_{\text{local improvement}} + \underbrace{q(v, \theta)}_{\text{feasibility}}.$$

155 The hope is to have an algorithm that generates a sequence $\{(v_k, \theta_k)\}_{k=0}^\infty$ that satisfies $\mathcal{K}(v_k, \theta_k) \to 0$
156 as $k \to +\infty$.

157 Intuitively, the first term in $\mathcal{K}(v, \theta)$ measures how much $\nabla f$ conflicts with $\nabla q$ (how much we
158 can decrease $f$ without increasing $q$), as it is equal to the squared $\ell_2$ norm of the solution to the
159 problem $\min_\delta \|\nabla f - \delta\|^2$ s.t. $\langle \nabla q, \delta \rangle \geq 0$. The second term in $\mathcal{K}$ measures how much the $\arg\min g$
160 constraint is satisfied.

## 4.2 Convergence with unimodal $g$

162 We first present the convergence rate when assuming $g(v, \cdot)$ has unique minimizer and satisfies the
163 Polyak-Łojasiewicz (PL) inequality for all $v$, which guarantees a linear convergence rate of the
164 gradient descent on the low level problem.

165 **Assumption 1** (PL-inequality). *Given any $v$, assume $g(v, \cdot)$ has a unique minimizer denoted as $\theta^*(v)$.*
166 *Also assume there exists $\kappa > 0$ such that for any $(v, \theta)$, $\|\nabla_\theta g(v, \theta)\|^2 \geq \kappa(g(v, \theta) - g(v, \theta^*(v)))$.*

167 The PL inequality gives a characterization on how a small gradient norm implies global optimality.
168 It is implied from, but weaker than strongly convexity. The PL-inequality is more appealing than
169 convexity because some modern over-parameterized deep neural networks have been shown to satisfy
170 the PL-inequality along the trajectory of gradient descent. See, for example, Frei and Gu [13], Song
171 et al. [48], Liu et al. [30] for more discussion.

172 **Assumption 2** (Smoothness). *$f$ and $g$ are differentiable, and $\nabla f$ and $\nabla g$ are $L$-Lipschitz w.r.t. the*
173 *joint inputs $(v, \theta)$ for some $L \in (0, +\infty)$.*

174 **Assumption 3** (Boundedness). *There exists a constant $M < \infty$ such that $\|\nabla g(v, \theta)\|$, $\|\nabla f(v, \theta)\|$,*
175 *$|f(v, \theta)|$ and $|g(v, \theta)|$ are all upper bounded by $M$ for any $(v, \theta)$.*

176 Assumptions 2 and 3 are both standard in optimization.

177 **Theorem 1.** *Consider Algorithm 1 with $\xi, \alpha \leq 1/L$, $\phi_k = \eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$, and $\eta > 0$. Suppose*
178 *that Assumptions 1, 2, and 3 hold. Then there exists a constant $c$ depending on $\alpha, \kappa, \eta, L$ such that*
179 *when $T \geq c$, we have for any $K \geq 0$,*

$$\min_{k \leq K} \mathcal{K}(v_k, \theta_k) = O\left(\sqrt{\xi} + \sqrt{\frac{q_0}{\xi K}} + \frac{1}{\xi K} + \exp(-bT)\right)$$

180 *where $q_0 = q(v_0, \theta_0)$, and $b > 0$ is a constant depending on $\kappa$, $L$, and $\alpha$.*

181 **Remark** Note that one of the dominant terms depends on the initial value $q_0 = q(v_0, \theta_0)$. Therefore,
182 we can obtain a better rate if we start from a $\theta_0$ with small $q_0$ (hence near the optimum of $g(v_0, \cdot)$). In
183 particular, when $q(v_0, \theta_0) = O(1)$, choosing $\xi = O(K^{-1/2})$ gives $\min_{k \leq K} \mathcal{K}(v_k, \theta_k) = O(K^{-1/4} +$
184 $\exp(-bT))$ rate. On the other hand, if we start from a better initialization such that $q(v_0, \theta_0) =$
185 $O((\xi K)^{-1})$, then choosing $\xi = O(K^{-2/3})$ gives $\min_{k \leq K} \mathcal{K}(v_k, \theta_k) = O(K^{-1/3} + \exp(-bT))$.

### 4.3 Convergence with multimodal $g$

The PL-inequality eliminates the possibility of having stationary points that are not global optimum. To study cases in which $g$ has multiple local optima, we introduce the notion of attraction points following gradient descent.

**Definition 2** (Attraction points). *Given any $(v, \theta)$, we say that $\theta^\diamond(v, \theta)$ is the attraction point of $(v, \theta)$ with step size $\alpha > 0$ if the sequence $\{\theta^{(t)}\}_{t=0}^\infty$ generated by gradient descent $\theta^{(t)} = \theta^{(t-1)} - \alpha \nabla_\theta g(v, \theta^{(t-1)})$ starting from $\theta^{(0)} = \theta$ converges to $\theta^\diamond(v, \theta)$.*

Assume the step size $\alpha \le 1/L$ where $L$ is the smoothness constant defined in Assumption 2, one can show the existence and uniqueness of attraction point of any $(v, \theta)$ using Proposition 1.1 of Traonmilin and Aujol [49]. Intuitively, the attraction of $(v, \theta)$ is where the gradient descent algorithm can not make improvement. In fact, when $\alpha \le 1/L$, one can show that $g(v, \theta) \le g(v, \theta^\diamond(v, \theta))$ is equivalent to the stationary condition $\nabla_\theta g(v, \theta) = 0$.

The set of $(v, \theta)$ that have the same attraction point forms an attraction basin. Our analysis needs to assume the PL-inequality within the individual attraction basins.

**Assumption 4** (Local PL-inequality within attraction basins). *Assume that for any $(v, \theta)$, $\theta^\diamond(v, \theta)$ exists. Also assume that there exists $\kappa > 0$ such that for any $(v, \theta)$ $\|\nabla_\theta g(v, \theta)\|^2 \ge \kappa(g(v, \theta) - g(v, \theta^\diamond(v, \theta)))$.*

We can also define local variants of $q$ and $\mathcal{K}$ as follows:

$$q^\diamond(v, \theta) = g(v, \theta) - g(v, \theta^\diamond(v, \theta)), \quad \mathcal{K}^\diamond(v, \theta) = \min_{\lambda \ge 0} \|\nabla f(v, \theta) + \lambda \nabla q^\diamond(v, \theta)\|^2 + q^\diamond(v, \theta).$$

Compared with Section 4.2, a key technical challenge is that $\theta^\diamond(v, \theta)$ and hence $q^\diamond(v, \theta)$ can be discontinuous w.r.t. $\theta$ when it is on the boundary of different attraction basins; $\mathcal{K}^\diamond$ is not well defined on these points. However, these boundary points are not stable stationary points, and it is possible to use arguments based on the stable manifold theorem to show that an algorithm with random initialization will almost surely not visit them [47, 27].

**Theorem 2.** *Consider Algorithm 1 with $\xi, \alpha \le 1/L$, $\phi_k = \eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$, and $\eta > 0$. Suppose that Assumptions 2, 3, and 4 hold and that $q^\diamond$ is differentiable on $(v_k, \theta_k)$ at every iteration $k \ge 0$. Then there exists a constant $c$ depending on $\alpha, \kappa, \eta, L$, such that when $T \ge c$, we have*

$$\min_{k \le K} \mathcal{K}^\diamond(v_k, \theta_k) = O\left(\sqrt{\xi} + \sqrt{\frac{1}{\xi K}} + \exp(-bT)\right),$$

*where $b$ is a positive constant depending on $\kappa$, $L$, and $\alpha$.*

Unlike Theorem 1, the rate does not improve when $q_0^\diamond := q^\diamond(v_0, \theta_0)$ is small because the attraction basin may change in different iterations, eliminating the benefit of starting from a good initialization. Choosing $\xi = O(K^{-1/2})$ gives $O(K^{-1/4} + \exp(-bT))$ rate of $\min_{k \le K} \mathcal{K}^\diamond(v_k, \theta_k)$.

## 5 Related Works

The value-function formulation (4) is a classical approach in bilevel optimization [41, 53, 7]. However, despite its attractive properties, it has been mostly used as a theoretical tool, and much less exploited for practical algorithms compared with the more widely known hypergradient approach (Section 2), especially for challenging nonconvex functions $f$ and $g$ such as those encountered in deep learning. One exception is Liu et al. [33], which proposes a BO method by solving the value-function formulation using an interior-point method combined with a smoothed approximation. This was improved later in a pessimistic trajectory truncation approach [35] and a sequential minimization approach [34] (BVFSM). Similar to our approach, these methods do not require computation of Hessians, thanks to the use of value function. However, as we observe in experiments (Section 6.2), BVFSM tends to be dominated by our method both in accuracy and speed, and is sensitive to some hyperparameters that are difficult to tune (such as the coefficients of the log-barrier function in interior point method). Theoretically, Liu et al. [33, 34, 35] provide only asymptotic analysis on the convergence of the smoothed and penalized surrogate loss to the target loss. They do not give an analysis for the algorithm that was actually implemented.

Our algorithm is build up on the dynamic control barrier method of Gong et al. [16], an elementary approach for constrained optimization. Gong et al. [16] also applied their approach to solve a lexico-graphical optimization of form $\min_\theta f(\theta)$ s.t. $\theta \in \arg\min_{\theta'} g(\theta')$, which is a bilevel optimization

Figure 1: Results on the toy coreset problem and mini-max problem. (a)-(c): the trajectories of $(v_k, \theta_k)$, and $f(v_k, v_k)$ and $\hat{q}_k(v_k, v_k)$ of BOME (our method), BSG-1 [15], BVFSM [34], Penalty [39] and Optimistic GD [5] (only for minimax problem). (d)-(e) trajectories of BOME with different choices of inner gradient step $T$ and the control coefficient $\eta$.

234 without an outer variable (known as *simple bilevel optimization* [8]). Our method is an extension of
235 their method to general bilevel optimization. We provide non-asymptotic analysis for our method,
236 going beyond the continuous time analysis in Gong et al. [16]. A key sophistication in the theoretical
237 analysis is that we need to control the approximation error of $\theta^*(v_k)$ with $\theta_k^{(T)}$ at each step, which
238 requires an analysis significantly different from that of Gong et al. [16].

239 Another recent body of theoretical works on BO focus on how to optimize when only stochastic
240 approximation of the objectives is provided [14, 19, 22, 51, 18, 4, 25]; there are also recent works on
241 the lower bounds and minimax optimal algorithms [21, 22]. These algorithms and analysis are based
242 on hypergradient descent and hence require Hessian-vector products in implementation.

## 6 Experiment

244 We conduct experiments (1) to study the correctness, basic properties, and robustness to hyperpa-
245 rameters of BOME, and (2) to test its performance and computational efficiency on challenging ML
246 applications, compared with state-of-the-art bilevel algorithms. In the following, we first list the
247 baseline methods and how we set the hyperparameters. Then we introduce the experiment problems
248 in Section 6.1, which includes 3 toy problems and 3 ML applications, and provide the experiment
249 results. Finally we summarize observations and findings in Section 6.2.

250 **Baselines** A comprehensive set of state-of-the-art BO methods are chosen as baseline methods. This
251 includes the *fully first-order* methods: BSG-1 [15] and BVFSM [34], ; a *stationary-seeking* method:
252 Penalty [39], *explicit/implicit* methods: ITD [22], AID-CG (using conjugate gradient), AID-FP
253 (using fixed point method) [17], reverse (using reverse auto-differentiation) [11] stocBiO [22], and
254 VRBO [51].

255 **Hyperparameters** Unless otherwise specified, BOME strictly follows Algorithm 1 with $\phi_k = $
256 $\eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$, $\eta = 0.5$, and $T = 10$. The inner stepsize $\alpha$ is set to be the same
257 as outer stepsize $\xi$. The stepsizes of all methods are set by a grid search from the set
258 $\{0.01, 0.05, 0.1, 0.5, 1, 5, 10, 50, 100, 500, 1000\}$. All toy problems adopt vanilla gradient descent
259 (GD) and applications on hyperparameter optimization adapts GD with a momentum of 0.9. Details
260 are provided in Appendix A.

### 6.1 Experiment Problems and Results

262 **Toy Coreset Problem** To validate the *convergence* property of BOME, we consider:

$$\min_{v,\theta} \|\theta - x_0\|^2 \quad s.t. \quad \theta \in \arg\min_\theta \|\theta - X\sigma(v)\|^2,$$

263 where $\sigma(v) = \exp(v)/\sum_{i=1}^4 \exp(v_i)$ is the softmax function, $v \in \mathbb{R}^4, \theta \in \mathbb{R}^2$, and $X = $
264 $[x_1, x_2, x_3, x_4] \in \mathbb{R}^{2 \times 4}$. The goal is to find the closest point to a target point $x_0$ within the convex
265 hull of $\{x_1, \ldots, x_4\}$. See Fig. 6.1 (upper row) for the illustration and results.

Figure 2: Result for hyperparameter optimization. **Top:** data hyper-cleaning on MNIST dataset. The solid black line is the model performance trained purely on the validation set and the dashed black line is the model performance trained on the validation set and on the part of training set that have correct labels. **Bottom:** learnable regularization on 20 Newsgroup dataset. The solid black line indicates the model performance without any regularization. All results are averaged on 5 random trials. See Appendix A.4 and A.5 for results on FashionMNIST and more details.

**Toy Mini-Max Game** Mini-max game is a special and challenging case of BO where $f$ and $g$ contradicts with each other completely (e.g., $f = -g$). We consider

$$\min_{v,\theta \in \mathbb{R}} v\theta \quad s.t. \quad \theta \in \arg\max_{\theta' \in \mathbb{R}} v\theta'. \tag{12}$$

The optimal solution is $v^* = \theta^* = 0$. Note that the naive gradient descent ascent algorithm diverges to infinity on this problem, and a standard alternative is to use optimistic gradient descent [5]. Figure 6.1 (lower row) shows that BOME works on this problem while other first-order BO methods fail.

**Degenerate Low Level Problem** Many existing BO algorithms require the low level singleton (LLS) assumption, which BOME does not require. To test this, we consider an example from Liu et al. [31]:

$$\min_{v \in \mathbb{R}, \theta \in \mathbb{R}^2} \|\theta - [v; 1]\|_2^2 \quad s.t. \quad \theta \in \arg\min_{(\theta'_1, \theta'_2) \in \mathbb{R}^2} (\theta'_1 - v)^2,$$

where $\theta = (\theta_1, \theta_2)$ and the solution is $v^* = 1, \theta^* = (1, 1)$. See Fig. 4 in Appendix A.3 for the result.

**Data Hyper-cleaning** We are given a noisy training set $\mathcal{D}_{\text{train}} := \{x_i, y_i\}_{i=1}^m$ and a clean validation set $\mathcal{D}_{\text{val}}$. The goal is to optimally weight the training data points so that the model trained on the weighted training set yields good performance on the validation set:

$$\min_{v,\theta} \ell^{\text{val}}(\theta), \quad s.t. \quad \theta = \arg\min_{\theta'} \left\{ \ell^{\text{train}}(\theta', v) + c \|\theta'\|^2 \right\},$$

where $\ell^{\text{val}}$ is the validation loss on $\mathcal{D}^{\text{val}}$, and $\ell^{\text{train}}$ is a weighted training loss: $\ell^{\text{train}} = \sum_{i=1}^m \sigma(v_i)\ell(x_i, y_i, \theta)$ with $\sigma(v) = \text{Clip}(v, [0,1])$ and $v \in \mathbb{R}^m$. We set $c = 0.001$. For the dataset, we use MNIST [9] (FashionMNIST [50]). We corrupt 50% of the training points by assigning them randomly sampled labels. See Fig. 2 (upper panel) for the results. (Results for FashionMNIST are reported in Appendix A.4.)

**Learnable Regularization** We apply bilevel optimization to learn the optimal regularization coefficient on the twenty newsgroup dataset:[2]

$$\min_{v,\theta} \ell^{\text{val}}(\theta) \quad s.t. \quad \theta \in \arg\min_{\theta'} \left\{ \ell^{\text{train}}(\theta') + \|W_v\theta'\|_2^2 \right\},$$

where $W_v$ is a matrix depending on $v$, e.g., $W_v = \text{diag}(\exp(v))$. See Fig. 2 (lower panel) for results.

**Continual Learning (CL)** CL studies how to learn on a sequence of tasks in an online fashion without catastrophic forgetting of previously learned tasks. We follow the setting of contextual transformation network (CTN) from Pham et al. [43], which trains a deep neural network consisting of a quickly updated backbone network (parameterized by $\theta$) and a slowly updated controller network (parameterized by $v$). When training the $\tau$-th task, we update $(v, \theta)$ by

$$\min_{v,\theta} \ell_{1:\tau}^{\text{val}}(v, \theta) \quad s.t. \quad \theta \in \arg\min_{\theta'} \ell_{1:\tau}^{\text{train}}(v, \theta'),$$

---

[2]Dataset from `https://scikit-learn.org/0.19/datasets/twenty_newsgroups.html`.

| Method | PMNIST | | | Split CIFAR | | |
|---|---|---|---|---|---|---|
| | ACC ($\uparrow$) | NBT ($\downarrow$) | FT ($\uparrow$) | ACC ($\uparrow$) | NBT ($\downarrow$) | FT ($\uparrow$) |
| Offline | $84.95 \pm 0.95$ | - | - | $74.11 \pm 0.66$ | - | - |
| MER | $76.59 \pm 0.74$ | $5.73 \pm 0.59$ | $82.32 \pm 0.34$ | $60.32 \pm 0.86$ | $8.91 \pm 0.86$ | $69.23 \pm 0.40$ |
| CTN (+ITD) | $78.40 \pm 0.28$ | $5.62 \pm 0.39$ | $84.02 \pm 0.29$ | $67.7 \pm 60.96$ | $4.88 \pm 0.77$ | $72.58 \pm 0.62$ |
| CTN (+BVFSM) | $77.78 \pm 0.32$ | $7.25 \pm 0.28$ | $\mathbf{85.03} \pm 0.28$ | $67.04 \pm 0.76$ | $6.97 \pm 0.62$ | $\mathbf{74.01} \pm 0.57$ |
| CTN (+BOME) | $\mathbf{80.70} \pm 0.26$ | $\mathbf{4.09} \pm 0.27$ | $84.79 \pm 0.25$ | $\mathbf{68.16} \pm 0.60$ | $\mathbf{4.72} \pm 0.75$ | $72.88 \pm 0.48$ |

Table 1: Results of continual learning as bilevel optimization. We compute the mean and standard error of each method's results over 5 independent runs. Best results are **bolded**. The full result with comparison against other methods are provided in Table 2 in the Appendix.

where $\ell_{1:\tau}^{\text{train}}$ and $\ell_{1:\tau}^{\text{val}}$ are the training and validation loss available up to task $\tau$. The goal is to update the controller such that the long term loss $\ell_{1:\tau}^{\text{val}}$ is minimized assuming $\theta$ is adapted to the available training loss when new tasks come. Assume the CL process terminates at time $t$. Denote by $a_\tau^s$ the test accuracy of task $s$ after training on task $\tau$. We measure the performance of CL by 1) the final mean accuracy on all seen tasks (ACC $= \frac{1}{t} \sum_{\tau \leq t} a_t^\tau$), 2) how much the model forgets as measured by negative backward transfer NBT $= \frac{1}{t} \sum_{\tau \leq t} (a_\tau^\tau - a_t^\tau)$, and 3) how fast the model learns on new tasks as measured by forward transfer FT $= \frac{1}{t} \sum_{\tau \leq t} a_t^\tau$. Note that FT $=$ ACC $+$ NBT.

We follow the setting of Pham et al. [43] closely, except replacing their bilevel optimizer (which is essentially ITD [22]) with BOME. See Appendix A.6 for experiment details. The results are shown in Table 1, where in addition to the bilevel algorithms, we also compare with a set of state-of-the-art CL algorithms, including MER [45], ER [3], and GEM [36]. Table 1 also includes an 'Offline" basline – learning $t$ tasks simultaneously using a single model (which is the upper bound on performance).

## 6.2 Observations

**BOME yields faster learning and better solutions at convergence** Figure 6.1-4 show that BOME converges to the optimum of the corresponding bilevel problems and work well on the mini-max optimization and the degenerate low level problem; see also Fig.3 in Appendix A.1. In comparison, the other methods like BSG-1, BVFSM, and Penalty fail to converge to the true optimum even with a grid search over their hyperparameters. Moreover, in all three toy examples, BOME guarantees that $\hat{q}$, which is a proxy for the optimality of the inner problem, decreases to 0. From Fig. 2, it is observed that BOME achieves comparable or better performance than the state-of-the-art bilevel methods for hyperparameter optimization. Moreover, BOME exhibits better computational efficiency (Fig. 2), especially on the twenty newsgroup dataset where the dimension of $\theta$ is large. In Table 1, we find that directly plugging in BOME to the CL problem yields a substantial performance boost.

**Robustness to parameter choices** Besides the standard step size $\xi$ in typical optimizers, BOME only has three parameters: control coefficient $\eta$, inner loop iteration $T$, and inner step size $\alpha$. We use the default setting of $\eta = 0.5$, $T = 10$ and $\alpha = \xi$ across the experiments. From Fig. 6.1 (d,e) and Fig. 2 (b,d), BOME is robust to the choice of $\eta$, $T$ and $\alpha$ as varying them results in almost identical performance. Specifically, $T = 1$ works well in many cases (see Figure 6.1 (e) and 2 (b)). The fact that BOME works well with a small $T$ empirically makes it computationally attractive in practice.

**Choice of control barrier** $\phi_k$ The control barrier is set as $\phi_k = \eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$ by default. Another option is to use $\phi_k = \eta \hat{q}(v_k, \theta_k)$. We test both options on the data hyper-cleaning and learnable regularization experiments in Fig. 2 (d), and observe no significant difference (we choose $\eta$ properly so that both choices of $\phi_k$ is on the same order). Hence we use $\phi_k = \eta \|\nabla \hat{q}(v_k, \theta_k)\|^2$ as the default.

**Comparison against BVFSM** The most relevant baseline to BOME is BVFSM, which similarly adopts the value-function reformulation of the bilevel problems. However, BOME consistently outperform BVFSM in both converged results and computational efficiency, across all experiments. More importantly, BOME has fewer hyperparameters and is robust to them, while we found BVFSM is sensitive to hyperparameters. This makes BOME a better fit for large practical bilevel problems.

## 7 Conclusion and Future Work

BOME, a simple fully first-order bilevel method, is proposed in this work with non-asymptotic convergence guarantee. While the current theory requires the inner loop iterations to scale in a logarithmic order w.r.t to the outer loop iterations, we do not observe this empirically. A further study to understand the mechanism is an interesting future direction.

# References

[1] Michael Arbel and Julien Mairal. Amortized implicit differentiation for stochastic bilevel optimization. *arXiv preprint arXiv:2111.14580*, 2021.

[2] Jonathan F Bard. *Practical bilevel optimization: algorithms and applications*, volume 30. Springer Science & Business Media, 2013.

[3] Arslan Chaudhry, Marcus Rohrbach, Mohamed Elhoseiny, Thalaiyasingam Ajanthan, Puneet K Dokania, Philip HS Torr, and Marc'Aurelio Ranzato. On tiny episodic memories in continual learning. *arXiv preprint arXiv:1902.10486*, 2019.

[4] Tianyi Chen, Yuejiao Sun, and Wotao Yin. A single-timescale stochastic bilevel optimization method. *arXiv preprint arXiv:2102.04671*, 2021.

[5] Constantinos Daskalakis, Andrew Ilyas, Vasilis Syrgkanis, and Haoyang Zeng. Training gans with optimism. *arXiv preprint arXiv:1711.00141*, 2017.

[6] Stephan Dempe. *Foundations of bilevel programming*. Springer Science & Business Media, 2002.

[7] Stephan Dempe and Alain Zemkoho. *Bilevel optimization*. Springer, 2020.

[8] Stephan Dempe, Nguyen Dinh, Joydeep Dutta, and Tanushree Pandit. Simple bilevel programming and extensions. *Mathematical Programming*, 188(1):227–253, 2021.

[9] Li Deng. The mnist database of handwritten digit images for machine learning research. *IEEE Signal Processing Magazine*, 29(6):141–142, 2012.

[10] Nguyen Dinh, B Mordukhovich, and Tran TA Nghia. Subdifferentials of value functions and optimality conditions for dc and bilevel infinite and semi-infinite programs. *Mathematical Programming*, 123(1):101–138, 2010.

[11] Luca Franceschi, Michele Donini, Paolo Frasconi, and Massimiliano Pontil. Forward and reverse gradient-based hyperparameter optimization. In *International Conference on Machine Learning*, pages 1165–1173. PMLR, 2017.

[12] Luca Franceschi, Paolo Frasconi, Saverio Salzo, Riccardo Grazzi, and Massimiliano Pontil. Bilevel programming for hyperparameter optimization and meta-learning. In *International Conference on Machine Learning*, pages 1568–1577. PMLR, 2018.

[13] Spencer Frei and Quanquan Gu. Proxy convexity: A unified framework for the analysis of neural networks trained by gradient descent. *Advances in Neural Information Processing Systems*, 34, 2021.

[14] Saeed Ghadimi and Mengdi Wang. Approximation methods for bilevel programming. *arXiv preprint arXiv:1802.02246*, 2018.

[15] Tommaso Giovannelli, Griffin Kent, and Luis Nunes Vicente. Bilevel stochastic methods for optimization and machine learning: Bilevel stochastic descent and darts. *arXiv preprint arXiv:2110.00604*, 2021.

[16] Chengyue Gong, Xingchao Liu, and Qiang Liu. Automatic and harmless regularization with constrained and lexicographic optimization: A dynamic barrier approach. *Advances in Neural Information Processing Systems*, 34, 2021.

[17] Riccardo Grazzi, Luca Franceschi, Massimiliano Pontil, and Saverio Salzo. On the iteration complexity of hypergradient computation. In *International Conference on Machine Learning*, pages 3748–3758. PMLR, 2020.

[18] Zhishuai Guo and Tianbao Yang. Randomized stochastic variance-reduced methods for stochastic bilevel optimization. *arXiv preprint arXiv:2105.02266*, 2021.

[19] Mingyi Hong, Hoi-To Wai, Zhaoran Wang, and Zhuoran Yang. A two-timescale framework for bilevel optimization: Complexity analysis and application to actor-critic. *arXiv preprint arXiv:2007.05170*, 2020.

[20] Robert Janin. Directional derivative of the marginal function in nonlinear programming. In *Sensitivity, Stability and Parametric Analysis*, pages 110–126. Springer, 1984.

[21] Kaiyi Ji and Yingbin Liang. Lower bounds and accelerated algorithms for bilevel optimization. *arXiv preprint arXiv:2102.03926*, 2021.

[22] Kaiyi Ji, Junjie Yang, and Yingbin Liang. Bilevel optimization: Convergence analysis and enhanced design. In *International Conference on Machine Learning*, pages 4882–4892. PMLR, 2021.

[23] Haoming Jiang, Zhehui Chen, Yuyang Shi, Bo Dai, and Tuo Zhao. Learning to defend by learning to attack. In *International Conference on Artificial Intelligence and Statistics*, pages 577–585. PMLR, 2021.

[24] Hamed Karimi, Julie Nutini, and Mark Schmidt. Linear convergence of gradient and proximal-gradient methods under the polyak-łojasiewicz condition. In *Joint European Conference on Machine Learning and Knowledge Discovery in Databases*, pages 795–811. Springer, 2016.

[25] Prashant Khanduri, Siliang Zeng, Mingyi Hong, Hoi-To Wai, Zhaoran Wang, and Zhuoran Yang. A near-optimal algorithm for stochastic bilevel optimization via double-momentum. *arXiv preprint arXiv:2102.07367*, 2021.

[26] Diederik P Kingma and Jimmy Ba. Adam: A method for stochastic optimization. *arXiv preprint arXiv:1412.6980*, 2014.

[27] Jason D Lee, Max Simchowitz, Michael I Jordan, and Benjamin Recht. Gradient descent only converges to minimizers. In *Conference on learning theory*, pages 1246–1257. PMLR, 2016.

[28] Junyi Li, Bin Gu, and Heng Huang. A fully single loop algorithm for bilevel optimization without hessian inverse. *arXiv preprint arXiv:2112.04660*, 2021.

[29] Renjie Liao, Yuwen Xiong, Ethan Fetaya, Lisa Zhang, KiJung Yoon, Xaq Pitkow, Raquel Urtasun, and Richard Zemel. Reviving and improving recurrent back-propagation. In *International Conference on Machine Learning*, pages 3082–3091. PMLR, 2018.

[30] Chaoyue Liu, Libin Zhu, and Mikhail Belkin. Loss landscapes and optimization in over-parameterized non-linear systems and neural networks. *Applied and Computational Harmonic Analysis*, 2022.

[31] Risheng Liu, Pan Mu, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang. A generic first-order algorithmic framework for bi-level programming beyond lower-level singleton. In *International Conference on Machine Learning*, pages 6305–6315. PMLR, 2020.

[32] Risheng Liu, Jiaxin Gao, Jin Zhang, Deyu Meng, and Zhouchen Lin. Investigating bi-level optimization for learning and vision from a unified perspective: A survey and beyond. *arXiv preprint arXiv:2101.11517*, 2021.

[33] Risheng Liu, Xuan Liu, Xiaoming Yuan, Shangzhi Zeng, and Jin Zhang. A value-function-based interior-point method for non-convex bi-level optimization. *arXiv preprint arXiv:2106.07991*, 2021.

[34] Risheng Liu, Xuan Liu, Shangzhi Zeng, Jin Zhang, and Yixuan Zhang. Value-function-based sequential minimization for bi-level optimization. *arXiv preprint arXiv:2110.04974*, 2021.

[35] Risheng Liu, Yaohua Liu, Shangzhi Zeng, and Jin Zhang. Towards gradient-based bilevel optimization with non-convex followers and beyond. *Advances in Neural Information Processing Systems*, 34, 2021.

[36] David Lopez-Paz and Marc'Aurelio Ranzato. Gradient episodic memory for continual learning. *Advances in neural information processing systems*, 30:6467–6476, 2017.

[37] Jonathan Lorraine, Paul Vicol, and David Duvenaud. Optimizing millions of hyperparameters by implicit differentiation. In *International Conference on Artificial Intelligence and Statistics*, pages 1540–1552. PMLR, 2020.

[38] Matthew MacKay, Paul Vicol, Jon Lorraine, David Duvenaud, and Roger Grosse. Self-tuning networks: Bilevel optimization of hyperparameters using structured best-response functions. *arXiv preprint arXiv:1903.03088*, 2019.

[39] Akshay Mehra and Jihun Hamm. Penalty method for inversion-free deep bilevel optimization. In *Asian Conference on Machine Learning*, pages 347–362. PMLR, 2021.

[40] Jorge Nocedal and Stephen J. Wright. *Numerical Optimization*. Springer Science & Business Media, second edition, 2006.

[41] Jivrí V Outrata. On the numerical solution of a class of stackelberg problems. *Zeitschrift für Operations Research*, 34(4):255–277, 1990.

[42] Fabian Pedregosa. Hyperparameter optimization with approximate gradient. In *International conference on machine learning*, pages 737–746. PMLR, 2016.

[43] Quang Pham, Chenghao Liu, Doyen Sahoo, and HOI Steven. Contextual transformation networks for online continual learning. In *International Conference on Learning Representations*, 2020.

[44] Aravind Rajeswaran, Chelsea Finn, Sham Kakade, and Sergey Levine. Meta-learning with implicit gradients. 2019.

[45] Matthew Riemer, Ignacio Cases, Robert Ajemian, Miao Liu, Irina Rish, Yuhai Tu, and Gerald Tesauro. Learning to learn without forgetting by maximizing transfer and minimizing interference. *arXiv preprint arXiv:1810.11910*, 2018.

[46] Amirreza Shaban, Ching-An Cheng, Nathan Hatch, and Byron Boots. Truncated backpropagation for bilevel optimization. In *The 22nd International Conference on Artificial Intelligence and Statistics*, pages 1723–1732. PMLR, 2019.

[47] Michael Shub. *Global stability of dynamical systems*. Springer Science & Business Media, 2013.

[48] Chaehwan Song, Ali Ramezani-Kebrya, Thomas Pethick, Armin Eftekhari, and Volkan Cevher. Subquadratic overparameterization for shallow neural networks. *Advances in Neural Information Processing Systems*, 34, 2021.

[49] Yann Traonmilin and Jean-Fran$\theta cois Aujol. The basins of attraction of the global minimizers of the non-convex sparse spike estimation problem. Inverse Problems$, $36(4):045003, 2020.$

[50] Han Xiao, Kashif Rasul, and Roland Vollgraf. Fashion-mnist: a novel image dataset for benchmarking machine learning algorithms. *arXiv preprint arXiv:1708.07747*, 2017.

[51] Junjie Yang, Kaiyi Ji, and Yingbin Liang. Provably faster algorithms for bilevel optimization. *arXiv preprint arXiv:2106.04692*, 2021.

[52] Zhuoran Yang, Yongxin Chen, Mingyi Hong, and Zhaoran Wang. Provably global convergence of actor-critic: A case for linear quadratic regulator with ergodic cost. 2019.

[53] JJ Ye and DL Zhu. Optimality conditions for bilevel programming problems. *Optimization*, 33 (1):9–27, 1995.

## Societal Impacts

This paper proposes a simple first order algorithm for bi-level optimization. Many specific instantiation of bi-level optimization such as adversarial learning and data attacking might be harmful to machine learning system in real world, as a general optimization algorithm for bi-level optimization, our method can be a tool in such process. We also develop a great amount of theoretical works and to our best knowledge, we do not observe any significant negative societal impact of our theoretical result.


## A  Experiment Details

We provide details about each experiment in this section. Regarding the implementation of baseline methods:

- BVFSM's implementation is adapted from `https://github.com/vis-opt-group/BVFSM`.

- Penalty's implementation is adapted from `https://github.com/jihunhamm/bilevel-penalty`.

- VRBO's implementation is adapted from `https://github.com/JunjieYang97/MRVRBO`.

- AID-CG and AID-FP implementations are adapted from `https://github.com/prolearner/hypertorch`.

- ITD implementation is adapted from `https://github.com/JunjieYang97/stocBiO`.

### A.1  Toy Coreset Problem

The problem is:

$$\min_{v,\theta} \|\theta - x_0\|^2 \quad s.t. \quad \theta \in \arg\min_{\theta} \|\theta - X\sigma(v)\|^2,$$

where $\sigma(v) = \exp(v)/\sum_{i=1}^{4}\exp(v_i)$ is the softmax function, $v \in \mathbb{R}^4, \theta \in \mathbb{R}^2$, and $X = [x_1, x_2, x_3, x_4] \in \mathbb{R}^{2\times4}$. where $\sigma(v) = \exp(v)/\sum_{i=1}^{4}\exp(v_i)$ is the softmax function. Here the outer objective $f$ pushes $\theta$ to towards $x_0$ while the inner objective $g$ ensures $\theta$ remains in the convex hull formed by 4 points in the 2D plane (e.g. $X = [x_1, x_2, x_3, x_4] \in \mathbb{R}^{2\times4}$). We choose $x_0 = (3, -2)$ and the four points $x_1 = (1, 3), x_2 = (3, 1), x_3 = (-2, 2)$ and $x_4 = (-3, 2)$. We set $v_0 = (0, 0, 0, 0)$ and $\theta_0 \in \{(0, 3), (-3, 1), (3.5, 1)\}$. For all methods, we fix both the inner stepsize $\alpha$ and the outer stepsize $\xi$ to be 0.05 and set $T = 10$. For BVFSM and Penalty, we grid search the best hyperparameters from $\{0.001, 0.01, 0.1\}$. For BOME, we choose $\phi = \eta \|\nabla\hat{q}\|^2$ and ablate over $\eta \in \{0.1, 0.5, 0.9\}$ and $T \in \{1, 10, 100\}$. The visualization of the optimization trajectories over the 3 initial points are plotted in Fig. 3.

Figure 3: Trajectories of $(v_k, \theta_k)$ on the toy coreset problem (6.1) obtained from BOME (blue) and three recent first-order bilevel methods: BSG-1 [15] (green), BVFSM [34] (orange), and Penalty [39] (red). The goal of the problem is to find the closet point (marked by opt.) to the goal $x_0$ within the convex envelop of the four vertexes. All methods start from 3 initial points (start 1-3), and the converged points are shown in darkblue. For BOME, we also plot the trajectory of $\{\hat{\theta}_k^T\}$ in cyan.

As shown, BOME successfully converges to the optimal solution regardless of the initial $\theta_0$, while BSG-1, BVFSM and Penalty methods converge to non-optimal points. We emphasize that for BVFSM and Penalty, the convergence point *depends on* the choice of hyperparameters.

### A.2  Toy Mini-max Game

The toy mini-max game we consider is:

$$\min_{v\in\mathbb{R}} v\theta^*(v) \quad s.t. \quad \theta^*(v) = \arg\max_{\theta\in\mathbb{R}} v\theta. \tag{13}$$

For BOME and BSG-1, BVFSM, and Penalty methods, we again set both the inner stepsize $\alpha$ and $\beta$ to be 0.05, as no significant difference is observed by varying the stepsizes. For all methods, we set the inner iteration $T = 10$. For BVFSM and Penalty, we grid search the best hyperparameters from $\{0.001, 0.01, 0.1\}$.

### A.3 Without LLS assumption

The toy example to validate whether BOME requires the low-level singleton assumption is borrowed from Liu et al. [31]:

$$\min_{v \in \mathbb{R}, \theta \in \mathbb{R}^2} \|\theta - [v; 1]\|_2^2 \quad s.t. \quad \theta \in \underset{(\theta_1', \theta_2') \in \mathbb{R}^2}{\arg\min} (\theta_1' - v)^2,$$

where $\theta = (\theta_1, \theta_2)$ and the optimal solution is $v^* = 1, \theta^* = (1, 1)$. Note that the inner objective has infinite many optimal solution $\theta^*(v)$ since it is degenerated. We set both the inner and outer stepsizes to 0.5 and $T = 10$ for all methods. For BVFSM and Penalty, we grid search the best hyperparameters from $\{0.001, 0.01, 0.1\}$. In Fig. 4, we provide the distance of $f(v_k, \theta_k), g(v_k, \theta_k), \theta_k, v_k$ to their corresponding optimal over training time in seconds. Note that BOME ensures that $\hat{q}(v_k, \theta_k) = g(v_k, \theta_k) - g(v^*, \theta^*)$ decreases to 0.

Figure 4: Results on Problem (A.3) which violates the low-level singleton (LLS). We compare BOME against BSG-1, BVFSM, and Penalty. $(v^*, \theta^*)$ denotes the true optimum and The four plots show how fast $f(v_k, \theta_k), g(v_k, \theta_k), \theta_k$ and $v_k$ to the corresponding optimal values w.r.t. the training time in seconds.

### A.4 Data Hyper-cleaning

The bilevel problem for data hyper-cleaning is

$$\min_{v, \theta} \ell^{\text{val}}(\theta), \quad s.t. \quad \theta = \underset{\theta}{\arg\min} \ell^{\text{train}}(\theta, v) + c \|\theta\|^2,$$

where $\ell^{\text{val}}$ is the validation loss on $\mathcal{D}^{\text{val}}$, and $\ell^{\text{train}}$ is a weighted training loss: $\ell^{\text{train}} = \sum_{i=1}^m \sigma(v_i)\ell(x_i, y_i, \theta)$ with $\sigma(v) = \text{Clip}(v, [0, 1])$ and $v \in \mathbb{R}^m$. The training data is of size $m = 50000$ and hence $w \in [0, 1]^{50000}$. The validation data is of size $m = 5000$. The model $\theta = (W, b)$ is a linear model with weight $W$ and bias $b$. Where $W \in \mathbb{R}^{10 \times 784}$ and $b \in \mathbb{R}^{10}$. For this problem, we set inner stepsize $\alpha = 0.01$ for both MNIST and FashionMNIST dataset for all methods as larger or smaller $\alpha$ result in worse performance. As we observe $v$'s gradient norm is much smaller than $\theta$'s in practice, we conduct a grid search over $\xi_v$ from $\{10.0, 50.0, 100.0, 500.0, 1000.0\}$ and also search whether to apply momentum for gradient descent, for all methods. The momentum is searched from $\{0.0, 0.9\}$. For BVFSM and Penalty methods, we also search for their best hyperparameters from $\{0.001, 0.01, 0.1, 1\}$. The model's initial parameter $\theta_0$ is initialized from a pretrained model learned only on the corrupted data. We split the dataset into 4 parts: train set, validation set 1, validation set 2, and the test set. For each method, the model is learned on the train set, and the hyperparameter $v$ is tuned using validation set 1. The best hyperparameter of any algorithm (e.g. stepsize, barrier coefficient, etc.) are then chosen based on the best validation performance on validation set 2. Then we report the final performance of the model on the test set. To conduct the ablation on $\alpha$ for BOME, we search for $\alpha \in \{0.25\xi, 0.5\xi, \xi, 2\xi\}$, where $\xi = 0.01$ is the best stepsize we found for BOME. Results on MNIST and FashionMNIST dataset are provided in Fig. 5. In the first column of Fig. 5, we compare BOME with ($T = 1$ and $T = 20$) with baseline methods whose $T$ is chosen based on best performance on the validation set 2.

Figure 5: Bilevel optimization for hyperparameter optimization. **Top:** data hyper-cleaning on MNIST dataset. The solid black line is the model performance trained purely on the validation set and the dashed black line is the model performance trained on the validation set and on the part of training set that have correct labels. **Middle:** data hyper-cleaning on FashionMNIST dataset. **Bottom:** Learnable regularization on 20 Newsgroup dataset. The solid black line indicates the model performance without any regularization. The results for each method is averaged over 5 independent runs.

**Remark:** In Fig. 2 (top row), we do not include the performance of BSG-1 as we fail to find a set of hyperparameters for BSG-1 to make it work on these data hyper-cleaning problems. VRBO's performance at convergence is tuned by hyparameter search. However, we observe that VRBO learns slowly in practice, as it requires multiple steps of Hessian vector products at each step. We notice that this is slightly inconsistent with the findings in the original paper [51]. We adapt the code from `https://github.com/JunjieYang97/MRVRBO` and find the original implementation is also slow. It is possible that a good set of hyperparameters can result in better performance.

## A.5  Learnable regularization

The bilevel optimization formulation of the learnable rgularization problem is:

$$\min_{v,\theta} \ell^{\text{val}}(\theta) \quad s.t. \quad \theta \in \arg\min_{\theta'} \ell^{\text{train}}(\theta') + \|W_v\theta'\|_2^2.$$

We use a linear model who's parameter $\theta$ is a matrix (e.g. $\theta \in \mathbb{R}^{20 \times 130107}$). Hence $v \in \mathbb{R}^{130107}$. For this experiment, the inner stepsize $\alpha$ of all methods are searched from $\{1, 10, 100, 1000\}$. The outer stepsize $\xi$ is searched from $\{0.5, 1, 5, 10, 50, 100, 500, 1000\}$. For BVFSM and Penalty methods, we also search for their best hyperparameters from $\{0.001, 0.01, 0.1, 1\}$. Similar to the experiment on Data Hyper-cleaning, we split the dataset into 4 parts: train set, validation set 1, validation set 2, and the test set. The initial model parameter $\theta_0$ is initialized from a pretrained model without any regularization (e.g. $v = 0$) to speed up the learning. To conduct the ablation on $\alpha$ for BOME, we search for $\alpha \in \{0.25\xi, 0.5\xi, \xi, 2\xi\}$, where $\xi = 100$ is the best stepsize we found for BOME. In the bottom left of Fig. 5, we compare BOME with ($T = 1$ and $T = 20$) with baseline methods whose $T$ is chosen based on best performance on the validation set 2.

**Remark:** In Fig. 2 (bottom row), we do not include the performance of VRBO as we fail to find a set of hyperparameters for VRBO that works well on the learnable regularization experiment.

## A.6  Continual Learning

Continual learning (CL) experiment follows closely to the setup in contextual transformation network (CTN) from Pham et al. [43], which trains a deep neural network consisting of a quickly updated backbone network (parameterized by $\theta$) a slowly updated controller network (parameterized by $\theta$). When training the $\tau$-th task, the update on $(v, \theta)$ is solved from a bilevel optimization:

$$\min_{v,\theta} \ell_{1:\tau}^{\text{val}}(v, \theta) \quad \text{s.t.} \quad \theta \in \arg\min_{\theta'} \ell_{1:\tau}^{\text{train}}(v, \theta').$$

More specifically,

$$\ell_{1:\tau}^{\text{val}}(v, \theta) = L^{\text{ctrl}}\big(\{\theta^*(v), v\}; \mathcal{M}_{<t+1}^{\text{sm}}\big), \quad \text{and} \quad \ell_{1:\tau}^{\text{train}}(v, \theta') = L^{\text{tr}}\big(\{\theta, v,\}, D_t \cup \mathcal{M}_{<t}^{\text{em}}\big). \quad (14)$$

Here, $\mathcal{M}_t^{\text{sm}}$ and $\mathcal{M}_t^{\text{em}}$ denote the semantic and episodic memory of task $t$ (e.g. they can be think of validation and training data) and $D_t$ is the training data of task $t$. Hence, the inner objective learns a backbone $\theta^*(v)$ that performs well on the training data which consists of the current task data $D_t$ as well as previous episodic memories $\mathcal{M}_{<t}^{\text{em}}$. Then, the outer objective encourages good generalization on the held out validation data, which consists of the semantic memory $\mathcal{M}_{<t+1}^{\text{sm}}$. All hyperparameters of BOME are set to the same as those of CTN. We choose $\phi_k = \eta \|\nabla \hat{q}_k(v_k, \theta_k)\|$ where $\eta = 2.0$, here $\eta$ is chosen from $\{0.1, 0.5, 1.0, 2.0\}$.

| Method | PMNIST | | | Split CIFAR | | |
|---|---|---|---|---|---|---|
| | ACC ($\uparrow$) | NBT ($\downarrow$) | FT ($\uparrow$) | ACC ($\uparrow$) | NBT ($\downarrow$) | FT ($\uparrow$) |
| Offline | $84.95 \pm 0.95$ | - | - | $74.11 \pm 0.66$ | - | - |
| MER | $76.59 \pm 0.74$ | $6.88 \pm 0.59$ | $82.32 \pm 0.34$ | $60.32 \pm 0.86$ | $11.80 \pm 0.86$ | $69.23 \pm 0.40$ |
| GEM | $72.74 \pm 0.91$ | $9.45 \pm 0.95$ | $80.53 \pm 0.28$ | $61.33 \pm 1.16$ | $9.21 \pm 0.94$ | $69.37 \pm 0.72$ |
| ER-Ring | $72.11 \pm 0.46$ | $9.53 \pm 0.23$ | $80.06 \pm 0.37$ | $61.96 \pm 1.22$ | $8.48 \pm 1.54$ | $69.14 \pm 0.87$ |
| CTN (+ITD) | $78.40 \pm 0.28$ | $6.25 \pm 0.41$ | $84.02 \pm 0.29$ | $67.7 \pm 60.96$ | $5.76 \pm 0.69$ | $72.58 \pm 0.62$ |
| CTN (+BVFSM) | $77.78 \pm 0.32$ | $7.86 \pm 0.32$ | $85.03 \pm 0.28$ | $67.04 \pm 0.76$ | $7.61 \pm 0.47$ | $\textbf{74.01} \pm 0.57$ |
| CTN (+Penalty) | $67.74 \pm 0.42$ | $10.10 \pm 0.48$ | $77.28 \pm 0.61$ | $47.41 \pm 2.93$ | $9.34 \pm 2.74$ | $55.76 \pm 1.64$ |
| CTN (+BOME) | $\textbf{80.70} \pm 0.26$ | $\textbf{4.73} \pm 0.23$ | $\textbf{84.79} \pm 0.25$ | $\textbf{68.16} \pm 0.60$ | $\textbf{5.32} \pm 0.76$ | $72.88 \pm 0.48$ |

Table 2: Results of continual learning as bilevel optimization. We compute the mean and standard error of each method's results over 5 independent runs. Best results are **bolded**.

## B  Proof of the Result in Section 4.1

We proof Proposition 1 using Proposition 6.3 (presented below using our notation) in Gong et al. [16] by checking all the assumptions required by Proposition 6.3 in Gong et al. [16] are satisfied. Specifically, it remains to show that for any $k$, $\lambda_k < \infty$, $\lim_{k\to\infty} \nabla q(v_k, \theta_k) = 0$ and $q$ is lower bounded (this is trivial as $q \geq 0$ by its definition), which we prove below.

Firstly, simple calculation shows that for any $k$,

$$\lambda_k = \left[ \frac{-\langle \nabla f(v_k, \theta_k), \nabla q(v_k, \theta_k) \rangle}{||\nabla q(v_k, \theta_k)||^2} \right]_+ \leq \frac{\sup_{v,\theta} ||\nabla f(v,\theta)||}{||\nabla q(v_k, \theta_k)||} < \infty.$$

Here the last inequality is by $||\nabla q(v_k, \theta_k)|| > 0$. Secondly, note that as we assume $\nabla q$ is continuous, this implies that

$$\lim_{k\to\infty} \nabla q(v_k, \theta_k) = \nabla q(v^*, \theta^*).$$

As $q(v^*, \theta^*) = 0$, we have $\nabla q(v^*, \theta^*) = 0$. Using Proposition 6.3 in Gong et al. [16] gives the desired result.

**Proposition B.1** (Proposition 6.3 in Gong et al. [16]). *Assume $f, q, \nabla q$ are continuously differentiable. Let $\{[v_k, \theta_k, \lambda_k] : k = 1, 2, ...\}$ be a sequence which satisfies $\lim_{k\to\infty} ||\nabla q(v_k, \theta_k)|| = 0$ and $\lim_{k\to\infty} ||\nabla f(v_k, \theta_k) + \lambda_k \nabla q(v_k, \theta_k)|| = 0$. Assume that $[v^*, \theta^*]$ is a limit point of $[v_k, \theta_k]$ as $k \to \infty$ and $[v^*, \theta^*]$ satisfies CRCQ with $\nabla_\theta q$, then there exists a vector-valued Lagrange multiplier $\omega^* \in \mathbb{R}^m$ (the same length as $\theta$) such that*

$$\nabla f(v^*, \theta^*) + \nabla(\nabla_\theta q(v^*, \theta^*))\omega^* = 0.$$

## C  Proof of the Result in Section 4.2

We define $L_q := 2L(L/\kappa + 1)$ and using Assumption 2 and 1, we are able to show that $q(v, \theta)$ is $L_q$-smooth (see Lemma 4 for details). For simplicity, we also assume that $\xi \leq 1$ throughout the proof. We use $b$ with some subscript to denote some general $O(1)$ constant and refer reader to section E for their detailed value.

Note that $\hat{q}$ defined in Section 3 changes in different iterations (as it depends on $\theta_k^{(T)}$) and so does $\nabla\hat{q}$. To avoid the confusion, we introduce several new notations. Firstly, given $v$ and $\theta$, $\theta^{(T)}$ denotes the results of $T$ steps of gradient of $g(v, \cdot)$ w.r.t. $\theta$ starting from $\theta$ with step size $\alpha$ (similar to the definition in (8)). Note that $\theta^{(T)}$ depends on $v, \theta$ and $\alpha$. Our notation does not reflects this dependency on $v, \alpha$ as we find it introduces no ambiguity while much simplifies the notation. Also note that when taking gradient on $\hat{q}$, the $\theta_k^{(T)}$ at iteration $k$ is treated as a constant and the gradient does not pass through it.

To be clear, we define $\hat{\nabla}q(v, \theta) = \nabla g(v, \theta) - \left[ \nabla_1^\top g(v, \theta^{(T)}), \mathbf{0}^\top \right]^\top$, where $\mathbf{0}$ denotes a zero vector with the same dimension as $\theta$. Using this definition, $\hat{\nabla}q(v_k, \theta_k) = \nabla\hat{q}(v_k, \theta_k)$ at iteration $k$. We also let $\lambda^*(v, \theta)$ be the solution of the dual problem of

$$\min_\delta ||\hat{\nabla}q(v, \theta) - \nabla f(v, \theta)||^2 \ s.t. \ \left\langle \hat{\nabla}q(v, \theta), \nabla f(v, \theta) \right\rangle \geq \eta ||\hat{\nabla}q(v, \theta)||^2. \tag{15}$$

That is

$$\lambda^*(v, \theta) = \begin{cases} \frac{[\eta||\hat{\nabla}q(v,\theta)||^2 - \langle \hat{\nabla}q(v,\theta), \nabla f(v,\theta) \rangle]_+}{||\hat{\nabla}q(v,\theta)||^2} & \text{when } ||\hat{\nabla}q(v, \theta)|| > 0 \\ 0 & \text{when } ||\hat{\nabla}q(v, \theta)|| = 0 \end{cases} \tag{16}$$

We might use $\lambda^*$ for $\lambda^*(v, \theta)$ when it introduces no confusion. Also, denote $\delta^*(v, \theta) = \lambda^*(v, \theta)\hat{\nabla}q(v, \theta) + \nabla f(v, \theta)$ and thus $\delta_k = \delta^*(v_k, \theta_k)$.

We start with several technical Lemmas showing some basic function properties.

### C.1  Technical Lemmas

**Lemma 1.** *Under Assumption 1, for any $v, \theta$, $g(v, \theta) - g(v, \theta^*(v)) \geq \frac{\kappa}{4}||\theta - \theta^*(v)||^2$.*

**Lemma 2.** *Under Assumption 1 and 2, we have* $||\nabla q(v,\theta) - \hat{\nabla} q(v,\theta)|| \leq L||\theta^{(T)} - \theta^*(v)||$ *for any* $v, \theta$. *Also, when* $||\hat{\nabla} q(v,\theta)|| = 0$, $q(v,\theta) = 0$.

**Lemma 3.** *Under Assumption 2, 1, we have* $||\theta^*(v_2) - \theta^*(v_1)|| \leq \frac{2L}{\kappa}||v_1 - v_2||$.

**Lemma 4.** *Under Assumption 2, we have* $||\nabla_\theta q(v, \theta_1) - \nabla_\theta q(v, \theta_2)|| \leq L||\theta_1 - \theta_2||$, *for any* $v$. *Further assume Assumption 1, we have*

$$||\nabla q(v_1, \theta_1) - \nabla q(v_2, \theta_2)|| \leq L_q ||[v_1, \theta_1] - [v_2, \theta_2]||,$$

*where* $L_q := 2L(L/\kappa + 1)$.

**Lemma 5.** *Under Assumption 1, 2 and assume that* $\alpha < 2/L$. *Given any* $v, \theta$, *suppose* $\theta^{(0)} = \theta$ *and* $\theta^{(t+1)} = \theta^{(t)} - \alpha \nabla_\theta q(v, \theta^{(t)})$, *then for any* $t$, *we have* $q(v, \theta^{(t)}) \leq \exp(-b_1(\alpha, L, \kappa)t)q(v, \theta)$, *where* $b_1(\alpha, L, \kappa) =$ *is some strictly positive constant that depends on* $\alpha$, $L$ *and* $\kappa$.

**Lemma 6.** *Under Assumption 3, for any* $[v, \theta]$, *we have* $||\delta^*(v, \theta)||, ||\nabla q(v, \theta)||, ||\hat{\nabla} q(v, \theta)|| \leq b_2(M, \eta)$, *where* $b_2(M, \eta) = (3 + \eta)M$.

**Lemma 7.** *Under Assumption 3, for any* $[v, \theta]$, *we have* $\lambda^* ||\hat{\nabla} q||^2 \leq \eta ||\hat{\nabla} q||^2 + M||\hat{\nabla} q||$, *where* $\lambda^*$ *are defined in (16).*

**Lemma 8.** *Under Assumption 1 and 2, we have* $||\nabla q(v, \theta)|| \leq 2\kappa^{-1/2} L_q q^{1/2}(v, \theta)$.

### C.1.1 Lemmas

Now we give several main lemmas that are used to prove the result in Section 4.2.

**Lemma 9.** *Under Assumption 1, 2 and 3, when* $||\hat{\nabla} q(v_k, \theta_k)|| > 0$, *we have*

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi\eta||\nabla q(v_k, \theta_k)||^2 + \xi\eta L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \left(L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2L_q||\theta_k - \theta^*(v_k)||\right)$$
$$+ \xi b_2 L||\theta_k^{(T)} - \theta^*(v_k)|| + L_q \xi^2 b_2^2/2.$$

*When* $||\hat{\nabla} q(v_k, \theta_k)|| = 0$, *we have* $q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq \xi^2 L_q b_2^2/2$.

**Lemma 10.** *Under Assumption 1, 2 and 3, choosing* $T \geq b_3(\eta, r, \kappa, L)$, *we have*

$$q(v_k, \theta_k) \leq \exp(-b_4 k)q(v_0, \theta_0) + \Delta,$$

*where* $b_4 = -\log(1 - \frac{\xi}{4}\eta\kappa)$ *is some strictly positive constant and* $\Delta = O(\exp(-b_1 T) + \xi)$.

**Lemma 11.** *Under Assumption 1, 2 and 3, we have*

$$\sum_{k=0}^{K-1} ||\nabla q(v_k, \theta_k)||^2 \leq \frac{b_5 q(v_0, \theta_0)}{\xi} + K\xi^2 b_6 \Delta,$$

*where* $b_5$ *is some constant depends on* $L_q, \eta, \kappa$; $b_6$ *is some constant depends on* $\kappa, L$ *and* $\Delta$ *is defined in Lemma 10.*

**Lemma 12.** *Under Assumption 1, 2 and 3, choosing* $T \geq b_3(\eta, r, \kappa, L)$ *and assume that* $r, \xi \leq 1/L$, *we have*

$$\sum_{k=0}^{K-1} \left[||\delta^*(v_k, \theta_k)||^2 + q(v_k, \theta_k)\right] = O(\xi^{-1} + K\exp(-b_1 T/2) + K\xi^{1/2} + \xi^{-1/2}K^{1/2}q^{1/2}(v_0, \theta_0)).$$

### C.2 Proof of Theorem 1

Using our definition of $\lambda^*$ in (16), we have

$$||\nabla f(v, \theta) + \lambda^*(v, \theta)\nabla q(v, \theta)|| \leq ||\nabla f(v, \theta) + \lambda^*(v, \theta)\hat{\nabla} q(v, \theta)|| + ||\lambda^*(v, \theta)(\hat{\nabla} q(v, \theta) - \nabla q(v, \theta))||$$
$$= ||\delta^*(v, \theta)|| + ||\lambda^*(v, \theta)(\hat{\nabla} q(v, \theta) - \nabla q(v, \theta))||.$$

669 Using Lemma 2, we know that when $||\hat{\nabla}q|| = 0$, we have $q = 0$ and thus $||\nabla q|| = 0$. In this case,
670 $||\lambda^*(\hat{\nabla}q - \nabla q)|| = 0$. When $||\hat{\nabla}q|| > 0$, some algebra shows that

$$
\begin{aligned}
||\lambda^*(\hat{\nabla}q - \nabla q)|| &\leq \left[\eta - \left\langle \nabla f, \hat{\nabla}q/||\hat{\nabla}q|| \right\rangle ||\hat{\nabla}q||^{-1}\right] ||\hat{\nabla}q - \nabla q|| \\
&\leq (\eta - \left\langle \nabla f, \hat{\nabla}q/||\hat{\nabla}q|| \right\rangle ||\hat{\nabla}q||^{-1})||\hat{\nabla}q - \nabla q||.
\end{aligned}
$$

671 Notice that

$$
\begin{aligned}
||\hat{\nabla}q(v,\theta) - \nabla q(v,\theta)|| &\leq L||\theta^{(T)} - \theta^*(v)|| \\
&\leq 2L\kappa^{-1/2}q^{1/2}(v,\theta^{(T)}) \\
&\leq 2L\kappa^{-1/2}\exp(-b_1 T/2)q^{1/2}(v,\theta) \\
&\leq 2L\kappa^{-1}\exp(-b_1 T/2)||\nabla q(v,\theta)||.
\end{aligned}
$$

672 Here the first inequality is by Lemma 2, the second inequality is by Lemma 1, the third inequality is by
673 Lemma 5 and the last inequality is by Assumption 1 (using $||\nabla q(v,\theta)|| \geq ||\nabla_\theta g(v,\theta)||$). Similarly,
674 under assumption that $T \geq \lceil -b_1^{-1}\log(\frac{1}{16}\kappa^2 L^{-2}) \rceil$, $L\kappa^{-1}\exp(-b_1 T/2) \leq 1/4$,

$$
\begin{aligned}
||\hat{\nabla}q(v,\theta)|| &= ||\hat{\nabla}q(v,\theta) - \nabla q(v,\theta) + \nabla q(v,\theta)|| \\
&\geq ||\nabla q(v,\theta)|| - ||\hat{\nabla}q(v,\theta) - \nabla q(v,\theta)|| \\
&\geq ||\nabla q(v,\theta)||(1 - (2L\kappa^{-1}\exp(-b_1 T/2))) \\
&\geq \frac{1}{2}||\nabla q(v,\theta)||.
\end{aligned}
$$

675 This implies that

$$
\frac{||\hat{\nabla}q - \nabla q||}{||\hat{\nabla}q||} \leq 2\frac{||\hat{\nabla}q - \nabla q||}{||\nabla q||} \leq 4L\kappa^{-1}\exp(-b_1 T/2).
$$

676 We thus have

$$
\begin{aligned}
||\lambda^*(v,\theta)(\hat{\nabla}q(v,\theta) - \nabla q(v,\theta))|| &\leq \eta||\hat{\nabla}q - \nabla q|| + \left\langle \nabla f, \frac{\hat{\nabla}q}{||\hat{\nabla}q||} \right\rangle \frac{||\hat{\nabla}q - \nabla q||}{||\hat{\nabla}q||} \\
&\leq 2L\kappa^{-1}\exp(-b_1 T/2)\left[\eta||\nabla q(v,\theta)|| + 2\left\langle \nabla f, \frac{\hat{\nabla}q}{||\hat{\nabla}q||} \right\rangle\right] \\
&\leq 2L\kappa^{-1}\exp(-b_1 T/2)(\eta + 2)b_2,
\end{aligned}
$$

677 where the last inequality is by Lemma 6. Combining all the results and using $||\nabla q(v_k,\theta_k)|| \leq$
678 $2\kappa^{-1/2}L_q q^{1/2}(v_k,\theta_k)$ by Lemma 8, we have

$$
\begin{aligned}
\mathcal{K}(v,\theta) &\leq ||\nabla f(v,\theta) + \lambda^*(v,\theta)\nabla q(v,\theta)||^2 + q(v,\theta) \\
&\leq 2||\nabla f(v,\theta) + \lambda^*(v,\theta)\hat{\nabla}q(v,\theta)||^2 + q(v,\theta) + 2||\lambda^*(v,\theta)(\hat{\nabla}q(v,\theta) - \nabla q(v,\theta))||^2 \\
&\leq 2||\delta^*(v,\theta)||^2 + q(v,\theta) + 8L^2\kappa^{-2}\exp(-b_1 T)(\eta + 2)^2 b_2^2.
\end{aligned}
$$

679 Using Lemma 12, we have

$$
\begin{aligned}
\min_k \mathcal{K}(v_k,\theta_k) &= O(\min_k(||\delta^*(v_k,\theta_k)||^2 + q(v_k,\theta_k)) + \exp(-b_1 T)) \\
&= O(\xi^{-1} + K\exp(-b_1 T/2) + K\xi^{1/2} + \xi^{-1/2}K^{1/2}q^{1/2}(v_0,\theta_0)).
\end{aligned}
$$

## C.3 Proof of Lemmas

### C.3.1 Proof of Lemma 9

When $||\hat{\nabla}q(v_k, \theta_k)|| > 0$, by Lemma 4, we know that $q$ is $L_q$-smoothness, we have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi \langle \nabla q(v_k, \theta_k), \delta^*(v_k, \theta_k) \rangle + \frac{L_q \xi^2}{2} ||\delta^*(v_k, \theta_k)||^2$$

$$\leq -\xi \left\langle \hat{\nabla}q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle - \xi \left\langle \nabla q(v_k, \theta_k) - \hat{\nabla}q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + L_q \xi^2 b_2^2/2$$

$$\leq -\xi\eta ||\hat{\nabla}q(v_k, \theta_k)||^2 - \xi \left\langle \nabla q(v_k, \theta_k) - \hat{\nabla}q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + L_q \xi^2 b_2^2/2$$

$$\leq -\xi\eta ||\hat{\nabla}q(v_k, \theta_k)||^2 + \xi b_2 ||\nabla q(v_k, \theta_k) - \hat{\nabla}q(v_k, \theta_k)|| + L_q \xi^2 b_2^2/2.$$

where the second and the last inequality is by Lemma 6 and the third inequality is ensured by
the constraint in the local subproblem ($\left\langle \nabla \hat{\nabla}q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle \geq \eta ||\hat{\nabla}q(v_k, \theta_k)^2||$.). And by
Lemma 2, we have $||\nabla q(v_k, \theta_k) - \hat{\nabla}q(v_k, \theta_k)|| \leq L||\theta_k^{(T)} - \theta^*(v_k)||$. Plug in the bound we have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi\eta ||\hat{\nabla}q(v_k, \theta_k)||^2 + \xi b_2 L ||\theta_k^{(T)} - \theta^*(v_k)|| .$$

Also notice that

$$\left| ||\hat{\nabla}q(v_k, \theta_k)||^2 - ||\nabla q(v_k, \theta_k)||^2 \right| \leq ||\hat{\nabla}q(v_k, \theta_k) - \nabla q(v_k, \theta_k)|| \, ||\hat{\nabla}q(v_k, \theta_k) + \nabla q(v_k, \theta_k)||$$

$$\leq ||\hat{\nabla}q(v_k, \theta_k) - \nabla q(v_k, \theta_k)|| \, (||\hat{\nabla}q(v_k, \theta_k) - \nabla q(v_k, \theta_k)|| + 2||\nabla q(v_k, \theta_k)||)$$

$$\leq L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \, (L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2||\nabla q(v_k, \theta_k)||)$$

$$= L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \, (L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2||\nabla q(v_k, \theta_k) - \nabla q(v_k, \theta^*(v_k))||)$$

$$\leq L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \, (L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2L_q ||\theta_k - \theta^*(v_k)||),$$

where the third inequality is by Lemma 2, the equality is by $\nabla q(v_k, \theta^*(v_k)) = 0$ and the last
inequality is by Lemma 4.

Using this bound, we further have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi\eta ||\nabla q(v_k, \theta_k)||^2 + \xi\eta \left| ||\hat{\nabla}q(v_k, \theta_k)||^2 - ||\nabla q(v_k, \theta_k)||^2 \right|$$

$$+ \xi b_2 ||\nabla q(v_k, \theta_k) - \hat{\nabla}q(v_k, \theta_k)|| + L_q \xi^2 b_2^2/2$$

$$\leq -\xi\eta ||\nabla q(v_k, \theta_k)||^2 + \xi\eta L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \, (L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2L_q ||\theta_k - \theta^*(v_k)||)$$

$$+ \xi b_2 L ||\theta_k^{(T)} - \theta^*(v_k)|| + L_q \xi^2 b_2^2/2.$$

When $||\hat{\nabla}q(v_k, \theta_k)|| = 0$, by Lemma 2, $q(v_k, \theta_k) = 0$ and hence $\nabla q(v_k, \theta_k) = 0$. We thus have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi \langle \nabla q(v_k, \theta_k), \delta^*(v_k, \theta_k) \rangle + \frac{L_q \xi^2}{2} ||\delta^*(v_k, \theta_k)||^2$$

$$= \frac{L_q \xi^2}{2} ||\delta^*(v_k, \theta_k)||^2$$

$$\leq \xi^2 L_q b_2^2/2.$$

### C.3.2 Proof of Lemma 10

By Lemma 9, when $||\hat{\nabla}q(v_k, \theta_k)|| > 0$, we have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi\eta ||\nabla q(v_k, \theta_k)||^2 + \xi\eta L_q ||\theta_k^{(T)} - \theta^*(v_k)|| \, (L_q ||\theta_k^{(T)} - \theta^*(v_k)|| + 2L_q ||\theta_k - \theta^*(v_k)||)$$

$$+ \xi b_2 L ||\theta_k^{(T)} - \theta^*(v_k)|| + L_q \xi^2 b_2^2/2.$$

By Lemma 1 and Lemma 5

$$||\theta_k^{(T)} - \theta^*(v_k)|| \leq 2\kappa^{-1/2} q^{1/2}(v_k, \theta_k^{(T)}) \leq 2\kappa^{-1/2} \exp(-b_1 T/2) q^{1/2}(v_k, \theta_k).$$

$$||\theta_k - \theta^*(v_k)|| \leq 2\kappa^{-1/2} q^{1/2}(v_k, \theta_k).$$

694    Using those bounds, we know that

$$L_q||\theta_k^{(T)} - \theta^*(v_k)|| \left(L_q||\theta_k^{(T)} - \theta^*(v_k)|| + 2L_q||\theta_k - \theta^*(v_k)||\right) \leq 12L_q^2\kappa^{-1}\exp(-b_1 T)q(v_k, \theta_k)$$

695    This implies that

$$\begin{aligned}
&q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \\
&\leq -\xi\eta||\nabla q(v_k, \theta_k)||^2 + 12\xi\eta L_q^2\kappa^{-1}\exp(-b_1 T)q(v_k, \theta_k) \\
&\quad + 2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2)q^{1/2}(v_k, \theta_k) + L_q\xi^2 b_2^2/2 \\
&\leq -\xi\eta\kappa q(v_k, \theta_k) + 12\xi\eta L_q^2\kappa^{-1}\exp(-b_1 T)q(v_k, \theta_k) \\
&\quad + 2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2)q^{1/2}(v_k, \theta_k) + L_q\xi^2 b_2^2/2.
\end{aligned}$$

696    Choosing $T$ such that $T \geq b_3(\eta, \alpha, \kappa, L)$ where

$$b_3(\eta, \alpha, \kappa, L) = \left\lceil -b_1^{-1}\log(\frac{\eta\kappa}{64\eta L_q^2}) \right\rceil,$$

697    we have

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\frac{3}{4}\xi\eta\kappa q(v_k, \theta_k) + 2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2)q^{1/2}(v_k, \theta_k) + L_q\xi^2 b_2^2/2.$$

698    This implies that when $\frac{64b_2^2 L^2}{\eta^2\kappa}\exp(-b_1 T) \leq q(v_k, \theta_k)$ and $\frac{2L_q\xi b_2^2}{\eta\kappa} \leq q(v_k, \theta_k)$,

$$q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\frac{1}{4}\xi\eta\kappa q(v_k, \theta_k).$$

699    Let $a = \max(\frac{64b_2^2 L^2}{\eta^2\kappa}\exp(-b_1 T), \frac{2L_q\xi b_2^2}{\eta\kappa})$. Also, when $q(v_k, \theta_k) < a$,

$$\begin{aligned}
q(v_{k+1}, \theta_{k+1}) &\leq q(v_k, \theta_k) + 2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2)q^{1/2}(v_k, \theta_k) + L_q\xi^2 b_2^2/2 \\
&< a + 2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2)\sqrt{a} + L_q\xi^2 b_2^2/2.
\end{aligned}$$

700    Note that

$$2\xi b_2 L\kappa^{-1/2}\exp(-b_1 T/2) \leq \frac{\xi\eta\kappa}{4}\sqrt{a}$$

$$L_q\xi^2 b_2^2/2 \leq \frac{\xi\eta\kappa}{4}a.$$

701    This gives that in the case of $q(v_k, \theta_k) < a$,

$$q(v_{k+1}, \theta_{k+1}) < (1 + \frac{\xi\eta\kappa}{4})a.$$

702    Define $k_0$ as the first iteration such that $q(v_k, \theta_k) < a$. This implies that, for any $k \leq k_0$,

$$q(v_k, \theta_k) \leq (1 - \frac{\xi}{4}\eta\kappa)^k q(v_0, \theta_0).$$

703    When any $k > k_0$, we show that $q(v_{k+1}, \theta_{k+1}) \leq (1 + \frac{\xi\eta\kappa}{4})a$. This can be proved by induction.
704    At $k = k_0 + 1$, if $q(v_k, \theta_k) < a$, we have $q(v_k, \theta_k) < (1 + \frac{\xi\eta\kappa}{4})a$. Else if at $k = k_0 + 1$,
705    $q(v_k, \theta_k) \geq a$, $q(v_{k+1}, \theta_{k+1}) \leq q(v_k, \theta_k) \leq a$. We thus have the conclusion that for any $k > k_0$,
706    $q(v_k, \theta_k) \leq (1 + \frac{\eta\kappa}{4})a$. Combining the result, we have

$$q(v_k, \theta_k) \leq (1 - \frac{\xi}{4}\eta\kappa)^k q(v_0, \theta_0) + \Delta,$$

707    where we denote

$$\Delta = (1 + \frac{\eta\kappa}{4})(\frac{64b_2^2 L^2}{\eta^2\kappa^3}\exp(-b_1 T) + \frac{2L_q\xi b_2^2}{\eta\kappa}) + L_q\xi^2 b_2^2/2 = O(\exp(-b_1 T) + \xi). \qquad (17)$$

708    Let $b_4(\eta, \kappa, \xi) = -\log(1 - \frac{\xi}{4}\eta\kappa)$, we have the desired result.

 ### C.3.3 Proof of Lemma 11

 By Lemma 8 and 10, we have

$$
\begin{aligned}
||\nabla q(v_k, \theta_k)||^2 &\le 2\kappa^{-1} L_q^2 q(v_k, \theta_k) \\
&\le 2\kappa^{-1} L_q^2 \left[ \exp(-b_4 k) q(v_0, \theta_0) + \Delta \right],
\end{aligned}
$$

 where $\Delta$ is defined in (17). Also notice that

$$
\begin{aligned}
||\hat{\nabla} q(v, \theta)|| &\le ||\hat{\nabla} q(v, \theta) - \nabla q(v, \theta)|| + ||\nabla q(v, \theta)|| \\
&\le L||\theta^{(T)} - \theta^*(v)|| + ||\nabla q(v, \theta)|| \\
&\le 2L\kappa^{-1/2} q^{1/2}(v, \theta^{(T)}) + ||\nabla q(v, \theta)|| \\
&\le 2L\kappa^{-1/2} \exp(-b_1 T/2) q^{1/2}(v, \theta) + ||\nabla q(v, \theta)|| \\
&\le (2L\kappa^{-1} \exp(-b_1 T/2) + 1)||\nabla q(v, \theta)|| \\
&\le (2L\kappa^{-1} + 1)||\nabla q(v, \theta)||
\end{aligned}
$$

 Here the first inequality is by triangle inequality, the second inequality is by Lemma 2, the third
 inequality is by Lemma 1, the forth inequality is by Lemma 5 and the fifth inequality is by Assumption
 1. Taking summation over iteration and using Lemma 10, we have

$$
\begin{aligned}
\sum_{k=0}^{K-1} ||\hat{\nabla} q(v, \theta)||^2 &\le (2L\kappa^{-1} + 1)^2 \sum_{k=0}^{K-1} ||\nabla q(v_k, \theta_k)||^2 \\
&\le (2L\kappa^{-1} + 1)^2 \left[ 2\kappa^{-1} L_q^2 q(v_0, \theta_0) \sum_{k=0}^{K-1} [\exp(-b_4 k)] + K\Delta \right] \\
&\le (2L\kappa^{-1} + 1)^2 \left[ \frac{2\kappa^{-1} L_q^2 q(v_0, \theta_0)}{1 - \exp(-b_4)} + K\Delta \right] \\
&= \frac{b_5 q(v_0, \theta_0)}{\xi} + K b_6 \Delta,
\end{aligned}
$$

 where we define $b_5(L_q, \eta, \kappa) = \frac{16 L_q^2}{\eta \kappa^2} (2L\kappa^{-1} + 1)^2$ and $b_6(\kappa, L) = (2L\kappa^{-1} + 1)^2$.

 ## C.4 Proof of Lemma 12

 Remind that by our definition of $\lambda^*$ in (16) and Assumption 2, we have

$$
\begin{aligned}
f(v_{k+1}, \theta_{k+1}) - f(v_k, \theta_k) &\le -\xi \left\langle \nabla f(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + \frac{L\xi^2}{2} ||\delta^*(v_k, \theta_k)||^2 \\
&= -\xi \left\langle \delta^*(v_k, \theta_k) - \lambda^*(v_k, \theta_k)\hat{\nabla} q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + \frac{L\xi^2}{2} ||\delta^*(v_k, \theta_k)||^2 \\
&= -(\xi - \frac{L\xi^2}{2})||\delta^*(v_k, \theta_k)||^2 + \xi\lambda^*(v_k, \theta_k) \left\langle \hat{\nabla} q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle \\
&\le -(\xi - \frac{L\xi^2}{2})||\delta^*(v_k, \theta_k)||^2 + \xi\eta\lambda^*(v_k, \theta_k)||\hat{\nabla} q(v_k, \theta_k)||^2 \\
&\le -\frac{\xi}{2}||\delta^*(v_k, \theta_k)||^2 + \xi\eta\lambda^*(v_k, \theta_k)||\hat{\nabla} q(v_k, \theta_k)||^2,
\end{aligned}
$$

 where the last inequality is by the assumption on $\xi \le 1/L$. To show the second inequality, we use the
 complementary slackness of Problem (15), that is

$$
\lambda^*(v_k, \theta_k) \left[ \left\langle \hat{\nabla} q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle - \eta ||\hat{\nabla} q(v_k, \theta_k)|| \right] = 0.
$$

By telescoping,

$$\sum_{k=0}^{K-1} f(v_{k+1}, \theta_{k+1}) - f(v_k, \theta_k) \leq -\frac{\xi}{2} \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 + \xi\eta \sum_{k=0}^{K-1} \lambda^*(v_k, \theta_k)||\hat{\nabla}q(v_k, \theta_k)||^2$$

$$\leq -\frac{\xi}{2} \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 + \xi\eta \sum_{k=0}^{K-1} (\eta||\hat{\nabla}q(v_k, \theta_k)||^2 + M||\hat{\nabla}q(v_k, \theta_k)||)$$

$$= -\frac{\xi}{2} \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 + \xi\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2 + \xi\eta M \sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||$$

$$\leq -\frac{\xi}{2} \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 + \xi\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2 + \xi\eta M\sqrt{K}\sqrt{\sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2},$$

where the second inequality is by Lemma 7 and the last inequality is by Holder's inequality. Since $\sum_{k=0}^{K-1} f(v_{k+1}, \theta_{k+1}) - f(v_k, \theta_k) = f(v_K, \theta_K) - f(v_0, \theta_0)$, rearrange the terms, we have

$$\xi \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 \leq 2(f(v_0, \theta_0) - f(v_K, \theta_K)) + 2\xi\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2 + 2\xi\eta M\sqrt{K}\sqrt{\sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2}.$$

This implies that

$$\xi \sum_{k=0}^{K-1} \left[||\delta^*(v_k, \theta_k)||^2 + q(v_k, \theta_k)\right] \leq 2(f(v_0, \theta_0) - f(v_K, \theta_K)) + 2\xi\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2$$

$$+ 2\xi\eta M\sqrt{K}\sqrt{\sum_{k=0}^{K-1} ||\hat{\nabla}q(v_k, \theta_k)||^2} + \xi \sum_{k=0}^{K-1} q(v_k, \theta_k).$$

Using Lemma 10, we know that

$$q(v_k, \theta_k) \leq (1 - \frac{\xi}{4}\eta\kappa)^k q(v_0, \theta_0) + \Delta.$$

This gives that

$$\xi \sum_{k=0}^{K-1} q(v_k, \theta_k) \leq \frac{4q(v_0, \theta_0)}{\eta\kappa} + \xi K\Delta.$$

Using Lemma 11, 10 and $\sqrt{x+y} \leq \sqrt{x} + \sqrt{y}$, we have

$$2\xi\eta^2 \sum_{k=1}^{K} ||\hat{\nabla}q(v_k, \theta_k)||^2 \leq 2\eta^2 b_5 q(v_0, \theta_0) + 2K\eta^2\xi b_6\Delta$$

$$2\xi\eta M\sqrt{K}\sqrt{\sum_{k=1}^{K} ||\hat{\nabla}q(v_k, \theta_k)||^2} \leq 2\xi^{1/2}K^{1/2}b_5^{1/2}\eta Mq^{1/2}(v_0, \theta_0) + 2K\xi b_6^{1/2}\eta M\Delta^{1/2}$$

This implies that

$$\xi \sum_{k=0}^{K-1} \left[||\delta^*(v_k, \theta_k)||^2 + q(v_k, \theta_k)\right]$$

$$\leq 2(f(v_0, \theta_0) - f(v_K, \theta_K)) + 2\eta^2 b_5 q(v_0, \theta_0) + 2K\eta^2\xi b_6\Delta + 2\xi^{1/2}K^{1/2}b_5^{1/2}\eta Mq^{1/2}(v_0, \theta_0)$$

$$+ 2K\xi b_6^{1/2}\eta M\Delta^{1/2} + \frac{4q(v_0, \theta_0)}{\eta\kappa} + \xi K\Delta$$

$$\leq 2(f(v_0, \theta_0) - f(v_K, \theta_K)) + (2\eta^2 b_5 + \frac{4}{\eta\kappa})q(v_0, \theta_0) + 2K\xi(b_6^{1/2}\eta M\Delta^{1/2} + (b_6\eta^2 + 1/2)\Delta)$$

$$+ 2\xi^{1/2}K^{1/2}b_5^{1/2}\eta Mq^{1/2}(v_0, \theta_0)$$

We thus have

$$\sum_{k=0}^{K-1} \left[ ||\delta^*(v_k, \theta_k)||^2 + q(v_k, \theta_k) \right]$$

$$= O(\xi^{-1} + K\Delta^{1/2} + \xi^{-1/2}K^{1/2}q^{1/2}(v_0, \theta_0))$$

$$= O(\xi^{-1} + K\exp(-b_1 T/2) + K\xi^{1/2} + \xi^{-1/2}K^{1/2}q^{1/2}(v_0, \theta_0)).$$

## C.5  Proofs of Technical Lemmas

### C.5.1  Proof of Lemma 1

Please see the proof of Theorem 2 in Karimi et al. [24].

### C.5.2  Proof of Lemma 2

Since $\nabla_2 g(v, \theta^*(v)) = 0$, we have $\nabla_v g(v, \theta^*(v)) = \nabla_1 g(v, \theta^*(v)) + \nabla_v \theta^*(v) \nabla_2 g(v, \theta^*(v)) = \nabla_1 g(v, \theta^*(v))$. Thus

$$\nabla q(v, \theta) = \left[ \begin{array}{c} \nabla_v g(v, \theta) - \nabla_v g(v, \theta^*(v)) \\ \nabla_\theta g(v, \theta) \end{array} \right] = \left[ \begin{array}{c} \nabla_v g(v, \theta) - \nabla_1 g(v, \theta^*(v)) \\ \nabla_\theta g(v, \theta) \end{array} \right].$$

Also note that

$$\hat{\nabla} q(v, \theta) = \left[ \begin{array}{c} \nabla_v g(v, \theta) - \nabla_1 g(v, \theta^{(T)}) \\ \nabla_\theta g(v, \theta) \end{array} \right].$$

This gives that

$$||\nabla q(v, \theta) - \hat{\nabla} q(v, \theta)|| = ||\nabla_1 g(v, \theta^{(T)}) - \nabla_1 g(v, \theta^*(v))||$$

$$\leq L||\theta^{(T)} - \theta^*(v)||.$$

Also when $0 = ||\hat{\nabla} q(v, \theta)|| = \sqrt{||\nabla_v g(v, \theta) - \nabla_1 g(v, \theta^{(T)})||^2 + ||\nabla_\theta g(v, \theta)||^2}$, we have $||\nabla_\theta g(v, \theta)|| = 0$. Under Assumption 1,

$$0 = ||\nabla_\theta g(v, \theta)|| \geq \kappa(g(v, \theta) - g(v, \theta^*(v))) = \kappa q(v, \theta).$$

### C.5.3  Proof of Lemma 3

Using Assumption 1 and $\nabla_2 g(v_1, \theta^*(v_1)) = 0$, we have

$$||\nabla_2 g(v_1, \theta^*(v_2))|| \geq \sqrt{\kappa(g(v_1, \theta^*(v_2)) - g(v_1, \theta^*(v_1)))}.$$

Also by Lemma 1, we have $g(v_1, \theta^*(v_2)) - g(v_1, \theta^*(v_1)) \geq \frac{1}{4}\kappa||\theta^*(v_2) - \theta^*(v_1)||^2$. These imply that

$$||\nabla_2 g(v_1, \theta^*(v_2))|| \geq \frac{1}{2}\kappa||\theta^*(v_2) - \theta^*(v_1)||.$$

Also

$$||\nabla_2 g(v_1, \theta^*(v_2))||$$

$$= ||\nabla_2 g(v_1, \theta^*(v_2)) - \nabla_\theta g(v_2, \theta^*(v_2))||$$

$$= ||\nabla_2 [g(v_1, \theta^*(v_2)) - g(v_2, \theta^*(v_2))]||$$

$$\leq ||\nabla_{[1,2]} [g(v_1, \theta^*(v_2)) - g(v_2, \theta^*(v_2))]||$$

$$\leq L||v_1 - v_2||,$$

where $\nabla_{[1,2]}$ denotes taking the derivative on both first and second variables. We thus conclude that

$$||\theta^*(v_2) - \theta^*(v_1)|| \leq \frac{2L}{\kappa}||v_1 - v_2||.$$

### C.5.4 Proof of Lemma 4

To prove the first property,

$$||\nabla_\theta q(v, \theta_1) - \nabla_\theta q(v, \theta_2)|| = ||\nabla_\theta g(v, \theta_1) - \nabla_\theta g(v, \theta_2)||$$
$$\leq L||\theta_1 - \theta_2||.$$

Also

$$||\nabla q(v_1, \theta_1) - \nabla q(v_2, \theta_2)|| = ||\nabla g(v_1, \theta_1) - \nabla g(v_2, \theta_2) - \nabla g(v_1, \theta^*(v_1)) + \nabla g(v_2, \theta^*(v_2))||$$
$$\leq ||\nabla g(v_1, \theta_1) - \nabla g(v_2, \theta_2)|| + ||\nabla_1 g(v_1, \theta^*(v_1)) - \nabla_1 g(v_2, \theta^*(v_2))||.$$

By Assumption 2 (Lipschitz continuity of $\nabla g$),

$$||\nabla_1 g(v_1, \theta^*(v_1)) - \nabla_1 g(v_2, \theta^*(v_2))|| \leq ||\nabla_{[1,2]} g(v_1, \theta^*(v_1)) - \nabla_{[1,2]} g(v_2, \theta^*(v_2))||$$
$$\leq L\sqrt{||\theta^*(v_1) - \theta^*(v_2)||^2 + ||v_1 - v_2||^2},$$

where $\nabla_{[1,2]}$ denotes taking the derivative on both first and second variable. Also By Lemma 3,

$$L\sqrt{||\theta^*(v_1) - \theta^*(v_2)||^2 + ||v_1 - v_2||^2} \leq L\sqrt{\frac{4L^2}{\kappa^2}||v_1 - v_2||^2 + ||v_1 - v_2||^2}$$
$$\leq L(\frac{2L}{\kappa} + 1)||v_1 - v_2||.$$

This gives that

$$||\nabla q(v_1, \theta_1) - \nabla q(v_2, \theta_2)|| \leq ||\nabla g(v_1, \theta_1) - \nabla g(v_2, \theta_2)|| + ||\nabla_1 g(v_1, \theta^*(v_1)) - \nabla_1 g(v_2, \theta^*(v_2))||$$
$$\leq L\sqrt{||v_1 - v_2||^2 + ||\theta_1 - \theta_2||^2} + ||\nabla_1 g(v_1, \theta^*(v_1)) - \nabla_1 g(v_2, \theta^*(v_2))||$$
$$\leq L\sqrt{||v_1 - v_2||^2 + ||\theta_1 - \theta_2||^2} + L(\frac{2L}{\kappa} + 1)||v_1 - v_2||$$
$$\leq L_q\sqrt{||v_1 - v_2||^2 + ||\theta_1 - \theta_2||^2},$$

where $L_q := 2L(L/\kappa + 1)$.

### C.5.5 Proof of Lemma 5

By Lemma 4, we have

$$q(v, \theta^{(t+1)}) - q(v, \theta^{(t)}) \leq -(\alpha - \frac{L\alpha^2}{2})||\nabla_\theta q(v, \theta^{(t)})||^2.$$

By Assumption 1, we have

$$||\nabla_\theta q(v, \theta^{(t)})||^2 = ||\nabla_2 g(v, \theta^{(t)})||^2 \geq \kappa(g(v, \theta^{(t)}) - g(v, \theta^*(v)) = \kappa q(v, \theta^{(t)}).$$

Plug-in, we have

$$q(v, \theta^{(t+1)}) \leq (1 - (\alpha - \frac{L\alpha^2}{2})\kappa)q(v, \theta^{(t)}).$$

Recursively apply this inequality, we have

$$q(v, \theta^{(t)}) \leq (1 - (\alpha - \frac{L\alpha^2}{2})\kappa)^t q(v, \theta).$$

Let $b_1(r, L, \kappa) = \log(1 - (\alpha - L\alpha^2/2)\kappa)$, we have the desired result.

### C.5.6 Proof of Lemma 6

Notice that $||\nabla q(v, \theta)|| \leq ||\nabla g(v, \theta)|| + ||\nabla g(v, \theta^*(v))|| \leq 2M$. $||\hat{\nabla} q(v, \theta)|| \leq ||\nabla_v g(v, \theta)|| + ||\nabla_1 g(v, \theta^{(T)})|| + ||\nabla_\theta g(v, \theta)|| \leq 3M$. When $||\hat{\nabla} q|| = 0$, $||\delta^*|| = ||\nabla f|| \leq M$. When $||\hat{\nabla} q|| > 0$,

$$||\delta^*|| = ||[\eta||\hat{\nabla} q||^2 - \langle \nabla f, \hat{\nabla} q \rangle]_+ / ||\hat{\nabla} q||^2 \hat{\nabla} q + \nabla f||$$
$$\leq \eta||\hat{\nabla} q|| + 2||\nabla f|| \leq (2 + \eta)M.$$

This concludes that $||\delta^*|| \leq (2 + \eta)M$.

 **C.5.7 Proof of Lemma 7**

763 In the case that $\left\langle \nabla f, \hat{\nabla} q \right\rangle < \eta ||\hat{\nabla} q||^2$, $\lambda^* ||\hat{\nabla} q||^2 = \eta ||\hat{\nabla} q||^2 - \left\langle \nabla f, \hat{\nabla} q \right\rangle$. In the other case,

764 $\lambda^* ||\hat{\nabla} q||^2 = 0$. Thus in all cases,

$$\lambda^* ||\hat{\nabla} q||^2 \leq \eta ||\hat{\nabla} q||^2 + ||\nabla f|| \, ||\hat{\nabla} q||$$
$$\leq \eta ||\hat{\nabla} q||^2 + M ||\hat{\nabla} q||.$$

765 **C.5.8 Proof of Lemma 8**

766 Notice that since $\nabla q(v, \theta^*(v)) = 0$, we have

$$||\nabla q(v, \theta)|| = ||\nabla q(v, \theta) - \nabla q(v, \theta^*(v))|| \leq L_q ||\theta - \theta^*(v)|| \leq 2\kappa^{-1/2} L_q q^{1/2}(v, \theta),$$

767 where the first inequality is by Lemma 4 and the second inequality is by Lemma 1.

# D   Proof of the Result in Section 4.3

769 We use $b$ with some subscript to denote some general $O(1)$ constant and refer reader to section E for
770 their detailed value.

771 For notation simplicity, given $v$ and $\theta$, $\theta^{(T)}$ denotes the results of $T$ steps of gradient of $g(v, \cdot)$
772 w.r.t. $\theta$ starting from $\theta$ using step size $\alpha$ (similar to the definition in (8)). And note that $\hat{\nabla} q(v, \theta) =$
773 $\nabla g(v, \theta) - \left[ \nabla_1^\top g(v, \theta^{(T)}), \mathbf{0}^\top \right]^\top$, where $\mathbf{0}$ denotes a zero vector with the same dimension as $\theta$. We
774 refer readers to the beginning of Appendix C for a discussion on the design of this extra notation and
775 how it relates to the notation we used in Section 3. For simplicity, we omit the superscript $\diamond$ in $q^\diamond$ and
776 simply use $q$ to denote $q^\diamond$ in the proof.

777 We start with the following two Lemmas.

778 **Lemma 13.** *Under Assumption 4 and assume $\alpha \leq 1/L$, for any $v, \theta$, $g(v, \theta) - g(v, \theta^\diamond(v, \theta)) \geq$*
779 $\frac{\kappa}{4} ||\theta - \theta^\diamond(v, \theta)||^2$.

780 *Proof.* It is easy to show that

$$g(v, \theta^{(t+1)}) \leq g(v, \theta^{(t)}) - (\alpha - \frac{L\alpha^2}{2}) ||\nabla_\theta g(v, \theta^{(t)})||^2 \leq g(v, \theta^{(t)}).$$

781 We thus have $g(v, \theta^\diamond(v, \theta)) \leq g(v, \theta)$. The result of the proof follows the proof of Theorem 2 in
782 Karimi et al. [24]. $\square$

783 **Lemma 14.** *Under Assumption 2 and 4, $||\theta^\diamond(v_2, \theta) - \theta^\diamond(v_1, \theta)|| \leq \frac{4L}{\kappa} ||v_1 - v_2||$ for any $v_1, v_2$.*

784 *Proof.* Notice that $\nabla q(v_2, \theta^\diamond(v_2, \theta)) = 0$, we have

$$||\nabla q(v_1, \theta^\diamond(v_2, \theta)) - \nabla q(v_2, \theta^\diamond(v_2, \theta))|| = ||\nabla q(v_1, \theta^\diamond(v_2, \theta))||.$$

785 By Assumption 4, we have $||\nabla q(v_1, \theta^\diamond(v_2, \theta))|| \geq \sqrt{\kappa(g(v_1, \theta^\diamond(v_2, \theta)) - g(v_1, \theta^\diamond(v_1, \theta)))}$. And by
786 Lemma 13, we have

$$g(v_1, \theta^\diamond(v_2, \theta)) - g(v_1, \theta^\diamond(v_1, \theta)) \geq \frac{\kappa}{4} ||\theta^\diamond(v_2, \theta) - \theta^\diamond(v_1, \theta)||^2.$$

787 Combing all bounds gives that

$$2L||v_1 - v_2|| \geq ||\nabla q(v_1, \theta^\diamond(v_2, \theta)) - \nabla q(v_2, \theta^\diamond(v_2, \theta))|| = ||\nabla q(v_1, \theta^\diamond(v_2, \theta))|| \geq \frac{\kappa}{2} ||\theta^\diamond(v_2, \theta) - \theta^\diamond(v_1, \theta)||.$$

788 This implies that $||\theta^\diamond(v_2, \theta) - \theta^\diamond(v_1, \theta)|| \leq \frac{4L}{\kappa} ||v_1 - v_2||$. $\square$

789 Now we proceed to give the proof of Theorem 2.

790 Note that

$$
\begin{aligned}
q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) &= [g(v_{k+1}, \theta_{k+1}) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))] - [g(v_k, \theta_k) - g(v_k, \theta^\diamond(v_k, \theta_k))] \\
&= [g(v_{k+1}, \theta_{k+1}) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k))] - [g(v_k, \theta_k) - g(v_k, \theta^\diamond(v_k, \theta_k))] \\
&\quad + [g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))] \\
&= [g(v_{k+1}, \theta_{k+1}) - g(v_k, \theta_k)] - [g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_k, \theta^\diamond(v_k, \theta_k))] \\
&\quad + [g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))].
\end{aligned}
$$

791 Note that

$$
g(v_{k+1}, \theta_{k+1}) - g(v_k, \theta_k) \leq -\xi \left\langle \nabla g(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + \frac{L\xi^2}{2} ||\delta^*(v_k, \theta_k)||^2
$$

$$
-[g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_k, \theta^\diamond(v_k, \theta_k))] \leq \left\langle \nabla_{[1,2]} g(v_k, \theta^\diamond(v_k, \theta_k)), [v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)] - [v_k, \theta^\diamond(v_k, \theta_k)] \right\rangle
$$

$$
+ \frac{L}{2} ||[v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)] - [v_k, \theta^\diamond(v_k, \theta_k)]||^2.
$$

792 Notice that as $\nabla_2 g(v_k, \theta^\diamond(v_k, \theta_k)) = 0$,

$$
\left\langle \nabla_{[1,2]} g(v_k, \theta^\diamond(v_k, \theta_k)), [v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)] - [v_k, \theta^\diamond(v_k, \theta_k)] \right\rangle = \xi \left\langle \nabla_{[1,2]} g(v_k, \theta^\diamond(v_k, \theta_k)), \delta^*(v_k, \theta_k) \right\rangle.
$$

793 Also using Lemma 14, we have

$$
||\theta^\diamond(v_{k+1}, \theta_k) - \theta^\diamond(v_k, \theta_k)|| \leq \frac{4L}{\kappa} ||v_{k+1} - v_k||.
$$

794 This implies that

$$
||[v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)] - [v_k, \theta^\diamond(v_k, \theta_k)]||^2 \leq (\frac{16L^2}{\kappa^2} + 1)||v_{k+1} - v_k||^2 \leq (\frac{16L^2}{\kappa^2} + 1)\xi^2 ||\delta^*(v_k, \theta_k)||^2.
$$

795 We thus have

$$
q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\xi \left\langle \nabla q(v_k, \theta_k), \delta^*(v_k, \theta_k) \right\rangle + L_q \xi^2 ||\delta^*(v_k, \theta_k)||^2 / 2 + \chi_k,
$$

796 where we define $L_q = (\frac{16L^2}{\kappa^2} + 2)$ and $\chi_k = [g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))]$.

797 Using the same argument in the proof of Lemma 10 and Lemma 11, we have

$$
\begin{aligned}
&q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \\
&\leq -\xi\eta ||\nabla q(v_k, \theta_k)||^2 + 12\xi\eta L_q^2 \kappa^{-1} \exp(-b_1 T) q(v_k, \theta_k) \\
&\quad + 2\xi b_2 L \kappa^{-1/2} \exp(-b_1 T/2) q^{1/2}(v_k, \theta_k) + L_q \xi^2 b_2^2 / 2 + \chi_k \\
&\leq -\xi\eta ||\nabla q(v_k, \theta_k)||^2 + 12\xi\eta L_q^2 \kappa^{-2} \exp(-b_1 T) ||\nabla q(v_k, \theta_k)||^2 \\
&\quad + 2\xi b_2 L \kappa^{-1} \exp(-b_1 T/2) ||\nabla q(v_k, \theta_k)|| + L_q \xi^2 b_2^2 / 2 + \chi_k.
\end{aligned}
$$

798 Here the second inequality is by Assumption 4. Choosing $T$ such that $T \geq b_7(\eta, \alpha, \kappa, L)$ where

$$
b_7(\eta, \alpha, \kappa, L) = \left\lceil -b_1^{-1} \log(\frac{\kappa^2}{48\eta L_q^2}) \right\rceil,
$$

799 we have

$$
q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\frac{3}{4}\xi\eta ||\nabla q(v_k, \theta_k)||^2 + 2\xi b_2 L \kappa^{-1} \exp(-b_1 T/2) ||\nabla q(v_k, \theta_k)|| + L_q \xi^2 b_2^2 / 2 + \chi_k.
$$

800 Using Young's inequality, given any $x > 0$,

$$
\exp(-b_1 T/2) ||\nabla q(v_k, \theta_k)|| \leq x \exp(-b_1 T) + \frac{1}{x} ||\nabla q(v_k, \theta_k)||^2.
$$

801 Choosing $x = \frac{4Lb_2}{\eta\kappa}$, we have

$$
q(v_{k+1}, \theta_{k+1}) - q(v_k, \theta_k) \leq -\frac{1}{4}\xi\eta ||\nabla q(v_k, \theta_k)||^2 + \Delta + \chi_k,
$$

where we denote $\Delta = \xi \frac{8L^2 b_2^2}{\eta \kappa^2} \exp(-b_1 T) + \frac{1}{2} L_q \xi^2 b_2^2$. This gives that

$$\frac{1}{4} \xi \eta \sum_{k=0}^{K} ||\nabla q(v_k, \theta_k)||^2 \leq q(v_0, \theta_0) - q(v_K, \theta_K) + K\Delta + \sum_{k=0}^{K-1} \chi_k.$$

Using the same argument in the proof of Lemma 11,

$$||\hat{\nabla} q(v, \theta)|| \leq (2L\kappa^{-1} + 1)||\nabla q(v, \theta)||.$$

We hence have

$$\sum_{k=0}^{K-1} ||\hat{\nabla} q(v_k, \theta_k)||^2 \leq (2L\kappa^{-1} + 1)^2 \sum_{k=0}^{K-1} ||\nabla q(v_k, \theta_k)||^2$$

$$\leq \frac{4(2L\kappa^{-1} + 1)^2}{\xi \eta} (q(v_0, \theta_0) - q(v_K, \theta_K) + K\Delta + \sum_{k=0}^{K-1} \chi_k).$$

Similar to the proof of Lemma 12,

$$\sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||^2 \leq \frac{2(f(v_0, \theta_0) - f(v_K, \theta_K))}{\xi} + 2\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla} q(v_k, \theta_k)||^2 + 2\eta M \sqrt{K} \sqrt{\sum_{k=0}^{K-1} ||\hat{\nabla} q(v_k, \theta_k)||^2}.$$

Using $\sqrt{x+y} \leq \sqrt{x} + \sqrt{y}$, we have

$$2\eta^2 \sum_{k=0}^{K-1} ||\hat{\nabla} q(v_k, \theta_k)||^2 \leq \frac{8\eta(2L\kappa^{-1} + 1)^2}{\xi}(q(v_0, \theta_0) - q(v_K, \theta_K) + K\Delta + \sum_{k=0}^{K-1} \chi_k)$$

$$2\eta M \sqrt{K} \sqrt{\sum_{k=0}^{K-1} ||\hat{\nabla} q(v_k, \theta_k)||^2} \leq \sqrt{K} \frac{4\eta^{1/2} M(2L\kappa^{-1} + 1)}{\xi^{1/2}} (\sqrt{q(v_0, \theta_0) - q(v_K, \theta_K)} + K^{1/2}\Delta^{1/2} + \sqrt{\left[\sum_{k=0}^{K-1} \chi_k\right]_+}).$$

Also notice that by Assumption 4,

$$\sum_{k=0}^{K-1} q(v_k, \theta_k) \leq \sum_{k=0}^{K-1} \frac{\xi}{\kappa} ||\nabla q(v_k, \theta_k)||^2$$

$$\leq \frac{4}{\eta \kappa \xi}(q(v_0, \theta_0) - q(v_K, \theta_K) + K\Delta + \sum_{k=0}^{K-1} \chi_k)$$

We hence have

$$\sum_{k=0}^{K-1} (||\delta^*(v_k, \theta_k)||^2 + q(v_k, \theta_k)) = O\left(\frac{1}{\xi} + \frac{K\Delta}{\xi} + \frac{K^{1/2}}{\xi^{1/2}} + \frac{K\Delta^{1/2}}{\xi^{1/2}} + K^{1/2} \left(\left[\sum_{k=0}^{K-1} \chi_k\right]_+\right)^{1/2}\right)$$

$$= O\left(\frac{1}{\xi} + K\exp(-b_1 T/2) + K\xi^{1/2} + \frac{K^{1/2}}{\xi^{1/2}} + \left(K\left[\sum_{k=0}^{K-1} \chi_k\right]_+\right)^{1/2}\right).$$

Using the same argument as the proof of Theorem 1, when $T \geq \lceil -b_1^{-1} \log(\frac{1}{16}\kappa^2 L^{-2}) \rceil$,

$$\mathcal{K}^\diamond(v, \theta) \leq 2||\delta^*(v, \theta)||^2 + q(v, \theta) + 8L^2 \exp(-b_1 T)\kappa^{-2}(\eta + 2)^2 b_2^2.$$

This implies that

$$\min_k \mathcal{K}^\diamond(v_k, \theta_k) \leq \frac{1}{K} \sum_{k=0}^{K-1} [2||\delta^*(v, \theta)||^2 + q(v, \theta)] + 8L^2 \exp(-b_1 T)\kappa^{-2}(\eta + 2)^2 b_2^2$$

$$= O\left(\frac{1}{\xi K} + \exp(-b_1 T/2) + \xi^{1/2} + \frac{1}{\xi^{1/2} K^{1/2}} + \left(\left[\frac{1}{K}\sum_{k=0}^{K-1} \chi_k\right]_+\right)^{1/2}\right).$$

811  Now we proceed to bound $\frac{1}{K}\sum_{k=0}^{K-1}\chi_k$. Notice that

$$\chi_k = g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))$$
$$= g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_k, \theta^\diamond(v_k, \theta_k)) + g(v_k, \theta^\diamond(v_k, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1})).$$

812  Notice that using Assumption 2 and Lemma 14

$$g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)) - g(v_k, \theta^\diamond(v_k, \theta_k)) \leq L||[v_{k+1}, \theta^\diamond(v_{k+1}, \theta_k)] - [v_k, \theta^\diamond(v_k, \theta_k)]||$$
$$\leq L(||v_{k+1} - v_k|| + ||\theta^\diamond(v_{k+1}, \theta_k) - \theta^\diamond(v_k, \theta_k)||)$$
$$\leq (L + \frac{4L}{\kappa})||v_{k+1} - v_k||$$
$$\leq (L + \frac{4L}{\kappa})\xi||\delta^*(v_k, \theta_k)||.$$

813  Note that using the same procedure as the proof of Lemma 6, $||\delta^*(v_k, \theta_k)|| \leq b_2$. We thus conclude
814  that

$$\sum_{k=0}^{K-1}\chi_k \leq \sum_{k=0}^{K-1} g(v_k, \theta^\diamond(v_k, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1}))$$

$$+ (L + \frac{4L}{\kappa})\xi \sum_{k=0}^{K-1} ||\delta^*(v_k, \theta_k)||$$

$$\leq \sum_{k=0}^{K-1} g(v_k, \theta^\diamond(v_k, \theta_k)) - g(v_{k+1}, \theta^\diamond(v_{k+1}, \theta_{k+1})) + (L + \frac{4L}{\kappa})b_2\xi K$$

$$= g(v_0, \theta^\diamond(v_0, \theta_0)) - g(v_K, \theta^\diamond(v_K, \theta_K)) + (L + \frac{4L}{\kappa})b_2\xi K.$$

815  We thus have $\frac{1}{K}\sum_{k=0}^{K-1}\chi_k = O(\frac{1}{K} + \xi)$.

# E    List of absolute constants used in the proofs

817  Here we summarize the absolute constant used in the proofs.

$$b_1(\alpha, L, \kappa) = \log(1 - (\alpha - L\alpha^2/2)\kappa)$$
$$b_2(M, \eta) = (3 + \eta)M$$
$$b_3(\eta, \alpha, \kappa, L) = \left\lceil -b_1^{-1}\log(\frac{\eta\kappa}{64\eta L_q^2}) \right\rceil$$
$$b_4(\eta, \kappa, \xi) = -\log(1 - \frac{\xi}{4}\eta\kappa)$$
$$b_5(L_q, \eta, \kappa) = \frac{16L_q^2}{\eta\kappa^2}(2L\kappa^{-1} + 1)^2$$
$$b_6(\kappa, L) = (2L\kappa^{-1} + 1)^2$$
$$b_7(\eta, \alpha, \kappa, L) = \left\lceil -b_1^{-1}\log(\frac{\kappa^2}{48\eta L_q^2}) \right\rceil$$