A Reinforcement Learning Framework for Time Dependent Causal Effects Evaluation in A/B Testing

Abstract

A/B testing, or online experiment is a standard business strategy to compare a new product with an old one in pharmaceutical, technological, and traditional industries. The aim of this paper is to introduce a reinforcement learning framework for carrying A/B testing in two-sided marketplace platforms, while characterizing the long-term treatment effects. Our proposed testing procedure allows for sequential monitoring and online updating. It is generally applicable to a variety of treatment designs in different industries. In addition, we systematically investigate the theoretical properties (e.g., size and power) of our testing procedure. Finally, we apply our framework to both synthetic data and a real-world data example obtained from a technological company to illustrate its advantage over the current practice.

1 Introduction

A/B testing, or online experiment is a business strategy to compare a new product with an old one in pharmaceutical, technological, and traditional industries (e.g., Google, Amazon, or Facebook). Most works in the literature focus on the setting, in which observations are independent across time (see e.g. Johari et al., 2015; Kharitonov et al., 2015; Johari et al., 2017; Abhishek & Mannor, 2017; Yang et al., 2017; Ju et al., 2019; Zhou et al., 2020, and the references therein). In many applications, however, there is only one unit (or system) that receives a sequence of treatments over time. The treatment at a given time can impact future outcomes. For instance, in a ride-sharing company (e.g., Uber), an order dispatching strategy not only affects its immediate income, but also impacts the spatial distribution of drivers in the future, thus affecting its future income. In medicine, it usually takes time for drugs to distribute to the site of action. The independence assumption is thus violated.

The focus of this paper is to test the difference in long-term treatment effects between two products in online experiments. There are three major challenges as follows. (i) The first one lies in modelling the temporal dependence between treatments and outcomes. (ii) Running each experiment takes a considerable time. The company wishes to terminate the experiment as early as possible in order to save both time and budget. (iii) Treatments are desired to be allocated in a manner to maximize the cumulative outcomes and to detect the alternative more efficiently. The testing procedure shall allow the treatment to be adaptively assigned.

We summarize our contributions as follows. First, we introduce a reinforcement learning (RL, see e.g., Sutton & Barto, 2018, for an overview) framework for A/B testing. In addition to the treatment-outcome pairs, it is assumed that there is a set of time-varying state confounding variables. We model the state-treatment-outcome triplet by using the Markov decision process (MDP, see e.g. Puterman, 1994) to characterize the association between treatments and outcomes across time. Specifically, at each time point, the decision maker selects a treatment based on the observed state variables. The system responds by giving the decision maker a corresponding outcome and moving into a new state in the next time step. In this way, past treatments will have an indirect influence on future rewards through its effect on future state variables. See Figure 1 for an illustration. In addition, the long-term treatment effects can be characterized by the value functions (see Section 3.1 for details) that measure the discounted cumulative gain from a given initial state. Under this framework, it suffices to evaluate the difference between two value functions to compare different treatments. This addresses the challenge mentioned in (i).

Figure 1: Causal diagram for MDP under settings where treatments depend on current states only. (S_t, A_t, Y_t) represents the state-treatment-outcome triplet. Solid lines represent causal relationships.

Second, we propose a novel sequential testing procedure for detecting the difference between two value functions. To the best of our knowledge, this is the first work on developing valid sequential tests in the RL framework. Our proposed test integrates temporal difference learning (see e.g., Precup et al., 2001; Sutton et al., 2008), the α-spending approach (Lan & DeMets, 1983) and bootstrap (Efron & Tibshirani, 1994) to allows for sequential monitoring and online updating. It is generally applicable to a variety of treatment designs, including the Markov design, the alternating-time-interval design and the adaptive design (see Section 4.2 for details). This addresses the challenges mentioned in (ii) and (iii).

Third, we systematically investigate the asymptotic properties of our testing procedure. We show that our test not only maintains the nominal type I error rate, but also has non-negligible powers against local alternatives. To our knowledge, these results have not been established in RL.

Finally, we introduce a potential outcome framework for MDP. We state all necessary conditions that guarantee that the value functions are estimable from the observed data.

2 RELATED WORK

There is a huge literature on RL such that various algorithms are proposed for an agent to learn an optimal policy and interact with an environment. Our work is closely related to the literature on off-policy evaluation, whose objective is to estimate the value of a new policy based on data collected by a different policy. Popular methods include Thomas et al. (2015); Jiang & Li (2016); Thomas & Brunskill (2016); Liu et al. (2018); Farajtabar et al. (2018); Kallus & Uehara (2019). Those methods required the treatment assignment probability (propensity score) to be bounded away from 0 and 1. As such, they are inapplicable to the alternating-time-interval design, which is the treatment allocation strategy in our real data application.

Our work is related to the temporal-difference learning method based on function approximation. Convergence guarantees of the value function estimators have been derived by Sutton et al. (2008) under the setting of independent noise and by Bhandari et al. (2018) for Markovian noise. However, uncertainty quantification of the resulting value function estimators have been less studied. Such results are critical for carrying out A/B testing. Luckett et al. (2019) outlined a procedure for estimating the value under a given policy. Shi et al. (2020c) developed a confidence interval for the value. However, these method do not allow for sequential monitoring or online updating.

In addition to the literature on RL, our work is also related to a line of research on evaluating time-varying causal effects (see e.g. Robins, 1986; Boruvka et al., 2018; Ning et al., 2019; Rambachan & Shephard, 2019; Viviano & Bradic, 2019; Bojinov & Shephard, 2020). However, none of the above cited works used an RL framework to characterize treatment effects. In particular, Bojinov & Shephard (2020) proposed to use importance sampling (IS) based methods to test the null hypothesis of no (average) temporal causal effects in time series experiments. Their causal estimand is different from ours since they focused on p lag treatment effects, whereas we consider the long-term effects characterized by the value function. Moreover, their method requires the propensity score to be bounded away from 0 and 1, and thus it is not valid for our applications.

Furthermore, our work is also related to the literature on sequential analysis (see e.g. Jennison & Turnbull, 1999, and the references therein). Most test statistics in classical sequential analysis have the canonical joint distribution (see Equation (3.1) in Jennison & Turnbull, 1999) and their associated stopping boundary can be recursively updated via numerical integration. However, in our setup, test statistics no longer have the canonical joint distribution. To resolve this issue, we propose a scalable bootstrap-assisted procedure to determine the stopping boundary.

In addition, we note that there is a line of research on bandit/RL with causal graphs (see e.g., Lee & Bareinboim, 2018; 2019). We remark that the problems considered and the solutions developed in this article are different from these works. Specifically, these works considered applying causal inference methods to deal with unmeasured confounders in bandit/RL settings whereas we apply the RL framework to evaluate time-dependant causal effects.

Finally, we relax several key conditions used in Ertefaie (2014) and Luckett et al. (2019) that presented a potential outcome framework for MDP (see Section 3.1 for details). Specifically, Ertefaie (2014) and Luckett et al. (2019) imposed the Markov conditions on the observed data rather than the potential outcomes, while assuming that the outcome at time t is a deterministic function of the state variables at time t, $t+1$ and the treatment at time t.

3 PROBLEM FORMULATION

3.1 A POTENTIAL OUTCOME FRAMEWORK FOR MDP

For simplicity, we assume that there are only two treatments (actions, products), coded as 0 and 1, respectively. For any $t \geq 0$, let $\bar{a}_t = (a_0, a_1, \cdots, a_t)^\top \in \{0,1\}^{t+1}$ denote a treatment history vector up to time t. Let $\mathbb{S}$ denote the support of state variables and S_0 denote the initial state variable. We assume $\mathbb{S}$ is a compact subset of $\mathbb{R}^d$. For any $(\bar{a}_{t-1}, \bar{a}_t)$, let $S_t^*(\bar{a}_{t-1})$ and $Y_t^*(\bar{a}_t)$ be the counterfactual state and counterfactual outcome, respectively, that would occur at time t had the agent followed the treatment history $\bar{a}_t$. The set of potential outcomes up to time t is given by

$$W_t^*(\bar{a}_t) = \{S_0, Y_0^*(a_0), S_1^*(a_0), \cdots, S_t^*(\bar{a}_{t-1}), Y_t^*(\bar{a}_t)\}.$$

Let $W^* = \cup_{t \geq 0, \bar{a}_t \in \{0,1\}^{t+1}} W_t^*(\bar{a}_t)$ be the set of all potential outcomes.

A deterministic policy π is a function that maps the space of state variables to the set of available actions. For any such π, let $\bar{\pi}_t$ denote the treatment history up to time t, assigned according to π. We use $S_t^*(\bar{\pi}_{t-1})$ and $Y_t^*(\bar{\pi}_t)$ to denote the associated potential state and outcome that would occur at time t had the agent followed π. The goodness of a policy π is measured by its value function,

$$V(\pi; s) = \sum_{t \geq 0} \gamma^t \mathrm{E}\{Y_t^*(\bar{\pi}_t)|S_0 = s\},$$

where $0 < \gamma < 1$ is a discounted factor that reflects the trade-off between immediate and future outcomes. Note that our definition of the value function is slightly different from those in the existing literature (see Sutton & Barto, 2018, for example). Specifically, $V(\pi; s)$ is defined through potential outcomes rather than the observed data. Similarly, we define the Q function by

$$Q(\pi; a, s) = \sum_{t \geq 0} \gamma^t \mathrm{E}\{Y_t^*(\bar{\pi}_t(a))|S_0 = s\},$$

where $\{\bar{\pi}_t(a)\}_{t \geq 0}$ denotes the treatment history where the initial action equals to a and all other actions are assigned according to π.

In our setup, we focus on two nondynamic policies that assign the same treatment at each time point. We use their value functions (denote by $V(1; \cdot)$ and $V(0; \cdot)$) to measure their long-term treatment effects. Meanwhile, our proposed method is equally applicable to the dynamic policy scenario as well. See Section B for details. To quantitatively compare the two policies, we introduce the Average Treatment Effect (ATE) based on their value functions which relates RL to causal inference.

Definition. For a given reference distribution function $\mathbb{G}$, ATE is defined by the integrated difference between two value function, i.e., $\text{ATE} = \int_s \{V(1; s) - V(0; s)\}\mathbb{G}(ds)$.

The focus of this paper is to test the following hypotheses:

$$H_0 : \tau_0 = \text{ATE} \leq 0 \quad \text{v.s} \quad H_1 : \tau_0 = \text{ATE} > 0.$$

When H_0 holds, the new product is no better than the old one.

3.2 IDENTIFIABILITY OF ATE

One of the most important question in causal inference is the identifiability of causal effects. In this section, we present sufficient conditions that guarantee the identifiability of the value function.

In practice, with the exception of S_0, the set W^* cannot be observed, whereas at time t, we observe the state-action-outcome triplet (S_t, A_t, Y_t). For any $t \geq 0$, let $\bar{A}_t = (A_0, A_1, \cdots, A_t)^\top$ denote

the observed treatment history. We first introduce two conditions that are commonly assumed in multi-stage decision making problems (see e.g. Murphy, 2003; Robins, 2004; Zhang et al., 2013; Kennedy, 2019).

(CA) Consistency assumption: $S_{t+1} = S_{t+1}^*(\bar{A}_t)$ and $Y_t = Y_t^*(\bar{A}_t)$ for all $t \geq 0$, almost surely.

(SRA) Sequential randomization: A_t is independent of W^* given S_t and $\{S_j, A_j, Y_j\}_{0 \leq j < t}$.

The SRA implies that there are no unmeasured confounders and it automatically holds in online experiments, in which the treatment assignment mechanism is pre-specified. In SRA, we allows A_t to depend on the observed data history $S_t, \{S_j, A_j, Y_j\}_{0 \leq j < t}$ and thus, the treatments can be adaptively chosen. We next introduce two conditions that are unique to the RL setting.

(MA) Markov assumption: there exists a Markov transition kernel $\mathcal{P}$ such that for any $t \geq 0$, $\bar{a}_t \in \{0,1\}^{t+1}$ and $\mathcal{S} \subseteq \mathbb{R}^d$, we have $\Pr\{S_{t+1}^*(\bar{a}_t) \in \mathcal{S}|W_t^*(\bar{a}_t)\} = \mathcal{P}(\mathcal{S}; a_t, S_t^*(\bar{a}_{t-1}))$.

(CMIA) Conditional mean independence assumption: there exists a function r such that for any $t \geq 0, \bar{a}_t \in \{0,1\}^{t+1}$, we have $\mathrm{E}\{Y_t^*(\bar{a}_t)|S_t^*(\bar{a}_{t-1}), W_{t-1}^*(\bar{a}_{t-1})\} = r(a_t, S_t^*(\bar{a}_{t-1}))$.

These two conditions are central to the empirical validity of reinforcement learning (RL). Specifically, under these two conditions, there exists an optimal policy π^* such that $V(\pi^*; s) \geq V(\pi; s)$ for any π and s. We observe that Ertefaie (2014) and Luckett et al. (2019) imposed the Markov conditions on the observed data rather than the potential outcomes. When CA and SRA hold, these assumptions are equivalent. When SRA is violated, their Markov assumptions could be violated as the treatment depends on unobserved confounders and the observed data process is no longer Markovian. CMIA requires past treatments to affect $Y_t^*(\bar{a}_t)$ only through its impact on $S_t^*(\bar{a}_{t-1})$. In other words, the state variables shall be chosen to include those that serve as important mediators between past treatments and current outcomes. Under MA, CMIA is automatically satisfied when $Y_t^*(\bar{a}_t)$ is a deterministic function of $(S_{t+1}^*(\bar{a}_t), a_t, S_t^*(\bar{a}_{t-1}))$ that measures the system's status at time $t+1$. The latter condition is commonly imposed in the reinforcement learning literature. As we have discussed, when CA and SRA hold, MA and CMIA hold on the observed data as well. The latter two conditions are thus testable. See the goodness-of-fit test developed by Shi et al. (2020a).

To conclude this section, we derive a version of Bellman equation for the Q function under the potential outcome framework. Specifically, for $a' \in \{0,1\}$, let $Q(a'; \cdot, \cdot)$ denote the Q function where treatment a' is repeatedly assigned after the initial decision.

Lemma 1 *Under MA, CMIA, CA and SRA, for any $t \geq 0$, $a' \in \{0,1\}$ and any function φ : $\mathbb{S} \times \{0,1\} \to \mathbb{R}$, we have $E[\{Q(a'; A_t, S_t) - Y_t - \gamma Q(a'; a', S_{t+1})\}\varphi(S_t, A_t)] = 0$.*

Lemma 1 implies that the Q-function is estimable from the observed data. As a result, $\tau_0 =$ATE is identifiable. Note that the positivity assumption is not needed in Lemma 1. Our procedure can thus handle the case where treatments are deterministically assigned. This is due to MA and CMIA that assume the system dynamics are invariant across time. To elaborate this, note that the discounted value function is completely determined by the transition kernel $\mathcal{P}$ and the reward function r. We remark that these quantities can be consistently estimated under certain conditions, regardless of whether the treatments are deterministically assigned or not.

4 TESTING PROCEDURE

4.1 AN OVERVIEW OF THE PROPOSAL

First, we estimate τ_0 based on a version of temporal difference learning. The idea is to apply basis function approximations to solve an estimating equation derived from Lemma 1. Specifically, let $\mathcal{Q} = \{\Psi^\top(s)\beta_{a',a} : \beta_{a',a} \in \mathbb{R}^q\}$ be a large linear approximation space for $Q(a'; a, s)$, where $\Psi(\cdot)$ is a vector containing q basis functions on $\mathbb{S}$. The dimension q is allowed to depend on the number of samples T to alleviate the effects of model misspecification. Let us suppose $Q \in \mathcal{Q}$ for a moment. By Lemma 1, there exists some $\boldsymbol{\beta}^* = (\beta_{0,0}^{*\top}, \beta_{0,1}^{*\top}, \beta_{1,0}^{*\top}, \beta_{1,1}^{*\top})^\top$ such that

$$\mathrm{E}[\{\Psi^\top(S_t)\beta_{a',a}^* - Y_t - \gamma\Psi^\top(S_{t+1})\beta_{a',a'}^*\}\Psi(S_t)\mathbb{I}(A_t = a)] = 0, \quad \forall a, a' \in \{0,1\},$$

where $\mathbb{I}(\cdot)$ denotes the indicator function. Let $\xi(s, a) = \{\Psi^\top(s)\mathbb{I}(a = 0), \Psi^\top(s)\mathbb{I}(a = 1)\}^\top$. The above equations can be rewritten as $\mathrm{E}(\boldsymbol{\Sigma}_t\boldsymbol{\beta}^*) = \mathrm{E}\boldsymbol{\eta}_t$, where $\boldsymbol{\Sigma}_t$ is a block diagonal matrix given by

$$\boldsymbol{\Sigma}_t = \left[\begin{array}{l} \xi(S_t, A_t)\{\xi(S_t, A_t) - \gamma\xi(S_{t+1}, 0)\}^\top \\ \qquad\qquad\qquad\qquad \xi(S_t, A_t)\{\xi(S_t, A_t) - \gamma\xi(S_{t+1}, 1)\}^\top \end{array} \right]$$

$$\text{and} \quad \boldsymbol{\eta}_t = \{\xi(S_t, A_t)^\top Y_t, \xi(S_t, A_t)^\top Y_t\}^\top.$$

Let $\widehat{\boldsymbol{\Sigma}}(t) = t^{-1} \sum_{j < t} \boldsymbol{\Sigma}_j$ and $\widehat{\boldsymbol{\eta}}(t) = t^{-1} \sum_{j < t} \boldsymbol{\eta}_j$. It follows that $\mathrm{E}\{\widehat{\boldsymbol{\Sigma}}(t)\beta^*\} = \mathrm{E}\{\widehat{\boldsymbol{\eta}}(t)\}$. This motivates us to estimate $\boldsymbol{\beta}^*$ by $\widehat{\boldsymbol{\beta}}(t) = \{\widehat{\beta}_{0,0}^\top(t), \widehat{\beta}_{0,1}^\top(t), \widehat{\beta}_{1,0}^\top(t), \widehat{\beta}_{1,1}^\top(t)\}^\top = \widehat{\boldsymbol{\Sigma}}^{-1}(t)\widehat{\boldsymbol{\eta}}(t)$. ATE can thus be estimated by the plug-in estimator $\widehat{\tau}(t) = \int_s \Psi^\top(s)\{\widehat{\beta}_{1,1}(t) - \widehat{\beta}_{0,0}(t)\}\mathbb{G}(ds)$.

Second, we use $\widehat{\tau}(t)$ to construct our test statistic at time t. We will show $\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}$ is asymptotically normal. Its variance can be consistently estimated by $\widehat{\sigma}^2(t) = \boldsymbol{U}^\top \widehat{\boldsymbol{\Sigma}}^{-1}(t)\widehat{\boldsymbol{\Omega}}(t)\{\widehat{\boldsymbol{\Sigma}}^{-1}(t)\}^\top \boldsymbol{U}$, as t grows to infinity, where the explicit forms of $\boldsymbol{U}$ and $\widehat{\boldsymbol{\Omega}}(t)$ are given in equation 3 and equation 4. This yields our test statistic $\sqrt{t}\widehat{\tau}(t)/\widehat{\sigma}(t)$, at time t. For a given significance level $\alpha > 0$, we reject H_0 when $\sqrt{t}\widehat{\tau}(t)/\widehat{\sigma}(t)$ exceeds the upper α-th quantile of a standard normal distribution.

Third, we integrate the α-spending approach with bootstrap to sequentially implement our test (see Section 4.3). The idea is to generate bootstrap samples that mimic the distribution of our test statistics, to specify the stopping boundary at each interim stage. Suppose that the interim analyses are conducted at time points $T_1 < \cdots < T_K = T$. To simplify the presentation, for each $1 \leq k < K$, we assume $T_k/T \to c_k$ for some constants $0 < c_1 < \cdots < c_{K-1} < 1$. To better understand our algorithm, we next investigate the limiting distribution of our test statistics at these interim stages.

4.2 ASYMPTOTIC PROPERTIES UNDER DIFFERENT TREATMENT DESIGNS

We consider three treatment allocation designs that can be handled by our procedure as follows:

D1. Markov design: $\Pr(A_t = 1 | S_t, \{S_j, A_j, Y_j\}_{0 \leq j < t}) = b^{(0)}(S_t)$ for some function $b^{(0)}(\cdot)$ uniformly bounded away from 0 and 1.

D2. Alternating-time-interval design: $A_{2j} = 0$, $A_{2j+1} = 1$ for all $j \geq 0$.

D3. Adaptive design (e.g., ϵ-greedy): For $T_k \leq t < T_{k+1}$ for some $k \geq 0$, $\Pr(A_t = 1 | S_t, \{S_j, A_j, Y_j\}_{0 \leq j < t}) = b^{(k)}(S_t)$ for some $b^{(k)}(\cdot)$ that depends on $\{S_j, A_j, Y_j\}_{0 \leq j < T_k}$.

More discussions on these designs are given in Appendix E to save space. To study the asymptotic properties of our test, we need to introduce assumptions C1-C3 and move them and their corresponding detailed discussions to Appendix F. In C1, we require the above mentioned Markov chains to be geometrically ergodic. Geometric ergodicity is weaker than the uniform ergodicity condition imposed in the existing reinforcement learning literature (see e.g. Bhandari et al., 2018; Zou et al., 2019; Mengersen & Tweedie, 1996). In C2, we impose conditions on the set of basis functions $\Psi(\cdot)$ such that $\Psi(\cdot)$ is chosen to yield a good approximation for the Q function. We present examples that satisfy these conditions in Appendix F.2. In C3, we impose some mild conditions on the temporal-difference error, requiring their variance to be non-degenerate.

Let $\{Z_1, \cdots, Z_K\}$ denote the sequence of our test statistics, where $Z_k = \sqrt{T_k}\widehat{\tau}(T_k)/\widehat{\sigma}(T_k)$. In the following, we study their joint asymptotic distributions. We also present an estimator of their covariance matrix that is consistent under all designs.

Theorem 1 (Limiting distributions) *Assume C1-C3, MA, CMIA, CA, and SRA hold. Assume all immediate rewards are uniformly bounded variables, the density function of S_0 is uniformly bounded on $\mathbb{S}$ and q satisfies $q = o(\sqrt{T}/\log T)$. Then under either D1, D2 or D3, we have*

- *$\{Z_k\}_{1 \leq k \leq K}$ are jointly asymptotically normal;*
- *their asymptotic means are non-positive under H_0;*
- *their covariance matrix can be consistently estimated by some $\widehat{\boldsymbol{\Xi}}$, whose (k_1, k_2)-th element $\widehat{\Xi}_{k_1,k_2}$ equals $\sqrt{T_{k_1}/T_{k_2}}\boldsymbol{U}^\top \widehat{\boldsymbol{\Sigma}}^{-1}(T_{k_1})\widehat{\boldsymbol{\Omega}}(T_{k_1})\{\widehat{\boldsymbol{\Sigma}}^{-1}(T_{k_2})\}^\top \boldsymbol{U}/\{\widehat{\sigma}(T_{k_1})\widehat{\sigma}(T_{k_2})\}$.*

4.3 SEQUENTIAL MONITORING AND ONLINE UPDATING

To sequentially monitor our test, we need to specify the stopping boundary $\{b_k\}_{1 \leq k \leq K}$ such that the experiment is terminated and H_0 is rejected when $Z_k > b_k$ for some k.

First, we use the α spending function approach to guarantee the validity of our test. It requires to specify a monotonically increasing function $\alpha(\cdot)$ that satisfies $\alpha(0) = 0$ and $\alpha(T) = \alpha$. Some popular choices of the α spending function include

$$\alpha_1(t) = 2 - 2\Phi\{\Phi^{-1}(1 - \alpha/2)\sqrt{T/t}\} \quad \text{and} \quad \alpha_2(t) = \alpha(t/T)^\theta \quad \text{for} \ \theta > 0, \tag{1}$$

where $\Phi(\cdot)$ denotes the normal cumulative distribution function. Adopting the α spending approach, we require b_k's to satisfy

$$\Pr(\cup_{j=1}^k \{Z_j > b_j\}) = \alpha(T_k) + o(1), \quad \forall 1 \le k \le K. \tag{2}$$

When the sequence $\{Z_k\}_{1 \le k \le K}$ satisfies the Markov property, the stopping boundary can be efficiently computed based on the numerical integration method detailed in Section 19.2 of Jennison & Turnbull (1999). However, in our setup, the Markov assumption might not hold when adaptive design is used. As a result, the numerical integration method is not applicable.

The proposed method is built upon the wild bootstrap (Wu et al., 1986). The idea is to generate bootstrap samples that have asymptotically the same joint distribution as the test statistics. However, we note that directly applying the wild bootstrap algorithms is time consuming. See Section D for detailed illustration. To facilitate the computation, we present a scalable bootstrap algorithm to determine $\{b_k\}_k$, based on Theorem 1. Let $\{e_k\}_{1 \le k \le K}$ be a sequence of i.i.d $N(0, I_{4q})$ random vectors, where I_J stands for a $J \times J$ identity matrix for any J. Let $\widehat{\Omega}(T_0)$ be an $4q \times 4q$ zero matrix. At the k-th stage, we compute

$$\widehat{Z}_k^* = \frac{\boldsymbol{U}^\top \widehat{\boldsymbol{\Sigma}}^{-1}(T_k)}{\sqrt{T_k}\widehat{\sigma}(T_k)} \sum_{j=1}^k \{T_j \widehat{\boldsymbol{\Omega}}(T_j) - T_{j-1}\widehat{\boldsymbol{\Omega}}(T_{j-1})\}^{1/2} e_j.$$

A key observation is that, conditional on the observed dataset, the covariance of $\widehat{Z}_{k_1}^*$ and $\widehat{Z}_{k_2}^*$ equals $\widehat{\Xi}_{k_1,k_2}$. By Theorem 1, the covariance matrices of $\{Z_k\}_k$ and $\{Z_k^*\}_k$ are asymptotically equivalent. In addition, the limiting distributions of $\{Z_k\}_k$ and $\{Z_k^*\}_k$ are multivariate normal with zero means. As such, the joint distribution of $\{Z_k\}_{1 \le k \le K}$ can be well approximated by that of $\{Z_k^*\}_{1 \le k \le K}$ conditional on the data. This forms the basis of our bootstrap algorithm. By the requirement on $\{b_k\}_k$ in equation 2, we obtain $\Pr(\max_{1 \le j < k}(Z_j - b_j) \le 0, Z_k > b_k) = \alpha(T_k) - \alpha(T_{k-1}) + o(1)$. To implement our test, we thus recursively calculate the threshold $\widehat{b}_k$ as follows,

$$\Pr^* \left\{ \max_{1 \le j < k}(Z_j^* - \widehat{b}_j) \le 0, Z_k^* > \widehat{b}_k \right\} = \alpha(T_k) - \alpha(T_{k-1}),$$

where $\Pr^*$ denotes the conditional probability given on the data, and reject H_0 when $Z_k^* > \widehat{b}_k$ for some k. In practice, the left-hand-side can be approximated via Monte carlo simulations.

Theorem 2 (Type-I error) *Suppose that the conditions of Theorem 1 hold and $\alpha(\cdot)$ is continuous. Then the proposed thresholds satisfy $Pr(\bigcup_{j=1}^k \{Z_j > \widehat{b}_j\}) \le \alpha(T_k) + o(1)$, for all $1 \le k \le K$ under H_0. The equality holds when $\tau_0 = 0$.*

Theorem 2 implies that the type-I error rate of the proposed test is well controlled. We next investigate the power property of our test.

Theorem 3 (Power) *Under the conditions of Theorem 2, assume $\tau_0 \gg T^{-1/2}$, then $Pr(\bigcup_{j=1}^k \{Z_j > \widehat{b}_j\}) \to 1$. Assume $\tau_0 = T^{-1/2}h$ for some $h > 0$. Then $\lim_{T \to \infty}[Pr(\cup_{j=1}^k \{Z_j > \widehat{b}_j\}) - \alpha(T_k)] > 0$.*

Combining Theorems 2 and 3 yields the consistency of our test. The second assertion in Theorem 3 implies that our test has non-negligible powers against local alternatives converging to H_0 at the $T^{-1/2}$ rate. Finally, it is worth mentioning that our test can be online updated as batches of observations arrive at the end of each interim stage. We summarize our procedure in Algorithm 1 (see Appendix A). Its time complexity is dominated by $O(Bq^3 + Tq^2)$.

5 NUMERICAL STUDIES

5.1 SYNTHETIC DATA

Simulated data of states and rewards was generated as follows,

$$S_{1,t} = (2A_{t-1} - 1)S_{1,(t-1)}/2 + S_{2,(t-1)}/4 + \delta A_{t-1} + \varepsilon_{1,t},$$
$$S_{2,t} = (2A_{t-1} - 1)S_{2,(t-1)}/2 + S_{1,(t-1)}/4 + \delta A_{t-1} + \varepsilon_{2,t}, \quad Y_t = 1 + (S_{1,t} + S_{2,t})/2 + \varepsilon_{3,t},$$

(a) The proposed test under H_1 and H_0 (from left plots to right plots)

(b) Two-sample t-test under H_1 and H_0 (from left plots to right plots)

Figure 2: Empirical rejection probabilities of our test and the two-sample t-test with $\alpha(\cdot) = \alpha_1(\cdot)$. Settings correspond to the alternating-time-interval, adaptive and Markov design, from top plots to bottom plots.

where the random errors $\{\varepsilon_{j,t}\}_{j=1,2,0 \leq t \leq T}$ are i.i.d $N(0, 0.5^2)$ and $\{\varepsilon_{3,t}\}_{0 \leq t \leq T}$ are i.i.d $N(0, 0.3^2)$. Let $S_t = (S_{1,t}, S_{2,t})^\top$ denote the state at time t. Under this model, treatments have delayed effects on the outcomes. Specifically, Y_t does not depend on A_t, but is affected by A_{t-1} through S_t. The parameter δ characterizes the degree of such carryover effects. When $\delta = 0$, $\tau_0 = 0$ and H_0 holds. When $\delta > 0$, H_1 holds. Moreover, τ_0 increases as δ increases.

We set $K = 5$ and $(T_1, T_2, T_3, T_4, T_5) = (300, 375, 450, 525, 600)$. The discounted factor γ is set to 0.6 and $\mathbb{G}$ is chosen as the initial state distribution. We consider three behavior policies, according to the designs D1-D3, respectively. For the behavior policy in D1, we set $b^{(0)}(s) = 0.5$ for any $s \in \mathbb{S}$. For the behavior policy in D3, we use an ϵ-greedy policy and set $b^{(k)}(s) = \epsilon/2 + (1 - \epsilon)\mathbb{I}(\Psi(s)^\top(\widehat{\beta}_{1,1}(T_k) - \widehat{\beta}_{0,0}(T_k)) > 0)$, with $\epsilon = 0.1$, for any $k \geq 1$ and $s \in \mathbb{S}$.

For each design, we further consider five choices of δ, corresponding to $0, 0.05, 0.1, 0.15$ and 0.2. The significance level α is set to 0.05 in all cases. To implement our test, we choose two α-spending functions, corresponding to $\alpha_1(\cdot)$ and $\alpha_2(\cdot)$ given in equation 1. The hyperparameter θ in $\alpha_2(\cdot)$ is set to 3. The number of bootstrap sample is set to 1000. In addition, we consider the following polynomial basis function, $\Psi(s) = \Psi(s_1, s_2) = (1, s_1, s_1^2, \cdots, s_1^J, s_2, s_2^2, \cdots, s_2^J)^\top$, with $J = 4$. We also tried some other values of J by setting J to 3 and 5. Results are reported in Figure 5 (see Appendix H). It can be seen that the resulting tests have very similar performance and is not sensitive to the choice of J.

All experiments run on a macbook pro with a dual-core 2.7 GHz processor. Implementing a single test takes one second. Figures 2 (a) and 4 (a) (see Appendix H) depict the empirical rejection probabilities of our test statistics at different interim stages under H_0 and H_1 with different combinations of δ, $\alpha(\cdot)$ and the designs. These rejection probabilities are aggregated over 500 simulations. We also plot $\alpha_1(\cdot)$ and $\alpha_2(\cdot)$ under H_0. Based on the results, it can be seen that under H_0, the Type-I error rate of our test is well-controlled and close to the nominal level at each interim stage. Under H_1, the power of our test increases as δ increases, showing the consistency of our test procedure.

5.2 BASELINE COMPARISON

To further evaluate our method, we compare it with the classical two-sample t-test that is commonly used in online experiments. We remark that the current practice of policy evaluation in most two-sided marketplace platforms is to employ classical two-sample t-test. Specifically, for each T_k, we apply the t-test to the data $\{A_t, Y_t\}_{0 \leq t \leq T_k}$ and plot the corresponding empirical rejection probabili-

Figure 3: Our test statistic (the orange line) and the rejection boundary (the black line) in the A/A (left plot) and A/B (right plot) experiments.

ties in Figures 2(b) and 4(b) (see Appendix H). As shown in these figures, the test fails to detect any carryover effects and does not have power at all.

We next explain why several other methods mentioned in the introduction cannot be used for comparison. First, a lot of causal effects evaluation methods did not consider early termination. Consequently, they are unsuitable to apply in our numerical studies. Second, standard temporal difference learning method did not study uncertainty quantification and asymptotic distribution of the resulting value function estimators. These results are critical for carrying out A/B testing. Finally, many methods proposed to use inverse propensity-score based methods. These methods require the treatment assignment probability to be strictly bounded away from 0 and 1, and thus it is not valid for the alternating-time-interval design.

5.3 REAL DATA APPLICATION

We apply the proposed test to a real dataset from a ride-sharing platform. Order dispatching is one of the most critical problems in online ride-hailing platforms to adapt the operation and management strategy to the dynamics in demand and supply. The purpose of this study is to compare the performance of a newly developed strategy with a standard control strategy used in the platform. The new strategy is expected to reduce the answer time of passengers and increase drivers income. For a given order, the new strategy will dispatch it to a nearby driver that has not yet finished their previous ride request, but almost. In comparison, the standard control assigns orders to drivers that have completed their ride requests.

The experiment is conducted at a given city from December 3rd to December 16th. Dispatch strategies are executed based on alternating half-hourly time intervals. We also apply our test to a data from an A/A experiment (which compares the baseline strategy against itself), conducted from November 12th to November 25th. We expect that our test will not reject H_0 when applied to the data from the A/A experiment, since the two strategies used are essentially the same.

Both experiments last for two weeks. Thirty-minutes is defined as one time unit. We set $T_k = 48 \times (k + 6)$ for $k = 1, \ldots, 8$. That is, the first interim analysis is performed at the end of the first week, followed by seven more at the end of each day during the second week. We choose the overall drivers' income in each time unit as the response. The new strategy is expected to reduce the answer time of passengers and increase drivers' income. Three time-varying variables are used to construct the state. The first two correspond to the number of requests (demand) and drivers' online time (supply) during each 30-minutes time interval. These factors are known to have large impact on drivers' income. The last one is the supply and demand equilibrium metric. This variable characterizes the degree that supply meets the demand and serves as an important mediator between past treatments and future outcomes.

To implement our test, we set $\gamma = 0.6$, $B = 1000$ and use a fourth-degree polynomial basis for $\Psi(\cdot)$, as in simulations. We use $\alpha_1(\cdot)$ as the spending function for interim analysis and set $\alpha = 0.05$. The test statistic and its corresponding rejection boundary at each interim stage are plotted in Figure 3. It can be seen that our test is able to conclude, at the end of the 12th day, that the new order dispatch strategy can significantly increase drivers' income, and meet more order requests. In addition, based on the dataset from the A/B experiment, we found that the new strategy reduces the answer time of orders by 2%, leading to almost 2% increment of drivers income. When applied to the data from the A/A experiment, we fail to reject H_0, as expected. For comparison, we also apply the two-sample t-test to the data collected from the A/B experiment. The corresponding p-value is 0.18. Similar to our findings in simulations, the t-test fails to detect the difference between the long-term effects of these two strategies.

REFERENCES

Vineet Abhishek and Shie Mannor. A nonparametric sequential test for online randomized experiments. In *Proceedings of the 26th International Conference on World Wide Web Companion*, pp. 610–616. International World Wide Web Conferences Steering Committee, 2017.

Jalaj Bhandari, Daniel Russo, and Raghav Singal. A finite time analysis of temporal difference learning with linear function approximation. *arXiv preprint arXiv:1806.02450*, 2018.

Iavor Bojinov and Neil Shephard. *Time series experiments and causal estimands: exact randomization tests and trading*, volume accepted. Taylor & Francis, 2020.

Audrey Boruvka, Daniel Almirall, Katie Witkiewitz, and Susan A. Murphy. Assessing time-varying causal effect moderation in mobile health. *J. Amer. Statist. Assoc.*, 113(523):1112–1121, 2018. ISSN 0162-1459. doi: 10.1080/01621459.2017.1305274.

Prabir Burman and Keh-Wei Chen. Nonparametric estimation of a regression function. *Ann. Statist.*, 17(4):1567–1596, 1989. ISSN 0090-5364. doi: 10.1214/aos/1176347382.

Xiaohong Chen and Timothy M. Christensen. Optimal uniform convergence rates and asymptotic normality for series estimators under weak dependence and weak conditions. *J. Econometrics*, 188(2):447–465, 2015. ISSN 0304-4076. doi: 10.1016/j.jeconom.2015.03.010.

Bradley Efron and Robert J Tibshirani. *An introduction to the bootstrap*. CRC press, 1994.

Ashkan Ertefaie. Constructing dynamic treatment regimes in infinite-horizon settings. *arXiv preprint arXiv:1406.0764*, 2014.

Mehrdad Farajtabar, Yinlam Chow, and Mohammad Ghavamzadeh. More robust doubly robust off-policy evaluation. In *International Conference on Machine Learning*, 2018.

Jianhua Z. Huang. Projection estimation in multiple regression with application to functional ANOVA models. *Ann. Statist.*, 26(1):242–272, 1998. ISSN 0090-5364. doi: 10.1214/aos/1030563984.

Christopher Jennison and Bruce W Turnbull. *Group sequential methods with applications to clinical trials*. Chapman and Hall/CRC, 1999.

Nan Jiang and Lihong Li. Doubly robust off-policy value evaluation for reinforcement learning. In *International Conference on Machine Learning*, pp. 652–661, 2016.

Ramesh Johari, Leo Pekelis, and David J Walsh. Always valid inference: Bringing sequential analysis to a/b testing. *arXiv preprint arXiv:1512.04922*, 2015.

Ramesh Johari, Pete Koomen, Leonid Pekelis, and David Walsh. Peeking at a/b tests: Why it matters, and what to do about it. In *Proceedings of the 23rd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, pp. 1517–1525. ACM, 2017.

Nianqiao Ju, Diane Hu, Adam Henderson, and Liangjie Hong. A sequential test for selecting the better variant: Online a/b testing, adaptive allocation, and continuous monitoring. In *Proceedings of the Twelfth ACM International Conference on Web Search and Data Mining*, pp. 492–500. ACM, 2019.

Nathan Kallus and Masatoshi Uehara. Efficiently breaking the curse of horizon: Double reinforcement learning in infinite-horizon processes. *arXiv preprint arXiv:1909.05850*, 2019.

Edward H Kennedy. Nonparametric causal effects based on incremental propensity score interventions. *Journal of the American Statistical Association*, 114(526):645–656, 2019.

Eugene Kharitonov, Aleksandr Vorobev, Craig Macdonald, Pavel Serdyukov, and Iadh Ounis. Sequential testing for early stopping of online experiments. In *Proceedings of the 38th International ACM SIGIR Conference on Research and Development in Information Retrieval*, pp. 473–482. ACM, 2015.

K. K. Gordon Lan and David L. DeMets. Discrete sequential boundaries for clinical trials. *Biometrika*, 70(3):659–663, 1983. ISSN 0006-3444. doi: 10.2307/2336502.

Sanghack Lee and Elias Bareinboim. Structural causal bandits: where to intervene? In *Advances in Neural Information Processing Systems*, pp. 2568–2578, 2018.

Sanghack Lee and Elias Bareinboim. Structural causal bandits with non-manipulable variables. In *Proceedings of the AAAI Conference on Artificial Intelligence*, volume 33, pp. 4164–4172, 2019.

Qiang Liu, Lihong Li, Ziyang Tang, and Dengyong Zhou. Breaking the curse of horizon: Infinite-horizon off-policy estimation. In *Advances in Neural Information Processing Systems*, pp. 5356–5366, 2018.

Daniel J Luckett, Eric B Laber, Anna R Kahkoska, David M Maahs, Elizabeth Mayer-Davis, and Michael R Kosorok. Estimating dynamic treatment regimes in mobile health using v-learning. *Journal of the American Statistical Association*, accepted, 2019.

D. L. McLeish. Dependent central limit theorems and invariance principles. *Ann. Probability*, 2: 620–628, 1974. ISSN 0091-1798. doi: 10.1214/aop/1176996608.

K. L. Mengersen and R. L. Tweedie. Rates of convergence of the Hastings and Metropolis algorithms. *Ann. Statist.*, 24(1):101–121, 1996. ISSN 0090-5364. doi: 10.1214/aos/1033066201.

S. A. Murphy. Optimal dynamic treatment regimes. *J. R. Stat. Soc. Ser. B Stat. Methodol.*, 65(2): 331–366, 2003. ISSN 1369-7412. doi: 10.1111/1467-9868.00389.

Bo Ning, Subhashis Ghosal, and Jewell Thomas. Bayesian method for causal inference in spatially-correlated multivariate time series. *Bayesian Anal.*, 14(1):1–28, 2019. ISSN 1936-0975. doi: 10.1214/18-BA1102.

Doina Precup, Richard S Sutton, and Sanjoy Dasgupta. Off-policy temporal-difference learning with function approximation. In *ICML*, pp. 417–424, 2001.

Martin L. Puterman. *Markov decision processes: discrete stochastic dynamic programming*. Wiley Series in Probability and Mathematical Statistics: Applied Probability and Statistics. John Wiley & Sons, Inc., New York, 1994. ISBN 0-471-61977-9. A Wiley-Interscience Publication.

B. Rabta and D. Aïssani. Perturbation bounds for Markov chains with general state space. *J. Math. Sci. (N.Y.)*, 228(5):510–521, 2018. ISSN 1072-3374. doi: 10.1007/s10958-017-3640-9.

Ashesh Rambachan and Neil Shephard. A nonparametric dynamic causal model for macroeconometrics. *Available at SSRN 3345325*, 2019.

James Robins. A new approach to causal inference in mortality studies with a sustained exposure period—application to control of the healthy worker survivor effect. volume 7, pp. 1393–1512. 1986. doi: 10.1016/0270-0255(86)90088-6. Mathematical models in medicine: diseases and epidemics, Part 2.

James M Robins. Optimal structural nested models for optimal sequential decisions. In *Proceedings of the second seattle Symposium in Biostatistics*, pp. 189–326. Springer, 2004.

Chengchun Shi, Runzhe Wan, Rui Song, Wenbin Lu, and Ling Leng. Does the markov decision process fit the data: Testing for the markov property in sequential decision making. *arXiv preprint arXiv:2002.01751*, 2020a.

Chengchun Shi, Sheng Zhang, Wenbin Lu, and Rui Rong. Statistical inference of the value function for reinforcement learning in infinite horizon settings. *arXiv preprint arXiv:2001.04515*, 2020b.

Chengchun Shi, Sheng Zhang, Wenbin Lu, and Rui Song. Statistical inference of the value function for reinforcement learning in infinite horizon settings. *arXiv preprint arXiv:2001.04515*, 2020c.

Charles J. Stone. Optimal global rates of convergence for nonparametric regression. *Ann. Statist.*, 10(4):1040–1053, 1982. ISSN 0090-5364.

Richard S. Sutton and Andrew G. Barto. *Reinforcement learning: an introduction*. Adaptive Computation and Machine Learning. MIT Press, Cambridge, MA, second edition, 2018. ISBN 978-0-262-03924-6.

Richard S Sutton, Csaba Szepesvári, and Hamid Reza Maei. A convergent o(n) algorithm for off-policy temporal-difference learning with linear function approximation. *Advances in neural information processing systems*, 21(21):1609–1616, 2008.

Philip Thomas and Emma Brunskill. Data-efficient off-policy policy evaluation for reinforcement learning. In *International Conference on Machine Learning*, pp. 2139–2148, 2016.

Philip S Thomas, Georgios Theocharous, and Mohammad Ghavamzadeh. High-confidence off-policy evaluation. In *Twenty-Ninth AAAI Conference on Artificial Intelligence*, 2015.

Davide Viviano and Jelena Bradic. Synthetic learner: model-free inference on treatments over time. *arXiv preprint arXiv:1904.01490*, 2019.

Chien-Fu Jeff Wu et al. Jackknife, bootstrap and other resampling methods in regression analysis. *the Annals of Statistics*, 14(4):1261–1295, 1986.

Fanny Yang, Aaditya Ramdas, Kevin G Jamieson, and Martin J Wainwright. A framework for multi-a (rmed)/b (andit) testing with online fdr control. In *Advances in Neural Information Processing Systems*, pp. 5957–5966, 2017.

Baqun Zhang, Anastasios A. Tsiatis, Eric B. Laber, and Marie Davidian. Robust estimation of optimal dynamic treatment regimes for sequential treatment decisions. *Biometrika*, 100(3):681–694, 2013.

Yifan Zhou, Yang Liu, Ping Li, and Feifang Hu. Cluster-adaptive network a/b testing: From randomization to estimation. *arXiv preprint arXiv:2008.08648*, 2020.

Shaofeng Zou, Tengyu Xu, and Yingbin Liang. Finite-sample analysis for sarsa with linear function approximation. In *Advances in Neural Information Processing Systems*, pp. 8665–8675, 2019.

Input: no. of basis functions q, no. of bootstrap samples B, an α spending function $\alpha(\cdot)$.
Initialize: $\mathcal{I} = \{1, \cdots, B\}$. **Set** $\widehat{\boldsymbol{\Omega}}$, $\widehat{\boldsymbol{\Sigma}}_0$, $\widehat{\boldsymbol{\Sigma}}_1$ to zero matrcies, and $\widehat{\boldsymbol{\eta}}$, $\widehat{S}_1, \cdots, \widehat{S}_B$ to zero vectors.
Compute U (see equation 3) using either Monte Carlo methods or numerical integration.
For $k = 1$ to K:
    Step 1. Online update of ATE.
    For $t = T_{k-1}$ to $T_k - 1$:
        $\widehat{\boldsymbol{\Sigma}}_a = (1 - t^{-1})\widehat{\boldsymbol{\Sigma}}_a + t^{-1}\xi(S_t, A_t)\{\xi(S_t, A_t) - \gamma\xi(S_{t+1}, a)\}^\top, a = 0, 1$;
        $\widehat{\boldsymbol{\eta}} = (1 - t^{-1})\widehat{\boldsymbol{\eta}} + t^{-1}\xi(S_t, A_t)Y_t$.
    Set $(\widehat{\beta}_{a,0}^\top, \widehat{\beta}_{a,1}^\top)^\top = \widehat{\boldsymbol{\Sigma}}_a^{-1}\widehat{\boldsymbol{\eta}}$ for $a \in \{0, 1\}$ and $\widehat{\tau} = U^\top\widehat{\boldsymbol{\beta}}$.
    Step 2. Online update of the variance estimator.
    Initialize $\widehat{\boldsymbol{\Omega}}^*$ to a zero matrix.
    For $t = T_{k-1}$ to $T_k - 1$:
        $\widehat{\varepsilon}_{t,a} = Y_t + \gamma\Psi^\top(S_{t+1})\widehat{\beta}_{a,a} - \Psi^\top(S_t)\widehat{\beta}_{a,A_t}$ for $a = 0, 1$;
        $\widehat{\boldsymbol{\Omega}}^* = \widehat{\boldsymbol{\Omega}}^* + (\xi(S_t, A_t)^\top\widehat{\varepsilon}_{t,0}, \xi(S_t, A_t)^\top\widehat{\varepsilon}_{t,1})^\top(\xi(S_t, A_t)^\top\widehat{\varepsilon}_{t,0}, \xi(S_t, A_t)^\top\widehat{\varepsilon}_{t,1})$.
    Set $\widehat{\boldsymbol{\Sigma}}$ to a block diagonal matrix by aligning $\widehat{\boldsymbol{\Sigma}}_0$ and $\widehat{\boldsymbol{\Sigma}}_1$ along the diagonal of $\widehat{\boldsymbol{\Sigma}}$;
    Set $\widehat{\boldsymbol{\Omega}} = T_k^{-1}(T_{k-1}\widehat{\boldsymbol{\Omega}} + \widehat{\boldsymbol{\Omega}}^*)$ and the variance estimator $\widehat{\sigma}^2 = U^\top\widehat{\boldsymbol{\Sigma}}^{-1}\widehat{\boldsymbol{\Omega}}\{\widehat{\boldsymbol{\Sigma}}^{-1}\}^\top U$.
    Step 3. Bootstrap test statistic.
    For $b = 1$ to B:
        Generate $e_k^{(b)} \sim N(0, I_{4q})$; $\widehat{S}_b = \widehat{S}_b + \widehat{\boldsymbol{\Omega}}^{*1/2}e_k^{(b)}$; $\widehat{Z}_b^* = T_k^{-1/2}\widehat{\sigma}^{-1}U^\top\widehat{\boldsymbol{\Sigma}}^{-1}\widehat{S}_b$;
    Set z to be the upper $\{\alpha(t) - |\mathcal{I}^c|/B\}/(1 - |\mathcal{I}^c|/B)$-th percentile of $\{\widehat{Z}_b^*\}_{b\in\mathcal{I}}$.
    Update $\mathcal{I}$ and $\widehat{\boldsymbol{\Omega}}$ as $\mathcal{I} \leftarrow \{b \in \mathcal{I} : \widehat{Z}_b^* \leq z\}$;
    Step 4. Reject or not?
    Reject the null if $\sqrt{T_k}\widehat{\sigma}^{-1}\widehat{\tau} > z$.

Algorithm 1: The testing procedure

A   MORE ON THE ALGORITHM

A pseudo algorithm summarizing our procedure is given in Algorithm 1. We next introduce some notations. The vector U is defined by

$$\left\{ -\int_{s\in\mathbb{S}} \Psi(s)^\top\mathbb{G}(ds), 0_q^\top, 0_q^\top, \int_{s\in\mathbb{S}} \Psi(s)^\top\mathbb{G}(ds) \right\}^\top, \tag{3}$$

here 0_q denotes a zero vector of size q. The matrix $\widehat{\boldsymbol{\Omega}}(t)$ is defined by

$$\widehat{\boldsymbol{\Omega}}(t) = \frac{1}{t}\sum_{j=0}^{t-1} \begin{pmatrix} \xi_j\widehat{\varepsilon}_{j,0} \\ \xi_j\widehat{\varepsilon}_{j,1} \end{pmatrix} \begin{pmatrix} \xi_j\widehat{\varepsilon}_{j,0} \\ \xi_j\widehat{\varepsilon}_{j,1} \end{pmatrix}^\top, \tag{4}$$

where $\widehat{\varepsilon}_{j,0}$ and $\widehat{\varepsilon}_{j,1}$ are the temporal difference errors defined in Algorithm 1.

B   EXTENSIONS TO DYNAMIC POLICIES

In this paper, we focus on comparing the long-term treatment effects between two nondynamic policies. The proposed method can be easily extended to handle dynamic policies as well. Specifically, consider two time-homogeneous policies π_1 and π_2 where each $\pi_j(s)$ measures the treatment assignment probability $\Pr(A_t = 1|S_t = s)$. Note that the integrated value difference function τ_0 can be represented by

$$\int_s \{V(\pi_1; s) - V(\pi_2; s)\}\mathbb{G}(ds) = \int_s [\{Q(\pi_1; 1, s) - Q(\pi_1; 0, s)\}\pi_1(s)$$
$$-\{Q(\pi_2; 1, s) - Q(\pi_2; 0, s)\}\pi_2(s) + Q(\pi_1; 0, s) - Q(\pi_2; 0, s)]\mathbb{G}(ds).$$

The Q-estimators can be similarly computed via temporal difference learning. More specifically, for a given policy π, let

$$\widehat{Q}_t(\pi; a, s) = \Psi^\top(s)\widehat{\boldsymbol{\Sigma}}_\pi^{-1}(t)\left\{ \frac{1}{t}\sum_{j<t} \xi(S_j, A_j)Y_j \begin{pmatrix} A_j \\ 1 - A_j \end{pmatrix} \right\}, \tag{5}$$

be the Q-estimator given the data $\{(S_j, A_j, Y_j)\}_{j<t}$ where $\widehat{\Sigma}_\pi(t) = t^{-1} \sum_{j<t} \Sigma_j$ where Σ_j is defined by

$$\left[\begin{array}{cc} \Psi(S_j)(1-A_j)\{\Psi(S_j) - \gamma\Psi(S_{j+1})(1-\pi(S_{j+1}))\}^\top & -\gamma\Psi(S_j)(1-A_j)\Psi^\top(S_{j+1})\pi(S_{j+1}) \\ -\gamma\Psi(S_j)A_j\Psi^\top(S_{j+1})\pi(S_{j+1}) & \Psi(S_j)A_j\{\Psi(S_j) - \gamma\Psi(S_{j+1})(1-\pi(S_{j+1}))\}^\top \end{array} \right].$$

We can plug-in the Q-estimator in equation 5 to estimate τ_0. The corresponding variance estimator and the resulting test statistic can be similarly derived. A bootstrap procedure can be similarly developed as in Section 4.3 for sequential testing. We omit the details for brevity.

C Potential outcomes under a random policy

We focus on the class of Markov policies that are functions of current state variables only. Following Luckett et al. (2019), we define $\{\xi_b^t(\cdot)\}_{t\geq 0}$ as a sequence of independent, binary-valued stochastic processes that satisfy $\Pr\{\xi_b^t(s) = 1\} = b_t(s)$ for any $t \geq 0$ and $s \in \mathbb{S}$, where $b_t(\cdot)$ denotes the policy used at time t. The potential outcomes $Y_t^*(\bar{b}_t)$ and $S_{t+1}^*(\bar{b}_t)$ can thus be recursively defined as

$$Y_t^*(\bar{b}_t) = \sum_{\bar{a}_t \in \{0,1\}^{t+1}} Y_t^*(\bar{a}_t)\mathbb{I}(\xi_b^t(S_t^*(\bar{a}_{t-1})) = \bar{a}_t),$$

$$S_{t+1}^*(\bar{b}_t) = \sum_{\bar{a}_t \in \{0,1\}^{t+1}} S_{t+1}^*(\bar{a}_t)\mathbb{I}(\xi_b^t(S_t^*(\bar{a}_{t-1})) = \bar{a}_t),$$

for $t = 0, 1, \cdots$.

D More on the wild bootstrap algorithm

The idea is to generate bootstrap samples $\{\widehat{Z}^{\mathrm{MB}}(t)\}_t$ that have asymptotically the same joint distribution as $\{\sqrt{t}\widehat{\sigma}^{-1}(t)(\widehat{\tau}(t) - \tau_0)\}_t$. Specifically, let $\{\zeta_t\}_{t\geq 0}$ be a sequence of i.i.d. random variables independent of the observed data. Define

$$\widehat{\boldsymbol{\beta}}^{\mathrm{MB}}(t) = \widehat{\Sigma}^{-1}(t)\left\{ \frac{1}{t}\sum_{j<t} \xi(S_j, A_j)\zeta_j \left(\begin{array}{c} \widehat{\varepsilon}_{j,0} \\ \widehat{\varepsilon}_{j,1} \end{array} \right) \right\}, \tag{6}$$

where $\widehat{\varepsilon}_{t,a}$ is the temporal difference error defined in Algorithm 1. Based on $\widehat{\boldsymbol{\beta}}^{\mathrm{MB}}(t)$, one can define the bootstrap sample $\widehat{Z}^{\mathrm{MB}}(t) = \sqrt{t}\widehat{\sigma}^{-1}(t)\boldsymbol{U}^\top\widehat{\boldsymbol{\beta}}^{\mathrm{MB}}(t)$.

We remark that although the wild bootstrap method is developed under the i.i.d. settings, it is valid under our setup as well. This is due to that under CMIA, $\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^*$ forms a martingale sequence with respect to the filtration $\{(S_j, A_j, Y_j) : j < t\}$. This guarantees that the covariance matrices of $\widehat{\boldsymbol{\beta}}^{\mathrm{MB}}(t)$ and $\widehat{\boldsymbol{\beta}}(t)$ are asymptotically equivalent. As such, the bootstrap approximation is valid.

However, calculating $\widehat{\boldsymbol{\beta}}^{\mathrm{MB}}(T_k)$ requires $O(T_k)$ operations. The time complexity of the resulting bootstrap algorithm is $O(BT_k)$ up to the k-th interim stage, where B is the total number of bootstrap samples. This can be time consuming when $\{T_k - T_{k-1}\}_{k=1}^K$ are large.

E More on the designs

In D3, we require $b^{(k)}$ to be strictly bounded between 0 and 1. Suppose an ϵ-greedy policy is used, i.e. $b^{(k)}(s) = \epsilon/2 + (1-\epsilon)\widehat{\pi}^{(k)}(s)$, where $\widehat{\pi}^{(k)}$ denotes some estimated optimal policy. It follows that $\epsilon/2 \leq b^{(k)}(s) \leq 1 - \epsilon/2$ for any s. Such a requirement is automatically satisfied.

For any behaviour policy b in D1-D3, define $S_t^*(\bar{b}_{t-1})$ and $Y_t^*(\bar{b}_t)$ as the potential outcomes at time t, where $\bar{b}_t$ denotes the action history assigned according to b. When b is a random policy as in D1 or D3, definitions of these potential outcomes are more complicated than those under a deterministic policy (see Appendix C for details). When b is a stationary policy, it follows from MA that $\{S_{t+1}^*(\bar{b}_t)\}_{t\geq -1}$ forms a time-homogeneous Markov chain. When b follows the alternating-time-interval design, both $\{S_{2t}^*(\bar{b}_{2t-1})\}_{t\geq 0}$ and $\{S_{2t+1}^*(\bar{b}_{2t})\}_{t\geq 0}$ form time-homogeneous Markov chains.

F   TECHNICAL CONDITIONS

To simplify the presentation, we assume all state variables are continuous.

F.1   CONDITION C1

C1 Suppose (i) holds. Assume (ii) holds under D1, (iii) holds under D2 and (ii), (iv) hold under D3.
(i) The transition kernel $\mathcal{P}$ is absolutely continuous and satisfies $\mathcal{P}(ds; a, s') = p(s; a, s')ds$ for some transition density function p. In addition, assume p is uniformly bounded away from 0 and ∞.
(ii) The Markov chain $\{S_t^*(\bar{b}_{t-1}^{(0)})\}_{t \geq 0}$ formed under the behaviour policy $b^{(0)}$ is geometrically ergodic, i.e. there exists some function M on $\mathbb{S}$, some constant $0 \leq \rho < 1$ and some probability density function Π such that $\int_{s \in \mathbb{S}} M(s)\Pi(ds) < +\infty$ and

$$\left\| \Pr(S_t^*(\bar{b}_{t-1}^{(0)}) \in \mathcal{S}|S_0 = s) - \Pi(\mathcal{S}) \right\|_{TV} \leq M(s)\rho^t, \quad \forall t \geq 0, s \in \mathbb{S}, \mathcal{S} \subseteq \mathbb{S},$$

where $\| \cdot \|_{TV}$ denotes the total variation norm.
(iii) The Markov chains $\{S_{2t}^*(\bar{b}_{2t})\}_{t \geq 0}$ and $\{S_{2t+1}^*(\bar{b}_{2t+1})\}_{t \geq 0}$ are geometrically ergodic.
(iv) For any $k = 1, \cdots, K - 1$, the following events occur with probability tending to 1: the Markov chain $\{S_t^*(\bar{b}_{t-1}^{(k)})\}_{t \geq 0}$ is geometrically ergodic; $\sup_{s \in \mathbb{S}} |b^{(k)}(s) - b^*(s)| \xrightarrow{P} 0$ for some $b^*(\cdot)$; the stationary distribution of $\{S_t^*(\bar{b}_{t-1}^{(k)})\}_{t \geq 0}$ will converge to some Π^* in total variation.

Remark: By C1(ii), Π is the stationary distribution of $\{S_t^*(\bar{b}_{t-1}^{(0)})\}_{t \geq 0}$. It follows that

$$\Pi(\mathcal{S}) = \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} \mathcal{P}(\mathcal{S}; a, s)\{ab^{(1)}(s) + (1-a)b^{(0)}(s)\}\Pi(ds),$$

for any $\mathcal{S} \subseteq \mathbb{S}$. By C1(i), we obtain

$$\Pi(\mathcal{S}) = \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} \int_{s' \in \mathcal{S}} [a\{1 - b^{(0)}(s)\} + (1-a)b^{(0)}(s)]p(s'; a, s)ds'\Pi(ds)$$

$$= \int_{s' \in \mathcal{S}} \underbrace{\sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} [a\{1 - b^{(0)}(s)\} + (1-a)b^{(0)}(s)]p(s'; a, s)\Pi(ds)}_{\mu(s')} \, ds'. \tag{7}$$

This implies that $\mu(\cdot)$ is the density function of Π. Since p is uniformly bounded away from 0 and ∞, so is μ.

Under C1(iv), for any $k \in \{1, \cdots, K - 1\}$, there exist some $M^{(k)}(\cdot)$, $\Pi^{(k)}(\cdot)$ and $\rho^{(k)}$ that satisfy $\int_{s \in \mathbb{S}} M^{(k)}(s)\Pi^{(k)}(ds) < +\infty$ and

$$\left\| \Pr(S_t^*(\bar{b}_{t-1}^{(k)}) \in \mathcal{S}|S_0 = s) - \Pi^{(k)}(\mathcal{S}) \right\|_{TV} \leq M^{(k)}(s)\{\rho^{(k)}\}^t, \quad \forall t \geq 0, s \in \mathbb{S}, \mathcal{S} \subseteq \mathbb{S}, \tag{8}$$

with probability tending to 1. Since $b^{(k)}$ is a function of the observe data history, so are $M^{(k)}(\cdot)$, $\Pi^{(k)}(\cdot)$ and $\rho^{(k)}$.

Suppose an ϵ-greedy policy is used, i.e. $b^{(k)}(s) = \epsilon/2 + (1 - \epsilon)\widehat{\pi}^{(k)}(s)$ where $\widehat{\pi}^{(k)}$ denotes some estimated optimal policy. Then the condition $\sup_{s \in \mathbb{S}} |b^{(k)}(s) - b^*(s)| \xrightarrow{P} 0$ requires $\widehat{\pi}^{(k)}$ to converge. The total variation distance between the one-step transition kernel under $\bar{b}^{(k)}$ and that under b^* can be bounded by

$$\sup_s |\Pr(S_1^*(b^{(k)}) \in \mathcal{S}|S_0 = s) - \Pr(S_1^*(b^*) \in \mathcal{S}|S_0 = s)| \leq \sup_s |b^{(k)}(s) - b^*(s)| \sup_{s,s',a} p(s'; a, s),$$

and converges to zero in probability. When the markov chain $\{S_t^*(\bar{b}_{t-1}^{(k)})\}_{t \geq 0}$ is uniformly ergodic, it follows from Theorems 2 and 3 of Rabta & Aïssani (2018) that $\|\Pi^{(k)} - \Pi^*\|_{TV} \to 0$ where Π^* corresponds to the stationary distribution of $\{S_t^*(\bar{b}_{t-1}^*)\}$. The last condition in C1(iv) is thus satisfied.

F.2 CONDITION C2

C2(i) Assume there exists some β^* such that

$$\sup_{a',a\in\{0,1\},s\in\mathbb{S}} |Q(a';a,s) - \Psi^\top(s)\beta^*_{a',a}| = o(T^{-1/2}).$$

(ii) Assume there exists some constant $\bar{c}^* \geq 1$ such that

$$(\bar{c}^*)^{-1} \leq \lambda_{\min}\left\{\int_{s\in\mathbb{S}} \Psi(s)\Psi^\top(s)ds\right\} \leq \lambda_{\max}\left\{\int_{s\in\mathbb{S}} \Psi(s)\Psi^\top(s)ds\right\} \leq \bar{c}^*, \tag{9}$$

and $\sup_s \|\Psi(s)\|_2 = O(\sqrt{q})$.
(iii) Assume $\liminf_q \|\int_{s\in\mathbb{S}} \Psi(s)\mathbb{G}(ds)\|_2 > 0$.

Remark: For any $a, a' \in \{0, 1\}$, suppose $Q(a'; a, s)$ is p-smooth as a function of s (see e.g. Stone, 1982, for the definition of p-smoothness). When tensor product B-splines or wavelet basis functions (see Section 6 of Chen & Christensen, 2015, for an overview of these bases) are used for $\Psi(\cdot)$, we have

$$\sup_{a',a\in\{0,1\},s\in\mathbb{S}} |Q(a';a,s) - \Psi^\top(s)\beta^*_{a',a}| = o(q^{-p/d}),$$

under certain mild conditions. See Section 2.2 of Huang (1998) for details. It follows that Condition C2(i) automatically holds when the number of basis functions q satisfies $q \gg T^{d/(2p)}$.

Condition C2(ii) is satisfied when tensor product B-splines or wavelet basis is used. For B-spline basis, the assertion in equation 9 follows from the arguments used in the proof of Theorem 3.3, Burman & Chen (1989). For wavelet basis, the assertion in equation 9 follows from the arguments used in the proof of Theorem 5.1, Chen & Christensen (2015). For both bases, the number of nonzero elements in $\Psi(\cdot)$ is bounded by some constant. Moreover, each basis function is uniformly bounded by $O(\sqrt{q})$. The condition $\sup_s \|\Psi(s)\|_2 = O(\sqrt{q})$ thus holds.

Condition C2(iii) automatically holds for tensor product B-splines basis. Notice that $\mathbf{1}^\top \Psi(s) = q^{1/2}$ for any $s \in \mathbb{S}$ where $\mathbf{1}$ denotes a vector of ones. It follows from Cauchy-Schwarz inequality that

$$\sqrt{q} \left\| \int_{s\in\mathbb{S}} \Psi(s)\mathbb{G}(ds) \right\|_2 \geq \left\| \int_{s\in\mathbb{S}} \mathbf{1}^\top \Psi(s)\mathbb{G}(ds) \right\|_2 = \sqrt{q}.$$

C2(iii) is thus satisfied.

F.3 CONDITION C3

Let $\varepsilon^*(a', a) = Y_0^*(a) + \gamma Q(a'; a', S_1^*(a)) - Q(a'; a, S_0)$.

C3 Assume $\inf_q \inf_{a',a\in\{0,1\},s\in\mathbb{S}} \text{Var}\{\varepsilon^*(a',a)|S_0 = s\} > 0$ and $\sup_q \sup_{a\in\{0,1\},s\in\mathbb{S}} \rho_\varepsilon(a,s) < 1$
where

$$\rho_\varepsilon(a,s) = \frac{\text{E}\{\varepsilon^*(0,a)\varepsilon^*(1,a)|S_0 = s\}}{\sqrt{\text{Var}(\varepsilon^*(0,a)|S_0 = s)\text{Var}(\varepsilon^*(1,a)|S_0 = s)}}.$$

Here, ρ_ε corresponds to the partial correlation of $\varepsilon^*(0, a)$ and $\varepsilon^*(1, a)$ given S_0.

G TECHNICAL PROOFS

G.1 PROOF OF LEMMA 1

To prove Lemma 1, we state the following lemma.

Lemma 2 *Under MA and CMIA, $Q(a'; a, s) = r(a, s) + \gamma \int_{s'} Q(a'; a', s')\mathcal{P}(ds'; a, s)$ for any (s, a).*

Proof of Lemma 2: For any $a, a' \in \{0, 1\}$, define the potential outcome $Y_t^*(a', a)$ and $S_t^*(a', a)$ as the reward and state variables that would occur at time t had the agent assigned Treatment a at the initial time point and Treatment a' afterwards.

Let $\mathcal{P}_{a'}^t(\mathbb{S}, a, s) = \Pr\{S_t^*(a', a) \in \mathbb{S} | S_0 = s\}$ for any $\mathbb{S} \subseteq \mathbb{S}, a, a' \in \{0, 1\}, s \in \mathbb{S}$ and $t \geq 0$. We break the proof into two parts. In Part 1, we show Lemma 2 holds when the following is satisfied:

$$\Pr\{S_{t+1}^*(a', a) \in \mathbb{S} | S_1^*(a) = s, S_0\} = \mathcal{P}_{a'}^t(\mathbb{S}, a', s), \tag{10}$$

In Part2, we show equation 10 holds.

Part 1: Under CMIA, we have

$$\begin{aligned}
\mathrm{E}\{Y_t^*(a', a) | S_0 = s\} &= \mathrm{E}[\mathrm{E}\{Y_t^*(a', a) | S_t^*(a', a), S_0 = s\} | S_0 = s] \\
&= \mathrm{E}\{r(\pi(S_t^*(a', a)), S_t^*(a', a)) | S_0 = s\}.
\end{aligned} \tag{11}$$

It follows that

$$Q(a'; a, s) = \sum_{t \geq 0} \gamma^t \mathrm{E}\{r(\pi(S_t^*(a', a)), S_t^*(a', a)) | S_0 = s\}. \tag{12}$$

Similar to equation 11, we can show

$$\begin{aligned}
\mathrm{E}\{Y_{t+1}^*(a', a) | S_0 = s\} &= \mathrm{E}\{r(\pi(S_{t+1}^*(a', a)), S_{t+1}^*(a', a)) | S_0 = s\} \\
&= \mathrm{E}[\mathrm{E}\{r(\pi(S_{t+1}^*(a', a)), S_{t+1}^*(a', a)) | S_1^*(a), S_0 = s\} | S_0 = s],
\end{aligned}$$

and hence

$$\sum_{t \geq 0} \gamma^t \mathrm{E}\{Y_{t+1}^*(a', a) | S_0 = s\} = \mathrm{E}\left[\sum_{t \geq 0} \gamma^t \, \mathrm{E}\{r(\pi(S_{t+1}^*(a', a)), S_{t+1}^*(a', a)) \big| S_1^*(a), S_0 = s\} | S_0 = s\right].$$

By equation 10, the conditional distribution of $S_{t+1}^*(a', a)$ given $S_1^*(a) = s$ and S_0 are the same as the conditional distribution of $S_t^*(a', a)$ given $S_0 = s$. It follows that from equation 12 that

$$\sum_{t \geq 0} \gamma^t \mathrm{E}\{Y_{t+1}^*(a', a) | S_0 = s\} = \mathrm{E}\{Q(a'; a, S_1^*(a)) | S_0 = s\}.$$

This together with the definition of Q function and CMIA yields

$$Q(a'; a, s) = r(a, s) + \gamma \left[\sum_{t \geq 0} \gamma^t \mathrm{E}\{Y_{t+1}^*(a', a) | S_0 = s\}\right] = r(a, s) + \gamma \mathrm{E}\{Q(a'; a, S_1^*(a)) | S_0 = s\}. \tag{13}$$

Under MA, we have

$$\mathrm{E}\{Q(a'; a, S_1^*(a)) | S_0 = s\} = \int_{s' \in \mathbb{S}} Q(a'; a, s') \mathcal{P}(ds'; a, s).$$

Combining this together with equation 13 yields the desired result.

Part 2: We use induction to prove equation 10. When $t = 0$, it trivially holds.

Suppose equation 10 holds for $t = k$. In the following, we show equation 10 holds for $t = k + 1$. Under MA, we have

$$\begin{aligned}
\Pr\{S_{k+2}^*(a', a) \in \mathbb{S} | S_1^*(a) = s, S_0\} &= \mathrm{E}[\Pr\{S_{k+2}^*(a', a) \in \mathbb{S} | S_{k+1}^*(a', a), S_1^*(a) = s, S_0\} | S_1^*(a) = s, S_0] \\
&= \mathrm{E}[\mathcal{P}(\mathbb{S}; a', S_{k+1}^*(a', a)) | S_1^*(a) = s, S_0].
\end{aligned}$$

Since we have shown equation 10 holds for $t = k$, it follows that

$$\Pr\{S_{k+2}^*(a', a) \in \mathbb{S} | S_1^*(a) = s, S_0\} = \int_{s' \in \mathbb{S}} \mathcal{P}(\mathbb{S}; a', s') \mathcal{P}_{a'}^k(ds', a', s).$$

Similarly, we can show

$$\mathcal{P}_{a'}^{k+1}(\mathbb{S}, a', s) = \Pr\{S_{k+1}^*(a', a') \in \mathbb{S} | S_0 = s\} = \int_{s' \in \mathbb{S}} \mathcal{P}(\mathbb{S}; a', s') \mathcal{P}_{a'}^k(ds', a', s).$$

The proof is hence completed.

Proof of Lemma 1: By CA, it is equivalent to show

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\}\varphi(A_t, S_t^*(\bar{A}_{t-1})) = 0.$$

Let $\mathbb{S}_0$ denote the support of S_0. For any $s_0 \in \mathbb{S}_0$, it suffices to show

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\varphi(A_t, S_t^*(\bar{A}_{t-1}))|S_0 = s_0\} = 0.$$

This is equivalent to show

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\varphi(A_t, S_t^*(\bar{A}_{t-1}))\mathbb{I}(A_0 = a_0)\}|S_0 = s_0] = 0,$$

for any $s_0 \in \mathbb{S}_0$, $a_0 \in \{0, 1\}$.

Let $\mathcal{A}_0(s_0) = \{a \in \{0, 1\} : \Pr(A_0 = a|S_0 = s) > 0\}$. It suffices to show for any $s_0 \in \mathbb{S}_0, a_0 \in \mathcal{A}_0(s_0)$,

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\varphi(A_t, S_t^*(\bar{A}_{t-1}))\mathbb{I}(A_0 = a_0)|S_0 = s_0\} = 0,$$

or equivalently,

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\varphi(A_t, S_t^*(\bar{A}_{t-1}))|S_0 = s_0, A_0 = a_0\} = 0. \quad (14)$$

Let $\bar{s}_j = (s_0, s_1, \cdots, s_j)^\top$, $\bar{y}_j = (y_0, y_1, \cdots, y_j)^\top$, $\bar{S}_j = (S_0, S_1, \cdots, S_j)^\top$ and $\bar{Y}_j = (Y_0, Y_1, \cdots, Y_j)^\top$. We can recursively define the sets $\mathcal{Y}_j(\bar{s}_j, \bar{a}_j, \bar{y}_{j-1})$, $\mathbb{S}_{j+1}(\bar{s}_j, \bar{a}_j, \bar{y}_j)$, $\mathcal{A}_{j+1}(\bar{s}_{j+1}, \bar{a}_j, \bar{y}_j)$ to be the supports of Y_j, S_{j+1}, A_{j+1} conditional on $(\bar{S}_j = \bar{s}_j, \bar{A}_j = \bar{a}_j, \bar{Y}_{j-1} = \bar{y}_{j-1})$, $(\bar{S}_j = \bar{s}_j, \bar{A}_j = \bar{a}_j, \bar{Y}_j = \bar{y}_j)$, $(\bar{S}_{j+1} = \bar{s}_{j+1}, \bar{A}_j = \bar{a}_j, \bar{Y}_j = \bar{y}_j)$ respectively, for $j \geq 0$. Similar to equation 14, it suffices to show

$$\mathrm{E}\{Q(a'; A_t, S_t^*(\bar{A}_{t-1})) - Y_t^*(\bar{A}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{A}_t))\varphi(A_t, S_t^*(\bar{A}_{t-1}))|\bar{S}_t = \bar{s}_t, \bar{A}_t = \bar{a}_t, \bar{Y}_{t-1} = \bar{y}_{t-1}\} = 0,$$

for any $s_0 \in \mathbb{S}_0, a_0 \in \mathcal{A}_0(s_0), y_0 \in \mathcal{Y}_0(s_0, a_0), \cdots, s_t \in \mathbb{S}_t(\bar{s}_{t-1}, \bar{a}_{t-1}, \bar{y}_{t-1}), a_t \in \mathcal{A}_t(\bar{s}_t, \bar{a}_{t-1}, \bar{y}_{t-1})$. This is equivalent to show

$$\mathrm{E}\{Q(a'; a_t, S_t^*(\bar{a}_{t-1})) - Y_t^*(\bar{a}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{a}_t))|\bar{S}_t = \bar{s}_t, \bar{A}_t = \bar{a}_t, \bar{Y}_{t-1} = \bar{y}_{t-1}\} = 0. \quad (15)$$

By construction, we have $\Pr(A_t = a_t|\bar{S}_t = \bar{s}_t, \bar{Y}_{t-1} = \bar{y}_{t-1}, \bar{A}_{t-1} = \bar{a}_{t-1}) > 0$. Under SRA, the left-hand-side (LHS) of equation 15 equals

$$\mathrm{E}\{Q(a'; a_t, S_t^*(\bar{a}_{t-1})) - Y_t^*(\bar{a}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{a}_t))|\bar{S}_t = \bar{s}_t, \bar{A}_{t-1} = \bar{a}_{t-1}, \bar{Y}_{t-1} = \bar{y}_{t-1}\}. \quad (16)$$

Notice that the conditioning event is the same as $\{S_t^*(\bar{a}_{t-1}) = s_t, Y_{t-1}^*(\bar{a}_{t-1}) = y_{t-1}, \bar{S}_{t-1} = \bar{s}_{t-1}, \bar{A}_{t-1} = \bar{a}_{t-1}, \bar{Y}_{t-2} = \bar{y}_{t-2}\}$. Under SRA, equation 16 equals

$$\mathrm{E}\{Q(a'; a_t, S_t^*(\bar{a}_{t-1})) - Y_t^*(\bar{a}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{a}_t))|S_t^*(\bar{a}_{t-1}) = s_t, Y_{t-1}^*(\bar{a}_{t-1}) = y_{t-1},$$
$$\bar{S}_{t-1} = \bar{s}_{t-1}, \bar{A}_{t-2} = \bar{a}_{t-2}, \bar{Y}_{t-2} = \bar{y}_{t-2}\}.$$

By recuisvely applying SRA, we can show the left-hand-side of equation 15 equals

$$\mathrm{E}[Q(a'; a_t, S_t^*(\bar{a}_{t-1})) - Y_t^*(\bar{a}_t) - \gamma Q(a'; a', S_{t+1}^*(\bar{a}_t))|\{S_j^*(\bar{a}_{j-1}) = s_j\}_{1 \leq j \leq t}, \{Y_j^*(\bar{a}_j) = y_j\}_{1 \leq j \leq t-1}].$$

This is equal to zero by MA, CMIA and Lemma 2. The proof is hence completed.

G.2   PROOF OF THEOREM 1

G.2.1   PROOF UNDER D1

We begin by providing an outline of the proof. The proof is divided into three steps. In the first step, we show for any $T_1 \leq t \leq T_k$, the estimator $\widehat{\boldsymbol{\beta}}(t)$ satisfies the following linear representation,

$$\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^* = \boldsymbol{\Sigma}^{-1}(t) \underbrace{\left\{ \frac{1}{t} \sum_{j=0}^{t-1} \left(\begin{array}{c} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{array} \right) \right\}}_{\zeta_1(t)} + o_p(t^{-1/2}), \quad (17)$$

where $\boldsymbol{\Sigma}(t) = \mathrm{E}\widehat{\boldsymbol{\Sigma}}(t)$ and $\varepsilon_{j,a} = Y_j + \gamma Q(a; a, S_{j+1}) - Q(a; A_j, S_j)$ for $a = 0, 1$. Based on this representation, in the second step, we show the asymptotic normality of $\widehat{\tau}(t)$. Specifically, we show

$$\frac{\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}}{\widehat{\sigma}(t)} \xrightarrow{d} N(0, 1).$$

In the last step, we prove Theorem 1.

Part 1: By definition, we have

$$
\begin{aligned}
\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^* &= \widehat{\boldsymbol{\Sigma}}^{-1}(t)\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j Y_j \\ \xi_j Y_j\end{pmatrix} - \widehat{\boldsymbol{\Sigma}}(t)\boldsymbol{\beta}^*\right\} = \widehat{\boldsymbol{\Sigma}}^{-1}(t)\left[\frac{1}{t}\sum_{j=0}^{t-1}\left\{\begin{pmatrix}\xi_j Y_j \\ \xi_j Y_j\end{pmatrix} - \boldsymbol{\Sigma}_j\boldsymbol{\beta}^*\right\}\right] \\
&= \widehat{\boldsymbol{\Sigma}}^{-1}(t)\left[\frac{1}{t}\sum_{j=0}^{t-1}\left\{\begin{array}{c}\xi_j\{Y_j - \Psi^\top(S_j)\beta_{0,A_j}^* + \gamma\Psi^\top(S_{j+1})\beta_{0,0}^*\} \\ \xi_j\{Y_j - \Psi^\top(S_j)\beta_{1,A_j}^* + \gamma\Psi^\top(S_{j+1})\beta_{1,1}^*\}\end{array}\right\}\right],
\end{aligned}
$$

where the last equality is due to the definition of $\boldsymbol{\Sigma}_j$. Let

$$r_{a,j} = \Psi^\top(S_j)\beta_{a,A_t}^* - \gamma\Psi^\top(S_{j+1})\beta_{a,0}^* - Q(a; A_j, S_j) + \gamma Q(a; a, S_{j+1}).$$

It follows that

$$\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^* = \widehat{\boldsymbol{\Sigma}}^{-1}(t)\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0} \\ \xi_j\varepsilon_{j,1}\end{pmatrix}\right\} - \widehat{\boldsymbol{\Sigma}}^{-1}(t)\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j r_{j,0} \\ \xi_j r_{j,1}\end{pmatrix}\right\},$$

and hence

$$
\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^* = \boldsymbol{\Sigma}^{-1}(t)\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0} \\ \xi_j\varepsilon_{j,1}\end{pmatrix}\right\} + \underbrace{\{\widehat{\boldsymbol{\Sigma}}^{-1}(t) - \boldsymbol{\Sigma}^{-1}(t)\}\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0} \\ \xi_j\varepsilon_{j,1}\end{pmatrix}\right\}}_{\zeta_2(t)}
$$

$$
\underbrace{-\widehat{\boldsymbol{\Sigma}}^{-1}(t)\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j r_{j,0} \\ \xi_j r_{j,1}\end{pmatrix}\right\}}_{\zeta_3(t)}.
$$

We first consider $\zeta_3(t)$. It can be upper bounded by

$$
\|\widehat{\boldsymbol{\Sigma}}^{-1}(t)\|_2 \left\|\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0} \\ \xi_j\varepsilon_{j,1}\end{pmatrix}\right\|_2 = \|\widehat{\boldsymbol{\Sigma}}^{-1}(t)\|_2 \max_{a\in\{0,1\}}\sup_{\|\boldsymbol{a}\|_2=1}\left|\boldsymbol{a}^\top\left(\frac{1}{t}\sum_{j=0}^{t-1}\xi_j r_{a,j}\right)\right|
$$

$$
\leq \|\widehat{\boldsymbol{\Sigma}}^{-1}(t)\|_2 \max_{a\in\{0,1\}}\sup_{\|\boldsymbol{a}\|_2=1}\sqrt{\frac{1}{t}\sum_{j=0}^{t-1}(\boldsymbol{a}^\top\xi_j)^2 r_{a,j}^2} \leq \max_{a,j}|r_{a,j}|\|\widehat{\boldsymbol{\Sigma}}^{-1}(t)\|_2\sqrt{\lambda_{\max}\left(\frac{1}{t}\sum_{j=0}^{t-1}\xi_j\xi_j^\top\right)},
$$

where the second follows from Cauchy-Schwarz inequality. Under Condition C2(i), we have for any $j \leq t \leq T_k$, $\max_j|r_{a,j}| = o(t^{-1/2})$. Suppose for now, we have shown

$$\|\widehat{\boldsymbol{\Sigma}}^{-1}(t)\|_2 = O_p(1) \quad \text{and} \quad \lambda_{\max}\left(\frac{1}{t}\sum_{j=0}^{t-1}\xi_j\xi_j^\top\right) = O_p(1). \tag{18}$$

It follows that

$$\zeta_3(t) = o_p(t^{-1/2}). \tag{19}$$

To bound $\zeta_2(t)$, notice that for any $a \in \{0, 1\}$,

$$\mathrm{E}\left\|\frac{1}{t}\sum_{j=0}^{t-1}\xi_j\varepsilon_{j,a}\right\|_2^2 = \frac{1}{t^2}\sum_{j=0}^{t-1}\mathrm{E}\xi_j^\top\xi_j\varepsilon_{j,a}^2 + \frac{1}{t^2}\sum_{j_1\neq j_2}\mathrm{E}\xi_{j_1}^\top\xi_{j_2}\varepsilon_{j_1,a}\varepsilon_{j_2,a}.$$

Similar to Lemma 1, we can show for any $\varphi(\cdot)$ that is a function of $\bar{A}_t, \bar{S}_t, \bar{Y}_{t-1}$ that

$$\mathrm{E}\{Q(a'; A_t, S_t) - Y_t - \gamma Q(a'; a', S_{t+1})\}\varphi(\bar{S}_t, \bar{A}_t, \bar{Y}_{t-1}) = 0. \tag{20}$$

This implies that $\mathrm{E}\xi_{j_1}^\top \xi_{j_2}\varepsilon_{j_1,a}\varepsilon_{j_2,a} = 0$ for any $j_1 \neq j_2$. It follows that

$$\mathrm{E}\left\|\frac{1}{t}\sum_{j=0}^{t-1}\xi_j\varepsilon_{j,a}\right\|_2^2 = \frac{1}{t^2}\sum_{j=0}^{t-1}\mathrm{E}\xi_j^\top \xi_j\varepsilon_{j,a}^2 \leq q\lambda_{\max}\left(\frac{1}{t^2}\sum_{j=0}^{t-1}\mathrm{E}\xi_j\xi_j^\top \varepsilon_{j,a}^2\right).$$

Since all immediate rewards are uniformly bounded, so is the Q function. As a result, $|\varepsilon_{j,a}|$'s are uniformly bounded. Suppose for now, we have shown

$$\lambda_{\max}\left(\frac{1}{t}\sum_{j=0}^{t-1}\mathrm{E}\xi_j\xi_j^\top\right) = O(1). \tag{21}$$

It follows that $\mathrm{E}\|t^{-1}\sum_{j=0}^{t-1}\xi_j\varepsilon_{j,a}\|_2^2 = O(q)$ and hence

$$\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0}\\ \xi_j\varepsilon_{j,1}\end{pmatrix} = O_p(t^{-1/2}\sqrt{q}).$$

Suppose

$$\|\widehat{\boldsymbol{\Sigma}}^{-1}(t) - \boldsymbol{\Sigma}^{-1}(t)\|_2 = o_p(q^{-1/2}). \tag{22}$$

It follows that

$$\|\zeta_2(t)\|_2 \leq \|\widehat{\boldsymbol{\Sigma}}^{-1}(t) - \boldsymbol{\Sigma}^{-1}(t)\|_2\left\|\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0}\\ \xi_j\varepsilon_{j,1}\end{pmatrix}\right\|_2 = o_p(t^{-1/2}).$$

This together with equation 19 yields equation 17.

It remains to show equation 18, equation 21 and equation 22 hold. We summarize these results in Lemma 3.

Lemma 3 *Under the given conditions, we have equation 18, equation 21 and equation 22 hold.*

Part 2: By definition, we have $\widehat{\tau}(t) - \tau_0 = \boldsymbol{U}^\top\{\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^*\} + \boldsymbol{U}^\top\boldsymbol{\beta}^* - \tau_0$. Define

$$\boldsymbol{\Omega}(t) = \mathrm{E}\left\{\frac{1}{t}\sum_{j=0}^{t-1}\begin{pmatrix}\xi_j\varepsilon_{j,0}\\ \xi_j\varepsilon_{j,1}\end{pmatrix}\begin{pmatrix}\xi_j\varepsilon_{j,0}\\ \xi_j\varepsilon_{j,1}\end{pmatrix}^\top\right\}$$

The asymptotic variance of $\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}$ is given by $\sigma^2(t) = \boldsymbol{U}^\top\boldsymbol{\Sigma}^{-1}(t)\boldsymbol{\Omega}(t)\{\boldsymbol{\Sigma}^{-1}(t)\}^\top\boldsymbol{U}$. We begin by providing a lower bound for $\sigma^2(t)$. Notice that

$$\sigma^2(t) \geq \lambda_{\min}\{\boldsymbol{\Omega}(t)\}\|\boldsymbol{U}^\top\boldsymbol{\Sigma}^{-1}(t)\|_2^2 \geq \lambda_{\min}\{\boldsymbol{\Omega}(t)\}\lambda_{\min}[\boldsymbol{\Sigma}^{-1}(t)\{\boldsymbol{\Sigma}^{-1}(t)\}^{-1}]\|\boldsymbol{U}\|_2^2. \tag{23}$$

Under C1(iii), we have $\liminf_q \|\boldsymbol{U}\|_2^2 > 0$.

In addition, notice that $\boldsymbol{\Sigma}^{-1}(t)\{\boldsymbol{\Sigma}^{-1}(t)\}^{-1}$ is positive semi-definite. It follows that $\lambda_{\min}[\boldsymbol{\Sigma}^{-1}(t)\{\boldsymbol{\Sigma}^{-1}(t)\}^{-1}] = 1/\lambda_{\max}[\boldsymbol{\Sigma}(t)\{\boldsymbol{\Sigma}(t)\}]$. Using similar arguments in showing $\|\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\|_2 = O(1)$ in the proof of Lemma 3, we can show $\sup_{t\geq 1}\|\boldsymbol{\Sigma}(t)\|_2 = O(1)$ and hence $\sup_{t\geq 1}\lambda_{\max}[\boldsymbol{\Sigma}(t)\{\boldsymbol{\Sigma}(t)\}] = O(1)$. This further yields

$$\inf_{t\geq 1}\lambda_{\min}[\boldsymbol{\Sigma}^{-1}(t)\{\boldsymbol{\Sigma}^{-1}(t)\}^{-1}] > 0. \tag{24}$$

Suppose $\boldsymbol{\Omega}(t)$ satisfies

$$\liminf_t \lambda_{\min}\{\boldsymbol{\Omega}(t)\} > 0. \tag{25}$$

It follows that $\sigma^2(t)$ is bounded away from zero, for sufficiently large t. Under Condition C2(i), we have $\boldsymbol{U}^\top \boldsymbol{\beta}^* - \tau_0 = o(T_k^{-1/2}) = o(t^{-1/2})$. It follows that

$$\frac{\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}}{\sigma(t)} = \frac{\sqrt{t}\boldsymbol{U}^\top\{\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^*\}}{\sigma(t)} + \frac{\sqrt{t}(\boldsymbol{U}^\top\boldsymbol{\beta}^* - \tau_0)}{\sigma(t)} = \frac{\sqrt{t}\boldsymbol{U}^\top\{\widehat{\boldsymbol{\beta}}(t) - \boldsymbol{\beta}^*\}}{\sigma(t)} + o(1).$$

Moreover, it follows from equation 23, equation 24 and equation 25 that $\sigma(t)/\|\boldsymbol{U}\|_2$ is uniformly bounded away from zero, for sufficiently large t. Combining this together with equation 17 yields

$$\frac{\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}}{\sigma(t)} = \frac{\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)}{\sigma(t)} + \frac{\sqrt{t}\boldsymbol{U}^\top R_t}{\sigma(t)},$$

where the remainder term satisfies $\|R_t\|_2 = o_p(t^{-1/2})$. It follows that the second term on the right-hand-side (RHS) of the above expression is bounded from above by $\sqrt{t}\|R_t\|_2\|\boldsymbol{U}\|_2/\sigma(t) = o_p(1)$ and hence

$$\frac{\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}}{\sigma(t)} = \frac{\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)}{\sigma(t)} + o_p(1). \tag{26}$$

Similar to the proof of Lemma 1, we can show for any $j \geq 0$, $a \in \{0, 1\}$,

$$\mathrm{E}(\xi_j\varepsilon_{j,a}|\{S_i, A_i, Y_i\}_{i<j}) = 0.$$

By the definition of $\zeta_1(t)$, $\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)/\sigma(t)$ forms a martingle with respect to the filtration $\sigma(\{S_j, A_j, Y_j\}_{j<t})$, i.e. the σ-algebra generated by $\{S_j, A_j, Y_j\}_{j<t}$. By the martingale central limit theorem, we can show $\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)/\sigma(t) \xrightarrow{d} N(0,1)$ (see Lemma 4 for details).

To complete the proof of Part 2, we need to show equation 25 holds and that $\widehat{\sigma}(t)/\sigma(t) \xrightarrow{P} 1$. The assertion $\widehat{\sigma}(t)/\sigma(t) \xrightarrow{P} 1$ can be similarly proven using arguments from Step 3 of the proof of Theorem 1, Shi et al. (2020b). We show the asymptotic normality of $\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)/\sigma(t)$ and that equation 25 holds in the following lemma.

Lemma 4 *Under the given conditions, we have equation 25 holds and that $\sqrt{t}\boldsymbol{U}^\top\zeta_1(t)/\sigma(t) \xrightarrow{d} N(0,1)$.*

Part 3: Results in Part 2 yield that $\sqrt{T_k}\{\widehat{\tau}(T_k) - \tau_0\}/\sigma(T_k) \xrightarrow{d} N(0,1)$ for each $1 \leq k \leq K$. In addition, for any K-dimensional vector $\boldsymbol{a} = (a_1, \cdots, a_K)^\top$, it follows from equation 26 that

$$\sum_{k=1}^{K} \frac{a_k\sqrt{T_k}\{\widehat{\tau}(T_k) - \tau_0\}}{\sigma(T_k)} = \sum_{k=1}^{K} \frac{a_k\sqrt{T_k}\boldsymbol{U}^\top\zeta_1(T_k)}{\sigma(T_k)} + o_p(1).$$

The leading term on the RHS can be rewritten as a weighted sum of $\{\xi_j\varepsilon_{j,0}, \xi_j\varepsilon_{j,1}\}_{0 \leq j < t}$. Similar to the proof in Part 2, we can show it forms a martingale with respect to the filtration $\sigma(\{S_j, A_j, Y_j\}_{j<t})$. We now derive its asymptotic normality for any $\boldsymbol{a}$, using the martingale central limit theorem for triangular arrays.

By Corollary 2 of McLeish (1974), we need to verify the following two conditions:

(a) $\max_{0 \leq j < t} |\sum_{k=1}^{K} a_k T_k^{-1/2}\boldsymbol{U}^\top\boldsymbol{\Sigma}^{-1}(T_k)(\xi_j^\top\varepsilon_{j,0}, \xi_j^\top\varepsilon_{j,1})^\top\{\sigma(T_k)\}^{-1}\mathbb{I}(j < T_k)| \xrightarrow{P} 0$;
(b) $\sum_{j=0}^{T-1} |\sum_{k=1}^{K} a_k T_k^{-1/2}\boldsymbol{U}^\top\boldsymbol{\Sigma}^{-1}(T_k)(\xi_j^\top\varepsilon_{j,0}, \xi_j^\top\varepsilon_{j,1})^\top\{\sigma(T_k)\}^{-1}\mathbb{I}(j < T_k)|^2$ converges to some constant in probability.

Since K is fixed, to verify (a), it suffices to show

$$\max_{1 \leq j < t, 1 \leq k \leq K} T_k^{-1/2}|\boldsymbol{U}^\top\boldsymbol{\Sigma}^{-1}(T_k)(\xi_j^\top\varepsilon_{j,0}, \xi_j^\top\varepsilon_{j,1})^\top\{\sigma(T_k)\}^{-1}| \xrightarrow{P} 0.$$

In Lemma 3, we have shown $\|\boldsymbol{\Sigma}^{-1}(t)\| = O(1)$. In Part 1 and Part 2 of the proof, we have shown $|\varepsilon_{j,a}|$'s are uniformly bounded and that $\sigma(t)/\|\boldsymbol{U}\|_2$ is bounded away from zero. Therefore, it suffices to show $T_1^{-1/2}\max_{0 \leq j < t}\|\xi_j\|_2 \xrightarrow{P} 0$. Under Condition C2(ii), we have $\sup_s\|\Psi(s)\|_2 = O(q^{1/2})$

and hence $\max_{0 \leq j < t} \|\xi_j\|_2 = O(q^{1/2})$. The assertion thus follows by noting that $T_1 = c_1 T$ and $q = o(T)$.

Using similar arguments in Step 3 of the proof of Shi et al. (2020b), we can show

$$\left\| \frac{1}{t} \sum_{j=0}^{t-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1}) - \mathbf{\Omega}(t) \right\|_2 \xrightarrow{P} 0, \tag{27}$$

as $t \to \infty$. This together with the facts $\|\mathbf{\Sigma}^{-1}(t)\| = O(1)$ and $\sigma(t)/\|\mathbf{U}\|_2$ is bounded away from zero implies that

$$\left| \frac{a_{k_1} a_{k_2}}{\sqrt{T_{k_1} T_{k_2}} \sigma^2(T_{k_1} \wedge T_{k_2})} \sum_{j=0}^{T_{k_1} \wedge T_{k_2}} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_{k_1})(\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})\{\mathbf{\Sigma}^{-1}(T_{k_2})\}^\top \mathbf{U} \right.$$

$$\left. - \frac{a_{k_1} a_{k_2}(T_{k_1} \wedge T_{k_2})}{\sqrt{T_{k_1} T_{k_2}} \sigma^2(T_{k_1} \wedge T_{k_2})} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_{k_1}) \mathbf{\Omega}(T_{k_1} \wedge T_{k_2})\{\mathbf{\Sigma}^{-1}(T_{k_2})\}^\top \mathbf{U} \right\|_2$$

$$\leq \frac{a_{k_1} a_{k_2}}{\sigma^2(T_{k_1} \wedge T_{k_2})} \|\mathbf{U}\|_2^2 \max_k \|\mathbf{\Sigma}^{-1}(T_k)\|_2^2 \left\| \frac{1}{T_{k_1} \wedge T_{k_2}} \sum_{j=0}^{T_{k_1} \wedge T_{k_2}-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1}) - \mathbf{\Omega}(t) \right|$$

$$\xrightarrow{P} 0,$$

where $a \wedge b = \min(a, b)$. It follows that

$$\left| \sum_{j=0}^{T-1} \left| \sum_{k=1}^{K} a_k T_k^{-1/2} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_k)(\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top \{\sigma(T_k)\}^{-1} \mathbb{I}(j < T_k) \right|^2 \right. \tag{28}$$

$$\left. - \sum_{k_1 \neq k_2} \frac{a_{k_1} a_{k_2}(T_{k_1} \wedge T_{k_2})}{\sqrt{T_{k_1} T_{k_2}} \sigma^2(T_{k_1} \wedge T_{k_2})} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_{k_1}) \mathbf{\Omega}(T_{k_1} \wedge T_{k_2})\{\mathbf{\Sigma}^{-1}(T_{k_2})\}^\top \mathbf{U} \right| = o_p(1).$$

In the proofs of Lemmas 3 and 4, we show that $\|\mathbf{\Sigma}^{-1}(t) - (\mathbf{\Sigma}^{(0)*})^{-1}\|_2 = O(t^{-1/2})$ and $\|\mathbf{\Omega}(t) - \mathbf{\Omega}^{(0)*}\|_2 = O(t^{-1/2})$ for some matrices $\mathbf{\Sigma}^{(0)*}$ and $\mathbf{\Omega}^{(0)*}$ that are invariant to t. Definitions of these two matrices can be found in Sections G.5 and G.6. Under C2(ii) and the condition that $q = o(\sqrt{T}/\log T)$, we can show $\|\mathbf{U}\|_2 = O(q^{1/2})$ and hence $\sigma^2(t) \xrightarrow{P} (\sigma^{(0)*})^2$ where $(\sigma^{(0)*})^2 = \mathbf{U}^\top (\mathbf{\Sigma}^{(0)*})^{-1} \mathbf{\Omega}^{(0)*}\{(\mathbf{\Sigma}^{(0)*})^{-1}\}^\top \mathbf{U}$. Similar to equation 28, we have

$$\sum_{k_1 \neq k_2} \frac{a_{k_1} a_{k_2}(T_{k_1} \wedge T_{k_2})}{\sqrt{T_{k_1} T_{k_2}} \sigma^2(T_{k_1} \wedge T_{k_2})} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_{k_1}) \mathbf{\Omega}(T_{k_1} \wedge T_{k_2})\{\mathbf{\Sigma}^{-1}(T_{k_2})\}^\top \mathbf{U} \tag{29}$$

$$\xrightarrow{P} \sum_{k_1 \neq k_2} \frac{a_{k_1} a_{k_2}(T_{k_1} \wedge T_{k_2})}{\sqrt{T_{k_1} T_{k_2}} (\sigma^{(0)*})^2} \mathbf{U}^\top \{\mathbf{\Sigma}^{(0)*}\}^{-1} \mathbf{\Omega}^{(0)*}\{(\mathbf{\Sigma}^{(0)*})^{-1}\}^\top \mathbf{U} \to \sum_{k_1 \neq k_2} \frac{a_{k_1} a_{k_2}(c_{k_1} \wedge c_{k_2})}{\sqrt{c_{k_1} c_{k_2}}},$$

where c_k's are defined in Section 4.2. This together with equation 28 yields that

$$\sum_{j=0}^{T-1} \left| \sum_{k=1}^{K} a_k T_k^{-1/2} \mathbf{U}^\top \mathbf{\Sigma}^{-1}(T_k)(\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top \{\sigma(T_k)\}^{-1} \mathbb{I}(j < T_k) \right|^2 \xrightarrow{P} \frac{a_{k_1} a_{k_2}(c_{k_1} \wedge c_{k_2})}{\sqrt{c_{k_1} c_{k_2}}}.$$

Conditions (a) and (b) are thus verified. By Lemma 4, we can show

$$\sum_{k=1}^{K} \frac{a_k \sqrt{T_k} \{\widehat{\tau}(T_k) - \tau_0\}}{\sigma(T_k)} = \sum_{k=1}^{K} \frac{a_k \sqrt{T_k} \{\widehat{\tau}(T_k) - \tau_0\}}{\widehat{\sigma}(T_k)} + o_p(1),$$

for any $(a_1, \cdots, a_K)$. This yields the joint asymptotic normality of our test statistics.

By equation 29, its covariance matrix is given by $\mathbf{\Xi}_0$ whose (k_1, k_2)-th entry is equal to $(c_{k_1} c_{k_2})^{-1/2} c_{k_1} \wedge c_{k_2}$. Using similar arguments in proving equation 28, equation 29 and Step 3 of the proof in Theorem 1, Shi et al. (2020b), we can show $\widehat{\mathbf{\Xi}}$ is a consistent estimator for $\mathbf{\Xi}_0$. This completes the proof of Theorem 1 under D1.

G.2.2 PROOF UNDER D2

The proof is very similar to that under D1. Suppose we can show equation 18, equation 21, equation 22 and equation 25 hold. Then similar to the proof under D1, we have

$$\frac{\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}}{\sigma(t)} = \frac{\sqrt{t}\boldsymbol{U}^\top \zeta_1(t)}{\sigma(t)} + o_p(1).$$

The following lemma shows these assertions hold under D2 as well.

Lemma 5 *Under the given conditions, we have equation 18, equation 21, equation 22 and equation 25 hold.*

It follows that for any K-dimensional vector $\boldsymbol{a} = (a_1, \cdots, a_K)^\top$,

$$\sum_{k=1}^{K} \frac{a_k \sqrt{T_k}\{\widehat{\tau}(T_k) - \tau_0\}}{\sigma(T_k)} = \sum_{k=1}^{K} \frac{a_k \sqrt{T_k}\boldsymbol{U}^\top \zeta_1(T_k)}{\sigma(T_k)} + o_p(1).$$

In the proof of Lemma 5, we show $\|\{\boldsymbol{\Sigma}(t)\}^{-1}\|_2 = O(1)$, $\|\boldsymbol{\Sigma}(t) - \boldsymbol{\Sigma}^*\|_2 = O(t^{-1/2})$ for some time-invariant matrix $\boldsymbol{\Sigma}^*$ that satisfies $\|(\boldsymbol{\Sigma}^*)^{-1}\|_2 = O(1)$. It follows that

$$\|\{\boldsymbol{\Sigma}(t)\}^{-1} - (\boldsymbol{\Sigma}^*)^{-1}\|_2 = \|\{\boldsymbol{\Sigma}(t)\}^{-1}(\boldsymbol{\Sigma}(t) - \boldsymbol{\Sigma}^*)(\boldsymbol{\Sigma}^*)^{-1}\|_2 \leq$$
$$\|\{\boldsymbol{\Sigma}(t)\}^{-1}\|_2 \|\boldsymbol{\Sigma}(t) - \boldsymbol{\Sigma}^*\|_2 \|(\boldsymbol{\Sigma}^*)^{-1}\|_2 = O(t^{-1/2}).$$

Similarly, we can show $\|\boldsymbol{\Omega}(t) - \boldsymbol{\Omega}^*\|_2 = O(t^{-1/2})$ for some matrix $\boldsymbol{\Omega}^*$.

In addition, using similar arguments in the proof of Lemma 5, we can show equation 27 holds under D2 as well. Now, the joint asymptotic normality of our test statistics follow using arguments from Part 3 of the proof under D1. Similarly, we can show $\widehat{\boldsymbol{\Xi}}$ is consistent. This completes the proof under D2.

G.2.3 PROOF UNDER D3

The proof under D1 indicates that equation 18, equation 21, equation 22 and equation 25 hold with $t = T_1$. It follows that

$$\frac{\sqrt{T_1}\{\widehat{\tau}(T_1) - \tau_0\}}{\sigma(T_1)} = \frac{\sqrt{T_1}\boldsymbol{U}^\top \zeta_1(T_1)}{\sigma(T_1)} + o_p(1). \tag{30}$$

The rest of the proof is divided into two parts. In the first part, we show for $k = 2, \cdots, K$,

$$\frac{\sqrt{T_k}\{\widehat{\tau}(T_k) - \tau_0\}}{\sigma^*(T_k)} = \frac{\sqrt{T_k}\boldsymbol{U}^\top \zeta_1^*(T_k)}{\sigma^*(T_k)} + o_p(1), \tag{31}$$

for some $\zeta_1^*(T_k)$ and $\sigma^*(T_k)$ defined below. In the second part, we show the assertion in Theorem 1 holds under D3.

Part 1: For any $1 \leq k \leq K$, consider the matrices

$$\boldsymbol{\Sigma}^{(k)} = \frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \mathrm{E}[\boldsymbol{\Sigma}_j | \{(S_t, A_t, Y_t)\}_{0 \leq t < T_{k-1}}] \quad \text{and} \quad \widehat{\boldsymbol{\Sigma}}^{(k)} = \frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \boldsymbol{\Sigma}_j.$$

We show in Lemma 6 below that for $k = 2, \cdots, K$,

$$\|\boldsymbol{\Sigma}^{(k)} - \widehat{\boldsymbol{\Sigma}}^{(k)}\|_2 = o_p(q^{-1/2}), \tag{32}$$

and

$$\|\{\overline{\boldsymbol{\Sigma}}^{(k)}\}^{-1}\|_2 = O_p(1). \tag{33}$$

where $\overline{\boldsymbol{\Sigma}}^{(k)} = T_k^{-1} \sum_{i=1}^{k} (T_i - T_{i-1}) \boldsymbol{\Sigma}^{(i)}$.

Lemma 6 *Under the given conditions, we have equation 32 and equation 33 hold.*

Notice that $(T_i - T_{i-1})/T_k \to (c_i - c_{i-1})/c_k$ and $\|\{c_k^{-1}\sum_{i=1}^{k}(c_i - c_{i-1})\boldsymbol{\Sigma}^{(i)}\}^{-1}\|_2 = O_p(1)$ where $c_0 = 0$. It follows from equation 32 that $\|c_k^{-1}\sum_{i=1}^{k}(c_i - c_{i-1})(\widehat{\boldsymbol{\Sigma}}^{(i)} - \boldsymbol{\Sigma}^{(i)})\|_2 = o_p(q^{-1/2})$ and hence

$$\left\| \frac{1}{T_k} \sum_{i=1}^{k} (T_i - T_{i-1})(\widehat{\boldsymbol{\Sigma}}^{(i)} - \boldsymbol{\Sigma}^{(i)}) \right\|_2 = o_p(q^{-1/2}), \quad \forall k = 2, \cdots, K.$$

Similar to the proof under D1, we can show

$$\left\| \left\{ \frac{1}{T_k} \sum_{i=1}^{k} (T_i - T_{i-1})\widehat{\boldsymbol{\Sigma}}^{(i)} \right\}^{-1} \right\|_2 = O_p(1) \quad \text{and} \quad \left\| \left\{ \frac{1}{T_k} \sum_{i=1}^{k} (T_i - T_{i-1})\widehat{\boldsymbol{\Sigma}}^{(i)} \right\}^{-1} - \{\overline{\boldsymbol{\Sigma}}^{(k)}\}^{-1} \right\|_2 = o_p(q^{-1/2}),$$

for $k = 2, \cdots, K$. Thus, equation 22 and the first assertion in equation 18 hold with $t = T_2, T_3, \cdots, T_K$ under D3.

In addition, similar to Lemma 6, we can show

$$\lambda_{\max} \left[\frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \mathrm{E}[\xi_j \xi_j^\top | \{(S_t, A_t, Y_t)\}_{0 \le t < T_{k-1}}] \right] = O_p(1), \tag{34}$$

and

$$\left\| \frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \left(\xi_j \xi_j^\top - \mathrm{E}[\xi_j \xi_j^\top | \{(S_t, A_t, Y_t)\}_{0 \le t < T_{k-1}}] \right) \right\|_2 = o_p(q^{-1/2}),$$

for $k = 2, \cdots, K$. This yields $\|(T_k - T_{k-1})^{-1} \sum_{j=T_{k-1}}^{T_k - 1} \xi_j \xi_j^\top \|_2 = O_p(1)$. As a result, the second assertion in equation 18 holds with $t = T_2, T_3, \cdots, T_K$.

Moreover, using similar arguments in showing $t^{-1} \sum_{j=0}^{t-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top = O_p(t^{-1/2}\sqrt{q})$ under D1, we have by equation 34 that $(T_k - T_{k-1})^{-1} \sum_{j=T_{k-1}}^{T_k - 1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top = O_p\{(T_k - T_{k-1})^{-1/2}\sqrt{q}\}$, for $k = 1, \cdots, K$. Under the given conditions on $\{T_k\}_k$, we obtain $t^{-1} \sum_{j=0}^{t-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top = O_p(t^{-1/2}\sqrt{q})$ for $t = T_2, T_3, \cdots, T_K$.

Based on these results, using similar arguments in Part 1 of the proof under D1, we can show

$$\sqrt{T_k}\{\widehat{\boldsymbol{\beta}}(T_k) - \boldsymbol{\beta}^*\} = \sqrt{T_k}\zeta_1^*(T_k) + o_p(1), \quad \forall k \in \{2, \cdots, K\}, \tag{35}$$

where

$$\zeta_1^*(T_k) = \frac{1}{T_k} \sum_{j=1}^{T_k} (\overline{\boldsymbol{\Sigma}}^{(k)})^{-1} \begin{pmatrix} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{pmatrix}.$$

For $1 \le k \le K$, define

$$\boldsymbol{\Omega}^{(k)} = \frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \mathrm{E}[(\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1}) | \{(S_t, A_t, Y_t)\}_{0 \le t < T_{k-1}}],$$

and $\overline{\boldsymbol{\Omega}}^{(k)} = T_k^{-1} \sum_{i=1}^{k}(T_i - T_{i-1})\boldsymbol{\Sigma}^{(i)}$. For any $2 \le k \le K$, we have $\lambda_{\min}(\overline{\boldsymbol{\Omega}}^{(k)}) \ge \lambda_{\min}(T_k^{-1}T_1\boldsymbol{\Omega}^{(1)})$. Since $T_k^{-1}T_1 \to c_k^{-1}c_1 > 0$ and $\lambda_{\min}(\boldsymbol{\Omega}^{(1)}) = \lambda_{\min}(\boldsymbol{\Omega}(T_1))$ is bounded away from zero, $\lambda_{\min}(\overline{\boldsymbol{\Omega}}^{(k)})$ is bounded away from zero for $k = 2, \cdots, K$ as well. Define

$$\{\sigma^*(T_k)\}^2 = \boldsymbol{U}^\top (\overline{\boldsymbol{\Sigma}}^{(k)})^{-1} \overline{\boldsymbol{\Omega}}^{(k)} \{(\overline{\boldsymbol{\Sigma}}^{(k)})^{-1}\}^\top \boldsymbol{U}.$$

It can be shown that $\sigma^*(T_k)/\|\boldsymbol{U}\|_2$ is bounded away from zero, for $k = 2, \cdots, K$. Using similar arguments in Part 2 of the proof under D1, we can show equation 31 holds. This completes the proof for Part 1.

Part 2: Let $\sigma^*(T_1) = \sigma(T_1)$. By equation 30 and equation 31, we have for any K-dimensional vector $\boldsymbol{a} = (a_1, \cdots, a_K)^\top$ that

$$\sum_{k=1}^{K} \frac{a_k \sqrt{T_k} \{\hat{\tau}(T_k) - \tau_0\}}{\sigma^*(T_k)} = \sum_{k=1}^{K} \frac{a_k \sqrt{T_k} \boldsymbol{U}^\top \zeta_1(T_k)}{\sigma^*(T_k)} + o_p(1). \tag{36}$$

In the following, we show the leading term on the RHS of equation 36 is asymptotically normal. Similar to the proof under D1, it suffices to verify the following conditions:

(a) $\max_{0 \le j < T} |\sum_{k=1}^{K} a_k T_k^{-1/2} \boldsymbol{U}^\top (\overline{\boldsymbol{\Sigma}}^{(k)})^{-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top \mathbb{I}(j < T_k)| \xrightarrow{P} 0$;

(b) $\sum_{j=0}^{T-1} |\sum_{k=1}^{K} a_k T_k^{-1/2} \boldsymbol{U}^\top (\overline{\boldsymbol{\Sigma}}^{(k)})^{-1} (\xi_j^\top \varepsilon_{j,0}, \xi_j^\top \varepsilon_{j,1})^\top \{\sigma^*(T_k)\}^{-1} \mathbb{I}(j < T_k)|^2$ converges to some constant in probability.

Condition (a) can be proven in a similar manner as in Part 3 of the proof under D1. Notice that for $k = 2, \cdots, K$, $\overline{\boldsymbol{\Sigma}}^{(k)}$, $\overline{\boldsymbol{\Omega}}^{(k)}$ and $\sigma^*(T_k)$ are random variables and depend on the observed data history. In the proof of Lemma 6, we show $\|(\overline{\boldsymbol{\Sigma}}^{(k)})^{-1} - (\boldsymbol{\Sigma}^{**})^{-1}\|_2 = O_p(T^{-1/2})$ for some deterministic matrix $\boldsymbol{\Sigma}^*$ and all $k \in \{2, \cdots, K\}$. Similarly, we can show $\|\overline{\boldsymbol{\Omega}}^{(k)} - \boldsymbol{\Omega}^{**}\|_2 = O_p(T^{-1/2})$ and $\|\{\sigma^*(T_k)\}^2 - (\sigma^{**})^2\|_2 = O_p(T^{-1/2})$ for some $\boldsymbol{\Sigma}^*, \sigma^{**}$ and all $k \in \{2, \cdots, K\}$. Moreover, using similar arguments in the proof of Lemma 6, we can show

$$\left\| \frac{1}{T_k - T_{k-1}} \sum_{j=T_{k-1}}^{T_k - 1} \begin{pmatrix} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{pmatrix} \begin{pmatrix} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{pmatrix}^\top - \boldsymbol{\Omega}^{(k)} \right\|_2 = o_p(q^{-1/2}), \quad \forall k = 2, \cdots, K.$$

This further implies that

$$\left\| \frac{1}{T_k} \sum_{j=0}^{T_k - 1} \begin{pmatrix} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{pmatrix} \begin{pmatrix} \xi_j \varepsilon_{j,0} \\ \xi_j \varepsilon_{j,1} \end{pmatrix}^\top - \boldsymbol{\Omega}^{**} \right\|_2 = o_p(q^{-1/2}), \quad \forall k = 2, \cdots, K.$$

Based on these results, using similar arguments in Part 3 of the proof of Lemma 3, we obtain (b). The joint asymptotic normality of $\sqrt{T_1}\{\hat{\tau}(T_1) - \tau_0\}/\sigma^*(T_1), \cdots, \sqrt{T_1}\{\hat{\tau}(T_1) - \tau_0\}/\sigma^*(T_K)$ thus follows.

Consistency of $\widehat{\boldsymbol{\Xi}}$ can be similarly proven. We omit the details for brevity.

G.3 PROOF OF THEOREM 2

As discussed in Section 4.3, $(Z_1^*, Z_2^*, \cdots, Z_K^*)^\top$ is jointly normal with mean zero and covariance matrix $\widehat{\boldsymbol{\Xi}}$, conditional on the observed data. By Theorem 1, we have $\widehat{\boldsymbol{\Xi}} \xrightarrow{P} \boldsymbol{\Xi}_0$ where $\boldsymbol{\Xi}_0$ is the asymptotic covariance matrix of $(Z_1, Z_2, \cdots, Z_K)^\top$. Let $\alpha^*(t) = \alpha(tT)$ for any $0 \le t \le 1$, we have $\alpha(T_k) \to \alpha^*(c_k)$ for any $1 \le k \le K$. Notice that $\{\hat{b}_k\}_{1 \le k \le K}$ is a continuous function of $\widehat{\boldsymbol{\Xi}}$ and $\{\alpha(T_k)\}_{1 \le k \le K}$, it follows that $\hat{b}_k \xrightarrow{P} b_{k,0}$ for $1 \le k \le K$, where $\{b_{k,0}\}_{1 \le k \le K}$ are recursively defined as follows:

$$\Pr\left\{ \max_{1 \le j < k} (Z_{j,0} - b_{j,0}) \le 0, Z_{k,0} > b_{k,0} \right\} = \alpha^*(c_k) - \alpha^*(c_{k-1}),$$

where $(Z_{1,0}, Z_{2,0}, \cdots, Z_{K,0})^\top$ is asymptotically normal with mean zero and covariance matrix $\boldsymbol{\Xi}_0$.

Theorem 1 implies that $(Z_1 - \sqrt{T_1}\tau_0/\hat{\sigma}(T_1), Z_2 - \sqrt{T_2}\tau_0/\hat{\sigma}(T_2), \cdots, Z_K - \sqrt{T_K}\tau_0/\hat{\sigma}(T_K))^\top \xrightarrow{d} (Z_{1,0}, Z_{2,0}, \cdots, Z_{K,0})^\top$. It follows that

$$\Pr\left(\bigcup_{j=1}^{k} \{Z_j > \hat{b}_j\} \right) \le \Pr\left(\bigcup_{j=1}^{k} \{Z_j - \sqrt{T_j}\tau_0/\hat{\sigma}(T_j) > \hat{b}_j\} \right) \to \Pr\left(\bigcup_{j=1}^{k} \{Z_{j,0} > b_{j,0}\} \right) = \alpha^*(c_k). \tag{37}$$

The proof is hence completed by noting that $\alpha(T_k) \to \alpha^*(c_k)$. When $\tau_0 = 0$, we have $\mathbb{E}Z_k = o(1)$. The rejection probability thus converges to the nominal level.

G.4 PROOF OF THEOREM 3

Suppose $\tau_0 = T^{-1/2}h$ for some $h > 0$. Based on the proof of Theorem 1, we can show $\widehat{\sigma}(T_k) \xrightarrow{P} \sigma_k^*$ for some $\sigma_k^* > 0$. It follows from equation 37 that

$$\Pr\left(\bigcup_{j=1}^{k}\{Z_j > \widehat{b}_j\}\right) = \Pr\left(\bigcup_{j=1}^{k}\{Z_j - \sqrt{T_j}\tau_0/\widehat{\sigma}(T_j) > \widehat{b}_j - h/\widehat{\sigma}(T_j)\}\right)$$

$$\rightarrow \Pr\left(\bigcup_{j=1}^{k}\{Z_{j,0} > b_{j,0} - h/\sigma_j^*\}\right) > \alpha^*(c_k).$$

The second assertion in Theorem 3 thus holds by noting that $\alpha(T_k) \rightarrow \alpha^*(c_k)$.

Let $h \rightarrow \infty$, we obtain

$$\Pr\left(\bigcup_{j=1}^{k}\{Z_j > \widehat{b}_j\}\right) = \Pr\left(\bigcup_{j=1}^{k}\{Z_{j,0} > b_{j,0} - h/\sigma_j^*\}\right) + o(1) \rightarrow 1.$$

The proof is hence completed.

G.5 PROOF OF LEMMA 3

Under the given conditions in C1(i), C1(ii) and C2(ii), equation 21 and the second assertion in equation 18 can be proven using similar arguments in the proof of Lemma E.2 and E.3 of Shi et al. (2020b). We omit the proof for brevity.

It remains to show equation 22 and the first assertion in equation 18 hold. Recall that μ is the density function of the stationary distribution Π (see the remark below Condition C1). In addition, μ is uniformly bounded away from 0 and ∞ under C1(i). For $a' \in \{0, 1\}$, define the matrix

$$\boldsymbol{\Sigma}^{(0)*}(a') = \int_{s,s' \in \mathbb{S}} \sum_{a \in \{0,1\}} \xi(s,a)\{\xi(s,a) - \gamma\xi(s',a')\}\mu(s)\{(1-a)b^{(0)}(s) + a(1 - b^{(0)}(s))\}p(s';a,s)dsds'.$$

Define

$$\boldsymbol{\Sigma}^{(0)*} = \left[\begin{array}{cc} \boldsymbol{\Sigma}^{(0)*}(0) & \\ & \boldsymbol{\Sigma}^{(0)*}(1) \end{array}\right].$$

The matrix $\boldsymbol{\Sigma}^{(0)*}$ is the population limit of $\widehat{\boldsymbol{\Sigma}}(t)$ under D1. To prove the first assertion in equation 18, we first show

$$\|(\boldsymbol{\Sigma}^{(0)*})^{-1}\|_2 = O(1). \tag{38}$$

By definition, this is equivalent to show

$$\|\{\boldsymbol{\Sigma}^{(0)*}(a)\}^{-1}\|_2 = O(1),$$

for $a \in \{0, 1\}$. The matrix $\boldsymbol{\Sigma}^{(0)*}(0)$ can be written as

$$\boldsymbol{\Sigma}^{(0)*}(0) = \left[\begin{array}{cc} \boldsymbol{\Sigma}_{1,1}^{(0)*}(0) & \\ \boldsymbol{\Sigma}_{2,1}^{(0)*}(0) & \boldsymbol{\Sigma}_{2,2}^{(0)*}(0) \end{array}\right],$$

where

$$\boldsymbol{\Sigma}_{1,1}^{(0)*}(0) = \int_{s,s' \in \mathbb{S}} \Psi(s)\{\Psi(s) - \gamma\Psi(s')\}^\top b^{(0)}(s)\mu(s)p(s';0,s)dsds',$$

$$\boldsymbol{\Sigma}_{2,1}^{(0)*}(0) = -\gamma\int_{s,s' \in \mathbb{S}} \Psi(s)\Psi^\top(s')(1 - b^{(0)}(s))\mu(s)p(s';0,s)dsds',$$

$$\boldsymbol{\Sigma}_{2,2}^{(0)*}(0) = \int_{s \in \mathbb{S}} \Psi(s)\Psi^\top(s)\mu(s)(1 - b^{(0)}(s))p(s';1,s)ds.$$

It follows that

$$\{\boldsymbol{\Sigma}^{(0)*}(0)\}^{-1} = \begin{bmatrix} \{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1} & \\ -\{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1} & \{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1} \end{bmatrix},$$

and hence

$$\|\{\boldsymbol{\Sigma}^{(0)*}(0)\}^{-1}\|_2 = \sup_{\|\boldsymbol{a}_1\|_2=1,\|\boldsymbol{a}_2\|_2=1} \left| \boldsymbol{a}_1^\top \begin{bmatrix} \{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1} & \\ -\{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1} & \{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1} \end{bmatrix} \boldsymbol{a}_2 \right|$$

$$\leq \sup_{\|\boldsymbol{a}_3\|_2=1,\|\boldsymbol{a}_4\|_2=1} |\boldsymbol{a}_3^\top \{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1}\boldsymbol{a}_4| + \sup_{\|\boldsymbol{a}_3\|_2=1,\|\boldsymbol{a}_4\|_2=1} |\boldsymbol{a}_3^\top \{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\boldsymbol{a}_4|$$

$$+ \sup_{\|\boldsymbol{a}_3\|_2=1,\|\boldsymbol{a}_4\|_2=1} |\boldsymbol{a}_3^\top \{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1}\boldsymbol{a}_4|$$

$$\leq \|\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1}\|_2 + \|\{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\|_2 + \|\{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\|_2\|\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\|_2\|\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1}\|_2.$$

Thus, to prove $\|\{\boldsymbol{\Sigma}^{(0)*}(0)\}^{-1}\|_2 = O(1)$, it suffices to show

$$\|\{\boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\}^{-1}\|_2 = O(1), \tag{39}$$

$$\|\{\boldsymbol{\Sigma}_{2,2}^{(0)*}(0)\}^{-1}\|_2 = O(1), \tag{40}$$

$$\|\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\|_2 = O(1). \tag{41}$$

We first consider equation 39. Using similar arguments in Part 1 of the proof of Lemma E.2, Shi et al. (2020b), it suffices to show

$$\boldsymbol{a}^\top \boldsymbol{\Sigma}_{1,1}^{(0)*}(0)\boldsymbol{a} \geq \bar{c}_1\|\boldsymbol{a}\|_2^2, \quad \forall \boldsymbol{a},$$

for some $\bar{c}_1 > 0$. Under D1, $b^{(0)}$ is strictly positive. Since μ is strictly positive, it suffices to show

$$\boldsymbol{a}^\top \int_{s,s'\in\mathbb{S}} \Psi(s)\{\Psi(s) - \gamma\Psi(s')\}^\top dsds'\boldsymbol{a} \geq \bar{c}_2\|\boldsymbol{a}\|_2^2, \quad \forall \boldsymbol{a}, \tag{42}$$

for some $\bar{c}_2 > 0$. Notice that LHS of equation 42 is equal to

$$\lambda(\mathbb{S}) \int_{s\in\mathbb{S}} \{\boldsymbol{a}^\top\Psi(s)\}^2 ds - \gamma \int_{s,s'\in\mathbb{S}} \{\boldsymbol{a}^\top\Psi(s)\}\{\boldsymbol{a}^\top\Psi(s')\}dsds',$$

where $\lambda(\mathbb{S})$ is the Lebesgue measure of $\mathbb{S}$. Since $\mathbb{S}$ is compact, we have $\lambda(\mathbb{S}) < +\infty$. By Cauchy-Schwarz inequality, LHS of equation 42 is greater than or equal to

$$\lambda(\mathbb{S}) \int_{s\in\mathbb{S}} \{\boldsymbol{a}^\top\Psi(s)\}^2 ds - \lambda(\mathbb{S}) \int_{s\in\mathbb{S}} \frac{\gamma}{2}\{\boldsymbol{a}^\top\Psi(s)\}^2 ds - \lambda(\mathbb{S}) \int_{s\in\mathbb{S}} \frac{\gamma}{2}\{\boldsymbol{a}^\top\Psi(s')\}ds'$$

$$\geq (1-\gamma)\lambda(\mathbb{S}) \int_{s\in\mathbb{S}} \{\boldsymbol{a}^\top\Psi(s)\}^2 ds.$$

This is directly implied by Condition C2(ii). The proof of equation 39 is hence completed. Similarly, we can prove equation 40. In addition, notice that

$$\|\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\|_2 \leq \sup_{\|\boldsymbol{a}_1\|_2=1,\|\boldsymbol{a}_2\|_2=1} \int_{s,s'\in\mathbb{S}} |\boldsymbol{a}_1^\top\Psi(s)||\boldsymbol{a}_2^\top\Psi(s')|\{1 - b^{(0)}(s)\mu(s)\}p(s';0,s)dsds'$$

$$\leq \sup_{\|\boldsymbol{a}_1\|_2=1,\|\boldsymbol{a}_2\|_2=1} \int_{s,s'\in\mathbb{S}} |\boldsymbol{a}_1^\top\Psi(s)||\boldsymbol{a}_2^\top\Psi(s')|\mu(s)p(s';0,s)dsds'.$$

Since the density function μ is uniformly bounded, we have

$$\|\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\|_2 \leq O(1) \sup_{\|\boldsymbol{a}_1\|_2=1,\|\boldsymbol{a}_2\|_2=1} \int_{s,s'\in\mathbb{S}} |\boldsymbol{a}_1^\top\Psi(s)||\boldsymbol{a}_2^\top\Psi(s')|dsds',$$

where $O(1)$ denotes the universal constant. By Cauchy-Schwarz inequality, we obtain

$$\|\boldsymbol{\Sigma}_{2,1}^{(0)*}(0)\|_2 \leq O(1)\lambda(\mathbb{S}) \sup_{\|\boldsymbol{a}\|_2=1} \int_{s\in\mathbb{S}} |\boldsymbol{a}^\top\Psi(s)|^2 ds \leq O(1)\lambda(\mathbb{S})\lambda_{\max}\left[\int_{s\in\mathbb{S}} \Psi(s)\Psi^\top(s)ds\right].$$

In view of C2(ii), we obtain equation 41.

To summarize, we have shown $\|\{\boldsymbol{\Sigma}^{(0)*}(0)\}^{-1}\|_2 = O(1)$. Similarly, we can prove $\|\{\boldsymbol{\Sigma}^{(0)*}(1)\}^{-1}\|_2 = O(1)$. Assertion equation 38 thus holds. Similar to Lemma E.5 of Shi et al. (2020b), we can show $\|\boldsymbol{\Sigma}(t) - \boldsymbol{\Sigma}^{(0)*}\|_2 = O(t^{-1/2})$. Using similar arguments in Part 1 of the proof of Lemma E.2, Shi et al. (2020b), this yields $\|\boldsymbol{\Sigma}^{-1}(t) - (\boldsymbol{\Sigma}^{(0)*})^{-1}\|_2 = o(t^{-1/2})$ and $\|\boldsymbol{\Sigma}^{-1}(t)\|_2 = O(1)$. Under the given conditions, equation 22 and the first assertion in equation 18 now follow from the arguments used in Part 2 and 3 of the proof of Lemma E.2, Shi et al. (2020b).

G.6 PROOF OF LEMMA 4

The asymptotic normality of $\sqrt{t}\{\widehat{\tau}(t) - \tau_0\}/\sigma(t)$ can be proven using similar arguments in Part 3 of the proof of Theorem 1. In the following, we focus on equation 25. Define the matrix

$$\boldsymbol{\Omega}^{(0)*} = \int_{s\in\mathbb{S}} \sum_{a\in\{0,1\}} \mathrm{E}\left\{ \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right) \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right)^\top \middle| S_0 = s, A_0 = a \right\} \mu(s)\{ab^{(0)}(s) + (1-a)(1-b^{(0)}(s))\}ds.$$

Similar to Lemma E.5 of Shi et al. (2020b), we can show $\|\boldsymbol{\Omega}^{(0)*} - \boldsymbol{\Omega}(t)\|_2 = O(t^{-1/2})$. Thus, it suffices to show $\inf_q \lambda_{\min}(\boldsymbol{\Omega}^{(0)*}) > 0$. Under CA and SRA, we have

$$\mathrm{E}\left\{ \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right) \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right)^\top \middle| S_0 = s, A_0 = a \right\}$$

$$= \mathrm{E}\left\{ \left(\begin{array}{c} \xi_0(a)\varepsilon^*(0,a) \\ \xi_0(a)\varepsilon^*(1,a) \end{array} \right) \left(\begin{array}{c} \xi_0(a)\varepsilon^*(0,a) \\ \xi_0(a)\varepsilon^*(1,a) \end{array} \right)^\top \middle| S_0 = s, A_0 = a \right\}$$

$$= \mathrm{E}\left\{ \left(\begin{array}{c} \xi_0(a)\varepsilon^*(0,a) \\ \xi_0(a)\varepsilon^*(1,a) \end{array} \right) \left(\begin{array}{c} \xi_0(a)\varepsilon^*(0,a) \\ \xi_0(a)\varepsilon^*(1,a) \end{array} \right)^\top \middle| S_0 = s \right\}$$

For any $2q$-dimensional vectors $\boldsymbol{a}_1, \boldsymbol{a}_2$ that satisfy $\|\boldsymbol{a}_1\|_2^2 + \|\boldsymbol{a}_2\|^2 = 1$, it follows that

$$(\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)\mathrm{E}\left\{ \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right) \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right)^\top \middle| S_0 = s, A_0 = a \right\}(\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)^\top$$

$$= \{\boldsymbol{a}_1^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(0,a)\}^2|S_0 = s] + \{\boldsymbol{a}_2^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s]$$

$$+ 2\{\boldsymbol{a}_1^\top\xi(s,a)\}\{\boldsymbol{a}_2^\top\xi(s,a)\}\mathrm{E}[\{\varepsilon^*(0,a)\varepsilon^*(1,a)\}|S_0 = s]$$

$$\geq \{\boldsymbol{a}_1^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(0,a)\}^2|S_0 = s] + \{\boldsymbol{a}_2^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s]$$

$$- 2\rho_\varepsilon|\boldsymbol{a}_1^\top\xi(s,a)\|\boldsymbol{a}_2^\top\xi(s,a)|\sqrt{\mathrm{E}[\{\varepsilon^*(0,a)\}^2|S_0 = s]\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s]}$$

$$= (1-\rho_\varepsilon)\{\boldsymbol{a}_1^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(0,a)\}^2|S_0 = s] + (1-\rho_\varepsilon)\{\boldsymbol{a}_2^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s]$$

$$+ \rho_\varepsilon\left| |\boldsymbol{a}_1^\top\xi(s,a)|\sqrt{\mathrm{E}\{\varepsilon^*(0,a)\}^2|S_0 = s\}} + |\boldsymbol{a}_2^\top\xi(s,a)|\sqrt{\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s]} \right|^2$$

$$\geq (1-\rho_\varepsilon)\{\boldsymbol{a}_1^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(0,a)\}^2|S_0 = s] + (1-\rho_\varepsilon)\{\boldsymbol{a}_2^\top\xi(s,a)\}^2\mathrm{E}[\{\varepsilon^*(1,a)\}^2|S_0 = s],$$

where $\rho_\varepsilon = \sup_q \sup_{a\in\{0,1\},s\in\mathbb{S}} \rho_\varepsilon(a,s)$. Under C3, we have $\rho_\varepsilon < 1$ and $\inf_q \inf_{a',a,s} \mathrm{E}[\{\varepsilon^*(a',a)\}^2|S_0 = s] > 0$. It follows that

$$(\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)\mathrm{E}\left\{ \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right) \left(\begin{array}{c} \xi_0\varepsilon_{0,0} \\ \xi_0\varepsilon_{0,1} \end{array} \right)^\top \middle| S_0 = s, A_0 = a \right\}(\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)^\top \geq \bar{c}[\{\boldsymbol{a}_1^\top\xi(s,a)\}^2 + \{\boldsymbol{a}_2^\top\xi(s,a)\}^2],$$

for some constant $\bar{c}_3 > 0$. Therefore,

$$\lambda_{\min}(\boldsymbol{\Omega}^{(0)*}) = \inf_{\|\boldsymbol{a}_1\|_2^2 + \|\boldsymbol{a}_2\|_2^2 = 1} (\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)\boldsymbol{\Omega}^{(0)*}(\boldsymbol{a}_1^\top, \boldsymbol{a}_2^\top)^\top$$

$$\geq \bar{c}_3 \inf_{\|\boldsymbol{a}_1\|_2^2 + \|\boldsymbol{a}_2\|_2^2 = 1} \int_{s\in\mathbb{S}} \sum_{a\in\{0,1\}} [\{\boldsymbol{a}_1^\top\xi(s,a)\}^2 + \{\boldsymbol{a}_2^\top\xi(s,a)\}^2]\mu(s)\{ab^{(0)}(s) + (1-a)(1-b^{(0)}(s))\}ds.$$

The strict positivity of $\mu(\cdot)$ and the condition that $b^{(0)}(\cdot)$ is uniformly bounded away from 0 and 1 yields

$$\lambda_{\min}(\boldsymbol{\Omega}^{(0)*}) \geq \bar{c}_4 \inf_{\|\boldsymbol{a}_1\|_2^2 + \|\boldsymbol{a}_2\|_2^2 = 1} \int_{s \in \mathbb{S}} \sum_{a \in \{0,1\}} [\{\boldsymbol{a}_1^\top \xi(s,a)\}^2 + \{\boldsymbol{a}_2^\top \xi(s,a)\}^2] ds, \qquad (43)$$

for some constant $\bar{c}_4 > 0$. With some calculation, we can show the RHS of equation 43 is equal to

$$\bar{c}_4 \lambda_{\min} \left\{ \int_{s \in \mathbb{S}} \Psi(s) \Psi^\top(s) \right\}.$$

By Condition C2(ii), it is strictly positive. This yields $\inf_q \lambda_{\min}(\boldsymbol{\Omega}^{(0)*}) > 0$. Thus, we obtain equation 25.

G.7 PROOF OF LEMMA 5

We begin by proving

$$\|\boldsymbol{\Sigma}^{-1}(t)\|_2 = O(1), \qquad (44)$$

under D2. For any matrices $\boldsymbol{M}_1$ and $\boldsymbol{M}_2$, denote by $\text{diag}[\boldsymbol{M}_1, \boldsymbol{M}_2]$ the block diagonal matrix

$$\begin{bmatrix} \boldsymbol{M}_1 & \\ & \boldsymbol{M}_2 \end{bmatrix}.$$

By MA and Condition C1(ii), the two Markov chains $\{S_{2t-1}\}_{t \geq 1}$, $\{S_{2t}\}_{t \geq 0}$ are geometrically ergodic. Let μ_1 and μ_2 denote the density function of their stationary distributions, respectively. Under C1(i), we can similarly show that they are uniformly bounded away from 0 and ∞. Define

$$\boldsymbol{\Sigma}_1^* = \int_{s,s' \in \mathbb{S}} \text{diag}[\xi(s,1)\{\xi(s,1) - \gamma\xi(s',0)\}^\top, \xi(s,1)\{\xi(s,1) - \gamma\xi(s',1)\}^\top] \mu_1(s) p(s';1,s) ds ds',$$

$$\boldsymbol{\Sigma}_2^* = \int_{s,s' \in \mathbb{S}} \text{diag}[\xi(s,0)\{\xi(s,0) - \gamma\xi(s',0)\}^\top, \xi(s,0)\{\xi(s,0) - \gamma\xi(s',1)\}^\top] \mu_2(s) p(s';0,s) ds ds'.$$

The matrix $(\boldsymbol{\Sigma}_1^* + \boldsymbol{\Sigma}_2^*)/2$ corresponds to the population limit of $\boldsymbol{\Sigma}(t)$. Similar to Lemma E.5 of Shi et al. (2020b), we can show $\|\boldsymbol{\Sigma}_1^* - (2t)^{-1} \sum_{j=0}^t \mathbb{E}\boldsymbol{\Sigma}_{2j+1}\|_2 = o(t^{-1/2})$ and $\|\boldsymbol{\Sigma}_2^* - (2t)^{-1} \sum_{j=0}^t \mathbb{E}\boldsymbol{\Sigma}_{2j}\|_2 = o(t^{-1/2})$. This further yields

$$\left\| \frac{\boldsymbol{\Sigma}_1^* + \boldsymbol{\Sigma}_2^*}{2} - \boldsymbol{\Sigma}(t) \right\|_2 = o(t^{-1/2}).$$

Similar to the proof of Lemma 3, in order to show equation 44, it suffices to show

$$\|(\boldsymbol{\Sigma}_1^* + \boldsymbol{\Sigma}_2^*)^{-1}\|_2 = O(1). \qquad (45)$$

Notice that $\boldsymbol{\Sigma}_1^* + \boldsymbol{\Sigma}_2^* = \text{diag}[\boldsymbol{\Sigma}^*(0), \boldsymbol{\Sigma}^*(1)]$ where

$$\boldsymbol{\Sigma}^*(a) = \int_{s,s' \in \mathbb{S}} [\xi(s,0)\{\xi(s,0) - \gamma\xi(s',a)\} \mu_2(s) p(s';0,s) + \xi(s,1)\{\xi(s,1) - \gamma\xi(s',a)\} \mu_1(s) p(s';1,s)] \, ds ds'.$$

The matrix $\boldsymbol{\Sigma}^*(0)$ can be further decomposed into

$$\boldsymbol{\Sigma}^*(0) = \begin{bmatrix} \boldsymbol{\Sigma}_{1,1}^*(0) & \\ \boldsymbol{\Sigma}_{2,1}^*(0) & \boldsymbol{\Sigma}_{2,2}^*(0) \end{bmatrix},$$

where

$$\boldsymbol{\Sigma}_{1,1}^*(0) = \int_{s,s' \in \mathbb{S}} \Psi(s)\{\Psi(s) - \gamma\Psi(s')\}^\top \mu_2(s) p(s';0,s) ds ds',$$

$$\boldsymbol{\Sigma}_{2,1}^*(0) = -\gamma \int_{s,s' \in \mathbb{S}} \Psi(s)\Psi^\top(s') \mu_1(s) p(s';1,s) ds ds',$$

$$\boldsymbol{\Sigma}_{2,2}^*(0) = \int_s \Psi(s)\Psi^\top(s) \mu_1(s) ds.$$

Similar to the proof of Lemma 3, we can show $\|\{\mathbf{\Sigma}_{1,1}^*(0)\}^{-1}\|_2 = O(1)$, $\|\{\mathbf{\Sigma}_{2,2}^*(0)\}^{-1}\|_2 = O(1)$ and $\|\mathbf{\Sigma}_{2,1}^*(0)\|_2 = O(1)$. It follows that $\|\{\mathbf{\Sigma}^*(0)\}^{-1}\|_2 = O(1)$. Similarly, we can show $\|\{\mathbf{\Sigma}^*(1)\}^{-1}\|_2 = O(1)$. This proves equation 45. Thus, we obtain equation 44.

Using similar arguments in Part 2 of the proof of Lemma E.2, Shi et al. (2020b), we can show $\|t^{-1}\sum_{j=0}^{t-1}\mathbf{\Sigma}_{2j} - \mathbf{\Sigma}_2^*\|_2 = O_p(t^{-1/2}\log t)$ and $\|t^{-1}\sum_{j=0}^{t-1}\mathbf{\Sigma}_{2j+1} - \mathbf{\Sigma}_1^*\|_2 = O_p(t^{-1/2}\log t)$. This further implies $\|\widehat{\mathbf{\Sigma}}(t) - (\mathbf{\Sigma}_1^* + \mathbf{\Sigma}_2^*)/2\| = O_p(t^{-1/2}\log t)$ and hence $\|\widehat{\mathbf{\Sigma}}(t) - \mathbf{\Sigma}(t)\|_2 = O_p(t^{-1/2}\log t)$. Combining these results together with equation 44, we can show equation 22 and the first assertion in equation 18 hold. equation 21 and the second assertion in equation 18 hold can be proven in a similar manner.

Finally, using similar arguments in the proof of Lemma 4, we can show equation 25 holds. We omit the details to save space.

G.8 PROOF OF LEMMA 6

Under C1(iv), we have equation 7 holds. Similar to equation 8, we can show $\Pi^{(k)}$ has a probability density function $\mu^{(k)}$ given by

$$\mu^{(k)}(s') = \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} [a\{1 - b^{(k)}(s)\} + (1-a)b^{(k)}(s)]p(s';a,s)\Pi^{(k)}(ds). \tag{46}$$

For $a' \in \{0,1\}$, define

$$\mathbf{\Sigma}^{(k)*}(a) = \int_{s,s' \in \mathbb{S}} \sum_{a \in \{0,1\}} \xi(s,a)\{\xi(s,a) - \gamma\xi(s',a')\}^\top \mu^{(k)}(s)\{a\{1 - b^{(k)}(s)\} + (1-a)b^{(k)}(s)\}p(s';a,s)dsds'.$$

Condition on $\{(S_j, A_j, Y_j)\}_{1 \le j < T_{k-1}}$, the matrix $\mathbf{\Sigma}^{(k)*}(a)$ is deterministic. Let $\mathbf{\Sigma}^{(k)} = \mathrm{diag}[\mathbf{\Sigma}^{(k)*}(0), \mathbf{\Sigma}^{(k)*}(1)]$. Similar to the proof of Lemma 3, we can show $\|\mathbf{\Sigma}^{(k)*} - \mathbf{\Sigma}^{(k)}\|_2 = o(1)$, conditional on $\{(S_j, A_j, Y_j)\}_{1 \le j < T_{k-1}}$, with probability tending to 1. This implies for any sufficiently small $\epsilon > 0$,

$$\Pr(\|\mathbf{\Sigma}^{(k)*} - \mathbf{\Sigma}^{(k)}\|_2 > \epsilon | \{(S_j, A_j, Y_j)\}_{1 \le j < T_{k-1}}) \xrightarrow{P} 0.$$

The above conditional probability is bounded between 0 and 1. Using bounded convergence theorem, we have

$$\Pr(\|\mathbf{\Sigma}^{(k)*} - \mathbf{\Sigma}^{(k)}\|_2 > \epsilon) = o(1), \tag{47}$$

and hence $\|\mathbf{\Sigma}^{(k)*} - \mathbf{\Sigma}^{(k)}\|_2 = o_p(1)$.

Notice that $\sup_s |b^{(k)}(s) - b^*(s)| \xrightarrow{P} 0$ and $\|\Pi^{(k)} - \Pi^*\|_{TV} \xrightarrow{P} 0$. Define

$$\mu^*(s) = \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} [a\{1 - b^*(s)\} + (1-a)b^*(s)]p(s';a,s)\Pi^*(ds).$$

It follows that

$$|\mu^{(k)}(s') - \mu^*(s')| \le \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} \{a|b^*(s) - b^{(k)}(s)| + (1-a)|b^*(s) - b^{(k)}(s)|\}p(s';a,s)\Pi^{(k)}(ds)$$

$$+ \sum_{a \in \{0,1\}} \int_{s \in \mathbb{S}} [a\{1 - b^*(s)\} + (1-a)b^*(s)]p(s';a,s)|\Pi^{(k)}(ds) - \Pi^*(ds)|,$$

and hence $\sup_s |\mu^{(k)}(s) - \mu^*(s)| \xrightarrow{P} 0$. With some calculations, we can show $\|\mathbf{\Sigma}^{(k)*}(a) - \mathbf{\Sigma}^*(a)\|_2 \xrightarrow{P} 0$ where

$$\mathbf{\Sigma}^*(a) = \int_{s,s' \in \mathbb{S}} \sum_{a \in \{0,1\}} \xi(s,a)\{\xi(s,a) - \gamma\xi(s',a')\}^\top \mu^*(s)\{a\{1 - b^*(s)\} + (1-a)b^*(s)\}p(s';a,s)dsds'.$$

(a) The proposed test under H_1 and H_0 (from left plots to right plots)

(b) Two-sample t-test under H_1 and H_0 (from left plots to right plots)

Figure 4: Empirical rejection probabilities of our test and the two-sample t-test with $\alpha(\cdot) = \alpha_2(\cdot)$. Settings correspond to the alternating-time-interval, adaptive and Markov design, from top plots to bottom plots.

Let $\boldsymbol{\Sigma}^* = \mathrm{diag}[\boldsymbol{\Sigma}^*(0), \boldsymbol{\Sigma}^*(1)]$, we obtain $\|\boldsymbol{\Sigma}^{(k)*} - \boldsymbol{\Sigma}^*\|_2 = o_p(1)$. Combining this together with equation 47, we obtain $\|\boldsymbol{\Sigma}^{(k)} - \boldsymbol{\Sigma}^*\|_2 = o_p(1)$. The proof of Lemma 3 yields $\|\boldsymbol{\Sigma}^{(1)} - \boldsymbol{\Sigma}^{(0)*}\|_2 = o(1)$. Thus, we have for any $2 \le k \le K$ that

$$\|\overline{\boldsymbol{\Sigma}}^{(k)} - T_k^{-1} T_1 \boldsymbol{\Sigma}^{(0)*} - T_k^{-1}(T_k - T_1)\boldsymbol{\Sigma}^*\|_2 = o_p(1). \tag{48}$$

Similar to the proof of Lemma 3, we can show $\mu^{(k)}$'s are uniformly bounded away from 0 and ∞. It follows that μ^* is uniformly bounded away from 0 and ∞. Using similar arguments in Lemma 3, we can show $\|\{T_k^{-1} T_1 \boldsymbol{\Sigma}^{(0)*} + T_k^{-1}(T_k - T_1)\boldsymbol{\Sigma}^*\}^{-1}\|_2 = O(1)$. Using similar arguments in Part 1 of the proof of Lemma E.2, Shi et al. (2020b), we have by equation 48 that $\|(\overline{\boldsymbol{\Sigma}}^{(k)})^{-1}\|_2 = O(1)$, with probability tending to 1. equation 33 is thus proven.

Assertion equation 32 now follows using similar arguments in Part 2 and Part 3 of the proof of Lemma E.2, Shi et al. (2020b).

H   ADDITIONAL FIGURES

We present some additional figures to report the simulation results in this section.

(a) The proposed test under H_1 and H_0 (from left plots to right plots). $J = 3, \alpha(\cdot) = \alpha_1(\cdot)$.

(b) The proposed test under H_1 and H_0 (from left plots to right plots). $J = 3, \alpha(\cdot) = \alpha_2(\cdot)$.

(c) The proposed test under H_1 and H_0 (from left plots to right plots). $J = 5, \alpha(\cdot) = \alpha_1(\cdot)$.

(d) The proposed test under H_1 and H_0 (from left plots to right plots). $J = 5, \alpha(\cdot) = \alpha_2(\cdot)$.

Figure 5: Empirical rejection probabilities of our test. Settings correspond to the alternating-time-interval, adaptive and Markov design, from top plots to bottom plots.