EDCDE - Extended Discovery of Closed-Form Differential Equations

Abstract

Understanding the mathematical connections between variables in a physical system, such as Ordinary Differential Equations (ODEs) is an essential part of the scientific method. This is where symbolic regression plays a key role in looking for closed-form functions given a dataset. We extend the results of the original Discovering Closed-Form ODEs from Observed Trajectories (D-CODE) by considering a generalized variational formulation that can work with most forms of ODEs. We conclude the paper with numerical results and applications.

1 Introduction

Recently Qian et al. (2021) introduced the D-CODE paper, where they studied first-order autonomous systems. However, they do comment on how one can transform non-autonomous and higher-order ODEs into systems of autonomous ODEs by including more variables. This can be useful in certain situations, such as when the system's behaviour is easier to analyze or when more efficient numerical algorithms can be developed to solve the system. However, this process can be challenging, especially if the system is complex or the equations that describe it are poorly understood. It may require significant mathematical analysis and modelling efforts to successfully reduce the order of an autonomous system. This can make it challenging to understand how the non-autonomous system is behaving and how it is likely to behave in the future. Moreover, the increase from n variables to a system of $2n + 1$ can be cumbersome. In this tiny paper, we showcase we can easily solve the original system of order n using a generalized variational formulation which is the first of its kind. We also provide numerical results to support our claim and consider extensions of higher degree higher order systems in the Appendix.

2 First Degree Higher Order ODEs

we consider the system with the system with $J \in \mathbb{N}^+$ variables defined as $\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t)) = f_j(\boldsymbol{x}(t)), \forall j = 1, \ldots, J$ where we use Newton's notation $\overset{n}{\dot{x}}_j(t)$ for the *nth* time derivative and $\mathcal{F}(x, \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t))$ to denote any combination of these derivative powers. More specifically $\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t)) = \sum_{i=0}^{n} \alpha_i \overset{i}{\dot{x}}_j(t)$ where α_i are just arbitrary constants and $\overset{0}{\dot{x}}_j(t) = x_j(t)$. The functions $f_j : \mathbb{R}^J \to \mathbb{R}$ will sometimes be directly referred to as the ODEs. We denote $T \in \mathbb{R}^+$ as the maximum time horizon we have the data for, and the trajectory $x_j : [0, T] \to \mathbb{R}$ is a function of time, whereas the state $x_j(t) \in \mathbb{R}, \forall t \in [0, T]$ is a point on the trajectory. We denote the state vector $:= [x_1(t), \ldots, x_J(t)]^\top \in \mathbb{R}^J$ and the vector-valued trajectory function $\boldsymbol{x} := [x_1, \ldots, x_J]$. Similarly, as in the paper, let f_j^*'s be the true but unknown ODEs to be uncovered, and $\boldsymbol{x}_i : [0, T] \to \mathbb{R}^j, i \leq N, N \in \mathbb{N}^+$ be the true trajectories that satisfy f_j^*'s. In practice, we only measure the true trajectories at discrete times and with noise. Denote the measurement of trajectory i at time t as $\boldsymbol{y}_i(t) \in \mathbb{R}^J$; we assume

$$\boldsymbol{y}_i(t) = \boldsymbol{x}_i(t) + \epsilon_i(t), \quad \forall i \leq N, t \in \mathcal{T}$$

where $\epsilon_i(t) \in \mathbb{R}^J$ is zero-mean noise with standard deviation σ. The measurements are made at time $t \in \mathcal{T} = \{t_1, t_2, \ldots, T\}$. We denote the dataset as $\mathcal{D} = \{\boldsymbol{y}_i(t) \mid i \leq N, t \in \mathcal{T}\}$.

2.1 GENERALIZED VARIATIONAL FORMULATION

The variational formulation provides a direct link between the trajectory $\boldsymbol{x}$ and the ODE f_j without involving any of the nth derivatives $\overset{n}{x}_j(t)$. In particular consider $J \in \mathbb{N}^+, T \in \mathbb{R}^+$, continuous functions $\boldsymbol{x} : [0,T] \to \mathbb{R}^J, f : \mathbb{R}^J \to \mathbb{R}$, and $g \in \mathcal{C}^n[0,T]$, where $\mathcal{C}^n$ is the set of n times continuously differentiable functions. We define the functionals

$$C_j(f, \boldsymbol{x}, g^n) := \int_0^T \left(f(\boldsymbol{x}(t))g(t)^n + x_j(t)\Phi(\overset{n}{g}(t)) \right) dt, \quad \forall j \in \{1, 2, \ldots, J\} \tag{1}$$

where $\Phi(\overset{n}{g})$ is a function of derivatives of g. To get the explicit representation of Φ we must do a Taylor series expansion of g^n and choose the *n-th* term in the series expansion for the corresponding $\overset{n}{g}$. We obtain a linear combination of the derivatives of $\dot{g}$. The functional C_j depends now on the testing function $g(t)$ and its n - derivatives $\overset{n}{g}(t)$ but not on any n - derivatives of $\overset{n}{x}_j(t)$. Then we can easily consider an extension of the proposition in the paper, which can be stated as follows.

Proposition 1: Consider $J \in \mathbb{N}^+, T \in \mathbb{R}^+$, a continuously differentiable function $\boldsymbol{x} : [0,T] \to \mathbb{R}^J$, and continuous functions $f_j : \mathbb{R}^J \times [0,T] \to \mathbb{R}$ for $j = 1, \ldots, J$. Then $\boldsymbol{x}$ is the solution to the system of ODEs in Equation 1 if and only if
$$C_j(f_j, \boldsymbol{x}, g^n) = 0, \forall g \in \mathcal{C}^n[0,T], g(0) = g(T) \equiv 0$$

In regard to the choice of the testing functions, we can consider $g_s(t) = \sqrt{(2/T)}\sin(s/T)$ which are an orthonormal basis for $L^2[0,T]$ and satisfy $g(0) = g(T) = 0$. Moreover, their derivatives are easy to compute and have a closed-form analytic solution such as for odd powers we have $\overset{2k+1}{\dot{g}_s}(t) = \dfrac{(-1)^k \sqrt{2}\,\pi^k s^k \cos\left(\frac{\pi sx}{T}\right)}{T^{\frac{2k+1}{2}}}$ and for even powers of n we have $\overset{2k}{\dot{g}_s}(t) = \dfrac{(-1)^k \sqrt{2}\,\pi^k s^k \sin\left(\frac{\pi sx}{T}\right)}{T^{\frac{2k+1}{2}}}$.
Finally the Pseudocode and proof of proposition 1 is in the appendix.

3 NUMERICAL RESULTS

We create generalized ODEs from the ODEs considered in the original paper. In particular, we consider the third-order generalized ODEs (Logistic and Gompertz) given by

$$\overset{3}{x}(t) = -\theta_1 x(t) \cdot \log(\theta_2 x(t)), \quad \overset{3}{x}(t) = \theta_1 x(t) \cdot \left(1 - x(t)^{\theta_2}\right), \ \theta_1, \theta_2 \in \mathbb{R}^+ \tag{2}$$

We have no benchmark to compare our EDCDE algorithm as it is difficult to estimate the $\overset{3}{x}$ and hence will present the closed-form ODEs discovered by the algorithm. Tables A.3 and A.3 showcase our results and some of the closed-form ODEs discovered by EDCDE. We fix the number of basis as 50, do 10 simulations for each experiment and set the noise level to 0 for simplicity.

4 APPLICATIONS

Like the original D-CODE paper, we demonstrate that our generalized algorithms can be used to discover most closed-form ODEs. Examples include

1. **Data Centric AI:** As the amount of data available to organizations has grown dramatically, advances in ML have made it possible to build more powerful and accurate models using this data. Using this data, we can use symbolic regression to analyze large, complex datasets and identify underlying ODE/PDE systems and relationships that may not immediately appear to human analysts.

2. **Theory discovery:** Closed-form Differential equations can be very useful because they provide a compact and interpretable representation of a system's behaviour and can be used to make predictions and understand the underlying relationships governing the system. It is possible to use symbolic regression to develop new theories about the system and make predictions about its behaviour.

URM STATEMENT

The authors acknowledge that the author of this work meets the URM criteria of ICLR 2023 Tiny Papers Track.

REFERENCES

Zhaozhi Qian, Krzysztof Kacprzyk, and Mihaela van der Schaar. D-code: Discovering closed-form odes from observed trajectories. In *International Conference on Learning Representations*, 2021.

A   APPENDIX

A.1   PROOF OF PROPOSITION 1

Lemma 1: If we had an ODE of order n, i.e. $\overset{n}{\dot{x}_1}$ then the following statements are true

1: $\overset{n}{\dot{x}_1}$ the corresponding Taylor series expansion should be computed until the $n + 1$ term, and the first n terms correspond to the boundary terms, which equal to 0 due to $g(0) = g(T) = 0$. The $n+1$ term is the corresponding integral term.

2: $\overset{n-1}{\dot{x}_1}$ the corresponding boundary terms equal to the first $n - 1$ terms and the n term is the integral. Continuing the process, we eventually get that for the $\dot{x}_1$ term, our boundary term is equal to the first term, and the integral is equal to the second term.

Proof 1: We now prove our proposition in full generality. The proof is straightforward, as in the original paper, with repeated integration by parts. We have from 1 that

$$\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}_j}(t)) = f_j(\boldsymbol{x}(t))$$

Now using the Fundamental lemma of the calculus of variations, we get the following two statements are equivalent.

$$\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}_j}(t)) - f_j(\boldsymbol{x}(t)) = 0 \ \forall t \in [0, T]$$

$$\int_0^T \left(\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}_j}(t)) - f_j(\boldsymbol{x}(t)) \right) g(t)dt = 0, \quad \forall g \in \mathcal{C}^n[0, T], g(0) = g(T) \equiv 0$$

Using linearity of integration, we get

$$= \int_0^T \left(\mathcal{F}(x_j(t), \ldots, \overset{n}{\dot{x}_j}(t)) \right) g(t)dt - \int_0^T f_j(\boldsymbol{x}(t))g(t)dt$$

$$= \int_0^T \left(\sum_{i=0}^n \alpha_i \overset{i}{\dot{x}_j}(t) \right) g(t)dt - \int_0^T (f_j(\boldsymbol{x}(t))) g(t)dt$$

$$= \sum_{i=0}^n \int_0^T \alpha_i \overset{i}{\dot{x}_j}(t)g(t)dt - \int_0^T (f_j(\boldsymbol{x}(t))) g(t)dt$$

Where the swap can be made due to the dominated convergence theorem (Again, we only have a finite sum, so it is valid mathematically). Now we only care about the first term above. Therefore we get (Avoiding the explicit notation of t)

$$= \sum_{i=0}^n \int_0^T \alpha_i \overset{i}{\dot{x}_j}(t)g(t)dt = \sum_{i=0}^n \left(\alpha_i \left(\sum_{k=1}^i \xi_k \{g^n(t)\} \overset{k}{\dot{x}} - \int_0^T \xi_{i+1} \{g^n(t)\} x_j dt \right) \right)$$

$$= \sum_{i=0}^n \left(- \int_0^T \xi_{i+1} \{g^n(t)\} x_j dt \right)$$

where in the last equality we used the fact that $g(0) = g(T) = 0$ and where $\xi_k \{g^n(t)$ is the $k - th$ term in the power series expansion of $g^n(t)$. That means that

$$= - \int_0^T \sum_{i=0}^n \xi_{i+1} \{g^n(t)x_j\} dt = \int_0^T \Phi(\overset{n}{\dot{g}}(t))x_j dt = \int_0^T \left(f(\boldsymbol{x}(t))g(t)^n + x_j(t)\Phi(\overset{n}{\dot{g}}(t)) \right) dt$$

This proves that (Let $\mathcal{J} = \{1, 2 \ldots J\}$ and $\forall j \in \mathcal{J}$)

$$\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t)) = f_j(\boldsymbol{x}(t)) \iff C_j(f_j, \boldsymbol{x}, g^n) = 0, \ \forall g \in \mathcal{C}^n[0, T], \ g(0) = g(T) = 0$$

A.2 PSEUDOCODE

Algorithm 1 EDCDE - ODE

   Input: Dataset $\mathcal{D} = \{\boldsymbol{G}_i(t) \mid i \leq N, t \in \mathcal{T}\}$
   Input: Smoothing Algo S, Optimization Algo O
   Input: Numerical Integration Algo I, Order n
   Input: Test functions $g_s, s \leq S$, Initial guesses $f_j, j \leq J$ $\widehat{x}_i = \mathcal{S}(\boldsymbol{x}_i(t_1), \boldsymbol{x}_i(t_2), \ldots), i \leq N$
   Φ = Symbolically differentiate $g(x)^n$
   for $i = 1$ **to** J **do**
      $\hat{f}_j = f_j$
      Converge = False
      while Not Converge **do**
         obj = $\sum_{i=1}^{N} \sum_{s=1}^{S} C_j(f_j, \widehat{x}_i, g_s^n)$ [Use Φ]
         $\hat{f}_j$, Converge = $\mathcal{O}(obj, \hat{f}_j)$
      end while
   end for
   Output: The discovered ODE's $f_j, \ j \leq J$

A.3 NUMERICAL RESULTS

Higher Order ODE Training Time		
ODE	Training Time	Order
Logistic	1540	1
Gompertz	1200	1
Logistic	1080	3
Gompertz	3000	3

Higher Order ODE Closed Form Discovery		
ODE	Order	ODE Discovered (Best)
Logistic	1	$\theta_1 x(t) - x(t)^{\theta_2}$
Gompertz	1	$-\theta_1 x(t) \cdot \log(\theta_2 x(t))$
Logistic	3	$x(t)^{\theta_1}(\theta_2 - x(t))$
Gompertz	3	$(-x(t) - 4\log x(t))(\theta_1 \log(\theta_2 x(t)))$

B CONVERGENCE TO DISTANCE

The convergence properties remain the same as in the original paper and the proof of the theorem follows the same procedure as the original paper but replaces g_s by g_s^n.

Theorem 2: Consider $J \in \mathbb{N}^+, j \in \{1, \ldots, J\}, T \in \mathbb{R}^+$. Let $f^* : \mathbb{R}^J \to \mathbb{R}$ be a continuous function, and let $\boldsymbol{x} : [0, T] \to \mathbb{R}^J$ be a continuously differentiable function satisfying $\dot{x}_j(t) = f^*(\boldsymbol{x}(t))$. Consider a sequence of functions $(\widehat{\boldsymbol{x}}_k)$, where $\widehat{\boldsymbol{x}}_k : [0, T] \to \mathbb{R}^J$ is a continuously differentiable function. If $(\widehat{\boldsymbol{x}}_k)$ converges to $\boldsymbol{x}$ in L^2 norm. Then for any Lipschitz continuous function f

$$\lim_{S \to \infty} \lim_{k \to \infty} \sum_{s=1}^{S} C_j(f, \widehat{\boldsymbol{x}}_k, g_s^n)^2 = d_{\boldsymbol{x}}(f, f^*)^2,$$ where $\{g_1, g_2, \ldots\}$ is a Hilbert (orthonormal) basis

for $L^2[0, T]$ such that $\forall i, g_i(0) = g_i(T) = 0$ and $g_i \in \mathcal{C}^n[0, T]$.

C NON - AUTONOMOUS ODES

We consider a Non-Autonomous system of ODEs as $\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t)) = f_j(\boldsymbol{x}(t), t), \forall j = 1, \ldots, J$. The only key difference here is to make one variable substitution i.e. $x_m = t$, and so $\dot{x}_m = 1$, and we solve the following ODE system

$$\mathcal{F}(x_j(t), \dot{x}_j(t), \ldots, \overset{n}{\dot{x}}_j(t)) = f_j(\boldsymbol{x}(t), t) \text{ and } \dot{x}_m = 1, \ m = J+1, \quad \forall j = 1, \ldots, J \quad (3)$$

Then we can easily use our generalized D-CODE algorithm given the measurements of x_m and x_j.