Score-Based Generative Models with Lévy Processes

Abstract

Time reversibility of stochastic processes is a primary cornerstone of the score-based generative models through stochastic differential equations (SDEs). While a broader class of Markov processes is reversible, previous continuous-time approaches restrict the range of noise processes to Brownian motion (BM) since the closed-form of the time reversal formula is only known for diffusion processes. In this paper, to expand the class of noise distribution, we propose a class of score-based probabilistic generative models, Lévy-Itō Model (LIM), which utilizes α-stable distribution for noise injection. To this end, we derive an approximate time reversal formula for the SDEs with Lévy processes that can allow discontinuous pure jump motion. Consequently, we advance the score-based generative models with a broad range of non-Gaussian Markov processes. Empirical results on MNIST, CIFAR-10, CelebA, and CelebA-HQ show that our approach is valid.

1 Introduction

The recent successes of score-based generative models [24, 26, 11] and their applications [18, 14, 6] draw huge attention from machine learning communities. Score-based generative models via stochastic differential equations (SDEs) [26] rely on the time reversal theory of diffusion processes, Anderson theorem [1], which shows that the time reversal of the diffusion process belongs to the class of diffusion processes again. One can interpret this result as solving a martingale problem which induces a weak solution to the reverse SDEs [9, 4]. Due to the advances in the SDE theory with jump Markov processes [13, 22, 5], one can desire a positive expectation for applying a class of non-Gaussian noise distribution to score-based generative models. However, since the closed form of the time reversal formula is only known for diffusion processes, whether a score-based method is feasible for a non-Gaussian Markov process other than a Brownian motion has been an open question in this field. To tackle the challenging problem, we propose the second-order approximate formula for the time reversal of SDEs with Lévy processes and a novel score-based generative method, Lévy-Itō Model (LIM), which utilizes a symmetric α-stable distribution as noise injection. We apply the proposed method to MNIST, CIFAR-10, CelebA, CelebA-HQ. Our approach and empirical results establish the bridge between probability theory and generative models.

2 Score-Based Generative Models with Lévy Processes

2.1 Background

Symmetric α-stable distribution   Let $\alpha \in (0, 2]$ be a characteristic exponent which determines the decay rate at which the tails of the distribution, and γ be a scale parameter. Symmetric α-Stable distributions $\mathcal{S}\alpha\mathcal{S}(\gamma)$ have the heavy-tail properties $P(X > x) \sim x^{-\alpha}$ and densities with unknown closed-form expressions, except for the Cauchy ($\alpha = 1$) and Gaussian ($\alpha = 2$) [1].

---

[1]When $\alpha = 2$, it holds $\mathcal{S}\alpha\mathcal{S}(\gamma) = \mathcal{N}(0, \sqrt{2}\gamma)$.

Figure 1: (a) PDF of Gaussian and α-stable distributions. α-stable distributions have heavier tails as alpha decreases. (b) Trajectories of Brownian motion and α-stable Lévy processes with different α. Lévy processes can have infinitely many discontinuous jumps, unlike Brownian motion.

Lévy process and α-stable Lévy motion   A stochastic process $L_t = (L_t)_{t \geq 0}$ with $L_0 = 0$ is called Lévy process if (i) L_t has independent increments, (ii) L_t has stationary increments, (iii) L_t is stochastically continuous. If the difference $(L_t - L_s)$ and L_{t-s} have the same distribution following $\mathcal{S}\alpha\mathcal{S}((t-s)^{1/\alpha})$ for $s < t$, then the Lévy process is called symmetric α-stable Lévy motion L_t^α. Due to the stochastic continuity (iii), Lévy processes have a countable number of discontinuous points (i.e. jumps) [31]. Notably, L_t^α is a prototypical pure jump process. The heavy-tail properties of α-stable distribution implies that the frequency of large jumps of L_t^α increases as α gets smaller (see Figure 1).

2.2   Lévy-Itō Model: Approximate Time-Reversal of SDEs driven by Lévy Processes

Due to Lévy-Ito decomposition [2, 16], we consider a family of SDEs driven by a Lévy process consisting of continuous Brownian motion part B_t and pure jump part L_t^α as follows:

$$d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_B(t)dB_t + \sigma_L(t)dL_t^\alpha, \quad t \in [0, 1] \tag{1}$$

The following approximate time-reversal formula is our main result.

Theorem 2.1 (Time-reversal formula of SDEs with Lévy Processes)**.** The reverse SDE of (1) is

$$d\overleftarrow{X}_t = \left(b(t, \overleftarrow{X}_t) - [\sigma_B^2(t) + C_\alpha\sigma_L^2(t)]\nabla_x \log p_t(\overleftarrow{X}_t)\right)dt + \sigma_B(t)d\bar{B}_t + \sigma_L(t)d\bar{L}_t^\alpha \tag{2}$$

where $C_\alpha = \frac{\Gamma(\alpha+1)}{\frac{\pi}{2}(2-\alpha)}\sin(\frac{\pi}{2}(2-\alpha))$. $\bar{B}_t, \bar{L}_t^\alpha$ is a corresponding backward Brownian motion and backward symmetric α-stable Lévy motion, respectively.

Here (2) uses the second-order approximation to obtain the explicit SDE representation. See Theorem A.10 for more details. We also remark that (2) recovers the result of [1] if $\alpha \to 2$. To shed light on the probabilistic approach with the jump Markov process, we propose **Lévy-Itō M**odel (LIM), a novel score-based generative model through SDE driven by symmetric α-stable motion only ($\sigma_B(t) \equiv 0$):

$$d\overleftarrow{X}_t = \left(b(t, \overleftarrow{X}_t) - C_\alpha\sigma_L^2\nabla_x \log p_t(\overleftarrow{X}_t)\right)dt + \sigma_L(t)d\bar{L}_t^\alpha \tag{3}$$

Considering a beta-scheduling version of (3), we obtain $d\overrightarrow{X}_t = -\frac{\beta(t)}{\alpha}\overrightarrow{X}_t + (\beta(t))^{1/\alpha}dL_t^\alpha$. Then the solution becomes $\overrightarrow{X}_t = a(t)\overrightarrow{X}_0 + \gamma(t)\epsilon$, $\epsilon \sim \mathcal{S}\alpha\mathcal{S}(1)$, where $a(t)$ is $\exp(-\int_0^t \frac{\beta(s)}{\alpha}ds)$ and the scale parameter $\gamma(t)$ is $(1 - a(t))^{1/\alpha}$ (see Lemma C.3). Due to the Euler-Maruyama method, we can induce a stochastic sampling of LIM, a sequence convergent to a data point in p_{data}(see Corollary C.3.1).

The Probability ODE   We can also derive the Probability ODE of $d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_L(t)dL_t^\alpha$:

$$d\overrightarrow{X}_t = \left(b(t, \overrightarrow{X}_t) - \frac{C_\alpha}{2}\sigma_L^2(t)\nabla_x \log p_t(\overrightarrow{X}_t)\right)dt \tag{4}$$

The proof of (4) can be found in Theorem B.3. Deterministic ODE sampling of LIM can be deduced from (4) by using the Euler-Maruyama method (see Corollary C.3.2).

Figure 2: Comparison between (a) Lévy score and (b) ReELS. (c) and (d) the synthetic data sampled by the SDE with (3) ($\alpha = 1.5$) trained with Lévy score and ReELS, respectively. There are divergent points as indicated in the red circle since the value of Lévy score decreases for large noise.

2.3 Score function for Lévy-Itō Model

Let q_α be the density function of $\mathcal{S}\alpha\mathcal{S}$. Recall the solution of beta-scheduling version of (3) is $\overrightarrow{X}_t = a(t)\overrightarrow{X}_0 + \gamma(t)\epsilon$ with the transition density function $p_t(x_t|x_0)$. The score function of $p_t(x_t|x_0)$ satisfies $\nabla \log p_t(x_t|x_0) = \nabla \log q_\alpha(\epsilon)/\gamma(t)$ (See Lemma D.1). We denote $S_\alpha = \nabla \log q_\alpha$. Figure 2.(a) shows that the score function of BM is linearly decreasing, while the Lévy score functions are not monotonic. For instance, the Lévy score towards to zero when $|x| \geq 2$ if $\alpha = 1.5$. Hence, if we train the score model to target the Lévy score, it is difficult to denoise the divergent large noise generated at the heavy tail (Figure 2.(b)). These phenomena worsen as α decreases.

Rectified Enhanced Lévy Score (ReELS) One can employ rejection sampling (i.e. clamping) to avoid the large noise, by reassigning a point inside the clamped range when a jump exceeds a certain size. However, rejection sampling requires exhaustive hyperparameter tuning during the training and sampling. To denoise the large noise at the heavy tail without losing the nature of the Lévy score function, we propose **Re**ctified **E**nhanced **L**évy **S**core (ReELS) as follows:

$$\text{ReELS}_\alpha(x) = \begin{cases} S_\alpha(x) & : x \in I_\alpha \\ -\text{sgn}(x)\hat{c}|x|^{\hat{\beta}} & : \text{otherwise} \end{cases}, \quad \hat{\beta}(\alpha) \in (0, 1) \tag{5}$$

Here we set the range I_α as the interval between two local optimum points of the given Lévy score. We find parameters $\hat{c}, \hat{\beta}$ in ReELS by fitting $-\text{sgn}(x)\hat{c}|x|^{\hat{\beta}}$ to the Lévy score inside I_α (see Figure 2.(b)). This procedure is equivalent to the fitting score function of a generalized Gaussian distribution to the Lévy score [17]. Note that ReELS_α recovers the BM score as α converges to 2. We remark that utilizing the BM score does not outperform ReELS for Lévy-driven SDEs because generalized Gaussian distributions have a score function more similar to the Lévy score function (see Table D.1, D.2, and D.3). The experiments on synthetic data (Mixture of Gaussian, Two-Moon, Swiss-Roll) demonstrate that LIM trained by ReELS converges to the true data distribution and performs better than using BM score for Lévy-driven SDEs.

2.4 Loss function

We use the U-net architecture [19] as in DDPM [11] and apply L_2-loss to train the model using $\text{ReELS}_\alpha(\epsilon)$ as a label through the Denoising Score Matching (DSM) [25] . For $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$ and $x_0 \sim p_\text{data}$, we let $x_t = a(t)x_0 + \gamma(t)\epsilon$ where $\beta(t) = \beta_0 + (\beta_1 - \beta_0)t$, $a(t) = \exp(-\frac{(\beta_1-\beta_0)}{2\alpha}t^2 - \frac{\beta_0}{\alpha}t)$, and $\gamma(t) = (1 - a(t)^\alpha)^{1/\alpha}$. Let $U(0, 1)$ denotes an uniform distribution. Then the loss with the relative weight $\gamma(t)$ is defined as

$$L(\theta; \gamma(t)) := \mathbb{E}_{t\sim U(0,1)}\mathbb{E}_{x_0\sim p_\text{data}}\mathbb{E}_{\epsilon\sim\mathcal{S}\alpha\mathcal{S}}\|\gamma(t)S_\theta(x_t, t) - \text{ReELS}_\alpha(\epsilon)\|_2^2 \tag{6}$$

(a) MNIST Stochastic sampling in varied α values

(b) Top: FID, Bottom: bits/dim

Figure 3: Generated MNIST images (a) by DM (Brownian motion, [26]) and LIM (α-stable Lévy motion for $\alpha = 1.8, 1.5, 1.2$), and corresponding plots (b) of FID score($\downarrow$) and bits per dimension ($\downarrow$) for different NFEs with various α. LIM shows a faster generation speed than DM.

3 Experiment

We empirically validate the proposed score-based generative model on image data including MNIST (Figure E.1), CIFAR10 (Figure E.2), CelebA (Figure E.4, E.5), and CelebA-HQ (Figure E.6). We adjust the model size of each dataset for training efficiency. For $\alpha \in \{1.8, 1.5, 1.2\}$, we train our model on MNIST for 1000 epochs with $\beta_0 = 0.1$, $\beta_1 = 5.0$ and use the noise clamping to control the large-scale noise to improve the sample quality. See Section E.1 to find the other configurations in different datasets. Figure 3.(a) shows that LIM with ReELS converges faster than DM (Diffusion Models [26], $\alpha = 2.0$). See Figure 3.(b) to compare FID scores [10] and bits per dimension (bits/dim) on the MNIST dataset for each number of function evaluations (NFE) with different α. To evaluate bits/dim, we use a uniformly dequantized test dataset with 5 iterations and compute log-likelihood by using an ODE solver (See Section E.2.2). LIM achieves competitive sample quality to DM at $\alpha = 1.5, 1.8$, and tends to converge quickly as lower α. Figure 4 shows ReELS can be adaptive to the

Figure 4: (a) SDE sampling results and (b) deterministic ODE sampling results with different NFEs. ODE shows a faster sampling speed than that of SDE.

probability ODE, which enables fast sampling than stochastic sampling. Although the large jump of the reverse process can be controlled in the reverse SDE sampling, it is challenging to control the jump size in the deterministic ODE sampling. Hence, the image quality may degrade due to the effect of uncontrolled large noise, which leads to the higher bits per dimension in LIM 3.17 ($\alpha = 1.8$) > 1.67 ($\alpha = 2.0$).

4 Conclusion

In this paper, we broaden the range of noise distribution used in score-based generative models by inducing an approximate time reversal formula for SDEs with Lévy processes and by proposing a novel score-based generative model, Lévy-Itō Model (LIM) with Rectified Enhanced Lévy Score (ReELS). Empirical results validate that the proposed approach works well with different ranges of symmetric α-stable Lévy distributions in synthetic datasets and various image data. Consequently, our study presents a feasible solution and demonstrates the potential for applying a broader class of non-Gaussian Markov processes to score-based generative models.

References

[1] Brian D.O. Anderson. Reverse-time diffusion equation models. volume 12, pages 313–326, 1982.

[2] David Applebaum. *Lévy processes and stochastic calculus*. Cambridge university press, 2009.

[3] Björn Böttcher. Feller evolution systems: Generators and approximation. *Stochastics and Dynamics*, 14(03):1350025, 2014.

[4] Patrick Cattiaux, Giovanni Conforti, Ivan Gentil, and Christian Léonard. Time reversal of diffusion processes under a finite entropy condition. *arXiv preprint arXiv:2104.07708*, 2021.

[5] Giovanni Conforti and Christian Léonard. Time reversal of markov processes with jumps under a finite entropy condition. 2021.

[6] Florinel-Alin Croitoru, Vlad Hondru, Radu Tudor Ionescu, and Mubarak Shah. Diffusion models in vision: A survey. *arXiv preprint arXiv:2209.04747*, 2022.

[7] John R Dormand and Peter J Prince. A family of embedded runge-kutta formulae. *Journal of computational and applied mathematics*, 6(1):19–26, 1980.

[8] Will Grathwohl, Ricky TQ Chen, Jesse Bettencourt, Ilya Sutskever, and David Duvenaud. Ffjord: Free-form continuous dynamics for scalable reversible generative models. *arXiv preprint arXiv:1810.01367*, 2018.

[9] U. G. Haussmann and E. Pardoux. Time Reversal of Diffusions. *The Annals of Probability*, 14(4):1188 – 1205, 1986.

[10] Martin Heusel, Hubert Ramsauer, Thomas Unterthiner, Bernhard Nessler, Günter Klambauer, and Sepp Hochreiter. Gans trained by a two time-scale update rule converge to a nash equilibrium. *arXiv preprint arXiv:1706.08500*, 12(1), 2017.

[11] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. In H. Larochelle, M. Ranzato, R. Hadsell, M.F. Balcan, and H. Lin, editors, *Advances in Neural Information Processing Systems*, volume 33, pages 6840–6851. Curran Associates, Inc., 2020.

[12] Michael F Hutchinson. A stochastic estimator of the trace of the influence matrix for laplacian smoothing splines. *Communications in Statistics-Simulation and Computation*, 18(3):1059–1076, 1989.

[13] Jean Jacod and Philip Protter. Time Reversal on Levy Processes. *The Annals of Probability*, 16(2):620 – 641, 1988.

[14] Jihoon Kim, Jiseob Kim, and Sungjoon Choi. Flame: Free-form language-based motion synthesis & editing. *arXiv preprint arXiv:2209.00349*, 2022.

[15] Thomas G. Kurtz. Equivalence of stochastic equations and martingale problems. 2011.

[16] Paul Lévy. Sur les intégrales dont les éléments sont des variables aléatoires indépendantes. *Annali della Scuola Normale Superiore di Pisa-Classe di Scienze*, 3(3-4):337–366, 1934.

[17] Shengyang Luan, Jiayuan Li, Yinrui Gao, Jinfeng Zhang, and Tianshuang Qiu. Generalized covariance-based esprit-like solution to direction of arrival estimation for strictly non-circular signals under alpha-stable distributed noise. *Digital Signal Processing*, 118:103214, 2021.

[18] Aditya Ramesh, Prafulla Dhariwal, Alex Nichol, Casey Chu, and Mark Chen. Hierarchical text-conditional image generation with clip latents. *arXiv preprint arXiv:2204.06125*, 2022.

[19] Olaf Ronneberger, Philipp Fischer, and Thomas Brox. U-net: Convolutional networks for biomedical image segmentation. In *International Conference on Medical image computing and computer-assisted intervention*, pages 234–241. Springer, 2015.

[20] Ludger Rueschendorf, Alexander Schnurr, and Viktor Wolf. Comparison of time-inhomogeneous markov processes, 2015.

[21] Tim Salimans and Durk P Kingma. Weight normalization: A simple reparameterization to accelerate training of deep neural networks. *Advances in neural information processing systems*, 29, 2016.

[22] Umut Şimşekli. Fractional langevin monte carlo: Exploring lévy driven stochastic differential equations for markov chain monte carlo. In *International Conference on Machine Learning*, pages 3200–3209. PMLR, 2017.

[23] John Skilling. *Maximum Entropy and Bayesian Methods: Cambridge, England, 1988*, volume 36. Springer Science & Business Media, 2013.

[24] Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. 2019.

[25] Yang Song and Stefano Ermon. Generative modeling by estimating gradients of the data distribution. In H. Wallach, H. Larochelle, A. Beygelzimer, F. d'Alché-Buc, E. Fox, and R. Garnett, editors, *Advances in Neural Information Processing Systems*, volume 32. Curran Associates, Inc., 2019.

[26] Yang Song, Jascha Sohl-Dickstein, Diederik P Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations. In *International Conference on Learning Representations*, 2020.

[27] Padmanabhan Sundar. Time reversal of solutions of equations driven by lévy processes. 1992.

[28] Isabelle Tristani. Fractional fokker-planck equation, 2013.

[29] Ashish Vaswani, Noam Shazeer, Niki Parmar, Jakob Uszkoreit, Llion Jones, Aidan N Gomez, Łukasz Kaiser, and Illia Polosukhin. Attention is all you need. *Advances in neural information processing systems*, 30, 2017.

[30] Yuxin Wu and Kaiming He. Group normalization. In *Proceedings of the European conference on computer vision (ECCV)*, pages 3–19, 2018.

[31] Umut Şimşekli. Fractional langevin monte carlo: Exploring lévy driven stochastic differential equations for markov chain monte carlo, 2017.

Appendix

A Time-reversal of SDE

In this chapter, given the generator of the general Markov process with jump kernel, we show that the reverse form can be deduced into a computational formula under certain conditions. Finally, we introduce the stochastic sampling and the deterministic ODE sampling of LIM. Let us outline some necessary lemmas before we move on to the proof. A homogeneous Markov process that corresponds to an inhomogeneous Markov process $(\overrightarrow{X}_t)$ always exists according to Lemma A.1. The existence of an SDE representation of a homogeneous Markov process with a particular generator is given in Lemma A.5. Lemma A.6 introduces the general reverse-time formula. Through the transformation of time-inhomogeneous Markov processes and a SDE representation of given a specific generator, we find the SDE representation that corresponds to the generator of the time-reverse process. From these lemmas, we deduce a reverse SDE representation, on which we also get a stochastic sampling and a deterministic ODE sampling based on probability ODE if a Markov process is provided as a solution to (1).

A.1 Time-Reversal of General Markov process with jump kernel

Definition A.1 (Evolution system). Let $(\overrightarrow{X}_t)$ be an $\mathbb{R}^d$-valued continuous time inhomogeneous Markov process on an probability space $(\Omega, \mathcal{A}, \mathbb{P})$. The evolution system is defined as

$$T(s,t)u(x) = \mathbb{E}(u(\overrightarrow{X}_t)|\overrightarrow{X}_s = x) \text{ for } s \leq t, s,t \in \mathbb{R} \tag{7}$$

and this operator is well-defined on the set of Borel measurable function u on $\mathbb{R}^d$, denoted by $B(\mathbb{R}^d)$. The operator is linear and positive preserving with $T(s,t)1 = 1$ and $T(s,t) = T(s,r)T(r,t)$ for $s \leq r \leq t$. This operator is also strongly continuous such that for each $v, w \in \mathbb{R}, v \leq w$ and $s \leq t$ $\lim_{(s,t)\to(v,w)} ||U(s,t)u - U(v,w)u||_\infty = 0$ where $||\cdot||_\infty$ is the supreme norm. For all $u \in C_\infty(\mathbb{R}^d)$, the set of a continuous function with vanishing at ∞, the generators of the evolution system is given by

$$\mathcal{L}_s u = \lim_{h \to 0} \frac{T(s,s+h)u - u}{h} \text{ for each } s \in \mathbb{R} \tag{8}$$

A family of linear operators $T(s,t)$ on C_∞ is a Feller evolution system if it is a strongly continuous, positive, contraction semigroup on C_∞.

Definition A.2 (Space-time process). Let $\mathcal{B}$ be a Borel algebra in $\mathbb{R}^d$ and an a state space $(\mathbb{R}_+ \times \mathbb{R}^d, \tilde{\mathcal{B}})$ with $\tilde{x} \in \mathbb{R}_+ \times \mathbb{R}^d$ and σ-algebra $\tilde{\mathcal{B}} = \{B \in \mathbb{R}_+ \times \mathbb{R}^d | B_s \in \mathcal{B}\}$, and a new sample space $(\tilde{\Omega}, \tilde{\mathcal{A}})$ with $\tilde{w} = (s, w) \in \mathbb{R}_+ \times \Omega = \tilde{\Omega}$ and $\tilde{\mathcal{A}} = \{A \subset \mathbb{R}_+ \times \Omega | A_s \in \mathcal{A}, \forall s \in \mathbb{R}_+\}$. A space-time process $(\tilde{X}_t)$ is defined by

$$\tilde{X}_t(\tilde{w}) = (s + t, \overrightarrow{X}_{s+t}(w)) \tag{9}$$

with the probability measure for $A \in \tilde{\mathcal{A}}$ and $\tilde{x} \in \mathbb{R}_+ \times \mathbb{R}^d$ such that $\tilde{P}_{\tilde{x}}(A) = \tilde{P}(A|\tilde{X}_0 = (s, x)) \doteq P(A_s|\overrightarrow{X}_s = x)$ and the transition probabilities are given by $\tilde{P}(\tilde{X}_t \in B|\tilde{X}_0 = \tilde{x}) = \tilde{P}(\tilde{X}_t \in B|\tilde{X}_0 = (s, x)) = P(\overrightarrow{X}_{s+t} \in B_{s+t}|\overrightarrow{X}_s = x)$ where $B \in \tilde{B}, \tilde{x} \in \mathbb{R}_+ \times \mathbb{R}^d$. The transition function is defined by $\tilde{P}(t, \tilde{x}, B) = P(s, x; s + t, B_{s+t})$.

Lemma A.1. Given a inhomogeneous Markov process (X_t), the space-time process $(\tilde{X}_t)$ on $(\tilde{\Omega}, \tilde{\mathcal{A}}, \tilde{P})$ is a homogeneous Markov process.

Proof. See Transformation 3.1 in [3]. □

Lemma A.2. Let $(\overrightarrow{X}_t)$ be the stochastic process with Feller evolution system $U(s,t)$ and the generator of $(\overrightarrow{X}_t)$ be $\mathcal{L}_s$. Let $\widetilde{X}_t$ be its space-time process with associated semigroup $T(t)$ by $T_t u(\tilde{x}) = \tilde{\mathbb{E}}(u(\tilde{X}_t)|\tilde{X}_0 = \tilde{x})$ for $\tilde{x} \in \mathbb{R}_+ \times \mathbb{R}^d$ and $u \in B_b(\mathbb{R}_+ \times \mathbb{R}^d)$. Then the extended generator $\tilde{\mathcal{L}}$ of T_t is given for all $u \in C_\infty([0,1] \times \mathbb{R}^d)$ satisfying some conditions,

$$\tilde{\mathcal{L}}u(\tilde{x}) = \frac{\partial}{\partial s}u(s,x) + \mathcal{L}_s u_s(x) \quad \text{where } \tilde{x} = (s,x) \text{ and } u_s(x) = u(s,x). \tag{10}$$

Proof. See Theorem 3.2 in [3]. □

A Markov process typically has a generator that takes the form

$$\mathcal{L}u(x) = \frac{1}{2}\sum_{i,j=1}^{d} a_{ij}(x)\frac{\partial^2}{\partial x_i \partial x_j}u(x) + b(x)\cdot\nabla u(x) \tag{11}$$

$$+ \int_{\mathbb{R}^d}\left(u(x+y) - u(x) - \mathbf{1}_{B_1}(y)y\cdot\nabla u(x)\right)\eta(x,dy) \tag{12}$$

where $b(x)$ is a locally bounded $\mathbb{R}^d$-valued function and (a_{ij}) is a locally bounded and $d \times m$ matrix-valued function, B_1 is the ball with a radius of one and a center of zero and η satisfies

$$\int 1 \wedge |y^2|\,\eta(x,dy) < \infty \tag{13}$$

Suppose there exist $\lambda : \mathbb{R}^d \times S \to [0,1], \hat{\gamma} : \mathbb{R}^d \times S \to \mathbb{R}^d$, and a σ-finite measure v on a measurable space $(S,\mathcal{S})$ such that

$$\eta(x,\Gamma) = \int_S \lambda(x,u)\mathbf{1}_\Gamma(\hat{\gamma}(x,u))\nu(du).$$

We decompose S into $S_1 \cup S_2$ such that $1_{S_1} = 1_{B_1}(\hat{\gamma}((s,x),y))$ and $1_{S_2} = 1_{B_1^c}(\hat{\gamma}((s,x),y))$. We can rewrite the form of the generator is

$$\mathcal{L}u(x) = \frac{1}{2}\sum_{i,j=1}^{d} a_{ij}(x)\frac{\partial^2}{\partial x_i \partial x_j}u(x)$$

$$+ b(x)\cdot\nabla u(x) + \int_S \lambda(x,u((f(x,\hat{\gamma}(x,u)) - f(x) - 1_{S_1}(u)\hat{\gamma}(x,u)\cdot\nabla f(x))\nu(du)$$

Lemma A.3. Let the generator $\mathcal{L}$ be the form of (11). Let ξ be a Poisson random measure on $[0,1] \times S \times [0,\infty)$ with mean measure $m \times \nu \times m$, and let $\tilde{\xi}(A) = \xi(A) - m \times \nu \times m(A)$. Let $(S_0, \mathcal{S}_0)$ be a measurable space, μ a σ_B-finite measure on $(S_0, \mathcal{S}_0)$ where $\sigma_B : \mathbb{R}^d \times S_0 \to \mathbb{R}^d$ satisfies $\int_{S_0} |\sigma_B(x,u)|^2\mu(du) < \infty$ and

$$a(x) = \int_{S_0} \sigma_B(x,u)\sigma_B^T(x,u)\mu(du) \tag{14}$$

Assume that for each compact $K \subset \mathbb{R}^d$,

$$\sup_{x \in K}\left(|b(x)| + \int_{S_0}|\sigma_B(x,u)|^2\mu(du) + \int_{S_1}\lambda(x,u)|\hat{\gamma}(x,u)|^2\nu(du).\right. \tag{15}$$

$$\left. + \int_{S_2}\lambda(x,u)|\hat{\gamma}(x,u)| \wedge 1\nu(du)\right) < \infty \tag{16}$$

Then $\vec{X}$ satisfies a stochastic differential equation of the form

$$\vec{X_t} = \vec{X_0} + \int_{S_0}\int_0^t \sigma_B(\vec{X_s},u)W(du \times ds) + \int_0^t b(\vec{X_s})ds \tag{17}$$

$$+ \int_{v=0}^{v=1}\int_{u\in S_1}\int_{s=0}^{s=t}\mathbf{1}_{[0,\lambda(\vec{X_s},u)]}(v)\hat{\gamma}(\vec{X_s},u)\tilde{\xi}(dv \times du \times ds) \tag{18}$$

$$+ \int_{v=0}^{v=1}\int_{u\in S_2}\int_{s=0}^{s=t}\mathbf{1}_{[0,\lambda(\vec{X_s},u)]}(v)\hat{\gamma}(\vec{X_s},u)\xi(dv \times du \times ds), \tag{19}$$

240 *Proof.* See Theorem 2.3 in [15]. □

241 **Lemma A.4.** Let $\lambda((s,x),y) = \frac{p_s(x+y)}{p_s(x)}\sigma_L^\alpha(s)$ for $\sigma_L(s) \geq 0$ and $\hat{\gamma}((s,x),y)$ be $(0,y)$ and $\nu(dy)$ be
242 a Lévy measure such that it is a Borel measure on $\mathbb{R}^d$ and $\nu(\{0\}) = 0$ and $\int_{\mathbb{R}^d}(|x|^2 \wedge 1)\nu(dx) < \infty$.
243 If $(\vec{X}_t)$ has the corresponding generator $\mathcal{L}_t$

$$\mathcal{L}_t u(x) = b(x) \cdot \nabla u(x) + \int_{\mathbb{R}^d} [u(x+y) - u(x) - y \cdot \nabla u(x)1_{S_1}(y)]\frac{p_t(x+y)}{p_t(x)}\sigma_L^\alpha(t)\nu(dy) \quad (20)$$

244 where $u \in B_b(\mathbb{R}^d)$. Then the corresponding generator $\tilde{\mathcal{L}}$ of the space-time process $\tilde{X}_t$ is

$$\tilde{\mathcal{L}}u(s,x) = (1,b(x)) \cdot \nabla u(s,x) \tag{21}$$

$$+ \int_{\mathbb{R}^d} [u((s,x) + \hat{\gamma}((s,x),y)) - u(s,x) - \gamma((s,x),y) \cdot \nabla u(s,x)1_{S_1}(y)]\lambda((s,x),y)\nu(dy)$$
$$\tag{22}$$

245 where $u \in C_\infty([0,1] \times \mathbb{R}^d)$.

Proof.

$$\tilde{\mathcal{L}}u(s,x) = \frac{\partial}{\partial s}u(s,x) + \mathcal{L}_s u_s(x) \quad \text{for } u_s(x) = u(s,x)$$

$$= \frac{\partial}{\partial s}u(s,x) + b(x) \cdot \nabla_x u_s(x) + \int [u_s(x+y) - u_s(x)$$

$$- y \cdot \nabla_x u_s(x)1_{S_1}(y)]\frac{p_t(x+y)}{p_t(x)}\sigma_L^\alpha(t)\nu(dy)$$

$$= (1,b(x)) \cdot \nabla u(s,x) + \int [u(s,x+y) - u(s,x) - (0,y) \cdot \nabla u(x)1_{S_1}(y)]\frac{p_t(x+y)}{p_t(x)}\sigma_L^\alpha(t)\nu(dy)$$

$$= (1,b(x)) \cdot \nabla u(s,x) + \int [u((s,x) + (0,y)) - u(s,x)$$

$$- (0,y) \cdot \nabla u(x)1_{S_1}(y)]\frac{p_t(x+y)}{p_t(x)}\sigma_L^\alpha(t)\nu(dy)$$

246 □

247 Now, we consider the Lévy measure ν of the symmetric α-stable distribution such that $\nu(dy) =$
248 $\frac{C}{|y|^{1+\alpha}}dy$ with $C = \frac{\Gamma(\alpha+1)\sin(\alpha\pi/2)}{\pi}$.
249 **Theorem A.5.** Let a generator $\mathcal{L}_t$ has a jump kernel driven by the symmetric α-Lévy process
250 represented by (20). Let $N(dt,dx)$ is another Poisson measure on $[0,\infty) \times \mathbb{R}^d$ with mean measure
251 $m \times \nu$. Let ξ be a Poisson random measure on $\mathbb{R}_+ \times \mathbb{R}^d \times [0,\infty)$ with $\mathbb{E}[\xi(dv \times dy \times ds)] =$
252 $dm \times \nu(dy) \times dm$ and $\tilde{\xi}(A) = \xi(A) - m \times \nu \times m(A)$. Then the SDE representation of the generator
253 $\tilde{\mathcal{L}}$ satisfies

$$\vec{X}_t = \vec{X}_0 + \int_0^t b(s,\vec{X}_s)ds + \int_{v=0}^{v=t}\int_{y\in S_1}\int_{s=0}^{s=\frac{p_s(y+\vec{X}_s)}{p_s(\vec{X}_s)}\sigma_L^\alpha(s)} y \cdot \tilde{\xi}(dv \times dy \times ds) \tag{23}$$

$$+ \int_{v=0}^{v=t}\int_{y\in S_2}\int_{s=0}^{s=\frac{p_s(y+\vec{X}_s)}{p_s(\vec{X}_s)}\sigma_L^\alpha(s)} y \cdot \xi(dv \times dy \times ds) \tag{24}$$

$$= \vec{X}_0 + \int_0^t b(s,\vec{X}_s)ds + \int_0^t \sigma_L(s)dL_s^\alpha \tag{25}$$

254 *Proof.* We know λ satisfies

$$\int_{\mathbb{R}} \lambda((s,x),y)1_{S_1}(y)|r((s,x),y))|^2 + 1_{S_2}(y)\nu(dy) \tag{26}$$

$$= \int_{S_1}\left[\frac{p_s(x+y)}{p_s(x)}\sigma_L^\alpha(s)|y|^2\nu(dy)\right]dy + \int_{S_2}\frac{p_s(x+y)}{p_s(x)}\sigma_L^\alpha(s)\nu(dy) < \infty \tag{27}$$

since $\int_{S_1} |y|^2 \nu(dy) < \infty$ and $\int_{S_2} \nu(dy) < \infty$ with $\sup_{x,y,s} \frac{p_s(x+y)}{p_s(y)} \sigma_L^\alpha(s) < \infty$. Since we set $a(s,x) = 0$, $\sigma_B((s,x),y)$ is 0. Therefore, for any compact set $K \subset \mathbb{R}^2$,

$$\sup_{(s,x) \in K} \Big(|b(x)| + \int_{S_1} \lambda((s,x),y)|r((s,x),y)|^2 \nu(du)$$
$$+ \int_{S_2} \int \lambda((s,x),y)|r((s,x),y)| \wedge \nu(du) \Big) < \infty$$

since $\int_{S_1} \lambda((s,x),y)|\hat{\gamma}((s,x),y)|^2 \nu(du) + \int_{S_2} \int \lambda((s,x),y)|\hat{\gamma}((s,x),y)| \wedge \nu(du))$ is well-defined and continuous with respect to (s,x) and $b(s,x)$ is locally bounded $\mathbb{R}$-valued function. We can apply Lemma $A.2$ to the transformed homogeneous generator $\tilde{\mathcal{L}}$ of the inhomogeneous generator $\mathcal{L}_t$ from Lemma $A.4$. Now, we define $Y_t = \int_{v=0}^{v=t} \int_{y \in S_1} \int_{s=0}^{s=\frac{p_s(y+\overrightarrow{X}_s)}{p_s(\overrightarrow{X}_s)}\sigma_L^\alpha(s)} y \cdot \tilde{\xi}(dv \times dy \times ds) + \int_{v=0}^{v=t} \int_{y \in S_2} \int_{s=0}^{s=\frac{p_s(y+\overrightarrow{X}_s)}{p_s(\overrightarrow{X}_s)}\sigma_L^\alpha(s)} y \cdot \xi(dv \times dy \times ds)$ and $Z_t = \int_0^t \sigma_L(s) dL_s^\alpha$. If we show $\mathbb{E}[\exp(i(u,Y_t))] = \mathbb{E}[\exp(i(u,Z_t))]$, then we can conclude $dX_t = b(t,X(t))dt + \sigma_L^\alpha(t)dL_t^\alpha$.

$$\mathbb{E}[\exp(i(u,Y_t))] = \mathbb{E}\Big[\exp(i(u, \int_{v=0}^{v=t} \int_{y \in S_1} \int_{s=0}^{s=\frac{p_s(y+\overrightarrow{X}_s)}{p_s(\overrightarrow{X}_s)}\sigma_L^\alpha(s)} y \cdot \tilde{\xi}(dv \times dy \times ds)$$
$$+ \int_{v=0}^{v=t} \int_{y \in S_2} \int_{s=0}^{s=\frac{p_s(y+\overrightarrow{X}_s)}{p_s(\overrightarrow{X}_s)}\sigma_L^\alpha(s)} y \cdot \xi(dv \times dy \times ds))\Big]$$
$$= \mathbb{E}\Big[\exp(i(u, \int_{v=0}^{v=t} \int_{y \in S_1} \int_{s=0}^{s=\sigma_L^\alpha(s)} y \cdot \tilde{\xi}(dv \times dy \times ds)$$
$$+ \int_{v=0}^{v=t} \int_{y \in S_2} \int_{s=0}^{s=\sigma_L^\alpha(s)} y \cdot \xi(dv \times dy \times ds))\Big]$$
$$= \mathbb{E}[\exp(i(u,Z_t))]$$

$\square$

So far, we have proven that SDE representations can be found for inhomogeneous Markov processes that satisfy certain conditions. Afterward, we will examine how the time reversal of a generator appears when a homogeneous Markov process is given. Then Lemma A.5 is used to obtain the score-based reverse formula of Lévy-driven SDE. We use the reverse formula of Theorem 5.7 in [5] to propose a new class of generative model, LIM.

Lemma A.6. Consider a Markov process $(\overrightarrow{X}_t)_{t \in [0,1]}$ with generator $\mathcal{L}_t$ defined for any function u in set of continuous functions with compact support $C_c^1(\mathbb{R}^d)$ such that $\mathcal{L}_t u(x) = b(t,x) \cdot \nabla u(x) + \int_{\mathbb{R}^n} [u(y) - u(x) - \nabla u(x) \cdot [y-x]^\delta] \overrightarrow{J}_{t,x}(dy)$, $(t,x) \in [0,T] \times \mathbb{R}^n$ for some $\delta > 0$, where $b(t,x)$ is a vector field, and the jump kernel is $\overrightarrow{J}_{t,x}(dy)$. Let $[x]^\delta \doteq 1_{|x| \le \delta}x$ Then, under some hypotheses, the Markov generator $\overleftarrow{\mathcal{L}}_t$ of the time-reversed process is

$$\overleftarrow{\mathcal{L}}_t u(x) = \overleftarrow{b}(t,x) \cdot \nabla u(x) + \int_{\mathbb{R}^n} \int [u(y) - u(x) - \nabla u(x) \cdot [y-x]^\delta \overleftarrow{J}_{t,x}(dy) \quad (28)$$

where $\mathbf{p}_t(dy)$ is a marginal distribution of $(\overrightarrow{X}_t)_{t \in [0,1]}$ such that it satisfies $\mathbf{p}_t(dy)\overleftarrow{J}_{t,x}(dx) = \mathbf{p}_t(dx)\overrightarrow{J}_{t,x}(dy)$ for almost every t and the backward drift $\overleftarrow{b}(t,x)$ is given by

$$b(t,x) + \overleftarrow{b}(t,x)(t,x) = \int_{\mathbb{R}^n} [y-x]^\delta (\overrightarrow{J}_{t,x} + \overleftarrow{J}_{t,x})(dy) \quad \mathbf{p_t} - \text{a.e.} \quad (29)$$

Proof. See Theorem 5.7 in [5] $\square$

Corollary A.6.1. If we assume the marginal distribution has the density function $p_t(x)$ such that $\mathbf{p}_t(dx) = p_t(x)dx$ and $\overrightarrow{J}_{t,x}(dy)$ is a symmetric kernel with $\overrightarrow{J}_{t,x}(dy) = v_t(y-x)dy$ for some Lévy measure v_t that is a Borel measure and $v_t(\{0\}) = 0$ and $\int 1 \wedge |y|^2 v_t(dy) < \infty$ for each t. Then $\overleftarrow{J}_{t,x}(dy) = \frac{p_t(x)}{p_t(y)}v_t(y-x)dy$. Therefore (29) satisfies $\int_{|y|\leq\delta} y \cdot \frac{p_t(x)}{p_t(y)}v_t(y)dy$.

The following introduces the necessary lemmas before using Corollary A.6.1.

Lemma A.7. For $1 \leq \alpha < 2$, $|p_t^{(n)}(x)| \leq \frac{((-1)^n+1)}{2\pi}\Gamma(n+1)$.

Proof.

$$\begin{aligned}
|p_t^{(n)}(x)| &\leq \int \left|\frac{e^{-|u|^\alpha}(-iu)^n}{2\pi}e^{-iux}\right| du = \frac{1}{2\pi}\int e^{-|u|^\alpha}u^n du \\
&= \frac{1}{2\pi}\int e^{-|u|^1}u^n du \quad \text{since } 1 \leq \alpha < 2 \\
&= \frac{(-1)^n+1}{2\pi}\Gamma(n+1)
\end{aligned}$$

$\square$

Lemma A.8. For $1 \leq \alpha < 2$, $n \in \mathbb{N}$, and $0 \leq \sup_t \sigma_L(t) < 1$

$$\sum_{n=3,\text{ odd}}^{\infty} \frac{2 \cdot p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1} = O(\sigma_L^4(t)) \tag{30}$$

Proof.

$$\begin{aligned}
&\left|\sum_{n=3,\text{ odd}}^{\infty} \frac{2 \cdot p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1}\right| \\
&\leq \sum_{n=3,\text{ odd}}^{\infty} \left|\frac{2\Gamma(n+1)}{n!\pi p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1}\right| \quad \text{by Lemma } A.7 \\
&= \sum_{n=3,\text{ odd}}^{\infty} \left|\frac{2}{\pi p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1}\right| \\
&= \sum_{k=1}^{\infty} \left|\frac{2}{\pi p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{2k+21}}{2k+2-\alpha}\sigma_L(t)^2\right| \\
&\leq \frac{2}{\pi}\frac{1}{p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\frac{1}{4-\alpha}\frac{\sigma_L^4(t)}{1-\sigma_L^2(t)} = O(\sigma_L^4(t))
\end{aligned}$$

$\square$

Lemma A.9. If a stochastic process $(\overrightarrow{X}_t)$ is a solution to $d\overrightarrow{X}_t = -\frac{\beta(t)}{\alpha}\overrightarrow{X}_t + (\beta(t))^{1/\alpha}dL_t^\alpha$ then the jump kernel of $\overrightarrow{X}_t$ is

$$\overrightarrow{J}_t(x,dy) = \frac{\Gamma(\alpha+1)\sin(\alpha\pi/2)}{\pi}\frac{\sigma_L^\alpha(t)dy}{|y-x|^{\alpha+1}} \tag{31}$$

Proof. See Lemma 4 in [20]

$\square$

Therefore, when $\overrightarrow{X}_t$ is a solution to $d\overrightarrow{X}_t = -\frac{\beta(t)}{\alpha}\overrightarrow{X}_t + (\beta(t))^{1/\alpha}dL_t^\alpha$, the Markov generator $\overleftarrow{\mathcal{L}}_t$ of the time-reversed process $(\overleftarrow{X}_t)$ is the form of (20). So, we can use Theorem A.5 to $\overleftarrow{\mathcal{L}}_t$.

Theorem A.10. If $d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_L(t)dL_t^\alpha$ then the reverse SDE with respect to backward integral is $d\overleftarrow{X}_t = -\overleftarrow{b}(t, \overleftarrow{X}_t)dt + \sigma(t)d\bar{L}_t^\alpha$ with $\overleftarrow{b}$ satisfying

$$\left(\overleftarrow{b}(t, x) + b(t, x)\right) = 2\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^2(t)}{(2-\alpha)}\nabla_x \log p_t(x) + O(\sigma_L^4(t)) \qquad (32)$$

Proof.

$$\left(b(t, x) + \overleftarrow{b}(t, x)\right) = \int_{|y|<\sigma_L(t)} y\left(1 + \frac{p_t(y+x)}{p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\frac{\sin(\alpha\pi/2)}{|y|^{\alpha+1}}\sigma_L(t)^\alpha\right)dy$$

$$= \frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\sigma_L^\alpha(t)\int_{|y|<\sigma_L(t)} y\frac{p_t(y+x)}{p_t(x)}\frac{dy}{|y|^{\alpha+1}}$$

$$= \frac{\Gamma(\alpha+1)}{\pi}\frac{\sin(\alpha\pi/2)}{p_t(x)}\sigma_L^\alpha(t)\int_{|y|<\alpha}\frac{y}{|y|^{\alpha+1}}\sum_{n=0}^\infty\frac{p_t^{(n)}(x)y^n}{n!}dy$$

$$= \sum_{n=0}^\infty\frac{p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\sigma_L^\alpha(t)\int_{|y|<\sigma_L(t)}\frac{y^{n+1}}{|y|^{\alpha+1}}dy$$

$$= \sum_{n=0,\text{odd}}^\infty\frac{2\cdot p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\sigma_L^\alpha(t)\int_0^{\sigma_L(t)} y^{n-\alpha}dy$$

$$= \sum_{n=0,\text{odd}}^\infty\frac{2\cdot p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1}$$

$$= \frac{2\cdot p_t^{(1)}(x)}{p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^2}{2-\alpha} + \sum_{n=3,\text{odd}}^\infty\frac{2\cdot p_t^{(n)}(x)}{n!p_t(x)}\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^{n+1}}{n-\alpha+1}$$

$$= 2\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^2(t)}{(2-\alpha)}\nabla_x \log p_t(x) + O(\sigma_L^4(t)) \quad \text{by Lemma 30}$$

$\square$

Theorem A.11 (The general reverse SDE). The reverse SDE of $d\overrightarrow{X}_t = b(t, x)dt + \sigma_B(t)dB_t + \sigma_L(t)dL_t^\alpha$ is

$$d\overleftarrow{X}_t = \left(b(t, \overleftarrow{X}_t) - [\sigma_B^2(t) + C_\alpha\sigma_L^2(t)]\nabla_x \log p_t(\overleftarrow{X}_t)\right)dt + \sigma_B(t)d\bar{B}_t + \sigma_L(t)d\bar{L}_t^\alpha \qquad (33)$$

where $C_\alpha = \frac{\Gamma(\alpha+1)}{\frac{\pi}{2}(2-\alpha)}\sin(\frac{\pi}{2}(2-\alpha))$ and $\bar{B}_t, \bar{L}_t^\alpha$ is a corresponding backward Brownian motion and backward symmetric α-stable Lévy motion, respectively.

Proof. We denote the generator of $d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_B(t)dB_t + \sigma_L(t)dL_t^\alpha$ as $\mathcal{L}_t$ and the generator can be divided into two generators $\mathcal{L}_t^1$ and $\mathcal{L}_t^2$, for any function u in $C_c^1(\mathbb{R}^d)$ such that $\mathcal{L}_t^1(x) = \frac{1}{2}\sigma_B^2(t)\Delta u(x)$ and $L_t^2(x) = b(t, x)\cdot\nabla u(x) + \int_{\mathbb{R}_*^n}[u(y) - u(x) - \nabla u(x)\cdot[y-x]^\delta]\overrightarrow{J}_{t,x}(dy)$ with $\mathcal{L}_t = \mathcal{L}_t^1 + \mathcal{L}_t^2$ since B_t and L_t^α is independent. The reverse of Lévy-driven SDE can be expressed as backward SDE again[27]. In that case, the SDE will be determined again by the drift term, diffusion term, and jump kernel. However, the jump kernel is fully determined from the reverse of $\mathcal{L}_t^2$. By the same logic, the diffusion term of reverse is fully determined by $\mathcal{L}_t^1$, and diffusion and jump are still independent when we reverse the SDE. Therefore, we can find the generator of $\overleftarrow{\mathcal{L}}_t$ by finding out the time reversal of generators $\mathcal{L}_t^1$ and $\mathcal{L}_t^2$, respectively, which mean $\overleftarrow{\mathcal{L}}_t = \overleftarrow{\mathcal{L}}_t^1 + \overleftarrow{\mathcal{L}}_t^2$. Thus, we just need

to know $\mathcal{L}_t^2$ as we already know the SDE of $\mathcal{L}_t^1$. From Theorem A.10,

$$-\overleftarrow{b}(t,x) = b(t,x) + 2\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^2(t)}{(2-\alpha)}\nabla_x \log p_t(x) + O(\sigma_L^4(t))$$

$$\approx b(t,x) + 2\frac{\Gamma(\alpha+1)}{\pi}\sin(\alpha\pi/2)\frac{\sigma_L^2(t)}{(2-\alpha)}\nabla_x \log p_t(x) \quad \text{for small } \sigma_L$$

$$= b(t,x) - 2\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{\sigma_L^2(t)}{(2-\alpha)}\nabla_x \log p_t(x)$$

$\square$

B Probability ODE

This chapter introduces the fractional Fokker-Planck equation, which is an extended version of the Fokker-Planck equation in BM into a symmetric α-stable distribution and obtains the existence of probability ODEs from the equation. By deriving the probability ODE with the fractional derivative, the computational formula is obtained by using first-order approximation. In order to prove the existence of probability ODE, we first define fractional calculus.

B.1 Fractional Calculus

Fractional calculus is a concept that extends the existing differentiation and has the characteristic that it satisfies (36) when Fourier transformation is performed.

Definition B.1 (Multivariate Grünwald-Letnikov Fractional Derivative). For $\alpha \in (0,2)$ and $x = (x_1, \ldots, x_d) \in \mathbb{R}^d$, we define multivariate Grünwald-Letnikov fractional derivative as follows:

$$\mathbb{D}_{x_i}^\alpha f(x) = \lim_{h\to 0} \frac{\Delta_{h,x_i}^\alpha f(x)}{h^\alpha} \tag{34}$$

$$\Delta_{h,x_i}^\alpha f(x) = \sum_{j=-\infty}^{\infty} (-1)^j \frac{\Gamma(\alpha+1)}{\Gamma\left(\frac{\alpha}{2}-j+1\right)\Gamma\left(\frac{\alpha}{2}+j+1\right)} f(x_1, \ldots, x_i - jh, \ldots, x_d) \tag{35}$$

Its Fourier transform is given by:

$$\mathcal{F}\left[(D_{c1}^\alpha f)(x_1, \ldots, x_j, \ldots, x_d)\right](k) = |k_i|^\alpha \hat{f}(k_1, \ldots, k_d) \text{ for } k = (k_1, \ldots, k_d) \tag{36}$$

Proposition B.1. Let $\partial_{|x|}^\alpha f(x) \equiv (D_{cx_1}^{\alpha_1} f)(x) + \cdots + (D_{cx_d}^{\alpha_d} f)(x)$ for $\alpha = (\alpha_1, \ldots, \alpha_d)$. Then $\mathcal{F}[(\partial_{|x|}^\alpha f)(x)] = \sum_{i=1}^d |k_i|^{\alpha_i} \hat{f}(k_1, \ldots, k_d)$. If $\alpha = (0, \ldots, \alpha, \ldots, 0)$ then $\partial_{|x|}^\alpha f(x_1, \ldots, x_d) = \partial_{|x_i|}^\alpha f(x_1, \ldots, x_d)$

Proposition B.2. $\partial_{|x_i|}^\alpha f(x) = -\partial_{x_i}^2 \partial_{|x_i|}^{\alpha-2} f(x) = -\partial_{x_i} \partial_{|x_i|}^{\alpha-2} \partial_{x_i} f(x)$

Proof.

$$\mathcal{F}(\partial_{|x_i|}^\alpha f(x)) = |k_i|^\alpha \hat{f}(k) = |k_i|^2 |k_i|^{\alpha-2} \hat{f}(k) = \mathcal{F}(-\partial_{x_i} \partial_{|x_i|}^{\alpha-2} \partial_{x_i} f(t,x)) \tag{37}$$

$\square$

B.2 Stochastic Calculus for Lévy-driven Stochastic Differential Equations

Lemma B.1 (Fractional Fokker-Planck equation). Given a Lévy-driven SDE, $d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma(t)dL_t^\alpha$ for $dL_t^\alpha = (dL_1^\alpha, \ldots, dL_d^\alpha)$ with set of independent symmetric alpha stable distributions $(L_i^\alpha)_{i=1}^d$. Then the marginal distribution $p_t(x)$ satisfies fractional-Fokker-Planck equation

$$\frac{\partial p_t(x_t)}{\partial t} = -\nabla \cdot [b(t,x)p_t(x_t)] - \sigma_L(t)^\alpha \partial_{|x_t|}^\alpha p_t(x_t) \tag{38}$$

Proof. See Proposition 1 in [28]

$\square$

Corollary B.1.1 (General Fractional Fokker-Planck equation). Given a Lévy-driven SDE and, $\overrightarrow{X}_t \in \mathbb{R}^d$ which satisfies

$$d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_B(t)dB_t + \sigma_L(t)dL_t^\alpha \tag{39}$$

where $dB_t = (dB_1, \ldots, dB_d)$ with set of independent BMs $(B_i)_{i=1}^d$, and $dL_t^\alpha = (dL_1^\alpha, \ldots, dL_d^\alpha)$ with set of independent symmetric alpha stable distributions $(L_i^\alpha)_{i=1}^d$. Then the marginal distribution $p_t(x)$ satisfies General fractional-Fokker-Planck equation

$$\frac{\partial p_t(x_t)}{\partial t} = -\nabla \cdot [b(t, x)p_t(x_t)] + \sigma_B^2(t) \sum_{i=1}^d \frac{\partial^2 p_t(x_t)}{\partial x_i{}^2} - \sigma_L^\alpha(t)\partial_{|x_t|}^\alpha p_t(x_t) \tag{40}$$

Theorem B.2 (Existence of Probability ΨDE). If $p_t(x)$ satisfies Fractional Fokker-Planck equation then $p_t(x)$ satisfies

$$\frac{\partial p_t(x_t)}{\partial t} = -\nabla \cdot [(b(t, x) - \sigma_L^\alpha(t)F(t, x_t))p_t(x_t)] \tag{41}$$

such that $F_i(t, x_t) = \frac{\partial_{|x_i|}^{\alpha-2}\partial_{x_i}p_t(x_t)}{p_t(x_t)}$. So X_t satisfies the ODE,

$$d\overrightarrow{X}_t = (b(t, x) - \sigma_L\alpha(t)^F(t, x_t))dt \tag{42}$$

Proof.

$$\frac{\partial p(t, x_t)}{\partial t} = -\sum_{i=1}^d \partial_{x_i}(b_i(t, x)p(t, x_t)) - \sum_{i=1}^d \sigma_L^\alpha(t)\partial_{|x_i|}^\alpha p(t, x_t) \tag{43}$$

$$= -\sum_{i=1}^d \left[\partial_{x_i}b_i(t, x)p(t, x_t) + \sigma_L^\alpha(t)\partial_{|x_i|}^\alpha p(t, x_t) \right] \tag{44}$$

$$= -\sum_{i=1}^d \left[\partial_{x_i}b_i(t, x)p(t, x_t) - \sigma_L^\alpha(t)\partial_{x_i}\partial_{|x_i|}^{\alpha-2}\partial_{x_i}p(t, x_t) \right] \tag{45}$$

$$= -\sum_{i=1}^n \partial_{x_i} \left(\left[b_i(t, x) - \sigma_L^\alpha(t)\frac{\partial_{|x_i|}^{\alpha-2}\partial_{x_i}p(t, x_t)}{p(t, x_t)} \right] p(t, x_t) \right) \tag{46}$$

$$= -\nabla \cdot [(b(t, x) - \sigma_L^\alpha(t)F(t, x))p(t, x_t)] \tag{47}$$

$\square$

Therefore, we proved the existence of the probability ODE given an SDE driven by Lévy process. A noteworthy feature here is that the newly added $F(t, x)$ depends only on the marginal distribution and the jump coefficient, $\sigma_L(t)$. However, we can not exactly calculate fractional derivatives. Therefore, based on (33) and Theorem B.2, we can obtain a computational probability ODE by first-order approximation of $F(t, x)$.

Theorem B.3 (The Probability ODE). The probability ODE of $d\overrightarrow{X}_t = b(t, \overrightarrow{X}_t)dt + \sigma_L(t)dL_t^\alpha$ is

$$d\overrightarrow{X}_t = \left(b(t, \overrightarrow{X}_t) - \nabla_x \log p_t(\overrightarrow{X}_t)\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{\sigma_L^2(t)}{2-\alpha} \right) \tag{48}$$

C General OU process

Given a SDE driven by a symmetric α-stable Lévy process $L_t^\alpha \sim S_\alpha(t^{1/\alpha}, 0, 0)$ such that

$$d\overrightarrow{X}_t = -\beta\overrightarrow{X}_tdt + (\alpha \cdot \beta)^{1/\alpha}dL_t^\alpha \tag{49}$$

the solution of the SDE is

$$\overrightarrow{X}_t = x_0 e^{-\beta t} + (\alpha \cdot \beta)^{1/\alpha}\int_0^t e^{-\beta(t-s)}dL_s^\alpha \tag{50}$$

350 Since the integral $\int_0^t e^{-\beta(t-s)} dL_s^\alpha$ is also a symmetric α-stable $\sim \mathcal{S}\alpha\mathcal{S}(\gamma(t))$ for some $\gamma(t)$ as L_t^α is

351 a symmetric α-stable. We want to find the scale parameter $\gamma(t)$ of $\int_0^t e^{-\beta(t-s)} dL_s^\alpha$ for each t.

Lemma C.1. Given α with $0 < \alpha < 2$ and f is a measurable function such that $f : [0, T] \to \mathbb{R}$ with

353 $\int_0^T |f(s)|^\alpha ds < \infty$. Let $\vec{X}_t = \int_0^t f(s) dL_s^\alpha$ then

$$\vec{X}_t \sim \mathcal{S}\alpha\mathcal{S} \left(\int_0^t |f(s)|^\alpha ds)^{1/\alpha} \right) \tag{51}$$

354 *Proof.* If $f(t) = \sum_{i=1}^N a_i \chi_{(t_{i-1}, t_i]}$ with $t_0 = 0, t_N = t$,

$$\vec{X}_t = \int_0^t \sum_{i=1}^N a_i \chi_{(t_{i-1}, t_i]}(s) dL_s^\alpha = \sum_{i=1}^n a_i [L_{t_i}^\alpha - L_{t_{i-1}}^\alpha] \overset{d}{=} \sum_{i=1}^N a_i L_{\Delta t_i}^\alpha, \quad \Delta t_i = t_i - t_{i-1} \tag{52}$$

355 Using the above equation,

$$\mathbb{E}[e^{(iu\vec{X}_t)}] = \mathbb{E}[e^{iu \sum_{i=1}^N a_i L_{\Delta t_i}^\alpha}] = \prod_{i=1}^N \mathbb{E}[e^{iu a_i L_{\Delta_{i}}^\alpha}] \tag{53}$$

$$= \prod_{i=1}^N e^{-|u|^\alpha |a_i|^\alpha \Delta t_i} = e^{-\sum_{i=1}^N |a_i|^\alpha \Delta t_i |u|^\alpha} = e^{-(\int_0^t |f(s)|^\alpha ds)|u|^\alpha} \tag{54}$$

356 $\Rightarrow \vec{X}_t \sim \mathcal{S}\alpha\mathcal{S} \left(\int_0^t |f(s)|^\alpha ds)^{1/\alpha} \right)$. Let us prove that f is not a simple function. With loss of

357 generality, assume $f(t) \geq 0$. If not, we decompose $f(t) = f^+(t) - f^-(t)$ such that f^+, f^- are

358 non-negative functions. Then we can construct an non-decreasing sequence of simple functions

359 f_n such that $\lim_{n\to\infty} f_n(t) = f(t)$ for all $t \in [0, T]$. So, $\sup_n f_n(t) \leq f(t)$ for all t. Define

360 $X_t^n \equiv \int_0^t f_n(s) dL_s^\alpha$. As $\int_0^T |f(s)|^\alpha ds < \infty$, we can use dominated convergence theorem so that

361 $\lim_{n\to\infty} \vec{X}_t^n(w) = \vec{X}_t(w)$ for all $w \in \Omega, t \in [0, T]$.

$$\mathbb{E}[e^{(iu\vec{X}_t)}] = \lim_{n\to\infty} \mathbb{E}[e^{(iuX_t^n)}] = \lim_{n\to\infty} e^{-(\int_0^t |f_n(s)|^\alpha ds)|u|^\alpha} = e^{-(\int_0^t |f(s)|^\alpha ds)|u|^\alpha} \tag{55}$$

362 $\therefore X_t \sim \mathcal{S}\alpha\mathcal{S} \left(\int_0^t |f(s)|^\alpha ds)^{1/\alpha} \right)$ when f is a measurable function. $\square$

Theorem C.2. If $a(t) = e^{-\beta t}, \gamma(t) = (1 - e^{-\alpha\beta t})^{1/\alpha} = (1 - (a(t))^\alpha)^{1/\alpha}$ and $\vec{X}_t = a(t)x_0 + \gamma(t)\epsilon$

364 for some $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$ then X_t is a solution to $d\vec{X}_t = -\beta\vec{X}_t dt + (\alpha \cdot \beta)^{1/\alpha} dL_t^\alpha$ and

$$\vec{X}_t \overset{d}{=} x_0 e^{-\beta t} + (\alpha \cdot \beta)^{1/\alpha} \int_0^t e^{-\beta(t-s)} dL_s^\alpha \tag{56}$$

365 *Proof.* Use Lemma C.1. $\square$

Lemma C.3. If $\vec{X}_t$ is a solution to $d\vec{X}_t = -\frac{\beta(t)}{\alpha}\vec{X}_t dt + \beta(t)^{1/\alpha} dL_t^\alpha$, then X_t can be represented

367 by

$$\vec{X}_t = e^{-\int_0^t \frac{\beta(s)}{\alpha} ds} \vec{X}_0 + \int_0^t e^{-\int_u^t \frac{\beta(s)}{\alpha} ds} \beta(u)^{1/\alpha} dL_u^\alpha \tag{57}$$

368 If we define $a(t) = e^{-\int_0^t \frac{\beta(s)}{\alpha} ds}$, then the scale parameter $\gamma(t)$ of $\int_0^t e^{-\int_u^t \frac{\beta(s)}{\alpha} ds} (\beta(u)^{1/\alpha} dL_t^\alpha$

369 satisfies $\gamma^\alpha(t) = (1 - a^\alpha(t))$. If $\beta(t) = \beta_0 + (\beta_1 - \beta_0)t$ then $\mathbb{E}[X_t] = e^{-\frac{(\beta_1-\beta_0)}{2\alpha}t^2 - \frac{\beta_0 t}{\alpha}} x_0 = a(t)x_0$,

370 with $\log a(t) = -\frac{(\beta_1-\beta_0)}{2\alpha}t^2 - \frac{\beta_0}{\alpha}t$.

Proof.

$$d(e^{\int_0^t \frac{\beta(s)}{\alpha} ds}) = e^{\int_0^t \frac{\beta(s)}{\alpha} ds} \cdot \frac{\beta(t)}{\alpha} dt + e^{\int_0^t \frac{\beta(s)}{\alpha} ds}\left(-\frac{\beta(t)}{\alpha}\vec{X}_t dt + (\beta(t))^{1/\alpha} dL_t^\alpha\right)$$

$$= e^{\int_0^t \frac{\beta(s)}{\alpha} ds}(\beta(t))^{1/\alpha} dL_t^\alpha$$

$\vec{X}_t = e^{-\int_0^t \frac{\beta(s)}{\alpha} ds} X_0 + \int_0^t e^{-\int_u^t \frac{\beta(s)}{\alpha} ds} \beta(u)^{1/\alpha} dL_t^\alpha$. If we set $a(t) = e^{-\int_0^t \frac{\beta(s)}{\alpha} ds}$ then $\frac{d}{dt}\log a(t) = -\frac{\beta(s)}{\alpha}$. And the scale parameter $\gamma(t)$ satisfies

$$\gamma^\alpha(t) = \int_0^t \frac{a(t)^\alpha}{a(u)^\alpha}(\beta(u)) du = \int_0^t \frac{a^\alpha(t)}{a^\alpha(u)}(-\alpha)\frac{d}{dt}\log a(u) du = a^\alpha(t)\int_0^t \frac{-\alpha}{a^\alpha(u)}\frac{a'(u)}{a(u)} du$$

$$= a^\alpha(t)\int_0^t (-\alpha)\frac{a'(u)}{a^{\alpha+1}(u)} du = a^\alpha(t)\int_0^t \frac{d}{du}(a^{-\alpha}(u)) du = a^\alpha(t)[a^{-\alpha}(t) - a^{-\alpha}(0)]$$

$$= (1 - a^\alpha(t)).$$

$\square$

Corollary C.3.1 (Stochastic sampling of LIM). When $t < s$, $\Delta t = s - t$

$$x(t) = (1 + \frac{\beta(s)}{\alpha}\Delta t)x(s) + (\beta(s)\Delta t)^{2/\alpha}(\Delta t)^{1-2/\alpha}C_\alpha \nabla_x \log p_s(x(s)) + (\beta(s)\Delta t)^{1/\alpha}\epsilon \quad (58)$$

where $C_\alpha = \frac{\sin(\frac{\pi}{2}(2-\alpha))}{\frac{\pi}{2}\cdot(2-\alpha)}\Gamma(\alpha+1)$ and $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$.

Proof. We know the reverse SDE of $dX_t = -\frac{\beta(t)}{\alpha}X_t + (\beta(t))^{1/\alpha}dL_t^\alpha$ is

$$d\vec{X}_t = (-\frac{\beta(t)}{\alpha}\vec{X}_t - 2\nabla_x \log p_t(\vec{X}_t)\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{\beta(t)}{2-\alpha}) + (\beta(t))^{1/\alpha})d\bar{L}_t^\alpha \quad (59)$$

We can apply Euler-Maruyama method to (59). When $t < s$,

$$x(s) = x(s) + \frac{\beta(s)\Delta t}{\alpha}x(s)) + 2\nabla_x \log p_t(x(s))\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{(\beta(s)\Delta t)^{2/\alpha}}{2-\alpha}(\Delta t)^{1-2/\alpha}$$
$$+ (\beta(s)\Delta t)^{1/\alpha}\epsilon$$

where $\Delta t = |t - s|$ and for some $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$. $\square$

Corollary C.3.2 (Deterministic ODE sampling of LIM). Using Euler-Maruyama method, the deterministic ODE sampling becomes $x(t) = (1 + \frac{\beta(s)}{\alpha}\Delta t)x(s) + \frac{1}{2}(\beta(s)\Delta t)^{2/\alpha}\cdot(\Delta t)^{1-2/\alpha}\cdot\frac{\sin(\frac{\pi}{2}(2-\alpha))}{\frac{\pi}{2}\cdot(2-\alpha)}\cdot\Gamma(\alpha+1)\nabla_x \log p_s(x(s))$ for $t < s$, $\Delta t = s - t$, $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$

Proof. From Theorem B.3, we know the ODE of $dX_t = -\frac{\beta(t)}{\alpha}X_t + (\beta(t))^{1/\alpha}dL_t^\alpha$ is

$$d\vec{X}_t \approx (-\frac{\beta(t)}{\alpha}\vec{X}_t - \nabla_x \log p_t(\vec{X}_t)\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{\beta(t)}{2-\alpha}) \quad (60)$$

We apply Euler-Maruyama method to (60). When $t < s$,

$$x(s) = x(s) + \frac{\beta(s)\Delta t}{\alpha}x(s)) + \nabla_x \log p_t(x(s))\frac{\Gamma(\alpha+1)}{\pi}\sin(\frac{\pi}{2}(2-\alpha))\frac{(\beta(s)\Delta t)^{2/\alpha}}{2-\alpha}(\Delta t)^{1-2/\alpha}$$

where $\Delta t = |t - s|$. $\square$

D Score Function for Lévy-Itō Models

Lemma D.1. Let q_α be the density function of $\mathcal{S}\alpha\mathcal{S}$ and the value of X_t satisfies $x_t = a(t)x_0 + \gamma(t)\epsilon$ for given x_0 and $\epsilon \sim \mathcal{S}\alpha\mathcal{S}$. Then the score function of the transition distribution satisfies $\nabla \log p_t(x_t|x_0) = \nabla \log q_\alpha(\epsilon)/\gamma(t)$.

Proof. Let $\vec{X}_t$ and Y be defined on a probability space $(\Omega, \mathcal{A}, \mathbb{P})$ where the transition density function of $\vec{X}_t$ is $p_t(x_t|x_0) = \frac{d\mathbb{P}(\vec{X}_t \leq x_t|\vec{X}_t = x_0)}{dx_t}$ and the density function of Y is $q_\alpha(y) = \frac{d\mathbb{P}(Y \leq y)}{dy}$. Let $\vec{X}_t = a(t)x_0 + \gamma(t)\epsilon$. Then

$$\mathbb{P}(\vec{X}_t \leq x_t|\vec{X}_0 = x_0) = \mathbb{P}(a(t)x_0 + \gamma(t)Y \leq x_t) \text{ since } \vec{X}_t = a(t)x_0 + \gamma(t)Y \tag{61}$$

$$= \mathbb{P}(Y \leq \frac{x_t - a(t)x_0}{\gamma(t)}) \tag{62}$$

$$= \mathbb{P}(Y \leq \epsilon) \tag{63}$$

Then $p_t(x_t|x_0) = \frac{d\mathbb{P}(\vec{X}_t \leq x_t|\vec{X}_0 = x_0)}{dx_t} = \frac{d\mathbb{P}(Y \leq \epsilon)}{d\epsilon}/\gamma(t) = q_\alpha(\epsilon)/\gamma(t)$. If we take logarithm of both sides then $\log p(x_t|x_0) = \log q_\alpha(\epsilon) - \log \gamma(t)$. Therefore, we can get $\nabla_{x_t} \log p(x_t|x_0) = \nabla_\epsilon \log q_\alpha(\epsilon)/\gamma(t)$. $\qquad\square$

Definition D.1 (Generalized Gaussian distribution)**.** The generalized Gaussian distribution is two families of parametric probability distributions with a continuous path on $\mathbb{R}$ with a shape parameter $\tilde{\beta}$ and scale parameter $\tilde{\sigma}$ such that

$$G_{\tilde{\sigma},\tilde{\beta}}(x) = \frac{\tilde{\beta}}{2\tilde{\sigma}\Gamma(\tilde{\beta} - 1)} \exp\left(-\frac{|x|^{\tilde{\beta}}}{\tilde{\sigma}^{\tilde{\beta}}}\right) \tag{64}$$

where $\Gamma(\cdot)$ is the Gamma function.

The score function of the Generalized Gaussian distribution is

$$\nabla_x \log G_{\tilde{\sigma},\tilde{\beta}} = -\frac{\tilde{\beta}}{\tilde{\sigma}^{\tilde{\beta}}} \text{sgn}(x))|x|^{\tilde{\beta}-1} \tag{65}$$

Which is the same form of

$$\text{ReELS}_\alpha(x) = -\text{sgn}(x)\hat{c}|x|^{\hat{\beta}}$$

when $\hat{\beta} = \tilde{\beta} - 1$ and $\hat{c} = \frac{\tilde{\beta}}{\tilde{\sigma}^{\tilde{\beta}}}$.

D.1 ReELS

The principle behind the ReELS approaches to approximate the Lévy score function of the α-stable distribution is similar to that provided in [17]. Because computing the Lévy score exactly requires higher computation complexity, score functions of generalized Gaussian distribution are employed [17] as an approximation technique. ReLES is employed with a similar concept to conduct enough denoising at large noise to allow data to converge while maintaining the heavy-tailed features. We empirically observe that $\hat{\beta}$ becomes an approximation with a value less than 1 when α is less than 2. This means that distribution with a score similar to each Lévy score is a generalized Gaussian distribution where $\tilde{\beta}$ is less than 2.

D.2 Stochastic sampling for synthetic data

In order to show that the results predicted by the theory are valid, the performance of the model trained with the BM score and ReLES with synthetic data is compared with FID. Additionally, it is tested with synthetic data if the synthetic data converges to the original distribution when using ReELS for LIM. The synthetic data used as the test were two mixtures of Gaussian, Two moons, and swiss-roll. In the case of two mixtures of Gaussian, the simplest MLP with a model depth of 3 was used, and for Two Moon and swiss-roll, MLP with a model depth of 6 was used. Detailed

experimental settings are described in (Table D.1, D.2, D.3). When using different score functions for each $\alpha = 1.2, 1.5, 1.8, 2$, the generation ability of LIM was compared with FID. As a result of the experiments, the mean value and variance of FID were low for the overall α in the case of using ReELS. The stochastic sampling according to the time step when different α is given can be seen in (Figure E.3, E.8, E.9).

FID $\downarrow \backslash \alpha$	1.2	1.5	1.8
Lévy Score	304192 ± 233	25410 ± 98	1966 ± 31
ReELS	**2.01** ± 0.08	**0.34** ± 0.02	**0.14** ± 0.01
BM Score	2.61 ± 0.09	0.71 ± 0.03	0.20 ± 0.01

Table D.1: FID score (mean ±95% CI) of stochastic sampling on synthetic data (Mixture of Gaussian). The mean values of the data distributions are (5,5), (-5,-5), respectively, and the covariance is $0.2I$. The training data is 5000 pieces and the test data is 5000 pieces. As a score model, an MLP model with a depth of 3 and a channel of [3,32,2] is used. β_0 is set to 0, β_1 is set to 10, and the clamp is set to 20. It can be seen that the FID is low when ReELS is used for all $\alpha = 1.2, 1.5$, and 1.8.

FID $\downarrow \backslash \alpha$	1.2	1.5	1.8
Lévy Score	22258 ± 182	3145 ± 62	316 ± 26
ReELS	**0.85** ± 0.013	**0.21** ± 0.0053	**0.11** ± 0.0032
BM Score	0.99 ± 0.023	0.34 ± 0.0083	0.16 ± 0.0038

Table D.2: FID score (mean ±95% CI) of stochastic sampling on synthetic data (Two-moon). The noise of two-moon synthetic data was set to 0.05. The training data is 5000 pieces and the test data is 5000 pieces. As a score model, an MLP model with a depth of 6 and a channel [3,32,64,64,32,2] is used. β_0 is set to 0, β_1 is set to 5, and the clamp is set to 20. It can be seen that the FID is low when ReELS is used for all $\alpha = 1.2, 1.5$, and 1.8.

FID $\downarrow \backslash \alpha$	1.2	1.5	1.8
Lévy Score	1486 ± 3.68	197.54 ± 0.73	17 ± 0.13
ReELS	1.16 ± 0.090	**0.210** ± 0.0087	**0.114** ± 0.0019
BM Score	**0.952** ± 0.11	0.44 ± 0.017	0.210 ± 0.0087

Table D.3: FID score (mean ±95% CI) of stochastic sampling on synthetic data (Swiss-roll). The noise of Swiss-roll synthetic data was set to 0.1. The training data is 5000 pieces and the test data is 5000 pieces. As a score model, an MLP model with a depth of 6 and a channel [3,32,64,64,32,2] is used. β_0 is set to 0, β_1 is set to 5, and the clamp is set to 20. It can be seen that the FID is low when ReELS is used for all $\alpha = 1.2, 1.5$, and 1.8.

D.3 Deterministic ODE sampling for synthetic data

In this chapter, we demonstrate the validity of the probability ODE (Theorem B.3) from the deterministic ODE sampling of LIM by showing the ability to generate synthetic data. We train score models by using three synthetic data such as Two mixtures of Gaussian, Two moons, and Swiss roll. The deterministic ODE sampling according to the time step when different α is given can be seen in (Figure E.10, E.11, E.12).

FID $\downarrow \setminus \alpha$	1.2	1.5	1.8	2.0
Two mixture	30.36 ± 0.094	2.39 ± 0.0076	2.39 ± 0.0076	0.21 ± 0.0021
Two moon	35.67 ± 0.170	0.99 ± 0.028	0.99 ± 0.028	0.41 ± 0.0026
Swiss roll	30.96 ± 0.584	0.77 ± 0.0097	0.15 ± 0.0021	0.39 ± 0.0039

Table D.4: FID (mean $\pm 95\%$ CI) of deterministic ODE sampling on synthetic data (Two mixture, Two moons, Swiss-roll). The mean values of the Mixture of Gaussian distributions are (5,5), (-5,-5), respectively, and the covariance is $0.2I$. The training data is 5000 pieces and the test data is 5000 pieces. β_0 is set to 0, β_1 is set to 10, and the clamp is set to 20. It can be seen that the FID is low when ReELS is used for all $\alpha = 1.2, 1.5$, and 1.8.

E Dataset Experiment

E.1 Implementation Detail

Our diffusion model is U-Net[19] following DDPM[11], which replaces weight normalization[21] with group normalization[30] for simple implementation. We set the model size suitable for dataset, such that MNIST (28×28) is [16, 32, 64], CIFAR10 (32×32) is [128, 256, 256, 256], CelebA (64×64) is [128, 256, 256, 256, 1024], and CelebA-HQ (256×256) is [128, 256, 256, 256, 1024, 1024], but fix the number of residual blocks with 2 in each resolution level, and add self-attention block only in 16×16 resolution level. Continuous diffusion time $t \in [0, 1)$ is injected into model through Transformer sinusoidal position embedding[29] after added with 0.0001, and we use swish function as activation function.

We train our MNIST model used in experiments for 1000 epochs with batch size 128, CIFAR10 model for 250 epochs with batch size 128, CelebA model for 140 epochs with batch size 128, and CelebA-HQ model for 160 epochs with batch size 32. All training and experiments are conducted on NVIDIA A100 GPU and NVIDIA GeForce RTX 3090, and we tune the batch size for sampling adjusted for computation resources. Because the target distribution of our model is α-stable distribution, sample quality is very sensitive to hyperparameter setting according to α scale. So we improve sample quality by tuning hyperparameters to be optimized for each dataset:

- Though DDPM[11] used linear noise schedule with fixed $\beta_0 = 0.1$, $\beta_1 = 20$, we tuned β_0, β_1 for each α because variance of α-stable distribution depends on α scale. For MNIST dataset, we fixed the β schedule as $\beta_0 = 0.1$, $\beta_1 = 5$ in all α values. In CIFAR10/CelebA/CelebA-HQ, we chose $\beta_1 = 20$ for $\alpha = 1.8$, and $\beta_1 = 15$ for $\alpha = 1.5$ to optimize convergence into sample space, and fix β_0 to 0.1.

- Different from Gaussian distribution, α-stable distribution can have large-scale noise at lower α values, which leads to sample quality degradation. To prevent this problem, we used noise clamping as a heuristic in the training and sampling phase. It consists of 3 clamps, clamp(training, init sample, SDE sample), and we adjusted the scale of clamps suitable for each dataset:

  - 10, 10, 10 at $\alpha = 1.8$, and 100, 50, 100 at $\alpha = 1.5, 1.2$ for MNIST.
  - 30, 30, 30 at $\alpha = 1.8$, and 100, 50, 100 at $\alpha = 1.5$ for CIFAR10.
  - 10, 10, 10 at $\alpha = 1.8$, and 50, 50, 50 at $\alpha = 1.5$ for CelebA and CelebA-HQ.
  - In the case of deterministic ODE sampling, we only used clamp(training, init sample), and set them 10, 5.

E.2 Evaluation Metric

E.2.1 FID(Fréchet Inception Distance) score

To evaluate generated sample quality, we choose the widely used FID score metric([10]), where a lower score means better sample quality. After computing both mean/variance of distributions in the training dataset and generated 50k samples by using the pre-trained Inception-V3 model, we calculate distances between two distributions as FID score.

E.2.2 Likelihood computation

Our ReELS method is adaptive to probability ODE(Figure 4), so we can compute the exact likelihood on any input data in the same way as [26]. By replacing the score $\nabla_x \log p_t(X_t)$ with score model $S_\theta(X_t, t)$, we can rewrite (48) as

$$d\vec{X}_t = \underbrace{\left(b(t, \vec{X}_t) - S_\theta(\vec{X}_t, t) \frac{\Gamma(\alpha+1)}{\pi} \sin(\frac{\pi}{2}(2-\alpha)) \frac{\sigma_L^2(t)}{2-\alpha} \right)}_{=:\tilde{f}_\theta(\vec{X}_t, t)} dt \tag{66}$$

Then we can compute the log-likelihood of $p_0(X_0)$ such that

$$\log p_0(\vec{X}_0) = \log p_T(\vec{X}_T) + \int_0^T \nabla \cdot \tilde{f}_\theta(\vec{X}_t, t) dt \tag{67}$$

where X_T is noise mapping to X_0 which can be obtained by solving the probability ODE in (66) with ODE solver. Because of expensive computation of $\nabla \cdot \tilde{f}_\theta(\vec{X}_t, t)$, we estimate it by using the Skilling-Hutchinson trace estimator([23], [12]), which is following [8].

To solve the integral term in (66), we choose the RK45 ODE solver[7] which can be used as `solve-ivp` function in `scipy.integrate` library. As same as [8], we also set parameters `atol=1e-5`, `rtol=1e-5`. We use a test dataset applied uniform dequantization, and take the average of the bits/dim values over 5 repeats for exact likelihood computation. By changing initial time t_0 of integral $\int_{t_0}^T \nabla \cdot \tilde{f}_\theta(\vec{X}_t, t) dt$ after adding 0.001, we compute bits/dim with varied number of function evaluations(NFE) like Figure 3(b).

E.3 Additional Samples

Additional sampling results on MNIST 28×28, CIFAR10 32×32, CelebA 64×64, CelebA-HQ 256×256 are reported in below figures.

$\alpha = 2.0$ (DM) $\alpha = 1.8$ (LIM) $\alpha = 1.5$ (LIM) $\alpha = 1.2$ (LIM)

Figure E.1: Samples generated by DM(Brownian motion, [26]), and LIM($\alpha = 1.8, 1.5, 1.2$) with on MNIST(28×28) dataset.

Figure E.2: Samples generated by LIM at $\alpha = 1.8$ on CIFAR10(32×32) dataset.

Figure E.3: Samples generated by LIM at $\alpha = 1.5$ on CIFAR10(32×32) dataset.

Figure E.4: Samples generated by LIM at $\alpha = 1.8$ on CelebA(64×64) dataset.

Figure E.5: Samples generated by LIM at $\alpha = 1.5$ on CelebA(64×64) dataset.

Figure E.6: Samples generated by LIM($\alpha = 1.8$) with 0, 500, 750, 825, 900, 1000 number of function evaluations(NFE) on CelebA-HQ(256×256) dataset.

Figure E.7: Stochastic sampling(C.3.1) of two mixtures of Gaussian using ReELS for (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. The orange color represents the original distribution of two moons.

Figure E.8: Stochastic sampling(C.3.1) of two moons using ReELS for (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. The orange color represents the original distribution of two moons.

Figure E.9: Stochastic sampling(C.3.1) of swiss roll using ReELS for (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. The orange color represents the original distribution of swiss roll.

Figure E.10: Deterministic ODE sampling(C.3.2s) of Mixture of Gaussian synthetic data for (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. Unlike stochastic sampling, there is a small number of points that do not converge to modes. The existence of these points is presumed to have occurred because ODE sampling cannot directly clamp the noise size in the middle of the reverse process.

Figure E.11: Deterministic ODE sampling(C.3.2s) of Two Moons synthetic data for (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. Unlike stochastic sampling, there is a small number of points that do not converge to modes. The existence of these points is presumed to have occurred because ODE sampling cannot directly clamp the noise size in the middle of the reverse process.

Figure E.12: Deterministic ODE sampling(C.3.2s) for Swiss Roll synthetic data when (a) $\alpha = 1.2$, (b) $\alpha = 1.5$, (c) $\alpha = 1.8$, and (d) BM-driven synthetic image. Unlike stochastic sampling, there is a small number of points that do not converge to modes. The existence of these points is presumed to have occurred because ODE sampling cannot directly clamp the noise size in the middle of the reverse process.