Beyond Thresholds: Learning Perioperative Risk from Intraoperative Physiological Trajectories

Sadaf Sohrabi [1] [2] **Shehan Hettiaratchy** [2] **Aldo A. Faisal** [3]

Abstract

Clinical risk scores for surgical patients are computed once from preoperative variables and consequently fail to capture the physiological dynamics that unfold during surgery. We ask whether intraoperative vital-sign time series, routinely recorded but rarely used for risk stratification, provide predictive information that static models miss. Using 130,960 operations from the INSPIRE perioperative dataset, we compare four modelling arms across progressively richer feature regimes (static baseline, physiological summaries, HMM-derived trajectory features, and their combination) and evaluate performance across four postoperative outcomes using grouped cross-validation. We find that the HMM-derived representations capture prognostic structure complementary to the threshold-based summaries for ICU-related outcomes, with combined models performing best. Predictive signal is detectable as early as 15 minutes after anaesthesia onset, and gains are largest in patients whose static risk profile underestimates true intraoperative vulnerability. These results indicate that the multivariate structure of intraoperative physiology carries prognostic information complementary to the threshold-based per-channel summaries evaluated here, with the largest gains from combining the two.

1. Introduction

Surgery is performed at massive scale, with over 300 million procedures worldwide each year (Weiser et al., 2015). While most patients have an uneventful perioperative course, a substantial minority develop serious complications in the hours and days that follow, including unplanned intensive care unit (ICU) admission and death. Identifying which patients will deteriorate remains a central and unresolved challenge in perioperative medicine. The European Surgical Outcomes Study found that 73% of patients who died after non-cardiac surgery had never been admitted to ICU (Pearse et al., 2012), indicating that adverse outcomes often arise from failures to recognise risk in time.

Current approaches to risk identification are fundamentally static. Preoperative tools such as the Surgical Outcome Risk Tool (SORT) (Protopapa et al., 2014) assign a scalar risk estimate before the patient enters the operating room (OR), based on American Society of Anesthesiologists (ASA) physical status grade, urgency, surgical severity, specialty, cancer status, and age. While these tools perform well on average, achieving AUROC of 0.90 for 30-day mortality (Wong et al., 2020), the resulting estimate is not designed to be updated once surgery begins. It does not reflect the patient's response to anaesthesia and subsequent surgical stress. Once the operation starts, risk assessment shifts to threshold-based alarm systems that flag individual vital signs when they breach pre-defined limits. These systems treat vital signs as independent scalar quantities, discarding the multivariate temporal context that encodes the patient's evolving physiological state. As a result, they provide no representation of the physiological trajectory over the course of the procedure.

During surgery, each patient generates a continuous stream of physiological data at minute-level resolution, yet this information is rarely incorporated into predictive models and is absent from formal risk estimates. Machine learning provides a natural framework for modelling multivariate physiological time series, where clinically relevant information is encoded in temporal patterns and cross-signal dependencies rather than in isolated measurements. Prior studies using gradient-boosted models, deep neural networks, and sequence models have shown promise for postoperative risk prediction from intraoperative data (Xue et al., 2021; Fritz et al., 2019; Shickel et al., 2023; Hao et al., 2025). However, it is unclear whether predictive gains arise from detecting threshold violations or from modelling the structure of physiological trajectories. More broadly, fundamental questions

---

[1]UKRI CDT in AI for Healthcare, Imperial College London [2]Imperial College Healthcare NHS Trust [3]Department of Computing, Imperial College London. Correspondence to: Sadaf Sohrabi <ss16225@ic.ac.uk>.

Proceedings of the Workshop on Structured Data for Health at the 43rd International Conference on Machine Learning, Seoul, South Korea.

remain about what feature representations capture the most prognostic information, when the signal becomes reliable, and for whom dynamic monitoring adds most value over static preoperative assessment.

We address these questions using the INSPIRE dataset (Lim et al., 2024), the largest publicly available perioperative dataset, which comprises 130,960 operations with linked intraoperative vital signs and postoperative outcomes. We make four contributions. (1) A structured comparison of static baseline risk, intraoperative physiological summaries, and hidden Markov model (HMM)-derived trajectory features, demonstrating that each adds incremental predictive value over the previous. (2) Evidence that HMM trajectory features capture complementary predictive structure to per-channel summaries, with the combined representation performing best, indicating that prognostic signal is multivariate and contextual, rather than reducible to threshold crossings of individual signals. (3) A demonstration that predictive signal is detectable as early as 15 minutes after anaesthesia onset. (4) A finding that gains concentrate in patients whose static risk profile underestimates true physiological vulnerability, a pattern not explained by any single preoperative variable.

2. Problem Setup and Methods

Dataset. INSPIRE (Lim et al., 2024) contains 130,960 surgical operations from a single academic centre (2011–2020), with continuously recorded intraoperative vital signs, including heart rate, peripheral oxygen saturation (SpO_2), non-invasive mean arterial pressure (NIBP MAP), invasive mean arterial pressure (arterial MAP), and inspired oxygen fraction (FiO_2), alongside postoperative outcomes. The dataset is publicly available on PhysioNet and provides sufficient scale and outcome prevalence for robust evaluation. Cohort characteristics and outcome incidence are summarised in Appendix A.

Outcomes. We evaluate four binary outcomes: prolonged ICU stay >48 h (prevalence 2.2%), ICU admission (9.6%), early ICU escalation within 6 h of OR exit (9.4%), and in-hospital death (1.2%). The primary outcome is prolonged ICU stay, capturing sustained postoperative deterioration and providing a more specific marker of clinically meaningful risk than ICU admission alone. ICU admission captures all postoperative ICU transfers within 24 hours of surgery, while ICU within 6 h captures only immediate escalations. The high overlap between these definitions explains their near-identical predictive performance. Class imbalance motivates reporting both AUROC and AUPRC.

Feature regimes. We compare three feature regimes (baseline, summaries, and trajectories) and their combination, evaluated across four modelling arms. *Baseline* uses static

preoperative variables (age, sex, BMI, ASA grade, emergency status, surgical specialty, anaesthesia type, procedure class). Anaesthesia and operation duration are excluded to maintain a strictly preoperative baseline. *Summaries* augment the baseline with per-channel statistics computed over the full intraoperative period (anaesthesia onset to OR exit, where anaesthesia onset is the timestamp of first anaesthetic agent administration and OR exit is the recorded departure time from the operating room), including the mean, SD, min, max, and time beyond clinical thresholds (SpO_2 <92%, MAP <65 mmHg, heart rate <40 or >120 bpm). *Trajectories* augment the baseline with features derived from a Gaussian HMM fit to 5-minute binned multivariate intraoperative time series, replacing per-channel summaries. The four modelling arms were baseline only (A), baseline plus summaries (B), baseline plus trajectories (C), and the combined model (D).

HMM design. Six channels were used: NIBP MAP, arterial MAP, SpO_2, heart rate, FiO_2, and a binary arterial line indicator. Signals were binned into 5-minute intervals (using the within-bin mean), then standardised via a `StandardScaler` fitted on training bins after median imputation. Short gaps of up to two consecutive bins (10 minutes) were forward-filled using the last observed value (LOCF). This procedure can introduce bias and is acknowledged as a limitation. Per-channel missingness varied. SpO_2 and heart rate had <2% missing bins, FiO_2 had 16%, and NIBP MAP and arterial MAP each had ~42–44% missing per bin. The combined MAP channel, preferring arterial MAP with NIBP fallback, achieved ~82% bin coverage before LOCF. Bins with no observations across all channels were excluded, as were operations with fewer than six valid bins (30 minutes). This left 124,992 operations for the four-arm predictive comparison.

The number of states K was selected by sweeping $K \in \{3, \ldots, 8\}$ and evaluating held-out log-likelihood with a Bayesian Information Criterion (BIC)-style complexity penalty and silhouette score on a grouped 80/20 training split. Improvements in held-out log-likelihood decelerated substantially after $K=6$, and silhouette reached a local maximum at $K=6$ among $K \geq 4$. We therefore selected $K=6$ to balance data fit and state separability (Table 1).

Operation-level features comprised (i) state occupancy fractions (fraction of bins spent in each state), (ii) transition entropy $H = -\sum_{i,j} \hat{P}_{ij} \log \hat{P}_{ij}$, where $\hat{P}_{ij}$ is the observed fraction of decoded transitions from state i to state j within that operation, and (iii) transitions into and out of physiologically unstable states, defined as states whose emission means breach standard clinical thresholds (MAP < 65 mmHg, SpO_2 < 92%, heart rate < 45 or > 110 bpm, or FiO_2 > 60%).

The inclusion of an arterial line indicator may partly reflect

Table 1. K-selection sweep for the 6-channel Gaussian HMM on a grouped 80/20 held-out split. Val LL/bin is the mean held-out log-likelihood per 5-minute bin (higher is better). ΔBIC-style ($\times 10^3$) is the decrease in a BIC-style criterion (held-out log-likelihood minus complexity penalty) relative to $K-1$ (larger = more improvement). Silhouette peaks at $K=6$; ΔBIC-style decelerates sharply at $K=8$.

K	Val LL/bin	Silhouette	ΔBIC-style ($\times 10^3$)	
3	9.07	0.172	–	
4	9.51	0.150	679	
5	11.63	0.096	3,298	
6	12.24	**0.132**	939	•
7	12.76	0.126	817	
8	12.93	0.114	250	

monitoring intensity rather than physiology alone, as higher-risk patients are more likely to receive invasive monitoring. To assess robustness, we repeated the analysis excluding this indicator and restricting inputs to the four universally available channels (NIBP MAP, SpO_2, heart rate, FiO_2; $K=7$) on the subset of operations with at least 30 minutes of valid data ($n=124{,}990$). Arm C AUROC was within 0.003–0.007 of the six-channel model across all outcomes (ICU admission 0.932 vs 0.936; prolonged ICU stay 0.944 vs 0.947; in-hospital death 0.876 vs 0.877), confirming that gains are not driven by the arterial line indicator.

Fold-safe evaluation. All models and preprocessing steps were fit within cross-validation folds using training operations only and applied to held-out data. HMMs were trained and decoded in a fold-safe manner. Cross-fold state alignment was performed via nearest-neighbour emission remapping, matching each fold's states to those from fold one by minimum Euclidean distance between emission means. All experiments use 5-fold grouped cross-validation (`GroupKFold`) by patient identifier to prevent leakage between multiple operations from the same patient.

Models. Logistic regression was used as the primary classifier to ensure that observed differences between arms reflect feature representations rather than model flexibility.

3. Results

3.1. Trajectory features complement summaries

Table 2 shows the predictive comparison across four arms: baseline (A), baseline plus summaries (B), baseline plus HMM trajectories (C), and the combined model (D). For prolonged ICU stay, AUPRC increases from 0.244 (baseline) to 0.305 (summaries), remains similar for HMM features alone (0.303), and reaches 0.339 when combined. HMM trajectory features outperform summaries on AUPRC for ICU admission (0.599 vs 0.582) and ICU within 6 h (0.594 vs 0.575), while summaries outperform trajectories for pro-

Table 2. Fold-safe predictive comparison. Arms: A = baseline, B = baseline + summaries, C = baseline + HMM trajectories, D = all combined. Best per outcome in bold.

Outcome	Arm	AUROC	AUPRC
ICU stay >48 h	A: Baseline	0.927	0.244
	B: + Summaries	0.944	0.305
	C: + HMM	0.947	0.303
	D: Combined	**0.954**	**0.339**
ICU admission	A: Baseline	0.903	0.520
	B: + Summaries	0.927	0.582
	C: + HMM	0.936	0.599
	D: Combined	**0.944**	**0.623**
ICU within 6 h	A: Baseline	0.903	0.513
	B: + Summaries	0.927	0.575
	C: + HMM	0.936	0.594
	D: Combined	**0.943**	**0.616**
In-hospital death	A: Baseline	0.863	0.151
	B: + Summaries	0.882	0.198
	C: + HMM	0.877	0.172
	D: Combined	**0.889**	**0.204**

longed ICU stay (0.305 vs 0.303) and in-hospital death (0.198 vs 0.172). This pattern is consistent with immediate ICU escalation being more sensitive to trajectory structure, whereas sustained deterioration and mortality may be more strongly associated with aggregate physiological burden. The combined model performs best across all outcomes, confirming that both representations carry partially independent prognostic information.

3.2. Latent states are prognostic beyond threshold violations

The six inferred states differ in multivariate combinations of physiological signals while remaining within clinically normal ranges (Figure 1). Despite the absence of overt abnormalities in any individual variable, state occupancy is strongly associated with postoperative outcomes (Appendix B). For prolonged ICU stay, operations with adverse outcomes spend 57.1% of operative time in State 1 compared with 19.7% for operations without the event, with the opposite pattern observed for State 3 (1.3% vs 59.6%). This separation is consistent across outcomes. These states do not reflect overt physiological instability. Instead, prognostic signal arises from multivariate structure across channels, indicating that clinically relevant information is encoded in combinations of physiological signals rather than isolated threshold violations.

3.3. Predictive signal emerges within 15 minutes

Predictive signal is detectable within 15 minutes of anaesthesia onset for all outcomes (Appendix C). As shown in Table 3, for prolonged ICU stay, ΔAUROC increases from

Figure 1. Emission means for the six HMM states across five continuous physiological channels. Colour encodes relative values within each channel (red = minimum, green = maximum). All states lie within standard clinical ranges, with no individual variable breaching typical instability thresholds. Differences between states arise from subtle multivariate patterns across channels rather than from isolated abnormalities. The binary arterial line indicator is omitted.

Table 3. ΔAUROC at intraoperative checkpoints for the combined model over baseline. Signal is detectable at 15 minutes for all outcomes and accumulates progressively for ICU-related outcomes.

Outcome	15 min	30 min	60 min	Full
ICU stay >48 h	+0.014	+0.015	+0.017	+0.029
ICU admission	+0.017	+0.022	+0.027	+0.042
ICU within 6 h	+0.017	+0.022	+0.027	+0.042
In-hospital death	+0.023	+0.024	+0.024	+0.027

+0.014 at 15 minutes to +0.029 at full duration. For ICU admission and early ICU escalation within 6 h, gains increase from +0.017 at 15 minutes to +0.042 at full duration. For in-hospital death, gains plateau after 30 minutes, with early physiological response to anaesthesia induction carrying disproportionate prognostic weight. For ICU-related outcomes, gains accumulate progressively, supporting the value of continuous rather than single-point monitoring. These results indicate that a monitoring-augmented model can flag elevated risk early in the procedure. We note, however, that this analysis is a retrospective simulation (the model is not deployed online) and that absolute gains of +0.014–0.023 AUROC at 15 minutes, while statistically reliable at this sample size, may not be sufficient to change individual clinical decisions without additional validation.

3.4. Gains concentrate where static models fail

The incremental value of dynamic physiology varies substantially across patients. Stratifying operations by tertiles of the composite preoperative model's predicted probability (Low, Medium, and High baseline risk strata, distinct from ASA grade strata) reveals marked heterogeneity. For prolonged ICU stay, ΔAUROC is +0.223 (95% CI +0.170 to +0.274) in the lowest-risk stratum compared with +0.106 in the medium-risk stratum (Appendix D). For in-hospital death, gains decrease monotonically from low to high baseline risk.

This heterogeneity is not explained by any single preoperative variable. When stratified by ASA physical status, ΔAUROC is nearly identical in ASA I–II and ASA III+ groups (difference <0.01 AUROC points). The differential benefit therefore reflects heterogeneity in the *joint* baseline risk profile, captured only by composite learned representations. Patient vulnerability is thus encoded in combinations of preoperative factors rather than in any individual measure.

4. Discussion

Three findings emerge. First, HMM-derived trajectory representations add complementary predictive value to per-channel summaries for ICU-related outcomes, indicating that prognostic structure is multivariate and contextual rather than reducible to threshold crossings. The core contribution of this work is a feature representation comparison demonstrating that HMM trajectory features and summary statistics carry partially independent prognostic information. The Gaussian HMM serves as a deliberately simple baseline. It makes strong parametric assumptions, uses fixed 5-minute bins, and cannot model long-range dependencies. That it nonetheless improves prediction suggests that multivariate structure in intraoperative physiology is genuinely informative, rather than an artefact of model flexibility. This leaves substantial room for richer sequence models. Transformers could capture long-range dependencies across the operative course, state space models could better handle irregular sampling and missingness, and neural process models could represent uncertainty over physiological trajectories.

Second, predictive signal is available within 15 minutes of anaesthesia onset. A model need not wait until the procedure ends to provide useful risk stratification. This raises a representation learning question: if early physiology carries substantial predictive information, what features of anaesthesia induction are most informative, and can they be learned in a self-supervised manner from unlabelled data? The scale of intraoperative monitoring data in hospital systems makes such pretraining a promising direction.

Third, gains concentrate in patients whose preoperative profile appears reassuring but whose physiological trajec-

tory reveals unexpected vulnerability, precisely where static models underestimate risk. ASA grade is a coarse variable, whereas the model integrates multiple baseline factors jointly and is more sensitive to where physiology adds value. This motivates learning-based approaches to identifying which patients benefit most from monitoring. Rather than relying on clinical proxies, a learned representation of the joint preoperative risk profile may better identify patients for whom intraoperative data is most informative. More broadly, this suggests that the value of a data stream is not fixed, but depends on what the model already knows, a principle relevant to any setting where dynamic observations complement static risk.

Taken together, the findings support the view that clinically relevant information lies in the multivariate combination of physiological variables, rather than exclusively in isolated threshold breaches. From a machine learning perspective, this setting motivates representation learning methods that capture multivariate structure in clinical time series. The INSPIRE dataset provides a reproducible benchmark for this agenda, with sufficient scale and clinically meaningful outcomes.

Limitations. Data originate from a single centre in South Korea, and case-mix, monitoring practices, and outcome rates may differ elsewhere. External validation at an independent institution is required before clinical translation. Temporal validation was precluded by date anonymisation in INSPIRE, and direct comparison with established preoperative risk scores such as SORT was prevented by missing variables (e.g. surgical urgency). Bootstrap confidence intervals were computed at the observation level for the stratified analysis and should be interpreted as approximate. Grouped resampling by patient would be more principled.

The two secondary ICU outcomes (ICU admission, 9.6%, and ICU within 6 h, 9.4%) may include planned ICU admissions or resource-use decisions unrelated to acute deterioration. This confounding is less problematic for the primary outcome (prolonged ICU stay >48 h, where planned admissions rarely persist beyond 48 h) but warrants caution in interpreting the secondary outcomes clinically.

The early 15-minute signal may partly reflect patient baseline tolerance to anaesthesia induction rather than intraoperative physiological dynamics per se. Whether this signal is recoverable earlier than surgery and could motivate modified preoperative assessment remains an open question. Additionally, the 5-minute binning discards sub-minute physiological dynamics, and LOCF forward-filling of short gaps may introduce correlation bias at the start of procedures.

Arm B (summaries) was constructed using a defined threshold set. Richer hand-engineered features such as trend, variability, or spectral indices could close the gap between

Arms B and C, so our claim is one of complementarity to the specific summaries used rather than a fundamental limitation of summary-based representations. Finally, model outputs represent one static risk score per surgery, a useful research formulation but not an online risk trajectory. Real-time deployment would require rolling-window recomputation and considerably more validation.

Broader impact. This work explores how intraoperative monitoring data could support dynamic risk stratification during surgery, enabling earlier recognition of patient deterioration and more timely escalation of care. By leveraging existing data streams, such approaches could improve decision-making without requiring changes to monitoring workflows. However, deploying predictive models in this setting carries risks. Automation bias may lead clinicians to over-rely on model outputs, particularly in high-pressure environments such as the operating room. Model errors or miscalibration could result in inappropriate escalation or missed deterioration. In addition, performance may vary across patient subgroups, raising concerns about fairness and generalisability. Careful evaluation, calibration, and integration into workflows will be essential to ensure such systems augment rather than replace clinical judgement.

Acknowledgements

Sadaf Sohrabi is supported by the UKRI AI Centre for Doctoral Training in Digital Healthcare [EP/Y030974/1]. Aldo A. Faisal acknowledges the support of the UKRI AI programme and the Engineering and Physical Sciences Research Council (EPSRC) for the AI Hub in Generative Models [EP/Y028805/1].

References

Fritz, B. A., Cui, Z., Zhang, M., He, Y., Chen, Y., Kronzer, A., Ben Abdallah, A., King, C. R., and Avidan, M. S. Deep-learning model for predicting 30-day postoperative mortality. *British Journal of Anaesthesia*, 123(5):688–695, 2019. doi: 10.1016/j.bja.2019.07.025.

Hao, X., Wang, Y., Li, K., Zhu, T., Herasevich, V., et al. Applying machine learning for perioperative adverse event prediction: a narrative review toward better clinical efficacy and usability. *Anesthesiology and Perioperative Science*, 3(51), 2025. doi: 10.1007/s44254-025-00136-z.

Lim, L., Lee, H., Jung, C.-W., Sim, D., Borrat, X., Pollard, T. J., Celi, L. A., Mark, R. G., Vistisen, S. T., and Lee, H.-C. INSPIRE, a publicly available research dataset for perioperative medicine. *Scientific Data*, 11(1):655, 2024. doi: 10.1038/s41597-024-03517-4.

Pearse, R. M., Moreno, R. P., Bauer, P., Pelosi, P., Metnitz, P., Spies, C., Vallet, B., Vincent, J.-L., Hoeft, A., Rhodes, A., and European Surgical Outcomes Study (EuSOS) group. Mortality after surgery in Europe: a 7 day cohort study. *The Lancet*, 380(9847):1059–1065, 2012. doi: 10.1016/S0140-6736(12)61148-9.

Protopapa, K. L., Simpson, J. C., Smith, N. C. E., and Moonesinghe, S. R. Development and validation of the Surgical Outcome Risk Tool (SORT). *British Journal of Surgery*, 101(13):1774–1783, 2014. doi: 10.1002/bjs.9638.

Shickel, B., Loftus, T. J., Ruppert, M., Upchurch, G. R., Ozrazgat-Baslanti, T., Rashidi, P., and Bihorac, A. Dynamic predictions of postoperative complications from explainable, uncertainty-aware, and multi-task deep neural networks. *Scientific Reports*, 13(1):1224, 2023. doi: 10.1038/s41598-023-27418-5.

Weiser, T. G., Haynes, A. B., Molina, G., Lipsitz, S. R., Esquivel, M. M., Uribe-Leitz, T., Fu, R., Azad, T., Chao, T. E., Berry, W. R., and Gawande, A. A. Estimate of the global volume of surgery in 2012: an assessment supporting improved health outcomes. *The Lancet*, 385(Suppl 2):S11, 2015. doi: 10.1016/S0140-6736(15)60806-6.

Wong, D. J. N., Harris, S., Sahni, A., Bedford, J. R., Cortes, L., Shawyer, R., Wilson, A. M., Lindsay, H. A., Campbell, D., Popham, S., Barneto, L. M., Myles, P. S., SNAP-2: EPICCS collaborators, and Moonesinghe, S. R. Developing and validating subjective and objective risk-assessment measures for predicting mortality after major surgery: an international prospective cohort study. *PLOS Medicine*, 17(10):e1003253, 2020. doi: 10.1371/journal.pmed.1003253.

Xue, B., Li, D., Lu, C., King, C. R., Wildes, T., Avidan, M. S., Kannampallil, T., and Abraham, J. Use of machine learning to develop and evaluate models using preoperative and intraoperative data to identify risks of postoperative complications. *JAMA Network Open*, 4(3):e212240, 2021. doi: 10.1001/jamanetworkopen.2021.2240.

A. Cohort Characteristics

Table A1. Baseline characteristics of the INSPIRE study cohort. Unit of analysis is the surgical operation ($n = 130{,}960$; 99,886 unique patients).

Characteristic	Value
Patient demographics	
Age, years (median [IQR])	60 (45–70)
Female sex	73,023 (55.8%)
BMI, kg/m^2 (median [IQR])	23.8 (21.5–26.0)
Surgical characteristics	
Emergency surgery	12,330 (9.4%)
General anaesthesia	102,790 (78.5%)
ASA I–II	115,139 (87.9%)
ASA III+	12,274 (9.4%)
ASA missing	3,547 (2.7%)
Intraoperative monitoring	
Operations with usable intraoperative physiology	130,888 (99.9%)
Measurements per operation (median [IQR])	69 (36–126)
Anaesthesia duration, min (median [IQR])	115 (70–195)
Outcomes	
ICU admission	12,540 (9.6%)
ICU within 6 h of OR exit	12,285 (9.4%)
ICU stay >48 h (primary)	2,871 (2.2%)
In-hospital death	1,555 (1.2%)

B. HMM State Occupancy by Outcome

Figure B1. Mean fraction of operative time in each of the six HMM states, stratified by event (dark blue) and no-event (light blue) across four postoperative outcomes. Operations followed by adverse events spend substantially more time in State 1 and less in State 3 than operations without adverse outcomes. No state breaches any standard clinical instability threshold, yet the occupancy pattern is strongly and consistently prognostic across all four outcomes.

C. Temporal Emergence of Predictive Signal

Figure C1. (A) AUROC for combined models across progressively longer observation windows for ICU-related outcomes. All three improve steadily with additional physiological data. (B) Incremental AUROC gains decomposed by phase across all four outcomes. For in-hospital death the early gain dominates. Negative bars at 15–120 min are attributable to cohort restriction to longer operations rather than loss of signal.

D. Stratified Benefit Analysis

Figure D1. ΔAUROC from adding intraoperative physiology across baseline risk strata. Strata are defined by tertiles of the baseline model's predicted probability (not by ASA grade). Gains are largest in the lowest-risk stratum for all ICU outcomes, indicating that dynamic physiology adds most value where static models underestimate true risk. Error bars: 95% bootstrap CIs (observation-level).