Bootstrapping the error of Oja's algorithm

Abstract

We consider the problem of quantifying uncertainty for the estimation error of
the leading eigenvector from Oja's algorithm for streaming Principal Component
Analysis, where the data are generated IID from some unknown distribution. By
combining classical tools from the U-statistics literature with recent results on
high-dimensional central limit theorems for quadratic forms of random vectors
and concentration of matrix products, we establish a Gaussian approximation
result for the $\sin^2$ distance between the population eigenvector and the output
of Oja's algorithm. Since estimating the covariance matrix associated with the
approximating distribution requires knowledge of unknown model parameters, we
propose a multiplier bootstrap algorithm that may be updated in an online manner.
We establish conditions under which the bootstrap distribution is close to the
corresponding sampling distribution with high probability, thereby establishing the
bootstrap as a consistent inferential method in an appropriate asymptotic regime.

1 Introduction

Since its discovery over a century ago [16], principal component analysis (PCA) has been a corner-
stone of data analysis. In many applications, dimension reduction is paramount and PCA offers an
optimal low-rank approximation of the original data. PCA is also highly interpretable as it projects
the dataset onto the directions that capture the most variance known as principal components.

Important applications of PCA include image and document analysis, where the largest few principal
components may be used to compress a large dimensional dataset to a manageable size without
incurring much loss; for a discussion of some other applications of PCA, see for example, [31]. In
these settings, the original dimensionality, which could be the number of pixels in an image or the
vocabulary size after removing stop-words, is in the tens of thousands. An offline computation of the
principal components would require the computation of eigenvectors of the sample covariance matrix.
However, in high-dimensional settings, storing the covariance matrix and subsequent eigen-analysis
can be challenging. Streaming PCA methods have gained significant traction owing to their ability to
iteratively update the principal components by considering one data-point at a time.

One of the most widely used algorithms for streaming PCA is Oja's algorithm, proposed in the
seminal work of [50]. Oja's algorithm involves the following update rule:

$$w_{t+1} - w_t = \eta(w_t^T X_t)X_t; \qquad w_{t+1}^T w_{t+1} = 1, \tag{1}$$

where $X_t \in \mathbb{R}^d$ is the t^{th} data point and w_t is the current estimate for the leading eigenvector of
$\Sigma = \mathbb{E}XX^T$ after t data-points have been seen. The parameter η can be thought of as a learning rate,
which can either be fixed or varied as a function of t. In this paper we fix the learning rate, similar
to [29].

Contribution In the present work, we consider the problem of uncertainty quantification for the
estimation error of the leading eigenvector from Oja's algorithm, which is one of the most commonly
used streaming PCA algorithms. Our contributions may be summarized as follows:

1. We derive a high-dimensional χ^2 approximation to the $\sin^2$ error for the leading eigenvector of Oja's algorithm. We recover the optimal convergence rate $O(1/n)$ while allowing d to be potentially exponentially large under suitable structural assumptions on the covariance matrix, matching state-of-the-art theoretical results for consistency of Oja's algorithm. Our result provides a distributional characterization of the $\sin^2$ distance for Oja's algorithm for the first time in the literature. The approximation holds for a wide range of step sizes.

2. Since the χ^2 approximation depends on unknown parameters, we propose an online bootstrap algorithm and establish conditions under which the bootstrap is consistent. Our bootstrap procedure allows the approximation of important quantities such as the quantiles of the error associated with Oja's algorithm for the first time.

Prior analysis of Oja's algorithm. While Oja's algorithm was invented in 1982 it was not until recently that the theoretical workings of Oja's algorithm have been understood. A number of papers in recent years have focused on proving guarantees of convergence of the iterative update in (1) toward the principal eigenvector of the (unknown) covariance matrix $\mathbb{E}XX^T$, which can be recast as projected Stochastic gradient descent on the quadratic objective function

$$\min_{\substack{w \\ w^T w = 1}} -\text{trace}(w^T \Sigma w), \qquad \Sigma = \mathbb{E}XX^T, \tag{2}$$

where projection is onto the non-convex unit sphere. We assume that the data-points are mean zero. Despite being non-convex and thus falling outside the framework for which stochastic gradient descent convergence is soundly established, the output of Oja's algorithm be viewed as a product of random matrices and shares similar structure to other important classes of non-convex problems, such as matrix completion [30, 32], matrix sensing [30], and subspace tracking [4]. Thus, this optimization problem serves as a natural first step toward understanding the behavior of Stochastic Gradient Descent in more general non-convex settings.

Let v_1 denote the principal eigenvector of Σ, and let $\hat{v}_1 = w_n$ be the solution to the stochastic iterative method applying Eq 1. Finally, let $\lambda_1 > \lambda_2$ be the first and second principal eigenvalues of Σ. Sharp rates of convergence for Oja's updates were recently established in [28]. They show that[1], with constant probability, the square of the sine of the angle between v_1 and w satisfies:

$$1 - (v_1^T \hat{v}_1)^2 = O\left(\frac{1}{n}\right) \tag{3}$$

where the O hides a constant which depends in the optimal way on the eigengap between the top two eigenvalues, and independent of n or d, improving on previous error bounds for Oja's algorithm [57, 22, 3, 58, 46, 2] which showed convergence rates which deteriorate with the ambient dimension d, and thus did not fully explain the efficiency of Oja's update. This sharp rate is remarkable, as it matches the error of the principal eigenvector of the sample covariance matrix, which is the batch or offline version of PCA. Other notable work include [36, 38] for unbounded X_i, analysis of Oja's algorithm for computing top k principal components [1, 27].

The Bootstrap. The bootstrap, proposed by [12], is one of the most widely used methods for uncertainty quantification in machine learning and statistics and accordingly has a vast literature. We refer the reader to [21, 61] for expositions on the classical theory of the bootstrap for IID data. Recently, since the groundbreaking work of [9, 10], the bootstrap has seen a renewed surge of interest in the context of high-dimensional data where d can be potentially exponentially larger than n. Of particular relevance to the present work are high-dimensional central limit theorems (CLTs) for quadratic forms, which have been studied by [53, 66, 18] . In particular, our CLT for the estimation error of Oja's algorithm invokes a modest adaptation of [66] to independent but non-identically distributed random variables. In machine learning, bootstrap methods have been used to estimate the uncertainty of randomized algorithms such as bagging and random forests [40], sketching for large scale singular value decomposition (SVD) [41], randomized matrix multiplication [43], randomized least squares [39], to name a few.

A standard notion of bootstrap consistency is that, conditioned on the data, the distribution of the suitably centered and scaled bootstrap functional approaches the true distribution with high probability in some norm on probability measures, typically the Kolmogorov distance, which is the supremum

---

[1]Assuming $\|X_i X_i^T - \Sigma\|$ are bounded almost surely

of the absolute pointwise difference between two CDFs. Bootstrap consistency is often established by deriving a Gaussian approximation for the sampling distribution and showing that the bootstrap distribution is close to the corresponding Gaussian approximation with high probability.

It may seem that if one knows that the limiting distribution of a statistic is Gaussian, this defeats the purpose of bootstrap. However, *for most statistics, the parameters of the limiting normal distribution depend on unknown model parameters, and have to be estimated if one intends to use the normal approximation*. Furthermore, the CLT only gives a *first-order* correct approximation of the target distribution, i.e. with $O(1/\sqrt{n})$ error. In contrast, the bootstrap of a suitably centered and scaled statistic has been shown to be higher order correct for many functionals [20, 21, 23].

Quantifying the Uncertainty for SGD Behind the recent success of neural networks in a wide range of sub-fields of machine learning, the workhorse algorithm has become Stochastic gradient descent (SGD) [51, 49, 55]. For establishing consistency of bootstrap, one requires to establish asymptotic normality [14, 52, 56, 47]. There has also been many works on uncertainty estimation of SGD [8, 37, 15, 60]. However, *all these works are for convex, and predominantly strongly convex loss functions.* Only recently, [67] has established asymptotic normality for nonconvex loss functions under dissipativity conditions and appropriate growth conditions on the gradient, which are weaker conditions than strong convexity but not significantly so.

Now, in Section 2 we present notation and do setup, present our main theoretical results in Section 3, followed by simulations in Section 4.

2   Notation and problem setup

We consider a row-wise IID triangular array, where the the random vectors $\{X_i\}$ in the n^{th} row take values in $\mathbb{R}^{d_n}$, with $E[X_i] = 0$ and $\mathrm{Var}(X_i) = \Sigma_n$. Note that the triangular array ensures that $\{X_1, \ldots, X_n\}$ come from a different distribution for each n. This is similar to traditional settings for the Lindeberg-Feller theorem [11]. Note that the case where d is fixed and n grows is a special case of this setting. For readability, we will drop the subscript n from Σ_n.

Expanding out the recursive definition in Eq 1, we see that Oja's iteration can be expressed as $w_{t+1} = (I_d + \eta X_t X_t^T) w_t$. Thus, after n iterations the vector can be written as a matrix-vector product, where the matrix is a product of n independent matrices. Expanding out the recursive definition, we get:

$$B_n := \prod_{i=1}^n (I_d + \eta X_i X_i^T) \qquad \hat{v}_1 = \frac{B_n u_0}{\|B_n u_0\|}, \tag{4}$$

where I_d is a $d \times d$ identity matrix. where u_0 is a random unit vector in d dimensions. In the scalar case, when $\eta = 1/n$, for large n, the numerator of Eq 4 behaves like $\exp(\sum_i X_i^2/n)$, which in turn converges to $\exp(E[X_1^2])$. For matrices, one hopes that, by independence, a result of the same flavor will hold. And in fact if it does hold, then for $\eta = \frac{\log n}{n}$, the numerator in Eq 4 will concentrate around $\exp(\log n\Sigma)$. The spectrum of this matrix is dominated by the principal eigenvector, i.e. the ratio of the first eigenvalue to the second one is $\exp(\log n(\lambda_1 - \lambda_2))$, where λ_i is the i^{th} eigenvalue of the covariance matrix Σ. This makes it clear that *Oja's algorithm is essentially a matrix vector product of this matrix exponential (suitably scaled) and a random unit vector.*

However, the intuition from the scalar case is nontrivial to generalize to matrices because of the non-commutativity. Limits of products of random matrices have been studied intensively in mathematics in the context of ergodic theory on Markov chains (see [17, 35, 5, 13] etc.). However, until recently, there has not been much work on quantifying the exact *rate of convergence*, or finite-sample large deviation bounds for how a random matrix product deviates from its expectation.

We reparametrize η as η_n/n, where η_n is chosen carefully to obtain a suitable error rate. Note that this is not a scheme where we decrease η over time as in [24], but hold it as a constant which is a function of the total number of data-points.

2.1   The Hoeffding decomposition

The Hoeffding decomposition, introduced in [25], is a key technical tool for studying the asymptotic properties of U-statistics, generalizes far beyond U-statistics [63]. Before explaining the Hoeffding

decomposition, we will first introduce the Hájek projection [63], which gives the best approximation of a general function of n IID random variables, by functions of single random variables X_i. If one can show that this projection onto sums of IID random variables incurs asymptotically negligible error, it immediately leads to an asymptotic CLT. The Hoeffding decomposition essentially improves on this approximation by projecting onto sums of *two or more* variables, i.e. a quadratic space, a cubic space of random variables, and so on. Each new space is chosen to be orthogonal to the previous space. Thus, the Hoeffding decomposition can be thought of as a tool for building approximations of increasing levels of complexity.

2.1.1   The matrix product

The Hoeffding decomposition motivates the following expansion for products of independent random matrices. Let $Y_i = X_i X_i^T - \Sigma$. Observe that B_n can be expressed as:

$$B_n = \sum_{k=0}^{n} T_k, \qquad T_k = \left(\frac{\eta_n}{n}\right)^k \sum_{S \in \mathcal{S}_k^{(n)}} \prod_{i=1}^{n} A_i^{(S)}. \tag{5}$$

where $A_i^{(S)}$ is given by: $A_i^{(S)} = \begin{cases} Y_i & \text{if } i \in S \\ I + \frac{\eta_n}{n}\Sigma & \text{otherwise} \end{cases}$.

The above expansion has favorable properties that facilitate second-moment calculations. In fact, we show in the Supplementary material that,

$$\mathbb{E}\left[\|B_n\|_F^2\right] = \sum_{k=0}^{n} \left(\frac{\eta_n}{n}\right)^{2k} \sum_{S \in \mathcal{S}_k^{(n)}} \mathbb{E}\left[\|\prod_{i=1}^{n} A_i^{(S)}\|_F^2\right]$$

$$\mathbb{E}\left[\|B_n x\|_2^2\right] = \sum_{k=0}^{n} \left(\frac{\eta_n}{n}\right)^{2k} \sum_{S \in \mathcal{S}_k^{(n)}} \mathbb{E}\left[\|\prod_{i=1}^{n} A_i^{(S)} x\|_2^2\right]$$

where the second statement holds for any $x \in \mathbb{R}^d$. In Section 3 we establish the asymptotic distribution of the $\sin^2$ distance between v_1 and the Oja vector in Eq 1. Here we will express that in terms of the matrix product in Eq 4 and the expansion introduced above. Denote by $\hat{v}_1$ the Oja vector, and by $V_\perp$ the $d \times d - 1$ matrix with $2, \ldots, d$ eigenvectors of Σ on its columns. Let $\lambda_1 \geq \cdots \geq \lambda_d$ denote the eigenvalues of Σ. We have the following representation of the $\sin^2$ error:

$$1 - \left(v_1^T \hat{v}_1\right)^2 = 1 - \frac{u_0^T B_n^T v_1 v_1^T B_n u_0}{u_0^T B_n^T B_n u_0} = \frac{u_0^T B_n^T (I - v_1 v_1^T) B_n u_0}{u_0^T B_n^T B_n u_0} = \frac{\|V_\perp V_\perp^T B_n u_0\|^2}{\|B_n u_0\|^2} \tag{6}$$

2.2   Online bootstrap for streaming PCA

The multiplier bootstrap is based on the multiplier CLT [62]. Also known as the wild bootstrap, or the weighted bootstrap, it is one of the most widely used resampling methods for measuring the significance of test or for constructing confidence intervals. For a better understanding of the theory of multiplier bootstraps and its wide applicability for example see [7, 9, 10, 45, 59].

The multiplier bootstrap involves perturbing the data by multiplier random variables that match the first two moments of the data. As observed by [9], a Gaussian multiplier random variable eliminates the need to establish a Gaussian approximation for the bootstrap since, conditional on the data, it is already Gaussian. For many inferential quantities, this bootstrap procedure results in a good approximation to the distribution of interest. It is not hard to see that this is a natural candidate for the online setting. It has been used for bootstrapping the stochastic gradient descent estimator [15].

We present our bootstrap in Algorithm 1. In our procedure, wew update $m + 1$ vectors at every iteration. The first one is $\hat{v}$, which will result in the final Oja estimate of the first principal component. The other vectors $\{v^{*(j)}, j = 1, \ldots m\}$ are obtained by perturbing the basic Oja update (Eq 1).

The W_i's are the multiplier random variables, which are scaled mean zero scaled Gaussians with variance $1/2$. The update of the $v^{*(j)}$ is novel because it preserves the mean and the variance of the original Oja estimator, while not requiring access to the full sample covariance matrix. Hence we can make our updates online, and thus, to get error estimates in addition to a pointwise estimate of the principal eigenvector, *we only have to increase computation and storage by a factor of m.*

Algorithm 1: Bootstrap for Oja's algorithm

Input: Datapoints $X_1, \ldots, X_n$, stepsize η, number of bootstrap resamples m, initial unit vector u_0

Output: Oja's solution $\hat{v}_1$ and m bootstrapped versions of it $v_1^{*(1)}, \ldots, v_1^{*(m)}$

Draw $g \sim N(0, I_p)$

Create unit vector $u_0 \leftarrow g/\|g\|$

Initialize $\hat{v}_1, v_1^{*(1)}, \ldots, v_1^{*(m)} \leftarrow u_0$

for $t=2,\ldots, n$ **do**

    Update $\hat{v}_1 \leftarrow \hat{v}_1 + \eta(X_t^T \hat{v}_1)\hat{v}_1$

    Normalize $\hat{v}_1$ to have unit norm;

    for $i=1{:}m$ **do**

        Draw $W_i \sim N(0, 1/2)$;

        Let $h^{(i)} \leftarrow (X_t^T v_1^{*(i)})X_t$;

        Let $g^{(i)} \leftarrow (X_{t-1}^T v_1^{*(i)})X_{t-1}$;

        Update $v_1^{*(i)} \leftarrow v_1^{*(i)} + \eta\left(Y_t + Z_i(h^{(i)} - g^{(i)})\right)$;

        Normalize $v_1^{*(i)}$ to have unit norm;

    end

end

---

3   Main Results

In this section we present our main contributions. We start by stating a CLT for the error of Oja's algorithm. To state this theorem, we will need to introduce some notation.

Let $\Lambda_\perp$ be a diagonal matrix with $\Lambda_\perp(i,i) = (1 + \eta_n \lambda_{i+1}/n)/(1 + \eta_n \lambda_1/n)$, $i = 1, \ldots, d-1$. Also let

$$\mathbb{M} := \mathbb{E}\left[V_\perp^T (X_1^T v_1)^2 X_1 X_1^T V_\perp\right]. \tag{7}$$

Now we define

$$\bar{\mathbb{V}}_n = \frac{\eta_n}{n} \sum_i \mathbb{E}[V_\perp \Lambda_\perp^{i-1} V_\perp^T (X_i X_i^T - \Sigma) v_1 v_1^T (X_i X_i^T - \Sigma) V_\perp \Lambda_\perp^{i-1} V_\perp^T]$$

$$= \frac{\eta_n}{n} V_\perp \left(\sum_i \Lambda_\perp^{i-1} \mathbb{M} \Lambda_\perp^{i-1}\right) V_\perp^T \tag{8}$$

Theorem 1. *Suppose that u_0 is drawn from the uniform distribution on $\mathcal{S}^{d-1}$, $\lambda_1 = O(1)$. Choose $\eta_n \to \infty$ such that $\sqrt{nd}\exp(-\eta_n(\lambda_1 - \lambda_2)) \to 0$, $\frac{(\eta_n \vee \log d)\,\eta_n^2 M_d^2}{n} \to 0$, where $M_d = \mathbb{E}[\|X_i X_i^T - \Sigma\|^2]$. Further, suppose that:*

$$\|\mathbb{M}\|_F \geq C > 0 \tag{9}$$

$$\frac{\mathbb{E}\left[\|V_\perp(X_1 X_1^T - \Sigma)v_1\|^6\right]}{\|\mathbb{M}\|_F^3} = o(n) \tag{10}$$

Then, there exists a sequence of Gaussian distributions $\{Z_n\}_{n \geq 1}$ with mean 0 and covariance matrix $\bar{\mathbb{V}}_n$ (see Eq 8) such that

$$\sup_{t \in \mathbb{R}} \left|P\left(n/\eta_n\, \sin^2(\hat{v}_1, v_1) \leq t\right) - P(Z_n^T Z_n \leq t)\right| \to 0 \tag{11}$$

Theorem 1 is very general. It uses a triangular array, and allows the dimensionality to grow with the number of observations, which is typically done in high dimensional bootstrap literature. Note that the case of fixed d and growing n is a special case of this setup.

We want to point out that while previous literature obtained sharp bounds on the $\sin^2$ error $1 - (v_1^T \hat{v}_1)^2$, we go a step further. *We establish the distribution of $n/\eta_n(1 - (v_1^T \hat{v}_1)^2)$.*

Remark 1 (Condition on norm). *For simplicity, we assume $\lambda_1 = O(1)$, which can be easily relaxed to growly wih n. We do not assume that the $\|X_i X_i^T - \Sigma\|_2$ is bounded almost surely. However, the norm of $X_i X_i^T - \Sigma$ comes into play implicitly via the assumption in Eq 10. Consider the case where X_i are drawn from some multivariate Gaussian distribution. Note that convergence to a χ^2 in this case is immediate. However we use this to build intuition about the assumptions in Eq 9 and 10. In this case, $X_1^T V_\perp$ is a gaussian of independent entries and thus $\mathbb{E}\left[\left\|V_\perp (X_1 X_1^T - \Sigma)v_1\right\|\right]^6 = \mathbb{E}\|X_1^T v_1\|^6 \mathbb{E}\left(\sum_{j>1}(X_j^T v_j)^2\right)^3$. Note that $\sum_{j>1}((X_j^T v_j)^2 - \lambda_j)$ is a sub-exponential random variable with parameters $(c_1 \sum_{j>1} \lambda_j, c_2)$. Furthermore, $\|\mathbb{M}\|_F^2 = \lambda_1 \sum_{i>1} \lambda_i$. Thus Eq 10 basically reduces to checking if*

$$\frac{\lambda_1^{3/2}(\sum_{j>1} \lambda_j)^3}{\left(\sum_i \lambda_i^2\right)^{3/2}} = o(n)$$

Under appropriate conditions on the marginal variances, which are often needed for high-dimensional PCA results [42, 6, 54, 44, 34, 33, 48, 64], this is a mild condition.

As a side product of our main theorem, we obtain the following error bound on the $\sin^2$ error.

Corollary 1. *Under the conditions in Theorem 1, we have*

$$\sin^2(\hat{v}_1, v_1) = \frac{\eta_n}{n}\left[\text{trace}(\bar{\mathbb{V}}_n) + O_P\left(\|\bar{\mathbb{V}}_n\|_F + d\exp\left\{-\eta_n(\lambda_1 - \lambda_2)\right\}\right)\right]$$

Remark 2 (Comparison with previous work). *Note that, the above rate is similar to the error bounds in previous work in the dependence on n and d. Also note that, the main error essentially comes from the limiting χ^2 distribution, whose mean is $\text{trace}(\bar{\mathbb{V}}_n)$ and variance is $\|\bar{\mathbb{V}}_n\|_F$.*

Consider a matrix $\mathbb{M}' = V_\perp^T \bar{\mathbb{V}}_n V_\perp$, and observe that the Frobenius norm and trace of $\mathbb{M}'$ are the same as that of $\bar{\mathbb{V}}_n$. Some calculations show that the k, ℓ^{th} entry of $\mathbb{M}'$:

$$\mathbb{M}'(k,\ell) = \frac{\mathbb{M}(k,\ell)(1 + O(\eta_n/n))}{2\lambda_1 - (\lambda_k + \lambda_\ell)}$$

Therefore, by Eq 8, we have

$$\text{trace}(\bar{\mathbb{V}}_n) \leq \frac{C\text{trace}(\mathbb{M})}{\lambda_1 - \lambda_2} \leq C\frac{M_d}{\lambda_1 - \lambda_2}$$
$$\|\bar{\mathbb{V}}_n\|_F \leq C'\frac{\|\mathbb{M}\|_F}{\lambda_1 - \lambda_2} \leq C'\frac{M_d}{\lambda_1 - \lambda_2}$$

If we set $\eta_n = c_1 \log n/(\lambda_1 - \lambda_2)$, for large enough c_1, the dominating term in the error is $O_P\left(\frac{M_d \log n}{n(\lambda_1 - \lambda_2)^2}\right)$ under mild conditions on d.

Remark 3 (Rate of Convergence in Kolmogorov Distance). *To simplify the theorem statement, we have stated Theorem 1 without giving an explicit rate of convergence in the Kolmogorov distance. Convergence rates depend on the rate of decay of the remainder terms, which are worked out in the Supplement, and the magnitude of the quantity in Eq 10. The contribution of the latter quantity to the rate is worked out in the IID case in [66].*

Remark 4 (Lower bound on norm). *While our rate matches the sharp bounds in literature and our assumptions on norm upper bounds are similar or weaker than previous work, we do assume a lower bound on the Frobenius norm of the covariance matrix as in Eq 9. Note that if indeed all X_i's were a scalar multiple of v_1, then the $\bar{\mathbb{V}}_n$ matrix in Eq 8 will be zero. This will lead to a perfect point estimate, but there will not be any variability from the data and hence there will be no non-degenerate approximation. Therefore, the lower bound on the norm is not resulting from loose analysis; similar lower bounds on the variance are imposed in the high-dimensional CLT literature [9, 10].*

Now we provide a proof sketch of Theorem 1 below.

Proof sketch for Theorem 1. Our goal is to show that $\frac{\|V_\perp V_\perp^T B_n u_0\|^2}{\|B_n u_0\|^2}$ is asymptotically equivalent to a χ^2 distribution.

1. Our first step is to show that $\|B_n u_0\|$ concentrates around its expectation $(1 + \eta_n \lambda_1/n)^n |v_1^T u_0|$. This replaces the denominator by a non-random quantity.

2. Next we establish that $\frac{\|V_\perp V_\perp B_n V_\perp V_\perp^T u_0\|_2}{\|B_n u_0\|}$ is $O_P\left(\sqrt{d} \cdot \exp\{-\eta_n(\lambda_1 - \lambda_2)\} + \sqrt{\frac{\eta_n^3 M_d^2 \log d}{n^2}}\right)$.

   This is achieved by using a similar recursive argument as in [28], but with the crucial observation that the residual or common difference term is of a lower order because it can be replaced by a matrix product minus its expectation.

3. Now we go back to the expansion in Eq 5.

$$(v_1^T u_0) V_\perp V_\perp^T B_n v_1 = (v_1^T u_0) \sum_k V_\perp V_\perp^T T_k v_1$$

   We see that because of orthogonality, since $T_0 = (I + \eta_n/n\Sigma)^n$, $V_\perp V_\perp^T T_0 v_1$ is the zero vector. Now we examine the $(v_1^T u_0) V_\perp V_\perp^T (B_n - T_1) v_1$ term. Here we use the structure of the higher order terms T_k. In particular, we use the fact that it is a matrix product interlaced with k $X_i X_i^T - \Sigma$ matrices. For example, for $k = 2$ we have

$$T_2 = \frac{\eta_n^2}{n^2} \sum_{i<j} \left(I + \frac{\eta_n}{n}\Sigma\right)^{i-1} Y_i \left(I + \frac{\eta_n}{n}\Sigma\right)^{j-i-1} Y_j \left(I + \frac{\eta_n}{n}\Sigma\right)^{n-j}$$

   We show that the norm of $(v_1^T u_0) V_\perp V_\perp^T (B_n - T_1) v_1$, normalized by the denominator, is $O(\eta_n^2 M_d^2/n^2)$. The fact that the summands of T_k are uncorrelated, and T_k and T_ℓ are uncorrelated, for $k \neq \ell$ makes this possible.

4. Finally, we are left with $V_\perp V_\perp^T T_1 v_1 (v_1^T u_0)$. Note that this is of the following form:

$$\frac{\eta_n}{n} \frac{(v_1^T u_0) V_\perp V_\perp^T T_1 v_1}{|v_1^T u_0|(1 + \lambda_1 \eta_n/n)^n} = \frac{\eta_n \text{sgn}(v_1^T u_0)}{n} \sum_{i=1}^n V_\perp \Lambda_\perp^{i-1} V_\perp^T (X_i X_i^T - \Sigma) v_1$$

   It is not hard to see that this is a sum of independent random vectors with covariance matrix $\eta_n/n\overline{\mathbb{V}}_n$ (see Eq 8).

5. We adapt a result of distributional convergence of squared norm of sums of IID random vectors in [66] to squared norm of sums of independent random vectors. Under the assumptions 10 and 9, the conditions of distributional convergence are satisfied.

6. Finally, all the error terms are combined along with standard anti-concentration argument for χ^2 to establish the final result. The full proof and accompanying lemmas are in Section A of the Supplement.

$\square$

Note that, given the exact form of the χ^2 distribution in Theorem 1, it is natural to wonder if one can just estimate the parameters of this distributions and directly do inference for the error of Oja's algorithm. However, this will require estimation of the eigenvalues of Σ as well as other population quantities. Instead, we propose a streaming bootstrap procedure that mimics the properties of the original Oja algorithm. While the similar structure leads to error terms that are similar to the CLT, the bootstrap presents its own technical challenges. In what follows let P^* denote the bootstrap measure, which is conditioned on the data, and let $\mathbb{E}^*[\cdot]$ denote the corresponding expectation operator.

A common strategy for establishing consistency of the Gaussian multiplier bootstrap is to invoke a Gaussian comparison lemma. Since the multipliers are themselves Gaussian and the data is treated as fixed, the idea is that one can use specialized results for comparing the distributions of two Gaussians (bootstrapped Z^* and limiting Z) that only depend on how close the covariance matrices $\mathbb{E}^* Z^* Z^{*T}$ and $\mathbb{E} ZZ^T$ are in an appropriate metric. For quadratic forms, various results in the literature exist [18, 19], of which we use the later, which upper bounds the Kolmogorov distance between the distributions of norms using the trace and Frobenius norm of the difference between the corresponding covariance matrices. We invoke the comparison lemma to establish closeness in probability between the covariance matrix of the main term in the bootstrap with that that of the Gaussian approximation. While the original covariance matrix is complicated, it turns out that this closeness can be stated succinctly:

Lemma 1. *[Bounding the difference between the bootstrap covariance and true covariance] Let:*

$$Z^* = sgn(v_1^T u_0)\sqrt{\frac{\eta_n}{n}} \sum_i W_i V_\perp \Lambda_\perp^{i-1} V_\perp^T (X_i X_i^T - X_{i-1} X_{i-1}^T) v_1. \tag{12}$$

Recall the definition of $\bar{\mathbb{V}}_n$ from Eq 8. We have,

$$|\text{trace}(\mathbb{E}^* Z^* Z^{*T} - \bar{\mathbb{V}}_n)|, \|\mathbb{E}^* Z^* Z^{*T} - \bar{\mathbb{V}}_n\|_F = O_P\left(\sqrt{\frac{\mathbb{E}\|X_1 X_1^T - \Sigma\|^4}{n(\lambda_1 - \lambda_2)}}\right)$$

With this lemma in hand, we are ready to state our bootstrap approximation result.

Theorem 2 (Bootstrap Consistency)**.** *Suppose that the conditions of Theorem 1 are satisfied. Furthermore, let α_n be a sequence such that $P(\mathcal{A}_n^c) \to 0$, where $\mathcal{A}_n$ is defined as $\mathcal{A}_n = \left\{\max_{i \le i \le n} \|X_i\|_2^2 \le \alpha_n\right\}$. Further suppose that $\frac{\alpha_n^4 \eta_n^3}{n} \to 0$, $\sqrt{\alpha_n}\eta_n \exp(-\eta_n(\lambda_1 - \lambda_2)) \to 0$, $\frac{\alpha_n \eta_n^{3/2} M_d}{n(\lambda_1-\lambda_2)} \to 0$, and $\frac{\mathbb{E}[\|X_i X_i^T - \Sigma\|^4]}{n(\lambda_1-\lambda_2)} \to 0$. Then,*

$$\sup_{t\in\mathbb{R}} \left|P^*(n/\eta_n \sin^2(\hat{v}_1, v_1) \le t) - P(n/\eta_n \sin^2(v_1^*, \hat{v}_1) \le t)\right| \xrightarrow{P} 0$$

Proof sketch of Theorem 2. The proof follows the same route as Theorem 2. We provide the detailed analysis in Supplementary Section B. A similar representation as Eq 5 Hoeffding decomposition is now obtained conditioned on the data. So in step one we have B_n^* replace B_n, where B_n^* is given by:

$$B_n^* = \prod_{i=1}^{n}\left(I + \eta_n/n(X_i X_i^T + W_i(X_i X_i^T - X_{i-1}X_{i-1}^T))\right)$$

So we work out Step 1 using concentration of matrix products [26]. For step 2-3, we see that T_k^* has the same structure as T_k with the difference that $(I + \eta_n \Sigma/n)^i$ gets replaced by its sample counter-part which is a product of i independent matrices of the form $I + \eta_n/n X_j X_j^T$. Concentration of these terms in operator norm are established with results from [26]. Finally for step 4, we see that the main term that approximates the bootstrap residual $\widehat{V}_\perp \widehat{V}_\perp^T B_n^* u_0$ is given by $\sqrt{\eta_n/n} Z^*$, where Z^* is given in Eq 12. Conditioned on the data, this is already a normal distribution because of the multiplier random variables W_i, albeit with a different covariance matrix $\mathbb{E}^* Z^* Z^{*T}$. So now we invoke Gaussian comparison results (e.g. Corollary 2.2, [19]) and 1 to obtain convergence to the χ^2 approximation. $\square$

In theorem above, we have stated the good set $\mathcal{A}_n$ in an abstract manner, but one may wonder how stringent the condition is in various problem settings. Below, we describe a general setup with sub-exponential entries of X_i in which α_n grows as $\log n$; under milder forms of various decay, all we need is for α_n to grow slowly with n. Here $\|\cdot\|_{\psi_1}$ is the sub-Exponential Orlicz norm and $\|\cdot\|_{\psi_2}$ is the sub-Gaussian Orlicz norm (see, for example [65]).

Proposition 1 (The effect of variance decay on the norm)**.** *For each $1 \le j \le p$, suppose that X_{1j} satisfies $\|X_{1j}\|_{\psi_2} \le \nu_j \sum_{j=1}^p \nu_j \le C_1 < \infty$. Then, for some universal constant $0 \le C_2 < \infty$, $\left\|\sum_{j=1}^p X_{1j}^2\right\|_{\psi_1} < C_2$ and there exists some $c_1, c_2 > 0$ such that:*

$$P\left(\max_{1\le i \le n} \|X_i\|_2^2 > c_1 \log n\right) \le \frac{c_2}{n}$$

We now present experimental validation of our bootstrap procedure below.

4 Experimental validation of the online multiplier bootstrap

We draw $Z_{ij} \overset{IID}{\sim} \text{Uniform}(-\sqrt{3}, \sqrt{3})$, for $i = 1, \ldots, n$ and $j = 1, \ldots d$. Consider a PSD matrix $K_{ij} = \exp(-|i-j|c)$ with $c = 0.01$. Now we create a covariate matrix such that $\Sigma_{ij} = K(i, j)\sigma_i\sigma_j$.

We use $\sigma_i = 5i^{-\beta}$. Now we transform our data to introduce dependence by using $X = Z(\Sigma^{1/2})^T$. Note that we have $\mathbb{E}X^T X = \Sigma$. Our goal is to simply demonstrate that the bootstrap distribution of $\sin^2$ errors closely match that of the sampling distribution. To this effect, we fix u_0 and draw 500 datasets and run streaming PCA on each and then construct an empirical CDF (F) from the $\sin^2$ error with the true v_1. This is the point of comparison for the bootstrap distribution (F^*), for which we fix a dataset X. We then invoke algorithm 1 to obtain 500 bootstrap replicates $\hat{v}_1^*$ as well as the Oja vector for the dataset $\hat{v}_1$. The bootstrap distribution is the empirical CDF of $1 - (\hat{v}_1^T \hat{v}_1^*)^2$. We use $\eta_n = \log n$. In Figure 1, we see that for $\beta = 0.2$ (see (A) and (B)), where the variance decay

Figure 1: Bootstrapped and sampling CDF for $n = 1000, d = 500$ in (A) and (C) and for $n = 10,000, d = 500$ in (B) and (D). (A) and (B) use $\beta = 0.2$ whereas (C) and (D) use $\beta = 1$.

is slow and therefore the error bounds of the residual terms are expected to be large, the quality of approximation is poorer compared to (C) and (D), where $\beta = 1$. However, even for $\beta = 0.2$, increasing n improves the performance. We also did these simulations for $X \sim N(0, \Sigma)$, where the results were qualitatively similar, leading us to believe that error from the residuals of higher order terms in the expansion 5 incur more error than the Gaussian approximation.

5   Discussion and Limitations

Modern tools in nonasymptotic random matrix theory have given rise to recent breakthroughs in establishing pointwise convergence rates for stochastic iterative methods in optimizing certain nonconvex objectives, including the classic Oja's algorithm for online principal component analysis. By synthesizing modern random matrix theory tools with classic results from the U-statistics literature and recently developed high-dimensional central limit theorems, we extend the error analysis of Oja's algorithm from pointwise convergence rates to *distributional* convergence and moreover establish an efficient online bootstrap method for Oja's algorithm to quantify the error on the fly. Our results are a first step toward incorporating uncertainty estimation into the general framework of stochastic optimization algorithms, but we acknowledge the present limitations of our analysis: new tools will be needed to extend the current analysis to estimating higher-dimensional principal subspaces, and additional tools will be needed to account for non-independent matrix products which appear beyond the setting of online PCA.

References

[1] Z. Allen-Zhu, Y. Li, R. Oliveira, and A. Wigderson. Much Faster Algorithms for Matrix Scaling. In *Proceedings of the 58th Symposium on Foundations of Computer Science*, FOCS '17, 2017. Full version available at http://arxiv.org/abs/1704.02315.

[2] M.-F. Balcan, S. S. Du, Y. Wang, and A. W. Yu. An improved gap-dependency analysis of the noisy power method. In V. Feldman, A. Rakhlin, and O. Shamir, editors, *29th Annual Conference on Learning Theory*, volume 49 of *Proceedings of Machine Learning Research*, pages 284–309, Columbia University, New York, New York, USA, 23–26 Jun 2016. PMLR.

[3] A. Balsubramani, S. Dasgupta, and Y. Freund. The fast convergence of incremental pca. In C. J. C. Burges, L. Bottou, M. Welling, Z. Ghahramani, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems 26*, pages 3174–3182. Curran Associates, Inc., 2013.

[4] L. Balzano, R. Nowak, and B. Recht. Online Identification and Tracking of Subspaces from Highly Incomplete Information. *arXiv e-prints*, page arXiv:1006.4046, June 2010.

[5] Y. Benoist and J.-F. Quint. *Random Walks on Reductive Groups*. Springer International Publishing, June 2016.

[6] F. Bunea, L. Xiao, et al. On the sample covariance matrix estimator of reduced effective rank population matrices, with applications to fpca. *Bernoulli*, 21(2):1200–1230, 2015.

[7] S. Chatterjee and A. Bose. Generalized bootstrap for estimating equations. *Ann. Statist.*, 33(1):414–436, 02 2005.

[8] X. Chen, J. D. Lee, X. T. Tong, and Y. Zhang. Statistical inference for model parameters in stochastic gradient descent. *Ann. Statist.*, 48(1):251–273, 02 2020.

[9] V. Chernozhukov, D. Chetverikov, and K. Kato. Gaussian approximations and multiplier bootstrap for maxima of sums of high-dimensional random vectors. *The Annals of Statistics*, 41(6):2786 – 2819, 2013.

[10] V. Chernozhukov, D. Chetverikov, and K. Kato. Central limit theorems and bootstrap in high dimensions. *Ann. Probab.*, 45(4):2309–2352, 07 2017.

[11] R. Durrett. Probability: Theory and examples, 2011.

[12] B. Efron and R. Tibshirani. Bootstrap methods for standard errors, confidence intervals, and other measures of statistical accuracy. *Statist. Sci.*, 1(1):54–75, 02 1986.

[13] J. Emme and P. Hubert. Limit laws for random matrix products. *arXiv e-prints*, page arXiv:1712.03698, Dec. 2017.

[14] V. Fabian. On asymptotic normality in stochastic approximation. *Ann. Math. Statist.*, 39(4):1327–1332, 08 1968.

[15] Y. Fang, J. Xu, and L. Yang. Online bootstrap confidence intervals for the stochastic gradient descent estimator. *J. Mach. Learn. Res.*, 19(1):3053–3073, Jan. 2018.

[16] K. P. F.R.S. Liii. on lines and planes of closest fit to systems of points in space. *The London, Edinburgh, and Dublin Philosophical Magazine and Journal of Science*, 2(11):559–572, 1901.

[17] A. Furman. Chapter 12 random walks on groups and random transformations. volume 1 of *Handbook of Dynamical Systems*, pages 931 – 1014. Elsevier Science, 2002.

[18] A. Giessing and J. Fan. Bootstrapping ℓ_p-statistics in high dimensions, 2020.

[19] F. Gʹotze, A. Naumov, V. Spokoiny, and V. Ulyanov. Large ball probabilities, Gaussian comparison and anti-concentration. *Bernoulli*, 25(4A):2538 – 2563, 2019.

[20] F. Götze and H. R. Künsch. Second-order correctness of the blockwise bootstrap for stationary observations. *Ann. Statist.*, 24(5):1914–1933, 10 1996.

[21] P. Hall. *The Bootstrap and Edgeworth Expansion*. Springer-Verlag, New York, 1992.

[22] M. Hardt and E. Price. The noisy power method: A meta algorithm with applications. In Z. Ghahramani, M. Welling, C. Cortes, N. D. Lawrence, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems 27*, pages 2861–2869. Curran Associates, Inc., 2014.

[23] R. Helmers. On the edgeworth expansion and the bootstrap approximation for a studentized u-statistic. *Ann. Statist.*, 19(1):470–484, 03 1991.

[24] A. Henriksen and R. Ward. AdaOja: Adaptive Learning Rates for Streaming PCA. *arXiv e-prints*, page arXiv:1905.12115, May 2019.

[25] W. Hoeffding. A class of statistics with asymptotically normal distribution. *Annals of Mathematical Statistics*, 19(3):293–325, 09 1948.

[26] D. Huang, J. Niles-Weed, J. A. Tropp, and R. Ward. Matrix concentration for products, 2020.

[27] D. Huang, J. Niles-Weed, and R. Ward. Streaming k-pca: Efficient guarantees for oja's algorithm, beyond rank-one updates, 2021.

[28] P. Jain, C. Jin, S. Kakade, P. Netrapalli, and A. Sidford. Streaming pca: Matching matrix bernstein and near-optimal finite sample guarantees for oja's algorithm. In *Proceedings of The 29th Conference on Learning Theory (COLT)*, June 2016.

[29] P. Jain, D. Nagaraj, and P. Netrapalli. SGD without Replacement: Sharper Rates for General Smooth Convex Functions. *arXiv e-prints*, page arXiv:1903.01463, Mar. 2019.

[30] P. Jain, P. Netrapalli, and S. Sanghavi. Low-rank matrix completion using alternating minimization. In *Proceedings of the Forty-Fifth Annual ACM Symposium on Theory of Computing*, STOC '13, page 665–674, New York, NY, USA, 2013. Association for Computing Machinery.

[31] I. T. Jolliffe and J. Cadima. Principal component analysis: a review and recent developments. *Phil. Trans. R. Soc. A.*, 2016.

[32] R. Keshavan, A. Montanari, and S. Oh. Matrix completion from a few entries. *Information Theory, IEEE Transactions on*, 56:2980 – 2998, 07 2010.

[33] V. Koltchinskii, M. Löffler, R. Nickl, et al. Efficient estimation of linear functionals of principal components. *The Annals of Statistics*, 48(1):464–490, 2020.

[34] V. Koltchinskii and K. Lounici. Concentration inequalities and moment bounds for sample covariance operators. *Bernoulli*, 23(1):110–133, 2017.

[35] F. Ledrappier. *Some asymptotic properties of random walks on free groups*, pages 117–152. 06 2001.

[36] C. J. Li, M. Wang, H. Liu, and T. Zhang. Near-optimal stochastic approximation for online principal component estimation. *Math. Program.*, 167(1):75–97, 2018.

[37] T. Li, L. Liu, A. Kyrillidis, and C. Caramanis. Statistical inference using SGD. *arXiv e-prints*, page arXiv:1705.07477, May 2017.

[38] X. Liang. On the optimality of the oja's algorithm for online pca, 2021.

[39] M. Lopes, S. Wang, and M. Mahoney. Error estimation for randomized least-squares algorithms via the bootstrap. volume 80 of *Proceedings of Machine Learning Research*, pages 3217–3226, Stockholmsmässan, Stockholm Sweden, 10–15 Jul 2018. PMLR.

[40] M. E. Lopes. Estimating the algorithmic variance of randomized ensembles via the bootstrap. *Ann. Statist.*, 47(2):1088–1112, 04 2019.

[41] M. E. Lopes, N. B. Erichson, and M. W. Mahoney. Error Estimation for Sketched SVD via the Bootstrap. In *Proceedings of the International Conference of Machine Learning*, 2020.

[42] M. E. Lopes, Z. Lin, and H.-G. Müller. Bootstrapping max statistics in high dimensions: Near-parametric rates under weak variance decay and application to functional and multinomial data. *The Annals of Statistics*, 48(2):1214 – 1229, 2020.

[43] M. E. Lopes, S. Wang, and M. W. Mahoney. A bootstrap method for error estimation in randomized matrix multiplication. *Journal of Machine Learning Research*, 20(39):1–40, 2019.

[44] K. Lounici. High-dimensional covariance matrix estimation with missing observations. *Bernoulli*, 20(3):1029 – 1058, 2014.

[45] E. Mammen. Bootstrap and wild bootstrap for high dimensional linear models. *Ann. Statist.*, 21(1):255–285, 03 1993.

[46] I. Mitliagkas, C. Caramanis, and P. Jain. Memory limited, streaming pca. In *Proceedings of the 26th International Conference on Neural Information Processing Systems - Volume 2*, NIPS'13, page 2886–2894, Red Hook, NY, USA, 2013. Curran Associates Inc.

[47] E. Moulines and F. R. Bach. Non-asymptotic analysis of stochastic approximation algorithms for machine learning. In J. Shawe-Taylor, R. S. Zemel, P. L. Bartlett, F. Pereira, and K. Q. Weinberger, editors, *Advances in Neural Information Processing Systems 24*, pages 451–459. Curran Associates, Inc., 2011.

[48] A. Naumov, V. Spokoiny, and V. Ulyanov. Bootstrap confidence sets for spectral projectors of sample covariance. *Probability Theory and Related Fields*, 174(3):1091–1132, 2019.

[49] A. Nemirovski, A. B. Juditsky, G. Lan, and A. Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM J. Optimization*, 19(4):1574–1609, 2009.

[50] E. Oja. Simplified neuron model as a principal component analyzer. *Journal of Mathematical Biology*, 15(3):267–273, Nov. 1982.

[51] B. T. Polyak and A. B. Juditsky. Acceleration of stochastic approximation by averaging. *SIAM J. Control Optim.*, 30(4):838–855, July 1992.

[52] B. T. Polyak and A. B. Juditsky. Acceleration of stochastic approximation by averaging. *SIAM Journal on Control and Optimization*, 30(4):838–855, 1992.

[53] D. Pouzo. Bootstrap consistency for quadratic forms of sample averages with increasing dimension. *Electronic Journal of Statistics*, 9(2):3046 – 3097, 2015.

[54] M. Reiß and M. Wahl. Nonasymptotic upper bounds for the reconstruction error of PCA. *The Annals of Statistics*, 48(2):1098 – 1123, 2020.

[55] H. Robbins and S. Monro. A stochastic approximation method. *Ann. Math. Statist.*, 22(3):400–407, 09 1951.

[56] D. Ruppert. Efficient estimations from a slowly convergent robbins-monro process. 1988.

[57] C. D. Sa, K. Olukotun, and C. Ré. Global convergence of stochastic gradient descent for some nonconvex matrix problems. *CoRR*, abs/1411.1134, 2014.

[58] O. Shamir. Convergence of stochastic gradient descent for pca. In *Proceedings of the 33rd International Conference on International Conference on Machine Learning - Volume 48*, ICML'16, page 257–265. JMLR.org, 2016.

[59] V. Spokoiny and M. Zhilova. Bootstrap confidence sets under model misspecification. *Ann. Statist.*, 43(6):2653–2675, 12 2015.

[60] W. J. Su and Y. Zhu. Uncertainty quantification for online learning and stochastic approximation via hierarchical incremental gradient descent, 2018.

[61] A. van der Vaart. *Asymptotic statistics*. Cambridge University Press, 2000.

[62] A. van der Vaart, A. van der Vaart, A. van der Vaart, and J. Wellner. *Weak Convergence and Empirical Processes: With Applications to Statistics*. Springer Series in Statistics. Springer, 1996.

[63] A. W. Van der Vaart. *Asymptotic statistics*, volume 3. Cambridge university press, 2000.

[64] R. Vershynin. Introduction to the non-asymptotic analysis of random matrices. *arXiv preprint arXiv:1011.3027*, 2010.

[65] R. Vershynin. *High-Dimensional Probability*. Cambridge University Press, Cambridge, UK, 2018.

[66] M. Xu, D. Zhang, and W. B. Wu. Pearson's chi-squared statistics: approximation theory and beyond. *Biometrika*, 106(3):716–723, 04 2019.

[67] L. Yu, K. Balasubramanian, S. Volgushev, and M. A. Erdogdu. An analysis of constant step size sgd in the non-convex regime: Asymptotic normality and bias, 2020.